

On quantitative uniqueness for parabolic equations

Igor Kukavica and Quinn Le

ABSTRACT. We consider the quantitative uniqueness properties for a parabolic type equation $u_t - \Delta u = w(x, t)\nabla u + v(x, t)u$, when $v \in L_t^{p_2} L_x^{p_1}$ and $w \in L_t^{q_2} L_x^{q_1}$, with a suitable range for exponents p_1, p_2, q_1 , and q_2 . We prove a strong unique continuation property and provide a pointwise in time observability estimate. November 17, 2021

CONTENTS

1. Introduction	1
2. The main result	2
3. Proof on the statement for quantitative uniqueness	4
4. Doubling property	14
5. The case \mathbb{R}^n	16
6. $L_t^q L_x^p$ case	17
Acknowledgments	18
References	18

1. Introduction

We address quantitative uniqueness for solutions of the parabolic equation

$$u_t - \Delta u = w(x, t)\nabla u + v(x, t)u, \quad (x, t) \in \Omega \times [T_0, T_0 + T],$$

where v and w belong to the mixed Lebesgue spaces $v \in L_t^{p_2} L_x^{p_1}$ and $w \in L_t^{q_2} L_x^{q_1}$, respectively. In particular, we obtain bounds on the order of vanishing which are algebraic in the corresponding norms of v and w . Additionally, we obtain pointwise in time doubling estimates, i.e., inequalities of the form

$$\|u(\cdot, t)\|_{L^2(\Omega)} \leq M_\delta \|u(\cdot, t)\|_{L^2(B_\delta(x_0))}$$

for an arbitrary $\delta > 0$ under the Lebesgue conditions on v and w .

The unique continuation for PDEs has a rich history (cf. the review papers by Kenig [K1, K2] and Vessella [V]), so we only mention several results pertaining to this paper. In [JK], Jerison and Kenig proved that the second order elliptic equation has the strong unique continuation property (i.e., is identically zero if it vanishes to an infinite order at a point) if $w = 0$ and $v \in L^{n/2}$, with a sufficiently small $L^{n/2}$ norm, which is a sharp result. The parabolic counterpart was obtained by Escoriaza and Vega [EV] (see also [E] for a previous unique continuation result when $v \in L^\infty L^p$ with $p > n/2$). The difficult case when w is nonzero was addressed by Koch and Tataru for the elliptic case in [KT1] and the parabolic case in [KT2]. In particular, they obtained the strong unique continuation for $v \in L_t^1 L_x^\infty + L_t^\infty L_x^{n/2}$ with the norm sufficiently small and $w \in L_{x,t}^{n+2}$. All the mentioned works rely on suitable Carleman type estimates and lead to observability type estimates on the space-time rectangles. For other works on the frequency approach to the unique continuation, see [A, AN, Ch, Ku1, Ku2], for the works related to Dirichlet quotients, see [A, CFNT, FS] and for other related works, cf. [AE, AMRV, An, AV, BC, B, BK, CRV, DF1, DF2, DZ1, DZ2, EF, EFV, EVe, H1, H2, K3, KSW, L, M, Z].

In this paper (cf. Theorems (2.1)–(2.3) below), we obtain explicit algebraic observability estimates for fixed time (i.e., not only on space-time rectangles) under the assumptions on the coefficients $v \in L^\infty L^p$ and $w \in L^\infty L^q$, where

$p > 2n/3$ and $q > 2n$. More general conditions $v \in L_t^{p_2} L_x^{p_1}$ and $w \in L_t^{q_2} L_x^{q_1}$ under certain range are addressed in Theorems 2.1–2.3. While the results cited in the second paragraph use the Carleman estimates, we rely on the frequency function approach, developed in [AI, GL] for the elliptic and [Kur, P] for the parabolic equations. The main idea of this approach for parabolic equations is the logarithmic convexity of the frequency function

$$Q(t) = \frac{|t| \int_{\mathbb{R}^n} |\nabla u(x, t)|^2 G(x, t) dx}{\int_{\mathbb{R}^n} u(x, t)^2 G(x, t) dx} \quad (1.1)$$

for the heat equation. In (1.1), G is the $(4\pi)^{n/2}$ -multiple of the backward Gaussian kernel, i.e.,

$$G(x, t) = \frac{1}{|t|^{n/2}} e^{|x|^2/4t}.$$

Another reformulation of the idea is to use the similarity variables (cf. (3.8) and (3.9) below) and obtain a logarithmic convexity of the unweighted norm [C, Ku3]. In [CK], the approach was used to obtain an estimate for an order of vanishing $C(\|w\|_{L^\infty}^2 + \|v\|_{L^\infty}^{2/3})$, which is, at least for complex valued coefficients, sharp [CKW1, CKW2]. Poon and Kurata showed that the frequency approach leads to unique continuation for $p > n$ and $q = \infty$.

In this paper, we show that this may be improved to $p > 2n/3$ and $q > 2n$. The improved range is obtained by three main devices. The first is to find the point in space where the frequency function is the smallest and translate the equation so it starts at that point (cf. Lemma 3.1 below); this idea has been introduced in [Ku4, CK]. The second device is to use the embedding theorems with Laplacian and use (3.24)–(3.25) below to bound the parts containing v and w . The third device is to use the finiteness of the integral in (3.47) below, which then allows us to show the convergence of the quantity under the integral. Note that we obtain an explicit algebraic bound on the order of vanishing, which is a constant multiple of

$$\|v\|_{L_t^\infty L_x^p}^{2/(3-2n/p)} + \|w\|_{L_t^\infty L_x^q}^{2/(1-2n/q)}.$$

When setting $p = q = \infty$, the estimate reduces to the sharp bound from [CK]. We also provide a pointwise estimate in time for a better understanding of the behavior of solution u . In particular, for all $\delta_0 \in (0, 1]$, we have

$$\|u(\cdot, t)\|_{L^2(\mathbb{R}^n)} \lesssim e^P \|u(\cdot, t)\|_{L^2(B_{\delta_0}(0))} \quad (1.2)$$

for all $t \in (T_0, T_0 + T)$, where P is a polynomial depending only on $n, q, p, \delta_0, \|v\|_{L^p}$ and $\|w\|_{L^q}$. The explicit formula for P can be found in Lemma 4.2 below. Note that the estimates of the type (1.2) are an essential ingredient when considering qualitative properties of solutions of evolutionary parabolic PDE. For instance, they are needed when considering the size of the zero set of solution at time t or, more generally, complexity of a graph of a function at a time t ([Ku3, Ku4]). Note finally that compared to [CK], we reduce the necessary regularity for the solution u to a simple boundedness, rather than differentiability.

The paper is structured as follows. In Section 2, we provide the setup of the problem and state the two main theorems on the order of vanishing with periodic boundary conditions and the analog for \mathbb{R}^n . For simplicity, the theorems address periodic boundary conditions. The case of \mathbb{R}^n under a natural non-growth condition on the solution is considered in Theorem 2.3 and proven in Section 4. Theorems 2.1–2.3 are concerned with the case when the Lebesgue exponent of t is infinity, while the general range is addressed in Theorem 6.1 in Section 6.

2. The main result

We address the quantitative uniqueness of a nontrivial solution $u \in L^\infty(I, L^2(\mathbb{T}^n)) \cap L^2(I, H^1(\mathbb{T}^n))$ of the equation

$$\begin{aligned} u_t - \Delta u &= w(x, t) \nabla u + v(x, t) u \\ u(x, T_0) &= u_0(t) \end{aligned} \quad (2.1)$$

defined for $(x, t) \in \mathbb{R}^n \times I$ where $I = [T_0, T_0 + T]$ is a given time interval and $n \geq 3$. The theorems are also valid for $n = 1, 2$ with minor changes; cf. Remark 2.4 below. We assume that u, v , and w are Ω -periodic, where $\Omega = \mathbb{T}^n = [-1/2, 1/2]^n$ and that they satisfy

$$\|v(\cdot, t)\|_{L^p(\Omega)} \leq M_0 \quad (2.2)$$

and

$$\|w(\cdot, t)\|_{L^q(\Omega)} \leq M_1, \quad (2.3)$$

for all $t \in I$. For simplicity, and since we are interested in the regime of M_1 and M_2 large, we assume $M_1, M_2 \geq 1$. Let $O_{(x_0, t_0)}(u)$ be the vanishing order of u at (x_0, t_0) , which is defined as the largest integer d such that

$$\|u\|_{L^2(Q_r(x_0, t_0))} = O(r^{d+(n+2)/2}) \quad \text{as } r \rightarrow 0^+$$

where

$$Q_r(x_0, t_0) = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} : |x - x_0| < r, -r^2 < t - t_0 < 0\} \quad (2.4)$$

stands for the parabolic cylinder centered at (x_0, t_0) with the radius r . For $t \in I$, denote by

$$q_D(t) = \frac{\|\nabla u(\cdot, t)\|_{L^2(\Omega)}^2}{\|u(\cdot, t)\|_{L^2(\Omega)}^2} \quad (2.5)$$

the Dirichlet quotient of u at time t . We assume that $\|u(\cdot, t)\|_{L^2(\Omega)}$ is nonzero for all $t \in I$. We also assume

$$q_0 = \sup_{t \in I} q_D(t) < \infty. \quad (2.6)$$

The following is the main result of this paper; cf. also Theorem 6.1 for $v \in L_t^{p_2} L_x^{p_1}(\Omega \times I)$ and $w \in L_t^{q_2} L_x^{q_1}(\Omega \times I)$.

THEOREM 2.1. *Let $u \in L_{x,t}^\infty(\mathbb{T}^n \times I)$ be a solution of (2.1) with v and w satisfying (2.2) and (2.3) such that*

$$p > \frac{2n}{3}$$

and

$$q > 2n.$$

Then for all $(x_0, t_0) \in \mathbb{T}^n \times I$, the vanishing order of u at (x_0, t_0) satisfies

$$O_{(x_0, t_0)}(u) \lesssim M_0^a + M_1^b, \quad (2.7)$$

where $a = 2/(3 - 2n/p)$ and $b = 2/(1 - 2n/q)$, with the implicit constant in (2.7) depending on q_0 and I .

We emphasize that the dimension is considered fixed, so we allow all constants to depend on n without mention. In the next statement, we provide a pointwise in time doubling property of u under the conditions of Theorem 2.1.

THEOREM 2.2. *Under the conditions of Theorem 2.1, we have*

$$\|u(\cdot, t)\|_{L^2(\Omega)} \lesssim e^{P(n, \delta_0, M_0, M_1, p, q)} \|u(\cdot, t)\|_{L^2(B(0, \delta_0))},$$

for all $t \in (T_0, T_0 + T)$, where P is a polynomial.

Now, consider u to be given on \mathbb{R}^n , instead of \mathbb{T}^n . Suppose that it satisfies the condition

$$\int_{\mathbb{R}^n} u(x, t)^2 dx \leq K \int_{B(0, 1)} u(x, t)^2 dx, \quad t \in [T_0, T_0 + T], \quad (2.8)$$

for some constant K . In this case, we obtain the following analogue of Theorems 2.1 and 2.2.

THEOREM 2.3. *Let $u \in L_t^\infty L_x^\infty$ be a solution of (2.1) such that the condition (2.8) holds with v and w satisfying (2.2) and (2.3). Assume additionally that*

$$q > 2n$$

and

$$p > \frac{2n}{3}.$$

Then for all $(x_0, t_0) \in \mathbb{R}^n \times I$, the vanishing order of u at (x_0, t_0) satisfies

$$O_{(x_0, t_0)}(u) \lesssim M_0^a + M_1^b, \quad (2.9)$$

where $a = 2/(3 - 2n/p)$ and $b = 2/(1 - 2n/q)$, with the implicit constant in (2.9) depending on q_0 , K , and I . Moreover, for $\delta_0 \in (0, 1]$, we have

$$\|u(\cdot, t)\|_{L^2(\Omega)} \lesssim e^{P(n, \delta_0, M_0, M_1, q, p)} \|u(\cdot, t)\|_{L^2(B(0, \delta_0))}, \quad (2.10)$$

for all $t \in (T_0, T_0 + T)$, where P is a polynomial.

The theorem is proven in Section 5 below.

REMARK 2.4. In the theorems above, we assumed $n \geq 3$. In this remark, we address the changes when $n = 1$ or $n = 2$. If $n = 2$, the results are exactly the same with small modifications in the proof; the modifications are pointed out in the proof of Theorem 2.1. For the case $n = 1$, we need to assume additionally that $p \geq 2$ and $q \geq 4$.

3. Proof on the statement for quantitative uniqueness

This section is devoted to the proof of Theorem 2.1. We first start with the case where v , w , and u_0 are smooth and then use approximation argument to prove the theorem under the boundedness assumption. Thus assume for now that v , w , and u_0 are smooth. By translation and rescaling, we may assume that $I = [-1, 0]$ and $(x_0, t_0) = (0, 0)$. Suppose additionally that u is periodic in x with respect to $\Omega = [-1/2, 1/2]^n$. For any starting time $-\epsilon$, we claim that we can choose a starting point x_ϵ that is controlled by q_0 .

LEMMA 3.1. *Let u be a solution of (2.1) that is periodic in x with respect to $\Omega = [-1/2, 1/2]^n$ where v and w satisfying (2.2) and (2.3). For any $\epsilon \in (0, T)$, there exist $x_\epsilon \in \Omega$*

$$\frac{\epsilon \int_{\mathbb{R}^n} |\Delta u(x_\epsilon + y, -\epsilon)|^2 G(y, -\epsilon) dy}{\int_{\mathbb{R}^n} u(x_\epsilon + y, -\epsilon)^2 G(y, -\epsilon) dy} \leq 2\epsilon q_D(-\epsilon),$$

where

$$G(x, t) = \frac{1}{|t|^{n/2}} e^{|x|^2/4t}, \quad x \in \mathbb{R}^n, \quad t < 0. \quad (3.1)$$

The lemma is proven in [CK, Ku4]; we provide a short argument for the sake of completeness. In the sequel, we reserve ϵ for the time provided in Lemma 3.1. We always tacitly assume that $\epsilon \in (0, T)$.

PROOF OF LEMMA 3.1. Assume that we have

$$2q_D(-\epsilon) \int_{\mathbb{R}^n} u(x + y, -\epsilon)^2 G(y, -\epsilon) dy \leq \int_{\mathbb{R}^n} |\nabla u(x + y, -\epsilon)|^2 G(y, -\epsilon) dy, \quad (3.2)$$

for all $x \in \Omega$. With a simple change of variables, (3.2) becomes

$$\int_{\mathbb{R}^n} u(y, -\epsilon) G(y - x, -\epsilon) dy \leq \frac{1}{2q_D(-\epsilon)} \int_{\mathbb{R}^n} |\nabla u(y, -\epsilon)|^2 G(y - x, -\epsilon) dy, \quad x \in \Omega. \quad (3.3)$$

Integrating (3.3) over Ω , we obtain

$$\begin{aligned} \int_{\Omega} \int_{\mathbb{R}^n} u(y, -\epsilon)^2 G(y, -\epsilon) dy dx &= \int_{\Omega} \sum_{j \in \mathbb{Z}^n} \int_{j+\Omega} u(y, -\epsilon)^2 G(y - x, -\epsilon) dy dx \\ &= \int_{\Omega} \sum_{j \in \mathbb{Z}^n} \int_{\Omega} u(y', -\epsilon)^2 G(y' + j - x, -\epsilon) dy' dx \\ &= \int_{\Omega} u(y', -\epsilon)^2 \left(\sum_{j \in \mathbb{Z}^n} \int_{\Omega} G(y' + j - x, -\epsilon) dx \right) dy' \\ &= \int_{\Omega} u(y', -\epsilon)^2 \int_{\mathbb{R}^n} G(y' - x, -\epsilon) dx dy' = (4\pi)^{n/2} \int_{\Omega} u(y', -\epsilon)^2 dy', \end{aligned} \quad (3.4)$$

where we used $\int_{\mathbb{R}^n} G(y - x, -\epsilon) dx = (4\pi)^{n/2}$ in the last equality. Similarly, we have

$$\int_{\Omega} \int_{\mathbb{R}^n} |\nabla u(y, -\epsilon)|^2 G(y - x, -\epsilon) dy dx = (4\pi)^{n/2} \int_{\Omega} |\nabla u(y', -\epsilon)|^2 dy'. \quad (3.5)$$

By (3.3), (3.4), and (3.5), we obtain

$$2q_D(-\epsilon) \int_{\Omega} u(y, -\epsilon)^2 dy \leq \int_{\Omega} |\nabla u(y, -\epsilon)|^2 dy = q_D(-\epsilon) \int_{\Omega} u(y, -\epsilon)^2 dy.$$

We have a contradiction since $\int_{\Omega} u(y, -\epsilon)^2 dy > 0$. Therefore, (3.2) cannot hold for all $x \in \Omega$. Thus Lemma 3.1 follows. \square

We proceed with a change of variables so that at the time $-\epsilon$, given by Lemma 3.1, the solution u starts at x_ϵ with x_ϵ also as in Lemma 3.1. In particular, we let

$$u(x, t) = \bar{u} \left(x + \frac{x_\epsilon}{\epsilon} t, t \right).$$

It is not difficult to check that \bar{u} solves the equation

$$\partial_t \bar{u} - \Delta \bar{u} = -\frac{1}{\epsilon} x_\epsilon \nabla \bar{u} + w \nabla \bar{u} + v \bar{u}. \quad (3.6)$$

Denoting $a = -x_\epsilon/\epsilon$ and writing u instead of \bar{u} , the equation (3.6) becomes

$$\partial_t u - \Delta u = a_j \partial_j u + w_j \partial_j u + vu. \quad (3.7)$$

We now proceed with a change of variable

$$u(x, t) = e^{|x|^2/8(-t)} U \left(\frac{x}{\sqrt{-t}}, -\log(-t) \right), \quad (3.8)$$

that is,

$$U(y, \tau) = e^{-|y|^2/8} u(ye^{-\tau/2}, -e^{-\tau}) \quad (3.9)$$

with $\tau = -\log(-t)$. Also, let

$$V(y, \tau) = v(ye^{-\tau/2}, -e^{-\tau}), \quad (y, \tau) \in \mathbb{R}^n \times [\tau_0, \infty)$$

and

$$W_j(y, \tau) = w_j(ye^{-\tau/2}, -e^{-\tau}), \quad (y, \tau) \in \mathbb{R}^n \times [\tau_0, \infty),$$

for $j = 1, \dots, n$, where

$$\tau_0 = \log(1/\epsilon). \quad (3.10)$$

Then

$$\partial_\tau U + HU = e^{-\tau/2} (a_j y_j U + a_j \partial_j U) + e^{-\tau/2} (y_j W_j(y, \tau) U + W_j(y, \tau) \partial_j U) + e^{-\tau} V(y, \tau) U, \quad (3.11)$$

where

$$HU = -\Delta U + \left(\frac{|y|^2}{16} - \frac{n}{4} \right) U$$

with the initial data

$$U(y, \tau_0) = U \left(y, \log \frac{1}{\epsilon} \right) = e^{-|y|^2/8} u \left(\frac{y}{\sqrt{\epsilon}}, -\epsilon \right).$$

Now, define the frequency function

$$Q(\tau) = \frac{(HU, U)_{L^2(\mathbb{R}^n)}}{\|U\|^2},$$

where we denote

$$\|\cdot\| = \|\cdot\|_{L^2(\mathbb{R}^n)} \quad (3.12)$$

throughout. Letting

$$A(\tau)U = HU - e^{-\tau/2} a_j y_j U \quad (3.13)$$

and

$$\bar{Q}(\tau) = \frac{(A(\tau)U, U)_{L^2(\mathbb{R}^n)}}{\|U\|^2} = Q(\tau) - \frac{e^{-\tau/2} a_j}{\|U\|^2} \int_{\mathbb{R}^n} y_j U^2 dy, \quad (3.14)$$

we may rewrite (3.11) as

$$\partial_\tau U + (A(\tau) - \bar{Q}(\tau)I)U + \bar{Q}(\tau)U = e^{-\tau/2} a_j \partial_j U + e^{-\tau/2} (y_j W_j U + W_j \partial_j U) + e^{-\tau} VU. \quad (3.15)$$

For simplicity of notation, denote

$$\tilde{U} = \frac{U}{\|U\|},$$

so that $\|\tilde{U}\| = 1$. We now show that the modified frequency function \bar{Q} is bounded.

LEMMA 3.2. *Under the assumptions of Theorem 2.1, for all $\tau \geq \tau_0$, the modified frequency function satisfies*

$$\bar{Q}(\tau) \lesssim M_0^a + M_1^b, \quad (3.16)$$

where $a = 2/(3 - 2n/p)$ and $b = 2/(1 - 2n/q)$.

PROOF OF LEMMA 3.2. Without loss of generality, assume that $M_0, M_1 \geq 1$. Dividing (3.15) by $\|U\|$ and then taking the inner product with $(A(\tau) - \bar{Q}(\tau)I)\tilde{U}$, we obtain, using $((A(\tau) - \bar{Q}(\tau)I)\tilde{U}, \tilde{U}) = 0$,

$$\frac{1}{\|\tilde{U}\|} (\partial_\tau \tilde{U}, (A(\tau) - \bar{Q}(\tau)I)\tilde{U}) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 = (F(\tilde{U}), (A(\tau) - \bar{Q}(\tau)I)\tilde{U}), \quad (3.17)$$

where $F(\tilde{U})$ denotes the right-hand side of (3.15) divided by $\|U\|$, i.e.,

$$F(\tilde{U}) = e^{-\tau/2} a_j \partial_j \tilde{U} + e^{-\tau/2} (y_j W_j \tilde{U} + W_j \partial_j \tilde{U}) + e^{-\tau} V \tilde{U}.$$

From (3.17), we get

$$\frac{1}{\|U\|} (\partial_\tau U, (A(\tau) - \bar{Q}(\tau)I)\tilde{U}) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 = (F(\tilde{U}), (A(\tau) - \bar{Q}(\tau)I)\tilde{U}). \quad (3.18)$$

By differentiating (3.14), we have

$$\bar{Q}'(\tau) = 2(\partial_\tau \tilde{U}, A(\tau)\tilde{U}) + (\tilde{U}, A'(\tau)\tilde{U}) - 2(\partial_\tau \tilde{U}, \tilde{U})(A(\tau)\tilde{U}, \tilde{U})$$

from where we use that $A(\tau)$ is a symmetric operator with the derivative $A'(\tau)U = \frac{1}{2}e^{-\tau/2}a_j y_j U$ to obtain

$$\begin{aligned} \frac{1}{2}\bar{Q}'(\tau) &= (\partial_\tau \tilde{U}, A(\tau)\tilde{U}) + \frac{1}{4}e^{-\tau/2}(\tilde{U}, a_j y_j \tilde{U}) - (\partial_\tau \tilde{U}, \tilde{U})(A(\tau)\tilde{U}, \tilde{U}) \\ &= (\partial_\tau \tilde{U}, (A(\tau) - \bar{Q}(\tau)I)\tilde{U}) + \frac{1}{4}e^{-\tau/2}(\tilde{U}, a_j y_j \tilde{U}). \end{aligned} \quad (3.19)$$

Substituting (3.18) into (3.19), we get

$$\begin{aligned} \frac{1}{2}\bar{Q}'(\tau) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 &= \frac{1}{4}e^{-\tau/2}(a_j y_j \tilde{U}, \tilde{U}) + e^{-\tau/2} a_j (\partial_j \tilde{U}, (A(\tau) - \bar{Q}(\tau)I)\tilde{U}) \\ &\quad + (e^{-\tau/2}(y_j W_j \tilde{U} + W_j \partial_j \tilde{U}) + e^{-\tau} V \tilde{U}, (A(\tau) - \bar{Q}(\tau)I)\tilde{U}). \end{aligned} \quad (3.20)$$

Note that $\int \tilde{U} \partial_j U = 0$ and $\int \Delta U \partial_j U = 0$ since v, w , and u_0 are assumed smooth. Integrating by parts the second term on the right-hand side of (3.20), we obtain

$$\begin{aligned} \frac{1}{2}\bar{Q}'(\tau) &= \frac{1}{4}e^{-\tau/2} a_j \int y_j \tilde{U}^2 dy - \frac{1}{16}e^{-\tau/2} a_j \int y_j \tilde{U}^2 dy + \frac{1}{2}e^{-\tau} |a|^2 \\ &\quad + e^{-\tau/2} \int (y_j W_j \tilde{U} + W_j \partial_j \tilde{U} + e^{-\tau/2} V \tilde{U})(A(\tau) - \bar{Q}(\tau)I)\tilde{U} dy \\ &= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Using the Cauchy-Schwarz inequality and $\|\tilde{U}\| = 1$, we have

$$I_1 \leq e^{-\tau/2} |a| \int |y| \tilde{U}^2 dy \leq e^{-\tau/2} |a| \left(\int |y|^2 \tilde{U}^2 dy \right)^{1/2},$$

and similarly

$$I_2 + I_3 \leq e^{-\tau/2} |a| \left(\int |y|^2 \tilde{U}^2 dy \right)^{1/2} + e^{-\tau} |a|^2.$$

In order to estimate I_4 , we first claim

$$\|D^2 U\|^2 \lesssim \|HU\|^2 + \|U\|^2. \quad (3.21)$$

To prove (3.21), we expand $\|HU\|^2$ as

$$\begin{aligned} \|HU\|^2 &= \int_{\mathbb{R}^n} \left(-\Delta U + \left(\frac{|y|^2}{16} - \frac{n}{4} \right) U \right)^2 dy \\ &= \int_{\mathbb{R}^n} (\Delta U)^2 + \left(\frac{|y|^2}{16} - \frac{n}{4} \right)^2 U^2 dy - 2 \int_{\mathbb{R}^n} \left(\frac{|y|^2}{16} - \frac{n}{4} \right) U \Delta U dy \\ &= \int_{\mathbb{R}^n} \left((\Delta U)^2 + \left(\frac{|y|^2}{16} - \frac{n}{4} \right)^2 U^2 \right) dy + 2 \int_{\mathbb{R}^n} \partial_j U \partial_j U \left(\frac{|y|^2}{16} - \frac{n}{4} \right) dy + \frac{1}{4} \int_{\mathbb{R}^n} y_j U \partial_j U dy, \end{aligned}$$

from where

$$\begin{aligned} \|HU\|^2 &= \int_{\mathbb{R}^n} \left((\Delta U)^2 + \left(\frac{|y|^2}{16} - \frac{n}{4} \right)^2 U^2 \right) dy + 2 \int_{\mathbb{R}^n} |\nabla U|^2 \left(\frac{|y|^2}{16} - \frac{n}{4} \right) dy - \frac{n}{8} \int_{\mathbb{R}^n} U^2 dy \\ &\geq \int_{\mathbb{R}^n} \left((\Delta U)^2 + \left(\frac{|y|^2}{16} - \frac{n}{4} \right)^2 U^2 \right) dy - \frac{n}{2} \int_{\mathbb{R}^n} |\nabla U|^2 dy - \frac{n}{8} \int_{\mathbb{R}^n} U^2 dy. \end{aligned}$$

Using $n/4 - |y|^2/16 \leq n/4$ in the third integral, we get

$$\|\Delta U\|^2 \lesssim \|HU\|^2 + \|U\|^2 + \|\nabla U\|^2.$$

By Sobolev's and the Cauchy-Schwarz inequalities, we get

$$\|\nabla U\|^2 \leq \|U\| \|\Delta U\| \leq \frac{\|U\|^2 + \|\Delta U\|^2}{2},$$

and then using $\|D^2 U\| \lesssim \|\Delta U\|$, we obtain (3.21).

We now estimate $\|V\tilde{U}\|$. By Hölder's and Sobolev's inequalities, we have

$$\|V\tilde{U}\| \lesssim \|V\|_{L^p(\mathbb{R}^n)} \|\tilde{U}\|_{L^{2p/(p-2)}(\mathbb{R}^n)} \lesssim \|V\|_{L^p(\mathbb{R}^n)} \|\tilde{U}\|^{1-\alpha} \|D^2 \tilde{U}\|^\alpha \quad (3.22)$$

with

$$\alpha = \frac{n}{2p}. \quad (3.23)$$

Note that we need $p > 2$ in (3.22). This automatically holds if $n \geq 3$ and $p > 2n/3$. If $n = 2$, p can be 2 since we only require $p > 2n/3 = 4/3$. In this case, we can simply replace $2p/(p-2)$ by ∞ and use Sobolev's inequalities for L^∞ . This estimate also works for the case $n = 1$ since we have assumed additionally that $p \geq 2$ if $n = 1$. By (3.21), we have

$$\|V\tilde{U}\| \lesssim \|V\|_{L^p(\mathbb{R}^n)} \|\tilde{U}\|^{1-\alpha} (\|H\tilde{U}\|^2 + \|\tilde{U}\|^2)^{\alpha/2} \lesssim \|V\|_{L^p(\mathbb{R}^n)} (\|H\tilde{U}\|^\alpha + 1). \quad (3.24)$$

In order to estimate the last expression, we write

$$\begin{aligned} \|H\tilde{U}\| &\leq \|H\tilde{U} - A(\tau)\tilde{U}\| + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| + \|\bar{Q}(\tau)\tilde{U}\| \\ &\lesssim e^{-\tau/2}|a|\|y\tilde{U}\| + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| + \bar{Q}(\tau), \end{aligned} \quad (3.25)$$

where we used (3.13) in the last step. Since

$$\|\nabla \tilde{U}\| + \|y\tilde{U}\| \lesssim \bar{Q}(\tau)_+^{1/2} + e^{-\tau/2}|a| + 1, \quad (3.26)$$

([Ku4, p.780]), as one may readily check, we get

$$\|H\tilde{U}\| \lesssim e^{-\tau/2}|a|(\bar{Q}(\tau)_+^{1/2} + e^{-\tau/2}|a| + 1) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| + |\bar{Q}(\tau)|. \quad (3.27)$$

Since $\|V\|_{L^p(\mathbb{R}^n)} \leq M_0 e^{\alpha\tau}$ by a simple change of variables, we obtain from (3.24) and (3.27) the inequality

$$\|V\tilde{U}\| \lesssim M_0 e^{\alpha\tau} \left(e^{-\alpha\tau/2}|a|^\alpha \left(\bar{Q}(\tau)_+^{\alpha/2} + e^{-\alpha\tau/2}|a|^\alpha + 1 \right) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\alpha + |\bar{Q}(\tau)|^\alpha + 1 \right). \quad (3.28)$$

Next, to estimate $\|y_j W_j \tilde{U} + W_j \partial_j \tilde{U}\|$, we first bound $\|W_j \partial_j \tilde{U}\|$ and then $\|y_j W_j \tilde{U}\|$. By Hölder's and Sobolev's inequalities again, we have, also using $\|W\|_{L^p(\mathbb{R}^n)} \lesssim M_1 e^{n\tau/2q} = M_1 e^{(\beta-1/2)\tau}$,

$$\|W_j \partial_j \tilde{U}\| \lesssim \|W\|_{L^q(\mathbb{R}^n)} \|\nabla \tilde{U}\|_{L^{2q/(q-2)}(\mathbb{R}^n)} \lesssim M_1 e^{(\beta-1/2)\tau} \|\tilde{U}\|^{1-\beta} \|D^2 \tilde{U}\|^\beta$$

with

$$\beta = \frac{n}{2q} + \frac{1}{2}. \quad (3.29)$$

Similarly as in (3.25), we have

$$\begin{aligned} \|W_j \partial_j \tilde{U}\| &\lesssim M_1 e^{(\beta-1/2)\tau} \left(\|H\tilde{U}\|^\beta + 1 \right) \\ &\lesssim M_1 e^{(\beta-1/2)\tau} \left(e^{-\beta\tau/2}|a|^\beta (\bar{Q}(\tau)_+^{\beta/2} + e^{-\beta\tau/2}|a|^\beta + 1) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\beta + \bar{Q}(\tau)^\beta + 1 \right). \end{aligned} \quad (3.30)$$

Next,

$$\begin{aligned}
\|y_j W_j \tilde{U}\| &\lesssim \|W\|_{L^q(\mathbb{R}^n)} \|y \tilde{U}\|_{L^{2q/(q-2)}(\mathbb{R}^n)} \lesssim M_1 e^{(\beta-1/2)\tau} \|y \tilde{U}^{1/2}\|_{L^4(\mathbb{R}^n)} \|\tilde{U}^{1/2}\|_{L^{4q/(q-4)}(\mathbb{R}^n)} \\
&= M_1 e^{(\beta-1/2)\tau} \|y^2 \tilde{U}\|^{1/2} \|\tilde{U}\|_{L^{2q/(q-4)}(\mathbb{R}^n)}^{1/2} \lesssim M_1 e^{(\beta-1/2)\tau} \|y^2 \tilde{U}\|^{1/2} \|\Delta \tilde{U}\|^{n/2q} \|\tilde{U}\|^{1/2-n/2q} \\
&\lesssim M_1 e^{(\beta-1/2)\tau} \|y^2 \tilde{U}\|^{1/2} \|\Delta \tilde{U}\|^{n/2q},
\end{aligned} \tag{3.31}$$

where we used (3.29) in the last inequality. Similarly to estimate $\|V \tilde{U}\|$, we need $q \geq 4$ in (3.31). This is obviously true if $n \geq 2$ and $q > 2n$. If $n = 1$, (3.31) still holds since we assumed $q \geq 4$. Recall that $HU = -\Delta U + (|y|^2/16 - n/4)U$, from where $\|y^2 U\| \lesssim \|HU\| + \|\Delta U\| + \|U\|$. Therefore,

$$\begin{aligned}
\|y_j W_j \tilde{U}\| &\lesssim M_1 e^{(\beta-1/2)\tau} \|\Delta \tilde{U}\|^{n/2q} (\|H \tilde{U}\|^{1/2} + \|\Delta \tilde{U}\|^{1/2} + 1) \\
&\lesssim M_1 e^{(\beta-1/2)\tau} (\|H \tilde{U}\|^{n/2q} + 1) (\|H \tilde{U}\|^{1/2} + 1) \\
&\lesssim M_1 e^{(\beta-1/2)\tau} (\|H \tilde{U}\|^\beta + 1),
\end{aligned}$$

where we used $\beta = n/2q + 1/2$. By (3.27) we then obtain

$$\|y_j W_j \tilde{U}\| \lesssim M_1 e^{(\beta-1/2)\tau} \left(e^{-\beta\tau/2} |a|^\beta (\bar{Q}(\tau)_+^{\beta/2} + e^{-\beta\tau/2} |a|^\beta + 1) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\beta + \bar{Q}(\tau)^\beta + 1 \right). \tag{3.32}$$

By (3.28), (3.30), and (3.32), we get an estimate for I_4 which reads

$$\begin{aligned}
I_4 &\lesssim \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| \left(e^{(\beta-1)\tau} M_1 (e^{-\beta\tau/2} |a|^\beta (\bar{Q}(\tau)_+^{\beta/2} + e^{-\beta\tau/2} |a|^\beta + 1) \right. \\
&\quad \left. + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\beta + \bar{Q}(\tau)^\beta + 1 \right) \\
&\quad + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| \left(e^{(\alpha-1)\tau} M_0 (e^{-\alpha\tau/2} |a|^\alpha (\bar{Q}(\tau)_+^{\alpha/2} + e^{-\alpha\tau/2} |a|^\alpha + 1) \right. \\
&\quad \left. + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\alpha + \bar{Q}(\tau)^\alpha + 1 \right).
\end{aligned}$$

Combining the bounds for I_1, I_2, I_3 , and I_4 , we get

$$\begin{aligned}
&\frac{1}{2} \bar{Q}'(\tau) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 \\
&\lesssim e^{-\tau/2} |a| (\bar{Q}(\tau)_+^{1/2} + e^{-\tau/2} |a| + 1) \\
&\quad + e^{(\beta-1)\tau} M_1 \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| \left(e^{-\beta\tau/2} |a|^\beta \bar{Q}(\tau)_+^{\beta/2} + e^{-\beta\tau} |a|^{2\beta} + e^{-\beta\tau/2} |a|^\beta \right. \\
&\quad \left. + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\beta + \bar{Q}(\tau)^\beta + 1 \right) \\
&\quad + e^{(\alpha-1)\tau} M_0 \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\| \left(e^{-\alpha\tau/2} |a|^\alpha \bar{Q}(\tau)_+^{\alpha/2} + e^{-\alpha\tau} |a|^{2\alpha} + e^{-\alpha\tau/2} |a|^\alpha \right. \\
&\quad \left. + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\alpha + \bar{Q}(\tau)^\alpha + 1 \right).
\end{aligned} \tag{3.33}$$

We now apply Young's inequality to terms involving $\|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|$ on the right-hand side so that we can absorb them into the second term in the left hand side of (3.33). Namely, we use

$$N \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^\gamma \leq \epsilon_0 \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 + C_{\epsilon_0} N^{2/(2-\gamma)},$$

where $\epsilon_0 \in (0, 1]$ is arbitrarily small, with $\gamma = 1, \alpha + 1, \beta + 1$ and corresponding expressions for N . Thus, we obtain

$$\begin{aligned}
&\bar{Q}'(\tau) + \|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 \\
&\lesssim e^{-\tau/2} |a| \bar{Q}(\tau)_+^{1/2} + e^{(\beta-2)\tau} M_1^2 |a|^{2\beta} \bar{Q}(\tau)_+^\beta + e^{(2\beta-2)\tau} M_1^2 \bar{Q}(\tau)_+^{2\beta} \\
&\quad + M_0^2 e^{(\alpha-2)\tau} |a|^{2\alpha} \bar{Q}(\tau)_+^\alpha + M_0^2 e^{(2\alpha-2)\tau} \bar{Q}(\tau)_+^{2\alpha} + e^{-\tau} |a|^2 + e^{-\tau/2} |a| \\
&\quad + M_1^2 e^{-2\tau} |a|^{4\beta} + M_1^2 e^{(\beta-2)\tau} |a|^{2\beta} + M_1^{2/(1-\beta)} e^{-2\tau} + M_1^2 e^{(2\beta-2)\tau} \\
&\quad + M_0^2 e^{-2\tau} |a|^{4\alpha} + M_0^2 e^{(\alpha-2)\tau} |a|^{2\alpha} + M_0^{2/(1-\alpha)} e^{-2\tau} + M_0^2 e^{(2\alpha-2)\tau}.
\end{aligned} \tag{3.34}$$

Since $|a| \lesssim 1/\epsilon$, the last inequality implies

$$\begin{aligned} \bar{Q}'(\tau) &\lesssim e^{-\tau/2} \epsilon^{-1} \bar{Q}(\tau)_+^{1/2} + e^{(\beta-2)\tau} M_1^2 \epsilon^{-2\beta} \bar{Q}(\tau)_+^\beta + e^{(2\beta-2)\tau} M_1^2 \bar{Q}(\tau)_+^{2\beta} \\ &\quad + M_0^2 e^{(\alpha-2)\tau} \epsilon^{-2\alpha} \bar{Q}(\tau)_+^\alpha + M_0^2 e^{(2\alpha-2)\tau} \bar{Q}(\tau)_+^{2\alpha} + e^{-\tau} \epsilon^{-2} + e^{-\tau/2} \epsilon^{-1} \\ &\quad + M_1^2 e^{-2\tau} \epsilon^{-4\beta} + M_1^2 e^{(\beta-2)\tau} \epsilon^{-2\beta} + M_1^{2/(1-\beta)} e^{-2\tau} + M_1^2 e^{(2\beta-2)\tau} \\ &\quad + M_0^2 e^{-2\tau} \epsilon^{-4\alpha} + M_0^2 e^{(\alpha-2)\tau} \epsilon^{-2\alpha} + M_0^{2/(1-\alpha)} e^{-2\tau} + M_0^2 e^{(2\alpha-2)\tau}. \end{aligned} \quad (3.35)$$

For the initial value, we have

$$\begin{aligned} \bar{Q}(\tau_0) &\lesssim Q(\tau_0) + \left| e^{-\tau_0/2} a_j \int y_j \tilde{U}^2 dy \right| \lesssim 2\epsilon q_D(-\epsilon) + \epsilon^{-1/2} \frac{(HU(\tau_0), U(\tau_0))^{1/2}}{(U(\tau_0), U(\tau_0))^{1/2}} \\ &\lesssim 2\epsilon q_D(-\epsilon) + \epsilon^{-1/2} (\epsilon q_D(-\epsilon))^{1/2} \lesssim \epsilon q_0 + q_0^{1/2}, \end{aligned} \quad (3.36)$$

which implies

$$\bar{Q}(\tau_0) \leq C_1,$$

where $C_1 = C_0(q_0 + q_0^{1/2})$ and $C_0 \geq 1$ is a constant from (3.36), which is further increased below. Denote by $C_2 \geq 1$ the implicit constant in the inequality (3.35) and choose

$$\epsilon = \frac{1}{2(M_0^a + M_1^b)}, \quad (3.37)$$

where

$$a = \frac{2}{3-4\alpha} \quad (3.38)$$

and

$$b = \frac{2}{3-4\beta}. \quad (3.39)$$

Note that $\epsilon \in (0, 1/2]$ since $M_0 \geq 1$. We claim that

$$\bar{Q}(\tau) < \frac{C_1 + C_2}{\epsilon}, \quad \tau \geq \tau_0 = -\log \epsilon \quad (3.40)$$

if we choose $C_0 \geq 1$ sufficiently large. Assume, contrary to the assertion, that there exists τ_1 such that $\bar{Q}(\tau_1) = (C_1 + C_2)/\epsilon$, and assume that τ_1 is the first such time. Integrating (3.35) between τ_0 and τ_1 , we arrive at

$$\begin{aligned} \bar{Q}(\tau_1) &\leq \bar{Q}(\tau_0) + 2C_2(C_1 + C_2)^{1/2} \epsilon^{-1} + M_1^2 C_2 (C_1 + C_2)^\beta \epsilon^{2-4\beta} + 2M_1^2 C_2 (C_1 + C_2)^{2\beta} \epsilon^{2-4\beta} \\ &\quad + M_0^2 C_2 (C_1 + C_2)^\alpha \epsilon^{2-4\alpha} + 2M_0^2 C_2 (C_1 + C_2)^{2\alpha} \epsilon^{2-4\alpha} + C_2 \epsilon^{-1} + 2C_2 \epsilon^{-1/2} \\ &\quad + M_1^2 C_2 \epsilon^{2-4\beta} + M_1^2 C_2 \epsilon^{2-3\beta} + M_1^{2/(1-\beta)} C_2 \epsilon^2 + 2M_1^2 C_2 \epsilon^{2-2\beta} + M_0^2 C_2 \epsilon^{2-4\alpha} \\ &\quad + M_0^2 C_2 \epsilon^{2-3\alpha} + M_0^{2/(1-\alpha)} C_2 \epsilon^2 + 2M_0^2 C_2 \epsilon^{2-2\alpha}, \end{aligned} \quad (3.41)$$

where we also used $e^{-\tau_0} = \epsilon$ and $0 \leq \alpha, \beta \leq 3/4$. The second term on the right-hand side satisfies

$$2C_2(C_1 + C_2)^{1/2} \epsilon^{-1} \leq \frac{C_1 + C_2}{100\epsilon},$$

given C_1 is sufficiently large compared to C_2 , which is achieved by increasing C_0 . For the third term, we have

$$M_1^2 C_2 (C_1 + C_2)^\beta \epsilon^{2-4\beta} = M_1^2 C_2 (C_1 + C_2)^\beta \epsilon^{3-4\beta} \epsilon^{-1} \leq C_2 (C_1 + C_2)^\beta \epsilon^{-1} \leq \frac{C_1 + C_2}{100\epsilon},$$

where in the second inequality we used $M_1^2 \epsilon^{3-4\beta} \leq 1$, and this results from (3.37). In the last step, we used that C_1 is sufficiently large compared to C_2 . We proceed similarly for the rest of the terms in (3.41) and obtain

$$\bar{Q}(\tau_1) \leq \bar{Q}(\tau_0) + \frac{C_1 + C_2}{4\epsilon} \leq C_1 + \frac{C_1 + C_2}{4\epsilon} < \frac{C_1 + C_2}{\epsilon},$$

where in the last step we used $\epsilon \leq 1/2$. This is a contradiction with a choice of τ_1 , and thus we conclude that (3.40) holds for all $\tau \geq \tau_0$. Finally, by (3.37) and (3.40), we get $\bar{Q}(\tau) \lesssim M_0^a + M_1^b$, as desired. \square

Now we show that the modified frequency function \bar{Q} controls the vanishing order of u . We first prove the following lemma, which shows the convergence of this function as $\tau \rightarrow \infty$ and the order of vanishing of $\int u^2(x, t) G(x, t) dx$, where G is defined in (3.1).

LEMMA 3.3. *Under the assumptions of Theorem 2.1, the modified frequency function satisfies $\bar{Q}(\tau) \rightarrow m/2$ as $\tau \rightarrow \infty$ for some $m \in \mathbb{N}$ such that $m \lesssim M_0^a + M_1^b$, where a and b are as in (3.38) and (3.39). Moreover, for all $\delta > 0$, there exist $\eta \in (\log(1/\epsilon), 0)$ and $A_1(\delta), A_2(\delta) > 0$ such that*

$$A_1(\delta)|t|^{m+\delta} \leq \int u^2(x, t)G(x, t) dx \leq A_2(\delta)|t|^{m-\delta}, \quad (3.42)$$

for all $t \in [-\eta, 0)$.

PROOF OF LEMMA 3.3. Taking the inner product of (3.15) with U , we obtain

$$\frac{1}{2} \frac{d}{d\tau} \|U\|^2 + (A(\tau)U, U) = e^{-\tau/2} (y_j W_j U, U) + e^{-\tau/2} (W_j \partial_j U, U) + e^{-\tau} (VU, U), \quad (3.43)$$

where we used $(e^{-\tau/2} a_j \partial_j U, U) = 0$. Denoting the right-hand side of (3.43) by $f(\tau)$, we obtain

$$\begin{aligned} f(\tau) &= e^{-\tau/2} \int y_j W_j U^2 dy + e^{-\tau/2} \int W_j U \partial_j U dy + e^{-\tau} \int VU^2 dy \\ &\lesssim e^{(\beta-1)\tau} M_1 \|yU^2\|_{L^{q/(q-1)}(\mathbb{R}^n)} + e^{(\beta-1)\tau} M_1 \|U \nabla U\|_{L^{q/(q-1)}(\mathbb{R}^n)} + e^{(\alpha-1)\tau} M_0 \|U^2\|_{L^{p/(p-1)}(\mathbb{R}^n)} \\ &\lesssim e^{(\beta-1)\tau} M_1 \|yU\| \|U\|_{L^{2q/(q-2)}(\mathbb{R}^n)} + e^{(\beta-1)\tau} M_1 \|U\|_{L^{2q/(q-2)}(\mathbb{R}^n)} \|\nabla U\| \\ &\quad + e^{(\alpha-1)\tau} M_0 \|U\|_{L^{2p/(p-1)}(\mathbb{R}^n)}^2. \end{aligned}$$

We now use the Gagliardo-Nirenberg inequality to get

$$\begin{aligned} f(\tau) &\lesssim e^{(\beta-1)\tau} M_1 \|yU\| \|U\|^{1-n/q} \|\nabla U\|^{n/q} + e^{(\beta-1)\tau} M_1 \|U\|^{1-n/q} \|\nabla U\|^{n/q+1} \\ &\quad + e^{(\alpha-1)\tau} M_0 \|U\|^{2-n/p} \|\nabla U\|^{n/p}. \end{aligned}$$

Therefore, by (3.26), we may estimate

$$\frac{f(\tau)}{\|U\|^2} \lesssim e^{(\beta-1)\tau} M_1 (\bar{Q}(\tau)_+ + e^{-\tau}|a|^2 + 1)^\beta + e^{(\alpha-1)\tau} M_0 (\bar{Q}(\tau)_+ + e^{-\tau}|a|^2 + 1)^\alpha, \quad (3.44)$$

where α and β are as in (3.23) and (3.29). For convenience, we denote

$$M = M_0^a + M_1^b. \quad (3.45)$$

Now, we integrate (3.44) from τ_0 to τ and obtain

$$\begin{aligned} \int_{\tau_0}^{\tau} \frac{f(s)}{\|U(\cdot, s)\|^2} ds &\lesssim M^\beta M_1 (e^{-(1-\beta)\tau_0} - e^{-(1-\beta)\tau}) + |a|^{2\beta} M_1 (e^{-\tau_0} - e^{-\tau}) + M_1 (e^{-(1-\beta)\tau_0} - e^{-(1-\beta)\tau}) \\ &\quad + M^\alpha M_0 (e^{-(1-\alpha)\tau_0} - e^{-(1-\alpha)\tau}) + |a|^{2\alpha} M_0 (e^{-\tau_0} - e^{-\tau}) + M_0 (e^{-(1-\alpha)\tau_0} - e^{-(1-\alpha)\tau}), \end{aligned}$$

where we used (3.16). On the other hand, if we integrate from τ_1 to τ the equation

$$\frac{1}{2} \frac{d}{d\tau} \|U\|^2 + \bar{Q}(\tau) = \frac{f(\tau)}{\|U\|^2},$$

we get

$$\begin{aligned} \frac{1}{2} \log \|U(\cdot, \tau)\|^2 - \frac{1}{2} \log \|U(\cdot, \tau_1)\|^2 &= - \int_{\tau_1}^{\tau} \bar{Q}(s) ds + \int_{\tau_1}^{\tau} \frac{f(s)}{\|U(\cdot, s)\|^2} ds \\ &\leq -\frac{m}{2}(\tau - \tau_1) - \int_{\tau_1}^{\tau} \left(\bar{Q}(s) - \frac{m}{2} \right) ds + C(e^{-\tau_1/4} + e^{-\tau/4}), \end{aligned} \quad (3.46)$$

where the constant depends on $|a|$, M (note that α and β are considered fixed). To show that $\bar{Q}(\tau) \rightarrow m/2$ as $\tau \rightarrow \infty$ for some $m \lesssim M$, we use $\bar{Q}(\tau) \lesssim M$ in (3.34) and integrate between τ_1 and τ , where $0 \leq \tau_1 \leq \tau$ obtaining

$$\bar{Q}(\tau) - \bar{Q}(\tau_1) + \int_{\tau_1}^{\tau} \|(A(s) - \bar{Q}(s)I)\tilde{U}\|^2 ds \lesssim e^{-\tau_1/4} + e^{-\tau/4}.$$

Therefore,

$$\int_{\tau_0}^{\tau} \|(A(s) - \bar{Q}(s)I)\tilde{U}\|^2 ds \lesssim M + e^{-\tau_1/4} + e^{-\tau/4}.$$

We also have

$$\|(A(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 \gtrsim \frac{1}{2} \|(H(\tau) - \bar{Q}(\tau)I)\tilde{U}\|^2 - e^{-\tau}|a|^2 \bar{Q} - e^{-\tau}|a|^2$$

by definitions of $A(\tau)$ and $H(\tau)$. Therefore,

$$\int_{\tau_0}^{\tau} \|(H(s) - \bar{Q}(s)I)\tilde{U}\|^2 ds \lesssim \int_{\tau_0}^{\tau} \|(A(s) - \bar{Q}(s)I)\tilde{U}\|^2 ds + e^{-\tau} \lesssim M + e^{-\tau_0/2} + e^{-\tau/2} < \infty, \quad (3.47)$$

where we used the boundedness of \bar{Q} and a in the first inequality. Combining (3.47) with $\|(H - \bar{Q}(\tau)I)\tilde{U}\| \geq \text{dist}(\bar{Q}(\tau), \text{sp}(H))$ and recalling that $\|\tilde{U}\| = 1$, we conclude that $\text{dist}(\bar{Q}(\tau), \text{sp}(H)) \rightarrow 0$ as $\tau \rightarrow \infty$. It is well-known that $\text{sp}(H) = \{m/2 : m \in \mathbb{N}_0\}$, (cf. [CK, p.664]). Thus we obtain $\bar{Q}(\tau) \rightarrow m/2$ as $\tau \rightarrow \infty$ for some $m \lesssim M_0^a + M_1^b$.

Now, from (3.46), we deduce that for all $\delta > 0$ there exists $\tau_1 > 0$ depending on δ such that

$$-\delta(\tau - \tau_1) \leq \log \|U(\tau)\|^2 - \log \|U(\tau_1)\|^2 + m(\tau - \tau_1) \leq \delta(\tau - \tau_1), \quad \tau \geq \tau_1.$$

Then,

$$e^{-\delta(\tau - \tau_1)} \leq \frac{e^{\tau m} \|U(\tau)\|^2}{e^{\tau_1 m} \|U(\tau_1)\|^2} \leq e^{\delta(\tau - \tau_1)}.$$

Therefore, there exists $A_1(\delta), A_2(\delta) > 0$ such that

$$A_1(\delta)e^{-\delta\tau} \leq e^{\tau m} \|U(\tau)\|^2 \leq A_2(\delta)e^{\delta\tau}. \quad (3.48)$$

Restating (3.48) in the original variable u , we obtain (3.42). \square

The following lemma provides control on $\int_{B(0,R)} Q(x,t)G(x,t) dx$, with Q a homogeneous polynomial of degree d . We use this lemma in our proof of Theorem 2.1.

LEMMA 3.4. *Let $Q = \sum_{|\mu|+2l=d} C_{\mu,l} x^\mu t^l$ be a homogeneous polynomial of degree $d \in \mathbb{N}$. Then,*

$$\int_{\mathbb{R}^n} Q(x,t)G(x,t) dx \lesssim |t|^{d/2}, \quad (3.49)$$

where the constant in (3.49) depends on the polynomial only. Moreover, if all the coordinates of $\mu = (\mu_1, \mu_2, \dots, \mu_n)$ are even, then for all $R > 0$,

$$\int_{B(0,R)} x^\mu t^l G(x,t) dx \lesssim |t|^{l+|\mu|/2},$$

as $t \rightarrow 0^-$. If μ_i is an odd integer for some $i \in \{1, \dots, n\}$, then

$$\int_{B(0,R)} x^\mu t^l G(x,t) dx = 0.$$

For the proof of Lemma 3.4, cf. [CK, p.670]).

PROOF OF THEOREM 2.1. Without loss of generality, let $I = [0, T]$ and $(x_0, t_0) = (0, 0)$. Let m be as in the statement of Lemma 3.3, and denote by d the vanishing order of u at $(0, 0)$. We claim that $d \leq m$. Since the degree of vanishing of u at $(0, 0)$ is d , we have

$$|u(x,t)| \lesssim (|x|^2 + |t|)^{d/2},$$

for all $(x,t) \in Q_1(0,0)$ with $Q_1(0,0)$ defined in (2.4). Also, for any $R > 0$, we have

$$\int_{\mathbb{R}^n \setminus B(0,R)} u^2(x,t)G(x,t) dx \leq \|u\|_{L^\infty(\Omega)}^2 \int_R^\infty \int_{\partial B(0,R)} \frac{e^{-r^2/4|t|}}{|t|^{n/2}} dS_y dr \lesssim \|u\|_{L^\infty(\Omega)}^2 \frac{|t|^{1/2}}{R} e^{-R^2/8|t|}.$$

By Lemma 3.3, there exists $\eta \in (0, \epsilon)$ and $A_1(\delta), A_2(\delta) > 0$ such that

$$A_1(\delta)|t|^{m+\delta} \leq \int_{B(0,R)} u^2(x,t)G(x,t) dx \leq A_2(\delta)|t|^{m-\delta}, \quad t \in [-\eta, 0]. \quad (3.50)$$

Moreover, by Lemma 3.4,

$$\int_{B(0,R)} u^2(x,t)G(x,t) dx \lesssim \int_{B(0,R)} (|x|^2 + |t|)^d G(x,t) dx \lesssim |t|^d, \quad (3.51)$$

where the constant in (3.51) depends on R . Combining (3.50) with (3.51), we get, for a fixed $\delta \in (0, 1]$,

$$A_1(\delta)|t|^{m+\delta} \lesssim |t|^d, \quad t \in [-\eta, 0].$$

Sending $t \rightarrow 0$, we obtain

$$d \leq m + \delta,$$

and then letting $\delta \rightarrow 0$, we conclude that $d \leq m \lesssim M_0^a + M_0^b$ as desired.

Therefore, we have proven Theorem 2.1 when v, w , and u_0 smooth. For the general case, we approximate v, w , and u_0 by smooth functions v^ϵ, w^ϵ , and u_0^ϵ in $L_t^\infty L_x^p, L_t^\infty L_x^q$, and $L_t^\infty L_x^\infty$, respectively. Let u^ϵ be a solution of the equation

$$\begin{aligned} \partial_t u^\epsilon - \Delta u^\epsilon &= w^\epsilon \partial_j u^\epsilon + v^\epsilon u^\epsilon \\ u^\epsilon(x, 0) &= u_0^\epsilon(x). \end{aligned} \quad (3.52)$$

Subtracting (3.52) from (2.1), we get

$$\begin{aligned} \partial_t \tilde{u} - \Delta \tilde{u} &= w^\epsilon \partial_j \tilde{u} + \tilde{w} \partial_j u + v^\epsilon \tilde{u} + \tilde{v} u \\ \tilde{u}(x, 0) &= u_0^\epsilon(x) - u_0(x), \end{aligned} \quad (3.53)$$

where $\tilde{u} = u^\epsilon - u, \tilde{v} = v^\epsilon - v$ and $\tilde{w} = w^\epsilon - w$. First, we have $u, \tilde{u} \in L^\infty L^2 \cap L^2 H^1$. Using $v \in L^\infty L^{\max\{2, n/2\}}$ and $w \in L^\infty L^{2n}$, bootstrapping, and the $W^{2,p}$ maximal regularity for parabolic equations, we get

$$D^2 u, D^2 \tilde{u} \in L^2 L^{2n}(\Omega \times (0, T)). \quad (3.54)$$

Taking the inner product of (3.53) with \tilde{u} , we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 + \|\nabla \tilde{u}\|_{L^2(\Omega)}^2 &= \int w^\epsilon \tilde{u} \partial_j \tilde{u} + \int \tilde{w} \tilde{u} \partial_j u + \int v^\epsilon \tilde{u}^2 + \int \tilde{v} u \tilde{u} \\ &\lesssim \|w^\epsilon\|_{L^q(\Omega)} \|\tilde{u}\|_{L^{2q/(q-2)}(\Omega)} \|\nabla \tilde{u}\|_{L^2(\Omega)} + \|\tilde{w}\|_{L^q(\Omega)} \|\tilde{u}\|_{L^{2q/(q-2)}(\Omega)} \|\nabla u\|_{L^2(\Omega)} \\ &\quad + \|v^\epsilon\|_{L^p(\Omega)} \|\tilde{u}\|_{L^{2p/(p-1)}(\Omega)}^2 + \|\tilde{v}\|_{L^p(\Omega)} \|u\|_{L^{2p/(p-1)}(\Omega)} \|\tilde{u}\|_{L^{2p/(p-1)}(\Omega)}. \end{aligned}$$

Using $\|\tilde{u}\|_{L^{2q/(q-2)}(\Omega)} \lesssim \|\tilde{u}\|_{L^2(\Omega)}^{1-n/q} \|\nabla \tilde{u}\|_{L^2(\Omega)}^{n/q}$ and $\|\tilde{u}\|_{L^{2p/(p-1)}(\Omega)} \lesssim \|\tilde{u}\|_{L^2(\Omega)}^{1-n/2p} \|\nabla \tilde{u}\|_{L^2(\Omega)}^{n/2p}$, with the same inequality for u , and applying Young's inequality, we get, for an arbitrary $\epsilon_0 \in (0, 1]$,

$$\begin{aligned} \frac{d}{dt} \|\tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 &\leq C \|w^\epsilon\|_{L^q(\Omega)}^{2q/(q-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + C_{\epsilon_0} \|\tilde{w}\|_{L^q(\Omega)}^{2q/(q-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 \\ &\quad + C \|v^\epsilon\|_{L^p(\Omega)}^{2p/(2p-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + C_{\epsilon_0} \|\tilde{v}\|_{L^p(\Omega)}^{2p/(2p-n)} \|\tilde{u}\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)} + \epsilon_0 \|\nabla u\|_{L^2(\Omega)}^2. \end{aligned} \quad (3.55)$$

Integrating (3.55) from 0 to t_0 , we obtain

$$\begin{aligned} &\|\tilde{u}(\cdot, t_0)\|_{L^2(\Omega)}^2 - \|\tilde{u}(\cdot, 0)\|_{L^2(\Omega)}^2 \\ &\leq C \int_0^{t_0} \|w^\epsilon\|_{L^q(\Omega)}^{2q/(q-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + C_{\epsilon_0} \int_0^{t_0} \|\tilde{w}\|_{L^q(\Omega)}^{2q/(q-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 \\ &\quad + C \int_0^{t_0} \|v^\epsilon\|_{L^p(\Omega)}^{2p/(2p-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + C_{\epsilon_0} \int_0^{t_0} \|\tilde{v}\|_{L^p(\Omega)}^{2p/(2p-n)} \|\tilde{u}\|_{L^2(\Omega)} \|u\|_{L^2(\Omega)} + \epsilon_0 \int_0^{t_0} \|\nabla u\|_{L^2(\Omega)}^2, \end{aligned}$$

for every $\epsilon_0 \in (0, 1]$. Sending $\epsilon \rightarrow 0^+$, it follows that

$$\limsup_{\epsilon \rightarrow 0} \|\tilde{u}(\cdot, t_0)\|_{L^2(\Omega)}^2 \leq \epsilon_0 \int_0^{t_0} \|\nabla u\|_{L^2(\Omega)}^2, \quad t_0 \geq 0,$$

for every $\epsilon_0 \in (0, 1]$. Sending then $\epsilon_0 \rightarrow 0$, we then obtain

$$\|u^\epsilon(\cdot, t_0)\|_{L^2(\Omega)} \rightarrow \|u(\cdot, t_0)\|_{L^2(\Omega)}, \quad t_0 \geq 0. \quad (3.56)$$

Now, we claim that $\nabla u^\epsilon \rightarrow \nabla u$ in $L_t^\infty L_x^2$. To prove this, we test the first equation in (3.53) with $-\Delta \tilde{u}$, which we may by (3.54), and write

$$\frac{d}{dt} \|\nabla \tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 + \|\Delta \tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 = - \int w^\epsilon \partial_j \tilde{u} \Delta \tilde{u} dx - \int \tilde{w} \partial_j u \Delta \tilde{u} dx - \int v^\epsilon \tilde{u} \Delta \tilde{u} dx - \int \tilde{v} u \Delta \tilde{u} dx.$$

By Hölder's and Sobolev's inequalities, we have

$$\begin{aligned}
& \frac{d}{dt} \|\nabla \tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 + \|\Delta \tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 \\
& \lesssim \|w^\epsilon\|_{L^q(\Omega)} \|\nabla \tilde{u}\|_{L^{2q/(q-2)}(\Omega)} \|\Delta \tilde{u}\|_{L^2(\Omega)} + \|\tilde{w}\|_{L^q(\Omega)} \|\nabla u\|_{L^{2q/(q-2)}(\Omega)} \|\Delta \tilde{u}\|_{L^2(\Omega)} \\
& \quad + \|v^\epsilon\|_{L^p(\Omega)} \|\tilde{u}\|_{L^{2p/(p-2)}(\Omega)} \|\Delta \tilde{u}\|_{L^2(\Omega)} + \|\tilde{v}\|_{L^p(\Omega)} \|u\|_{L^{2p/(p-2)}(\Omega)} \|\Delta \tilde{u}\|_{L^2(\Omega)} \\
& \lesssim \|w^\epsilon\|_{L^q(\Omega)} \|\tilde{u}\|_{L^2(\Omega)}^{1/2-n/2q} \|\Delta \tilde{u}\|_{L^2(\Omega)}^{3/2+n/2q} + \|\tilde{w}\|_{L^q(\Omega)} \|u\|_{L^2(\Omega)}^{1/2-n/2q} \|\Delta u\|_{L^2(\Omega)}^{1/2+n/2q} \|\Delta \tilde{u}\|_{L^2(\Omega)} \\
& \quad + \|v^\epsilon\|_{L^p(\Omega)} \|\tilde{u}\|_{L^2(\Omega)}^{1-n/2p} \|\Delta \tilde{u}\|_{L^2(\Omega)}^{n/2p+1} + \|\tilde{v}\|_{L^p(\Omega)} \|u\|_{L^2(\Omega)}^{1-n/2p} \|\Delta u\|_{L^2(\Omega)}^{n/2p} \|\Delta \tilde{u}\|_{L^2(\Omega)}.
\end{aligned} \tag{3.57}$$

Applying Young's inequality on all four terms and absorbing the factors containing $\Delta \tilde{u}$, we get

$$\begin{aligned}
\frac{d}{dt} \|\nabla \tilde{u}(\cdot, t)\|_{L^2(\Omega)}^2 & \lesssim \|w^\epsilon\|_{L^q(\Omega)}^{4q/(q-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + \|\tilde{w}\|_{L^q(\Omega)}^2 \|u\|_{L^2(\Omega)}^{1-n/q} \|\Delta u\|_{L^2(\Omega)}^{1+n/q} \\
& \quad + \|v^\epsilon\|_{L^p(\Omega)}^{4p/(2p-n)} \|\tilde{u}\|_{L^2(\Omega)}^2 + \|\tilde{v}\|_{L^p(\Omega)}^2 \|u\|_{L^2(\Omega)}^{2-n/p} \|\Delta u\|_{L^2(\Omega)}^{n/p}.
\end{aligned} \tag{3.58}$$

For any $t_0 \geq 0$, we integrate (3.58) from 0 to t_0 to obtain

$$\begin{aligned}
& \|\nabla \tilde{u}(\cdot, t_0)\|_{L^2(\Omega)}^2 - \|\nabla \tilde{u}(\cdot, 0)\|_{L^2(\Omega)}^2 \\
& \lesssim t_0 \|w^\epsilon\|_{L^\infty L^q(\Omega \times I)} \|\tilde{u}\|_{L^\infty L^2(\Omega \times I)}^2 + \|\tilde{w}\|_{L^\infty L^q(\Omega \times I)}^2 \|u\|_{L^\infty L^2(\Omega \times I)}^{1-n/q} \int_0^{t_0} \|\Delta \tilde{u}\|_{L^2(\Omega)}^{1+n/q} \\
& \quad + t_0 \|v^\epsilon\|_{L^\infty L^p(\Omega \times I)}^{4p/(2p-n)} \|\tilde{u}\|_{L^\infty L^2(\Omega \times I)}^2 + \|\tilde{v}\|_{L^\infty L^p(\Omega \times I)}^2 \|u\|_{L^2 L^{2p/(p-2)}(\Omega \times I)}^{2-n/p} \int_0^{t_0} \|\Delta u\|_{L^2 L^2(\Omega \times I)}^{n/p},
\end{aligned}$$

where $\|\Delta \tilde{u}\|_{L^2 L^2(\Omega \times I)} < \infty$ due to (3.54). Letting $\epsilon \rightarrow 0^+$, we obtain

$$\|\nabla u^\epsilon(\cdot, t_0)\|_{L^2(\Omega)} \rightarrow \|\nabla u(\cdot, t_0)\|_{L^2(\Omega)}, \quad t_0 \geq 0. \tag{3.59}$$

Denote by $\bar{Q}^\epsilon(\tau)$ and $\bar{Q}(\tau)$ the modified frequency functions corresponding to u^ϵ and u , respectively. Recalling that

$$\bar{Q}(\tau) = \frac{(HU, U)}{\|U\|^2} - \frac{e^{-\tau/2} a_j}{\|U\|^2} \int y_j U^2 dy,$$

we now claim that $\bar{Q}^\epsilon(\tau) \rightarrow \bar{Q}(\tau)$ as $\epsilon \rightarrow 0$ for all τ . Note that

$$\|U^\epsilon(\tau)\|^2 - \|U(\tau)\|^2 = \int (u_\epsilon^2 - u^2) G(x, t) dx \lesssim \|\tilde{u}\|_{L^2(\Omega)} \|(u^\epsilon + u)\|_{L^2(\Omega)} \rightarrow 0, \tag{3.60}$$

as $\epsilon \rightarrow 0^+$ by (3.56). For $(HU_\epsilon, U_\epsilon) - (HU, U)$, we have

$$(HU_\epsilon, U_\epsilon) - (HU, U) = |t| \int (|\nabla u_\epsilon|^2 - |\nabla u|^2) G(x, t) dx \lesssim |t| \|\nabla \tilde{u}\|_{L^2(\Omega)} \|\nabla(u^\epsilon + u)\|_{L^2(\Omega)} \rightarrow 0,$$

as $\epsilon \rightarrow 0^+$ by (3.59). Lastly, we estimate $\int y_j U_\epsilon^2 dy - \int y_j U^2 dy$ as

$$\begin{aligned}
& \int y_j U_\epsilon^2 dy - \int y_j U^2 dy = \int y_j (U_\epsilon^2 - U^2) \lesssim \|y(U_\epsilon - U)\| \|U_\epsilon + U\| \\
& \lesssim ((H(U_\epsilon - U), U_\epsilon - U)^{1/2} + \|U_\epsilon - U\|) \|U^\epsilon + U\| \\
& \lesssim (|t| \|\nabla(u^\epsilon - u)\| \|\nabla(u^\epsilon - u)G\|^{1/2} + \|U_\epsilon - U\|) \|U_\epsilon + U\| \rightarrow 0,
\end{aligned} \tag{3.61}$$

as $\epsilon \rightarrow 0^+$ where we used $\|yU\|^2 \lesssim (HU, U) + \|U\|^2$ in the second inequality. By (3.60)–(3.61), we obtain $\bar{Q}^\epsilon(\tau) \rightarrow \bar{Q}(\tau)$ as $\epsilon \rightarrow 0$ for all τ . Note that we have

$$\bar{Q}^\epsilon(\tau) \lesssim M_0^a + M_1^b, \tag{3.62}$$

uniformly in τ and $\epsilon \in (0, 1]$. Passing to the limit, we get

$$\bar{Q}(\tau) \lesssim M_0^a + M_1^b, \quad \tau \geq 0. \tag{3.63}$$

Next, using $\text{dist}(\bar{Q}^\epsilon(\tau), \text{sp}(H)) \lesssim \|(H - \bar{Q}(\tau)I)\tilde{U}\|$ and (3.47), we obtain, for all $\epsilon \in (0, 1]$

$$\int_{\tau_0}^\tau \text{dist}(\bar{Q}^\epsilon(\tau), \text{sp}(H))^2 \lesssim M_0^a + M_1^b + e^{-\tau_0/2} + e^{-\tau/2} \lesssim M_0^a + M_1^b = K, \tag{3.64}$$

where we used $M_0, M_1 \geq 1$. Observe that by (3.62) we have $Q^\epsilon(\tau) \in [0, \bar{M}]$, where \bar{M} is independent of ϵ and τ . Thus letting $\epsilon \rightarrow 0$ in (3.64) yields

$$\int_{\tau_0}^{\tau} \text{dist}(\bar{Q}(\tau), \text{sp}(H))^2 \leq K. \quad (3.65)$$

Combining (3.63), (3.65), and $\text{sp}(H) = \{m/2 : m \in \mathbb{N}_0\}$, we get $\bar{Q}(\tau) \rightarrow m/2$ for some $m \lesssim M_0^a + M_1^b$ with a and b as in (2.7). The rest of the proof is similar to the case when v, w , and u_0 are smooth. \square

4. Doubling property

In the proof of Theorem 2.2, we need the following observability lemmas.

LEMMA 4.1. *Let $K \geq 0$ and $\delta_0 \in (0, 1/2]$. If a Ω -periodic function u satisfies*

$$\|u\|_{L^2(\Omega)}^2 \leq e^K \int_{\mathbb{R}^n} u(x)^2 G(x, -\delta^2) dx$$

with $\delta \in (0, \delta_0]$ so small that

$$\frac{1}{\delta^2} \geq \frac{C}{\delta_0^2} \log \frac{1}{\delta} + \frac{C(K+1)}{\delta_0^2},$$

for a sufficiently large constant $C > 0$, then

$$\|u\|_{L^2(\Omega)}^2 \leq \frac{Ce^K}{\delta^n} \|u\|_{L^2(B_{\delta_0})}^2,$$

where $\Omega = [0, 1]^n$.

For the proof, cf. [Ku4, p.775].

LEMMA 4.2. *Under assumptions of Theorem 2.1, let $t_0 < 0$ such that $|t_0| < \epsilon$ with ϵ defined in (3.37). Then, for any $\delta_0 \in (0, 1/2]$, we have*

$$\|u(\cdot, t)\|_{L^2(\Omega)}^2 \lesssim e^{(n+1)\delta_0^2/\delta^2} \|u(\cdot, t)\|_{L^2(B_{\delta_0})}^2, \quad t \in [-\delta^2, 0] \quad (4.1)$$

with $\delta > 0$ satisfying

$$\frac{1}{\delta^2} \geq \frac{\log 1/\delta}{\delta_0^2} + \frac{\gamma(\delta) + 1}{\delta_0^2} \quad (4.2)$$

and

$$\delta < \min\{\sqrt{\epsilon}, \delta_0\}, \quad (4.3)$$

where

$$\gamma(\delta) = M_1^2 + M_0 + M_1 M^{2\beta-1} + M_0 M^{2\alpha-1} + M \log(1/\delta^2), \quad (4.4)$$

with M defined in (3.45). Moreover,

$$\delta = \min \left\{ \sqrt{\epsilon}, \frac{\delta_0}{\log(1/\delta_0)}, \frac{\delta_0^2}{M^2}, \frac{\delta_0}{M_1 + M_0^{1/2} + M^{\beta-1/2} M_0^{1/2} + M_0^{1/2} M^{\alpha-1/2}} \right\}$$

satisfies the conditions (4.2) and (4.3). Therefore, $(n+1)\delta_0^2/\delta^2$ can be replaced by a polynomial P with respect δ_0 and P depends on n, M_0, M_1, p , and q .

PROOF OF LEMMA 4.2. From (3.44), we see that

$$\frac{f(\tau)}{\|U\|^2} \gtrsim -e^{(\beta-1)\tau} M_1 (\bar{Q}(\tau)_+ + e^{-\tau}|a|^2 + 1)^\beta - e^{(\alpha-1)\tau} M_0 (\bar{Q}(\tau)_+ + e^{-\tau}|a|^2 + 1)^\alpha,$$

where α and β are as in (3.23) and (3.29). Therefore, by replacing τ_1 by τ_0 in (3.46), we have

$$\begin{aligned} \frac{1}{2} \log \|U(\cdot, \tau)\|^2 - \frac{1}{2} \log \|U(\cdot, \tau_0)\|^2 &= - \int_{\tau_0}^{\tau} \bar{Q}(s) ds + \int_{\tau_0}^{\tau} \frac{f(s)}{\|U(\cdot, s)\|^2} ds \\ &\gtrsim -(\tau - \tau_0) \frac{2C_1 + C_2}{\epsilon} - e^{(\beta-1)\tau_0} M_1 \left(\frac{2C_1 + C_2}{\epsilon} + e^{-\tau_0}|a|^2 + 1 \right)^\beta \\ &\quad - e^{(\alpha-1)\tau_0} M_0 \left(\frac{2C_1 + C_2}{\epsilon} + e^{-\tau_0}|a|^2 + 1 \right)^\alpha, \end{aligned}$$

where we used $\bar{Q}(\tau) \lesssim (2C_1 + C_2)/\epsilon$ for all $\tau \geq \tau_0$ in the last inequality. Since $\epsilon = e^{-\tau_0}$, we have

$$\log \|U(\cdot, \tau)\|^2 - \log \|U(\cdot, \tau_0)\|^2 \gtrsim -(\tau - \tau_0)\epsilon^{-1} - \epsilon^{1-\beta} M_1 (\epsilon^{-1} + \epsilon|a|^2 + 1)^\beta - \epsilon^{1-\alpha} M_0 (\epsilon^{-1} + \epsilon|a|^2 + 1)^\alpha. \quad (4.5)$$

Let $q_D(t) = \|\nabla u\|_{L^2(\Omega)}^2 / \|u\|_{L^2(\Omega)}^2$. We now derive a similar estimate for $\|u(\cdot, t)\|_{L^2(\Omega)}^2 / \|u(\cdot, -\epsilon)\|_{L^2(\Omega)}^2$. Taking the inner product of (3.7) with u and using $\|\nabla u\|_{L^2(\Omega)}^2 = q_D(t)\|u\|_{L^2(\Omega)}^2$, we obtain

$$\begin{aligned} & \frac{d}{dt} \|u(\cdot, t)\|_{L^2(\Omega)}^2 + q_D(t) \|u(\cdot, t)\|_{L^2(\Omega)}^2 \\ & \lesssim \|w(\cdot, t)\|_{L^q(\Omega)} \|u(\cdot, t)\|_{L^{2q/(q-2)}(\Omega)} \|u(\cdot, t)\|_{L^2(\Omega)} q_D(t)^{1/2} + \|v(\cdot, t)\|_{L^p(\Omega)} \|u(\cdot, t)\|_{L^{2p/(p-1)}(\Omega)}^2 \\ & \lesssim M_1 \|u(\cdot, t)\|_{L^2(\Omega)}^{2-2\beta} \|\nabla u(\cdot, t)\|_{L^2(\Omega)}^{2\beta-1} \|u(\cdot, t)\|_{L^2(\Omega)} q_D(t)^{1/2} + M_0 \|u(\cdot, t)\|_{L^2(\Omega)}^{2-2\alpha} \|\nabla u(\cdot, t)\|_{L^2(\Omega)}^{2\alpha} \end{aligned} \quad (4.6)$$

since $\int_\Omega a_j u \partial_j u \, dx$ vanishes. Dividing both sides of (4.6) by $\|u(\cdot, t)\|_{L^2(\Omega)}^2$, we get

$$\begin{aligned} & \frac{1}{\|u(\cdot, t)\|_{L^2(\Omega)}^2} \frac{d}{dt} \|u(\cdot, t)\|_{L^2(\Omega)}^2 + q_D(t) \lesssim M_1 \frac{\|\nabla u(\cdot, t)\|_{L^2(\Omega)}^{2\beta-1}}{\|u(\cdot, t)\|_{L^2(\Omega)}^{2\beta-1}} q_D(t)^{1/2} + M_0 \frac{\|\nabla u(\cdot, t)\|_{L^2(\Omega)}^{2\alpha}}{\|u(\cdot, t)\|_{L^2(\Omega)}^{2\alpha}} \\ & \lesssim M_1 q_0^{\beta-1/2} q_D(t)^{1/2} + M_0 q_0^\alpha \lesssim M_1 q_D(t)^{1/2} + M_0 \end{aligned} \quad (4.7)$$

if we allow the last implicit constant to depend on q_0 . Applying Young's inequality to the right-hand side of (4.7), we obtain

$$\frac{1}{\|u(\cdot, t)\|_{L^2(\Omega)}^2} \frac{d}{dt} \|u(\cdot, t)\|_{L^2(\Omega)}^2 \lesssim M_1^2 + M_0. \quad (4.8)$$

Integrating (4.8) from $-\epsilon$ to t leads to

$$\log \frac{\|u(\cdot, t)\|_{L^2(\Omega)}^2}{\|u(\cdot, -\epsilon)\|_{L^2(\Omega)}^2} \lesssim (M_1^2 + M_0)(t + \epsilon) \lesssim (M_1^2 + M_0)\epsilon, \quad t \in (\epsilon, 0). \quad (4.9)$$

By definition of U in (3.9), we have

$$\frac{\|U(\cdot, \tau_0)\|^2}{\|u(\cdot, -\epsilon)\|_{L^2(\Omega)}^2} = \frac{\int_{\mathbb{R}^n} u(x, -\epsilon)^2 e^{-|x|^2/4\epsilon} \, dx}{\epsilon^{n/2} \int_\Omega u(x, -\epsilon)^2 \, dx} \geq \frac{e^{-n/4\epsilon}}{\epsilon^{n/2}} \quad (4.10)$$

since $|x| \leq \sqrt{n}$ for $x \in \Omega$. We then take the logarithm of (4.10) to get

$$\log \frac{\|u(\cdot, -\epsilon)\|_{L^2(\Omega)}^2}{\|U(\cdot, \tau_0)\|^2} \leq -\frac{n}{2} \log \frac{1}{\epsilon} + \frac{n}{4\epsilon} \lesssim \frac{1}{\epsilon}. \quad (4.11)$$

Using (4.5), (4.9), and (4.11), we obtain

$$\begin{aligned} \log \frac{\|u(\cdot, t)\|_{L^2(\Omega)}^2}{\|U(\cdot, \tau)\|^2} & \lesssim (M_1^2 + M_0)\epsilon + \epsilon^{-1} + (\tau - \log \epsilon^{-1})\epsilon^{-1} \\ & \quad + \epsilon^{1-\beta} M_1 (\epsilon^{-1} + \epsilon|a|^2 + 1)^\beta + \epsilon^{1-\alpha} M_0 (\epsilon^{-1} + \epsilon|a|^2 + 1)^\alpha \\ & \lesssim M_1^2 + M_0 + \epsilon^{-1} + \tau\epsilon^{-1} + M_1\epsilon^{1-2\beta} + M_0\epsilon^{1-2\alpha}, \end{aligned}$$

from where, after an short computation,

$$\log \frac{\|u(\cdot, t)\|_{L^2(\Omega)}^2}{\|U(\cdot, \tau)\|^2} \lesssim M_1^2 + M_0 + M\tau + M_1 M^{2\beta-1} + M_0 M^{2\alpha-1}, \quad (4.12)$$

with $M = M_0^{2/(3-4\alpha)} + M_1^{2/(3-4\beta)}$ as defined in (3.45). Fix $\delta_0 \in (0, 1/2]$. Let $t = -\delta^2$, i.e., $\tau = 2 \log(1/\delta)$. If $\delta \in (0, \sqrt{\epsilon})$, then

$$\log \frac{\|u(\cdot, -\delta^2)\|_{L^2(\Omega)}^2}{\|U(\cdot, \log 1/\delta^2)\|^2} \lesssim \gamma(\delta)$$

with $\gamma(\delta)$ defined in (4.4), which equals to the right-hand side of (4.12). Therefore,

$$\|u(\cdot, -\delta)\|_{L^2(\Omega)}^2 \lesssim e^{\gamma(\delta)} \int_{\mathbb{R}^n} u(x, -\delta^2)^2 G(x, -\delta^2) \, dx.$$

By Lemma 4.1, we conclude that

$$\|u(\cdot, -\delta^2)\|_{L^2(\Omega)}^2 \lesssim \frac{e^{\gamma(\delta)}}{\delta^n} \int_{B_{\delta_0}} u(x, -\delta^2)^2 dx.$$

Our assumption on δ in (4.2) gives $\gamma(\delta) \lesssim \delta_0^2/\delta^2$ and $\log(1/\delta) \leq \delta_0^2/\delta^2$. So we obtain (4.1) for $t = -\delta^2$. To get (4.1) for $t \in (-\delta^2, 0]$, we simply translate in time. Therefore, we conclude that

$$\|u(\cdot, t)\|_{L^2(\Omega)}^2 \lesssim e^{(n+1)\delta_0^2/\delta^2} \|u(\cdot, t)\|_{L^2(B_{\delta_0})}^2, \quad t \in [-\delta^2, 0],$$

as desired. \square

PROOF OF THEOREM 2.2. We first assume that u_0, v , and w are smooth as in the proof of Theorem 2.1. Fix any $t \in (0, T)$. For all $\delta_0 \in (0, 1/2]$, by restating Lemma 4.2 to address positive time instead of negative, we claim that

$$\|u(\cdot, s)\|_{L^2(\Omega)}^2 \lesssim e^{P(n, \delta_0, M_0, M_1, p, p)} \|u(\cdot, s)\|_{L^2(B_{\delta_0})}^2, \quad (4.13)$$

for all $s \in [0, \delta^2]$, where P is determined by Lemma 4.2. Choose $t' \in (0, t)$ such that $t - t' < \epsilon$ with ϵ as in Lemma 4.2. By (4.13), we have

$$\|u(\cdot, t')\|_{L^2(\Omega)}^2 \lesssim e^{P(n, \delta_0, M_0, M_1, q, p)} \|u(\cdot, t')\|_{L^2(B_{\delta_0})}^2.$$

Therefore, (2.10) holds for all $t \in (0, T)$. Thus we have proven Theorem 2.2 under the assumption that u_0, v , and w are smooth. For the general case, we approximate u_0, v , and w by smooth functions u_0^ϵ, v^ϵ , and w^ϵ respectively. We then have $\|u^\epsilon(\cdot, t)\|_{L^2(\Omega)}^2 \rightarrow \|u(\cdot, t)\|_{L^2(\Omega)}^2$ and $\|u^\epsilon(\cdot, t)\|_{L^2(B_{\delta_0})}^2 \rightarrow \|u(\cdot, t)\|_{L^2(B_{\delta_0})}^2$ as $\epsilon \rightarrow 0^+$ by (3.57). Thus (2.10) holds for all $t \in (0, T)$, as desired. \square

5. The case \mathbb{R}^n

In this section, we prove the theorem concerning the case of \mathbb{R}^n .

PROOF OF THEOREM 2.3. Note that we now assume the growth condition (2.8) instead of periodicity. The proof is similar to the periodic case with small modifications. With $\epsilon \in (0, T)$ such that $\epsilon \leq 1/4$, we have

$$\frac{\epsilon \int_{\mathbb{R}^n} |\nabla u(x_\epsilon + y, -\epsilon)|^2 G(y, -\epsilon) dy}{\int_{\mathbb{R}^n} u(x_\epsilon + y, -\epsilon)^2 G(y, -\epsilon) dy} \leq 4K\epsilon q_D(-\epsilon), \quad (5.1)$$

for some $x_\epsilon \in B_2$. This was proven in [CK], but since the argument is short, we present it here. To see why (5.1) holds, we assume that

$$\lambda \int_{\mathbb{R}^n} u(x + y, -\epsilon)^2 G(y, -\epsilon) dy \leq \int_{\mathbb{R}^n} |\nabla u(x + y, -\epsilon)|^2 G(y, -\epsilon) dy, \quad x \in B_2, \quad (5.2)$$

where $\lambda = 4Kq_D(-\epsilon)$. Integrating (5.2) over B_2 and proceeding similarly as in (3.1), we have

$$\begin{aligned} \int_{\mathbb{R}^n} G(y, -\epsilon) dy \int_{B_2} u(x + y, -\epsilon)^2 dx &\leq \frac{1}{\lambda} \int_{\mathbb{R}^n} G(y, -\epsilon) dy \int_{B_2} |\nabla u(x + y, -\epsilon)|^2 dx \\ &\leq \frac{1}{\lambda} \|\nabla u(\cdot, -\epsilon)\|^2 \int_{\mathbb{R}^n} G(y, -\epsilon) dy \leq \frac{(2\pi)^{n/2}}{\lambda} \|\nabla u(\cdot, -\epsilon)\|^2, \end{aligned} \quad (5.3)$$

where we continue using the convention (3.12). On the other hand, we have the lower bound for the left hand side of (5.3)

$$\begin{aligned} \int_{\mathbb{R}^n} G(y, -\epsilon) dy \int_{B_2} u(x + y, -\epsilon)^2 dx &= \int_{\mathbb{R}^n} G(y, -\epsilon) dy \int_{B_2(y)} u(x, -\epsilon)^2 dx \\ &= \int_{B_{1/2}} G(y, -\epsilon) dy \int_{B_2(y)} u(x, -\epsilon)^2 dx + \int_{\mathbb{R}^n \setminus B_{1/2}} G(y, -\epsilon) dy \int_{B_2(y)} u(x, -\epsilon)^2 dx \\ &\geq \int_{B_{1/2}} G(y, -\epsilon) dy \int_{B_2(y)} u(x, -\epsilon)^2 dx \geq \int_{B_{1/2}} G(y, -\epsilon) dy \int_{B_{1/2}} G(y, -\epsilon) dy \int_{B_1} u(x, -\epsilon)^2 dx \\ &\geq \frac{1}{2} (2\pi)^{n/2} \int_{B_1} u(x, -\epsilon)^2 dx \geq \frac{1}{2K} (2\pi)^{n/2} \int_{\mathbb{R}^n} u(x, -\epsilon)^2 dx, \end{aligned}$$

where the last inequality holds by the doubling condition (2.8). Thus we obtain

$$\|u(x, -\epsilon)\|^2 \leq \frac{2K}{\lambda} \|\nabla u(\cdot, -\epsilon)\|^2 \leq \frac{1}{2q_D(-\epsilon)} \|\nabla u(\cdot, -\epsilon)\|^2.$$

We have a contradiction since $\|\nabla u(\cdot, -\epsilon)\|^2/\|u(\cdot, -\epsilon)\|^2 \leq q_D(-\epsilon)$. Therefore, (5.2) does not hold for all $x \in B_2$, i.e., there exists $x_\epsilon \in B_2$ such that (5.1) holds true. Theorem 2.3 then follows as in the proofs of Theorems 2.1 and 2.2. \square

6. $L_t^q L_x^p$ case

The method we used also works for the case where $v \in L_t^{p_2} L_x^{p_1}$ and $w \in L_t^{q_2} L_x^{q_1}$ when p_2 and q_2 are finite, as shown in the next statement. When limiting $p_2 \rightarrow \infty$ and $q_2 \rightarrow \infty$, the theorem reduces to the results in Section 2.

THEOREM 6.1. *Let $n \geq 2$. Assume that $u \in L_{x,t}^\infty(\Omega \times I)$ is a solution of (2.1) with $v \in L_t^{p_2} L_x^{p_1}(\Omega \times I)$ and $w \in L_t^{q_2} L_x^{q_1}(\Omega \times I)$ are such that $p_1 > 2n/3$ and $q_1 > 2n$ with*

$$p_2 > \max \left\{ \frac{2}{3-4\alpha}, \frac{2}{1-\alpha} \right\} \quad (6.1)$$

and

$$q_2 > \max \left\{ \frac{2}{3-4\beta}, \frac{2}{1-\beta} \right\},$$

where $\alpha = n/2p_1$ and $\beta = 1/2 + n/2q_1$. Then for all $(x_0, t_0) \in \mathbb{T}^n \times I$, the vanishing order of u at (x_0, t_0) satisfies

$$O_{(x_0, t_0)}(u) \lesssim \|v\|_{L_t^{p_2} L_x^{p_1}(\Omega \times I)}^a + \|w\|_{L_t^{q_2} L_x^{q_1}(\Omega \times I)}^b + 1$$

where

$$a = \frac{2}{3-2/p_2-4\alpha} \quad (6.2)$$

and

$$b = \frac{2}{3-2/q_2-4\beta}.$$

PROOF OF THEOREM 6.1. Without loss of generality, we can assume $I = [-1, 0]$ and $(x_0, t_0) = (0, 0)$. For simplicity, we only consider the case $w = 0$ and $v \in L_t^{p_2} L_x^{p_1}$, as for a nonzero w the proof is similar. By a simple change of variables, we obtain

$$\int_{-1}^0 \left(\int |v(x, t)|^{p_1} dx \right)^{p_2/p_1} dt = \int_0^\infty \left(\int |V(y, \tau)|^{p_1} e^{-n\tau/2} dy \right)^{p_2/p_1} e^{-\tau} d\tau = \int_0^\infty K_0(\tau)^{p_2} e^{-\tau} d\tau,$$

where $K_0(\tau) = \|V\|_{L^{p_1}} e^{-n\tau/2p_1}$. Therefore,

$$\int_0^\infty K_0(\tau)^{p_2} e^{-\tau} d\tau \leq M_0^{p_2},$$

where $M_0 = \|v\|_{L_t^{p_2} L_x^{p_1}(\Omega \times I)} + 1$. Note that $K_0(\tau)$ plays the role of M_0 in the proof of Theorem 2.1, except that here K_0 is a function of τ . As in the proof of Theorem 2.1, we have

$$\begin{aligned} \bar{Q}'(\tau) &\lesssim e^{-\tau/2} \epsilon^{-1} \bar{Q}_+^{1/2} + K_0(\tau)^2 e^{(\alpha-2)\tau} \epsilon^{-2\alpha} \bar{Q}_+^\alpha + K_0(\tau)^2 e^{(2\alpha-2)\tau} \bar{Q}_+^{2\alpha} + e^{-\tau} \epsilon^{-2} + e^{-\tau/2} \epsilon^{-1} \\ &\quad + K_0(\tau)^2 e^{-2\tau} \epsilon^{-4\alpha} + K_0(\tau)^2 e^{(\alpha-2)\tau} \epsilon^{-2\alpha} + K_0(\tau)^2 e^{(1-\alpha)\tau} e^{-2\tau} + K_0(\tau)^2 e^{(2\alpha-2)\tau}, \end{aligned} \quad (6.3)$$

where $\alpha = n/2p_1$ and $\epsilon = M_0^{-a}/K$ with a as in (6.2) and K sufficiently large. Denote by $C_2 \geq 1$ the implicit constant in (6.3). Under the condition (6.1), we claim that

$$\bar{Q}_+(\tau) < \frac{C_1 + C_2}{\epsilon},$$

where $C_1 = C_0(q_0 + q_0^{1/2})$ and $C_0 \geq 1$ is sufficiently large. To prove this claim, we use a barrier argument. Assume, contrary to the assertion, that there exists τ_1 such that $\bar{Q}(\tau_1) = (C_1 + C_2)/\epsilon$, and suppose that τ_1 is the first time with

this property. We have

$$\begin{aligned}\bar{Q}'(\tau) &\leq C_2(C_1 + C_2)^{1/2}e^{-\tau/2}\epsilon^{-3/2} + C_2K_0(\tau)^2e^{(\alpha-2)\tau}\epsilon^{-3\alpha}(C_1 + C_2)^\alpha \\ &\quad + C_2K_0(\tau)^2e^{(2\alpha-2)\tau}(C_1 + C_2)^{2\alpha}\epsilon^{-2\alpha} + C_2e^{-\tau}\epsilon^{-2} + C_2K_0(\tau)^2e^{-2\tau}\epsilon^{-4\alpha} \\ &\quad + C_2K_0(\tau)^2e^{(\alpha-2)\tau}\epsilon^{-2\alpha} + C_2K_0(\tau)^{2/(1-\alpha)}e^{-2\tau} + C_2K_0(\tau)^2e^{(2\alpha-2)\tau}.\end{aligned}$$

We integrate (6.3) in τ from τ_0 to τ_1 , where τ_0 is given in (3.10), to obtain

$$\begin{aligned}\bar{Q}(\tau_1) &\leq \bar{Q}(\tau_0) + 2C_2(C_1 + C_2)^{1/2}\epsilon^{-1} + C_2(C_1 + C_2)^\alpha\epsilon^{-3\alpha}\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{(\alpha-2)\tau} d\tau \\ &\quad + C_2(C_1 + C_2)^{2\alpha}\epsilon^{-2\alpha}\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{(2\alpha-2)\tau} d\tau + C_2\epsilon^{-1} + C_2\epsilon^{-4\alpha}\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{-2\tau} d\tau \\ &\quad + C_2\epsilon^{-2\alpha}\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{(\alpha-2)\tau} d\tau + C_2\int_{\tau_0}^{\tau_1} K_0(\tau)^{2/(1-\alpha)}e^{-2\tau} d\tau + C_2\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{(2\alpha-2)\tau} d\tau.\end{aligned}\tag{6.4}$$

By Hölder's inequality, the third term on the right-hand side satisfies

$$\begin{aligned}C_2(C_1 + C_2)^\alpha\epsilon^{-3\alpha}\int_{\tau_0}^{\tau_1} K_0(\tau)^2e^{(\alpha-2)\tau} d\tau &\leq C_2(C_1 + C_2)^\alpha\epsilon^{-3\alpha}\|K_0(\tau)^2e^{-2\tau/p_2}\|_{L^{p_2/2}(0,\infty)}\|e^{(\alpha-2+2/p_2)\tau}\|_{L^{p_2/(p_2-2)}(0,\infty)} \\ &\leq C_1(C_1 + C_2)^\alpha\epsilon^{-3\alpha}\left(\int_0^\infty K_0(\tau)^{p_2}e^{-\tau} d\tau\right)^{2/p_2}\epsilon^{2-\alpha-2/p_2} \leq C_1(C_1 + C_2)^\alpha\epsilon^{3-4\alpha-2/p_2}M_0^2\epsilon^{-1} \\ &\leq C_1(C_1 + C_2)^\alpha\frac{\epsilon^{-1}}{K^{3-4\alpha-2/p_2}} \leq \frac{C_1 + C_2}{100\epsilon},\end{aligned}\tag{6.5}$$

for sufficiently large C_0 and K , where we used $\epsilon^{3-4\alpha-2/p_2}M_0^2 = 1/K^{3-4\alpha-2/p_2}$ since $a = 2/(3 - 2/p_2 - 4\alpha)$ in the second last inequality. In the last inequality, noting that $3 - 4\alpha - 2/p_2 > 0$ by condition (6.1), we have $1/K^{3-4\alpha-2/p_2} < 1/100$ for sufficiently large K . We proceed similarly as in (6.5) for every term involving $K_0(\tau)$ on the right-hand side of (6.4) to obtain

$$\bar{Q}(\tau_1) \leq \bar{Q}(\tau_0) + \frac{C_1 + C_2}{4\epsilon} < \frac{C_1 + C_2}{\epsilon}$$

with C_0 and K sufficiently large. Note that the condition $p_2 > 2/(1 - \alpha)$ is needed in the tenth term on the right side of (6.4). This is a contradiction, showing that

$$\bar{Q}(\tau) \leq \frac{C_1 + C_2}{\epsilon}.$$

Therefore, $\bar{Q}(\tau) \lesssim M_0^\alpha$, as desired. The case $v = 0$ and $w \in L_t^{q_2}L_x^{q_1}$ is similar with M_0 , α , and K_0 being replaced by M_1 , β , and K_1 respectively. The general case, when both $v \in L_t^{p_2}L_x^{p_1}$ and $w \in L_t^{q_2}L_x^{q_1}$ are present, follows analogously. \square

Acknowledgments

The authors were supported in part by the NSF grant DMS-1907992.

References

- [A] S. Agmon, *Unicité et convexité dans les problèmes différentiels*, Séminaire de Mathématiques Supérieures, No. 13 (Été, 1965), Les Presses de l'Université de Montréal, Montreal, Que., 1966.
- [AN] S. Agmon and L. Nirenberg, *Lower bounds and uniqueness theorems for solutions of differential equations in a Hilbert space*, Comm. Pure Appl. Math. **20** (1967), 207–229.
- [AE] G. Alessandrini and L. Escauriaza, *Null-controllability of one-dimensional parabolic equations*, ESAIM Control Optim. Calc. Var. **14** (2008), no. 2, 284–293.
- [AMRV] G. Alessandrini, A. Morassi, E. Rosset, and S. Vessella, *On doubling inequalities for elliptic systems*, J. Math. Anal. Appl. **357** (2009), no. 2, 349–355.
- [AI] F.J. Almgren, Jr., *Dirichlet's problem for multiple valued functions and the regularity of mass minimizing integral currents*, Minimal submanifolds and geodesics (Proc. Japan-United States Sem., Tokyo, 1977), North-Holland, Amsterdam-New York, 1979, pp. 1–6.
- [An] S. Angenent, *The zero set of a solution of a parabolic equation*, J. Reine Angew. Math. **390** (1988), 79–96.

- [AV] G. Alessandrini and S. Vessella, *Local behaviour of solutions to parabolic equations*, Comm. Partial Differential Equations **13** (1988), no. 9, 1041–1058.
- [BC] L. Bakri and J.-B. Casteras, *Quantitative uniqueness for Schrödinger operator with regular potentials*, Math. Methods Appl. Sci. **37** (2014), no. 13, 1992–2008.
- [B] L. Bakri, *Carleman estimates for the Schrödinger operator. Applications to quantitative uniqueness*, Comm. Partial Differential Equations **38** (2013), no. 1, 69–91.
- [BK] J. Bourgain and C.E. Kenig, *On localization in the continuous Anderson-Bernoulli model in higher dimension*, Invent. Math. **161** (2005), no. 2, 389–426.
- [CRV] B. Canuto, E. Rosset, and S. Vessella, *Quantitative estimates of unique continuation for parabolic equations and inverse initial-boundary value problems with unknown boundaries*, Trans. Amer. Math. Soc. **354** (2002), no. 2, 491–535.
- [C] T. Carleman, *Sur un problème d'unicité pur les systèmes d'équations aux dérivées partielles à deux variables indépendantes*, Ark. Mat., Astr. Fys. **26** (1939), no. 17, 9.
- [Ch] X.-Y. Chen, *A strong unique continuation theorem for parabolic equations*, Math. Ann. **311** (1998), no. 4, 603–630.
- [CFNT] P. Constantin, C. Foias, B. Nicolaenko, and R. Temam, *Spectral barriers and inertial manifolds for dissipative partial differential equations*, J. Dynam. Differential Equations **1** (1989), no. 1, 45–73.
- [CK] G. Camliyurt and I. Kukavica, *Quantitative unique continuation for a parabolic equation*, Indiana Univ. Math. J. **67** (2018), no. 2, 657–678.
- [CKW1] G. Camliyurt, I. Kukavica, and F. Wang, *On quantitative uniqueness for elliptic equations*, Math. Z. **291** (2019), no. 1-2, 227–244.
- [CKW2] G. Camliyurt, I. Kukavica, and F. Wang, *On localization and quantitative uniqueness for elliptic partial differential equations*, Partial differential equations in fluid mechanics, London Math. Soc. Lecture Note Ser., vol. 452, Cambridge Univ. Press, Cambridge, 2018, pp. 68–96.
- [DZ1] B. Davey and J. Zhu, *Quantitative uniqueness of solutions to second order elliptic equations with singular potentials in two dimensions*, Calc. Var. Partial Differential Equations **57** (2018), no. 3, Paper No. 92, 27.
- [DZ2] B. Davey and J. Zhu, *Quantitative uniqueness of solutions to second-order elliptic equations with singular lower order terms*, Comm. Partial Differential Equations **44** (2019), no. 11, 1217–1251.
- [DF1] H. Donnelly and C. Fefferman, *Nodal sets of eigenfunctions on Riemannian manifolds*, Invent. Math. **93** (1988), no. 1, 161–183.
- [DF2] H. Donnelly and C. Fefferman, *Nodal sets for eigenfunctions of the Laplacian on surfaces*, J. Amer. Math. Soc. **3** (1990), no. 2, 333–353.
- [EF] L. Escauriaza and F.J. Fernández, *Unique continuation for parabolic operators*, Ark. Mat. **41** (2003), no. 1, 35–60.
- [E] L. Escauriaza, *Carleman inequalities and the heat operator*, Duke Math. J. **104** (2000), no. 1, 113–127.
- [EFV] L. Escauriaza, F. J. Fernández, and S. Vessella, *Doubling properties of caloric functions*, Appl. Anal. **85** (2006), no. 1-3, 205–223.
- [EV] L. Escauriaza and L. Vega, *Carleman inequalities and the heat operator. II*, Indiana Univ. Math. J. **50** (2001), no. 3, 1149–1169.
- [EVe] L. Escauriaza and S. Vessella, *Optimal three cylinder inequalities for solutions to parabolic equations with Lipschitz leading coefficients*, Inverse problems: theory and applications (Cortona/Pisa, 2002), Contemp. Math., vol. 333, Amer. Math. Soc., Providence, RI, 2003, pp. 79–87.
- [FS] C. Foias and J.-C. Saut, *Asymptotic behavior, as $t \rightarrow \infty$, of solutions of the Navier-Stokes equations*, Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. IV (Paris, 1981/1982), Res. Notes in Math., vol. 84, Pitman, Boston, MA, 1983, pp. 74–86.
- [GL] N. Garofalo and F.-H. Lin, *Monotonicity properties of variational integrals, A_p weights and unique continuation*, Indiana Univ. Math. J. **35** (1986), no. 2, 245–268.
- [H1] Q. Han, *Schauder estimates for elliptic operators with applications to nodal sets*, J. Geom. Anal. **10** (2000), no. 3, 455–480.
- [H2] Q. Han, *On the Schauder estimates of solutions to parabolic equations*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **27** (1998), no. 1, 1–26 (1999).
- [JK] D. Jerison and C.E. Kenig, *Unique continuation and absence of positive eigenvalues for Schrödinger operators*, Ann. of Math. (2) **121** (1985), no. 3, 463–494, With an appendix by E. M. Stein.
- [K1] C.E. Kenig, *Restriction theorems, Carleman estimates, uniform Sobolev inequalities and unique continuation*, Harmonic analysis and partial differential equations (El Escorial, 1987), Lecture Notes in Math., vol. 1384, Springer, Berlin, 1989, pp. 69–90.
- [K2] C.E. Kenig, *Some recent applications of unique continuation*, Recent developments in nonlinear partial differential equations, Contemp. Math., vol. 439, Amer. Math. Soc., Providence, RI, 2007, pp. 25–56.
- [K3] C.E. Kenig, *Quantitative unique continuation, logarithmic convexity of Gaussian means and Hardy's uncertainty principle*, Perspectives in partial differential equations, harmonic analysis and applications, Proc. Sympos. Pure Math., vol. 79, Amer. Math. Soc., Providence, RI, 2008, pp. 207–227.

- [KSW] C. Kenig, L. Silvestre, and J.-N. Wang, *On Landis' conjecture in the plane*, Comm. Partial Differential Equations **40** (2015), no. 4, 766–789.
- [Ku1] I. Kukavica, *Quantitative uniqueness for second-order elliptic operators*, Duke Math. J. **91** (1998), no. 2, 225–240.
- [Ku2] I. Kukavica, *Quantitative uniqueness and vortex degree estimates for solutions of the Ginzburg-Landau equation*, Electron. J. Differential Equations (2000), No. 61, 15 pp. (electronic).
- [Ku3] I. Kukavica, *Self-similar variables and the complex Ginzburg-Landau equation*, Comm. Partial Differential Equations **24** (1999), no. 3-4, 545–562.
- [Ku4] I. Kukavica, *Length of vorticity nodal sets for solutions of the 2D Navier-Stokes equations*, Comm. Partial Differential Equations **28** (2003), no. 3-4, 771–793.
- [Kur] K. Kurata, *On a backward estimate for solutions of parabolic differential equations and its application to unique continuation*, Spectral and scattering theory and applications, Adv. Stud. Pure Math., vol. 23, Math. Soc. Japan, Tokyo, 1994, pp. 247–257.
- [KT1] H. Koch and D. Tataru, *Carleman estimates and unique continuation for second-order elliptic equations with nonsmooth coefficients*, Comm. Pure Appl. Math. **54** (2001), no. 3, 339–360.
- [KT2] H. Koch and D. Tataru, *Carleman estimates and unique continuation for second order parabolic equations with nonsmooth coefficients*, Comm. Partial Differential Equations **34** (2009), no. 4-6, 305–366.
- [L] F.-H. Lin, *Nodal sets of solutions of elliptic and parabolic equations*, Comm. Pure Appl. Math. **44** (1991), no. 3, 287–308.
- [M] V.Z. Meshkov, *On the possible rate of decrease at infinity of the solutions of second-order partial differential equations*, Mat. Sb. **182** (1991), no. 3, 364–383.
- [P] C.-C. Poon, *Unique continuation for parabolic equations*, Comm. Partial Differential Equations **21** (1996), no. 3-4, 521–539.
- [V] S. Vessella, *Carleman estimates, optimal three cylinder inequalities and unique continuation properties for parabolic operators*, Progress in analysis, Vol. I, II (Berlin, 2001), World Sci. Publ., River Edge, NJ, 2003, pp. 485–492.
- [Z] J. Zhu, *Quantitative uniqueness of solutions to parabolic equations*, J. Funct. Anal. **275** (2018), no. 9, 2373–2403.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089
Email address: kukavica@usc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089
Email address: ntle@usc.edu