

ON CATEGORY OF LIE ALGEBRAS

P. G. ROMEO AND ARJUN. S. N

ABSTRACT. In this paper we describe the the category of Lie algebras of group algebras and the category of Plesken Lie algebras and explore the categorical relations between them. Further we provide the examples of the Lie algebra of the group algebra of subgroups of Heisenberg group and the Plesken Lie algebra of subgroups of Heisenberg group.

1. INTRODUCTION

The concept of Lie algebras was introduced by Sophus Lie to solve problems in Lie groups with some ease. Here we introduce some class of Lie algebras like Lie algebras of group algebras and Plesken Lie algebras of groups. In recent times category theory established itself as a practical tool in dealing with mathematical structures. In particular a categorical approach enables to extract more insight into the interactions between Lie groups and Lie algebras. Here we discuss the category of some class of Lie algebras, functorial relation that exists between them with some examples.

2. PRELIMINARIES

In the following we briefly recall all basic definitions and the elementary concepts needed in the sequel. In particular we recall the definitions of Lie algebra, group algebra, Plesken Lie algebra, categories, functors and discusses some interesting properties of these structures.

Definition 1. (cf.[5]) *A category \mathcal{C} consists of the following data:*

- (1) *A class called the class of vertices or objects $\nu\mathcal{C}$.*
- (2) *A class of disjoint sets $\mathcal{C}(a, b)$ one for each pair $(a, b) \in \nu\mathcal{C} \times \nu\mathcal{C}$. An element $f \in \mathcal{C}$ is called a morphism from a to b , written $f : a \rightarrow b$; $a = \text{dom } f$ called the domain of f and $b = \text{cod } f$ called the codomain of f .*

2020 *Mathematics Subject Classification.* 20M10.

Key words and phrases. Lie algebra, Plesken Lie algebra, Group algebra, Category, Functor.

- (3) For $a, b, c, \in \nu\mathcal{C}$, a map
 $\circ : \mathcal{C}(a, b) \times \mathcal{C}(b, c) \rightarrow \mathcal{C}(a, c) \quad (f, g) \rightarrow g \circ f$
 \circ is called the composition of morphisms in \mathcal{C} .
- (4) for each $a \in \nu\mathcal{C}$, a unique $1_a \in \mathcal{C}(a, a)$ is called the identity morphism on a .

These must satisfy the following axioms :

- (cat1) The composition is associative : for $f \in \mathcal{C}(a, b), g \in \mathcal{C}(b, c)$ and $h \in \mathcal{C}(c, d)$, we have

$$f \circ (g \circ h) = (f \circ g) \circ h$$

- (cat 2) for each $a \in \nu\mathcal{C}, f \in \mathcal{C}(a, b)$ and $g \in \mathcal{C}(c, a)$,

$$1_a \circ f = f \quad \text{and} \quad g \circ 1_a = g$$

Clearly $\nu\mathcal{C}$ can be identify as a subclass of \mathcal{C} and with this identification it is possible to regard categories in terms of morphisms alone. The category \mathcal{C} is said to be small if the class \mathcal{C} is a set. A morphism $f \in \mathcal{C}(a, b)$ is said to be an isomorphism if there exists $f^{-1} \in \mathcal{C}(b, a)$ such that $ff^{-1} = 1_a = e_a$, domain identity and $f^{-1}f = 1_b = e_b$, range identity.

Example 1. A group G can be regarded as a category \mathcal{C} with the object set of \mathcal{C} say $\nu\mathcal{C} = G$, and morphisms $\mathcal{C}(G, G) = G$ and composition in \mathcal{C} is the binary operation in G . Identity element in the group will be the identity morphism on the vertex G .

Definition 2. [1] For categories \mathcal{C} and \mathcal{D} a functor $T : \mathcal{C} \rightarrow \mathcal{D}$ with domain \mathcal{C} and codomain \mathcal{D} consists of two functions: the object function T , which assigns to each object c of \mathcal{C} an object Tc of \mathcal{D} and the arrow function which assigns to each arrow $f : c \rightarrow c'$ of \mathcal{C} an arrow $Tf : Tc \rightarrow Tc'$ of \mathcal{D} , in such a way that $T(1_c) = 1_{Tc}$ and $T(g \circ f) = Tg \circ Tf$.

Definition 3. [3] A vector space L over a field \mathbb{F} , with an operation $L \times L \rightarrow L$, denoted by $(x, y) \mapsto [x, y]$ for x and y in L and satisfying the following axioms:

- (1) The bracket operation is bilinear. For $x, y, z \in L, a, b \in \mathbb{F}$

$$\begin{aligned} [ax + by, z] &= a[x, z] + b[y, z] \\ [x, ay + bz] &= a[x, y] + b[x, z] \end{aligned}$$

- (2) $[x, x] = 0$ for all $x \in L$

- (3) Jacobi identity :

$$[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0 \text{ for all } x, y, z \in L$$

is called a bracket product and $(L, [., .])$ is called a Lie algebra over \mathbb{F} .

Example 2. $End(V)$, the set of all linear transformations on a finite dimensional vector space V over a field \mathbb{F} is a Lie algebra with Lie bracket $[x, y] = xy - yx$ for $x, y \in End(V)$.

A subspace K of a Lie algebra L is called a Lie subalgebra if $[x, y] \in K$ whenever $x, y \in K$.

Definition 4. Let G be a group and let \mathbb{F} be \mathbb{R} or \mathbb{C} . Define a vector space over \mathbb{F} with elements of G as a basis, and denote it by $\mathbb{F}G$. That is; $\mathbb{F}G = \{ \sum_{i=0}^n a_i g_i : a_i \in \mathbb{F} \text{ for all } i \}$. The addition and scalar multiplication in $\mathbb{F}G$ are defined by; for

$$\alpha = \sum_{i=0}^n a_i g_i \text{ and } \beta = \sum_{i=0}^n b_i g_i$$

in $\mathbb{F}G$ and $k \in \mathbb{F}$,

$$\alpha + \beta = \sum_i (a_i + b_i) g_i \text{ and } k\alpha = \sum_i (ka_i) g_i$$

The vector space $\mathbb{F}G$, with multiplication defined by

$$\left(\sum_i a_i g_i \right) \left(\sum_j b_j g_j \right) = \sum_{i,j} a_i b_j (g_i g_j), \quad a_i, b_j \in \mathbb{F}$$

is called the group algebra of G over \mathbb{F} .

The group algebra of a finite group G is a vector space of dimension $|G|$ which also carries extra structure involving the the product operation on G .

Example 3. $G = C_3 = \langle a : a^3 = e \rangle$ where e is the identity in G . Then

$$\mathbb{C}G = \{ \lambda_1 e + \lambda_2 a + \lambda_3 a^2 : \lambda_i \in \mathbb{C} \text{ for } i = 1, 2, 3 \}$$

$\mathbb{C}G$ is a group algebra with usual addition and multiplication of series.

3. LIE ALGEBRAS OF GROUP ALGEBRAS AND PLESKEN LIE ALGEBRAS

Let $\mathbb{F}G$ be a group algebra over \mathbb{F} . Then $\mathbb{F}G$ can be regarded as a Lie algebra by defining the Lie bracket $[., .] : \mathbb{F}G \times \mathbb{F}G \rightarrow \mathbb{F}G$ as follows :

$$[\alpha, \beta] = \alpha\beta - \beta\alpha \text{ for } \alpha = \sum_i a_i g_i \text{ and } \beta = \sum_i b_i g_i \text{ in } \mathbb{F}G$$

Clearly $\mathbb{F}G$ is a Lie algebra with respect to the given Lie bracket and is called the Lie algebra of the group algebra $\mathbb{F}G$ and we denote it by $L_{\mathbb{F}G}$.

A linear map between $L_{\mathbb{F}G}$ and $L_{\mathbb{F}H}$ which preserves the Lie bracket is a *homomorphism* of Lie algebras of group algebras.

Proposition 1. *Let $f : G \rightarrow H$ be a group homomorphism. Then $\bar{f} : L_{\mathbb{F}G} \rightarrow L_{\mathbb{F}H}$ defined by*

$$\bar{f}(\alpha) = \bar{f}\left(\sum_i a_i g_i\right) = \sum_i a_i f(g_i)$$

is a homomorphism between Lie algebras of group algebras.

Proof. Let $G = \{g_1, g_2, g_3, \dots\}$ and $L_{\mathbb{F}G} = \{\sum_i a_i g_i : a_i \in \mathbb{F} \text{ for all } i\}$.

Then for $\alpha = \sum_i a_i g_i, \beta = \sum_j b_j g_j \in L_{\mathbb{F}G}$,

$$\begin{aligned} \bar{f}([\alpha, \beta]) &= \bar{f}(\alpha\beta - \beta\alpha) \\ &= \bar{f}\left(\sum_i a_i g_i \sum_j b_j g_j - \sum_j b_j g_j \sum_i a_i g_i\right) \\ &= \bar{f}\left(\sum_{i,j} a_i b_j g_i g_j - \sum_{j,i} b_j a_i g_j g_i\right) \\ &= \sum_{i,j} a_i b_j f(g_i g_j) - \sum_{j,i} b_j a_i f(g_j g_i) \\ &= \sum_{i,j} a_i b_j f(g_i) f(g_j) - \sum_{j,i} b_j a_i f(g_j) f(g_i) \\ &= \sum_i a_i f(g_i) \sum_j b_j f(g_j) - \sum_j b_j f(g_j) \sum_i a_i f(g_i) \\ &= [\bar{f}(\alpha), \bar{f}(\beta)] \end{aligned}$$

Hence, \bar{f} is a homomorphism between Lie algebras of group algebras. \square

Next we proceed to describe Plesken Lie algebra. Let G be a group and $\mathbb{F}G$ be its group algebra over \mathbb{F} , then for each $g \in G$, $g - g^{-1} \in \mathbb{F}G$, denote it by \hat{g} , then the linear span of \hat{g} admits a Lie algebra structure as explained below.

Definition 5. [2] *Plesken Lie algebra $\mathcal{L}(G)$ of a group G over \mathbb{F} is the linear span of elements $\hat{g} \in \mathbb{F}G$ together with the Lie bracket*

$$[\hat{g}, \hat{h}] = \hat{g}\hat{h} - \hat{h}\hat{g}$$

That is, for any group $G = \{g_1, g_2, g_3, \dots\}$, $\{\sum_i a_i \hat{g}_i : a_i \in \mathbb{F} \text{ for all } i\}$ together with the Lie bracket defined above is the Plesken Lie algebra $\mathcal{L}(G)$.

Lemma 1. *The Plesken Lie algebra $\mathcal{L}(G)$ over \mathbb{F} is a Lie subalgebra of the Lie algebra $L_{\mathbb{F}G}$.*

Proof. Let $G = \{g_1, g_2, g_3, \dots\}$ be a group. Then the Lie algebra of the group algebra $\mathbb{F}G$ is $L_{\mathbb{F}G} = \{\sum_i \lambda_i g_i : \lambda_i \in \mathbb{F} \text{ for all } i\}$ and the Plesken

Lie algebra is $\mathcal{L}(G) = \{\sum_i a_i \hat{g}_i : a_i \in \mathbb{F} \text{ for all } i\}$. Since $\mathcal{L}(G)$ is the linear span of \hat{g} and $\hat{g} \in L_{\mathbb{F}G}$, $\mathcal{L}(G)$ is a subset of $L_{\mathbb{F}G}$.

Let $\hat{\alpha} = \sum_i a_i \hat{g}_i$ and $\hat{\beta} = \sum_i b_i \hat{g}_i$ in $\mathcal{L}(G)$,

$$\begin{aligned} \hat{\alpha} + \hat{\beta} &= \sum_i a_i \hat{g}_i + \sum_i b_i \hat{g}_i \\ &= \sum_i (a_i + b_i) \hat{g}_i \in \mathcal{L}(G) \end{aligned}$$

and $k(\hat{\alpha}) = k \sum_i a_i \hat{g}_i = \sum_i (ka_i) \hat{g}_i \in \mathcal{L}(G)$.

Thus, $\mathcal{L}(G)$ is a subspace of $L_{\mathbb{F}G}$.

Let $\hat{g}, \hat{h} \in \mathcal{L}(G)$, then

$$\begin{aligned} [\hat{g}, \hat{h}] &= \hat{g}\hat{h} - \hat{h}\hat{g} \\ &= (g - g^{-1})(h - h^{-1}) - (h - h^{-1})(g - g^{-1}) \\ &= \widehat{gh} - \widehat{gh^{-1}} - \widehat{g^{-1}h} + \widehat{g^{-1}h^{-1}} \end{aligned}$$

Thus Lie bracket is closed in $\mathcal{L}(G)$. Hence, $\mathcal{L}(G)$ is a Lie subalgebra of $L_{\mathbb{F}G}$. \square

Example 4. *Consider the symmetric group S_3 , then*

$$\begin{aligned} L(S_3) &= \text{span}\{\sigma - \sigma^{-1} : \sigma \in S_3\} \\ &= \{a_1((1) - (1)) + a_2((1\ 2) - (1\ 2)) + a_3((1\ 3) - (1\ 3)) + a_4((2\ 3) - (2\ 3)) \\ &\quad + a_5((1\ 2\ 3) - (1\ 3\ 2)) + a_6((1\ 3\ 2) - (1\ 2\ 3)) : a_i \in \mathbb{C}\} \\ &= \{a((1\ 2\ 3) - (1\ 3\ 2)) : a \in \mathbb{C}\} \end{aligned}$$

is a one dimensional Plesken Lie algebra over \mathbb{C} with Lie bracket

$$[a(\widehat{1\ 2\ 3}), b(\widehat{1\ 2\ 3})] = 0$$

A linear map between two Plesken Lie algebras $\mathcal{L}(G)$ and $\mathcal{L}(H)$ is a *Plesken Lie algebra homomorphism* if it preserves the Lie bracket.

Proposition 2. *Let $f : G \rightarrow H$ be a group homomorphism. Then $\hat{f} : \mathcal{L}(G) \rightarrow \mathcal{L}(H)$ defined by*

$$\hat{f}\left(\sum_i a_i \hat{g}_i\right) = \sum_i a_i \widehat{f(g_i)}$$

is a Plesken Lie algebra homomorphism.

Proof. For $\sum_i a_i \hat{g}_i, \sum_i b_i \hat{g}_i \in \mathcal{L}(G)$ and $k \in \mathbb{F}$,

$$\begin{aligned} \hat{f}\left(\sum_i a_i \hat{g}_i + k \sum_i b_i \hat{g}_i\right) &= \hat{f}\left(\sum_i (a_i + kb_i) \hat{g}_i\right) \\ &= \sum_i (a_i + kb_i) \widehat{f(g_i)} \\ &= \hat{f}\left(\sum_i a_i \hat{g}_i\right) + k \hat{f}\left(\sum_i b_i \hat{g}_i\right) \end{aligned}$$

thus \hat{f} is linear.

For $\sum_i a_i \hat{g}_i, \sum_j b_j \hat{g}_j \in \mathcal{L}(G)$,

$$\begin{aligned} \hat{f}\left([\sum_i a_i \hat{g}_i, \sum_j b_j \hat{g}_j]\right) &= \hat{f}\left(\sum_{i,j} (a_i b_j - b_i a_j) \widehat{g_i g_j} - \sum_{i,j} (a_i b_j - b_i a_j) \widehat{g_i g_j^{-1}}\right. \\ &\quad \left. - \sum_{i,j} (a_i b_j - b_i a_j) \widehat{g_i^{-1} g_j} + \sum_{i,j} (a_i b_j - b_i a_j) \widehat{g_i^{-1} g_j^{-1}}\right) \\ &= \sum_{i,j} (a_i b_j - b_i a_j) f(\widehat{g_i}) f(\widehat{g_j}) - \sum_{i,j} (a_i b_j - b_i a_j) f(\widehat{g_i}) f(\widehat{g_j^{-1}}) \\ &\quad - \sum_{i,j} (a_i b_j - b_i a_j) f(\widehat{g_i^{-1}}) f(\widehat{g_j}) + \sum_{i,j} (a_i b_j - b_i a_j) f(\widehat{g_i^{-1}}) f(\widehat{g_j^{-1}}) \\ &= \left[\sum_i a_i \widehat{f(g_i)}, \sum_j b_j \widehat{f(g_j)}\right] \end{aligned}$$

Hence \hat{f} preserves Lie bracket and is a Plesken Lie algebra homomorphism. \square

4. CATEGORY OF LIE ALGEBRAS

Here we describe the category of Lie algebras over \mathbb{F} whose objects are Lie algebras over \mathbb{F} and morphisms Lie algebra homomorphisms. Let L and L' be two Lie algebras over a field \mathbb{F} . If f and g are two morphisms, then $f \circ g$ exists only when $f \in \text{hom}(L, L')$ and $g \in \text{hom}(L', L'')$. Also the identity in $\text{hom}(L, L)$ is the morphism 1_L .

Example 5. Consider a group G and all its subgroups H_i . The group algebras $\mathbb{F}H_i$ of each subgroup H_i of G together with a Lie bracket

$[x, y] = xy - yx$ for $x = \sum_i a_i h_i, y = \sum_j b_j h_j \in \mathbb{F}H_i$ is the Lie algebra $L_{\mathbb{F}H_i}$. The collection of all such Lie algebras of group algebras of subgroups of G form the category $L_{\mathbb{F}G}$ whose morphisms are $\{ \bar{f}_{ij} : L_{\mathbb{F}H_i} \rightarrow L_{\mathbb{F}H_j} \mid \bar{f}(\sum_i a_i g_i) = \sum_i a_i f(g_i) \text{ where } f : H_i \rightarrow H_j \text{ is the group homomorphism} \}$.

Example 6. Consider the Heisenberg group $H(\mathbb{R}) = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in \mathbb{R} \right\}$ and its non-isomorphic subgroups :

$$H_1 = \{ I_{3 \times 3}, \text{ the identity matrix} \}, H_2 = \left\{ \begin{pmatrix} 1 & 0 & b \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} : b \in \mathbb{R} \right\}$$

$$H_3 = \left\{ \begin{pmatrix} 1 & 0 & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} : b, c \in \mathbb{R} \right\}, H_4 = H(\mathbb{R})$$

The group algebras $\mathbb{F}H_i$ over \mathbb{F} of each subgroups H_i of $H(\mathbb{R})$ together with a Lie bracket $[X, Y] = XY - YX$ where $X = \sum_i \lambda_i A_i, Y = \sum_i \mu_i B_i \in \mathbb{F}H_i$ is the Lie algebra $L_{\mathbb{F}H_i}$ of the group algebra $\mathbb{F}H_i$. Consider $L_{\mathbb{F}H(\mathbb{R})}$ whose objects are

$L_{\mathbb{F}H_1} = \{ \lambda I : \lambda \in \mathbb{F} \} = \text{the Lie algebra of scalar matrices}$

$$L_{\mathbb{F}H_2} = \left\{ \sum_i \lambda_i A_i : A_i \in H_2, \lambda_i \in \mathbb{F} \right\} = \left\{ \begin{pmatrix} \sum_i \lambda_i & 0 & \sum_i \lambda_i b_i \\ 0 & \sum_i \lambda_i & 0 \\ 0 & 0 & \sum_i \lambda_i \end{pmatrix} \right\}$$

$$L_{\mathbb{F}H_3} = \left\{ \sum_i \lambda_i A_i : A_i \in H_3, \lambda_i \in \mathbb{F} \right\} = \left\{ \begin{pmatrix} \sum_i \lambda_i & 0 & \sum_i \lambda_i b_i \\ 0 & \sum_i \lambda_i & \sum_i \lambda_i c_i \\ 0 & 0 & \sum_i \lambda_i \end{pmatrix} \right\}$$

$$L_{\mathbb{F}H_4} = \left\{ \sum_i \lambda_i A_i : A_i \in H_4, \lambda_i \in \mathbb{F} \right\} = \left\{ \begin{pmatrix} \sum_i \lambda_i & \sum_i \lambda_i a_i & \sum_i \lambda_i b_i \\ 0 & \sum_i \lambda_i & \sum_i \lambda_i c_i \\ 0 & 0 & \sum_i \lambda_i \end{pmatrix} \right\}$$

and morphisms : $\text{hom}(L_{\mathbb{F}H_i}, L_{\mathbb{F}H_j}) = \{ \bar{f}_{ij} : L_{\mathbb{F}H_i} \rightarrow L_{\mathbb{F}H_j} \mid \bar{f}(\sum_i a_i g_i) = \sum_i a_i f(g_i) \text{ where } f : H_i \rightarrow H_j \text{ is the group homomorphism} \}$ is the category $L_{\mathbb{F}(H(\mathbb{R}))}$ of Lie algebras of the group algebras of subgroups of $H(\mathbb{R})$.

Next we consider the category whose objects are Plesken Lie algebras $\mathcal{L}(G)$ over a field \mathbb{F} and morphisms are Plesken Lie algebra homomorphisms which we denote it by \mathcal{C}_{PLG} .

Example 7. Consider the Heisenberg group $H(\mathbb{R}) = \left\{ \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} : a, b, c \in \mathbb{R} \right\}$ and its non-isomorphic subgroups H_1, H_2, H_3 and H_4 as given in Example.6. For each $A = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} \in H_i$,

$$\hat{A} = A - A^{-1} = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix} - \begin{pmatrix} 1 & -a & ac - b \\ 0 & 1 & -c \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 2a & 2b - ac \\ 0 & 0 & 2c \\ 0 & 0 & 0 \end{pmatrix}$$

Thus $\mathcal{L}(H_i) = \left\{ \sum_i \lambda_i \hat{A}_i : A_i \in H_i, \lambda_i \in \mathbb{F} \right\}$ and so

$$\mathcal{L}(H_1) = \{0\}, \quad \mathcal{L}(H_2) = \left\{ \begin{pmatrix} 0 & 0 & 2 \sum_i \lambda_i b_i \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\}$$

$$\mathcal{L}(H_3) = \left\{ \begin{pmatrix} 0 & 0 & 2 \sum_i \lambda_i b_i \\ 0 & 0 & 2 \sum_i \lambda_i c_i \\ 0 & 0 & 0 \end{pmatrix} \right\}, \quad \mathcal{L}(H_4) = \left\{ \begin{pmatrix} 0 & 2 \sum_i \lambda_i a_i & \sum_i \lambda_i (2b_i - a_i c_i) \\ 0 & 0 & 2 \sum_i \lambda_i c_i \\ 0 & 0 & 0 \end{pmatrix} \right\}$$

The category whose objects are $\mathcal{L}(H_i), i = 1, 2, 3, 4$ and morphisms are $\text{hom}(\mathcal{L}(H_i), \mathcal{L}(H_j)) = \{ \hat{f} : \mathcal{L}(H_i) \rightarrow \mathcal{L}(H_j) \mid \hat{f}(\sum_i \lambda_i \hat{A}_i) = \sum_i \lambda_i \widehat{f(A_i)} \}$

where $f : H_i \rightarrow H_j$ is the group homomorphism } is the category of Plesken Lie algebras of subgroups of $H(\mathbb{R})$.

Theorem 1. *If $L_{\mathbb{F}G}$ is the category of Lie algebras of group algebras and \mathcal{C}_{PLG} is the category of Plesken Lie algebras, then there exists a functor from $L_{\mathbb{F}G}$ to \mathcal{C}_{PLG} .*

Proof. Define $T : L_{\mathbb{F}G} \rightarrow \mathcal{C}_{PLG}$ such that $\nu T : L_{\mathbb{F}G_i} \rightarrow \mathcal{L}(G_i)$ is defined by

$$\nu T\left(\sum_i a_i g_i\right) = \sum_i (a_i - a'_i)(g_i - g_i^{-1})$$

and on morphisms, for a group homomorphism $f : G_i \rightarrow G_j$, $\bar{f} : L_{\mathbb{F}G_i} \rightarrow L_{\mathbb{F}G_j}$ is given by

$$\bar{f}\left(\sum_i a_i g_i\right) = \sum_i a_i f(g_i)$$

and $T\bar{f} : T(L_{\mathbb{F}G_i}) \rightarrow T(L_{\mathbb{F}G_j})$ is defined by

$$T\bar{f} = \hat{f}$$

and $\hat{f} : \mathcal{L}(G_i) \rightarrow \mathcal{L}(G_j)$ is given by

$$\hat{f}\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) = \sum_i (a_i - a'_i)(f(g_i) - f(g_i)^{-1})$$

where \hat{f} is a Plesken Lie algebra homomorphism (Proposition 2).

Let $\bar{f}_1 : L_{\mathbb{F}G_i} \rightarrow L_{\mathbb{F}G_j}$ and $\bar{f}_2 : L_{\mathbb{F}G_j} \rightarrow L_{\mathbb{F}G_k}$ be two homomorphisms between Lie algebras of group algebras which are induced from the group homomorphisms $f_1 : G_i \rightarrow G_j$ and $f_2 : G_j \rightarrow G_k$. Then $T(\bar{f}_1) : T(L_{\mathbb{F}G_i}) \rightarrow T(L_{\mathbb{F}G_j})$ and $T(\bar{f}_2) : T(L_{\mathbb{F}G_j}) \rightarrow T(L_{\mathbb{F}G_k})$ are Plesken Lie algebra homomorphisms. Then their composition $T(\bar{f}_2) \circ T(\bar{f}_1) : T(L_{\mathbb{F}G_i}) \rightarrow T(L_{\mathbb{F}G_k})$ is also a Plesken Lie algebra homomorphism. Moreover,

$$\begin{aligned} (1) \quad (T(\bar{f}_2) \circ T(\bar{f}_1))\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) &= (\hat{f}_2 \circ \hat{f}_1)\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) \\ &= \hat{f}_2\left(\sum_{i=1} (a_i - a'_i)(f_1(g_i) - f_1(g_i)^{-1})\right) \\ &= \sum_i (a_i - a'_i)((f_2 \circ f_1)(g_i) - ((f_2 \circ f_1)(g_i))^{-1}) \end{aligned}$$

and

$$\begin{aligned} (2) \quad T(\bar{f}_2 \circ \bar{f}_1)\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) &= (\widehat{f_2 \circ f_1})\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) \\ &= \sum_i (a_i - a'_i)((f_2 \circ f_1)(g_i) - ((f_2 \circ f_1)(g_i))^{-1}) \end{aligned}$$

(1) and (2) shows that $T(\bar{f}_2 \circ \bar{f}_1) = T(\bar{f}_2) \circ T(\bar{f}_1)$.

The identity Lie algebra homomorphism, $1_{L_{\mathbb{F}G_i}} : L_{\mathbb{F}G_i} \rightarrow L_{\mathbb{F}G_i}$ is induced from the identity group homomorphism $1_{G_i} : G_i \rightarrow G_i$. Then $T(1_{L_{\mathbb{F}G_i}}) : T(L_{\mathbb{F}G_i}) \rightarrow T(L_{\mathbb{F}G_i})$ is a Plesken Lie algebra homomorphism and

$$\begin{aligned} T(1_{L_{\mathbb{F}G_i}})\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) &= \sum_i (a_i - a'_i)(1_{G_i}(g_i) - (1_{G_i}(g_i))^{-1}) \\ &= \sum_i (a_i - a'_i)(g_i - g_i^{-1}) \\ &= 1_{T(L_{\mathbb{F}G_i})}\left(\sum_i (a_i - a'_i)(g_i - g_i^{-1})\right) \end{aligned}$$

That is, $T(1_{L_{\mathbb{F}G_i}}) = 1_{T(L_{\mathbb{F}G_i})}$, hence T is a functor. \square

Corollary 1. *The functor $T : L_{\mathbb{F}G} \rightarrow \mathcal{C}_{PLG}$ in Theorem 1 is a full functor.*

Proof. For the functor $T : L_{\mathbb{F}G} \rightarrow \mathcal{C}_{PLG}$ define a map $T_{L_{\mathbb{F}G_i}, L_{\mathbb{F}G_j}} : \text{hom}(L_{\mathbb{F}G_i}, L_{\mathbb{F}G_j}) \rightarrow \text{hom}(TL_{\mathbb{F}G_i}, TL_{\mathbb{F}G_j})$ by

$$T_{L_{\mathbb{F}G_i}, L_{\mathbb{F}G_j}}(\bar{f}) = \hat{f}$$

Now it is enough to prove that $T_{L_{\mathbb{F}G_i}, L_{\mathbb{F}G_j}}$ is surjective. For $\hat{f} \in \text{hom}(TL_{\mathbb{F}G_i}, TL_{\mathbb{F}G_j})$, a Plesken Lie algebra homomorphism, define $\bar{f} : L_{\mathbb{F}G_i} \rightarrow L_{\mathbb{F}G_j}$ by

$$\bar{f}\left(\sum_i a_i g_i\right) = \sum_i a_i f(g_i)$$

is a homomorphism between Lie algebras of group algebras and $T\bar{f} : TL_{\mathbb{F}G_i} \rightarrow TL_{\mathbb{F}G_j}$ given by,

$$T\bar{f}\left(\sum_i a_i \hat{g}_i\right) = \sum_i a_i \widehat{f(g_i)} = \hat{f}\left(\sum_i a_i \hat{g}_i\right)$$

That is, $T\bar{f} = \hat{f}$. Hence T is full. \square

However, it should be noted that the functor defined above need not be faithful as illustrated in the following example.

Example 8. *Let $K_4 = \{e, a, b, c\}$ be the Klein 4- group and $L_{\mathbb{F}K_4}$ Lie algebra of the group algebra $\mathbb{F}K_4$. Consider the identity map $1_{L_{\mathbb{F}K_4}}$ and $\bar{f} : L_{\mathbb{F}K_4} \rightarrow L_{\mathbb{F}K_4}$ given by*

$$\bar{f}\left(\sum_i a_i g_i\right) = \sum_i a_i f(g_i)$$

where $g_i \in K_4$ and f is the trivial homomorphism. Clearly both $T\bar{f}$ and $T1_{L_{\mathbb{F}K_4}}$ are zeroes for the functor $T : L_{\mathbb{F}G} \rightarrow \mathcal{C}_{PLG}$.

REFERENCES

- [1] Saunders Mac Lane, *Categories for the working Mathematician*, Second edition, Springer, 1997
- [2] Arjeh M. Cohen and D. E. Taylor, *On a Certain Lie algebra Defined by a Finite Group*. The American Mathematical Monthly, pp. 633 - 639, Aug. - Sep., 2007.
- [3] James E. Humphreys, *Introduction to Lie Algebras and Representation Theory*. Springer - Verlag, 1972.
- [4] Brian C. Hall, Lie groups, *Lie Algebras, and Representations An Elementary Introduction*, Springer, 2003
- [5] K.S.S. Nambooripad, *Theory of Regular Semigroups*. Sayahna Foundation, Trivandrum, 2018.
- [6] Gordon James and Martin Liebeck, *Representations and Characters of Groups*. Second Edition, Cambridge University Press, 2001.

DEPT. OF MATHEMATICS, COCHIN UNIVERSITY OF SCIENCE AND TECHNOLOGY, KOCHI, KERALA, INDIA.

Email address: romeo-parackal@yahoo.com, arjunsnmaths1996@gmail.com