

HESSIAN EQUATIONS OF KRYLOV TYPE ON KÄHLER MANIFOLDS

LI CHEN

ABSTRACT. In this paper, we consider a Hessian equation with its structure as a combination of elementary symmetric functions on closed Kähler manifolds. We provide a sufficient and necessary condition for the solvability of this equation, which generalizes the results of Hessian equation and Hessian quotient equation. As a consequence, we can solve a complex Monge-Ampère type equation proposed by Chen in the case that one of the coefficients is negative. The key to our argument is a clever use of the special properties of the Hessian quotient operator $\frac{\sigma_k}{\sigma_{k-1}}$.

1. INTRODUCTION

Let (M, ω) be a closed Kähler manifold of complex dimension $n \geq 2$ and fix a real smooth closed $(1, 1)$ -form χ_0 on M . For any C^2 function $u : M \rightarrow \mathbb{R}$, we can obtain a new real $(1, 1)$ -form

$$\chi_u = \chi_0 + \frac{\sqrt{-1}}{2} \partial \bar{\partial} u.$$

We consider the following Hessian type equation on (M, ω)

$$(1.1) \quad \chi_u^k \wedge \omega^{n-k} = \sum_{l=0}^{k-1} \alpha_l(z) \chi_u^l \wedge \omega^{n-l}, \quad 2 \leq k \leq n,$$

where $\alpha_0(z), \dots, \alpha_{k-1}(z)$ are real smooth functions on M . This equation includes some of the most partial differential equations in complex geometry and analysis.

- For $k = n$, $\alpha_1 = \dots = \alpha_{k-1} = 0$, (1.1) is the complex Monge-Ampère equation $\chi_u^n = \alpha_0(z) \omega^n$ which was famously solved on compact Kähler manifolds by Yau in his resolution of Calabi conjecture [56], and on compact Hermitian manifolds by Tosatti-Weinkove [50] with some earlier work by Cherrier [12], Hanani [26], Guan-Li [22] and Zhang [58].
- For $k = n$, $\alpha_0 = \dots = \alpha_{k-2} = 0$ and the constant α_{n-1} , (1.1) is the $(n, n-1)$ -quotient equation $\chi_u^n = \alpha_{n-1} \chi_u^{n-1} \wedge \omega$ which appears

2010 *Mathematics Subject Classification.* 35J96, 52A20, 53C44.

Key words and phrases. Complex Hessian equations; Cone condition; Kähler manifolds.

in a problem proposed by Donaldson [15] in the setting of moment maps. After some progresses made in [10, 52, 53, 54], it was solved by Song-Weinokove [41] via J -flow.

- For $k = n$ and $\alpha_0, \dots, \alpha_{n-1} \in \mathbb{R}$, (1.1) was proposed by Chen (see Conjecture/Question 4 in [9]) in the study of Mabuchi energy. A special case $\chi_u^n = \alpha_l \chi_u^l \wedge \omega^{n-l}$ ($1 < l < n$) was solved by Fang-Lai-Ma [17] by generalizing the result of Song-Weinokove [41] and non constant α_l on Hermitian manifolds were considered by [23, 24, 43]. Chen's problem was solved by Collins and Székelyhidi [13] for $\alpha_i \geq 0$ ($0 \leq i \leq k-1$) as was conjectured by Fang-Lai-Ma. Moreover, a generalized equation of Chen's problem was studied by Sun [47, 48], Pingali [33, 34, 35] and Phong-Tô [38].
- The Hessian equation $\chi_u^k \wedge \omega^{n-k} = \alpha_0(z)$ and the (k, l) -quotient equation $\chi_u^k \wedge \omega^{n-k} = \alpha_l(z) \chi_u^l \wedge \omega^{n-l}$ for $1 \leq l < k < n$ are also the special cases of (1.1). The first one was solve by Dinew-Kolodziej [14] (combined with the estimate of Hou-Ma-Wu [28]) on Kähler manifolds and by Sun [45] on Hermitian manifolds (see also Zhang [57]). The second one was solved by Székelyhidi [40] for constant α_l and non constant α_l was settled by Sun [45].

All previous results on the equation (1.1) require all coefficients $\alpha_i \geq 0$ for $0 \leq i \leq k-1$. Thus, it is interesting to ask:

Question 1.1. Can we solve the equation (1.1) for some $\alpha_i(z)$ which change sign or are negative? In particular, can we solve Chen's problem for some $\alpha_i < 0$?

In this paper, we make some progress on this direction. To statement our main theorem, we first need to introduce the cone condition.

Definition 1.1. Following [41, 17, 24, 45], we set

$$(1.2) \quad \mathcal{C}_k(\omega) : = \left\{ [\chi] : \exists \bar{\chi} \in [\chi] \cap \Gamma_{k-1}(M), \text{ such that} \right. \\ \left. k\bar{\chi}^{k-1} \wedge \omega^{n-k} > \sum_{l=1}^{k-1} l\alpha_l \bar{\chi}^{l-1} \wedge \omega^{n-l} \right\}.$$

where $\alpha_i(z)$ with $0 \leq i \leq k-1$ are real smooth functions on M and $[\chi] = \{\chi + \frac{\sqrt{-1}}{2} \partial \bar{\partial} u : u \in C^2(M)\}$. If $[\chi] \in \mathcal{C}_k(\omega)$, we say that χ satisfies the cone condition for (1.1).

Next, we make the following assumption on the coefficient functions $\alpha_i(z)$ ($0 \leq i \leq k-1$).

Assumption 1.1. Suppose that $\alpha_0(z), \alpha_1(z), \dots, \alpha_{k-1}(z)$ are real smooth functions on M satisfying

- either $\alpha_l(z) > 0$ or $\alpha_l(z) \equiv 0$ for $0 \leq l \leq k-2$ and all $z \in M$;
- $\sum_{l=0}^{k-2} \alpha_l(z) > 0$ for all $z \in M$;

(iii) $\alpha_l(z) \geq c_{k,l}$ for $0 \leq l \leq k-1$ and

$$(1.3) \quad \int_M \chi_0^k \wedge \omega^{n-k} \leq \sum_{l=0}^{k-1} c_{k,l} \int_M \chi_0^l \wedge \omega^{n-l},$$

where $c_{k,0}, \dots, c_{k,k-1}$ are constants.

Remark 1.1. We make no sign requirement for $\alpha_{k-1}(z)$ in Assumption 1.1.

Based on Lemma 2.2, it is easy to see the condition $[\chi_0] \in \mathcal{C}_k(\omega)$ is necessary for the solvability of (1.1). We show such condition is also sufficient.

Theorem 1.1. *Let (M, ω) be a closed Kähler manifold of complex dimension $n \geq 2$ and χ_0 be a smooth closed real $(1,1)$ -form on M . Assume that $\alpha_0(z), \alpha_1(z), \dots, \alpha_{k-1}(z)$ satisfies Assumption 1.1. Then, there exist a unique (up to a constant) smooth function u and $a \in \mathbb{R}$ satisfies*

$$\chi_u^k \wedge \omega^{n-k} = \sum_{l=0}^{k-2} \alpha_l(z) \chi_u^l \wedge \omega^{n-l} + [\alpha_{k-1}(z) + a] \chi_u^{k-1} \wedge \omega^{n-k+1}$$

provided that $\chi_0 \in \mathcal{C}_k(\omega)$.

In particular, if $\alpha_0, \alpha_1, \dots, \alpha_{k-1}$ are constants, the equation (1.1) was proposed by Chen (see Conjecture/Question 4 in [9]) for $k = n$ and it was solved by Collins and Székelyhidi [13] for $\alpha_i \geq 0$ ($0 \leq i \leq n-1$). As a corollary of Theorem 1.1, we can solve Chen's problem for $\alpha_{k-1} < 0$.

Theorem 1.2. *Let (M, ω) be a closed Kähler manifold of complex dimension $n \geq 2$ and χ_0 be a smooth closed real $(1,1)$ -form on M . Assume $\alpha_{k-1} \in \mathbb{R}$ and $\alpha_0, \alpha_1, \dots, \alpha_{k-2}$ are nonnegative real numbers satisfying $\sum_{i=0}^{k-2} \alpha_i > 0$ and*

$$(1.4) \quad \int_M \chi_0^k \wedge \omega^{n-k} = \sum_{l=0}^{k-2} \alpha_l \int_M \chi_0^l \wedge \omega^{n-l},$$

Then, there exist a unique (up to a constant) smooth function u satisfies (1.1) if and only if $[\chi_0] \in \mathcal{C}_k(\omega)$.

Remark 1.2. Integrating both sides of (1.1) on M , it is easy to see the equality (1.4) is a necessary condition for the solvability of (1.1) for $\alpha_i \in \mathbb{R}$ ($0 \leq i \leq k-1$).

The equation (1.1) is just a Hessian type of equation with its structure as a combination of elementary symmetric functions (see Section 2.1 for details):

$$(1.5) \quad \sigma_k(\chi_u) = \sum_{l=0}^{k-1} \beta_l(z) \sigma_l(\chi_u),$$

where $\beta_l(z) = \frac{C_n^k}{C_n^l} \alpha_l(z) > 0$. Such type of equation has been firstly considered by Krylov about twenty years ago. In [30], he considered the Dirichlet

problem of the following degenerate equation in a $(k-1)$ -convex domain D in \mathbb{R}^n ,

$$\sigma_k(D^2u) = \sum_{l=0}^{k-1} \beta_l(x) \sigma_l(D^2u)$$

with all coefficient $\beta_l(x) \geq 0$ for $0 \leq l \leq k-1$. Recently, Krylov's equation was extended by Guan-Zhang [25] to the case without the sign requirement for the coefficient function $\beta_{k-1}(x)$ and the corresponding Neumann problems of Krylov's equation in real and complex spaces were studied by the author with his collaborators in [5, 6]. Moreover, such type of equations with its structure as a combination of elementary symmetric functions arise naturally from many important geometric problems, such as the so-called Fu-Yau equation arising from the study of the Hull-Strominger system in theoretical physics, see Fu-Yau [19, 18] and Phong-Picard-Zhang [37, 36, 39]. Furthermore, the special Lagrangian equations introduced by Harvey and Lawson [27] can also be written as the alternative combinations of elementary symmetric functions. Lastly, the Krylov type of equations in prescribed curvature problems were also studied by the author with his collaborators in [7, 8].

Compared with Hessian equations or Hessian quotient equations, the ellipticity and concavity of (1.5) is not so easy to see. Moreover, we even have several ways to make (1.5) elliptic and concave. In [13, 47], the authors rewrite the equation (1.5) as the form

$$(1.6) \quad - \sum_{l=0}^{k-1} \beta_l(z) \frac{\sigma_l(\chi_u)}{\sigma_k(\chi_u)} = -1.$$

The disadvantage of the form (1.6) is that the ellipticity and concavity of (1.6) require all $\alpha_i \geq 0$ for $0 \leq i \leq k-1$. In this paper, we follow the viewpoint of Guan-Zhang [25] to rewrite the equation (1.5) as the form

$$(1.7) \quad \frac{\sigma_k(\chi_u)}{\sigma_{k-1}(\chi_u)} - \sum_{l=0}^{k-2} \beta_l(z) \frac{\sigma_l(\chi_u)}{\sigma_{k-1}(\chi_u)} = \beta_{k-1}(z).$$

Then, we have the following very important fact (see Lemma 2.5 for details) which was first observed by Guan-Zhang [25] for Krylov' equation.

Fact 1.1. Assume $\beta_l(z) \geq 0$ for $0 \leq l \leq k-2$ and $\chi_u \in \Gamma_{k-1}(M)$, then the equation (1.6) is elliptic and concave. Compared with the form (1.6), its advantages lie in two aspects: (i) we make no sign requirement for $\beta_{k-1}(z)$ and (ii) the proper admissible set of solutions is Γ_{k-1} which is larger than Γ_k .

Fact 1.1 is based on the special properties of the Hessian quotient operator $\frac{\sigma_k}{\sigma_{k-1}}$ which are key to our argument. In details, Propositions 4 and 5, and Lemma 2.2 declare that if $l = k-1$, some properties of Hessian quotient $\frac{\sigma_k}{\sigma_l}$

hold true in a larger cone Γ_{k-1} . The root of these special properties goes back to the Newton inequality (2.2).

In view of Fact 1.1, using the equation (1.7), we first prove a C^0 estimate, generalizing the approach of [45] in the case of Hessian quotient equations. On the one hand, we improve the inequalities (2.11) and (2.12) in [45] to get the inequalities (3.4) and (3.5) such that they do not depend on the lower bound of α_i ($0 \leq i \leq k-1$). This improvement makes sure Theorem 1.1 even holds true for α_{k-1} with indefinite sign and some $\alpha_i(z) \equiv 0$ ($0 \leq i \leq k-2$). On the other hand, in order to control the integrals of Hessian operators with lower degree in Lemma 3.3, we need to get a new inequality (3.12) by iterating the inequality (3.15) in [45].

Then, we follow a technique of Hou-Ma-Wu [28] in the case of Hessian equation (which is in turn based on ideas developed for the real Hessian equation by Chou-Wang [11]) to derive a second derivative bound of the form

$$(1.8) \quad \sup_M |\partial\bar{\partial}u| \leq C(\sup_M |\nabla u|^2 + 1).$$

To achieve such estimate, the main task is to control a third order derivative term of the form

$$(1.9) \quad \frac{F^{i\bar{i}}|\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2}$$

in the case of Hessian equations or Hessian quotient equations. This term can be controlled with the help of the positive term

$$(1.10) \quad -F^{i\bar{j},r\bar{s}}\chi_{i\bar{j};1}\chi_{r\bar{s};\bar{1}}.$$

However, in the case of the equation (1.7), there are an extra third order derivative term of the form

$$(1.11) \quad \sum_{l=0}^{k-2} \left(\nabla_{\bar{1}}\beta_l \cdot F_l^{i\bar{j}}\chi_{i\bar{j};1} + \nabla_1\beta_l \cdot F_l^{i\bar{j}}\chi_{i\bar{j};\bar{1}} \right).$$

Our first contribution is to extract a part of the term (1.10) to cancel the term (1.11) and meanwhile make sure the rest of the term (1.10) is enough to help us control the (1.9). Thus, the extra third order derivative term makes our argument significantly more delicate. Our another contribution is the inequalities in Lemma 2.8 and Lemma 2.9 which are very important for our C^2 estimate, while similar equalities for the case of Hessian equations or Hessian quotient equations are trivial.

A C^1 -bound is derived by combing a second derivative bound (1.8), with a blow-up argument and Liouville-type theorem due to Dinew-Kolodziej [14]. The gradient bound combined with (1.8) then bound $|\partial\bar{\partial}u|$.

Once we establish the a priori estimates for solutions to (1.1), we can use the continuity method to prove Theorem 1.1. Although the method is very standard in the study of elliptic PDEs, it is not easy to carry out this

method for (1.1) on closed complex manifold. On the one hand, since the essential cone condition depends on $\alpha_0, \dots, \alpha_{k-1}$, we can not obtain a priori estimates for the whole family of solutions in the method of continuity. To overcome this difficulty, we use the idea in [43, 45] to technically set up an intermediate step and apply the method twice. On the other hand, it is very hard to make the inverse function theorem work. To resolve the issue, we further develop the approach in [50]. In this method, a new Hermitian metric is constructed, which is a crucial new technique in implementing the method of continuity.

2. PRELIMINARIES

In this section, we give some basic properties of elementary symmetric functions, which could be found in [31, 42], and establish some key lemmas.

2.1. Basic properties of elementary symmetric functions. For $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$, the k -th elementary symmetric function is defined by

$$\sigma_k(\lambda) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k}.$$

We also set $\sigma_0 = 1$ and denote by $\sigma_k(\lambda|i)$ the k -th symmetric function with $\lambda_i = 0$. Recall that the Gårding's cone is defined as

$$\Gamma_k = \{\lambda \in \mathbb{R}^n : \sigma_i(\lambda) > 0, \forall 1 \leq i \leq k\}.$$

Proposition 1. *Let $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n$ and $k = 1, \dots, n$, then*

$$(2.1) \quad \sigma_k(\lambda) = \sigma_k(\lambda|i) + \lambda_i \sigma_{k-1}(\lambda|i), \quad \forall 1 \leq i \leq n,$$

$$\sum_{i=1}^n \lambda_i \sigma_{k-1}(\lambda|i) = k \sigma_k(\lambda),$$

$$\sum_{i=1}^n \sigma_k(\lambda|i) = (n - k) \sigma_k(\lambda).$$

The Newton inequality and the generalized Newton-MacLaurin inequality are as follows, which will be used later.

Proposition 2. *For $1 \leq k \leq n - 1$ and $\lambda \in \mathbb{R}^n$, we have*

$$(2.2) \quad (n - k + 1)(k + 1) \sigma_{k-1}(\lambda) \sigma_{k+1}(\lambda) \leq k(n - k) \sigma_k^2(\lambda).$$

In particular, we have

$$(2.3) \quad \sigma_{k-1}(\lambda) \sigma_{k+1}(\lambda) \leq \sigma_k^2(\lambda).$$

Proposition 3. *For $\lambda \in \Gamma_k$ and $n \geq k > l \geq 0$, $r > s \geq 0$, $k \geq r$, $l \geq s$, we have*

$$(2.4) \quad \left[\frac{\sigma_k(\lambda)/C_n^k}{\sigma_l(\lambda)/C_n^l} \right]^{\frac{1}{k-l}} \leq \left[\frac{\sigma_r(\lambda)/C_n^r}{\sigma_s(\lambda)/C_n^s} \right]^{\frac{1}{r-s}}.$$

Proposition 4. For $\lambda \in \Gamma_k$, we have if $0 \leq l < k \leq n$

$$\frac{\partial[\frac{\sigma_k}{\sigma_l}](\lambda)}{\partial\lambda_i} > 0, \quad \forall 1 \leq i \leq n.$$

But,

$$\frac{\partial[\frac{\sigma_k}{\sigma_{k-1}}](\lambda)}{\partial\lambda_i} \geq 0, \quad \forall 1 \leq i \leq n,$$

holds for $\lambda \in \Gamma_{k-1}$.

Proposition 5. For any $n \geq k > l \geq 0$, we have

$$\left[\frac{\sigma_k(\lambda)}{\sigma_l(\lambda)} \right]^{\frac{1}{k-l}}$$

is a concave function in Γ_k . But, for any $k \geq 1$, we have

$$\frac{\sigma_k(\lambda)}{\sigma_{k-1}(\lambda)}$$

is a concave function in Γ_{k-1} .

We recall the Gårding's inequality (see [4]).

Lemma 2.1. If $\lambda, \mu \in \Gamma_k$, then for any $k \in \{1, 2, \dots, n\}$

$$\sum_{i=1}^n \mu_i \sigma_{k-1}(\lambda|i) \geq k [\sigma_k(\mu)]^{\frac{1}{k}} [\sigma_k(\lambda)]^{1-\frac{1}{k}}.$$

In particular,

$$(2.5) \quad \sum_{i=1}^n \mu_i \sigma_{k-1}(\lambda|i) > 0.$$

Lemma 2.2. If $\lambda \in \Gamma_k$, then for any $i \in \{1, 2, \dots, n\}$ and $1 \leq l < k \leq n$ we have

$$(2.6) \quad \frac{\sigma_{k-1}(\lambda|i)}{\sigma_{l-1}(\lambda|i)} > \frac{\sigma_k(\lambda)}{\sigma_l(\lambda)}.$$

In particular, if $\lambda \in \Gamma_{k-1}$, we have

$$(2.7) \quad \frac{\sigma_{k-1}(\lambda|i)}{\sigma_{k-2}(\lambda|i)} \geq \frac{\sigma_k(\lambda)}{\sigma_{k-1}(\lambda)}.$$

Proof. We get from (2.1)

$$h(\lambda_i) := \frac{\sigma_k(\lambda)}{\sigma_l(\lambda)} = \frac{\lambda_i \sigma_{k-1}(\lambda|i) + \sigma_k(\lambda|i)}{\lambda_i \sigma_{l-1}(\lambda|i) + \sigma_l(\lambda|i)}.$$

A direct computation implies

$$\frac{d}{d\lambda_i} h(\lambda_i) = \frac{\sigma_{k-1}(\lambda|i) \sigma_l(\lambda|i) - \sigma_k(\lambda|i) \sigma_{l-1}(\lambda|i)}{\sigma_l^2(\lambda)}.$$

If $\sigma_k(\lambda|i) \leq 0$, it is easy to see $\frac{d}{d\lambda_i}h(\lambda_i) > 0$. Otherwise, $\sigma_k(\lambda|i) > 0$. Thus, λ with $\lambda_i = 0$ belongs to Γ_k . So, using the generalized Newton-MacLaurin inequality (2.4), we have

$$\frac{\sigma_k(\lambda|i)}{\sigma_{k-1}(\lambda|i)} < \frac{k(n-l)}{l(n-k)} \frac{\sigma_k(\lambda|i)}{\sigma_{k-1}(\lambda|i)} \leq \frac{\sigma_l(\lambda|i)}{\sigma_{l-1}(\lambda|i)}.$$

Thus, $\frac{d}{d\lambda_i}h(\lambda_i) > 0$. Therefore,

$$\frac{\sigma_k(\lambda)}{\sigma_l(\lambda)} < \lim_{\lambda_i \rightarrow +\infty} h(\lambda_i) = \frac{\sigma_{k-1}(\lambda|i)}{\sigma_{l-1}(\lambda|i)}.$$

Hence, we obtain (2.6). For the special case $l = k - 1$, we have $\frac{d}{d\lambda_i}h(\lambda_i) \geq 0$ by the Newton inequality (2.3). Thus, the inequality (2.7) follows as before. \square

At last, we list the following well-known result (See [1]).

Lemma 2.3. *If $A = (a_{ij})$ is a Hermitian matrix, $\lambda_i = \lambda_i(A)$ is one of its eigenvalues ($i = 1, \dots, n$) and $F = F(A) = f(\lambda(A))$ is a symmetric function of $\lambda_1, \dots, \lambda_n$, then for any Hermitian matrix $B = (b_{ij})$, we have the following formulas:*

$$(2.8) \quad \frac{\partial^2 F}{\partial a_{ij} \partial a_{st}} b_{ij} b_{st} = \frac{\partial^2 f}{\partial \lambda_p \partial \lambda_q} b_{pp} b_{qq} + 2 \sum_{p < q} \frac{\frac{\partial f}{\partial \lambda_p} - \frac{\partial f}{\partial \lambda_q}}{\lambda_p - \lambda_q} b_{pq}^2.$$

In addition, if f is a concave function and

$$\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n,$$

then we have

$$(2.9) \quad f_1 \geq f_2 \geq \dots \geq f_n,$$

where $f_i = \frac{\partial f}{\partial \lambda_i}$.

Let $\lambda(a_{i\bar{j}})$ denote the eigenvalues of a Hermitian matrix $(a_{i\bar{j}})$. Define $\sigma_k(a_{i\bar{j}}) = \sigma_k(\lambda(a_{i\bar{j}}))$. This definition can be naturally extended to Kähler manifolds (and more generally, Hermitian manifolds). Let $\mathcal{A}^{1,1}(M)$ be the space of real smooth $(1, 1)$ -forms on (M, ω) . For any $\chi \in \mathcal{A}^{1,1}(M)$, in a local normal coordinate system of M with respect with ω , we have

$$\chi = \frac{\sqrt{-1}}{2} \chi_{i\bar{j}} dz^i \wedge d\bar{z}^j.$$

Definition 2.1. We define $\sigma_k(\chi)$ with respect to ω as

$$\sigma_k(\chi) = \sigma_k(\chi_{i\bar{j}}) = \sigma_k(\lambda(\chi_{i\bar{j}})).$$

The definition of σ_k is independent of the choice of local normal coordinate system. In fact, σ_k can be defined without the use of local normal coordinate by

$$\sigma_k(\chi) = C_n^k \frac{\chi^k \wedge \omega^{n-k}}{\omega^n},$$

where $C_n^k = \frac{n!}{(n-k)!k!}$. We also define the Gårding's cone on M by

$$\Gamma_k(M) = \{\chi \in \mathcal{A}^{1,1}(M) : \sigma_i(\chi) > 0, \forall 1 \leq i \leq k\}.$$

Using the above notation, we can rewrite equation (1.1) as the following local form:

$$(2.10) \quad \sigma_k(\chi_u) = \sum_{l=0}^{k-1} \beta_l(z) \sigma_l(\chi_u),$$

where $\beta_l(z) = \frac{C_n^k}{C_n^l} \alpha_l(z)$.

Moreover, we also have a local version of the cone condition (1.2).

Lemma 2.4. $\chi \in \mathcal{C}_k(\omega)$ is equivalent to

$$(2.11) \quad \sigma_{k-1}(\chi|i) - \sum_{l=1}^{k-1} \beta_l \sigma_{l-1}(\chi|i) > 0$$

for any $i \in \{1, 2, \dots, n\}$, where $(\chi|i)$ denotes the matrix obtained by deleting the i -th row and i -th column of χ , and $\beta_l(z) = \frac{C_n^k}{C_n^l} \alpha_l(z)$ is denoted as before.

Proof. The equality (2.11) follows directly if we observe that coefficient of $(n-1, n-1)$ form $\prod_{j=1, j \neq i}^n dz^j d\bar{z}^j$ in $\chi^{l-1} \wedge \omega^{n-l}$ is

$$(l-1)!(n-l)! \sigma_{l-1}(\chi|i) = \frac{1}{l} \frac{n!}{C_n^l} \sigma_{l-1}(\chi|i).$$

□

2.2. The ellipticity and concavity. To make the equation (2.10) elliptic and concave, we follow an important observation by Guan-Zhang [25] to rewrite (2.10) as the form:

$$(2.12) \quad F(u_{i\bar{j}}, z) := \frac{\sigma_k(\chi_u)}{\sigma_{k-1}(\chi_u)} - \sum_{l=0}^{k-1} \beta_l(z) \frac{\sigma_l(\chi_u)}{\sigma_{k-1}(\chi_u)} = \beta_{k-1}(z).$$

Then, as a corollary of Propositions 4 and 5, we obtain (see also Proposition 2.2 in [25]):

Lemma 2.5. *If u is a C^2 function with $\chi_u \in \Gamma_{k-1}(M)$, and $\beta_l(z)$ ($0 \leq l \leq k-2$) are nonnegative, then the operator F is elliptic and concave. Moreover, if $\sum_{l=0}^{k-2} \beta_l(z) > 0$ for all $z \in M$, the operator F is strictly elliptic.*

2.3. Some key lemmas. In this subsection, we prove some inequalities and lemmas which will play an important role in the establishment of the a priori estimates. In the following, for $\lambda(z) = (\lambda_1(z), \dots, \lambda_n(z)) \in C^0(M, \mathbb{R}^n)$, we set

$$(2.13) \quad f(\lambda(z), z) := \frac{\sigma_k(\lambda(z))}{\sigma_{k-1}(\lambda(z))} - \sum_{l=0}^{k-1} \beta_l(z) \frac{\sigma_l(\lambda(z))}{\sigma_{k-1}(\lambda(z))} = \beta_{k-1}(z)$$

and denote by $f_i(\lambda) = \frac{\partial f}{\partial \lambda_i}(\lambda)$.

Lemma 2.6. Let $\beta_0(z), \dots, \beta_{k-2}(z)$ be nonnegative functions on M . Assume that $\lambda = (\lambda_1, \dots, \lambda_n) \in \Gamma_{k-1}$ and $\mu = (\mu_1, \dots, \mu_n) \in \Gamma_{k-1}$, then we have

$$(2.14) \quad \sum_i f_i(\lambda) \mu_i \geq f(\mu) + (k-1) \sum_{l=0}^{k-1} \beta_l \frac{\sigma_l(\lambda)}{\sigma_{k-1}(\lambda)}.$$

Proof. Since f is concave in Γ_{k-1} (see Lemma 2.5), we have

$$f(\mu) \leq \sum_i f_i(\lambda) (\mu_i - \lambda_i) + f(\lambda),$$

which implies

$$\sum_i f_i(\lambda) \mu_i \geq f(\mu) + (k-1) \sum_{l=0}^{k-1} \beta_l \frac{\sigma_l(\lambda)}{\sigma_{k-1}(\lambda)},$$

where we used the fact

$$\sum_i f_i(\lambda) \lambda_i = f(\lambda) + \sum_{l=0}^{k-1} (k-l) \beta_l \frac{\sigma_l(\lambda)}{\sigma_{k-1}(\lambda)}.$$

So, the proof is complete. \square

Lemma 2.7. Let $\beta_0(z), \dots, \beta_{k-2}(z)$ be nonnegative continuous functions on M and $\beta_{k-1}(z)$ be continuous functions on M . Assume $\mu(z) = (\mu_1(z), \dots, \mu_n(z)) \in \Gamma_{k-1}$ for any $z \in M$ and μ satisfy

$$(2.15) \quad \frac{\sigma_{k-1}(\mu|i)}{\sigma_{k-2}(\mu|i)} - \sum_{l=1}^{k-2} \beta_l(z) \frac{\sigma_{l-1}(\mu|i)}{\sigma_{k-2}(\mu|i)} > \beta_{k-1}(z).$$

for arbitrary $z \in M$ and $i \in \{1, 2, \dots, n\}$. Then there are constants $N, \theta > 0$ depending on $|\mu|_{C^0(M)}$ and $|\beta_i|_{C^0(M)}$ with $0 \leq i \leq k-1$ such that for any $z \in M$, if

$$\max_{1 \leq i \leq n} \{\lambda_i(z)\} \geq N,$$

we have at z

$$(2.16) \quad \sum_i f_i(\lambda) (\lambda_i - \mu_i) \leq -\theta - \theta \sum_i f_i(\lambda)$$

or

$$(2.17) \quad f_1(\lambda) \lambda_1 \geq \theta.$$

Proof. For any $z \in M$, without loss of generality, we may assume $\lambda_1(z) = \max_{1 \leq i \leq n} \{\lambda_i(z)\}$. If $\lambda_1 \gg -\mu_1 + \epsilon$, we have

$$\begin{aligned}
& \sum_i f_i(\lambda)(\lambda_i - \mu_i) \\
&= \sum_{i \geq 2} f_i(\lambda)(\lambda_i - (\mu_i - \epsilon)) - \epsilon \sum_i f_i(\lambda) + f_1(\lambda)(\lambda_1 - \mu_1 + \epsilon) \\
(2.18) \quad &\leq \sum_{i \geq 2} f_i(\lambda)(\lambda_i - (\mu_i - \epsilon)) - \epsilon \sum_i f_i(\lambda) + 2f_1(\lambda)\lambda_1.
\end{aligned}$$

Choosing ϵ small enough which is independent of z such that $(\mu_1 - \epsilon, \dots, \mu_n - \epsilon) \in \Gamma_{k-1}$ satisfies (2.15). So we can choose λ_1 large enough which is independent of z such that $\tilde{\mu} = (\lambda_1, \mu_2 - \epsilon, \dots, \mu_n - \epsilon) \in \Gamma_{k-1}$ and

$$(2.19) \quad f(\tilde{\mu}) \geq \beta_{k-1} + \tilde{\epsilon}$$

in view that

$$\lim_{\lambda_1 \rightarrow +\infty} f(\tilde{\mu}) = \frac{\sigma_{k-1}(\mu')}{\sigma_{k-2}(\mu')} - \sum_{l=1}^{k-2} \beta_l \frac{\sigma_{l-1}(\mu')}{\sigma_{k-2}(\mu')} > \beta_{k-1}.$$

where $\mu' = (\mu_2 - \epsilon, \dots, \mu_n - \epsilon)$ and we use the inequality (2.15). Now, we rewrite (2.18) as

$$\begin{aligned}
& \sum_i f_i(\lambda)(\lambda_i - \mu_i) \\
(2.20) \quad &\leq \sum_i f_i(\lambda)(\lambda_i - \tilde{\mu}_i) - \epsilon \sum_i f_i(\lambda) + 2f_1(\lambda)\lambda_1.
\end{aligned}$$

Since f is concave in Γ_{k-1} (see Lemma 2.5), there is

$$f(\tilde{\mu}) \leq f_i(\lambda)(\tilde{\mu}_i - \lambda_i) + f(\lambda),$$

which implies together with (2.20)

$$\sum_i f_i(\lambda)(\lambda_i - \mu_i) \leq f(\lambda) - f(\tilde{\mu}) - \epsilon \sum_i f_i(\lambda) + 2f_1(\lambda)\lambda_1.$$

Thus, when λ_1 is large enough, we have by (2.19)

$$\sum_i f_i(\lambda)(\lambda_i - \mu_i) \leq -\tilde{\epsilon} - \epsilon \sum_i f_i(\lambda) + 2f_1(\lambda)\lambda_1.$$

If $f_1(\lambda)\lambda_1 \leq \frac{\tilde{\epsilon}}{4}$, we obtain

$$\sum_i f_i(\lambda)(\lambda_i - \mu_i) \leq -\frac{\tilde{\epsilon}}{2} - \epsilon \sum_i f_i(\lambda).$$

Then, we complete the proof if we choose $\theta = \min\{\frac{\tilde{\epsilon}}{4}, \epsilon\}$. □

Lemma 2.8. Assume $\lambda \in \Gamma_{k-1}$ satisfies (2.13) and $\beta_l(z) \geq 0$ ($0 \leq l \leq k-2$), then we have

$$(2.21) \quad 0 < \frac{\sigma_l(\lambda)}{\sigma_{k-1}(\lambda)} \leq C, \quad 0 \leq l \leq k-2,$$

where the constant C depends on $n, k, \inf \beta_l, \sup |\beta_{k-1}|$. Moreover, we have

$$(2.22) \quad -\sup |\beta_{k-1}| \leq \frac{\sigma_k(\lambda)}{\sigma_{k-1}(\lambda)} \leq C,$$

where the constant C depends on $n, k, \inf \beta_l$ for $0 \leq l \leq k-2$, and $\sum_{i=0}^{k-1} |\beta_i|$.

Proof. On the one hand, if $\frac{\sigma_k}{\sigma_{k-1}} \leq 1$, then we get from the equation (2.13)

$$\beta_l \frac{\sigma_l}{\sigma_{k-1}} \leq \frac{\sigma_k}{\sigma_{k-1}} - \beta_{k-1} \leq 1 + \sup |\beta_{k-1}|, \quad 0 \leq l \leq k-2.$$

On the other hand, if $\frac{\sigma_k}{\sigma_{k-1}} > 1$, i.e. $\frac{\sigma_{k-1}}{\sigma_k} < 1$. We can get for $0 \leq l \leq k-2$ by the Newton-MacLaurin inequality (2.4),

$$\frac{\sigma_l}{\sigma_{k-1}} \leq \frac{(C_n^k)^{k-1-l} C_n^l}{(C_n^{k-1})^{k-l}} \left(\frac{\sigma_{k-1}}{\sigma_k} \right)^{k-1-l} \leq \frac{(C_n^k)^{k-1-l} C_n^l}{(C_n^{k-1})^{k-l}} \leq C(n, k).$$

So, we get (2.21). Then, it follows that

$$-\sup |\beta_{k-1}| \leq \frac{\sigma_k}{\sigma_{k-1}} = \sum_{l=0}^{k-2} \beta_l \frac{\sigma_l}{\sigma_{k-1}} + \beta_{k-1} \leq C \sum_{i=0}^n |\beta_i|.$$

Thus, the proof is complete. \square

Lemma 2.9. Assume $\lambda \in \Gamma_{k-1}$ and $\beta_l(z) \geq 0$ ($0 \leq l \leq k-2$), then

$$(2.23) \quad \sum_{i=1}^n f_i \geq \frac{n-k+1}{k}.$$

Proof. By direct computations, we can get by Propositions 4, 1 and 3

$$\begin{aligned} \sum_{i=1}^n f_i &\geq \sum_{i=1}^n \frac{\partial \left(\frac{\sigma_k}{\sigma_{k-1}} \right)}{\partial \lambda_i} = \sum_{i=1}^n \frac{\sigma_{k-1}(\lambda|i) \sigma_{k-1} - \sigma_k \sigma_{k-2}(\lambda|i)}{\sigma_{k-1}^2} \\ &= \frac{(n-k+1) \sigma_{k-1}^2 - (n-k+2) \sigma_k \sigma_{k-2}}{\sigma_{k-1}^2} \\ &\geq \frac{n-k+1}{k}, \end{aligned}$$

where we get the last inequality from Newton' inequality (2.2). Hence (2.23) holds. \square

3. C^0 ESTIMATE

In this section, we follow the idea of Sun [45] to derive C^0 estimate directly from the cone condition. Moreover, we expect that our estimate holds true for α_{k-1} with indefinite sign and does not depend on the lower bound of α_i ($0 \leq i \leq k-2$).

3.1. Some lemmas. Since $\chi_0 \in \Gamma_{k-1}(M)$, we may assume that there is a uniform constant $\tau > 0$ such that

$$(3.1) \quad \chi_0 - \tau\omega \in \Gamma_{k-1}(M) \quad \text{and} \quad \omega - \tau\chi_0 \in \Gamma_{k-1}(M).$$

Then, we can succeed in extending Lemma 2.3 in [45] to our case.

Lemma 3.1. *Let (M, ω) be a Kähler manifold of complex dimension $n \geq 2$ and $\alpha_0(z), \dots, \alpha_{k-1}(z)$ be continuous functions on M which satisfy*

$$\alpha_i(z) \geq 0, \quad 0 \leq i \leq k-2 \quad \text{and} \quad \sum_{l=0}^{k-2} \alpha_l(z) > 0 \quad \text{for all } z \in M.$$

Suppose that $\chi_0 \in \Gamma_{k-1}(M)$ satisfies (3.1). If $u \in C^2(M)$ satisfies $\chi_u \in \Gamma_{k-1}(M)$, then we have the following pointwise inequalities:

(1) for $0 \leq t \leq 1$ and $1 \leq l \leq k-1$

$$(3.2) \quad \chi_{tu}^{l-1} \wedge \omega^{n-l} \geq (1-t)^{l-1} \tau^{l-1} \omega^{n-1}.$$

(2) for $0 < t \leq 1$ and $1 \leq l \leq k-1$,

$$(3.3) \quad \chi_u^l \wedge \omega^{n-l} \leq \frac{1}{t^l} \chi_{tu}^l \wedge \omega^{n-l}.$$

Moreover, if u is a solution to the equation (1.1) and χ_0 satisfies the cone condition (1.2), then there exists a uniform constant $C > 0$ such that for $0 \leq t \leq 1$

$$(3.4) \quad \begin{aligned} & k\chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l\alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \\ & > C(1-t)\chi_{tu}^{k-2} \wedge \omega^{n-k+1}, \end{aligned}$$

and consequently

$$(3.5) \quad \begin{aligned} & k\chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l\alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \\ & > C\tau^{k-2}(1-t)^{k-1} \omega^{n-1}. \end{aligned}$$

Proof. For $1 \leq l \leq k-1$ and $1 \leq i \leq n$, $\sigma_l^{\frac{1}{l}}(\lambda)$ and $\sigma_{l-1}^{\frac{1}{l-1}}(\lambda|i)$ are concave in Γ_l and Γ_{l-1} respectively. Thus,

$$(3.6) \quad \sigma_l^{\frac{1}{l}}(\chi_{tu}) \geq (1-t)\sigma_l^{\frac{1}{l}}(\chi_0) + t\sigma_l^{\frac{1}{l}}(\chi_u) \geq t\sigma_l^{\frac{1}{l}}(\chi_u),$$

$$\begin{aligned}
\sigma_{l-1}^{\frac{1}{l-1}}(\chi_{tu}|i) &\geq (1-t)\sigma_{l-1}^{\frac{1}{l-1}}(\chi_0|i) + t\sigma_{l-1}^{\frac{1}{l-1}}(\chi_u|i) \\
(3.7) \qquad \qquad \qquad &\geq (1-t)\sigma_{l-1}^{\frac{1}{l-1}}(\chi_0|i),
\end{aligned}$$

and

$$(3.8) \quad \sigma_{l-1}^{\frac{1}{l-1}}(\chi_0|i) \geq \sigma_{l-1}^{\frac{1}{l-1}}(\chi_0 - \tau\omega|i) + \tau\sigma_{l-1}^{\frac{1}{l-1}}(\omega|i) \geq \tau\sigma_{l-1}^{\frac{1}{l-1}}(\omega|i).$$

Combining (3.7) and (3.8) gives

$$\sigma_{l-1}(\chi_{tu}|i) \geq (1-t)^{l-1}\tau^{l-1}\sigma_{l-1}(\chi_0|i),$$

which is just the local form of (3.2). Similarly, (3.3) is a consequence of (3.6). Now we only need to prove (3.4) and (3.5). Since

$$f(\chi|i) := \frac{\sigma_{k-1}(\chi|i)}{\sigma_{k-2}(\chi|i)} - \sum_{l=1}^{k-2} \beta_l \frac{\sigma_{l-1}(\chi|i)}{\sigma_{k-2}(\chi|i)}$$

is concave in Γ_{k-2} (see Lemma 2.5), we can obtain

$$(3.9) \quad f(\chi_{tu}|i) \geq (1-t)f(\chi_0|i) + tf(\chi_u|i).$$

Moreover, we have by Lemma 2.2

$$\begin{aligned}
f(\chi_u|i) &= \frac{\sigma_{k-1}(\chi_u|i)}{\sigma_{k-2}(\chi_u|i)} - \sum_{l=1}^{k-2} \beta_l \frac{\sigma_{l-1}(\chi_u|i)}{\sigma_{k-2}(\chi_u|i)} \\
&> \frac{\sigma_k(\chi_u)}{\sigma_{k-1}(\chi_u)} - \sum_{l=1}^{k-2} \beta_l \frac{\sigma_l(\chi_u)}{\sigma_{k-1}(\chi_u)} \\
&= \frac{\beta_0}{\sigma_{k-1}(\chi_u)} + \beta_{k-1} \\
(3.10) \qquad \qquad \qquad &\geq \beta_{k-1}.
\end{aligned}$$

In addition, since χ_0 satisfies the cone condition (1.2), there exists some uniform constant $\delta > 0$ which is independent of z such that

$$(3.11) \quad f(\chi_0|i) > \beta_{k-1}(z) + \delta,$$

where we write the cone condition in a local version as Lemma 2.4. Substituting (3.10) and (3.11) into (3.9), we get

$$\frac{\sigma_{k-1}(\chi_{tu}|i)}{\sigma_{k-2}(\chi_{tu}|i)} - \sum_{l=1}^{k-2} \beta_l \frac{\sigma_{l-1}(\chi_{tu}|i)}{\sigma_{k-2}(\chi_{tu}|i)} > (1-t)(\beta_{k-1} + \delta) + t\beta_{k-1},$$

this is to say

$$\sigma_{k-1}(\chi_{tu}|i) - \sum_{l=1}^{k-1} \beta_l \sigma_{l-1}(\chi_{tu}|i) > \delta(1-t)\sigma_{k-2}(\chi_{tu}|i),$$

which is just the local form of (3.4). Then, (3.4) follows by substituting (3.2) into it. \square

Next, we derive the following important inequality by iterating the inequality (3.15) in [45].

Lemma 3.2. *Let (M, ω) be a Kähler manifold of complex dimension $n \geq 2$. Suppose that $\chi_0 \in \Gamma_{k-1}(M)$ satisfies (3.1). If $u \in C^2(M)$ satisfies $\chi_u \in \Gamma_{k-1}(M)$, then we have the following inequalities for $l < k$:*

$$(3.12) \quad \begin{aligned} & \frac{k-1}{l-1} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{k-2} \wedge \omega^{n-k+1} \\ & \geq \tau^{k-l} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1}. \end{aligned}$$

Proof. Using integration by parts and Garding's inequality (2.5), it yields

$$\begin{aligned} & \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-1} \wedge \omega^{n-l} \\ & \geq \tau \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1} \\ & \quad + \frac{1}{l-1} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge t \frac{d}{dt} (\chi_{tu}^{l-1} \wedge \omega^{n-l}) \\ & \geq \tau \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1} \\ & \quad - \frac{1}{l-1} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-1} \wedge \omega^{n-l}. \end{aligned}$$

Thus,

$$\begin{aligned} & \frac{l}{l-1} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-1} \wedge \omega^{n-l} \\ & \geq \tau \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1}. \end{aligned}$$

So, we obtain by iteration

$$\begin{aligned} & \frac{k-1}{l-1} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{k-2} \wedge \omega^{n-k+1} \\ & \geq \tau^{k-l} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1}, \end{aligned}$$

which complete the proof. \square

Once we have established the following inequality (3.13), we can use it to derive C^0 estimate (3.14) by the standard argument in [49, 50] without using the equation (1.1).

Lemma 3.3. *Let (M, ω) be a closed Kähler manifold of complex dimension $n \geq 2$ and $\alpha_0(z), \dots, \alpha_{k-1}(z)$ be C^2 functions on M which satisfy*

$$\alpha_i(z) \geq 0, \quad 0 \leq i \leq k-2 \quad \text{and} \quad \sum_{l=0}^{k-2} \alpha_l(z) > 0 \quad \text{for all } z \in M.$$

Suppose that $\chi_0 \in \Gamma_{k-1}(M)$ is closed and satisfies (3.1). If $u \in C^2(M)$ satisfies $\chi_u \in \Gamma_{k-1}(M)$, then there exists uniform constants C and p_0 such that for $p \geq p_0$ the following inequality holds

$$(3.13) \quad \int_M |\partial e^{-\frac{p}{2}u}|_g^2 \omega^n \leq Cp \int_M e^{-pu} \omega^n.$$

Thus, there exists a uniform constant C depends on the given data M, ω, χ_0 , and $|\alpha_i|_{C^0(M)}$ ($0 \leq i \leq k-1$) such that

$$(3.14) \quad |u|_{C^0(M)} \leq C \quad \text{with} \quad \sup_M u = 0.$$

Proof. Since

$$\begin{aligned} & (\chi_u^k \wedge \omega^{n-k} - \chi_0^k \wedge \omega^{n-k}) - \sum_{l=0}^{k-1} \alpha_l (\chi_u^l \wedge \omega^{n-l} - \chi_0^l \wedge \omega^{n-l}) \\ &= \int_0^1 \sqrt{-1} \partial \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) dt, \end{aligned}$$

we have

$$\begin{aligned} & \int_M e^{-pu} \left[(\chi_u^k \wedge \omega^{n-k} - \chi_0^k \wedge \omega^{n-k}) - \sum_{l=0}^{k-1} \alpha_l (\chi_u^l \wedge \omega^{n-l} - \chi_0^l \wedge \omega^{n-l}) \right] \\ &= p \int_0^1 dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) \\ & \quad - \sum_{l=1}^{k-1} \frac{l}{p} \int_0^1 dt \int_M e^{-pu} \sqrt{-1} \partial \bar{\partial} \alpha_l \wedge \chi_{tu}^{l-1} \wedge \omega^{n-l} \\ &\geq p \int_0^1 dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) \\ (3.15) \quad & - \sum_{l=1}^{k-1} \frac{C_l}{p} \int_0^1 dt \int_M e^{-pu} \chi_{tu}^{l-1} \wedge \omega^{n-l+1}. \end{aligned}$$

Notice that

$$\begin{aligned}
(3.16) \quad & \int_M e^{-pu} \left[(\chi_u^k \wedge \omega^{n-k} - \chi_0^k \wedge \omega^{n-k}) - \sum_{l=0}^{k-1} \alpha_l (\chi_u^l \wedge \omega^{n-l} - \chi_0^l \wedge \omega^{n-l}) \right] \\
&= \int_M e^{-pu} \left(-\chi_0^k \wedge \omega^{n-k} + \sum_{l=0}^{k-1} \alpha_l \chi_0^l \wedge \omega^{n-l} \right) \\
&\leq C \int_M e^{-pu} \omega^n.
\end{aligned}$$

Using the inequality (3.3), we obtain

$$\begin{aligned}
(3.17) \quad & \int_0^1 dt \int_M e^{-pu} \chi_{tu}^{l-1} \wedge \omega^{n-l+1} \\
&\leq 2^{l-1} \int_0^1 dt \int_M e^{-pu} \chi_{\frac{tu}{2}}^{l-1} \wedge \omega^{n-l+1} \\
&\leq 2^l \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \chi_{tu}^{l-1} \wedge \omega^{n-l+1}.
\end{aligned}$$

Plugging the inequalities (3.16) and (3.17) into (3.15), it yields

$$\begin{aligned}
(3.18) \quad & p \int_0^1 dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) \\
&\leq \sum_{l=1}^{k-1} \frac{2^l C_l}{p} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \chi_{tu}^{l-1} \wedge \omega^{n-l+1} + C \int_M e^{-pu} \omega^n.
\end{aligned}$$

We deal with the first term on the right side of the inequality (3.18) according to the inequality (3.12)

$$\begin{aligned}
(3.19) \quad & \frac{1}{p} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \chi_{tu}^{l-1} \wedge \omega^{n-l+1} \\
&= (l-1) \int_0^{\frac{1}{2}} dt \int_0^t ds \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{su}^{l-2} \wedge \omega^{n-l+1} \\
&\quad + \frac{1}{2p} \int_M e^{-pu} \chi_0^{l-1} \wedge \omega^{n-l+1} \\
&\leq \frac{l-1}{2} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{l-2} \wedge \omega^{n-l+1} \\
&\quad + \frac{1}{2p} \int_M e^{-pu} \chi_0^{l-1} \wedge \omega^{n-l+1} \\
&\leq \frac{k-1}{\tau^{k-l}} \int_0^{\frac{1}{2}} dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{k-2} \wedge \omega^{n-k+1} \\
&\quad + \frac{C}{2p} \int_M e^{-pu} \omega^n.
\end{aligned}$$

To cancel the first term in the right hand of (3.19), we will use part of the left term in (3.18). In details, we can get the following positive term for $0 \leq t \leq \frac{1}{2}$ from the inequality (3.4)

$$(3.20) \quad \begin{aligned} & p e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=1}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) \\ & \geq C p e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \chi_{tu}^{k-2} \wedge \omega^{n-k+1}. \end{aligned}$$

Thus, if we choose p sufficiently large, the integral of the term (3.20) on M can kill the first term in the right hand of (3.19). Then, (3.18) becomes

$$\begin{aligned} & \frac{p}{2} \int_0^1 dt \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \left(k \chi_{tu}^{k-1} \wedge \omega^{n-k} - \sum_{l=0}^{k-1} l \alpha_l \chi_{tu}^{l-1} \wedge \omega^{n-l} \right) \\ & \leq C \int_M e^{-pu} \omega^n, \end{aligned}$$

which implies in view of (3.5)

$$p \int_M e^{-pu} \sqrt{-1} \partial u \wedge \bar{\partial} u \wedge \omega^{n-1} \leq C \int_M e^{-pu} \omega^n.$$

So, our proof is completed. \square

4. C^2 ESTIMATE

4.1. Notations and some lemmas. In local complex coordinates (z^1, \dots, z^n) , the subscripts of a function u always denote the covariant derivatives of u with respect to ω in the directions of the local frame $\frac{\partial}{\partial z^1}, \dots, \frac{\partial}{\partial z^n}$. Namely,

$$u_i = \nabla_{\frac{\partial}{\partial z^i}} u, \quad u_{i\bar{j}} = \nabla_{\frac{\partial}{\partial \bar{z}^j}} \nabla_{\frac{\partial}{\partial z^i}} u, \quad u_{i\bar{j}k} = \nabla_{\frac{\partial}{\partial z^k}} \nabla_{\frac{\partial}{\partial \bar{z}^j}} \nabla_{\frac{\partial}{\partial z^i}} u.$$

But, the covariant derivatives of a $(1, 1)$ -form χ with respect to ω will be denoted by indices with semicolons, e.g.,

$$\chi_{i\bar{j};k} = \nabla_{\frac{\partial}{\partial z^k}} \chi \left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j} \right), \quad \chi_{i\bar{j};k\bar{l}} = \nabla_{\frac{\partial}{\partial \bar{z}^l}} \nabla_{\frac{\partial}{\partial z^k}} \chi \left(\frac{\partial}{\partial z^i}, \frac{\partial}{\partial \bar{z}^j} \right).$$

We recall the following commutation formula on Kähler manifolds (M, ω) [28, 22, 23].

Lemma 4.1. *For $u \in C^4(M)$, we have*

$$u_{i\bar{j}l} = u_{i\bar{j}l} - u_p R_{l\bar{j}i}^p, \quad u_{p\bar{j}m} = u_{p\bar{j}m}, \quad u_{i\bar{q}l} = u_{l\bar{q}i},$$

$$u_{i\bar{j}l\bar{m}} = u_{l\bar{m}i\bar{j}} + u_{p\bar{j}} R_{l\bar{m}i}^p - u_{p\bar{m}} R_{i\bar{j}l}^p,$$

where R is the curvature tensor of (M, ω) .

For the convenience of notations, we will denote

$$F_k(\chi) := \frac{\sigma_k(\chi)}{\sigma_{k-1}(\chi)}, \quad F_l(\chi) := -\frac{\sigma_l(\chi)}{\sigma_{k-1}(\chi)}, \quad 0 \leq l \leq k-2,$$

$$F(\chi, z) := F_k(\chi) + \sum_{l=0}^{k-2} \beta_l(z) F_l(\chi),$$

and

$$F^{i\bar{j}} := \frac{\partial F}{\partial \chi_{i\bar{j}}}, \quad F^{i\bar{j}, k\bar{l}} := \frac{\partial^2 F}{\partial \chi_{i\bar{j}} \partial \chi_{r\bar{s}}}, \quad F_l^{i\bar{j}} := \frac{\partial F_l}{\partial \chi_{i\bar{j}}}, \quad F_l^{i\bar{j}, r\bar{s}} := \frac{\partial^2 F_l}{\partial \chi_{i\bar{j}} \partial \chi_{r\bar{s}}},$$

where $1 \leq i, j, r, s \leq n$ and $0 \leq l \leq k-2$.

Lemma 4.2. *Let $u \in C^4(M)$ be a solution to the equation (2.12) on (M, ω) and $\chi \in \Gamma_{k-1}(M)$. For any $z \in M$, we choose a normal coordinate such that at this point $\omega = \frac{\sqrt{-1}}{2} \delta_{ij} dz^i \wedge d\bar{z}^j$. Then, we have for any $\delta < 1$ at z*

$$(4.1) \quad \begin{aligned} F^{i\bar{j}} \chi_{i\bar{j}; p\bar{p}} &\geq -(1 - \delta^2) F^{i\bar{j}, r\bar{s}} \chi_{i\bar{j}; p} \chi_{r\bar{s}; \bar{p}} + \nabla_{\bar{p}} \nabla_p \beta_{k-1} \\ &+ \sum_{l=0}^{k-2} \frac{1}{1 + \frac{1}{k+1-l}} \frac{|\nabla_p \beta_l|^2}{\delta^2 \beta_l} F_l - \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l. \end{aligned}$$

Proof. Differentiating the equation (2.12) once, we have

$$\nabla_p \beta_{k-1} = F^{i\bar{j}} \chi_{i\bar{j}; p} + \sum_{l=0}^{k-2} \nabla_p \beta_l \cdot F_l.$$

Differentiating the equation (2.12) again, we obtain

$$\begin{aligned} \nabla_{\bar{p}} \nabla_p \beta_{k-1} &= F^{i\bar{j}, r\bar{s}} \chi_{i\bar{j}; p} \chi_{r\bar{s}; \bar{p}} + F^{i\bar{j}} \chi_{i\bar{j}; p\bar{p}} + \sum_{l=0}^{k-2} \left(\nabla_{\bar{p}} \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j}; p} + \nabla_p \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j}; \bar{p}} \right) \\ &+ \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l. \end{aligned}$$

Moreover, since the operator $(\frac{\sigma_{k-1}}{\sigma_l})^{\frac{1}{k-1-l}}$ is concave for $0 \leq l \leq k-2$ (see Proposition 5), we have

$$(4.2) \quad -F_l^{i\bar{j}, r\bar{s}} \chi_{i\bar{j}; p} \chi_{r\bar{s}; \bar{p}} \geq -\left(1 + \frac{1}{k-1-l}\right) F_l^{-1} F_l^{i\bar{j}} F_l^{r\bar{s}} \chi_{i\bar{j}; p} \chi_{r\bar{s}; \bar{p}}.$$

Since F_k is concave in Γ_{k-1} , throwing away the negative term

$$\delta^2 \frac{\partial^2 F_k}{\partial \chi_{i\bar{j}} \partial \chi_{r\bar{s}}} \chi_{i\bar{j}; p} \chi_{r\bar{s}; \bar{p}}$$

gives

$$\begin{aligned}
& \nabla_{\bar{p}} \nabla_p \beta_{k-1} - (1 - \delta^2) F^{i\bar{j}, r\bar{s}} \chi_{i\bar{j};p} \chi_{r\bar{s};\bar{p}} \\
& \leq \sum_{l=0}^{k-2} \delta^2 \beta_l F_l^{i\bar{j}, r\bar{s}} \chi_{i\bar{j};p} \chi_{r\bar{s};\bar{p}} + F^{i\bar{j}} \chi_{i\bar{j};p\bar{p}} + \sum_{l=0}^{k-2} \left(\nabla_{\bar{p}} \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j};p} + \nabla_p \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j};\bar{p}} \right) \\
& \quad + \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l \\
& \leq \delta^2 \sum_{l=0}^{k-2} \beta_l \left(1 + \frac{1}{k-1-l} \right) F_l^{-1} |F_l^{i\bar{j}} \chi_{i\bar{j};p}|^2 + F^{i\bar{j}} \chi_{i\bar{j};p\bar{p}} \\
& \quad + \sum_{l=0}^{k-2} \left(\nabla_{\bar{p}} \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j};p} + \nabla_p \beta_l \cdot F_l^{i\bar{j}} \chi_{i\bar{j};\bar{p}} \right) + \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l \\
& = \frac{\delta^2 (k-l)}{k-1-l} \sum_{l=1}^{k-2} \beta_l F_l^{-1} \left| F_l^{i\bar{j}} \chi_{i\bar{j};p} + \frac{1}{1 + \frac{1}{k-1-l}} \frac{\nabla_p \beta_l}{\delta^2 \beta_l} F_l \right|^2 \\
& \quad - \sum_{l=0}^{k-2} \frac{1}{1 + \frac{1}{k-1-l}} \frac{|\nabla_p \beta_l|^2}{\delta^2 \beta_l} F_l + F^{i\bar{j}} \chi_{i\bar{j};p\bar{p}} + \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l \\
& \leq - \sum_{l=0}^{k-2} \frac{1}{1 + \frac{1}{k-1-l}} \frac{|\nabla_p \beta_l|^2}{\delta^2 \beta_l} F_l + F^{i\bar{j}} \chi_{i\bar{j};p\bar{p}} + \sum_{l=0}^{k-2} \nabla_{\bar{p}} \nabla_p \beta_l \cdot F_l,
\end{aligned}$$

where we used the inequality (4.2) to get the second inequality. So, our proof is completed. \square

4.2. C^2 estimate.

Lemma 4.3. *Let (M, ω) be a closed Kähler manifold of complex dimension $n \geq 2$. Suppose that χ_0 satisfies the cone condition (1.2) and $\alpha_0(z), \alpha_1(z), \dots, \alpha_{k-1}(z)$ are real C^2 functions on M satisfying (i) and (ii) in Assumption 1.1. If $u \in C^4(M)$ is a solution to the equation (2.12) with $\chi_u \in \Gamma_{k-1}(M)$, we have*

$$\sup_M |\partial\bar{\partial}u| \leq C(\sup_M |\nabla u|^2 + 1),$$

where the constant C depends on $M, \omega, \chi_0, |u|_{C^0}, \inf_M \alpha_l$ ($0 \leq l \leq k-2$) and $|\alpha_i|_{C^2}$ ($0 \leq i \leq k-1$).

Proof. For convenience, in the following argument, we write $\chi = \chi_u$ for short (suppressing the subscript u). Following the work of Hou-Ma-Wu [28], we define for $z \in M$ and a unit vector $\xi \in T_z^{(1,0)} M$

$$W(z, \xi) = \ln(\chi_{ij} \xi^i \xi^j) + \varphi(|\nabla u|^2) + \psi(u),$$

where

$$\varphi(s) = -\frac{1}{2} \log \left(1 - \frac{s}{2K} \right) \quad \text{for } 0 \leq s \leq K-1$$

and

$$\psi(t) = -A \log \left(1 + \frac{t}{2L} \right) \quad \text{for} \quad -L + 1 \leq t \leq 0.$$

Here, we set

$$K := \sup_M |\nabla u|^2 + 1, \quad L := \sup_M |u| + 1, \quad A := 2L\Lambda,$$

and Λ is a large constant that we will choose later. Clearly, φ satisfies

$$\frac{1}{2K} \geq \varphi' \geq \frac{1}{4K}, \quad \varphi'' = 2(\varphi')^2 > 0,$$

and ψ satisfies the bounds

$$2\Lambda \geq -\psi' \geq \Lambda, \quad \psi'' \geq \frac{2\varepsilon}{1-\varepsilon} (\psi')^2 \quad \text{for all} \quad \varepsilon \leq \frac{1}{2A+1}.$$

The function W must achieve its maximum at some point z in some unit direction of η . Around z , we choose a normal chart such that χ is diagonal and

$$(4.3) \quad \chi_{\eta\bar{\eta}} = \chi_{1\bar{1}} \geq \dots \geq \chi_{n\bar{n}}.$$

It follows from (2.9)

$$(4.4) \quad F^{1\bar{1}} \leq \dots \leq F^{n\bar{n}}.$$

Furthermore, we arrive at z

$$(4.5) \quad W_i = \frac{\chi_{1\bar{1};i}}{\chi_{1\bar{1}}} + \varphi' \nabla_i (|\nabla u|^2) + \psi' u_i = 0$$

and

$$(4.6) \quad \begin{aligned} W_{i\bar{i}} &= \frac{\chi_{1\bar{1};i\bar{i}}}{\chi_{1\bar{1}}} - \frac{|\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} + \varphi'' |\nabla_i (|\nabla u|^2)|^2 + \varphi' \nabla_{i\bar{i}} (|\nabla u|^2) \\ &+ \psi'' |u_i|^2 + \psi' u_{i\bar{i}} \leq 0. \end{aligned}$$

Commuting derivatives, we have the identity from Lemma 4.1

$$\chi_{1\bar{1};i\bar{i}} = \chi_{i\bar{i};1\bar{1}} + \chi_{1\bar{1}} R_{1\bar{1}i\bar{i}} - \chi_{i\bar{i}} R_{i\bar{i}1\bar{1}}.$$

Using the above inequality, we get from (4.1)

$$\begin{aligned} F^{i\bar{j}} \chi_{1\bar{1};i\bar{j}} &\geq -(1-\delta^2) F^{i\bar{j},r\bar{s}} \chi_{i\bar{j};1} \chi_{r\bar{s};\bar{1}} - C \chi_{1\bar{1}} \sum_i F^{i\bar{i}} - C F_k - C \sum_{l=0}^{k-1} |F_l| \\ &+ \nabla_{\bar{1}} \nabla_1 \beta_{k-1} - \sum_{l=0}^{k-2} \frac{1}{1 + \frac{1}{k+1-l}} \frac{|\nabla_1 \beta_l|^2}{\delta^4 \beta_l} F_l - \sum_{l=0}^{k-2} \nabla_{\bar{1}} \nabla_1 \beta_l \cdot F_l \\ &\geq -(1-\delta^2) F^{i\bar{j},r\bar{s}} \chi_{i\bar{j};1} \chi_{r\bar{s};\bar{1}} - C \chi_{1\bar{1}} \sum_i F^{i\bar{i}} - C, \end{aligned}$$

where we used the equality

$$F^{i\bar{i}}\chi_{i\bar{i}} = F_k + \sum_{l=0}^{k-2} (l-k+1)\beta_l F_l$$

and the fact that F_k and F_l ($0 \leq l \leq k-2$) are bounded (see (2.21) and (2.22)).

Multiplying (4.6) by $F^{i\bar{i}}$ and summing it over index i , we can know from the above inequality

$$\begin{aligned} 0 &\geq -\frac{F^{i\bar{i}}|\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - (1-\delta^2)\frac{1}{\chi_{1\bar{1}}}F^{i\bar{j},r\bar{s}}\chi_{i\bar{j};1}\chi_{r\bar{s};\bar{1}} \\ &\quad + \varphi''F^{i\bar{i}}|\nabla_i(|\nabla u|^2)|^2 + \varphi'F^{i\bar{i}}\nabla_{\bar{i}}\nabla_i(|\nabla u|^2) + \psi''F^{i\bar{i}}|u_i|^2 + \psi'F^{i\bar{i}}u_{i\bar{i}} \\ (4.7) \quad &-C\sum_i F^{i\bar{i}} - \frac{C}{\chi_{1\bar{1}}}. \end{aligned}$$

To proceed, we need the following calculation

$$\nabla_i(|\nabla u|^2) = \sum_j (-u_j\chi_{0i\bar{j}} + u_{j\bar{i}}u_{\bar{j}}) + u_i\chi_{ii}$$

and

$$\begin{aligned} \nabla_{\bar{i}}\nabla_i(|\nabla u|^2) &= \sum_j (\chi_{0j\bar{i}}\chi_{0i\bar{j}} + u_{j\bar{i}}u_{\bar{j}} - \chi_{0i\bar{i};j}u_{\bar{j}} - u_j\chi_{0i\bar{i};\bar{j}}) + \sum_{j,k} R_{i\bar{i}j\bar{k}}u_k u_{\bar{j}} \\ &\quad + \chi_{i\bar{i}}^2 - 2\chi_{0i\bar{i}}\chi_{i\bar{i}} + 2\sum_j \operatorname{Re}\{\chi_{i\bar{i};j}u_{\bar{j}}\} \\ &\geq -2\sum_j \operatorname{Re}\{\chi_{0i\bar{i};j}u_{\bar{j}}\} + \sum_{j,k} R_{i\bar{i}j\bar{k}}u_k u_{\bar{j}} + \frac{1}{2}\chi_{i\bar{i}}^2 - 2\chi_{0i\bar{i}}^2 \\ &\quad + 2\sum_j \operatorname{Re}\{\chi_{i\bar{i};j}u_{\bar{j}}\}, \end{aligned}$$

where $\operatorname{Re}\{z\}$ denotes the real part of the complex number z . Thus, we can estimate the term

$$\begin{aligned} &\varphi' \sum_i F^{i\bar{i}}\nabla_{\bar{i}}\nabla_i(|\nabla u|^2) \\ &\geq -C\frac{|\nabla u| + |\nabla u|^2}{K} \sum_i F^{i\bar{i}} + \frac{1}{2}\varphi' \sum_i F^{i\bar{i}}\chi_{i\bar{i}}^2 - 2\varphi' \sum_i F^{i\bar{i}}\chi_{0i\bar{i}}^2 \\ &\quad - 2\varphi' \sum_j \operatorname{Re}\left\{\sum_{l=0}^{k-1} (\nabla_j\beta_l \cdot F_l)u_{\bar{j}}\right\} \\ (4.8) \quad &\geq -C\frac{1 + |\nabla u| + |\nabla u|^2}{K} \sum_i F^{i\bar{i}} + \frac{1}{2}\varphi' \sum_i F^{i\bar{i}}\chi_{i\bar{i}}^2. \end{aligned}$$

where we used the following equality

$$(4.9) \quad F^{i\bar{j}} \chi_{i\bar{j};p} = - \sum_{l=0}^{k-1} \nabla_p \beta_l \cdot F_l$$

to get the first inequality and (4.9) can be directly deduced by (2.12).

Taking the inequality (4.8) into (4.7), it yields

$$(4.10) \quad \begin{aligned} 0 &\geq -\frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - (1 - \delta^2) \frac{1}{\chi_{1\bar{1}}} F^{i\bar{j}, r\bar{s}} \chi_{i\bar{j};1} \chi_{r\bar{s};\bar{1}} \\ &\quad + \varphi'' F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 + \frac{\varphi'}{2} F^{i\bar{i}} \chi_{i\bar{i}}^2 + \psi'' F^{i\bar{i}} |u_i|^2 + \psi' F^{i\bar{i}} u_{i\bar{i}} \\ &\quad - C \sum_i F^{i\bar{i}} - C. \end{aligned}$$

Now, we divide our proof into two cases separately, depending on whether $\chi_{n\bar{n}} < -\delta \chi_{1\bar{1}}$ or not, for a small δ to be chosen later.

Case 1. $\chi_{n\bar{n}} < -\delta \chi_{1\bar{1}}$. In this case, it follows that $\chi_{1\bar{1}}^2 \leq \frac{1}{\delta^2} \chi_{n\bar{n}}^2$. So, we only need to bound $\chi_{n\bar{n}}^2$. Clearly, we can obtain if we throw some positive terms in (4.10)

$$(4.11) \quad \begin{aligned} 0 &\geq -\frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} + \varphi'' F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 + \frac{\varphi'}{2} F^{i\bar{i}} \chi_{i\bar{i}}^2 + \psi' F^{i\bar{i}} u_{i\bar{i}} \\ &\quad - C \sum_i F^{i\bar{i}} - C. \end{aligned}$$

Note that we get from (2.14)

$$(4.12) \quad \begin{aligned} \psi' \sum_i F^{i\bar{i}} u_{i\bar{i}} &= \psi' \sum_i F^{i\bar{i}} (\chi_{i\bar{i}} - (\chi_{0i\bar{i}} - \tau) - \tau) \\ &= \psi' \left(F_k + \sum_{l=0}^{k-1} (l - k + 1) \beta_l F_l - \sum_i F^{i\bar{i}} (\chi_{0i\bar{i}} - \tau) - \tau \sum_i F^{i\bar{i}} \right) \\ &\geq -C\Lambda + \tau\Lambda \sum_i F^{i\bar{i}} \end{aligned}$$

Plugging (4.12) into (4.11), we get by choosing Λ large enough

$$(4.13) \quad \varphi'' F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 + \frac{\varphi'}{2} F^{i\bar{i}} \chi_{i\bar{i}}^2 \leq \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} + C(1 + \Lambda).$$

In addition, we have from (4.5)

$$(4.14) \quad \begin{aligned} \sum_i \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} &= \sum_i F^{i\bar{i}} |\varphi' \nabla_i (|\nabla u|^2) + \psi' u_i|^2 \\ &\leq 2(\varphi')^2 \sum_i F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 + 8\Lambda^2 K \sum_i F^{i\bar{i}}. \end{aligned}$$

Moreover, we obtain in view of (4.3) and (4.4)

$$(4.15) \quad \sum_i F^{i\bar{i}} \chi_{i\bar{i}}^2 \geq F^{n\bar{n}} \chi_{n\bar{n}}^2 \geq \frac{1}{n} \chi_{n\bar{n}}^2 \sum_i F^{i\bar{i}}.$$

Substituting (4.14) and (4.15) into (4.13), we get

$$\frac{1}{K} \chi_{n\bar{n}}^2 \sum_i F^{i\bar{i}} \leq 8\Lambda^2 K \sum_i F^{i\bar{i}} + C(1 + \Lambda).$$

Then, it follows that by (2.23)

$$\chi_{1\bar{1}} \leq CK.$$

Case 2. $\chi_{n\bar{n}} \geq -\delta\chi_{1\bar{1}}$. Define the set

$$I = \left\{ i \in \{1, 2, \dots, n\} : F^{i\bar{i}} > \delta^{-1} F^{1\bar{1}} \right\}.$$

It follows from (2.8)

$$(4.16) \quad \begin{aligned} -\frac{1}{\chi_{1\bar{1}}} F^{i\bar{j}, r\bar{s}} \chi_{i\bar{j};1} \chi_{r\bar{s};\bar{1}} &\geq \frac{1-\delta}{1+\delta} \frac{1}{\chi_{1\bar{1}}^2} \sum_{i \in I} F^{i\bar{i}} |\chi_{i\bar{1};1}|^2 \\ &\geq \frac{1-\delta}{1+\delta} \frac{1}{\chi_{1\bar{1}}^2} \sum_{i \in I} F^{i\bar{i}} \left(|\chi_{1\bar{1};i}|^2 + 2\operatorname{Re}\{\chi_{1\bar{1};i} \bar{b}_i\} \right), \end{aligned}$$

where $b_i = \chi_{0i\bar{1};1} - \chi_{0\bar{1};i}$.

Using (4.5), we have in view of the fact $\varphi'' = 2(\varphi')^2$

$$(4.17) \quad \begin{aligned} &\varphi'' \sum_{i \in I} F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 \\ &\geq 2 \sum_{i \in I} F^{i\bar{i}} \left(\delta \left| \frac{\chi_{1\bar{1};i}}{\chi_{1\bar{1}}} \right|^2 - \frac{\delta}{1-\delta} |\psi' u_i|^2 \right). \end{aligned}$$

Choosing $\delta \leq \frac{1}{2A+1}$, we have $\psi'' \geq \frac{2\delta}{1-\delta}(\psi')^2$. Then, we get by combining (4.16) and (4.17)

$$\begin{aligned}
& - \sum_{i \in I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - (1-\delta^2) \frac{1}{\chi_{1\bar{1}}} F^{i\bar{j},r\bar{s}} \chi_{i\bar{j};1} \chi_{r\bar{s};\bar{1}} \\
& + \varphi'' \sum_{i \in I} F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 + \psi'' \sum_{i \in I} F^{i\bar{i}} |u_i|^2 \\
\geq & \delta^2 \sum_{i \in I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} + \frac{2(1-\delta^2)(1-\delta)}{1+\delta} \frac{1}{\chi_{1\bar{1}}^2} \sum_{i \in I} F^{i\bar{i}} \operatorname{Re}\{\chi_{1\bar{1};i} \bar{b}_i\} \\
\geq & \delta^2 \sum_{i \in I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - 2(1-\delta)^2 \frac{1}{\chi_{1\bar{1}}^2} \sum_{i \in I} F^{i\bar{i}} \operatorname{Re}\{\chi_{1\bar{1};i} \bar{b}_i\} \\
\geq & \frac{\delta^2}{2} \sum_{i \in I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - C \frac{4(1-\delta)^4}{\delta^2} \frac{1}{\chi_{1\bar{1}}^2} \sum_{i \in I} F^{i\bar{i}} \\
(4.18) \geq & \frac{\delta^2}{2} \sum_{i \in I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} - C \sum_{i \in I} F^{i\bar{i}},
\end{aligned}$$

where we choose $\chi_{1\bar{1}}$ large enough to get the last inequality.

For the terms without an index in I , we have by (4.14)

$$\begin{aligned}
& - \sum_{i \notin I} \frac{F^{i\bar{i}} |\chi_{1\bar{1};i}|^2}{\chi_{1\bar{1}}^2} + \varphi'' \sum_{i \notin I} F^{i\bar{i}} |\nabla_i (|\nabla u|^2)|^2 \\
\geq & -8\Lambda^2 K \sum_{i \notin I} F^{i\bar{i}} \\
(4.19) \geq & -\frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}},
\end{aligned}$$

where we used the fact $1 \notin I$ to get the last inequality.

Substituting (4.18) and (4.19) into (4.10)

$$\begin{aligned}
& C + C \sum_i F^{i\bar{i}} + \frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}} \\
(4.20) \geq & \frac{1}{8K} \sum_i F^{i\bar{i}} \chi_{i\bar{i}}^2 + \psi' \sum_i F^{i\bar{i}} u_{i\bar{i}}.
\end{aligned}$$

Suppose that $\chi_{1\bar{1}} \geq N$. Since the cone condition (1.2) is equivalent to the condition (2.11) for μ being the eigenvalues of χ_0 , Lemma 2.7 works. Thus we will divide the following argument into two cases:

Case A: If (2.16) hold, we have

$$\sum_i F^{i\bar{i}} u_{i\bar{i}} \leq -\theta - \theta \sum_i F^{i\bar{i}},$$

which implies

$$(4.21) \quad \psi' \sum_i F^{i\bar{i}} u_{i\bar{i}} \geq \Lambda \theta (1 + \sum_i F^{i\bar{i}}).$$

Substituting (4.21) into (4.20)

$$\begin{aligned} & C + C \sum_i F^{i\bar{i}} + \frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}} \\ & \geq \frac{1}{8K} \sum_i F^{i\bar{i}} \chi_{i\bar{i}}^2 + \Lambda \theta (1 + \sum_i F^{i\bar{i}}), \end{aligned}$$

which implies if we choose Λ large enough

$$\frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}} \geq \frac{1}{8K} \sum_i F^{i\bar{i}} \chi_{i\bar{i}}^2 \geq \frac{1}{8K} F^{1\bar{1}} \chi_{1\bar{1}}^2.$$

Thus,

$$\chi_{1\bar{1}}^2 \leq \frac{64n\Lambda^2 K^2}{\delta}.$$

Case B: Otherwise, we have by (2.17)

$$(4.22) \quad F^{1\bar{1}} \chi_{1\bar{1}} \geq \theta.$$

Thus, we get from (4.12)

$$(4.23) \quad \begin{aligned} & C(1 + \Lambda) + C \sum_i F^{i\bar{i}} + \frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}} \\ & \geq \frac{1}{8K} F^{1\bar{1}} \chi_{1\bar{1}}^2 + \tau \Lambda \sum_i F^{i\bar{i}}. \end{aligned}$$

Choosing Λ large enough, it follows from (4.23)

$$(4.24) \quad C(1 + \Lambda) + \frac{8n\Lambda^2 K}{\delta} F^{1\bar{1}} \geq \frac{1}{8K} F^{1\bar{1}} \chi_{1\bar{1}}^2.$$

Then, using (4.22), we choose $\chi_{1\bar{1}}$ large enough such that

$$\frac{1}{16K} F^{1\bar{1}} \chi_{1\bar{1}}^2 \geq \frac{1}{16K} \theta \chi_{1\bar{1}} \geq C(1 + \Lambda).$$

Taking the above inequality into (4.24), we arrive

$$\chi_{1\bar{1}} \leq \sqrt{\frac{128n}{\theta}} \Lambda K.$$

So, we complete the proof. \square

5. THE GRADIENT ESTIMATE

We will derive the gradient estimate using a blow-up argument and Liouville-type theorem due to Dinew-Kolodziej [14]. Suppose the gradient estimate fails. Then, there exists a sequence solutions $u_m \in C^4(M)$ to the equation (2.12) such that

$$N_m = \sup_M |\nabla u_m| \rightarrow +\infty, \quad \text{as } m \rightarrow +\infty.$$

Clearly,

$$(5.1) \quad \chi_{u_m}^k \wedge \omega^{n-k} = \sum_{l=0}^{k-1} \alpha_l(z) \chi_{u_m}^l \wedge \omega^{n-l}, \quad \sup_M u_m = 0.$$

For each m , we assume that $|\nabla u_m|$ attains its maximum value at $z_m \in M$. Then, after passing a subsequence, z_m converges some point $z \in M$. Choosing a normal coordinate chart around z , which we identify with an open set in \mathbb{C}^n with coordinates (z^1, \dots, z^n) , and such that $\omega(0) = \omega_0 := \frac{\sqrt{-1}}{2} \delta_{ij} dz^i \wedge d\bar{z}^j$. Without loss of generality, we may assume that the open set contains $B_1(0)$. Set

$$v_m(z) = u\left(\frac{z}{N_m}\right), \quad z \in B_{N_m}(0).$$

Then, we know from C^2 estimate

$$|v_m|_{C^2(B_{N_m}(0))} \leq C.$$

By passing to a subsequence again, we can assume v_m is $C^{1,\alpha}$ convergent to a limit function $v \in C^{1,\alpha}$ with $\nabla v(0) = 1$. Then, we have from (5.1)

$$\left[\chi\left(\frac{z}{N_m}\right) \right]^k \wedge \left[\omega\left(\frac{z}{N_m}\right) \right]^{n-k} = \sum_{l=0}^{k-1} \alpha_l\left(\frac{z}{N_m}\right) \left[\chi\left(\frac{z}{N_m}\right) \right]^l \wedge \left[\omega\left(\frac{z}{N_m}\right) \right]^{n-l},$$

which results in

$$\begin{aligned} & \left[\chi_0\left(\frac{z}{N_m}\right) + N_m^2 \frac{\sqrt{-1}}{2} \partial\bar{\partial}v_m \right]^k \wedge \left[\omega\left(\frac{z}{N_m}\right) \right]^{n-k} \\ &= \sum_{l=0}^{k-1} \alpha_l\left(\frac{z}{N_m}\right) \left[\chi_0\left(\frac{z}{N_m}\right) + N_m^2 \frac{\sqrt{-1}}{2} \partial\bar{\partial}v_m \right]^l \wedge \left[\omega\left(\frac{z}{N_m}\right) \right]^{n-l}. \end{aligned}$$

Thus,

$$\begin{aligned} & N_m^{2k} \left[O\left(\frac{1}{N_m^2}\right) \omega_0 + \frac{\sqrt{-1}}{2} \partial\bar{\partial}v_m \right]^k \wedge \left[\omega_0 + O\left(\frac{|z|^2}{N_m^2}\right) \omega_0 \right]^{n-k} \\ &= \sum_{l=0}^{k-1} \alpha_l\left(\frac{z}{N_m}\right) N_m^{2l} \left[O\left(\frac{1}{N_m^2}\right) \omega_0 + \frac{\sqrt{-1}}{2} \partial\bar{\partial}v_m \right]^l \wedge \left[\omega_0 + O\left(\frac{|z|^2}{C_m^2}\right) \omega_0 \right]^{n-l}. \end{aligned}$$

Therefore, we have

$$(\sqrt{-1} \partial\bar{\partial}v)^k \wedge \omega_0^{n-k} = 0,$$

which is in the pluripotential sense. Moreover, we have for any $1 \leq l \leq k$ by a similar reasoning

$$(\sqrt{-1}\partial\bar{\partial}v)^l \wedge \omega_0^{n-l} \geq 0.$$

Thus, the limiting function v is k -subharmonic. A result of Blocki [2] tell us that v is a maximal k -subharmonic function in \mathbb{C}^n . Then the Liouville theorem in [14] implies that v is a constant, which contradicts the fact $\nabla v(0) = 1$.

6. THE PROOF OF MAIN THEOREM

To get higher-order estimates, we need to show the equation (2.12) is uniformly elliptic. First, we recall the following lemma.

Lemma 6.1. *Under the assumption in Theorem 1.1, we have*

$$(6.1) \quad \sigma_{k-1}(\chi_u) \geq C > 0.$$

Proof. Without loss of generality, we assume there exists $0 \leq l_0 \leq k-2$ such that $\beta_{l_0}(z) > 0$. Then, we get from the inequality (2.21)

$$0 < \frac{\sigma_{l_0}(\chi_u)}{\sigma_{k-1}(\chi_u)} \leq C.$$

Thus, we complete the proof by using the generalized Newton-MacLaurin inequality (2.4) □

Thus, using (6.1) and the uniform estimates up to second order by Lemmas in the previous sections, we can deduce that the equation (2.12) is uniformly elliptic. Then, higher-order estimates follow from the Evans-Krylov theorem [16, 29] and the Schauder estimate (see also [51]). Next, we shall give a proof of Theorem 1.1 by the continuity method. The obstacle is that we have to find a uniform cone condition for the solution flow of the continuity method.

we define $\gamma(z)$ by

$$\chi_0^k \wedge \omega^{n-k} - \sum_{l=1}^{k-2} \alpha_l \chi_0^l \wedge \omega^{n-l} = \gamma(z) \chi_0^{k-1} \wedge \omega^{n-k+1}$$

It is easy to see from (2.6) and (2.7)

$$(6.2) \quad k\chi_0^{k-1} \wedge \omega^{n-k} > \sum_{l=1}^{k-2} l\alpha_l \chi_0^{l-1} \wedge \omega^{n-l} + (k-1)\gamma\chi_0^{k-2} \wedge \omega^{n-k+1}.$$

Set

$$\tilde{\alpha}_{k-1}(z) = \max\{\alpha_{k-1}(z), \gamma(z)\}.$$

Then, χ_0 satisfies the cone condition by combining (6.2) and (1.2)

$$(6.3) \quad k\chi_0^{k-1} \wedge \omega^{n-k} > \sum_{l=1}^{k-2} l\alpha_l \chi_0^{l-1} \wedge \omega^{n-l} + (k-1)\tilde{\alpha}_{k-1}\chi_0^{k-2} \wedge \omega^{n-k+1}.$$

We will apply the continuous method introduced in [49, 43, 45] to complete my proof. First, we consider the family of the equations for $t \in [0, 1]$

$$(6.4) \quad \chi_{u_t}^k \wedge \omega^{n-k} = \sum_{l=0}^{k-2} \alpha_l(z) \chi_{u_t}^l \wedge \omega^{n-l} + \left[(1-t)\gamma(z) + t\tilde{\alpha}_{k-1}(z) + a_t \right] \chi_{u_t}^{k-1} \wedge \omega^{n-k+1},$$

where $\chi_{u_t} \in \Gamma_{k-1}(M)$ and a_t is a constant for each t . We consider

$$\mathcal{T}_1 = \{s \in [0, 1] : \exists u_t \in C^{2,\alpha}(M) \text{ and } a_t \text{ solving (6.4) for } t \in [0, s]\}$$

Clearly, $0 \in \mathcal{T}_1$ and $a_0 = 0$.

Lemma 6.2. \mathcal{T}_1 is open.

Proof. Rewriting the equations (6.4) as the form

$$\begin{aligned} \log H(u_t, z, t) &:= \frac{\sigma_k(\chi_{u_t})}{\sigma_{k-1}(\chi_{u_t})} - \sum_{l=0}^{k-2} \beta_l \frac{\sigma_l(\chi_{u_t})}{\sigma_{k-1}(\chi_{u_t})} \\ &= \frac{n-k+1}{k} \left[(1-t)\gamma(z) + t\tilde{\alpha}_{k-1}(z) + a_t \right]. \end{aligned}$$

In the following, we will write $H(u_t) = H(u_t, z, t)$ for short (suppressing the subscript z, t). Now we follow the idea in [49, 43] to show \mathcal{T}_1 is open. Assume $\hat{t} \in \mathcal{T}_1$, we need to show that there exists small $\varepsilon > 0$ such that $t \in \mathcal{T}_1$ for any $t \in [\hat{t}, \hat{t} + \varepsilon)$.

Define a new Hermitian metric corresponding to u_t

$$\Omega_t := \frac{\sqrt{-1}}{2} H(u_t) H_{i\bar{j}}(u_t) dz^i \wedge dz^{\bar{j}}$$

and

$$\hat{\Omega} = \Omega_{\hat{t}} := \frac{\sqrt{-1}}{2} H(u_{\hat{t}}) H_{i\bar{j}}(u_{\hat{t}}) dz^i \wedge dz^{\bar{j}},$$

where $H^{i\bar{j}} = \frac{\partial H}{\partial u_{i\bar{j}}}$ and $\{H_{i\bar{j}}\}$ is the inverse of $H^{i\bar{j}}$. Using Gauduchon's theorem [20] to $\hat{\Omega}$, there exists a function \hat{f} such that $\hat{\Omega}_G = e^{\hat{f}} \hat{\Omega}$ is Gauduchon, i.e., $\partial\bar{\partial}(\hat{\Omega}_G^{n-1}) = 0$. We may assume that by adding a constant to \hat{f}

$$\int_M e^{(n-1)\hat{f}} \hat{\Omega}^n = 1.$$

We will show that we can find $u_t \in C^{2,\alpha}(M)$ for $t \in [\hat{t}, \hat{t} + \varepsilon)$ solving

$$(6.5) \quad \frac{H(u_t)}{H(u_{\hat{t}})} = \left(\int_M \frac{H(u_t)}{H(u_{\hat{t}})} e^{(n-1)\hat{f}\widehat{\Omega}^n} \right) e^{\frac{n-k+1}{k} \left[(\hat{t}-t)\gamma(z) + (t-\hat{t})\tilde{\alpha}_{k-1}(z) + a_t - a_{\hat{t}} \right]},$$

where a_t is chosen such that

$$\int_M e^{\frac{n-k+1}{k} \left[(\hat{t}-t)\gamma(z) + (t-\hat{t})\tilde{\alpha}_{k-1}(z) + a_t - a_{\hat{t}} \right]} e^{(n-1)\hat{f}\widehat{\Omega}^n} = 1.$$

If u_t is a solution to (6.5), $u_t + c$ is also for any $c \in \mathbb{R}$. Thus, we normalized u_t such that

$$\int_M u_t e^{(n-1)\hat{f}\widehat{\Omega}^n} = 0.$$

Moreover, we define two Banach manifolds \mathcal{B}_1 and \mathcal{B}_2 by

$$\mathcal{B}_1 := \left\{ \xi \in C^{2,\alpha}(M) : \int_M \xi e^{(n-1)\hat{f}\widehat{\Omega}^n} = 0 \right\}$$

and

$$\mathcal{B}_2 := \left\{ \eta \in C^\alpha(M) : \int_M e^\eta e^{(n-1)\hat{f}\widehat{\Omega}^n} = 1 \right\}.$$

It is easy to see the tangent spaces of \mathcal{B}_1 and \mathcal{B}_2 at 0:

$$T_0\mathcal{B}_1 = \mathcal{B}_1 \quad \text{and} \quad T_0\mathcal{B}_2 = \left\{ \rho \in C^\alpha(M) : \int_M \rho e^{(n-1)\hat{f}\widehat{\Omega}^n} = 0 \right\}.$$

Define a linear operator $L : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ by

$$L(\xi) := \log H(u_{\hat{t}} + \xi) - \log H(u_{\hat{t}}) - \log \left(\int_M \frac{H(u_{\hat{t}} + \xi)}{H(u_{\hat{t}})} e^{(n-1)\hat{f}\widehat{\Omega}^n} \right).$$

Observe that $L(0) = 0$. By the inverse function theorem, to get openness of the equation (6.5) at $\xi = 0$ we just need to show that

$$(DL)_0 : T_0\mathcal{B}_1 \rightarrow T_0\mathcal{B}_2$$

is invertible. Compute

$$\begin{aligned} (DL)_0(\xi) &= \frac{1}{H(u_{\hat{t}})} H^{i\bar{j}}(u_{\hat{t}}) \xi_{ij} - \int_M \frac{1}{H(u_{\hat{t}})} H^{i\bar{j}}(u_{\hat{t}}) \xi_{ij} e^{(n-1)\hat{f}\widehat{\Omega}^n} \\ &= \Delta_{\widehat{\Omega}} \xi - n \int_M e^{(n-1)\hat{f}\widehat{\Omega}^n} \wedge \partial\bar{\partial}\xi \\ &= \Delta_{\widehat{\Omega}} \xi. \end{aligned}$$

where we are using the fact that $\partial\bar{\partial}(\widehat{\Omega}_G^{n-1}) = 0$. Recall that we can solve the equation $\Delta_{\widehat{\Omega}_G} u = v$ as long as $\int_M v \widehat{\Omega}_G^n = 0$ (see e.g. [3]). Given $\rho \in T_0\mathcal{B}_2$

we see that

$$\int_M \rho e^{(n-1)\hat{f}} \hat{\Omega}^n = \int_M \rho e^{-\hat{f}} \hat{\Omega}_G^n = 0.$$

Then there exists η solves

$$e^{-\hat{f}} \Delta_{\hat{\Omega}} \eta = \Delta_{\hat{\Omega}_G} \eta = \rho e^{-\hat{f}},$$

which implies

$$\Delta_{\hat{\Omega}} \eta = \rho,$$

as required. This shows that we can find u_t solving (6.5) for $t \in [\hat{t}, \hat{t} + \varepsilon)$. Thus \mathcal{T}_1 is open. \square

Lemma 6.3. \mathcal{T}_1 is closed.

Proof. It suffices to show (i) the uniform bound for a_t and (ii) the uniform C^∞ estimates for all u_t . In fact, at the maximum point of u_t , using the concavity of $(\frac{\sigma_k}{\sigma_l})^{\frac{1}{k-l}}$ (see Proposition 5), we have

$$\frac{\chi_0^k \wedge \omega^{n-k}}{\chi_0^l \wedge \omega^{n-l}} \geq \frac{\chi_{u_t}^k \wedge \omega^{n-k}}{\chi_{u_t}^l \wedge \omega^{n-l}}, \quad 0 \leq l \leq k-1.$$

Thus, we get from (6.2) and (6.3)

$$\begin{aligned} \chi_0^k \wedge \omega^{n-k} &\geq \sum_{l=0}^{k-2} \alpha_l(z) \chi_0^l \wedge \omega^{n-l} + \left[(1-t)\gamma(z) + t\tilde{\alpha}_0(z) + a_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1} \\ &\geq \sum_{l=0}^{k-2} \alpha_l(z) \chi_0^l \wedge \omega^{n-l} + \left[\gamma(z) + a_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1}. \end{aligned}$$

Thus, $a_t \leq 0$. Similarly, at the minimum point of u_t , we have

$$\begin{aligned} \chi_0^k \wedge \omega^{n-k} &\leq \sum_{l=0}^{k-2} \alpha_l(z) \chi_0^l \wedge \omega^{n-l} + \left[(1-t)\gamma(z) + t\tilde{\alpha}_0(z) + a_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1} \\ &\leq \sum_{l=1}^{k-1} \alpha_l(z) \chi_0^l \wedge \omega^{n-l} + \left[\tilde{\alpha}_0(z) + a_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1}. \end{aligned}$$

Therefore, a_t is uniformly bounded from below. Thus, the requirement (i) is met. For the requirement (ii), it is sufficient to show the cone condition

$$\begin{aligned} k\chi_0^{k-1} \wedge \omega^{n-k} &> \sum_{l=1}^{k-2} l\alpha_l \chi_0^{l-1} \wedge \omega^{n-l} \\ &\quad + (k-1) \left[(1-t)\gamma(z) + t\tilde{\alpha}_0(z) + a_t \right] \chi_0^{k-2} \wedge \omega^{n-k+1}. \end{aligned}$$

is uniform for the equations (6.4). This fact can be easily deduced in view of $a_t \leq 0$, (6.2) and (6.3). \square

Therefore, there exists a solution to the equation

$$\chi_v^k \wedge \omega^{n-k} = \sum_{l=0}^{k-2} \alpha_l(z) \chi_v^l \wedge \omega^{n-l} + [\tilde{\alpha}_{k-1}(z) + a_1] \chi_0^{k-1} \wedge \omega^{n-k+1}.$$

Second, we consider the family of equations

$$(6.6) \chi_{u_t}^k \wedge \omega^{n-k} = \sum_{l=0}^{k-2} \alpha_l(z) \chi_{u_t}^l \wedge \omega^{n-l} + \left[(1-t)\tilde{\alpha}_{k-1}(z) + t\alpha_{k-1}(z) + b_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1}.$$

where $\chi_{u_t} \in \Gamma_{k-1}(M)$ and b_t is a constant for each t . We consider

$$\mathcal{T}_2 = \{s \in [0, 1] : \exists u_t \in C^{2,\alpha}(M) \text{ and } b_t \text{ solving (6.6) for } t \in [0, s]\}$$

Clearly, $0 \in \mathcal{T}_2$ with $u_0 = v$ and $b_0 = a_1$. The openness of \mathcal{T}_2 and uniform bound for b_t can be similarly deduced by the previous argument for (6.4). Moreover, from the argument for \mathcal{T}_1 , we know it suffices to show the cone condition holds for the equations (6.6) in order to guarantee that the continuity method works.

Integrating (6.6) on M , we get from the condition (1.3)

$$\begin{aligned} & \int_M \chi_0^k \wedge \omega^{n-k} \\ &= \int_M \chi_{tu}^k \wedge \omega^{n-k} \\ &= \sum_{l=0}^{k-2} \int_M \alpha_l \chi_{tu}^l \wedge \omega^{n-l} + \int_M \left[(1-t)\tilde{\alpha}_{k-1} + t\alpha_{k-1} + b_t \right] \chi_0^{k-1} \wedge \omega^{n-k+1} \\ &\geq \sum_{l=0}^{k-2} c_{k,l} \int_M \chi_0^l \wedge \omega^{n-l} + (c_{k,k-1} + b_t) \int_M \chi_0^{k-1} \wedge \omega^{n-k+1} \\ &\geq \int_M \chi_0^k \wedge \omega^{n-k} + b_t \int_M \chi_0^{k-1} \wedge \omega^{n-k+1}. \end{aligned}$$

Thus, $b_t \leq 0$. So the cone condition holds for the equations (6.6). Therefore, we complete the proof of Theorem 1.1.

REFERENCES

- [1] B. Andrews: Contraction of convex hypersurfaces in Euclidean space. *Calc. Var. Partial Differ. Equ.* 2, 151-171 (1994)
- [2] Z. Blocki: Weak solutions to the complex Hessian equation. *Ann. Inst. Fourier (Grenoble)* 55 (2005), no. 5, 1735-1756.
- [3] N. Buchdahl: On compact Kähler surfaces. *Ann. Inst. Fourier (Grenoble)* 49 (1999), no.1, 287-302.
- [4] L. Caffarelli, L. Nirenberg, J. Spruck: Dirichlet problem for nonlinear second order elliptic equations III, Functions of the eigenvalues of the Hessian. *Acta Math.*, 155(1985), 261-301.

- [5] C. Q. Chen, L. Chen, X. Q. Mei, N. Xiang: The Classical Neumann Problem for a class of mixed Hessian equations. To appear in *Studies in Applied Mathematics* (2021).
- [6] C. Q. Chen, L. Chen, X. Q. Mei, N. Xiang: The Neumann problem for a class of mixed complex Hessian equations. Preprint, 2019.
- [7] L. Chen, X. Guo, Y. He: A class of fully nonlinear equations arising in conformal geometry. To appear in *Int. Math. Res. Not.*, (2021), <https://doi.org/10.1093/imrn/rnaa253>.
- [8] L. Chen, A. G. Shang, Q. Tu: A class of prescribed Weingarten curvature equations in Euclidean space. *Comm. Partial Differential Equations* 46 (2021), no. 7, 1326-1343.
- [9] X. X. Chen: On the lower bound of the Mabuchi energy and its application. *Int. Math. Res. Not.* 12, 607-623 (2000)
- [10] X. X. Chen: A new parabolic flow in Kähler manifolds. *Comm. Anal. Geom.* 12 (2004), no. 4, 837-852.
- [11] K. S. Chou, X.J. Wang, A variational theory of the Hessian equation, *Commun. Pure Appl. Math.* 54 (9) (2001) 1029-1064.
- [12] P. Cherrier: équations de Monge-Ampère sur les variétés Hermitiennes compactes. *Bull. Sci. Math.* (2), 111 (1987), 343-385.
- [13] T. Collins, G. Székelyhidi: Convergence of the J -flow on toric manifolds. *J. Differential Geom.* 107 (2017) no. 1, 47-81.
- [14] S. Dinew, S. Kolodziej: Liouville and Calabi-Yau type theorems for complex Hessian equations. *Amer. J. Math.*, 139(2), 403-415 (2017).
- [15] S. Donaldson: Moment maps and diffeomorphisms. *Asian J. Math.* 3 (1999), no. 1, 1-15.
- [16] L. C. Evans: Classical solutions of fully nonlinear, convex, second-order elliptic equations. *Comm. Pure Appl. Math.* 35, no. 3(1982), 333-363.
- [17] H. Fang, M. J. Lai, X. N. Ma: On a class of fully nonlinear flows in Kähler geometry. *J. Reine Angew. Math.*, 653 (2011), 189-220.
- [18] J. X. Fu, S. T. Yau: The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampère equation. *J. Differential Geom.*, Vol 78, Number 3 (2008), 369-428.
- [19] J. X. Fu, S.T. Yau: A Monge-Ampère type equation motivated by string theory. *Comm. Anal. Geom.* 15 (2007), no. 1, 29-76.
- [20] P. Gauduchon: Le théorème de l'excentricité nulle. *C. R. Acad. Sci. Paris* 285, 387-390 (1977).
- [21] D. Gilbarg, N.S. Trudinger: Elliptic partial differential equations of second order. Reprint of the 1998 edition. *Classics in Mathematics*. Springer-Verlag, Berlin, 2001.
- [22] B. Guan, Q. Li: Complex Monge-Ampère equations and totally real submanifolds. *Adv. Math.* 225 (2010), no. 3, 1185-1223.
- [23] B. Guan, Q. Li: A Monge-Ampère type fully nonlinear equation on Hermitian manifolds. *Discrete Contin. Dyn. Syst. Ser. B* 17 (2012) 1991-1999.
- [24] B. Guan, W. Sun: On a class of fully nonlinear elliptic equations on Hermitian manifolds. *Calc. Var. Partial Differential Equations* 54 (2015), no. 1, 901-916.
- [25] P. F. Guan, X. W. Zhang: A class of curvature type equations. *Pure and Applied Math Quarterly*, Vol. 17, No. 3 (2021), 865-907. [arXiv:1909.03645](https://arxiv.org/abs/1909.03645).
- [26] A. Hanani, Équations du type de Monge-Ampère sur les variétés Hermitiennes compactes. *J. Funct. Anal.*, 137 (1996), 49-75.
- [27] R. Harvey, H.B. Lawson: Calibrated geometries. *Acta Math.*, 148 (1982), 47-157.
- [28] Z. Hou, X.N. Ma, D. M. Wu: A second order estimate for complex Hessian equations on a compact Kähler manifold. *Math. Res. Lett.* 17 (2010), no. 3, 547-561.
- [29] N. V. Krylov: Boundedly inhomogeneous elliptic and parabolic equations in a domain. *Izv. Akad. Nauk SSSR Ser. Mat.* 47,no 1 (1983), 75-108.

- [30] N. V. Krylov: On the general notion of fully nonlinear second order elliptic equation. Trans. Amer. Math. Soc. 347 (3),(1995), 857-895.
- [31] G. Lieberman: Second order parabolic differential equations. World Scientific, 1996.
- [32] M. Lin, N. S. Trudinger: On some inequalities for elementary symmetric functions. Bull. Aust. Math. Soc. 50(1994), 317-326.
- [33] Vamsi P. Pingali: A fully nonlinear generalized Monge-Ampère PDE on a torus. Electron. J. Differential Equations 2014, No. 211, 8 pp.
- [34] Vamsi P. Pingali: A generalised Monge-Ampère equation. J. Partial Differ. Equ. 27 (2014), no. 4, 333-346.
- [35] Vamsi P. Pingali: A priori estimates for a generalized Monge-Ampère PDE on some compact Kähler manifolds. Complex Var. Elliptic Equ. 61 (2016), no. 8, 1037-1051.
- [36] D. H. Phong, S. Picard, X. W. Zhang: On estimates for the Fu-Yau generalization of a Strominger system. J. Reine Angew. Math. (Crelle's Journal), Vol. 2019, Issue 751 (2019), 243-274
- [37] D. H. Phong, S. Picard, X. W. Zhang: The Fu-Yau equation with negative slope parameter. Invent. Math., Vol. 209, No. 2 (2017), 541-576.
- [38] D. H. Phong, D. T. Tô: Fully non-linear parabolic equations on compact Hermitian manifolds. arXiv:1711.10697
- [39] D. H. Phong, S. Picard, X. W. Zhang: Fu-Yau Hessian Equations. arXiv:1801.09842, to appear in J. Differential Geometry.
- [40] G. Székelyhidi: Fully non-linear elliptic equations on compact Hermitian manifolds. J. Differ. Geom. 109(2), 337-378 (2018).
- [41] J. Song, B. Weinkove: On the convergence and singularities of the J -flow with applications to the Mabuchi energy. Comm. Pure Appl. Math., 61 (2008), 210-229.
- [42] J. Spruck: Geometric aspects of the theory of fully nonlinear elliptic equations. Clay Mathematics Proceedings, volume 2, 2005, 283-309.
- [43] W. Sun: On a class of fully nonlinear elliptic equations on closed Hermitian manifolds. J. Geom. Anal. 26(3), 2459-2473 (2016)
- [44] W. Sun: Parabolic complex Monge-Ampère type equations on closed Hermitian manifolds. Calc. Var. PDE 54(4), 3715-3733 (2015)
- [45] W. Sun: On a class of fully nonlinear elliptic equations on closed Hermitian manifolds II: L^∞ estimate. Commun. Pure Appl. Math. 70(1), 172-199 (2017)
- [46] W. Sun: The Parabolic Flows for Complex Quotient Equations. J. Geom. Anal. 29:1520-1545 (2019)
- [47] W. Sun, Generalized complex Monge-Ampère type equations on closed Hermitian manifolds, preprint, arXiv:1412.8192.
- [48] W. Sun, Parabolic Flow for Generalized complex Monge-Ampère type equations preprint, arXiv:1501.04255.
- [49] V. Tosatti, B. Weinkove: Estimates for the complex Monge-Ampère equation on Hermitian and balanced manifolds. Asian J. Math. 14, 19-40 (2010)
- [50] V. Tosatti, B. Weinkove: The complex Monge-Ampère equation on compact hermitian manifolds. J. Amer. Math. Soc., 23 (2010), 1187-1195.
- [51] V. Tosatti, Y. Wang, B. Weinkove, X. K. Yang: $C^{2,\alpha}$ estimates for nonlinear elliptic equations in complex and almost complex geometry. Calc. Var. Partial Differential Equations 54 (2015), no. 1, 431-453.
- [52] B. Weinkove: Convergence of the J -flow on Kähler surfaces. Comm. Anal. Geom. 12 (2004), no. 4, 949-965.
- [53] B. Weinkove: The J -flow, the Mabuchi energy, the Yang-Mills flow and multiplier ideal sheaves. Doctoral dissertation, Columbia University, New York, 2004.
- [54] B. Weinkove: On the J -flow in higher dimensions and the lower boundedness of the Mabuchi energy. J. Differential Geom. 73 (2006), no. 2, 351-358.
- [55] N. S. Trudinger: The Dirichlet problem for the prescribed curvature equations. Arch. Rational Mech. Anal. 111(2) (1990), 153-179.

- [56] S.T. Yau: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I. *Comm. Pure Appl. Math.*, 31 (1978), 339-411.
- [57] D. K. Zhang: Hessian equations on closed Hermitian manifolds. *Pac. J. Math.* 291(2), 485-510 (2017)
- [58] X. W. Zhang: A priori estimates for complex Monge-Ampère equation on Hermitian manifolds. *Int. Math. Res. Not.* 2010, no. 19, 3814-3836.

FACULTY OF MATHEMATICS AND STATISTICS, HUBEI KEY LABORATORY OF APPLIED
MATHEMATICS, HUBEI UNIVERSITY, WUHAN 430062, P.R. CHINA
Email address: chenli@hubu.edu.cn