# Exponential stability of linear systems under a class of Desch-Schappacher perturbations

S. El Alaoui and M. Ouzahra

M2PA Laboratory, University of Sidi Mohamed Ben Abdellah P.O. Box 5206, Bensouda, Fès, Morocco mohamed.ouzahra@usmba.ac.ma

#### Abstract

In this paper we investigate the uniform exponential stability of the system  $\frac{dx(t)}{dt} = Ax(t) - \rho Bx(t)$ ,  $(\rho > 0)$ , where the unbounded operator A is the infinitesimal generator of a linear  $C_0$ -semigroup of contractions S(t) in a Hilbert space X and B is a Desch-Schappacher operator. Then we give sufficient conditions for exponential stability of the above system. The obtained stability result is then applied to show the uniform exponential stabilization of bilinear partial differential equations.

**Keywords**: Exponential stabilization, linear system, bilinear control, Desch-Schappacher operator, unbounded control operator.

#### I. INTRODUCTION

Consider the following abstract system

$$\begin{cases} \dot{x}(t) = Ax(t) - \rho Bx(t), \ t > 0\\ x(0) = x_0, \end{cases}$$
(1)

where the state x(.) takes values in a Hilbert state space X endowed with an inner product  $\langle ., . \rangle_X$  with associate norm  $\|.\|_X$ , the unbounded operator (A, D(A)) generates a  $C_0$ -semigroup S(t) on X. Here, B is an unbounded linear operator of X in the sense that it is bounded from X to some extrapolating space of X. In the case of various real problems, the modeling may lead to mathematical model of the form (1) with an operator B which is of type Desch-Schappacher. Such a perturbation operator B appears for instance in case of control actions exercised through the boundary or at a point of the geometrical domain of partial differential equations, and also in many other situations of internal control (see [4, 7, 8] and the references therein). Due to the unbounded aspect of the operator B, the solution of (1) does not exist, in general, with values in X. Thus, to confront this difficulty the concept of admissibility is needed, which requires the introduction of interpolating and extrapolating spaces of the state space X.

Our goal in this paper is to investigate the uniform exponential stability of the system (1). This consists on looking for a set of parameters  $\rho$  for which there exists a global X-valued mild solution x(t) of (1) and is such that  $||x(t)|| \le Ke^{-\sigma t}||x_0||$ ,  $\forall t \ge 0$  for some constants  $K, \sigma > 0$ . As an application, one can consider the stabilization of bilinear systems by means of switching controllers, which leads to a closed-loop system like (1). This problem has been considered in [11]

for a bounded operator *B*. The case of a Miyadera-Voigt type operator has been investigated in [15]. Moreover, in [2] the case of 1–admissibility in Banach space has been considered. However, the 1–admissibility assumption prevents us to consider the case of Hilbert state space as in this case the operator *B* will be necessary bounded (see [18]). In other words, the 1–admissibility condition excludes several applications that are also available in Hilbert space. Moreover, in [2] it was assumed that  $D((A_{-1} - \rho B)|X) = D(A_{-1}) \cap D(B|X)$ , which played an essential role in the proofs of the stabilization results (in a technical point of view). Unfortunately, there are several examples in which this domain condition is not fulfilled (see e.g Examples 1&2). In this paper, we will rather use the *p*–admissibility property with  $p \ge 1$ . Then we introduce new sufficient conditions for uniform exponential stability of system (1), which are easily checkable. In the sequel, we proceed as follows: The main results of this paper are contained in Section 2. In Section 3, we provide applications to feedback stabilization of bilinear heat and transport equations.

#### II. EXPONENTIAL STABILITY

In this section, we state and prove our two main stabilization results. We start by introducing the necessary tools regarding the notion of admissibility in connection with the generation results and then provide some a priori estimations of the solution.

### i. Preliminary results

As pointed out in the introduction, the unbounded aspect of the operator B do not guarantee the existence of an X-valued solution x(t) of (1). However, one may extend the system at hand in a larger (extrapolating) space  $X_{-1}$  of the state space X in which the existence of the solution x(t) is ensured and then give the required admissibility conditions of B, so that the solution x(t) lies in X. Classically, the spaces  $X_1$  and  $X_{-1}$  are defined as follows:  $X_1 := (D(A), \|\cdot\|_1)$ , where  $\|x\|_1 :=$  $\|(\lambda I - A)x\|_X$ ,  $x \in D(A)$  for some  $\lambda$  in the resolvent set  $\rho(A)$  of A, and  $X_{-1}$  is the completion of X with respect to the norm  $||x||_{-1} := ||(\lambda I - A)^{-1}x||_X$ ,  $x \in X$ . These spaces are independent of the choice of  $\lambda$  and are related by the following continuous and dense embedding:  $X_1 \hookrightarrow$  $X \hookrightarrow X_{-1}$ . That way the unbounded operator *B* becomes bounded from *X* to the extrapolating space  $X_{-1}$ , i.e.,  $B \in \mathcal{L}(X, X_{-1})$ . Thus, in order to give a meaning to solutions of (1), we have to use the fact that the semigroup S(t) can be extended to a  $C_0$ -semigroup  $S_{-1}(t)$  on  $X_{-1}$ , whose generator  $A_{-1}$  has  $D(A_{-1}) = X$  as domain and is such that  $A_{-1}x = Ax$ , for any  $x \in D(A)$ . By definition,  $(X_{-1}, \|.\|_{-1})$  is a Banach space and one can easily show that the operator A is bounded from  $(D(A), \|.\|_X)$  into the Banach space  $(X_{-1}, \|.\|_{-1})$ , so that it can be extended to an operator  $A_{-1} \in \mathcal{L}(X, X_{-1})$ , generating a  $C_0$ -semigroup  $S_{-1}(t)$  on  $X_{-1}$  and which is given by  $S_{-1}(t) =$  $(\lambda - A_{-1})S(t)(\lambda - A_{-1})^{-1}$ . Indeed, it clear that  $S_{-1}(t)$  is a  $C_0$ -semigroup on  $X_{-1}$ . Moreover, to show that its generator is  $A_{-1}$ , we show that for all  $x \in X$ ,  $\lim_{t\to 0} \frac{S_{-1}(t)x-x}{t} = A_{-1}x$ . We first show this for  $x \in D(A)$ . This case follows from the fact that  $\lim_{t\to 0} \frac{S_{-1}(t)x-x}{t} = \lim_{t\to 0} \frac{S(t)x-x}{t} = Ax$ . Now for  $x \in X$ , we use the fact that  $R(\lambda, A)x \in D(A)$  to deduce that  $R(\lambda, A_{-1}) \lim_{t \to 0} \frac{\dot{S}_{-1}(t)x - x}{t} = C(A)$  $R(\lambda, A_{-1})A_{-1}x$ , which gives the desired result. Moreover, if the semigroup S(t) is a contraction, then so is  $S_{-1}(t)$ . Indeed, having in mind that for all  $\lambda \in \rho(A_{-1})$  and for all  $x \in X_{-1}$ , we have  $(\lambda I - A_{-1})^{-1}x \in X$ , and  $(\lambda I - A_{-1})^{-1}x = (\lambda I - A)^{-1}x$ ,  $\forall x \in X$ , we can see that for all  $x \in X$ we have

$$||S_{-1}(t)x||_{-1} = ||(\lambda I - A)^{-1}S(t)x||_X \le ||(\lambda I - A)^{-1}x||_X = ||x||_{-1},$$

thus, by density of X in  $X_{-1}$ , we conclude that the semigroup  $S_{-1}(t)$  is a contraction on  $X_{-1}$ . Recall that for any given initial state  $x_0 \in X$ , a mild solution of (1) is an X-valued continuous function x on [0, T] satisfying the following variation of parameters formula:

$$x(t) = S(t)x_0 - \rho \int_0^t S_{-1}(t-s)Bx(s)ds, \ \forall t \ge 0,$$

which always makes sense in  $X_{-1}$ . The system (1) can be rewritten in the large space  $X_{-1}$  in the following abstract form:

$$\begin{cases} \dot{x}(t) = A_{-1}x(t) - \rho Bx(t), \\ x(0) = x_0. \end{cases}$$
(2)

which is well-posed in *X* whenever  $A - \rho B$  is the generator of a  $C_0$ -semigroup on *X* (cf. [6], Section II.6). The well-posedeness of systems like (1) has been studied in many works using different approaches (see e.g. [1, 3, 5, 9, 13, 18]).

The next result provides sufficient conditions on a Desch-Schappacher perturbation B to guarantee the existence and uniqueness of the mild solution of (1) (see [1] & ([6], p. 183)).

**Theorem II.1** Let A be the generator of a  $C_0$ -semigroup S(t) on X and let  $B \in \mathcal{L}(X, X_{-1})$  be p-admissible for some  $1 \le p < \infty$ , i.e., there is a T > 0 such that

$$\int_0^T S_{-1}(T-t)Bu(s)ds \in X, \quad \forall u \in L^p(0,T;X).$$
(3)

Then for any  $\rho$ , the operator  $(A_{-1} - \rho B)|_X$  defined on the domain  $D((A_{-1} - \rho B)|_X) := \{x \in X : (A_{-1} - \rho B)x \in X\}$  by

$$(A_{-1} - \rho B)|_X x := A_{-1} x - \rho B x, \quad \forall x \in D((A_{-1} - \rho B)|_X)$$
(4)

is the generator of a  $C_0$ -semigroup  $(T(t))_{t>0}$  on X, which verifies the variation of parameters formula

$$T(t)x = S(t)x - \rho \int_0^t S_{-1}(t-s)BT(s)xds, \ \forall t \ge 0, \ \forall x \in D((A_{-1} - \rho B)|_X).$$

An operator  $B \in \mathcal{L}(X, X_{-1})$  satisfying the condition (3) is called a Desch-Schappacher operator or perturbation. Moreover, the operator defined by (4) is the part  $(A_{-1} - \rho B)|_X$  of  $(A_{-1} - \rho B)$  on X (see ([16], p. 39) and ([6], p. 147)).

**Remark 1** Notice that since  $W^{1,p}(0,T;X)$  is dense in  $L^p(0,T;X)$ , the range condition (3) is equivalent to the existence of some M > 0 such that

$$\left\| \int_{0}^{T} S_{-1}(T-s) Bu(s) ds \right\|_{X} \le M \|u\|_{L^{p}(0,T;U)}, \quad \forall u \in W^{1,p}(0,T;X),$$

$$with \|u\|_{L^{p}(0,T;X)} = \left( \int_{0}^{T} \|u(t)\|_{X}^{p} dt \right)^{\frac{1}{p}}.$$
(5)

**Remark 2** Note that if the operator  $B \in \mathcal{L}(X, X_{-1})$  is p-admissible in [0, T], then it is so in [0, t] for any  $t \in [0, T]$ . In other words, if (5) holds then for all  $t \in [0, T]$  we have the following inequality

$$\left\|\int_{0}^{t} S_{-1}(t-r)Bu(r)dr\right\|_{X} \le M\|u\|_{L^{p}(0,t;X)}.$$
(6)

3

Indeed, if for all  $t \in [0, T]$  and  $u \in L^p(0, T; X)$ , we define  $u_t \in L^p(0, T; X)$  by

$$u_t(r) := \begin{cases} 0, & \text{for } r \in [0, T-t] \\ \\ u(r+t-T), & \text{for } r \in (T-t, T] \end{cases}$$

then, observing that  $\int_0^t S_{-1}(t-r)Bu(r)dr = \int_0^T S_{-1}(T-r)Bu_t(r)dr \in X$ , it comes from the *p*-admissibility of *B* that  $\left\|\int_0^t S_{-1}(t-r)Bu(r)dr\right\|_X \leq M \|u_t\|_{L^p(0,T;X)}$ . Thus for all  $t \in [0,T]$ , we have  $\|\int_0^t S_{-1}(t-r)Bu(r)dr\|_X \leq M \|u\|_{L^p(0,t;X)}$ .

Let us now show the following lemma that will be needed in the sequel.

**Lemma II.2** Let assumptions of Theorem II.1 hold. Then for any  $0 < \rho < \frac{1}{T^{\frac{1}{p}}M}$ , the mild solution x(t) of the system (1) satisfies the following estimate

$$\|x(.)\|_{L^{p}(0,T;X)} \leq \frac{T^{\frac{1}{p}}}{1 - \rho T^{\frac{1}{p}}M} \|x_{0}\|_{X}, \, \forall x_{0} \in X$$
(7)

and

$$\left\| \int_{0}^{t} S_{-1}(t-s) Bx(s) ds \right\|_{X} \le M_{\rho} \|x_{0}\|_{X}, \ \forall t \in [T, 2T], \ \forall x_{0} \in X,$$
with  $M_{\rho} := \frac{MT^{\frac{1}{p}}}{1 - \sigma T^{\frac{1}{p}}M} \left( 2 + \rho MT^{\frac{1}{p}} \right).$ 

**Proof 1** Let  $x_0 \in D((A_{-1} - \rho B)|_X)$ . From Theorem II.1, we know that the system (1) admits a unique mild solution x(t) which is given by

$$x(t) = S(t)x_0 - \rho \int_0^t S_{-1}(t-s)Bx(s)ds, \quad \forall t \ge 0.$$
(8)

Let us estimate  $||x(\cdot)||_{L^p(0,T;X)}$ . From (8), we get via Minkowski's inequality

$$\|x(.)\|_{L^{p}(0,T;X)} \leq \left(\int_{0}^{T} \|S(t)x_{0}\|_{X}^{p} dt\right)^{\frac{1}{p}} + \rho \left(\int_{0}^{T} \left\|\int_{0}^{t} S_{-1}(t-s)Bx(s)ds\right\|_{X}^{p}\right)^{\frac{1}{p}} dt$$

Then from Remark 2 we derive

$$\|x(\cdot)\|_{L^{p}(0,T;X)} \leq T^{\frac{1}{p}} \|x_{0}\|_{X} + \rho T^{\frac{1}{p}} M \|x(.)\|_{p},$$

where  $||x(\cdot)||_{L^{p}(0,T;X)} := \left(\int_{0}^{T} ||x(\tau)||_{X}^{p} d\tau\right)^{\frac{1}{p}}$ , which gives the estimate (7) for any  $0 < \rho < \frac{1}{T^{\frac{1}{p}}M}$ . Now, since the mapping  $x_{0} \mapsto x(t)$  defines a  $C_{0}$ -semigroup T(t) on X, the mapping  $x_{0} \mapsto x(\cdot) = T(\cdot)x_{0}$  is continuous from X to  $L^{p}(0,T;X)$ . Then the estimate (7) holds by density for any  $x_{0} \in X$ . Let  $x_{0} \in X$ , and let us write for any  $t \in [T,2T]$ ,

$$\int_0^t S_{-1}(t-s)Bx(s)ds = \int_0^T S_{-1}(t-s)Bx(s)ds + \int_T^t S_{-1}(t-s)Bx(s)ds$$
  
:=  $L_1 + L_2$ 

Then we consider the two terms of the sum separately. For the first one, the admissibility of B together with the contraction property of  $S_{-1}(t)$  yields

$$\|L_1\|_X = \left\|S_{-1}(t-T)\int_0^T S_{-1}(T-s)Bx(s)ds\right\|_X \le M\|x(\cdot)\|_{L^p(0,T;X)}$$
(9)

For the second term, observing that  $L_2 = \int_0^{t-T} S_{-1}(t-T-\tau)Bx(\tau+T)d\tau$ , we obtain again from the admissibility of B

$$||L_2||_X \le M ||x(.+T)||_{L^p(0,T;X)}$$

Based on the V.C.F (8), it follows directly from Lemma (7) that for all  $t \ge 0$ , we have

$$\|x(t)\|_{X} \le \left(1 + \frac{\rho M T^{\frac{1}{p}}}{1 - T^{\frac{1}{p}} \rho M}\right) \|x_{0}\|_{X}, \forall x_{0} \in X$$
(10)

for any  $0 < \rho < \frac{1}{T^{\frac{1}{p}}M}$ . Using the last estimate, we derive the following inequalities:

$$\|x(.+T)\|_{L^{p}(0,T;X)} \leq T^{\frac{1}{p}} \left(1 + \frac{\rho M^{2} T^{\frac{1}{p}}}{1 - \rho M T^{\frac{1}{p}}}\right) \|x_{0}\|_{X}.$$
(11)

Combining (9) and (11), we obtain the desired estimate:

$$\left\|\int_0^t S_{-1}(t-s)Bx(s)ds\right\|_X \le M_\rho \|x_0\|_X$$

## ii. A direct approach

**Theorem II.3** Let  $B \in \mathcal{L}(X, X_{-1})$  and let A be the infinitesimal generator of a linear  $C_0$ -semigroup of contractions S(t) on X, and assume that for some T > 0, we have

- (i) there exists  $1 such that for all <math>u \in L^p(0,T;X)$ , we have  $\int_0^T S_{-1}(T-s)Bu(s)ds \in X,$
- (ii) for some  $\delta > 0$  we have

$$\int_0^T \operatorname{Re} \left\langle S(t)x, B^*S(t)x \right\rangle_X dt \ge \delta \|S(T)x\|_X^2, \quad \forall x \in X.$$
(12)

Then there is a  $\rho_1 > 0$  such that for all  $\rho \in (0, \rho_1)$ , the system (1) is exponentially stable on X.

**Proof 2** Foe any  $\rho > 0$ , we set  $A_{\rho B} := (A_{-1} - \rho B)|_X$ . According to assumption (i), we deduce from Theorem 1 that the system (1) admits a unique mild solution which is given, for  $x_0 \in D(A_{\rho B})$ , by the variation of parameters formula (see [5]):

$$x(t) = S(t)x_0 - \rho \int_0^t S_{-1}(t-s)Bx(s)ds, \quad \forall t \ge 0.$$
(13)

For  $\lambda \in \rho(A)$  ( $\rho(A)$  is the resolvent set of A), we consider the system (1) with  $B_{\lambda} := \lambda R(\lambda, A_{-1})B$ instead of B. Observing that the operator  $B_{\lambda}$  is bounded, we deduce that the corresponding system admits a unique mild solution denoted by  $x_{\lambda}$ , which satisfies the following formula

$$x_{\lambda}(t) = S(t)x_0 - \rho \int_0^t S_{-1}(t-s)B_{\lambda}x_{\lambda}(s)ds, \quad \forall t \ge 0.$$
(14)

5

*We claim that*  $x_{\lambda}(t)$  *converges to* x(t) *as*  $\lambda \to +\infty$ *. Indeed, for all* t > 0*, we have* 

$$\begin{aligned} x_{\lambda}(t) - x(t) &= \rho \int_{0}^{t} S_{-1}(t-s) B_{\lambda} x_{\lambda}(s) - \rho \int_{0}^{t} S_{-1} B x(s) ds \\ &= \rho \int_{0}^{t} S_{-1}(t-s) B_{\lambda}(x_{\lambda}(s) - x(s)) ds + \rho \int_{0}^{t} S_{-1}(t-s) B_{\lambda} x(s) ds - \rho \int_{0}^{t} S_{-1}(t-s) B x(s) ds. \end{aligned}$$

*Then, using (5), this yields for all*  $t \in [0, T]$ 

$$\|x_{\lambda}(t) - x(t)\|_{X} \leq \rho M \|x_{\lambda}(.) - x(.)\|_{L^{p}(0,t;X)} + \rho \left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda}x(s) ds - \int_{0}^{t} S_{-1}(t-s) Bx(s) ds \right\|_{X},$$

which by integrating gives for all  $t \in [0, T]$ 

$$\|x_{\lambda}(t) - x(t)\|_{L^{p}(0,T;X)}^{p} \leq T(2\rho M)^{p} \|x_{\lambda}(.) - x(.)\|_{L^{p}(0,T;X)}^{p} + (2\rho)^{p} \left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda}x(s) ds - \int_{0}^{t} S_{-1}(t-s) B_{X}(s) ds \right\|_{X}^{p}.$$

It follows that

$$\|x_{\lambda}(.) - x(.)\|_{L^{p}(0,T;X)}^{p} \leq \rho C_{\rho} \int_{0}^{T} \left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda}x(s) ds - \int_{0}^{t} S_{-1}(t-s) Bx(s) ds \right\|_{X}^{p} dt,$$

with  $C_{\rho} := \frac{(2\rho)^p}{1-T(2\rho M)^p}$ . It is clear that  $\lim_{\lambda \to \infty} \int_0^t S_{-1}(t-s) B_{\lambda} x(s) ds = \int_0^t S_{-1}(t-s) B x(s) ds$  in X and we have

$$\left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda} x(s) ds - \int_{0}^{t} S_{-1}(t-s) B_{\lambda}(s) ds \right\|_{X} \le \left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda} x(s) ds \right\|_{X} + \left\| \int_{0}^{t} S_{-1}(t-s) B_{\lambda}(s) ds \right\|_{X}.$$

Moreover, by the admissibility assumption we get

$$\left\|\int_0^t S_{-1}(t-s)B_{\lambda}x(s)ds - \int_0^t S_{-1}(t-s)Bx(s)ds\right\|_X \le 2M\|x(.)\|_{L^p(0,t;X)}.$$

Then, according to the dominated convergence theorem we have

$$\begin{split} \lim_{\lambda \to \infty} \|x_{\lambda}(.) - x(.)\|_{L^{p}(0,T;X)}^{p} &\leq \lim_{\lambda \to \infty} \int_{0}^{T} \left\| \rho \int_{0}^{t} S_{-1}(t-s) B_{\lambda}x(s) ds - \rho \int_{0}^{t} S_{-1}(t-s) Bx(s) ds \right\|_{X} dt \\ &= \int_{0}^{T} \lim_{\lambda \to \infty} \left\| \rho \int_{0}^{t} S_{-1}(t-s) B_{\lambda}x(s) ds - \rho \int_{0}^{t} S_{-1}(t-s) Bx(s) ds \right\|_{X} dt \\ &= 0. \end{split}$$

Let  $x_0 \in D(A_{\rho B})$  be fixed. Thus for all t > 0 we have

$$\frac{d}{dt} \|x_{\lambda}(t)\|_{X}^{2} \leq -2\rho \mathcal{R}e \langle B_{\lambda}x_{\lambda}(t), x_{\lambda}(t) \rangle_{X}, \quad \forall t > 0.$$
(15)

For all t > 0, we have the following equality

$$\langle B_{\lambda}S(t)x_{0}, S(t)x_{0} \rangle_{X} = \langle B_{\lambda}S(t)x_{0}, S(t)x_{0} - x_{\lambda}(t) \rangle_{X} + \langle B_{\lambda}S(t)x_{0} - B_{\lambda}x_{\lambda}(t), x_{\lambda}(t) \rangle_{X} + \langle B_{\lambda}x_{\lambda}(t), x_{\lambda}(t) \rangle_{X}$$

which gives

$$\begin{split} \langle S(t)x_0, B^*\lambda R^*(\lambda, A)S(t)x_0\rangle_X &= \langle S(t)x_0, B^*\lambda R^*(\lambda, A)\left(S(t)x_0 - x_\lambda(t)\right)\rangle_X \\ &+ \langle S(t)x_0 - x_\lambda(t), B^*\lambda R^*(\lambda, A)x_\lambda(t)\rangle_X + \langle B_\lambda x_\lambda(t), x_\lambda(t)\rangle_X \end{split}$$

Let us estimate each term of this last expression.

We deduce from (i) that for some constant M > 0 and for all  $u \in L^{p}(0,T;X)$ , we have

$$\left\|\int_{0}^{T} S_{-1}(T-s) Bu(s) ds\right\|_{X} \le M \|u\|_{L^{p}(0,T;X)}$$
(16)

The formula (14) combined with the estimate (6), gives

$$\|S(t)x_0 - x_{\lambda}(t)\|_{X} = \rho \left\| \int_0^t S_{-1}(t-s)Bx_{\lambda}(s)ds \right\|_{X} \le \rho M \|x_{\lambda}(.)\|_{L^p(0,T;X)}, \forall t \in [0,T].$$

Then, according to Lemma II.2, we conclude that

$$\|S(t)x_0 - x_{\lambda}(t)\|_X \le \frac{\rho M T^{\frac{1}{p}}}{1 - \rho T^{\frac{1}{p}} M} \|x_0\|_X$$
(17)

For every t > 0, we have

$$\begin{aligned} \mathcal{R}e \left\langle S(t)x_{0}, B^{*}\lambda R^{*}(\lambda, A)S(t)x_{0}\right\rangle_{X} &\leq C\|S(t)x_{0}\|_{X}\|B^{*}\|_{\mathcal{L}(X_{-1}, X)}\|\lambda R^{*}(\lambda, A)\|_{\mathcal{L}(X)}\|S(t)x_{0} - x_{\lambda}(t)\|_{X} \\ &+ C\|B^{*}\|_{\mathcal{L}(X_{-1}, X)}\|\lambda R^{*}(\lambda, A)\|_{\mathcal{L}(X)}\|x(t)\|_{X}\|S(t)x_{0} - x(t)\|_{X} \\ &+ \mathcal{R}e \left\langle B_{\lambda}x_{\lambda}(t), x_{\lambda}(t)\right\rangle_{X}, \end{aligned}$$

where *C* is a positive constant.

Using the fact that S(t) is a contraction, it comes

$$\mathcal{R}e \left\langle S(t)x_0, B^*\lambda R^*(\lambda, A)S(t)x_0 \right\rangle_X \leq C \|B^*\|_{\mathcal{L}(X_{-1}, X)} \|x_0\|_X \|S(t)x_0 - x_\lambda(t)\|_X \\ + C \|B^*\|_{\mathcal{L}(X_{-1}, X)} \|x(t)\|_X \|S(t)x_0 - x(t)\|_X + \mathcal{R}e \left\langle B_\lambda x_\lambda(t), x_\lambda(t) \right\rangle_X$$

*Using* (10) *and* (17)*, we deduce that for all*  $t \in (0, T]$  *we have* 

$$\begin{aligned} \mathcal{R}e\left\langle S(t)x_{0},B^{*}\lambda R^{*}(\lambda,A)S(t)x_{0}\right\rangle_{X} &\leq \frac{\rho CLMT^{\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M}\|x_{0}\|_{X}^{2} \\ &+ \frac{\rho CLMT^{\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M}\left(1+\frac{\rho MT^{\frac{1}{p}}}{1-T^{\frac{1}{p}}\rho M}\right)\|x_{0}\|_{X}^{2}+\mathcal{R}e\left\langle B_{\lambda}x_{\lambda}(t),x_{\lambda}(t)\right\rangle_{X},\end{aligned}$$

with  $L := \|B^*\|_{\mathcal{L}(X_{-1},X)}$ . Then, integrating the last inequality and using (15) we get for all  $t \in [0, t]$ 

$$\begin{aligned} 2\rho \int_{0}^{T} \mathcal{R}e \left\langle S(t)x_{0}, B^{*}\lambda R^{*}(\lambda, A)S(t)x_{0}\right\rangle_{X} &\leq \frac{2\rho^{2}CLMT^{\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M} \|x_{0}\|_{X}^{2} \\ &+ \frac{2\rho^{2}CLMT^{\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M} \left(1+\frac{\rho MT^{\frac{1}{p}}}{1-T^{\frac{1}{p}}\rho M}\right) \|x_{0}\|_{X}^{2} + \|x_{0}\|_{X}^{2} - \|x_{\lambda}(T)\|_{X}^{2}, \end{aligned}$$

Thus, letting  $\lambda \to +\infty$ , we drive

$$2\mathcal{R}e\int_{0}^{T} \mathcal{R}e \langle S(t)x_{0}, B^{*}S(t)x_{0} \rangle_{X} dt \leq 2\rho^{2}C_{1} \|x_{0}\|_{X}^{2} + \|x_{0}\|_{X}^{2} - \|x(T)\|_{X}^{2}$$
$$= \frac{MCLT^{1+\frac{1}{p}}}{2} \left(2 + \frac{\rho MT^{\frac{1}{p}}}{2}\right)$$

with  $C_1 = \frac{MCLT}{1-\rho T^{\frac{1}{p}}M} \left(2 + \frac{\rho MT}{1-\rho T^{\frac{1}{p}}M}\right)$ . Applying the inequality (12), it follows that

$$2\rho\delta\|S(T)x_0\|_X^2 - 2\rho^2 C_1\|x_0\|_X^2 \le \|x_0\|_X^2 - \|x(T)\|_X^2$$
(18)

*Using Lemma II.2, we deduce via the variation of constants formula (13) that for all*  $t \in [T, 2T]$ *, we have* 

$$\|x(t)\|_{X} \leq \|S(T)x_{0}\|_{X} + \rho \|\int_{0}^{t} S_{-1}(t-s)Bx(s)ds\|_{X}$$

$$\leq \|S(T)x_0\|_X + \rho M_{\rho}\|x_0\|_X.$$

By reiterating the processes for  $t \in [kT, (k+1)T], k \ge 1$ , we deduce that

$$||x(t)||_{X} \le ||S(T)x(kT)||_{X} + \rho M_{\rho} ||x(kT)||_{X}.$$

*Then for all*  $k \ge 1$ *, we have* 

$$\|x((k+1)T)\|_{X}^{2} \leq 2\|S(T)x(kT)\|_{X}^{2} + 2\rho M_{\rho}\|x(kT)\|_{X}^{2}.$$
(19)

Moreover, (18) becomes

$$2\rho\delta\|S(T)(kT)\|_X^2 - 2\rho^2 C_1 \|x(kT)\|_X^2 \le \|x(kT)\|_X^2 - \|x((k+1)T)\|_X^2$$
(20)

This together with (19) implies

$$\rho\delta\bigg(\|x((k+1)T)\|_X^2 - 2\rho M_\rho \|x(kT)\|_X^2\bigg) - 2C_1\rho^2 \|x(kT)\|_X^2 \le \|x(kT)\|_X^2 - \|x((k+1)T)\|_X^2.$$

Hence

with  $C_1$ 

$$(1+\rho\delta)\|x((k+1)T)\|_X^2 \le \left(2\delta\rho^2 M_\rho + 2C_1\rho^2 + 1\right)\|x(kT)\|_X^2, \ k \ge 0$$

This implies

$$\|x((k+1)T)\|_X^2 \le C_2 \|x(kT)\|_X^2$$

where  $C_2 = \frac{2\rho^2(\delta M_{\rho}+C_1)+1}{1+\rho\delta}$ , which is in (0, 1) for  $\rho \to 0^+$ . Since  $||x(t)||_X$  decreases, we get for  $k = E\left(\frac{t}{T}\right)$  (where E(.) is the integer part function).

$$||x(t)||_X^2 \le (C_2)^k ||x_0||_X^2$$

which gives the following exponential decay

$$\|x(t)\|_X \le Ke^{-\sigma t} \|x_0\|, \ \forall t \ge 0$$

where  $K = (C_2)^{-\frac{1}{2}}$  and  $\sigma = \frac{-ln(C_2)}{2T}$ . This estimate extends by density to all  $x_0 \in X$ . Hence the uniform exponential stability hold for any  $0 < \rho < \rho_1$ , where  $\rho_1$  is such that  $0 < \rho_1 < \frac{1}{T^{\frac{1}{p}}M}$  and  $\frac{2\rho^2(\delta M_\rho+C_1)+1}{1+\rho\delta}\in(0,1).$ 

#### Remark 1

In the case where  $Range(BS(t)) \subset X$ ,  $\forall t > 0$ , the condition (12) is equivalent to the conventional one (see [2, 15]):

$$\int_0^T \operatorname{Re} \left\langle BS(t)x, S(t)x \right\rangle_X dt \ge \delta \|S(T)x\|_X^2, \ \forall x \in X.$$
(21)

Moreover, the condition (21) can be weakened if an appropriate decomposition of Range(B) is available. This is the aim of the next section.

#### iii. A range decomposition method

Let  $\mathfrak{X} \oplus \mathfrak{X}_{-1}$  be a direct sum in  $X_{-1}$ , where  $\mathfrak{X} = i(X)$  (*i* being the canonical injection of X in  $X_{-1}$ ), so we can write  $\mathfrak{X} = X$ . Then for any  $C \in \mathcal{L}(X, X_{-1})$  such that  $rg(C) \subset \mathfrak{X} \oplus \mathfrak{X}_{-1}$ , we set  $_XC =: P_{\mathfrak{X}}C$ , where  $P_{\mathfrak{X}}$  is the projection of  $\mathfrak{X}$  according to  $\mathfrak{X} \oplus \mathfrak{X}_{-1}$ . Now, given a pair of operators  $(K, L) \in \mathcal{L}(X, X_{-1}) \times \mathcal{L}(X, X_{-1})$ , the decomposition  $\mathfrak{X} \oplus \mathfrak{X}_{-1}$  is said to be admissible for (K, L) if the three following properties hold:

(a)  $rg(K) \subset \mathfrak{X} \oplus \mathfrak{X}_{-1}$  and  $rg(L) \subset \mathfrak{X} \oplus \mathfrak{X}_{-1}$ ,

(b)  $_{X}K$  is dissipative on  $D((K+L)|_{X}) := \{x \in X : Kx + Lx \in X\}$ ,

(c)  $_{X}L \in \mathcal{L}(X).$ 

For our stabilization problem, we will be interested with admissible decompositions for the pairs  $(A_{-1}, -\rho B)$  with  $\rho > 0$  small enough. Note that if the domain of the operator  $(A_{\rho B})|_X$  is independent of  $\rho > 0$  (small enough), which is equivalent to  $D((A_{\rho B})|_X) = D(A) \cap D(B|_X)$ , then for the sum  $\mathfrak{X} \oplus \mathfrak{X}_{-1}$  to be admissible for the pairs  $(A_{-1}, -\rho B)$ ,  $\rho > 0$ , it suffices to be admissible for the pair  $(A_{-1}, B)$ .

We are ready to state our second main result.

**Theorem II.4** Let A be the infinitesimal generator of a linear  $C_0$ -semigroup of contractions S(t) on X and let  $B \in \mathcal{L}(X, X_{-1})$ . Let  $\mathfrak{X} \oplus \mathfrak{X}_{-1}$  be an admissible decomposition for the pair  $(A_{-1}, -\rho B)$  for any  $\rho > 0$  small enough, and assume that for some T > 0, the operator B is p-admissible for some 1 and satisfies the estimate:

$$\int_0^T \operatorname{Re}\left\langle_X BS(t)x, S(t)x\right\rangle_X dt \ge \delta \|S(T)x\|_X^2, \ \forall x \in X,$$
(22)

for some  $T, \delta > 0$ .

Then there is a  $\rho_1 > 0$  such that the system (1) is exponentially stable on X for all  $\rho \in (0, \rho_1)$ .

**Proof 3** Let  $0 < \rho < \frac{1}{T^{\frac{1}{p}}M}$ , and let x(t) be the unique mild solution of the system (1) given for  $x_0 \in D((A_{\rho B})|_X)$  by the formula (13).

*The admissibility assumption on B together with Lemma II.2 implies the following estimate for*  $t \in [0, T]$  :

$$\|x(t) - S(t)x_0\|_X \le \frac{\rho M T^{\frac{1}{p}}}{1 - \rho T^{\frac{1}{p}} M} \|x_0\|_X$$
(23)

Moreover, observing that  $A_{\rho B}x(t) =_X (A_{\rho B})x(t)$ , we can write

$$\frac{d}{dt}\|x(t)\|_X^2 = 2\mathcal{R}e\langle_X(A_{-1})x(t) - \rho_X Bx(t), x(t)\rangle_X, \ \forall t > 0$$

5	L		

Integrating this last equality and using the dissipativeness of  $_X(A_{-1})$  gives

$$2\rho \int_{s}^{t} \mathcal{R}e\langle_{X}Bx(\tau), x(\tau)\rangle_{X}d\tau \le \|x(s)\|_{X}^{2} - \|x(t)\|_{X}^{2}, \ t \ge s \ge 0.$$
(24)

We have the following equality

$$\langle {}_{X}BS(t)x_{0},S(t)x_{0}\rangle_{X} = \langle {}_{X}BS(t)x_{0} - {}_{X}Bx(t),S(t)x_{0}\rangle_{X} + \langle {}_{X}Bx(t),S(t)x_{0} - x(t)\rangle_{X} + \langle {}_{X}Bx(t),x(t)\rangle_{X}$$

Then using the fact that the operator  $_XB$  is bounded, it comes

$$\mathcal{R}e \langle_X BS(t)x_0, S(t)x_0 \rangle_X \leq \|_X B\|_{\mathcal{L}(X)} \|x_0\|_X \|S(t)x_0 - x(t)\|_X + \|_X B\|_{\mathcal{L}(X)} \|x(t)\|_X \|S(t)x_0 - x(t)\|_X + \mathcal{R}e \langle_X Bx(t), x(t) \rangle_X$$

The estimate (23) combined with (10), implies

$$\begin{aligned} \mathcal{R}e\left\langle_{X}BS(t)x_{0},S(t)x_{0}\right\rangle_{X} &\leq \|_{X}B\|_{\mathcal{L}(X)}\frac{\rho MT^{\frac{1}{p}}}{1-T^{\frac{1}{p}}\rho M}\|x_{0}\|_{X}^{2} \\ &+ \|_{X}B\|_{\mathcal{L}(X)}\frac{\rho MT^{\frac{1}{p}}}{1-T^{\frac{1}{p}}\rho M}\left(1+\frac{\rho MT^{\frac{1}{p}}}{1-T^{\frac{1}{p}}\rho M}\right)\|x_{0}\|_{X}^{2} \\ &+ \mathcal{R}e\left\langle_{X}Bx(t),x(t)\right\rangle_{X}, \,\forall t \in [0,T]. \end{aligned}$$

Integrating this inequality and using the inequality (22), we deduce that

$$\delta \|S(T)x(kT)\|_{X}^{2} - \rho C_{1} \|x(kT)\|_{X}^{2} \leq \int_{kT}^{(k+1)T} \mathcal{R}e \langle_{X}Bx(s), x(s)\rangle_{X} ds$$
with  $C_{1} = \frac{MT^{1+\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M} \|_{X}B\|_{\mathcal{L}(X)} \left(2 + \frac{\rho MT^{\frac{1}{p}}}{1-\rho T^{\frac{1}{p}}M}\right).$ 

By using Lemma II.2, we derive

$$\rho \delta \left( \|x((k+1)T)\|_X^2 - 2\rho M_\rho \|x(kT)\|_X^2 \right) - 2C_1 \rho^2 \|x(kT)\|_X^2 \le \|x(kT)\|_X^2 - \|x((k+1)T)\|_X^2,$$

or equivalently

$$||x((k+1)T)||_X^2 \le C_2 ||x(kT)||_X^2,$$

where  $C_2 = \frac{2\rho^2(\delta M_{\rho}+C_1)+1}{1+\rho\delta}$ , which lies in (0,1) for  $\rho \to 0^+$ . Hence, using the decreasing of  $||x(t)||_X$ , we deduce the following exponential decay

$$||x(t)||_X \le Ke^{-\sigma t} ||x_0||, \ \forall t \ge 0,$$

where  $K = (C_2)^{-\frac{1}{2}}$  and  $\sigma = \frac{-ln(C_2)}{2T}$ . This estimate extends by density to all  $x_0 \in X$ . Thus, taking  $\rho_1 > 0$  such that  $0 < \rho_1 < \frac{1}{T^{\frac{1}{p}}M}$  and  $C_2 \in (0,1)$ , we get the result of the theorem.

## III. EXAMPLES

**Example 1** Let  $\Omega$  be an open and bounded subset of  $\mathbb{R}^d$ ,  $d \ge 1$ , and let us consider the following bilinear equation of diffusion type

$$\begin{cases} \frac{\partial}{\partial t}x = \Delta x + gx + \nu(t)(-\Delta)^{\frac{1}{2}}x & \text{in } \Omega \times (0,\infty), \\ x(t) = 0 & \text{on } \partial\Omega \times (0,\infty), \\ x(0) = x_0 & \text{in } \Omega. \end{cases}$$
(25)

where  $g \in L^{\infty}(\Omega)$ , v is a real valued bilinear control and  $x(t) = x(\zeta, t) \in L^{2}(\Omega)$  is the state. The system (25) is an example of fractional equation of diffusion equations type, and may describe transport processes in complex systems which are slower than the Brownian diffusion. As practical situations displaying such anomalous behaviour, let us mention the charge carrier transport in amorphous semiconductors, the nuclear magnetic resonance diffusometry in percolative and porous media etc (see [3, 10, 13, 12]). Here, we aim to show the exponential stabilization of (25). Let us observe that system (25) can be written in the form of (1) if we close it by the switching feedback control  $v(t) = -\rho \mathbf{1}_{\{t \ge 0 / x(t) \neq 0\}}$ . This is because we have  $\mathbf{1}_{\{t \ge 0 / x(t) \neq 0\}}(-\Delta)^{\frac{1}{2}}x(t) = (-\Delta)^{\frac{1}{2}}x(t), \forall t \ge 0$ . Let us take the state space  $X = L^{2}(\Omega)$  (endowed with its natural scalar product  $\langle \cdot, \cdot \rangle_{X}$ ), and consider the control operator  $B = (-\Delta)^{\frac{1}{2}}$  and the system's operator  $A = \Delta + gI$  with  $D(A) = H^{2}(\Omega) \cap H_{0}^{1}(\Omega)$ . The operator A generates an analytic semigroup S(t) on X(see [6], p. 107 and p. 176) which is given by the following variation of constants formula:

$$S(t)x = S_0(t)x + \int_0^t S_0(t-s)g(\xi)S(s)xds, \ t \ge 0,$$

where  $S_0(t)$  is the semigroup generated by A with g = 0. Let us verify the assumptions of Theorem 2. In order to make the computation easier, we restrict our self to the mono-dimension case, thus we consider  $\Omega = (0, 1)$ . In this case the semigroup  $(S_0(t))$  is is given by

$$S_0(t)x = \sum_{j\geq 1} e^{-lpha_j t} \langle x, \phi_j \rangle_X \phi_j, \ \forall x \in L^2(\Omega)$$

with  $\alpha_j = j^2 \pi^2$ ,  $j \ge 1$  is the set of eigenvalues of  $-\Delta$  with the corresponding orthonormal basis of  $L^2(\Omega)$ :  $\phi_j(x) = \sqrt{2} \sin(j\pi x)$ . Moreover, the semigroup S(t) is a contraction if in addition

$$\int_{\Omega} g(\xi) y^2(\xi) d\xi \leq \|y\|_{H^1_0(\Omega)'}^2, \ \forall y \in H^1_0(\Omega).$$

Thus, in the sequel we suppose this condition satisfied. Then the operator B can be expressed as

$$Bx = \sum_{j\geq 1} \alpha_j^{\frac{1}{2}} \langle x, \phi_j \rangle_X \phi_j, \ x \in L^2(\Omega).$$

Here, B is unbounded on  $L^2(\Omega)$  and it is bounded from  $L^2(\Omega)$  onto the space  $X_{-1}$  defined as the completion of  $L^2(\Omega)$  for the norm  $||y|| = \left(\sum_{j\geq 1} \frac{1}{\alpha_j} \langle y, \phi_j \rangle^2\right)^{\frac{1}{2}}$ ,  $\forall y \in L^2(\Omega)$ , which can be also interpreted as the dual space of  $D((-\Delta)^{\frac{1}{2}})$  with respect to the  $L^2(\Omega)$ -topology (the space  $L^2(\Omega)$  being the pivot space). Note also that the space  $D((-\Delta)^{\frac{1}{2}})$  can be doted with the norm  $||x||_{D((-\Delta)^{\frac{1}{2}})} = \left(\sum_{j\geq 1} \alpha_j |\langle x, \phi_j \rangle_X|^2\right)^{\frac{1}{2}}$ . Let p > 1, T > 0 and let  $u \in L^p(0, T; X)$ . It follows from the fact that  $(-\Delta)^{\frac{1}{2}} \in \mathcal{L}(X, X_{-1})$ , that the  $X_{-1}$ -valued integral  $\int_0^T S_{-1}(T-s)(-\Delta)^{\frac{1}{2}}u(s)ds$  is well-defined (see [6], Theorem 5.34). Moreover, since the semigroup  $(S(t))_{t\geq 0}$  is analytic, then so is  $((S_{-1}(t))_{t\geq 0})$ . This implies that  $S_{-1}(\frac{T-s}{2})(-\Delta)^{\frac{1}{2}}u(s) \in X$ ,  $\forall s \in [0,T)$  (see [6], p. 101). Then we have  $\int_0^T S_{-1}(T-s)(-\Delta)^{\frac{1}{2}}u(s)ds \in X$ , which gives the *p*-admissibility of B (see [14], Prop. 3.3 and [17], Lemma. 4.3.9). For all  $x \in L^2(\Omega)$ ,  $t \geq 0$  and  $j \geq 1$ , we have

$$\begin{aligned} |\langle \int_0^t S_0(t-s)gS(s)xds,\phi_j\rangle_X| &= \int_0^t \langle S_0(t-s)gS(s)x,\phi_j\rangle ds = |\int_0^t \langle gS(s)x,e^{-\alpha_j(t-s)}\phi_j\rangle_X ds| \\ &\leq \|g\|_{L^{\infty}(\Omega)}\|x\|_X \frac{1-e^{-\alpha_j t}}{\alpha_j}. \end{aligned}$$

We deduce that

$$|\langle S(t)x,\phi_j\rangle_X| \le e^{-\alpha_j t} \|x\|_X + \|g\|_{L^{\infty}(\Omega)} \frac{1 - e^{-\alpha_j t}}{\alpha_j} \|x\|_X, \ t \ge 0, \ j \ge 1.$$
(26)

*Now for any*  $x \in X$ *, we have* 

$$BS(t)x = \sum_{j\geq 1} \alpha_j^{\frac{1}{2}} \left\langle S(t)x, \phi_j \right\rangle_X \phi_j.$$

This combined with (26) implies that  $BS(t)x \in X$  for all  $x \in X$  and for any t > 0. Now, using the series expansion of BS(t)x for  $x \in X$ , we get

$$\langle BS(t)x, S(t)x \rangle_X = \sum_{j \ge 1} \alpha_j^{\frac{1}{2}} \langle S(t)x, \phi_j \rangle_X^2$$

$$\geq \|S(t)x\|_X^2$$

$$\geq \|S(T)x\|_X^2, \, \forall t \in [0,T].$$

It follows that the assumption (12) is fulfilled.

We conclude by Theorem II.3 that for  $\rho > 0$  small enough, the control  $\nu(t) = -\rho \mathbf{1}_{\{t \ge 0, x(t) \neq 0\}}$  guarantees the uniform exponential stability of the system (25).

**Example 2** Consider the following system

$$(S_0) \begin{cases} \frac{\partial}{\partial t}(\zeta,t) = \frac{\partial}{\partial \zeta}x(\zeta,t) - \alpha x(\zeta,t) + \nu(t)h(\zeta)x(\zeta,t) & \text{ in } (0,1) \times (0,\infty), \\ x(1,t) = 0 & \text{ in } (0,\infty), \\ x(\cdot,0) = x_0 \in L^2(0,1) \end{cases}$$

where  $X = L^2(0, 1)$ ,  $\alpha > 0$  and  $h \in L^{\infty}(0, 1)$  is such that  $h \ge c > 0$ , for some positive constant c. Here we can take  $A = \frac{d}{d\zeta} - \alpha$  id with domain  $D(A) := \{x \in H^1(0, 1) : x(1) = 0\}$ . The operator A is the generator of a contraction semigroup  $(S(t)_{t\ge 0})$  given by

$$(S(t)x)(\zeta) = \begin{cases} e^{-\alpha t}x(\zeta+t) & \text{if } \zeta+t \leq 1, \\ 0 & \text{else.} \end{cases}$$

According to previous theorems, the system  $(S_0)$  is exponentially stabilizable by the switching feedback control  $v(t) = -\rho \mathbf{1}_{\{t>0/x(t)\neq 0\}}$ . Indeed, here the semigroup S(t) is a contraction (so that ||S(t)|| is

decreasing) and the linear bounded operator  $B_1 := h$  id is a bounded linear operator  $(h \in L^{\infty})$  and satisfies the observation condition (since  $h \ge c > 0$ ). Let us now consider the following system

$$(S_1)\begin{cases} \dot{x}(t) = x_{\zeta}(t) - \alpha x(\zeta, t) + \nu(t)h(\zeta)x(t) & in(0, 1) \times (0, \infty) \\ x(1, t) + \epsilon \psi(x(t)) = 0 & in(0, \infty) \end{cases}$$

where  $\psi : X \to \mathbf{R}$  is a non null linear functional of X. This may be seen as a perturbed version of  $(S_0)$  on its boundary conditions. According to Riesz representation, one can assume that  $\psi(x) = \int_0^1 f(s)x(s)ds$ ,  $\forall x \in X$  for some  $f \in X - (0)$ .

We aim to show that under small valuers of  $\epsilon > 0$ , this system is still exponentially stabilizable. The system  $(S_1)$  can be reformulated as:

$$(S_2)\begin{cases} \dot{x}(t) = \mathcal{A}x(t) + \nu(t)h(\zeta)x(t) & in(0,1) \times (0,\infty) \\ x(0) = x_0 & in(0,1) \end{cases}$$

where  $\mathcal{A} : D(\mathcal{A}) \subset X \to X$  is defined by:

$$\mathcal{A}x := Ax - \epsilon hx, \ \forall x \in D(\mathcal{A}) := \left\{ x \in H^1(0,1), x(1) + \epsilon \psi(x) = 0 \right\}.$$

We claim that A is the generator of a strongly continuous semigroup on X. In order to verify this assertion, we will consider A as a perturbation of the generator A.

In order to write the system  $(S_2)$  in the form (1), let us consider the function  $\theta(\zeta) = \mathbf{1}(\zeta) := 1, \zeta \in X$ , which is such that  $A_m \theta = 0$ , and  $\theta(1) = 1$ , where  $A_m := \frac{d}{d\zeta}$  with domain  $D(A_m) := H^1(0, 1)$ . Let us introduce the following operator

$$Bx = hx - \psi(x)A_{-1}\theta, \ \forall x \in X$$

which is a one to one operator since we have  $\theta \notin D(A)$ .

In the sequel, we will verify the assumptions of Theorem II.4 and then conclude the stabilization of the perturbed system  $(S_1)$ .

• From the boundary conditions of  $(S_2)$ , we can see that

$$\forall x \in X, x \in D(\mathcal{A}) \Leftrightarrow x \in H^1(0,1) \text{ and } x + \epsilon \psi(x) \theta \in D(\mathcal{A})$$
.

*This together with the definition of*  $\theta$  *implies that for*  $x \in D(\mathcal{A})$ *, we have* 

$$X \ni \mathcal{A}x = A_m x - \epsilon h x$$
  
=  $A_m (x + \epsilon \psi(x)\theta) - \epsilon h x$   
=  $A (x + \epsilon \psi(x)\theta) - \epsilon h x$   
=  $A_{-1} (x + \epsilon \psi(x)\theta) - \epsilon h x$   
=  $A_{-1} x - \epsilon B x$   
=  $(A_{-1} - \epsilon B)|_X x$ 

Moreover, for all  $x \in D((A_{-1} - \epsilon B)|_X)$ , we have  $A_{-1}(x + \epsilon \psi(x)\theta) \in X$ , i.e.  $x + \epsilon \psi(x)\theta \in D(A) \subset H^1(0,1)$  which implies that  $x \in H^1(0,1)$ . Then we have  $(A_{-1} - \epsilon B)|_X x = Ax$ . In other words,

$$(\mathcal{A}, D(\mathcal{A})) = ((A_{-1} - \epsilon B)|_X, D(A_{-1} - \epsilon B)|_X)).$$

• The operator  $(A_{-1} - \epsilon B)_X$  is a generator if we can show that

$$\int_0^1 S_{-1}(1-r)\psi(u(r))A_{-1}\theta dr \in X,$$

or, equivalently

$$\int_0^1 S_{-1}(1-r)\mathbf{1}(.)\psi(u(r))dr \in D(A), \ \forall u \in L^2(0,1;X).$$

We have

$$\int_0^1 S_{-1}(1-r)\mathbf{1}(.)\psi(u(r))dr = \int_0^1 \psi(u(r))S(1-r)\mathbf{1}(.)dr$$
  
=  $\int_{.}^1 e^{-\alpha(1-r)}\psi(u(r))dr := g(.)$ 

Since  $\psi ou \in L^2(0,1)$ , this implies that  $g \in H^1(0,1)$  and g(1) = 0. In other words,  $g \in D(A)$ . Hence, for  $\epsilon > 0$  small enough, the system  $(S_1)$  is well-posed.

• Here we can take  $\mathfrak{X}_{-1} = span(A_{-1}\theta)$ , so we obtain an admissible decomposition for the pair  $(A_{-1}, -\epsilon B)$ . Indeed, it is clear that  $_{X}Bx = hx$ ,  $x \in X$ , so  $_{X}B$  is a bounded operator from X to X. Moreover, for all  $x \in D((A_{-1} - \epsilon B)|_{X}) = D((A_{-1} + \epsilon \psi(.)A_{-1}\theta)|_{X})$ , we have

$$A_{-1}x = A_{-1}(x + \epsilon \psi(x)\theta) - \epsilon \psi(x)A_{-1}\theta = A(x + \epsilon \psi(x)\theta) - \epsilon \psi(x)A_{-1}\theta,$$

from which it comes that

$$_{\mathbf{X}}(A_{-1})x = A(x + \epsilon \psi(x)\theta), \ \forall x \in D((A_{-1} - \epsilon B)|_{\mathbf{X}}).$$

where

$$D((A_{-1} - \epsilon B)|_X) = \{ x \in L^2(0, 1) / x + \epsilon \psi(x)\theta \in D(A) \}$$

Then for  $x \in D((A_{-1} - \epsilon B)|_X)$ , we have  $(A_{-1} - \epsilon B)x \in X$  or equivalently  $x + \epsilon \psi(x)\theta \in D(A)$ , and

$$\begin{aligned} \langle_{\mathbf{x}}(A_{-1})\mathbf{x},\mathbf{x}\rangle &= \langle A(\mathbf{x} + \epsilon \psi(\mathbf{x})\theta), \mathbf{x}\rangle \\ &= \langle A_m(\mathbf{x} + \epsilon \psi(\mathbf{x})\theta), \mathbf{x}\rangle \\ &= \langle A_m \mathbf{x}, \mathbf{x}\rangle \\ &= \int_0^1 \mathbf{x}'(s)\mathbf{x}(s)ds - \alpha \|\mathbf{x}\|^2 \\ &\leq \left(\frac{\epsilon^2 \|f\|^2}{2} - \alpha\right) \|\mathbf{x}\|^2 - \frac{1}{2}\mathbf{x}^2(0) \end{aligned}$$

Thus the operator  $_{X}(A_{-1})$  is dissipative in  $D((A_{-1} - \epsilon B)|_{X})$  for every  $0 < \epsilon \le \frac{(2\alpha)^{1/2}}{\|f\|}$ . • Finally, the observation estimate follows from the fact that  $h \ge c > 0$  and that for any  $x \in X$ , the

mapping  $t \mapsto ||S(t)x||$  is decreasing. We conclude by Theorem II.4 that for  $\epsilon > 0$  small enough, the control  $\nu(t) = -\epsilon \mathbf{1}_{\{t \ge 0: x(t) \neq 0\}}$  guarantees the exponentially stabilization of the system  $(S_1)$ .

# IV. CONCLUSION

In this paper we have shown that it is possible for a linear system with dissipative dynamic, to be exponentially stable under small Desch-Schapacher perturbations of the dynamic. The main assumptions of sufficiency are formulated in terms of admissibility and observability assumptions

of unbounded linear operators. An explicit decay rate of the stabilized state is given. The previous research on this problem concerned either bounded or Miyadera's type perturbations [11, 15]. The main stabilization result is further applied to show the uniform exponential stabilization of unbounded bilinear reaction diffusion and transport equations using a bang controller.

#### CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

# References

- Adler, M., Bombieri, M., & Engel, K. J. (2014). On Perturbations of Generators of-Semigroups. In Abstract and Applied Analysis (Vol. 2014). Hindawi.
- [2] Ammari, K., El Alaoui, S., & Ouzahra, M. (2020). Feedback stabilization of linear and bilinear unbounded systems in Banach space. SCL, 2021.
- [3] Berrahmoune, L. (2009). A note on admissibility for unbounded bilinear control systems. Bulletin of the Belgian Mathematical Society-Simon Stevin, 16(2), 193-204.
- [4] Boulouz, A., Bounit, H., & Hadd, S. (2021). Well-posedness and exponential stability of boundary control systems with dynamic boundary conditions. Systems & Control Letters, 147, p. 104825.
- [5] Desch, W., & Schappacher, W. (1989). Some generation results for perturbed semigroup, Semigroup Theory and Applications (Clémnet, Invernizzi, Mitidieri, and Vrabie, eds.). Lect. Notes Pure Appl. Math, 116, 125-152.
- [6] Engel, K. J., & Nagel, R. (2001, June). One-parameter semigroups for linear evolution equations. In Semigroup forum (Vol. 63, No. 2). Springer-Verlag.
- [7] G. Greiner, Perturbing the boundary conditions of a generator, Houston Journal of Mathematics, 13 (1987), 213–229.
- [8] Hadd, S., Manzo, R., & Rhandi, A. (2015). Unbounded perturbations of the generator domain. Discrete & Continuous Dynamical Systems-A, 35(2), 703.
- [9] Idrissi, A. (2003). On the unboundedness of control operators for bilinear systems. Quaestiones Mathematicae, 26(1), 105-123.
- [10] Kilbas, A. A., Srivastava, H. M., & Trujillo, J. J. (2003). Fractional differential equations: A emergent field in applied and mathematical sciences. In Factorization, singular operators and related problems (pp. 151-173). Springer, Dordrecht.
- [11] Liu, K., Liu, Z., & Rao, B. (2001). Exponential stability of an abstract nondissipative linear system. SIAM journal on control and optimization, 40(1), 149-165.
- [12] Metzler, R., & Klafter, J. (2000). The random walk's guide to anomalous diffusion: a fractional dynamics approach. Physics reports, 339(1), 1-77.
- [13] Maragh, F., Bounit, H., Fadili, A., & Hammouri, H. (2014). On the admissible control operators for linear and bilinear systems and the Favard spaces. Bulletin of the Belgian Mathematical Society-Simon Stevin, 21(4), 711-732.

- [14] Nagel, R., and E. Sinestrari. Inhomogeneous Volterra integrodifferential equations for Hille-Yosida operators, Functional Analysis Proc. Essen Conference (KD Bierstedt and A. Pietsch and WM Ruess and D. Vogt, eds.). (1994): 51-70.
- [15] Ouzahra, M. (2017). Exponential stability of nondissipative linear system in Banach space and application to unbounded bilinear systems. Systems & Control Letters, 109, 53-62.
- [16] A. Pazy, Semi-groups of linear operators and applications to partial differential equations, Springer Verlag, New York, 1983.
- [17] Van Neerven, J. (1992). The adjoint semigroup. In The Adjoint of a Semigroup of Linear Operators (pp. 1-18). Springer, Berlin, Heidelberg.
- [18] Weiss, G. (1989). Admissibility of unbounded control operators. SIAM Journal on Control and Optimization, 27(3), 527-545.