

Natural dynamical reduction of the three-body problem

Barak Kol

Racah Institute of Physics, Hebrew University, Jerusalem 91904, Israel

barak.kol@mail.huji.ac.il

ABSTRACT: The three-body problem is a fundamental long-standing open problem, with applications in all branches of physics, including astrophysics, nuclear physics and particle physics. In general, conserved quantities allow to reduce the formulation of a mechanical problem to fewer degrees of freedom, a process known as dynamical reduction. However, extant reductions are either non-general, or hide the problem's symmetry or include unexplained definitions. This paper presents a dynamical reduction that avoids these issues, and hence is general and natural.

Any three-body configuration defines a triangle, and its orientation in space. Accordingly, we decompose the dynamical variables into the geometry (shape + size) and orientation of the triangle. The geometry variables are shown to describe the motion of an abstract point in a curved 3d space, subject to a potential-derived force and a magnetic-like force with a monopole charge. The orientation variables are shown to obey a dynamics analogous to the Euler equations for a rotating rigid body, only here the moments of inertia depend on the geometry variables, rather than being constant.

The reduction rests on a novel symmetric solution to the center of mass constraint inspired by Lagrange's solution to the cubic. The formulation of the orientation variables is novel and rests on a little known generalization of the Euler-Lagrange equations to non-coordinate velocities. Applications to special exact solutions and to the statistical solution are described or discussed. Moreover, a generalization to the four-body problem is presented.

Contents

1	Introduction	1
2	Reduction	3
2.1	Translations and the complex position vector \vec{w}	3
2.2	Rotations and the orientation–geometry decomposition	5
2.3	Rotations within the plane and geometry space	6
2.4	Geometry of geometry space	9
2.5	Angular momentum variables	12
2.6	Discussion	16
3	Applications to planar motion	17
4	Four-body problem	19
A	Lagrange’s solution to the cubic	20
B	Bi-complex numbers and the isotropic oscillator	21
C	Equations of motion in terms of non-coordinate velocities and the rotating rigid body	23

To Joseph-Louis Lagrange

1 Introduction

The three body problem involves the motion of three point masses influenced by mutual forces which depend on their separations.

While the two-body problem is integrable, the three-body problem is in general non-integrable and chaotic. Its solution has been a puzzle ever since it was studied by Newton in an attempt to understand the motion of the moon [1]. The system is known to be rather rich, and along the centuries its study deeply influenced several scientific fields including perturbation theory, chaos and topology.

It is interesting to seek a dynamical reduction of the system: a simplified formulation, where the dynamical variables are chosen such that the equations of motion simplify. More specifically, the existence of conserved quantities allows to reduce the order of the system of differential equations. While a dynamical reduction is often applied to integrable systems, it could also apply to a non-integrable system, in which case the reduced system remains non-integrable.

This work originated from the recent flux-based reduction of the statistical solution to the three-body problem in Newtonian gravity [2], see also progress on the statistical solution through non-flux-based methods [3, 4]. The flux-based theory involves the (regularized) phase space volume as a function of the conserved charges. It is defined by a multi-dimensional integral, yet it was found that a certain change of variables enables analytic integration over numerous variables [2, 5]. This suggested the possibility that certain dynamical reductions may exist, involving the new variables, thereby providing a deeper reason for the success of the analytic integrations.

Over the centuries much research was devoted to the dynamical reduction of the three-body problem (3BP). Lagrange (1772) [6] reduced the equations' order through replacing the position vectors by variables describing the triangle geometry and additional variables. Jacobi (1842) [7] showed how to perform one additional reduction of order, known as the elimination of the nodes. Levi-Civita (1915) [8] was the first to discuss the reduction at the level of the action, namely using a Lagrangian or a Hamiltonian, carrying out the reduction to the instantaneous three-body plane. Murnaghan (1936) [9] performed a reduction of the planar 3BP at the level of the action, choosing side lengths as the dynamical variables. However, the side length variables become singular for collinear configurations. Van Kampen and Wintner (1937) [10] generalized [9] to 3d by adding the average azimuthal angle as a fourth coordinate. Lemaître (1952) [11] introduced a different set of symmetric coordinates, which remain smooth in the collinear limit, and presented a Hamiltonian for the general, non-planar case. More recently, Moeckel and Montgomery (2013) [12] introduced the term shape sphere, for a set of symmetric coordinates for the planar 3BP, see also [13, 14]. These coordinates are essentially equivalent to those introduced by Lemaître, though it is not immediately apparent. In addition, [12] recast the analysis into modern mathematical terminology.

The quantum version of the three body problem was studied in the context of the Helium atom (nucleus + 2 electrons = 3 bodies). Interestingly, some works on the quantum version reached new definitions of the dynamical variables and new forms for the quantum Hamiltonian, which in hindsight, are related with novel formulations of the non-quantum problem. Hylleraas (1929) [15] presented a Hamiltonian in terms of side lengths for the $\vec{J} = 0$ sector and furthermore setting two electron masses to be equal and the the nucleus mass to be infinite. Gronwall (1932) [16] introduced 3d variables which made the kinetic term conformally Euclidean. Fock (1954) [17] introduced 4d variables that reduce to Gronwall's variables after imposing the reduction to $\vec{J} = 0$. Finally, Tkachenko (1978) [18] presented a Hamiltonian for non-zero \vec{J} .

Despite this venerable history, certain features of the reduction are still lacking. First, the choice of coordinates that satisfy the center of mass constraint appears to violate the permutation symmetry (see section 2.1), which is unnatural. Secondly and relatedly, the shape sphere variables display somewhat surprising and undeserved properties (see last paragraph of section 2.6), which suggest that a more direct and natural route to their introduction might exist, which would make them clearer. Thirdly and lastly, extant formulations do not incorporate the angular momentum components as canonical variables, which would be natural given their central standing as conserved quantities.

In this paper, the reduction process is described in section 2. It starts by introducing a solution to the center of mass constraints, which transforms nicely under permutations, and hence is natural. It involves complex numbers in a novel way, and it is inspired by Lagrange’s solution to the cubic. Next, the orientation of the plane defined by the three bodies is separated from motion within the plane simply by using a rotating body frame. We proceed to the separation of rotations within the plane from the geometry of the triangle formed by the three bodies. This requires to define certain spherical coordinates on \mathbb{C}^2 and to handle correctly one of the angles, resulting in a famous quotient of it. Finally, we Legendre transform from angular velocities to angular momenta and obtain their Poisson brackets from a general theory for non-coordinate velocities and their conjugate momenta. This leads us to our final result, (2.49). We end the section with an illustration of the resulting formulation and a discussion of it.

The formulation can be applied to study both special exact solutions and the statistical solution. In section 3, we provide a novel derivation of the uniformly rotating solutions, and sketch a novel derivation of the linear stability region for the equilateral solution. The section ends with an outlook on applications to the statistical solution and more.

This work has interesting implications for our-body problem, and a sketchy generalization appears in section 4.

Appendices provide background material on Lagrange’s solution to the cubic, on bi-complex numbers and the isotropic oscillator and finally on the equations of motion in terms of non-coordinate velocities and the rotating rigid body.

2 Reduction

Setup. Consider three point masses, m_1, m_2, m_3 . Let us take the bodies’ position vectors, \vec{r}_a , $a = 1, 2, 3$, to be our initial dynamical variables. The kinetic energy is given by

$$T := \sum_{a=1}^3 \frac{1}{2} m_a \dot{\vec{r}}_a^2 \quad (2.1)$$

For concreteness, take the potential to be given by the Newtonian gravitational potential

$$V := -\frac{G m_1 m_2}{r_{12}} - \frac{G m_1 m_3}{r_{13}} - \frac{G m_2 m_3}{r_{23}} \quad (2.2)$$

where G is Newton’s gravitational constant, and $r_{ab} = |\vec{r}_a - \vec{r}_b|$, are the inter-body distances. We note that the whole reduction procedure is independent of the choice of V as long as it depends on r_{ab} only, namely $V = V(\{r_{ab}\})$. Finally, the system can be defined by the Lagrangian

$$\mathcal{L} \left(\{\vec{r}_a, \dot{\vec{r}}_a\}_{a=1}^3 \right) := T - V . \quad (2.3)$$

2.1 Translations and the complex position vector \vec{w}

The system is invariant under translations and hence the total linear momentum is conserved, the center of mass moves in uniform motion, and henceforth we work in the center

of mass frame

$$0 = \vec{r}_{CM} := \frac{1}{M} \sum_a m_a \vec{r}_a \quad (2.4)$$

where $M := m_1 + m_2 + m_3$ is the total mass.

The center of mass constraint reduces the configuration space, reducing the number of degrees of freedom from 9 to 6. This requires a choice coordinates on the reduced configuration space. We shall see that this choice creates a tension with the bodies' permutation symmetry, which will affect the whole reduction process, and hence we shall describe it in some detail.

The coordinates are chosen to be translation-invariant vectors: vectors, in order to preserve the transformation properties under rotations, and translation-invariant, in order to decouple them from the center of mass coordinates in the expression for the kinetic energy.

In the two-body problem, one chooses the relative coordinate to be

$$\vec{r}_1 - \vec{r}_2 \quad (2.5)$$

It is a translation-invariant vector, and under a $1 \leftrightarrow 2$ exchange, it changes only by a sign.

In the three-body problem, there are three relative vectors

$$\vec{r}_{12} := \vec{r}_1 - \vec{r}_2, \quad \vec{r}_{23} := \vec{r}_2 - \vec{r}_3, \quad \vec{r}_{31} := \vec{r}_3 - \vec{r}_1 \quad (2.6)$$

They satisfy the constraint $0 = \vec{r}_{12} + \vec{r}_{23} + \vec{r}_{31}$.

Any pair of linear combinations of the relative vectors can serve as coordinates in the center of mass frame. There are two popular choices for that. The first set consists of the planetary coordinates

$$\vec{r}_{13}, \quad \vec{r}_{23} \quad (2.7)$$

They are useful for the case where one of the masses, say m_3 , is much heavier than the other two: $m_1, m_2 \ll m_3$. The second set consists of the lunar coordinates

$$\vec{r}_{12}, \quad \frac{1}{m_1 + m_2} (m_1 \vec{r}_1 + m_2 \vec{r}_2) - \vec{r}_3 \quad (2.8)$$

These are useful for the hierarchical case where the magnitude of the first vector is much smaller than the second, namely bodies 1 and 2 are much closer to each other, relative to their distance from the third body. These coordinates are commonly called Jacobi coordinates, because they appeared in [7], but they are a rather natural choice, and they appeared already in [6].

Both coordinate choices break the symmetry between the bodies. Hence, they are not optimal for the non-hierarchical case where all distances and all masses are comparable.

Can we define natural, namely symmetric, coordinates in the center of mass frame? A solution is suggested by Lagrange's solution to the cubic equation. Lagrange realized that the key to a solution of a polynomial equation is to study expressions in terms of the roots that display a certain measure of symmetry among them. For more see appendix A. Inspired by this, we define

$$\vec{w} := \vec{r}_1 + \eta \vec{r}_2 + \bar{\eta} \vec{r}_3 \quad (2.9)$$

where η is the cubic root of unity, namely $\eta = \exp(2\pi j/3)$, and we denote here the imaginary unit by j , $j^2 = -1$, since i will have a different use below.

The definition (2.9) is one of the central results of this paper. Let us discuss it. By construction \vec{w} is a complex vector. Naturally, it is composed of 2 real vectors, and so it has the correct number of components to serve as coordinates in the center of mass frame. It is translation-invariant as a result of the identity $1 + \eta + \bar{\eta} = 0$. Under a cyclic rotation it is merely multiplied by a $2\pi/3$ phase, while the exchange $2 \leftrightarrow 3$ transforms it into its complex conjugate. *These simple transformations under permutations render \vec{w} a natural coordinate.*

Transformed Lagrangian Given \vec{w} , the relative positions can be expressed by

$$\vec{r}_{23} = \frac{1}{j\sqrt{3}} (\vec{w} - \vec{\bar{w}}) \quad (2.10)$$

and similarly, replacing $w \rightarrow \eta w$ in this expression gives \vec{r}_{12} , and $w \rightarrow \bar{\eta} w$ gives \vec{r}_{31} .

The kinetic energy in the center of mass frame is given by

$$\begin{aligned} T_{CM} &:= T - \frac{M}{2} \dot{\vec{r}}_{CM}^2 = \frac{1}{2M} \left(m_2 m_3 \dot{\vec{r}}_{23}^2 + cyc. \right) = \\ &= \frac{1}{6M} \left(m_2 m_3 \left\| \frac{d}{j dt} (\vec{w} - \vec{\bar{w}}) \right\|^2 + cyc. \right) \end{aligned} \quad (2.11)$$

where the cyclic transformation denotes now $w \rightarrow \bar{\eta} w$, in addition to $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$.

2.2 Rotations and the orientation–geometry decomposition

One can naturally divide the system's coordinates into orientation coordinates and coordinates that describe the configuration up to rotations, namely, geometry coordinates. At any given moment, the positions of the three bodies define an instantaneous plane. We define a rotating body frame whose z axis is normal to the instantaneous plane, and 2d vectors $\vec{\rho}_a$ specify the positions within the rotating (x, y) plane. As usual, inertial frame velocities are given by

$$\left. \frac{d}{dt} \right|_{inert} \vec{\rho}_{ab} = \dot{\vec{\rho}}_{ab} + \vec{\omega} \times \vec{\rho}_{ab} \quad (2.12)$$

where $\vec{\omega}$ is the angular velocity vector that describes the rotation of the body frame, and ω is reserved for later use.

Substituting, the kinetic energy (2.11) becomes

$$T = \frac{1}{2} I_{ij} \tilde{\omega}^i \tilde{\omega}^j + \vec{L}_w \cdot \vec{\omega} + T_w \quad (2.13)$$

where I_{ij} , $i, j = 1, 2, 3$ denotes the inertia tensor in the rotating (and center of mass) frame

$$I_{ij} = \frac{1}{M} \left[m_2 m_3 \left(\delta_{ij} \vec{\rho}_{23}^2 - \rho_{23}^i \rho_{23}^j \right) + cyc. \right] ; \quad (2.14)$$

\vec{L}_w denotes the the angular momentum due to the $\vec{\rho}_a$ motion

$$\begin{aligned} \vec{L}_w &= L_w \hat{z} \\ L_w &= \frac{1}{M} \left(m_2 m_3 \vec{\rho}_{23} \wedge \dot{\vec{\rho}}_{23} + cyc. \right) , \end{aligned} \quad (2.15)$$

where the wedge product between any 2d vectors \vec{u}, \vec{v} is defined by $\vec{u} \wedge \vec{v} := (\vec{u} \times \vec{v}) \cdot \hat{z}$; and finally, the kinetic energy due to the $\vec{\rho}_a$ motion

$$T_w := \frac{1}{M} \left(m_2 m_3 \dot{\rho}_{23}^2 + \text{cyc.} \right) . \quad (2.16)$$

We note several properties of I^{ij} . Since the mass distribution is confined to the $z = 0$ plane one has

$$\begin{aligned} 0 &= I_{13} = I_{23} \\ I &:= I_{33} = I_{11} + I_{22} \end{aligned} \quad (2.17)$$

In addition, since the mass distribution consists of only 3 point masses, one has

$$I_{11} I_{22} - (I_{12})^2 = \frac{m_1 m_2 m_3}{M} \Delta^2 \quad (2.18)$$

where

$$\Delta := \det [x_1, y_1, 1; x_2, y_2, 1; x_3, y_3, 1] \quad (2.19)$$

is twice the (signed) area of the triangle formed by the bodies. In other words, the 2×2 determinant of $I_{\alpha\beta}$, $\alpha, \beta = 1, 2$ factors into a part that depends only on positions, and a part which depends only on the masses.

2.3 Rotations within the plane and geometry space

The orientation–geometry decomposition is not done yet. To see that, let us count the number of generalized coordinates. We know that the 3d problem has 6 degrees of freedom (d.o.f) in the center of mass frame (CM). The orientation is specified by 3 angles, such as the Euler angles, and the $\vec{\rho}_a$ at CM (or equivalently the associated 2d \vec{w}) provide 4 other d.o.f., so altogether we have 7 generalized coordinates. This means that these coordinates are redundant by a single degree of freedom.

Rotations within the (x, y) plane are the origin of this redundancy: since a rotation of the body frame around the z -axis is equivalent to a rotation of the 2d vectors, the configuration depends on the two associated angles only through their sum.

Bi-complex w . In order to account for plane rotations, it is useful to complexify the plane, so that rotations would be represented by a phase multiplication. By complexification one means that any 2 vector $\vec{\rho}$ is mapped onto a complex number $\vec{\rho} \rightarrow \rho := \vec{\rho} \cdot (\hat{x} + i \hat{y})$. Similarly, the 2d j -complex vector \vec{w} , defined in (2.9), is mapped onto a so-called bi-complex number

$$\vec{w} \rightarrow w = \vec{w} \cdot (\hat{x} + i \hat{y}) \quad (2.20)$$

which is of the form $w = a + b i + c j + d i j$. The imaginary unit i represents a quarter rotation in the (x, y) plane. Hence, i, j are commuting imaginary units, namely $i^2 = j^2 = -1$ and $i j = j i$. For more on the algebraic structure of bi-complex numbers, and their application to the isotropic oscillator of mechanics, see appendix B.

Let us introduce a natural basis for the w space. Evaluating the w variable (2.20) for a right-handed and a left-handed equilateral triangles of unit sides motivates the definitions

$$e_R := \frac{\sqrt{3}}{2} (1 + ij) \quad e_L := \frac{\sqrt{3}}{2} (1 - ij) , \quad (2.21)$$

where for concreteness, the equilateral triangles are chosen to be oriented such that the height from edge 2—3 to vertex 1 is pointed along $+\hat{x}$ within the (x, y) plane.

Spherical coordinates for w . Considering w space as a 2d complex vector space over complex numbers involving i , namely $\mathbb{C}[i]^2$, one can expand a general state w in the e_R, e_L basis as follows

$$w = r e^{i\psi_0} \left[e^{i\phi/2} \cos \frac{\theta}{2} e_R + e^{-i\phi/2} \sin \frac{\theta}{2} e_L \right] \quad (2.22)$$

In this expression, r sets the overall triangle scale, $\psi \sim \psi + 2\pi$ is an overall rotation angle, $\theta \in [0, \pi]$ determines the relative magnitudes of the right and left components, and finally $\phi \sim \phi + 2\pi$ is the relative phase of the right and left components. Altogether r, θ, ϕ, ψ are spherical coordinates for $w \in \mathbb{C}[i]^2$.

We define G to be the quotient space of planar three-body configurations, up to rotations, where G stands for geometry. In other words, G is the space of equivalence classes of congruent triangles. We can write

$$G = \mathbb{C}[i]^2 / U(1) \quad (2.23)$$

where the $U(1)$ acts by overall phase rotations. We shall later see how to parameterize G in terms of quadratics in w . In coordinates, G is given by the w variable up to ψ -shifts, and it is parameterized by r, θ, ϕ . G includes information on both size and shape: the r coordinate determines the triangle size, while the θ, ϕ coordinates parameterize a sphere that describes triangles up to similarity, and is known as the *shape sphere*. Coordinates that are equivalent to r, θ, ϕ appeared in [11] in Eqs. (1,2,14), and the term “shape sphere” appeared in [12].

Substituting (2.22) into (2.10), the relative position vector r_{23} becomes

$$\rho_{23} = i r e^{i\psi_0} \left[e^{i\phi/2} \cos \frac{\theta}{2} - e^{-i\phi/2} \sin \frac{\theta}{2} \right] . \quad (2.24)$$

In particular,

$$r_{23}^2 \equiv |\rho_{23}|^2 = r^2 (1 - \sin \theta \cos \phi) . \quad (2.25)$$

Similarly, r_{31}, r_{12} are given by the same expression after the substitutions $\phi \rightarrow \phi - 2\pi/3$, and $\phi \rightarrow \phi + 2\pi/3$, respectively. This means that expressions are invariant under a cyclic permutation combined with a third of a revolution in ϕ .

The definition of ψ_0 can be changed into $\psi = \psi_0 + \chi(\theta, \phi)$, which will be seen to be a gauge transformation. ψ_0 , defined in (2.22), is symmetric between the R and L hemispheres, $0 \leq \theta \leq \pi/2$ and $\pi/2 \leq \theta \leq \pi$, respectively. However, ψ_0 is singular for both R and L poles. A gauge that is regular at the R pole is given by

$$\psi_+ = \psi_0 + \phi/2 \quad (2.26)$$

Kinetic energy. In spherical coordinates, and using the ψ_+ gauge (2.26), T_w (2.16) becomes

$$T_w = \frac{1}{2} I \left(\dot{\psi}_+ + \frac{L_G}{I} \right)^2 + T_G \quad (2.27)$$

where the largest principal moment of inertia (for rotations within the plane) (2.14), is given by

$$I := I_{33} = \frac{r^2}{M} [m_2 m_3 (1 - \sin \theta \cos \phi) + cyc.] \quad (2.28)$$

L_G denotes the angular momentum in the geometry space G and it is given by

$$L_G = -\frac{r^2}{2M} \left[m_2 m_3 \left((1 - \cos \theta - \sin \theta \cos \phi) \dot{\phi} - \sin \phi \dot{\theta} \right) + cyc. \right] \quad (2.29)$$

where *cyc.* denotes $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ together with $\phi \rightarrow \phi - 2\pi/3$, just like the comment after (2.25). L_G is related to L_w , defined in (2.15), by

$$L_w = I \dot{\psi}_+ + L_G \quad (2.30)$$

T_G denotes the kinetic energy in G space, and is given by

$$T_G := \frac{1}{8I} \dot{I}^2 + \frac{3M_3}{8M} \frac{r^4}{I} \left(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2 \right) \quad (2.31)$$

Finally, M, M_2, M_3 denote the elementary symmetric functions of the masses

$$\begin{aligned} M &:= m_1 + m_2 + m_3 \\ M_2 &:= m_2 m_3 + m_3 m_1 + m_1 m_2 \\ M_3 &:= m_1 m_2 m_3 \end{aligned} \quad (2.32)$$

The derivation proceeds by projecting $\dot{\rho}_{23}$ along the radial and the tangential directions with respect to ρ_{23} , performed in the r, θ, ϕ coordinates, and similarly for the other relative velocities. It will be detailed in a later version.

The kinetic energy (2.27) is in a form of a dimensional reduction (also known as Kaluza-Klein reduction) over the coordinate ψ . Therefore, the expression for L_G depends on the choice of gauge for ψ , while the kinetic metric T_G is gauge-independent, and represents the metric on the quotient space (2.23). The expression for T_G appeared essentially in [12].

Fixing the ψ gauge. Let us return to the issue of coordinate redundancy discussed at the beginning of this subsection. This redundancy can be removed by fixing a gauge for the ψ coordinate, and we choose to set

$$\psi = 0 \quad (2.33)$$

This can always be achieved through a choice of the x – y body axes, which in turn fixes the value of $\tilde{\omega}_z$. Clearly, the effect of this gauge is directly related to the choice of definition of ψ .

Relatedly, the kinetic energy depends on $\tilde{\omega}_z, \dot{\psi}$ only through their sum. Indeed, the relevant terms in T are (2.13,2.27)

$$\begin{aligned} T &\supset \frac{1}{2}I(\tilde{\omega}_3)^2 + L_w \tilde{\omega}_3 + \frac{1}{2}I\left(\dot{\psi} + \frac{L_G}{I}\right)^2 = \\ &= \frac{1}{2}I\left(\tilde{\omega}_3 + \dot{\psi} + \frac{L_G}{I}\right)^2 = \frac{1}{2}I\left(\omega_3 + \frac{L_G}{I}\right)^2 \end{aligned} \quad (2.34)$$

where in passing to the second line, we have used (2.30), and in the last equality we have used $\vec{\omega}$ defined by

$$\vec{\omega} = \vec{\tilde{\omega}} + \dot{\psi} \hat{z} . \quad (2.35)$$

Altogether, the expression for the kinetic energy in the center of mass frame becomes

$$T[\vec{\omega}, r, \theta, \phi] = T_G + \frac{1}{2}I\left(\omega_3 + \frac{L_G}{I}\right)^2 + \frac{1}{2}I_{\alpha\beta}\omega^\alpha\omega^\beta \quad (2.36)$$

where $T_G, I, L_G, I_{\alpha\beta}$ $\alpha, \beta = 1, 2$ depend on the G -space variables and were defined in (2.31,2.28,2.29,2.14) respectively.

2.4 Geometry of geometry space

Let us study further the geometry of geometry space, namely three-body configurations up to rotations (2.23).

Invariants. So far, geometry space was described by the coordinates r, θ, ϕ . Alternatively, we can employ the invariants of the quotient. This approach will clarify the geometry at the origin of geometry space.

We define the quadratic invariants

$$Q_s := \begin{bmatrix} w_1^* & w_2^* \end{bmatrix} \tau_s \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \quad s = 0, 1, 2, 3, 4 \quad (2.37)$$

where $w_1, w_2 \in \mathbb{C}[i]$ denote the j -real and imaginary parts of the bi-complex w , up to normalization, defined as follows

$$w = \sqrt{\frac{3}{2}}(w_1 + j w_2) \quad (2.38)$$

and the Pauli-like matrices τ_r are given by

$$\tau_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \tau_1 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \tau_2 = \frac{1}{2} \begin{bmatrix} -1 & -\sqrt{3} \\ -\sqrt{3} & 1 \end{bmatrix}, \quad \tau_3 = \frac{1}{2} \begin{bmatrix} -1 & \sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix}, \quad \tau_4 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \quad (2.39)$$

Since Q_s are of the form $\sim w^* w$ they are manifestly invariant under the rotations $w \rightarrow \exp(i\psi) w$.

The quadratic invariants Q_0, \dots, Q_4 are not all independent, but rather satisfy the relations

$$0 = Q_1 + Q_2 + Q_3 \quad (2.40a)$$

$$Q_0^2 = \frac{2}{3} (Q_1^2 + Q_2^2 + Q_3^2) + Q_4^2 \quad (2.40b)$$

Geometrically, relation (2.40a) means that Q_1, Q_2, Q_3 describe a flat 2d space, and relation (2.40b) means that the quotient is a (light) cone in a 3+1 space. The cone singularity at the origin corresponds to a triple collision configuration.

We note that the Q_0, \dots, Q_4 variables are closely related to the Stokes parameters S_0, \dots, S_3 [20], which are used to describe states of polarization of light and are reviewed in appendix B. More precisely, Q_0, Q_4 are identical to the S_0, S_3 while Q_1, Q_2, Q_3 up to the relation (2.40a) are analogous to S_1, S_2 . However, the Q variables are distinguished by being compatible with a symmetry of order 3.

In terms of the r, θ, ϕ coordinates, substitution of (2.22) shows that the invariants are given by

$$Q_0 = r^2 \quad (2.41a)$$

$$Q_a = r^2 \sin \theta \cos \phi_a, \quad a = 1, 2, 3 \quad (2.41b)$$

$$Q_4 = r^2 \cos \theta \quad (2.41c)$$

where

$$\phi_a := \phi - (a - 1)2\pi/3 . \quad (2.42)$$

In this form the invariants are seen to be closely related to Cartesian coordinates associated with the spherical coordinates r^2, θ, ϕ .

Alternatively, one can define Cartesian coordinates on G -space, by considering r^2, θ, ϕ to be spherical coordinates of a vector \vec{G} , namely $G_1 = r^2 \sin \theta \cos \phi$, $G_2 = r^2 \sin \theta \sin \phi$, $G_3 = r^2 \cos \theta$. The three components of \vec{G} are independent coordinates, which solve the relations on the Q variables (2.40). Similarly, we define the vector \vec{g} to be the Cartesian coordinates associated with r, θ, ϕ (r instead of r^2).

Using (2.25), the triangle edge lengths can now be expressed as

$$r_{23}^2 = Q_0 - Q_1 , \quad (2.43)$$

and similarly for r_{31}^2, r_{12}^2 . Summing all three, and using (2.40a, 2.41a) we find

$$r^2 = \frac{1}{3} (r_{23}^2 + r_{31}^2 + r_{12}^2) . \quad (2.44)$$

This means that the geometric interpretation of the radial coordinate r is the root mean square of the side lengths.

We note that one could take a wider perspective, and rather than study the planar configuration of the bi-complex w up to phase shifts, one could study the 3d complex position vector \vec{w} (2.9) up to identification by 3d rotations, namely $\vec{w}/SO(3)$. The quadratic

invariants Q_0, \dots, Q_3 can be expressed in terms of scalar products among \vec{w}, \vec{w} , and hence are invariants from the 3d perspective. On the other hand, Q_4 cannot be described in this way, and hence while it is an invariant of $\mathbb{C}^2/U(1)$, only Q_4^2 is an invariant of $\vec{w}/SO(3)$.

In addition, we note that Q_4 is proportional to the triangle area, more precisely

$$Q_4 = \frac{2}{\sqrt{3}} \Delta \quad (2.45)$$

where Δ is twice the (signed) triangle area, and was defined in (2.19). This can be shown by following the definition of Q_4 and using translation invariance to set $\vec{p}_1 = 0$.

Shape sphere. Geometry up to size is known as shape. For this reason, the unit sphere in geometry space, namely $r = 1$, is known as shape sphere [12].

For triangles, shape is the same as classification up to similarity, which is well known to be classified by the values of the three angles which are constrained to sum to π . This means that shape space should be a 2d surface. As we have seen, it turns out that this surface has the topology of sphere. More precisely, the shape sphere is the space of shapes of triangles with labeled vertices.

The shape sphere is shown in fig. 1. The geometrical interpretation of various locations on the shape sphere is known [12], and will be repeated here for convenience. The vertical Q_4 coordinate is proportional to the triangle area, as stated in (2.45). Hence, the $Q_4 = 0$ equator corresponds to collinear configurations. Within the collinear equator, the point $Q_1 = 1$ implies $w_2 = 0$ and hence it corresponds to the collision of the 2,3 vertices and it is denoted by $C1$. Similarly for $C2, C3$. Going away from the equator, the $Q_4 = 1$ pole is associated with $w = 1 + ij$, which corresponds to a right equilateral triangle, namely, such the 1,2,3 vertices are oriented in the positive mathematical direction (counter-clockwise). Similarly, the $Q_4 = -1$ pole corresponds to a left equilateral triangle.

Just as the Q invariants were noted above to be closely related to the Stokes parameters and the Pauli matrices (Pauli, 1927) [22], the shape sphere is closely related to the Poincaré sphere, the sphere of polarizations that was introduced in Poincaré's lectures on optics [21]. These notions are also closely related to the Hopf fibration (Hopf, 1931) [23] and the Bloch sphere (Bloch 1946) [24]. The Pauli matrices are used to describe quantum operators on the states of a spin-half particle. The Hopf fibration represents the 3-sphere as a circle fibration over the 2-sphere. The Bloch sphere describes the states of a two-state quantum system, through an analogy with the spin-half system. Mathematically, the $\mathbb{C}^2/U(1)$ quotient is at the root of all of the above-mentioned topics.

We record the form that several quantities, which appear in the motion potentials, obtain after transformation to the Q variables

$$\begin{aligned} I &= \frac{1}{M} [m_2 m_3 (Q_0 - Q_1) + cyc.] \\ L_G &= - \frac{[m_2 m_3 ((Q_1 - Q_0 - Q_4) \frac{d}{dt}(Q_2 - Q_3) - (Q_2 - Q_3) \frac{d}{dt}(Q_1 - Q_0 - Q_4)) + cyc.]}{2\sqrt{3} M (Q_0 + Q_4)} \end{aligned} \quad (2.46)$$




Figure 1. The shape sphere describes the space of all possible labeled triangles in a plane up to similarity. See text for the description of various locations on it. The vertical direction represents Q_4 , while the horizontal directions represent Q_1, Q_2, Q_3 constrained by (2.40a).

2.5 Angular momentum variables

In order to incorporate into the reduction the conservation of angular momenta, we transform from angular velocity variables into the conjugate momenta, which are non other than the angular momenta

$$J_i = \frac{\partial T}{\partial \omega^i} \quad i = 1, 2, 3. \quad (2.47)$$

We have arrived at the final set of dynamical variables, namely

$$\vec{J}, \vec{g} \quad (2.48)$$

where \vec{g} is a location in geometry space, which usually would be represented by the spherical coordinates r, θ, ϕ .

Performing a Legendre transform over the Lagrangian (2.3) we obtain

$$\mathcal{L}_J := \mathcal{L} - \vec{\omega} \cdot \vec{J} = \mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2. \quad (2.49)$$

\mathcal{L}_J is a function of the dynamical variables (2.48) (together with the generalized velocities associated with \vec{g}). The expression for \mathcal{L}_J is organized into three parts $\mathcal{L}_0, \mathcal{L}_1, \mathcal{L}_2$ according to powers of \vec{J} , and their values are detailed below.

The transform that defined \mathcal{L}_J was taken only with respect to part of the velocity variables, and therefore \mathcal{L}_J is a hybrid of Lagrangian and a Hamiltonian (sometimes known as Routhian). Later, we shall see that the equations of motion can be derived from \mathcal{L}_J . It is useful to have a term that refers to any kind of such function, whether it is a Lagrangian, a Hamiltonian or hybrid. In mechanics, the standard potential function encodes the forces

through derivatives. In thermodynamics, one uses one of several *thermodynamic potentials*, such as the energy, the free-energy, the Gibbs free energy, all related among themselves by Legendre transforms. Similarly, we shall use the term *motion potential* to refer to any function from which the equations of motion can be derived. From this perspective, the standard potential can be distinguished by the term *force potential*.

The first term appeared already in (2.2,2.31) and is repeated here for convenience. It depends only on the geometry space variables

$$\mathcal{L}_0 := \frac{1}{8I} \dot{I}^2 + \frac{3M_3 r^4}{8M I} \left(\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2 \right) - \left(\frac{Gm_2 m_3}{r_{23}} + cyc. \right) \quad (2.50)$$

where

$$I = \frac{1}{M} (m_2 m_3 r_{23}^2 + cyc.) \quad (2.51)$$

and

$$\begin{aligned} r_{23}^2 &= r^2 (1 - \sin \theta \cos \phi) \\ r_{31}^2 &= r^2 (1 - \sin \theta \cos(\phi - 2\pi/3)) \\ r_{12}^2 &= r^2 (1 - \sin \theta \cos(\phi - 4\pi/3)) \end{aligned} \quad (2.52)$$

The first two terms of \mathcal{L}_0 specify the kinetic energy on geometry space – in the radial and angular directions, respectively. The last term is minus the potential, and for concreteness we present the case of a Newtonian gravitational potential.

The \mathcal{L}_1 term couples geometry space and the rotating body, and it is given by

$$\mathcal{L}_1 := J_3 \frac{L_G}{I} \quad (2.53)$$

where L_G is given by

$$L_G = -\frac{r^2}{2M} \left[m_2 m_3 \left((1 - \cos \theta - \sin \theta \cos \phi) \dot{\phi} - \sin \phi \dot{\theta} \right) + cyc. \right] \quad (2.54)$$

cyc. denotes $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ together with $\phi \rightarrow \phi - 2\pi/3$. The expression for L_G is given in the ψ_+ gauge (2.26) and a gauge transformation with a gauge-function χ could shift it by $\Delta \mathcal{L}_1 = -J_3 \dot{\chi}$. In geometry space, \mathcal{L}_1 is analogous in form to a coupling of the motion in G -space to a vector potential, as noticed by [11]. This means that the motion in G -space experiences a velocity-dependent force, which is magnetic-like in form, and can be thought to originate from a Coriolis force.

The magnetic-like two-form is given by

$$B := d \left(J_3 \frac{L_G dt}{I} \right) = \frac{3}{2} M M_3 J_3 \frac{\sin \theta d\theta d\phi}{[m_2 m_3 (1 - \sin \theta \cos \phi) + cyc.]^2} \quad (2.55)$$

It can be seen that this 2-form does not involve the radial coordinate r . It carries non-zero magnetic monopole charge, proportional to J_3 and located at $\vec{g} = 0$. The monopole charge is a consequence of the intrinsic charge associated with the Hopf fibration, which is used

to reduce over ψ rotations. The author is not familiar with previous appearances of the expression for the magnetic field.

On the side of the rotating body, \mathcal{L}_1 will be understood to generate a precession of \vec{J} around the z -axis.

The \mathcal{L}_2 term also couples geometry space and the rotating body, only this term is quadratic in \vec{J} and is given by

$$-\mathcal{L}_2 := \frac{1}{2I} J_3^2 + \frac{M}{2M_3} \frac{1}{\Delta^2} \bar{I}^{\alpha\beta} J_\alpha J_\beta \quad (2.56)$$

where Δ is twice the triangle's area, which was defined around (2.18), and it can be expressed in G -variables through

$$\Delta = \frac{3}{4} r^4 \cos^2 \theta, \quad (2.57)$$

and where $\bar{I}^{\alpha\beta}$ is the inverse of the $I_{\alpha\beta}$ matrix and is given by

$$\bar{I}^{\alpha\beta} := \frac{1}{M} \left(m_2 m_3 \rho_{23}^\alpha \rho_{23}^\beta + \text{cyc.} \right) \quad (2.58)$$

where ρ_{23} is given within the ψ_0 gauge by

$$\rho_{23} = i r \left[\cos \frac{\theta}{2} - e^{-i\phi} \sin \frac{\theta}{2} \right]. \quad (2.59)$$

In geometry space, minus \mathcal{L}_2 is interpreted as the *centrifugal potential*. It can be thought to generalize the familiar 2-body centrifugal potential

$$V_{cent,2\text{ body}} = \frac{L^2}{2\mu r^2} \quad (2.60)$$

where \vec{L} is the system's angular momentum, and μ is its reduced mass. Indeed, 2-body motion is necessarily planar and hence $J_1 = J_2 = 0$. Moreover, $I = \mu r^2$, thereby the 3-body expression for the centrifugal potential reduces to that of the 2-body.

As usual, a coupling in the motion potential, implies several terms in the equations of motion. In this case, in addition to a centrifugal force acting on the geometric variables \vec{g} , we shall see that \mathcal{L}_2 also implies the Euler equations for the rotating body.

Equations of motion. We performed several natural changes of variables in the kinetic energy in order to re-formulate the problem, and thereby re-phrase the equations of motion. One possibility to achieve the equations of motion would be to express $\vec{\omega}$ in (2.36) in terms of a set of frame orientation angles, such as the Euler angles, which would be used as the fundamental dynamical variables. However, this procedure requires to make an arbitrary choice of the Euler-like angles, and obscures the rotational symmetry.

Alternatively, it would be nice to derive the equations of motion using $\vec{\omega}$ as fundamental velocities variables. However, the standard Euler-Lagrange equations would not produce the correct equations of motion, as can be seen for the example of the rigid body in

appendix C. Indeed, if we write $\omega^i = \beta_j^i \dot{q}^j$, where q^i are generalized coordinates, then $\beta^i \equiv \beta_j^i dq^j$ are 1-forms over G -space, such that $d\beta^i \neq 0$ and hence β^i cannot be expressed as a differential of any generalized coordinates. In other words, β^i define a non-coordinate basis of differential forms, also known as a non-holonomic basis.

In order to derive the equations of motion from a Lagrangian expressed in terms of non-coordinate velocities, we rediscovered the appropriate generalization of the Euler-Lagrange equations. This little-known generalization was originally found by Poincare in 1901 [19] and it is described in Appendix C. Here we only state the result for the case at hand.

In a Lagrangian formulation, one takes

$$\mathcal{L}(\vec{\omega}, \vec{g}) = T(\vec{\omega}, \vec{g}) - V(\vec{g}) \quad (2.61)$$

where $T(\vec{\omega}, \vec{g})$ is given by (2.36) and $V(\vec{g})$ is an arbitrary potential. The above-mentioned differential 1-forms have the following non-zero exterior differentials

$$d\beta^i = \frac{1}{2} \epsilon_{ijk} \beta^k \beta^j \quad (2.62)$$

and hence the equations of motion read

$$\begin{aligned} \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \omega^i} \right) + \epsilon_{ijk} \omega^j \frac{\partial \mathcal{L}}{\partial \omega^k} &= 0 \\ \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}^r} \right) &= \frac{\partial \mathcal{L}}{\partial q^r} \end{aligned} \quad (2.63)$$

where $q^r = (r, \theta, \phi)$, the spherical coordinates in geometry space. The first line describes 3 generalized equations of motion which originate from variations with respect to ω , while the second line describes 3 standard Euler-Lagrange equations of motion that originate from variation with respect to the G -space variables.

In terms of the partial Legendre transform \mathcal{L}_J (2.49), the equations of motion are given by

$$\begin{aligned} \frac{d}{dt} J_i &= -\{J_i, \mathcal{L}_J\} \\ \frac{d}{dt} \left(\frac{\partial \mathcal{L}_J}{\partial \dot{q}^r} \right) &= \frac{\partial \mathcal{L}_J}{\partial q^r} \end{aligned} \quad (2.64)$$

where the Poisson brackets among the dynamic variables originate from the non-coordinate nature of the corresponding velocities according to the general rule (C.9), and in our case are given by

$$\{J_i, J_j\} = -\epsilon_{ijk} J_k \quad (2.65)$$

After a solution $\vec{J} = \vec{J}(t)$, $\vec{g} = \vec{g}(t)$ is found, one can further integrate to obtain the frame orientation angles.

Note that \vec{J}^2 is conserved as a result of (2.49, 2.64). Hence phase space is essentially 8 dimensional: 6 dimensions for \vec{g} and its generalized velocities and 2 additional dimensions for \vec{J} that lies on a sphere of fixed \vec{J}^2 .

We comment that one could obtain a formulation where the vector potential is replaced by the gauge-invariant field-strength by performing a Legendre transform on the remaining velocity variables, and employing “covariant momenta” (in analogy with the covariant derivative).

2.6 Discussion

The motion potential $\mathcal{L}_J = \mathcal{L}_J(\vec{J}, \vec{g})$ given by (2.49) defines a reformulation of the three-body problem, and it is a central result of this paper. The dynamic variables \vec{J}, \vec{g} are compatible with the conserved charges in such a way as to reduce the number of effective degrees of freedom.

The employed dynamical variables decompose into two sets. The first set consists of the \vec{J} variables that describe the angular momentum in the body frame. They are the momenta canonically conjugate to $\vec{\omega}$, the angular velocity vector in the body frame. Hence, \vec{J} describes the rotational motion of the configuration triangle. The second set of variables consists of \vec{g} or in spherical coordinates (r, θ, ϕ) . These describe the geometry of the triangle, namely its shape and size, see fig. 1. The two sets are coupled as the geometry of the triangle determines its moment of inertia, which affects its rotation motion.

The corresponding decomposition of the mechanics into a rotational motion and a motion in 3d geometry space is illustrated in figure 2.

Let us look for the origin of the chaotic nature of the system in terms of these two components. Already in the limit of planar motion, the rotational component of the motion becomes trivial, yet the system is chaotic. Indeed, the motion in geometry space is non-integrable since it has 3 degrees of freedom and only a single conserved charge, namely the energy. Coupling to a rotating body cannot change this non-integrable nature. Moreover, given that the motion of a rotating body is integrable in the rigid body limit, *geometry space can be considered to be the core chaotic component in the three-body system.*

Various ingredients of this formulation have already appeared in the literature. The r, θ, ϕ coordinates together with the vector potential appeared in [11]. The kinetic term in shape space, which is conformal to the round sphere, appeared in [12] together with the origin of geometry space from a quotient of a 4d space. In hindsight, these elements partially appeared also in the Helium atom context: the 3d geometry space and its conformal equivalence to the sphere in [16], and the 4d predecessor in [17].

The current paper includes two main novelties. First, the definition of the complex position vector \vec{w} that solves the center of mass constraint (2.9). Secondly, the formulation in terms of \vec{J} , the angular momentum components in the rotating system, see subsection 2.5. In comparing to previous work, [12] was limited to planar motion, while we address the full 3d problem. \vec{J} has not been used as dynamical variables before, and in particular neither in [11] nor in [12]. While in hindsight, [18] can be recognized to have some relation to a formulation in terms of \vec{J} within the Helium atom context, it is at most partial and special due to the special values of the masses and the distinct quantum context.

Finally, the definition of \vec{w} naturally explains certain features that were previously observed yet deemed surprising. [11] introduces the spherical coordinates in geometry space without explanation. It starts with the equal mass case, where the correct definitions are




Figure 2. The natural dynamical reduction into orientation and geometry. Right - the three bodies define a triangle. Its dynamics is decomposed into its orientation and its geometry (shape + size). The dynamical variables for the orientation are taken to be the components of the total angular momentum within the rotating system.

Left: Mechanics in geometry space. Geometry space is a 3-dimensional space in which each point describes the shape and size of a triangle formed by the three bodies. r, θ, ϕ are considered to be spherical coordinates in this space. The (yellow) surface shaped like a pipe joint is a surface of constant effective potential (shown for $\vec{J} = 0$), so that motion is confined to be within it. The 3 solid (black) lines are sources of an attractive potential which originates with the Newtonian gravitational potential. Finally the (blue) arrows show the radial magnetic-field field, which originates from a Coriolis force.

easier to guess, before generalizing to the general case of unequal masses. [12] arrives at these coordinates after a choice of coordinates on $\mathbb{C}\mathbb{P}^1$ and comments that “Remarkably, it turns out that if we put the binary collisions at the third roots of unity... then the equilateral points are automatically moved to the north and south poles.” The definition of \vec{w} provides the missing link and makes natural the transition to the geometry space coordinates and their properties.

3 Applications to planar motion

This section describes an application of current formulation to planar three-body motion, and in particular to uniformly rotating configurations.

If the initial velocities are within the three-body plane then the motion remains planar throughout. For planar motion, J_1, J_2 vanish identically, while J_3 is conserved. Hence, the rotating body motion is trivial, and \mathcal{L}_2 specializes to

$$\mathcal{L}_{2,2d} = -\frac{1}{2I} J_3^2 \quad (3.1)$$

where $\mathcal{L}_0, \mathcal{L}_1$ are defined in (2.50,2.53).

Equilibria. Equilibria of the reduced planar motion describe rigidly rotating solutions.

The effective potential in geometry space $V_{eff}^G = V_{eff}^G(r, \theta, \phi)$ is given by

$$V_{eff}^G = V + \frac{J_3^2}{2I} \quad (3.2)$$

where the second term is the centrifugal potential.

If V is a power-law in the inter-body distance, one can solve the equilibrium potential with respect to r . In the remainder of this section, we assume the Newtonian gravitational potential. Solving the equation $0 = \partial V_{eff}^G / \partial r$ for r and substituting back into V_{eff}^G one gets a potential that may be called the effective potential in shape space $V_{eff}^S = V_{eff}^S(\theta, \phi)$. It is given by

$$V_{eff}^S = -\frac{1}{2J_3^2} I V^2 \quad (3.3)$$

Note that $I V^2$ is indeed r -independent, on account of the r -scaling properties of I and V .

To proceed towards the equilibria, it remains to differentiate V_{eff}^S with respect to θ, ϕ . It turns out to be convenient to use the side-lengths, r_{12}, r_{23}, r_{31} , rather than θ, ϕ . We require

$$\begin{aligned} 0 &= \frac{\partial}{\partial r_{23}} \log V_{eff}^S = \frac{\partial}{\partial r_{23}} (\log I + 2 \log(-V)) = \\ &= \frac{2 m_2 m_3 r_{23}}{M I} - \frac{2 G m_2 m_3}{(-V) r_{23}^2} \end{aligned} \quad (3.4)$$

where we have used the expressions for V, I from (2.2, 2.51). Solving for r_{23} the factors of $m_2 m_3$ cancel out on account of the equality of gravitational and inertial masses, and one finds that the side lengths are all equal

$$r_{12}^3 = r_{23}^3 = r_{31}^3 = G M \frac{I}{(-V)} \quad (3.5)$$

These are the *equilateral solutions found by Lagrange* [6]. Thus, we presented a *derivation of them through the reduced Lagrangian in geometry space*.

Note that the angular velocity associated with an equilateral with sides a is given by

$$\Omega^2 \equiv \left(\frac{J_3}{I} \right)^2 = \frac{GM}{a^3} \quad (3.6)$$

which is consistent with Kepler's third law and the fact that within the equilateral orbit each body moves as though attracted toward the center of mass (for this reason, these configurations are known as central configurations).

We add some comments.

Collinear solutions. V_{eff}^S is invariant under a space reflection, which when translated to geometry space becomes a reflection through the equator. Hence, the gradient of V_{eff}^S must lie within the equator. By restricting V_{eff}^S to the equator one finds the three collinear solutions found by Euler [29]. These solutions were not found above while differentiating

with respect to the r_{ab} variables, since the transformation between them and the r, θ, ϕ variables is singular at the equator.

Type of equilibrium. By noting the behavior of V_{eff}^S near the equator, one concludes that the extremum associated with equilateral motion is in fact a maximum.

Stability of Lagrange’s equilateral solutions. The stability of the equilibria found in the previous subsection are closely related to the stability of the associated uniformly rotating solutions. In this subsection, we focus on the stability of Lagrange’s equilateral solutions with uniform rotation, and we sketch the determination of the associated domain of linear stability.

Given m_1, m_2, m_3 and J_3 this solution is represented by the equilibrium point at the Right or Left pole on the shape sphere. Without loss of generality, let us consider the Right pole. Radial displacements can be shown to result in a stable normal mode, which is analogous to quasi-circular motion in the two-body problem.

For displacement in the plane tangent to the shape sphere, one has to take into account also the magnetic-like field (2.55). Qualitatively, one finds a negative (unstable) effective spring constant, yet the magnetic-like field can stabilize it. More specifically, the equation for the eigenfrequency ω is

$$0 = x^2 - x + \frac{27}{4} \frac{M_2}{M^2} \quad (3.7)$$

where $x := \omega/\Omega$ is the dimensionless frequency of oscillation, and Ω is the frequency of the equilateral solution (3.6).

This equation has real solutions, implying linear stability, whenever

$$\frac{M_2}{M^2} \leq \frac{1}{27} \quad (3.8)$$

This stability bound was found by Gascheau (1843) [25], see also [26] since I was not able to locate the former reference.

The current derivation is novel, reducing the dynamical stability analysis to a static one in geometry space.

Outlook. A future version will include the necessary details for the preceding discussion of stability.

In addition to the application of the formulation to the special exact solutions, we anticipate applications to the statistical solution. In a work in preparation, [5], the formulation would be shown to be essential to the determination of the system’s regularized phase-volume. A future version will also apply it to a visualization of the system through allowed regions of motion (analogous to the Hill region).

4 Four-body problem

In the three-body problem, we introduced symmetric vector coordinates in the center of mass frame, inspired by Lagrange’s solution to the cubic (2.9). The quartic equation also has a general solution, which suggests a generalization to the four-body problem.

Denoting the masses by m_a , $a = 1, 2, 3, 4$ and the positions by \vec{r}_a we define

$$\begin{aligned}\vec{s}_1 &= \vec{r}_1 - \vec{r}_2 - \vec{r}_3 + \vec{r}_4 \\ \vec{s}_2 &= -\vec{r}_1 + \vec{r}_2 - \vec{r}_3 + \vec{r}_4 \\ \vec{s}_3 &= -\vec{r}_1 - \vec{r}_2 + \vec{r}_3 + \vec{r}_4\end{aligned}\tag{4.1}$$

These variables are translation-invariant vectors. They contain 9 degrees of freedom, which are necessary to cover the configuration space at the center of mass frame. As the labels 1, 2, 3, 4 are permuted, the \vec{s} vectors transform nicely – they permute among themselves and/or change signs.

In order to proceed and decompose the variables into orientation and geometrical variables, the invariants are the 6 scalar products $Q_{rt} = \vec{s}_r \cdot \vec{s}_t$, $r, t = 1, 2, 3$ and the gauge-field and associated field-strength become $SO(3)$ -valued, and so, non-Abelian.

Acknowledgments

I thank Yogesh Dandekar, Lior Lederer and Subhajit Mazumdar for collaboration on a related project. This research, as well as [2], was performed during the times of the COVID-19 pandemic, and benefitted from the associated isolation.

I dedicate this work to the memory of Ami Nathan, 15.6.1934–24.10.2020, my father in law and a dedicated family man.

A Lagrange’s solution to the cubic

In this appendix, we provide a bit more information on Lagrange’s solution to the cubic.

A cubic is an equation of the form

$$a x^3 + b x^2 + c x + d = 0 .\tag{A.1}$$

Denoting the roots by x_1, x_2, x_3 Lagrange defines [27]

$$\begin{aligned}s_1 &:= x_1 + \eta x_2 + \bar{\eta} x_3 \\ s_2 &:= x_1 + \bar{\eta} x_2 + \eta x_3\end{aligned}\tag{A.2}$$

where η is the cubic root of unity.

Under cyclic permutations of x_a , $a = 1, 2, 3$, the variables s_1, s_2 transform by a third of a revolution in phase. Hence, s_1^3 and s_2^3 are invariant under cyclic permutations. Moreover, any exchange permutation of x_a , $a = 1, 2, 3$, exchanges s_1^3 and s_2^3 . Hence $S := s_1^3 + s_2^3$ is a fully symmetric function of x_a , $a = 1, 2, 3$ and so is $P := s_1 s_2$. Therefore, by the fundamental theorem of symmetric polynomials, both S, P can be expressed in terms of $b/a, c/a, d/a$. Now $z = s_1^3, s_2^3$ are the roots of the quadratic

$$z^2 - S z + P^3 = 0 .\tag{A.3}$$

After solving this quadratic, one takes cubic roots to obtain s_1, s_2 . Finally, (A.2) together with $-b/a = s_0 := x_1 + x_2 + x_3$ is a system of linear equations which can be inverted

to produce the three roots $x_a, a = 1, 2, 3$. These transformations between the x and s variables are known as a discrete Fourier transform.

In this way, Lagrange's method relies on the symmetry properties of expressions in the roots in order to solve the cubic. This is a crucial insight on the road to Galois theory.

The s_1, s_2 variables are invariant under a shift in x and have nice transformation properties under permutations. In this way, they inspired the definition of \vec{z} (2.9).

B Bi-complex numbers and the isotropic oscillator

A bi-complex number w is an expression of form

$$w = a + bi + cj + dij \tag{B.1}$$

where a, b, c, d are real numbers.

Bi-complex numbers can be added and multiplied using the relations

$$\begin{aligned} -1 &= i^2 = j^2 \\ ij &= ji \end{aligned} \tag{B.2}$$

These relations define an algebraic structure known as a commutative ring. Note that bi-complex numbers are distinct from Quaternions: while both are dimension 4 over the Reals, the latter is non-commutative, e.g. $ij = -ji = k$.

Not all non-zero bi-complex numbers have an inverse. This is true of zero-divisors, namely non-zero elements u, v such that $u \cdot v = 0$. The zero-divisors can be generated from the elementary zero divisors

$$R := 1 + ij \quad L := 1 - ij \tag{B.3}$$

The notation R, L stands for right and left: for the isotropic oscillator these states correspond to right/left circular motion, while for the three-body problem they correspond to right/left equilateral triangles.

Indeed, $R \cdot L = 0$. Moreover, all other zero divisors can be generated from these through multiplication by a bi-complex number. R, L satisfy the relations

$$\begin{aligned} iR &= -jR \\ iL &= +jL \end{aligned} \tag{B.4}$$

We note that after multiplication R, L obtain the following simple form: $iR = i - j$, $iL = i + j$.

The isotropic oscillator. The isotropic oscillator is the mechanical system consisting of a point mass m , free to move in the plane under the influence of a harmonic restoring force $\vec{F} = -m\omega^2 \vec{r}$. It can be defined through the Hamiltonian

$$H = \frac{1}{2m} \vec{p}^2 + \frac{1}{2} m \omega^2 \vec{r}^2 . \tag{B.5}$$

One complexifies phase space through the definition of a complex vector

$$\vec{a} := \frac{1}{\sqrt{2}} \left(\sqrt{m\omega} \vec{r} + \frac{i}{\sqrt{m\omega}} \vec{p} \right) \quad (\text{B.6})$$

In terms of these variables, one has

$$H = \omega \vec{a} \cdot \vec{a}^* \quad (\text{B.7})$$

where $*$ denotes complex conjugation, and the Poisson brackets

$$\{a_\alpha, a_\beta^*\} = -i \delta_{\alpha\beta} \quad (\text{B.8})$$

— all other Poisson brackets among the complex variables vanish.

It is natural to further complexify the plane. Since plane rotations commute with phase space rotations we use a different imaginary unit j which commutes with i and define a bi-complex number through

$$\vec{a} \rightarrow a := \vec{a} \cdot (\hat{x} + j \hat{y}) \quad (\text{B.9})$$

Since time evolution is given by

$$\vec{a}(t) = \exp(i\omega t) \vec{a}(0) \quad (\text{B.10})$$

the conserved charges can be expressed in terms of the Stokes parameters S_0, S_1, S_2, S_3 defined by

$$\begin{aligned} S_0 + i j S_3 &:= a \cdot \bar{a}^* \\ S_1 + j S_2 &= a \cdot a^* , \end{aligned} \quad (\text{B.11})$$

where a bar denotes conjugation with respect to j , and a $*$ continues to denote conjugation with respect to i . These expressions are equivalent to $S_0 = \vec{a} \cdot \vec{a}^*$, $S_i = \sigma_{\alpha\beta}^{i-1} a_\alpha^* a_\beta$, $i = 1, 2, 3$, $\alpha, \beta = 1, 2$ where $\sigma^i, i = 1, 2, 3$ are the Pauli matrices. If (a_1, a_2) undergo a unitary transformation, $\vec{S} \equiv (S_1, S_2, S_3)$ undergoes an orthogonal transformation.

The Stokes parameters satisfy the relation

$$S_0^2 = \vec{S}^2 . \quad (\text{B.12})$$

In addition, their Poisson brackets are given by

$$\begin{aligned} \{S_i, S_j\} &= \epsilon_{ijk} S_k \\ \{S_i, S_0\} &= 0 . \end{aligned} \quad (\text{B.13})$$

C Equations of motion in terms of non-coordinate velocities and the rotating rigid body

Consider an n -dimensional configuration space \mathcal{M} , described by some generalized coordinates $(q^i)_{i=1}^n$, as well as non-coordinate velocities (also known as quasi velocities [28])

$$v^a = \beta_i^a \dot{q}^i, \quad a = 1, \dots, n \quad (\text{C.1})$$

where $\beta_i^a = \beta_i^a(q^j)$ defines a changes of basis from the coordinate velocities \dot{q}^i to v^a . As an example, we shall see below that the angular velocity components of a rotating rigid body, are non-coordinate velocities.

Equivalently, the velocities define a basis of 1-forms over \mathcal{M}

$$\beta^a := \beta_i^a dq^i \quad (\text{C.2})$$

A basis of 1-forms is called a coordinate basis, or a holonomic basis, if the 1-forms can all be expressed as differentials of some functions. The non-coordinate nature of a basis is captured by the torsion coefficients $C_{ab}^c = C_{ab}^c(q^i)$ defined by

$$d\beta^c = \frac{1}{2} C_{ab}^c \beta^b \beta^a. \quad (\text{C.3})$$

The dual basis of vector fields e_a is defined by $\beta^a \cdot e_b = \delta_b^a$. In components $e_b = e_b^i \partial_i$ and e_b^i is the inverse matrix for β_i^a . In terms of e_a , the structure constants can be defined equivalently through the Lie brackets $[e_a, e_b] = C_{ab}^c e_c$.

Suppose the Lagrangian is given in terms of the generalized coordinates and the velocities v^a

$$\mathcal{L} = \mathcal{L}(q^i, v^a). \quad (\text{C.4})$$

The generalization of the Euler-Lagrange equations of motion to a non-coordinate basis are

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial v^a} \right) + C_{ab}^c v^b \frac{\partial \mathcal{L}}{\partial v^c} = \partial_a \mathcal{L} \quad (\text{C.5})$$

where $\partial_a \equiv e_a^i \partial_i$ is a derivative with respect to the vector field e_a . This generalization was found by Poincaré (1901) [19, 28].

For coordinate velocities $C_{ab}^c = 0$ and hence the term $C_{ab}^c v^b \frac{\partial \mathcal{L}}{\partial v^c}$ vanishes and the equations reduce to the Euler-Lagrange equations.

Derivation. The derivation of the Euler-Lagrange equations relies on the commutation of the variation and the time derivative. For non-coordinate velocities, this no longer holds, and a correction term arises. Indeed

$$\begin{aligned} \delta v^a &\equiv \delta \left(\beta_i^a \frac{d}{dt} q^i \right) = \frac{d}{dt} (\beta_i^a \delta q^i) + 2 (d\beta^a)_{ij} \delta q^i \dot{q}^j = \\ &= \frac{d}{dt} (\beta_i^a \delta q^i) + C_{bc}^a \delta q^c \dot{q}^b \end{aligned} \quad (\text{C.6})$$

where the last equality uses (C.3).

Now the variation of the action can be performed

$$\begin{aligned}
\delta S &= \int \left[\frac{\partial \mathcal{L}}{\partial q^i} \delta q^i + \frac{\partial \mathcal{L}}{\partial v^a} \delta v^a \right] = \\
&= \int \left[\frac{\partial \mathcal{L}}{\partial q^i} \delta q^i + \frac{\partial \mathcal{L}}{\partial v^a} \frac{d}{dt} (\beta_i^a \delta q^i) + \frac{\partial \mathcal{L}}{\partial v^c} C^c_{ba} \delta q^a \dot{q}^b \right] = \\
&= \int \delta q^a \left[\partial_a \mathcal{L} - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial v^a} \right) - C^c_{ab} v^b \frac{\partial \mathcal{L}}{\partial v^c} \right]
\end{aligned} \tag{C.7}$$

where in passing to the second line, we have used (C.6), and in passing to the third, we have used $q^i = e_a^i q^a$ and $C^c_{ba} = -C^c_{ab}$. The last line implies the modified equations of motion (C.5) — QED.

Momentum space. This procedure can be generalized so that it applies to Hamiltonians as well (though I have not seen this in the literature). Define the momenta conjugate to v^a

$$p_a = \frac{\partial \mathcal{L}}{\partial v^a} \tag{C.8}$$

These momenta can be expressed in terms of momenta conjugate to \dot{q}^i through $p_a = e_a^i p_i$, where e_a^i were defined below (C.3). This implies non-zero Poisson brackets for p_a

$$\{p_a, p_b\} = -C^a_{bc} p_c \tag{C.9}$$

From here one proceeds as usual to define the Hamiltonian through the Legendre transform, and to obtain the equations of motion for any dynamical variable $A = A(q^i, p_a)$, through $dA/dt = \{A, \mathcal{H}\}$.

Lagrangian for Euler's rigid body equations. Let us see how this formulation allows us to formulate Euler's equations for a rotating rigid body within a Lagrangian formulation.

A freely rotating rigid body is described by Euler's equations

$$0 = \frac{d}{dt} \Big|_{inert} \vec{S} \equiv \dot{\vec{S}} + \vec{\omega} \times \vec{S} . \tag{C.10}$$

Equivalently, in coordinate notation

$$0 = \frac{d}{dt} \Big|_{inert} I_1 \omega_1 := I_1 \dot{\omega}_1 + (I_3 - I_2) \omega_2 \omega_3 \tag{C.11}$$

and similarly 2 other equation by cyclic permutations. $\vec{\omega}$ is the angular velocity vector in a body frame whose axes are the body's principal axes, I_i , $i = 1, 2, 3$ are the principal moments of inertia, and $\frac{d}{dt} \Big|_{inert}$ denotes a time derivative in the inertial frame.

The kinetic energy of the body is given by $T = \frac{1}{2} I_{ij} \omega^i \omega^j$. A free body has no potential energy and hence the Lagrangian is defined to be

$$\mathcal{L} := T = \frac{1}{2} I_{ij} \omega^i \omega^j . \tag{C.12}$$

If one applies the standard Euler-Lagrange equations carelessly, one finds $(d/dt) (\partial\mathcal{L}/\partial\omega^i) = 0$. This is different from the Euler equations of motion (C.11). However, we notice that ω^i are non-coordinate velocities with torsion coefficients

$$C_{jk}^i = \epsilon_{ijk} \quad (\text{C.13})$$

where ϵ_{ijk} is the Levi-Civita tensor. This can be shown by expressing ω^i in terms of Euler angles,¹ and we note that the coefficients are constant over \mathcal{M} and coincide with the structure constants of $SO(3)$. The modified equations of motion (C.5) read

$$0 = \dot{S}^i + \epsilon_{ij}^k \omega^j S_k = \dot{S}^i + (\vec{\omega} \times \vec{S})^i \quad (\text{C.15})$$

which now fully agree with Euler's equations.

We can now proceed to a Hamiltonian formulation. The conjugate momenta

$$S_i = \frac{\partial\mathcal{L}}{\partial\omega^i} = I_{ij} \omega^j \quad (\text{C.16})$$

are the coordinates of the spin (angular momentum) vector in body coordinates. By (C.9, C.13) their Poisson brackets are given by

$$\{S_i, S_j\} = -\epsilon_{ijk} S_k \quad (\text{C.17})$$

The Hamiltonian is given by

$$\mathcal{H} = \omega^i S_i - \mathcal{L} = \frac{1}{2} (I^{-1})^{ij} S_i S_j \quad (\text{C.18})$$

where I^{-1} is in the inverse matrix for the moment of inertia tensor. Finally, the equations of motion are given by

$$\frac{d}{dt} S^i = \{S^i, \mathcal{H}\} = \{S^i, S^j\} \frac{\partial\mathcal{H}}{\partial S^j} = -\epsilon_{ijk} \omega^j S^k \quad (\text{C.19})$$

which reproduce the Euler equations.

¹E.g.

$$\begin{aligned} \omega^x &= -\sin\theta_n \cos\psi_n \dot{\phi}_n + \sin\psi_n \dot{\theta}_n \\ \omega^y &= \sin\theta_n \sin\psi_n \dot{\phi}_n + \cos\psi_n \dot{\theta}_n \\ \omega^z &= \cos\theta_n \dot{\phi}_n + \dot{\psi}_n \end{aligned} \quad (\text{C.14})$$

where θ_n, ϕ_n, ψ_n are Euler-like angles. These expressions are of the form (C.1).

References

- [1] I. Newton, “Philosophiæ Naturalis Principia Mathematica” (1687).
- [2] B. Kol, “Flux-based statistical prediction of three-body outcomes,” *Celestial Mech. Dyn. Astron.*, Volume **133**, 17 (2021) doi:10.1007/s10569-021-10015-x [arXiv:2002.11496 [gr-qc]].
- [3] N. C. Stone and N. W. C. Leigh, “A statistical solution to the chaotic, non-hierarchical three-body problem,” *Nature* **576**, no. 7787, 406 (2019). doi:10.1038/s41586-019-1833-8
- [4] Y. B. Ginat and H. B. Perets, “An Analytical, Statistical Approximate Solution for Dissipative and non-Dissipative Binary-Single Stellar Encounters,” arXiv:2011.00010 [astro-ph.GA].
- [5] “Regularized phase-volume for the three-body problem”, to appear.
- [6] J. L. Lagrange, “Essai sur le Problème des Trois Corps,” Prix de l’Académie Royale des Sciences de Paris, tome IX (1772), in vol. 6 of *Oeuvres*.
- [7] C. G. J. Jacobi, “Sur l’élimination des noeuds dans le problème des trois corps,” *Astron. Nachr.* **462**, 81 (1842).
- [8] T. Levi-Civita, “Sulla riduzione del problem dei tre corpi,” *Atti Ist. Veneto di Sc., lett. ed art* 907 **LXXIV** (1914-15).
- [9] F. D. Murnaghan, “A symmetric reduction of the planar three-body problem,” *Am. J. Math.* **58**, 829 (1936).
- [10] E. R. van Kampen and A. Wintner, “On a symmetrical canonical reduction of the problem of three bodies,” *Am. J. Math.* **59** 153 (1937).
- [11] G. Lemaître, “Coordonnées symétriques dans le problème des trois corps,” *Ext. Bull. Acad. roy. Belg.* **120**, 582 (1952).
- [12] R. Moeckel and R. Montgomery, “Symmetric regularization, reduction and blow-up of the planar three-body problem,” *Pac. J. Math.* **262**, 129 (2013).
- [13] R. Montgomery, “The three-body problem and the shape sphere,” *Am. Math. Month.* **122**, 299 (2015).
- [14] R. Montgomery, “The three-body problem,” *Sci. Am.* **321**, 2, 66 (August 2019). doi:10.1038/scientificamerican0819-66. See also scientificamerican.com.
- [15] E. A. Hylleraas, “Neue Berechnung der Energie des Heliums im Grundzustande, sowie des tiefsten Terms von Ortho-Helium,” *Z. Physik* **54**, 347 (1929).
- [16] T. H. Gronwall, “A Special Conformally Euclidean Space of Three Dimensions Occurring in Wave Mechanics,” *Ann. Math.* **33**, 279 (1932).
- [17] V. Fock, “On the Schrodinger equation of the Helium atom,” *Izv. AN* **18**, 161 (1954). *Det Kongelige Norske Videnskabers Selskabs Forhandling* **B31**, 138 (1958) - Author’s English version.
- [18] V. K. Tkachenko, “Schrödinger equation for the Helium atom,” *Teor. Mat. Fiz.* **34**, 142 (1978).
- [19] H. Poincare, “Sur une form nouvelle des équations de la mécanique,” *CR Acad. Sci.* **132**, 369 (1901).
- [20] G. G. Stokes, “On the composition and resolution of streams of polarized light from different

sources,” *Trans. Cambridge Phil. Soc.* **9**, 399 (1852).

Reprinted in *Mathematical and Physical Papers* [by George Gabriel Stokes], Vol. III pp. 233-258, Cambridge University Press. doi:10.1017/CBO9780511702266.010

- [21] H. Poincaré, “Théorie mathématique de la lumière II,” G. Carré éditeur, Paris (1892).
- [22] W. Pauli, “Zur Quantenmechanik des magnetischen Elektrons,” *Z. Physik* **43**, 601 (1927). doi:10.1007/BF01397326
- [23] H. Hopf, “Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche,” *Math. Ann.* **104**, 637 (1931). doi:10.1007/BF01457962
- [24] F. Bloch, “Nuclear induction,” *Phys. Rev.* **70**, 460 (1946). doi:10.1103/physrev.70.460
- [25] G. Gascheau, “Examen d’une classe d’équations différentiel et applications à un cas particulier du problème des trois corps,” *C. R. Acad. Sci.* **16** 393 (1843).
- [26] E. J. Routh, “On Laplace’s three particles, with a supplement on the stability of steady motion,” *Proc. London Math. Soc.* **6**, 86 (1875) (repr. in *Stability of Motion*, Fuller, A.T. (ed.) [London: Taylor & Francis], Chap. 3 [1975]).
- [27] J. L. Lagrange, “Réflexions sur la résolution algébrique des équations,” *Nouveaux Mémoires de l’Académie royale des Science et Belles-Lettres de Berlin* (1771), in vol. 3 of *Oeuvres* (p. 205).
- [28] V.I. Arnold, V.V. Kozlov and A.I. Neishtadt, *Mathematical Aspects of Classical and Celestial Mechanics in: Encyclopaedia of Mathematical Science, Vol. III, Dynamical Systems* (Springer-Verlag, Berlin, 1988).
- [29] L. Euler, “De motu rectilineo trium corporum se mutuo attrahentium,” *Novi commentarii acadmiæ scientarum Petropolitanæ* 11, pp. 144-151 (1767), in *Oeuvres, Seria Secunda tome XXV Commentationes Astronomicæ* (p. 286).