

HYPERSURFACES OF PRESCRIBED NULL EXPANSION

XIAOXIANG CHAI

ABSTRACT. We study hypersurfaces with prescribed null expansion in an initial data set. We propose a notion of stability and prove a topology theorem. Eichmair's Perron approach toward the existence of marginally outer trapped surface adapts to the settings of hypersurfaces with prescribed null expansion with only minor modifications.

Gromov [Gro18, Gro21] studied stable μ -bubbles in the study of scalar curvature which leads to many interesting results in positive scalar curvature. See [Gro21]. The use of stable μ -bubble dates back to minimal surface techniques used by Schoen-Yau settled the positive mass conjecture [SY79b] and [SY79a]. The analog of minimal surfaces in initial data sets is a *marginally outer trapped surface* (in short, *MOTS*; See [AMS05]). An initial data set (M, g, p) is a Riemannian manifold (M, g) equipped with an extra symmetric 2-form p . A MOTS arises boundaries of blow up sets of Jang equation [SY81]. It was applied to establish the spacetime positive mass theorem [EHLS16].

Definition 1. *Given a function h on (M, g) , a hypersurface Σ is called a hypersurface of prescribed null expansion if*

$$(0.1) \quad \theta := H_\Sigma + \text{tr}_\Sigma p = h$$

where h is a function on (M, g) and θ is the null expansion.

The case $h \equiv 0$ gives the definition of MOTS. Motivated by the stability of MOTS [AMS05] and the stability of the μ -bubble, we define the following.

Definition 2. *We say that Σ is stable if there exists a vector field $X = \varphi\nu$ with nonzero $\varphi \geq 0$ such that*

$$(0.2) \quad \delta_X(H + \text{tr}_\Sigma p - h) \geq 0.$$

It is desirable to find a physical interpretation of such hypersurfaces. By [AEM10, (10)],

$$\delta_X \theta = -\Delta \varphi + 2\langle W, \nabla \varphi \rangle + (\text{div } W - W + Q)\varphi - \frac{1}{2}\theta(\theta - 2\text{tr } p)$$

where W is the vector field tangential to Σ which is dual to $p(\nu, \cdot)$ and

$$Q = \frac{1}{2}R_\Sigma - \mu - J(\nu) - \frac{1}{2}|p_\Sigma + A|^2,$$

and $\mu = \frac{1}{2}R_g - |p|_g^2 + (\text{tr}_g p)^2$, $J = \text{div}_g p - d(\text{tr}_g p)$, A is the second fundamental form of Σ in M and we write in short $\chi = p_\Sigma + A$ the null second fundamental form. With $X(h) = \varphi\langle Dh, \nu \rangle$ and $\theta = h$, we have that

$$\begin{aligned} & L\varphi \\ &= -\Delta \varphi + 2\langle W, \nabla \varphi \rangle + (\text{div } W - W + Q)\varphi \\ &\quad - \frac{1}{2}(h^2 - 2h \text{tr } p + 2\nu(h))\varphi. \end{aligned}$$

Equivalently, Σ is stable if the first eigenvalue to the eigen problem $L\varphi = \lambda\varphi$ is real and nonnegative by Krein-Rutman theorem (See [AMS05]). Note that $h \equiv 0$ recovers the stability operator of a MOTS and $p = 0$ recovers the stability operator of a μ -bubble.

Remark 0.1. By requiring that $\partial M \neq \emptyset$ and Σ does not change intersecting angle along the deformation induced by X , one can define a capillary version of (0.2) (See [ALY20]). More generally, one can prescribe arbitrary angle on ∂M as well, the boundary stability operator is no different from that of the case $p \equiv 0$. See [Gro21] for a derivation of boundary operator for the case $p \equiv 0$.

Assuming for some h defined on M ,

$$(0.3) \quad \mu - |J| + \frac{1}{2}\left(\frac{n}{n-1}h^2 - 2h \operatorname{tr} p - 2|Dh|\right) \geq 0.$$

The condition (0.3) when $h \equiv 0$ is called *dominant energy condition*. When $p \equiv 0$, it reduces to the relaxed positivity of scalar curvature [Gro21]. We have the following theorem asserting the Yamabe type of a stable hypersurface of null expansion $\theta = h$.

Theorem 0.2. *Assume that Σ is a stable hypersurface of prescribed null expansion h in an initial data set (M, g, p) satisfying (0.3), then Σ is of positive Yamabe type unless Σ is Ricci flat, χ^0 vanishes and $\mu - |J| + \frac{1}{2}\left(\frac{n}{n-1}h^2 - 2h \operatorname{tr} p + 2\nu(h)\right)$ vanishes along Σ .*

Proof of Theorem 0.2. Consider the eigenvalue problem

$$-\Delta\varphi + [Q - \frac{1}{2}(h^2 - 2h \operatorname{tr} p + 2\nu(h))]\varphi = \lambda\varphi.$$

Since Σ is stable, it follows the same lines as in [GS06] that the first eigenvalue $\lambda_1 \geq 0$ and the associated eigenfunction f can be chosen positive everywhere on Σ . Consider the metric $\tilde{h} = f^{\frac{2}{n-2}}h$, we have that the scalar curvature \tilde{S} of Σ with respect to \tilde{h} is

$$(0.4) \quad \begin{aligned} & \tilde{S} \\ &= f^{-\frac{n}{n-2}}(-2\Delta f + Sf + \frac{n-1}{n-2}f^{-1}|\nabla f|^2) \\ &= f^{-\frac{2}{n-2}}[2\lambda_1 + 2(\mu + J(\nu)) \\ & \quad + (h^2 - 2h \operatorname{tr} p + 2\langle Dh, \nu \rangle) + |\chi|^2 + \frac{n-1}{n-2}f^{-1}|\nabla f|^2] \\ &= f^{-\frac{2}{n-2}}[2\lambda_1 + 2(\mu + J(\nu)) \\ & \quad + (\frac{n}{n-1}h^2 - 2h \operatorname{tr} p + 2\langle Dh, \nu \rangle) + |\chi^0|^2 + \frac{n-1}{n-2}f^{-1}|\nabla f|^2], \end{aligned}$$

where $\chi^0 = \chi - \frac{1}{n-1}\theta g_\Sigma$ denotes the trace free part of the null second fundamental form.

By the condition (0.3), we have that $\tilde{S} \geq 0$. If $\tilde{S} > 0$ somewhere, then it is well known that Σ carries a metric of strict positive scalar curvature. If \tilde{S} vanishes identically, then $\lambda_1 = 0$, χ^0 vanishes along Σ , $\theta = h$ along Σ ,

$$2(\mu + J(\nu)) + (\frac{n}{n-1}h^2 - 2h \operatorname{tr} p + 2\langle Dh, \nu \rangle)$$

vanishes along Σ and f is a positive constant. We see from (0.4) that (Σ, h) is of vanishing scalar curvature. By a result of Bourguignon (see [KW75]), Σ carries a metric of positive scalar curvature unless Σ is Ricci flat. \square

Eichmair [Eic09] and Andersson-Metzger [AM09] established the existence of a marginally outer trapped surface in domains bounded by two boundaries with certain convexity conditions. In addition, Eichmair obtained the solution to the Plateau problem via Perron methods. We note that the Eichmair’s approach works for surfaces of prescribed null expansion with only minor modifications. In particular analogs of Theorem 1.1 and Theorem 3.1 of [Eic09] could be established via a study of the Jang equation,

$$\operatorname{div}_g\left(\frac{Du}{\sqrt{1+|Du|^2}}\right) + \left(g^{ij} - \frac{D^i u D^j u}{1+|Du|^2}\right)p_{ij} - \tau u - h = 0,$$

here $\tau \geq 0$ is a constant. The following theorem is an analog of [Eic09, Theorem 3.1]. We leave the proof and the solution of the Plateau problem to interested readers.

Theorem 0.3. *Let (M^n, g, p) be an initial data set and let U be a connected bounded open subset with smooth embedded boundary ∂U . Assume that the boundary consists of two non-empty closed hypersurfaces $\partial_1 U$ and $\partial_2 U$ so that*

$$H_{\partial_1 U} + (\operatorname{tr}_{\partial_1 U} p - h) > 0 \text{ and } H_{\partial_2 U} - (\operatorname{tr}_{\partial_2 U} p - h) > 0,$$

where the mean curvature is computed with respect to the normal pointing out of U . Then there exists a closed boundary Σ^{n-1} homologous to $\partial_1 U$ which is C -almost minimizing in U for a constant $C = C(|p|_{C(\bar{U})}, |h|_{C(\bar{U})})$. (Hence, Σ has a singular set of Hausdorff dimension less than $n - 8$ which satisfies (0.1) distributionally. In particular, if $2 \leq n \leq 7$, Σ is non-empty, closed and embedded.) If $3 \leq n \leq 7$ and if the data satisfies (0.3), the Σ has non-negative Yamabe type.

Acknowledgment Research of Xiaoxiang Chai is supported by KIAS Grants under the research code MG074402.

REFERENCES

[AEM10] Lars Andersson, Michael Eichmair, and Jan Metzger. Jang’s equation and its applications to marginally trapped surfaces. *ArXiv:1006.4601 [gr-qc]*, 2010.

[ALY20] Aghil Alaei, Martin Lesourd, and Shing-Tung Yau. Stable Surfaces and Free Boundary Marginally Outer Trapped Surfaces. *ArXiv:2009.07933 [gr-qc, physics:math-ph]*, 2020.

[AM09] Lars Andersson and Jan Metzger. The area of horizons and the trapped region. *Comm. Math. Phys.*, 290(3):941–972, 2009.

[AMS05] Lars Andersson, Marc Mars, and Walter Simon. Local existence of dynamical and trapping horizons. *Phys. Rev. Lett.*, 95:111102, Sep 2005.

[EHLS16] Michael Eichmair, Lan-Hsuan Huang, Dan A. Lee, and Richard Schoen. The space-time positive mass theorem in dimensions less than eight. *J. Eur. Math. Soc. (JEMS)*, 18(1):83–121, 2016.

[Eic09] Michael Eichmair. The Plateau problem for marginally outer trapped surfaces. *Journal of Differential Geometry*, 83(3):551–584, 2009.

[Gro18] Misha Gromov. Metric inequalities with scalar curvature. *Geom. Funct. Anal.*, 28(3):645–726, 2018.

[Gro21] Misha Gromov. Four Lectures on Scalar Curvature. *ArXiv:1908.10612 [math]*, 2021.

[GS06] Gregory J. Galloway and Richard Schoen. A generalization of Hawking’s black hole topology theorem to higher dimensions. *Comm. Math. Phys.*, 266(2):571–576, 2006.

[KW75] Jerry Kazdan and F Warner. Prescribing curvatures. In *Proceedings of Symposia in Pure Mathematics*, volume 27, pages 309–319. 1975.

[SY79a] R. Schoen and Shing Tung Yau. Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature. *Ann. of Math. (2)*, 110(1):127–142, 1979.

[SY79b] Richard Schoen and Shing Tung Yau. On the proof of the positive mass conjecture in general relativity. *Comm. Math. Phys.*, 65(1):45–76, 1979.

- [SY81] Richard Schoen and Shing Tung Yau. Proof of the positive mass theorem. ii. *Communications in Mathematical Physics*, 79(2):231–260, 1981.

KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 02455, SOUTH KOREA
Email address: `xxchai@kias.re.kr`