

Lecture notes for pseudodifferential operators and microlocal analysis

Ma, Shiqi

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ
Email address: mashiqi01@gmail.com, shiqi.s.ma@jyu.fi

arXiv:2107.12839v1 [math.AP] 27 Jul 2021

ABSTRACT. This is an introductory course focusing on some basic notions in pseudodifferential operators (Ψ DOs) and microlocal analysis. We start these lecture notes with some notations and necessary preliminaries. Then the notion of symbols and Ψ DOs are introduced. In Chapter 3 we define the oscillatory integrals of different types. Chapter 4 is devoted to the stationary phase lemmas. One of the features of the lecture is that the stationary phase lemmas are proved for not only compactly supported functions but also for more general functions with certain order of smoothness and certain order of growth at infinity. We build the results on the stationary phase lemmas. Chapters 5, 6 and 7 cover the main results in Ψ DOs and the proofs are heavily built on the results in Chapter 4. Some aspects of the semi-classical analysis are similar to that of microlocal analysis. In Chapter 8 we finally introduce the notion of wavefront, and Chapter 9 focuses on the propagation of singularities of solutions of partial differential equations. Important results are circled in black boxes and some key steps are marked in red color. Exercises are provided at the end of each chapter.

Version: May, 2021.

Contents

Chapter 1. Preliminaries	5
1.1. Notations	5
1.2. Schwartz Space and tempered distributions	6
1.3. Fourier transforms	9
Exercise	11
Chapter 2. Pseudodifferential operators	12
2.1. Symbols	12
2.2. Pseudodifferential operators	15
2.3. Kernels	19
2.4. Pseudolocal property	20
Exercise	21
Chapter 3. Oscillatory integrals	22
3.1. Oscillatory integrals - Type I	22
3.2. Oscillatory integrals - Type II	27
Exercise	30
Chapter 4. Stationary phase lemmas	31
4.1. A simple case	31
4.2. Lemma Statements	33
4.3. Proofs of the results	35
Exercise	41
Chapter 5. Symbolic calculus of Ψ DOs	42
5.1. Composition of Ψ DOs	42
5.2. Reduction of variables	45
5.3. The Adjoint and transpose	47
Exercise	48
Chapter 6. Parametrix and Boundedness of Ψ DOs	49
6.1. Parametrix	49
6.2. The L^2 boundedness	52
6.3. Gårding's inequalities	55
Exercise	60
Chapter 7. Semi-classical Ψ DOs and its symbolic calculus	61
7.1. Semi-classical Ψ DOs	61
7.2. Composition of the standard quantizations	65
7.3. Composition of the Weyl quantizations	68
7.4. Applications in Carleman estimates	74
Exercise	75
Chapter 8. The wavefront set	76

8.1. Basic facts	76
8.2. Wavefront set of product of distributions	79
8.3. The wavefront sets of Fourier integral operators	85
8.4. Applications	90
Exercise	94
Chapter 9. Propagation of the singularities	95
9.1. Microlocal parametrix	95
9.2. Bicharacteristics	98
9.3. Propagation of singularities	101
9.4. Cauchy problems of hyperbolic PDEs	105
Exercise	107
Bibliography	108
Index	110

CHAPTER 1

Preliminaries

A good reference for is [26, Chapters 1-5].

1.1. Notations

\mathbb{R}^n : the Euclidean space. For $x \in \mathbb{R}^n$, $|x| := \sqrt{x_1^2 + \cdots + x_n^2}$, and the inner product $x \cdot y := \sum_{j=1}^n x_j y_j$. The notation $\langle x \rangle := (1 + |x|^2)^{1/2}$ will be frequently used throughout the lecture. For two quantities \mathcal{A} and \mathcal{B} , we write $\mathcal{A} \lesssim \mathcal{B}$ to signify $\mathcal{A} \leq C\mathcal{B}$, and write $\mathcal{A} \simeq \mathcal{B}$ to signify $C_1\mathcal{B} \leq \mathcal{A} \leq C_2\mathcal{B}$, for some generic positive constants C , C_1 and C_2 . It can be checked that $\langle x \rangle \simeq 1 + |x|$.

LEMMA 1.1. *For any $s \in \mathbb{R}$ and any multi-index α , there exists a constant C independent of x such that*

$$|\partial^\alpha (\langle x \rangle^s)| \leq C \langle x \rangle^{s-|\alpha|}, \quad \text{when } |x| \geq 1.$$

The proof is left as an exercise.

$C^m(\mathbb{R}^n; \mathbb{C})$ is the set of complex-valued functions that has continuous derivative up to order m . $C_c^\infty(\mathbb{R}^n)$ is comprised of C^∞ functions with compact support.

The Fourier and inverse Fourier transforms of f are denoted as $\mathcal{F}f$ (also \hat{f}) and $\mathcal{F}^{-1}f$ (also \check{f}):

$$\begin{aligned} \hat{f}(\xi) &= \mathcal{F}f(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} f(x) \, dx, \\ \check{f}(\xi) &= \mathcal{F}^{-1}f(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix \cdot \xi} f(x) \, dx. \end{aligned}$$

$\langle f, g \rangle := \int_{\mathbb{R}^n} f(x)g(x) \, dx$, $(f, g) := \int_{\mathbb{R}^n} f(x)\overline{g(x)} \, dx$, where $\overline{g(x)}$ is the complex conjugation of $g(x)$.

$\partial_j := \frac{\partial}{\partial x_j}$, $D_j := \frac{1}{i}\partial_j$, where i is the imaginary unit.

Multi-index: in \mathbb{R}^n , a multi-index is $\alpha = (\alpha_1, \dots, \alpha_n)$ where α_j are non-negative integers. $D^\alpha := D_1^{\alpha_1} \cdots D_n^{\alpha_n}$, $\partial^\alpha := \partial_1^{\alpha_1} \cdots \partial_n^{\alpha_n}$, and $x^\alpha := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, and the length of α is $|\alpha| := \alpha_1 + \cdots + \alpha_n$.

LEMMA 1.2. *Assume $x \in \mathbb{R}^n$ and α is a multi-index. Then*

$$|x^\alpha| \leq |x|^{|\alpha|}.$$

PROOF. We have

$$\begin{aligned} |x^\alpha| &= |x_1^{\alpha_1} \cdots x_n^{\alpha_n}| = |x_1^{\alpha_1}| \cdots |x_n^{\alpha_n}| = |x_1|^{\alpha_1} \cdots |x_n|^{\alpha_n} \\ &\leq |x|^{\alpha_1} \cdots |x|^{\alpha_n} = |x|^{\alpha_1 + \cdots + \alpha_n} = |x|^{|\alpha|}. \end{aligned}$$

□

More on multi-index:

- $\beta \leq \alpha$ means $\beta_j \leq \alpha_j$ for $j = 1, \dots, n$;

- the notion $\alpha - \beta$ is valid only when $\beta \leq \alpha$, and $\alpha - \beta := (\alpha_1 - \beta_1, \dots, \alpha_n - \beta_n)$;
- $\alpha! := \alpha_1! \cdots \alpha_n!$;
- when $\beta \leq \alpha$, $\binom{\alpha}{\beta} := \frac{\alpha!}{\beta!(\alpha-\beta)!} = \binom{\alpha_1}{\beta_1} \cdots \binom{\alpha_n}{\beta_n}$, where $\binom{\alpha_j}{\beta_j} = \frac{\alpha_j!}{\beta_j!(\alpha_j-\beta_j)!}$;

A typical form of a linear differential operator is $\sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha$. If we denote a polynomial $p(x, \xi) := \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$ where $\xi \in \mathbb{R}^n$, then

$$\sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha = p(x, D).$$

LEMMA 1.3. *Assume $f, g \in C^\infty(\mathbb{R}^n)$ and α is a multi-index. Then*

$$D^\alpha(fg) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (D^{\alpha-\beta} f)(D^\beta g).$$

The proof is left as an exercise.

1.2. Schwartz Space and tempered distributions

DEFINITION 1.4 (Schwartz Space). Let $\varphi \in C^\infty(\mathbb{R}^n)$. For multi-indices α and β , we define the semi-norm $|\cdot|_{\alpha, \beta}$ of φ as

$$|\varphi|_{\alpha, \beta} := \sup_{x \in \mathbb{R}^n} |x^\alpha D^\beta \varphi(x)| < +\infty. \quad (1.1)$$

We call φ a *Schwartz function* when $|\varphi|_{\alpha, \beta} < +\infty$ for any α and β . The set

$$\{\varphi \in C^\infty(\mathbb{R}^n); |\varphi|_{\alpha, \beta} < +\infty, \forall \alpha, \beta\}$$

together with the topology induced by the set of semi-norms $|\cdot|_{\alpha, \beta}$ is called the *Schwartz space*, denoted as $\boxed{\mathcal{S}(\mathbb{R}^n)}$.

The topology \mathcal{T} is induced by $\{|\cdot|_{\alpha, \beta}\}$ is defined as follows. Choose

$$N(\alpha, \beta; \epsilon) := \{\varphi \in \mathcal{S}(\mathbb{R}^n); |\varphi|_{\alpha, \beta} < \epsilon\}$$

to be open neighborhoods of point $0 \in \mathcal{S}(\mathbb{R}^n)$. Choose

$$\mathcal{N} := \{N(\alpha, \beta; \epsilon); \alpha, \beta \text{ are multi-index, } \epsilon > 0\}$$

to be a open neighborhood basis of 0, and $\varphi + \mathcal{N}$ the open neighborhood basis of $\varphi \in \mathcal{S}(\mathbb{R}^n)$. Then the topology \mathcal{T} is generated by these open neighborhood basis, see [16, §1.8] for more details.

DEFINITION 1.5 (Convergence in Schwartz space). A sequence of functions $\{\varphi_j\}_j \subset \mathcal{S}(\mathbb{R}^n)$ is said to *converge to zero in $\mathcal{S}(\mathbb{R}^n)$* if

$$\forall \alpha, \beta, |\varphi_j|_{\alpha, \beta} \rightarrow 0 \quad j \rightarrow +\infty, \quad (1.2)$$

denoted as $\varphi_j \rightarrow 0$ in $\mathcal{S}(\mathbb{R}^n)$.

LEMMA 1.6. *We have*

$$\mathcal{F}\mathcal{S}(\mathbb{R}^n) = \mathcal{S}(\mathbb{R}^n), \quad \partial^\alpha \mathcal{S}(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n).$$

The space $\mathcal{S}(\mathbb{R}^n)$ are often be used as test functions set. There is also another commonly used test functions set: $C_c^\infty(\mathbb{R}^n)$. In Fourier analysis the set $\mathcal{S}(\mathbb{R}^n)$ is more commonly used than $C_c^\infty(\mathbb{R}^n)$, and one of the reason is that $\mathcal{S}(\mathbb{R}^n)$ is closed for the Fourier transform \mathcal{F} . The uncertainty principle claims that the Fourier transform of any compactly supported function is impossible to be compactly supported, namely, $\mathcal{F}C_c^\infty(\mathbb{R}^n) \neq C_c^\infty(\mathbb{R}^n)$.

LEMMA 1.7. *$C_c^\infty(\mathbb{R}^n)$ is dense in $\mathcal{S}(\mathbb{R}^n)$.*

PROOF. Fix a function $\phi \in C_c^\infty(\mathbb{R}^n)$ satisfying $\phi \equiv 1$ when $|x| \leq 1$, and $\phi \equiv 0$ when $|x| \geq 2$, and $0 \leq \phi(x) \leq 1$.

For any $\varphi \in \mathcal{S}(\mathbb{R}^n)$, denote $\varphi_\epsilon(x) := \varphi(x)\phi(\epsilon x)$, then $\{\varphi_\epsilon\}_{\epsilon>0}$ is a sequence in $C_c^\infty(\mathbb{R}^n)$. Then for any multi-index α and β , we have

$$\begin{aligned} |\phi - \phi_\epsilon|_{\alpha,\beta} &= \sup_{x \in \mathbb{R}^n} |x^\alpha \partial^\beta (\varphi(x)[1 - \phi(\epsilon x)])| = \sup_{|x| \geq 1/\epsilon} |x^\alpha \partial^\beta (\varphi(x)[1 - \phi(\epsilon x)])| \\ &\leq \sup_{1/\epsilon \leq |x| \leq 2/\epsilon} |x^\alpha \partial^\beta (\varphi(x)[1 - \phi(\epsilon x)])| + \sup_{|x| \geq 2/\epsilon} |x^\alpha \partial^\beta \varphi(x)| \\ &= \sup_{1/\epsilon \leq |x| \leq 2/\epsilon} |x^\alpha \partial^\beta \varphi(x)| |1 - \phi(\epsilon x)| + \mathcal{O}(\epsilon) + \sup_{|x| \geq 2/\epsilon} |x^\alpha \partial^\beta \varphi(x)| \\ &\leq \sup_{1/\epsilon \leq |x| \leq 2/\epsilon} |x^\alpha \partial^\beta \varphi(x)| + \mathcal{O}(\epsilon) + \sup_{|x| \geq 2/\epsilon} |x^\alpha \partial^\beta \varphi(x)| \\ &\leq 2 \sup_{|x| \geq 1/\epsilon} |x^\alpha \partial^\beta \varphi(x)| + \mathcal{O}(\epsilon). \end{aligned}$$

Because $\sup_{\mathbb{R}^n} |x_j^2 x^\alpha \partial^\beta \varphi(x)| < +\infty$, we have that $|x_j^2 x^\alpha \partial^\beta \varphi(x)|$ is bounded in \mathbb{R}^n , so $|x|^2 |x^\alpha \partial^\beta \varphi(x)|$ is bounded in \mathbb{R}^n , thus $|x^\alpha \partial^\beta \varphi(x)| \leq C_{\alpha,\beta} \langle x \rangle^{-2}$ for certain constant $C_{\alpha,\beta}$. Therefore,

$$|\phi - \phi_\epsilon|_{\alpha,\beta} \leq 2C_{\alpha,\beta} \sup_{|x| \geq 1/\epsilon} \langle x \rangle^{-2} + \mathcal{O}(\epsilon) \rightarrow 0, \quad \epsilon \rightarrow 0.$$

The proof is complete. \square

LEMMA 1.8. *Let $f \in \mathcal{S}(\mathbb{R}^n)$. Then $\forall s \in \mathbb{R}$, we have*

$$(1 + |x|^2)^s f(x) \in \mathcal{S}(\mathbb{R}^n).$$

PROOF. Let α be a multi-index. Then

$$D^\alpha [(1 + |x|^2)^s f(x)] = \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} D^\delta ((1 + |x|^2)^s) \cdot (D^{\alpha-\delta} f)(x).$$

We should notice that $|D^\delta ((1 + |x|^2)^s)|$ can be always controlled by $(1 + |x|^2)^{t_\delta}$ for $t_\delta \in \mathbb{R}$ large enough:

$$|D^\delta ((1 + |x|^2)^s)| \leq (1 + |x|^2)^{t_\delta}, \quad \forall x \in \mathbb{R}^n.$$

So for any non-negative integer k and multi-index α , we have

$$\begin{aligned} |(1 + |x|^2)^k \cdot D^\alpha [(1 + |x|^2)^s f(x)]| &\leq \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} (1 + |x|^2)^{t_\delta} \cdot |(D^{\alpha-\delta} f)(x)| \\ &\leq \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} |(1 + |x|^2)^{t_\delta+k} (D^{\alpha-\delta} f)(x)| \\ &\leq \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} C_{k,\alpha,\delta} = C_{k,\alpha} < +\infty. \end{aligned}$$

We proved the conclusion. \square

Schwartz functions are these who decay fast enough. Now we introduce another type of functions which grow at infinity, but with a mild speed. These functions are called tempered functions.

DEFINITION 1.9 (Tempered functions). Let f be a measurable function defined on \mathbb{R}^n such that

$$\sup_{x \in \mathbb{R}^n} |(1 + |x|)^{-m} f(x)| < +\infty$$

for some positive integer m . Then we call f a *tempered function*. If f is continuous, then we call it *continuous tempered function*.

LEMMA 1.10. *Assume f is a smooth function such that $\partial^\alpha f$ are tempered functions $\forall \alpha$, then we have $f \cdot \mathcal{S}(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n)$.*

The proof is left as an exercise.

DEFINITION 1.11 (Tempered Distributions). A linear functional T is called a *tempered distribution* if for any sequence $\{\varphi_j\}_j$ of functions in $\mathcal{S}(\mathbb{R}^n)$ converging to zero in $\mathcal{S}(\mathbb{R}^n)$, we have

$$T(\varphi_j) \rightarrow 0, \quad (j \rightarrow +\infty).$$

It can be checked that the set of tempered distribution, denoted as $\boxed{\mathcal{S}'(\mathbb{R}^n)}$, is the dual of Schwartz Space $\mathcal{S}(\mathbb{R}^n)$.

Recall the semi-norm $|\cdot|_{\alpha,\beta}$ defined in (1.1). We define a new norm $|\cdot|_m$ as

$$|\varphi|_m := \sum_{|\alpha|, |\beta| \leq m} |\varphi|_{\alpha,\beta},$$

It can be seen that $|\varphi|_m \leq |\varphi|_{m+1}$.

LEMMA 1.12. *“ $T \in \mathcal{S}'(\mathbb{R}^n)$ ” is equivalent to the following statement:*

there exists a constant C such that $\exists m \in \{0\} \cup \mathbb{N}^+$ s.t. $\boxed{|T(\varphi)| \leq C|\varphi|_m}$, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$.

PROOF. (\Leftarrow) Assume $\exists m \in \{0\} \cup \mathbb{N}^+$ s.t. $|T(\varphi)| \leq C|\varphi|_m$, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$. Then for every sequence $\{\varphi_k\}_k \subset \mathcal{S}(\mathbb{R}^n)$ satisfying $\varphi_k \rightarrow 0$ ($k \rightarrow +\infty$), we have $|\varphi_k|_m \rightarrow 0$ ($k \rightarrow +\infty$). So $|T(\varphi_k)| \leq C|\varphi_k|_m \rightarrow 0$ ($k \rightarrow +\infty$). This means $T \in \mathcal{S}'(\mathbb{R}^n)$.

(\Rightarrow) Assume $T \in \mathcal{S}'(\mathbb{R}^n)$. Suppose that the claim is not true, then for every positive integer M , and every $m \in \{0\} \cup \mathbb{N}^+$, there exists $\varphi_{M,m} \in \mathcal{S}(\mathbb{R}^n)$ such that

$$|T(\varphi_{M,m})| > M|\varphi_{M,m}|_m, \quad \text{so} \quad |T(\varphi_{M,m}/(M|\varphi_{M,m}|_m))| > 1.$$

Let $\phi_{M,m} = \varphi_{M,m}/(M|\varphi_{M,m}|_m)$, then $|\phi_{M,m}|_m = \frac{1}{M}$ and $|T(\phi_{M,m})| \geq 1$. Further, we denote $\phi_M := \phi_{M,M}$, then $|\phi_M|_M = \frac{1}{M}$ and

$$|T(\phi_M)| \geq 1, \quad \forall M \in \mathbb{N}^+. \tag{1.3}$$

Now for every $m \in \{0\} \cup \mathbb{N}^+$, when j is large enough, we have

$$|\phi_j|_m \leq |\phi_j|_j = \frac{1}{j} \rightarrow 0 \quad (j \rightarrow +\infty).$$

So according to the Definition 1.5, we have $\phi_j \rightarrow 0$ in $\mathcal{S}(\mathbb{R}^n)$, so according to the definition of tempered Distributions we shall have $|T(\phi_j)| \rightarrow 0$. But this is contradictory with (1.3).

The proof is complete. \square

The notion of “tempered function” and “tempered distribution” are closely related. Every tempered function f defines a tempered distribution $T_f \in \mathcal{S}'(\mathbb{R}^n)$ by the following way:

$$T_f(\varphi) := \int_{\Omega} f(x)\varphi(x) dx, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n).$$

At the first glance, the definition of tempered distribution is not a generalization of the definition of tempered function. But the following theorem will characterize tempered distributions through tempered functions.

THEOREM 1.13 (Schwartz representation Theorem). *Every $T \in \mathcal{S}'(\mathbb{R}^n)$ can be represented as a sum of certain order of derivative of continuous tempered functions in $\mathcal{S}'(\mathbb{R}^n)$, i.e. for every $T \in \mathcal{S}'(\mathbb{R}^n)$, there exist a finite collection $T_{\alpha,\beta}$ of bounded continuous functions such that*

$$T = \sum_{|\alpha|+|\beta| \leq m} x^\alpha D^\beta T_{\alpha,\beta}$$

See §1.4 in <https://math.mit.edu/~rbm/iml/Chapter1.pdf>

THEOREM 1.14. *For $1 \leq p \leq +\infty$, there holds $\mathcal{S}'(\mathbb{R}^n) \subset L^p(\mathbb{R}^n) \subset \mathcal{S}'(\mathbb{R}^n)$.*

1.3. Fourier transforms

DEFINITION 1.15 (Fourier Transform on $\mathcal{S}'(\mathbb{R}^n)$). Let $f \in \mathcal{S}'(\mathbb{R}^n)$, then the *Fourier transform* of f is defined as

$$(\mathcal{F}f)(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} f(x) dx, \quad \forall \xi \in \mathbb{R}^n,$$

where $x \cdot \xi = \sum_{i=1}^n x_i \xi_i$. We also denote the Fourier transform of f as \hat{f} .

DEFINITION 1.16 (Inverse Fourier Transform). Let $f \in \mathcal{S}'(\mathbb{R}^n)$, then the *inverse Fourier transform* of f is defined as

$$(\mathcal{F}^{-1}f)(x) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix \cdot \xi} f(\xi) d\xi, \quad \forall \xi \in \mathbb{R}^n.$$

We also denote the inverse Fourier transform of f as \check{f} .

LEMMA 1.17. *For every $f, g \in \mathcal{S}'(\mathbb{R}^n)$, we have:*

- (1) $\mathcal{F}, \mathcal{F}^{-1}: \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ are linear bijection;
- (2) $\langle \hat{f}, g \rangle = \langle f, \hat{g} \rangle$;
- (3) $(f, g) = (\hat{f}, \hat{g})$. (Parseval's Relation);
- (4) $\mathcal{F}(f * g) = (2\pi)^{n/2} \hat{f} \cdot \hat{g}$;
- (5) $\mathcal{F}(f \cdot g) = (2\pi)^{-n/2} \hat{f} * \hat{g}$.

Let's define an operator

$$\mathcal{R}: f(x) \in \mathcal{S}'(\mathbb{R}^n) \mapsto (\mathcal{R}f)(x) = f(-x) \in \mathcal{S}'(\mathbb{R}^n).$$

Then these four operators $\{I, \mathcal{R}, \mathcal{F}, \mathcal{F}^{-1}\}$ act very like $\{1, -1, i, -i\}$. Denote a multiplication operation X_j as $X_j \varphi(x) := x_j \varphi(x)$. We have the following relations:

PROPOSITION 1.18.

- | | |
|--|---|
| (1.a) $\mathcal{R}\mathcal{F} = \mathcal{F}\mathcal{R} = \mathcal{F}^{-1}$; | $((-1) \cdot i = i \cdot (-1) = -i)$ |
| (1.b) $\mathcal{R}\mathcal{F}^{-1} = \mathcal{F}^{-1}\mathcal{R} = \mathcal{F}$; | $((-1) \cdot (-i) = (-i) \cdot (-1) = i)$ |
| (1.c) $\mathcal{F}\mathcal{F} = \mathcal{F}^{-1}\mathcal{F}^{-1} = \mathcal{R}$; | $(i \cdot i = (-i) \cdot (-i) = -1)$ |
| (1.d) $\mathcal{R}\mathcal{R} = I$. | $((-1) \cdot (-1) = 1)$ |
| (2.a) $\mathcal{F}D_j = X_j\mathcal{F}, \quad \mathcal{F}X_j = -D_j\mathcal{F}$, | |
| (2.b) $\mathcal{F}^2D_j = -D_j\mathcal{F}^2, \quad \mathcal{F}^2X_j = -X_j\mathcal{F}^2$. | |

THEOREM 1.19 (Plancherel Theorem). \mathcal{F} and \mathcal{F}^{-1} defined on $\mathcal{S}'(\mathbb{R}^n)$ can be extended uniquely to a unitary operator on $L^2(\mathbb{R}^n)$.

PROOF. $C_c^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n) \subset L^2(\mathbb{R}^n)$ and $\overline{C_c^\infty(\mathbb{R}^n)}^{\|\cdot\|_{L^2(\mathbb{R}^n)}} = L^2(\mathbb{R}^n)$, So $\mathcal{S}(\mathbb{R}^n)$ is dense in $L^2(\mathbb{R}^n)$ with respect to the L^2 norm.

For any $f \in L^2(\mathbb{R}^n)$, let $\{\varphi_n\}_n \subset \mathcal{S}(\mathbb{R}^n)$ such that $\|\varphi_n - f\|_{L^2(\mathbb{R}^n)} \rightarrow 0$ ($n \rightarrow +\infty$), then

$$\|\mathcal{F}\varphi_m - \mathcal{F}\varphi_n\|_{L^2(\mathbb{R}^n)} = \|\mathcal{F}(\varphi_m - \varphi_n)\|_{L^2(\mathbb{R}^n)}$$

and by the Parseval's Relation we can continue

$$\|\mathcal{F}\varphi_m - \mathcal{F}\varphi_n\|_{L^2(\mathbb{R}^n)} = \|\varphi_m - \varphi_n\|_{L^2(\mathbb{R}^n)} \rightarrow 0, \quad (m, n \rightarrow +\infty).$$

Therefore $\{\mathcal{F}\varphi_n\}_n$ is a Cauchy sequence in $L^2(\mathbb{R}^n)$ and has a limit. We denote the limit as $\mathcal{F}f$ and assign it to f as the Fourier transform of f . \square

The Fourier transform and inverse Fourier transform can also be uniquely extended on $\mathcal{S}'(\mathbb{R}^n)$.

DEFINITION 1.20 (Fourier Transform on $\mathcal{S}'(\mathbb{R}^n)$). Let $T \in \mathcal{S}'(\mathbb{R}^n)$, then the *Fourier transform* and *inverse Fourier transform* of T are defined to be the linear functionals $\mathcal{F}T$ on $\mathcal{S}(\mathbb{R}^n)$ given by

$$\begin{aligned} (\mathcal{F}T)(\varphi) &:= T(\hat{\varphi}), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n), \\ (\mathcal{F}^{-1}T)(\varphi) &:= T(\check{\varphi}), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

THEOREM 1.21. For every $\varphi \in \mathcal{S}(\mathbb{R}^n)$, $T \in \mathcal{S}'(\mathbb{R}^n)$, we have:

- (1) $\mathcal{F}, \mathcal{F}^{-1}: \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ are linear continuous bijection.
- (2) $\mathcal{F}(\varphi * T) = (2\pi)^{n/2} \hat{\varphi} \cdot \hat{T}$;
- (3) $\mathcal{F}(\varphi \cdot T) = (2\pi)^{-n/2} \hat{\varphi} * \hat{T}$.

The Proposition 1.18 also holds on $\mathcal{S}'(\mathbb{R}^n)$. The operator \mathcal{R} for $T \in \mathcal{S}'(\mathbb{R}^n)$ is defined as:

$$(\mathcal{R}T)(\varphi) := T(\mathcal{R}\varphi), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n).$$

The Fourier transform are both $(1, +\infty)$ -type and $(2, 2)$ -type bounded. And we have $L^p(\mathbb{R}^n) \subset L^1(\mathbb{R}^n) + L^2(\mathbb{R}^n)$ when $1 < p < 2$. Therefore we can define the Fourier transform \mathcal{F} on $L^1(\mathbb{R}^n) + L^2(\mathbb{R}^n)$ by (1.4) and then study the boundedness of Fourier transform on $L^p(\mathbb{R}^n)$ with $1 < p < 2$. For details about these $L^1(\mathbb{R}^n) + L^2(\mathbb{R}^n)$ things, please Google "Riesz–Thorin theorem". Therefore according to the Marcinkiewicz interpolation theorem (see [25, Appendix B]), we have the following result.

$$\left\{ \begin{array}{l} f = f_1 + f_2 \in L^p(\mathbb{R}^n), \\ f_1 \in L^1(\mathbb{R}^n), f_2 \in L^2(\mathbb{R}^n), \\ \mathcal{F}_1: \text{Fourier transform from } L^1(\mathbb{R}^n) \text{ to } L^1(\mathbb{R}^n), \\ \mathcal{F}_2: \text{Fourier transform from } L^2(\mathbb{R}^n) \text{ to } L^2(\mathbb{R}^n), \\ \mathcal{F}(f) := \mathcal{F}_1(f_1) + \mathcal{F}_2(f_2). \end{array} \right. \quad (1.4)$$

THEOREM 1.22 (Hausdorff-Young inequality). Define the Fourier transform on $L^1(\mathbb{R}^n) + L^2(\mathbb{R}^n)$ by (1.4), then there exists a constant C_p such that for all $f \in L^p(\mathbb{R}^n)$, ($1 \leq p \leq 2$), we have

$$\|\hat{f}\|_{L^{p'}(\mathbb{R}^n)} \leq C \|f\|_{L^p(\mathbb{R}^n)},$$

where $1 \leq p \leq 2$ and $1/p + 1/p' = 1$.

PROOF. We know that \mathcal{F} and \mathcal{F}^{-1} are bounded from $L^1(\mathbb{R}^n)$ to $L^\infty(\mathbb{R}^n)$ and $L^2(\mathbb{R}^n)$ to $L^2(\mathbb{R}^n)$. So according to Marcinkiewicz interpolation theorem, $\forall t \in [0, 1]$, \mathcal{F} and \mathcal{F}^{-1} are bounded from $L^{t_1}(\mathbb{R}^n)$ to $L^{t_2}(\mathbb{R}^n)$, where

$$\begin{cases} t_1 = \left(t \cdot \frac{1}{2} + (1-t) \cdot 1 \right)^{-1} = \frac{2}{2-t} \\ t_2 = \left(t \cdot \frac{1}{2} + (1-t) \cdot 0 \right)^{-1} = \frac{2}{t} \end{cases} .$$

Let $p = t_1$ $p' = t_2$, then we proved the theorem. □

Exercise

EXERCISE 1.1. Prove Lemma 1.1.

EXERCISE 1.2. Prove Lemma 1.3.

EXERCISE 1.3. Prove Lemma 1.10.

CHAPTER 2

Pseudodifferential operators

In this chapter we introduce the pseudodifferential operators, and in most of the place we abbreviate it as Ψ DOs. First, we introduce symbols and its asymptotics. Then the Ψ DOs are its kernels are defined. Finally, we prove an important property of Ψ DOs—the pseudolocal property. Other references are [26, Chapter 6], [11, §1 & §3].

2.1. Symbols

Recall the general form $p(x, D) = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha$ of the linear differential operators mentioned is §1.1. For a test function φ , we have

$$\begin{aligned} p(x, D)\varphi(x) &= \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha \varphi(x) = \sum_{|\alpha| \leq m} a_\alpha(x) \mathcal{F}^{-1} \{ \widehat{D^\alpha \varphi} \}(x) \\ &= \sum_{|\alpha| \leq m} a_\alpha(x) \mathcal{F}^{-1} \{ \xi^\alpha \hat{\varphi} \}(x) \\ &= \sum_{|\alpha| \leq m} a_\alpha(x) (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \xi^\alpha \hat{\varphi}(\xi) \, d\xi \\ &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha \hat{\varphi}(\xi) \, d\xi \\ &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix \cdot \xi} p(x, \xi) \hat{\varphi}(\xi) \, d\xi. \end{aligned}$$

This observation encourages us to define operators by functions $p(x, \xi)$.

DEFINITION 2.1 (Kohn-Nirenberg symbol). Let $m \in (-\infty, +\infty)$. Then we define S^m to be the set of all functions $\sigma(x, \xi) \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{C})$ such that for any two multi-indices α and β , there is a positive constant $C_{\alpha, \beta}$, independent of (x, ξ) , such that

$$\boxed{|(D_x^\alpha D_\xi^\beta \sigma)(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - |\beta|}, \quad \forall x, \xi \in \mathbb{R}^n}$$

holds. We call any function σ in S^m a *symbol* of order m . We write $S^{-\infty} = \bigcap_{m \in \mathbb{R}} S^m$ and $S^{+\infty} = \bigcup_{m \in \mathbb{R}} S^m$.

EXAMPLE 2.2. Here we give some examples of symbols.

- $\sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$ is a symbol of order m when $a_\alpha \in \mathcal{S}(\mathbb{R}^n)$;
- $\mathcal{S}(\mathbb{R}^n) \subset S^{-\infty}$;
- Fix a bounded $\psi \in C^\infty(\mathbb{R}^n)$, then $\psi(x) \langle \xi \rangle^m$ is a symbol of order m ;
- Fix a $\phi \in C_c^\infty(\mathbb{R}^n)$ with $\phi(0) = 1$, then $(1 - \phi(\xi))(1 + |\xi|)^m$ is a symbol of order m .

LEMMA 2.3. Assume $\sigma_j \in S^{m_j}$ ($j = 1, 2$), then $\sigma_1 \sigma_2 \in S^{m_1 + m_2}$. $\partial^\alpha \sigma_1 \in S^{m_1 - |\alpha|}$.

The proof is left as an exercise.

One can also define a more general symbol which the effect of x is taken into consideration, and the dimension of x variable and ξ variable can be different.

DEFINITION 2.4. Let $m \in (-\infty, +\infty)$ and $0 \leq \delta < \rho \leq 1$. Then we define $S_{\rho, \delta}^m$ to be the set of all functions $\sigma(x, \xi) \in C^\infty(\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}; \mathbb{C})$ such that for any two multi-indices α and β , there is a positive constant $C_{\alpha, \beta}$, depending on α and β only, for which

$$\boxed{|(D_x^\alpha D_\xi^\beta \sigma)(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\beta| + \delta|\alpha|}, \quad \forall x \in \mathbb{R}^{n_1}, \xi \in \mathbb{R}^{n_2}}$$

holds. We also call any function σ in $S_{\rho, \delta}^m$ a *symbol*.

The Kohn-Nirenberg symbol $S^m = S_{1, 0}^m$. In what follows, we only focus on S^m , and the situations for $S_{\rho, \delta}^m$ shall be followed in similar manners.

Now we introduce an important notion: the asymptotic expansion of symbols.

DEFINITION 2.5 (Asymptotics). Let symbol $a \in S^m$ and $a_j \in S^{m_j}$ ($j = 0, 1, \dots$) where the orders m_j satisfies

$$m = m_0 > m_1 > \dots > m_j > m_{j+1} \rightarrow -\infty, \quad j \rightarrow \infty.$$

If

$$a - \sum_{j=0}^N a_j \in S^{m_{N+1}},$$

holds for every integer N , we write

$$a \sim \sum_j a_j \text{ in } S^m,$$

and we call $\{a_j\}$ an asymptotics of a . The a_0 is called the *principal symbol* of a .

We often write $a = b + S^m$ as a shorthand of $a = b + r$ for some $r \in S^m$. Then we can summarize Definition 2.5 as follows,

$$\boxed{a \sim \sum_j a_j \text{ in } S^{m_{N+1}} \Leftrightarrow a = \sum_{j=0}^N a_j + S^{m_{N+1}}.}$$

Now let's randomly pick up some m, m_j that satisfy the requirement in Definition 2.5, and randomly pick up $a_j \in S^{m_j}$. A natural question is to ask, does there exist $a \in S^m$ such that $a \sim \sum_j a_j$ in S^m ? The answer is yes.

THEOREM 2.6. For any m and m_j satisfying

$$m = m_0 > m_1 > \dots > m_j > m_{j+1} \rightarrow -\infty, \quad j \rightarrow \infty,$$

and for any $a_j \in S^{m_j}$, there exists a symbol (not unique) $a \in S^m$ such that $a \sim \sum_j a_j$ in S^m .

When $x \rightarrow +\infty$, $\frac{1}{1-1/\langle x \rangle} = 1 + 1/\langle x \rangle + 1/\langle x \rangle^2 + \mathcal{O}(1/\langle x \rangle^3)$. Arbitrarily pick up α_j , is there a function $f(x)$ such that in $[1, +\infty)$,

$$f(x) = \sum_{0 \leq j \leq N} \alpha_j \langle x \rangle^j + \mathcal{O}(1/\langle x \rangle^{N+1}), \quad x \rightarrow +\infty, \quad (2.1)$$

holds for all $N \in \mathbb{N}$? The answer is no and an example is $\alpha_j := j!$ (the convergence radius goes to infinity as N grows). The problem is that $\alpha_j/\langle x = 1 \rangle^j$ will be too big when $j \rightarrow +\infty$. However, we can fix this problem by cutoff, so that there exist a function f (not unique!) such that (2.1) holds on intervals $[A_j, +\infty)$ where the A_j is in accordance with a_j , $A_1 < A_2 < \dots$. The key step is to choose a cutoff function χ_j to cutoff term $\alpha_j/\langle x \rangle^j$ such that

$$\chi_j(x) \alpha_j / \langle x \rangle^j \leq 1/2^j, \text{ or } 1/3^j \text{ etc.} \quad (2.2)$$

The following function satisfies the requirement:

$$\begin{cases} \chi_j \in C^\infty(\mathbb{R}), \\ \chi_j \equiv 0, |x| \leq 2(\alpha_j)^{1/j}, \\ \chi_j \equiv 1, |x| \geq 1 + 2(\alpha_j)^{1/j}, \\ 0 \leq \chi_j \leq 1, \text{ otherwise.} \end{cases} \quad (2.3)$$

The requirement (2.3) can also be realized by fix some χ satisfying

$$\begin{cases} \chi \in C^\infty(\mathbb{R}), 0 \leq \chi(x) \leq 1 \\ \chi \equiv 0 \text{ when } |x| \leq 1, \chi \equiv 1 \text{ when } |x| \geq 2. \end{cases} \quad (2.4)$$

and then set $\chi_j(x) := \chi(\epsilon_j x)$, where the ϵ_j shall be chosen according to (2.3).

SKETCH OF THE PROOF OF THEOREM 2.6. Choose suitable coefficients ϵ_j and define

$$a(x, \xi) := \sum_{j \geq 0} \chi(\epsilon_j \xi) a_j(x, \xi), \quad x, \xi \in \mathbb{R}^n.$$

It can be checked that $\chi(\epsilon_j \xi) \in S^0$. For any fixed (x_0, ξ_0) , there is only finitely many terms in $\sum_{j \geq 0} \chi(\epsilon_j \xi_0) a_j(x_0, \xi_0)$ which are non-zero, so $a(x, \xi) \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n; \mathbb{C})$. Moreover, we need to show first a is a symbol, and second a is an asymptotics of a_j .

First, we show that $a \in S^m$. It can be checked that for any multi-index β ,

$$|D_\xi^\beta(\chi(\epsilon_j \xi))| \leq C_\beta \langle \xi \rangle^{-\beta}, \quad \forall \xi \in \mathbb{R}^n,$$

where the constant C_β is independent of ϵ_j . we notice that every term $\chi(\epsilon_j \xi) a_j(x, \xi)$ is in $S^{0+m_j} = S^{m_j}$, so

$$\begin{aligned} D_x^\alpha D_\xi^\beta(\chi(\epsilon_j \xi) a_j(x, \xi)) &= \chi(2\epsilon_j \xi) D_x^\alpha D_\xi^\beta(\chi(\epsilon_j \xi) a_j(x, \xi)) \\ &= \chi(2\epsilon_j \xi) \sum_{\beta' \leq \beta} \binom{\beta}{\beta'} D_\xi^{\beta'}(\chi(\epsilon_j \xi)) \cdot (D_x^\alpha D_\xi^{\beta-\beta'} a_j)(x, \xi) \\ &\leq \chi(2\epsilon_j \xi) \sum_{\beta' \leq \beta} \binom{\beta}{\beta'} C_{\beta'} \langle \xi \rangle^{-|\beta'|} \cdot C_{\alpha, \beta, \beta'} \langle \xi \rangle^{m_j - |\beta| + |\beta'|} \\ &= \chi(2\epsilon_j \xi) \sum_{\beta' \leq \beta} \binom{\beta}{\beta'} C_{\beta'} C_{\alpha, \beta, \beta'} \langle \xi \rangle^{m_j - |\beta|} = \chi(2\epsilon_j \xi) C_{\alpha, \beta} \langle \xi \rangle^{m_j - |\beta|} \\ &= C_{\alpha, \beta} \langle \xi \rangle^{m_j - m} \chi(2\epsilon_j \xi) \cdot \langle \xi \rangle^{m - |\beta|} \\ &\leq C_{\alpha, \beta} (2\epsilon_j)^{m - m_j} \chi(2\epsilon_j \xi) \cdot \langle \xi \rangle^{m - |\beta|} \\ &\leq C_{\alpha, \beta} (2\epsilon_j)^{m - m_j} \cdot \langle \xi \rangle^{m - |\beta|}, \end{aligned}$$

where the change " $\langle \xi \rangle^{m_j - m} \rightarrow (2\epsilon_j)^{m - m_j}$ " is due to the presence of $\chi(2\epsilon_j \xi)$. Hence,

$$|D_x^\alpha D_\xi^\beta a(x, \xi)| \leq \langle \xi \rangle^{m - |\beta|} \cdot \sum_{j \geq 0} C_{\alpha, \beta} (2\epsilon_j)^{m - m_j}.$$

We choose ϵ_j to decrease fast enough such that $\sum_{j \geq 0} C_{\alpha, \beta} (2\epsilon_j)^{m - m_j}$ is finite for every α, β (see [26, Theorem 6.10] for details). We proved $a \in S^m$.

Second, to show $a \sim \sum_j a_j$ in S^m , we see

$$\begin{aligned} a - \sum_{0 \leq j \leq N} a_j &= \sum_{0 \leq j \leq N} [\chi(\epsilon_j \xi) - 1] a_j(x, \xi) + \sum_{j \geq N+1} \chi(\epsilon_j \xi) a_j(x, \xi) \\ &\in S^{-\infty} + S^{m_{N+1}} = S^{m_{N+1}}. \end{aligned}$$

The proof is complete. \square

2.2. Pseudodifferential operators

2.2.1. Some basics about the Ψ DOs. Based on the notion of symbols, we introduce the pseudodifferential operators.

DEFINITION 2.7 (Pseudodifferential operator, Ψ DO). Let σ be a symbol. Then the *pseudo-differential operator* T_σ , defined on $\mathcal{S}(\mathbb{R}^n)$ and associated with σ , is defined as

$$\begin{aligned} (T_\sigma\varphi)(x) &:= (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix\cdot\xi} \sigma(x, \xi) \hat{\varphi}(\xi) \, d\xi \\ &= \boxed{(2\pi)^{-n} \iint_{\mathbb{R}^n \times \mathbb{R}^n} e^{i(x-y)\cdot\xi} \sigma(x, \xi) \varphi(y) \, dy \, d\xi}, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

We denote the set of Ψ DOs of order m as Ψ^m . We write $\Psi^{-\infty} = \bigcap_{m \in \mathbb{R}} \Psi^m$ and $\Psi^{+\infty} = \bigcup_{m \in \mathbb{R}} \Psi^m$.

EXAMPLE 2.8. Here we give some examples of Ψ DOs:

- $-\Delta \in \Psi^2$, with symbol $|\xi|^2$;
- $\sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha \in \Psi^m$, with symbol $\sum_{|\alpha| \leq m} a_\alpha(x) \xi^\alpha$;
- $(I - \Delta)^{m/2} \in \Psi^m$, which is defined by the symbol $\langle \xi \rangle^m = (I + |\xi|^2)^{m/2}$;
- The DtN map of the Calderón problem is a Ψ DO living on the boundary, see [18].

It is an interesting question to ask for the symbol when given a certain Ψ DO.

EXAMPLE 2.9. Some simple Ψ DOs whose symbol are also simple:

- $D \mapsto \xi$;
- $-\Delta = D \cdot D$, so $-\Delta \mapsto |\xi|^2$.

Similar to Lemma 2.3, we have the following claim, whose proof will be provided in Theorem 5.3.

LEMMA 2.10. *Assume $\sigma_j \in S^{m_j}$ ($j = 1, 2$), then $T_{\sigma_1} \circ T_{\sigma_2} \in \Psi^{m_1+m_2}$.*

We show that the map $\sigma \mapsto T_\sigma$ is a bijection.

LEMMA 2.11. *map $\sigma \mapsto T_\sigma \in \mathcal{L}(\mathcal{S}(\mathbb{R}^n), \mathcal{S}(\mathbb{R}^n))$ is a bijection.*

PROOF. The Ψ DO T_σ is defined by σ , so the surjectivity is obvious. The injectivity amounts to prove $T_\sigma = T_\tau \Rightarrow \sigma = \tau$.

Let's assume σ and τ are two symbols and $T_\sigma = T_\tau$, then

$$\int e^{ix\cdot\xi} [\sigma(x, \xi) - \tau(x, \xi)] \hat{\varphi}(\xi) \, d\xi = 0$$

holds for any $x \in \mathbb{R}^n$ and any $\varphi \in \mathcal{S}(\mathbb{R}^n)$. Replace φ by its inverse Fourier transform, and fix x to some x_0 , we can see

$$\int e^{ix_0\cdot\xi} [\sigma(x_0, \xi) - \tau(x_0, \xi)] \varphi(\xi) \, d\xi = 0.$$

The arbitrary of φ gives

$$e^{ix_0\cdot\xi} [\sigma(x_0, \xi) - \tau(x_0, \xi)] = 0, \quad \forall x_0 \in \mathbb{R}^n.$$

And the arbitrary of x_0 gives $\sigma(x_0, \xi) = \tau(x_0, \xi)$ for $\forall x_0 \in \mathbb{R}^n$. The injectivity is proved. We arrive at the conclusion. \square

REMARK 2.12. For people who the first time encounter the form

$$T_\sigma \varphi(x) \simeq \int_{\mathbb{R}^n} e^{ix \cdot \xi} \sigma(x, \xi) \hat{\varphi}(\xi) \, d\xi,$$

one may think that $T_\sigma \varphi(x)$ is just the inverse Fourier transform of $\sigma(x, \xi) \hat{\varphi}(\xi)$, and consequently, $\sigma(x, \xi) \hat{\varphi}(\xi)$ can be recovered by taking the Fourier transform of $T_\sigma \varphi(x)$. Unfortunately this is not true. The function $\sigma(x, \xi) \hat{\varphi}(\xi)$ depends not only ξ but also x , so that is not a Fourier transform anymore. When a symbol a is independent of x , we have

$$\mathcal{F}\{T_\sigma \varphi\}(\xi) = a(\xi) \hat{\varphi}(\xi).$$

But when a depends on x , the Fourier transform of $(T_\sigma \varphi)(x)$ is generally *NOT* $a(x, \xi) \hat{\varphi}(\xi)$. In generally, we cannot use the expression above to get the symbol a :

$$\boxed{a(x, \xi) \hat{\varphi}(\xi) \neq \mathcal{F}\{(T_\sigma \varphi)(\cdot)\}(\xi).}$$

Similar to generalizing the Fourier transform from functions to distributions, the notion of Ψ DOs can also extend to $\mathcal{S}'(\mathbb{R}^n)$ by using duality arguments. Formally speaking, we have the following computation,

$$\begin{aligned} (T_\sigma u, \varphi) &= \int (T_\sigma u) \overline{\varphi} \, dx = \int [(2\pi)^{-n/2} \int e^{ix \cdot \xi} \sigma(x, \xi) \hat{u}(\xi) \, d\xi] \overline{\varphi(x)} \, dx \\ &= \int [(2\pi)^{-n} \int e^{i(x-y) \cdot \xi} \sigma(x, \xi) u(y) \, dy \, d\xi] \overline{\varphi(x)} \, dx \\ &= \int u(y) \overline{[(2\pi)^{-n} \int e^{i(y-x) \cdot \xi} \overline{\sigma(x, \xi)} \varphi(x) \, dx \, d\xi]} \, dy \\ &\sim \int u(y) \overline{[(2\pi)^{-n} \int e^{i(y-x) \cdot \xi} \sum_{\alpha} \frac{(x-y)^\alpha}{\alpha!} \partial_y^\alpha \overline{\sigma(y, \xi)} \varphi(x) \, dx \, d\xi]} \, dy \quad (\text{Taylor's}) \\ &\sim \int u(y) \overline{[(2\pi)^{-n} \int (-D)_\xi^\alpha (e^{i(y-x) \cdot \xi}) \sum_{\alpha} \frac{1}{\alpha!} \partial_y^\alpha \overline{\sigma(y, \xi)} \varphi(x) \, dx \, d\xi]} \, dy \\ &\sim \int u(y) \overline{[(2\pi)^{-n} \int e^{i(y-x) \cdot \xi} \sum_{\alpha} \frac{1}{\alpha!} \partial_y^\alpha D_\xi^\alpha \overline{\sigma(y, \xi)} \varphi(x) \, dx \, d\xi]} \, dy \\ &= \int u(y) \overline{[(2\pi)^{-n/2} \int e^{iy \cdot \xi} \sum_{\alpha} \frac{1}{\alpha!} \partial_y^\alpha D_\xi^\alpha \overline{\sigma(y, \xi)} \hat{\varphi}(\xi) \, d\xi]} \, dy \\ &= \int u(y) \overline{T_{\sigma^*} \varphi(y)} \, dy, \quad \sigma^*(y, \xi) := \sum_{\alpha} \frac{1}{\alpha!} \partial_y^\alpha D_\xi^\alpha \overline{\sigma(y, \xi)} \\ &= (u, T_{\sigma^*} \varphi). \end{aligned} \tag{2.5}$$

The computation (2.5) implies the existence of the adjoint of T_σ (denoted as T_σ^*), and we leave the rigorous proof of the existence of T_σ^* to §5.3. Now, by assuming the existence of T_σ^* , we extend the domain of T_σ from $\mathcal{S}(\mathbb{R}^n)$ to $\mathcal{S}'(\mathbb{R}^n)$ as follows.

DEFINITION 2.13 (Pseudodifferential operators in \mathcal{S}'). Let σ be a symbol. For every $u \in \mathcal{S}'(\mathbb{R}^n)$, we can define the *pseudo-differential operator* T_σ acting on u as

$$\boxed{(T_\sigma u, \varphi) := (u, T_\sigma^* \varphi), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n).}$$

where T_σ^* is the adjoint of T_σ and the bracket (\cdot, \cdot) signifies the pair of distributions with test functions.

LEMMA 2.14. Let σ be a symbol, and denote its corresponding Ψ DO as T_σ . Then $T_\sigma(\mathcal{S}(\mathbb{R}^n)) \subset \mathcal{S}(\mathbb{R}^n)$. And also, $T_\sigma(\mathcal{S}'(\mathbb{R}^n)) \subset \mathcal{S}'(\mathbb{R}^n)$.

SKETCH OF THE PROOF. Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$, we need to show $x^\alpha D^\beta(T_\sigma\varphi)$ are bounded in \mathbb{R}^n . We can show

$$x^\alpha D^\beta(T_\sigma\varphi)(x) \simeq \sum_{\alpha', \beta'} \int e^{ix \cdot \xi} \tilde{\sigma}_{\alpha', \beta'}(x, \xi) \mathcal{F}\{x^{\alpha'} D^{\beta'} \varphi\}(\xi) d\xi$$

where $\tilde{\sigma}_{\alpha', \beta'}$ are also symbols of certain orders, say, m . Because $\varphi \in \mathcal{S}(\mathbb{R}^n)$, we know that the Fourier transform $\mathcal{F}\{x^{\alpha'} D^{\beta'} \varphi\} \in \mathcal{S}(\mathbb{R}^n)$, so

$$|\mathcal{F}\{x^{\alpha'} D^{\beta'} \varphi\}(\xi)| \lesssim \langle \xi \rangle^{-M}$$

for any positive integer M . Hence,

$$|x^\alpha D^\beta(T_\sigma\varphi)(x)| \lesssim \sum_{\alpha', \beta'} \int \langle \xi \rangle^m \langle \xi \rangle^{-M} d\xi < +\infty$$

when we take M to be large enough.

For the second conclusion, from $(T_\sigma u, \varphi) := (u, T_\sigma^* \varphi)$, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$ we have

$$|(T_\sigma u, \varphi)| \leq \|u\| \|T_\sigma^* \varphi\| \lesssim \|u\| |\varphi|_m$$

where m is the order of σ . Then by Lemma 1.12 we can conclude $T_\sigma u \in \mathcal{S}'(\mathbb{R}^n)$. \square

In conclusion, there holds

$$T_\sigma: \begin{cases} \mathcal{S} \rightarrow \mathcal{S}, \\ \mathcal{S}' \rightarrow \mathcal{S}', \end{cases} \quad (2.6)$$

where \mathcal{S} is a shorthand for $\mathcal{S}(\mathbb{R}^n)$. Space \mathcal{S} represents functions which are extremely smooth (good), while \mathcal{S}' represents “functions” which are extremely rough (bad). To quantize the goodness and the badness, we introduce the potential spaces.

2.2.2. Sobolev spaces.

DEFINITION 2.15 (Sobolev spaces). We denote

$$\boxed{H^{s,p}(\mathbb{R}^n) := \{f \in \mathcal{S}'(\mathbb{R}^n); (I - \Delta)^{s/2} f \in L^p(\mathbb{R}^n)\}},$$

and define the norm $\|f\|_{H^{s,p}} := \|(I - \Delta)^{s/2} f\|_{L^p(\mathbb{R}^n)}$. Write $H^s(\mathbb{R}^n) := H^{s,2}(\mathbb{R}^n)$.

LEMMA 2.16. *The normed vector space $(H^{s,p}(\mathbb{R}^n), \|\cdot\|_{H^{s,p}})$ is a Banach space, and $(H^s(\mathbb{R}^n), \|\cdot\|_{H^s})$ is a Hilbert space.*

THEOREM 2.17. *Let $\sigma \in S^m$ and denote its corresponding Ψ DO as T_σ . Then the mapping $T_\sigma: H^s(\mathbb{R}^n) \rightarrow H^{s-m}(\mathbb{R}^n)$ is bounded.*

The proof of Theorem 2.17 is based on the L^2 boundedness of Ψ DOs of order 0. Formally speaking,

$$\|T_\sigma f\|_{H^{s-m}} = \|(I - \Delta)^{(s-m)/2} \circ T_\sigma f\|_{L^2} \lesssim \|(I - \Delta)^{s/2} f\|_{L^2} = \|f\|_{H^s}.$$

THEOREM 2.18. *For a fixed constant $s \in \mathbb{R}$, $\forall r, t: r \leq s \leq t$, $\forall C > 0$, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$, we have:*

$$\boxed{\|\varphi\|_{H^s}^2 \leq \frac{1}{C^{t-s}} \|\varphi\|_{H^t}^2 + C^{s-r} \|\varphi\|_{H^r}^2.} \quad (2.7)$$

REMARK 2.19. We know that when $t > s$, $\|\varphi\|_{H^s}$ can be controlled by $\|\varphi\|_{H^t}$:

$$\|\varphi\|_{H^s} \leq 1 \cdot \|\varphi\|_{H^t}.$$

The key point of Theorem 2.18 is that $\|\varphi\|_{H^s}$ can even be “controlled” by $c \cdot \|\varphi\|_{H^t}$ with $0 < c < 1$. But we need to pay for this: being dominated only by $c \cdot \|\varphi\|_{H^t}$ is not enough. Due to the fact that c is less than 1, certain “byproduct” should cost to compensate the advantage, and this so-called “byproduct” is $\|\varphi\|_{H^r}$.

PROOF OF THEOREM 2.18. When $0 < C \leq 1$, it is trivial. When $C \geq 1$, we have:

$$\begin{aligned} \|\varphi\|_{H^s}^2 &= \|\mathcal{F}^{-1}\sigma_{-s}\mathcal{F}\varphi\|_2^2 = \|\sigma_{-s}\hat{\varphi}\|_2^2 = \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\{\langle \xi \rangle \leq \sqrt{C}\}} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi + \int_{\{\langle \xi \rangle > \sqrt{C}\}} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\{\langle \xi \rangle \leq \sqrt{C}\}} \langle \xi \rangle^{2s-2r} \cdot \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi + \int_{\{\langle \xi \rangle^{-1} < \frac{1}{\sqrt{C}}\}} (\langle \xi \rangle^{-1})^{2t-2s} \cdot \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \\ &\leq \int_{\{\langle \xi \rangle \leq \sqrt{C}\}} \sqrt{C}^{2s-2r} \cdot \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi + \int_{\{\langle \xi \rangle^{-1} < \frac{1}{\sqrt{C}}\}} \left(\frac{1}{\sqrt{C}}\right)^{2t-2s} \cdot \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \\ &\leq \sqrt{C}^{2s-2r} \int_{\mathbb{R}^n} \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi + \left(\frac{1}{\sqrt{C}}\right)^{2t-2s} \int_{\mathbb{R}^n} \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \\ &= C^{s-r} \|\varphi\|_{H^r}^2 + C^{s-t} \|\varphi\|_{H^t}^2. \end{aligned}$$

This completes the proof. \square

REMARK 2.20. In the proof of Theorem 2.18, when φ is compactly supported and $s = 0$, and if we replace $\langle \xi \rangle$ by $|\xi|$ and choose $r = 0$ and C to be small enough and use the fact that

$$\int_{\{|\xi| \leq \sqrt{C}\}} |\hat{\varphi}(\xi)|^2 d\xi \lesssim C^{n/2} \sup_{|\xi| \leq \sqrt{C}} |\hat{\varphi}(\xi)|^2 \lesssim C^{n/2} \|\varphi\|_{L^1}^2 \leq C^{n/2} \sqrt{|\text{supp } \varphi|} \|\varphi\|_{L^2},$$

we can prove the Poincaré’s inequality $\|\varphi\|_{L^2} \lesssim \|(-\Delta)^{t/2}\varphi\|_{L^2}$ for the set of functions with uniformly compact support.

Noticing that $\langle \xi \rangle \geq 1$, we can further extend Theorem 2.18 to more generalized situation.

THEOREM 2.21. *For a fixed constant $s \in \mathbb{R}$, $\forall t \geq s$ and $\forall r \in \mathbb{R}^1$, $\forall \epsilon > 0$, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$, there exists a constant $C_{r,s,t,\epsilon}$ such that:*

$$\boxed{\|\varphi\|_{H^s}^2 \leq \epsilon \|\varphi\|_{H^t}^2 + C_{r,s,t,\epsilon} \|\varphi\|_{H^r}^2.} \quad (2.8)$$

As mentioned in Remark 2.19, Theorem 2.21 expresses the same information, in addition that the “byproduct” can be $\|\varphi\|_{H^r}$ with any $r \in \mathbb{R}^1$.

PROOF OF THEOREM 2.21. We pick up some constant $C > 1$ first, and then we decide its value later.

$$\begin{aligned} \|\varphi\|_{H^s}^2 &= \|\mathcal{F}^{-1}\sigma_{-s}\mathcal{F}\varphi\|_2^2 = \|\sigma_{-s}\hat{\varphi}\|_2^2 = \int_{\mathbb{R}^n} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\{1 \leq \langle \xi \rangle \leq \sqrt{C}\}} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi + \int_{\{\langle \xi \rangle > \sqrt{C}\}} \langle \xi \rangle^{2s} |\hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\{1 \leq \langle \xi \rangle \leq \sqrt{C}\}} \langle \xi \rangle^{2s-2r} \cdot \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi + \int_{\{\langle \xi \rangle^{-1} < \frac{1}{\sqrt{C}}\}} (\langle \xi \rangle^{-1})^{2t-2s} \cdot \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \end{aligned}$$

$$\begin{aligned}
&\leq \max\{1, \sqrt{C}^{2s-2r}\} \cdot \int_{\{1 \leq \langle \xi \rangle \leq \sqrt{C}\}} \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi \\
&\quad + \int_{\{\langle \xi \rangle^{-1} < \frac{1}{\sqrt{C}}\}} \left(\frac{1}{\sqrt{C}}\right)^{2t-2s} \cdot \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \\
&\leq \max\{1, \sqrt{C}^{2s-2r}\} \cdot \int_{\mathbb{R}^n} \langle \xi \rangle^{2r} |\hat{\varphi}(\xi)|^2 d\xi + \left(\frac{1}{\sqrt{C}}\right)^{2t-2s} \int_{\mathbb{R}^n} \langle \xi \rangle^{2t} |\hat{\varphi}(\xi)|^2 d\xi \\
&= \max\{1, \sqrt{C}^{2s-2r}\} \cdot \|\varphi\|_{H^r}^2 + C^{s-t} \|\varphi\|_{H^t}^2.
\end{aligned}$$

Now let $C = \epsilon^{t-s}$ and let $C_{r,s,t,\epsilon} = \max\{1, \sqrt{C}^{2s-2r}\}$, then we completes the proof. \square

2.2.3. Other phases. Beside the phase $(x-y) \cdot \xi$ in the expression

$$(T_\sigma \varphi)(x) = (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} \sigma(x, y, \xi) \varphi(y) dy d\xi$$

in Definition 2.7, it is possible to use more general functions as the phase functions and the corresponding operators are still Ψ DOs, i.e.,

$$P\varphi(x) = (2\pi)^{-n} \int e^{i\phi(x,y,\xi)} \sigma(x, y, \xi) \varphi(y) dy d\xi$$

will still be a Ψ DO is the ϕ satisfies certain conditions. See [24, §3.2] for details.

2.3. Kernels

The expression in Definition 2.7 can also be represented as

$$\boxed{(T_\sigma \varphi)(x) = \int_{\mathbb{R}^n} K(x, y) \varphi(y) dy,} \quad (2.9)$$

where $K(x, y)$ is called the kernel of T_σ ,

$$K(x, y) := (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} \sigma(x, \xi) d\xi,$$

and the integration shall be understood as an oscillatory integral (see Definition 3.8).

Differential operators such as $P = \sum_{j=1}^n x_j \partial_j$ maps $\mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$, $C_c^\infty(\mathbb{R}^n) \rightarrow C_c^\infty(\mathbb{R}^n)$, and $\mathcal{E}(\mathbb{R}^n) \rightarrow \mathcal{E}(\mathbb{R}^n)$, and so by duality argument, we know differential operators P maps $\mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$, $\mathcal{D}'(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}^n)$, and $\mathcal{E}'(\mathbb{R}^n) \rightarrow \mathcal{E}'(\mathbb{R}^n)$:

$$P: \begin{cases} \mathcal{E}'(\mathbb{R}^n) \rightarrow \mathcal{E}'(\mathbb{R}^n) \\ \mathcal{D}'(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}^n) \end{cases}$$

But for pseudo-differential operators T_σ , generally speaking, we only have $T_\sigma: \mathcal{E}'(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}^n)$.

$$T_\sigma: \begin{cases} \mathcal{E}'(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}^n) \\ \mathcal{E}'(\mathbb{R}^n) \not\rightarrow \mathcal{E}'(\mathbb{R}^n) \\ \mathcal{D}'(\mathbb{R}^n) \not\rightarrow \mathcal{D}'(\mathbb{R}^n). \end{cases}$$

A Ψ DO which maps \mathcal{E}' to \mathcal{E}' is called properly supported. In fact any Ψ DO can be divided into a properly supported part and a C^∞ -smooth part.

LEMMA 2.22. *Assume $m \in \mathbb{R}$ and $\sigma \in S^m$ is a symbol, and $K(x, y)$ is the kernel of T_σ . Then for any $\epsilon > 0$, there exists two symbols $\sigma_1 \in S^m$ and $\sigma_2 \in S^{-\infty}$ such that*

$\sigma = \sigma_1 + \sigma_2$, T_{σ_1} is properly supported, T_{σ_2} is smooth, and their kernels K_1, K_2 has the following properties:

$$\text{supp } K_1 \subset \{(x, y) \in \mathbb{R}^{2n}; |x - y| \leq \epsilon\}, \quad (\text{properly supported})$$

$$\text{supp } K_2 \subset \{(x, y) \in \mathbb{R}^{2n}; |x - y| \geq \epsilon/2\}. \quad (\text{smooth})$$

IDEA OF THE PROOF. The proof needs Theorem 5.6.

Fix a cutoff function $\chi \in C_c^\infty(\mathbb{R})$ such that $\chi(t) = 1$ when $|t| \leq \epsilon/2$ and $\chi(t) = 0$ when $|t| \geq \epsilon$. We have

$$T_\sigma \varphi(x) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) \varphi(y) dy d\xi = T' \varphi(x) + T'' \varphi(x),$$

where

$$\begin{cases} T' \varphi(x) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \chi(|x-y|^2) \sigma(x, \xi) \varphi(y) dy d\xi, \\ T'' \varphi(x) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} (1 - \chi(|x-y|^2)) \sigma(x, \xi) \varphi(y) dy d\xi. \end{cases} \quad (2.10)$$

By Theorem 5.6 we see that there exist $\sigma_1, \sigma_2 \in S^m$ such that $T' = T_{\sigma_1}$ and $T'' = T_{\sigma_2}$, so $T_\sigma = T_{\sigma_1} + T_{\sigma_2} = T_{\sigma_1 + \sigma_2}$. By Lemma 2.11 we know $\sigma = \sigma_1 + \sigma_2$.

The fact $\sigma_2 \in S^{-\infty}$ can be seen when using the asymptotics in Theorem 5.6, namely,

$$\sigma_2(x, \xi) = \sum_{|\alpha| \leq N} \frac{1}{\alpha!} D_y^\alpha \partial_\eta^\alpha ((1 - \chi(|x-y|^2)) \sigma(x, \eta))|_{(y, \eta) = (x, \xi)} + S^{m-N-1} = S^{m-N-1}$$

holds for $\forall N \in \mathbb{N}$, so $\sigma_2 \in S^{-\infty}$.

From (2.10) we can see

$$\begin{cases} K_1(x, y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \chi(|x-y|^2) \sigma(x, \xi) d\xi, \\ K_2(x, y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} (1 - \chi(|x-y|^2)) \sigma(x, \xi) d\xi. \end{cases}$$

which implies T_1 is properly supported. And the requirements for the $\text{supp } K_1$ and $\text{supp } K_2$ can be seen from the expression above. The proof is complete. \square

2.4. Pseudolocal property

We talk about singular support and pseudolocal property.

DEFINITION 2.23 (Singular support). For a distribution $u \in \mathcal{D}'$, we define its *singular support* to be the complement of the set $\bigcap \{O \subset \mathbb{R}^n; O \text{ is open and } A \subset O\}$ where

$$A = \{x \in \mathbb{R}^n; u \text{ is } C^\infty \text{ at } x\}.$$

We denote the singular support of a distribution $u \in \mathcal{D}'$ as $\text{sing supp } u$.

It is obvious that $\text{sing supp } u$ is as closed set and

$$\text{sing supp } u \subset \text{supp } u.$$

We know a differential operator doesn't increase the support of a distribution, but this is not true for a Ψ DO. More specifically, if a distribution u is supported in Ω , then Tu might not be supported in a domain Ω anymore. Instead, Ψ DOs have another property, called pseudolocal property, which means Ψ DOs don't increase the singular support of a distribution.

THEOREM 2.24 (Pseudolocal property). Assume T is a Ψ DO, then

$$\boxed{\text{sing supp}(Tu) \subset \text{sing supp } u.}$$

PROOF. Assume $x_0 \notin \text{sing supp } u$. Because $\text{sing supp } u$ is closed, we can find $\epsilon > 0$ such that u is C^∞ in $B(x_0, \epsilon)$. According to Lemma 2.22, we can divided T into T_1 and T_2 such that T_2 is C^∞ -smooth and the kernel K_1 of T_1 satisfies

$$\text{supp } K_1 \subset \{(x, y) \in \mathbb{R}^{2n}; |x - y| \leq \epsilon/4\}.$$

Hence, for $\forall x \in B(x_0, \epsilon/4)$, we have $K(x, y) = 0$ when $|y - x_0| \geq \epsilon/2$.

Fix a function $\chi \in C_c^\infty$ such that $\chi(y) = 1$ when $|y - x_0| \leq \epsilon/2$ and $\text{supp } \chi = B(x_0, \epsilon)$, then

$$\forall x \in B(x_0, \epsilon/4), \quad T_1 u(x) = \int K(x, y)u(y) dy = \int K(x, y)\chi(y)u(y) dy = T_1(\chi u)(x).$$

Note that $\chi u \in C_c^\infty \subset \mathcal{S}$ due to the fact that u is C^∞ in $\text{supp } \chi$, so $T_1(\chi u) \in \mathcal{S}$. Because $T_1 u = T_1(\chi u) \in \mathcal{S}$ on $B(x_0, \epsilon/4)$, we conclude that $T_1 u \in C^\infty(B(x_0, \epsilon/4))$. Also, $T_2 u \in C^\infty$ because T_2 is C^∞ -smooth. In total, Tu is C^∞ -smooth in a small neighborhood of x_0 , so $x_0 \notin \text{sing supp}(Tu)$.

We obtain $(\text{sing supp } u)^c \subset (\text{sing supp}(Tu))^c$. The proof is complete. \square

Exercise

EXERCISE 2.1. Prove $\mathcal{S}(\mathbb{R}^n) \subset S^{-\infty}$, namely, $\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$, $\varphi(\xi) \in S^{-\infty}$.

EXERCISE 2.2. Prove Lemma 2.3.

EXERCISE 2.3. Prove Lemma 2.14. See [26, Prop. 6.7] for reference.

Oscillatory integrals

In §2.1 we encountered the notion of kernel of a Ψ DO,

$$K(x, y) := (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) \, d\xi.$$

which might not be integral in the Lebesgue sense (e.g. when $\sigma(x, \xi) = 1$). However, if we look back to the original definition of a Ψ DO,

$$(T_\sigma\varphi)(x) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{ix\cdot\xi} \sigma(x, \xi) \hat{\varphi}(\xi) \, d\xi,$$

the integral above is always well-defined in the Lebesgue sense, because $\hat{\varphi}$ is rapidly decaying. Specifically, for any $m \in \mathbb{R}$ and any $\sigma \in S^m$, we have

$$|(T_\sigma\varphi)(x)| \lesssim \int |\sigma(x, \xi) \hat{\varphi}(\xi)| \, d\xi \lesssim \int \langle \xi \rangle^m \langle \xi \rangle^{-m-n-1} \, d\xi < +\infty.$$

The problems emerges when we expand the Fourier transform $\hat{\varphi}$ (by a variable y) and **exchange** the integration order of y and ξ :

$$\begin{aligned} (T_\sigma\varphi)(x) &= (2\pi)^{-n} \int \left(\int e^{i(x-y)\cdot\xi} \sigma(x, \xi) \varphi(y) \, dy \right) \, d\xi, \\ \int_{\mathbb{R}^n} K(x, y) \varphi(y) \, dy &= (2\pi)^{-n} \int \left(\int e^{i(x-y)\cdot\xi} \sigma(x, \xi) \, d\xi \right) \varphi(y) \, dy. \end{aligned} \tag{3.1}$$

According to Fubini's theorem, this exchange is valid only when all of the integrals involved are absolutely integrable. From time to time we will encounter integrals of the form (3.1), but also more general than that. A rigorous framework is appealing for making the integrals of these type always well-defined.

3.1. Oscillatory integrals - Type I

Generally speaking, for any $u \in \mathcal{S}(\mathbb{R}^n)$ and $\sigma \in S^m(\mathbb{R}_x^n \times R_\xi^N)$, one is interested in the following integral

$$I(u) := \int e^{i\varphi(x, \xi)} \sigma(x, \xi) u(x) \, dx \, d\xi \tag{3.2}$$

where φ is a phase function defined as follows.

DEFINITION 3.1 (Phase function). Function φ is called a *phase function (of order μ)* if it satisfies

- (1) $\varphi \in C^\infty(\mathbb{R}_x^n \times (\mathbb{R}_\xi^N \setminus \{0\}); \mathbb{R})$ is real-valued;
- (2) φ is homogeneous w.r.t. ξ of order $\mu > 0$, i.e. $\varphi(x, t\xi) = t^\mu \varphi(x, \xi)$;
- (3) $|\nabla_{(x, \xi)} \varphi(x, \xi)| \neq 0$ for $\forall (x, \xi) \in \mathbb{R}_x^n \times (\mathbb{R}_\xi^N \setminus \{0\})$.

There are different ways to define the notion of phase functions, see [13, §7.8], and we don't pursue diversity here. Note that n might not equal N , and most of the results in §2.1 holds also for the case $n \neq N$. Here we consider phases of order μ , instead of just

order 1, because in §5 and §7 we do encounter phases of order 2. The condition $\mu > 0$ is indispensable.

In contrast with (3.1), in (3.2) it is not sure that integrating first w.r.t x (or ξ) can guarantee it's integrable. Instead, we study

$$\lim_{\epsilon \rightarrow 0^+} I_\epsilon(u) := \lim_{\epsilon \rightarrow 0^+} \int e^{i\varphi(x,\xi)} \sigma(x,\xi) \chi(\epsilon\xi) u(x) dx d\xi, \quad (3.3)$$

where χ is a function in $C_c^\infty(\mathbb{R}^n)$ with $\chi(0) = 1$. We show that the limit (3.3) exists and its value is independent of the choice of χ .

THEOREM 3.2. *Assume $m \in \mathbb{R}$, $\sigma \in S^m(\mathbb{R}_x^n \times \mathbb{R}_\xi^N)$ and φ is a phase function of order μ . Fix a function $\chi \in C_c^\infty(\mathbb{R}^n)$ with $\chi(0) = 1$. Assume either $u \in C_c^\infty(\mathbb{R}^n)$, or $u \in \mathcal{S}(\mathbb{R}^n)$ and $\partial_x^\alpha \varphi(x,\xi)$ is tempered w.r.t. x for any α . Then the limit (3.3) exists and its value is independent of the choice of χ , and it equals to*

$$\int e^{i\varphi(x,\xi)} L^T(\sigma(x,\xi)u(x)) dx d\xi,$$

when integer $T > (m + N)/\mu$ where L is given in Lemma 3.4.

The proof Theorem 3.2, we first do some preparation.

LEMMA 3.3. *Assume $\chi \in C_c^\infty(\mathbb{R}^n)$ and let $\epsilon \in \mathbb{R}$. There exists a constant C independent of ϵ such that*

$$\boxed{|\partial_\xi^\alpha(\chi(\epsilon\xi))| \leq C \langle \xi \rangle^{-|\alpha|}.}$$

PROOF. Because $\chi \in C_c^\infty(\mathbb{R}^n)$, there exists a fixed constant C such that $\chi(\epsilon\xi) \equiv 0$ when $|\epsilon\langle \xi \rangle| \geq C$. When $|\epsilon\langle \xi \rangle| \geq C$, $\chi(\epsilon\xi) = 0$ so $\partial_\xi^\alpha(\chi(\epsilon\xi)) = 0$; when $|\epsilon\langle \xi \rangle| \leq C$, we have

$$|\partial_\xi^\alpha(\chi(\epsilon\xi))| = |\epsilon^{|\alpha|} \partial_\xi^\alpha \chi(\epsilon\xi)| \leq (C \langle \xi \rangle)^{-|\alpha|} \sup_{\mathbb{R}^n} |\partial_\xi^\alpha \chi| \simeq C \langle \xi \rangle^{-|\alpha|}.$$

We arrive at the conclusion. \square

LEMMA 3.4. *Assume φ is a phase function φ of order μ , and $\partial_x^\alpha \varphi(x,\xi)$ is tempered w.r.t. x for any α . Then there exists a first order linear differential operator*

$$L = a_j(x,\xi) \partial_{x_j} + b_j(x,\xi) \partial_{\xi_j} + c(x,\xi)$$

such that ${}^t L(e^{i\varphi(x,\xi)}) = e^{i\varphi(x,\xi)}$, and for any fixed x_0 , $a_j(x_0, \cdot) \in S^{-\mu}$, $b_j(x_0, \cdot) \in S^{1-\mu}$, $c(x_0, \cdot) \in S^{-\mu}$, and a_j, b_j, c are tempered functions of x variable.

Here $\langle {}^t L f, g \rangle := \langle f, L g \rangle$, where the integral is w.r.t. (x,ξ) , and $f, g \in C_0^\infty$. ${}^t L$ is call the transpose of L , e.g. ${}^t(\nabla_\xi) = -\nabla_\xi$.

PROOF OF LEMMA 3.4. We write $\nabla_x \varphi = \varphi_x$ and $\nabla_\xi \varphi = \varphi_\xi$ for short. Fix a $\chi \in C_c^\infty(\mathbb{R}^N)$ with $\chi \equiv 1$ in a neighborhood of 0. Construct

$$M := (1 - \chi(\xi)) \frac{\varphi_x \cdot D_x + \langle \xi \rangle^2 \varphi_\xi \cdot D_\xi}{|\varphi_x|^2 + \langle \xi \rangle^2 |\varphi_\xi|^2} + \chi(\xi).$$

We mention several facts about M :

- First, M is well-defined. Note that the denominator $|\varphi_x|^2 + \langle \xi \rangle^2 |\varphi_\xi|^2 \geq |\varphi_x|^2 + |\varphi_\xi|^2 \neq 0$ when $(x,\xi) \in \mathbb{R}_x^n \times (\mathbb{R}_\xi^N \setminus \{0\})$, and the point $\xi = 0$ has been cutoff by $1 - \chi$, so M is always well-defined;
- Second, away from $\xi = 0$, $\varphi_x(x_0, \cdot) \in S^\mu$, $\varphi_\xi(x_0, \cdot) \in S^{\mu-1}$, the denominator $(x_0, \cdot) \in S^{2\mu}$;
- Third, $M e^{i\varphi(x,\xi)} = (1 - \chi) e^{i\varphi(x,\xi)} + \chi e^{i\varphi(x,\xi)} = e^{i\varphi(x,\xi)}$.

The transpose of M is the desired operator. Indeed, it can be checked that, when x is fixed,

$${}^tM = (1 - \chi)S^{-\mu}\partial_x + (1 - \chi)S^{1-\mu}\partial_\xi + (1 - \chi)S^{-\mu} + \chi(\xi). \quad (3.4)$$

The proof is complete. \square

REMARK 3.5. Here $S^{-\mu}\partial_x$ is a shorthand of $a \in C^\infty$ such that $a_j\partial_{x_j}$ for some $a(x_0, \cdot) \in S^{-\mu}$. Readers should note that (3.4) is somehow misleading because the coefficients might not be bounded w.r.t. x , e.g. ${}^tM = x_1\partial_{x_1}$. However, they must be tempered, and these tempered growth will be balanced by the rapid decay of u , The notations in (3.4) wouldn't hurt.

PROOF OF THEOREM 3.2. Choose L according to Lemma 3.4, then

$$\begin{aligned} I_\epsilon(u) &= \int ({}^tL)^T(e^{i\varphi(x,\xi)}) \sigma(x, \xi) \chi(\epsilon\xi) u(x) dx d\xi \\ &= \int e^{i\varphi(x,\xi)} L^T(\sigma(x, \xi) \chi(\epsilon\xi) u(x)) dx d\xi. \end{aligned} \quad (3.5)$$

Readers should note that the transpose of L is realized by the **classical integration by parts** (nothing fancy here), and it is the presence of $\chi(\epsilon\xi)$ that cancels the boundary terms and makes the integration by parts applicable.

The conditions “ $a_j \in S^{-\mu}$, $b_j \in S^{1-\mu}$, $c \in S^{-\mu}$ ” in Lemma 3.4 give us

$$\begin{aligned} &L^T(\sigma(x, \xi) \chi(\epsilon\xi) u(x)) \\ &= ((a_j(x, \xi)\partial_{x_j} + b_j(x, \xi)\partial_{\xi_j} + c(x, \xi))^T(\sigma(x, \xi) \chi(\epsilon\xi) u(x)) \\ &= \sum_{|\alpha+\beta|\leq T} S^{-\mu|\alpha|+(1-\mu)|\beta|-\mu(T-|\alpha|-\beta)} \partial_x^\alpha \partial_\xi^\beta (\sigma(x, \xi) \chi(\epsilon\xi) u(x)) \\ &= \sum_{|\alpha+\beta|\leq T} S^{|\beta|-\mu T} \partial_x^\alpha \partial_\xi^\beta (\sigma(x, \xi) \chi(\epsilon\xi) u(x)) \\ &= \sum_{|\alpha+\beta|\leq T} \sum_{\beta'+\beta''=\beta} C_{\beta',\beta''} S^{|\beta|-\mu T} \partial_x^\alpha (\partial_\xi^{\beta'} \sigma(x, \xi) \partial_\xi^{\beta''} [\chi(\epsilon\xi)] u(x)) \\ &= \sum_{|\alpha+\beta|\leq T} \sum_{\beta'+\beta''=\beta} C S^{|\beta|-\mu T} \partial_x^\alpha (\partial_\xi^{\beta'} \sigma(x, \xi) \partial_\xi^{\beta''} [\chi(\epsilon\xi)] u(x)) \\ &= \sum_{|\alpha+\beta|\leq T} \sum_{\beta'+\beta''=\beta} \sum_{\alpha'+\alpha''=\alpha} C S^{|\beta|-\mu T} \partial_x^{\alpha'} \partial_\xi^{\beta'} \sigma(x, \xi) \partial_\xi^{\beta''} [\chi(\epsilon\xi)] \partial_x^{\alpha''} u(x). \end{aligned}$$

The term $\partial_\xi^{\beta''} [\chi(\epsilon\xi)]$ is the only term that depends on ϵ . Hence, by Lemma 3.3 we can have

$$\begin{aligned} &|L^T(\sigma(x, \xi) \chi(\epsilon\xi) u(x))| \\ &\leq \sum_{|\alpha+\beta|\leq T} \sum_{\beta'+\beta''=\beta} \sum_{\alpha'+\alpha''=\alpha} C \langle \xi \rangle^{|\beta|-\mu T} C \langle \xi \rangle^{m-|\beta'|} C \langle \xi \rangle^{-|\beta''|} \mathcal{S}(\mathbb{R}_x^n) \\ &\leq C \langle \xi \rangle^{m-\mu T} \mathcal{S}(\mathbb{R}_x^n), \end{aligned}$$

where the constant C is independent of ϵ . Then T is chosen to be larger than $(m + N)/\mu$, the integrand in (3.5) is bounded by a absolutely integral function. Therefore, according to LDCT, the limit $\lim_{\epsilon \rightarrow 0^+} I_\epsilon(u)$ exists. Readers may think where we used the condition $\mu > 0$.

We also show that the limit is independent of χ . Fix a $T > (m + N)/\mu$, then by (3.5) we have

$$\begin{aligned} \lim_{\epsilon \rightarrow 0^+} I_\epsilon(u) &= \lim_{\epsilon \rightarrow 0^+} \int e^{i\varphi(x,\xi)} L^T(\sigma(x,\xi)\chi(\epsilon\xi)u(x)) \, dx \, d\xi \\ &= \int e^{i\varphi(x,\xi)} \lim_{\epsilon \rightarrow 0^+} L^T(\sigma(x,\xi)\chi(\epsilon\xi)u(x)) \, dx \, d\xi \quad (\text{thanks to LDCT}) \\ &= \int e^{i\varphi(x,\xi)} L^T(\sigma(x,\xi)u(x)) \, dx \, d\xi, \end{aligned} \quad (3.6)$$

which implies $\lim_{\epsilon \rightarrow 0^+} I_\epsilon(u)$ is independent of χ . The proof is complete. \square

Readers may think about if the framework can be generalized to symbols in $S_{\rho,\delta}^m$.

Now let's summarize the definition of oscillatory integrals.

DEFINITION 3.6 (Oscillatory integral). For any $m \in \mathbb{R}$, any $\sigma \in S^m(\mathbb{R}_x^n \times \mathbb{R}_\xi^N)$, and any phase function φ of order μ , and either $u \in C_c^\infty(\mathbb{R}^n)$, or $u \in \mathcal{S}(\mathbb{R}^n)$ and $\partial_x^\alpha \varphi(x,\xi)$ is tempered w.r.t. x for any α , the integral

$$I(u) = \int e^{i\varphi(x,\xi)} \sigma(x,\xi) u(x) \, dx \, d\xi$$

is defined as

$$I(u) := \lim_{\epsilon \rightarrow 0^+} \int e^{i\varphi(x,\xi)} \sigma(x,\xi) \chi(\epsilon\xi) u(x) \, dx \, d\xi, \quad (3.7)$$

where the result is independent of χ , as long as $\chi \in C_c^\infty$ and $\chi(0) = 1$. The limit equals

$$I(u) = \int e^{i\varphi(x,\xi)} L^T(\sigma(x,\xi)u(x)) \, dx \, d\xi$$

when integer $T > (m + N)/\mu$, where L is given in Lemma 3.4.

In many cases we will meet oscillatory integrals involving parameters.

LEMMA 3.7. For any $\sigma \in S^m(\mathbb{R}_x^{n_1} \times \mathbb{R}_y^{n_2} \times \mathbb{R}_\xi^N)$, and any phase function φ of order μ , and for either $u \in C_c^\infty(\mathbb{R}_x^{n_1} \times \mathbb{R}_y^{n_2})$, or $u \in \mathcal{S}(\mathbb{R}_x^{n_1} \times \mathbb{R}_y^{n_2})$ and $\partial_{(x,y)}^\alpha \varphi(x,\xi)$ is tempered w.r.t. (x,y) for any α , the integral

$$I(u)(y) := \int e^{i\varphi(x,y,\xi)} \sigma(x,y,\xi) u(x,y) \, dx \, d\xi \quad (3.8)$$

is a well-defined oscillatory integral, and $I: \mathcal{S}(\mathbb{R}^{n_1}) \rightarrow \mathcal{S}(\mathbb{R}^{n_2})$ bounded. Moreover, we have

$$\begin{aligned} \frac{\partial}{\partial y} (I(u)(y)) &= \int \frac{\partial}{\partial y} (e^{i\varphi(x,y,\xi)} \sigma(x,y,\xi) u(x,y)) \, dx \, d\xi, \\ \int I(u)(y) \, dy &= \int e^{i\varphi(x,y,\xi)} \sigma(x,y,\xi) u(x,y) \, dx \, dy \, d\xi. \end{aligned}$$

We omit the proof. The take-home message of Lemma 3.7 is that oscillatory integrals can have parameters, and there are much freedom to put operations w.r.t. y inside the integration $I(u)(y)$.

Now we go back to Ψ DOs and its kernel. We have intuitively claimed that the kernel of T_σ is of the form

$$K(x,y) = \int e^{i(x-y)\cdot\xi} \sigma(x,\xi) \, d\xi.$$

Note that this object has variables (x, y) , so a proper candidate of test functions should be $w(x, y) \in \mathcal{S}$. We choose $w(x, y) = u(y)v(x)$ where $u, v \in \mathcal{S}$, then formally we should have

$$\langle K, v \otimes u \rangle \simeq \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) u(y) v(x) dx dy d\xi \simeq \langle T_\sigma u, v \rangle.$$

The integral above is exactly an example of Lemma 3.7, so it is a well-defined oscillatory integral. Now the kernel of a Ψ DO can be defined.

DEFINITION 3.8 (Kernel). Assume $m \in \mathbb{R}$ and $\sigma \in S^m$, and T_σ is the corresponding Ψ DO. The *kernel* of T_σ is defined as a map:

$$K_\sigma : w \in \mathcal{S}(\mathbb{R}^{2n}) \mapsto \langle K_\sigma, w \rangle := (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) w(x, y) dx dy d\xi \in \mathbb{C}.$$

When $x \neq y$, we write $K_\sigma(x, y)$ as

$$K_\sigma(x, y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) d\xi. \quad (3.9)$$

The well-definedness of Definition 3.8 is guaranteed by Lemma 3.7.

REMARK 3.9. From (3.9), we know that when $m < -N$, K_σ is a well-defined bounded function for any (x, y) because

$$|K_\sigma| \lesssim \int |\sigma(x, \xi)| d\xi \lesssim \int \langle \xi \rangle^m d\xi \leq C < +\infty.$$

This implies when the order of σ is small enough, there should hold some type of boundedness for T_σ , and we will cover this in §6. However when $m \geq -N$, only when $x \neq y$ the kernel K_σ can be expressed as (3.9).

LEMMA 3.10. *Under the assumption of Definition 3.8, we have*

$$\langle K_\sigma, u(y)v(x) \rangle = \langle T_\sigma u, v \rangle.$$

We omit the proof.

LEMMA 3.11. *Under the assumption of Definition 3.8, when $x \neq y$, K_σ is C^∞ smooth. Moreover, for T large enough, we have*

$$|K_\sigma(x, y)| \leq C_T |x - y|^{-T}, \quad |x - y| \geq 1.$$

PROOF. For any fixed point (x, y) with $x \neq y$, we show that K_σ is C^∞ at this point. Fix a function $\chi \in C^\infty$ satisfying $\chi \equiv 0$ in a small neighborhood U of the diagonal $\{x = y\}$ and $\chi \equiv 1$ in the interior of the complement of U . We can shrink U such that for any $x \neq y$, $\chi(x, y) = 1$. For any $w \in \mathcal{S}$, we have

$$\langle \partial_x^\alpha (\chi K_\sigma), w \rangle \simeq \int e^{i(x-y)\cdot\xi} \xi^\alpha \sigma(x, \xi) w(x, y) dx dy d\xi.$$

Apply the operator L to $e^{i(x-y)\cdot\xi}$ and integrate by parts, we obtain

$$\langle \partial_x^\alpha \chi K_\sigma, w \rangle = \langle (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \left(\frac{-(x-y) \cdot D_\xi}{|x-y|^2} \right)^T (\xi^\alpha \sigma(x, \xi)) d\xi, w \chi \rangle,$$

which implies

$$\partial_x^\alpha (\chi K_\sigma)(x, y) \simeq \int e^{i(x-y)\cdot\xi} \left(\frac{-(x-y) \cdot D_\xi}{|x-y|^2} \right)^T (\xi^\alpha \sigma(x, \xi)) d\xi.$$

It can be checked that when T is large enough, the integral above will be absolutely integrable, and is of the order $|x-y|^{-T}$ for T large enough. Therefore, $K_\sigma \in C^\infty(\mathbb{R}^{2n} \setminus \{x = y\})$, and K satisfies the desired decay. \square

By Lemma 3.11, we see that $K(x, y)$ behave nicely when off the diagonal, thanks to the notion of oscillatory integrals. However, on the diagonal, $K(x, y)$ might still be ill-defined. See Example 3.12.

EXAMPLE 3.12. The kernel corresponding to the identity operator (symbol = 1) is the distribution $\delta(x - y)$. This is because

$$\begin{aligned}\langle K, v \otimes u \rangle &= \langle T_1 u, v \rangle = \langle u, v \rangle = \int u(x)v(x) dx \\ &= \int \delta(x - y)u(y)v(x) dx dy \\ &= \langle \delta(x - y), v \otimes u \rangle.\end{aligned}$$

which gives $K(x, y) = \delta(x - y)$.

EXAMPLE 3.13. The kernel corresponding to D_1 (symbol = ξ_1) is $D_1\delta(x - y)$. This is because

$$\begin{aligned}\langle K, v \otimes u \rangle &= \langle D_1 u, v \rangle = \int D_1 u(x)v(x) dx = \int \delta(x - y)D_1 u(y)v(x) dx dy \\ &= - \int D_{y_1}(\delta(x - y))u(y)v(x) dx dy = \int D_1 \delta(x - y)u(y)v(x) dx dy \\ &= \langle D_1 \delta(x - y), v \otimes u \rangle.\end{aligned}$$

which gives $K(x, y) = D_1\delta(x - y)$. Besides, readers may also have tried another way to compute the kernel and get a zero result: when $x \neq y$

$$\begin{aligned}K(x, y) &= (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \xi_1 d\xi = (2\pi)^{-n} \int \left(\frac{(x-y)\cdot D\xi}{|x-y|^2}\right)^2 (e^{i(x-y)\cdot\xi}) \xi_1 d\xi \\ &= (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \left(\frac{-(x-y)\cdot D\xi}{|x-y|^2}\right)^2 (\xi_1) d\xi = 0.\end{aligned}$$

This result is technically correct (because $D_1\delta(x - y) = 0$ when $x \neq y$), but is not complete: it cannot speak about the behavior of K on the diagonal. This example told us, none of the methods is the best one to get most accurate expression for a kernel, sometimes we need to do complicated and delicate computations.

3.2. Oscillatory integrals - Type II

We know the Fourier transform of a constant function is the δ function and so the inverse Fourier transform of the δ function should be the constant, namely,

$$\int e^{i(x-y)\cdot\xi} dx d\xi \simeq 1. \quad (3.10)$$

This integral can be regarded as the $I(u)$ defined in Lemma 3.7 where the symbol and the u are both constant 1. However, this is not covered by Lemma 3.7 because $1 \notin \mathcal{S}$. The map I in Lemma 3.7 is defined on \mathcal{S} . Now by using duality arguments we shall generalize it from \mathcal{S} to $S^{+\infty}$.

Let $\sigma \in S^m$ and $u \in \mathcal{S}$, so $T_\sigma u \in \mathcal{S}$. Let $f \in S^{+\infty}$, then f is a smooth tempered function, so $\langle T_\sigma u, f \rangle$ is meaningful and

$$\langle T_\sigma u, f \rangle = \lim_{\epsilon \rightarrow 0^+} \langle T_\sigma u, \chi(\epsilon \cdot) f \rangle$$

holds for any $\chi \in C_c^\infty$. Expanding the integral, we have

$$\langle T_\sigma u, f \rangle = \lim_{\epsilon \rightarrow 0^+} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) u(y) f(x) \chi(\epsilon x) dx dy d\xi$$

$$\begin{aligned}
&= \lim_{\epsilon \rightarrow 0^+} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) \chi(\epsilon\xi) u(y) f(x) \chi(\epsilon x) dx dy d\xi \\
&= \lim_{\epsilon \rightarrow 0^+} \int u(y) \left(\int e^{i(x-y)\cdot\xi} \sigma(x, \xi) f(x) \chi(\epsilon x) \chi(\epsilon\xi) dx d\xi \right) dy.
\end{aligned}$$

This inspired us to define (3.10) as

$$\lim_{\epsilon \rightarrow 0^+} \int e^{i(x-y)\cdot\xi} \chi(\epsilon x) \chi(\epsilon\xi) dx d\xi.$$

More generally, we can generalize Definition 3.6 as follows.

DEFINITION 3.14 (Oscillatory integral). For any $m \in \mathbb{R}$ and $\rho > 0$, any $\sigma \in S_\rho^m(\mathbb{R}^N)$, and any phase function $\varphi \in C^\infty(\mathbb{R}^N \setminus \{0\})$ of order μ (real-valued, $\varphi(\theta) = t^\mu \varphi(\theta)$, and $\nabla_\theta \varphi(\theta) \neq 0$ when $\theta \neq 0$) satisfying

$$\boxed{\rho + \mu > 1}, \quad (3.11)$$

the integral

$$I = \int e^{i\varphi(\theta)} \sigma(\theta) d\theta$$

is defined as

$$\boxed{I := \lim_{\epsilon \rightarrow 0^+} \int e^{i\varphi(\theta)} \sigma(\theta) \chi(\epsilon\theta) d\theta}, \quad (3.12)$$

where the result is independent of χ , as long as $\chi \in C_c^\infty$ and $\chi(0) = 1$. The limit equals

$$\boxed{I = \int e^{i\varphi(\theta)} L^T(\sigma(\theta)) d\theta} \quad (3.13)$$

when $T > (m + N)/(\rho + \mu - 1)$, where L is given in Lemma 3.16 below.

REMARK 3.15. The space $S_\rho^m(\mathbb{R}^N)$ is defined as

$$S_\rho^m(\mathbb{R}^N) := \{\varphi \in C^\infty(\mathbb{R}^N); |\partial^\alpha \varphi(\theta)| \lesssim \langle \theta \rangle^{m-\rho|\alpha|}\}.$$

See [1, §I.8.1] for more details. Also, the formula (3.9) in the definition of the kernel is meaningful now.

LEMMA 3.16. *Under the condition in Definition 3.14, there exists a first order linear differential operator*

$$L = b_j(\theta) \partial_{\theta_j} + c(\theta)$$

such that ${}^tL(e^{i\varphi(\theta)}) = e^{i\varphi(\theta)}$, $0 \notin \text{supp } b_j$, and $b_j \in S^{1-\mu}$, $c \in S^{-\mu}$, and in a small neighborhood of $\theta = 0$ there holds $b \equiv 0$ and $c \equiv 1$.

PROOF. We write $\nabla_\theta \varphi = \varphi_\theta$ for short. Fix a $\chi \in C_c^\infty(\mathbb{R}^n)$ with $\chi \equiv 1$ in a neighborhood of 0. Construct

$$M := (1 - \chi(\theta)) \frac{\varphi_\theta \cdot D_\theta}{|\varphi_\theta|^2} + \chi(\theta).$$

We mention several facts about M :

- First, M is well-defined. Note that the denominator $|\varphi_\theta|^2 \neq 0$ when $\theta \in \mathbb{R}^N \setminus \{0\}$, and the point $\theta = 0$ has been cutoff by $1 - \chi$, so M is always well-defined;
- Second, away from $\theta = 0$, $\varphi_\theta \in S^{\mu-1}$;
- Third, $M e^{i\varphi(\theta)} = (1 - \chi) e^{i\varphi(\theta)} + \chi e^{i\varphi(\theta)} = e^{i\varphi(\theta)}$.

The transpose of M is the desired operator. Indeed, it can be checked that, when x is fixed,

$${}^tM = (1 - \chi) S^{1-\mu} \partial_\theta + (1 - \chi) S^{-\mu} + \chi(\theta).$$

The proof is complete. \square

Readers may compare Lemma 3.16 with Lemma 3.4.

LEMMA 3.17. *Assume L is chosen as in Lemma 3.16, and $\sigma \in S_\rho^m(\mathbb{R}^N)$, then*

$$|L^T(\sigma(\theta))| \lesssim \langle \theta \rangle^{m-(\rho+\mu-1)T}.$$

The proof is left as an exercise. Combining Lemmas 3.16 and 3.17, we obtain a result similar to Theorem 3.2.

THEOREM 3.18. *Assume $m \in \mathbb{R}$ and $\rho > 0$, $\sigma \in S_\rho^m(\mathbb{R}^N)$ and $\varphi \in C^\infty(\mathbb{R}^N \setminus \{0\})$ is a phase function of order μ . Fix a function $\chi \in C_c^\infty(\mathbb{R}^n)$ with $\chi(0) = 1$. Then the limit (3.12) exists and its value is independent of the choice of χ , and it equals to*

$$\int e^{i\varphi(\theta)} L^T(\sigma(\theta)) \, d\theta,$$

when $T > (m + N)/(\rho + \mu - 1)$ where T is given in Lemma 3.16.

The proof is similar to that of Theorem 3.2. The generalized definition of the oscillatory integral can handle more cases. One of the examples is as follows.

LEMMA 3.19. *The following equality holds in oscillatory sense,*

$$\int_{\mathbb{R}^{2n}} e^{\pm ix \cdot \xi} \, dx \, d\xi = (2\pi)^n.$$

PROOF. We shall regard (x, ξ) as the θ in Definition 3.14, then this phase function $x \cdot \xi$ is of order 2. One can also check that $\nabla_{(x, \xi)}(x \cdot \xi) \neq 0$ when $(x, \xi) \neq 0$. We choose the cutoff function as $\chi(\epsilon x)\chi(\epsilon \xi)$ where $\chi \in C_c^\infty$ with $\chi(0) = 1$, then

$$\begin{aligned} \int e^{ix \cdot \xi} \, dx \, d\xi &:= \lim_{\epsilon \rightarrow 0^+} \int e^{ix \cdot \xi} \chi(\epsilon x) \chi(\epsilon \xi) \, dx \, d\xi = \lim_{\epsilon \rightarrow 0^+} \int \left(\int e^{ix \cdot \xi} \chi(x) \, dx \right) \cdot \chi(\epsilon^2 \xi) \, d\xi \\ &= (2\pi)^{n/2} \lim_{\epsilon \rightarrow 0^+} \int \widehat{\chi}(-\xi) \chi(\epsilon^2 \xi) \, d\xi = (2\pi)^{n/2} \int \widehat{\chi}(-\xi) \, d\xi \quad (\text{LDCT}) \\ &= (2\pi)^n \chi(0) = (2\pi)^n. \end{aligned}$$

The case for $e^{-ix \cdot \xi}$ is similar. Note that all of the integrals above are usual integral besides the first one on the LHS. \square

The following result will be useful.

LEMMA 3.20. *The following equality holds in oscillatory sense,*

$$\int_{\mathbb{R}^{2n}} e^{\pm ix \cdot \xi} x^\alpha \xi^\beta \, dx \, d\xi = (\pm i)^{|\alpha|} (2\pi)^n \alpha! \delta^{\alpha\beta}.$$

PROOF. We have

$$\begin{aligned} \int e^{\pm ix \cdot \xi} x^\alpha \xi^\beta \, dx \, d\xi &= \int (\pm D_\xi)^\alpha (e^{\pm ix \cdot \xi}) \xi^\beta \, dx \, d\xi \\ &= \int e^{\pm ix \cdot \xi} (\mp D_\xi)^\alpha (\xi^\beta) \, dx \, d\xi = (\pm i)^{|\alpha|} \int e^{\pm ix \cdot \xi} \partial_\xi^\alpha (\xi^\beta) \, dx \, d\xi. \end{aligned}$$

It can be check that $\partial_\xi^\alpha (\xi^\beta) = \beta! / (\beta - \alpha)! \xi^{\beta - \alpha}$ when $\alpha \leq \beta$, and $= 0$ otherwise. Hence, when $\alpha \leq \beta$ we have

$$\int e^{\pm ix \cdot \xi} x^\alpha \xi^\beta \, dx \, d\xi = (\pm i)^{|\alpha|} \beta! / (\beta - \alpha)! \int e^{\pm ix \cdot \xi} \xi^{\beta - \alpha} \, dx \, d\xi. \quad (3.14)$$

When $\alpha \neq \beta$, we can continue

$$\int e^{\pm ix \cdot \xi} \xi^{\beta - \alpha} \, dx \, d\xi \simeq \int D_x^{\beta - \alpha} (e^{\pm ix \cdot \xi}) \, dx \, d\xi = 0 \simeq \int e^{\pm ix \cdot \xi} D_x^{\beta - \alpha} (1) \, dx \, d\xi = 0.$$

Therefore, when $\alpha \neq \beta$, we have

$$\int_{\mathbb{R}^{2n}} e^{\pm ix \cdot \xi} x^\alpha \xi^\beta dx d\xi = 0. \quad (3.15)$$

When $\alpha = \beta$, by Lemma 3.19 we can continue (3.14) as follows,

$$\int_{\mathbb{R}^{2n}} e^{\pm ix \cdot \xi} x^\alpha \xi^\beta dx d\xi = (\pm i)^{|\alpha|} \alpha! \int e^{\pm ix \cdot \xi} dx d\xi = (\pm i)^{|\alpha|} \alpha! (2\pi)^n.$$

The proof is complete. \square

REMARK 3.21. The space $S_\rho^m(\mathbb{R}^N)$ is defined as

$$S_\rho^m(\mathbb{R}^N) := \{\varphi \in C^\infty(\mathbb{R}^N); |\partial^\alpha \varphi(\theta)| \lesssim \langle \theta \rangle^{m-\rho|\alpha|}\}.$$

See [1, §I.8.1] for more details. Also, the formula (3.9) in the definition of the kernel is meaningful now.

We can generalize Lemma 3.19

LEMMA 3.22. Assume $m \in \mathbb{R}$ and $f \in S_\rho^m(\mathbb{R}^n)$ (see Remark 3.15) with $\rho + 2 > 1$. The following equality holds in oscillatory sense,

$$\int_{\mathbb{R}^{2n}} e^{\pm ix \cdot \xi} f(x) dx d\xi = (2\pi)^n f(0).$$

REMARK 3.23. Lemma 3.22 indicates the the “inverse Fourier transform” is indeed the inverse of “Fourier transform”.

PROOF. We only show the case $+ix \cdot \xi$. The condition on f guarantees the integral is well-defined, see Definition 3.14. By Taylor’s expansion we have

$$f(x) = f(0) + \sum_{j=1}^n x_j g_j(x), \quad \text{where } g_j(x) := \int_0^1 \partial_{x_j} f(tx), dt.$$

so

$$\begin{aligned} \int e^{ix \cdot \xi} f(x) dx d\xi &= \int e^{ix \cdot \xi} [f(0) + \sum_{j=1}^n x_j g_j(x)] dx d\xi \\ &= f(0) \int e^{ix \cdot \xi} dx d\xi + \sum_{j=1}^n \int D_{\xi_j} (e^{ix \cdot \xi}) g_j(x) dx d\xi \\ &= (2\pi)^n f(0) - \sum_{j=1}^n \int e^{ix \cdot \xi} D_{\xi_j} (g_j(x)) dx d\xi \\ &= (2\pi)^n f(0). \end{aligned}$$

The proof is complete. \square

Exercise

EXERCISE 3.1. Check that the tM given in (3.4) satisfies (3.4). Hint: utilize the second fact about M to facilitate the derivation.

EXERCISE 3.2. Prove Lemma 3.17.

Stationary phase lemmas

The stationary phase lemmas is a useful tools for computing certain asymptotics. Some useful references are [8, §5], [9, §19.3], [13, §7.7], [28, §3].

From time to time we will encounter oscillatory integrals of the form

$$I(\lambda) = \int e^{i\lambda\varphi(x)} a(x) dx \quad (4.1)$$

where φ is a phase function of some order and a is a symbol ($|\partial^\beta a(\xi)| \lesssim \langle \xi \rangle^{m-|\beta|}$). In §3 we have introduced schemes to make $I(\lambda)$ well-defined. Now we focus on the asymptotics of $I(\lambda)$ w.r.t. $\lambda \rightarrow +\infty$ when φ satisfies certain conditions.

When φ is linear, i.e. $\varphi(x) = p \cdot x$ for certain fixed vector $p \neq 0$, there is no critical point of φ ($|\nabla\varphi| = |p| \neq 0$). In this case we call φ *non-stationary*. For the non-stationary case, the asymptotics of I is straightforward:

$$\begin{aligned} I(\lambda) &= \int e^{i\lambda p \cdot x} a(x) dx = \int \left(\frac{p \cdot D_x}{\lambda |p|^2} \right)^N (e^{i\lambda p \cdot x}) a(x) dx \\ &= \lambda^{-N} \int e^{i\lambda p \cdot x} \left(\frac{-p \cdot D_x}{|p|^2} \right)^N (a(x)) dx \lesssim |p|^{-N} \lambda^{-N} \int \sum_{|\beta|=N} C_\beta \partial^\beta a(x) dx \\ &\lesssim |p|^{-N} \lambda^{-N} \int \langle x \rangle^{m-N} dx \lesssim \lambda^{-N}, \end{aligned}$$

provided N is large enough. This means that $\int e^{i\lambda p \cdot x} a(x) dx$ is of rapid decay w.r.t. λ .

The gradient of φ has been put in the denominator in the derivation above, so the method will not be applicable when the phase function contains critical points. In this case we call the phase *stationary*. In this chapter we devote ourselves into the stationary case.

4.1. A simple case

To help readers understand the method of stationary phase, we start with a simple case where the phase is stationary. To that end, we need to do some preparations.

4.1.1. Preliminaries. We need the Taylor's expansion. Suppose $f \in C^{N+1}(\mathbb{R}^n; \mathbb{C})$, then we have that

$$\begin{aligned} f(x) &= \sum_{|\delta| \leq N} \frac{1}{\delta!} (\partial^\delta f)(x_0) \cdot (x - x_0)^\delta \\ &\quad + (N+1) \sum_{|\delta|=N+1} \frac{(x - x_0)^\delta}{\delta!} \int_0^1 (1-t)^N (\partial^\delta f)(x_0 + t(x - x_0)) dt. \end{aligned} \quad (4.2)$$

The proof of (4.2) can be found in most of the calculus textbook and we omit it here.

Secondly, for a measurable function u in \mathbb{R}^n , as long as $\partial^\alpha u \in L^1(\mathbb{R}^n)$ for $|\alpha| \geq n+1$, then \hat{u} exists and there exists a constant C such that

$$\|\hat{u}\|_{L^1(\mathbb{R}^n)} \leq C \sum_{|\alpha| \leq n+1} \|\partial^\alpha u\|_{L^1(\mathbb{R}^n)}. \quad (4.3)$$

PROOF. We have

$$\begin{aligned} \int |\hat{u}(\xi)| d\xi &= \int \langle \xi \rangle^{-n-1} |\langle \xi \rangle^{n+1} \hat{u}(\xi)| d\xi \leq C \sup_{\mathbb{R}^n} |\langle \xi \rangle^{n+1} \hat{u}(\xi)| \\ &\leq C \sum_{|\alpha| \leq n+1} C_\alpha \sup_{\mathbb{R}^n} |\xi^\alpha \hat{u}(\xi)| \leq \sum_{|\alpha| \leq n+1} C_\alpha \sup_{\mathbb{R}^n} |\mathcal{F}\{\partial^\alpha u\}(\xi)| \\ &\leq \sum_{|\alpha| \leq n+1} C_\alpha \|\partial^\alpha u\|_{L^1(\mathbb{R}^n)}. \end{aligned}$$

We arrive at the conclusion. \square

We also need the following transformation. For a fixed non-degenerate, symmetric, real-valued square matrix Q , we have

$$\mathcal{F}\{e^{\pm i \langle Q \cdot, \cdot \rangle / 2}\}(\xi) = \frac{e^{\pm i \frac{\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} e^{\mp i \langle Q^{-1} \xi, \xi \rangle / 2}. \quad (4.4)$$

Here the non-degeneracy condition of Q means $\det Q \neq 0$.

PROOF. We have

$$\begin{aligned} \mathcal{F}\{e^{\pm i |\cdot|^2 / 2}\}(\xi) &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} e^{\pm i |x|^2 / 2} dx = (2\pi)^{-n/2} e^{\mp i |\xi|^2 / 2} \int_{\mathbb{R}^n} e^{\pm i |x \mp \xi|^2 / 2} dx \\ &= \pi^{-n/2} e^{\mp i |\xi|^2 / 2} \int_{\mathbb{R}^n} e^{\pm i |x|^2} dx. \end{aligned}$$

By standard Cauchy's integral theorem we can have

$$\int_{-\infty}^{+\infty} e^{\pm i x^2} dx = \sqrt{\pi} e^{\pm i \pi / 4},$$

so we can continue,

$$\mathcal{F}\{e^{\pm i |\cdot|^2 / 2}\}(\xi) = \pi^{-n/2} e^{\mp i |\xi|^2 / 2} (\sqrt{\pi} e^{\pm i \frac{\pi}{4}})^n = e^{\pm i \frac{\pi}{4} n} e^{\mp i |\xi|^2 / 2}. \quad (4.5)$$

We left the computation from (4.5) to (4.4) as an exercise. \square

4.1.2. A simple case. We study the quadratic case in \mathbb{R}^1 .

LEMMA 4.1. *Assume $a \in C_c^\infty(\mathbb{R})$ with $a(0) \neq 0$. Fix an arbitrary integer $N \in \mathbb{N}$. Then for the integral $I(\lambda)$:*

$$I(\lambda) = \int_{\mathbb{R}} e^{i\lambda x^2 / 2} a(x) dx,$$

there holds

$$I(\lambda) = \left(\frac{2\pi}{\lambda}\right)^{1/2} e^{i\frac{\pi}{4}} \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{i}{2}\right)^j a^{(2j)}(0) + \mathcal{O}(\lambda^{-\frac{1}{2}-N-1} \sum_{j \leq 2N+4} \sup |a^{(j)}|), \quad (4.6)$$

where $a^{(j)}$ signifies $\frac{d^j a}{dx^j}$.

PROOF. By the Plancherel theorem (which claims $(f, g) = (\hat{f}, \hat{g})$) we have

$$\begin{aligned} I(\lambda) &= \int e^{-i\lambda x^2 / 2} a(x) dx = \int (\lambda)^{-1/2} e^{-\frac{i\pi}{4}} e^{\frac{i}{2\lambda} \xi^2} \cdot \hat{a}(\xi) d\xi \\ &= \lambda^{-1/2} e^{\frac{i\pi}{4}} \int e^{-\frac{i\xi^2}{2} h} \hat{a}(\xi) d\xi, \quad h := \lambda^{-1}. \end{aligned}$$

Using (4.2) to expand $I(\lambda)$ w.r.t. h at $h = 0$, we obtain

$$\begin{aligned}
\int e^{-\frac{i\xi^2}{2}h}\hat{a}(\xi) \, d\xi &= \sum_{j=0}^N \int h^j \left(-\frac{i\xi^2}{2}\right)^j / j! \hat{a}(\xi) \, d\xi \\
&\quad + \left| \frac{h^{N+1}}{N!} \int_0^1 \int (1-t)^N \left(-\frac{i\xi^2}{2}\right)^{N+1} e^{-\frac{i\xi^2}{2}th} \hat{a}(\xi) \, d\xi \, dt \right| \\
&= \sum_{j=0}^N h^j \frac{\left(\frac{i}{2}\right)^j}{j!} \int \mathcal{F}\{a^{(2j)}\}(\xi) \, d\xi + \mathcal{O}(h^{N+1} \int |\xi^{2N+2} \hat{a}(\xi)| \, d\xi) \\
&= \sum_{j=0}^N \frac{(2\pi)^{1/2} \left(\frac{i}{2}\right)^j h^j}{j!} a^{(2j)}(0) + \mathcal{O}(h^{N+1} \|\mathcal{F}\{a^{(2N+2)}\}\|_{L^1}) \tag{4.7} \\
&= \sum_{j=0}^N \frac{(2\pi)^{1/2} \left(\frac{i}{2}\right)^j h^j}{j!} a^{(2j)}(0) + \mathcal{O}(h^{N+1} \sum_{j \leq 2N+4} \|a^{(j)}\|_{L^1(\mathbb{R}^n)}).
\end{aligned}$$

Note that we used (4.3). Combining the computations and changing h back to λ^{-1} , we arrive at

$$I(\lambda) = \left(\frac{2\pi}{\lambda}\right)^{1/2} e^{\frac{i\pi}{4}} \sum_{j=0}^N \frac{\lambda^{-j}}{j!} \left(\frac{i}{2}\right)^j a^{(2j)}(0) + \mathcal{O}(\lambda^{-\frac{1}{2}-N-1} \sum_{j \leq 2N+4} \sup |a^{(j)}|).$$

The proof is complete. \square

From this short proof, we see extract several main steps:

- (1) use Plancherel theorem to turn λ into λ^{-1} in the exponent;
- (2) expand the integral w.r.t. $h := \lambda^{-1}$ at $h = 0$ with integral remainder;
- (3) estimate the remainder with (4.3).

4.2. Lemma Statements

THEOREM 4.2. *Let $n \in \mathbb{N}^+$ be the dimension. We consider the oscillatory integral $I(\lambda)$:*

$$I(\lambda) = \int_{\mathbb{R}^n} e^{i\lambda \langle Q(x-x_0), x-x_0 \rangle / 2} a(x; \lambda) \, dx,$$

where $\langle Qx, y \rangle$ signifies $(Qx)^T y$ as matrix multiplication. Fix two arbitrary integers $M, N \in \mathbb{N}$, and we assume

- Q is a non-degenerate, symmetric, real-valued matrix;
- for each λ , $a(\cdot; \lambda) \in C^{n+2N+3}(\mathbb{R}^n; \mathbb{C})$;
- for each λ , $a(\cdot; \lambda) \in C^M(\mathbb{R}^n; \mathbb{C})$, and $\forall \alpha : |\alpha| \leq M$, there exists λ -dependent constants $C_{M,\alpha}(\lambda) > 0$ such that $\forall x \in \mathbb{R}^n$ there holds

$$|\partial_x^\alpha a(x - x_0; \lambda)| < C_{M,\alpha}(\lambda) \langle x \rangle^{2M-n-1-|\alpha|}.$$

Then the integral $I(\lambda)$ is well-defined in the oscillatory integral sense, and as $\lambda \rightarrow +\infty$ we have

$$\begin{aligned}
 I(\lambda) = & \left(\frac{2\pi}{\lambda} \right)^{n/2} \frac{e^{i\frac{\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1}D, D \rangle}{2i} \right)^j (a(x; \lambda))|_{x=x_0} \\
 & + \mathcal{O}(\lambda^{-\frac{n}{2}-N-1} \times \sum_{|\alpha| \leq n+2N+3} \sup_{B(x_0,1)} |\partial^\alpha a(\cdot; \lambda)|) \\
 & + \mathcal{O}(\lambda^{-M} \times \sum_{|\alpha| \leq M} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(x-x_0; \lambda)|}{\langle x \rangle^{2M-n-1-|\alpha|}}),
 \end{aligned} \tag{4.8}$$

where $B(x_0, 1)$ stands for the ball centered at x_0 with radius 1, and $\operatorname{sgn} Q$ stands for the difference between the number of positive eigenvalues and the number of negative eigenvalues of the matrix Q .

REMARK 4.3. In contrast to many other versions of the stationary phase lemma, here we don't require a to be compactly supported. Instead, some other boundedness conditions are required, which makes the oscillatory integral well-defined.

REMARK 4.4. Eq. (4.8) is not an asymptotics w.r.t. j , but is rather w.r.t. λ . To get enough terms w.r.t. j , one could choose M be large enough first, and then check if a satisfies the requirements.

REMARK 4.5. The integers M and N in Theorem 4.2 shall be chosen properly to serve for your own purposes. For example, if one cares more about the decaying behavior w.r.t. λ , then the M can be set to $\lceil n/2 \rceil + N + 1$. However, if one is dealing with these functions a which doesn't have good decaying behavior at the infinity, then one could set M to be large enough such that $\langle x \rangle^{2M-n-1-|\alpha|}$ can dominate $\partial^\alpha a$, with the cost that we should demand more smoothness of $a(x)$.

REMARK 4.6. The unit ball $B(x_0, 1)$ involved in the term $\sup_{B(x_0,1)} |\partial^\alpha a|$ can be changed to other bounded domain containing x_0 . But one should be careful that when the domain used has a very small radius, the underlying coefficients of the $\mathcal{O}(\cdot)$ term will be very large accordingly.

REMARK 4.7. The function a is allowed to be dependent on λ , hence the expression (4.8) is an asymptotic expansion only when $\partial_x^\alpha a(x; \lambda)$ doesn't increase significantly when $\lambda \rightarrow +\infty$.

If chosen $M = n + 2N + 3$, Theorem 4.2 will be simplified as follows.

PROPOSITION 4.8. *Let $n \in \mathbb{N}^+$ be the dimension. We consider the oscillatory integral $I(\lambda)$:*

$$I(\lambda) = \int_{\mathbb{R}^n} e^{i\lambda \langle Q(x-x_0), x-x_0 \rangle / 2} a(x; \lambda) dx,$$

where $\langle Qx, y \rangle$ signifies $(Qx)^T y$ as matrix multiplication. Fix an integer $N \in \mathbb{N}$, and we assume

- Q is a non-degenerate, symmetric, real-valued matrix;
- for each λ , $a(\cdot; \lambda) \in C^{n+2N+3}(\mathbb{R}^n; \mathbb{C})$, and $\forall \alpha : |\alpha| \leq n + 2N + 3$, there exists λ -dependent constants $C_{N,n,\alpha}(\lambda) > 0$ such that $\forall x \in \mathbb{R}^n$ there holds

$$|\partial_x^\alpha a(x-x_0; \lambda)| < C_{N,n,\alpha}(\lambda) \langle x \rangle^{2N+2}. \tag{4.9}$$

Then the integral $I(\lambda)$ is well-defined in the oscillatory integral sense, and as $\lambda \rightarrow +\infty$ we have

$$I(\lambda) = \left(\frac{2\pi}{\lambda}\right)^{n/2} \frac{e^{i\frac{\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1}D, D \rangle}{2i}\right)^j (a(x; \lambda))|_{x=x_0} + \mathcal{O}\left(\lambda^{-\frac{n}{2}-N-1} \sum_{|\alpha| \leq n+2N+3} \sup_{x \in \mathbb{R}^n} \frac{|\partial^\alpha a(x-x_0; \lambda)|}{\langle x \rangle^{n+4N+5-|\alpha|}}\right). \quad (4.10)$$

Proposition 4.8 can be extended to a more general case where the phase function is not quadratic.

THEOREM 4.9 (Stationary phase lemma [9]). *We consider the oscillatory integral $I(\lambda)$:*

$$I(\lambda) = \int_{\mathbb{R}^n} e^{i\lambda\varphi(x)} a(x) dx.$$

For an arbitrary integer $N \in \mathbb{N}$, assume

- $a \in C^{n+2N+3}(\mathbb{R}^n; \mathbb{C})$ with $\sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} |\partial^\alpha a| < +\infty$;
- $\varphi \in C^{n+2N+6}(\mathbb{R}^n; \mathbb{R})$ with $\sum_{|\alpha| \leq n+2N+6} \sup_{\mathbb{R}^n} |\partial^\alpha \varphi| < +\infty$;
- x_0 is the only critical point of $\varphi(x)$ on $\operatorname{supp} a(x)$, i.e., $\varphi(x_0) = \nabla\varphi(x_0) = 0$, $\varphi_x(x) \neq 0$ for $x \neq x_0$;
- the Hessian $\varphi_{xx}(x_0) := [\frac{\partial^2 \varphi}{\partial x_j \partial x_k}(x_0)]_{j,k=1}^n$ satisfies $\det \varphi_{xx}(x_0) \neq 0$.

Then the integral $I(\lambda)$ is well-defined in the oscillatory integral sense, and as $\lambda \rightarrow +\infty$ we have

$$I(\lambda) = \left(\frac{2\pi}{\lambda}\right)^{n/2} \frac{e^{i\lambda\varphi(x_0) + i\frac{\pi}{4} \operatorname{sgn} \varphi_{xx}(x_0)}}{|\det \varphi_{xx}(x_0)|^{1/2}} \left(a(x_0) + \sum_{j=1}^N a_j(x_0) \lambda^{-j}\right) + \mathcal{O}\left(\lambda^{-\frac{n}{2}-N-1} \times \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} |\partial^\alpha a| \times \sum_{|\alpha| \leq n+2N+6} \sup_{\mathbb{R}^n} |\partial^\alpha \varphi|\right), \quad (4.11)$$

for some functions a_j , $1 \leq j \leq N$.

REMARK 4.10. Proposition 4.8 is a special case of Theorem 4.9 where

$$\varphi(x) = \langle Q(x-x_0), x-x_0 \rangle / 2,$$

which guarantees $\varphi_{xx}(x_0) = Q$. Theorem 4.9 is not a generalization of Theorem 4.2 because unlike the quadratic phase function in Theorem 4.2, the phase function φ in Theorem 4.9 is not assumed to possess the property that $|\nabla\varphi(x)| \simeq |x|$ as $|x|$ is large. However if φ is a homeomorphism of \mathbb{R}^n , it is possible to generalize Theorem 4.2.

In one-dimensional case, explicit expressions for these $a_j(x_0)$ are given in [28, (3.4.11)], and the details are given in [28, Second proof of Theorem 3.11]. For explicit expressions for these $a_j(x_0)$ in higher dimension, readers may refer to [13, Theorem 7.7.5] for details. In [9, Lemma 19.3] there is also another routine to prove the stationary phase lemma. [8, Chapter 5] by Dimassi and Sjöstrand is also a good reference. See also [27, §2.3 & §6.4].

4.3. Proofs of the results

We first prove the quadratic case.

PROOF OF THEOREM 4.2. We omit notationally the dependence of a on λ until related clarifications are needed. Without loss of generality we assume $x_0 = 0$, and $a(0) = 1$. For readers' convenient we rewrite the expression of I here: $I(\lambda) = \int_{\mathbb{R}^n} e^{i\lambda\langle Qx, x \rangle/2} a(x) dx$.

Step 1: cutoff singularity of the phase function. According to the assumption on Q , we know there exists a decomposition $Q = P\Lambda P^T$ where P is an orthogonal matrix and $\Lambda := (\alpha_j)_{j=1, \dots, n}$ is a diagonal matrix. Make the change of variable $y = P^T x$, we can have

$$\begin{aligned} I &= \int_{\mathbb{R}^n} a(Py) e^{i\lambda \sum_{j=1}^n \alpha_j y_j^2/2} dy \\ &= \int_{\mathbb{R}^n} e^{i\lambda \sum_{j=1}^n \alpha_j y_j^2/2} (1 - \chi(y)) f(y) dy + \int_{\mathbb{R}^n} e^{i\lambda \sum_{j=1}^n \alpha_j y_j^2/2} \chi(y) f(y) dy \quad (\det P = 1) \\ &=: J_1 + J_2, \end{aligned} \tag{4.12}$$

where $f(y) := a(Py)$ and $\chi \in C_c^\infty(\mathbb{R}^n)$ is a cutoff function satisfying $0 \leq \chi \leq 1$ and $\chi \equiv 1$ in a neighborhood of the origin. We will see:

J_1 is rapidly decaying and J_2 gives the desired asymptotics.

Step 2: J_1 is rapidly decaying. Noting that neighborhoods of the origin is not included in the support of the integrand in J_1 , we can estimate J_1 by using integration by parts (in the oscillatory integral sense). For any integer $M \in \mathbb{N}$ we have

$$\begin{aligned} |J_1| &= \left| \int_{\mathbb{R}^n} \left(\frac{\sum_j \alpha_j^{-1} y_j \partial_j}{i\lambda |y|^2} \right)^M (e^{i\lambda \sum_{j=1}^n \alpha_j y_j^2/2}) [(1 - \chi(y)) f(y)] dy \right| \\ &\lesssim \lambda^{-M} \int_{\mathbb{R}^n} \left| \left(\sum_j \partial_j \circ (y_j |y|^{-2}) \right)^M [(1 - \chi(y)) f(y)] \right| dy \\ &\lesssim \lambda^{-M} \int_{\text{supp}(1-\chi)} \sum_{|\alpha| \leq M} C_{M;\alpha} |y|^{|\alpha|-2M} |\partial^\alpha((1 - \chi)f(y))| dy \end{aligned} \tag{4.13}$$

$$\lesssim \lambda^{-M} \left(C \sum_{|\alpha| \leq M} \sup_{\{0 < \chi < 1\}} |\partial^\alpha f| + \sum_{|\alpha| \leq M} C_{M;\alpha} \int_{\{\chi=0\}} |y|^{|\alpha|-2M} |\partial^\alpha f(y)| dy \right) \tag{4.14}$$

$$\lesssim \lambda^{-M} \sum_{|\alpha| \leq M} \left(\sup_{\{0 < \chi < 1\}} |\partial^\alpha a| + \int_{\{\chi=0\}} |y|^{-n-1} \langle y \rangle^{|\alpha|-2M+n+1} |\partial^\alpha a(y)| dy \right)$$

$$\lesssim \lambda^{-M} \sum_{|\alpha| \leq M} \left(\sup_{\{0 < \chi < 1\}} |\partial^\alpha a| + \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y)|}{\langle y \rangle^{2M-n-1-|\alpha|}} \right)$$

$$\lesssim \lambda^{-M} \sum_{|\alpha| \leq M} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y)|}{\langle y \rangle^{2M-n-1-|\alpha|}}. \tag{4.15}$$

The inequality (4.13) is due to the fact that

$$\left(\sum_{1 \leq j \leq n} \partial_j \circ (y_j |y|^{-2}) \right)^M \varphi = \sum_{|\alpha| \leq M} C_{M;\alpha} |y|^{|\alpha|-2M} \partial^\alpha \varphi$$

which can be derived by induction and we omit the details. Inequality (4.14) is due to the fact that $\partial^\alpha((1 - \chi)f) = \partial^\alpha f$ in $\{\chi = 0\}$.

Step 3: J_2 and Plancherel theorem. We turn to J_2 . Keep in mind that $f(y) = a(Py)$ and χf is compactly support and in $C_c^{n+2N+3}(\mathbb{R}^n)$. Here we analyze I by borrowing idea from [28, First proof of Theorem 3.11]. First we use Plancherel theorem (which states $(f, g) = (\hat{f}, \hat{g})$),

$$J_2 = \int_{\mathbb{R}^n} \overline{e^{-i\lambda \sum_{j=1}^n \alpha_j y_j^2/2}} \chi f(y) dy = \int_{\mathbb{R}^n} \overline{\mathcal{F}\{e^{-i\lambda \sum_{j=1}^n \alpha_j (\cdot)^2/2}\}(\xi)} \cdot \widehat{\chi f}(\xi) d\xi$$

$$\begin{aligned}
&= \int_{\mathbb{R}^n} \overline{(\lambda)^{-n/2} \frac{e^{-\frac{i\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} e^{\frac{i}{2\lambda} \alpha_j^{-1} \xi_j^2} \cdot \widehat{\chi f}(\xi)} \, d\xi \quad (\text{by (4.4)}) \\
&=: \left(\frac{2\pi}{\lambda}\right)^{n/2} \frac{e^{\frac{i\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} J(1/\lambda, 1/\lambda, \chi f), \tag{4.16}
\end{aligned}$$

where we ignored the summation notation over j and the function J is defined by

$$J(h_1, h_2, \chi f) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{\frac{\xi_j^2 h_1}{i2\alpha_j}} \cdot \widehat{\chi f}(\xi; 1/h_2) \, d\xi. \tag{4.17}$$

Note that in (4.17) we put emphasize on the dependence of f on h_2 (i.e. dependence of a on λ). The smoothness of J w.r.t. h_1 is guaranteed by the L^1 of derivatives of f , namely, we have the following claim whose justification will be clear in (4.20),

$$\forall m \in \mathbb{N}, \quad \max_{|\alpha| \leq n+2m+1} \|\partial^\alpha f\|_{L^1(\mathbb{R}^n)} < +\infty \Rightarrow J(\cdot, h_2, f) \in C^m(\mathbb{R}).$$

Step 4: Taylor's expansion. We abbreviate $\partial_{h_1} J$ as $\partial_1 J$. Expand J w.r.t. h_1 ,

$$\begin{aligned}
\partial_1^k J(0, h_2, \chi f) &= (2\pi)^{-n/2} \int_{\mathbb{R}^n} \partial_1^k (e^{\frac{\xi_j^2 h_1}{i2\alpha_j}})|_{h_1=0} \cdot \widehat{\chi f}(\xi) \, d\xi \\
&= (2\pi)^{-n/2} \int_{\mathbb{R}^n} \left(\sum_j \frac{\xi_j^2}{i2\alpha_j}\right)^k \cdot \widehat{\chi f}(\xi) \, d\xi \\
&= (2\pi)^{-n/2} \int_{\mathbb{R}^n} \widehat{T^k \chi f}(\xi) \, d\xi = T^k f(0) = T^k(a(Py))|_{y=0} \\
&= \left(\frac{i}{2} P^{lj} P^{kj} \alpha_j^{-1} \partial_{kl}\right)^k a(0) = A^k a(0; 1/h_2),
\end{aligned}$$

where $T = \frac{i}{2} \sum_j \frac{\partial_j^2}{\alpha_j}$ and $A = \frac{i}{2} (Q^{-1})^{jl} \partial_{jl} = \frac{1}{2i} \langle Q^{-1} D, D \rangle$ (recall that $D = \frac{1}{i} \nabla$ is vertical). We expand J using Taylor series (i.e. (4.2)),

$$\begin{aligned}
J(h, h_2, \chi f) &= \sum_{k \leq N} \frac{h^k}{k!} \partial_h^k J(0, h_2, \chi f) + \frac{h^{N+1}}{N!} \int_0^1 (1-t)^N \cdot \partial_1^{N+1} J(th, h_2, \chi f) \, dt \\
&= \sum_{0 \leq k \leq N} \frac{(hA)^k}{k!} a(0; h_2) + \frac{h^{N+1}}{N!} \int_0^1 (1-t)^N \cdot \partial_1^{N+1} J(th, h_2, \chi f) \, dt. \tag{4.18}
\end{aligned}$$

Step 5: The remainder term. By invoking (4.17), the remainder term in (4.18) can be estimated as

$$\begin{aligned}
& \left| \frac{h^{N+1}}{N!} \int_0^1 (1-t)^N \cdot \partial_1^{N+1} J(th, h_2, \chi f) \, dt \right| \\
& \leq C_N h^{N+1} \int_{\mathbb{R}^n} \left| \left(\frac{-i}{4\alpha_j} \xi_j^2\right)^{N+1} \cdot \widehat{\chi f}(\xi; 1/h_2) \right| \, d\xi \\
& \leq C_N h^{N+1} \sum_{|\beta| \leq 2N+2} C_\beta \|(\partial_x^\beta (\chi f(\cdot; 1/h_2)))^\wedge\|_{L^1(\mathbb{R}^n)}.
\end{aligned}$$

By using (4.17) and (4.3), we can continue

$$\left| \frac{h^{N+1}}{N!} \int_0^1 (1-t)^N \cdot \partial_1^{N+1} J(th, h_2, \chi f) \, dt \right|$$

$$\leq C_N h^{N+1} \sum_{\substack{|\alpha| \leq n+1 \\ |\beta| \leq 2N+2}} \|\partial_x^{\alpha+\beta}(\chi f(\cdot; 1/h_2))\|_{L^1(\mathbb{R}^n)} \quad (4.19)$$

$$\begin{aligned} &\leq C_N h^{N+1} \sum_{|\alpha| \leq n+2N+3} \|\partial_x^\alpha(\chi f(\cdot; 1/h_2))\|_{L^1(\mathbb{R}^n)} \\ &\leq C_N h^{N+1} \sum_{|\alpha| \leq n+2N+3} \sup_{\text{supp } \chi} |\partial^\alpha a(\cdot; 1/h_2)|. \end{aligned} \quad (4.20)$$

Letting $h = h_2 = 1/\lambda$ and combining (4.16), (4.17), (4.18) and (4.20), we obtain

$$\begin{aligned} J_2 &= \left(\frac{2\pi}{\lambda}\right)^{n/2} \frac{e^{\frac{i\pi}{4}\text{sgn } Q}}{|\det Q|^{1/2}} \sum_{j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1}D, D \rangle}{2i}\right)^j a(0; \lambda) \\ &\quad + C_N \lambda^{-\frac{n}{2}-N-1} \sum_{|\alpha| \leq n+2N+3} \sup_{\text{supp } \chi} |\partial^\alpha a(\cdot; \lambda)|. \end{aligned} \quad (4.21)$$

Combining (4.21) with (4.12), (4.15), we have

$$\begin{aligned} I &= \left(\frac{2\pi}{\lambda}\right)^{n/2} \frac{e^{\frac{i\pi}{4}\text{sgn } Q}}{|\det Q|^{1/2}} \sum_{j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1}D, D \rangle}{2i}\right)^j a(0; \lambda) \\ &\quad + \mathcal{O}\left(\lambda^{-\frac{n}{2}-N-1} \sum_{|\alpha| \leq n+2N+3} \sup_{\text{supp } \chi} |\partial^\alpha a(\cdot; \lambda)|\right) + \mathcal{O}\left(\lambda^{-M} \sum_{|\alpha| \leq M} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{2M-n-1-|\alpha|}}\right), \end{aligned}$$

which is (4.8). The proof is complete. \square

PROOF OF PROPOSITION 4.8. The statement of Proposition 4.8 is almost the same as Theorem (4.2), except that M is set to be $n + 2N + 3$. Hence, we set $M := n + 2N + 3$, then we have $-M \leq -n/2 - N - 1$, so

$$\begin{aligned} \lambda^{-M} \sum_{|\alpha| \leq M} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{2M-n-1-|\alpha|}} &= \lambda^{-M} \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{2(n+2N+3)-n-1-|\alpha|}} \\ &\leq \lambda^{-n/2-N-1} \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{n+4N+5-|\alpha|}}. \end{aligned}$$

Also, for the first remainder term in (4.8) we have

$$\begin{aligned} \sum_{|\alpha| \leq n+2N+3} \sup_{B(x_0, 1)} |\partial^\alpha a(\cdot; \lambda)| &\leq C_{x_0, n, N} \sum_{|\alpha| \leq n+2N+3} \sup_{y \in B(x_0, 1)} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{n+4N+5-|\alpha|}} \\ &\leq C_{x_0, n, N} \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} \frac{|\partial^\alpha a(y; \lambda)|}{\langle y \rangle^{n+4N+5-|\alpha|}}. \end{aligned}$$

Combining these with (4.8), we arrive at (4.10). The proof is complete. \square

Based on Theorem 4.2, now we prove the more general case.

PROOF OF THEOREM 4.9. Without loss of generality we assume $x_0 = 0$, $\varphi(0) = 0$ and $a(0) = 1$. Hence by Taylor's expansion (4.2) we have

$$\varphi(x) = \sum_{j, k \leq n} x_j x_k \int_0^1 (1-t) \partial_{jk} \varphi(tx) dt = x^T \cdot \int_0^1 (1-t) \varphi_{xx}(tx) dt \cdot x.$$

Note that $|\varphi_{xx}(0)| \neq 0$ and $|\varphi_{xx}(x)|$ is continuous on x ($\varphi \in C^2$), thus there exists a positive constant \boxed{r} such that $|\varphi_{xx}(x)| > |\varphi_{xx}(0)|/2 > 0$ for all $x \in B(0, r)$. Fix a cutoff function $\chi \in C_c^\infty(\mathbb{R}^n)$ such that $\text{supp } \chi \subset B(0, r)$ and $\chi \equiv 1$ in $B(0, r/2)$. Hence:

- on $B(0, r)$, matrix φ_{xx} is non-degenerate;
- on $\text{supp } a \setminus B(0, r)$, $|\nabla\varphi(x)|$ is uniformly bounded away from 0.

Step 1: cutoff singularity of the phase function. We divide I into two parts

$$I(\lambda) = \int_{\mathbb{R}^n} (1 - \chi(x))a(x)e^{i\lambda\varphi(x)} dx + \int_{\mathbb{R}^n} \chi(x)a(x)e^{i\lambda\varphi(x)} dx := I_1 + I_2, \quad (4.22)$$

and we will show that I_1 is rapidly decreasing w.r.t. λ while I_2 can be analyzed by using Theorem 4.2.

Step 2: I_1 is rapidly decaying. For I_1 , denote $L = \sum_{j=1}^n \frac{\varphi_{x_j}}{|\nabla\varphi|^2} \partial x_j$, where φ_{x_j} is short for $\partial_{x_j}\varphi$. Then $\frac{1}{i\lambda}Le^{i\lambda\varphi} = e^{i\lambda\varphi}$ and ${}^tLf = \sum_{j=1}^n \partial_{x_j}(\frac{\varphi_{x_j}}{|\nabla\varphi|^2})$. For any $N \in \mathbb{N}^+$, I_1 can be easily estimated as follows (which requires $a \in C^{n+N+1}(\mathbb{R}^n)$ and $\varphi \in C^{n+N+2}(\mathbb{R}^n)$)

$$\begin{aligned} I_1 &= \int_{\mathbb{R}^n} (1 - \chi)a \cdot ((i\lambda)^{-n-N-1} L^{N+1} e^{i\lambda\varphi(x)}) dx \\ &= (i\lambda)^{-n-N-1} \int_{\mathbb{R}^n} ({}^tL)^{n+N+1} ((1 - \chi)a) \cdot e^{i\lambda\varphi(x)} dx \\ &= \mathcal{O}(\lambda^{-n-N-1} \sum_{|\alpha| \leq n+N+1} \|\partial^\alpha a\|_{L^1(\mathbb{R}^n)}), \quad \lambda \rightarrow \infty. \end{aligned} \quad (4.23)$$

As mentioned before, due to the presence of $1 - \chi$, the denominator $|\nabla\varphi|^2$ in L keeps a positive distance away from 0, guaranteeing that $({}^tL)^N((1 - \chi)a)$ is bounded and compactly supported.

Step 3: Turn I_2 into quadratic phase form (e.g. “ J_2 ”). Now we turn to I_2 . Because $\varphi \in C^2(\mathbb{R}^n)$, $\varphi_{xx}(x)$ is symmetric and thus there exist orthogonal matrix $P(x)$ and diagonal matrix $\Lambda(x) = (\alpha_j(x))_{j=1, \dots, n}$ such that

$$2 \int_0^1 (1 - t)\varphi_{xx}(tx) dt = P(x)\Lambda(x)P^T(x).$$

Especially we have $P(0)\Lambda(0)P^T(0) = \varphi_{xx}(0)$. Denote $\alpha_j = \alpha_j(0)$ and $n \times n$ diagonal matrix $\Lambda := (\alpha_j)_{j=1, \dots, n}$ for short. Thus

$$\Lambda(x) = \left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot \Lambda \cdot \left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n}.$$

Note that we can choose the support of χ to be small enough such that, on $\text{supp } \chi$, $\alpha_j(x)$ doesn't change sign, so $\alpha_j(x)/\alpha_j$ will always be positive on $\text{supp } \chi$. This grants the use of the square root operation.

Make the change of variable:

$$y = \Phi(x) := \left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot P^T(x) \cdot x. \quad (4.24)$$

Note that

$$\varphi \in C^{n+2N+6} \Rightarrow \Phi \in C^{n+2N+4}. \quad (4.25)$$

We have

$$\varphi(x) = \frac{1}{2}x^T \cdot \left[2 \int_0^1 (1 - t)\varphi_{xx}(tx) dt \right] \cdot x = \frac{1}{2}x^T \cdot [P(x)\Lambda(x)P^T(x)] \cdot x$$

$$\begin{aligned}
&= \frac{1}{2} [P^T(x) \cdot x]^T \cdot \left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot \Lambda \cdot \left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot [P^T(x) \cdot x] \\
&= \frac{1}{2} \left[\left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot P^T(x) \cdot x \right]^T \cdot \Lambda \cdot \left[\left(\sqrt{\frac{\alpha_j(x)}{\alpha_j}} \right)_{j=1, \dots, n} \cdot P^T(x) \cdot x \right] \\
&= \frac{1}{2} \langle \Lambda y, y \rangle.
\end{aligned}$$

We have $\Phi(0) = 0$. It is easy to check that $\frac{\partial \Phi}{\partial x}(0) = P^T(0)$. From (4.24) it is clear that there exists inverse of Φ , i.e. $\phi = \Phi^{-1}$. Note that $x = \phi(\Phi(x))$ and

$$\Phi \in C^{n+2N+4} \Rightarrow \phi \in C^{n+2N+4}. \quad (4.26)$$

We have

$$\begin{aligned}
I_2 &= \int_{\mathbb{R}^n} \chi(\phi(y)) a(\phi(y)) \cdot e^{i\lambda \langle \Lambda y, y \rangle / 2} d\phi(y) \\
&= \int_{\mathbb{R}^n} \chi(\phi(y)) a(\phi(y)) |\det \nabla_y \phi(y)| \cdot e^{i\lambda \langle \Lambda y, y \rangle / 2} dy \\
&= \int_{\mathbb{R}^n} f(y) e^{i\lambda \langle \Lambda y, y \rangle / 2} dy
\end{aligned}$$

where

$$f(y) = \chi(\phi(y)) \cdot a(\phi(y)) \cdot |\det \nabla_y \phi(y)|,$$

Note that

$$\phi \in C^{n+2N+4}, a \in C^{n+2N+2} \Rightarrow f \in C^{n+2N+2}. \quad (4.27)$$

Now we can conclude from (4.27), (4.27) and (4.27) that

$$\varphi \in C^{n+2N+6}, a \in C^{n+2N+3} \Rightarrow f \in C^{n+2N+3}. \quad (4.28)$$

Step 4: Apply Theorem 4.2. By using Theorem 4.2, we can obtain

$$\begin{aligned}
I_1(\lambda) &= \left(\frac{2\pi}{\lambda} \right)^{n/2} \frac{e^{i\frac{\pi}{4} \operatorname{sgn} \Lambda}}{|\det \Lambda|^{1/2}} \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle \Lambda^{-1} D, D \rangle}{2i} \right)^j f(0) \\
&\quad + \mathcal{O}(\lambda^{-\frac{n}{2} - N - 1}) \times \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} |\partial^\alpha f| \\
&= \left(\frac{2\pi}{\lambda} \right)^{n/2} \frac{e^{i\frac{\pi}{4} \operatorname{sgn} \Lambda}}{|\det \Lambda|^{1/2}} \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle \Lambda^{-1} D, D \rangle}{2i} \right)^j f(0) \\
&\quad + \mathcal{O}(\lambda^{-\frac{n}{2} - N - 1}) \times \sum_{|\alpha| \leq n+2N+3} \sup_{\mathbb{R}^n} |\partial^\alpha a| \times \sum_{|\alpha| \leq n+2N+6} \sup_{\mathbb{R}^n} |\partial^\alpha \varphi|. \quad (4.29)
\end{aligned}$$

It can be checked that $\operatorname{sgn} \Lambda = \operatorname{sgn} \varphi_{xx}(0)$ and $\det \Lambda = \det \varphi_{xx}(0)$.

Step 5: The leading term. We are now almost arrive at (4.11) except for the explicit computation of the leading term in (4.11) and (4.29). From equality $x = \phi(\Phi(x))$ we know $I = \nabla_y \phi(\Phi(x)) \cdot \nabla_x \Phi(x)$. Formula (4.24) implies $\Phi(0) = 0$ and $\nabla_x \Phi(0) = P^T(0)$, hence $\det \nabla_y \phi(0) = \det \nabla_y \phi(\Phi(0)) = (\det \nabla_x \Phi(0))^{-1} = (\det P^T(0))^{-1} = 1$. Therefore,

$$f(0) = \chi(\phi(0)) \cdot a(\phi(0)) \cdot |\det \nabla_y \phi(0)| = \chi(0) \cdot a(0) = a(0). \quad (4.30)$$

Combining (4.22), (4.23), (4.29) and (4.30), we arrive at the conclusion. \square

Exercise

EXERCISE 4.1. Use (4.5) to derive (4.4).

EXERCISE 4.2. Show details about how to derive (4.7) from $\int e^{-\frac{i\xi^2}{2}h} \widehat{a}(\xi) d\xi$.

EXERCISE 4.3. (optional) In (4.17), if we instead set

$$J(h, \chi f) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{\frac{\xi_j^2 h}{i2\alpha_j}} \cdot \widehat{\chi f}(\xi; 1/h) d\xi,$$

and later on expand J w.r.t. h at $h = 0$, will the computations following (4.17) still give the desired result? Explain the reason briefly.

EXERCISE 4.4. Assume $a \in C_c^\infty(\mathbb{R}^{2n})$ and denote a Lebesgue integral

$$I(y, \eta; \lambda) := (2\pi)^{-n} \int_{\mathbb{R}^{2n}} e^{i\lambda x \cdot \xi} a(x + y, \xi + \eta) dx d\xi.$$

- (1) fix y and η , and use Proposition 4.8 to find the asymptotic expansion of I w.r.t. λ as $\lambda \rightarrow +\infty$;
- (2) write down the first $1 + n$ terms (the leading term + the first order terms) of the asymptotic expansion.

Hint: $x \cdot \xi = \frac{1}{2} \langle Q(x, \xi), (x, \xi) \rangle$ with $Q = \begin{pmatrix} 0 & I_{n \times n} \\ I_{n \times n} & 0 \end{pmatrix}$, where (x, ξ) is treated as a vertical vector.

EXERCISE 4.5. Assume symbol $a \in S^m(\mathbb{R}^n \times \mathbb{R}^n)$ and denote an oscillatory integral

$$I(y, \eta) := (2\pi)^{-n} \int_{\mathbb{R}^{2n}} e^{ix \cdot \xi} a(x + y, \xi + \eta) dx d\xi.$$

- (1) is I well-defined? If it is, should the cutoff function χ (cf. (3.12)) be chosen to cutoff ξ alone using $\chi(\epsilon\xi)$, or cutoff x alone using $\chi(\epsilon x)$, or cutoff both x and ξ together using $\chi(\epsilon x, \epsilon\xi)$?
- (2) use Proposition 4.8 to find the asymptotic expansion of I w.r.t. $\langle \eta \rangle$ as $|\eta| \rightarrow +\infty$;
- (3) write down the first $1 + n$ terms (the leading term + the first order terms) of the asymptotic expansion.
- (4) compare with the result in Exercise 4.4, and revise Remark 4.7.

Hint: Perform the change of variable $\xi \rightarrow \langle \eta \rangle \xi$.

Symbolic calculus of Ψ DOs

In this chapter we show certain symbolic calculus of Ψ DOs. We need some preparations.

LEMMA 5.1. *Assume $a, b \in \mathbb{R}$ such that $|a| \geq 1$ and $|b| \geq 1$, then for every $m \in \mathbb{R}$ there exists a constant independent of a, b such that*

$$\langle ab \rangle^m \leq C_m \langle a \rangle^m \langle b \rangle^m, \quad |a| \geq 1, \quad |b| \geq 1.$$

PROOF. When $m \geq 0$, we have

$$\langle ab \rangle^m \simeq (1 + |ab|)^m \leq (1 + |a|)^m (1 + |b|)^m \leq \langle a \rangle^m \langle b \rangle^m.$$

When $m < 0$, because $|a|, |b| \geq 1$, we have

$$\begin{aligned} \langle ab \rangle^m &\simeq \frac{1}{(1 + |ab|)^{|m|}} < \frac{1}{|ab|^{|m|}} = \langle a \rangle^m \langle b \rangle^m \left(\frac{\langle a \rangle}{|a|} \frac{\langle b \rangle}{|b|} \right)^{|m|} \\ &\lesssim \langle a \rangle^m \langle b \rangle^m. \end{aligned}$$

We proved the result. \square

LEMMA 5.2 (Peetre's inequality). *For $\forall a, b \in \mathbb{R}^n$ and $\forall m \in \mathbb{R}$, there exists a constant C_m independent of a and b such that*

$$\langle a \pm b \rangle^m \leq C_m \langle a \rangle^m \langle b \rangle^{|m|}.$$

PROOF. For any $a, b \in \mathbb{R}^n$, we have

$$1 + |a - b| \leq 1 + |a| + |b| \leq (1 + |a|) \cdot (1 + |b|).$$

Note that $\langle a \rangle \simeq 1 + |a|$, so we can conclude Lemma 5.2 for the case where $m \geq 0$.

When $m < 0$, we use the fact:

$$\begin{aligned} 1 + |a| &\leq 1 + |a - b| + |b| \leq (1 + |a - b|) \cdot (1 + |b|) \\ &\Rightarrow (1 + |a - b|) \geq (1 + |a|) \cdot (1 + |b|)^{-1}. \end{aligned}$$

Now assume $m < 0$, we have

$$(1 + |a - b|)^m \leq (1 + |a|)^m \cdot (1 + |b|)^{-m} = (1 + |a|)^m (1 + |b|)^{|m|}.$$

The proof is complete. \square

5.1. Composition of Ψ DOs

Assume $a \in S^{m_1}$ and $b \in S^{m_2}$. For notational convenience we denote $T = T_a \circ T_b$, thus for any $\varphi \in \mathcal{S}$, we have

$$\begin{aligned} T\varphi &= (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} a(x, \xi) T_b \varphi(y) dy d\xi \\ &= (2\pi)^{-n} \int e^{i(x-z)\cdot\eta} \left((2\pi)^{-n} \int e^{i(x-y)\cdot(\xi-\eta)} a(x, \xi) b(y, \eta) dy d\xi \right) \varphi(z) dz d\eta \\ &= (2\pi)^{-n} \int e^{i(x-z)\cdot\eta} \left((2\pi)^{-n} \int e^{-iy\cdot\xi} a(x, \eta + \xi) b(x + y, \eta) dy d\xi \right) \varphi(z) dz d\eta \end{aligned}$$

$$= (2\pi)^{-n} \int e^{i(x-z)\cdot\eta} c(x, \eta) \varphi(z) dz d\eta, \quad (5.1)$$

where c is defined as the oscillatory integral

$$c(x, \eta) := (2\pi)^{-n} \int e^{-iy\cdot\xi} a(x, \eta + \xi) b(x + y, \eta) dy d\xi. \quad (5.2)$$

If we could show $c \in S^m$ for certain m , then it implies the composition of Ψ DOs is still a Ψ DO. We use the stationary phase lemma under oscillatory integrals developed in §4 to show this expectation.

To show $c \in S^m$, the task boils down to show the asymptotics of c and its derivatives w.r.t. $|\eta|$, thus we set $\lambda := \langle \eta \rangle$, so

$$c(x, \eta) = (2\pi)^{-n} \lambda^n \int e^{-i\lambda y \cdot \xi} a(x, \lambda(\tilde{\eta} + \xi)) b(x + y, \eta) dy d\xi, \quad \text{where } \tilde{\eta} := \eta / \langle \eta \rangle.$$

To make better correspondence with the notations in §4, we set

$$c_{x,\eta}(y, \xi) := a(x, \lambda(\tilde{\eta} + \xi)) b(x + y, \eta),$$

thus

$$c(x, \eta) = (2\pi)^{-n} \lambda^n \int_{\mathbb{R}^{2n}} e^{i\lambda \langle Q(y, \xi), (y, \xi) \rangle / 2} c_{x,\eta}(y, \xi) d(y, \xi), \quad (5.3)$$

where (y, ξ) is treated as a $2n$ -dim vertical vector and

$$Q = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix} \quad (\Rightarrow Q^{-1} = Q, \text{sgn } Q = 0, \text{ and } \det Q = \pm 1).$$

In $c_{x,\eta}(y, \xi)$, we regard (x, η) as irrelevant parameters make the following correspondence:

	function	variable	fixed point	in total	dimension
In Prop. 4.8	a	x	x_0	$a(x - x_0)$	n
at here	$c_{x,\eta}$	(y, ξ)	$(y_0, \xi_0) = 0$	$c_{x,\eta}(y, \xi)$	$2n$

To use Proposition 4.8, the only thing left to check is (4.9), namely, to check

$$\forall \alpha, \beta : |\alpha| + |\beta| \leq 2n + 2N + 3, \quad |\partial_y^\alpha \partial_\eta^\beta (c_{x,\eta}(y, \xi))| \lesssim C_{N,n,\alpha,\beta}(\lambda) \langle (y, \xi) \rangle^{2N+2}. \quad (5.4)$$

For $|\xi| \geq 2$, we have

$$\begin{aligned} |\partial_y^\alpha \partial_\xi^\beta (c_{x,\eta}(y, \xi))| &= |\partial_y^\alpha \partial_\xi^\beta [a(x, \lambda(\tilde{\eta} + \xi)) b(x + y, \eta)]| \\ &\leq C_{\alpha,\beta} \lambda^{|\beta|} |\partial_\xi^\beta a(x, \lambda(\tilde{\eta} + \xi))| \cdot |\partial_x^\alpha b(x + y, \eta)| \\ &\leq C_{\alpha,\beta} \lambda^{|\beta|} \langle \lambda(\tilde{\eta} + \xi) \rangle^{m_1 - |\beta|} \langle \eta \rangle^{m_2}. \end{aligned}$$

Because $|\tilde{\eta}| < 1$, when $|\xi| \geq 2$ we can have $|\tilde{\eta} + \xi| \geq 1$. Recall that $\lambda \geq 1$. Hence when $|\xi| \geq 2$, we can use Lemma 5.1 to continue the computation as follows,

$$\begin{aligned} |\partial_y^\alpha \partial_\xi^\beta (c_{x,\eta}(y, \xi))| &\leq C_{\alpha,\beta} \lambda^{|\beta|} \langle \lambda \rangle^{m_1 - |\beta|} \langle \tilde{\eta} + \xi \rangle^{m_1 - |\beta|} \langle \eta \rangle^{m_2} \\ &\leq C_{\alpha,\beta} \lambda^{m_1} \langle \tilde{\eta} + \xi \rangle^{m_1 - |\beta|} \lambda^{m_2} \quad (\lambda = \langle \eta \rangle \Rightarrow \lambda \simeq \langle \lambda \rangle) \\ &\leq C_{\alpha,\beta} \lambda^{m_1 + m_2} \langle \xi \rangle^{m_1 - |\beta|} \langle \tilde{\eta} \rangle^{|m_1 - |\beta||} \quad (\text{Lemma 5.2}) \\ &\leq C_{\alpha,\beta} \lambda^{m_1 + m_2} \langle \xi \rangle^{m_1 - |\beta|}. \end{aligned} \quad (5.5)$$

We emphasize that (5.5) holds when $|\xi| \geq 2$, and the constant $C_{\alpha,\beta}$ is uniform for x, y, η . Then, due to the continuity, (5.5) actually holds for all ξ . Hence, the condition (5.4) is satisfied when $2N + 2 > m_1$, with $C_{N,n,\alpha,\beta}(\lambda) = C_{\alpha,\beta} \lambda^{m_1 + m_2}$, so we can use Proposition 4.8 directly on (5.3) to obtain

$$c(x, \eta) = (2\pi)^{-n} \lambda^n \int_{\mathbb{R}^{2n}} e^{i\lambda \langle Q(y, \xi), (y, \xi) \rangle / 2} c_{x,\eta}(y, \xi) d(y, \xi)$$

$$\begin{aligned}
&= (2\pi)^{-n} \lambda^n \times \left(\frac{2\pi}{\lambda} \right)^n \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1} D_{(y,\xi)}, D_{(y,\xi)} \rangle}{2i} \right)^j c_{x,\eta}(0,0) \\
&\quad + \lambda^n \times \mathcal{O}(\lambda^{-n-N-1}) \sum_{|\alpha|+|\beta| \leq 2n+2N+3} \sup_{(y,\xi) \in \mathbb{R}^{2n}} \frac{|\partial_y^\alpha \partial_\xi^\beta (c_{x,\eta}(y,\xi))|}{\langle (y,\xi) \rangle^{2n+4N+5-|\alpha|-|\beta|}} \\
&= \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} (D_y \cdot \nabla_\xi)^j c_{x,\eta}(0,0) + \mathcal{O}(\lambda^{-N-1+m_1+m_2}) \\
&= \sum_{|\alpha| \leq N} \frac{\lambda^{-|\alpha|}}{\alpha!} D_y^\alpha \partial_\xi^\alpha (c_{x,\eta}(y,\xi))|_{(y,\xi)=(0,0)} + \mathcal{O}(\lambda^{-N-1+m_1+m_2}) \tag{5.6}
\end{aligned}$$

$$\begin{aligned}
&= \sum_{|\alpha| \leq N} \frac{\lambda^{-|\alpha|}}{\alpha!} \lambda^{|\alpha|} \partial_\eta^\alpha a(x, \lambda \tilde{\eta}) D_x^\alpha b(x, \eta) + \mathcal{O}(\lambda^{-N-1+m_1+m_2}) \\
&= \sum_{|\alpha| \leq N} \frac{1}{\alpha!} \partial_\eta^\alpha a(x, \eta) D_x^\alpha b(x, \eta) + \mathcal{O}(\lambda^{-N-1+m_1+m_2}). \tag{5.7}
\end{aligned}$$

In (5.6) we used

$$\boxed{(D_y \cdot \nabla_\xi)^j = (D_{y_1} \partial_{\xi_1} + \cdots + D_{y_n} \partial_{\xi_n})^j = \sum_{|\alpha|=j} \frac{j!}{\alpha!} D_y^\alpha \partial_\xi^\alpha.} \tag{5.8}$$

By letting N to be large enough, (5.7) implies the following inequality

$$|\partial_x^\alpha \partial_\eta^\beta c(x, \eta)| \lesssim \langle \eta \rangle^{m_1+m_2-|\beta|} \tag{5.9}$$

holds when $|\alpha| = |\beta| = 0$. To show the case when α and/or β are nonzero, we compute

$$\begin{aligned}
\partial_x^\alpha \partial_\eta^\beta c(x, \eta) &= (2\pi)^{-n} \lambda^n \partial_x^\alpha \partial_\eta^\beta \int e^{-i\lambda y \cdot \xi} a(x, \lambda(\tilde{\eta} + \xi)) b(x+y, \eta) dy d\xi \\
&\simeq \lambda^n \sum_{\alpha, \beta} \partial_\eta^\beta \int e^{-i\lambda y \cdot \xi} \partial_x^{\alpha'} \partial_\eta^{\beta'} a(x, \lambda(\tilde{\eta} + \xi)) \partial_x^{\alpha''} \partial_\eta^{\beta''} b(x+y, \eta) dy d\xi. \tag{5.10}
\end{aligned}$$

Note that $\lambda \tilde{\eta} = \eta$. Then we repeat the long computation (with the help of Proposition 4.8) as in (5.7), and this can give (5.9) for all nonzero α and β . The rigorous computation is left as an exercise. Therefore, $c \in S^{m_1+m_2}$.

By letting N to be large enough, (5.7) implies

$$c(x, \eta) \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\eta^\alpha a(x, \eta) D_x^\alpha b(x, \eta),$$

We proved the following result:

THEOREM 5.3. *Assume $m_1, m_2 \in \mathbb{R}$, $a \in S^{m_1}$ and $b \in S^{m_2}$. Then $T_a \circ T_b \in \Psi^{m_1+m_2}$. Denote the symbol of $T_a \circ T_b$ as $\boxed{a \# b}$, then $a \# b \in S^{m_1+m_2}$ and*

$$\boxed{a \# b(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} \partial_\eta^\alpha (a(x, \eta))|_{\eta=\xi} D_y^\alpha (b(y, \xi))|_{y=x}.}$$

REMARK 5.4. We deliberately write $\partial_\eta^\alpha (a(x, \eta))|_{\eta=\xi}$ instead of $\partial_\xi^\alpha a(x, \xi)$, to avoid possible computation mistakes. The same for b .

REMARK 5.5. When symbol a is of the form $a(x, \xi) = \sum_{|\alpha| \leq m_1} d_\alpha(x) \xi^\alpha$ where $d_\alpha \in C^\infty$ are all bounded, or when symbol $b(x, \xi)$ is independent of x -variable, the asymptotics in Theorem (5.3) stops in finite term and the asymptotic is “exact”: we can replace ‘ \sim ’ by ‘ $=$ ’. This can be seen from the expression (5.2) of $c(x, \eta)$. See also Exercise 5.3 and [21, Remark 2.6.9].

From Theorem 5.3 we know, if $a \in S^{m_1}$ and $b \in S^{m_2}$, then

$$a \# b = ab + S^{m_1+m_2-1} = ab + \frac{1}{i} \nabla_\xi a \cdot \nabla_x b + S^{m_1+m_2-2} = ab + \frac{1}{i} \{a, b\} + S^{m_1+m_2-2}. \quad (5.11)$$

5.2. Reduction of variables

As we have seen in (2.5) that

$$(T_\sigma u, \varphi) = (u, (2\pi)^{-n} \int e^{i(y-x) \cdot \xi} \bar{\sigma}(x, \xi) \varphi(x) dx d\xi).$$

In practice we may encounter Ψ DOs of the form

$$\int e^{i(x-y) \cdot \xi} a(x, y, \xi) \varphi(y) dy d\xi$$

where the symbol a depends not only on x but also on y , e.g. in §5.3 we shall see Ψ DOs possessing this type of symbols. We have the following result.

THEOREM 5.6. *Assume $a \in S^m(\mathbb{R}_x^n \times \mathbb{R}_y^n \times \mathbb{R}_\xi^n)$, then there exists symbol $a' \in S^m(\mathbb{R}_x^n \times \mathbb{R}_\xi^n)$ such that*

$$T_{a'} \varphi(x) = (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} a(x, y, \xi) \varphi(y) dy d\xi, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n), \quad (5.12)$$

and this $T_{a'}$ takes the following as its kernel:

$$K(x, y) := (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} a(x, y, \xi) d\xi.$$

Moreover, a' has the asymptotics

$$\boxed{a'(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} D_y^\alpha \partial_\eta^\alpha (a(x, y, \eta))|_{(y, \eta)=(x, \xi)}}.$$

If (5.12) holds, we will have

$$\int e^{i(x-y) \cdot \xi} a'(x, \xi) \varphi(y) dy d\xi = \int e^{i(x-y) \cdot \xi} a(x, y, \xi) \varphi(y) dy d\xi$$

and so we can expect

$$\int e^{i(x-y) \cdot \xi} a'(x, \xi) d\xi = \int e^{i(x-y) \cdot \xi} a(x, y, \xi) d\xi$$

to hold in the oscillatory integral sense. By changing y to $y+x$, we see the LHS is a Fourier transform,

$$\mathcal{F}_\xi \{a'(x, \xi)\}(y) = (2\pi)^{-n/2} \int e^{-iy \cdot \xi} a(x, y+x, \xi) d\xi,$$

so

$$\begin{aligned} a'(x, \eta) &= (2\pi)^{-n} \int e^{iy \cdot \eta} dy \cdot \int e^{-iy \cdot \xi} a(x, y+x, \xi) d\xi \\ &= (2\pi)^{-n} \int e^{-iy \cdot (\xi - \eta)} a(x, y+x, \xi) dy d\xi \end{aligned}$$

$$\begin{aligned}
&= (2\pi)^{-n} \int e^{-iy \cdot \xi} a(x, y + x, \xi + \eta) dy d\xi \\
&= (2\pi)^{-n} \lambda^n \int e^{-i\lambda y \cdot \xi} a(x, y + x, \lambda(\xi + \tilde{\eta})) dy d\xi,
\end{aligned}$$

where again $\lambda := \langle \eta \rangle$ and $\tilde{\eta} := \eta / \langle \eta \rangle$. The rigorous proof we go by first set a' as in this way, and then prove a' is a symbol of order m .

PROOF OF THEOREM 5.6. We set

$$a'(x, \eta) = (2\pi)^{-n} \lambda^n \int e^{-i\lambda y \cdot \xi} a(x, y + x, \lambda(\xi + \tilde{\eta})) dy d\xi,$$

where $\lambda := \langle \eta \rangle$ and $\tilde{\eta} := \eta / \langle \eta \rangle$. Following the arguments preceding this proof, we can show that a' satisfies (5.12). It's left to show a' satisfies the asymptotics, which will automatically show $a' \in S^m$.

To show a' satisfies the asymptotics, we use the stationary phase lemma in a similar manner as in §5.1. We set

$$a_{x,\eta}(y, \xi) := a(x, x + y, \lambda(\xi + \tilde{\eta})),$$

thus

$$a'(x, \eta) = (2\pi)^{-n} \lambda^n \int_{\mathbb{R}^{2n}} e^{i\lambda \langle Q(y,\xi), (y,\xi) \rangle / 2} a_{x,\eta}(y, \xi) d(y, \xi), \quad (5.13)$$

where y and ξ is treated as horizontal vector and

$$Q = \begin{pmatrix} 0 & -I \\ -I & 0 \end{pmatrix} \quad (\Rightarrow Q^{-1} = Q, \text{sgn } Q = 0, \text{ and } \det Q = \pm 1).$$

For $|\xi| \geq 2$, we have

$$\begin{aligned}
|\partial_y^\alpha \partial_\xi^\beta (a_{x,\eta}(y, \xi))| &= |\partial_y^\alpha \partial_\xi^\beta [a(x, x + y, \lambda(\xi + \tilde{\eta}))]| \\
&\leq C_{\alpha,\beta} \lambda^{|\beta|} |(\partial_y^\alpha \partial_\xi^\beta a)(x, x + y, \lambda(\xi + \tilde{\eta}))| \\
&\leq C_{\alpha,\beta} \lambda^{|\beta|} \langle \lambda(\xi + \tilde{\eta}) \rangle^{m-|\beta|} \\
&\leq C_{\alpha,\beta} \lambda^{|\beta|} \langle \lambda \rangle^{m-|\beta|} \langle \xi + \tilde{\eta} \rangle^{m-|\beta|} \quad (\text{Lemma 5.1}) \\
&\lesssim C_{\alpha,\beta} \lambda^m \langle \xi \rangle^{m-|\beta|} \langle \tilde{\eta} \rangle^{|m-|\beta||} \quad (\lambda \simeq \langle \lambda \rangle, \text{ Lemma 5.2}) \\
&\leq C_{\alpha,\beta} \lambda^m \langle \xi \rangle^{m-|\beta|}.
\end{aligned}$$

Hence, the condition (4.9) is satisfied when $2N + 2 > m$, with $C_{N,n,\alpha,\beta}(\lambda) = C_{\alpha,\beta} \lambda^m$, so we can use Proposition 4.8 directly on (5.13) to obtain

$$\begin{aligned}
a'(x, \eta) &= (2\pi)^{-n} \lambda^n \int_{\mathbb{R}^{2n}} e^{i\lambda \langle Q(y,\xi), (y,\xi) \rangle / 2} c_{x,\eta}(y, \xi) d(y, \xi) \\
&= (2\pi)^{-n} \lambda^n \times \left(\frac{2\pi}{\lambda} \right)^n \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} \left(\frac{\langle Q^{-1} D_{(y,\xi)}, D_{(y,\xi)} \rangle}{2i} \right)^j c_{x,\eta}(0, 0) \\
&\quad + \lambda^n \times \mathcal{O}(\lambda^{-n-N-1} \sum_{|\alpha|+|\beta| \leq 2n+2N+3} \sup_{(y,\xi) \in \mathbb{R}^{2n}} \frac{|\partial_y^\alpha \partial_\xi^\beta (c_{x,\eta}(y, \xi))|}{\langle (y, \xi) \rangle^{2n+4N+5-|\alpha|-|\beta|}}) \\
&= \sum_{0 \leq j \leq N} \frac{\lambda^{-j}}{j!} (D_y \cdot \nabla_\xi)^j c_{x,\eta}(0, 0) + \mathcal{O}(\lambda^{-N-1+m}) \\
&= \sum_{|\alpha| \leq N} \frac{\lambda^{-|\alpha|}}{\alpha!} D_y^\alpha \partial_\xi^\alpha (c_{x,\eta}(y, \xi))|_{(y,\xi)=(0,0)} + \mathcal{O}(\lambda^{-N-1+m})
\end{aligned}$$

$$\begin{aligned}
&= \sum_{|\alpha| \leq N} \frac{\lambda^{-|\alpha|}}{\alpha!} D_y^\alpha \partial_\xi^\alpha (a(x, y+x, \lambda\xi + \eta))|_{(y,\xi)=(0,0)} + \mathcal{O}(\lambda^{-N-1+m}) \\
&= \sum_{|\alpha| \leq N} \frac{1}{\alpha!} D_y^\alpha \partial_\xi^\alpha (a(x, y+x, \xi + \eta))|_{(y,\xi)=(0,0)} + \mathcal{O}(\lambda^{-N-1+m}) \\
&= \sum_{|\alpha| \leq N} \frac{1}{\alpha!} D_y^\alpha \partial_\xi^\alpha (a(x, y, \xi))|_{(y,\xi)=(x,\eta)} + \mathcal{O}(\lambda^{-N-1+m}). \tag{5.14}
\end{aligned}$$

Due to the same logic as in (5.9)-(5.10), we can let N to be large enough, and by doing so, (5.14) can implies $c \in S^m$ and

$$c(x, \eta) \sim \sum_{\alpha} \frac{1}{\alpha!} D_y^\alpha \partial_\xi^\alpha (a(x, y, \xi))|_{(y,\xi)=(x,\eta)},$$

The proof is complete. \square

Theorem 5.6 completes the proof of Lemma 2.22.

5.3. The Adjoint and transpose

We define the adjoint and transpose of the Ψ DO T_a acting on Schwartz functions as follows,

$$\begin{aligned}
\text{adjoint } T_a^* &: (T_a^* u, v) := (u, T_a v), \\
\text{transpose } {}^t T_a &: \langle {}^t T_a u, v \rangle := \langle u, T_a v \rangle,
\end{aligned} \tag{5.15}$$

where $u, v \in \mathcal{S}$.

THEOREM 5.7. *Assume $a(x, \xi) \in S^m$. The T_a^* and ${}^t T_a$ defined in (5.15) exist uniquely, and both are Ψ DOs. There exist symbols a^* and ${}^t a$ of the same order as a such that $T_a^* = T_{a^*}$ and ${}^t T_a = T_{{}^t a}$. Moreover, we have the asymptotics*

$$\begin{aligned}
&\boxed{a^*(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} D_x^\alpha \partial_\xi^\alpha \bar{a}(x, \xi),} \\
&\boxed{{}^t a(x, \xi) \sim \sum_{\alpha} \frac{(-1)^{|\alpha|}}{\alpha!} D_x^\alpha \partial_\xi^\alpha a(x, -\xi).}
\end{aligned}$$

REMARK 5.8. The computation (2.5) gives an very efficient intuitive way to compute the asymptotics of a^* .

PROOF. Here we only show the proof for a^* , and that of ${}^t a$ is left as an exercise.

Step 1. Existence. As explained at the beginning of §5.2, for $u, v \in \mathcal{S}$ we have

$$(u, T_a v) = ((2\pi)^{-n} \int e^{i(y-x)\cdot\xi} \bar{a}(x, \xi) u(x) dx d\xi, v)$$

so if we define a mapping T as

$$Tu(y) := (2\pi)^{-n} \int e^{i(y-x)\cdot\xi} \bar{a}(x, \xi) u(x) dx d\xi,$$

then $(Tu, v) := (u, T_a v)$. Also, this T is of the form (5.12), so by Theorem 5.6 we know T is a Ψ DO.

Step 2. Uniqueness. Assume there are two adjoint of T , and we denote them as T_1 and T_2 , respectively. Then for any $u, v \in \mathcal{S}$ we can conclude

$$(T_1 u, v) = (u, T_a v) = (T_2 u, v) \Rightarrow ((T_1 - T_2)u, v) = 0.$$

Hence, $(T_1 - T_2)u = 0$ for any $u \in \mathcal{S}$ and so $T_1 = T_2$.

Step 3. Asymptotics. Theorem 5.6 suggests that the symbol of T , denoted as a^* , satisfies the asymptotics:

$$a^*(x, \xi) \sim \sum_{\alpha} \frac{1}{\alpha!} D_y^\alpha \partial_\eta^\alpha (\bar{a}(y, \eta))|_{(y, \eta)=(x, \xi)} = \sum_{\alpha} \frac{1}{\alpha!} D_x^\alpha \partial_\xi^\alpha \bar{a}(x, \xi).$$

The proof is complete. \square

Exercise

EXERCISE 5.1. Use stationary phase lemmas to complete the estimate in $\partial_x^\alpha \partial_\eta^\beta c(x, \eta)$ in (5.10). Hint: mimic the computations in (5.7).

EXERCISE 5.2. Assume $a \in S^{m_1}$ and $b \in S^{m_2}$. Utilize Theorem 5.3 to show that $[T_a, T_b] \in \Psi^{m_1+m_2-1}$, where $[T_a, T_b] := T_a T_b - T_b T_a$ is called the *commutator* of T_a and T_b , and $T_a T_b$ is a shorthand of the composition $T_a \circ T_b$.

EXERCISE 5.3. Prove the statement in Remark 5.5. In Theorem 5.3, assume $a(x, \xi) = \sum_{|\alpha| \leq m_1} d_\alpha(x) \xi^\alpha$ where $d_\alpha \in C^\infty$ are all bounded, or assume $b = b(\xi)$, then show that

$$c(x, \eta) = \sum_{|\alpha| \leq N} \frac{1}{\alpha!} \partial_\eta^\alpha a(x, \eta) D_x^\alpha b(x, \eta)$$

for some finite integer N . Hint: substitute the expressions of a or b into (5.2) and use Lemma 3.22.

EXERCISE 5.4. Mimic the proof for a^* in Theorem 5.7 to prove the result for ${}^t a$.

EXERCISE 5.5. Let T_1, T_2 be two Ψ DOs. Show that $(T_1^*)^* = T_1$ and $(T_1 T_2)^* = T_2^* T_1^*$. Here “ T^* ” stands for taking the adjoint of T .

Parametrix and Boundedness of Ψ DOs

In this chapter we investigate the parametrix and boundedness of Ψ DOs, both of which heavily utilize the symbolic calculus. The notion of parametrix can be understood as the approximate inverse, or the inverse modulo C^∞ an operator. For a homogeneous polynomial $T(\xi) := \sum_{|\alpha|=m} a_\alpha \xi^\alpha$, its corresponding operator $T := T(D)$ is a Ψ DO.

To find the inverse, a typical idea is to design $S(\xi) := 1/T(\xi)$ and let $S := S(D)$. Inaccurately this seems to give us $ST = I$ where I is the identity operator, which is (inaccurately) because by Theorem 5.3 (and Remark 5.5) we have

$$\text{symbol of } ST = \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} (T(\xi)) D_x^{\alpha} (S(\xi)) = T(\xi) S(\xi) = 1.$$

Unfortunately, this is wrong, because $1/T(\xi)$ has singularities when $T(\xi) = 0$. And due to this reason, S may not be a Ψ DO so Theorem 5.3 is not applicable here.

However, the S can be saved if we cutoff the singularity. Specifically, fix a $\chi \in C_c^\infty$ with $\chi(0) = 1$ and we re-design S as $S(\xi) := (1 - \chi(\xi))/T(\xi)$ and once again let $S := S(D)$. It is straightforward that this new $S(\xi)$ is a symbol and so S is a Ψ DO. Again, by Theorem 5.3 (and Remark 5.5) we have

$$\begin{aligned} \text{symbol of } ST &= \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} (T(\xi)) D_x^{\alpha} (S(\xi)) = T(\xi) S(\xi) \\ &= T(\xi) (1 - \chi(\xi)) / T(\xi) \\ &= 1 - \chi(\xi). \end{aligned}$$

It is also true that the symbol of $TS = 1 - \chi(\xi)$. Note that $\chi(D) \in \Psi^{-\infty}$, so we conclude

$$ST = I + \Psi^{-\infty}, \quad TS = I + \Psi^{-\infty}.$$

This inspires us to introduce the notion of parametrix.

6.1. Parametrix

In what follows we use I to signify the identity operator unless otherwise stated.

DEFINITION 6.1 (Parametrix). Assume $m \in \mathbb{R}$ and $T \in \Psi^m$. If there exists a Ψ DO S such that $ST - I \in \Psi^{-\infty}$, we call S a *left parametrix* of T . If $TS - I \in \Psi^{-\infty}$, we call S a *right parametrix* of T . We call S a *parametrix* of T if it is both a left and a right parametrix.

The notion of left and right parametrix is somewhat redundant.

LEMMA 6.2. *Assume both S and T both Ψ DOs. If S is a left (right) parametrix of T , and T has a right (left) parametrix, then S is also a right (left) parametrix of T .*

PROOF. We only prove the left-case. There exists S' such that $TS' = I + \Psi^{-\infty}$. From $ST = I + \Psi^{-\infty}$ we have $(ST)S' = S' + \Psi^{-\infty} = S(TS')$, so $S' + \Psi^{-\infty} = S(I + \Psi^{-\infty})$, which gives $S = S' + \Psi^{-\infty}$. Therefore,

$$TS = T(S' + \Psi^{-\infty}) = TS' + \Psi^{-\infty} = I + \Psi^{-\infty} + \Psi^{-\infty} = I + \Psi^{-\infty},$$

which implies S is a right parametrix of T . □

The parametrix of a Ψ DO is not always exists. And in contrast with the notion of inverse of an operator, when parametrices exist, they are not unique.

LEMMA 6.3. *Assume S is a parametrix of T , and $R \in \Psi^{-\infty}$, then $S + R$ is also a parametrix of T .*

The proof is left as an exercise. One of the condition that guarantees the existence of parametrix is the ellipticity.

DEFINITION 6.4 (Ellipticity). Assume $m \in \mathbb{R}$ and $a \in S^m$. We call a and also its corresponding Ψ DO T_a *elliptic* when there exist fixed positive constants C and R such that

$$\boxed{|a(x, \xi)| \geq C\langle \xi \rangle^m, \quad \text{when } x \in \mathbb{R}^n, |\xi| \geq R.}$$

There is an equivalent definition for the ellipticity of a symbol.

LEMMA 6.5. *Assume $m \in \mathbb{R}$ and $a \in S^m$. The ellipticity condition for a is equivalent to the fact that there exist two positive constants C and D such that*

$$\boxed{|a(x, \xi)| \geq C\langle \xi \rangle^m - D\langle \xi \rangle^{m-1}, \quad \forall x, \xi \in \mathbb{R}^n.} \quad (6.1)$$

PROOF. Assume $a \in S^m$ is elliptic, then there are constants $C, R > 0$ such that

$$|a(x, \xi)|/\langle \xi \rangle^m \geq C, \quad \forall |\xi| \geq R,$$

so for any positive constant D we have

$$|a(x, \xi)|/\langle \xi \rangle^m \geq C - D\langle \xi \rangle^{-1}, \quad (6.2)$$

for $\forall |\xi| \geq R$. If we set $D := C\langle R \rangle$, then

$$\forall |\xi| \leq R, \quad C\langle \xi \rangle \leq D \Rightarrow C - D\langle \xi \rangle^{-1} \leq 0,$$

so (6.2) holds for both $|\xi| \geq R$ and $|\xi| \leq R$. This gives (6.1).

On the other hand, from (6.1) it is easy to see a is elliptic. \square

We will show that

$$\boxed{\text{Ellipticity} \Leftrightarrow \exists \text{ parametrix.}}$$

First, we show the ellipticity condition gives the existence of parametrices.

THEOREM 6.6 (Ellipticity \Rightarrow parametrix). *Assume $m \in \mathbb{R}$ and $a \in S^m$ and a is elliptic, then T_a has a parametrix.*

PROOF. Here we use the notation $\sigma(T)$ to represent the symbol of a Ψ DO T , the well-definedness of the mapping σ is guaranteed by Lemma 2.11. We denote T_a as A for simplicity. Fix a cutoff function $\chi \in C_c^\infty(\mathbb{R}^n)$ such that $\chi(\xi) = 1$ when $|\xi| \leq R$ and $\chi(\xi) = 0$ when $|\xi| \geq R + 1$, where the R is given in Definition 6.4.

Step 1. Define $b_0(x, \xi) := (1 - \chi(\xi))/a(x, \xi)$ and $B_0 := T_{b_0}$, then b_0 is well-defined because the denominator is nonzero in the support of $1 - \chi$. Also, it can be checked that b_0 is a symbol of order $-m$ (see Exercise 6.2). Then according to Theorem 5.3, we have

$$\sigma(AB_0) = a(1 - \chi)/a - r_1 = 1 - \chi - r_1, \quad \text{for some } r_1 \in S^{-1}.$$

Step 2. Define $b_1(x, \xi) := (1 - \chi(\xi))/a(x, \xi) \cdot r_1(x, \xi) \in S^{-m-1}$ and $B_1 := T_{b_1}$. Again, according to Theorem 5.3, we have

$$\begin{aligned} \sigma(A(B_0 + B_1)) &= \sigma(AB_0) + \sigma(AB_1) = 1 - \chi - r_1 + a(1 - \chi)/ar_1 - r_2 \\ &= 1 - (1 + r_1)\chi - r_2, \quad \text{for some } r_2 \in S^{-2}. \end{aligned}$$

Step 3. Define recursively $b_j(x, \xi) := (1 - \chi(\xi))/a(x, \xi) \cdot r_j(x, \xi) \in S^{-m-j}$ and $B_j := T_{b_j}$. According to Theorem 5.3, we have

$$\begin{aligned} \sigma(A(B_0 + \cdots + B_j)) &= \sigma(A(B_0 + \cdots + B_{j-1})) + \sigma(AB_j) \\ &= [1 - (1 + r_1 + \cdots + r_{j-1})\chi - r_j] + a(1 - \chi)/ar_j - r_{j+1} \\ &= 1 - (1 + r_1 + \cdots + r_j)\chi - r_{j+1}, \quad \text{for some } r_{j+1} \in S^{-j-1}. \end{aligned}$$

Step 4. According to Theorem 2.6, there exists $b \in S^{-m}$ such that $b \sim \sum_j b_j$. Denote $B = T_b$, so for any $N \in \mathbb{N}$ there holds $B = B_0 + \cdots + B_N + \Psi^{-m-N-1}$. Hence we can compute the symbol of AB as follows,

$$\begin{aligned} \sigma(AB) &= \sigma(A(B_0 + \cdots + B_N + \Psi^{-m-N-1})) \\ &= \sigma(A(B_0 + \cdots + B_N)) + \sigma(A\Psi^{-m-N-1}) \\ &= 1 - (1 + r_1 + \cdots + r_N)\chi - r_{N+1} + S^{-N-1} = 1 + S^{-N-1}, \end{aligned} \quad (6.3)$$

where the last equal sign is due to $\chi \in S^{-\infty}$ and $r_{N+1} \in S^{-N-1}$. Due to the arbitrariness of N , (6.3) implies that

$$AB - I \in \Psi^{-\infty},$$

so B is right parametrix of A . By repeating steps 1-4 we can also show A has a right parametrix, so by Lemma 6.2 we conclude that B is a parametrix of A . \square

Second, we show the existence of parametrices gives the ellipticity.

THEOREM 6.7 (Parametrix \Rightarrow ellipticity). *Assume $m \in \mathbb{R}$ and $a \in S^m$ and T_a has either a right parametrix or a left parametrix, then T_a is elliptic.*

PROOF. Assume T_b is the right parametrix, then b is necessarily a symbol of order $-m$, so

$$\sigma(T_a T_b) = ab + S^{-1}, \quad \text{and} \quad \sigma(T_a T_b) = \sigma(I + \Psi^{-\infty}) = 1 + S^{-\infty},$$

thus

$$ab = 1 + S^{-1} + S^{-\infty} = 1 + S^{-1}.$$

Therefore, when $|\xi|$ is large enough

$$\forall (x, \xi) \in \mathbb{R}^{2n}, \quad |a(x, \xi)b(x, \xi) - 1| \leq C\langle \xi \rangle^{-1}.$$

Therefore, $\langle \xi \rangle \geq C/2$ is large enough, we can conclude

$$|a(x, \xi)b(x, \xi)| \geq 1/2 \quad \Rightarrow \quad |a(x, \xi)| \geq 1/(2|b(x, \xi)|).$$

This gives

$$|a(x, \xi)| \geq \langle \xi \rangle^m / 2 \quad \text{when} \quad \langle \xi \rangle \geq C/2,$$

so a is elliptic.

The proof for the left-case is similar. \square

From Theorems 6.6 & 6.7, we see that the condition “ T has a right (left) parametrix” in Lemma 6.2 can be lifted.

PROPOSITION 6.8. *Assume both S and T are Ψ DOs. If S is a left (right) parametrix of T , then S is also a right (left) parametrix of T .*

PROOF. If S is a left (right) parametrix of T , then by Theorem 6.7 we know that T is elliptic, so by Theorem 6.6 we know T has a right (left) parametrix. Then Lemma 6.2 tells us S is a right (left) parametrix of T . \square

We recall that when T is a Ψ DO, T doesn't increase the singular support of a distribution (see Theorem 2.24). Now if we know T is also elliptic, then T doesn't decrease the singular support.

LEMMA 6.9. *Assume T is an elliptic Ψ DO and $u \in \mathcal{E}'$, then*

$$\boxed{\text{sing supp}(Tu) = \text{sing supp } u.}$$

Readers may compare Lemma 6.9 with Theorem 2.24.

PROOF. Denote $Tu = f$, then Theorem 2.24 implies

$$\text{sing supp}(Tu) \subset \text{sing supp } u.$$

Theorem 6.6 implies T possesses parametrices. Let S be a parametrix of T . Then we have $Sf = STu = (I + \Psi^{-\infty})u = u + C^\infty(\mathbb{R}^n)$, so

$$\text{sing supp } u = \text{sing supp}(Sf) \subset \text{sing supp } f = \text{sing supp}(Tu).$$

The proof is done. \square

We will revisit the notion of parametrix and ellipticity in §9.1.

6.2. The L^2 boundedness

LEMMA 6.10 (Schur estimate). *Assume $K \in L^1_{loc}(\mathbb{R}^{2n})$ and for $\varphi \in L^1_{loc}(\mathbb{R}^n)$ we denote $T\varphi(x) := \int_{\mathbb{R}^n} K(x, y)\varphi(y) dy$. Also, denote*

$$L := \sup_{x \in \mathbb{R}^n} \int_{\mathbb{R}^n} |K(x, y)| dy, \quad R := \sup_{y \in \mathbb{R}^n} \int_{\mathbb{R}^n} |K(x, y)| dx. \quad (6.4)$$

When $L, R < +\infty$, for $\forall p \in [1, +\infty]$ and $\varphi \in L^p(\mathbb{R}^n)$ we have

$$\|T\varphi\|_{L^p} \leq L^{1-1/p} R^{1/p} \|\varphi\|_{L^p}.$$

PROOF. When $p = +\infty$ is trivial, we have

$$\begin{aligned} \|T\varphi\|_{L^\infty} &= \text{ess sup}_x \left| \int K(x, y)\varphi(y) dy \right| \leq \text{ess sup}_x \int |K(x, y)| dy \cdot \text{ess sup}_y |\varphi(y)| \\ &= L \|\varphi\|_{L^\infty}. \end{aligned}$$

When $p = 1$, we have

$$\begin{aligned} \|T\varphi\|_{L^1} &= \left\| \int K(x, y)\varphi(y) dy \right\|_{L^1} \leq \int \|K(\cdot, y)\|_{L^1} |\varphi(y)| dy \\ &\leq R \int |\varphi(y)| dy = R \|\varphi\|_{L^1}. \end{aligned}$$

Now we assume $1 < p < +\infty$. Let $p' = p/(p-1)$, so $1 = 1/p + 1/p'$. We have

$$\begin{aligned} |T\varphi(x)| &\leq \int |K(x, y)\varphi(y)| dy = \int |K(x, y)|^{1/p'} |K(x, y)|^{1/p} |\varphi(y)| dy \\ &\leq \left(\int |K(x, y)| dy \right)^{1/p'} \left(\int |K(x, y)| |\varphi(y)|^p dy \right)^{1/p} \quad (\text{by Hölder's ineq.}) \\ &\leq L^{1/p'} \left(\int |K(x, y)| |\varphi(y)|^p dy \right)^{1/p}. \end{aligned}$$

Hence,

$$\begin{aligned} \|T\varphi\|_{L^p} &\leq L^{1/p'} \left(\iint |K(x, y)| |\varphi(y)|^p dy dx \right)^{1/p} \leq L^{1/p'} \left(R \int |\varphi(y)|^p dy \right)^{1/p} \\ &\leq L^{1/p'} R^{1/p} \|\varphi\|_{L^p}. \end{aligned}$$

The proof is complete. \square

As already mentioned in Remark 3.9, when the order m is small enough, T_σ possesses certain types of boundedness.

LEMMA 6.11. *In \mathbb{R}^n , we assume $m < -n$ and $\sigma \in S^m(\mathbb{R}_x^n \times \mathbb{R}_\xi^n)$, then the Ψ DO $T_\sigma: L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ is bounded.*

PROOF. Denote the kernel of T_σ as K , so

$$K(x, y) = (2\pi)^{-n} \int e^{i(x-y)\cdot\xi} \sigma(x, \xi) d\xi.$$

Because $\sigma \in S^m$ with $m < -n$, we know that integral above is absolutely integrable. This means that K is a well-defined function in \mathbb{R}^{2n} , especially, K is well-defined on the diagonal $\{(x, x); x \in \mathbb{R}^n\}$. However, we remind the readers that the condition “ $m < -n$ ” doesn’t guarantee that K is also C^∞ on the diagonal (recall that Lemma 3.11 tells us K is C^∞ off diagonal). The value of K on \mathbb{R}^{2n} is uniformly bounded, because

$$|K(x, y)| \leq (2\pi)^{-n} \int |\sigma(x, \xi)| d\xi \leq (2\pi)^{-n} \int C\langle\xi\rangle^m d\xi \leq C.$$

Because K is well-defined and uniformly bounded on \mathbb{R}^{2n} , we can define the corresponding L and R of it as in (6.4), and we can also enhance the estimate in Lemma 3.11 as follows,

$$|K(x, y)| \leq C\langle x - y \rangle^{-n-1}, \quad \forall x, y \in \mathbb{R}^n,$$

which implies both L and R are finite. Because $T_\sigma\varphi(x) = \int_{\mathbb{R}^n} K(x, y)\varphi(y) dy$, we can use Lemma 6.10 to conclude $\|T_\sigma\varphi\|_{L^p} \lesssim \|\varphi\|_{L^p}$. The proof is complete. \square

THEOREM 6.12 (L^2 boundedness). *Assume symbol $a \in S^0$, then $T_a: L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bounded.*

PROOF. Recall the definition for “ $a\#b$ ” in Theorem 5.3. To prove the result, it amounts to find a suitable positive constant M such that for $\forall\varphi \in \mathcal{S}$,

$$\|T_a\varphi\|_{L^2} \leq M\|\varphi\|_{L^2} \quad \Leftrightarrow \quad ((M - T_a^*T_a)\varphi, \varphi) \geq 0.$$

Our strategy is: we try to find such a M so that $M - T_a^*T_a$ can be represented as B^*B for some B so that

$$((M - T_a^*T_a)\varphi, \varphi) = (B^*B\varphi, \varphi) = (B\varphi, B\varphi) \geq 0.$$

Step 1. Symbolic calculus. Because $a \in S^0$, we know $|a(x, \xi)| \leq C$ uniformly for some C . Let

$$M = M_1 + M_2, \quad \text{where} \quad M_1 := 2 \sup_{\mathbb{R}^{2n}} |a(x, \xi)|^2 + 1, \quad (6.5)$$

and M_2 shall be determined later, and define

$$b(x, \xi) := \sqrt{M_1 - |a(x, \xi)|^2}.$$

It can be checked that $b \in S^0$. We use $\sigma(T)$ to signify the symbol of T . Then by Theorems 5.3 & 5.7 we have

$$\sigma(T_b^*T_b) = |b|^2 + S^{-1} = M_1 - |a(x, \xi)|^2 + S^{-1},$$

and also

$$\sigma(M_1 - T_a^*T_a) = M_1 - \sigma(T_a^*T_a) = M_1 - (|a(x, \xi)|^2 + S^{-1}).$$

Hence

$$\sigma(M - T_a^*T_a) = M_2 + \sigma(M_1 - T_a^*T_a) = M_2 + \sigma(T_b^*T_b) + S^{-1},$$

which implies

$$M - T_a^* T_a = M_2 + T_b^* T_b - R, \quad \text{for some } \Psi\text{DO } R \in \Psi^{-1}.$$

Therefore it is equivalent to prove

$$((M_2 + T_b^* T_b - R)\varphi, \varphi) \geq 0,$$

so we only need to prove

$$(R\varphi, \varphi) \leq M_2 \|\varphi\|_{L^2}^2. \quad (6.6)$$

Step 2. To prove (6.6), we can do the following derivations:

$$\begin{aligned} (R\varphi, \varphi) \leq M_2 \|\varphi\|_{L^2}^2 &\Leftrightarrow |(R\varphi, \varphi)| \leq M_2 \|\varphi\|_{L^2}^2 \\ &\Leftrightarrow \|R\varphi\|_{L^2} \|\varphi\|_{L^2} \leq M_2 \|\varphi\|_{L^2}^2 \\ &\Leftrightarrow \|R\varphi\|_{L^2} \leq M_2 \|\varphi\|_{L^2} \\ &\Leftrightarrow (R^* R\varphi, \varphi) \leq M_2^2 \|\varphi\|_{L^2}^2 \\ &\Leftrightarrow |(R^* R\varphi, \varphi)| \leq M_2^2 \|\varphi\|_{L^2}^2 \\ &\Leftrightarrow \|R^* R\varphi\|_{L^2} \leq M_2^2 \|\varphi\|_{L^2}. \end{aligned}$$

We observe that

$$\|R^* R\varphi\|_{L^2} \leq M_2^2 \|\varphi\|_{L^2} \Rightarrow \|R\varphi\|_{L^2} \leq M_2 \|\varphi\|_{L^2}. \quad (6.7)$$

Step 3. Using (6.7) iteratively, we can obtain $(R^* R)^* R^* R$, $((R^* R)^* R^* R)^* (R^* R)^* R^* R$, etc, and each time the order of the corresponding ΨDO decreases by at least 1. We will end up with a ΨDO of order less than $-n$ in finite time. And by Lemma 6.11, that operator is L^2 -bounded. Then we use (6.7) to bring the boundedness back to R , so we arrive at

$$\|R\varphi\|_{L^2} \leq M_2 \|\varphi\|_{L^2}, \quad \forall \varphi \in \mathcal{S}.$$

This gives (6.6). The proof is complete. \square

As a corollary of Theorem 6.12, we have the following H^m boundedness for any $T \in \Psi^m$.

COROLLARY 6.13 (H^m boundedness). *Assume $T \in \Psi^m$, then for any $s \in \mathbb{R}$, the mapping $T: H^{s+m}(\mathbb{R}^n) \rightarrow H^s(\mathbb{R}^n)$ is bounded.*

PROOF. Denote $J := (I - \Delta)^{1/2}$. Because $T \in \Psi^m$, we have $J^s T J^{-s-m} \in \Psi^0$. Hence for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$ we have

$$\begin{aligned} \|T\varphi\|_{H^s} &= \|J^s T\varphi\|_{L^2} = \|J^s T J^{-s-m} J^{s+m}\varphi\|_{L^2} \\ &\leq C \|J^{s+m}\varphi\|_{L^2} = C \|\varphi\|_{H^{s+m}}. \end{aligned}$$

By a density argument we can extend the result to any $\varphi \in H^{s+m}$. The proof is done. \square

Theorem 6.12 can be generalized to a more general case. The L^2 -boundedness results are given in [2, 3, 12]. Then A. Calderón and R. Vaillancourt generalized their own result [2] in [3]. We comment that [14] gives an elementary proof of the results in [3]. Here we restate the main results in [2, 3] as follows. Recall the symbol space $S_{\rho, \delta}^m$ defined in Definition 2.4.

THEOREM 6.14 (Calderón-Vaillancourt Theorem [2]). *Assume $a \in S_{0,0}^0(\mathbb{R}_x^n \times \mathbb{R}_\xi^n; \mathbb{C})$, then the ΨDO T_a is bounded in $L^2(\mathbb{R}^n)$, and there exist constants C, N such that*

$$\forall \varphi \in L^2(\mathbb{R}^n), \quad \|T_a \varphi\|_{L^2} \leq C \max_{|\alpha+\beta| \leq N} \|\partial_x^\alpha \partial_\xi^\beta a\|_{L^\infty} \|\varphi\|_{L^2}. \quad (6.8)$$

THEOREM 6.15 (Generalized Calderón-Vaillancourt Theorem [3]). *Let $a \in C^\infty(\mathbb{R}_x^n \times \mathbb{R}_y^n \times \mathbb{R}_\xi^n; \mathbb{C})$, and $0 \leq \rho \leq \delta_j < 1$ ($j = 1, 2$) and $M/n \geq \frac{1}{2}(\delta_1 + \delta_2) - \rho$. If there exists a constant \mathcal{C} such that $\forall (x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n$,*

$$|\partial_x^{\alpha_1} \partial_\xi^\beta a(x, y, \xi)| \leq \mathcal{C} \langle \xi \rangle^{-M + \delta_1 |\alpha_1| - \rho |\beta|}, \quad |\partial_y^{\alpha_2} \partial_\xi^\beta a(x, y, \xi)| \leq \mathcal{C} \langle \xi \rangle^{-M + \delta_2 |\alpha_2| - \rho |\beta|}$$

holds for all $0 \leq |\beta| \leq 2\lceil n/2 \rceil + 2$ and $0 \leq |\alpha_j| \leq 2m_j$ ($j = 1, 2$) with m_j being the least integer satisfying $m_j(1 - \delta_j) \geq 5n/4$, then the linear operator T_a defined as

$$T_a \varphi(x) := (2\pi)^{-n} \iint_{\mathbb{R}^n \times \mathbb{R}^n} e^{i(x-y) \cdot \xi} a(x, y, \xi) \varphi(y) \, dy \, d\xi$$

is bounded from $L^2(\mathbb{R}^n)$ to $L^2(\mathbb{R}^n)$ and $\|T_a\|_{L^2 \rightarrow L^2} \leq C_{\delta_1, \delta_2, n} \mathcal{C}$ for some constant $C_{\delta_1, \delta_2, n}$.

For simplicity, we summarize a easy-to-use L^2 -boundedness result as follows,

$$\boxed{a \in S_{0,0}^0 \Rightarrow \|T_a\|_{L^2 \rightarrow L^2} < \infty.}$$

For the L^p -boundedness ($1 < p < +\infty$) result, readers may refer to [6] (in French) and [15].

6.3. Gårding's inequalities

We use notation $\Re f$ to signify the real-valued part of any object f . Recall the Sobolev spaces $H^{s,p}(\mathbb{R}^n)$ defined in Definition 2.15, and the corresponding Sobolev norms $\|\cdot\|_{H^{m,p}}$ and $\|\cdot\|_{H^m}$. We denote $J^m := (I - \Delta)^{m/2}$ and $J := J^1$, namely, J^m takes $\langle \xi \rangle^m$ as its symbol. It can be checked that $J^m \in \Psi^m$, $J^{m_1} J^{m_2} = J^{m_1 + m_2}$, J is self-adjoint, and J^0 is the identity operator.

6.3.1. Gårding's Inequality.

DEFINITION 6.16 (Strongly elliptic). Let $m \in \mathbb{R}$. A symbol a is said to be *strongly elliptic* of order $2m$, if $a \in S^{2m}$ and if there exist fixed positive constants C, R such that

$$\boxed{\Re a(x, \xi) \geq C \langle \xi \rangle^{2m}, \quad \forall |\xi| \geq R,}$$

holds.

Similar to Lemma 6.5, there is an equivalent definition for the strong ellipticity of a symbol.

LEMMA 6.17. *Assume $m \in \mathbb{R}$ and $a \in S^{2m}$. The strong ellipticity condition for a is equivalent to the fact that there exist two positive constants C and D such that*

$$\boxed{\Re a(x, \xi) \geq C \langle \xi \rangle^{2m} - D \langle \xi \rangle^{2m-1}, \quad \forall x, \xi \in \mathbb{R}^n.} \quad (6.9)$$

PROOF. Assume $a \in S^{2m}$ is strongly elliptic, then there are constants $C, R > 0$ such that

$$\Re a(x, \xi) / \langle \xi \rangle^{2m} \geq C, \quad \forall |\xi| \geq R,$$

so for any positive constant D we have

$$\Re a(x, \xi) / \langle \xi \rangle^{2m} \geq C - D \langle \xi \rangle^{-1}, \quad (6.10)$$

for $\forall |\xi| \geq R$. Also, because a is a symbol of order $2m$, for some $M > 0$ we have,

$$|\Re a(x, \xi) / \langle \xi \rangle^{2m}| \leq |a(x, \xi) / \langle \xi \rangle^{2m}| \leq M \Rightarrow \Re a(x, \xi) / \langle \xi \rangle^{2m} \geq -M, \quad \forall \xi \in \mathbb{R}^n.$$

We set D to be large enough such that

$$-M \geq \sup_{|\xi| \leq R} (C - D \langle \xi \rangle^{-1}), \quad \text{e.g. } D := (C + M) \langle R \rangle,$$

then (6.10) holds for both $|\xi| \geq R$ and $|\xi| \leq R$. This gives (6.9).

On the other hand, from (6.9) it is easy to see a is strongly elliptic. \square

We are ready for the Gårding's Inequality.

THEOREM 6.18 (Gårding's inequality). *Assume $m \in \mathbb{R}$ and the symbol $a \in S^{2m}$ is strongly elliptic. Then we can find a positive constant C and a positive constant C_s for every real numbers $s \geq \frac{1}{2}$ such that*

$$\Re(T_a \varphi, \varphi) \geq C \|\varphi\|_{H^m}^2 - C_s \|\varphi\|_{H^{m-s}}^2, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n). \quad (6.11)$$

REMARK 6.19. When $a(x, \xi) = \langle \xi \rangle^{2m}$, then a is strongly elliptic and

$$\Re(T_a \varphi, \varphi) = \Re(J^{2m} \varphi, \varphi) = \Re(J^m \varphi, J^m \varphi) = \|\varphi\|_{H^m}^2,$$

which implies (6.11). Theorem 6.18 implies that even if a symbol is not of the form $\langle \xi \rangle^{2m}$ but is only strongly elliptic of order $2m$, then T_a still possesses some positiveness.

PROOF OF THEOREM 6.18. Let's denote the symbol of T_a^* as a^* , then it can be checked that

$$\Re(T_a \varphi, \varphi) = (T_{\frac{1}{2}(a+a^*)} \varphi, \varphi).$$

Step 1. When $m = 0$. Because a is strongly elliptic and $m = 0$, by Lemma 6.17 and Theorem 5.7 we have

$$\frac{1}{2}(a + a^*) = \Re a + r \geq C - D \langle \xi \rangle^{-1} + r = C - r,$$

where r is a generic symbol in S^{-1} . This makes it legal to define a symbol¹ $b \in S^0$ as follows,

$$b(x, \xi) := \left(\frac{1}{2}(a + a^*) - \frac{1}{2}C + r \right)^{1/2} \quad (6.12)$$

Then by Theorems 5.3 & 5.7 we have (symbolic calculus)

$$b^* \# b = (\bar{b} + S^{-1})b + S^{-1} = b^2 + S^{-1} = \frac{1}{2}(a + a^*) - \frac{1}{2}C + r + S^{-1},$$

so

$$\frac{1}{2}(a + a^*) = b^* \# b + \frac{1}{2}C + r,$$

where r is a generic symbol in S^{-1} . Therefore,

$$\begin{aligned} \Re(T_a \varphi, \varphi) &= (T_{\frac{1}{2}(a+a^*)} \varphi, \varphi) = (T_b^* T_b \varphi, \varphi) + \frac{C}{2} \|\varphi\|_{L^2}^2 + (R\varphi, \varphi) \\ &= \|T_b \varphi\|_{L^2}^2 + \frac{C}{2} \|\varphi\|_{L^2}^2 + (R\varphi, \varphi) \geq \frac{C}{2} \|\varphi\|_{L^2}^2 + (R\varphi, \varphi), \end{aligned} \quad (6.13)$$

for some $R \in \Psi^{-1}$. We have

$$\begin{aligned} |(R\varphi, \varphi)| &= |(R J^{1/2} J^{-1/2} \varphi, J^{1/2} J^{-1/2} \varphi)| = |(J^{1/2} R J^{1/2} (J^{-1/2} \varphi), J^{-1/2} \varphi)| \\ &\leq \|J^{1/2} R J^{1/2} (J^{-1/2} \varphi)\|_{L^2} \|\varphi\|_{H^{-1/2}} \leq C' \|J^{-1/2} \varphi\|_{L^2} \|\varphi\|_{H^{-1/2}} \end{aligned} \quad (6.14)$$

$$= C' \|\varphi\|_{H^{-1/2}}^2. \quad (6.15)$$

Note that in (6.14) we used the facts $J^{1/2} R J^{1/2} \in \Psi^0$ and operators in Ψ^0 are L^2 -bounded. Combining (6.15) with (6.13) we arrive at

$$\Re(T_a \varphi, \varphi) \geq \frac{C}{2} \|\varphi\|_{L^2}^2 - C' \|\varphi\|_{H^{-1/2}}^2. \quad (6.16)$$

¹See Exercise 6.3.

Step 2. When $m \neq 0$. Let $T_{a'} = J^{-m}T_a J^{-m}$ for certain $a' \in S^0$. Then it can be checked that there exist $C, D > 0$ so that

$$\Re a'(x, \xi) \geq C - D\langle \xi \rangle^{-1}, \quad \forall x, \xi \in \mathbb{R}^n. \quad (6.17)$$

Hence according to Lemma 6.17, a' is strongly elliptic, so by using the result in **Step 1** we can have

$$\begin{aligned} \Re(T_a \varphi, \varphi) &= \Re(J^m T_{a'} J^m \varphi, \varphi) = \Re(T_{a'} J^m \varphi, J^m \varphi) \\ &\geq \frac{C}{2} \|J^m \varphi\|_{L^2}^2 - C' \|J^m \varphi\|_{H^{-1/2}}^2 \\ &= \frac{C}{2} \|\varphi\|_{H^m}^2 - C' \|\varphi\|_{H^{m-1/2}}^2. \end{aligned} \quad (6.18)$$

Step 3. From Theorem 2.18 we have

$$\|\varphi\|_{H^{m-1/2}}^2 \leq \frac{1}{D^{1/2}} \|\varphi\|_{H^m}^2 + D^{s-1/2} \|\varphi\|_{H^{m-s}}^2,$$

for any $D > 0$. Set D to be small enough and substitute the inequality above into (6.16) and (6.18), we arrive at the conclusion. \square

Gårding's Inequality is used for giving the existence and uniqueness of the following type equation:

$$(T_a + \lambda I)u = f.$$

Let $m \geq 1/2$ and $s = m$, and assume $a \in S^{2m}$ is strongly elliptic symbol, then

$$C \|\varphi\|_{H^m}^2 - \lambda_0 \|\varphi\|_{L^2}^2 \leq \Re(T_a \varphi, \varphi), \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n)$$

for some constant $\lambda_0 > 0$, then for all $\lambda > \lambda_0$, we can conclude

$$C \|\varphi\|_{L^2}^2 \leq C \|\varphi\|_{H^m}^2 \leq \Re((T_a + \lambda)\varphi, \varphi) = \Re(\varphi, (T_a^* + \lambda)\varphi) \leq \|\varphi\|_{L^2} \|(T_a^* + \lambda)\varphi\|_{L^2},$$

which leads to a coercive condition:

$$C \|\varphi\|_{L^2} \leq \|(T_a^* + \lambda)\varphi\|_{L^2}.$$

Combining this with the Lax-Milgram theorem we can conclude that:

COROLLARY 6.20. *Assume $m \geq 1/2$ and $a \in S^{2m}$ is strongly elliptic. There exists a constant λ_0 such that when any $\lambda > \lambda_0$, for any $f \in L^2(\mathbb{R}^n)$ there exists a unique weak solution $u \in L^2(\mathbb{R}^n)$ satisfying the equation*

$$(T_a + \lambda)u = f.$$

6.3.2. Sharp Gårding's Inequality. In the proof of Theorem 6.18 later on, we see that having a strictly positive lower bound for $\Re a$ is critical, and the method in that proof will fail if the lower bound reduces to zero. However, when $\Re a \geq 0$, one can still obtain some lower bound of $\Re(T_a \varphi, \varphi)$ and that result is called sharp Gårding's inequality.

THEOREM 6.21 (Sharp Gårding's Inequality). *For a symbol $a \in S^{2m}$ satisfying*

$$\boxed{\Re a(x, \xi) \geq 0, \quad \forall |\xi| \geq R,} \quad (6.19)$$

we can find a positive constant C such that

$$\boxed{\Re(T_a \varphi, \varphi) \geq -C \|\varphi\|_{H^{m-1/2}}^2, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n).}$$

To prove (6.21), we introduce the wave packet transform. The wave packet transform $W : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n \times \mathbb{R}^n)$ is defined as (see [4, Theorem 4.2.3])

$$Wu(z, \xi) := c_n \langle \xi \rangle^{n/4} (e^{i(\cdot)\xi - \langle \xi \rangle |\cdot|^2} *_z u), \quad (6.20)$$

and its conjugate in terms of the L^2 -inner product is given by,

$$W^*F(x) = c_n \int \langle \xi \rangle^{n/4} (e^{i(\cdot)\xi - \langle \xi \rangle |\cdot|^2} *_x F(\cdot, \xi)) d\xi, \quad (6.21)$$

where the constant $c_n = 2^{-n/4} \pi^{-3n/4}$ and $(f *_x g)$ signifies $\int f(x-y)g(y) dy$.

LEMMA 6.22. *The wave packet transform W defined in (6.20) is a bounded linear operator.*

PROOF. The linearity is obvious.

To show the boundedness, we compute $\|Wu\|_{L^2(\mathbb{R}^n \times \mathbb{R}^n)}^2$,

$$\begin{aligned} \|Wu\|_{L^2(\mathbb{R}^n \times \mathbb{R}^n)}^2 &\simeq \iint |\langle \xi \rangle^{n/4} (e^{i(\cdot)\xi - \langle \xi \rangle |\cdot|^2} *_z u)|^2 dz d\xi \\ &= \int \langle \xi \rangle^{n/2} \int |\mathcal{F}\{(e^{i(\cdot)\xi - \langle \xi \rangle |\cdot|^2}\}(\eta))|^2 \cdot |\hat{u}(\eta)|^2 d\eta d\xi \quad (\text{Plancherel theorem}) \\ &= \int \left(\int \langle \xi \rangle^{n/2} |\mathcal{F}\{(e^{i(\cdot)\xi - \langle \xi \rangle |\cdot|^2}\}(\eta))|^2 d\xi \right) \cdot |\hat{u}(\eta)|^2 d\eta \\ &\simeq \int \left(\int \langle \xi \rangle^{n/2} |\langle \xi \rangle^{-n/2} e^{-\langle \xi \rangle^{-1} |\eta - \xi|^2/4}|^2 d\xi \right) \cdot |\hat{u}(\eta)|^2 d\eta \\ &= \int \left(\int \langle \xi \rangle^{-n/2} e^{-\langle \xi \rangle^{-1} |\eta - \xi|^2/2} d\xi \right) \cdot |\hat{u}(\eta)|^2 d\eta \\ &\leq \int \left(\int e^{-|\eta - \xi|^2/2} d\xi \right) \cdot |\hat{u}(\eta)|^2 d\eta \lesssim \int |\hat{u}(\eta)|^2 d\eta \\ &= \|u\|_{L^2(\mathbb{R}^n)}^2. \end{aligned}$$

The proof is complete. \square

PROOF OF THEOREM 6.21. similar to Proof of Theorem 6.18, the general cases w.r.t. m stem from the special case where $m = 1/2$. Let's assume $m = 1/2$ for the time being and try to show $\Re(T_a \varphi, \varphi) \gtrsim -\|\varphi\|_{-1/2, 2}^2$.

The condition (6.19) can be replaced by " $\Re a(x, \xi) \geq 0, \forall \xi \in \mathbb{R}^n$ ", and this is because we can fix some $\chi \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$ (thus $\chi \in S^{-\infty}$) satisfying $\chi(\xi) \geq \sup_{(x, \xi)} \Re a(x, \xi)$ when $\{|\xi| \leq R\}$, and then we can obtain $\Re a(x, \xi) + \chi(\xi) \geq 0, \forall \xi \in \mathbb{R}^n$. Note that $\Re(\chi \varphi, \varphi) \gtrsim -\|\varphi\|_{L^2}^2$ because $\chi \in S^{-\infty}$. Therefore, from now on we assume $\Re a(x, \xi) \geq 0, \forall \xi \in \mathbb{R}^n$.

Denote as $b(x, \xi)$ the symbol of $W^* \Re a W$. The operator $T_b := W^* \Re a W$ is defined by $T_b \varphi(x) = W^*(\Re a \cdot W \varphi)(x)$ for $\varphi \in \mathcal{S}(\mathbb{R}^n)$. We have

$$\begin{aligned} T_b \varphi(x) &\simeq \int \langle \xi \rangle^{n/4} \int e^{i(x-y)\xi - \langle \xi \rangle |x-y|^2} \Re a(y, \xi) W \varphi(y, \xi) dy d\xi \\ &\simeq \int \langle \xi \rangle^{n/4} \int e^{i(x-y)\xi - \langle \xi \rangle |x-y|^2} \Re a(y, \xi) \langle \xi \rangle^{n/4} \int e^{i(y-z)\xi - \langle \xi \rangle |y-z|^2} \varphi(z) dz dy d\xi \\ &\simeq \iint e^{i(x-z)\xi} \left(\int e^{-\langle \xi \rangle (|y-z|^2 + |x-y|^2)} \langle \xi \rangle^{n/2} \Re a(y, \xi) dy \right) \varphi(z) dz d\xi \\ &= \iint e^{i(x-z)\xi} \tilde{a}(x, z, \xi) \varphi(z) dz d\xi, \end{aligned} \quad (6.22)$$

where $\tilde{a}(x, z, \xi) := \langle \xi \rangle^{n/2} \int e^{-\langle \xi \rangle (|y-z|^2 + |x-y|^2)} \Re a(y, \xi) dy$. It can be checked that $\tilde{a} \in S_{1,1/2}^1$. By [4, Theorem 2.4.1], we have the asymptotic expansion

$$b(x, \xi) = \tilde{a}(x, x, \xi) + \sum_{j=1}^n e_j(x, \xi) + r, \quad r \in S_{1,1/2}^0, \quad (6.23)$$

where $e_j(x, \xi) = \partial_{z_j} D_{\xi_j} \tilde{a}(x, z, \xi)|_{z=x}$. Note that e_j belongs to $S_{1,1/2}^{1/2}$, not $S_{1,1/2}^0$, and this is why we expand \tilde{a} to the second order. The symbols e_j is purely imaginary because \tilde{a} is real.

For $\tilde{a}(x, x, \xi)$, we have

$$\begin{aligned} \tilde{a}(x, x, \xi) &= \langle \xi \rangle^{n/2} \Re \int e^{-2\langle \xi \rangle |x-y|^2} a(y, \xi) dy \\ &= \langle \xi \rangle^{n/2} \Re \int e^{-2\langle \xi \rangle |x-y|^2} [a(x, \xi) + \sum_j (y-x)^{(j)} \partial_{x_j} a(x, \xi) \\ &\quad + \sum_{|\alpha|=2} (y-x)^\alpha \partial_x^\alpha a(\rho x + (1-\rho)y, \xi)/2] dy \\ &\simeq \Re a(x, \xi) + \langle \xi \rangle^{n/2} \Re \partial_{x_j} a(x, \xi) \sum_j \int e^{-2\langle \xi \rangle |x-y|^2} (y-x)^{(j)} dy \\ &\quad + \langle \xi \rangle^{n/2} \sum_{|\alpha|=2} \Re \int e^{-2\langle \xi \rangle |x-y|^2} (y-x)^\alpha \cdot \partial_x^\alpha a(\rho x + (1-\rho)y, \xi)/2 dy \\ &= \Re a(x, \xi) + \langle \xi \rangle^{n/2} \sum_{|\alpha|=2} \Re \int e^{-2\langle \xi \rangle |x-y|^2} (y-x)^\alpha \cdot \partial_x^\alpha a(\rho x + (1-\rho)y, \xi)/2 dy \\ &= \Re a(x, \xi) + r', \quad r' \in S^0. \end{aligned} \quad (6.24)$$

The last equal sign in (6.24) is due to the following computation,

$$\begin{aligned} &|\langle \xi \rangle^{n/2} \int e^{-2\langle \xi \rangle |x-y|^2} (y-x)^\alpha \cdot \partial_x^\alpha a(\rho x + (1-\rho)y, \xi)/2 dy| \\ &\lesssim \langle \xi \rangle^{n/2} \int e^{-2\langle \xi \rangle |x-y|^2} |(y-x)^\alpha| \cdot \langle \xi \rangle dy \\ &= \int e^{-2|\langle \xi \rangle^{1/2} y|^2} |(\langle \xi \rangle^{1/2} y)^\alpha| d(\langle \xi \rangle^{1/2} y) \\ &= \int e^{-2|y|^2} |y^\alpha| dy \leq C \end{aligned}$$

for some positive constant C .

Combining (6.23) and (6.24), we obtain

$$b(x, \xi) = \Re a(x, \xi) + \sum_{j=1}^n e_j(x, \xi) + r, \quad e_j \in S_{1,1/2}^{1/2}, \quad r \in S_{1,1/2}^0, \quad (6.25)$$

and thus

$$(a + a^*)/2 = b(x, \xi) - \sum_{j=1}^n e_j(x, \xi) - r, \quad r \in S_{1,1/2}^0. \quad (6.26)$$

The r in (6.25) and (6.26) are different from each other and are also different from the r in (6.23). Now we have

$$\Re(T_a \varphi, \varphi) = [(T_a \varphi, \varphi) + \overline{(T_a \varphi, \varphi)}]/2 = [(T_a \varphi, \varphi) + (\varphi, T_a \varphi)]/2 = (T_{(a+a^*)/2} \varphi, \varphi)$$

$$\begin{aligned}
&= (T_b\varphi, \varphi) - (T_{e_j}\varphi, \varphi) - (T_r\varphi, \varphi) \\
&= (\Re a \cdot W\varphi, W\varphi) - (T_{e_j}\varphi, \varphi) - (T_r\varphi, \varphi) \\
&\geq -(T_{e_j}\varphi, \varphi) - (T_r\varphi, \varphi) \quad (\text{because } \Re a \geq 0) \\
&= -\Re(T_{e_j}\varphi, \varphi) - \Re(T_r\varphi, \varphi) = -(T_{(e_j+e_j^*)/2}\varphi, \varphi) - \Re(T_r\varphi, \varphi) \\
&\geq -(T_{(e_j+e_j^*)/2}\varphi, \varphi) - C\|\varphi\|_{L^2}^2, \tag{6.27}
\end{aligned}$$

for some positive constant C . The L^2 -boundedness of operators whose symbol come from $S_{1,1/2}^0$ can be proved in a similar manner as in the proof of that of S^0 , cf. [4, Theorem 4.1.1] and [1, Theorem 5.1].

Recall that $e_j \in S_{1,1/2}^{1/2}$ and e_j is purely imaginary, thus the principal symbol of $T_{(e_j+e_j^*)/2}$ equals to zero and hence $(e_j + e_j^*)/2 \in S_{0,1/2}^{1/2}$. Therefore,

$$(T_{(e_j+e_j^*)/2}\varphi, \varphi) \leq C\|\varphi\|_{L^2}^2 \tag{6.28}$$

for some positive constant C .

Combining (6.27) and (6.28), we arrive at the conclusion for the case $m = 1/2$. Based on the result regarding $m = 1/2$, the proof of general cases become trivial. \square

Exercise

EXERCISE 6.1. Prove Lemma 6.3.

EXERCISE 6.2. Assume $m \in \mathbb{R}$ and $a \in S^m$ and a is elliptic. Fix a cutoff function $\chi \in C^\infty(\mathbb{R}^n)$ such that $\chi(\xi) = R$ when $|\xi| \leq 1$ and $\chi(\xi) = 0$ when $|\xi| \geq R + 1$, where the R comes from the definition of the ellipticity of a . Define $r_0(x, \xi) := (1 - \chi(\xi))/a(x, \xi)$. Prove that $r_0 \in S^{-m}$.

EXERCISE 6.3. Prove the $b(x, \xi)$ defined in (6.12) is indeed a symbol and is of order 0. Hint: use [1, Lemma 2.1.1] or [26, Lemma 17.2].

EXERCISE 6.4. Prove (6.17) is true.

EXERCISE 6.5. Assume the symbols a and b are elliptic. Show that $T_a T_b$ and T_a^* are also elliptic. Hint: utilize Lemma 6.5.

Semi-classical Ψ DOs and its symbolic calculus

Semiclassical analysis shares lots of features with Ψ DO theory, while also keeping some of its own specialties. One of the application of semiclassical analysis is Carleman estimates.

7.1. Semi-classical Ψ DOs

7.1.1. Symbol classes.

DEFINITION 7.1 (Order function). A measurable function $m: \mathbb{R}^{2n} \rightarrow \mathbb{R}^+$ is call an *order function* if there exist constants $C > 0$ and $N \in \mathbb{N}$ such that

$$m(z_1 - z_2) \leq C \langle z_1 \rangle^N m(z_2), \quad \forall z_1, z_2 \in \mathbb{R}^{2n}.$$

The integer N is called the order of m .

For any $a, b \in \mathbb{R}$, $m(x, \xi) = \langle x \rangle^a \langle \xi \rangle^b$ are an order functions with $N = 2 \max\{|a|, |b|\}$. If m_1, m_2 are order functions, so does $m_1 m_2$.

DEFINITION 7.2 (Semiclassical symbol class). Let $h \in (0, 1)$, $\delta \in [0, \frac{1}{2}]$ and m be an order function with order N . For $a(\cdot; h) \in C^\infty(\mathbb{R}^{2n})$, we say $a \in S_\delta(m)$ with order N if

$$|\partial_z^\alpha a(z; h)| \leq C_\alpha h^{-\delta|\alpha|} m(z), \quad \forall z \in \mathbb{R}^{2n}.$$

Define a family of seminorms

$$|a(\cdot; h)|_{S_\delta(m), \alpha} = |a|_\alpha := \sup_{z \in \mathbb{R}^{2n}} \frac{|\partial_z^\alpha a(z; h)|}{h^{-\delta|\alpha|} m(z)},$$

and so the semiclassical symbol class $S_\delta(m)$ is given by

$$S_\delta(m) := \{a(z; h) \in C^\infty; \forall \text{ multi-index } \alpha, |a|_\alpha < +\infty\}.$$

We abbreviate $S_\delta(1)$ as S_δ and $S_0(1)$ as S .

Note that in contrast to the Kohn-Nirenberg symbol (cf. Definitions 2.1 & 2.4), the semiclassical symbol doesn't gain decay w.r.t. its arguments after being differentiated.

We write $a(\cdot; h) = \mathcal{O}_{S_\delta(m)}(f(h))$ if for every multi-index α , there exist h_0 and a such that $|a|_\alpha \leq C_\alpha f(h)$ holds for all $h \in (0, h_0)$, namely,

$$a(\cdot; h) = \mathcal{O}_{S_\delta(m)}(f(h)) \Leftrightarrow |\partial^\alpha a(\cdot; h)| \lesssim f(h) h^{-\delta|\alpha|} m.$$

It can be checked that $\forall a \in S_\delta(m)$, we have $f(h)a = \mathcal{O}_{S_\delta(m)}(f(h))$. For $S_\delta(m)$ and $a_j \in S_\delta(m)$, we write $a \sim \sum_j h^j a_j$ if $a - \sum_{j=0}^N h^j a_j = \mathcal{O}_{S_\delta(m)}(h^{N+1})$.

LEMMA 7.3. Assume $0 \leq \delta \leq \frac{1}{2}$ and $a \in S_\delta(m)$ and $a_1 \in S_\delta(m_1)$, $a_2 \in S_\delta(m_2)$. Then $hD_j a \in S_\delta(m)$ and $a_1 a_2 \in S_\delta(m_1 m_2)$.

PROOF. We can compute

$$\begin{aligned} |\partial^\alpha(hD_j a)| &= h|\partial^{\alpha+e_j} a| \leq C_\alpha h h^{-\delta(|\alpha+e_j|)} m = C_\alpha h h^{-\delta|\alpha|-\delta} m = C_\alpha h^{1-\delta} h^{-\delta|\alpha|} m \\ &\leq h_0 C_\alpha h^{-\delta|\alpha|} m. \end{aligned}$$

Hence $hD_j a \in S_\delta(m)$. We omit the rest of the proof. \square

DEFINITION 7.4 (Asymptotics). For symbol a , $a_j \in S_\delta(m)$ ($j = 0, 1, \dots$), we write $a \sim \sum_j h^{(1-2\delta)j} a_j$ in $S_\delta(m)$ if $a - \sum_{j=0}^N h^{(1-2\delta)j} a_j = \mathcal{O}_{S_\delta(m)}(h^{(1-2\delta)(N+1)})$ holds for every $n \in \mathbb{N}$, namely,

$$a \sim \sum_j h^{(1-2\delta)j} a_j \text{ in } S_\delta(m) \quad \Leftrightarrow \quad a = \sum_{j=0}^N h^{(1-2\delta)j} a_j + h^{(1-2\delta)(N+1)} S_\delta(m).$$

Here $h^{(1-2\delta)(N+1)} S_\delta(m)$ means $h^{(1-2\delta)(N+1)} r$ for some $r \in S_\delta(m)$. The a_0 is called the *principal symbol* of a .

The asymptotics is more about h than the Kohn-Nirenberg symbol which is more about ξ . To avoid confusion, we would like to comment in advance that even though the definition of asymptotics semiclassical symbol is in the form $a \sim a_0 + h^{1-2\delta} a_1 + h^{(1-2\delta)2} a_2 + \dots$, but later we may see a be expressed as $b_0 + h b_1 + h^2 b_2 + \dots$, e.g. in (7.15). The difference is that it is $h^{2\delta j} b_j$ rather than b_j itself that is in $S_\delta(m)$.

THEOREM 7.5. For $\forall a_j \in S_\delta(m)$ ($j = 0, 1, \dots$), there always exists $a \in S_\delta(m)$ such that $a \sim \sum_j h^{(1-2\delta)j} a_j$.

PROOF. We choose a cutoff function $\chi \in C_c^\infty(\mathbb{R})$ satisfying $\chi \equiv 1$ in $(-1, 1)$, $0 \leq \chi \leq 1$, χ is decreasing in the interval $(1, 2)$ and $\text{supp } \chi \subset (-2, 2)$. Note that we define χ on the whole real axis but will only use its definition on the positive real axis.

Step 1. Define

$$a := \sum_{j \geq 0} \chi(\lambda_j h) h^{(1-2\delta)j} a_j$$

for some λ_j which shall be determined. Our scheme is to choose $\lambda_j > 0$ properly (grows fast enough) such that a will be well-defined at each point and satisfies Definition 7.4. From the construction of χ it can be checked that

$$\forall h \geq 0, \forall k \geq 0, \quad \chi(h) h^{(1-2\delta)k} \leq 2^{(1-2\delta)k}. \quad (7.1)$$

Hence,

$$\begin{aligned} |a| &= \left| \sum_{j \geq 0} \chi(\lambda_j h) h^{(1-2\delta)j} a_j \right| = \left| \sum_{j \geq 0} \chi(\lambda_j h) (\lambda_j h)^{(1-2\delta)j} (\lambda_j h)^{-(1-2\delta)j} h^{(1-2\delta)j} a_j \right| \\ &\leq \sum_{j \geq 0} (2/\lambda_j)^{(1-2\delta)j} |a_j|, \end{aligned}$$

and similarly,

$$|\partial^\alpha a| \leq \sum_{j \geq 0} (2/\lambda_j)^{(1-2\delta)j} |\partial^\alpha a_j| \leq \sum_{j \geq 0} C_{j,\alpha} (2/\lambda_j)^{(1-2\delta)j} \cdot h^{-\delta|\alpha|} m.$$

Step 2. For a specific α , we only need to choose $\{\lambda_{j,\alpha}\}_{j \geq 0}$ grow fast enough such that

$$\sum_{j \geq 0} C_{j,\alpha} (2/\lambda_{j,\alpha})^{(1-2\delta)j}$$

is finite, and one example is $\lambda_{j,\alpha} = 3C_{j,\alpha}^{1/[(1-2\delta)j]}$. Then using diagonal arguments we could choose a suitable set $\{\lambda_j\}$ from $\{\lambda_{j,\alpha}\}_{j \geq 0}$. However, we want $\{\lambda_j\}$ to grow even more faster for our later use; particularly, to guarantee (7.5) is finite. To that end, for each fixed multi-index α and non-negative integer M , we first choose $\{\lambda_{j,\alpha,M}\}_{j \geq 0}$ to grow fast enough w.r.t. j such that

$$\begin{cases} \sum_{j \geq 0} C_{j+M,\alpha} (2/\lambda_{j,\alpha,M})^{(1-2\delta)j} < +\infty, \\ \lambda_{j,\alpha,M'} \geq \lambda_{j,\alpha,M} \text{ when } M' \geq M, \\ \lambda_{j,\alpha',M} \geq \lambda_{j,\alpha,M} \text{ when } \alpha' \geq \alpha, \end{cases} \quad (7.2)$$

then we choose

$$\lambda_j := \lambda_{j,(j,j,\dots,j),j}$$

where (j,j,\dots,j) stands for the multi-index of which the value of every component is j . By doing so, we are guaranteed that the sum $\sum_{j \geq 0} C_{j+M,\alpha} (2/\lambda_j)^{(1-2\delta)j}$ is finite for every α and M . Back to the estimate of $|\partial^\alpha a|$, we are guaranteed that a is well-defined and $a \in S_\delta(m)$. It remains to show $a \sim \sum_j h^{(1-2\delta)j} a_j$ in $S_\delta(m)$.

Step 3. To analyze $a - \sum_{j=0}^N h^j a_j$, we use another trick similar to (7.1),

$$\forall h \geq 0, \forall k \geq 0, \quad |\chi(h) - 1| h^{-k} \leq 1. \quad (7.3)$$

The verification of (7.3) is left as an exercise. By (7.1) and (7.3), for $h \in (0, h_0)$ where $h_0 < 1$ we have

$$\begin{aligned} & |\partial^\alpha (a - \sum_{j=0}^N h^{(1-2\delta)j} a_j)| \\ & \leq \sum_{j=0}^N |\chi(\lambda_j h) - 1| h^{(1-2\delta)j} |\partial^\alpha a_j| + \sum_{j \geq N+1} \chi(\lambda_j h) h^{(1-2\delta)j} |\partial^\alpha a_j| \\ & \leq \sum_{j=0}^N |\chi(\lambda_j h) - 1| (\lambda_j h)^{-(1-2\delta)(N+1-j)} \cdot (\lambda_j h)^{(1-2\delta)(N+1-j)} h^{(1-2\delta)j} |\partial^\alpha a_j| \\ & \quad + \sum_{j \geq 0} \chi(\lambda_j h) \left(\frac{\lambda_j h}{2}\right)^{(1-2\delta)j} \cdot C_{j+N+1,\alpha} \left(\frac{2}{\lambda_j h}\right)^{(1-2\delta)j} h^{(1-2\delta)(j+N+1)} h^{-\delta|\alpha|} m \end{aligned} \quad (7.4)$$

$$\begin{aligned} & \leq \left[\sum_{j=0}^N \lambda_j^{(1-2\delta)(N+1-j)} C_{j,\alpha} + \sum_{j \geq 0} C_{j+N+1,\alpha} \left(\frac{2}{\lambda_j}\right)^{(1-2\delta)j} \right] h^{(1-2\delta)(N+1)} h^{-\delta|\alpha|} m \\ & \leq \tilde{C}_{N,\alpha,h_0} h^{(1-2\delta)(N+1)} h^{-\delta|\alpha|} m. \end{aligned} \quad (7.5)$$

Here in (7.4) we used $\chi(\lambda_j h) \leq \chi(\lambda_{j-1} h)$. Hence,

$$a - \sum_{j=0}^N h^{(1-2\delta)j} a_j = \mathcal{O}_{S_\delta(m)}(h^{(1-2\delta)(N+1)}).$$

The proof is complete. \square

7.1.2. Semiclassical pseudodifferential operators. Just as Kohn-Nirenberg symbols, every semiclassical symbol produces an operator, and in semiclassical situation, these operators are also described as quantizations of the corresponding symbols.

DEFINITION 7.6 (Quantization). We quantize the symbol $a(x, \xi)$ by means of (7.8) for $\forall t \in [0, 1]$. And we also denote *Standard quantization* and *Weyl quantization* as in (7.6)-(7.7),

$$\text{(Standard quant.:)} \quad a(x, hD)u := (2\pi h)^{-n} \int e^{i(x-y)\cdot\xi/h} a(x, \xi)u(y) dy d\xi, \quad (7.6)$$

$$\text{(Weyl quant.:)} \quad a^w(x, hD)u := (2\pi h)^{-n} \int e^{i(x-y)\cdot\xi/h} a\left(\frac{x+y}{2}, \xi\right)u(y) dy d\xi, \quad (7.7)$$

$$\text{(General quant.:)} \quad a_t(x, hD)u := (2\pi h)^{-n} \int e^{i(x-y)\cdot\xi/h} a(tx + (1-t)y, \xi)u(y) dy d\xi. \quad (7.8)$$

These operators defined above are called *semiclassical pseudodifferential operators* (abbreviated as $S\Psi$ DOs). We denote the set of $S\Psi$ DO with symbols coming from $S_\delta(m)$ as $\text{Op}_h(S_\delta(m))$.

According to Definition 7.6 we know that $a^w(x, hD) = a_{\frac{1}{2}}(x, hD)$ and $a(x, hD) = a_1(x, hD)$. It is trivial to see

$$a(x, \xi) = f(x)\xi_j \quad \Leftrightarrow \quad a(x, hD) = f(x)hD_j.$$

We introduce the h -dependent Fourier transform.

DEFINITION 7.7. The *semiclassical Fourier transform* \mathcal{F}_h and its inverse \mathcal{F}_h^{-1} are defined as

$$\mathcal{F}_h u(\xi) := (2\pi h)^{-n/2} \int_{\mathbb{R}^n} e^{-ix\cdot\xi/h} u(x) dx, \quad (7.9)$$

$$\mathcal{F}_h^{-1} u(x) := (2\pi h)^{-n/2} \int_{\mathbb{R}^n} e^{ix\cdot\xi/h} u(\xi) d\xi. \quad (7.10)$$

It can be checked that

$$\boxed{a(x, hD)u := \mathcal{F}_h^{-1}\{a(x, \cdot)\mathcal{F}_h u(\cdot)\}.} \quad (7.11)$$

Formula (7.11) is one of the reason why the semiclassical Fourier transform shall defined as in Definition 7.7.

LEMMA 7.8. Assume $\delta \in \mathbb{R}$ and $a \in S_\delta(m)$. Then for $\forall t \in [0, 1]$, we have that the operator a_t satisfies $a_t(x, hD): \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$ and $a_t(x, hD): \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$, and the mappings are bounded with norm depending on δ and h , but uniformly on t .

PROOF. Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$. We have

$$a_t(x, hD)\varphi(x) = (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} a(tx + (1-t)y, \xi; h)\varphi(y) dy d\xi.$$

The integrability of y is not a problem because $\varphi(y)$ is rapidly decay. For ξ , we should use integration by parts to gain enough decay on ξ . Notice that $\frac{1-\xi\cdot hD_y}{\langle \xi \rangle^2} (e^{i(x-y)\cdot\xi/h}) = e^{i(x-y)\cdot\xi/h}$, we denote $L_1 = \frac{1-\xi\cdot hD_y}{\langle \xi \rangle^2}$, then act L_1^{n+1} on $e^{i(x-y)\cdot\xi/h}$ and use integration by parts, we will end up in a integrand of order $\langle \xi \rangle^{-n-1}$ on ξ and rapidly decay on y , thus integrable. Hence we proved that $a_t(x, hD): \mathcal{S}(\mathbb{R}^n) \rightarrow L^\infty(\mathbb{R}^n)$ for $a \in S_\delta(m)$. Adopt similar arguments on $x^\alpha \partial^\beta a_t(x, hD)$, we can obtain $x^\alpha \partial^\beta a_t(x, hD): \mathcal{S}(\mathbb{R}^n) \rightarrow L^\infty(\mathbb{R}^n)$ for $\forall \alpha, \beta$. Therefore $a_t(x, hD): \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n)$. And the continuity of the operator can also be seen from the arguments above.

The second result holds due to duality arguments. \square

LEMMA 7.9. *If symbol a is independent of ξ , i.e. $a(x, \xi) = a(x)$, then*

$$a_t(x, hD)u(x) = a(x)u(x), \quad \forall t \in [0, 1].$$

PROOF. It is enough to prove for $u \in \mathcal{S}$. When $t = 1$, we have

$$\begin{aligned} a_1(x, hD)u(x) &= (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} a(x, \xi)u(y) \, dy \, d\xi \\ &= (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} a(x)u(y) \, dy \, d\xi \\ &= a(x)u(x). \end{aligned}$$

We have

$$\begin{aligned} \partial_t(a_t(x, hD)u(x)) &= (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} \partial_t(a(tx + (1-t)y))u(y) \, dy \, d\xi \\ &= (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} (x-y) \cdot \nabla a(tx + (1-t)y)u(y) \, dy \, d\xi \\ &\simeq (2\pi h)^{-n} \iint \nabla_\xi e^{i(x-y)\cdot\xi/h} \cdot \nabla a(tx + (1-t)y)u(y) \, dy \, d\xi \\ &\simeq (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} \operatorname{div}_\xi(\nabla a(tx + (1-t)y)u(y)) \, dy \, d\xi \\ &= 0. \end{aligned}$$

We arrive at the conclusion. \square

From Lemma 7.9 and (7.8) we know that if $a(x, \xi)$ is either independent of ξ or independent of x , the quantized operator $a_t(x, hD)$ will be independent of t . Hence, for fixed $x^*, \xi^* \in \mathbb{R}^n$, and denote $l(x, \xi) := x^* \cdot x + \xi^* \cdot \xi$, then $l_t(x, hD)$ is independent of t , i.e.,

$$\boxed{l_t(x, hD) = x^* \cdot x + \xi^* \cdot hD.} \quad (7.12)$$

7.2. Composition of the standard quantizations

For a non-degenerate, symmetric, real-valued $n \times n$ matrix Q , the quantization of the exponential of quadratic forms is defined as the standard quantization (cf. (7.6)),

$$e^{\frac{i}{2h}\langle QhD, hD \rangle} \varphi(x) = (2\pi h)^{-n} \iint_{\mathbb{R}^n \times \mathbb{R}^n} e^{i(x-y)\cdot\xi/h} e^{\frac{i}{2h}\langle Q\xi, \xi \rangle} \varphi(y) \, dy \, d\xi. \quad (7.13)$$

Readers may compare (7.13) with (7.31). The following lemma shows how to express $e^{\frac{i}{2h}\langle QhD, hD \rangle}$. The $e^{\frac{i}{2h}\langle QhD, hD \rangle}$ can be expanded by using stationary phase lemmas.

LEMMA 7.10. *Assume Q is a non-degenerate, symmetric, real-valued $n \times n$ matrix. We have $e^{\frac{i}{2h}\langle QhD, hD \rangle} : \mathcal{S} \rightarrow \mathcal{S}$ continuously. And when $0 \leq \delta \leq \frac{1}{2}$, we have that $e^{\frac{i}{2h}\langle QhD, hD \rangle} : S_\delta(m) \rightarrow S_\delta(m)$, and the expression is*

$$e^{\frac{i}{2h}\langle QhD, hD \rangle} a(x) = \frac{e^{i\frac{\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2} (2\pi h)^{n/2}} \int_{\mathbb{R}^n} e^{\frac{-i}{2h}\langle Q^{-1}y, y \rangle} a(x+y) \, dy. \quad (7.14)$$

The integral (7.14) is defined in oscillatory sense. Moreover, when $0 \leq \delta < \frac{1}{2}$, for $a \in S_\delta(m)$ we have the asymptotics

$$\boxed{e^{\frac{i}{2h}\langle QhD, hD \rangle} a = \sum_{j=0}^N \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a + h^{(1-2\delta)(N+1)} S_\delta(m).} \quad (7.15)$$

PROOF. For a non-degenerate, symmetric, real-valued $n \times n$ matrix Q we have (see [20])

$$\mathcal{F}\{e^{\frac{i}{2}\langle Qx,x \rangle}\}(\xi) = \frac{e^{\frac{i\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} e^{-\frac{i}{2}\langle Q^{-1}\xi,\xi \rangle}. \quad (7.16)$$

For any measurable function $a \in \mathcal{S}$, as long as the right-hand-side of (7.14) is definable, by the definition (7.13) we have

$$\begin{aligned} e^{\frac{i}{2h}\langle QhD,hD \rangle} a(x) &= (2\pi h)^{-n} \iint e^{i(x-y)\cdot\xi/h} e^{\frac{i}{2h}\langle Q\xi,\xi \rangle} a(y) dy d\xi \\ &= (2\pi h)^{-n/2} \int [(2\pi h)^{-n/2} \int e^{i(x-y)\cdot\xi/h} e^{\frac{i}{2h}\langle Q\xi,\xi \rangle} d\xi] a(y) dy \\ &= (2\pi h)^{-n/2} \int [(2\pi)^{-n/2} \int e^{-i(y-x)/\sqrt{h}\cdot\xi} e^{\frac{i}{2}\langle Q\xi,\xi \rangle} d\xi] a(y) dy \\ &= (2\pi h)^{-n/2} \int \mathcal{F}\{e^{\frac{i}{2}\langle Q\xi,\xi \rangle}\}((y-x)/\sqrt{h}) \cdot a(y) dy \\ &= (2\pi h)^{-n/2} \int \frac{e^{\frac{i\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}} e^{-\frac{i}{2h}\langle Q^{-1}(y-x),y-x \rangle} a(y) dy \quad \text{by (7.16)} \\ &= \frac{e^{\frac{i\pi}{4} \operatorname{sgn} Q}}{|\det Q|^{1/2}(2\pi h)^{n/2}} \int e^{\frac{i}{h}\langle -Q^{-1}y,y \rangle/2} a(x+y) dy. \end{aligned} \quad (7.17)$$

We arrive at (7.14). From (7.17) and (4.8) it is easy to see that $x^\alpha \partial^\beta e^{\frac{i}{2h}\langle QhD,hD \rangle} : \mathcal{S} \rightarrow L^\infty$ for $\forall \alpha, \beta$, hence $e^{\frac{i}{2h}\langle QhD,hD \rangle} : \mathcal{S} \rightarrow \mathcal{S}$ continuously.

Now we use Proposition 4.8, to estimate (7.17) and confirm that $e^{\frac{i}{2h}\langle QhD,hD \rangle}$ indeed maps $S_\delta(m)$ into itself. Denote the order of the symbol a as \tilde{N} . Choose the N in Proposition 4.8 to be $N \geq \tilde{N}/2 - 1$. The constants $C_{N,n,\alpha}$ in Proposition 4.8 satisfy $C_{N,n,\alpha} = h^{-\delta|\alpha|}$. From (7.17) and Proposition 4.8 we have

$$\begin{aligned} &e^{\frac{i}{2h}\langle QhD,hD \rangle} a(x) \\ &= \sum_{0 \leq j \leq N} \frac{h^j}{j!} \left(\frac{\langle (-Q^{-1})^{-1}D, D \rangle}{2i} \right)^j a(x) + \mathcal{O}(h^{N+1} \times \sum_{|\alpha| \leq n+2N+3} \sup_{y \in \mathbb{R}^n} \frac{|\partial^\alpha a(x+y; h)|}{\langle y \rangle^{n+4N+5-|\alpha|}}) \\ &= \sum_{0 \leq j \leq N} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \mathcal{O}(h^{N+1} \times \sum_{|\alpha| \leq n+2N+3} \sup_{y \in \mathbb{R}^n} \frac{h^{-\delta|\alpha|} \langle y \rangle^{\tilde{N}} m(x)}{\langle y \rangle^{2N+2}}) \\ &= \sum_{0 \leq j \leq N} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \mathcal{O}(h^{N+1-\delta(n+2N+3)} \sup_{y \in \mathbb{R}^n} \langle y \rangle^{\tilde{N}-2N-2} m(x)) \\ &= \sum_{0 \leq j \leq N} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + h^{(1-2\delta)(N+1)} \mathcal{O}(h^{-\delta(n+1)} m(x)). \end{aligned} \quad (7.18)$$

Now $e^{\frac{i}{2h}\langle QhD,hD \rangle} : S_\delta(m) \rightarrow S_\delta(m)$ is justified by (7.18) and similar arguments work on $\partial^\alpha (e^{\frac{i}{2h}\langle QhD,hD \rangle} a)$. It can be checked that $h^j \langle QD, D \rangle^j a = h^{(1-2\delta)j} S_\delta(m)$. Here $f = h^{(1-2\delta)j} S_\delta(m)$ means there exists a symbol $g \in S_\delta(m)$ such that $f = h^{(1-2\delta)j} g$. Hence, these leading terms matched with the stipulation in Definition 7.4. From (7.18) it seems we didn't obtain the expansion because the remainder term may surpass some leading terms. However, when $\delta < \frac{1}{2}$, from (7.18) we see that the order of the remainder term goes higher as N goes larger (while when $\delta = \frac{1}{2}$ this doesn't happen), and when we set N to be larger enough, these leading terms in front of the remainder term can exposed themselves from the

remainder and will not be surpassed by the remainder. For example, if we want to expand (7.18) up to N' , we first choose N such that $(1-2\delta)(N+1) - \delta(2n+1) \geq (1-2\delta)(N'+1)$, then (7.18) can be continued as

$$\begin{aligned}
(*) &= \sum_{0 \leq j \leq N} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \mathcal{O}(h^{(1-2\delta)(N+1) - \delta(2n+1)} m(x)) \\
&= \sum_{0 \leq j \leq N'} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \sum_{N'+1 \leq j \leq N} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) \\
&\quad + \mathcal{O}(h^{(1-2\delta)(N+1) - \delta(2n+1)} m(x)) \\
&= \sum_{0 \leq j \leq N'} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \sum_{N'+1 \leq j \leq N} h^{(1-2\delta)j} S_\delta(m) + \mathcal{O}(h^{(1-2\delta)(N'+1)} m(x)) \\
&= \sum_{0 \leq j \leq N'} \frac{(ih)^j}{j!} \left(\frac{\langle QD, D \rangle}{2} \right)^j a(x) + \mathcal{O}_{S_\delta(m)}(h^{(1-2\delta)(N'+1)}). \tag{7.19}
\end{aligned}$$

Note that here we omitted the investigation of $|\partial^\alpha (e^{\frac{i}{2\hbar} \langle QhD, hD \rangle} a)|$, but the prove shall almost the same as above. We proved (7.15). \square

THEOREM 7.11 (Composition of standard quantizations). *Let $a \in S_\delta(m_1)$, $b \in S_\delta(m_2)$. Denote*

$$(a\#b)(x, hD) = a(x, hD) \circ b(x, hD),$$

then $a\#b \in S_\delta(m_1 m_2)$, and

$$a\#b(x, \eta) = e^{\frac{i}{2\hbar} \langle QhD_{(y, \xi)}, hD_{(y, \xi)} \rangle} (a(x, \eta + \xi) b(x + y, \eta))|_{y=0, \xi=0}, \tag{7.20}$$

where $Q = \begin{pmatrix} 0 & I_{n \times n} \\ I_{n \times n} & 0 \end{pmatrix}$. Moreover, when $h \rightarrow 0^+$ we have the semiclassical asymptotics,

$$\begin{aligned}
a\#b(x, \eta) &= \sum_{j=0}^N \frac{(ih)^j}{j!} (D_y \cdot D_\xi)^j (a(x, \eta + \xi) b(x + y, \eta))|_{y=0, \xi=0} + h^{(1-2\delta)(N+1)} S_\delta(m_1 m_2) \\
&= \sum_{|\alpha| \leq N} \frac{(-ih)^{|\alpha|}}{\alpha!} \partial_\eta^\alpha a(x, \eta) \partial_x^\alpha b(x, \eta) + h^{(1-2\delta)(N+1)} S_\delta(m_1 m_2).
\end{aligned}$$

(7.21)

REMARK 7.12. When either $a(x, \xi)$ or $b(x, \xi)$ is polynomial of ξ , the expansion (7.21) will be finite, i.e. when N is large enough the remainders $h^{(1-2\delta)(N+1)} S_\delta(m_1 m_2)$ will be exactly zero, see also [21, Remark 2.6.9]. This can be seen by directly working in the stationary phase lemma.

PROOF OF THEOREM 7.11. For a test function $\varphi \in \mathcal{S}(\mathbb{R}^n)$, we have

$$\begin{aligned}
(a\#b)(x, hD)\varphi &= (2\pi h)^{-2n} \int e^{i(x-z)\cdot\eta/h} \left(\int e^{i(x-y)\cdot(\xi-\eta)/h} a(x, \xi) b(y, \eta) dy d\xi \right) \varphi(z) dz d\eta \\
&= (2\pi h)^{-2n} \int e^{i(x-z)\cdot\eta/h} \left(\int e^{-iy\cdot\xi/h} a(x, \xi + \eta) b(y + x, \eta) dy d\xi \right) \varphi(z) dz d\eta \\
&= (2\pi h)^{-n} \int e^{i(x-z)\cdot\eta/h} \cdot [\dots] \cdot \varphi(z) dz d\eta,
\end{aligned}$$

$$\text{where } [\dots] = (2\pi h)^{-n} \int e^{\frac{-i}{2\hbar} \langle Q(y, \xi)^T, (y, \xi)^T \rangle} a(x, \eta + \xi) b(x + y, \eta) dy d\xi,$$

and $Q = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}$. Note that $Q^{-1} = Q$, $\text{sgn } Q = 0$ and $\det Q = 1$ or -1 . From (7.14) and (7.15), we see

$$\begin{aligned} [\dots] &= e^{\frac{i}{2\hbar} \langle Q^{-1} hD_{(y,\xi)}, hD_{(y,\xi)} \rangle} (a(x, \eta + \xi) b(x + y, \eta))|_{y=0, \xi=0} \\ &\sim \sum_j \frac{(ih)^j}{j!} \left(\frac{\langle QD_{(y,\xi)}, D_{(y,\xi)} \rangle}{2} \right)^j (a(x, \eta + \xi) b(x + y, \eta))|_{y=0, \xi=0}, \quad \text{in } S_\delta(m_1 m_2) \\ &\sim \sum_j \frac{(ih)^j}{j!} (D_y \cdot D_\xi)^j (a(x, \eta + \xi) b(x + y, \eta))|_{y=0, \xi=0}, \quad \text{in } S_\delta(m_1 m_2) \\ &= \sum_{|\alpha| \leq N} \frac{(-ih)^{|\alpha|}}{\alpha!} \partial_\eta^\alpha a(x, \eta) \partial_x^\alpha b(x, \eta) + \mathcal{O}_{S_\delta(m_1 m_2)}(h^{(1-2\delta)(N+1)}). \end{aligned}$$

We obtain (7.20) and (7.21). Note that we have used (5.8). The proof is complete. \square

Readers may compare Theorem 5.3 with (7.21). The asymptotics in Theorem 5.3 is in terms of the decay of $|\xi|$, but the asymptotics in (7.21) is about the order of h . The first two leading terms in Theorem 5.3 is $ab - i\nabla_\xi a \cdot \nabla_x b$ (no h), while that of (7.21) is $ab - ih\nabla_\xi a \cdot \nabla_x b$.

COROLLARY 7.13. *The first two leading terms of $a\#b$ is $ab - ih\nabla_\xi a \cdot \nabla_x b$. Assume $a \in S_\delta(m_1)$ and $b \in S_\delta(m_2)$, then the symbol of the commutator of $a(x, hD)$ and $b(x, hD)$ is*

$$\frac{\hbar}{i} \{a, b\} - \frac{\hbar^2}{2} \text{tr}(\nabla_\xi^2 a \cdot \nabla_x^2 b - \nabla_x^2 a \cdot \nabla_\xi^2 b) + \hbar^{3(1-2\delta)} S_\delta(m_1 m_2),$$

where $\{a, b\}$ is the Poisson bracket of a of b , and $\nabla_\xi^2 a \cdot \nabla_x^2 b$ is the product of two Hessian matrices, and tr is the trace.

The proof is left as an exercise. Finally, we also have symbolic calculus for the adjoint.

THEOREM 7.14 (Adjoint of standard quantizations). *Let $a \in S_\delta(m)$. Denote*

$$(a(x, hD)u, v) = (u, a^*(x, hD)v),$$

then $a^* \in S_\delta(m)$, and when $h \rightarrow 0^+$ we have the semiclassical asymptotics,

$$a^*(x, \xi) = \sum_{|\alpha| \leq N} \frac{\hbar^{|\alpha|}}{\alpha!} D_x^\alpha \partial_\xi^\alpha \bar{a}(x, \xi) + \hbar^{(1-2\delta)(N+1)} S_\delta(m). \quad (7.22)$$

We omit the proof.

7.3. Composition of the Weyl quantizations

The composition of the Weyl quantizations are more peculiar than that of the standard ones, and we explain this in §7.3.2. Before that, we make some preparation first.

7.3.1. Symplectic 2-form. We define the symplectic product.

DEFINITION 7.15 (Symplectic product). The *symplectic product* is defined as

$$\sigma: \mathbb{R}^{2n} \times \mathbb{R}^{2n} \rightarrow \mathbb{R}, \quad \sigma((x, \xi), (y, \eta)) := \xi \cdot y - x \cdot \eta.$$

REMARK 7.16. The underlying space \mathbb{R}^{2n} in Definition 7.15 can be generalized to be a tangent bundle. When \mathbb{R}^{2n} is replaced by a tangent bundle TM (or T^*M) where M is n -dimensional (hence TM is locally homeomorphic to \mathbb{R}^{2n}), σ can be generalized as a bilinear form on $T_p(TM) \times T_p(TM)$ in the following way. For any $p \in TM$ and $(u_x, u_\xi), (v_x, v_\xi) \in T_p(TM)$, we define

$$\sigma: T(TM) \times T(TM) \rightarrow \mathbb{R}, \quad \sigma|_p((u_x, u_\xi), (v_x, v_\xi)) := u_\xi \cdot v_x - u_x \cdot v_\xi$$

Locally speaking, when imposed a local coordinates system $\{x^j\}$ on M and the corresponding coordinates $\{\xi_j\}$ on the fiber, it can be checked that $\sigma = d\xi_j \wedge dx^j$ (Einstein summation convention invoked) and it is invariant w.r.t. the coordinates systems. This σ is a 2-form on the tangent bundle and is called the *symplectic 2-form*.

In what follows, we only work on \mathbb{R}^{2n} rather than on general manifolds. If without otherwise stated, we will use the following notations,

$$z = (x, \xi)^T \in \mathbb{R}^{2n}, \quad w = (y, \eta)^T \in \mathbb{R}^{2n}, \quad \zeta = (x, \xi, y, \eta)^T = (z^T, w^T)^T \in \mathbb{R}^{4n}. \quad (7.23)$$

Note that all of z, w and ζ are vertical vectors. Definition 7.15 is equivalent to

$$\sigma(z, w) = z^T \cdot \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \cdot w = z^T \cdot \sigma \cdot w = \langle \sigma^T z, w \rangle \quad (7.24)$$

where I is the identity $n \times n$ matrix and $\sigma = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$. Note that σ is non-degenerate and anti-symmetric, i.e. $\sigma^{-1} = \sigma^T = -\sigma$.

We note that (7.24) is homogeneous of degree 2 of ζ (i.e. $\sigma(h\zeta) = h^2\sigma(\zeta)$), but not in a quadratic form of ζ under a symmetric matrix (σ is not symmetric). We can achieve this by

$$\sigma(\zeta) = \sigma(z, w) = z^T \cdot \sigma \cdot w = \frac{1}{2}\zeta^T \cdot \begin{pmatrix} 0 & \sigma \\ \sigma^T & 0 \end{pmatrix} \cdot \zeta = \frac{1}{2}\zeta^T \cdot \Sigma \cdot \zeta,$$

where

$$\Sigma = \begin{pmatrix} 0 & \sigma \\ \sigma^T & 0 \end{pmatrix} \quad (7.25)$$

is a $4n \times 4n$ matrix. Note that Σ is non-degenerate and symmetric satisfying $\Sigma^{-1} = \Sigma^T = \Sigma$, $\det \Sigma = 1$ and $\text{sgn} \Sigma = 0$. In summary, we have

$$\sigma(z, w) = z^T \cdot \sigma \cdot w = \langle \sigma^T z, w \rangle = \frac{1}{2}\zeta^T \cdot \Sigma \cdot \zeta = \frac{1}{2}\langle \Sigma \zeta, \zeta \rangle. \quad (7.26)$$

7.3.2. The composition. If we mimic the proof of Theorem 7.11, we would have

$$\begin{aligned} & a^w(x, hD) \circ b^w(x, hD) \varphi \\ &= (2\pi h)^{-2n} \int e^{i(x-z)\cdot\eta/h} \left(\int e^{i(x-y)\cdot(\xi-\eta)/h} a\left(\frac{x+y}{2}, \xi\right) b\left(\frac{y+z}{2}, \eta\right) dy d\xi \right) \varphi(z) dz d\eta \\ &= (2\pi h)^{-2n} \int e^{i(x-z)\cdot\eta/h} \left(\int e^{-iy\cdot\xi/h} a\left(\frac{y}{2} + x, \xi + \eta\right) b\left(\frac{y}{2} + \frac{x+z}{2}, \eta\right) dy d\xi \right) \varphi(z) dz d\eta \\ &= (2\pi h)^{-n} \int e^{i(x-z)\cdot\eta/h} \cdot c\left(\frac{x+z}{2}, \eta\right) \cdot \varphi(z) dz d\eta = c^w(x, hD) \varphi \end{aligned}$$

where the c should satisfy

$$c\left(\frac{x+z}{2}, \eta\right) = (2\pi h)^{-n} \int e^{\frac{i}{2h}\langle Q(y, \xi), (y, \xi) \rangle} a\left(\frac{y}{2} + x, \xi + \eta\right) b\left(\frac{y}{2} + \frac{x+z}{2}, \eta\right) dy d\xi.$$

However, this argument doesn't work, because there is an additional x on the RHS.

Instead, from $a^w(x, hD) \circ b^w(x, hD) = c^w(x, hD)$ we can proceed as follows,

$$\begin{aligned} & a^w(x, hD) \circ b^w(x, hD) \varphi \\ &= (2\pi h)^{-2n} \int e^{i[(x-y)\cdot\xi + (y-z)\cdot\eta]/h} a\left(\frac{x+y}{2}, \xi\right) b\left(\frac{y+z}{2}, \eta\right) \varphi(z) \, dy \, dz \, d\xi \, d\eta \\ &= (2\pi h)^{-n} \int e^{i(x-z)\cdot\zeta/h} c\left(\frac{x+z}{2}, \zeta\right) \varphi(z) \, dz \, d\zeta, \end{aligned}$$

which, due to the arbitrary of φ , suggests

$$\int e^{i(x-z)\cdot\zeta/h} c\left(\frac{x+z}{2}, \zeta\right) \, d\zeta = (2\pi h)^{-n} \int e^{i[(x-y)\cdot\xi + (y-z)\cdot\eta]/h} a\left(\frac{x+y}{2}, \xi\right) b\left(\frac{y+z}{2}, \eta\right) \, dy \, d\xi \, d\eta.$$

Readers may note that the LHS is an inverse Fourier transform. We make the following change of variable before we perform the Fourier transform:

$$\begin{cases} \frac{x-z}{2} = s \\ \frac{x+z}{2} = t \end{cases} \Rightarrow \begin{cases} x = t + s \\ z = t - s \end{cases}$$

so

$$\begin{aligned} & \int e^{i2s\cdot\zeta/h} c(t, \zeta) \, d\zeta \\ &= (2\pi h)^{-n} \int e^{i[(t+s-y)\cdot\xi + (y-t+s)\cdot\eta]/h} a\left(\frac{t+s+y}{2}, \xi\right) b\left(\frac{y+t-s}{2}, \eta\right) \, dy \, d\xi \, d\eta, \end{aligned}$$

and

$$\begin{aligned} & c(t, \zeta) \\ &= (2\pi h)^{-2n} \int e^{-i2s\cdot\zeta/h} e^{i[(t+s-y)\cdot\xi + (y-t+s)\cdot\eta]/h} a\left(\frac{t+s+y}{2}, \xi\right) b\left(\frac{y+t-s}{2}, \eta\right) \, dy \, d\xi \, d\eta \, d(2s) \\ &= 2^n (2\pi h)^{-2n} \int e^{i[(y-t)\cdot(\eta-\xi) + s\cdot(\xi+\eta-2\zeta)]/h} a\left(\frac{t+y+s}{2}, \xi\right) b\left(\frac{t+y-s}{2}, \eta\right) \, dy \, ds \, d\xi \, d\eta \\ &= 2^n (2\pi h)^{-2n} \int e^{i[y\cdot(\eta-\xi) + s\cdot(\xi+\eta)]/h} a\left(\frac{y+s}{2} + t, \xi + \zeta\right) b\left(\frac{y-s}{2} + t, \eta + \zeta\right) \, dy \, ds \, d\xi \, d\eta \\ &= 2^n (2\pi h)^{-2n} \int e^{i[(y+s)\cdot\eta - (y-s)\cdot\xi]/h} a\left(\frac{y+s}{2} + t, \xi + \zeta\right) b\left(\frac{y-s}{2} + t, \eta + \zeta\right) \, dy \, ds \, d\xi \, d\eta \\ &= 2^n (2\pi h)^{-2n} \int e^{i(2y'\cdot\eta - 2s'\cdot\xi)/h} a(y' + t, \xi + \zeta) b(s' + t, \eta + \zeta) 2^n \, dy' \, ds' \, d\xi \, d\eta \\ &= (\pi h)^{-2n} \int e^{i2(y\cdot\eta - s\cdot\xi)/h} a(y + t, \xi + \zeta) b(s + t, \eta + \zeta) \, dy \, d\xi \, ds \, d\eta \\ &= (\pi h)^{-2n} \int e^{ih^{-1}\langle -2\Sigma(y, \xi, s, \eta), (y, \xi, s, \eta) \rangle/2} a(y + t, \xi + \zeta) b(s + t, \eta + \zeta) \, dy \, d\xi \, ds \, d\eta, \end{aligned} \quad (7.27)$$

where the $4n \times 4n$ matrix Σ is defined in (7.25), and we used Exercise 7.2.

Recall that $\Sigma^{-1} = \Sigma$, $\det \Sigma = 1$ and $\text{sgn } \Sigma = 0$. Now we apply Proposition 4.8 to (7.27) and obtain

$$\begin{aligned} c(t, \zeta) &\sim \frac{(\pi h)^{-2n} (2\pi h)^{2n}}{|\det(2\Sigma)|^{1/2}} \sum_j \frac{h^j}{j!} \left(\frac{\langle (-2\Sigma)^{-1} D_{(y, \xi, s, \eta)}, D_{(y, \xi, s, \eta)} \rangle}{2i} \right)^j (a(y + t, \xi + \zeta) \\ &\quad \times b(s + t, \eta + \zeta))|_{y=s=\xi=\eta=0} \end{aligned}$$

$$\begin{aligned}
&= \sum_j \frac{(h/(2i))^j}{j!} \left(\langle -\frac{1}{2} \Sigma D_{(y,\xi,s,\eta)}, D_{(y,\xi,s,\eta)} \rangle \right)^j (a(y,\xi)b(s,\eta))|_{y=s=t, \xi=\eta=\zeta} \\
&= \sum_j \frac{(ih/2)^j}{j!} (D_s \cdot D_\xi + \nabla_y \cdot \nabla_\eta)^j (a(y,\xi)b(s,\eta))|_{y=s=t, \xi=\eta=\zeta},
\end{aligned}$$

where we used Exercise 7.2. Here for simplicity we omitted the analysis of the remainder terms, and for the detailed analysis of the remainder, readers may refer to [28]. Noticing that

$$\begin{aligned}
(D_s \cdot D_\xi + \nabla_y \cdot \nabla_\eta)^j &= \sum_{0 \leq k \leq j} \binom{j}{k} (D_s \cdot D_\xi)^k (\nabla_y \cdot \nabla_\eta)^{j-k} \\
&= \sum_{0 \leq k \leq j} \binom{j}{k} \sum_{|\alpha|=k} \frac{k!}{\alpha!} D_s^\alpha D_\xi^\alpha \sum_{|\beta|=j-k} \frac{(j-k)!}{\beta!} \nabla_y^\beta \nabla_\eta^\beta \quad (\text{by (5.8)}) \\
&= \sum_{0 \leq k \leq j} \sum_{|\alpha|=k} \sum_{|\beta|=j-k} \frac{j!}{\alpha! \beta!} D_s^\alpha D_\xi^\alpha \partial_y^\beta \partial_\eta^\beta \\
&= \sum_{|\alpha|+|\beta|=j} \frac{j!}{\alpha! \beta!} D_s^\alpha D_\xi^\alpha \partial_y^\beta \partial_\eta^\beta,
\end{aligned}$$

we can continue

$$\begin{aligned}
c(t, \zeta) &\sim \sum_j \sum_{|\alpha|+|\beta|=j} \frac{(ih/2)^j}{j!} \frac{j!}{\alpha! \beta!} D_s^\alpha D_\xi^\alpha \partial_y^\beta \partial_\eta^\beta (a(y,\xi)b(s,\eta))|_{y=s=t, \xi=\eta=\zeta} \\
&= \sum_{\alpha, \beta} \frac{(ih/2)^{|\alpha|+|\beta|}}{\alpha! \beta!} [D_x^\beta \partial_\xi^\alpha (a(x,\xi)) D_x^\alpha \partial_\xi^\beta (b(x,\xi))] |_{x=t, \xi=\zeta}.
\end{aligned}$$

We have just proved the following result:

THEOREM 7.17 (Composition of quantizations of semiclassical symbols). *Assume that $a \in S_\delta(m_1)$, $b \in S_\delta(m_2)$. Denote*

$$(a \#^w b)^w(x, hD) = a^w(x, hD) \circ b^w(x, hD),$$

then $a \#^w b \in S_\delta(m_1 m_2)$ and

$$a \#^w b(x, \eta) = e^{ihA(D)} (a(x, \xi)b(y, \eta))|_{y=x, \xi=\eta}, \quad (7.28)$$

where $A(D) = \frac{1}{2} \sigma((D_x, D_\xi), (D_y, D_\eta)) = \frac{1}{2} (D_y \cdot D_\xi - D_x \cdot D_\eta)$ and the σ is defined in (7.23). Moreover, when $h \rightarrow 0^+$ we have the semiclassical asymptotics,

$$\begin{aligned}
a \#^w b(x, \eta) &= \sum_{j=0}^N \frac{(ih/2)^j}{j!} (D_y \cdot D_\xi + \nabla_x \cdot \nabla_\eta)^j (a(x, \xi)b(y, \eta))|_{y=x, \xi=\eta} \\
&\quad + h^{(1-2\delta)(N+1)} S_\delta(m_1 m_2) \\
&= \sum_{|\alpha|+|\beta| \leq N} \frac{(ih/2)^{|\alpha|+|\beta|}}{\alpha! \beta!} \partial_x^\beta D_\eta^\alpha a(x, \eta) D_x^\alpha \partial_\eta^\beta b(x, \eta) \\
&\quad + h^{(1-2\delta)(N+1)} S_\delta(m_1 m_2).
\end{aligned} \quad (7.29)$$

REMARK 7.18. Similar to Remark 7.12, the expansion (7.29) will be finite when either $a(x, \xi)$ or $b(x, \xi)$ is polynomial of ξ .

Readers may refer to [28, §4.11] for another proof of Theorem 7.17.

COROLLARY 7.19. *The first two leading terms of $a\#^w b$ is $\boxed{ab - ih\{a, b\}/2}$. Assume $a \in S_\delta(m_1)$ and $b \in S_\delta(m_2)$, then the commutator of $a^w(x, hD)$ and $b^w(x, hD)$ is*

$$\boxed{[a^w(x, hD), b^w(x, hD)] = \frac{h}{i}\{a, b\}^w(x, hD) + h^{3(1-2\delta)}\text{Op}_h(S_\delta(m_1 m_2))},$$

where $\{a, b\}$ is the Poisson bracket of a of b .

The remainder in the commutator expression looks out of expectation; it is of order $h^{3(1-2\delta)}$ rather than $h^{2(1-2\delta)}$. This is because the second order leading term is in fact zero.

PROOF. From (7.29) we have

$$\begin{aligned} a\#^w b &= ab + \frac{ih}{2}D_\xi a \cdot D_x b + \frac{ih}{2}\nabla_x a \cdot \nabla_\xi b \\ &\quad + \frac{(ih/2)^2}{2!}[(\nabla_x \cdot \nabla_\eta)^2 + (\nabla_\xi \cdot \nabla_y)^2 - 2(\nabla_x \cdot \nabla_\eta)(\nabla_\xi \cdot \nabla_y)](ab) \\ &\quad + h^{3(1-2\delta)}S_\delta(m_1 m_2) \\ &= ab - ih\{a, b\}/2 \\ &\quad + \frac{(ih/2)^2}{2!}[\text{tr}(\nabla_x^2 a \cdot \nabla_\eta^2 b) + \text{tr}(\nabla_\xi^2 a \cdot \nabla_y^2 b) - 2\text{tr}(\nabla_{(x,\xi)}^2 a \cdot \nabla_{(y,\eta)}^2 b)]|_{y=x, \xi=\eta} \\ &\quad + h^{3(1-2\delta)}S_\delta(m_1 m_2) \\ &= ab - ih\{a, b\}/2 + \frac{(ih/2)^2}{2!}[\text{tr}(\nabla_x^2 a \cdot \nabla_\eta^2 b) + \text{tr}(\nabla_x^2 b \cdot \nabla_\eta^2 a) - 2\text{tr}(\nabla_{(x,\eta)}^2 a \cdot \nabla_{(x,\eta)}^2 b)] \\ &\quad + h^{3(1-2\delta)}S_\delta(m_1 m_2), \end{aligned}$$

hence

$$a\#^w b - b\#^w a = -ih\{a, b\} + h^{3(1-2\delta)}S_\delta(m_1 m_2).$$

The proof is complete. \square

7.3.3. Specialties of Weyl quantization.

LEMMA 7.20. *For $u, v \in \mathcal{S}(\mathbb{R}^n)$, we have*

$$(a^w(x, hD)u, v) = (u, \bar{a}^w(x, hD)v).$$

The proof is left as an exercise.

The Weyl quantization is the correct generalization of a solution operator of an ODE. It is straightforward to check that $v(x, t) = e^{tf(x)}u(x)$ is the solution of an ODE

$$\begin{cases} \partial_t v(x, t) = f(x)v(x, t), & t \in \mathbb{R}, \\ v(x, 0) = u(x). \end{cases}$$

Recall the linear form $l(x, \xi) = x^* \cdot x + \xi^* \cdot \xi$. Now we would like to generalize the aforementioned idea by replacing $f(x)$ with an operator $\frac{i}{h}l(x, hD)$ and define $e^{\frac{it}{h}l(x, hD)}u$ as the unique solution of the corresponding ODE. But in order to avoid notational confusion between “ $e^{\frac{it}{h}l(x, hD)}u$ ” and “ $(e^{\frac{it}{h}l})(x, hD)u$ ” defined in (7.6), we deprecate the use of $e^{\frac{it}{h}l(x, hD)}u$. We will see from the following result that the correct generalization will be the Weyl quantization $(e^{\frac{it}{h}l})^w(x, hD)u$ instead of the standard quantization $(e^{\frac{it}{h}l})(x, hD)u$.

LEMMA 7.21. Let $l(x, \xi) = x^* \cdot x + \xi^* \cdot \xi$ for fixed $x^*, \xi^* \in \mathbb{R}^n$. For every $u \in \mathcal{S}$, the Weyl quantization $(e^{\frac{it}{\hbar}l})^w(x, hD)u$ is the unique solution of the ODE

$$\begin{cases} \partial_t v(x, t) = \frac{i}{\hbar} l(x, hD)v(x, t), & t \in \mathbb{R}, \\ v(x, 0) = u(x). \end{cases} \quad (7.30)$$

$$\boxed{e^{\frac{it}{\hbar}l(x, hD)} = (e^{\frac{it}{\hbar}l})^w(x, hD)}. \quad (7.31)$$

Specifically, we have

$$\boxed{(e^{\frac{it}{\hbar}l})^w(x, hD)u = e^{\frac{i}{\hbar}[(x^* \cdot x)t + (x^* \cdot \xi^*)t^2/2]}u(x + \xi^*t)}. \quad (7.32)$$

And we have the composition relation

$$\boxed{e^{\frac{it}{\hbar}l(x, hD)}e^{\frac{it}{\hbar}m(x, hD)} = e^{\frac{it^2}{2\hbar}\sigma(l, m)}e^{\frac{it}{\hbar}(l+m)(x, hD)}, \quad (7.33)$$

where the σ is given in Definition 7.15.

PROOF. First, we solve (7.30). By this ODE we have $\partial_t v(x, t) = \frac{i}{\hbar}(x^* \cdot x + \xi^* \cdot hD)v(x, t)$, which gives a transport equation $(\partial_t - \xi^* \cdot \nabla)v(x, t) = \frac{i}{\hbar}x^* \cdot xv(x, t)$. Let $\gamma: t \in \mathbb{R} \mapsto (x - \xi^*t, t) \in \mathbb{R}^{n+1}$ be a curve, then we can obtain

$$\frac{d}{dt}(v(\gamma(t))) = (\partial_t - \xi^* \cdot \nabla)v(\gamma(t)) = \frac{i}{\hbar}x^* \cdot (x - \xi^*t)v(\gamma(t)).$$

This is a one-dimensional ODE and the solution is straightforward,

$$v(\gamma(t)) = e^{\frac{i}{\hbar}[(x^* \cdot x)t - (x^* \cdot \xi^*)t^2/2]}v(\gamma(0)),$$

which is equivalent to

$$v(x - \xi^*t, t) = e^{\frac{i}{\hbar}[(x^* \cdot x)t - (x^* \cdot \xi^*)t^2/2]}v(x, 0).$$

By replacing x with $x + \xi^*t$ and substituting the boundary condition $v(x, 0) = u(x)$ into the solution above, we obtain

$$\begin{aligned} v(x, t) &= v((x + \xi^*t) - \xi^*t, t) \\ &= e^{\frac{i}{\hbar}[x^* \cdot (x + \xi^*t)t - (x^* \cdot \xi^*)t^2/2]}u(x + \xi^*t) \\ &= e^{\frac{i}{\hbar}[(x^* \cdot x)t + (x^* \cdot \xi^*)t^2/2]}u(x + \xi^*t) \\ \Rightarrow e^{\frac{it}{\hbar}l(x, hD)}u &= e^{\frac{i}{\hbar}[(x^* \cdot x)t + (x^* \cdot \xi^*)t^2/2]}u(x + \xi^*t). \end{aligned} \quad (7.34)$$

Second, we compute $(e^{\frac{it}{\hbar}l})^w(x, hD)u$. We have

$$\begin{aligned} (e^{\frac{it}{\hbar}l})^w(x, hD)u &= (2\pi\hbar)^{-n} \iint e^{i(x-y) \cdot \xi/h} e^{\frac{it}{\hbar}[x^* \cdot (\frac{x+y}{2}) + \xi^* \cdot \xi]} u(y) dy d\xi \\ &= e^{\frac{it}{\hbar}(x^* \cdot \frac{x}{2})} (2\pi\hbar)^{-n} \iint e^{i(x + \xi^*t - y) \cdot \xi/h} \cdot e^{\frac{it}{\hbar}(x^* \cdot \frac{y}{2})} u(y) dy d\xi \\ &= e^{\frac{it}{\hbar}(x^* \cdot \frac{x}{2})} \int \delta(x + \xi^*t - y) e^{\frac{it}{\hbar}(x^* \cdot \frac{y}{2})} u(y) dy \quad ((2\pi\hbar)^{-n} \int e^{ia \cdot \xi/h} d\xi = \delta(a)) \\ &= e^{\frac{it}{\hbar}(x^* \cdot \frac{x}{2})} e^{\frac{it}{\hbar}x^* \cdot \frac{(x + \xi^*t)}{2}} u(x + \xi^*t) = e^{\frac{i}{\hbar}[(x^* \cdot x)t + (x^* \cdot \xi^*)t^2/2]} u(x + \xi^*t), \end{aligned}$$

which is (7.32). From (7.34) and (7.32) we arrive at the first equality in the theorem.

For the composition relation, let $l(x, \xi) = x_1^* \cdot x + \xi_1^* \cdot \xi$ and $m(x, \xi) = x_2^* \cdot x + \xi_2^* \cdot \xi$, then from (7.34) we have

$$e^{\frac{it}{\hbar}(l+m)(x, hD)}u = e^{\frac{i}{\hbar}[(x_1^* + x_2^*) \cdot xt + (x_1^* + x_2^*) \cdot (\xi_1^* + \xi_2^*)t^2/2]}u(x + (\xi_1^* + \xi_2^*)t),$$

and

$$\begin{aligned}
e^{\frac{i\hbar}{\hbar}l(x,hD)}e^{\frac{i\hbar}{\hbar}m(x,hD)}u &= (e^{\frac{i\hbar}{\hbar}l})^w(x,hD) \circ e^{\frac{i\hbar}{\hbar}[(x_2^* \cdot x) + (x_2^* \cdot \xi_2^*)t^2/2]}u(x + \xi_2^*t) \\
&= e^{\frac{i\hbar}{\hbar}[(x_1^* \cdot x) + (x_1^* \cdot \xi_1^*)t^2/2]}e^{\frac{i\hbar}{\hbar}[(x_2^* \cdot (x + \xi_1^*t)) + (x_2^* \cdot \xi_2^*)t^2/2]}u(x + \xi_1^*t + \xi_2^*t) \\
&= e^{\frac{i\hbar}{\hbar}[(x_1^* + x_2^*) \cdot xt + (x_1^* \cdot \xi_1^* + 2x_2^* \cdot \xi_1^* + x_2^* \cdot \xi_2^*)t^2/2]}u(x + (\xi_1^* + \xi_2^*)t) \\
&= e^{\frac{i\hbar}{\hbar}[(x_1^* + x_2^*) \cdot xt + (x_1^* + x_2^*)(x_2^* + \xi_2^*)t^2/2]}u(x + (\xi_1^* + \xi_2^*)t) \cdot e^{\frac{i\hbar}{\hbar}[(x_2^* \cdot \xi_1^* - x_1^* \cdot \xi_2^*)t^2/2]} \\
&= e^{\frac{i\hbar}{2\hbar}\sigma((x_1^*, \xi_1^*), (x_2^*, \xi_2^*))}e^{\frac{i\hbar}{\hbar}(l+m)(x,hD)}u.
\end{aligned}$$

Readers should note that the σ here is the symplectic product defined in Definition 7.15.

The proof is complete. \square

By using (7.31), we can represent the corresponding Weyl quantization of a symbol by its Fourier transform.

LEMMA 7.22 (Fourier decomposition of a^w). *For any $a(x, \xi) \in \mathcal{S}(\mathbb{R}^{2n})$, we have*

$$a^w(x, hD) = (2\pi\hbar)^{-n} \int_{\mathbb{R}^{2n}} \mathcal{F}_h a(l) e^{\frac{i\hbar}{\hbar}l(x,hD)} dl,$$

where $l = (x^*, \xi^*) \in \mathbb{R}^{2n}$ and $l(x, hD)$ is defined as (7.12). This can directly generalize to the case where $a(x, \xi) \in \mathcal{S}'(\mathbb{R}^{2n})$ and then $\langle a^w(x, hD)u, v \rangle = (2\pi\hbar)^{-2n} \hat{a}(\langle e^{\frac{i\hbar}{\hbar}l(x,hD)}u, v \rangle)$ for $\forall u, v \in \mathcal{S}(\mathbb{R}^n)$.

REMARK 7.23. With the help of Lemma 7.22, every Weyl quantization can be represented by means of operators of the form $e^{\frac{i\hbar}{\hbar}l(x,hD)}$ where l is a linear form. Therefore, quantizations of the form $e^{\frac{i\hbar}{\hbar}l(x,hD)}$ plays an important role in semiclassical analysis.

PROOF OF LEMMA 7.22. When $a \in \mathcal{S}$, we have

$$a(x, \xi) = (2\pi\hbar)^{-n} \int_{\mathbb{R}^{2n}} e^{\frac{i\hbar}{\hbar}l(x,\xi)} \mathcal{F}_h a(l) dl,$$

thus by (7.31) we arrive at the statement. The case where $a(x, \xi) \in \mathcal{S}'(\mathbb{R}^{2n})$ is left as an exercise. \square

7.4. Applications in Carleman estimates

One of the examples of Carleman estimates is of the following

$$\tau^3 \|e^{\tau\phi}u\|^2 \lesssim \|e^{\tau\phi}Pu\|^2. \quad (7.35)$$

To prove it, we set $h = \tau^{-1}$, $v(x) = e^{\phi(x)/h}u(x)$ and denote an operator P_ϕ as

$$P_\phi: f \rightarrow e^{\phi/h}h^2P(e^{-\phi/h}f),$$

then (7.35) is equivalent to

$$\|P_\phi v\|^2 \gtrsim h\|v\|^2. \quad (7.36)$$

We assume

$$\sigma_{\text{scl}}(P_\phi) \in S(m) \quad (7.37)$$

for some order function m . Here we use $\sigma_{\text{scl}}(A)$ to signify the semiclassical symbol of A .

Set $A = (P_\phi + P_\phi^*)/2$ and $B = (P_\phi - P_\phi^*)/(2i)$, and denote

$$\sigma = \sigma_{\text{scl}}(ih^{-1}[A, B]),$$

then we can conclude

$$\|P_\phi v\|_{L^2}^2 = (P_\phi v, P_\phi v) = \|Av\|^2 + \|Bv\|^2 + (i[A, B]v, v) \geq h(\sigma(x, hD)v, v), \quad (7.38)$$

$$\sigma(x, hD) \text{ is self-adjoint, i.e. } \sigma(x, hD)^* = \sigma(x, hD). \quad (7.39)$$

Using (7.38), inequality (7.36) will be true if the following is true:

$$(\sigma(x, hD)v, v) \gtrsim \|v\|^2, \quad (7.40)$$

so (7.40) implies (7.35). It's left to prove (7.40).

To prove (7.40), we compute

$$\begin{aligned} \sigma &= \sigma_{\text{scl}}(ih^{-1}[A, B]) = \sigma_{\text{scl}}(ih^{-1}[(P_\phi + P_\phi^*)/2, (P_\phi - P_\phi^*)/(2i)]) \\ &= \frac{1}{2h} \sigma_{\text{scl}}([P_\phi^*, P_\phi]) = \frac{1}{2h} \left[\frac{h}{i} \{p_\phi^*, p_\phi\} + h^2 S(m^2) \right] \quad (\text{Corollary 7.13}) \\ &= \frac{1}{2i} \{ \overline{p_\phi} + hS(m), p_\phi \} + hS(m^2) \quad (\text{Theorem 7.14}) \\ &= \frac{1}{2i} \{ \overline{p_\phi}, p_\phi \} + hS(m^2) = \{ \Re p_\phi, \Im p_\phi \} + hS(m^2). \end{aligned} \quad (7.41)$$

The remainder term $S(m^2)$ comes from the assumption (7.37) and the fact $\sigma_{\text{scl}}(P_\phi^*) \in S(m)$. If

$$\boxed{\{ \Re p_\phi, \Im p_\phi \}(x, \xi) \geq Cm(x, \xi)^2, \quad (m(x, \xi) \geq 1)} \quad (7.42)$$

holds, from (7.41) we know when h is small enough we will have

$$|\sigma(x, \xi)| \gtrsim m(x, \xi)^2. \quad (7.43)$$

By combining (7.43), (7.39) and [28, Theorem 4.19], we can conclude (7.40). In summary, we have the following theorem.

THEOREM 7.24 (Carleman estimates). *Let P be a SΨDO and $\phi \in C^\infty(\mathbb{R}^n; \mathbb{R})$ and denote $P_\phi := e^{\phi/h} \circ h^2 P \circ e^{-\phi/h}$ and $p_\phi := \sigma_{\text{scl}}(P_\phi)$. Assume*

$$\begin{cases} \sigma_{\text{scl}}(P_\phi) \in S(m) \text{ for some order function } m(x, \xi) \geq 1 \\ \{ \Re p_\phi, \Im p_\phi \}(x, \xi) \geq Cm(x, \xi)^2, \end{cases}$$

then there exist positive constants C and τ_0 such that for $\forall \tau \geq \tau_0, \forall u \in \mathcal{S}(\mathbb{R}^n)$, there holds

$$\boxed{\tau^3 \|e^{\tau\phi} u\|^2 \leq C \|e^{\tau\phi} P u\|^2.}$$

Exercise

EXERCISE 7.1. Prove Corollary 7.13.

EXERCISE 7.2. Assume $y, \xi, s, \eta \in \mathbb{R}^n$, and define the $4n \times 4n$ matrix Σ by (7.25). Check

$$\begin{aligned} \frac{1}{2} \langle \Sigma(y, \xi, s, \eta), (y, \xi, s, \eta) \rangle &= s \cdot \xi - y \cdot \eta, \\ \frac{1}{2} \langle \Sigma D_{(y, \xi, s, \eta)}, D_{(y, \xi, s, \eta)} \rangle &= D_s \cdot D_\xi + \nabla_y \cdot \nabla_\eta. \end{aligned}$$

EXERCISE 7.3. Prove Lemma 7.20.

CHAPTER 8

The wavefront set

In this chapter we follow closely [4, §3].

8.1. Basic facts

Recall the Peetre's inequality (cf Lemma 5.2):

$$\langle a - b \rangle^m \leq \langle a \rangle^m \cdot \langle b \rangle^{|m|}, \quad \forall a, b \in \mathbb{R}^n \text{ and } \forall m \in \mathbb{R}. \quad (8.1)$$

We also need an angular separation inequality, which states that

$$|a - b| \geq C(|a| + |b|), \quad \forall a \in V_1, \forall b \in V_2, \quad (8.2)$$

provided that V_1 and V_2 are two cone in \mathbb{R}^n separating each other by a positive angle, and the positive constant C depends on this angle. One example is that $V_1 = \{\rho(\cos \alpha, \sin \alpha) \in \mathbb{R}^2; \rho \geq 0, 0 \leq \alpha \leq \pi/4\}$ and $V_2 = \{\rho(\cos \alpha, \sin \alpha) \in \mathbb{R}^2; \rho \geq 0, 3\pi/4 \leq \alpha \leq \pi\}$. From (8.2) we can easily derive

$$\langle a - b \rangle^{-m_1 - m_2} \leq C \langle a \rangle^{-m_1} \cdot \langle b \rangle^{-m_2}, \quad \forall a \in V_1, \forall b \in V_2, \forall m_1, m_2 \geq 0. \quad (8.3)$$

PROOF OF (8.3). From $|a - b| \gtrsim |a| + |b|$ we have $(1 + |a - b|)^{-1} \lesssim (1 + |a|)^{-1}$ and $(1 + |a - b|)^{-1} \lesssim (1 + |b|)^{-1}$, so $(1 + |a - b|)^{-m_1 - m_2} \lesssim (1 + |a|)^{-m_1} (1 + |b|)^{-m_2}$, which is equivalent to (8.3). \square

These inequalities are frequently used in microlocal analysis and sometimes play key role in the proofs of microlocal analysis. We use the notation $T^*\mathbb{R}^n \setminus 0$ to stand for the cotangent bundle with the zero section excluded. We deliberately exclude the zero section for some purpose, see Remark 8.13. We introduce the notion of conic sets, the smooth direction and the wavefront set as follows.

DEFINITION 8.1 (Conic set). A set $\Gamma \subset T^*\mathbb{R}^n \setminus 0$ is called a *conic set* if $\Gamma = \omega \times V$ for some $\omega \subset \mathbb{R}^n$ and some set $V \subset \mathbb{R}^n \setminus 0$, where the set V is conic in \mathbb{R}^n , i.e. if $\xi \in V$ then $t\xi \in V$ for all $t > 0$.

DEFINITION 8.2 (Smo). Let $m \in \mathbb{R}$ and $a \in S^m$, and A is the Ψ DO of a . Let $\Gamma \subset T^*\mathbb{R}^n \setminus 0$ be an open conic set. If for every integer N there exists a constant $C_{\Gamma, N}$ such that

$$|a(x, \xi)| \leq C_{\Gamma, N} \langle \xi \rangle^{-N}, \quad \forall (x, \xi) \in \Gamma, \quad (\text{or equivalently } \boxed{a \in S^{-\infty} \text{ in } \Gamma.}) \quad (8.4)$$

holds, we say Γ is a smooth direction set of a (and of A). We write $\boxed{\text{Smo}(A) := \bigcup \mathcal{F}}$ where $\mathcal{F} = \{\Gamma; \Gamma \text{ is a smooth direction set of } A\}$.

It can be checked that $\text{Smo}(a)$ is always open in $T^*\mathbb{R}^n \setminus 0$. We can also extend the $\text{Smo}(a)$ to a which is in $S^m(\mathbb{R}_x^n \times \mathbb{R}_\xi^N)$ (n and N need not to be the same). The idea of the smooth direction is that, for any symbol $a \in S^m$, no matter what the value of m is, there are chances that there exists some directions in ξ such that a decays at infinite speed in these direction.

LEMMA 8.3. Assume $A, B \in \Psi^{+\infty}$, then $\text{Smo}(A) \cup \text{Smo}(B) \subset \text{Smo}(A \circ B)$.

The proof is left as an exercise.

DEFINITION 8.4 (Wavefront set). Assume $\Omega \subset \mathbb{R}^n$ is a domain. For any given distribution $u \in \mathcal{D}'(\Omega)$, the *wavefront set* $\boxed{\text{WF}(u)} \subset T^*\Omega \setminus 0$ of u is defined as a closed subset such that, for any $(x_0, \xi_0) \notin \text{WF}(u)$, there exists a neighborhood ω of x_0 , a function $\varphi \in C_c^\infty(\Omega)$ with $\varphi(x_0) \neq 0$ and $\text{supp } \varphi \subset \omega$, and a cone neighborhood V of ξ_0 such that

$$|\widehat{\varphi u}(\xi)| \leq C_{N, \omega, V} \langle \xi \rangle^{-N}, \quad \forall N \in \mathbb{N}, \forall \xi \in V \quad (8.5)$$

holds for some positive constant $C_{N, \omega, V}$ depending on N , ω and specially on V .

EXAMPLE 8.5. Compute the wavefront set of $u(x_1, x_2) := H(x_1)$, where H is the Heaviside function. Fix a point (\bar{x}_1, \bar{x}_2) . When $\bar{x}_1 \neq 0$, we can always find a cutoff function $\varphi \in C_c^\infty(\mathbb{R}^2)$ such that $\varphi u \in C_c^\infty(\mathbb{R}^2)$, so $\widehat{\varphi u}$ will be rapidly decaying. This implies that

$$\{(\bar{x}_1, \bar{x}_2; \xi_1, \xi_2); \bar{x}_1 \neq 0\} \cap \text{WF}(u) = \emptyset, \quad (8.6)$$

so $\text{WF}(u)$ is made of points of the form $(0, x_2; \xi_1, \xi_2)$, thus in what follows we assume $\bar{x}_1 = 0$.

Fix cutoff functions $\varphi_1, \varphi_2 \in C_c^\infty(\mathbb{R}^1)$, such that φ_1 is supported in the neighborhood of 0 and φ_2 in the neighborhood of \bar{x}_2 , and denote $\varphi(x_1, x_2) = \varphi_1(x_1)\varphi_2(x_2)$, then

$$|\widehat{\varphi u}(\xi_1, \xi_2)| \simeq \left| \int_0^\infty e^{-ix_1\xi_1} \varphi_1(x_1) dx_1 \right| \cdot \left| \int e^{-ix_2\xi_2} \varphi_2(x_2) dx_2 \right|. \quad (8.7)$$

When $\xi_2 \neq 0$, we can continue (8.7) as

$$|\widehat{\varphi u}(\xi_1, \xi_2)| \simeq \int |\varphi_1(x_1)| dx_1 \cdot \langle \xi_2 \rangle^{-\infty} \lesssim \langle \xi_2 \rangle^{-\infty}. \quad (8.8)$$

For any cone $V_C := \{(\xi_1, \xi_2); |\xi_1| \leq C|\xi_2|\}$ where $C > 0$, we have $|\xi_2| \leq |x_1| + |\xi_2| \lesssim |\xi_2|$, which implies $\langle (\xi_1, \xi_2) \rangle \simeq \langle \xi_2 \rangle$. Hence, (8.8) becomes $|\widehat{\varphi u}(\xi_1, \xi_2)| \lesssim \langle \xi \rangle^{-\infty}$. Hence, for any constant $C > 0$, we have

$$\{(0, \bar{x}_2; \xi_1, \xi_2); \bar{x}_2 \in \mathbb{R}, |\xi_1| \leq C|\xi_2|\} \cap \text{WF}(u) = \emptyset. \quad (8.9)$$

Combining (8.6) and (8.9), we see that

$$\text{WF}(u) \subset \{(0, \bar{x}_2; \xi_1, 0); \bar{x}_2 \in \mathbb{R}\}. \quad (8.10)$$

Finally, we show

$$\text{WF}(u) \supset \{(0, \bar{x}_2; \xi_1, 0); \bar{x}_2 \in \mathbb{R}\}. \quad (8.11)$$

Fix $\bar{x}_2 \in \mathbb{R}$. For any $\varphi \in C_c^\infty(\mathbb{R}^2)$ supported in the neighborhood of $(0, \bar{x}_2)$, we have

$$\begin{aligned} \widehat{\varphi u}(\xi_1, \xi_2) &\simeq \int e^{-ix_1\xi_1} e^{-ix_2\xi_2} \varphi(x_1, x_2) H(x_1) dx_1 dx_2 \\ &= \int_0^\infty e^{-ix_1\xi_1} \tilde{\varphi}(x_1, \xi_2) dx_1 = i\xi_1^{-1} \tilde{\varphi}(0, \xi_2) + \xi_1^{-1} \int_0^\infty e^{-ix_1\xi_1} D_{x_1} \tilde{\varphi}(x_1, \xi_2) dx_1 \\ &= i\xi_1^{-1} \tilde{\varphi}(0, \xi_2) + \xi_1^{-1} (i\xi_1^{-1} D_{x_1} \tilde{\varphi}(0, \xi_2) + \xi_1^{-1} \int_0^\infty e^{-ix_1\xi_1} D_{x_1}^2 \tilde{\varphi}(x_1, \xi_2) dx_1) \\ &= i\xi_1^{-1} \tilde{\varphi}(0, \xi_2) + \mathcal{O}(|\xi_1|^{-2}), \end{aligned} \quad (8.12)$$

where $\tilde{\varphi}(x_1, \xi_2) = \int_{\mathbb{R}} e^{-ix_2\xi_2} \varphi(x_1, x_2) dx_2$. We know $\tilde{\varphi}(0, \xi_2) = \int_{\mathbb{R}} e^{-ix_2\xi_2} \varphi(0, x_2) dx_2$ is not compactly supported due to the uncertainty principle, so there is $\bar{\xi}_2 \neq 0$ such that $\tilde{\varphi}(0, \bar{\xi}_2) \neq 0$. For any cone $V'_C := \{(\xi_1, \xi_2); |\xi_2| \leq C|\xi_1|\}$ where $C > 0$, when $|\xi_1|$ is large enough we always have $(\xi_1, \xi_2) \in V'_C$. Hence, (8.12) means that in any cone V'_C , we have

$$|\widehat{\varphi u}(\xi_1, \bar{\xi}_2)| \simeq |\xi_1|^{-1} |\tilde{\varphi}(0, \bar{\xi}_2)| + \mathcal{O}(|\xi_1|^{-2}) \simeq |\xi_1|^{-1},$$

so φu is not rapidly decaying in any cone V'_C which contains $\{(\xi_1, 0)\}$ as their common part. By the definition of the wavefront set we can conclude

$$(0, \bar{x}_2; \xi_1, 0) \in \text{WF}(u),$$

which implies (8.11). Combining (8.10) with (8.11), we obtain

$$\underline{\text{WF}(u) = \{(0, x_2; \xi_1, 0); x_2 \in \mathbb{R}, \xi_1 \neq 0\}}, \quad \text{where } u(x_1, x_2) = H(x_1).$$

It is easy to see from the definition that $(\text{WF } u)^c$ is an open set, so every wavefront set is closed. In fact, we can relax the restriction on the function φ in the Definition (8.4) as follows.

LEMMA 8.6. *Using the same notation in Definition 8.4, if $(x_0, \xi_0) \notin \text{WF}(u)$, then there exists a another neighborhood $\omega' \subset \omega$ of x_0 , such that for any $\varphi \in C_c^\infty(\omega')$, which doesn't necessarily satisfy $\varphi(x_0) \neq 0$, the estimates (8.5) holds, with the constant $C_{N, \omega, V}$ now depends also on φ .*

PROOF. We call for the result (2) in Theorem 8.7 below in advance. Assume that $(x_0, \xi_0) \notin \text{WF}(u)$, then there exists a neighborhood ω of x_0 , a function $\varphi_0 \in C_c^\infty(\mathbb{R}^n)$ with $\varphi_0(x_0) \neq 0$ and a cone neighborhood V of ξ_0 such that (8.5) holds. Because $\varphi_0(x_0) \neq 0$ and φ_0 is continuous, there exists another neighborhood $\omega' \subset \omega$ of x_0 such that $|\varphi_0(x)| \geq |\varphi_0(x_0)/2| > 0$ for all $x \in \omega'$, and thus $1/\varphi_0(x)$ is well-defined in ω' ; the denominator keeps a positive distance from 0 in ω' . Now for any $\phi \in C_c^\infty(\omega')$, we know $\phi/\varphi_0 \in C_c^\infty(\omega')$, hence

$$\begin{aligned} |\widehat{\phi u}(\xi)| &= |(\phi/\varphi_0 \cdot \varphi_0 u)^\wedge(\xi)| \simeq \left| \int \widehat{\varphi_0 u}(\xi - \eta) \cdot \widehat{\phi/\varphi_0}(\eta) d\eta \right| \\ &\leq \int |\widehat{\varphi_0 u}(\xi - \eta)| \cdot |\widehat{\phi/\varphi_0}(\eta)| d\eta \lesssim \int \langle \xi - \eta \rangle^{-N} \cdot \langle \eta \rangle^{-N-n-1} d\eta \\ &\leq \langle \xi \rangle^{-N} \int \langle \eta \rangle^N \cdot \langle \eta \rangle^{-N-n-1} d\eta \lesssim \langle \xi \rangle^{-N}, \quad \forall N \in \mathbb{N}. \end{aligned}$$

Note that we used Peetre's inequality (8.1). The proof is complete. \square

The wavefront set possesses some simple facts [4].

THEOREM 8.7. *Assume that $u, v \in \mathcal{D}'(\Omega)$ and $a \in C_c^\infty(\Omega)$, then we have*

- (1) $\text{WF}(u + v) \subseteq \text{WF}(u) \cup \text{WF}(v)$;
- (2) $\text{WF}(au) \subseteq \text{WF}(u)$;
- (3) $\text{WF}(D^\alpha u) \subseteq \text{WF}(u)$.

PROOF. For (1). Assume that $(x_0, \xi_0) \notin \text{WF}(u) \cup \text{WF}(v)$, then $(x_0, \xi_0) \in (\text{WF}(u))^c \cap (\text{WF}(v))^c$, so there exists neighborhoods ω_1 and ω_2 of x_0 and cone neighborhoods V_1 and V_2 of ξ_0 such that

$$\begin{aligned} |\widehat{\varphi_{x_0} u}(\xi)| &\leq C \langle \xi \rangle^{-N}, \quad \forall \xi \in V_1, \forall \varphi_{x_0} \in \mathcal{D}(\omega_1) \text{ with } \varphi_{x_0}(x_0) \neq 0, \forall N \in \mathbb{N}, \\ |\widehat{\varphi_{x_0} v}(\xi)| &\leq C \langle \xi \rangle^{-N}, \quad \forall \xi \in V_2, \forall \varphi_{x_0} \in \mathcal{D}(\omega_2) \text{ with } \varphi_{x_0}(x_0) \neq 0, \forall N \in \mathbb{N}. \end{aligned}$$

Thus, we have

$$|\widehat{\varphi_{x_0} w}(\xi)| \leq C \langle \xi \rangle^{-N}, \quad \forall \xi \in V_1 \cap V_2, \forall \varphi_{x_0} \in \mathcal{D}(\omega_1 \cap \omega_2) \text{ with } \varphi_{x_0}(x_0) \neq 0, \forall N \in \mathbb{N},$$

where $w = u$ or v , so $(x_0, \xi_0) \notin \text{WF}(u + v)$. We can conclude (1).

For (2). Assume $(x_0, \xi_0) \notin \text{WF}(u)$, then there exists a neighborhood ω of x_0 , a function $\varphi \in \mathcal{D}(\mathbb{R}^n)$ with $\varphi(x_0) \neq 0$ and a cone neighborhood V of ξ_0 such that for all $\xi \in V$,

$$|\widehat{\varphi a u}(\xi)| = |(a \cdot \varphi u)^\wedge(\xi)| \simeq \left| \int \widehat{\varphi u}(\xi - \eta) \cdot \widehat{a}(\eta) d\eta \right|$$

$$\begin{aligned}
&\leq \int |\widehat{\varphi}u(\xi - \eta)| \cdot |\widehat{a}(\eta)| \, d\eta \lesssim \int \langle \xi - \eta \rangle^{-N} \cdot \langle \eta \rangle^{-N-n-1} \, d\eta \\
&\leq \langle \xi \rangle^{-N} \int \langle \eta \rangle^N \cdot \langle \eta \rangle^{-N-n-1} \, d\eta \lesssim \langle \xi \rangle^{-N}, \quad \forall N \in \mathbb{N}.
\end{aligned}$$

Therefore $(x_0, \xi_0) \notin \text{WF}(au)$. We can conclude (2).

For (3). Assume $(x_0, \xi_0) \notin \text{WF}(u)$. For any $\varphi \in \mathcal{D}(\omega')$ where the ω' is as in Lemma 8.6, we have

$$\begin{aligned}
\widehat{\phi D^\alpha u}(\xi) &\simeq \int e^{-ix \cdot \xi} \phi(x) D^\alpha u(x) \, dx \simeq \int D^\alpha (e^{-ix \cdot \xi} \phi(x)) u(x) \, dx \\
&= \int \sum_{|\beta| \leq |\alpha|} \binom{\alpha}{\beta} D^\beta (e^{-ix \cdot \xi}) D^{\alpha-\beta} \phi(x) u(x) \, dx \\
&= \sum_{|\beta| \leq |\alpha|} \xi^\beta \binom{\alpha}{\beta} \int e^{-ix \cdot \xi} (D^{\alpha-\beta} \phi \cdot u)(x) \, dx \\
&= \sum_{|\beta| \leq |\alpha|} \xi^\beta \binom{\alpha}{\beta} (D^{\alpha-\beta} \phi \cdot u)^\wedge(\xi).
\end{aligned}$$

Thus, by Lemma 8.6,

$$|\widehat{\phi D^\alpha u}(\xi)| \leq \sum_{|\beta| \leq |\alpha|} \xi^\beta \binom{\alpha}{\beta} |(D^{\alpha-\beta} \phi \cdot u)^\wedge(\xi)| \lesssim \sum_{|\beta| \leq |\alpha|} \binom{\alpha}{\beta} \langle \xi \rangle^{|\beta|} \langle \xi \rangle^{-N-|\alpha|} \lesssim \langle \xi \rangle^{-N},$$

for any $N \in \mathbb{N}$. Therefore $(x_0, \xi_0) \notin \text{WF}(D^\alpha u)$. We can conclude (3).

The proof is complete. \square

8.2. Wavefront set of product of distributions

In this section we deal with some more sophisticated cases of the computations of the wavefront sets.

8.2.1. Direct product. The first theorem is about the wavefront of the direct product $u \otimes v$. For $u: \mathcal{D}(\Omega_x) \rightarrow \mathbb{C}$ and $v: \mathcal{D}(\Omega_y) \rightarrow \mathbb{C}$, we define the *direct product* $u \otimes v$ of u and v as a distribution on $\mathcal{D}(\Omega_x \times \Omega_y)$ that maps $\varphi(x, y) \in \mathcal{D}(\Omega_x \times \Omega_y)$ to $\langle u, \langle v, \varphi(x, y) \rangle_y \rangle_x$,

$$\langle u \otimes v, \varphi(x, y) \rangle := \langle u, \langle v, \varphi(x, y) \rangle_y \rangle_x.$$

THEOREM 8.8. *For any given distributions $u \in \mathcal{D}'(\Omega_x)$ and $v \in \mathcal{D}'(\Omega_y)$, the wavefront set of the direct product $u \otimes v$ satisfies*

$$\boxed{\text{WF}(u \otimes v) \subseteq (\text{WF}(u) \times \text{WF}(v)) \cup (\text{WF}(u) \times \text{supp}_0 v) \cup (\text{supp}_0 u \times \text{WF}(v))}, \quad (8.13)$$

where $\text{supp}_0 u := \{(x, 0); x \in \text{supp } u\}$, $\text{supp}_0 v := \{(y, 0); y \in \text{supp } v\}$.

PROOF. Assume that $(x_0, y_0; \xi_0, \eta_0)$ doesn't belong to the right-hand-side of (8.13).

For the case where $\xi_0 \neq 0$ and $\eta_0 \neq 0$, we know $(x_0; \xi_0) \notin \text{WF}(u)$ and $(y_0; \eta_0) \notin \text{WF}(v)$, so the Fourier transform $(\varphi_{(x_0, y_0)} u \otimes v)^\wedge(\xi_0, \eta_0)$ cannot have the decay of the order $\langle (\xi, \eta) \rangle^{-N}$ for any $N \in \mathbb{N}$. Therefore, $(x_0, y_0; \xi_0, \eta_0) \notin \text{WF}(u \otimes v)$.

For the case where $\xi_0 = 0$ and $\eta_0 \neq 0$, if $x_0 \notin \text{supp } u$, obviously we can conclude $(x_0, y_0; \xi_0, \eta_0) \notin \text{WF}(u \otimes v)$, so we suggest that $x_0 \in \text{supp } u$, thus we must have $(y_0; \eta_0) \notin \text{WF}(v)$. Choose $\varphi(x, y) = \varphi_1(x) \varphi_2(y)$ as the cutoff function where $\varphi_1 \in \mathcal{D}(\omega_1)$ and ω_1 is some neighborhood of x_0 . So does φ_2 accordingly. Thus we have

$$(\varphi u \otimes v)^\wedge(\xi, \eta) = (\varphi_1 u)^\wedge(\xi) \cdot (\varphi_2 v)^\wedge(\eta).$$

We have that $(\varphi_2 v)^\wedge(\eta)$ is rapidly decaying and $(\varphi_1 u)^\wedge(\xi)$ grows in polynomial order of ξ in a cone neighborhood of $(0, \eta)$. It's easy to check that, in such a cone neighborhood, we have $\langle(\xi, \eta)\rangle \lesssim \langle\eta\rangle \lesssim \langle(\xi, \eta)\rangle$. Therefore,

$$\begin{aligned} |(\varphi u \otimes v)^\wedge(\xi, \eta)| &= |(\varphi_1 u)^\wedge(\xi)| \cdot |(\varphi_2 v)^\wedge(\eta)| \lesssim \langle\eta\rangle^{-N+l} \cdot \langle\xi\rangle^{-l} \\ &\lesssim \langle(\xi, \eta)\rangle^{-N+l} \cdot \langle(\xi, \eta)\rangle^{-l} = \langle(\xi, \eta)\rangle^{-N}, \end{aligned}$$

for any $N \in \mathbb{N}$. Therefore, $(x_0, y_0; \xi_0, \eta_0) \notin \text{WF}(u \otimes v)$.

The case where $\xi_0 \neq 0$ and $\eta_0 = 0$ is similar to the case where $\xi_0 = 0$ and $\eta_0 \neq 0$.

The proof is complete. \square

8.2.2. Product. Next, we investigate the product of two distributions. In contrast to the product of functions, the product of two distributions is not always well-defined. Under certain conditions, the product of two distributions can be defined, at least locally. We know that if $\varphi \in C_c^\infty(\Omega)$ and $u \in \mathcal{D}'(\Omega)$, we have $\varphi u \in \mathcal{E}'(\Omega)$ and thus the Fourier transform $\widehat{\varphi u}$ is well-defined and can be estimated of polynomial order at infinity. Thus we might have chance to define the product by using convolution,

$$(\varphi^2 uv)^\wedge(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} (\varphi u)^\wedge(\xi - \eta) \cdot (\varphi v)^\wedge(\eta) d\eta, \quad (8.14)$$

as long as the convolution (8.14) is integrable in the Lebesgue sense and grows under polynomial order in terms of $\langle\xi\rangle$ at infinity, which implies $\varphi^2 uv \in \mathcal{E}'(\Omega)$. This leads to the following result.

THEOREM 8.9 (Product Theorem). *For any given distributions $u, v \in \mathcal{D}'(\Omega)$, when*

$$(\text{WF}(u) + \text{WF}(v)) \cap O_x = \emptyset, \quad (8.15)$$

where $\text{WF}(u) + \text{WF}(v) := \{(x, \xi_1 + \xi_2); (x, \xi_1) \in \text{WF}(u), (x, \xi_2) \in \text{WF}(v)\}$, and $O_x := \{(x, 0); x \in \Omega\}$, the product “ uv ” can be well-defined in the sense of (8.14) and its wavefront set satisfies

$$\boxed{\text{WF}(uv) \subseteq (\text{WF}(u) + \text{WF}(v)) \cup \text{WF}(u) \cup \text{WF}(v)}. \quad (8.16)$$

PROOF. We partially follow [10, Proposition 11.2.3]. The proof is divided into two parts: first, we show that under condition (8.15) the convolution (8.14) can be controlled at polynomial of ξ ; second, we show the relation (8.16).

Step 1. For any open cone neighborhood V_3 of $\text{WF}(u) + \text{WF}(v)$, there exists open cone neighborhoods V_1' and V_2' of $\text{WF}(u)$ and $\text{WF}(v)$, respectively, such that $V_1' + V_2' \subset V_3$. Also, there must exist open cone neighborhoods V_1 and V_2 such that

$$\begin{cases} \text{WF}(u) \subsetneq V_1 \subsetneq V_1' \\ \text{WF}(v) \subsetneq V_2 \subsetneq V_2' \\ \text{WF}(u) + \text{WF}(v) \subsetneq V_1 + V_2 \subsetneq V_1' + V_2' \subsetneq V_3 \\ (V_1 + V_2) \cap O_x = \emptyset \end{cases} \quad (8.17)$$

The V_1' and V_2' will be utilized in **Step 2**.

Fix some $x_0 \in \Omega$, we can find some $\varphi \in \mathcal{D}(\Omega)$ with $\varphi(x_0) \neq 0$ and also φ guarantees φu and φv that (8.5) hold. For any fixed $\xi_0 \in \mathbb{R}^n \setminus \{0\}$, the integral (8.14) can be divided into four parts,

$$\begin{aligned} &\int_{\mathbb{R}^n} (\varphi u)^\wedge(\xi_0 - \eta) \cdot (\varphi v)^\wedge(\eta) d\eta \\ &= \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \notin V_2\}}} (\varphi u)^\wedge(\xi_0 - \eta) \cdot (\varphi v)^\wedge(\eta) d\eta + \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \in V_2\}}} (\varphi u)^\wedge(\xi_0 - \eta) \cdot (\varphi v)^\wedge(\eta) d\eta \end{aligned}$$

$$\begin{aligned}
& + \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \in V_1 \\ (x_0, \eta) \notin V_2\}}} (\varphi u)^\wedge(\xi_0 - \eta) \cdot (\varphi v)^\wedge(\eta) \, d\eta + \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \in V_1 \\ (x_0, \eta) \in V_2\}}} (\varphi u)^\wedge(\xi_0 - \eta) \cdot (\varphi v)^\wedge(\eta) \, d\eta \\
& =: I_1 + I_2 + I_3 + I_4.
\end{aligned} \tag{8.18}$$

The condition (8.15) will (only) be used to estimate I_4 .

According to Definition 8.4 and Peetre's inequality, we can estimate I_1 as

$$\begin{aligned}
|I_1| & \leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \notin V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| \, d\eta \\
& \lesssim \int_{\mathbb{R}^n} \langle \xi_0 - \eta \rangle^{-N} \cdot \langle \eta \rangle^{-N-n-1} \, d\eta \lesssim \langle \xi_0 \rangle^{-N} \int_{\mathbb{R}^n} \langle \eta \rangle^N \cdot \langle \eta \rangle^{-N-n-1} \, d\eta \\
& \lesssim \langle \xi_0 \rangle^{-N}, \quad \forall N \in \mathbb{N}.
\end{aligned} \tag{8.19}$$

For I_2 , we know that $\varphi v \in \mathcal{E}'(\Omega)$, so $|(\varphi v)^\wedge(\eta)|$ can be dominated by $\langle \eta \rangle^l$ for some $l \in \mathbb{N}$, thus

$$\begin{aligned}
|I_2| & \leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \in V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| \, d\eta \\
& \lesssim \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \in V_2\}}} \langle \xi_0 - \eta \rangle^{-l-n-1} \cdot \langle \eta \rangle^l \, d\eta \\
& \lesssim \langle \xi_0 \rangle^{l+n+1} \int_{\mathbb{R}^n} \langle \eta \rangle^{-l-n-1} \cdot \langle \eta \rangle^l \, d\eta \quad (\text{Peetre's inequality}) \\
& \lesssim \langle \xi_0 \rangle^{l+n+1}.
\end{aligned} \tag{8.20}$$

The estimation of I_3 is similar to that of I_2 ,

$$\begin{aligned}
|I_3| & \leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \in V_1 \\ (x_0, \eta) \notin V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| \, d\eta \\
& = \int_{\substack{\{\eta; (x_0, \gamma) \in V_1 \\ (x_0, \xi_0 - \gamma) \notin V_2\}}} |(\varphi u)^\wedge(\gamma)| \cdot |(\varphi v)^\wedge(\xi_0 - \gamma)| \, d\gamma \quad (\gamma = \xi_0 - \eta) \\
& \lesssim \int_{\substack{\{\eta; (x_0, \gamma) \in V_1 \\ (x_0, \xi_0 - \gamma) \notin V_2\}}} \langle \gamma \rangle^{l'} \cdot \langle \xi_0 - \gamma \rangle^{-l'-n-1} \, d\gamma \\
& \lesssim \langle \xi_0 \rangle^{l'+n+1} \int_{\mathbb{R}^n} \langle \gamma \rangle^{l'} \cdot \langle \gamma \rangle^{-l'-n-1} \, d\gamma \quad (\text{Peetre's inequality}) \\
& \lesssim \langle \xi_0 \rangle^{l'+n+1}.
\end{aligned} \tag{8.21}$$

For I_4 , we can show that the domain of integration $\{\eta; (x_0, \xi_0 - \eta) \in V_1, (x_0, \eta) \in V_2\}$ is bounded. We temporarily use $\hat{\eta}$ to mean the direction of η , $\hat{\eta} = \eta/|\eta|$. Therefore the direction of the vector $\xi_0 - \eta$ is parallel to $\xi_0/|\eta| - \hat{\eta}$, thus when $|\eta|$ is large enough, $(x_0, \xi_0 - \eta)$ will be in $-V_2 := \{(x, -\eta); (x, \eta) \in V_2\}$. We know $(x_0, \xi_0 - \eta) \in V_1$, so the set $\{(x_0, \gamma); (x_0, \gamma) \in V_1, (x_0, -\gamma) \in V_2\}$ is not empty. This contradict with $(V_1 + V_2) \cap O_x = \emptyset$ in (8.17). Therefore, when $|\eta|$ is large enough, the conditions $(x_0, \xi_0 - \eta) \in V_1$ and $(x_0, \eta) \in V_2$ cannot be satisfies simultaneously, which implies the set $\{\eta; (x_0, \xi_0 - \eta) \in V_1, (x_0, \eta) \in V_2\}$ is bounded. Therefore,

$$|I_4| \leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \notin V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| \, d\eta$$

$$\begin{aligned}
&\lesssim \int_{\{\eta; |\eta| \text{ bounded}\}} \langle \xi_0 - \eta \rangle^{l'} \cdot \langle \eta \rangle^l d\eta \\
&\lesssim \langle \xi_0 \rangle^{l'} \int_{\{\eta; |\eta| \text{ bounded}\}} \langle \eta \rangle^{l'} \cdot \langle \eta \rangle^l d\eta \quad (\text{Peetre's inequality}) \\
&\lesssim \langle \xi_0 \rangle^{l'}.
\end{aligned} \tag{8.22}$$

From (8.18)-(8.22), we conclude that the convolution (8.14) is Lebesgue integrable and grows with polynomial order in terms of $\langle \xi_0 \rangle$, thus $\varphi^2 uv \in \mathcal{E}'(\Omega)$. Now $uv \in \mathcal{D}'(\Omega)$ is well-defined.

Step 2. Under condition (8.15), we study the wavefront set of uv . Assume that

$$(x_0, \xi_0) \notin V_3 \cup V_1' \cup V_2', \tag{8.23}$$

Again, the condition (8.15) will (only) be used to estimate I_4 . Note the particular arrangements of the V_1, V_1' and V_2, V_2' in (8.23) and (8.18). We will utilize these arrangements combining with condition (8.17) to estimates I_2 and I_3 .

We estimate I_1 the same way as in **Step 1**, i.e. as in (8.19).

For I_2 , to get the rapid decay w.r.t. ξ_0 , we shall adapt different strategy. We know that $\varphi v \in \mathcal{E}'(\Omega)$, so $|(\varphi v)^\wedge(\eta)|$ can be dominated by $\langle \eta \rangle^l$ for some $l \in \mathbb{N}$. Thanks to the condition (8.23), we know $(x_0, \xi_0) \notin V_2'$ and now $(x_0, \eta) \in V_2$. Because $V_2 \subsetneq V_2'$, we know that V_2 and V_2' are separated with a positive angle, so the inequality (8.3) can apply to $\langle \xi_0 - \eta \rangle$,

$$\begin{aligned}
|I_2| &\leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \in V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| d\eta \\
&\lesssim \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \notin V_1 \\ (x_0, \eta) \in V_2\}}} \langle \xi_0 - \eta \rangle^{-N-l-n-1} \cdot \langle \eta \rangle^l d\eta \\
&\lesssim \langle \xi_0 \rangle^{-N} \int_{\mathbb{R}^n} \langle \eta \rangle^{-l-n-1} \cdot \langle \eta \rangle^l d\eta \quad (\text{by (8.3)}) \\
&\lesssim \langle \xi_0 \rangle^{-N}, \quad \forall N \in \mathbb{N}.
\end{aligned} \tag{8.24}$$

The estimation of I_3 is similar to (8.24),

$$\begin{aligned}
|I_3| &\leq \int_{\substack{\{\eta; (x_0, \xi_0 - \eta) \in V_1 \\ (x_0, \eta) \notin V_2\}}} |(\varphi u)^\wedge(\xi_0 - \eta)| \cdot |(\varphi v)^\wedge(\eta)| d\eta \\
&= \int_{\substack{\{\eta; (x_0, \gamma) \in V_1 \\ (x_0, \xi_0 - \gamma) \notin V_2\}}} |(\varphi u)^\wedge(\gamma)| \cdot |(\varphi v)^\wedge(\xi_0 - \gamma)| d\gamma \quad (\gamma = \xi_0 - \eta) \\
&\lesssim \int_{\substack{\{\eta; (x_0, \gamma) \in V_1 \\ (x_0, \xi_0 - \gamma) \notin V_2\}}} \langle \gamma \rangle^{l'} \cdot \langle \xi_0 - \gamma \rangle^{-N-l'-n-1} d\gamma \\
&\lesssim \langle \xi_0 \rangle^{-N} \int_{\mathbb{R}^n} \langle \gamma \rangle^{l'} \cdot \langle \gamma \rangle^{-l'-n-1} d\gamma \quad (\text{by (8.3)}) \\
&\lesssim \langle \xi_0 \rangle^{-N}, \quad \forall N \in \mathbb{N}.
\end{aligned} \tag{8.25}$$

Now we work on I_4 . From (8.15), (8.17) and (8.23), we know that $\xi_0 \notin V_1 + V_2$, thus the set $\{\eta; (x_0, \xi_0 - \eta) \in V_1, (x_0, \eta) \in V_2\}$ is empty. Therefore $I_4 = 0$. Combining this fact with (8.18), (8.19), (8.24) and (8.25), we arrive at

$$|(\varphi^2 uv)^\wedge(\xi)| \leq C_N \langle \xi_0 \rangle^{-N}, \quad \forall N \in \mathbb{N},$$

for $(x_0, \xi_0) \notin V_3 \cup V_1' \cup V_2'$. This implies $\text{WF}(u + v) \subset V_3 \cup V_1' \cup V_2'$. The sets V_3 , V_1' and V_2' can be close to $\text{WF}(u) + \text{WF}(v)$, $\text{WF}(u)$ and $\text{WF}(v)$, respectively, as close as possible, so we arrive at (8.16). The proof is complete. \square

8.2.3. Convolution. We define

$$\begin{cases} \text{WF}'(K) := \{(x, y; \xi, -\eta); (x, y; \xi, \eta) \in \text{WF}(K)\}, \\ \text{WF}_x(K) := \{(x; \xi); \exists y \text{ s.t. } (x, y; \xi, 0) \in \text{WF}(K)\}, \\ A \circ B := \{(x, \xi); \exists (y, \eta) \in B \text{ s.t. } (x, y; \xi, \eta) \in A\}, \\ O_x := \{(x, 0); x \in \Omega\}. \end{cases} \quad (8.26)$$

We need the following lemma.

LEMMA 8.10. *Assume $f \in \mathcal{D}'(\Omega \times \Omega)$, and there is a compact set $\mathcal{K} \subset \Omega$ such that $\text{supp } f \subset \Omega \times \mathcal{K}$. Then*

$$\boxed{\text{WF} \left(\int f(x, y) dy \right) = \text{WF}_x(f)}. \quad (8.27)$$

PROOF. Step 1. (\supset). Assume $(x_0, \xi_0) \notin \text{WF} \left(\int f(x, y) dy \right)$, then there exists $\chi_{x_0} \in C_c^\infty(\Omega)$ such that

$$\int e^{-ix_0 \cdot \xi_0} \chi_{x_0}(x) f(x, y) dy dx = \mathcal{O}(\langle \xi_0 \rangle^{-\infty}) = \mathcal{O}(\langle (\xi_0, 0) \rangle^{-\infty}).$$

which gives

$$\forall \bar{y} \in \mathcal{K}, \int e^{-i(x_0, \bar{y}) \cdot (\xi_0, 0)} \chi_{x_0}(x) \chi(y) f(x, y) d(x, y) = \mathcal{O}(\langle (\xi_0, 0) \rangle^{-\infty}),$$

where $\chi \in C_c^\infty(\Omega)$ with $\chi \equiv 1$ on \mathcal{K} . This means $(x_0, \bar{y}; \xi_0, 0) \notin \text{WF}(f)$ for $\forall \bar{y} \in \mathcal{K}$, so $(x_0, \xi_0) \notin \text{WF}_x(f)$. Hence,

$$\text{WF} \left(\int f(x, y) dy \right) \supset \text{WF}_x(f).$$

Step 2. (\subset). Assume $(x_0, \xi_0) \notin \text{WF}_x(f)$, then for $\forall \bar{y} \in \Omega$ we have $(x_0, \bar{y}; \xi_0, 0) \notin \text{WF}(f)$. Therefore, for $\forall \bar{y} \in \Omega$, there is a neighborhood of \bar{y} such that

$$\int e^{-i(x_0, \bar{y}) \cdot (\xi_0, 0)} \chi_{x_0}(x) \chi(y) f(x, y) d(x, y) = \mathcal{O}(\langle (\xi_0, 0) \rangle^{-\infty}), \quad (8.28)$$

for $\chi \equiv 1$ in that neighborhood. Because \mathcal{K} is compact, so by using partition of unity technique, we can remove the term $\chi(y)$ in (8.28), and obtain

$$\int e^{-ix_0 \cdot \xi_0} \chi_{x_0}(x) \left(\int f(x, y) dy \right) dx = \mathcal{O}(\langle \xi_0 \rangle^{-\infty}),$$

which gives $(x_0, \xi_0) \notin \text{WF}(\int f(x, y) dy)$. Hence,

$$\text{WF} \left(\int f(x, y) dy \right) \subset \text{WF}_x(f).$$

The proof is done. \square

THEOREM 8.11. *Assume $u \in \mathcal{E}'(\Omega)$, and $K \in \mathcal{D}'(\Omega \times \Omega)$. When $(\text{WF}'(K) \circ \text{WF}(u)) \cap O_x = \emptyset$, the distribution*

$$w(x) := \langle K(x, y), u(y) \rangle_y$$

is well-defined in the sense that

$$\forall \varphi \in \mathcal{D}(\Omega), \quad w(\varphi) := \langle K(x, y), u(y) \otimes \varphi(x) \rangle,$$

and we have the following *canonical relation*:

$$\boxed{\text{WF}(w) \subseteq (\text{WF}'(K) \circ \text{WF}(u)) \cup \text{WF}_x(K)}. \quad (8.29)$$

REMARK 8.12. Note that the $\text{supp } u$ should be contained in Ω , otherwise the w may be ill-defined.

PROOF. **Step 1.** Turn into product. Denote $\tilde{u}(x, y) = 1(x) \otimes u(y)$ where $1(x)$ is the constant function. The wavefront set of the function $1(x)$ is empty, so by Theorem 8.8 we have

$$\begin{aligned} \text{WF}(\tilde{u}) &\subset (\text{WF}(1) \times \text{WF}(u)) \cup (\text{WF}(1) \times \text{supp}_0 u) \cup (\text{supp}_0 1 \times \text{WF}(u)) \\ &= \emptyset \cup \emptyset \cup (\text{supp}_0 1 \times \text{WF}(u)) \\ &= \{(x, y; 0, \eta); x \in \text{supp } \Omega, (y, \eta) \in \text{WF}(u)\}. \end{aligned} \quad (8.30)$$

The $w(x)$ can be written as

$$w(x) = \langle K(x, y), u(y) \rangle_y = \int K(x, y) \cdot \tilde{u}(x, y) dy = \int K \tilde{u}(x, y) dy.$$

where $K \tilde{u}$ stands for the product of K and \tilde{u} . By Theorem 8.9, to guarantee the product $K \tilde{u}$ is well-defined, we need to check if the prerequisite

$$(\text{WF}(K) + \text{WF}(\tilde{u})) \cap O_{x,y} = \emptyset \quad (8.31)$$

is true. It can be shown that the condition $(\text{WF}'(K) \circ \text{WF}(u)) \cap O_x = \emptyset$ guarantees (8.31) (see Exercise 8.2), so $K \tilde{u}$ is well-defined.

Because $u \in \mathcal{E}'(\Omega)$, we see that for $\forall x \in \Omega$, $\text{supp } K \tilde{u}(x, \cdot)$ is uniformly compact, so by Lemma 8.10 we have $\text{WF}(\int K \tilde{u}(x, y) dy) = \text{WF}_x(K \tilde{u})$, so,

$$\text{WF}(w) = \text{WF}\left(\int K \tilde{u}(x, y) dy\right) = \text{WF}_x(K \tilde{u}) = \text{WF}(K \tilde{u}) \circ O_y, \quad (8.32)$$

where we used the fact that for general distribution $f \in \mathcal{D}'(\Omega \times \Omega)$,

$$\text{WF}_x(f) = \text{WF}(f) \circ O_y.$$

Step 2. Use product Theorem. Combining (8.32) with Theorem 8.9, we can have

$$\begin{aligned} \text{WF}(w) &= \text{WF}(K \tilde{u}) \circ O_y \subset \left((\text{WF}(K) + \text{WF}(\tilde{u})) \cup \text{WF}(K) \cup \text{WF}(\tilde{u}) \right) \circ O_y \\ &= M_1 \cup M_2 \cup M_3, \end{aligned} \quad (8.33)$$

where

$$\begin{cases} M_1 := (\text{WF}(K) + \text{WF}(\tilde{u})) \circ O_y, \\ M_2 := \text{WF}(K) \circ O_y, \\ M_3 := \text{WF}(\tilde{u}) \circ O_y. \end{cases}$$

The set $\text{WF}(K) + \text{WF}(\tilde{u})$ can be expressed as

$$\begin{aligned} &\text{WF}(K) + \text{WF}(\tilde{u}) \\ &= \{(x, y; \xi, \eta); \xi = \xi_1 + \xi_2, \eta = \eta_1 + \eta_2, (x, y; \xi_1, \eta_1) \in \text{WF}(K), (x, y; \xi_2, \eta_2) \in \text{WF}(\tilde{u})\} \\ &= \{(x, y; \xi, \eta); \eta = \eta_1 + \eta_2, (x, y; \xi, \eta_1) \in \text{WF}(K), (y, \eta_2) \in \text{WF}(u)\}. \quad (\text{by (8.30)}) \end{aligned}$$

Thus,

$$\begin{aligned} M_1 &= (\text{WF}(K) + \text{WF}(\tilde{u})) \circ O_y = \{(x, \xi); (x, y; \xi, -\eta) \in \text{WF}(K), (y, \eta) \in \text{WF}(u)\} \\ &= \{(x, \xi); (x, y; \xi, \eta) \in \text{WF}'(K), (y, \eta) \in \text{WF}(u)\} = \text{WF}'(K) \circ \text{WF}(u). \end{aligned} \quad (8.34)$$

By (8.30) it can also be checked that

$$M_2 = \text{WF}_x(K), \quad M_3 = \emptyset. \quad (8.35)$$

Combining (8.34), (8.35) with (8.33), we obtain (8.29). The proof is complete. \square

REMARK 8.13. In Theorem 8.11, if we know in advance that

$$\text{WF}(K) \subset (T^*\Omega_x \setminus 0) \times (T^*\Omega_y \setminus 0),$$

then $\text{WF}_x(K) = \emptyset$ and (8.29) can be reduced to

$$\boxed{\text{WF}(w) \subseteq \text{WF}'(K) \circ \text{WF}(u)}. \quad (8.36)$$

The set $\text{WF}'(K)$ is called the *twist* of $\text{WF}(K)$, and the operation “ $\text{WF}'(K) \circ$ ” is called *canonical relation* of the operator:

$$u(y) \mapsto w(x) := \langle K(x, y), u(y) \rangle$$

which takes K as its kernel. These can be generalized to the theory of *Fourier integral operators*.

8.3. The wavefront sets of Fourier integral operators

Recall the notion of phase function given in Definition 3.1.

THEOREM 8.14. *Assume $\varphi \in C^\infty(\mathbb{R}^n \times \mathbb{R}^N)$ is a phase function of order 1, and $a \in S^m$ is a symbol. Define $A(x)$ as*

$$A(x) := \int e^{i\varphi(x, \theta)} a(x, \theta) d\theta, \quad (8.37)$$

where the integral is understood as an oscillatory integral. Then A induces a distribution (also denoted as A) $A \in \mathcal{D}'(\Omega)$ for any domain $\Omega \subset \mathbb{R}^n$, i.e. $A: u \in \mathcal{D}(\Omega) \mapsto I_\varphi(au)$ by

$$A(u) := I_\varphi(au) = \langle A, u \rangle = \int e^{i\varphi(x, \theta)} a(x, \theta) u(x) dx d\theta$$

in oscillatory integral sense. The wavefront set of A satisfies

$$\boxed{\text{WF}(A) \subset \{(x, \varphi_x(x, \theta)); \varphi_\theta(x, \theta) = 0, (x, \theta) \notin \text{Smo}(a)\}}. \quad (8.38)$$

REMARK 8.15. When the following conditions are satisfied, the inclusion “ \subset ” in (8.38) can be improved to “ $=$ ” (see contexts preceding [7, Theorem 3.9], [5, Theorem 3.9]):

- (1) the phase function φ is non-degenerate on $C_\varphi := \{(x, \theta); \varphi_\theta(x, \theta) = 0, (x, \theta) \notin \text{Smo}(a)\}$, i.e. the $N \times (n + N)$ matrix $d\varphi_\theta$ is full rank on C_φ , here

$$d\varphi_\theta(x, \theta) = (\phi_{\theta x}(x, \theta) \quad \phi_{\theta\theta}(x, \theta));$$

- (2) the map $(x, \theta) \mapsto (x, \varphi_x(x, \theta))$ is injective when restricted to C_φ .

Readers may distinguish the A appeared in Theorem 8.14 with the operator B defined as

$$Bu(x) := \int e^{i\varphi(x, y, \theta)} a(x, y, \theta) u(y) dy d\theta.$$

The A is a distribution while the B just defined is an operator, namely, A maps a function to a scalar while B maps a function to another function.

However, A is a generalization of B , because

$$\begin{aligned} \langle Bu(x), v(x) \rangle &= \int e^{i\varphi(x, y, \theta)} a(x, y, \theta) u(y) v(x) dx dy d\theta \\ &= \left\langle \int e^{i\varphi(x, y, \theta)} a(x, y, \theta) d\theta, (v \otimes u)(x, y) \right\rangle \end{aligned}$$

$$= \langle K_B, v \otimes u \rangle.$$

where \tilde{B} is defined as

$$K_B(x, y) := \int e^{i\varphi(x, y, \theta)} a(x, y, \theta) d\theta.$$

Hence the operator B can be turned into a form of (8.37).

Moreover, we have $Bu(x) = \langle K_B(x, y), u(y) \rangle$, so by combining Theorems 8.14 and 8.11, hopefully we can obtain $\text{WF}(Bu)$.

SHORT PROOF OF THEOREM 8.14. This short proof is for summarizing the key idea of proving this theorem and thus the details may not be rigorously correct. After this short proof, we also present a formal proof of Theorem 8.14.

According to the Definition 8.4, we fix a cutoff function ϕ with $\phi(x_0) \neq 0$ and compute

$$\widehat{\phi A}(\xi) = A(\phi e^{-ix \cdot \xi}) = (2\pi)^{-n/2} \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) a(x, \theta) dx d\theta,$$

and the basic idea is to use N times (with N large enough) the operator $L := \frac{(\varphi_x(x, \theta) - \xi) \cdot \nabla_x}{i|\varphi_x(x, \theta) - \xi|^2}$ acting on $e^{i(\varphi(x, \theta) - x \cdot \xi)}$ and the fact (8.3) to get the desired estimate. But in order to do so, one needs to first address some singularities in the oscillatory integral. We have

$$\begin{aligned} \widehat{\phi A}(\xi) &\simeq \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} a(x, \theta) d\theta \right) dx \\ &\simeq \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\mathbb{R}_\theta^n} \left(\frac{\varphi_\theta(x, \theta) \cdot \nabla_\theta}{|\varphi_\theta(x, \theta)|^2} \right)^N e^{i\varphi(x, \theta)} a(x, \theta) d\theta \right) dx \\ &\sim \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} |\varphi_\theta(x, \theta)|^{-N} \partial_\theta^N a(x, \theta) d\theta \right) dx. \end{aligned} \quad (8.39)$$

Then as N be large enough, the $\partial_\theta^N a(x, \theta)$ will be integrable w.r.t. θ . But we notice that $|\varphi_\theta(x, \theta)|^{-1}$ has singularity at $\theta = 0$, so we first exclude the neighborhood of the origin of θ by using a cutoff function χ with $\chi(0) \neq 0$ as follows

$$\begin{aligned} \widehat{\phi A}(\xi) &\simeq \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) \chi(\theta) a(x, \theta) dx d\theta + \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \\ &= \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} \chi(\theta) a(x, \theta) d\theta \right) dx \\ &\quad + \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\Gamma_x} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx \\ &\quad + \int_{\theta \notin \Gamma_x} e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \\ &=: I_1(\xi) + I_2(\xi) + I_3(\xi), \end{aligned}$$

where

$$\Gamma_x := \{\theta; a(x, \theta) = \mathcal{O}(|\theta|^{-\infty})\}. \quad (8.40)$$

The $I_1(\xi)$ and $I_2(\xi)$ are $O(|\xi|^{-\infty})$ as $|\xi| \rightarrow +\infty$, because these terms $\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} \chi(\theta) a(x, \theta) d\theta$ and $\int_{\Gamma_x} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta$ are smooth in terms of x (for the first term, it is because the actual integral domain is compact, i.e. is contained in $\text{supp } \theta$; for the second term, it is because the integrand decays at infinity order). Then we can compute I_3 as follows,

$$I_3(\xi) \simeq \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\varphi_\theta(x, \theta) \neq 0 \text{ and } \theta \notin \Gamma_x} \left(\frac{\varphi_\theta(x, \theta) \cdot \nabla_\theta}{|\varphi_\theta(x, \theta)|^2} \right)^N e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx$$

$$\begin{aligned}
& + \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\varphi_\theta(x, \theta) = 0 \text{ and } \theta \notin \Gamma_x} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx \\
& =: I_4(\xi) + I_5(\xi).
\end{aligned}$$

Now here comes the key point: to obtain $\langle \xi \rangle^{-N}$, for I_4 we differentiate $e^{-ix \cdot \xi}$ w.r.t. x , and for I_5 we differentiate $e^{i(\varphi - x \cdot \xi)}$ w.r.t. x .

We can estimate I_4 by using the computation as in (8.39),

$$\begin{aligned}
I_4(\xi) & \simeq \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\varphi_\theta(x, \theta) \neq 0 \text{ and } \theta \notin \Gamma_x} \left(\frac{\varphi_\theta(x, \theta) \cdot \nabla_\theta}{|\varphi_\theta(x, \theta)|^2} \right)^N e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx \\
& \sim \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\varphi_\theta(x, \theta) \neq 0 \text{ and } \theta \notin \Gamma_x} e^{i\varphi(x, \theta)} |\varphi_\theta(x, \theta)|^{-N} \partial_\theta^N ((1 - \chi)a) d\theta \right) dx,
\end{aligned}$$

where the integer N can be arbitrary. And hence we have $I_4(\xi) = \mathcal{O}(|\xi|^{-\infty})$ for the same reason as I_1 .

It is the I_5 which finally decides $\text{WF}(A)$. For $\xi \neq \varphi_x(x, \theta)$, we can have I_3 as follows,

$$I_5(\xi) = \int_{\substack{\varphi_\theta(x, \theta) = 0, \\ (x, \theta) \notin \text{Smo}(a)}} e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \quad (8.41)$$

$$\begin{aligned}
& = \int_{\substack{\varphi_\theta(x, \theta) = 0, \\ (x, \theta) \notin \text{Smo}(a)}} \left[\left(\frac{(\varphi_x(x, \theta) - \xi) \cdot \nabla_x}{i|\varphi_x(x, \theta) - \xi|^2} \right)^{N_1 + N_2} e^{i(\varphi(x, \theta) - x \cdot \xi)} \right] \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \\
& \quad (8.42)
\end{aligned}$$

$$\begin{aligned}
& \lesssim \int_{\substack{\varphi_\theta(x, \theta) = 0, \\ (x, \theta) \notin \text{Smo}(a)}} \langle \varphi_x(x, \theta) - \xi \rangle^{-N_1 - N_2} \phi(x) |a(x, \theta)| dx d\theta \\
& \lesssim \int_{\substack{\varphi_\theta(x, \theta) = 0, \\ (x, \theta) \notin \text{Smo}(a)}} \langle \xi \rangle^{-N_1} \langle \varphi_x(x, \theta) \rangle^{-N_2} \phi(x) \langle \theta \rangle^m dx d\theta \\
& \simeq \langle \xi \rangle^{-N_1} \int_{\substack{\varphi_\theta(x, \theta) = 0, \\ (x, \theta) \notin \text{Smo}(a)}} \phi(x) \langle \theta \rangle^{m - N_2} dx d\theta \simeq \langle \xi \rangle^{-N_1},
\end{aligned}$$

To guarantee the derivation from (8.41) to (8.42), we need $\xi \neq \varphi_x(x, \theta)$ for these (x, θ) which satisfy $\varphi_x(x, \theta) = 0$ and $(x, \theta) \notin \text{Smo}(a)$. We finished the proof. \square

FORMAL PROOF OF THEOREM 8.14. We do some preparation first. Define \mathcal{A} as the collection of subsets Ω in $\mathbb{R}_x^n \times (\mathbb{R}_\xi^n \setminus \{0\})$ where $(x, \xi) \in \Omega \Rightarrow (x, t\xi) \in \Omega$ for any $t > 0$, and \mathcal{B} as the collection of subsets in $\mathbb{R}_x^n \times \mathbb{S}_\xi^{n-1}$. Then there is a one-to-one correspondence between \mathcal{A} and \mathcal{B} , and we denote the one-to-one mapping as S ,

$$S: \Omega \in \mathcal{A} \mapsto S\Omega = \{(x, \eta); \exists \xi \in \mathbb{R}^n \text{ s.t. } \eta = \xi/|\xi| \text{ and } (x, \xi) \in \Omega\} \in \mathcal{B}.$$

Let $T: (x, \theta) \mapsto (x, \varphi_x(x, \theta))$. Note that $ST = TS$. For any positive integer k , denote

$$V_k := \{(x, \theta) \in \mathbb{R}_x^n \times \mathbb{R}_\xi^n; |\varphi_\theta(x, \theta)| \leq 1/k\}. \quad (8.43)$$

It can be checked that

- $\{V_k\}_k$ and $\{TV_k\}_k$ are decreasing in terms of k ,
- V_k and TV_k are closed in $\mathbb{R}_x^n \times (\mathbb{R}_\xi^n \setminus \{0\})$,
- $V_k, TV_k \in \mathcal{A}$,
- SV_k and $STV_k (= TSV_k)$ are also closed in $\mathbb{R}_x^n \times \mathbb{S}_\xi^{n-1}$.

Now let's assume

$$(x_0, \xi_0) \notin TV_k, \quad (8.44)$$

then $(x_0, \xi_0/|\xi_0|)$ is not in STV_k , which is a closed set. Therefore, there exists $\epsilon_1 > 0$ such that

$$\{(x, \xi/|\xi|); |x - x_0| \leq \epsilon_1, |\xi/|\xi| - \xi_0/|\xi_0| \leq \epsilon_1\} \cap STV_k = \emptyset. \quad (8.45)$$

Because $\nabla_{(x,\theta)}\varphi(x, \theta)$ is always assumed to be nonzero, the number

$$\inf_{|x-x_0| \leq \epsilon_1, \theta \in \mathbb{S}^{n-1}} (|\varphi_x(x, \theta)| + |\varphi_\theta(x, \theta)|)$$

exists and is positive and we denote it as ϵ_2 ,

$$\epsilon_2 := \inf_{|x-x_0| \leq \epsilon_1, \theta \in \mathbb{S}^{n-1}} (|\varphi_x(x, \theta)| + |\varphi_\theta(x, \theta)|) > 0. \quad (8.46)$$

Let k_0 be any positive integer such that

$$k_0 > 2[1/\epsilon_2]. \quad (8.47)$$

Now, for any $(x, \theta) \in SV_{k_0}$, we know $(x, \theta) \in V_{k_0}$, so (8.43) gives $|\varphi_\theta(x, \theta)| \leq 1/k_0 < \epsilon_2/2$, so from (8.46) we can conclude that

$$|\varphi_x(x, \theta)| > \epsilon_2/2 > 0 \quad \text{in } W := \{(x, \theta) \in SV_{k_0}; |x - x_0| \leq \epsilon_1\}. \quad (8.48)$$

Fix some $\phi \in C_c^\infty(B(x_0, \epsilon_1))$. And $\chi \in C_c^\infty(\mathbb{R}^n)$ is a cut-off function with support containing the origin. Now we estimate $\widehat{\phi A}(\xi)$. We have

$$\begin{aligned} \widehat{\phi A}(\xi) &= A(\phi e^{-ix \cdot \xi}) = (2\pi)^{-n/2} \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) a(x, \theta) dx d\theta \\ &\simeq \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) \chi(\theta) a(x, \theta) dx d\theta + \int e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \\ &= \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} \chi(\theta) a(x, \theta) d\theta \right) dx \\ &\quad + \int_{(x, \theta) \in \text{Smo}'(a)} e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) d\theta dx \\ &\quad + \int_{(x, \theta) \notin \text{Smo}'(a)} e^{i(\varphi(x, \theta) - x \cdot \xi)} \phi(x) (1 - \chi(\theta)) a(x, \theta) dx d\theta \\ &=: I_1(\xi) + I_2(\xi) + I_3(\xi), \end{aligned} \quad (8.49)$$

where $\text{Smo}'(a)$ is an arbitrary subset of $\text{Smo}(a)$ such that for every fixed x , the projection of the intersect $S((x, \mathbb{R}_\xi^n) \cap \text{Smo}(a))$ is a compact subset of the sphere \mathbb{S}^{n-1} . The (x, \mathbb{R}_ξ^n) means $\{(x, \xi) \in \mathbb{R}_x^n \times \mathbb{R}_\xi^n; \xi \in \mathbb{R}^n\}$. The term I_1 is easy to estimate. The $\int_{\mathbb{R}_\theta^n} e^{i\varphi(x, \theta)} \chi(\theta) a(x, \theta) d\theta$ in I_1 is C^∞ -smooth in terms of x , so by using integration by parts we can have

$$|I_1(\xi)| = \left| \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \mathcal{C}(x) dx \right| \leq C_\alpha \xi^{-\alpha}, \quad \forall \xi, \forall \text{ multi-index } \alpha. \quad (8.50)$$

where \mathcal{C} is some function in $C_c^\infty(\mathbb{R}^n)$. The estimation (8.50) gives

$$|I_1(\xi)| \leq C_N \langle \xi \rangle^{-N}, \quad \forall \xi, \forall N \in \mathbb{N}. \quad (8.51)$$

And I_2 can be estimated as follows,

$$\begin{aligned} |I_2(\xi)| &= \left| \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \phi(x) \left(\int_{\{\theta; (x, \theta) \in \text{Smo}'(a)\}} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx \right| \\ &= \left| \int_{\mathbb{R}_x^n} [(i^{|\alpha|} \xi^{-\alpha} \partial_x^\alpha) e^{-ix \cdot \xi}] \phi(x) \left(\int_{\{\theta; (x, \theta) \in \text{Smo}'(a)\}} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) dx \right| \\ &= \xi^{-\alpha} \left| \int_{\mathbb{R}_x^n} e^{-ix \cdot \xi} \partial_x^\alpha \left[\phi(x) \left(\int_{\{\theta; (x, \theta) \in \text{Smo}'(a)\}} e^{i\varphi(x, \theta)} (1 - \chi(\theta)) a(x, \theta) d\theta \right) \right] dx \right|. \end{aligned}$$

It is easy to check that the term in $[\dots]$ are C^∞ -smooth and compactly supported w.r.t. x , thus it is integrable. Therefore,

$$|I_2(\xi)| \leq C_N \langle \xi \rangle^{-N}, \quad \forall \xi, \forall N \in \mathbb{N}. \quad (8.52)$$

Then we move on to I_3 ,

$$\begin{aligned} I_3(\xi) &= \int_{(\text{Smo}'(a))^c} e^{i(\varphi(x,\theta)-x\cdot\xi)} \phi(x)(1-\chi(\theta))a(x,\theta) dx d\theta \\ &= \int_{(\text{Smo}'(a))^c \cap (V_{k_0})^c} e^{i(\varphi(x,\theta)-x\cdot\xi)} \phi(x)(1-\chi(\theta))a(x,\theta) dx d\theta \\ &\quad + \int_{(\text{Smo}'(a))^c \cap V_{k_0}} e^{i(\varphi(x,\theta)-x\cdot\xi)} \phi(x)(1-\chi(\theta))a(x,\theta) dx d\theta \\ &=: I_4(\xi) + I_5(\xi), \end{aligned} \quad (8.53)$$

where $(\text{Smo}'(a))^c$ signifies the complementary set of $\text{Smo}'(a)$. Note that in $(V_{k_0})^c$, the $|\varphi_\theta(x,\theta)|$ is no less than $1/k_0$ (c.f. (8.43)), thus no singularity will accrue when $|\varphi_\theta(x,\theta)|$ appears in the denominator. Hence, for I_4 we have

$$\begin{aligned} I_4(\xi) &= \int_{(\text{Smo}'(a))^c \cap (V_{k_0})^c} e^{i(\varphi(x,\theta)-x\cdot\xi)} \phi(x)(1-\chi(\theta))a(x,\theta) dx d\theta \\ &= \int e^{-ix\cdot\xi} \phi(x) \left(\int e^{i\varphi(x,\theta)} (1-\chi(\theta))a(x,\theta) d\theta \right) dx \\ &= \int [(i^{|\alpha|} \xi^{-\alpha} \partial_x^\alpha) e^{-ix\cdot\xi}] \phi(x) \left(\int e^{i\varphi(x,\theta)} (1-\chi(\theta))a(x,\theta) d\theta \right) dx \\ &\simeq \xi^{-\alpha} \int e^{-ix\cdot\xi} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \partial_x^{\alpha-\beta} \phi(x) \cdot \partial_x^\beta \left(\int e^{i\varphi(x,\theta)} (1-\chi(\theta))a(x,\theta) d\theta \right) dx \\ &\simeq \xi^{-\alpha} \int e^{-ix\cdot\xi} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \partial_x^{\alpha-\beta} \phi(x) \\ &\quad \cdot \partial_x^\beta \left(\int \frac{-i\varphi_\theta(x,\theta) \cdot \nabla_\theta}{|\varphi_\theta(x,\theta)|^2} (e^{i\varphi(x,\theta)}) (1-\chi(\theta))a(x,\theta) d\theta \right) dx \\ &\lesssim \xi^{-\alpha} \sum_{\beta \leq \alpha} \int e^{-ix\cdot\xi} C_{\alpha,\beta} \partial_x^{\alpha-\beta} \phi(x) \left(\int \langle \theta \rangle^{m+|\beta|-N} d\theta \right) dx \\ &\leq C_\alpha \xi^{-\alpha}, \quad \forall \xi, \forall \text{ multi-index } \alpha. \end{aligned} \quad (8.54)$$

Now for the estimation of I_5 , we need some constraints on the direction of ξ . **It is this term that determines** $\text{WF}(A)$. Because $(x_0, \xi_0) \notin TV_{k_0}$ (see (8.44)), according to (8.45), there is a cone $W \subset \mathbb{R}_\xi^n \setminus \{0\}$ such that

$$\xi_0 \in W \quad \text{and} \quad |\nabla_x \varphi(x,\theta) - \xi| \geq C(|\nabla_x \varphi(x,\theta)| + |\xi|), \quad \forall \xi \in W. \quad (8.55)$$

Define $L := \frac{-i(\partial_x \varphi(x,\theta) - \xi) \cdot \nabla_x}{|\nabla_x \varphi(x,\theta) - \xi|^2}$. By using the fact that

$$|\partial^\alpha L f(x)| \lesssim |\nabla_x \varphi(x,\theta) - \xi|^{-1} \sum_\beta |\partial^\beta f(x)|,$$

we can have, for all $\xi \in W$,

$$I_5(\xi) = \int_{(\text{Smo}'(a))^c \cap V_{k_0}} e^{i(\varphi(x,\theta)-x\cdot\xi)} \phi(x)(1-\chi(\theta))a(x,\theta) dx d\theta$$

$$\begin{aligned}
&= \int_{(\text{Smo}'(a))^c \cap V_{k_0}} L^N (e^{i(\varphi(x,\theta)-x \cdot \xi)}) \phi(x) (1 - \chi(\theta)) a(x, \theta) \, dx \, d\theta \\
&= \int_{(\text{Smo}'(a))^c \cap V_{k_0}} e^{i(\varphi(x,\theta)-x \cdot \xi)} {}^t L^N (\phi(x) (1 - \chi(\theta)) a(x, \theta)) \, dx \, d\theta \\
&\lesssim \int_{\text{supp } \phi \times \mathbb{R}_\theta^n} (1 + |\varphi_x(x, \theta) - \xi|)^{-N} \langle \theta \rangle^m \, dx \, d\theta \\
(N_1 + N_2 = N) \quad &\lesssim \langle \xi \rangle^{-N_1} \int_{\text{supp } \phi \times \mathbb{R}_\theta^n} \langle |\theta| \cdot \varphi_x(x, \theta/|\theta|) \rangle^{-N_2} \langle \theta \rangle^m \, dx \, d\theta \\
(\text{by (8.48)}) \quad &\lesssim \langle \xi \rangle^{-N_1} \int_{\text{supp } \phi \times \mathbb{R}_\theta^n} \langle \theta \rangle^{-N_2} \langle \theta \rangle^m \, dx \, d\theta \\
&\lesssim \langle \xi \rangle^{-N_1} \quad \forall N_1 \in \mathbb{N}. \tag{8.56}
\end{aligned}$$

Combining (8.44), (8.47), (8.49), (8.51), (8.52), (8.53), (8.54) and (8.56), we arrive at

$$|\widehat{\phi A}(\xi)| \lesssim \langle \xi \rangle^{-N}, \quad \xi \in W, \quad \forall N \in \mathbb{N}, \tag{8.57}$$

where the W is a cone containing ξ_0 , and $(x_0, \xi_0) \neq (x, \varphi_x(x, \theta))$ for these (x, θ) satisfying

$$(x, \theta) \in (\text{Smo}'(a))^c \cap V_{k_0}.$$

Therefore, for any $k_0 > 2[1/\epsilon_2]$, there holds

$$\left(\{(x, \varphi_x(x, \theta)); |\varphi_\theta(x, \theta)| \leq 1/k_0, (x, \theta) \in (\text{Smo}'(a))^c\} \right)^c \subset \left(\text{WF}(A) \right)^c,$$

thus

$$\text{WF}(A) \subset \{(x, \varphi_x(x, \theta)); |\varphi_\theta(x, \theta)| \leq 1/k_0, (x, \theta) \notin \text{Smo}'(a)\}.$$

Finally, let k_0 goes to zero and choose $\text{Smo}'(a)$ to be arbitrarily close to $\text{Smo}(a)$, we arrive at the conclusion (8.38). \square

8.4. Applications

Now we are ready to apply those results.

8.4.1. Microlocality of Ψ DOs.

PROPOSITION 8.16. *Assume $a \in S^{+\infty}(\mathbb{R}_x^n \times \mathbb{R}_y^n \times \mathbb{R}_\xi^n)$ is symbol and K is the kernel of the corresponding Ψ DO of a , then*

$$\boxed{\text{WF}(K) \subset \{(x, x; \xi, -\xi); (x, x, \xi) \notin \text{Smo}(a)\}.} \tag{8.58}$$

PROOF. Denote the corresponding Ψ DO as A , then

$$\begin{aligned}
\langle K(x, y), u \otimes v(x, y) \rangle &= \langle K(x, y), u(x) \otimes v(y) \rangle = \langle \langle K(x, y), u(x) \rangle_x, v(y) \rangle_y \\
&= \langle Au(y), v(y) \rangle = (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} a(x, y, \xi) u(x) v(y) \, dx \, dy \, d\xi \\
&= (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} a(x, y, \xi) u \otimes v(x, y) \, dx \, dy \, d\xi.
\end{aligned}$$

Therefore, in the oscillatory integral sense,

$$K(x, y) = (2\pi)^{-n} \int e^{i(x-y) \cdot \xi} a(x, y, \xi) \, d\xi.$$

According to Theorem 8.14, we have

$$\text{WF}(K) \subset \{(x, y; \varphi_{x,y}(x, y, \xi)); \varphi_\xi(x, y, \xi) = 0, (x, y, \xi) \notin \text{Smo}(a)\},$$

where $\varphi(x, y, \xi) = (x - y) \cdot \xi$. Hence,

$$\begin{aligned} \text{WF}(K) &\subset \{(x, y; \xi, -\xi); x - y = 0, (x, y, \xi) \notin \text{Smo}(a)\} \\ &= \{(x, x; \xi, -\xi); (x, x, \xi) \notin \text{Smo}(a)\}. \end{aligned}$$

The proof is complete. \square

THEOREM 8.17. *Assume that A is a Ψ DO, then for $u \in \mathcal{E}'$ we have*

$$\text{microlocality: } \boxed{\text{WF}(Au) \subseteq \text{WF}(u) \setminus \text{Smo}(A)}. \quad (8.59)$$

Moreover, if A is elliptic in the sense of Definition 6.4, then $\text{Smo}(A) = \emptyset$ and

$$\boxed{\text{WF}(Au) = \text{WF}(u)}. \quad (8.60)$$

PROOF. We have $Au(x) = \langle K(x, y), u(y) \rangle$ where K is its kernel, so according to Theorem 8.11 and Proposition 8.16, we can conclude

$$\begin{aligned} \text{WF}(Au) &\subseteq (\text{WF}'(K) \circ \text{WF}(u)) \cup \text{WF}_x(K) \\ &= (\{(x, x; \xi, \xi); (x, \xi) \notin \text{Smo}(a)\} \circ \text{WF}(u)) \cup \emptyset \\ &= \text{WF}(u) \setminus \text{Smo}(a) = \text{WF}(u) \setminus \text{Smo}(A). \end{aligned}$$

If A is elliptic, then according to the definition, we have

$$|a(x, \xi)| \geq C \langle \xi \rangle^m, \quad \text{when } x \in \mathbb{R}^n, |\xi| \geq R,$$

for some constants $m \in \mathbb{R}$, $C > 0$ and $R > 0$, so it is obvious that $\text{Smo}(A) = \emptyset$, thus by (8.59),

$$\text{WF}(Au) \subseteq \text{WF } u.$$

Also, because A is elliptic, then by Theorem 6.6 we know A has a parametrix B such that $R := BA - I \in \Psi^{-\infty}$, so

$$\text{WF}(u) = \text{WF}(BAu - Ru) \subset \text{WF}(BAu) \cup \text{WF}(Ru) \subset \text{WF}(Au).$$

In total, $\text{WF}(Au) = \text{WF } u$. The proof is complete. \square

LEMMA 8.18. *Assume a is a symbol and $u \in \mathcal{E}'(\Omega)$. Denote the corresponding Ψ DO of a as A , then*

$$\boxed{\text{WF}(Au) \cap \text{Smo}(a) = \emptyset}. \quad (8.61)$$

PROOF. This is a straight forward outcome of (8.59). \square

8.4.2. Pull-back of distributions.

THEOREM 8.19. *Let Ω_1 and Ω_2 be two domain in \mathbb{R}^n , and $\psi: \Omega_1 \rightarrow \Omega_2$ is an diffeomorphism. Then for any $u \in \mathcal{D}'(\Omega_2)$, we have $\psi^*u \in \mathcal{D}'(\Omega_1)$, and*

$$\boxed{\text{WF}(\psi^*u) = \{(x, {}^t\psi'|_x\eta); (\psi(x), \eta) \in \text{WF}(u)\}}, \quad (8.62)$$

where ψ' signifies the matrix whose $(i\text{-row}, j\text{-column})$ element is $\partial_{x_i}\psi^j$, and $({}^t\psi')^{-1}|_x$ is the inverse of transpose of the matrix ψ' evaluated at x , and $({}^t\psi')^{-1}|_x\eta$ stands for the matrix multiplication of the matrix $({}^t\psi')^{-1}|_x$ and the vertical vector η .

The mapping in (8.62),

$$(x, {}^t\psi'|_x\eta) \mapsto (\psi(x), \eta)$$

is invariant on the cotangent bundle (see [19, for Theorem 18.1.17]).

FIRST PROOF OF THEOREM 8.19. It is obvious that $\psi^*u \in \mathcal{D}'(\Omega_1)$.

For (y_0, ξ_0) , because ψ is a diffeomorphism, we can find (x_0, η_0) such that $x_0 = \psi^{-1}(y_0)$ and $\xi_0 = (\psi^{-1})'|_{y_0} \cdot \eta_0$. Assume that $(y_0, \xi_0) \notin \text{WF}(u)$. For a smooth cutoff function φ_{x_0} satisfying $\varphi_{x_0}(x_0) \neq 1$, we have

$$\begin{aligned}
(\varphi_{x_0}\psi^*u)^\wedge(\eta) &\simeq \int e^{-ix \cdot \eta} \varphi_{x_0}(x) \psi^*u(x) dx = \int e^{-ix \cdot \eta_0} \varphi_{x_0}(x) u(\psi(x)) dx \\
&= \int e^{-i\psi^{-1}(y) \cdot \eta} \varphi_{x_0}(\psi^{-1}(y)) u(y) d\psi^{-1}(y) \quad (y = \psi(x)) \\
&= \int e^{-i{}^t\psi^{-1}(y)\eta} \cdot \varphi_{x_0}(\psi^{-1}(y)) \left| \frac{\partial \psi^{-1}}{\partial y}(y) \right| \phi^{-1}(y) \cdot (\phi u)(y) dy \\
&= \int e^{-i{}^t\psi^{-1}(y)\eta} \cdot \tilde{\varphi}_{y_0}(y) \cdot (\phi u)(y) dy = \int e^{-i({}^t\psi^{-1}(y)\eta - y \cdot \xi)} \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&= \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} e^{-i({}^t\psi^{-1}(y)\eta - y \cdot \xi)} \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&\quad + \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} e^{-i{}^t\psi^{-1}(y)\eta} e^{iy \cdot \xi} \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&=: I_1 + I_2,
\end{aligned}$$

where $\hat{\xi} := \xi/|\xi|$ and the same for $\hat{\xi}_0$. For convenience we have written $\psi^{-1}(y) \cdot \eta$ as ${}^t\psi^{-1}(y)\eta$, where tM signifies the transpose operation for any matrix M . By doing so it will be more straightforward when we make derivatives. Here $\tilde{\varphi}_{y_0}(y)$ is a generic function which is C^∞ -smooth and is compactly supported w.r.t. y and whose precise definition may varies from line to line.

For I_1 , because $|\hat{\xi}_0 - \hat{\xi}| > 1$ and $(y_0, \xi_0) \notin \text{WF}(u)$, we have $|({}^t\psi^{-1})'|_y \eta - \xi| \geq \frac{1}{2}$ and $|\widehat{\phi u}(\xi)| \lesssim \langle \xi \rangle^{-N_0}$ for some integer N_0 . The number N_0 comes from the fact that ϕu is a compactly supported distribution so its Fourier transform has (at most) polynomial growth. Hence,

$$\begin{aligned}
I_1 &= \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} \left(\frac{1 + \langle ({}^t\psi^{-1})'|_y \eta - \xi \rangle \cdot \nabla_y}{\langle ({}^t\psi^{-1})'|_y \eta - \xi \rangle^2} \right)^N (e^{-i({}^t\psi^{-1}(y)\eta - y \cdot \xi)}) \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&\simeq \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} \langle ({}^t\psi^{-1})'|_y \eta - \xi \rangle^{-N} e^{-i({}^t\psi^{-1}(y)\eta - y \cdot \xi)} \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&\lesssim \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} \langle ({}^t\psi^{-1})'|_y \eta \rangle^{-N_1} \langle \xi \rangle^{-N_2} \cdot |\tilde{\varphi}_{y_0}(y)| \cdot |\widehat{\phi u}(\xi)| d\xi dy \quad (N = N_1 + N_2) \\
&\lesssim \langle \eta \rangle^{-N_1} \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} \langle \xi \rangle^{-N_2} \cdot \left(\int |\tilde{\varphi}_{y_0}(y)| dy \right) \cdot |\widehat{\phi u}(\xi)| d\xi \\
&\lesssim \langle \eta \rangle^{-N_1} \int_{|\hat{\xi}_0 - \hat{\xi}| > 1} \langle \xi \rangle^{-N_2} \langle \xi \rangle^{N_0} d\xi \lesssim \langle \eta \rangle^{-N_1},
\end{aligned}$$

provided that $N_2 - N_0 > \text{the dimension of } \xi$.

For I_2 , we have

$$\begin{aligned}
I_2 &= \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} \left(\frac{1 + \langle ({}^t\psi^{-1})'|_y \eta \rangle \cdot \nabla_y}{\langle ({}^t\psi^{-1})'|_y \eta \rangle^2} \right)^N (e^{-i\psi^{-1}(y) \cdot \eta}) e^{iy \cdot \xi} \cdot \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) d\xi dy \\
&\simeq \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} \langle ({}^t\psi^{-1})'|_y \eta \rangle^{-N} e^{-i\psi^{-1}(y) \cdot \eta} \cdot (\nabla_y)^N (e^{iy \cdot \xi} \cdot \tilde{\varphi}_{y_0}(y)) \cdot \widehat{\phi u}(\xi) d\xi dy
\end{aligned}$$

$$\begin{aligned}
&\simeq \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} \langle \eta \rangle^{-N} e^{-i\psi^{-1}(y) \cdot \eta} \cdot \langle \xi \rangle^N \tilde{\varphi}_{y_0}(y) \cdot \widehat{\phi u}(\xi) \, d\xi \, dy \\
&\lesssim \langle \eta \rangle^{-N} \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} \langle \xi \rangle^N \cdot \left(\int |\tilde{\varphi}_{y_0}(y)| \, dy \right) \cdot |\widehat{\phi u}(\xi)| \, d\xi \\
&\lesssim \langle \eta \rangle^{-N} \int_{|\hat{\xi}_0 - \hat{\xi}| \leq 1} \langle \xi \rangle^N \langle \xi \rangle^{-N-n-1} \, d\xi \lesssim \langle \eta \rangle^{-N}.
\end{aligned}$$

In total, we have $|(\varphi_{x_0} \psi^* u)^\wedge(\eta)| \lesssim \langle \eta \rangle^{-N}$ for any integer N if $\hat{\eta}$ is in a small neighborhood of $\hat{\eta}_0$ where $(\psi(x_0), ({}^t\psi^{-1})'|_{y_0} \eta_0) \notin \text{WF}(u)$, namely,

$$(\psi(x_0), ({}^t\psi^{-1})'|_{\psi(x_0)} \eta_0) \notin \text{WF}(u) \Rightarrow (x_0, \eta_0) \notin \text{WF}(\psi^* u).$$

Therefore,

$$(x_0, \eta_0) \in \text{WF}(\psi^* u) \Rightarrow (\psi(x_0), ({}^t\psi^{-1})'|_{\psi(x_0)} \eta_0) \in \text{WF}(u),$$

so

$$\text{WF}(\psi^* u) \subset \{(x, \eta); (\psi(x), ({}^t\psi^{-1})'|_{\psi(x)} \eta) \in \text{WF}(u)\}.$$

Because ψ is invertible, we can obtain the opposite inclusion by looking at $\psi^{-1*}(\psi^* u)$.

It can be shown that $(\psi^{-1})'|_{\psi(x)} = (\psi'(x))^{-1}$. Indeed, by differentiating $x = \psi^{-1}(\psi(x))$ w.r.t. x we obtain $I = (\psi^{-1})'|_{\psi(x)} \cdot \psi'(x)$, so $(\psi^{-1})'|_{\psi(x)} = (\psi'(x))^{-1}$, and by taking transpose we obtain

$$({}^t\psi^{-1})'|_{\psi(x)} = ({}^t\psi'(x))^{-1},$$

so

$$\begin{aligned}
\text{WF}(\psi^* u) &= \{(x, \eta); (\psi(x), ({}^t\psi')^{-1}|_x \eta) \in \text{WF}(u)\} \\
&= \{(x, {}^t\psi'|_x \xi); (\psi(x), \xi) \in \text{WF}(u)\}.
\end{aligned}$$

The proof is complete. \square

There is also another proof for Theorem 8.19. As in Remark 8.15, we can use Theorems 8.14 and 8.11 to obtain $\text{WF}(\psi^* u)$.

SECOND PROOF OF THEOREM 8.19. The pull-back ψ^* has a kernel: for $f \in C^\infty(\Omega_2)$ and $g \in C^\infty(\Omega_1)$, we have

$$\begin{aligned}
\langle \psi^* f, g \rangle &= \int \psi^* f(x) g(x) \, dx = (2\pi)^{-n} \int e^{i(\psi(x)-y) \cdot \eta} f(y) g(x) \, dx \, dy \, d\eta \\
&\simeq \left\langle \int e^{i(\psi(x)-y) \cdot \eta} \, d\eta, (g \otimes f)(x, y) \right\rangle \\
&= \langle K, g \otimes f \rangle, \quad \text{where } K(x, y) = \int e^{i({}^t\psi(x)-{}^t y) \cdot \eta} \, d\eta.
\end{aligned}$$

and $\psi^* u(x) = \langle K(x, y), u(y) \rangle$. By Theorem 8.14 we have

$$\text{WF}(K) \subset \{(x, \psi(x); {}^t\psi'|_x \eta, -\eta)\},$$

and then by Theorem 8.11 we have

$$\text{WF}(\psi^* u) \subseteq (\text{WF}'(K) \circ \text{WF}(u)) \cup \text{WF}_x(K) = \{(x, {}^t\psi'|_x \eta); (\psi(x), \eta) \in \text{WF}(u)\}.$$

The opposite inclusion can be obtained by looking at $\psi^{-1*}(\psi^* u)$. We obtain (8.62). \square

Exercise

EXERCISE 8.1. Prove Lemme 8.3.

EXERCISE 8.2. Prove that the condition $(\text{WF}'(K) \circ \text{WF}(u)) \cap O_x = \emptyset$ guarantees (8.31).

EXERCISE 8.3. Show the details in the computations (8.34) and (8.35).

Propagation of the singularities

9.1. Microlocal parametrix

To study the microlocal parametrix, we recall notion of conic sets and the smooth direction “Smo” given in Definitions 8.1 & 8.2, and $T^*\mathbb{R}^n \setminus 0$ stands for the cotangent bundle with the zero section excluded. Now we generalize Definitions 6.1 & 6.4, microlocally, as follows.

DEFINITION 9.1 (Microlocal parametrix). Assume $m \in \mathbb{R}$ and $T \in \Psi^m$. We call a Ψ DO S a *left (resp. right) microlocal parametrix* of T if there exists a nonempty open conic set $\Gamma \subset T^*\mathbb{R}^n \setminus 0$ such that

$$\text{Smo}(ST - I) = \Gamma \quad (\text{resp. } \text{Smo}(TS - I) = \Gamma).$$

We call S a *microlocal parametrix* of T if it is both a left and a right microlocal parametrix under the same set Γ .

DEFINITION 9.2 (Microlocal ellipticity). Assume $m \in \mathbb{R}$ and $a \in S^m$, and A is the Ψ DO of a . Let $\Gamma \subset T^*\mathbb{R}^n \setminus 0$ be a open conic set. We say a (and A) is *microlocally elliptic in Γ* if for some constants $C_\Gamma > 0$, $R > 0$,

$$|a(x, \xi)| \geq C_\Gamma \langle \xi \rangle^m, \quad \forall (x, \xi) \in \Gamma, \quad |\xi| \geq R.$$

We write $\text{Char } A := (\bigcup \mathcal{F})^c$, where $\mathcal{F} = \{\Gamma; A \text{ is microlocally elliptic in } \Gamma\}$, and the notation Ω^c stands for the complement of Ω in $T^*\mathbb{R}^n \setminus 0$.

From Definition 9.2, it is obvious that $\text{Char } A$ is always closed. The following claim is trivial.

LEMMA 9.3. $\text{Char } A = \emptyset$ if and only if A is elliptic in the sense of Definition 6.4.

LEMMA 9.4. Assume P is the Ψ DO with principal symbol $p_m(x, \xi)$ homogeneous in ξ , then $\text{Char } P = p_m^{-1}(0)$, where $p_m^{-1}(0)$ signifies the set $\{(x, \xi) \in T^*\mathbb{R}^n \setminus 0; p_m(x, \xi) = 0\}$.

The proof is left as an exercise. The $\text{Char } A$ and $\text{Smo}(A)$ is closely related. Results in §6.1 can be modified to a microlocal version.

THEOREM 9.5 (Microlocal ellipticity \Leftrightarrow Microlocal parametrix). Let $m \in \mathbb{R}$ and $A \in \Psi^m$. Assume A is microlocally elliptic in Γ , where $\Gamma = (\text{Char } A)^c$ is non-empty. Then A has a microlocal parametrix B . Moreover, they satisfies

$$(\text{Char } A)^c \subset \text{Smo}(I - BA). \tag{9.1}$$

Conversely, if A has either a right or left microlocal parametrix, then A is microlocally elliptic.

PROOF. (\Rightarrow) Fix $(x_0, \xi_0) \in \Gamma$. In the proof of Theorem 6.6, we modify the function $\chi(\xi)$ to $\chi(x, \xi)$ which is given as $\chi(x, \xi) := \chi(x - x_0)\chi(\xi/|\xi| - \xi_0/|\xi_0|)$. Define $b_j(x, \xi) :=$

$(1 - \chi(x, \xi))/a(x, \xi) \cdot r_j(x, \xi)$ ($j \geq 0$) the same way, with $r_0 \equiv 1$, and follow the same steps as in the proof of Theorem 6.6 we can obtain $\forall N \in \mathbb{N}$,

$$\sigma(AB) = 1 - (1 + r_1 + \cdots + r_N)\chi - r_{N+1} + S^{-N-1} = 1 + S^{-N-1} \text{ in } \subset \Gamma,$$

as in (6.3), so

$$\sigma(I - AB) \in S^{-\infty} \text{ in } \subset \Gamma.$$

Due to the arbitrary of χ , we conclude

$$\sigma(I - AB) \in S^{-\infty} \text{ in } \Gamma.$$

Hence, by Definition 8.2 we obtain $\Gamma \subset \text{Smo}(I - AB)$.

(\Leftarrow) Assume B is the right parametrix of A , then there exists a nonempty conic open set $\Gamma \subset T^*\mathbb{R}^n \setminus 0$ such that $\text{Smo}(AB - I) \subset \Gamma$, which means $a\#b = 1 + S^{-\infty}$ in Γ . where a and b are symbols of A and B , respectively. Similar to the proof of Theorem 6.7, we can prove that

$$|a(x, \xi)| \geq \langle \xi \rangle^m / 2, \quad \text{when } (x, \xi) \in \Gamma, \langle \xi \rangle \geq C/2.$$

Therefore, a is microlocally elliptic in Γ . The proof for the left-case is similar.

The proof is complete. \square

COROLLARY 9.6. *For any $A \in \Psi^{+\infty}$ and any $u \in \mathcal{S}'$, there holds*

$$\boxed{\text{WF}(Au) \subset \text{WF } u \subset \text{WF}(Au) \cup \text{Char } A.}$$

PROOF. The “ $\text{WF}(Au) \subset \text{WF } u$ ” is from Theorem Theorem 8.17.

When $\text{Char } A = T^*\mathbb{R}^n \setminus 0$, the claim is trivial. When $\text{Char } A \subsetneq T^*\mathbb{R}^n \setminus 0$, $(\text{Char } A)^c$ is non-empty, so by Theorem 9.5, there exists a microlocal parametrix B of A , so we can have

$$\begin{aligned} \text{WF } u &= \text{WF}(BAu + (I - BA)u) \\ &\subset \text{WF}(BAu) \cup \text{WF}((I - BA)u) \\ &\subset \text{WF}(BAu) \cup (\text{WF } u \setminus \text{Smo}(I - BA)) \quad (\text{by (8.59)}) \\ &= \text{WF}(BAu) \cup (\text{WF } u \cap (\text{Smo}(I - BA))^c) \\ &\subset \text{WF}(BAu) \cup (\text{WF } u \cap \text{Char } A) \quad (\text{by Theorem 9.5}) \\ &= (\text{WF}(BAu) \cup \text{WF } u) \cap (\text{WF}(BAu) \cup \text{Char } A) \\ &\subset \text{WF } u \cap (\text{WF}(BAu) \cup \text{Char } A), \quad (\text{by (8.59)}) \end{aligned}$$

which gives

$$\text{WF } u \subset \text{WF}(BAu) \cup \text{Char } A \subset \text{WF}(Au) \cup \text{Char } A.$$

The proof is complete. \square

REMARK 9.7. Combining Corollary 9.6 and Lemma 9.3, we have

$$\begin{cases} A \text{ is microlocally elliptic: } \text{WF}(Au) \subset \text{WF } u \subset \text{WF}(Au) \cup \text{Char } A, \\ A \text{ is elliptic: } \text{WF}(Au) \subset \text{WF } u \subset \text{WF}(Au) \Leftrightarrow \text{WF}(Au) = \text{WF } u. \end{cases}$$

Hence, Corollary 9.6 can be viewed as a generalization of (8.60).

The following result is important.

THEOREM 9.8. *Assume $u \in \mathcal{S}'$, then*

$$\boxed{\text{WF}(u) = \bigcap_{A \in \Psi^{+\infty}, Au \in C^\infty} \text{Char } A.}$$

PROOF. By Corollary 9.6, we have $\text{WF } u \subset \text{WF}(Au) \cup \text{Char } A$, so

$$\text{WF}(u) \subset \bigcap_{Au \in C^\infty} \text{Char } A.$$

For the another direction, assume $(x_0, \xi_0) \notin \text{WF}(u)$, then we shall construct a suitable Ψ DO A such that

$$Au \in C^\infty, \quad \text{and} \quad (x_0, \xi_0) \notin \text{Char}(A), \quad (9.2)$$

which gives $(x_0, \xi_0) \notin \bigcap_{Au \in C^\infty} \text{Char } A$, and so

$$(\text{WF}(u))^c \subset \left(\bigcap_{Au \in C^\infty} \text{Char } A \right)^c \Rightarrow \bigcap_{Au \in C^\infty} \text{Char } A \subset \text{WF}(u),$$

and the proof will be finished.

It remains to construct such an operator A , and we present two ways to do it.

Method 1. Because $\text{WF}(u)$ is closed and $(x_0, \xi_0) \notin \text{WF}(u)$, so there exist bounded open neighborhoods ω, ω' of x_0 and conic open neighborhoods V, V' of ξ_0 such that

$$\begin{cases} \omega \subsetneq \omega', & V \subsetneq V', \\ \omega' \times V' \cap \text{WF}(u) = \emptyset. \end{cases}$$

We denote $\Gamma := \omega \times V$ and $\Gamma' := \omega' \times V'$, then $\Gamma \subset \Gamma'$ and $\Gamma' \cap \text{WF}(u) = \emptyset$, so

$$\text{WF}(u) \subset \Gamma^c. \quad (9.3)$$

Choose $a \in C^\infty(\mathbb{R}^{2n})$ such that

$$\begin{cases} \text{supp } a \subset \Gamma, & \text{and} & a(x_0, \xi_0) = 1, \\ a(x, \xi) = a(x, \xi/|\xi|) & \text{when } |\xi| \geq 1. \end{cases} \quad (9.4)$$

It can be shown that $a \in S^0$ (see Exercise 9.2) and $(x_0, \xi_0) \notin \text{Char}(T_a)$. Moreover, because $a \equiv 0$ in Γ^c and Γ^c is a conic set, we can conclude

$$\Gamma^c \subset \text{Smo}(T_a) \Rightarrow (\text{Smo}(T_a))^c \subset \Gamma. \quad (9.5)$$

Hence by Theorem 8.17 we have

$$\begin{aligned} \text{WF}(T_a u) &\subset \text{WF}(u) \setminus \text{Smo}(T_a) = \text{WF}(u) \cap (\text{Smo}(T_a))^c \subset \text{WF}(u) \cap \Gamma \quad (\text{by } (9.5)) \\ &= \text{WF}(u) \setminus \Gamma^c = \emptyset, \quad (\text{by } (9.3)) \end{aligned}$$

which implies $T_a u \in C^\infty$. Condition (9.2) is satisfied.

Method 2. Because $(x_0, \xi_0) \notin \text{WF}(u)$, there exists $\phi \in C_c^\infty(\mathbb{R}^n)$ such that $\widehat{\phi u}(\xi)$ is rapidly decaying when $\xi/|\xi|$ and $\xi_0/|\xi_0|$ are close enough, say, $|\xi/|\xi| - \xi_0/|\xi_0|| \leq \epsilon$ for certain $\epsilon > 0$. Hence we choose $\psi \in C^\infty(\mathbb{S}^{n-1})$ such that $\psi(\xi_0/|\xi_0|) = 1$ and $\psi(\eta) = 0$ when $|\eta - \xi_0/|\xi_0|| > \epsilon$ where $\eta \in \mathbb{S}^{n-1}$. Choose $\chi \in C_c^\infty(\mathbb{R}^n)$ such that $\chi(\xi) = 1$ when $|\xi| \geq 1/10$ and $\chi(\xi) = 0$ when $|\xi| \geq 1/5$. Now we define a operator A as follows

$$A\varphi(x) := (2\pi)^{-n} \int_{\mathbb{R}^{2n}} e^{i(x-y)\cdot\xi} \phi(y) (1 - \chi(\xi)) \psi(\xi/|\xi|) \varphi(y) dy d\xi, \quad \varphi \in \mathcal{S}(\mathbb{R}^n).$$

The purpose of the term “ $1 - \chi(\xi)$ ” is to cutoff the singularity near $\xi = 0$. By Theorem 5.6 we see A is a Ψ DO of order 0 with symbol

$$a(x, \xi) = \phi(x) (1 - \chi(\xi)) \psi(\xi/|\xi|) + S^{-1}, \quad (9.6)$$

By (9.6) we can show $(x_0, \xi_0) \notin \text{Char } A$, see Exercise 9.3. We can extend A from \mathcal{S} to \mathcal{S}' , and we have

$$\widehat{Au}(\xi) = (1 - \chi(\xi)) \psi(\xi/|\xi|) \widehat{\phi u}(\xi),$$

so \widehat{Au} is rapidly decaying, which means $Au \in C^\infty$. Condition (9.2) is satisfied.

The proof is complete. \square

9.2. Bicharacteristics

To prove a main result, we first introduce the notion of bicharacteristics. The Hamiltonian H_p of p is defined as:

$$H_p := \nabla_\xi p \cdot \nabla_x - \nabla_x p \cdot \nabla_\xi. \quad (9.7)$$

By Theorem 5.3 we can see

$$\begin{aligned} \sigma([P, Q]) &= \nabla_\xi p_m \cdot \nabla_x q_m - \nabla_x p_m \cdot \nabla_\xi q_m + S^{m_1+m_2-2} \\ &= H_{p_m} q_m + S^{m_1+m_2-2}, \end{aligned}$$

where m_1, m_2 are the order of P and Q , and p_m, q_m are principal symbols of P and Q , respectively.

In what follows we use the notation $T^*\mathbb{R}^n \setminus 0 := \{(x, \xi) \in T^*\mathbb{R}^n; \xi \neq 0\}$. We introduce the notion of bicharacteristic. For more details on the Hamiltonian flows, see [22, §2].

DEFINITION 9.9 (Null bicharacteristic). Assume $p \in C^1(T^*\mathbb{R}^n \setminus 0; \mathbb{R})$, and $I (\ni 0)$ is an open connected subset of \mathbb{R} . Let a curve $\gamma_{x_0, \xi_0}: s \in I \mapsto (x(s), \xi(s)) \in T^*\mathbb{R}^n \setminus 0$ satisfies

$$\begin{cases} \dot{x}(s) = \nabla_\xi p(x(s), \xi(s)), \quad \dot{\xi}(s) = -\nabla_x p(x(s), \xi(s)), \\ (x(0), \xi(0)) = (x_0, \xi_0) \in T^*\mathbb{R}^n \setminus 0. \end{cases} \quad (9.8)$$

We call $\boxed{\gamma_{x_0, \xi_0}(t)}$ a *bicharacteristic* of p . Furthermore, if $p(x_0, \xi_0) = 0$, then we shall have $p(x(s), \xi(s)) = 0$ for $\forall s$ and call $\gamma_{x_0, \xi_0}(t) := (x(t), \xi(t))$ a *null bicharacteristic* of p .

Note that in Definition 9.9, the function p is assumed to be real-valued. Without this assumption, we cannot guarantee $(x(s), \xi(s))$ are coordinates.

LEMMA 9.10. *For small enough $\epsilon > 0$, there exists a unique solution $\gamma: (-\epsilon, \epsilon) \mapsto (x(s), \xi(s)) \in T^*\mathbb{R}^n \setminus 0$ for the Hamiltonian equation (9.8). Moreover, assume either*

- (1) $\nabla_{(x, \xi)} p$ is uniformly bounded in $\{(x(s), \xi(s))/|\xi(s)|; s \in (-\epsilon, \epsilon)\}$, p is homogeneous of order 1;
- (2) or $\nabla_x p$ is uniformly bounded in $\{(x(s), \xi(s)); s \in (-\epsilon, \epsilon)\}$, p is homogeneous of order $\mu \geq 1$, and $\nabla_\xi p$ is bounded in uniformly bounded in $\{(x(s), \xi(s))/|\xi(s)|; s \in (-\epsilon, \epsilon)\}$;

then the domain of definition of γ can be extended from $(-\epsilon, \epsilon)$ to \mathbb{R} .

PROOF. Part 1: local solution. We use the Banach fixed-point theorem to show the existence of local solution. For simplicity denote $\eta_0 := (x_0, \xi_0)$ and $\eta(s) := (x(s), \xi(s))$ and $F(\eta(s)) := (\nabla_\xi p(\eta(s)), -\nabla_x p(\eta(s)))$, and we define a mapping \mathcal{F} :

$$\mathcal{F}: \eta \in C(I, T^*\mathbb{R}^n \setminus 0) \mapsto \eta_0 + \int_0^s F(\eta(\tau)) d\tau \in C(I, T^*\mathbb{R}^n \setminus 0).$$

Fix $\epsilon \leq (2\|\nabla F\|)^{-1}$, and let $I = (-\epsilon, \epsilon)$. Then for any $\eta_1, \eta_2 \in C(I, T^*\mathbb{R}^n \setminus 0)$, we have

$$\begin{aligned} \|\mathcal{F}\eta_1 - \mathcal{F}\eta_2\|_{C(I)} &= \left\| \int_0^s [F(\eta_1(\tau)) - F(\eta_2(\tau))] d\tau \right\|_{C(I)} \\ &\leq \epsilon \|F(\eta_1) - F(\eta_2)\|_{C(I)} \leq \epsilon \|\nabla F\| \|\eta_1 - \eta_2\|_{C(I)} \\ &\leq \frac{1}{2} \|\eta_1 - \eta_2\|_{C(I)}. \end{aligned}$$

The Banach fixed-point theorem can be applied, and we can find a fixed point η of \mathcal{F} such that

$$\eta(s) = \eta_0 + \int_0^s F(\eta(\tau)) d\tau, \quad \forall s \in I \quad \Rightarrow \quad \eta \text{ satisfies (9.8).}$$

We proved the existence.

For the uniqueness, assume η_1, η_2 solve (9.8). Because $\eta_1(0) = \eta_2(0)$, if there are not equal, their derivatives must be differ at a point, but this violates the first two equations in (9.8). The first part of the claim is proven.

Part 2: global solution (cf [22, §2]). To obtain the global solution, we can extend the local solution from $(-\epsilon, \epsilon)$ to $[-\epsilon, \epsilon]$, and then just paste local solutions on $[-\epsilon, \epsilon]$, $[\epsilon - \epsilon', \epsilon + \epsilon']$, $[\epsilon + \epsilon' - \epsilon'', \epsilon + \epsilon' + \epsilon'']$, etc. Now we show the endpoints extensions can be done. Assume γ is a local solution on $I = (-\epsilon, \epsilon)$ as given in **Part 1**.

Assume $\nabla_{(x,\xi)} p$ is uniformly bounded in $\{(x(s), \xi(s)/|\xi(s)|); s \in (-\epsilon, \epsilon)\}$, and p is homogeneous of order 1. From (9.8) we can have

$$\frac{d}{ds} (|\xi(s)|^2) = 2\xi(s) \cdot \dot{\xi}(s) = -2\xi(s) \cdot \nabla_x p(x(s), \xi(s)) = m(s)|\xi(s)|^2, \quad (9.9)$$

where $m(s) := -2\hat{\xi}(s) \cdot \nabla_x p(x(s), \hat{\xi}(s))$ with $\hat{\xi}(s) := \xi(s)/|\xi(s)|$. Here because $\xi(0) = \xi_0 \neq 0$, and $\xi(s)$ is continuous on s , so we can choose the interval $I = (-\epsilon, \epsilon)$ to be small enough such that $\xi(s) \neq 0$ for $\forall s \in I$, and this can make $\hat{\xi}(s)$ always well-defined. Solve (9.9) we obtain

$$|\xi(s)| = e^{-\int_0^s \hat{\xi}(\tau) \cdot \nabla_x p(x(\tau), \hat{\xi}(\tau)) d\tau} |\xi(0)|,$$

so

$$e^{-\epsilon M_1} |\xi(0)| \leq |\xi(s)| \leq e^{\epsilon M_1} |\xi(0)|$$

where $M_1 = \sup_{\{(x(s), \xi(s)/|\xi(s)|); s \in (-\epsilon, \epsilon)\}} |\nabla_x p|$, so $\{\xi(s); s \in (-\epsilon, \epsilon)\}$ is contained in a bounded domain. Similarly, for $x(s)$ we have

$$|\dot{x}(s)| = |\nabla_{\xi} p(x(s), \xi(s))| = |\nabla_{\xi} p(x(s), \hat{\xi}(s))| \leq \sup_{\mathbb{R}_x^n \times \mathbb{S}^{n-1}} |\nabla_{\xi} p|$$

where $M_2 = \sup_{\{(x(s), \xi(s)/|\xi(s)|); s \in (-\epsilon, \epsilon)\}} |\nabla_{\xi} p|$. Note that we used the homogeneity of p again. This gives

$$|x(s) - x(0)| \leq sn^{1/2} M_2.$$

Or, if $\nabla_x p$ is uniformly bounded in $\{(x(s), \xi(s)); s \in (-\epsilon, \epsilon)\}$, p is homogeneous of order $\mu \geq 1$, and $\nabla_{\xi} p$ is uniformly bounded in $\{(x(s), \xi(s)/|\xi(s)|); s \in (-\epsilon, \epsilon)\}$, by $|\dot{\xi}(s)| = |\nabla_x p(x(s), \xi(s))|$ we can have

$$|\xi(s)| = |\xi(0) + \int_0^s \dot{\xi}(\tau) d\tau| \leq |\xi(0)| + \int_0^s |\nabla_x p| d\tau \leq |\xi(0)| + sM \leq |\xi(0)| + \epsilon M'_1.$$

where $M'_1 = \sup_{\{(x(s), \xi(s)); s \in (-\epsilon, \epsilon)\}} |\nabla_x p|$. And similarly, for $x(s)$ we have

$$|\dot{x}(s)| = |\nabla_{\xi} p(x(s), \xi(s))| = |\xi(s)|^{\mu-1} |\nabla_{\xi} p(x(s), \hat{\xi}(s))| \leq (|\xi(0)| + \epsilon M'_1) M'_2,$$

where $M'_2 = \sup_{\{(x(s), \xi(s)/|\xi(s)|); s \in (-\epsilon, \epsilon)\}} |\nabla_{\xi} p|$.

Therefore, in both two cases the $(x(s), \xi(s))$ lives in a bounded domain when $s \in (-\epsilon, \epsilon)$, thus due to the continuity of $x(s)$ and $\xi(s)$ we can extend the domain of definition of γ from $(-\epsilon, \epsilon)$ to $[-\epsilon, \epsilon]$.

After extension, we set new initial value (x_0, ξ_0) to be $(x(\epsilon), \xi(\epsilon))$ and by **Part 1** we can get a local solution on $(\epsilon - \epsilon', \epsilon + \epsilon')$ for some small enough ϵ' . By doing this repeatedly, we can obtain a solution defined in \mathbb{R} . The proof is complete. \square

LEMMA 9.11. *Let a symbol p be homogeneous, i.e. $p(x, \lambda\xi) = \lambda p(x, \xi)$ for $\lambda \in \mathbb{R}_+$. Then we have $\boxed{\gamma_{x_0, \lambda\xi_0}(t) = (x(t), \lambda\xi(t))}$ for $\forall \lambda \in \mathbb{R}_+$.*

PROOF. Because p is homogeneous, from (9.8) we have

$$\begin{cases} \dot{x}(s) = \nabla_{\xi} p(x(s), \xi(s)) = \nabla_{\xi} p(x(s), \lambda\xi(s)), \\ \lambda \dot{\xi}(s) = -\lambda \nabla_x p(x(s), \xi(s)) = -\nabla_x p(x(s), \lambda\xi(s)), \\ p(x_0, \lambda\xi_0) = \lambda p(x_0, \xi_0), \end{cases}$$

so $(x(t), \lambda\xi(t))$ is also a solution of (9.8), with $(x(0), \lambda\xi(0)) = (x_0, \lambda\xi_0)$. The proof is done. \square

LEMMA 9.12. *Let $T > 0$. Assume a real-valued symbol $p \in S^1$ is homogeneous of order 1, and $F \in C^\infty([0, T] \times (T^*\mathbb{R}^n \setminus 0))$ and $\phi \in C^\infty(T^*\mathbb{R}^n \setminus 0)$. Then there exists a unique solution $q \in C^\infty(\mathbb{R} \times (T^*\mathbb{R}^n \setminus 0))$ satisfying*

$$\begin{cases} (\partial_t + H_p)q(t, x, \xi) = F(t, x, \xi), \\ q(0, x, \xi) = \phi(x, \xi), \end{cases}$$

where H_p is the Hamiltonian of p . The solution is given by

$$\forall (x_0, \xi_0) \in T^*\mathbb{R}^n, \quad q(t, \gamma_{x_0, \xi_0}(t)) = \phi(x_0, \xi_0) + \int_0^t F(\tau, \gamma_{x_0, \xi_0}(\tau)) d\tau.$$

More, when F and ϕ are homogeneous (with ξ) of order $m \in \mathbb{R}$, then q is also homogeneous (with ξ) of order $m \in \mathbb{R}$.

PROOF. Let $\gamma_{x_0, \xi_0}(t) = (x(t), \xi(t))$ be the bicharacteristic of p starting from (x_0, ξ_0) . The existence of γ_{x_0, ξ_0} is guaranteed by Lemma 9.10. Then we have

$$\begin{aligned} (\partial_t + H_p)q(t, x(t), \xi(t)) &= (\partial_t + \nabla_{\xi} p \cdot \nabla_x - \nabla_x p \cdot \nabla_{\xi})q(t, x(t), \xi(t)) \\ &= (\partial_t + \dot{x}(t) \cdot \nabla_x + \dot{\xi}(t) \cdot \nabla_{\xi})q(t, x(t), \xi(t)) \\ &= \frac{d}{dt}(q(t, x(t), \xi(t))), \end{aligned}$$

so $\frac{d}{dt}(q(t, x(t), \xi(t))) = F(t, x(t), \xi(t))$, which gives

$$\begin{aligned} q(t, x(t), \xi(t)) &= q(0, x(0), \xi(0)) + \int_0^t F(\tau, x(\tau), \xi(\tau)) d\tau \\ &= \phi(x_0, \xi_0) + \int_0^t F(\tau, x(\tau), \xi(\tau)) d\tau. \end{aligned}$$

For the homogeneity, fix $(x, \xi) \in T^*\mathbb{R}^n \setminus 0$, we solve the Hamiltonian equation with initial point (x, ξ) and we can obtain a bicharacteristic $\gamma_{x, \xi}$. Fix $t \in \mathbb{R}$, we set $(x_0, \xi_0) := \gamma_{x, \xi}(-t)$, so reversely we represent (x, ξ) as $\gamma_{x_0, \xi_0}(t) = (x(t), \xi(t))$. Because p is homogeneous of order 1, by Lemma 9.11 we have $(x(t), \lambda\xi(t)) = \gamma_{x_0, \lambda\xi_0}(t)$, so

$$\begin{aligned} q(t, x, \lambda\xi) &= q(t, x(t), \lambda\xi(t)) = q(t, \gamma_{x_0, \lambda\xi_0}(t)) = \phi(x_0, \lambda\xi_0) + \int_0^t F(\tau, \gamma_{x_0, \lambda\xi_0}(\tau)) d\tau \\ &= \phi(x_0, \lambda\xi_0) + \int_0^t F(\tau, (x(t), \lambda\xi(\tau))) d\tau \\ &= \lambda^m [\phi(x_0, \xi_0) + \int_0^t F(\tau, x(\tau), \xi(\tau)) d\tau] = \lambda^m q(t, x(t), \xi(t)). \end{aligned}$$

The proof is done. \square

9.3. Propagation of singularities

For other literature on the topic, [17, §10] is a good reference for this section. See [23, A.1.3], [11, §8] for different proofs. Now we are ready for the main result.

THEOREM 9.13. *Assume $m \in \mathbb{R}$ and $P \in \Psi^m$ is classical Ψ DO of real principal type, and denote its principal symbol as $p_m(x, \xi)$. We assume either*

- $u \in \mathcal{S}'(\mathbb{R}^n)$, or,
- P is properly supported and $u \in \mathcal{D}'(\mathbb{R}^n)$.

Let $Pu \in C^\infty$ and $p_m(x_0, \xi_0) = 0$. If $(x_0, \xi_0) \notin \text{WF}(u)$, then $\gamma_{x_0, \xi_0} \cap \text{WF}(u) = \emptyset$ where the γ_{x_0, ξ_0} is a null bicharacteristic of p_m defined in Definition 9.9. In other words, for a null bicharacteristic γ , it holds either $\gamma \subset \text{WF}(u)$ or $\gamma \cap \text{WF}(u) = \emptyset$.

PROOF. Step 1: change to Ψ^1 . Choose an elliptic $T_a \in \Psi^{1-m}$ with $a(x, \xi) > 0$ and $a(x, \xi)$ be real-valued, then

$$\text{WF}(T_a Pu) = \text{WF}(Pu) \quad \Rightarrow \quad T_a Pu \in C^\infty \text{ if and only if } Pu \in C^\infty,$$

namely, T_a doesn't change the wavefront set.

Also, we can show T_a doesn't change null bicharacteristics of the principal symbols as follows. Assume $(x(s), \xi(s))$ solves (9.8) with $p_m(x(0), \xi(0)) = 0$. **Let's assume we can find a function $f(s)$ such that**

$$f(s) := \int_0^s a(x(f(r)), \xi(f(r))) dr.$$

This is possible because it amounts to find a fix point of the transform $\mathcal{I} \circ a \circ \gamma: C^\infty(\mathbb{R}; \mathbb{R}) \rightarrow C^\infty(\mathbb{R}; \mathbb{R})$ where $\mathcal{I}g(s) := \int_0^s g(r) dr$ and $\gamma(s) := (x(s), \xi(s))$.

After obtained such an f , we can see f is a bijection because $f' = a > 0$. Denote

$$\tilde{x}(s) := x(f(s)), \quad \tilde{\xi}(s) := \xi(f(s)).$$

If $(x(s), \xi(s))$ is defined on a interval I , then we say $(\tilde{x}(s), \tilde{\xi}(s))$ is defined on a interval $I' := f^{-1}(I)$, so $p_m(\tilde{x}(s), \tilde{\xi}(s)) = 0$ for $s \in I'$, and we can have

$$\begin{aligned} \dot{\tilde{x}}(s) &= f'(s)\dot{x}(f(s)) = a(x(f(s)), \xi(f(s)))\nabla_\xi p_m(x(f(s)), \xi(f(s))) \\ &= a(\tilde{x}(s), \tilde{\xi}(s))\nabla_\xi p_m(\tilde{x}(s), \tilde{\xi}(s)) \\ &= a(\tilde{x}(s), \tilde{\xi}(s))\nabla_\xi p_m(\tilde{x}(s), \tilde{\xi}(s)) + p_m(\tilde{x}(s), \tilde{\xi}(s))\nabla_\xi a(\tilde{x}(s), \tilde{\xi}(s)) \\ &= \nabla_\xi (ap_m)(\tilde{x}(s), \tilde{\xi}(s)). \end{aligned}$$

Similarly, we have

$$\dot{\tilde{\xi}}(s) = -\nabla_x (ap_m)(\tilde{x}(s), \tilde{\xi}(s)).$$

These mean the null bicharacteristic $(x(s), \xi(s))$ of p_m , after a reparametrization, is also a null bicharacteristic of ap_m . Note that ap_m is the principal symbol of $T_a P$. Hence, to prove the claim for $P \in \Psi^m$ is equivalent to prove the claim for $P \in \Psi^1$, so, in the rest of the proof we assume $P \in \Psi^1$ of real principal type.

Step 2: find a t -dependent $Q = Q(t, x, D)$ such that

$$Qu|_{t=0} \in C^\infty. \tag{9.10}$$

Our plan is to construct a sequence of t -dependent Ψ DOs $Q_j = Q_j(t, x, D) \in \Psi^{-j}$ ($j \geq 0$) having classical symbol q_j , and set $Q \sim \sum_j Q_j$. Here $Q_j(t, x, D) \in \Psi^{-j}$ means its symbol $q_j(t, x, \xi)$ is in $S^{-j}([0, T] \times \mathbb{R}_x^n \times \mathbb{R}_\xi^n)$, i.e.,

$$|\partial_t^{\alpha'} \partial_x^{\alpha''} \partial_\xi^\beta q_j(t, x, \xi)| \lesssim \langle \xi \rangle^{-j-|\beta|},$$

see Definition 2.4.

Because $(x_0, \xi_0) \notin \text{WF}(u)$, we have $(x_0, t\xi_0) \notin \text{WF}(u)$ for $\forall t > 0$, and we can find a open conic neighborhood ω of (x_0, ξ_0) such that $\omega \cap \text{WF}(u) = \emptyset$. Choose a function $\chi(x, \xi) \in C^\infty$ satisfying

$$\begin{cases} \chi(x, \xi) \in C^\infty, \text{ supp } \chi \subset \omega, \chi(x, \lambda\xi) = \chi(x, \xi) \ (\forall \lambda > 0), \\ \chi \equiv 1 \text{ in a sufficiently small open conic neighborhood } \tilde{\omega} \text{ of } (x_0, \xi_0). \end{cases} \quad (9.11)$$

Set $q_0(0, x, \xi) := \chi(x, \xi)$, then $\omega^c \subset \text{Smo}(Q_0|_{t=0})$ where ω^c signifies the complement of the set ω in $T^*\mathbb{R}^n$, so by Theorem 8.17,

$$\begin{aligned} \text{WF}(Q_0u|_{t=0}) &= \text{WF}((Q_0|_{t=0})u) \subset \text{WF}(u) \setminus \text{Smo}(Q_0|_{t=0}) \\ &= \text{WF}(u) \cap (\text{Smo}(Q_0|_{t=0}))^c \\ &\subset \text{WF}(u) \cap \omega = \emptyset, \end{aligned}$$

so $\underline{Q_0u|_{t=0} \in C^\infty}$. For Q_j ($j \geq 1$), we set their symbol at $t = 0$ as zero, i.e.,

$$q_0(0, x, \xi) := \chi(x, \xi), \quad q_j(0, \cdot, \cdot) := 0 \ (j \geq 1), \quad (9.12)$$

then $Q_j|_{t=0} \in \Psi^{-\infty}$, so $\underline{Q_ju|_{t=0} \in C^\infty}$ for $j \geq 1$. By Theorem 2.6 we can find a Q satisfying $Q \sim \sum_j Q_j$ (thus Q is also t -dependent)¹. Q is of order 0. We can conclude (9.10).

Step 3: to make Q satisfy

$$(D_t + P)(Qu) \in C^\infty. \quad (9.13)$$

To achieve (9.13) is equivalent to achieve

$$[D_t + P, Q] \in \Psi^{-\infty} \quad (9.14)$$

because

$$\begin{aligned} (D_t + P)(Qu) &= [D_t + P, Q]u + Q(D_t + P)u \\ &= [D_t + P, Q]u + QPu = [P, Q]u + C^\infty. \end{aligned}$$

Here we used $\text{WF}(QPu) \subset \text{WF}(Pu) = \emptyset$ so $QPu \in C^\infty$. The fact $QD_tu = 0$ is because u is independent of t . Readers may note that in **Step 2** we only determined q_j on $\{t = 0\}$, while q_j on $\{t > 0\}$ hasn't been fixed yet. Here we design $q_j|_{t>0}$ to achieve (9.14).

We use the notation $\sigma(A)$ to signify the symbol of A . Because P is classical, we can expand $\sigma(P)$ as $\sum_k p_k$ for some homogeneous symbols $p_k \in S^{1-k}$. Recall **Step 1**, we see the integral curve of H_{p_m} is the same as H_{p_1} . Then by Theorem 5.3 and Remark 5.5, we have

$$\begin{aligned} \sigma([D_t, Q]) &\sim \sum_\alpha \frac{(-i)^{|\alpha|}}{\alpha!} (\partial_\tau^\alpha \tau) \partial_t^\alpha \left(\sum_{j \geq 0} q_j \right) - \sum_\alpha \frac{(-i)^{|\alpha|}}{\alpha!} (\partial_t^\alpha \tau) \partial_\tau^\alpha \left(\sum_{j \geq 0} q_j \right) \\ &= \tau \sum_{j \geq 0} q_j + (-i) \partial_t \sum_{j \geq 0} q_j - \tau \sum_{j \geq 0} q_j = \frac{1}{i} \sum_{j \geq 0} \partial_t q_j, \end{aligned} \quad (9.15)$$

and

$$\begin{aligned} \sigma([P, Q]) &\sim \sum_\alpha \frac{(-i)^{|\alpha|}}{\alpha!} \partial_\xi^\alpha \left(\sum_{k \geq 0} p_k \right) \partial_x^\alpha \left(\sum_{j \geq 0} q_j \right) - \sum_\alpha \frac{(-i)^{|\alpha|}}{\alpha!} \partial_x^\alpha \left(\sum_{k \geq 0} p_k \right) \partial_\xi^\alpha \left(\sum_{j \geq 0} q_j \right) \\ &= \sum_{k \geq 0} \sum_{j \geq 0} \sum_\alpha \frac{(-i)^{|\alpha|}}{\alpha!} \left((\partial_\xi^\alpha p_k) \partial_x^\alpha - (\partial_x^\alpha p_k) \partial_\xi^\alpha \right) q_j \end{aligned}$$

¹Note that such Q is not unique.

$$\begin{aligned}
&= \sum_{\ell \geq 0} \sum_{j+k+|\alpha|=\ell} \frac{(-i)^{|\alpha|}}{\alpha!} ((\partial_\xi^\alpha p_k) \partial_x^\alpha - (\partial_x^\alpha p_k) \partial_\xi^\alpha) q_j \\
&= \sum_{\substack{\ell \geq 1 \\ |\alpha| \geq 1}} \sum_{j+k+|\alpha|=\ell} L_{j,k,\alpha} q_j, \quad (\text{it can be checked that } L_{j,k,\alpha} q_j \in S^{1-\ell})
\end{aligned}$$

where the linear differential operator $L_{j,k,\alpha} := \frac{(-i)^{|\alpha|}}{\alpha!} ((\partial_\xi^\alpha p_k) \partial_x^\alpha - (\partial_x^\alpha p_k) \partial_\xi^\alpha)$. Note that p_0 is the principal symbol of P so $\underline{p_0}$ is real-valued. Also note that the restriction $\ell \geq 1$ and $|\alpha| \geq 1$ come from the fact that when $|\alpha| = 0$, $L_{j,k,\alpha} = p_k - p_k = 0$. It can be checked

$$|\alpha| = 1 \Rightarrow \sum_{|\alpha|=1} L_{j,k,\alpha} q_j = \frac{1}{i} H_{p_k} q_j. \quad (9.16)$$

We can further compute $\sigma([P, Q])$ as

$$\begin{aligned}
\sigma([P, Q]) &\sim \frac{1}{i} H_{p_0} q_0 + \sum_{\ell \geq 2} \sum_{\substack{j+k+|\alpha|=\ell \\ |\alpha| \geq 1}} L_{j,k,\alpha} q_j \\
&= \frac{1}{i} H_{p_0} q_0 + \sum_{\ell \geq 2} \left(\sum_{\substack{j+k+|\alpha|=\ell \\ |\alpha| \geq 1, j=\ell-1}} L_{j,k,\alpha} q_j + \sum_{\substack{j+k+|\alpha|=\ell \\ |\alpha| \geq 1, j < \ell-1}} L_{j,k,\alpha} q_j \right) \\
&= \frac{1}{i} H_{p_0} q_0 + \sum_{\ell \geq 2} \left(\sum_{|\alpha|=1} L_{j=\ell-1, k=0, \alpha} q_j + \sum_{\substack{j+k+|\alpha|=\ell \\ |\alpha| \geq 1, j < \ell-1}} L_{j,k,\alpha} q_j \right) \\
&= \frac{1}{i} H_{p_0} q_0 + \sum_{\ell \geq 2} \left(\frac{1}{i} H_{p_0} q_{\ell-1} + \sum_{\substack{j+k+|\alpha|=\ell \\ |\alpha| \geq 1, j < \ell-1}} L_{j,k,\alpha} q_j \right) \quad (\text{by (9.16)}) \\
&= \frac{1}{i} H_{p_0} q_0 + \sum_{\ell \geq 1} \left(\frac{1}{i} H_{p_0} q_\ell + \sum_{\substack{j+k+|\alpha|=\ell+1 \\ |\alpha| \geq 1, j < \ell}} L_{j,k,\alpha} q_j \right) \quad (\ell \rightarrow \ell - 1). \quad (9.17)
\end{aligned}$$

Combining (9.15) with (9.17), we obtain

$$\sigma([D_t + P, Q]) \sim \frac{1}{i} (\partial_t + H_{p_0}) q_0 + \sum_{\ell \geq 1} \left(\frac{1}{i} (\partial_t + H_{p_0}) q_\ell + \sum_{\substack{j+k+|\alpha|=\ell+1 \\ |\alpha| \geq 1, j < \ell}} L_{j,k,\alpha} q_j \right). \quad (9.18)$$

The requirement (9.14) thus amounts to require $\sigma([D_t + P, Q]) \in S^{-\infty}$, namely,

$$\begin{cases} (\partial_t + H_{p_0}) q_0 = 0, \\ \frac{1}{i} (\partial_t + H_{p_0}) q_\ell = - \sum_{\substack{j+k+|\alpha|=\ell+1 \\ |\alpha| \geq 1, j < \ell}} L_{j,k,\alpha} q_j, \quad \ell \geq 1. \end{cases} \quad (9.19)$$

Combining (9.19) with initial condition (9.12), these q_j ($j \geq 0$) can be solved iteratively in $[0, T] \times T^* \mathbb{R}^n$ by using see Lemma 9.12 (recall that p_0 is real-valued), and gives, $\forall (x, \xi) \in T^* \mathbb{R}^n \setminus 0$,

$$\begin{cases} q_0(t, \gamma_{x,\xi}(t)) = \chi(x, \xi), \\ q_\ell(t, \gamma_{x,\xi}(t)) = -i \int_0^t \sum_{\substack{j+k+|\alpha|=\ell+1 \\ |\alpha| \geq 1, j < \ell}} L_{j,k,\alpha} q_j(\tau, \gamma_{x,\xi}(\tau)) d\tau, \quad \ell \geq 1. \end{cases} \quad (9.20)$$

And they guarantee $\sigma([D_t + P, Q]) \in S^{-\infty}$, so (9.14) is achieved, thus (9.13) is satisfied.

By iteration we can show the RHS of (9.20) is of order $-\ell$, so the second conclusion in Lemma 9.12 implies q_ℓ is homogeneous of order $-\ell$, so they are all classical symbols.

Step 4: apply a hyperbolic PDE result. Combining (9.10) and (9.13), we can conclude

$$\begin{cases} (D_t + P)(Qu) = F \text{ in } \mathbb{R}_+ \times \mathbb{R}^n, \\ Qu|_{t=0} = \varphi \text{ on } \mathbb{R}^n, \end{cases} \quad (9.21)$$

for some $F \in C^\infty(\mathbb{R}_+ \times \mathbb{R}^n)$ and $\varphi \in C^\infty(\mathbb{R}^n)$. Now we use Lemma 9.17 in advance to conclude $Qu \in C([0, T], C^\infty(\mathbb{R}^n))$.

Step 5: conclusion. From $Qu \in C([0, T], C^\infty(\mathbb{R}^n))$ we see $Q|_t u \in C^\infty$ for each $t \in [0, T]$, where $Q|_t$ is an abbreviation of $Q(t, x, D)$. From (9.20) we see $q_0(t, \gamma_{x_0, \xi_0}(t)) = \chi(x_0, \xi_0)$. By Lemma 9.11, the homogeneity of p_0 gives $(x(t), \lambda\xi(t)) = \gamma_{x_0, \lambda\xi_0}(t)$, so

$$\begin{aligned} \forall \lambda > 0, \quad q_0(t, x(t), \lambda\xi(t)) &= q_0(t, \gamma_{x_0, \lambda\xi_0}(t)) && \text{(by Lemma 9.11)} && (9.22) \\ &= \chi(x_0, \lambda\xi_0) = \chi(x_0, \xi_0) && \text{(by (9.20), (9.11))} \\ &\neq 0. && \text{(by (9.11))} \end{aligned}$$

This means Q is elliptic at $(x(t), \xi(t)) = \gamma_{x_0, \xi_0}(t)$, i.e. $\gamma_{x_0, \xi_0}(t) \notin \text{Char}(Q|_t)$. Therefore, by Corollary 9.6,

$$\text{WF}(u) \subset \text{WF}(Q|_t u) \cup \text{Char}(Q|_t) = \text{Char}(Q|_t),$$

so

$$\gamma_{x_0, \xi_0}(t) \notin \text{WF}(u), \text{ for any } t \in [0, T].$$

which means $\gamma_{x_0, \xi_0} \cap \text{WF}(u) = \emptyset$. The proof is complete. \square

REMARK 9.14. The condition that P is of real principal type is used in the following ways:

- real-valued: in **Step 3**, in order to use Lemma 9.12, p_m has to be real-valued; Also, when p_m is real-valued, then $R := iP + (iP)^*$ is of order 0. This is used in **Step 4** which calls for Lemma 9.17;
- $|\nabla_\xi p_m(x, \xi)| \neq 0$ when $p(x, \xi) = 0$: **related to the solvability of (9.20)? Every $(x, \xi) \in T^*\mathbb{R}^n \setminus 0$ shall be reachable;**
- homogeneity: the condition “ $p_m(x, \lambda\xi) = \lambda p_m(x, \xi)$ ” is used at (9.22) to guarantee $(x(t), \lambda\xi(t)) = \gamma_{x_0, \lambda\xi_0}(t)$.

Theorem 9.13 can be interpreted by the following claim.

COROLLARY 9.15. *Assume $m \in \mathbb{R}$ and $P \in \Psi^m$ is classical Ψ DO of real principal type, and denote its symbol as $p(x, \xi)$. Assume Pu is well-defined and $Pu \in C^\infty$. Then $\text{WF}(u)$ is made of null bicharacteristic curves $\gamma_{x, \xi}$ for some $(x, \xi) \in p_m^{-1}(0)$.*

PROOF. We see that P is a Ψ DO with principal symbol $p_m(x, \xi)$ homogeneous in ξ , so we can apply Lemma 9.4 to conclude $\text{Char} P = p_m^{-1}(0)$. Also, when $Pu \in C^\infty$, by Corollary 9.6 we have $\text{WF}(u) \subset \text{Char} P$, so

$$\text{WF}(u) \subset p_m^{-1}(0). \quad (9.23)$$

For any $(x, \xi) \in \text{WF}(u)$, by (9.23) we know $(x, \xi) \in p_m^{-1}(0)$. Denote as $\gamma_{x, \xi}$ the null bicharacteristic of p_m passing through (x, ξ) , then $\gamma_{x, \xi} \subset p_m^{-1}(0)$ because the value of p_m is constant in bicharacteristics. According to Theorem 9.13, we can conclude $\gamma_{x, \xi} \subset \text{WF}(u)$. In summary, for every $(x, \xi) \in \text{WF}(u)$ we have $\gamma_{x, \xi} \subset \text{WF}(u)$ and $\gamma_{x, \xi}$ is a null bicharacteristic, so $\text{WF}(u)$ is made of null bicharacteristic curves. \square

9.4. Cauchy problems of hyperbolic PDEs

LEMMA 9.16. *Assume $T > 0$ and $s \in \mathbb{R}$, $P \in \Psi^1$ has a real-valued principal symbol. Denote $L = D_t + P$. There exists a constant $\lambda_0 > 0$ such that for any*

$$u \in C^1([0, T], H^s(\mathbb{R}^n)) \cap C([0, T], H^{s+1}(\mathbb{R}^n)),$$

we have

$$\sup_{t \in [0, T]} e^{-\lambda t} \|u(t, \cdot)\|_{H^s} \leq \|u(0, \cdot)\|_{H^s} + 2 \int_0^T e^{-\lambda t} \|Lu(t, \cdot)\|_{H^s} dt. \quad (9.24)$$

PROOF. Denote $Q = iP$ and $L' = iL = \partial_t + Q$. Then

$$\sigma(Q + Q^*) = \sigma(iP + (iP)^*) = i\sigma(P - P^*) = i[\sigma(P) + S^0 - \overline{\sigma(P)} - S^0] \in S^0,$$

because the principal symbol of P is real-valued. We denote $R = Q + Q^*$, then $R \in \Psi^0$ and thus is bounded in L^2 .

We prove the case $s = 0$ first. Denote $f(t) := \|e^{-\lambda t} u(t, \cdot)\|_{L^2}^2$, then

$$\begin{aligned} f'(t) &= 2e^{-2\lambda t} \Re(\partial_t u, u) - 2\lambda f(t) = 2e^{-2\lambda t} \Re((L' - Q)u, u) - 2\lambda f(t) \\ &= 2e^{-2\lambda t} \Re(L'u, u) + e^{-2\lambda t} (-Ru, u) - 2\lambda f(t) \\ &\leq 2e^{-2\lambda t} \|Lu(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2} + e^{-2\lambda t} \|Ru(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2} - 2\lambda f(t) \\ &\leq 2e^{-2\lambda t} \|Lu(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2} - (2\lambda - \|R\|) \|e^{-\lambda t} u(t, \cdot)\|_{L^2}^2 \\ &\leq 2e^{-2\lambda t} \|Lu(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2}, \quad (\text{when } \lambda > \|R\|/2), \end{aligned}$$

where $\|R\|$ is the L^2 operator norm. Hence, for any $t \in [0, T]$,

$$\begin{aligned} e^{-2\lambda t} \|u(t, \cdot)\|_{L^2}^2 &\leq \|u(0, \cdot)\|_{L^2}^2 + 2 \int_0^t e^{-2\lambda s} \|Lu(s, \cdot)\|_{L^2} \|u(s, \cdot)\|_{L^2} ds \\ &\leq \|u(0, \cdot)\|_{L^2}^2 + 2 \int_0^T e^{-2\lambda t} \|Lu(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2} dt. \end{aligned}$$

By denoting $M := \sup_{t \in [0, T]} e^{-\lambda t} \|u(t, \cdot)\|_{H^s}$, we can continue

$$\begin{aligned} M^2 &\leq \|u(0, \cdot)\|_{L^2}^2 + 2 \int_0^T e^{-2\lambda t} \|Lu(t, \cdot)\|_{L^2} \|u(t, \cdot)\|_{L^2} dt \\ &\leq M \|u(0, \cdot)\|_{L^2} + 2 \int_0^T e^{-\lambda t} \|Lu(t, \cdot)\|_{L^2} M dt \\ &\leq M (\|u(0, \cdot)\|_{L^2} + 2 \int_0^T e^{-\lambda t} \|Lu(t, \cdot)\|_{L^2} dt). \end{aligned}$$

We arrive at the conclusion for $s = 0$.

For $s \neq 0$, we can do something similar as in **Step 2** of the proof of Theorem 6.18. This completes the proof. \square

Based on the energy estimate in Lemma 9.16, we can obtain the following result.

LEMMA 9.17. *Assume $T > 0$ and $s \in \mathbb{R}$, $P \in \Psi^1$ has a real-valued principal symbol. Let $f \in L^1((0, T), H^s(\mathbb{R}^n))$ and $\phi \in H^s(\mathbb{R}^n)$. Then there is a unique solution $u \in C([0, T], H^s(\mathbb{R}^n))$ of the PDE*

$$\begin{cases} (D_t + P)u = f & \text{in } (0, T) \times \mathbb{R}^n, \\ u|_{t=0} = \phi & \text{on } \mathbb{R}^n, \end{cases} \quad (9.25)$$

PROOF. **Step 1:** variational formulation. Denote

$$\begin{cases} X := \{\varphi \in C^\infty([0, T] \times \mathbb{R}^n); \varphi(T, \cdot) \equiv 0\} \\ \ell(\varphi) := \int_0^T (f, \varphi) dt + \frac{1}{i}(\phi, \varphi). \end{cases}$$

We say $u \in \mathcal{S}'([0, T] \times \mathbb{R}^n)$ is a weak solution of (9.25) if u satisfies

$$\int_0^T (u, (D_t + P^*)\varphi) dt = \ell(\varphi), \quad \forall \varphi \in X. \quad (9.26)$$

To find a $u \in L^\infty((0, T), H^s)$ satisfying (9.26), we are to show $|\ell(\varphi)| \leq \|(D_t + P^*)\varphi\|_{L^1 H^{-s}}$, and then call for the Hahn-Banach theorem. Here $\|f\|_{L^1 H^{-s}}$ is a shorthand for $\int_0^T \|f\|_{H^{-s}} dt$.

Step 2: energy estimate. Because $P \in \Psi^1$ has a real-valued principal symbol, we see $-P^* \in \Psi^1$ and $-P^*$ also has a real-valued principal symbol. Apply Lemma 9.16 to $D_t + (-P^*)$ and $\varphi(T - t, x)$ we obtain

$$\sup_{t \in [0, T]} e^{-\lambda t} \|\varphi(T - t, \cdot)\|_{H^{-s}} \leq \|\varphi(T, \cdot)\|_{H^{-s}} + 2 \int_0^T e^{-\lambda t} \|(D_t + (-P^*))(\varphi(T - t, \cdot))\|_{H^{-s}} dt,$$

which gives

$$\sup_{t \in [0, T]} e^{\lambda t} \|\varphi(t, \cdot)\|_{H^{-s}} \leq 2 \int_0^T e^{\lambda t} \|(D_t + P^*)\varphi(t, \cdot)\|_{H^{-s}} dt.$$

so

$$\forall s \in \mathbb{R}, \quad \sup_{t \in [0, T]} \|\varphi(t, \cdot)\|_{H^{-s}} \leq 2e^{\lambda T} \|(D_t + P^*)\varphi\|_{L^1 H^{-s}}. \quad (9.27)$$

This means the map $\varphi \mapsto (D_t + P^*)\varphi$ is injective. (9.27) can be understood as a coercive condition.

Step 3: Hahn-Banach theorem. By using (9.27), we can estimate ℓ as follows,

$$\begin{aligned} |\ell(\varphi)| &\leq \int_0^T |(f, \varphi)| dt + |(\phi, \varphi)| \leq \int_0^T \|f\|_{H^s} \|\varphi\|_{H^{-s}} dt + \|\phi\|_{H^s} \|\varphi\|_{H^{-s}} \\ &\leq \left(\int_0^T \|f\|_{H^s} dt + \|\phi\|_{H^s} \right) \sup_{t \in [0, T]} \|\varphi(t, \cdot)\|_{H^{-s}} \\ &\leq C \left(\int_0^T \|f\|_{H^s} dt + \|\phi\|_{H^s} \right) \|(D_t + P^*)\varphi\|_{L^1 H^{-s}}. \end{aligned}$$

Therefore, the linear functional $\ell(\varphi)$ is also a linear functional for $(D_t + P^*)\varphi \in X$ under the norm $L^1((0, T), H^{-s})$. Because the dual space of $L^1((0, T), H^{-s})$ is $L^\infty((0, T), H^s)$, by the Hahn-Banach theorem, there exists a $u \in L^\infty((0, T), H^s)$ such that

$$\ell(\varphi) = (u, (D_t + P^*)\varphi)_{t,x}, \quad \forall \varphi \in X,$$

which is (9.26). This u is a weak solution.

Step 4: weak to strong solution. Because u is a distribution, on $(0, T)$ we have

$$D_t u + P u = f.$$

Because $u \in L^\infty((0, T), H^s)$, $P u \in L^\infty((0, T), H^{s-1})$.

Let f, ϕ be Schwartz, then $f \in L^\infty([0, T], H^s)$, so $D_t u = f - P u \in L^\infty((0, T), H^{s-1})$, which implies

$$u \in C([0, T], H^{s-1}).$$

Again, $f \in C([0, T], H^{s-2})$ and $P u \in C([0, T], H^{s-2})$, so $D_t u = f - P u \in C((0, T), H^{s-2})$, which implies

$$u \in C^1([0, T], H^{s-2}) \cap C([0, T], H^{s-1}) \text{ with } u(0) = \phi.$$

Due to the arbitrary of s , we can conclude

$$u \in C^1([0, T], H^s) \cap C([0, T], H^{s+1}) \text{ with } u(0) = \phi. \quad (9.28)$$

Therefore, $(u, (D_t + P^*)\varphi)_{t,x}$ can be legally write as $((D_t + P)u, \varphi)_{t,x}$, which implies u is a strong solution of (9.25).

Step 5: density arguments for f, ϕ . (9.28) is true when f and ϕ are Schwartz. For general $f \in L^1((0, T), H^s(\mathbb{R}^n))$ and $\phi \in H^s(\mathbb{R}^n)$, due to the density, we can find $\{f_k\} \subset \mathcal{S}$ and $\{\phi_k\} \subset \mathcal{S}$ such that

$$\begin{aligned} f_k &\rightarrow f \text{ in } L^1((0, T), H^s(\mathbb{R}^n)), \quad \phi_k \rightarrow \phi \text{ in } H^s(\mathbb{R}^n), \quad \text{and} \\ (D_t + P)u_k &= f_k, \quad u|_{t=0} = \phi_k, \quad u_k \in C^1([0, T], H^s) \cap C([0, T], H^{s+1}). \end{aligned} \quad (9.29)$$

From (9.29) and Lemma 9.16 we can obtain

$$e^{-\lambda T} \|u_k - u_{k'}\|_{C([0, T], H^s)} \leq \|\phi_k - \phi_{k'}\|_{H^s} + 2\|(f_k - f_{k'})(t, \cdot)\|_{L^1((0, T), H^s)},$$

so $\{u_k\}$ is Cauchy in $C([0, T], H^s)$ and the limit $u \in C([0, T], H^s)$ is a desired solution.

Step 6: uniqueness. By the energy estimate (9.24) it is easy to show the uniqueness of u .

The proof is complete. □

Exercise

EXERCISE 9.1. Proof Lemma 9.4.

EXERCISE 9.2. Prove the function a constructed in (9.4) is in S^0 .

EXERCISE 9.3. Show that a defined in (9.6) gives $(x_0, \xi_0) \notin \text{Char } T_a$. Hint: to borrow ideas from Lemma 6.17.

Bibliography

- [1] S. Alinhac and P. Gérard, *Pseudo-differential operators and the Nash-Moser theorem*, Graduate Studies in Mathematics, vol. 82, American Mathematical Society, Providence, RI, 2007. Translated from the 1991 French original by Stephen S. Wilson. MR2304160
- [2] A. P. Calderón and R. Vaillancourt, *On the boundedness of pseudo-differential operators*, J. Math. Soc. Japan **23** (1971), 374–378. MR284872
- [3] A. P. Calderón and R. Vaillancourt, *A class of bounded pseudo-differential operators*, Proc. Nat. Acad. Sci. U.S.A. **69** (1972), 1185–1187. MR298480
- [4] S. Chen, *Pseudodifferential operators (Chinese)*, Higher Education Press, 2017. ISBN: 9787040186758.
- [5] Q. Chou, S. Chen, J. Shi, J. Liu, and L. Jiang, *Fourier integral operators theory and its applications (Chinese)*, Science Press, 1997. ISBN: 9787030059925.
- [6] R. R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque, vol. 57, Société Mathématique de France, Paris, 1978. With an English summary. MR518170
- [7] M. V. de Hoop, S. Holman, and G. Uhlmann, *Microlocal analysis of seismic body waves and linearized inverse problems*, 202X. in preparation.
- [8] M. Dimassi and J. Sjöstrand, *Spectral asymptotics in the semi-classical limit*, London Mathematical Society Lecture Note Series, vol. 268, Cambridge University Press, Cambridge, 1999. MR1735654
- [9] G. Eskin, *Lectures on linear partial differential equations*, Graduate Studies in Mathematics, vol. 123, American Mathematical Society, Providence, RI, 2011. MR2809923
- [10] F. G. Friedlander, *Introduction to the theory of distributions*, Second, Cambridge University Press, Cambridge, 1998. With additional material by M. Joshi. MR1721032
- [11] A. Grigis and J. Sjöstrand, *Microlocal analysis for differential operators*, London Mathematical Society Lecture Note Series, vol. 196, Cambridge University Press, Cambridge, 1994. An introduction. MR1269107
- [12] L. Hörmander, *On the L^2 continuity of pseudo-differential operators*, Comm. Pure Appl. Math. **24** (1971), 529–535. MR281060
- [13] L. Hörmander, *The analysis of linear partial differential operators. I-V*, Classics in Mathematics, Springer-Verlag, Berlin, 2003. MR1996773
- [14] I. L. Hwang, *The L^2 -boundedness of pseudodifferential operators*, Trans. Amer. Math. Soc. **302** (1987), no. 1, 55–76. MR887496
- [15] I. L. Hwang and R. B. Lee, *L^p -boundedness of pseudo-differential operators of class $S_{0,0}$* , Trans. Amer. Math. Soc. **346** (1994), no. 2, 489–510. MR1264147
- [16] Z. Jiang and S. Sun, *Functional analysis*, China Higher Education Press, 2006.
- [17] M. Joshi, *Introduction to pseudo-differential operators*, arXiv preprint math.AP (1999).
- [18] J. M. Lee and G. Uhlmann, *Determining anisotropic real-analytic conductivities by boundary measurements*, Comm. Pure Appl. Math. **42** (1989), no. 8, 1097–1112. MR1029119
- [19] S. Ma, *Study note to Hormander’s book* (2020July). in preparation.
- [20] S. Ma, *Supplementary materials* (2020July). in preparation.
- [21] A. Martinez, *An introduction to semiclassical and microlocal analysis*, Universitext, Springer-Verlag, New York, 2002. MR1872698
- [22] M. Salo, *Stability for solutions of wave equations with $C^{1,1}$ coefficients*, Inverse Probl. Imaging **1** (2007), no. 3, 537–556. MR2308978
- [23] M. A. Shubin, *Pseudodifferential operators and spectral theory*, Second, Springer-Verlag, Berlin, 2001. Translated from the 1978 Russian original by Stig I. Andersson. MR1852334
- [24] C. D. Sogge, *Fourier integrals in classical analysis*, Second, Cambridge Tracts in Mathematics, vol. 210, Cambridge University Press, Cambridge, 2017. MR3645429
- [25] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J., 1970. MR0290095
- [26] M. W. Wong, *An introduction to pseudo-differential operators*, Third, Series on Analysis, Applications and Computation, vol. 6, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2014. MR3222682

- [27] R. Wong, *Asymptotic approximations of integrals*, Computer Science and Scientific Computing, Academic Press, Inc., Boston, MA, 1989. MR1016818
- [28] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics, vol. 138, American Mathematical Society, Providence, RI, 2012. MR2952218

Index

- adjoint, 47
- adjoint of quantizations, 68
- asymptotics, 13

- bicharacteristic, 98

- canonical relation, 84, 85
- Carleman estimates, 75
- commutator, 48
- composition of Ψ DOs, 44
- composition of quantizations, 67, 71
- conic set, 76

- direct product, 79

- ellipticity, 50

- Fourier integral operators, 85
- Fourier Transform, 9

- Gårding's inequality, 56

- kernel, 19, 26

- microlocal ellipticity, 95
- microlocal parametrix, 95

- null bicharacteristic, 98

- oscillatory integral I, 25
- oscillatory integral II, 28

- parametrix, 49
- Parseval's Relation, 9

- phase function, 22
- Plancherel Theorem, 9
- product of distributions, 80
- propagation of the singularities, 101
- properly supported, 19
- pseudodifferential operators, 15
- pseudolocal property, 20

- Schur estimate, 52
- Schwartz function, 6
- Schwartz Space, 6
- sharp Gårding's Inequality, 57
- singular support, 20
- smooth direction, Smo, 76
- Sobolev spaces, 17
- stationary phase lemma, 33
- strong ellipticity, 55
- symbols, 12
- symplectic 2-form, 69
- symplectic product, 68

- Taylor's expansion, 31
- tempered Distributions, 8
- tempered functions, 7
- transpose, 47
- twist, 85

- wavefront set, 77