

TITCHMARSH THEOREMS, HAUSDORFF-YOUNG-PALEY INEQUALITY AND $L^p - L^q$ BOUNDEDNESS OF FOURIER MULTIPLIERS ON HARMONIC NA GROUPS

VISHVESH KUMAR AND MICHAEL RUZHANSKY

ABSTRACT. In this paper we extend classical Titchmarsh theorems on the Fourier transform of Hölder-Lipschitz functions to the setting of harmonic NA groups, which relate smoothness properties of functions to the growth and integrability of their Fourier transform. We prove a Fourier multiplier theorem for L^2 -Hölder-Lipschitz spaces on Harmonic NA groups. We also derive conditions and a characterisation of Dini-Lipschitz classes on Harmonic NA groups in terms of the behaviour of their Fourier transform. Then, we shift our attention to the spherical analysis on Harmonic NA group. Since the spherical analysis on these groups fits well in the setting of Jacobi analysis we prefer to work in the Jacobi setting. We prove L^p - L^q boundedness of Fourier multipliers by extending a classical theorem of Hörmander to the Jacobi analysis setting. On the way to accomplish this classical result we prove Paley-type inequality and Hausdorff-Young-Paley inequality. We also establish L^p - L^q boundedness of spectral multipliers of the Jacobi Laplacian.

CONTENTS

1. Introduction	2
2. Essentials about harmonic NA groups	6
3. Titchmarsh theorems for (Helgason) Fourier transform of Hölder-Lipschitz functions on harmonic NA groups	10
3.1. Growth properties of Fourier transform of Hölder-Lipschitz functions on harmonic NA groups	10
3.2. Applications	12
3.3. An integrability result for Fourier transform of Hölder-Lipschitz functions on harmonic NA groups	13
4. Helgason transform of Dini-Lipschitz functions on harmonic NA groups	15
5. $L^p - L^q$ boundedness of multipliers for Jacobi transform for $1 < p \leq 2 \leq q < \infty$	20
5.1. Paley-type inequality for Jacobi transform	21
5.2. Hausdorff-Young-Paley inequality for Jacobi transform	24
Acknowledgment	27

2010 *Mathematics Subject Classification.* Primary 43A85 Secondary 22E30.

Key words and phrases. Titchmarsh theorems, Hölder-Lipschitz, Harmonic NA groups, Helgason transform, Paley-type inequality, Hausdorff-Young-Paley inequality, L^p - L^q boundedness, Fourier multipliers, Jacobi transform, Spectral multipliers

1. INTRODUCTION

Harmonic NA groups (also known as Damek-Ricci spaces) form a class of solvable Lie groups (non-unimodular), equipped with a left invariant Riemannian metric, called “harmonic extension of H -type groups”, is a natural generalisation of the Iwasawa NA groups of the real rank-one simple Lie groups. In particular, as Riemannian manifolds, the class of harmonic NA groups contains rank-one symmetric spaces of non-compact type as a very small subclass. Although, the rank one noncompact Riemannian symmetric spaces are the most discernible prototypes of harmonic NA groups, they have many contrasts with these. From the geometric point of view, Harmonic NA groups, in general, are not symmetric which can be seen through the fact that the geodesic inversion is not an isometry. On the other hand, these spaces cannot have Kunze-Stein property for being noncompact amenable groups.

The studies of relations between the smoothness of the functions and the growth and (or) the integrability of their Fourier coefficient are among the classical and fundamental problems in Fourier analysis. These types of studies include the classical theorems of Fourier transform like Riemann-Lebesgue Lemma relating the integrability of a functions and decay of its Fourier transform, and the Hausdorff-Young inequality relating the integrability of a function and its Fourier transform. Indeed, the Hausdorff-Young inequality on \mathbb{R} states that for $f \in L^p(\mathbb{R})$, $1 \leq p \leq 2$, its Fourier transform $\widehat{f} \in L^{p'}(\mathbb{R})$ with $\frac{1}{p} + \frac{1}{p'} = 1$. Titchmarsh [46] studies the problem of how much this fact can be strengthened if f additionally satisfies a Lipschitz condition. Indeed, in this vain, Titchmarsh [46] showed the this can be significantly improved. The first Titchmarsh theorem we deal with [46, Theorem 37] in this paper is recalled below: We define Hölder-Lipschitz space $\text{Lip}_{\mathbb{R}}(\alpha, p)$ for $0 < \alpha \leq 1$ and $1 < p < \infty$ by

$$\text{Lip}_{\mathbb{R}}(\alpha, p) := \{f \in L^p(\mathbb{R}) : \|f(t + \cdot) - f(\cdot)\|_{L^p(\mathbb{R})} = O(t^\alpha) \text{ as } t \rightarrow 0\}.$$

Theorem A. Let $0 < \alpha \leq 1$ and $1 < p \leq 2$. If $f \in \text{Lip}_{\mathbb{R}}(\alpha, p)$, then $\widehat{f} \in L^\beta(\mathbb{R})$ for

$$\frac{p}{p + \alpha p - 1} < \beta \leq p', \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

To discuss the improvement in the conclusion $\widehat{f} \in L^\beta(\mathbb{R})$ in Theorem A above in comparison to the Hausdorff-Young inequality, for instance we can take $\alpha = \frac{1}{2}$ and $p = 2$ then $\frac{p}{p + \alpha p - 1} = 1$, therefore $\widehat{f} \in L^\beta(\mathbb{R})$ for all $\beta \in (1, 2]$. The above theorem has been investigated in several different settings, e.g. on compact symmetric spaces of rank one, noncompact symmetric spaces of rank one, compact homogeneous manifolds, and the Euclidean space \mathbb{R}^n [42, 43, 27, 12, 50, 51]. Notably, our motivation is the work of Bray [12] on \mathbb{R}^n where he extends Theorem A in higher dimenstions, and a recent work of the third author [27] on compact homogeneous spaces, as well as the work of Ray and Sarkar [44] where they proved a version of the Hausdorff-Young inequality for Damek-Ricci spaces. In the context of Dini-Lipschitz spaces, Theorem A has been studied in the setting on \mathbb{R} by Younis [49]. In Theorem A' and Theorem 4.3 we extend the

Theorem A to the setting of Damek-Ricci spaces. It is worth noting that Theorem 4.3 is already new in rank one noncompact symmetric space setting.

The second Titchmarsh theorem [46, Theorem 85] which is also of our interest is stated below.

Theorem B. Let $0 < \alpha \leq 1$ and $f \in L^2(\mathbb{R})$. Then $f \in \text{Lip}_{\mathbb{R}}(\alpha, 2)$ if and only if

$$\int_{|\xi| > \frac{1}{t}} |\widehat{f}(\xi)|^2 d\xi = O(t^{2\alpha}) \quad \text{as } t \rightarrow 0.$$

This theorem was extended by Bray [12] to higher dimensional Euclidean spaces in a more general setting using multipliers by modifying the technique given in the seminal paper of Platonov [42] in the case of rank one noncompact symmetric spaces. For an overview of extensions of this theorem in different settings we refer to [27, 50, 51]. In Theorem B' and Theorem 4.1 we extend this theorem to the setting of Helgason Fourier transform on Damek-Ricci spaces.

To state our main results briefly, we first need to define the Hölder-Lipschitz spaces $\text{Lip}_S(\alpha, p)$ in a harmonic NA group S . We define $\text{Lip}_S(\alpha, p)$ using the spherical mean operator M_t , $t \in \mathbb{R}_+$, on Damek-Ricci spaces (see Section 2 for definition and more details). The use of the spherical mean operator is inspired by the work Platonov [42, 43] and Bray [12]. In the setting of Damek-Ricci spaces, the spherical mean operator was used in [36] to answer some questions initially posed by Bray and Pinsky [13] regarding the growth of Fourier transform. Thus, the Hölder-Lipschitz space $\text{Lip}_S(\alpha, p)$ is defined as

$$\text{Lip}_S(\alpha, p) := \{f \in L^p(S) : \|M_t f - f\|_p = O(t^\alpha) \quad \text{as } t \rightarrow 0\}.$$

Now, we state both of our results which are the suitable analogues of Titchmarsh theorems in the context of harmonic NA groups. We denote by \widetilde{f} the (Helgason) Fourier transform of f on the harmonic NA group S (see [9]). Also, we set $\gamma_p = \frac{2}{p} - 1$ and so $\gamma_{p'} = \frac{2}{p'} - 1 = -\gamma_p$, where p' is the Lebesgue conjugate of p .

Theorem A'. Let S be a harmonic NA group of dimension d . Let $0 < \alpha \leq 1$, $1 < p \leq 2$, and let p' be such that $\frac{1}{p} + \frac{1}{p'} = 1$. Let $f \in \text{Lip}_S(\alpha; p)$. Then the function F defined by

$$F(\lambda) := \left(\int_N |\widetilde{f}(\lambda + i\gamma_{p'} \rho, n)|^{p'} dn \right)^{\frac{1}{p'}}, \quad 1 < p \leq 2,$$

belongs to $L^\beta((0, \infty), |c(\lambda)|^{-2} d\lambda)$ provided that

$$\frac{dp}{dp + \alpha p - d} < \beta \leq p'.$$

Theorem B'. Let $0 < \alpha \leq 1$ and $f \in L^2(S)$. Then $f \in \text{Lip}_S(\alpha; 2)$ if and only if

$$\int_{\lambda > \frac{1}{t}} \int_N |\widetilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda = O(t^{2\alpha}) \quad \text{as } t \rightarrow 0.$$

We also prove the aforementioned theorem for Dini-Lipschitz spaces on harmonic NA groups. As an application of the above characterisation we show a Fourier multiplier theorem for Hölder-Lipschitz spaces on Damek-Ricci spaces.

Corollary 1.1. *Let $0 \leq \gamma < 1$ and let h be an even bounded measurable function on \mathbb{R} such that*

$$|h(\lambda)| \leq C \langle \lambda \rangle^{-\gamma},$$

where $\langle \lambda \rangle = (\lambda^2 + \frac{Q^2}{2})^{\frac{1}{2}}$, and Q denotes the homogeneous dimension of N . Let A be the Fourier multiplier with symbol h , i.e., given by $\widetilde{A}f(\lambda, n) = h(\lambda)\widetilde{f}(\lambda, n)$ for all $\lambda \in \mathbb{R}_+$ and $n \in N$. Then

$$A : Lip_S(\alpha; 2) \rightarrow Lip_S(\alpha + \gamma; 2)$$

is bounded for all α such that $0 < \alpha < 1 - \gamma$.

The multiplier theorem in Corollary 1.1 complements the other known multipliers theorems on Damek-Ricci spaces [6].

Now, we would like to divert our attention to the second part of the paper where we establish an L^p - L^q multiplier theorem for the Jacobi transform. In particular these results are true for the radial multipliers on Damek-Ricci spaces as the spherical analysis on Damek-Ricci spaces fits well in the setting of Jacobi analysis as observed in [6]. Here we deal with L^p - L^q multipliers as opposed to the L^p -multipliers for which theorems of Mihlin-Hörmander or Marcinkiewicz type provide results for Fourier multiplier in different settings based on the regularity of the symbol. We cite here [34, 33, 4, 5, 6, 20, 18, 19, 21, 45, 28, 29, 8, 7, 41, 10] to mention a few of them. In [6], several radial L^p -multipliers theorems have been established including the Mihlin-Hörmander multiplier theorem on Damek-Ricci spaces. The Paley-type inequality describes the growth of the Fourier transform of a function in terms of its L^p -norm. Interpolating the Paley-inequality with the Hausdorff-Young inequality one can obtain the following Hörmander's version of the Hausdorff-Young-Paley inequality,

$$\left(\int_{\mathbb{R}^n} |(\mathcal{F}f)(\xi)\phi(\xi)^{\frac{1}{r}-\frac{1}{p'}}|^r \right)^{\frac{1}{r}} \leq \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 < p \leq r \leq p' < \infty, \quad 1 < p < 2. \quad (1)$$

Also, as a consequence of the Hausdorff-Young-Paley inequality, Hörmander [33, page 106] proves that the condition

$$\sup_{t>0} t^b \{ \xi \in \mathbb{R}^n : m(\xi) \geq t \} < \infty, \quad \frac{1}{p} - \frac{1}{q} = \frac{1}{b}, \quad (2)$$

where $1 < p \leq 2 \leq q < \infty$, implies the existence of a bounded extension of a Fourier multiplier T_m with symbol m from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n)$. Recently, the third author with his collaborator R. Akylzhanov studied Hörmander classical results for unimodular locally compact groups and homogeneous spaces [2, 3]. In [3], the key idea behind the extension of Hörmander theorem is the reformulation of this theorem as follows:

$$\|T_m\|_{L^p(\mathbb{R}^n) \rightarrow L^q(\mathbb{R}^n)} \lesssim \sup_{s>0} s \left(\int_{\{\xi \in \mathbb{R}^n : m(\xi) \geq s\}} d\xi \right)^{\frac{1}{p}-\frac{1}{q}} \simeq \|m\|_{L^{r,\infty}(\mathbb{R}^n)} \simeq \|T_m\|_{L^{r,\infty}(\text{VN}(\mathbb{R}^n))},$$

where $\frac{1}{r} = \frac{1}{p} - \frac{1}{q}$, $\|m\|_{L^{r,\infty}(\mathbb{R}^n)}$ is the Lorentz norm of m , and $\|T_m\|_{L^{r,\infty}(\text{VN}(\mathbb{R}^n))}$ is the norm of the operator T_m in the Lorentz space on the group von Neumann algebra $\text{VN}(\mathbb{R}^n)$ of \mathbb{R}^n . They use the Lorentz spaces and group von Neumann algebra techniques for extending it to general locally compact unimodular groups. The unimodularity assumption has its own advantages such as existence of the canonical

trace on the group von Neumann algebra and consequently, Plancherel formula and the Hausdorff-Young inequality. They also pointed out that the unimodularity can be avoided by using the Tomita-Takesaki modular theory and the Haagerup reduction technique. In our case, Harmonic NA groups are nonunimodular groups but we are dealing with the Helgason Fourier transform and Harish Chandra transform instead of the group Fourier transform. So, the groups von Neumann algebra techniques are more intricate to implement. Untill now, there does not exist a suitable and optimal version of the Hausdorff-Young inequality for the (Helgason) Fourier transform for Damek-Ricci space, in particular, for rank one non-compact symmetric spaces, although several attempts has been made for it and consequently, different version of the Hausdorff-Young inequality were proved, for example, see [44, 40]. On the other hand, spherical analysis on Damek-Ricci spaces ([6]) fits perfectly in the well-established Jacobi analysis setup [37, 31, 32]. Therefore, we choose to work in the setting of Jacobi analysis, in particular, the spherical analysis on Damek-Ricci spaces. In this paper, we prove Paley-type inequality, Hausdorff-Young-Paley inequality and Hörmander multiplier theorem for Jacobi transform on half line \mathbb{R}_+ . For rest of the section, we assume that $\alpha \neq -1, -2, \dots$ and $\alpha \geq \beta > \frac{-1}{2}$. We set $A_{\alpha, \beta}(t) = (2 \sinh t)^{2\alpha+1} (2 \cosh t)^{2\beta+1}$, $t > 0$ and denote by $c(\lambda)$, a multiple of the meromorphic Harish-Chandra function given by the formula

$$c(\lambda) := \frac{2^{\rho-i\lambda} \Gamma(\alpha+1) \Gamma(i\lambda)}{\Gamma(\frac{1}{2}(\rho+i\lambda)) \Gamma(\frac{1}{2}(\rho+i\lambda) - \beta)}.$$

We refer to Section 5 for more details and all the notation used here. The following theorem is an analogue of the Paley inequality.

Theorem 1.2 (Paley-type inequality). *Suppose that ψ is a positive function on \mathbb{R}_+ satisfying the condition*

$$M_\psi := \sup_{t>0} t \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) > t}} |c(\lambda)|^{-2} d\lambda < \infty. \quad (3)$$

Then for $f \in L^p(\mathbb{R}_+, A_{\alpha, \beta}(t) dt)$, $1 < p \leq 2$, we have

$$\left(\int_0^\infty |\widehat{f}(\lambda)|^p \psi(\lambda)^{2-p} |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p}} \lesssim M_\psi^{\frac{2-p}{p}} \|f\|_{L^p(\mathbb{R}_+, A_{\alpha, \beta}(t) dt)}. \quad (4)$$

By interpolating the Hausdorff-Young inequality and Paley-type inequality we get the following Hausdorff-Young-Paley inequality for the Jacobi transform.

Theorem 1.3 (Hausdorff-Young-Paley inequality). *Let $1 < p \leq 2$, and let $1 < p \leq b \leq p' \leq \infty$, where $p' = \frac{p}{p-1}$. If $\psi(\lambda)$ is a positive function on \mathbb{R}_+ such that*

$$M_\psi := \sup_{t>0} t \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) > t}} |c(\lambda)|^{-2} d\lambda \quad (5)$$

is finite, then for every $f \in L^p(\mathbb{R}_+, A_{\alpha, \beta}(t) dt)$ we have

$$\left(\int_{\mathbb{R}_+} \left(|\widehat{f}(\lambda)| \psi(\lambda)^{\frac{1}{b} - \frac{1}{p'}} \right)^b |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{b}} \lesssim M_\psi^{\frac{1}{b} - \frac{1}{p'}} \|f\|_{L^p(\mathbb{R}_+, A_{\alpha, \beta}(t) dt)}. \quad (6)$$

Finally we establish the following L^p - L^q boundedness result for multipliers of Jacobi transform.

Theorem 1.4. *Let $1 < p \leq 2 \leq q < \infty$. Suppose that T is a Jacobi-Fourier multiplier with symbol h , that is,*

$$\widehat{Tf}(\lambda) = h(\lambda)\widehat{f}(\lambda), \quad \lambda \in \mathbb{R}_+,$$

where h is an bounded measurable even function on \mathbb{R} . Then we have

$$\|T\|_{L^p(\mathbb{R}_+, A_{\alpha, \beta}(t)dt) \rightarrow L^q(\mathbb{R}_+, A_{\alpha, \beta}(t)dt)} \lesssim \sup_{s>0} s \left[\int_{\{\lambda \in \mathbb{R}_+ : |h(\lambda)| > s\}} |c(\lambda)|^{-2} d\lambda \right]^{\frac{1}{p} - \frac{1}{q}}.$$

Now, we apply Theorem 1.4 to prove the L^p - L^q boundedness of spectral multipliers for Jacobi Laplacian $L := -\mathcal{L}_{\alpha, \beta} = -\frac{d^2}{dt^2} - ((2\alpha + 1) \coth t (2\beta + 1) \tanh t) \frac{d}{dt}$. Set $\rho = \alpha + \beta + 1$.

Theorem 1.5. *Let $1 < p \leq 2 \leq q < \infty$ and let φ be a monotonically decreasing continuous function on $[\rho^2, \infty)$ such that $\lim_{u \rightarrow \infty} \varphi(u) = 0$. Then we have*

$$\|\varphi(L)\|_{\text{op}} \lesssim \sup_{u > \rho^2} \varphi(u) \begin{cases} (u - \rho^2)^{(\frac{1}{p} - \frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} \leq 1, \\ (u - \rho^2)^{(\alpha+1)(\frac{1}{p} - \frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} > 1, \end{cases}$$

where $\|\cdot\|_{\text{op}}$ denotes the operator norm from $L^p(\mathbb{R}_+, A_{\alpha, \beta}(t)dt)$ to $L^q(\mathbb{R}_+, A_{\alpha, \beta}(t)dt)$.

2. ESSENTIALS ABOUT HARMONIC NA GROUPS

For basics of harmonic NA groups and Fourier analysis on them, one can refer to papers [22, 23, 24, 25, 6, 9, 17, 44, 36, 35, 7, 8]. However, we recall necessary definitions, notation and terminology that we shall use in this paper.

Let \mathfrak{n} be a two-step nilpotent Lie algebra, equipped with an inner product $\langle \cdot, \cdot \rangle$. Denote by \mathfrak{z} the center of \mathfrak{n} and by \mathfrak{v} the orthogonal complement of \mathfrak{z} in \mathfrak{n} with respect to the inner product of \mathfrak{n} . We assume that dimensions of \mathfrak{v} and \mathfrak{z} are m and l respectively as real vector spaces. The Lie algebra \mathfrak{n} is H -type algebra if for every $Z \in \mathfrak{z}$, the map $J_Z : \mathfrak{v} \rightarrow \mathfrak{v}$ defined by

$$\langle J_Z X, Y \rangle = \langle Z, [X, Y] \rangle, \quad X, Y \in \mathfrak{v}, Z \in \mathfrak{z},$$

satisfies the condition $J_Z^2 = -\|Z\|^2 I_{\mathfrak{v}}$, where $I_{\mathfrak{v}}$ is the identity operator on \mathfrak{v} . In [35], Kaplan proved that for $Z \in \mathfrak{z}$ with $\|Z\| = 1$ one has $J_Z^2 = -I_{\mathfrak{v}}$; that is, J_Z induced a complex structure on \mathfrak{v} and hence $m = \dim(\mathfrak{v})$ is always even. A connected and simply connected Lie group N is called H -type if its Lie algebra is of H -type. The exponential map is a diffeomorphism as N is nilpotent, we can parametrize the element of $N = \exp \mathfrak{n}$ by (X, Z) , for $X \in \mathfrak{v}$ and $Z \in \mathfrak{z}$. The multiplication on N follows from the Campbell-Baker-Hausdorff formula given by

$$(X, Z)(X', Z') = (X + X', Z + Z' + \frac{1}{2}[X, X']).$$

The group $A = \mathbb{R}_+^*$ acts on N by nonisotropic dilations as follows: $(X, Y) \mapsto (a^{\frac{1}{2}}X, aZ)$. Let $S = N \ltimes A$ be the semidirect product of N with A under the aforementioned action. The group multiplication on S is defined by

$$(X, Z, a)(X', Z', a') = (X + a^{\frac{1}{2}}X', Z + aZ' + \frac{1}{2}a^{\frac{1}{2}}[X, X'], aa').$$

Then S is a solvable (connected and simply connected) Lie group with Lie algebra $\mathfrak{s} = \mathfrak{z} \oplus \mathfrak{v} \oplus \mathbb{R}$ and Lie bracket

$$[(X, Z, \ell), (X', Z', \ell')] = (\frac{1}{2}\ell X' - \frac{1}{2}\ell' X, \ell Z' - \ell' Z + [X, X'], 0).$$

The group S is equipped with the left-invariant Riemannian metric induced by

$$\langle (X, Z, \ell), (X', Z', \ell') \rangle = \langle X, X' \rangle + \langle Z, Z' \rangle + \ell \ell'$$

on \mathfrak{s} . The homogeneous dimension of N is equal to $\frac{m}{2} + l$ and will be denoted by Q . At times, we also use symbol ρ for $\frac{Q}{2}$. Hence $\dim(\mathfrak{s}) = m + l + 1$, denoted by d . The associated left Haar measure dx on S is given by $a^{-Q-1}dXdZda$, where dX , dZ and da are the Lebesgue measures on \mathfrak{v} , \mathfrak{z} and \mathbb{R}_+^* respectively. The element of A will be identified with $a_t = e^t$, $t \in \mathbb{R}$. The group S can be realized as the unit ball $B(\mathfrak{s})$ in \mathfrak{s} using the Cayley transform $C : S \rightarrow B(\mathfrak{s})$ (see [6]).

To define (Helgason) Fourier transform on S we need to introduce the notion of the Poisson kernel ([9]). The Poisson Kernel $\mathcal{P} : S \times N \rightarrow \mathbb{R}$ is defined by $\mathcal{P}(na_t, n') = P_{a_t}(n'^{-1}n)$, where

$$P_{a_t}(n) = P_{a_t}(X, Z) = Ca_t^Q \left(\left(a_t + \frac{|X|^2}{4} \right)^2 + |Z|^2 \right)^{-Q}, \quad n = (X, Z) \in N.$$

The value of C is suitably adjusted so that $\int_N P_a(n)dn = 1$ and $P_1(n) \leq 1$. The Poisson kernel satisfies several useful properties (see [36, 44, 9]), we list here a few of them. For $\lambda \in \mathbb{C}$, the complex power of the Poisson kernel is defined as

$$\mathcal{P}_\lambda(x, n) = \mathcal{P}(x, n)^{\frac{1}{2} - \frac{i\lambda}{Q}}.$$

It is known ([44, 9]) that for each fixed $x \in S$, $\mathcal{P}_\lambda(x, \cdot) \in L^p(N)$ for $1 \leq p \leq \infty$ if $\lambda = i\gamma_p\rho$, where $\gamma_p = \frac{2}{p} - 1$. A very special feature of $\mathcal{P}_\lambda(x, n)$ is that it is constant on the hypersurfaces $H_{n, a_t} = \{n\sigma(a_t n') : n' \in N\}$. Here σ is the geodesic inversion on S , that is an involutive, measure-preserving, diffeomorphism which can be explicitly given by [17]:

$$\sigma(X, Z, a_t) = \left(\left(e^t + \frac{|V|^2}{4} \right)^2 + |Z|^2 \right)^{-1} \left(\left(- \left(e^t + \frac{|X|^2}{4} \right) + J_Z \right) X, -Z, a_t \right).$$

Let Δ_S be the Laplace-Beltrami operator on S . Then for every fixed $n \in N$, $\mathcal{P}_\lambda(x, n)$ is an eigenfunction of Δ_S with eigenvalue $-(\lambda^2 + \frac{Q^2}{4})$ (see [9]). For a measurable function f on S , the (Helgason) Fourier transform is defined as

$$\tilde{f}(\lambda, n) = \int_S f(x) \mathcal{P}_\lambda(x, n) dx$$

whenever the integral converges. For $f \in C_c^\infty(S)$, the following inversion formula holds ([9, Theorem 4.4]):

$$f(x) = C \int_{\mathbb{R}} \int_N \tilde{f}(\lambda, n) \mathcal{P}_{-\lambda}(x, n) |c(\lambda)|^{-2} d\lambda dn.$$

The authors also proved that the (Helgason) Fourier transform extends to an isometry from $L^2(S)$ onto the space $L^2(\mathbb{R}_+ \times N, |c(\lambda)|^{-2} d\lambda dn)$. In fact they have the precise value of constants, we refer the reader to [9]. The following estimates for the function $|c(\lambda)|$ hold:

$$|c(\lambda)|^{-2} = \begin{cases} |s|^2 & |s| \leq 1 \\ |s|^{d-1} & |s| > 1 \end{cases}$$

for all $\lambda \in \mathbb{R}$ (e. g. see [47, Theorem 1.14]). In [44, Theorem 4.6], the authors proved the following version of the Hausdorff-Young inequality: For $1 \leq p \leq 2$ we have

$$\left(\int_{\mathbb{R}} \int_N |\tilde{f}(\lambda + i\gamma_{p'} \rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p'}} \leq C_p \|f\|_p. \quad (7)$$

Let $\tilde{\mu}$ be the metric induced by the canonical left invariant Riemannian structure on S and let e be the identity element of S . A function f on S is called *radial* if for all $x, y \in S$, $f(x) = f(y)$ if $\tilde{\mu}(x, e) = \tilde{\mu}(y, e)$. Note that radial functions on S can be identified with the functions $f = f(r)$ of the geodesic distance $r = \tilde{\mu}(x, e) \in [0, \infty)$ to the identity. It is clear that $\tilde{\mu}(a_t, e) = |t|$ for $t \in \mathbb{R}$. At times, for any radial function f we use the notation $f(a_t) = f(t)$. For any function space $\mathcal{F}(S)$ on S , the subspace of radial functions will be denoted by $\mathcal{F}(S)^\#$. The elementary spherical function $\phi_\lambda(x)$ is defined by

$$\phi_\lambda(x) := \int_N \mathcal{P}_\lambda(x, n) \mathcal{P}_{-\lambda}(x, n) dn.$$

It follows ([6, 9]) that ϕ_λ is a radial eigenfunction of the Laplace-Beltrami operator Δ_S of S with eigenvalue $-(\lambda^2 + \frac{Q^2}{4})$ such that $\phi_\lambda(x) = \phi_{-\lambda}(x)$, $\phi_\lambda(x) = \phi_\lambda(x^{-1})$ and $\phi_\lambda(e) = 1$. In [6], the authors showed that the radial part (in geodesic polar coordinates) of the Laplace-Beltrami operator Δ_S given by

$$\text{rad } \Delta_S = \frac{\partial^2}{\partial t^2} + \left\{ \frac{m+l}{2} \coth \frac{t}{2} + \frac{k}{2} \tanh \frac{t}{2} \right\} \frac{\partial}{\partial t},$$

is (by substituting $r = \frac{t}{2}$) equal to $\frac{1}{4} \mathcal{L}_{\alpha, \beta}$ with indices $\alpha = \frac{m+l+1}{2}$ and $\beta = \frac{l-1}{2}$, where $\mathcal{L}_{\alpha, \beta}$ is the Jacobi operator studied by Koornwinder [37] in detail. It is worth noting that we are in the ideal situation of the Jacobi analysis with $\alpha > \beta > \frac{-1}{2}$. In fact, the Jacobi functions $\phi_\lambda^{\alpha, \beta}$ and elementary spherical functions ϕ_λ are related as ([6]): $\phi_\lambda(t) = \phi_{2\lambda}^{\alpha, \beta}(\frac{t}{2})$. As consequence of this relation, the following estimates for the elementary spherical functions hold true (see [42]).

Lemma 2.1. *The following inequalities are valid for spherical functions $\phi_\lambda(t)$ ($t, \lambda \in \mathbb{R}_+$) :*

- $|\phi_\lambda(t)| \leq 1$.
- $|1 - \phi_\lambda(t)| \leq \frac{t^2}{2} (4\lambda^2 + \frac{Q^2}{4})$.
- *There exists a constant $c > 0$, depending only on λ , such that $|1 - \phi_\lambda(t)| \geq c$ for $\lambda t \geq 1$.*

Now, we define the spherical (or Harish Chandra) transform of an integrable radial function f on S by

$$\mathcal{H}(f)(\lambda) := \int_S f(x) \phi_\lambda(x) dx.$$

For $f \in C_c^\infty(S)^\#$, the following inversion formula holds:

$$\mathcal{H}^{-1}(f)(x) := c_S \int_0^\infty \mathcal{H}f(\lambda) \phi_\lambda(x) |c(\lambda)|^{-2} d\lambda,$$

where c_S depends only on m and l , and $c(\lambda)$ is the Harish-Chandra function. Moreover, the following Plancherel formula holds:

$$\int_S |f(x)|^2 dx = c_S \int_{\mathbb{R}_+} |\mathcal{H}f(\lambda)|^2 |c(\lambda)|^{-2} d\lambda.$$

The spherical Fourier transform extends to an isometry from $L^2(S)^\#$ to $L^2(\mathbb{R}_+, c_S |c(\lambda)|^{-2} d\lambda)$. For details on the spherical analysis on harmonic NA group we refer to [6].

Let σ_t be the normalized surface measure of the geodesic sphere of radius t . Then σ_t is a nonnegative radial measure. The spherical mean operator M_t on a suitable function space on S is defined by $M_t f := f * \sigma_t$. It can be noted that $M_t f(x) = \mathcal{R}(f^x)(t)$, where f^x denotes the right translation of function f by x and \mathcal{R} is the radialization operator defined, for suitable function f , by

$$\mathcal{R}f(x) = \int_{S_\nu} f(y) d\sigma_\nu(y),$$

where $\nu = r(x) = \mu(C(x), 0)$, here C is the Cayley transform, and $d\sigma_\nu$ is the normalized surface measure induced by the left invariant Riemannian metric on the geodesic sphere $S_\nu = \{y \in S : \mu(y, e) = \nu\}$. It is easy to see that $\mathcal{R}f$ is a radial function and for any radial function f , $\mathcal{R}f = f$. Consequently, for a radial function f , $M_t f$ is the usual translation of f by t . In [36], the authors proved that M_t is a bounded linear operator on $L^p(S)$ and for a suitable function f on S , $\widetilde{M_t f}(\lambda, n) = \widetilde{f}(\lambda, n) \phi_\lambda(t)$ whenever both make sense. Also, $M_t f$ converges to f as $t \rightarrow 0$. The following result was proved by Kumar et al. in [36, Theorem 4.7 (a)] which was initially conjectured by Bray and Pinsky [13, Conjecture 16].

Theorem 2.2. *Let $1 < p \leq 2$ and $p \leq q \leq p'$. Then for $f \in L^p(S)$*

$$\left(\int_{\mathbb{R}} \min\{1, (\lambda t)^{2p'}\} \left(\int_N |\widetilde{f}(\lambda + i\gamma_q \rho, n)|^q dn \right)^{p'/q} |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p'}} \leq C_{p,q} \|M_t f - f\|_p.$$

For small $|\lambda|$, with $q = p'$, we have

$$\int_{|\lambda| < \frac{1}{t}} |\lambda|^{2p'} \int_N |\widetilde{f}(\lambda + i\gamma_{p'} \rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda \leq C_p^{p'} \left(\frac{\|M_t f - f\|_p}{t^2} \right)^{p'}. \quad (8)$$

For notational convenience, we take integral over an empty set to be zero.

3. TITCHMARSH THEOREMS FOR (HELGASON) FOURIER TRANSFORM OF HÖLDER-LIPSCHITZ FUNCTIONS ON HARMONIC NA GROUPS

This section is devoted to the investigation of Titchmarsh theorems on Hölder-Lipschitz functions on harmonic NA groups. In this plan of action we first study the growth property of Fourier transform of L^2 -Lipschitz function. Then we present some applications of it in the context of multiplier theorems and embedding theorems of Lipschitz-Sobolev spaces. After that, we prove an analogue of the second Titchmarsh theorem improving the range of Hausdorff-Young inequality for Hölder-Lipschitz functions on harmonic NA groups.

3.1. Growth properties of Fourier transform of Hölder-Lipschitz functions on harmonic NA groups. The following theorem generalizes one result found in [42] for rank one noncompact symmetric spaces. Recently, this result was extended by Bray [12] in higher dimensional Euclidean spaces, which dates back to Titchmarsh [46, Theorem 17] in the one dimensional setting (see Theorem B).

Theorem 3.1. *Let $0 < \alpha \leq 1$ and $f \in L^2(S)$. Then $f \in \text{Lip}_S(\alpha; 2)$ if and only if*

$$\int_{\lambda > \frac{1}{t}} \int_N |\tilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda = O(t^{2\alpha}) \quad \text{as } t \rightarrow 0.$$

Proof. Let us assume that $f \in \text{Lip}_S(\alpha; 2)$, i.e.,

$$\|M_t f - f\|_2 = O(t^\alpha) \quad \text{as } t \rightarrow 0. \quad (9)$$

By the Plancherel theorem and the fact that $\widetilde{M_t f}(\lambda, n) = \phi_\lambda(a_t) \tilde{f}(\lambda, n)$ we conclude that

$$\begin{aligned} \|M_t f - f\|_2^2 &= \|\widetilde{M_t f} - f\|_{L^2(\mathbb{R}_+ \times N, |c(\lambda)|^{-2} d\lambda dn)}^2 \\ &= \int_0^\infty \int_N |1 - \phi_\lambda(a_t)|^2 |\tilde{f}(\lambda, n)|^2 |c(\lambda)|^{-2} d\lambda dn \\ &= \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda, \end{aligned}$$

where $G(\lambda) = \int_N |\tilde{f}(\lambda, n)|^2 dn$. Now, by (9) it follows that

$$\int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda = O(t^{2\alpha}) \quad \text{as } t \rightarrow 0.$$

Now, if $\lambda \in [\frac{1}{t}, \frac{2}{t}]$ then $\lambda t \geq 1$ and by Lemma 2.1(iii) it follows that there exists a constant $c > 0$ such that $\frac{|1 - \phi_\lambda(a_t)|^2}{c^2} \geq 1$. Using this, we have

$$\begin{aligned} \int_{\frac{1}{t}}^{\frac{2}{t}} G(\lambda) |c(\lambda)|^{-2} d\lambda &\leq \frac{1}{c^2} \int_{\frac{1}{t}}^{\frac{2}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &\leq \frac{1}{c^2} \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda = O(t^{2\alpha}) \quad \text{as } t \rightarrow 0. \end{aligned}$$

By setting $r = \frac{1}{t}$, we may rewrite the above inequality as

$$\int_r^{2r} G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C' (r^{-2\alpha}).$$

Consequently, we have

$$\begin{aligned}
\int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda &= \left[\int_r^{2r} + \int_{2r}^{4r} + \int_{4r}^{8r} + \dots \right] G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&\leq C' (r^{-2\alpha} + (2r)^{-2\alpha} + (4r)^{-2\alpha} + \dots) \\
&\leq C' r^{-2\alpha} [1 + 2^{-2\alpha} + (2^{-2\alpha})^2 + (2^{-2\alpha})^3 + \dots] \\
&= C' (1 - 2^{-2\alpha})^{-1} r^{-2\alpha}.
\end{aligned}$$

Therefore, we have

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda = O(r^{-2\alpha}) \quad \text{as } r \rightarrow \infty.$$

Now we will prove the converse implication. Assume that

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^2 dn d\lambda = O(r^{-2\alpha}) \quad \text{as } r \rightarrow \infty. \quad (10)$$

As earlier we set $G(\lambda) = \int_N |\tilde{f}(\lambda, n)|^2 dn$. Then we get

$$\int_r^{2r} G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C r^{-2\alpha}.$$

As a consequence, it follows that

$$\int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda = \sum_{j=0}^\infty \int_{2^j r}^{2^{j+1} r} G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C \sum_{j=0}^\infty (2^{-2\alpha})^j r^{-2\alpha} \leq C r^{-2\alpha},$$

where C is a positive constant. Further, we have

$$\begin{aligned}
\|M_t f - f\|_{L^2(S)}^2 &= \|\widetilde{M_t f - f}\|_{L^2(\mathbb{R}_+ \times N, |c(\lambda)|^{-2} d\lambda dn)}^2 \\
&= \int_0^\infty \int_N |1 - \phi_\lambda(a_t)|^2 |\tilde{f}(\lambda, n)|^2 |c(\lambda)|^{-2} d\lambda dn \\
&= \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&= \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda + \int_{\frac{1}{t}}^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&= I_1 + I_2.
\end{aligned}$$

For second integral I_2 , by using $|\phi_\lambda(a_t)| \leq 1$ for $\lambda \in \mathbb{R}_+$ we have, with $r = \frac{1}{t}$,

$$I_2 = \int_{\frac{1}{t}}^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \leq 4 \int_{\frac{1}{t}}^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda = 4C r^{-2\alpha}$$

and so

$$I_2 = O(r^{-2\alpha}). \quad (11)$$

Next, to estimate I_1 , we will use the facts that $|\phi_\lambda(a_t)| \leq 1$ and $|1 - \phi_\lambda(a_t)| \leq t^2(\lambda^2 + \frac{Q^2}{4})$ for $\lambda, t \in \mathbb{R}_+$ (see Lemma 2.1), so that

$$I_1 = \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda = \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)| |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda$$

$$\begin{aligned}
&\leq \int_0^{\frac{1}{t}} (1 + |\phi_\lambda(a_t)|) |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&\leq 2 \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda \leq 2t^2 \int_0^{\frac{1}{t}} \left(\lambda^2 + \frac{Q^2}{4} \right) G(\lambda) |c(\lambda)|^{-2} d\lambda.
\end{aligned}$$

For a while, we put $F(r) = \int_r^\infty G(\lambda) d\mu(\lambda)$, where $d\mu(\lambda) = |c(\lambda)|^{-2} d\lambda$. Then we have, as $0 \leq \lambda \leq \frac{1}{t} = r$, that

$$I_1 \leq 2t^2 \int_0^{\frac{1}{t}} - \left(r^2 + \frac{Q^2}{4} \right) F'(r) dr \leq 2t^2 \int_0^{\frac{1}{t}} -r^2 F'(r) dr \quad (12)$$

$$= 2t^2 \left(\frac{-1}{t^2} F\left(\frac{1}{t}\right) + 2 \int_0^{\frac{1}{t}} r F(r) dr \right) = -2F\left(\frac{1}{t}\right) + 4t^2 \int_0^{\frac{1}{t}} r F(r) dr \quad (13)$$

$$\leq 4t^2 \int_0^{\frac{1}{t}} r F(r) dr. \quad (14)$$

Since $F(r) = \int_r^\infty G(\lambda) d\mu(\lambda) = O(r^{-2\alpha})$ we get

$$I_1 \leq Ct^2 \int_0^{\frac{1}{t}} r (r^{-2\alpha}) dr \leq Ct^{2\alpha},$$

where C is a positive constant. Hence, by combining I_1 and I_2 we get

$$\|M_t f - f\|_{L^2(S)}^2 = (t^{2\alpha}) \text{ as } t \rightarrow 0.$$

Hence, $f \in \text{Lip}_S(\alpha, 2)$. □

3.2. Applications. This subsection is devoted to the applications of the result presented in Subsection 3.1. Specifically, we formulate the following corollary for the regularity of Fourier multipliers of Hölder-Lipschitz spaces on harmonic NA groups. This corollary complements the other results available in the literature (e.g. see [6]) for boundedness of Fourier multipliers on harmonic NA groups.

Corollary 3.2. *Let $0 \leq \gamma < 1$ and let h be an even measurable function on \mathbb{R} such that*

$$|h(\lambda)| \leq C \langle \lambda \rangle^{-\gamma},$$

where $\langle \lambda \rangle = (\lambda^2 + \frac{Q^2}{2})^{\frac{1}{2}}$. Let A be the Fourier multiplier with symbol h , i.e., given by $\widetilde{A}f(\lambda, n) = h(\lambda) \widetilde{f}(\lambda, n)$ for all $\lambda \in \mathbb{R}_+$ and $n \in N$. Then

$$A : \text{Lip}_S(\alpha; 2) \rightarrow \text{Lip}_S(\alpha + \gamma; 2)$$

is bounded for all α such that $0 < \alpha < 1 - \gamma$.

Proof. Let $f \in \text{Lip}_S(\alpha; 2)$. Then by Theorem 3.1 we have

$$\begin{aligned}
\int_{\lambda > \frac{1}{t}} \int_N |\widetilde{A}f(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda &= \int_{\lambda > \frac{1}{t}} \int_N |h(\lambda) \widetilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda \\
&\leq C \int_{\lambda > \frac{1}{t}} \int_N \langle \lambda \rangle^{-2\gamma} |\widetilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda \\
&\leq Ct^{2\gamma} \int_{\lambda > \frac{1}{t}} \int_N |\widetilde{f}(\lambda, n)|^2 dn |c(\lambda)|^{-2} d\lambda
\end{aligned}$$

$$= O(t^{2(\alpha+\gamma)}) \text{ as } t \rightarrow 0.$$

Again by Theorem 3.1 this implies that $A : \text{Lip}_S(\alpha; 2) \rightarrow \text{Lip}_S(\alpha + \gamma; 2)$ is bounded for all $\alpha > 0$ such that $\alpha + \gamma < 1$. \square

As an example we consider the Lipschitz regularity for the Laplace-Beltrami operator $(-\Delta_S)^{\frac{-\gamma}{2}}$. First we observe that if $A := (-\Delta_S)^{\frac{-\gamma}{2}}$ with $0 \leq \gamma < 1$, by Corollary 3.2 we get

$$\|(-\Delta_S)^{\frac{-\gamma}{2}} f\|_{\text{Lip}_S(\alpha+\gamma; 2)} \leq C \|f\|_{\text{Lip}_S(\alpha; 2)}$$

for all α such that $0 < \alpha < 1 - \gamma$. Hence,

$$\|f\|_{\text{Lip}_S(\alpha+\gamma; 2)} \leq C \|(-\Delta_S)^{\frac{\gamma}{2}} f\|_{\text{Lip}_S(\alpha; 2)}. \quad (15)$$

Now, we can introduce the Sobolev-Lipschitz space $H^\gamma \text{Lip}_S(\alpha; 2)$ for every $0 \leq \gamma < 1$ and $0 < \alpha < 1 - \gamma$ by

$$H^\gamma \text{Lip}_S(\alpha; 2) := \left\{ f \in \mathcal{D}'(S) : (-\Delta_S)^{\frac{\gamma}{2}} f \in \text{Lip}_S(\alpha; 2) \right\}.$$

From (15) we have the following corollary:

Corollary 3.3. *For every $0 \leq \gamma < 1$ and $0 < \alpha < 1 - \gamma$ we have the continuous embedding*

$$H^\gamma \text{Lip}_S(\alpha; 2) \hookrightarrow \text{Lip}_S(\alpha + \gamma; 2).$$

3.3. An integrability result for Fourier transform of Hölder-Lipschitz functions on harmonic NA groups. The following result is a general integrability theorem for the function F on $[0, \infty)$ defined by

$$F(\lambda) := \left(\int_N |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn \right)^{\frac{1}{p'}}, \quad 1 < p \leq 2,$$

where $\gamma_{p'} = \frac{2}{p'} - 1 = -\gamma_p$. This theorem is an extension of a result of Bray ([12]) on \mathbb{R}^n to harmonic NA groups, in particular to rank one noncompact symmetric spaces. We prove this result by applying the similar technique as in [12]. In our case, an inequality (8) obtained as consequence of a result (Theorem 2.2) proved by Kumar et al. [36] plays a pivotal role.

Theorem 3.4. *Let S be a harmonic NA group of dimension d . Let $0 < \alpha \leq 1$, $1 < p \leq 2$, and let p' be such that $\frac{1}{p} + \frac{1}{p'} = 1$. Let $f \in \text{Lip}_S(\alpha; p)$. Then the function F defined by*

$$F(\lambda) := \left(\int_N |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn \right)^{\frac{1}{p'}}, \quad 1 < p \leq 2,$$

belongs to $L^\beta((0, \infty), |c(\lambda)|^{-2} d\lambda)$ provided that

$$\frac{dp}{dp + \alpha p - d} < \beta \leq p'. \quad (16)$$

Proof. For $f \in \text{Lip}_S(\alpha; p)$ we have that

$$\|M_t f - f\|_p = O(t^\alpha)$$

as $t \rightarrow 0$. Therefore, by setting $\Lambda = \frac{1}{t}$ from inequality (8) we get that

$$\int_1^\Lambda \lambda^{2p'} F(\lambda)^{p'} |c(\lambda)|^{-2} d\lambda \leq C_p^{p'} \left(\frac{\|M_t f - f\|_p}{t^2} \right)^{p'} \leq C \Lambda^{(2-\alpha)p'},$$

since $\|M_t f - f\|_p = O(t^\alpha)$ we get, by putting $t = \frac{1}{\Lambda}$, that

$$\left(\frac{\|M_t f - f\|_p}{t^2} \right)^{p'} = (O(\Lambda^{-\alpha}) O(\Lambda^2))^{p'} = O(\Lambda^{(2-\alpha)p'}).$$

Take $\beta < p'$ and let

$$G(\Lambda) = \int_1^\Lambda \lambda^{2\beta} F(\lambda)^\beta |c(\lambda)|^{-2} d\lambda.$$

By Hölder's inequality we conclude that by using the estimate $|c(\lambda)|^{-2} \lesssim |\lambda|^{d-1}$ as $|\lambda| > 1$,

$$\begin{aligned} G(\Lambda) &= \int_1^\Lambda [\lambda^{2\beta} F(\lambda)]^\beta |c(\lambda)|^{-2(\frac{\beta}{p'})} |c(\lambda)|^{-2(1-\frac{\beta}{p'})} d\lambda \\ &\leq C \left(\int_1^\Lambda \lambda^{2p'} F(\lambda)^{p'} |c(\lambda)|^{-2} d\lambda \right)^{\frac{\beta}{p'}} \left(\int_1^\Lambda |c(\lambda)|^{-2} d\lambda \right)^{1-\frac{\beta}{p'}} \\ &\leq C \left(\int_1^\Lambda \lambda^{2p'} F(\lambda)^{p'} |c(\lambda)|^{-2} d\lambda \right)^{\frac{\beta}{p'}} \left(\int_1^\Lambda |\lambda|^{d-1} d\lambda \right)^{1-\frac{\beta}{p'}} \\ &\leq C \Lambda^{(2-\alpha)\beta} \Lambda^{d(1-\frac{\beta}{p'})} = O(\Lambda^{2\beta+d-\alpha\beta-\frac{d\beta}{p'}}), \end{aligned} \tag{17}$$

where C is a generic constant which can be different at each step.

Now, integration by parts yield the following,

$$\begin{aligned} \int_1^\Lambda F(\lambda)^\beta |c(\lambda)|^{-2} d\lambda &= \int_1^\Lambda \lambda^{-2\beta} G'(\lambda) d\lambda \\ &= \Lambda^{-2\beta} G(\Lambda) + 2\beta \int_1^\Lambda \lambda^{-2\beta-1} G(\lambda) d\lambda = I_1 + I_2. \end{aligned}$$

The first term I_1 is bounded under the condition on β by using the above estimate of G . Indeed

$$I_1 = \Lambda^{-2\beta} G(\Lambda) \leq \Lambda^{-2\beta} C \Lambda^{(2-\alpha)\beta} \Lambda^{d(1-\frac{\beta}{p'})} = C \Lambda^{-\alpha\beta+d-\frac{d\beta}{p'}} = O(1)$$

since $-\alpha\beta + d - \frac{d\beta}{p'} = \alpha\beta + d - d\beta(1 - \frac{1}{p}) = \frac{dp - \beta(dp + \alpha p - d)}{p}$ which is negative by using the condition of β , i.e., $\beta > \frac{dp}{dp + \alpha p - d}$.

Similarly for I_2 , we get, using the estimate of $G(\lambda)$,

$$I_2 = \int_1^\Lambda \lambda^{-2\beta-1} G(\lambda) d\lambda \leq C \int_1^\Lambda \lambda^{-\alpha\beta+d-\frac{d\beta}{p'}-1} d\lambda = O(1) \tag{18}$$

because $-\alpha\beta + d - \frac{d\beta}{p'} - 1 = \frac{dp - \beta(dp + \alpha p - d) - p}{p} < 0$ by using the fact that $\beta > \frac{dp}{dp + \alpha p - d}$, which is equivalent to $dp - \beta(dp + \alpha p - d) < 0$. Therefore,

$$\int_1^\Lambda F(\lambda)^\beta |c(\lambda)|^{-2} d\lambda = O(1)$$

provided that $d - \alpha\beta - d\beta + \frac{d\beta}{p} < 0$ and hence $F \in L^\beta((0, \infty), |c(\lambda)|^{-2}d\lambda)$, since the above $O(1)$ are independent of Λ . The proof is complete as conditions on β are equivalent to (16). \square

4. HELGASON TRANSFORM OF DINI-LIPSCHITZ FUNCTIONS ON HARMONIC NA GROUPS

In this section, we discuss the result related with the growth and integrability of Fourier transform of Dini-Lipschitz functions on harmonic NA groups. This kind of results were first studied by Younis [49] in the context of one dimensional Euclidean space. These results were extended by many researchers in different settings [50, 51, 30, 27]. In particular, for compact homogeneous manifolds it was shown in [27]. In this direction, we state our first result below.

Theorem 4.1. *Let S be a harmonic NA group of dimension d and let $\alpha, \beta > 0$. Then the conditions*

$$\|M_t f - f\|_{L^2(S)} = O\left(\frac{t^\alpha}{(\log \frac{1}{t})^\beta}\right) \text{ as } t \rightarrow 0 \quad (19)$$

and

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^2 dn d\lambda = O\left(\frac{r^{-2\alpha-d+1}}{(\log r)^{2\beta}}\right) \text{ as } r \rightarrow \infty \quad (20)$$

are equivalent.

Proof. Let us assume that (19) holds. By the Plancherel theorem and the fact that

$$\widetilde{M_t f}(\lambda, n) = \phi_\lambda(a_t) \tilde{f}(\lambda, n)$$

we get

$$\begin{aligned} \|M_t f - f\|_2^2 &= \|\widetilde{M_t f - f}\|_{L^2(\mathbb{R}_+ \times N, |c(\lambda)|^{-2} d\lambda dn)}^2 \\ &= \int_0^\infty \int_N |1 - \phi_\lambda(a_t)|^2 |\tilde{f}(\lambda, n)|^2 |c(\lambda)|^{-2} d\lambda dn \\ &= \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda, \end{aligned}$$

where $G(\lambda) = \int_N |\tilde{f}(\lambda, n)|^2 dn$. Now, by (19) it follows that

$$\int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda = O\left(\frac{t^{2\alpha}}{(\log \frac{1}{t})^{2\beta}}\right) \text{ as } t \rightarrow 0.$$

Now, if $\lambda \in [\frac{1}{t}, \frac{2}{t}]$ then $|\lambda t| \geq 1$ and by Lemma 2.1(iii) it follows that there exists a constant $c > 0$ such that

$$\frac{|1 - \phi_\lambda(a_t)|^2}{c^2} \geq 1.$$

Using this, we have

$$\begin{aligned} \int_{\frac{1}{t}}^{\frac{2}{t}} G(\lambda) |c(\lambda)|^{-2} d\lambda &\leq \frac{1}{c^2} \int_{\frac{1}{t}}^{\frac{2}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &\leq \frac{1}{c^2} \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &= O\left(\frac{t^{2\alpha}}{(\log \frac{1}{t})^{2\beta}}\right) \text{ as } t \rightarrow 0. \end{aligned}$$

By setting $r = \frac{1}{t}$, we may write the above inequality as

$$\int_r^{2r} G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C' \left(\frac{r^{-2\alpha}}{(\log r)^{2\beta}} \right),$$

where $C' > 0$ is a generic constant which can be different for different inequalities below. Consequently, we have

$$\begin{aligned} \int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda &= \left[\int_r^{2r} + \int_{2r}^{4r} + \int_{4r}^{8r} + \dots \right] G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &\leq C' \left(\frac{r^{-2\alpha}}{(\log r)^{2\beta}} + \frac{(2r)^{-2\alpha}}{(\log 2r)^{2\beta}} + \frac{(4r)^{-2\alpha}}{(\log 4r)^{2\beta}} + \dots \right) \\ &\leq C' \frac{r^{-2\alpha}}{(\log r)^{2\beta}} [1 + 2^{-2\alpha} + (2^{-2\alpha})^2 + (2^{-2\alpha})^3 + \dots] \\ &= C' (1 - 2^{-2\alpha})^{-1} \frac{r^{-2\alpha}}{(\log r)^{2\beta}}. \end{aligned}$$

By using the estimate $|c(\lambda)|^{-2} \geq c_1 \lambda^{d-1}$ for $\lambda \rightarrow \infty$ we get

$$\int_r^\infty G(\lambda) d\lambda = c_1 r^{-d+1} \int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda = c_1 C' (1 - 2^{-2\alpha})^{-1} r^{-d+1} \frac{r^{-2\alpha}}{(\log r)^{2\beta}}$$

and therefore, we have

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^2 dn d\lambda = O \left(\frac{r^{-2\alpha-d+1}}{(\log r)^{2\beta}} \right) \text{ as } r \rightarrow \infty. \quad (21)$$

Now we will prove the converse implication. Assume that (20) holds, i.e.,

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^2 dn d\lambda = O \left(\frac{r^{-2\alpha-d+1}}{(\log r)^{2\beta}} \right) \text{ as } r \rightarrow \infty. \quad (22)$$

As earlier we set $G(\lambda) = \int_N |\tilde{f}(\lambda, n)|^2 dn$. Then by using the estimate $|c(\lambda)|^{-2} \leq c_2 |\lambda|^{d-1}$ we get

$$\begin{aligned} \int_r^{2r} G(\lambda) |c(\lambda)|^{-2} d\lambda &\leq c_2 \int_r^{2r} G(\lambda) \lambda^{d-1} d\lambda \leq c_2 (2r)^{d-1} \int_r^{2r} G(\lambda) d\lambda \\ &\leq C r^{d-1} \int_r^\infty G(\lambda) d\lambda \leq C \frac{r^{-2\alpha}}{(\log r)^{2\beta}}. \end{aligned}$$

As a consequence, it follows that

$$\begin{aligned} \int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda &= \sum_{j=0}^\infty \int_{2^j r}^{2^{j+1} r} G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &\leq C \sum_{j=0}^\infty \frac{(2^{-2\alpha})^j r^{-2\alpha}}{(\log r)^{2\beta}} \leq C \frac{r^{-2\alpha}}{(\log r)^{2\beta}}, \end{aligned}$$

where C is a positive constant. Further, we have

$$\begin{aligned} \|M_t f - f\|_{L^2(S)}^2 &= \|\widetilde{M_t f - f}\|_{L^2(\mathbb{R}_+ \times N, |c(\lambda)|^{-2} d\lambda dn)}^2 \\ &= \int_0^\infty \int_N |1 - \phi_\lambda(a_t)|^2 |\tilde{f}(\lambda, n)|^2 |c(\lambda)|^{-2} d\lambda dn \\ &= \int_0^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \end{aligned}$$

$$\begin{aligned}
&= \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda + \int_{\frac{1}{t}}^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&= I_1 + I_2.
\end{aligned}$$

For the second integral I_2 , by using $|\phi_\lambda(a_t)| \leq 1$ for $\lambda \in \mathbb{R}_+$, we have

$$\begin{aligned}
I_2 &= \int_{\frac{1}{t}}^\infty |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&\leq 4 \int_{\frac{1}{t}}^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda = 4C \frac{r^{-2\alpha}}{(\log r)^{2\beta}}
\end{aligned}$$

and so

$$I_2 = O\left(\frac{r^{-2\alpha}}{(\log r)^{2\beta}}\right). \quad (23)$$

Next, to estimate I_1 , we will use the facts that $|1 - \phi_\lambda(t)| \leq 1 + |\phi_\lambda(t)| \leq 2$ and $|1 - \phi_\lambda(t)| \leq t^2(\lambda^2 + \frac{Q^2}{4})$ for $\lambda, t \in \mathbb{R}_+$ (see Lemma 2.1), to obtain

$$\begin{aligned}
I_1 &= \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)|^2 G(\lambda) |c(\lambda)|^{-2} d\lambda = \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)| |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&\leq \int_0^{\frac{1}{t}} (1 + |\phi_\lambda(a_t)|) |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda \\
&\leq 2 \int_0^{\frac{1}{t}} |1 - \phi_\lambda(a_t)| G(\lambda) |c(\lambda)|^{-2} d\lambda \leq 2t^2 \int_0^{\frac{1}{t}} (\lambda^2 + \frac{Q^2}{4}) G(\lambda) |c(\lambda)|^{-2} d\lambda.
\end{aligned}$$

For a while, we put $F(r) = \int_r^\infty G(\lambda) d\mu(\lambda)$, where $d\mu(\lambda) = |c(\lambda)|^{-2} d\lambda$. Then we have

$$I_1 = 2t^2 \int_0^{\frac{1}{t}} -(r^2 + \frac{Q^2}{4}) F'(r) dr \leq 2t^2 \int_0^{\frac{1}{t}} -r^2 F'(r) dr \quad (24)$$

$$= 2t^2 \left(\frac{-1}{t^2} F\left(\frac{1}{t}\right) + 2 \int_0^{\frac{1}{t}} r F(r) dr \right) = -F\left(\frac{1}{t}\right) + 4t^2 \int_0^{\frac{1}{t}} r F(r) dr \quad (25)$$

$$\leq 4t^2 \int_0^{\frac{1}{t}} r F(r) dr. \quad (26)$$

Since $F(r) = \int_r^\infty G(\lambda) d\mu(\lambda) = O\left(\frac{r^{-2\alpha}}{(\log r)^{2\beta}}\right)$ we get

$$I_2 \leq Ct^2 \int_0^{\frac{1}{t}} r \left(\frac{r^{-2\alpha}}{(\log r)^{2\beta}} \right) \leq C \frac{t^{2\alpha}}{(\log \frac{1}{t})^{2\beta}},$$

where C is a positive constant. Hence, by combining I_1 and I_2 we get

$$\|M_t f - f\|_{L^2(S)}^2 = \left(\frac{t^{2\alpha}}{(\log \frac{1}{t})^{2\beta}} \right) \text{ as } t \rightarrow 0$$

proving (19). \square

This following Theorem can be proved in a similar way as Theorem 4.1 by using the following version of the Hausdorff-Young inequality [44, Theorem 4.6]

$$\left(\int_{\mathbb{R}} \int_N |\tilde{f}(\lambda + \gamma_{p'} \rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p'}} \leq C_p \|f\|_p \quad (27)$$

and the fact that $\widetilde{M_t f}(z, n) = \tilde{f}(z, n) \phi_z(a_t)$ for $t \in \mathbb{R}_+$ and $z \in S_p$.

Theorem 4.2. *Let S be a harmonic NA group of dimension d and let $\alpha, \beta > 0$. Let $f \in L^p(S)$, $1 < p \leq 2$ with $\frac{1}{p} + \frac{1}{p'} = 1$, be such that*

$$\|M_t f - f\|_{L^p(S)} = O\left(\frac{t^\alpha}{(\log \frac{1}{t})^\beta}\right) \text{ as } t \rightarrow 0. \quad (28)$$

Then

$$\int_r^\infty \int_N |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda = O(r^{-p'\alpha}(\log r)^{-p'\beta}) \text{ as } r \rightarrow \infty. \quad (29)$$

Proof. Suppose that (28) holds. If $\lambda \in [\frac{1}{t}, \frac{2}{t}]$, $(\lambda, t \in \mathbb{R}_+)$ then $\lambda t \geq 1$ and so from Lemma 2.1 we have

$$\frac{1}{c^{p'}} |1 - \phi_\lambda(a_t)|^{p'} \geq 1.$$

Set $G(\lambda) = \int_N |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn$. Now by using (27) we get

$$\begin{aligned} \int_{\frac{1}{t}}^{\frac{2}{t}} G(\lambda) |c(\lambda)|^{-2} d\lambda &\leq \frac{1}{c^{p'}} \int_{\frac{1}{t}}^{\frac{2}{t}} |1 - \phi_\lambda(a_t)|^{p'} G(\lambda) |c(\lambda)|^{-2} d\lambda \\ &\leq \frac{1}{c^{p'}} \int_{\frac{1}{t}}^{\frac{2}{t}} \int_N |1 - \phi_\lambda(a_t)|^{p'} |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda \\ &\leq \frac{1}{c^{p'}} \int_{\mathbb{R}} \int_N |1 - \phi_\lambda(a_t)|^{p'} |\tilde{f}(\lambda + i\gamma_{p'}\rho, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda \\ &\leq \frac{C_p}{c^{p'}} \|M_t f - f\|_{p'}^{p'} \\ &\leq C \|M_t f - f\|_{L^p(S)}^{p'} = O\left(\frac{t^{\alpha p'}}{(\log \frac{1}{t})^{\beta p'}}\right). \end{aligned}$$

Therefore, by putting $r = \frac{1}{t}$, we have

$$\int_r^{2r} G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C \left(\frac{r^{-\alpha p'}}{(\log r)^{\beta p'}} \right).$$

Now, as in Theorem 4.1 we get

$$\int_r^\infty G(\lambda) |c(\lambda)|^{-2} d\lambda \leq C_\alpha \frac{r^{-\alpha p'}}{(\log r)^{\beta p'}} \text{ as } r \rightarrow \infty,$$

with $C_\alpha = C(1 - 2^{p'\alpha})^{-1}$. Consequently, we have

$$\int_r^\infty \int_N |\tilde{f}(\lambda, n)|^{p'} dn |c(\lambda)|^{-2} d\lambda = O(r^{-p'\alpha}(\log r)^{-p'\beta}) \text{ as } r \rightarrow \infty,$$

proving (29). □

The following theorem is an analogue of Theorem 3.4 for Dini-Lipschitz functions on harmonic NA groups. This theorem was proved by Younis [49] on circle and extended in [27] to the setting of general compact homogeneous spaces. This theorem is new even in the case of rank one noncompact symmetric spaces.

Theorem 4.3. *Let S be a harmonic NA group of dimension d . Let $0 < \alpha \leq 1$, $\gamma > 0$, $1 < p \leq 2$, and let p' be such that $\frac{1}{p} + \frac{1}{p'} = 1$. Let $f \in L^p(S)$ be such that*

$$\|M_t f - f\|_{L^p(S)} = O\left(\frac{t^\alpha}{(\log \frac{1}{t})^\gamma}\right) \text{ as } t \rightarrow 0. \quad (30)$$

Then the function F defined by

$$F(\lambda) := \left(\int_N |\tilde{f}(\lambda + i\gamma_{p'} \rho, n)|^{p'} dn \right)^{\frac{1}{p'}},$$

belongs to $L^\beta((0, \infty), |c(\lambda)|^{-2} d\lambda)$ provided that

$$\frac{dp}{dp + \alpha p - d} < \beta \leq p'. \quad (31)$$

Proof. Arguing as in the proof of Theorem 3.4, under the assumption of Theorem 4.3 instead of (17) we get

$$G(\Lambda) = O(\Lambda^{2\beta + d - \alpha\beta - \frac{d\beta}{p'}} (\log \Lambda)^{-\beta\gamma}). \quad (32)$$

As a consequence, we have the following

$$\int_1^\Lambda F(\lambda)^\beta |c(\lambda)|^{-2} d\lambda = \Lambda^{-2\beta} G(\Lambda) + 2\beta \int_1^\Lambda \lambda^{-2\beta-1} G(\lambda) d\lambda = I_1 + I_2.$$

The term I_1 is bounded under the condition on β by using the above estimate of G . Indeed

$$I_1 = \Lambda^{-2\beta} G(\Lambda) = O(\Lambda^{-\alpha\beta + d - \frac{d\beta}{p'}} (\log \Lambda)^{-\beta\gamma}).$$

The right hand side of the last estimate is bounded as $\Lambda \rightarrow \infty$ as we must have $-\alpha\beta + d - \frac{d\beta}{p'} = \alpha\beta + d - d\beta(1 - \frac{1}{p}) = \frac{dp - \beta(dp + \alpha p - d)}{p} < 0$ which is always the case by using the condition of β , i.e., $\beta > \frac{dp}{dp + \alpha p - d}$ which also holds according to our assumption.

Similarly for I_2 , we get, using the estimate of $G(\lambda)$,

$$I_2 = \int_1^\Lambda \lambda^{-2\beta-1} G(\lambda) d\lambda = O(\Lambda^{-\alpha\beta + d - \frac{d\beta}{p'}} (\log \Lambda)^{-\beta\gamma}). \quad (33)$$

The right hand side of the above estimate is bounded because $-\alpha\beta + d - \frac{d\beta}{p'} = \frac{dp - \beta(dp + \alpha p - d)}{p} < 0$ by using the assumption that $\beta > \frac{dp}{dp + \alpha p - d}$, which is equivalent to $dp - \beta(dp + \alpha p - d) < 0$.

Therefore,

$$\int_1^\Lambda F(\lambda)^\beta |c(\lambda)|^{-2} d\lambda = O(1)$$

provided that $d - \alpha\beta - d\beta + \frac{d\beta}{p} < 0$. Hence $F \in L^\beta((0, \infty), |c(\lambda)|^{-2} d\lambda)$. The proof is complete as conditions on β are equivalent to (31). \square

5. $L^p - L^q$ BOUNDEDNESS OF MULTIPLIERS FOR JACOBI TRANSFORM FOR $1 < p \leq 2 \leq q < \infty$

The purpose of this section is to extend a classical result of Hörmander [34] on L^p - L^q boundedness of Fourier multipliers with the use of inequalities of Paley type [38, 39] and Hausdorff-Young-Paley type for the Jacobi transform [31, 37]. We have seen earlier in Section 2, initially noted in Anker et al. [6], that the spherical (radial) analysis on harmonic NA groups perfectly fits into the Jacobi analysis setting studied by Koornwinder and his coauthors [37, 31, 32]. So, instead of working in the setting of radial analysis on harmonic NA we choose to work in the broad setting of Jacobi analysis. In this section we prove the Hausdorff-Young-Paley inequality and apply it to prove $L^p - L^q$ boundedness of Fourier multipliers in this setting. We first briefly describe the harmonic analysis related to the Jacobi differential operators can be found in papers [37, 31, 32].

The hypergeometric function ${}_2F_1(a, b; c, z)$ is defined by

$${}_2F_1(a, b; c, z) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k k!} z^k, \quad |z| < 1,$$

where $(a)_k$ is the Pochhammer symbol of a given by $(a)_0 = 1$ and $(a)_k = a(a+1)\dots(a+k-1)$. The function $z \mapsto {}_2F_1(a, b; c, z)$ is the unique solution of the differential equation

$$z(1-z)u''(z) + (c - (a+b+1)z)u'(z) - abu(z) = 0,$$

which is regular at 0 and equals to 1 there. The Jacobi function with parameters (α, β) and $\lambda \in \mathbb{C}$ is defined by $\phi_{\lambda}^{\alpha, \beta}(t) = {}_2F_1\left(\frac{\alpha+\beta+1-i\lambda}{2}, \frac{\alpha+\beta+1+i\lambda}{2}; \alpha+1, -\sinh^2 t\right)$. It is well-known that the family $\{\phi_{\lambda}^{\alpha, \beta}\}_{\lambda \geq 0}$ forms an orthonormal system in \mathbb{R}_+ with respect to the weight $A_{\alpha, \beta}(t) = (2 \sinh t)^{2\alpha+1} (2 \cosh t)^{2\beta+1}$, $t > 0$ if $|\beta| < \alpha + 1$. In this section we assume that $\alpha \neq -1, -2, \dots$ and $\alpha \geq \beta > -\frac{1}{2}$. At times, for convenience we omit superscript and subscript (α, β) from $\phi_{\lambda}^{\alpha, \beta}$ and $A_{\alpha, \beta}(t)$ respectively and prefer to write them as ϕ_{λ} and $A(t)$ alone. It is worth noting that the following behavior for the weight $A_{\alpha, \beta}(t)$ holds true:

$$A_{\alpha, \beta}(t) \asymp \begin{cases} t^{2\alpha+1} & 0 < t < 1, \\ e^{2\rho t} & t \geq 1. \end{cases}$$

Also, we denote $\alpha + \beta + 1$ by ρ . The Jacobi Laplacian is given by

$$\mathcal{L}_{\alpha, \beta} = \frac{d^2}{dt^2} + ((2\alpha+1) \coth t + (2\beta+1) \tanh t) \frac{d}{dt},$$

where $\tanh t := \frac{e^t - e^{-t}}{e^t + e^{-t}}$ and $\coth t := \frac{1}{\tanh t}$. The Jacobi functions can be alternatively characterized as the unique solutions of

$$\mathcal{L}_{\alpha, \beta} \phi_{\lambda}^{\alpha, \beta} + (\lambda^2 + \rho^2) \phi_{\lambda}^{\alpha, \beta} = 0$$

on \mathbb{R}_+ such that $\phi_{\lambda}^{\alpha, \beta}(0) = 1$ and $(\phi_{\lambda}^{\alpha, \beta})'(0) = 0$. Therefore, the function $t \mapsto \phi_{\lambda}^{\alpha, \beta}(t)$ is analytic for $t \geq 0$. The function $\phi_{\lambda}^{\alpha, \beta}$ can be estimated as

$$|\phi_{\lambda}^{\alpha, \beta}(t)| \leq \begin{cases} 1 & \text{if } |\operatorname{Im}(\lambda)| \leq \rho \\ e^{(\operatorname{Im}(\lambda) - \rho)t} & \text{if } |\operatorname{Im}(\lambda)| > \rho \end{cases}$$

for $t \in \mathbb{R}_+$. It is also known that $\phi_\lambda^{\alpha, \beta}$ is bounded if and only if $|\operatorname{Im}(\lambda)| \leq \rho$.

Denote the space of all even smooth functions on \mathbb{R} with compact support by $C_c^\infty(\mathbb{R})^\#$ and the space consisting of restriction of functions in $C_c^\infty(\mathbb{R})^\#$ to \mathbb{R}_+ by $C_c^\infty(\mathbb{R}_+)^\#$. For a function $f \in C_c^\infty(\mathbb{R})^\#$, the Jacobi transform $\widehat{f}(\lambda)$, $\lambda \in \mathbb{C}$, is defined by

$$\widehat{f}(\lambda) := \int_0^\infty f(t) \phi_\lambda(t) A(t) dt.$$

The following inversion formula holds for functions in $C_c^\infty(\mathbb{R})^\#$:

$$f(t) = \frac{1}{2\pi} \int_0^\infty \widehat{f}(\lambda) \phi_\lambda(t) |c(\lambda)|^{-2} d\lambda, \quad t \in \mathbb{R},$$

where $c(\lambda)$ is a multiple of the meromorphic Harish-Chandra function given by the formula

$$c(\lambda) = \frac{2^{\rho-i\lambda} \Gamma(\alpha+1) \Gamma(i\lambda)}{\Gamma(\frac{1}{2}(\rho+i\lambda)) \Gamma(\frac{1}{2}(\rho+i\lambda) - \beta)}.$$

If we denote the measure $A(t)dt$ by $d\mu(t)$ then the space $L^p(\mathbb{R}_+, d\mu)$, $1 \leq p < \infty$, associated with the measure $d\mu$, at times, denoted by $L^p(\mu)$, is defined as the space of all even functions on \mathbb{R} such that

$$\|f\|_{L^p(\mu)} := \left(\int_0^\infty |f(t)|^p A(t) dt \right)^{\frac{1}{p}} = \left(\int_0^\infty |f(t)|^p d\mu(t) \right)^{\frac{1}{p}} < \infty, \quad 1 \leq p < \infty.$$

It is well-known that the Jacobi transform $f \mapsto \widehat{f}$ extends to an isometry from $L^2(\mu)$ onto $L^2(\mathbb{R}_+, \frac{1}{2\pi} |c(\lambda)|^{-2} d\lambda)$.

Let $S_p := \{\lambda \in \mathbb{C} : |\operatorname{Im}(\lambda)| < (1 - \frac{2}{p})\rho\}$. Then, for $f \in L^p(\mu)$, $1 \leq p < 2$ with $\frac{1}{p} + \frac{1}{p'} = 1$, the Jacobi transform $\widehat{f}(\lambda)$ is well-defined and holomorphic in S_p and for all $\lambda \in S_p$, we have

$$|\widehat{f}(\lambda)| \leq \|f\|_{L^p(\mu)} \|\phi_\lambda\|_p.$$

In particular, if $p = 1$, $\widehat{f}(\lambda)$ is continuous on S_1 and for all $\lambda \in S_1$, we have

$$|\widehat{f}(\lambda)| \leq \|f\|_{L^1(\mu)}.$$

For our convenience we denote the measure $\frac{1}{2\pi} |c(\lambda)|^{-2} d\lambda$ by $d\kappa(\lambda)$.

The following theorem is the Hausdorff-Young inequality for Jacobi transform (see [26]).

Theorem 5.1. *Let $1 \leq p \leq 2$ with $\frac{1}{p} + \frac{1}{p'} = 1$. If $f \in L^p(\mathbb{R}_+, d\mu)$ then we have*

$$\|\widehat{f}\|_{L^{p'}(\mathbb{R}_+, d\kappa(\lambda))} \leq \|f\|_{L^p(\mu)}.$$

5.1. Paley-type inequality for Jacobi transform. The following inequality can be regarded as a Paley-type inequality which plays an important in proving the Hausdorff-Young-Paley inequality.

Theorem 5.2. *Suppose that ψ is a positive function on \mathbb{R}_+ satisfying the condition*

$$M_\psi := \sup_{t>0} t \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) > t}} |c(\lambda)|^{-2} d\lambda < \infty. \quad (34)$$

Then for $f \in L^p(\mu)$, $1 < p \leq 2$, we have

$$\left(\int_0^\infty |\widehat{f}(\lambda)|^p \psi(\lambda)^{2-p} |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p}} \lesssim M_\psi^{\frac{2-p}{p}} \|f\|_{L^p(\mu)}. \quad (35)$$

Proof. We follow the proof in [2, 3]. Let us consider a measure ν on \mathbb{R}_+ given by

$$\nu(\lambda) = \psi(\lambda)^2 d\kappa(\lambda), \quad (36)$$

where $d\kappa = \frac{1}{2\pi}|c(\lambda)|^{-2}d\lambda$. We define the corresponding $L^p(\mathbb{R}_+, \nu)$ -space, $1 \leq p < \infty$, as the space of all complex-valued function f defined by \mathbb{R}_+ such that

$$\|f\|_{L^p(\mathbb{R}_+, \nu)} := \left(\int_{\mathbb{R}_+} |f(\lambda)|^p \psi(\lambda)^2 d\kappa(\lambda) \right)^{\frac{1}{p}} < \infty.$$

We define a sublinear operator T for $f \in L^p(\mu)$ by

$$Tf(\lambda) = \frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} \in L^p(\mathbb{R}_+, \nu).$$

We will show that T is well-defined and bounded from $L^p(\mu)$ to $L^p(\mathbb{R}_+, \nu)$ for any $1 < p \leq 2$. In other words, we claim the following estimate:

$$\|Tf\|_{L^p(\mathbb{R}_+, \nu)} = \left(\int_{\mathbb{R}_+} \frac{|\widehat{f}(\lambda)|^p}{\psi(\lambda)^p} \psi(\lambda)^2 d\kappa(\lambda) \right)^{\frac{1}{p}} \lesssim M_{\psi}^{\frac{2-p}{p}} \|f\|_{L^p(\mu)}, \quad (37)$$

which will give us the required inequality (35) with $M_{\psi} := \sup_{t>0} \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) > t}} d\kappa(\lambda)$. We will show that T is weak-type $(2, 2)$ and weak-type $(1, 1)$. More precisely, with the distribution function,

$$\nu(y; Tf) = \int_{\substack{\lambda \in \mathbb{R}_+ \\ \frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} \geq y}} \psi(\lambda)^2 d\kappa(\lambda),$$

where ν is give by formula (36), we show that

$$\nu(y; Tf) \leq \left(\frac{M_2 \|f\|_2}{y} \right)^2 \quad \text{with norm } M_2 = 1, \quad (38)$$

$$\nu(y; Tf) \leq \frac{M_1 \|f\|_1}{y} \quad \text{with norm } M_1 = M_{\psi}. \quad (39)$$

Then the estimate (37) follows from the Marcinkiewicz interpolation Theorem. Now, to show (38), using Plancherel identity we get

$$\begin{aligned} y^2 \nu(y; Tf) &\leq \sup_{y>0} y^2 \nu(y; Tf) = \|Tf\|_{L^{2,\infty}(\mathbb{R}_+, \nu)}^2 \leq \|Tf\|_{L^2(\mathbb{R}_+, \nu)}^2 \\ &= \int_{\mathbb{R}_+} \left(\frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} \right)^2 \psi(\lambda)^2 d\kappa(\lambda) \\ &= \int_{\mathbb{R}_+} |\widehat{f}(\lambda)|^2 d\kappa(\lambda) = \|f\|_2^2. \end{aligned}$$

Thus, T is type $(2, 2)$ with norm $M_2 \leq 1$. Further, we show that T is of weak type $(1, 1)$ with norm $M_1 = M_{\psi}$; more precisely, we show that

$$\nu \left\{ \lambda \in \mathbb{R}_+ : \frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} > y \right\} \lesssim M_{\psi} \frac{\|f\|_1}{y}. \quad (40)$$

The left hand side is an integral $\int \psi(\lambda) d\kappa(\lambda)$ taken over all those $\lambda \in \mathbb{R}_+$ for which $\frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} > y$. Since $|\widehat{f}(\lambda)| \leq \|f\|_1$ for all $\lambda \in \mathbb{R}_+$ we have

$$\left\{ \lambda \in \mathbb{R}_+ : \frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} > y \right\} \subset \left\{ \lambda \in \mathbb{R}_+ : \frac{\|f\|_1}{\psi(\lambda)} > y \right\},$$

for any $y > 0$ and, therefore,

$$\nu \left\{ \lambda \in \mathbb{R}_+ : \frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} > y \right\} \leq \nu \left\{ \lambda \in \mathbb{R}_+ : \frac{\|f\|_1}{\psi(\lambda)} > y \right\}.$$

Now by setting $w := \frac{\|f\|_1}{y}$, we have

$$\nu \left\{ \lambda \in \mathbb{R}_+ : \frac{\|f\|_1}{\psi(\lambda)} > y \right\} \leq \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) \leq w}} \psi(\lambda)^2 d\kappa(\lambda). \quad (41)$$

Now we claim that

$$\int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) \leq w}} \psi(\lambda)^2 d\kappa(\lambda) \lesssim M_\psi w. \quad (42)$$

Indeed, first we notice that

$$\int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) \leq w}} \psi(\lambda)^2 d\kappa(\lambda) = \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) \leq w}} d\kappa(\lambda) \int_0^{\psi(\lambda)^2} d\tau.$$

By interchanging the order of integration we get

$$\int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) \leq w}} d\kappa(\lambda) \int_0^{\psi(\lambda)^2} d\tau = \int_0^{w^2} d\tau \int_{\substack{\lambda \in \mathbb{R}_+ \\ \tau^{\frac{1}{2}} \leq \psi(\lambda) \leq w}} d\kappa(\lambda).$$

Further, by making substitution $\tau = t^2$, it gives

$$\begin{aligned} \int_0^{w^2} d\tau \int_{\substack{\lambda \in \mathbb{R}_+ \\ \tau^{\frac{1}{2}} \leq \psi(\lambda) \leq w}} d\kappa(\lambda) &= 2 \int_0^w t dt \int_{\substack{\lambda \in \mathbb{R}_+ \\ t \leq \psi(\lambda) \leq w}} d\kappa(\lambda) \\ &\leq 2 \int_0^w t dt \int_{\substack{\lambda \in \mathbb{R}_+ \\ t \leq \psi(\lambda)}} d\kappa(\lambda). \end{aligned}$$

Since

$$t \int_{\substack{\lambda \in \mathbb{R}_+ \\ t \leq \psi(\lambda)}} d\kappa(\lambda) \leq \sup_{t>0} t \int_{\substack{\lambda \in \mathbb{R}_+ \\ t \leq \psi(\lambda)}} d\kappa(\lambda) = M_\psi$$

is finite by assumption $M_\psi < \infty$, we have

$$2 \int_0^w t dt \int_{\substack{\lambda \in \mathbb{R}_+ \\ t \leq \psi(\lambda)}} d\kappa(\lambda) \lesssim M_\psi w.$$

This establishes our claim (42) and eventually proves (40). So, we have proved (38) and (39). Then by using the Marcinkiewicz interpolation theorem with $p_1 = 1$ and $p_2 = 2$ and $\frac{1}{p} = 1 - \theta + \frac{\theta}{2}$ we now obtain

$$\left(\int_{\mathbb{R}_+} \left(\frac{|\widehat{f}(\lambda)|}{\psi(\lambda)} \right)^p \psi(\lambda)^2 d\kappa(\lambda) \right)^{\frac{1}{p}} = \|Tf\|_{L^p(\mathbb{R}_+, \nu)} \lesssim M_\psi^{\frac{2-p}{p}} \|f\|_{L^p(\mu)}.$$

This completes the proof of the theorem. \square

5.2. Hausdorff-Young-Paley inequality for Jacobi transform. The following theorem [11] is useful to obtain one of our crucial result.

Theorem 5.3. *Let $d\mu_0(x) = \omega_0(x)d\mu'(x)$, $d\mu_1(x) = \omega_1(x)d\mu'(x)$, and write $L^p(\omega) = L^p(\omega d\mu')$ for the weight ω . Suppose that $0 < p_0, p_1 < \infty$. Then*

$$(L^{p_0}(\omega_0), L^{p_1}(\omega_1))_{\theta, p} = L^p(\omega),$$

where $0 < \theta < 1$, $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and $\omega = \omega_0^{\frac{p(1-\theta)}{p_0}} \omega_1^{\frac{p\theta}{p_1}}$.

The following corollary is immediate.

Corollary 5.4. *Let $d\mu_0(x) = \omega_0(x)d\mu'(x)$, $d\mu_1(x) = \omega_1(x)d\mu'(x)$. Suppose that $0 < p_0, p_1 < \infty$. If a continuous linear operator A admits bounded extensions, $A : L^p(Y, \mu) \rightarrow L^{p_0}(\omega_0)$ and $A : L^p(Y, \mu) \rightarrow L^{p_1}(\omega_1)$, then, we there exists a bounded extension $A : L^p(Y, \mu) \rightarrow L^b(\omega)$ of A , where $0 < \theta < 1$, $\frac{1}{b} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and $\omega = \omega_0^{\frac{b(1-\theta)}{p_0}} \omega_1^{\frac{b\theta}{p_1}}$.*

Using the above corollary we now present the Hausdorff-Young-Paley inequality.

Theorem 5.5 (Hausdorff-Young-Paley inequality). *Let $1 < p \leq 2$, and let $1 < p \leq b \leq p' < \infty$, where $p' = \frac{p}{p-1}$. If $\psi(\lambda)$ is a positive function on \mathbb{R}_+ such that*

$$M_\psi := \sup_{t>0} t \int_{\substack{\lambda \in \mathbb{R}_+ \\ \psi(\lambda) > t}} d\kappa(\lambda) \quad (43)$$

is finite then for every $f \in L^p(\mu)$ we have

$$\left(\int_{\mathbb{R}_+} \left(|\widehat{f}(\lambda)| \psi(\lambda)^{\frac{1}{b} - \frac{1}{p'}} \right)^b d\kappa(\lambda) \right)^{\frac{1}{b}} \lesssim M_\psi^{\frac{1}{b} - \frac{1}{p'}} \|f\|_{L^p(\mu)}. \quad (44)$$

This naturally reduced to Hausdorff-Young inequality when $b = p'$ and Paley inequality (35) when $b = p$.

Proof. From Theorem 5.2, the operator defined by

$$Af(\lambda) = \widehat{f}(\lambda), \quad \lambda \in \mathbb{R}_+$$

is bounded from $L^p(\mu)$ to $L^p(\mathbb{R}_+, \omega_0 d\mu')$, where $d\mu'(\lambda) = d\kappa(\lambda)$ and $\omega_0(\lambda) = \psi(\lambda)^{2-p}$. From Theorem 5.1, we deduce that $A : L^p(\mu) \rightarrow L^{p'}(\mathbb{R}_+, \omega_1 d\mu')$ with $d\mu'(\lambda) = d\kappa(\lambda)$ and $\omega_1(\lambda) = 1$ admits a bounded extension. By using the real interpolation (Corollary 5.4 above) we will prove that $A : L^p(\mu) \rightarrow L^b(\mathbb{R}_+, \omega d\mu')$, $p \leq b \leq p'$, is bounded, where the space $L^p(\mathbb{R}_+, \omega d\mu')$ is defined by the norm

$$\|\sigma\|_{L^p(\mathbb{R}, \omega d\mu')} := \left(\int_{\mathbb{R}_+} |\sigma(\lambda)|^p w(\lambda) d\mu'(\lambda) \right)^{\frac{1}{p}} = \left(\int_{\mathbb{R}_+} |\sigma(\lambda)|^p w(\lambda) d\kappa(\lambda) \right)^{\frac{1}{p}}$$

and $\omega(\lambda)$ is positive function over \mathbb{R}_+ to be determined. To compute ω , we can use Corollary 5.4, by fixing $\theta \in (0, 1)$ such that $\frac{1}{b} = \frac{1-\theta}{p} + \frac{\theta}{p'}$. In this case $\theta = \frac{p-b}{b(p-2)}$, and

$$\omega = \omega_0^{\frac{p(1-\theta)}{p-2}} \omega_1^{\frac{p\theta}{p-2}} = \psi(\lambda)^{1-\frac{b}{p'}}. \quad (45)$$

Thus we finish the proof. \square

The following theorem is the main result of this section which is an analogue of Hörmander multiplier theorem [33] for Jacobi transform.

Theorem 5.6. *Let $1 < p \leq 2 \leq q < \infty$. Suppose that T is a Jacobi-Fourier multiplier with symbol h , that is,*

$$\widehat{Tf}(\lambda) = h(\lambda)\widehat{f}(\lambda), \quad \lambda \in \mathbb{R}_+,$$

where h is an bounded measurable even function on \mathbb{R} . Then we have

$$\|T\|_{L^p(\mu) \rightarrow L^q(\mu)} \lesssim \sup_{s>0} s \left[\int_{\{\lambda \in \mathbb{R}_+ : |h(\lambda)| > s\}} d\kappa(\lambda) \right]^{\frac{1}{p} - \frac{1}{q}}.$$

Proof. Let us first assume that $p \leq q'$, where $\frac{1}{p} + \frac{1}{q'} = 1$. Since $q' \leq 2$, the Hausdorff-Young inequality gives that

$$\|Tf\|_{L^q(\mu)} \leq \|\widehat{Tf}\|_{L^{q'}(\mu)} = \|h\widehat{f}\|_{L^{q'}(\mu)}.$$

The case $q' \leq (p')' = p$ can be reduced to the case $p \leq q'$ as follows. Using the duality of L^p -spaces we have $\|T\|_{L^p(\mu) \rightarrow L^q(\mu)} = \|T^*\|_{L^{q'}(\mu) \rightarrow L^{p'}(\mu)}$. The symbol of adjoint operator T^* is equal to \check{h} , which equal to h and obviously we have $|\check{h}| = |h|$ (see Proposition 4.10 in [6]). Now, we are in a position to apply Theorem 5.5. Set $\frac{1}{p} - \frac{1}{q} = \frac{1}{r}$. Now, by applying Theorem 5.5 with $\psi = |h|^r$ with $b = q'$ we get

$$\|h\widehat{f}\|_{L^{q'}(\mu)} \lesssim \left(\sup_{s>0} s \int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)|^r > s}} d\kappa(\lambda) \right)^{\frac{1}{r}} \|f\|_{L^p(\mu)}$$

for all $f \in L^p(\mu)$, in view of $\frac{1}{p} - \frac{1}{q} = \frac{1}{q'} - \frac{1}{p'} = \frac{1}{r}$. Thus, for $1 < p \leq 2 \leq q < \infty$, we obtain

$$\|Tf\|_{L^q(\mu)} \lesssim \left(\sup_{s>0} s \int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)|^r > s}} d\kappa(\lambda) \right)^{\frac{1}{r}} \|f\|_{L^p(\mu)}.$$

Further, the proof follows from the following inequality:

$$\begin{aligned} \left(\sup_{s>0} s \int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)|^r > s}} d\kappa(\lambda) \right)^{\frac{1}{r}} &= \left(\sup_{s>0} s \int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)| > s^{\frac{1}{r}}}} d\kappa(\lambda) \right)^{\frac{1}{r}} \\ &= \left(\sup_{s>0} s^r \int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)| > s}} d\kappa(\lambda) \right)^{\frac{1}{r}} \end{aligned}$$

$$= \sup_{s>0} s \left(\int_{\substack{\lambda \in \mathbb{R}_+ \\ |h(\lambda)| > s}} d\kappa(\lambda) \right)^{\frac{1}{r}},$$

proving Theorem 5.6. \square

As an immediate consequence of above Theorem 5.6 we have the following multiplier theorem on harmonic NA groups.

Corollary 5.7. *Let S be a harmonic NA group and let $1 < p \leq 2 \leq q < \infty$. Suppose that h is a bounded measurable even function on \mathbb{R} . Then the Fourier multiplier defined by $T_h f := \mathcal{H}^{-1}(h\mathcal{H}f)$ for $f \in C_c^\infty(S)^\#$ is bounded from $L^p(S)^\#$ to $L^q(S)^\#$ provided that*

$$\sup_{s>0} s \int_{\{\lambda \in \mathbb{R}_+ : |h(\lambda)|^{\frac{1}{p}-\frac{1}{q}} > s\}} |c(\lambda)|^{-2} d\lambda < \infty.$$

Moreover,

$$\|T_h\|_{L^p(S)^\# \rightarrow L^q(S)^\#} \lesssim \sup_{s>0} s \left(\int_{\{\lambda \in \mathbb{R}_+ : |h(\lambda)| > s\}} |c(\lambda)|^{-2} d\lambda \right)^{\frac{1}{p}-\frac{1}{q}}.$$

Now, we apply Theorem 5.6 to prove the L^p - L^q boundedness of spectral multipliers for operator $L := -\mathcal{L}_{\alpha,\beta}$. If $\varphi \in L^\infty(\mathbb{R}_+, d\mu)$, the spectral multiplier $\varphi(L)$, defined by φ coincides with the Jacobi Fourier multiplier T_h with $h(\lambda) = \varphi(\lambda^2 + \rho^2)$ for $\lambda \in \mathbb{R}_+$. The L^p -boundedness of spectral multipliers has been proved by several authors in many different setting, e.g., Bessel transform [14], Dunkl harmonic oscillator [48]. The L^p - L^q boundedness of spectral multipliers for compact Lie groups, Heisenberg groups, graded Lie groups have been proved by R. Akyzhanov and the second author [3]. M. Chatzakou and the second author recently studied L^p - L^q boundedness of spectral multiplier for the anharmonic oscillator [16] (see also [15]).

Theorem 5.8. *Let $1 < p \leq 2 \leq q < \infty$ and let φ be a monotonically decreasing continuous function on $[\rho^2, \infty)$ such that $\lim_{u \rightarrow \infty} \varphi(u) = 0$. Then we have*

$$\|\varphi(L)\|_{\text{op}} \lesssim \sup_{u>\rho^2} \varphi(u) \begin{cases} (u - \rho^2)^{\frac{3}{2}(\frac{1}{p}-\frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} \leq 1, \\ (u - \rho^2)^{(\alpha+1)(\frac{1}{p}-\frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} > 1, \end{cases} \quad (46)$$

where $\|\cdot\|_{\text{op}}$ denotes the operator norm from $L^p(\mu)$ to $L^q(\mu)$.

Proof. Since $\varphi(L)$ is a Jacobi-Fourier multiplier with the symbol $\varphi(\lambda^2 + \rho^2)$, as an application of Theorem 5.6, we get

$$\begin{aligned} \|\varphi(L)\|_{\text{op}} &\lesssim \sup_{s>0} s \left[\int_{\{\lambda \in \mathbb{R}_+ : \varphi(\lambda^2 + \rho^2) \geq s\}} |c(\lambda)|^{-2} d\lambda \right]^{\frac{1}{p}-\frac{1}{q}} \\ &= \sup_{0 < s < \varphi(\rho^2)} s \left[\int_{\{\lambda \in \mathbb{R}_+ : \varphi(\lambda^2 + \rho^2) \geq s\}} |c(\lambda)|^{-2} d\lambda \right]^{\frac{1}{p}-\frac{1}{q}}, \end{aligned}$$

since $\varphi \leq \varphi(\rho^2)$. Now, as $s \in (0, \varphi(\rho^2)]$ we can write $s = \varphi(u)$ for some $u \in [\rho^2, \infty)$ and, therefore, we have

$$\|\varphi(L)\|_{\text{op}} \lesssim \sup_{\varphi(u) < \varphi(\rho^2)} \varphi(u) \left[\int_{\{\lambda \in \mathbb{R}_+ : \varphi(\lambda^2 + \rho^2) \geq \varphi(u)\}} |c(\lambda)|^{-2} d\lambda \right]^{\frac{1}{p} - \frac{1}{q}}.$$

Since φ is monotonically decreasing we get

$$\|\varphi(L)\|_{\text{op}} \lesssim \sup_{u > \rho^2} \varphi(u) \left[\int_{\{\lambda \in \mathbb{R}_+ : \lambda \leq (u - \rho^2)^{\frac{1}{2}}\}} |c(\lambda)|^{-2} d\lambda \right]^{\frac{1}{p} - \frac{1}{q}}.$$

Now, we use the estimate of the c -function to get

$$\begin{aligned} \|\varphi(L)\|_{\text{op}} &\lesssim \sup_{u > \rho^2} \varphi(u) \begin{cases} \left[\int_0^{(u - \rho^2)^{\frac{1}{2}}} \lambda^2 d\lambda \right]^{\frac{1}{p} - \frac{1}{q}} & \text{if } (u - \rho^2)^{\frac{1}{2}} \leq 1, \\ \left[\int_0^1 \lambda^2 d\lambda + \int_1^{(u - \rho^2)^{\frac{1}{2}}} \lambda^{2\alpha+1} d\lambda \right]^{\frac{1}{p} - \frac{1}{q}} & \text{if } (u - \rho^2)^{\frac{1}{2}} > 1 \end{cases} \\ &= \sup_{u > \rho^2} \varphi(u) \begin{cases} (u - \rho^2)^{\frac{3}{2}(\frac{1}{p} - \frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} \leq 1, \\ (u - \rho^2)^{(\alpha+1)(\frac{1}{p} - \frac{1}{q})} & \text{if } (u - \rho^2)^{\frac{1}{2}} > 1, \end{cases} \end{aligned}$$

proving Theorem 5.8. □

ACKNOWLEDGMENT

The authors thank Jean-Philippe Anker for discussions. Vishvesh Kumar thanks William O. Bray for his suggestions. VK and MR are supported by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations and by the Methusalem programme of the Ghent University Special Research Fund (BOF) (Grant number 01M01021). MR is also supported by EPSRC grant EP/R003025/2.

REFERENCES

1. Akylzhanov, R., Nursultanov, E., Ruzhansky, M. Hardy-Littlewood-Paley inequalities and Fourier multipliers on $SU(2)$. *Studia Math.* 234 (2016), no. 1, 1-29.
2. Akylzhanov, R., Ruzhansky, M., Nursultanov, E. Hardy-Littlewood, Hausdorff-Young-Paley inequalities, and $L^p - L^q$ Fourier multipliers on compact homogeneous manifolds. *J. Math. Anal. Appl.* 479 (2019), no. 2, 1519-1548.
3. Akylzhanov, R., Ruzhansky, M. $L^p - L^q$ multipliers on locally compact groups, *J. Func. Anal.*, 278(3) (2019), DOI: <https://doi.org/10.1016/j.jfa.2019.108324>
4. Anker, J.-P. L^p Fourier multipliers on Riemannian symmetric spaces of the noncompact type. *Ann. of Math. (2)* 132 (1990), no. 3, 597-628.
5. Anker, J.-P. Sharp estimates for some functions of the Laplacian on noncompact symmetric spaces. *Duke Math. J.* 65 (1992), no. 2, 257-297.
6. Anker, J.-P., Damek, E., Yacoub, C. Spherical analysis on harmonic AN groups. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) 23 (1996), no. 4, 643-679.
7. Astengo, F. A class of L^p convolutors on harmonic extensions of H-type groups. *J. Lie Theory* 5 (1995), no. 2, 147-164.
8. Astengo, F. Multipliers for a distinguished Laplacean on solvable extensions of H-type groups. *Monatsh. Math.* 120 (1995), no. 3-4, 179-188.

9. Astengo, F., Camporesi, R., Di Blasio, B. The Helgason Fourier transform on a class of nonsymmetric harmonic spaces. *Bull. Austral. Math. Soc.* 55 (1997), no. 3, 405-424.
10. Bloom, W. R., Xu, Z. Fourier multipliers for L_p on Chébli-Trimèche hypergroups. *Proc. London Math. Soc.* (3) 80 (2000), no. 3, 643-664.
11. Bergh, J., Lofstrom, J., *Interpolation spaces*, Grundlehren der mathematischen Wissenschaften, (1976).
12. Bray, W. O. Growth and integrability of Fourier transforms on Euclidean space. *J. Fourier Anal. Appl.* 20 (2014), no. 6, 1234-1256.
13. Bray, W. O., Pinsky, M. A. Growth properties of Fourier transforms via moduli of continuity. *J. Funct. Anal.* 255 (2008), no. 9, 2265-2285.
14. Betancor, J. J., Castro, A. J., Curbelo, J. Spectral multipliers for multidimensional Bessel operators, *J. Fourier Anal. Appl.* 17(5) (2011), 932-975.
15. Cardona, D., Kumar, V., Ruzhansky, M., Tokmagambetov, N. L^p - L^q boundedness of pseudo-differential operators on smooth manifolds and its applications to nonlinear equations, (2020). <https://arxiv.org/abs/2005.04936>
16. Chatzakou, M., Kumar M., L^p - L^q boundedness of Fourier multipliers associated with the anharmonic Oscillator, (2020). <https://arxiv.org/abs/2004.07801>
17. Cowling, M., Dooley A., Koranyi, A., Ricci, F. H -type groups and Iwasawa decompositions, *Adv. Math.* 87 (1991), no. 1, 1-41.
18. Cowling, M., Giulini, S., Meda, S. L^p - L^q estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. I. *Duke Math. J.* 72 (1993), no. 1, 109-150.
19. Cowling, M., Giulini, S., Meda, S. L^p - L^q -estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. II. *J. Lie Theory* 5 (1995), no. 1, 1-14.
20. Cowling, M., Giulini, S., Meda, S. L^p - L^q estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces. III. *Ann. Inst. Fourier (Grenoble)* 51 (2001), no. 4, 1047-1069.
21. Cowling, M., Giulini, S., Meda, S. Oscillatory multipliers related to the wave equation on noncompact symmetric spaces. *J. London Math. Soc.* (2) 66 (2002), no. 3, 691-709.
22. Damek, E. The geometry of a semidirect extension of a Heisenberg type nilpotent group. *Colloq. Math.* 53 (1987), no. 2, 255-268.
23. Damek, E., Ricci, F. A class of nonsymmetric harmonic Riemannian spaces. *Bull. Amer. Math. Soc.* 27 (1992), no. 1, 139-142.
24. Damek, E., Ricci, F. Harmonic analysis on solvable extensions of H -type groups. *J. Geom. Anal.* 2 (1992), no. 3, 213-248.
25. Di Blasio, B. Paley-Wiener type theorems on harmonic extensions of H -type groups. *Monatsh. Math.* 123 (1997), no. 1, 21-42.
26. Degenfeld-Schonburg, S. On the Hausdorff-Young theorem for commutative hypergroups, *Colloq. Math.*, 131(2) (2013) 219-231.
27. Daher, R., Delgado, J., Ruzhansky, M. Titchmarsh theorems for Fourier transforms of Hölder-Lipschitz functions on compact homogeneous manifolds. *Monatsh. Math.* 189 (2019), no. 1, 23-49.
28. Delgado, J., Ruzhansky, M. Fourier multipliers, symbols, and nuclearity on compact manifolds. *J. Anal. Math.* 135 (2018), no. 2, 757-800.
29. Johansen, T. R. Non-integrable multipliers for the Jacobi transform. *Integral Transforms Spec. Funct.* 24 (2013), no. 6, 425-440.
30. Fahlaoui, S., Boujeddaine, M., El Kassimi, M. Fourier transforms of Dini-Lipschitz functions on rank 1 symmetric spaces. *Mediterr. J. Math.* 13 (2016), no. 6, 4401-4411.

31. Flensted-Jensen, M., Koornwinder, Tom H. The convolution structure for Jacobi function expansions. *Ark. Mat.* 11 (1973), 245-262.
32. Flensted-Jensen, M., Koornwinder, Tom H. Jacobi functions: the addition formula and the positivity of the dual convolution structure. *Ark. Mat.* 17 (1979), no. 1, 139-151.
33. Hörmander, L. Estimates for translation invariant operators in L^p spaces. *Acta Math.*, 104:93-140, 1960.
34. Hörmander, L. The analysis of linear partial differential operators, volume 3, Springer-Verlag, Berlin, 1985.
35. Kaplan, A. Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms. *Trans. Amer. Math. Soc.* 258 (1980), no. 1, 147-153.
36. Kumar, P., Ray, S. K., Sarkar, R. P. The role of restriction theorems in harmonic analysis on harmonic NA groups. *J. Funct. Anal.* 258 (2010), no. 7, 2453-2482.
37. Koornwinder, Tom H. Jacobi functions and analysis on noncompact semisimple Lie groups. Special functions: group theoretical aspects and applications, 1-85, *Math. Appl.*, Reidel, Dordrecht, (1984).
38. Littlewood, J.E., Paley, R. E. A. Theorems on Fourier series and power series, *J. London Math. Soc.* 6 (1931), 230-233
39. Littlewood, J.E., Paley, R. E. A. Theorems on Fourier series and power series (II). *Proc. London Mat. Soc.*, 42 (1937), 52-89.
40. Mohanty, P., Ray, S. K., Sarkar, R. P., Sitaram, A. The Helgason-Fourier transform for symmetric spaces. II. *J. Lie Theory* 14 (2004), no. 1, 227-242.
41. Müller, D., Peloso, M. M., Ricci, F. L^p -spectral multipliers for the Hodge Laplacian acting on 1-forms on the Heisenberg group. *Geom. Funct. Anal.* 17 (2007), no. 3, 852-886.
42. Platonov, S.S. The Fourier transform of functions satisfying a Lipschitz condition on symmetric spaces of rank 1. *Sibirsk. Mat. Zh.* 46(6), 1374-1387 (2005).
43. Platonov, S. S. The Fourier transform of functions satisfying a Lipschitz condition on symmetric spaces of rank 1. (Russian) *Sibirsk. Mat. Zh.* 46 (2005), no. 6, 1374-1387; translation in *Siberian Math. J.* 46 (2005), no. 6, 1108-1118
44. Ray, S. K., Sarkar, R. P. Fourier and Radon transform on harmonic NA groups. *Trans. Amer. Math. Soc.* 361 (2009), no. 8, 4269-4297.
45. Ruzhansky, M., Wirth, J. L^p Fourier multipliers on compact Lie groups. *Math. Z.* 280 (2015), no. 3-4, 621-642.
46. Titchmarsh E. C. *Introduction to the Theory of Fourier Integral*, Oxford University Press, Oxford (1937).
47. Vallarino, M. *Analysis on harmonic extensions of H-type groups*, Ph.D. Thesis, Università degli Studi di Milano-Bicocca, 2005. arXiv:1504.00329.
48. Wróbel, B. Multivariate spectral multipliers for the Dunkl transform and the Dunkl harmonic oscillator, *Forum Math.* 27(4) (2015), 2301-2322.
49. Younis, M. S. Fourier transforms of Dini-Lipschitz functions. *Internat. J. Math. Math. Sci.* 9 (1986), no. 2, 301-312.
50. Younis, M. S. Fourier transforms of Lipschitz functions on certain Lie groups. *Int. J. Math. Math. Sci.* 27 (2001), no. 7, 439-448.
51. Younis, M. S. Fourier transforms of Dini-Lipschitz functions on Vilenkin groups. *Internat. J. Math. Math. Sci.* 15 (1992), no. 3, 609-612.

VISHVESH KUMAR

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS

GHENT UNIVERSITY, BELGIUM

Email address: vishveshmishra@gmail.com, Vishvesh.Kumar@UGent.be

MICHAEL RUZHANSKY

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS

GHENT UNIVERSITY, BELGIUM

AND

SCHOOL OF MATHEMATICAL SCIENCES

QUEEN MARY UNIVERSITY OF LONDON

UNITED KINGDOM

E-mail address michael.ruzhansky@ugent.be