

SHARP SUPERLEVEL SET ESTIMATES FOR SMALL CAP DECOUPLINGS OF THE PARABOLA

YUQIU FU, LARRY GUTH, AND DOMINIQUE MALDAGUE

ABSTRACT. We prove sharp bounds for the size of superlevel sets $\{x \in \mathbb{R}^2 : |f(x)| > \alpha\}$ where $\alpha > 0$ and $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ is a Schwartz function with Fourier transform supported in an R^{-1} -neighborhood of the truncated parabola \mathbb{P}^1 . These estimates imply the small cap decoupling theorem for \mathbb{P}^1 from [DGW20] and the canonical decoupling theorem for \mathbb{P}^1 from [BD15]. New (ℓ^q, L^p) small cap decoupling inequalities also follow from our sharp level set estimates.

In this paper, we further develop the high/low frequency proof of decoupling for the parabola [GMW20] to prove sharp level set estimates which recover and refine the small cap decoupling results for the parabola in [DGW20]. We begin by describing the problem and our results in terms of exponential sums. The main results in full generality are in §1.

For $N \geq 1$, $R \in [N, N^2]$, and $2 \leq p$, let $D(N, R, p)$ denote the smallest constant so that

$$(1) \quad |Q_R|^{-1} \int_{Q_R} \left| \sum_{\xi \in \Xi} a_\xi e((x, t) \cdot (\xi, \xi^2)) \right|^p dx dt \leq D(N, R, p) N^{p/2}$$

for any collection $\Xi \subset [-1, 1]$ with $|\Xi| \sim N$ consisting of $\sim \frac{1}{N}$ -separated points, $a_\xi \in \mathbb{C}$ with $|a_\xi| \sim 1$, and any cube $Q_R \subset \mathbb{R}^2$ of sidelength R .

A corollary of the small cap decoupling theorem for the parabola in [DGW20] is that if $2 \leq p \leq 2 + 2s$ for $R = N^s$, then

$$(2) \quad D(N, R, p) \leq C_\varepsilon N^\varepsilon.$$

This estimate is sharp, up to the $C_\varepsilon N^\varepsilon$ factor, which may be seen by Khintchine's inequality. The range $2 \leq p \leq 2 + 2s$ is the largest range of p for which $D(N, R, p)$ may be bounded by sub-polynomial factors in N . The case $R = N^2$ of (2) follows from the canonical ℓ^2 decoupling theorem of Bourgain and Demeter for the parabola [BD15]. For $R < N^2$ and the subset $\Xi = \{k/N\}_{k=1}^N$, the inequality (1) is an estimate for the moments of exponential sums over subsets smaller than the full domain of periodicity (i.e. N^2 in the t -variable). Bourgain investigated examples of this type of inequality in [Bou17b, Bou17a].

By a pigeonholing argument (see Section 5 of [GMW20]), (2) follows from upper bounds for superlevel sets U_α defined by

$$U_\alpha = \{(x, t) \in \mathbb{R}^2 : \left| \sum_{\xi \in \Xi} a_\xi e((x, t) \cdot (\xi, \xi^2)) \right| > \alpha\}.$$

In particular, (2) is equivalent, up to a $\log N$ factor, to proving that for any $\alpha > 0$ and for $R = N^s$,

$$(3) \quad \alpha^{2+2s} |U_\alpha \cap Q_R| \leq C_\varepsilon R^\varepsilon N^{1+s} R^2$$

Date: July 29, 2021.

when Ξ , a_ξ satisfy the hypotheses following (1). In this paper, we improve the above superlevel set estimate for all $\alpha > 0$ strictly between $N^{1/2}$ and N .

Theorem 1. *Let $R \in [N, N^2]$. For any $\varepsilon > 0$, there exists $C_\varepsilon < \infty$ such that*

$$|U_\alpha \cap Q_R| \leq C_\varepsilon N^\varepsilon \begin{cases} \frac{N^2 R}{\alpha^4} \sum_{\xi \in \Xi} |a_\xi|^2 & \text{if } \alpha^2 > R \\ \frac{N^2 R^2}{\alpha^6} \sum_{\xi \in \Xi} |a_\xi|^2 & \text{if } N \leq \alpha^2 \leq R \\ R^2 & \text{if } \alpha^2 < N. \end{cases}$$

whenever $\Xi \subset [-1, 1]$ is a $\gtrsim \frac{1}{N}$ -separated subset, $|a_\xi| \leq 1$ for each $\xi \in \Xi$, and $Q_R \subset \mathbb{R}^2$ is a cube of sidelength R .

Our superlevel set estimates are essentially sharp, which follows from analyzing the function $F(x, t) = \sum_{n=1}^N e((x, t) \cdot (\frac{n}{N}, \frac{n^2}{N^2}))$. It is not known whether the implicit constant in the upper bound of (2) goes to infinity with N except in the case that $p = 6$ and $s = 2$, when the same example $F(x, t) = \sum_{n=1}^N e((x, t) \cdot (\frac{n}{N}, \frac{n^2}{N^2}))$ shows that $D(N, N^2, 6) \gtrsim (\log N)$ [Bou93]. Roughly, the argument is that for each dyadic value $\alpha \in [N^{3/4}, N]$, one can show by counting the ‘‘major arcs’’ that

$$\alpha^6 \{(x, t) \in Q_{N^2} : |F(x, t)| \sim \alpha\} \gtrsim N^4 \cdot N^3.$$

Since there are $\sim \log N$ values of α , the lower bound for $\int_{Q_{N^2}} |F|^6$ follows. Theorem 1 implies that the corresponding superlevel set estimates (3) are not sharp for $1 \leq s < 2$, unless $\alpha \sim N$ or $\alpha^2 \sim N$, which leads to the following conjecture.

Conjecture 2. *Let $s \in [1, 2)$ and $2 \leq p \leq 2 + 2s$. There exists $C(s) > 0$ so that*

$$D(N, N^s, p) \leq C(s).$$

A more refined version of Theorem 1 leads to the following essentially sharp (ℓ^q, L^p) small cap decoupling theorem, stated here for general exponential sums.

Corollary 1. *Let $\frac{3}{p} + \frac{1}{q} \leq 1$, and let $R \in [N, N^2]$. Then for each $\varepsilon > 0$, there exists $C_\varepsilon < \infty$ so that*

$$\left\| \sum_{\xi \in \Xi} a_\xi e((x, t) \cdot (\xi, \xi^2)) \right\|_{L^p(B_R)} \leq C_\varepsilon N^\varepsilon (N^{1-\frac{1}{p}-\frac{1}{q}} R^{\frac{1}{p}} + N^{\frac{1}{2}-\frac{1}{q}} R^{\frac{2}{p}}) \left(\sum_{\xi} |a_\xi|^q \right)^{1/q}.$$

In the above corollary, the assumptions are that Ξ is a $\gtrsim \frac{1}{N}$ -separated subset of $[-1, 1]$ and that $a_\xi \in \mathbb{C}$.

1. MAIN RESULTS

We state our main results in the more general set-up for decoupling. Let \mathbb{P}^1 denote the truncated parabola

$$\{(t, t^2) : |t| \leq 1\}$$

and write $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ for the R^{-1} -neighborhood of \mathbb{P}^1 in \mathbb{R}^2 , where $R \geq 2$. For a partition $\{\gamma\}$ of $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ into almost rectangular blocks, an (ℓ^2, L^p) decoupling inequality is

$$(4) \quad \|f\|_{L^p(B_R)} \leq D(R, p) \left(\sum_{\gamma} \|f_\gamma\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2}$$

in which $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ is a Schwartz function with $\text{supp } \hat{f} \subset \mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ and f_γ means the Fourier projection onto γ , defined precisely below. When we refer to canonical caps or to

canonical decoupling, we mean that γ are approximately $R^{-1/2} \times R^{-1}$ blocks corresponding to the ℓ^2 -decoupling paper of [BD15]. In this paper, we allow γ to be approximate $R^{-\beta} \times R^{-1}$ blocks, where $\frac{1}{2} \leq \beta \leq 1$. This is the “small cap” regime studied in [DGW20]. We also consider (ℓ^q, L^p) decoupling for small caps, which replaces $(\sum_{\gamma} \|f_{\gamma}\|_p^2)^{1/2}$ by $(\sum_{\gamma} \|f_{\gamma}\|_p^q)^{1/q}$ in the decoupling inequality above (see Corollary 5).

To precisely discuss the collection $\{\gamma\}$, fix a $\beta \in [\frac{1}{2}, 1]$. Let $\mathcal{P} = \mathcal{P}(R, \beta) = \{\gamma\}$ be the partition of $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ given by

$$(5) \quad \bigsqcup_{|k| \leq \lceil R^{\beta} \rceil - 2} \{(x, t) \in \mathcal{N}_{R^{-1}}(\mathbb{P}^1) : k \lceil R^{\beta} \rceil^{-1} \leq x < (k+1) \lceil R^{\beta} \rceil^{-1}\}$$

and the two end pieces

$$\{(x, t) \in \mathcal{N}_{R^{-1}}(\mathbb{P}^1) : x < -1 + \lceil R^{\beta} \rceil^{-1}\} \sqcup \{(x, t) \in \mathcal{N}_{R^{-1}}(\mathbb{P}^1) : 1 - \lceil R^{\beta} \rceil^{-1} \leq x\}.$$

For a Schwartz function $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ with $\text{supp} \hat{f} \subset \mathcal{N}_{R^{-1}}(\mathbb{P}^1)$, define for each $\gamma \in \mathcal{P}(R, \beta)$

$$f_{\gamma}(x) := \int_{\gamma} \hat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi.$$

For $a, b > 0$, the notation $a \lesssim b$ means that $a \leq Cb$ where $C > 0$ is a universal constant whose definition varies from line to line, but which only depends on fixed parameters of the problem. Also, $a \sim b$ means $C^{-1}b \leq a \leq Cb$ for a universal constant C .

Let $U_{\alpha} := \{x \in \mathbb{R}^2 : |f(x)| \geq \alpha\}$. In Section 5 of [GMW20], through a wave packet decomposition and series of pigeonholing steps, bounds for $D(R, p)$ in (4) follow (with an additional power of $(\log R)$) from bounds on the constant $C(R, p)$ in

$$\alpha^p |U_{\alpha}| \leq C(R, p) (\#\{\gamma : f_{\gamma} \neq 0\})^{\frac{p}{2}-1} \sum_{\gamma} \|f_{\gamma}\|_2^2$$

for any $\alpha > 0$ and under the additional assumptions that $\|f_{\gamma}\|_{\infty} \lesssim 1$, $\|f_{\gamma}\|_p^p \sim \|f_{\gamma}\|_2^2$ for each γ . Thus decoupling bounds follow from upper bounds on the superlevel set $|U_{\alpha}|$. In this paper, we consider the question: given $\alpha > 0$ and a partition $\{\gamma\}$, how large can $|U_{\alpha}|$ be, varying over functions f satisfying $\|f_{\gamma}\|_{\infty} \lesssim 1$ for each γ ? We answer this question in the following theorem.

Theorem 3. *Let $\beta \in [\frac{1}{2}, 1]$, $R \geq 2$. Let $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ be a Schwartz function with Fourier transform supported in $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ satisfying $\|f_{\gamma}\|_{\infty} \leq 1$ for all $\gamma \in \mathcal{P}(R, \beta)$. Then for any $\alpha > 0$,*

$$|U_{\alpha} \cap [-R, R]^2| \leq C_{\varepsilon} R^{\varepsilon} \begin{cases} \frac{R^{2\beta-1}}{\alpha^4} \sum_{\gamma} \|f_{\gamma}\|_{L^2(\mathbb{R}^2)}^2 & \text{if } \alpha^2 > R \\ \frac{R^{2\beta}}{\alpha^6} \sum_{\gamma} \|f_{\gamma}\|_{L^2(\mathbb{R}^2)}^2 & \text{if } R^{\beta} \leq \alpha^2 \leq R \\ R^2 & \text{if } \alpha^2 < R^{\beta}. \end{cases}$$

Each bound in Theorem 3 is sharp, up to the $C_{\varepsilon} R^{\varepsilon}$ factor, which we show in §2.

Define notation for a distribution function for the Fourier support of a Schwartz function f with Fourier transform supported in $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ as follows. For each $0 \leq s \leq 2$, let

$$\lambda(s) = \sup_{\omega(s)} \#\{\gamma : \gamma \cap \omega(s) \neq \emptyset, f_{\gamma} \neq 0\}$$

where $\omega(s)$ is any arc of \mathbb{P}^1 with projection onto the ξ_1 -axis equal to an interval of length s . The following theorem implies Theorem 3 and replaces factors of R^{β} in the upper bounds

from Theorem 3 by expressions involving $\lambda(\cdot)$, which see the actual Fourier support of the input function f .

Theorem 4. *Let $\beta \in [\frac{1}{2}, 1]$, $R \geq 2$. For any f with Fourier transform supported in $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ satisfying $\|f_\gamma\|_\infty \lesssim 1$ for each $\gamma \in \mathcal{P}(R, \beta)$,*

$$|U_\alpha| \leq C_\varepsilon R^\varepsilon \begin{cases} \frac{1}{\alpha^4} \max_s \lambda(s^{-1} R^{-1}) \lambda(s) \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha^2 > \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)} \\ \frac{\lambda(1)^2}{\alpha^6} \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha^2 \leq \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)} \end{cases}$$

in which the maxima are taken over dyadic s , $R^{-\beta} \leq s \leq R^{-1/2}$.

Corollary 5 ((l^q, L^p) small cap decoupling). *Let $\frac{3}{p} + \frac{1}{q} \leq 1$. Then*

$$\|f\|_{L^p(B_R)} \leq C_\varepsilon R^\varepsilon (R^{\beta(1-\frac{1}{q})-\frac{1}{p}(1+\beta)} + R^{\beta(\frac{1}{2}-\frac{1}{q})}) (\sum_\gamma \|f_\gamma\|_{L^p(\mathbb{R}^2)}^q)^{1/q}$$

whenever f is a Schwartz function with Fourier transform supported in $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$.

The powers of R in the upper bound come from considering two natural sharp examples for the ratio $\|f\|_{L^p(B_R)}^p / (\sum_\gamma \|f_\gamma\|_p^q)^{p/q}$. The first is the square root cancellation example, where $|f_\gamma| \sim \chi_{B_R}$ for all γ and $f = \sum_\gamma e_\gamma f_\gamma$ where e_γ are ± 1 signs chosen (using Khintchine's inequality) so that $\|f\|_{L^p(B_R)}^p \sim R^{\beta p/2} R^2$.

$$\|f\|_p^p / (\sum_\gamma \|f_\gamma\|_p^q)^{p/q} \gtrsim (R^{\beta p/2} R^2) / (R^{\beta p/q} R^2) \sim R^{\beta p(\frac{1}{2}-\frac{1}{q})}.$$

The second example is the constructive interference example. Let $f_\gamma = R^{1+\beta} \check{\eta}_\gamma$ where η_γ is a smooth bump function approximating χ_γ . Since $|f| = |\sum_\gamma f_\gamma|$ is approximately constant on unit balls and $|f(0)| \sim R^\beta$, we have

$$\|f\|_p^p / (\sum_\gamma \|f_\gamma\|_p^q)^{p/q} \gtrsim (R^{\beta p}) / (R^{\beta p/q} R^{1+\beta}) \sim R^{\beta p(1-\frac{1}{q})-1-\beta}.$$

There is one more example which may dominate the ratio: The block example is $f = R^{1+\beta} \sum_{\gamma \subset \theta} \check{\eta}_\gamma$ where θ is a canonical $R^{-1/2} \times R^{-1}$ block. Since $f = f_\theta$ and $|f_\theta|$ is approximately constant on dual $\sim R^{1/2} \times R$ blocks θ^* , we have

$$\frac{\alpha^p |U_\alpha|}{(\#\gamma)^{\frac{p}{q}} \|f_\gamma\|_2^2} \gtrsim \frac{R^{(\beta-\frac{1}{2})p} R^{\frac{3}{2}}}{R^{(\beta-\frac{1}{2})\frac{p}{q} R^{1+\beta}}} = R^{(\beta-\frac{1}{2})p(1-\frac{1}{q})+\frac{1}{2}-\beta}.$$

One may check that the constructive interference examples dominate the block example when $\frac{3}{p} + \frac{1}{q} \leq 1$. We do not investigate (l^q, L^p) small cap decoupling in the range $\frac{3}{p} + \frac{1}{q} > 1$ in the present paper.

The paper is organized as follows. In §2, we demonstrate that Theorem 3 is sharp using an exponential sum example. In §3, we show how Theorem 3 follows easily from Theorem 4 and how after some pigeonholing steps, so does Corollary 5. Then in §4, we develop the multi-scale high/low frequency tools we use in the proof of Theorem 4. These tools are very similar to those developed in [GMW20]. It appears that a more careful version of the proof of Theorem 4 could also replace the $C_\varepsilon R^\varepsilon$ factor by a power of $(\log R)$, as is done for canonical decoupling in [GMW20]. Finally, in §5, we prove a bilinear version of Theorem 4 and then reduce to the bilinear case to finish the proof.

LG is supported by a Simons Investigator grant. DM is supported by the National Science Foundation under Award No. 2103249.

2. A SHARP EXAMPLE

Because we will show that Theorem 4 implies Theorem 3, it suffices to show that Theorem 3 is sharp, which we mean up to a $C_\varepsilon R^\varepsilon$ factor. Write $N = \lceil R^\beta \rceil$. The function achieving the sharp bounds is

$$F(x_1, x_2) = \sum_{k=1}^N e\left(\frac{k}{N}x_1 + \frac{k^2}{N^2}x_2\right)\eta(x_1, x_2),$$

where η is a Schwartz function satisfying $\eta \sim 1$ on $[-R, R]^2$ and $\text{supp } \widehat{\eta} \subset B_{R^{-1}}$. We will bound the set

$$U_\alpha = \{(x_1, x_2) \in [-R, R]^2 : |F(x_1, x_2)| \geq \alpha\}.$$

Case 1: $R < \alpha^2$.

Suppose that $\alpha \sim N$ and note that $F(0, 0) = N$ and $|F(0, 0)| \sim N$ when $|(x_1, x_2)| < \frac{1}{10^3}$. Using periodicity in the x_1 variable, there are $\sim R/N$ many other heavy balls where $|F(x)| \sim N$ in $[-R, R]^2$. For α in the range suppose that $R < \alpha^2 < N^2$, we will show that U_α is dominated by larger neighborhoods of the heavy balls.

Let $r = N^2/\alpha^2$ and assume without loss of generality that r is in the range $R^\varepsilon < r < N^2/R \sim R^{2\beta-1} \ll N$. The upper bound for $|U_\alpha|$ in Theorem 3 for this range is

$$|U_\alpha| \leq C_\varepsilon R^\varepsilon \frac{N^2}{\alpha^4 R} \sum_\gamma \|F_\gamma\|_2^2 \sim C_\varepsilon R^\varepsilon \frac{N^2}{\alpha^4 R} N R^2.$$

To demonstrate that this inequality is sharp, by periodicity in x_1 , it suffices to show that $|U_\alpha \cap B_r| \gtrsim r^2$. Let $\phi_{r^{-1}}$ be a nonnegative bump function supported in $B_{r^{-1}/2}$ with $\phi_{r^{-1}} \gtrsim 1$ on $B_{r^{-1}/4}$. Let $\eta_r = r^4(\phi_{r^{-1}} * \phi_{r^{-1}})^\vee$ and analyze the L^2 norm $\|F\|_{L^2(\eta_r)}$. By Plancherel's,

$$\begin{aligned} \|F\|_{L^2(\eta_r)}^2 &= \int |F|^2 \eta_r \sim \int \left| \sum_{k=1}^N e\left(\frac{k}{N}x_1 + \frac{k^2}{N^2}x_2\right) \right|^2 \eta_r(x_1, x_2) \\ &= \sum_{k=1}^N \sum_{k'=1}^N \widehat{\eta}_r\left(\xi\left(\frac{k-k'}{N}, \frac{k^2-(k')^2}{N^2}\right)\right) \sim N \cdot N/r \cdot r^2 = rN^2. \end{aligned}$$

Next we bound $\|F\|_{L^4(B_{R^\varepsilon r})}$ above. It follows from the local linear restriction statement (see [Dem20] Theorem 1.14, Prop 1.27, and Exercise 1.32)

$$\|f\|_{L^4(B_{R^\varepsilon r})}^4 \lesssim C_\varepsilon R^{O(\varepsilon)} r^{-3} \|\widehat{f}\|_{L^4(\mathbb{R}^2)}^4$$

that

$$\begin{aligned} \|F\|_{L^4(B_{R^\varepsilon r})}^4 &\sim \left\| \sum_{k=1}^N e\left(\frac{k}{N}x_1 + \frac{k^2}{N^2}x_2\right) \eta_r(x_1, x_2) \right\|_{L^4(B_{R^\varepsilon r})}^4 \\ &\lesssim C_\varepsilon R^\varepsilon r^{-3} \left\| \sum_{k=1}^N \widehat{\eta}_r\left(\xi - \left(\frac{k}{N}, \frac{k^2}{N^2}\right)\right) \right\|_{L^4(\mathbb{R}^2)}^4. \end{aligned}$$

The L^4 norm on the right hand side is bounded above by

$$\begin{aligned} \int_{B_2} \left| \sum_{k=1}^N \widehat{\eta}_r \left(\xi - \left(\frac{k}{N}, \frac{k^2}{N^2} \right) \right) \right|^4 d\xi &\lesssim (Nr^{-1})^3 \int_{B_2} \sum_{k=1}^N |\widehat{\eta}_r \left(\xi - \left(\frac{k}{N}, \frac{k^2}{N^2} \right) \right)|^4 d\xi \\ &\lesssim (Nr^{-1})^3 (r^2)^3 \int_{B_2} \sum_{k=1}^N |\widehat{\eta}_r \left(\xi - \left(\frac{k}{N}, \frac{k^2}{N^2} \right) \right)| d\xi \sim N^4 r^3. \end{aligned}$$

This leads to the upper bound $\|F\|_{L^4(B_{R^{\varepsilon r}})}^4 \lesssim (\log R) N^4$.

Finally, by dyadic pigeonholing, there is some $\lambda \in [R^{-1000}, N]$ so that $\|F\|_{L^2(\eta_r)}^2 \lesssim (\log R) \lambda^2 |\{x \in B_{R^{\varepsilon r}} : |F(x)| \sim \lambda\}| + C_\varepsilon R^{-2000}$. The lower bound for $\|F\|_{L^2(\eta_r)}^2$ and the upper bound for $\|F\|_{L^4(B_{R^{\varepsilon r}})}^4$ tell us that

$$\begin{aligned} \lambda^2 r N^2 \sim \lambda^2 \|F\|_{L^2(\eta_r)}^2 &\lesssim (\log R) \lambda^4 |\{x \in B_{R^{\varepsilon r}} : |F(x)| \sim \lambda\}| + C_\varepsilon \lambda^4 R^{-2000} \\ &\lesssim (\log R) \|F\|_{L^4(B_{R^{\varepsilon r}})}^4 + C_\varepsilon \lambda^4 R^{-2000} \lesssim C_\varepsilon R^\varepsilon N^4 + C_\varepsilon \lambda^4 R^{-2000}. \end{aligned}$$

Conclude that $\lambda^2 \lesssim C_\varepsilon R^\varepsilon N^2 / r \sim C_\varepsilon R^\varepsilon \alpha^2$. Assuming R is sufficiently large depending on ε ,

$$r N^2 \sim (\log R) \lambda^2 |\{x \in B_{R^{\varepsilon r}} : |F(x)| \sim \lambda\}| \lesssim C_\varepsilon R^\varepsilon (N^2 / r) |\{x \in B_{R^{\varepsilon r}} : |F(x)| \sim \lambda\}|,$$

so $|\{x \in B_{R^{\varepsilon r}} : |F(x)| \sim \lambda\}| \gtrsim C_\varepsilon^{-1} R^{-\varepsilon} r^2$ and $\lambda^2 \gtrsim C_\varepsilon^{-1} R^{-\varepsilon} N^2 / r \sim C_\varepsilon^{-1} R^{-\varepsilon} \alpha^2$.

Case 2: $R^\beta < \alpha^2 \leq R$. Let q, a , and b be integers satisfying

$$(6) \quad q \text{ odd}, \quad 1 \leq b \leq q \leq N^{2/3}, \quad (b, q) = 1, \quad \text{and} \quad 0 \leq a \leq q.$$

Define the set $M(q, a, b)$ to be

$$M(q, a, b) := \{(x_1, x_2) \in [0, N] \times [0, N^2] : |x_1 - \frac{a}{q}N| \leq \frac{1}{10^{10}}, \quad |x_2 - \frac{b}{q}N^2| \leq \frac{1}{10^{10}}\}.$$

Lemma 6. *For each $(q, a, b) \neq (q', a', b')$, both tuples satisfying (6), $M(q, a, b) \cap M(q', a', b') = \emptyset$.*

Proof. If $\frac{b}{q} = \frac{b'}{q'}$, then using the relatively prime part of (6), $b = b'$ and $q = q'$. Then we must have $a \neq a'$, meaning that if x_1 is the first coordinate of a point in $M(q, a, b) \cap M(q, a', b')$, then

$$\frac{2}{10^{10}} \geq |x_1 - \frac{a}{q}N| + |x_1 - \frac{a'}{q}N| \geq \frac{|a - a'|N}{q} \geq N^{1/3}$$

which is clearly a contradiction. The alternative is that $\frac{b}{q} \neq \frac{b'}{q'}$ in which case for x_2 the second coordinate of a point in $M(q, a, b) \cap M(q', a', b')$,

$$\frac{2}{10^{10}} \geq |x_2 - \frac{b}{q}N^2| + |x_2 - \frac{b'}{q'}N^2| \geq \frac{|b'q - bq'|N^2}{qq'} \geq \frac{N^2}{qq'} \geq N^{2/3},$$

which is another contradiction. \square

Lemma 7. *For each $(x_1, x_2) \in M(q, a, b)$, $|F(x_1, x_2)| \sim \frac{N}{q^{1/2}}$, here meaning within a factor of 4.*

Proof. This follows from Proposition 13.4 in [Dem20]. \square

Proposition 8. *Let $R^\beta < \alpha^2 \leq R$ be given. There exists $v \in [0, N^2]$ satisfying*

$$|\{(x_1, x_2) \in [0, R]^2 : |F(x_1, x_2 + v)| \geq \alpha\}| \gtrsim \frac{R^2 N^3}{\alpha^6}.$$

Proof. First note that by N -periodicity in x_1 ,

$$|\{(x_1, x_2) \in [0, R]^2 : |F(x_1, x_2 + v)| \geq \alpha\}| \gtrsim \frac{R}{N} |\{(x_1, x_2) \in ([0, N] \times [0, R]) : |F(x_1, x_2 + v)| \geq \alpha\}|.$$

The function F is N^2 periodic in x_2 , but $R < N^2$ so we need to find $v \in [0, N^2]$ making the set in the lower bound above largest.

By Lemma 7, it suffices to count the tuples (q, a, b) satisfying (6), $q \leq N^2/(16\alpha^2)$, and $|\frac{b}{q}N^2 - v| \leq R$, where v is to be determined. Begin by considering the distribution of points $\frac{b}{q}$ in $[0, 1]$, where $1 \leq b \leq q \sim \frac{N^2}{\alpha^2}$, $(b, q) = 1$. As in the proof of Lemma 6, if $\frac{b}{q} \neq \frac{b'}{q'}$, then $|\frac{b}{q} - \frac{b'}{q'}| \gtrsim \frac{\alpha^2}{N^4}$. Fix b_0, q_0 and consider the set $\{\frac{b}{q} : \frac{b}{q} = \frac{b_0}{q_0}, 1 \leq b \leq q \sim N^2/\alpha^2\}$. Let q_m be maximal such that for some $1 \leq b_m \leq q_m \sim N^2/\alpha^2$ and $(b_m, q_m) = 1$, $\frac{b_m}{q_m} = \frac{b_0}{q_0}$. Then $q_0 = q_m - k$ for some integer k and $b_m(q_m - k) = b_0 q_m$. Rearrange to get $q_m(1 - \frac{b_0}{b_m}) = k$. Thus $q_0 = q_m \frac{b_0}{b_m} \sim N^2/\alpha^2$, which implies that $\frac{b_0}{b_m} \sim 1$. Conclude that there are $\gtrsim \sum_{q \sim N^2/\alpha^2} \varphi(q)$ many unique points $\frac{b}{q}$ in $[0, 1]$ satisfying our prescribed conditions for φ denoting the Euler totient function. Use Theorem 3.7 in [Apo76] to estimate $\sum_{q \sim N^2/\alpha^2} \varphi(q) \sim N^4/\alpha^4$, as long as N/α is larger than some absolute constant. By the pigeonhole principle, there exists some R/N^2 interval $I \subset [0, 1]$ containing $\sim \lceil \frac{N^4}{\alpha^4} \frac{R}{N^2} \rceil$ many points $\frac{b}{q}$ with $1 \leq b \leq q \sim N^2/\alpha^2$, $(b, q) = 1$. There are also $\sim N^2/\alpha^2$ many choices for a to complete the tuple (q, a, b) satisfying (6). Let c denote the center of I and take $v = cN^2$ in the proposition statement and conclude that

$$|\{(x_1, x_2) \in ([0, N] \times [0, R]) : |F(x_1, x_2 + v)| \geq \alpha\}| \gtrsim \frac{RN^4}{\alpha^6}$$

to finish the proof. \square

Note that Proposition 8 shows the sharpness of Theorem 3 in the range $R^\beta < \alpha \leq R$ since

$$\frac{R^{2\beta}}{\alpha^6} \sum_{\gamma} \|F_{\gamma}\|_2^2 \sim \frac{R^{2\beta}}{\alpha^6} R^{\beta} R^2 = \frac{N^3 R^2}{\alpha^6}.$$

The sharpness of the trivial estimate $|U_{\alpha} \cap [-R, R]^2| \lesssim R^2$ in the range $\alpha^2 < R^\beta$ follows from Case 2 since for $\alpha^2 < R^\beta$,

$$|U_{\alpha} \cap [-R, R]^2| \geq |U_{R^{\beta/2}} \cap [-R, R]^2| \gtrsim \frac{R^{2\beta}}{(R^{\beta/2})^6} \sum_{\gamma} \|F_{\gamma}\|_2^2 \sim R^2.$$

3. IMPLICATIONS OF THEOREM 4

Proof of Theorem 3 from Theorem 4. First suppose that $\alpha^2 > \frac{\lambda(1)^2}{\max_s \lambda(s^{-1}R^{-1})\lambda(s)}$. Then

$$\begin{aligned} \max_s \lambda(s^{-1}R^{-1})\lambda(s) &\lesssim \max_s (s^{-1}R^{-1}R^{\beta})(sR^{\beta}) = R^{2\beta-1} \\ &\leq \begin{cases} R^{2\beta-1} & \text{if } \alpha^2 > R \\ \frac{R^{2\beta}}{\alpha^2} & \text{if } R^{\beta} \leq \alpha^2 \leq R \end{cases} \end{aligned}$$

Now suppose that $\alpha^2 \leq \frac{\lambda(1)^2}{\max_s \lambda(s^{-1}R^{-1})\lambda(s)}$. Then

$$\frac{\lambda(1)^2}{\alpha^2} \lesssim \begin{cases} R^{2\beta-1} & \text{if } \alpha^2 > R \\ \frac{R^{2\beta}}{\alpha^2} & \text{if } R^{\beta} \leq \alpha^2 \leq R \end{cases}$$

□

Proof of Corollary 5 from Theorem 4. To see how this corollary follows from Theorem 4, first use an analogous series of pigeonholing steps as in Section 5 of [GMW20] to reduce to the case where $\|f_\gamma\|_\infty \lesssim 1$ for all γ and there exists $C > 0$ so that $\|f_\gamma\|_p^p$ is either 0 or comparable to C for all γ . Split the integral

$$\int |f|^p = \sum_{R^{-1000} \leq \alpha \lesssim R^\beta} \int_{U_\alpha} |f|^p + \int_{|f| < R^{-1000}} |f|^p$$

where $U_\alpha = \{x : |f(x)| \sim \alpha\}$ and assume via dyadic pigeonholing that

$$\int |f|^p \lesssim \alpha^p |U_\alpha|$$

(ignoring the case that the set where $|f| \leq R^{-1000}$ dominates the integral which may be handled trivially). The result of all of the pigeonholing steps is that the statement of Corollary 5 follows from showing that

$$\alpha^p |U_\alpha| \leq C_\varepsilon R^\varepsilon (R^{\beta p(1-\frac{1}{q})-(1+\beta)} + R^{\beta p(\frac{1}{2}-\frac{1}{q})}) \lambda(1)^{\frac{p}{q}-1} \sum_\gamma \|f_\gamma\|_2^2$$

where f satisfies the hypotheses of Theorem 4. The full range $\frac{3}{p} + \frac{1}{q} \leq 1$ follows from p in the critical range $4 \leq p \leq 6$, which we treat first.

$4 \leq p \leq 6$: There are two cases depending on which upper bound is larger in Theorem 4. First we assume the L^4 bound holds, in which case

$$\begin{aligned} \alpha^p |U_\alpha| &\leq C_\varepsilon R^\varepsilon \alpha^{p-4} \max_s \lambda(s^{-1} R^{-1}) \lambda(s) \sum_\gamma \|f_\gamma\|_2^2 \\ &\sim C_\varepsilon R^\varepsilon \frac{\alpha^{p-4}}{\lambda(1)^{\frac{p}{q}-1}} \max_s \lambda(s^{-1} R^{-1}) \lambda(s) \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}} \\ &\lesssim C_\varepsilon R^\varepsilon \frac{\lambda(1)^{p-4}}{\lambda(1)^{\frac{p}{q}-1}} \max_s (R^\beta s^{-1} R^{-1}) (R^\beta s) \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}} \\ &\lesssim C_\varepsilon R^\varepsilon \lambda(1)^{p(1-\frac{1}{q})-3} R^{2\beta-1} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}}. \end{aligned}$$

Since $p(1 - \frac{1}{q}) - 3 \geq 0$, we may use the bound $\lambda(1) \lesssim R^\beta$ to conclude that

$$\lambda(1)^{p(1-\frac{1}{q})-3} R^{2\beta-1} \leq R^{\beta p(1-\frac{1}{q})-3\beta+2\beta-1} = R^{\beta p(1-\frac{1}{q})-(1+\beta)}.$$

The other case is that the L^6 bound holds in Theorem 4. We may also assume that $\alpha^2 > \lambda(1)$ since otherwise we trivially have

$$\alpha^p |U_\alpha| \leq \lambda(1)^{\frac{p}{2}-1} \sum_\gamma \|f_\gamma\|_2^2 \sim \lambda(1)^{\frac{p}{2}-1+1-\frac{p}{q}} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}} \lesssim R^{\beta p(\frac{1}{2}-\frac{1}{q})} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}}$$

where we used that $q \geq 2$ since $4 \leq p \leq 6$ and $\frac{3}{p} + \frac{1}{q} \leq 1$. Now using the assumptions $\alpha^2 > \lambda(1)$ and $p \leq 6$, we have

$$\begin{aligned} \alpha^p |U_\alpha| &\leq C_\varepsilon R^\varepsilon \alpha^{p-6} \lambda(1)^2 \lambda(1)^{1-\frac{p}{q}} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}} \\ &\sim C_\varepsilon R^\varepsilon \lambda(1)^{p(\frac{1}{2}-\frac{1}{q})} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}} \lesssim C_\varepsilon R^\varepsilon R^{\beta p(\frac{1}{2}-\frac{1}{q})} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}}. \end{aligned}$$

$3 \leq p < 4$: Suppose that $\alpha < R^{\beta/2}$. Then using L^2 -orthogonality,

$$\alpha^p |U_\alpha| \leq R^{\frac{\beta}{2}(p-2)} \sum_\gamma \|f_\gamma\|_2^2 \sim R^{\frac{\beta}{2}(p-2)} \lambda(1)^{1-\frac{p}{q}} \left(\sum_\gamma \|f_\gamma\|_p^q \right)^{\frac{p}{q}}.$$

Since in this subcase, $1-\frac{p}{q} \geq 1-(p-3) > 0$, we are done after noting that $R^{\frac{\beta}{2}(p-2)} \lambda(1)^{1-\frac{p}{q}} \leq R^{\beta p(\frac{1}{2}-\frac{1}{q})}$. Now assume that $\alpha \geq R^{\beta/2}$ and use the $p = 4$ case above (noting that $R^{4\beta(1-\frac{1}{q})-(1+\beta)} \leq R^{4\beta(\frac{1}{2}-\frac{1}{q})}$) to get

$$\begin{aligned} \alpha^p |U_\alpha| &\leq \frac{\alpha^4}{(R^{\beta/2})^{4-p}} |U_\alpha| \leq R^{-\frac{\beta}{2}(4-p)} C_\varepsilon R^\varepsilon R^{4\beta(\frac{1}{2}-\frac{1}{q})} \lambda(1)^{\frac{4}{q}-1} \sum_\gamma \|f_\gamma\|_2^2 \\ &\leq C_\varepsilon R^\varepsilon R^{\beta p(\frac{1}{2}-\frac{1}{q})} \lambda(1)^{\frac{p}{q}-1} \sum_\gamma \|f_\gamma\|_2^2. \end{aligned}$$

$6 < p$: In this range, we use the trivial bound $\alpha \leq \lambda(1)$ and the $p = 6$ case above (noting that $R^{6\beta(\frac{1}{2}-\frac{1}{q})} \leq R^{6\beta(1-\frac{1}{q})-(1+\beta)}$) to get

$$\begin{aligned} \alpha^p |U_\alpha| &\leq \lambda(1)^{p-6} \alpha^6 |U_\alpha| \leq \lambda(1)^{p-6} C_\varepsilon R^\varepsilon R^{6\beta(1-\frac{1}{q})-(1+\beta)} \lambda(1)^{\frac{6}{q}-1} \sum_\gamma \|f_\gamma\|_2^2 \\ &= \left(\frac{\lambda(1)}{R^\beta} \right)^{(p-6)(1-\frac{1}{q})} C_\varepsilon R^\varepsilon R^{p\beta(1-\frac{1}{q})-(1+\beta)} \lambda(1)^{\frac{p}{q}-1} \sum_\gamma \|f_\gamma\|_2^2 \\ &\leq C_\varepsilon R^\varepsilon R^{p\beta(1-\frac{1}{q})-(1+\beta)} \lambda(1)^{\frac{p}{q}-1} \sum_\gamma \|f_\gamma\|_2^2. \end{aligned}$$

□

4. TOOLS TO PROVE THEOREM 4

The proof of Theorem 4 follows the high/low frequency decomposition and pruning approach from [GMW20]. In this section, we introduce notation for different scale neighborhoods of \mathbb{P}^1 , a pruning process for wave packets at various scales, some high/low lemmas which are used to analyze the high/low frequency parts of square functions, and a version of a bilinear restriction theorem for \mathbb{P}^1 .

Begin by fixing some notation, as above. Let $\beta \in [\frac{1}{2}, 1]$ and $R \geq 2$. The parameter $\alpha > 0$ describes the superlevel set

$$U_\alpha = \{x \in \mathbb{R}^2 : |f(x)| \geq \alpha\}.$$

For $\varepsilon > 0$, we analyze scales $R_k = R^{k\varepsilon}$, noting that $R^{-1/2} \leq R_k^{-1/2} \leq 1$. Let N distinguish the index so that R_N is closest to R . Since R and R_N differ at most by a factor of R^ε , we will ignore the distinction between R_N and R in the rest of the argument.

Define the following collections, each of which partitions a neighborhood of \mathbb{P} into approximate rectangles.

- (1) $\{\gamma\}$ is a partition of $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ by approximate $R^{-\beta} \times R^{-1}$ rectangles, described explicitly in (5).
- (2) $\{\theta\}$ is a partition of $\mathcal{N}_{R^{-1}}(\mathbb{P}^1)$ by approximate $R^{-1/2} \times R^{-1}$ rectangles. In particular, let each θ be a union of adjacent γ .
- (3) $\{\tau_k\}$ is a partition of $\mathcal{N}_{R_k^{-1}}(\mathbb{P}^1)$ by approximate $R_k^{-1/2} \times R_k^{-1}$ rectangles. Assume the additional property that $\gamma \cap \tau_k = \emptyset$ or $\gamma \subset \tau_k$.

4.1. A pruning step. We will define wave packets at each scale τ_k , and prune the wave packets associated to f_{τ_k} according to their amplitudes.

For each τ_k , fix a dual rectangle τ_k^* which is a $2R_k^{1/2} \times 2R_k$ rectangle centered at the origin and comparable to the convex set

$$\{x \in \mathbb{R}^2 : |x \cdot \xi| \leq 1 \quad \forall \xi \in \tau_k\}.$$

Let \mathbb{T}_{τ_k} be the collection of tubes T_{τ_k} which are dual to τ_k , contain τ_k^* , and which tile \mathbb{R}^2 . Next, we will define an associated partition of unity $\psi_{T_{\tau_k}}$. First let $\varphi(\xi)$ be a bump function supported in $[-\frac{1}{4}, \frac{1}{4}]^2$. For each $m \in \mathbb{Z}^2$, let

$$\psi_m(x) = c \int_{[-\frac{1}{2}, \frac{1}{2}]^2} |\check{\varphi}|^2(x - y - m) dy,$$

where c is chosen so that $\sum_{m \in \mathbb{Z}^2} \psi_m(x) = c \int_{\mathbb{R}^2} |\check{\varphi}|^2 = 1$. Since $|\check{\varphi}|$ is a rapidly decaying function, for any $n \in \mathbb{N}$, there exists $C_n > 0$ such that

$$\psi_m(x) \leq c \int_{[0,1]^2} \frac{C_n}{(1 + |x - y - m|^2)^n} dy \leq \frac{\tilde{C}_n}{(1 + |x - m|^2)^n}.$$

Define the partition of unity $\psi_{T_{\tau_k}}$ associated to τ_k to be $\psi_{T_{\tau_k}}(x) = \psi_m \circ A_{\tau_k}$, where A_{τ_k} is a linear transformation taking τ_k^* to $[-\frac{1}{2}, \frac{1}{2}]^2$ and $A_{\tau_k}(T_{\tau_k}) = m + [-\frac{1}{2}, \frac{1}{2}]^2$. The important properties of $\psi_{T_{\tau_k}}$ are (1) rapid decay off of T_{τ_k} and (2) Fourier support contained in τ_k .

To prove upper bounds for the size of U_α , we will actually bound the sizes of $\sim \varepsilon^{-1}$ many subsets which will be denoted $U_\alpha \cap \Omega_k$, $U_\alpha \cap H$, and $U_\alpha \cap L$. The pruning process sorts between important and unimportant wave packets on each of these subsets, as described in Lemma 12 below.

Partition $\mathbb{T}_\theta = \mathbb{T}_\theta^g \sqcup \mathbb{T}_\theta^b$ into a “good” and a “bad” set as follows. Let $\delta > 0$ be a parameter to be chosen in §5.2 and set

$$T_\theta \in \mathbb{T}_\theta^g \quad \text{if} \quad \|\psi_{T_\theta} f_\theta\|_{L^\infty(\mathbb{R}^2)} \leq R^{M\delta} \frac{\lambda(1)}{\alpha}$$

where $M > 0$ is a universal constant we will choose in the proof of Proposition 1.

Definition 1 (Pruning with respect to τ_k). *For each θ and τ_{N-1} , define the notation $f_\theta^N = \sum_{T_\theta \in \mathbb{T}_\theta^g} \psi_{T_\theta} f_\theta$ and $f_{\tau_{N-1}}^N = \sum_{\theta \subset \tau_{N-1}} f_\theta^N$. For each $k < N$, let*

$$\begin{aligned} \mathbb{T}_{\tau_k}^g &= \{T_{\tau_k} \in \mathbb{T}_{\tau_k} : \|\psi_{T_{\tau_k}} f_{\tau_k}^{k+1}\|_{L^\infty(\mathbb{R}^2)} \leq R^{M\delta} \frac{\lambda(1)}{\alpha}\}, \\ f_{\tau_k}^k &= \sum_{T_{\tau_k} \in \mathbb{T}_{\tau_k}^g} \psi_{T_{\tau_k}} f_{\tau_k}^{k+1} \quad \text{and} \quad f_{\tau_{k-1}}^k = \sum_{\tau_k \subset \tau_{k-1}} f_{\tau_k}^k. \end{aligned}$$

For each k , define the k th version of f to be $f^k = \sum_{\tau_k} f_{\tau_k}^k$.

Lemma 9 (Properties of f^k). (1) $|f_{\tau_k}^k(x)| \leq |f_{\tau_k}^{k+1}(x)| \leq \#\gamma \subset \tau_k$.

(2) $\|f_{\tau_k}^k\|_{L^\infty} \leq C_\varepsilon R^{O(\varepsilon)} R^{M\delta} \frac{\lambda(1)}{\alpha}$.

(3) $\text{supp} \widehat{f_{\tau_k}^k} \subset 2\tau_k$.

(4) $\text{supp} \widehat{f_{\tau_{k-1}}^k} \subset (1 + (\log R)^{-1})\tau_{k-1}$.

Proof. The first property follows because $\sum_{T_{\tau_k} \in \mathbb{T}_{\tau_k}} \psi_{T_{\tau_k}}$ is a partition of unity, and

$$f_{\tau_k}^k = \sum_{T_{\tau_k} \in \mathbb{T}_{\tau_k}^h} \psi_{T_{\tau_k}} f_{\tau_k}^{k+1}.$$

Furthermore, by definition of $f_{\tau_k}^{k+1}$ and iterating, we have

$$\begin{aligned} |f_{\tau_k}^k| &\leq |f_{\tau_k}^{k+1}| \leq \sum_{\tau_{k+1} \subset \tau_k} |f_{\tau_{k+1}}^{k+1}| \leq \dots \leq \sum_{\tau_N \subset \tau_k} |f_{\tau_N}^N| \\ &\leq \sum_{\theta \subset \tau_k} |f_\theta| \leq \sum_{\gamma \subset \tau_k} |f_\gamma| \lesssim \#\gamma \subset \tau_k \end{aligned}$$

where we used the assumption $\|f_\gamma\|_\infty \lesssim 1$ for all γ . Now consider the L^∞ bound in the second property. We write

$$f_{\tau_k}^k(x) = \sum_{\substack{T_{\tau_k} \in \mathbb{T}_{\tau_k}^h, \\ x \in R^\varepsilon T_{\tau_k}}} \psi_{T_{\tau_k}} f_{\tau_k}^{k+1} + \sum_{\substack{T_{\tau_k} \in \mathbb{T}_{\tau_k, \lambda}, \\ x \notin R^\varepsilon T_{\tau_k}}} \psi_{T_{\tau_k}} f_{k+1, \tau_k}.$$

The first sum has at most $CR^{2\varepsilon}$ terms, and each term has norm bounded by $R^{M\delta} \frac{\lambda(1)}{\alpha}$ by the definition of $\mathbb{T}_{\tau_k}^h$. By the first property, we may trivially bound $f_{\tau_k}^{k+1}$ by $R \max_\gamma \|f_\gamma\|_\infty$. But if $x \notin R^\varepsilon T_{\tau_k}$, then $\psi_{T_{\tau_k}}(x) \leq R^{-1000}$. Thus

$$\left| \sum_{\substack{T_{\tau_k} \in \mathbb{T}_{\tau_k}^h, \\ x \notin R^\varepsilon T_{\tau_k}}} \psi_{T_{\tau_k}} f_{\tau_k}^{k+1} \right| \leq \sum_{\substack{T_{\tau_k} \in \mathbb{T}_{\tau_k}^h, \\ x \notin R^\varepsilon T_{\tau_k}}} R^{-500} \psi_{T_{\tau_k}}^{1/2}(x) \|f_{\tau_k}^{k+1}\|_\infty \leq R^{-250} \max_\gamma \|f_\gamma\|_\infty.$$

Since $\alpha \lesssim |f(x)| \lesssim \sum_\gamma \|f_\gamma\|_\infty \lesssim \lambda(1)$, (recalling the assumption that each $\|f_\gamma\|_\infty \lesssim 1$), we note $R^{-250} \leq CR^{2\varepsilon} \frac{\lambda(1)}{\alpha}$.

The fourth and fifth properties depend on the Fourier support of $\psi_{T_{\tau_k}}$, which is contained in $\frac{1}{2}\tau_k$. Initiate a 2-step induction with base case $k = N$: f_θ^N has Fourier support in 2θ because of the above definition. Then

$$f_{\tau_{N-1}}^N = \sum_{\theta \subset \tau_{N-1}} f_\theta^N$$

has Fourier support in $\bigcup_{\theta \subset \tau_{N-1}} 2\theta$, which is contained in $(1 + (\log R)^{-1})\tau_{N-1}$. Since each $\psi_{T_{\tau_{N-1}}}$ has Fourier support in $\frac{1}{2}\tau_{N-1}$,

$$f_{\tau_{N-1}}^{N-1} = \sum_{T_{\tau_{N-1}} \in \mathbb{T}_{\tau_{N-1}, \lambda}} \psi_{T_{\tau_{N-1}}} f_{\tau_{N-1}}^N$$

has Fourier support in $\frac{1}{2}\tau_{N-1} + (1 + (\log R)^{-1})\tau_{N-1} \subset 2\tau_{N-1}$. Iterating this reasoning until $k = 1$ gives (3) and (4).

□

Definition 2. For each τ_k , let w_{τ_k} be the weight function adapted to τ_k^* defined by

$$w_{\tau_k}(x) = w_k \circ R_{\tau_k}(x)$$

where

$$w_k(x, y) = \frac{c}{\left(1 + \frac{|x|^2}{R_k^2}\right)^{10} \left(1 + \frac{|y|^2}{R_k^2}\right)^{10}}, \quad \|w\|_1 = 1,$$

and $R_{\tau_k} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the rotation taking τ_k^* to $[-R_k^{1/2}, R_k^{1/2}] \times [-R_k, R_k]$. For each $T_{\tau_k} \in \mathbb{T}_{\tau_k}$, let $w_{T_{\tau_k}} = w_{\tau_k}(x - c_{T_{\tau_k}})$ where $c_{T_{\tau_k}}$ is the center of T_{τ_k} . For $s > 0$, we also use the notation w_s to mean

$$(7) \quad w_s(x) = \frac{c'}{\left(1 + |x|^2/s^2\right)^{10}}, \quad \|w_s\|_1 = 1.$$

The weights w_{τ_k} , $w_\theta = w_{\tau_N}$, and w_s are useful when we invoke the locally constant property. By locally constant property, we mean generally that if a function f has Fourier transform supported in a convex set A , then for a bump function $\varphi_A \equiv 1$ on A , $f = f * \widetilde{\varphi_A}$. Since $|\widetilde{\varphi_A}|$ is an L^1 -normalized function which is positive on a set dual to A , $|f| * |\widetilde{\varphi_A}|$ is an averaged version of $|f|$ over a dual set A^* . We record some of the specific locally constant properties we need in the following lemma.

Lemma 10 (Locally constant property). For each τ_k and $T_{\tau_k} \in \mathbb{T}_{\tau_k}$,

$$\|f_{\tau_k}\|_{L^\infty(T_{\tau_k})}^2 \lesssim |f_{\tau_k}|^2 * w_{\tau_k}(x) \quad \text{for any } x \in T_{\tau_k}.$$

For any collection of $\sim s^{-1} \times s^{-2}$ blocks θ_s partitioning $\mathcal{N}_{s^{-2}}(\mathbb{P}^1)$ and any s -ball B ,

$$\left\| \sum_{\theta_s} |f_{\theta_s}|^2 \right\|_{L^\infty(B)} \lesssim \sum_{\theta_s} |f_{\theta_s}|^2 * w_s(x) \quad \text{for any } x \in B.$$

Because the pruned versions of f and f_{τ_k} have essentially the same Fourier supports as the unpruned versions, the locally constant lemma applies to the pruned versions as well.

Proof of Lemma 10. Let ρ_{τ_k} be a bump function equal to 1 on τ_k and supported in $2\tau_k$. Then using Fourier inversion and Hölder's inequality,

$$|f_{\tau_k}(y)|^2 = |f_{\tau_k} * \widetilde{\rho_{\tau_k}}(y)|^2 \leq \|\widetilde{\rho_{\tau_k}}\|_1 |f_{\tau_k}|^2 * |\widetilde{\rho_{\tau_k}}|(y).$$

Since ρ_{τ_k} may be taken to be an affine transformation of a standard bump function adapted to the unit ball, $\|\widetilde{\rho_{\tau_k}}\|_1$ is a constant. The function $\widetilde{\rho_{\tau_k}}$ decays rapidly off of τ_k^* , so $|\widetilde{\rho_{\tau_k}}| \lesssim w_{\tau_k}$. Since for any $T_{\tau_k} \in \mathbb{T}_{\tau_k}$, $w_{\tau_k}(y)$ is comparable for all $y \in T_{\tau_k}$, we have

$$\begin{aligned} \sup_{x \in T_{\tau_k}} |f_{\tau_k}|^2 * w_{\tau_k}(x) &\leq \int |f_{\tau_k}|^2(y) \sup_{x \in T_{\tau_k}} w_{\tau_k}(x - y) dy \\ &\sim \int |f_{\tau_k}|^2(y) w_{\tau_k}(x - y) dy \quad \text{for all } x \in T_{\tau_k}. \end{aligned}$$

For the second part of the lemma, repeat analogous steps as above, except begin with ρ_{θ_s} which is identically 1 on a ball of radius $2s^{-1}$ containing θ_s . Then

$$\sum_{\theta_s} |f_{\theta_s}(y)|^2 = \sum_{\theta_s} |f_{\theta_s} * \widetilde{\rho_{\theta_s}}(y)|^2 \lesssim \sum_{\theta_s} |f_{\theta_s}|^2 * |\widetilde{\rho_{\theta_s}}|(y),$$

where we used that each ρ_{θ_s} is a translate of a single function $\rho_{s^{-1}}$. The rest of the argument is analogous to the first part. □

Definition 3 (Auxiliary functions). *Let $\varphi(x) : \mathbb{R}^2 \rightarrow [0, \infty)$ be a radial, smooth bump function satisfying $\varphi(x) = 1$ on B_1 and $\text{supp}\varphi \subset B_2$.*

$$\varphi(2^{J+1}\xi) + \sum_{j=-2}^J [\varphi(2^j\xi) - \varphi(2^{j+1}\xi)]$$

where J is defined by $2^J \leq [R^\beta] < 2^{J+1}$. Then for each dyadic $s = 2^j$, let

$$\eta_{\sim s}(\xi) = \varphi(2^j\xi) - \varphi(2^{j+1}\xi)$$

and let

$$\eta_{< [R^\beta]-1}(\xi) = \varphi(2^{J+1}\xi).$$

Finally, for $k = 1, \dots, N-1$, define

$$\eta_k(\xi) = \varphi(R_{k+1}^{1/2}x).$$

Definition 4. *Let $G(x) = \sum_\theta |f_\theta|^2 * w_\theta$, $G^\ell(x) = G * \check{\eta}_{< [R^\beta]-1}$, $G^h(x) = G(x) - G^\ell(x)$. For $k = 1, \dots, N-1$, let*

$$g_k(x) = \sum_{\tau_k} |f_{\tau_k}^{k+1}|^2 * w_{\tau_k}, \quad g_k^\ell(x) = g_k * \check{\eta}_k, \quad \text{and} \quad g_k^h(x) = g_k - g_k^\ell.$$

Definition 5. *Define the high set*

$$H = \{x \in B_R : G(x) \leq 2|G^h(x)|\}.$$

For each $k = 1, \dots, N-1$, let

$$\Omega_k = \{x \in B_R \setminus H : g_k \leq 2|g_k^h|, g_{k+1} \leq 2|g_{k+1}^\ell|, \dots, g_N \leq 2|g_N^\ell|\}$$

and for each $k = 1, \dots, N$. Define the low set

$$L = \{x \in B_R \setminus H : g_1 \leq 2|g_1^\ell|, \dots, g_N \leq 2|g_N^\ell|, G(x) \leq 2|G^\ell(x)|\}.$$

4.2. High/low frequency lemmas.

Lemma 11 (Low lemma). *For each x , $|G^\ell(x)| \lesssim \lambda(1)$ and $|g_k^\ell(x)| \lesssim g_{k+1}(x)$.*

Proof. For each θ , by Plancherel's theorem,

$$\begin{aligned} |f_\theta|^2 * \check{\eta}_{< [R^\beta]-1}(x) &= \int_{\mathbb{R}^2} |f_\theta|^2(x-y) \check{\eta}_{< [R^\beta]-1}(y) dy \\ &= \int_{\mathbb{R}^2} \widehat{f}_\theta * \widehat{f}_\theta(\xi) e^{-2\pi i x \cdot \xi} \eta_{< [R^\beta]-1}(\xi) d\xi \\ &= \sum_{\gamma, \gamma' \subset \theta} \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} \widehat{f}_\gamma * \widehat{f}_{\gamma'}(\xi) \eta_{< [R^\beta]-1}(\xi) d\xi. \end{aligned}$$

The integrand is supported in $(\gamma \setminus \gamma') \cap B_{2[R^\beta]-1}$. This means that the integral vanishes unless γ is within $CR^{-\beta}$ of γ' for some constant $C > 0$, in which case we write $\gamma \sim \gamma'$. Then

$$\sum_{\gamma, \gamma' \subset \theta} \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} \widehat{f}_\gamma * \widehat{f}_{\gamma'}(\xi) \eta_{< [R^\beta]-1}(\xi) d\xi = \sum_{\substack{\gamma, \gamma' \subset \theta \\ \gamma \sim \gamma'}} \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} \widehat{f}_\gamma * \widehat{f}_{\gamma'}(\xi) \eta_{< [R^\beta]-1}(\xi) d\xi$$

Use Plancherel's theorem again to get back to a convolution in x and conclude that

$$\begin{aligned} |G * \check{\eta}_{<[R^\beta]^{-1}}(x)| &= \left| \sum_{\theta} \sum_{\substack{\gamma, \gamma' \subset \theta \\ \gamma \sim \gamma'}} (f_\gamma \bar{f}_{\gamma'}) * w_\theta * \check{\eta}_{<[R^\beta]^{-1}}(x) \right| \\ &\lesssim \sum_{\theta} \sum_{\gamma \subset \theta} |f_\gamma|^2 * w_\theta * |\check{\eta}_{<[R^\beta]^{-1}}(x)| \lesssim \sum_{\gamma} \|f_\gamma\|_\infty^2 \lesssim \lambda(1). \end{aligned}$$

By an analogous argument as above, we have that

$$|g_k^\ell(x)| \lesssim \sum_{\tau_{k+1}} |f_{\tau_{k+1}}^{k+1}|^2 * w_{\tau_k} * |\check{\eta}_k|(x)$$

where for each summand, w_{τ_k} corresponds to the τ_k containing τ_{k+1} . By definition, $|f_{\tau_{k+1}}^{k+1}| \leq |f_{\tau_{k+1}}^k|$. By the locally constant property, $|f_{\tau_{k+1}}^k|^2 \lesssim |f_{\tau_{k+1}}|^2 * w_{\tau_{k+1}}$. It remains to note that

$$w_{\tau_{k+1}} * w_{\tau_k} * |\check{\eta}_k|(x) \lesssim w_{\tau_{k+1}}(x)$$

since $\tau_k^* \subset \tau_{k+1}^*$ and $\check{\eta}_k$ is an L^1 -normalized function that is rapidly decaying away from $B_{R_{k+1}^{1/2}}(0)$. □

Lemma 12 (Pruning lemma). *For any τ ,*

$$\begin{aligned} \left| \sum_{\tau_k \subset \tau} f_{\tau_k} - \sum_{\tau_k \subset \tau} f_{\tau_k}^{k+1}(x) \right| &\leq C_\varepsilon R^{-M\delta} \alpha \quad \text{for all } x \in \Omega_k \\ \text{and} \quad \left| \sum_{\tau_1 \subset \tau} f_{\tau_1} - \sum_{\tau_1 \subset \tau} f_{\tau_1}^1(x) \right| &\leq C_\varepsilon R^{-M\delta} \alpha \quad \text{for all } x \in L. \end{aligned}$$

Proof. By the definition of the pruning process, we have

$$f_\tau = f_\tau^N + (f_\tau - f_\tau^N) = \dots = f_\tau^{k+1}(x) + \sum_{m=k+1}^N (f_\tau^{m+1} - f_\tau^m)$$

with the understanding that $f^{N+1} = f$ and formally, the subscript τ means $f_\tau = \sum_{\gamma \subset \tau} f_\gamma$ and $f_\tau^m = \sum_{\tau_m \subset \tau} f_{\tau_m}^m$. We will show that each difference in the sum is much smaller than α . For each $m \geq k+1$ and τ_m ,

$$\begin{aligned} |f_{\tau_m}^m(x) - f_{\tau_m}^{m+1}(x)| &= \left| \sum_{T_{\tau_m} \in \mathbb{T}_{\tau_m}^b} \psi_{T_{\tau_m}}(x) f_{\tau_m}^{m+1}(x) \right| = \sum_{T_{\tau_m} \in \mathbb{T}_{\tau_m}^b} |\psi_{T_{\tau_m}}^{1/2}(x) f_{\tau_m}^{m+1}(x)| \psi_{T_{\tau_m}}^{1/2}(x) \\ &\lesssim \sum_{T_{\tau_m} \in \mathbb{T}_{\tau_m}^b} R^{-M\delta} \frac{\alpha}{\lambda(1)} \lambda^{-1} \|\psi_{T_{\tau_m}} f_{\tau_m}^{m+1}\|_{L^\infty(\mathbb{R}^2)} \|\psi_{T_{\tau_m}}^{1/2} f_{\tau_m}^{m+1}\|_{L^\infty(\mathbb{R}^2)} \psi_{T_{\tau_m}}^{1/2}(x) \\ &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} \sum_{T_{\tau_m} \in \mathbb{T}_{\tau_m}^b} \|\psi_{T_{\tau_m}}^{1/2} f_{\tau_m}^{m+1}\|_{L^\infty(\mathbb{R}^2)}^2 \psi_{T_{\tau_m}}^{1/2}(x) \\ &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} \sum_{T_{\tau_m} \in \mathbb{T}_{\tau_m}^b} \sum_{\tilde{T}_{\tau_m}} \|\psi_{T_{\tau_m}} |f_{\tau_m}^{m+1}|^2\|_{L^\infty(\tilde{T}_{\tau_m})} \psi_{T_{\tau_m}}^{1/2}(x) \\ &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} \sum_{T_{\tau_m}, \tilde{T}_{\tau_m} \in \mathbb{T}_{\tau_m}} \|\psi_{T_{\tau_m}}\|_{L^\infty(\tilde{T}_{\tau_m})} \| |f_{\tau_m}^{m+1}|^2 \|_{L^\infty(\tilde{T}_{\tau_m})} \psi_{T_{\tau_m}}^{1/2}(x). \end{aligned}$$

Let $c_{\tilde{T}_{\tau_m}}$ denote the center of \tilde{T}_{τ_m} and note the pointwise inequality

$$\sum_{T_{\tau_m}} \|\psi_{T_{\tau_m}}\|_{L^\infty(\tilde{T}_{\tau_m})} \psi_{T_{\tau_m}}^{1/2}(x) \lesssim R_m^{3/2} w_{\tau_m}(x - c_{\tilde{T}_{\tau_m}}),$$

which means that

$$\begin{aligned} |f_{\tau_m}^m(x) - f_{\tau_m}^{m+1}(x)| &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} R_m^{3/2} \sum_{\tilde{T}_{\tau_m} \in T_{\tau_m}} w_{\tau_m}(x - c_{\tilde{T}_{\tau_m}}) \| |f_{\tau_m}^{m+1}|^2 \|_{L^\infty(\tilde{T}_{\tau_m})} \\ &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} R_m^{3/2} \sum_{\tilde{T}_{\tau_m} \in T_{\tau_m}} w_{\tau_m}(x - c_{\tilde{T}_{\tau_m}}) |f_{\tau_m}^{m+1}|^2 * w_{\tau_m}(c_{\tilde{T}_{\tau_m}}) \\ &\lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} |f_{\tau_m}^{m+1}|^2 * w_{\tau_m}(x). \end{aligned}$$

where we used the locally constant property in the second to last inequality and the pointwise relation $w_{\tau_m} * w_{\tau_m} \lesssim w_{\tau_m}$ for the final inequality. Then

$$\left| \sum_{\tau_m \subset \tau} f_{\tau_m}^m(x) - f_{\tau_m}^{m+1}(x) \right| \lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} \sum_{\tau_m \subset \tau} |f_{\tau_m}^{m+1}|^2 * w_{\tau_m}(x) \lesssim R^{-M\delta} \frac{\alpha}{\lambda(1)} g_m(x).$$

By the definition of Ω_k and Lemma 11, $g_m(x) \leq 2|g_m^\ell(x)| \leq 2Cg_{m+1}(x) \leq \dots \leq (2C)^{\varepsilon^{-1}} G(x) \lesssim (2C)^{\varepsilon^{-1}} r$. Conclude that

$$\left| \sum_{\tau_m \subset \tau} f_{\tau_m}^m(x) - f_{\tau_m}^{m+1}(x) \right| \lesssim (2C)^{\varepsilon^{-1}} R^{-M\delta} \alpha.$$

The claim for L follows immediately from the above argument, using the low-dominance of g_k for all k . \square

Definition 6. Call the distribution function λ associated to a function f (R, ε) -normalized if for any τ_k, τ_m ,

$$\#\{\tau_k \subset \tau_m : f_{\tau_k} \neq 0\} \leq 100 \frac{\lambda(R_m^{-1/2})}{\lambda(R_k^{-1/2})}.$$

Lemma 13 (High lemma I). Assume that f has an (R, ε) -normalized distribution function $\lambda(\cdot)$. For each dyadic s , $R^{-\beta} \leq s \leq R^{-1/2}$,

$$\int_{\mathbb{R}^2} |G * \check{\eta}_{\sim s}|^2 \lesssim C_\varepsilon R^{2\varepsilon} \lambda(s^{-1} R^{-1}) \lambda(s) \sum_\gamma \|f_\gamma\|_2^2.$$

Proof. Organize the $\{\gamma\}$ into subcollections $\{\theta_s\}$ in which each θ_s is a union of γ which intersect the same $\sim s$ -arc of \mathbb{P}^1 , where here for concreteness, $\sim s$ means within a factor of 2. Then by Plancherel's theorem, since $\check{\eta}_{\sim s} = \check{\eta}_{\sim s}$, we have for each θ

$$\begin{aligned} |f_\theta|^2 * \check{\eta}_{\sim s}(x) &= \int_{\mathbb{R}^2} |f_\theta|^2(x-y) \check{\eta}_{\sim s}(y) dy \\ &= \int_{\mathbb{R}^2} \widehat{f}_\theta * \widehat{f}_\theta(\xi) e^{-2\pi i x \cdot \xi} \eta_{\sim s}(\xi) d\xi \\ (8) \quad &= \sum_{\theta_s, \theta'_s \subset \theta} \int_{\mathbb{R}^2} e^{-2\pi i x \cdot \xi} \widehat{f}_{\theta_s} * \widehat{f}_{\theta'_s}(\xi) \eta_{\sim s}(\xi) d\xi. \end{aligned}$$

The support of $\widehat{f}_{\theta'_s}(\xi) = \int e^{-2\pi i x \cdot \xi} \overline{f_{\theta'_s}}(x) dx = \overline{f_{\theta'_s}}(-\xi)$ is contained in $-\theta'_s$. This means that the support of $\widehat{f}_{\theta_s} * \widehat{f}_{\theta'_s}(\xi)$ is contained in $\theta_s - \theta'_s$. Since the support of $\eta_{\sim s}(\xi)$ is contained in the ball of radius $2s$, for each $\theta_s \subset \theta$, there are only finitely many $\theta'_s \subset \theta$ so that the integral in (8) is nonzero. Thus we may write

$$G * \check{\eta}_{\sim s}(x) = \sum_{\theta} |f_{\theta}|^2 * w_{\theta} * \check{\eta}_{\sim s}(x) = \sum_{\theta} \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (f_{\theta_s} \overline{f_{\theta'_s}}) * w_{\theta} * \check{\eta}_{\sim s}(x).$$

where the second sum is over $\theta_s, \theta'_s \subset \theta$ with $\text{dist}(\theta_s, \theta'_s) < 2s$. Using the above pointwise expression and then Plancherel's theorem, we have

$$\begin{aligned} \int_{\mathbb{R}^2} |G * \check{\eta}_{\sim s}|^2 &= \int_{\mathbb{R}^2} \left| \sum_{\theta} \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (f_{\theta_s} \overline{f_{\theta'_s}}) * w_{\theta} * \check{\eta}_{\sim s} \right|^2 \\ &= \int_{\mathbb{R}^2} \left| \sum_{\theta} \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (\widehat{f_{\theta_s}} * \widehat{f_{\theta'_s}}) \widehat{w_{\theta}} \eta_{\sim s} \right|^2 \end{aligned}$$

For each θ , $\sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (\widehat{f_{\theta_s}} * \widehat{f_{\theta'_s}})$ is supported in $\theta - \theta$, since each summand is supported in $\theta_s - \theta'_s$ and $\theta_s, \theta'_s \subset \theta$. For each $\xi \in \mathbb{R}^2$, $|\xi| > \frac{1}{2}r$, the maximum number of $\theta - \theta$ containing ξ is bounded by the maximum number of θ intersecting an $R^{-1/2} \cdot s^{-1} R^{-1/2}$ -arc of the parabola. Using that $\lambda(\cdot)$ is (R, ε) -normalized, this number is bounded above by $C_{\varepsilon} R^{\varepsilon} \frac{\lambda(s^{-1} R^{-1})}{\lambda(R^{-1/2})}$. Since $\eta_{\sim s}$ is supported in the region $|\xi| > \frac{1}{2}r$, by Cauchy-Schwarz

$$\begin{aligned} \int_{\mathbb{R}^2} \left| \sum_{\theta} \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (\widehat{f_{\theta_s}} * \widehat{f_{\theta'_s}}) \widehat{w_{\theta}} \eta_{\sim s} \right|^2 &\lesssim C_{\varepsilon} R^{\varepsilon} \frac{\lambda(r^{-1} R^{-1})}{\lambda(R^{-1/2})} \sum_{\theta} \int_{\mathbb{R}^2} \left| \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (\widehat{f_{\theta_s}} * \widehat{f_{\theta'_s}}) \widehat{w_{\theta}} \eta_{\sim s} \right|^2 \\ &= C_{\varepsilon} R^{\varepsilon} \frac{\lambda(r^{-1} R^{-1})}{\lambda(R^{-1/2})} \sum_{\theta} \int_{\mathbb{R}^2} \left| \sum_{\substack{\theta_s, \theta'_s \subset \theta \\ \theta_s \sim \theta'_s}} (f_{\theta_s} \overline{f_{\theta'_s}}) * w_{\theta} * \check{\eta}_{\sim s} \right|^2 \\ &\lesssim C_{\varepsilon} R^{\varepsilon} \frac{\lambda(r^{-1} R^{-1})}{\lambda(R^{-1/2})} \sum_{\theta} \int_{\mathbb{R}^2} \left| \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}| \right|^2. \end{aligned}$$

It remains to analyze each of the integrals above:

$$\int_{\mathbb{R}^2} \left| \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}| \right|^2 \lesssim \left\| \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}| \right\|_{\infty} \int_{\mathbb{R}^2} \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}|.$$

Bound the L^{∞} norms using the assumption that $\|f_{\gamma}\|_{\infty} \lesssim 1$ for all γ :

$$\left\| \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}| \right\|_{\infty} \lesssim \sum_{\theta_s \subset \theta} \|f_{\theta_s}\|_{\infty}^2 \lesssim \sum_{\theta_s \subset \theta} \sum_{\gamma \subset \theta_s} \|f_{\gamma}\|_{\infty}^2 \lesssim \lambda(R^{-1/2}) \lambda(s).$$

Finally, using Young's convolution inequality and the L^2 -orthogonality of the f_{γ} , we have

$$\int_{\mathbb{R}^2} \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 * w_{\theta} * |\check{\eta}_{\sim s}| \lesssim \int_{\mathbb{R}^2} \sum_{\theta_s \subset \theta} |f_{\theta_s}|^2 = \sum_{\gamma \subset \theta} \|f_{\gamma}\|_2^2.$$

□

Lemma 14 (High lemma II). *For each k ,*

$$\int_{\mathbb{R}^2} |g_k^h|^2 \lesssim R^{3\varepsilon} \sum_{\tau_k} \int_{\mathbb{R}^2} |f_{\tau_{k+1}}^{k+1}|^4.$$

Proof. By Plancherel's theorem, we have

$$\begin{aligned} \int_{\mathbb{R}^2} |g_k^h|^2 &= \int_{\mathbb{R}^2} |g_k - g_k^\ell|^2 \\ &= \int_{\mathbb{R}^2} \left| \sum_{\tau_k} (\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k} - \sum_{\tau_k} (\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k} \eta_k \right|^2 \\ &\leq \int_{|\xi| > cR_{k+1}^{-1/2}} \left| \sum_{\tau_k} (\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k} \right|^2 \end{aligned}$$

since $(1 - \eta_k)$ is supported in the region $|\xi| > cR_{k+1}^{-1/2}$ for some constant $c > 0$. For each τ_k , $\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}$ is supported in $2\tau_k - 2\tau_k$, using property (4) of Lemma 9. The maximum overlap of the sets $\{2\tau_k - 2\tau_k\}$ in the region $|\xi| \geq cR_{k+1}^{-1/2}$ is bounded by $\sim \frac{R_k^{-1/2}}{R_{k+1}^{-1/2}} \lesssim R^\varepsilon$. Thus using Cauchy-Schwarz,

$$\begin{aligned} \int_{|\xi| > cR_{k+1}^{-1/2}} \left| \sum_{\tau_k} (\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k} \right|^2 &\lesssim R^\varepsilon \sum_{\tau_k} \int_{|\xi| > cR_{k+1}^{-1/2}} |(\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k}|^2 \\ &\leq R^\varepsilon \sum_{\tau_k} \int_{\mathbb{R}^2} |(\widehat{f_{\tau_k}^{k+1}} * \widehat{f_{\tau_k}^{k+1}}) \widehat{w}_{\tau_k}|^2 \\ &= R^\varepsilon \sum_{\tau_k} \int_{\mathbb{R}^2} |f_{\tau_k}^{k+1}|^2 * w_{\tau_k}|^2 \leq R^{3\varepsilon} \sum_{\tau_{k+1}} \int_{\mathbb{R}^2} |f_{\tau_{k+1}}^{k+1}|^4 \end{aligned}$$

where we used Young's inequality with $\|w_{\tau_k}\|_1 \lesssim 1$ and $f_{\tau_k}^{k+1} = \sum_{\tau_{k+1} \subset \tau_k} f_{\tau_{k+1}}^{k+1}$ with Cauchy-Schwarz again in the last line. \square

4.3. Bilinear restriction. We will use the following version of a local bilinear restriction theorem, which follows from a standard Córdoba argument [Cor77] included here for completeness.

Theorem 15. *Let $S \geq 4$, $\frac{1}{2} \geq D \geq S^{-1/2}$, and $X \subset \mathbb{R}^2$ be any Lebesgue measurable set. Suppose that τ and τ' are D -separated subsets of $\mathcal{N}_{S^{-1}}(\mathbb{P}^1)$. Then for a partition $\{\theta_S\}$ of $\mathcal{N}_{S^{-1}}(\mathbb{P}^1)$ into $\sim S^{-1/2} \times S^{-1}$ -blocks, we have*

$$\int_X |f_\tau|^2(x) |f_{\tau'}|^2(x) dx \lesssim D^{-2} \int_{\mathcal{N}_{S^{1/2}}(X)} \left| \sum_{\theta_S} |f_{\theta_S}|^2 * w_{S^{1/2}}(x) \right|^2 dx.$$

In the following proof, the exact definition of the $\sim S^{-1} \times S^{-1}$ blocks θ_S is not important. However, by f_τ and $f_{\tau'}$, we mean more formally $f_\tau = \sum_{\theta_S \cap \tau \neq \emptyset} f_{\theta_S}$ and $f_{\tau'} = \sum_{\theta_S \cap \tau' \neq \emptyset} f_{\theta_S}$.

Proof. Let B be a ball of radius $S^{1/2}$ centered at a point in X . Let φ_B be a smooth function satisfying $\varphi_B \gtrsim 1$ in B , φ_B decays rapidly away from B , and $\widehat{\varphi_B}$ is supported in the $S^{-1/2}$ neighborhood of the origin. Then

$$\int_{X \cap B} |f_\tau|^2 |f_{\tau'}|^2 \lesssim \int_{\mathbb{R}^2} |f_\tau|^2 |f_{\tau'}|^2 \varphi_B.$$

Since S is a fixed parameter and θ_S are fixed $\sim S^{-1/2} \times S^{-1}$ blocks, simplify notation by dropping the S . Expand the squared terms in the integral above to obtain

$$\int_{\mathbb{R}^2} |f_\tau|^2 |f_{\tau'}|^2 \varphi_B = \sum_{\substack{\theta_i \cap \tau \neq \emptyset \\ \theta'_i \cap \tau' \neq \emptyset}} \int_{\mathbb{R}^2} f_{\theta_1} \bar{f}_{\theta_2} f_{\theta'_2} \bar{f}_{\theta'_1} \varphi_B.$$

By Placherel's theorem, each integral vanishes unless

$$(9) \quad (\theta_1 - \theta_2) \cap \mathcal{N}_{S^{-1/2}}(\theta'_1 - \theta'_2) \neq \emptyset.$$

Next we check that the number of tuples $(\theta_1, \theta_2, \theta'_1, \theta'_2)$ (with θ_1, θ_2 having nonempty intersection with τ and θ'_1, θ'_2 having nonempty intersection with τ') satisfying (9) is $O(D^{-1})$. Indeed, suppose that $\xi < \xi' < \xi'' < \xi'''$ satisfy

$$(\xi, \xi^2) \in \theta_1, \quad (\xi', (\xi')^2) \in \theta_2, \quad (\xi'', (\xi'')^2) \in \theta'_1, \quad (\xi''', (\xi''')^2) \in \theta'_2$$

and

$$\xi - \xi' = \xi'' - \xi''' + O(S^{-1/2}).$$

Then by the mean value theorem, $\xi^2 - (\xi')^2 = 2\xi_1(\xi - \xi')$ for some $\xi < \xi_1 < \xi'$ and $(\xi''')^2 - (\xi'')^2 = 2\xi_2(\xi'' - \xi''')$ for some $\xi'' < \xi_2 < \xi'''$. Since $(\xi_1, \xi_1^2) \in \tau$ and $(\xi_2, \xi_2^2) \in \tau'$, we also know that $|\xi_1 - \xi_2| \geq D$. Putting everything together, we have

$$\begin{aligned} |\xi^2 - (\xi')^2 - ((\xi'')^2 - (\xi''')^2)| &= 2|\xi_1(\xi - \xi') - \xi_2(\xi'' - \xi''')| \\ &\geq 2|\xi_1 - \xi_2||\xi - \xi'| - cS^{-1/2} \geq (2C - c)S^{-1/2} \end{aligned}$$

if either $\text{dist}((\xi, \xi^2), (\xi', (\xi')^2))$ or $\text{dist}((\xi'', (\xi'')^2), (\xi''', (\xi''')^2))$ is larger than $CD^{-1}S^{-1/2}$. Thus for a suitably large C , the heights will have difference larger than the allowed $O(S^{-1/2})$ -neighborhood imposed by (9). The conclusion is that

$$\begin{aligned} \sum_{\substack{\theta_i \cap \tau \neq \emptyset \\ \theta'_i \cap \tau' \neq \emptyset}} \int_{\mathbb{R}^2} f_{\theta_1} \bar{f}_{\theta_2} f_{\theta'_2} \bar{f}_{\theta'_1} \varphi_B &= \sum_{\substack{\theta_1 \cap \tau \neq \emptyset \\ \theta'_1 \cap \tau' \neq \emptyset}} \sum_{\substack{d(\theta_1, \theta_2) \leq CD^{-1}S^{-1/2} \\ d(\theta'_1, \theta'_2) \leq CD^{-1}S^{-1/2}}} \int_{\mathbb{R}^2} f_{\theta_1} \bar{f}_{\theta_2} f_{\theta'_2} \bar{f}_{\theta'_1} \varphi_B \\ &\lesssim D^{-2} \int_{\mathbb{R}^2} (\sum_{\theta} |f_\theta|^2)^2 \varphi_B. \end{aligned}$$

Using the locally constant property and summing over a finitely overlapping cover of \mathbb{R}^2 by $S^{1/2}$ -balls B' with centers $c_{B'}$, we have

$$\begin{aligned}
\int_{\mathbb{R}^2} \left(\sum_{\theta} |f_{\theta}|^2 \right)^2 \varphi_B &\leq \sum_{B'} |B| \left\| \sum_{\theta} |f_{\theta}|^2 \right\|_{L^\infty(B')}^2 \|\varphi_B\|_{L^\infty(B')} \\
&\leq |B| \left(\sum_{B'} \left\| \sum_{\theta} |f_{\theta}|^2 \right\|_{L^\infty(B')} \|\varphi_B^{1/2}\|_{L^\infty(B')} \right)^2 \\
&\lesssim |B| \left(\sum_{B'} \sum_{\theta} |f_{\theta}|^2 * w_{S^{1/2}}(c_{B'}) \|\varphi_B^{1/2}\|_{L^\infty(B')} \right)^2 \\
&\lesssim |B|^{-1} \left(\int_{\mathbb{R}^2} \sum_{\theta} |f_{\theta}|^2 * w_{S^{1/2}}(y) \varphi_B^{1/2}(y) dy \right)^2 \\
&\lesssim |B|^{-1} \left(\int_B \sum_{\theta} |f_{\theta}|^2 * w_{S^{1/2}}(y) dy \right)^2 \\
&\leq \int_B \left(\sum_{\theta} |f_{\theta}|^2 * w_{S^{1/2}} \right)^2
\end{aligned}$$

where we used that $w_{S^{1/2}} * \varphi_B^{1/2}(y) \lesssim w_{S^{1/2}} * \chi_B(y)$ in the second to last line. \square

5. PROOF OF THEOREM 4

Theorem 4 follows from the following proposition and a broad-narrow argument in §5.2. First we prove a version of Theorem 4 where U_α is replaced by a “broad” version of U_α .

5.1. The broad version of Theorem 4. Let $\delta > 0$ be a parameter we will choose in the broad/narrow analysis. The notation $\ell(\tau) = s$ means that τ is an approximate $s \times s^2$ block which is part of a partition of $\mathcal{N}_{s^2}(\mathbb{P}^1)$. For two non-adjacent blocks τ, τ' satisfying $\ell(\tau) = \ell(\tau') = R^{-\delta}$, define the broad version of U_α to be

$$(10) \quad \text{Br}_\alpha(\tau, \tau') = \{x \in \mathbb{R}^2 : \alpha \sim |f_\tau(x) f_{\tau'}(x)|^{1/2}, (|f_\tau(x)| + |f_{\tau'}(x)|) \leq R^{O(\delta)} \alpha\}.$$

Proposition 1. *Suppose that f satisfies the hypotheses of Theorem 4 and has an (R, ε) -normalized distribution function $\lambda(\cdot)$. Then*

$$|\text{Br}_\alpha(\tau, \tau')| \leq C_{\varepsilon, \delta} R^\varepsilon R^{O(\delta)} \begin{cases} \frac{1}{\alpha^4} \max_s \lambda(s^{-1} R^{-1}) \lambda(s) \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha^2 > \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)} \\ \frac{\lambda(1)^2}{\alpha^6} \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha^2 \leq \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)} \end{cases}.$$

Proof of Proposition 1. Bounding $|\text{Br}_\alpha(\tau, \tau') \cap H|$: Using bilinear restriction, given here by Theorem 15, we have

$$\alpha^4 |\text{Br}_\alpha(\tau, \tau') \cap H| \lesssim \sum_{\substack{\ell(\tau) = \ell(\tau') = R^{-\delta} \\ d(\tau, \tau') \gtrsim R^{-\delta}}} \int_{U_\alpha \cap H} |f_\tau|^2 |f_{\tau'}|^2 \lesssim R^{O(\delta)} \int_{\mathcal{N}_{R^{1/2}}(\text{Br}_\alpha(\tau, \tau') \cap H)} \left(\sum_{\theta} |f_{\theta}|^2 * w_{R^{1/2}} \right)^2.$$

By the locally constant property and the pointwise inequality $w_{R^{1/2}} * w_\theta \lesssim w_\theta$ for each θ , we have that $\sum_\theta |f_\theta|^2 * w_{R^{1/2}} \lesssim G(x)$. Then

$$(11) \quad \int_{\mathcal{N}_{R^{1/2}}(\text{Br}_\alpha(\tau, \tau') \cap H)} |G(x)|^2 dx \leq \sum_{\substack{Q_{R^{1/2}}: \\ Q_{R^{1/2}} \cap (\text{Br}_\alpha(\tau, \tau') \cap H) \neq \emptyset}} |Q_{R^{1/2}}| \|G\|_{L^\infty(Q_{R^{1/2}} \cap (\text{Br}_\alpha(\tau, \tau') \cap H))}^2$$

For each $x \in H$, $G(x) \leq 2|G^h(x)|$. Also note the equality $G^h(x) = \sum_s G * \check{\eta}_{\sim s}(x)$ where the sum is over dyadic s in the range $[R^\beta]^{-1} \lesssim s \lesssim R^{-1/2}$. This is because the Fourier support of G^h is contained in $\cup_\theta(\theta - \theta) \setminus B_{c[R^\beta]^{-1}}$ for a sufficiently small $c > 0$. By dyadic pigeonholing, there is some dyadic s , $[R^\beta]^{-1} \lesssim s \lesssim R^{-1/2}$, so that the upper bound in (11) is bounded by

$$(\log R) \sum_{\substack{Q_{R^{1/2}}: \\ Q_{R^{1/2}} \cap (\text{Br}_\alpha(\tau, \tau') \cap H) \neq \emptyset}} |Q_{R^{1/2}}| \|G * \check{\eta}_{\sim s}\|_{L^\infty(Q_{R^{1/2}} \cap (\text{Br}_\alpha(\tau, \tau') \cap H))}^2.$$

By the locally constant property, the above displayed expression is bounded by

$$(\log R) \sum_{\substack{Q_{R^{1/2}}: \\ Q_{R^{1/2}} \cap (\text{Br}_\alpha(\tau, \tau') \cap H) \neq \emptyset}} \int_{\mathbb{R}^2} |G * \check{\eta}_{\sim s}|^2 w_{Q_{R^{1/2}}} \lesssim (\log R) \int_{\mathbb{R}^2} |G * \check{\eta}_{\sim s}|^2.$$

Use Lemma 13 to upper bound the above integral to finish bounding $|\text{Br}_\alpha(\tau, \tau') \cap H|$.

Bounding $|\text{Br}_\alpha(\tau, \tau') \cap \Omega_k|$: First write the trivial inequality

$$\alpha^4 |\text{Br}_\alpha(\tau, \tau') \cap \Omega_k| \leq \sum_{\substack{\ell(\tau) = \ell(\tau') = R^{-\delta} \\ d(\tau, \tau') \gtrsim R^{-\delta}}} \int_{\text{Br}_\alpha(\tau, \tau') \cap \Omega_k \cap \{|f_\tau f_{\tau'}|^{1/2} \sim \alpha\}} |f_\tau|^2 |f_{\tau'}|^2.$$

By the definition of $\text{Br}_\alpha(\tau, \tau') \cap \Omega_k$ and Lemma 12, for each $x \in \text{Br}_\alpha(\tau, \tau') \cap \Omega_k$ we have

$$\begin{aligned} |f_\tau(x) f_{\tau'}(x)| &\leq |f_\tau(x)| |f_{\tau'}(x) - f_{\tau'}^{k+1}(x)| + |f_\tau(x) - f_\tau^{k+1}(x)| |f_{\tau'}^{k+1}(x)| + |f_\tau^{k+1}(x) f_{\tau'}^{k+1}(x)| \\ &\lesssim C_\varepsilon R^{O(\delta)} R^{-M\delta} \alpha^2 + |f_\tau^{k+1}(x) f_{\tau'}^{k+1}(x)|. \end{aligned}$$

For M large enough in the definition of pruning (depending on the implicit universal constant from the broad/narrow analysis which determines the set $\text{Br}_\alpha(\tau, \tau')$) so that $R^{O(\delta)} R^{-M\delta} \leq R^{-\delta}$ and for R large enough depending on ε and δ , we may bound each integral by

$$\int_{\{\text{Br}_\alpha(\tau, \tau') \cap \Omega_k \cap \{|f_\tau f_{\tau'}|^{1/2} \sim \alpha\}\}} |f_\tau|^2 |f_{\tau'}|^2 \lesssim \int_{\text{Br}_\alpha(\tau, \tau') \cap \Omega_k} |f_\tau^{k+1}|^2 |f_{\tau'}^{k+1}|^2.$$

Repeat analogous bilinear restriction, high-dominated from the definition of Ω_k , and locally-constant steps from the argument bounding $\text{Br}_\alpha(\tau, \tau') \cap H$ to obtain

$$\alpha^4 |\text{Br}_\alpha(\tau, \tau') \cap \Omega_k| \lesssim R^{O(\delta)} \int_{\mathbb{R}^2} |g_k^h|^2.$$

Use Lemma 14 and Lemma 9 to bound the above integral, obtaining

$$\begin{aligned} \alpha^4 |\text{Br}_\alpha(\tau, \tau') \cap \Omega_k| &\lesssim (\log R)^4 \int_{\mathbb{R}^2} |g_k^h|^2 \\ &\lesssim R^{O(\delta)} R^{O(\varepsilon)} \frac{\lambda(1)^2}{\alpha^2} \sum_{\tau_{k+1}} \int_{\mathbb{R}^2} |f_{\tau_{k+1}}^{k+1}|^2. \end{aligned}$$

Use L^2 -orthogonality and that $|f_{\tau_m}^m| \leq |f_{\tau_m}^{m+1}|$ for each m to bound each integral above:

$$\int_{\mathbb{R}^2} |f_{\tau_{k+1}}^{k+1}|^2 \leq \int_{\mathbb{R}^2} |f_{\tau_{k+1}}^{k+2}|^2 \leq C \sum_{\tau_{k+2} \subset \tau_{k+1}} \int_{\mathbb{R}^2} |f_{\tau_{k+2}}^{k+2}|^2 \leq \dots \leq C^{\varepsilon^{-1}} \sum_{\gamma \subset \tau_{k+1}} \int_{\mathbb{R}^2} |f_\gamma|^2.$$

We are done with this case because

$$\frac{\lambda(1)^2}{\alpha^2} \leq \begin{cases} \max_s \lambda(s^{-1}R^{-1})\lambda(s) & \text{if } \alpha^2 > \frac{\lambda(1)^2}{\max_s \lambda(s^{-1}R^{-1})\lambda(s)} \\ \frac{\lambda(1)^2}{\alpha^2} & \text{if } \alpha^2 \leq \frac{\lambda(1)^2}{\max_s \lambda(s^{-1}R^{-1})\lambda(s)} \end{cases}.$$

Bounding $|\text{Br}_\alpha(\tau, \tau') \cap L|$: Repeat the pruning step from the previous case to get

$$\alpha^6 |\text{Br}_\alpha(\tau, \tau') \cap L| \lesssim \sum_{\substack{\ell(\tau)=\ell(\tau')=R^{-\delta} \\ d(\tau, \tau') \gtrsim R^{-\delta}}} \int_{\text{Br}_\alpha(\tau, \tau') \cap L \cap \{|f_\tau f_{\tau'}|^2 \sim \alpha\}} |f_\tau^1 f_{\tau'}^1|^2 |f_\tau f_{\tau'}|.$$

Use Cauchy-Schwartz and the locally constant lemma for the bound $|f_\tau^1 f_{\tau'}^1| \lesssim R^{O(\varepsilon)} G_0$ and recall that by Lemma 11, $G_0 \leq C_\varepsilon R^\varepsilon \lambda(1)$. Then

$$R^{O(\varepsilon)} \sum_{\substack{\ell(\tau)=\ell(\tau')=R^{-\delta} \\ d(\tau, \tau') \gtrsim R^{-\delta}}} \int_{\text{Br}_\alpha(\tau, \tau') \cap L} |G_0|^2 |f_\tau f_{\tau'}| \leq R^{O(\varepsilon)} \lambda(1)^2 \sum_{\ell(\tau)=R^{-\delta}} \int_{\mathbb{R}^2} |f_\tau|^2 \lesssim R^{O(\varepsilon)} \lambda(1)^2 \sum_{\gamma} \|f_\gamma\|_2^2.$$

Using the same upper bound for $\frac{\lambda(1)^2}{\alpha^2}$ as in the previous case finishes the proof. \square

5.2. Bilinear reduction. We will present a broad/narrow analysis to show that Proposition 1 implies Theorem 4. In order to apply Proposition 1, we must reduce to the case that f has an (R, ε) -normalized distribution function $\lambda(\cdot)$. We demonstrate this through a series of pigeonholing steps.

Proposition 1 implies Theorem 4. We will pigeonhole the f_γ so that roughly, for any s -arc ω of the parabola, the number

$$\#\{\gamma : \gamma \cap \omega \neq \emptyset, \quad f_\gamma \neq 0\}$$

is either 0 or relatively constant among s -arcs ω . For the initial step, write

$$\{\tau_N : \exists \gamma \text{ s.t. } f_\gamma \neq 0, \gamma \subset \tau_N\} = \sum_{1 \leq \lambda \lesssim R^\beta R^{-\varepsilon}} \Lambda_N(\lambda)$$

where λ is a dyadic number, $\{\tau_N : \#\gamma \subset \tau_N \sim \lambda\}$, $\#\gamma \subset \tau_N$ means $\#\{\gamma \subset \tau_N : f_\gamma \neq 0\}$, and $\#\gamma \subset \tau_N \sim \lambda$ means $\lambda \leq \#\gamma \subset \tau_N < 2\lambda$. Since there are $\lesssim \log R$ many λ in the sum, there exists some λ_N such that

$$|\{x : |f(x)| > \alpha\}| \leq C(\log R) |\{x : C(\log R) \sum_{\tau_N \in \Lambda_N(\lambda_N)} |f_{\tau_N}(x)| > \alpha\}|.$$

Continuing in this manner, we have

$$\{\tau_k : \exists \tau_{k+1} \in \Lambda_{k+1}(\lambda_{k+1}) \text{ s.t. } \tau_{k+1} \subset \tau_k\} = \sum_{1 \leq \lambda \leq r_k} \Lambda_k(\lambda)$$

where $\Lambda_k(\lambda) = \{\tau_k : \exists \tau_{k+1} \in \Lambda_{k+1}(\lambda_{k+1}) \text{ s.t. } \tau_{k+1} \subset \tau_k \text{ and } \#\gamma \subset \tau_k \sim \lambda\}$ and for some λ_k ,

$$\begin{aligned} & |\{x : (C(\log R))^{N-k} \sum_{\tau_{k+1} \in \Lambda_{k+1}(\lambda_{k+1})} f_{\tau_{k+1}}(x) | > \alpha\}| \\ & \leq C(\log R) |\{x : (C(\log R))^{N-k+1} \sum_{\tau_k \in \Lambda_k(\lambda_k)} f_{\tau_k}(x) | > \alpha\}|. \end{aligned}$$

Continue this process until we have found τ_1, λ_1 so that

$$|\{x : |f(x)| > \alpha\}| \leq C^{\varepsilon^{-1}} (\log R)^{O(\varepsilon^{-1})} |\{x : C^{\varepsilon^{-1}} (\log R)^{O(\varepsilon^{-1})} \sum_{\tau_1 \in \Lambda_1(\lambda_1)} f_{\tau_1}(x) | > \alpha\}|.$$

The function $\sum_{\tau_1 \in \Lambda_1(\lambda_1)} f_{\tau_1}$ now satisfies the hypotheses of Theorem 4 and the property that $\#\gamma \subset \tau_k \sim \lambda_k$ or $\#\gamma \subset \tau_k = 0$ for all k, τ_k . It follows that the associated distribution function $\lambda(\cdot)$ of $\sum_{\tau_1 \in \Lambda_1(\lambda_1)} f_{\tau_1}$ is (R, ε) -normalized since

$$\lambda_m \sim \#\gamma \subset \tau_m = \sum_{\tau_k \subset \tau_m} \#\gamma \subset \tau_k \sim (\#\tau_k \subset \tau_m)(\lambda_k)$$

where we only count the γ or τ_k for which f_γ or f_{τ_k} is nonzero. Now we may apply Proposition 1. Note that since $\log R \leq \varepsilon^{-1} R^\varepsilon$ for all $R \geq 1$, the accumulated constant from this pigeonholing process satisfies $C^{\varepsilon^{-1}} (\log R)^{O(\varepsilon^{-1})} \leq C_\varepsilon R^\varepsilon$. It thus suffices to prove Theorem 4 assuming that f is (R, ε) -normalized.

Now we present a broad-narrow argument adapted to our set-up. Write $K = R^\delta$ for some $\delta > 0$ which will be chosen later. Since $|f(x)| \leq \sum_{\ell(\tau)=K^{-1}} |f_\tau(x)|$, there is a universal constant $C > 0$ so that $|f(x)| > K^C \max_{\substack{\ell(\tau)=\ell(\tau')=K^{-1} \\ \tau, \tau' \text{ nonadj.}}} |f_\tau(x) f_{\tau'}(x)|^{1/2}$ im-

plies $|f(x)| \leq C \max_{\ell(\tau)=K^{-1}} |f_\tau(x)|$. If $|f(x)| \leq K^C \max_{\substack{\ell(\tau)=\ell(\tau')=K^{-1} \\ \tau, \tau' \text{ nonadj.}}} |f_\tau(x) f_{\tau'}(x)|^{1/2}$ and

$K^C \max_{\substack{\ell(\tau)=\ell(\tau')=K^{-1} \\ \tau, \tau' \text{ nonadj.}}} |f_\tau(x) f_{\tau'}(x)|^{1/2} \leq C \max_{\ell(\tau)=K^{-1}} |f_\tau(x)|$, then $|f(x)| \leq C \max_{\ell(\tau)=K^{-1}} |f_\tau(x)|$.

Using this reasoning, we obtain the first step in the broad-narrow inequality

$$\begin{aligned} |f(x)| & \leq C \max_{\ell(\tau)=K^{-1}} |f_\tau(x)| + K^C \max_{\substack{\ell(\tau)=\ell(\tau')=K^{-1} \\ \tau, \tau' \text{ nonadj.}}} |f_\tau(x) f_{\tau'}(x)|^{1/2}. \\ & \leq C \max_{\ell(\tau_0)=K^{-1}} |f_{\tau_0}(x)| \leq K^C |f_\tau(x) f_{\tau'}(x)|^{1/2} \end{aligned}$$

Iterate the inequality m times (for the first term) where $K^m \sim R^{1/2}$ to bound $|f(x)|$ by

$$\begin{aligned} |f(x)| & \lesssim C^m \max_{\ell(\tau)=R^{-1/2}} |f_\tau(x)| \\ & \quad + C^m K^C \sum_{\substack{R^{-1/2} < \Delta < 1 \\ \Delta \in K^{\mathbb{N}}}} \max_{\ell(\tilde{\tau}) \sim \Delta} \max_{\substack{\ell(\tau)=\ell(\tau') \sim K^{-1} \Delta \\ \tau, \tau' \subset \tilde{\tau}, \text{ nonadj.}}} |f_\tau(x) f_{\tau'}(x)|^{1/2}. \\ & \quad \leq C \max_{\substack{\ell(\tau_0)=K^{-1} \Delta \\ \tau_0 \subset \tilde{\tau}}} |f_{\tau_0}(x)| \leq K^C |f_\tau(x) f_{\tau'}(x)|^{1/2} \end{aligned}$$

Recall that our goal is to bound the size of the set

$$U_\alpha = \{x \in \mathbb{R}^2 : \alpha \leq |f(x)|\}.$$

By the triangle inequality and using the notation θ for blocks τ with $\ell(\tau) = R^{-1/2}$

$$(12) \quad |U_\alpha| \leq |\{x \in \mathbb{R}^2 : \alpha \lesssim C^m \max_\theta |f_\theta(x)|\}| + \sum_{\substack{R^{-1/2} < \Delta < 1 \\ \Delta \in K^{\mathbb{N}}}} \sum_{\substack{\ell(\tilde{\tau}) \sim \Delta \\ \ell(\tau) = \ell(\tau') \sim K^{-1}\Delta \\ \tau, \tau' \subset \tilde{\tau}, \text{ nonadj.}}} |U_\alpha(\tau, \tau')|$$

where $U_\alpha(\tau, \tau')$ is the set

$$\{x \in \mathbb{R}^2 : \alpha \lesssim (\log R) C^m K^C |f_\tau(x) f_{\tau'}(x)|^{1/2}, C(|f_\tau(x)| + |f_{\tau'}(x)|) \leq K^C |f_\tau(x) f_{\tau'}(x)|^{1/2}\}.$$

The first term in the upper bound from (12) is bounded trivially by $\frac{\lambda(R^{-1/2})^2}{\alpha^4} \sum_\gamma \|f_\gamma\|_2^2$. By the assumption that $\|f_\gamma\|_\infty \lesssim 1$ for every γ , we know that $|f_\tau| \lesssim R^\beta$ for any τ . Also assume without loss of generality that $\alpha > 1$ (otherwise Theorem 4 follows from L^2 -orthogonality). This means that there are $\sim \log R$ dyadic values of α' between α and R^β so by pigeonholing, there exists $\alpha' \in [\alpha/(C^m K^C), R^\beta]$ so that

$$|U_\alpha(\tau, \tau')| \lesssim (\log R + \log(C^m K^C)) |\text{Br}_{\alpha'}(\tau, \tau')|$$

where the set $\text{Br}_{\alpha'}(\tau, \tau')$ is defined in (10). By parabolic rescaling, there exists an affine transformation T so that $f_\tau \circ T = g_{\underline{\tau}}$ and $f_{\tau'} \circ T = g_{\underline{\tau}'}$ where $\underline{\tau}$ and $\underline{\tau}'$ are $\sim K^{-1}$ -separated blocks in $\mathcal{N}_{\Delta^{-2}R^{-1}}(\mathbb{P}^1)$. Note that the functions $g_{\underline{\tau}}$ and $g_{\underline{\tau}'}$ inherit the property of being $(\Delta^2 R, \varepsilon)$ -normalized in the sense required to apply Proposition 1 in each of the following cases.

Case 1: Suppose that for some $\beta' \in [\frac{1}{2}, 1]$, $\Delta^{-1}R^{-\beta} = (\Delta^2 R)^{-\beta'}$. Then for each $\gamma \in \mathcal{P}(R, \beta)$, $f_\gamma \circ T = g_{\underline{\gamma}}$ for some $\underline{\gamma} \in \mathcal{P}(\Delta^2 R, \beta')$. Applying Proposition 1 with functions $g_{\underline{\tau}}$ and $g_{\underline{\tau}'}$ and level set parameter α' leads to the inequality

$$|\text{Br}_{\alpha'}(\tau, \tau')| \leq K^C \alpha' \} \leq C_{\varepsilon, \delta} R^\varepsilon C^m K^{O(1)} \times \begin{cases} \frac{1}{(\alpha')^4} \max_{R^{-\beta} < s < R^{-1/2}} \lambda(s^{-1}R^{-1}) \lambda(s) \sum_{\gamma \subset \tilde{\tau}} \|f_\gamma\|_2^2 & \text{if } (\alpha')^2 > \frac{\lambda(\Delta)^2}{\max_s \lambda(s^{-1}R^{-1}) \lambda(s)} \\ \frac{\lambda(\Delta)^2}{(\alpha')^6} \sum_{\gamma \subset \tilde{\tau}} \|f_\gamma\|_2^2 & \text{if } (\alpha')^2 \leq \frac{\lambda(\Delta)^2}{\max_s \lambda(s^{-1}R^{-1}) \lambda(s)} \end{cases}$$

Case 2: Now suppose that $\Delta^{-1}R^{-\beta} < (\Delta^2 R)^{-1}$. Let $\tilde{\theta}$ be $\Delta^{-1}R^{-1} \times R^{-1}$ blocks and let $\tilde{\theta}$ be $(\Delta^2 R)^{-1} \times (\Delta^2 R)^{-1}$ blocks so that $f_{\tilde{\theta}} \circ T = g_{\tilde{\theta}}$. Let $B = \max_{\tilde{\theta}} |f_{\tilde{\theta}}|$ and divide everything by B in order to satisfy the hypotheses $\|g_{\tilde{\theta}}\|_\infty / B \leq 1$ for all $\tilde{\theta}$. Let $\tilde{\lambda}(s) := \lambda(\Delta s) / \lambda(\Delta^{-1}R^{-1})$ count the number of $\tilde{\theta}$ intersecting an s -arc. In the case $(\alpha')^2 > \frac{\tilde{\lambda}(1)B^2}{\max_s \tilde{\lambda}(s^{-1}(\Delta^2 R)^{-1}) \tilde{\lambda}(s)}$ (with the maximum taken over $(\Delta^2 R)^{-1} < s < (\Delta^2 R)^{-1/2}$), use Proposition 1 with functions $g_{\underline{\tau}}/B$ and $g_{\underline{\tau}'}/B$ and level set parameter α'/B to get the inequality

$$|\text{Br}_{\alpha'}(\tau, \tau')| \leq C_{\varepsilon, \delta} R^\varepsilon C^m K^{O(1)} \frac{B^4}{(\alpha')^4} \max_{(\Delta^2 R)^{-1} < s < (\Delta^2 R)^{-1/2}} \tilde{\lambda}(s^{-1}(\Delta^2 R)^{-1}) \tilde{\lambda}(s) \sum_{\tilde{\theta} \subset \tilde{\tau}} \|f_{\tilde{\theta}}\|_2^2 / B^2.$$

Note that since $B \leq \lambda(\Delta^{-1}R^{-1})$,

$$B^2 \max_{(\Delta^2 R)^{-1} < s < (\Delta^2 R)^{-1/2}} \tilde{\lambda}(s^{-1}(\Delta^2 R)^{-1}) \tilde{\lambda}(s) \leq \max_{\Delta^{-1}R^{-1} < s < R^{-1/2}} \lambda(s^{-1}R^{-1}) \lambda(s)$$

and

$$\frac{\tilde{\lambda}(1)^2 B^2}{\max_s \tilde{\lambda}(s^{-1}(\Delta^2 R)^{-1}) \tilde{\lambda}(s)} \leq \frac{\lambda(\Delta)^2 \lambda(\Delta^{-1} R^{-1})^2}{\max_{\Delta^{-1} R^{-1} < s < R^{-1/2}} \lambda(s^{-1} R^{-1}) \lambda(s)} \leq \lambda(\Delta^{-1} R^{-1}) \lambda(\Delta).$$

Then in the case $(\alpha')^2 \leq \frac{\tilde{\lambda}(1) B^2}{\max_s \tilde{\lambda}(s^{-1}(\Delta^2 R)^{-1}) \tilde{\lambda}(s)}$, compute directly that

$$\begin{aligned} & (\alpha')^4 |\{x \in \mathbb{R}^2 : \alpha' \sim |f_\tau(x) f_{\tau'}(x)|^{1/2}, (|f_\tau(x)| + |f_{\tau'}(x)|) \leq K^C \alpha'\}| \\ & \lesssim \lambda(\Delta^{-1} R^{-1}) \lambda(\Delta) \int_{\mathbb{R}^2} (|f_\tau|^2 + |f_{\tau'}|^2) \lesssim \max_{\Delta^{-1} R^{-1} < s < R^{-1/2}} \lambda(s^{-1} R^{-1}) \lambda(s) \sum_{\gamma \subset \bar{\tau}} \|f_\gamma\|_2^2. \end{aligned}$$

Using also that $\sum_{\hat{\theta} \subset \bar{\tau}} \|f_{\hat{\theta}}\|_2^2 \leq \sum_{\gamma \subset \bar{\tau}} \|f_\gamma\|_2^2$, the bound for Case 2 is

$$\begin{aligned} & |\{x \in \mathbb{R}^2 : \alpha' \sim |f_\tau(x) f_{\tau'}(x)|^{1/2}, (|f_\tau(x)| + |f_{\tau'}(x)|) \leq K^C \alpha'\}| \\ & \leq C_{\varepsilon, \delta} R^\varepsilon C^m K^{O(1)} \frac{1}{(\alpha')^4} \max_{R^{-\beta} < s < R^{-1/2}} \lambda(s^{-1}(\Delta^2 R)^{-1}) \lambda(s) \sum_{\gamma \subset \bar{\tau}} \|f_\gamma\|_2^2. \end{aligned}$$

It follows from (12) and the combined Case 1 and Case 2 arguments above that

$$|U_\alpha| \leq C_{\varepsilon, \delta} R^\varepsilon C^m K^{O(1)} \times \begin{cases} \frac{1}{\alpha^4} \max_{R^{-\beta} < s < R^{-1/2}} \lambda(s^{-1} R^{-1}) \lambda(s) \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha > \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)} \\ \frac{\lambda(1)^2}{\alpha^6} \sum_\gamma \|f_\gamma\|_2^2 & \text{if } \alpha^2 \leq \frac{\lambda(1)^2}{\max_s \lambda(s^{-1} R^{-1}) \lambda(s)}. \end{cases}$$

Recall that $K^m \sim R^{-1/2}$ and $K = R^\delta$ so that $C_{\varepsilon, \delta} R^\varepsilon C^m K^{O(1)} \leq C_{\varepsilon, \delta} R^\varepsilon C^{O(\delta-1)} R^{O(1)\delta}$. Choosing δ small enough so that $R^{O(1)\delta} \leq R^\varepsilon$ finishes the proof. \square

REFERENCES

- [Apo76] Tom M. Apostol. *Introduction to analytic number theory*. Undergraduate Texts in Mathematics. Springer-Verlag, New York-Heidelberg, 1976.
- [BD15] Jean Bourgain and Ciprian Demeter. The proof of the l^2 decoupling conjecture. *Ann. of Math. (2)*, 182(1):351–389, 2015.
- [Bou93] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations. *Geom. Funct. Anal.*, 3(2):107–156, 1993.
- [Bou17a] J. Bourgain. Decoupling, exponential sums and the Riemann zeta function. *J. Amer. Math. Soc.*, 30(1):205–224, 2017.
- [Bou17b] Jean Bourgain. Decoupling inequalities and some mean-value theorems. *J. Anal. Math.*, 133:313–334, 2017.
- [Cor77] Antonio Cordoba. The Keakeya maximal function and the spherical summation multipliers. *American Journal of Mathematics*, 99(1):1–22, 1977.
- [Dem20] Ciprian Demeter. *Fourier restriction, decoupling, and applications*, volume 184 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2020.
- [DGW20] Ciprian Demeter, Larry Guth, and Hong Wang. Small cap decouplings. *Geom. Funct. Anal.*, 30(4):989–1062, 2020. With an appendix by D. R. Heath-Brown.
- [GMW20] Larry Guth, Dominique Maldague, and Hong Wang. Improved decoupling for the parabola. *arXiv preprint arXiv:2009.07953*, 2020.