

Shadowless rapidly rotating yet not ultraspinning Kerr-AdS₄ and Kerr-Newman-AdS₄ black holes

Di Wu^{1,2,*}, Puxun Wu^{2,†} and Hongwei Yu^{2,‡}

¹*College of Physics and Space Science, China West Normal University,
Nanchong, Sichuan 637002, People's Republic of China*

²*Department of Physics and Synergetic Innovation Center for Quantum Effect and Applications,
Hunan Normal University, Changsha, Hunan 410081, People's Republic of China*

(Dated: December 22, 2021)

We find that the Kerr-(Newman)-AdS₄ black hole will be shadowless if its rotation parameter is larger than a critical value and the shadowless-ness may be related to the appearance of the null hypersurface caustics (NHC) both inside the Cauchy horizon and outside the event horizon for the black hole with the rotation parameter beyond the critical value. Our studies also further confirm that whether an ultraspinning black hole is superentropic or not is unrelated to the existence of the NHC outside the event horizon.

The black hole, as one of the most remarkable and fascinating objects in nature, has always been a subject of extensive studies both theoretically and observationally. On the observational side, recent years have witnessed exciting achievements in “hearing” the gravitational waves from the coalescence of black holes [1–6] as well as “seeing” its shadow by the event horizon telescope [7–14]. On the theoretical side, the study of asymptotically anti de-Sitter (AdS) black holes has shed light on the nature of gravity through gauge-gravity dualities [15–17]. About ten years ago, it was conjectured that an AdS black hole should satisfy the reverse isoperimetric inequality (RII) [18, 19]

$$\mathcal{R} = \left[\frac{(d-1)V}{\mathcal{A}_{d-2}} \right]^{1/(d-1)} \left(\frac{\mathcal{A}_{d-2}}{A} \right)^{1/(d-2)} \geq 1, \quad (1)$$

where V is the thermodynamic volume of the black hole, $\mathcal{A}_{d-2} = 2\pi^{[(d-1)/2]}/\Gamma[(d-1)/2]$ the area of the unit $(d-2)$ -sphere with d being the number of spacetime dimensions and A the area of the outer horizon. Equality is attained for the Schwarzschild-AdS black hole, which means that the Schwarzschild-AdS black hole has the maximum entropy or it occupies the least volume for a specified entropy.

The first counterexample of an AdS black hole, which does not satisfy the RII, is the ultraspinning black hole [20], which is constructed by boosting one of the angular velocities of the rotating AdS black hole to the speed of light [20]. Such constructed ultraspinning black hole has a finite horizon area but noncompact event horizon topology. Since the ultraspinning black hole exceeds the maximum entropy bound [21–23], it is dubbed “superentropic”. Soon afterward, a lot of new ultraspinning AdS black hole solutions [24–30] have been generated successfully from the corresponding known rotating AdS black holes. And various aspects of the ultraspinning black holes, including thermodynamic properties [20, 24–27, 30–34], horizon geometry [21, 24, 25], Kerr/CFT correspondence [25, 26, 35] and geodesic motion [36], etc, have been explored widely. However, not all ultraspinning black holes are superentropic since a few new ultraspinning AdS black holes

[24, 28, 29], i.e. the ultraspinning Kerr-Sen-AdS₄ (dyonic) black holes, violate the RII only in certain parameter spaces of the solutions. Therefore, ultraspinning is not a sufficient condition for a black hole to be superentropic.

So, a question arises naturally as to what the physical condition is for an ultraspinning black hole to be superentropic. Recently, it was suggested that the superentropic-ness may be related to the presence of the null hypersurface caustics (NHC) outside the event horizon of the black hole. This is because the NHC exists outside the event horizon for the (high dimensional) Kerr-(Newman)-AdS superentropic black hole [37, 38]. The existence of the NHC is thought previously to mean that the causal structure of spacetime has some pathologies. However, it was recently found that the spacetime with the NHC is free of closed timelike curve [39].

For the ultraspinning Kerr-Sen-AdS₄ black hole, whether it is superentropic or not, the NHC always appears both out and inside of the horizon [39], which seems to indicate that the existence of the NHC outside the event horizon is related to ultraspinning rather than superentropicity. The ultraspinning means that the rotational angular velocity of the black hole is boosted to the speed of light. Then a further issue, which we are going to address, is whether the rotational angular velocity reaching the speed of light is a necessary requirement for existence of the NHC outside the event horizon.

Let us start with the four-dimensional Kerr-Newman-AdS black hole solution [40], whose metric has the form [41]

$$ds^2 = -\frac{\Delta_r}{\Sigma} \left(dt - \frac{a}{\Xi} \sin^2 \theta d\phi \right)^2 + \frac{\Sigma}{\Delta_r} dr^2 + \frac{\Sigma}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{\Sigma} \left(a dt - \frac{r^2 + a^2}{\Xi} d\phi \right)^2 \quad (2)$$

in the Boyer-Lindquist coordinates, where

$$\Delta_r = (r^2 + a^2)(1 + r^2/l^2) - 2mr + q^2, \quad \Xi = 1 - a^2/l^2, \\ \Delta_\theta = 1 - a^2 \cos^2 \theta / l^2, \quad \Sigma = r^2 + a^2 \cos^2 \theta,$$

a is the rotation parameter, m the mass parameter, q the electric charge parameter and l the AdS radius. The horizon is determined by equation $\Delta_r = 0$. When $q = 0$, the metric in Eq. (2) reduces to that of the Kerr-AdS₄ black hole. The RN-AdS₄ black hole is obtained if $a = 0$ in Eq. (2). In Fig. 1, we plot the rescaled horizon radius \bar{r} as a function of the

* wdcwnu@163.com

† pxwu@hunnu.edu.cn

‡ hwyu@hunnu.edu.cn




FIG. 1. The rescaled mass vs the rescaled horizon radius, where $\bar{m} = m/l$, $\bar{a} = a/l$, $\bar{q} = q/l$, $\bar{r} = r/l$. The red, orange, green, blue, purple and brown lines represent the Schwarzschild-AdS₄, Kerr-AdS₄, ultraspinning Kerr-AdS₄, RN-AdS₄, Kerr-Newman-AdS₄ and ultraspinning Kerr-Newman-AdS₄ black holes, respectively.

scaled mass \bar{m} . It is easy to see that if the black hole mass is larger than a critical value both the RN-AdS₄ black hole and the Kerr-(Newman)-AdS₄ black hole have the Cauchy and event horizons. When the black hole mass equals to the critical value, the Cauchy and event horizons coincide. While the Schwarzschild-AdS₄ black hole, which corresponds to the case with both $a = 0$ and $q = 0$, only has the event horizon.

To derive the condition for existence of the NHC for the Kerr-Newman-AdS₄ black hole, we use the method given in Refs. [37–39, 42, 43]. After introducing the outgoing and ingoing Eddington-Finkelstein coordinates defined in terms of the “generalized tortoise coordinate” $r_*(r, \theta)$: $u = t - r_*(r, \theta)$, $v = t + r_*(r, \theta)$, the null hypersurfaces can be described by $u = \text{const}$, $v = \text{const}$, which are dubbed the outgoing and ingoing null congruences of the hypersurfaces, respectively. It is easy to obtain that the null hypersurfaces defined by $u = \text{const}$ or $v = \text{const}$ satisfy the equation

$$g^{\mu\nu} \partial_\mu(t + r_*) \partial_\nu(t + r_*) = g^{tt} + g^{rr}(\partial_r r_*)^2 + g^{\theta\theta}(\partial_\theta r_*)^2 = 0. \quad (3)$$

By using the contravariant components g^{tt} , g^{rr} , and $g^{\theta\theta}$ of the metric (2), Eq. (3) can be re-expressed as

$$\Delta_r (\partial_r r_*)^2 - \Xi^2 \frac{(r^2 + a^2)^2}{\Delta_r} = -\Delta_\theta (\partial_\theta r_*)^2 - \Xi^2 \frac{a^2 \sin^2 \theta}{\Delta_\theta}. \quad (4)$$

This equation can be solved by the method of separation of variables. Introducing the so-called “constant of separation” $a^2 \lambda$ (hereinafter, λ is referred to as the separation constant) for Eq. (4), we have

$$(\partial_r r_*)^2 = \frac{Q^2(r)}{\Delta_r^2}, \quad (\partial_\theta r_*)^2 = \frac{P^2(\theta)}{\Delta_\theta^2}. \quad (5)$$




FIG. 2. Contours of the NHC, the horizon and the photon sphere in the (\bar{r}, \bar{m}) plane with a fixed \bar{q} for the Kerr-Newman-AdS₄ black hole, where $\bar{m} = m/l$, $\bar{a} = a/l$, $\bar{q} = q/l$, $\bar{r} = r/l$.

where

$$Q^2(r) = \Xi^2 \left[(r^2 + a^2)^2 - a^2 \lambda \Delta_r \right], \quad (6)$$

$$P^2(\theta) = \Xi^2 a^2 (\lambda \Delta_\theta - \sin^2 \theta).$$

In [37, 43], λ is chosen as $\lambda < 1$ and it was then found that the NHC cannot exist outside the event horizon of the Kerr-(Newman)-AdS black hole. However, this choice is incorrect, since when $\theta = \pi/2$, $P^2(\theta) = \Xi^2 a^2 (\lambda - 1) \geq 0$ must be satisfied. Thus, the allowed region of λ should be $\lambda \geq 1$.

After some complicated computations, which are presented in detail in the Appendix, we find that $Q(r) = 0$ is a sufficient condition for the existence of the NHC, which means that

$$(r^2 + a^2)^2 - a^2 \lambda \Delta_r = 0. \quad (7)$$

Figure 2 shows the numerical results of the NHC condition (7). In addition, the photon sphere condition, which is governed by the equation $(r^2 + a^2)\Delta'_r - 4r\Delta_r = 0$ [44, 45], and the horizon condition ($\Delta_r = 0$), are plotted, where a prime indicates the derivative with respect to r . In this figure, two intersection points between the dashed black line and the blue/purple line represent that the black hole has the Cauchy horizon and the event horizon. One can see that the NHC exists only inside the Cauchy horizon when $\bar{a} = 0.25$, and it appears both inside the Cauchy horizon and outside the event horizon of the black hole when $\bar{a} = 0.75$. This suggests that the NHC can appear outside the event horizon of the black hole for a large enough a which is not necessarily equal to l . The photon sphere line indicates that the NHC, which appears outside of the event horizon, locates between the event horizon and the photon sphere of the black hole.

How large is the value of the rotational angular velocity required for existence of the NHC outside the event horizon?

After a detailed calculation, we find that when the rotation parameter a satisfies

$$a \geq \frac{l}{\sqrt{\lambda}}, \quad (8)$$

the NHC exists both outside the event horizon and inside the Cauchy horizon of the Kerr-(Newman)-AdS₄ black hole. Otherwise, there is the NHC only inside the Cauchy horizon. Thus, if the rotation parameter is larger than a critical value $a_c = l/\sqrt{\lambda}$, the NHC appears outside the event horizon. Apparently, this critical value is determined by l and λ . When λ equals unity, the NHC exists outside the event horizon only in the case of the ultraspinning Kerr-(Newman)-AdS₄ black hole ($a \rightarrow l$). If $\lambda > 1$, a large enough a , which is not necessarily equal to l , can lead to the existence of the NHC outside the event horizon of the black hole. Therefore, the value of λ determines how large a has to be to ensure the existence of the NHC outside the event horizon.

Now we demonstrate that when the rotation parameter is larger than a critical value, in addition to the existence of the NHC outside the event horizon, the Kerr-(Newman)-AdS black hole's shadow cannot be obtained and thus it should be shadowless. Here, we take the Kerr-AdS₄ black hole as an example and investigate its shadows. The results are expected to be valid for the Kerr-Newman-AdS₄ case. Figure 3 gives the numerical results in the equatorial plane $\theta = \pi/2$. With the increase of the rotational angular velocity, the black hole shadows change from an ellipse to a "D" shape. Once $a > 0.741l$, the shadow of the Kerr-AdS₄ black hole disappears. Clearly, both the shadowless-ness and the existence of the NHC outside the event horizon of the Kerr-(Newman)-AdS black hole require that the black hole rotates rapidly enough with the rotation parameter larger than a critical value. This seems to suggest that the existence of the HNC outside the event horizon might be the cause of the shadowless-ness of a rapidly rotating Kerr-(Newman)-AdS black hole. Therefore, we can obtain the value of λ from analyzing the shadow of the Kerr-(Newman)-AdS₄ black hole. From the inequality (8), we find that the value of λ should be chosen to be about $\lambda \sim 1.821$ when $\bar{m} = 1$.

It is easy to obtain that when the rotation parameter of the black hole is larger than the critical value a_c , i.e. $a_c \simeq 0.741l$ for the Kerr-AdS₄ black hole with $\bar{m} = 1$, the NHC will appear outside the event horizon of the black hole. Apparently, the ultraspinning limit ($a \rightarrow l$) of the black hole is not a necessary but sufficient condition for the appearance of the NHC outside the event horizon of black hole. Thus, we further confirm that the superentropic-ness of the black hole is unrelated to the presence of the NHC outside the event horizon of the rotating AdS black hole.

To summarize, by investigating the NHC of the Kerr-(Newman)-AdS₄ black hole, we find that there exists a critical value a_c for the rotation parameter a . When the rotation parameter of the black hole is smaller than this critical value, the NHC only exists inside the Cauchy horizon. Once a is larger

than this critical value, the Kerr-(Newman)-AdS₄ black hole will have the NHC both inside the Cauchy horizon and outside the event horizon. Our results demonstrate that the NHC




FIG. 3. The shadows for the Kerr-AdS₄ black hole for different rescaled rotation parameter \bar{a} with $\bar{m} = 1$, where $\bar{m} = m/l$, $\bar{a} = a/l$.

can exist outside the event horizon of the black hole which is not ultraspinning, in sharp contrast to the conclusion that the Kerr-(Newman)-AdS₄ black hole has the NHC outside the event horizon only when it is ultraspinning [37, 43]. The fact that the shadows of the rotating AdS black holes cannot be obtained if the rotation parameter is larger than a critical value seems to suggest that the appearance of the NHC outside the event horizon leads to the disappearance of the black hole shadow. Thus, for the rapidly rotating yet not ultraspinning the Kerr-(Newman)-AdS₄ black holes, we will not see their shadows neither. Our results confirm that the superentropic-ness of ultraspinning black holes is unrelated to the presence of the NHC outside the event horizon. Finally, we believe that our work can be generalized to other rotating AdS black holes, and for these black holes there is also a critical value for the rotation parameter.

ACKNOWLEDGMENTS

We thank Professor Shuang-Qing Wu and Mr. Yuebing Zhou for helpful discussions. This work is supported by the National Natural Science Foundation of China under Grants No. 11675130, No. 11775077, No. 11690034, No. 12075084, by the National Key Research and Development Program of China Grant No. 2020YFC2201502, and by the Science and Technology Innovation Plan of Hunan province under Grant No. 2017XK2019.

-
- [1] B.P. Abbott et al. (LIGO Scientific and Virgo Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [2] B.P. Abbott et al. (LIGO Scientific and Virgo Collaborations), GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence, *Phys. Rev. Lett.* **116**, 241103 (2016).
- [3] B.P. Abbott et al. (LIGO Scientific and Virgo Collaborations), GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, *Phys. Rev. X* **9**, 031040 (2019).
- [4] R. Abbott et al. (LIGO Scientific and Virgo Collaborations), GW190521: A Binary Black Hole Merger with a Total Mass of $150M_{\odot}$, *Phys. Rev. Lett.* **125**, 101102 (2020).
- [5] R. Abbott et al. (LIGO Scientific and Virgo Collaborations), GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run, *Phys. Rev. X* **11**, 021053 (2021).
- [6] R. Abbott et al. (LIGO Scientific and Virgo and KAGRA Collaborations), Observation of gravitational waves from two neutron star-black hole coalescences, *Astrophys. J.* **915**, L5 (2021).
- [7] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole, *Astrophys. J.* **875**, L1 (2019).
- [8] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. II. Array and Instrumentation, *Astrophys. J.* **875**, L2 (2019).
- [9] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. III. Data Processing and Calibration, *Astrophys. J.* **875**, L3 (2019).
- [10] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. IV. Imaging the Central Supermassive Black Hole, *Astrophys. J.* **875**, L4 (2019).
- [11] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. V. Origin of the Asymmetric Ring, *Astrophys. J.* **875**, L5 (2019).
- [12] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. VI. The Shadow and Mass of the Central Black Hole, *Astrophys. J.* **875**, L6 (2019).
- [13] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. VII. Polarization of the Ring, *Astrophys. J.* **910**, L12 (2021).
- [14] The Event Horizon Telescope Collaboration, First M87 Event Horizon Telescope Results. VIII. Magnetic Field Structure near The Event Horizon, *Astrophys. J.* **910**, L13 (2021).
- [15] J.M. Maldacena, The Large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2**, 231 (1998); *Int. J. Theor. Phys.* **38**, 1113 (1999).
- [16] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Gauge theory correlators from noncritical string theory, *Phys. Lett. B* **428**, 105 (1998).
- [17] E. Witten, Anti-de Sitter space and holography, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
- [18] M. Cvetič, G.W. Gibbons, D. Kubizňák, and C.N. Pope, Black hole enthalpy and an entropy inequality for the thermodynamic volume, *Phys. Rev. D* **84**, 024037 (2011).
- [19] B.P. Dolan, D. Kastor, D. Kubizňák, R.B. Mann, and J. Traschen, Thermodynamic volumes and isoperimetric inequalities for de Sitter black holes, *Phys. Rev. D* **87**, 104017 (2013).
- [20] R.A. Hennigar, R.B. Mann, and D. Kubizňák, Entropy Inequality Violations from Ultraspinning Black Holes, *Phys. Rev. Lett.* **115**, 031101 (2015).
- [21] D. Klemm, Four-dimensional black holes with unusual horizons, *Phys. Rev. D* **89**, 084007 (2014).
- [22] A. Gneccchi, K. Hristov, D. Klemm, C. Toldo, and O. Vaughan, Rotating black holes in 4d gauged supergravity, *J. High Energy Phys.* **01** (2014) 127.
- [23] M. Appels, L. Cuspinera, R. Gregory, P. Krtouš, and D. Kubizňák, Are “Superentropic” black holes superentropic?, *J. High Energy Phys.* **02** (2020) 195.
- [24] R.A. Hennigar, D. Kubizňák, R.B. Mann, and N. Musoke, Ultraspinning limits and super-entropic black holes, *J. High Energy Phys.* **06** (2015) 096.
- [25] S.M. Noorbakhsh and M. Ghominejad, Ultra-spinning gauged supergravity black holes and their Kerr/CFT correspondence, *Phys. Rev. D* **95**, 046002 (2017).
- [26] S.M. Noorbakhsh and M. Ghominejad, Higher dimensional charged AdS black holes at ultra-spinning limit and their 2d CFT duals, *arXiv:1702.03448 [hep-th]*.
- [27] S.M. Noorbakhsh and M.H. Vahidinia, Extremal vanishing horizon Kerr-AdS black holes at ultraspinning limit, *J. High Energy Phys.* **01** (2018) 042.
- [28] D. Wu, P. Wu, H. Yu, and S.-Q. Wu, Are ultraspinning Kerr-Sen-AdS₄ black holes always superentropic?, *Phys. Rev. D* **102**, 044007 (2020).
- [29] D. Wu, S.-Q. Wu, P. Wu, and H. Yu, Aspects of the dyonic Kerr-Sen-AdS₄ black hole and its ultraspinning version, *Phys. Rev. D* **103**, 044014 (2021).
- [30] D. Wu, Ultraspinning Chow’s black holes in six-dimensional gauged supergravity and their thermodynamical properties, *arXiv:2106.14218 [hep-th]*.
- [31] C.V. Johnson, Instability of super-entropic black holes in extended thermodynamics, *Mod. Phys. Lett. A* **35**, 2050098 (2020).
- [32] C.V. Johnson, V.L. Martin, and A. Svesko, A microscopic description of thermodynamic volume in extended black hole thermodynamics, *Phys. Rev. D* **101**, 086006 (2020).
- [33] D. Wu, P. Wu, H. Yu, and S.-Q. Wu, Notes on the thermodynamics of superentropic AdS black holes, *Phys. Rev. D* **101**, 024057 (2020).
- [34] Z.M. Xu, The correspondence between thermodynamic curvature and isoperimetric theorem from ultraspinning black hole, *Phys. Lett. B* **807**, 135529 (2020).
- [35] M. Sinamuli and R.B. Mann, Super-entropic black holes and the Kerr-CFT correspondence, *J. High Energy Phys.* **08** (2016) 148.
- [36] K. Flathmann and N. Wassermann, Geodesic equations for particles and light in the black spindle spacetime, *J. Math. Phys.* **61**, 122504 (2020).
- [37] M.T.N. Imseis, A.A. Balushi, and R.B. Mann, Null hypersurfaces in Kerr-Newman-AdS black hole and super-entropic black hole spacetimes, *Classical Quantum Gravity* **38**, 045018 (2021).
- [38] D. Wu and P. Wu, Null hypersurface caustics for high-dimensional superentropic black holes, *Phys. Rev. D* **103**, 104020 (2021).
- [39] S. Noda and Y.C. Ong, Null hypersurface caustics, closed null curves, and super entropy, *Phys. Rev. D* **103**, 024053 (2021).
- [40] B. Carter, Hamilton-Jacobi and Schrodinger separable solutions of Einstein’s equations, *Commun. Math. Phys.* **10**, 280 (1968).
- [41] S.W. Hawking, C.J. Hunter, and M.M. Taylor-Robinson, Rotation and the AdS-CFT correspondence, *Phys. Rev. D* **59**,

064005 (1999).

- [42] F. Pretorius and W. Israel, Quasi-spherical light cones of the Kerr geometry, *Classical Quantum Gravity* **15**, 2289 (1998).
- [43] A.A. Balushi and R.B. Mann, Null hypersurfaces in Kerr-(A)dS spacetimes, *Classical Quantum Gravity* **36**, 245017 (2019).
- [44] A. Belhaj, M. Benali, A. El Balali, W. El Hadri, and H. El Moumni, Shadows of charged and rotating black holes with a cosmological constant, [arXiv:2007.09058 \[gr-qc\]](https://arxiv.org/abs/2007.09058).
- [45] A. Grenzebach, V. Perlick, and C. Lämmerzahl, Photon regions and shadows of Kerr-Newman-NUT black holes with a cosmological constant, *Phys. Rev. D* **89**, 124004 (2014).

Appendix A

Using the relations given in (5), one can express the total differential $dr_* = \partial_r r_* dr + \partial_\theta r_* d\theta$ as

$$dr_* = \frac{Q(r)}{\Delta_r} dr + \frac{P(\theta)}{\Delta_\theta} d\theta. \quad (\text{A1})$$

By treating λ as a variable, the exact differential Eq. (A1) can be generalized to be

$$dr_* = \frac{Q(r, \lambda)}{\Delta_r} dr + \frac{P(\theta, \lambda)}{\Delta_\theta} d\theta + c_1 F(r, \theta, \lambda) d\lambda, \quad (\text{A2})$$

where c_1 is an arbitrary constant and $F(r, \theta, \lambda)$ is an arbitrary function. Since Eqs. (A1) and (A2) are functionally equivalent, the condition

$$F(r, \theta, \lambda) = 0 \quad (\text{A3})$$

should be satisfied, which indicates $dF(r, \theta, \lambda) = 0$. Thus, we arrive at

$$[\partial_\lambda F(r, \theta, \lambda)] d\lambda + [\partial_r F(r, \theta, \lambda)] dr + [\partial_\theta F(r, \theta, \lambda)] d\theta = 0. \quad (\text{A4})$$

From the Poincaré lemma in the external differentiation theory, namely $d(dr_*) = 0$, one can obtain the following integrable conditions

$$\frac{\partial_\lambda Q(r, \lambda)}{\Delta_r} = c_1 \partial_r F(r, \theta, \lambda), \quad \frac{\partial_\lambda P(\theta, \lambda)}{\Delta_\theta} = c_1 \partial_\theta F(r, \theta, \lambda). \quad (\text{A5})$$

From the definitions of $Q(r, \lambda)$ and $P(\theta, \lambda)$ given in (6), one finds straightforwardly

$$\partial_\lambda Q(r, \lambda) = -\frac{\Xi^2 a^2 \Delta_r}{2Q(r, \lambda)}, \quad \partial_\lambda P(\theta, \lambda) = \frac{\Xi^2 a^2 \Delta_\theta}{2P(\theta, \lambda)}, \quad (\text{A6})$$

and one can rewrite Eq. (A4) as

$$v d\lambda = -\frac{dr}{Q(r, \lambda)} + \frac{d\theta}{P(\theta, \lambda)}, \quad (\text{A7})$$

after choosing the constant c_1 to be $\Xi^2 a^2 / 2$ and defining $v = -\partial_\lambda F(r, \theta, \lambda)$. By making use of Eq. (A1) and Eq. (A7), the metric of the Kerr-Newman-AdS₄ metric given in (2) can be re-expressed in terms of the coordinates $(t, r_*, \theta, \lambda)$

$$ds^2 = \frac{\Delta_r \Delta_\theta}{\Xi^4 R^2} (dr_*^2 - dt^2) + R^2 \sin^2 \theta (d\phi - \Omega dt)^2 + \frac{v^2 P^2(\theta, \lambda) Q^2(r, \lambda)}{\Xi^4 R^2} d\lambda^2, \quad (\text{A8})$$

where

$$R^2 = \frac{g_{\phi\phi}}{\sin^2 \theta} = \frac{\Delta_\theta (r^2 + a^2)^2 - \Delta_r a^2 \sin^2 \theta}{\Xi^2 \Sigma}, \quad (\text{A9})$$

$$\Omega = -\frac{g_{t\phi}}{g_{\phi\phi}} = \frac{a [\Delta_\theta (r^2 + a^2) - \Delta_r]}{\Xi \Sigma R^2}. \quad (\text{A10})$$

Then, using Eqs. (A1) and (A7), one has

$$dr = \frac{Q(r, \lambda) \Delta_r}{\Xi^4 R^2 \Sigma} [\Delta_\theta dr_* - P^2(\theta, \lambda) v d\lambda], \quad (\text{A11})$$

$$d\theta = \frac{P(\theta, \lambda) \Delta_\theta}{\Xi^4 R^2 \Sigma} [\Delta_r dr_* + Q^2(r, \lambda) v d\lambda]. \quad (\text{A12})$$

Since the outgoing and ingoing null congruences are defined to be $du = dv = 0$, which means $dr_*^2 = dt^2$, the metric of the Kerr-Newman-AdS₄ black hole on the null hypersurface is reduced to

$$dh^2 = R^2 \sin^2 \theta (d\phi - \Omega dt)^2 + \frac{v^2 P^2(\theta, \lambda) Q^2(r, \lambda)}{\Xi^4 R^2} d\lambda^2. \quad (\text{A13})$$

Because the volume element of a null hypersurface is the square root of the determinant of the induced metric, the points where the induced metric determinant goes to zero correspond to the NHC. Thus, the condition of the NHC is

$$v P(\theta, \lambda) Q(r, \lambda) \sin \theta = 0, \quad (\text{A14})$$

which is the determinant of the induced metric (A13) of the Kerr-Newman-AdS₄ black hole. Let us consider the ingoing null hypersurface case (same for the outgoing null hypersurface case), namely, $\lambda = \text{const}$ and a decreasing r , as an example to study the NHC. We need to analyze each factor in Eq. (A14). Since Eq. (A7) gives that the decreasing of θ will lead to the decreasing of r for a fixed λ , we have $P(\theta, \lambda) > 0$. Therefore, $Q(r, \lambda) = 0$ is a sufficient condition for the existence of the NHC.