ON A DEFORMATION OF GLUING STABILITY CONDITIONS

KOTARO KAWATANI

ABSTRACT. On a triangulated category **D** equipped with a semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$, Collins and Polishchuk develop a gluing construction of stability condition on **D**. The gluing construction gives a stability condition on **D** from these on \mathbf{D}_1 and \mathbf{D}_2 . We study a deformation of gluing stability conditions on for a nice semiorthogonal decomposition. As a consequence, we construct a continuous family of stability conditions by showing a deformation property introduced by Bridgeland's original paper. Here the deformation property is weaker than the support property which is the standard solution for the continuousness. After proving the continuousness of the family, we show that each stability condition in the family satisfies the support property via specialization. More precisely we find a stability condition with support property at the boundary of the family. Finally applying these results, we study the space of stability conditions on the category of morphisms in a triangulated category.

1. INTRODUCTION

Motivated by Douglas's work for Π -stability, Bridgeland [Bri07] introduced the notion of stability conditions on a triangulated category. Let **Stab D** be the set of locally finite stability conditions on a triangulated category **D**. By virtue of [Bri07], **Stab D** is not only a set, but also a topological space. Moreover each connected component of **Stab D** is a complex manifold if the rank of Grothendieck group $K_0(\mathbf{D})$ is finite. Thus **Stab D** is well understood locally and geometric properties of **Stab D** is expected to reflect categorical properties of **D** (see also [BB17], [Bri08]).

On the other hand, global description of $\operatorname{Stab} \mathbf{D}$ is difficult. For instance the connectedness, which is the most fundamental topological property, is widely open. If \mathbf{D} is the triangulated category of A_n -singularity, then Ishii, Ueda and Uehara showed that $\operatorname{Stab} \mathbf{D}$ is connected (the details are in [IUU10]). However for the other Kleinian singularities, the connectedness is still open. To discuss other cases, suppose that \mathbf{D} is the bounded derived category $\mathbf{D}^b(X)$ of coherent sheaves on a smooth projective variety X. If dim X = 1, then $\operatorname{Stab} \mathbf{D}^b(X)$ is non-empty and connected by [Bri07] and [Mac07]. If dim X = 2, then $\operatorname{Stab} \mathbf{D}^b(X)$ is non-empty by [AB13] but the connectedness is open. If dim X = 3, then the non-emptiness of $\operatorname{Stab} \mathbf{D}^b(X)$ follows from the generalized Bogomolov inequality proposed in [BMT14] but the connectedness is widely open.

The global description of Stab D is difficult but the space Stab D is expected to be contractible (unless it is empty), that is, the homotopy type of Stab D is the simplest in non-empty topological spaces. Motivated by the expectation, the author studied the space of stability conditions on the category of morphisms in a triangulated category D in the previous paper [Kaw19]. Unfortunately the category of morphisms in a triangulated category D is the homotopy category h(\mathcal{C}) of a stable infinity category \mathcal{C} . Then the infinity category \mathcal{C}^{Δ^1} of morphisms in the infinity category \mathcal{C}^{Δ^1} is triangulated. Thus h(\mathcal{C}^{Δ^1}) is a reasonable candidate of a triangulated category of morphisms in $\mathbf{D} = \mathbf{h}(\mathcal{C})$.

Although not directly related to the theme of this article, we would like to introduce some interesting aspects of the category of morphisms. Suppose that \mathcal{C} is the derived (infinity) category of the affine scheme Spec **k** of a field **k**. Then the category $h(\mathcal{C}^{\Delta^1})$ is equivalent to the derived

Date: September 15, 2021.

²⁰²⁰ Mathematics Subject Classification. 18G80, 32G10.

Key words and phrases. Stability conditions, Triangulated categories, Deformation, Support property.

category of the representation of A_2 -quiver by [Kaw19, Corollary 6.2]. Hence the category $h(\mathcal{C}^{\Delta^1})$ for a stable infinity category \mathcal{C} can be regarded as a generalization of the derived category of the quiver representations. In addition, recalling that the objects in the derived category $\mathbf{D}^b(X)$ of a projective variety represent (topological) D-branes, one can expect the category $h(\mathcal{C}^{\Delta^1})$ relates to open strings in string theory (for instance see [ABC⁺09]).

Our basic motivation is to understand a relation between $\operatorname{Stab} h(\mathcal{C})$ and $\operatorname{Stab} h(\mathcal{C}^{\Delta^1})$. Before recalling our previous results, let us recall basic notation. Since any object $f \in \mathcal{C}^{\Delta^1}$ is a morphism in \mathcal{C} , the object f is written as $[f: x \to y]$ for objects x and y in \mathcal{C} . Under these notation, we have three natural functors (as infinity categories) between \mathcal{C} and \mathcal{C}^{Δ^1}

(1.1)
$$\mathbb{C} \xrightarrow{d_0}_{s \to c} \mathbb{C}^{\Delta^1}; d_0 \dashv s \dashv d_1,$$

where d_0 sends the object $[f: x \to y] \in \mathbb{C}^{\Delta^1}$ to $y \in \mathbb{C}$, d_1 sends $f \in \mathbb{C}^{\Delta^1}$ to $x \in \mathbb{C}$, and s sends an object $z \in \mathbb{C}$ to the object $\mathrm{id}_z \in \mathbb{C}^{\Delta^1}$ which represents the identity morphism $[\mathrm{id}: z \to z]$ in \mathbb{C} . These functors give adjoint pairs of functors. Precisely d_0 is a left adjoint of s, and d_1 is a right adjoint of s.

In [Kaw19], we have shown the following:

Theorem 1.1 ([Kaw19, Theorem 1.2]). Let \mathcal{C} be a stable infinity category with rank $K_0(h(\mathcal{C})) < \infty$.

(1) If there exists a reasonable stability condition on h(C) then there exists a reasonable stability condition on h(C^{Δ1}). Moreover both functors d₀ and d₁ induce continuous and injective maps d₀^{*} and d₁^{*} from the space Stab^r h(C) of reasonable stability conditions on h(C) to that of h(C^{Δ1}):

 d_0^*, d_1^* : Stab^r h(C) \longrightarrow Stab^r h(C^{Δ^1}).

- (2) Both images $\operatorname{Im} d_0^*$ and $\operatorname{Im} d_1^*$ are closed in $\operatorname{Stab}^r h(\mathbb{C}^{\Delta^1})$ and do not intersect each other.
- (3) A stability condition σ is full if and only if $d_0^*\sigma$ (or $d_1^*\sigma$) is full.

The category $h(\mathcal{C}^{\Delta^1})$ has two semiorthogonal decompositions $\langle h(\mathcal{C}), h(\mathcal{C}_{/0}) \rangle$ and $\langle h(\mathcal{C}_{0/}), h(\mathcal{C}) \rangle$ (see also (8.1) and (8.2)) corresponds to the adjoint pairs $d_0 \dashv s$ and $s \dashv d_1$. Collins-Polishchuk [CP10] constructed a "gluing" stability condition from a semiorthogonal decomposition of a triangulated category. A reasonable stability condition introduced by [CP10] is necessary for the gluing construction. The construction of continuous maps d_1 and d_0 is based on the gluing construction. Hence we had to focus on reasonable stability conditions. Since a full stability condition is reasonable, $\operatorname{Stab} h(\mathcal{C}^{\Delta^1})$ is non-empty when there exists a full stability condition on $h(\mathcal{C})$ by Theorem 1.1.

In this article, we would like to understand the relation between $\mathsf{Stab}\,h(\mathfrak{C})$ and $\mathsf{Stab}\,h(\mathfrak{C}^{\Delta^1})$ deeper. Several problems might be considered for the sake and the following is one of these:

Problem 1.2. Let \mathcal{C} be a stable infinity category and σ a stability condition on $h(\mathcal{C})$. Is $d_0^*\sigma$ path connected to $d_1^*\sigma$?

Once recalling that the space of stability conditions is expected to be contractible, one can guess that both $\mathsf{Stab}\,h(\mathcal{C})$ and $\mathsf{Stab}\,h(\mathcal{C}^{\Delta^1})$ are contractible, in particular, they are homotopy equivalent to each other. In fact, if $h(\mathcal{C})$ is the derived category of an affine Noetherian scheme, $\mathsf{Stab}\,h(\mathcal{C})$ and $\mathsf{Stab}\,h(\mathcal{C}^{\Delta^1})$ are homotopy equivalent by the author [Kaw20]. However it seems hard to show the homotopy equivalence in general, because of the difficulty of the global description.

Now if the answer to Problem 1.2 is negative, $\operatorname{Stab} h(\mathbb{C})$ and $\operatorname{Stab} h(\mathbb{C}^{\Delta^1})$ might not be homotopy equivalent. Recalling that there is no example of **D** whose spaces of full stability conditions is non-connected, it is natural to expect to an affirmative answer to Problem 1.2 at least for a full stability condition on $h(\mathbb{C})$. Moreover, by the previous paper [Kaw19], the answer is positive when $h(\mathcal{C})$ is the derived category $\mathbf{D}^{b}(\mathbb{P}^{1})$ of the projective line \mathbb{P}^{1} over a field \mathbf{k} . The aim of this article is giving a positive answer to Problem 1.2.

Theorem 1.3 (=Theorem 8.5). Let \mathcal{C} be a stable infinity category with rank $K_0(h(\mathcal{C})) < \infty$. If a stability condition $\sigma \in \mathsf{Stab}\,h(\mathcal{C})$ is rational, then $d_0^*\sigma$ is path connected to $d_1^*\sigma$. In particular, the restriction of d_0^* and d_1^* to the space $\mathsf{Stab}^{\mathsf{fl}}\,h(\mathcal{C})$ of full stability conditions on $h(\mathcal{C})$ gives the same map at the level of connected components $\pi_0(\mathsf{Stab}^{\mathsf{fl}}\,h(\mathcal{C})) \to \pi_0(\mathsf{Stab}^{\mathsf{fl}}\,h(\mathcal{C}^{\Delta^1}))$.

To prove Theorem 1.3, it is necessary to study an explicit deformation of gluing stability conditions on $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$. Similarly to the case of K3 surfaces (cf. [Bri08]), we construct a complex one dimensional family of stability conditions which will be denoted by $\mathcal{S}(\epsilon_1, \epsilon_2) =$ $\{\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} \mid (\beta,\omega) \in \mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)\}$ where $\sigma_i \in \mathsf{Stab} \mathbf{D}_i$ with nice gluing property (cf. Definition 2.18).

The construction is divided into mainly four steps:

- (Step 1) An observation of a common property of two semiorthogonal decomposition $h(\mathcal{C}^{\Delta^1}) = \langle h(\mathcal{C}), h(\mathcal{C}_{/0}) \rangle$ and $h(\mathcal{C}^{\Delta^1}) = \langle h(\mathcal{C}_{0/}), h(\mathcal{C}) \rangle$.
- (Step 2) An introduction of a stability on the gluing heart which is an analogy of the slope stability of vector bundles on K3 surfaces (or more generally smooth projective variety X with $\dim X > 1$).
- (Step 3) Construction of a continuous family of stability conditions on $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$.
- (Step 4) Specialization of the semiorthogonal decompositions.

We first introduce some conditions for semiorthogonal decompositions (cf. Definition 2.18). These conditions are motivated by two semiorthogonal decompositions of $h(\mathcal{C}^{\Delta^1})$

Recall that the tilting construction on K3 surfaces is based on the slope stability of vector bundles (cf. [Bri08]). Similarly to the case of K3 surfaces, we introduce a stability on the heart of a gluing stability condition $gl(\sigma_1, \sigma_2)$ on a triangulated category **D** equipped with the semiorthogonal decomposition $\langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ discussed in Step 1 above. Then such a slope like stability satisfies the support property which is an analogy of the support property of the slope stability.

Then, we introduce a torsion pair on the heart of a gluing stability condition. It is not hard to construct $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ for rational numbers $(\beta,\omega) \in \mathbb{Q} \times \mathbb{Q}_{>0}$. The hardest part is to show the continuousness of the family. As mentioned in [BMS16, Appendix], if the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ satisfies the support property one could deform locally and show the continuousness of the family. However the support property is very hard in our situation since the category $h(\mathcal{C}^{\Delta^1})$ is not geometric but too general. In stead of it, we show a weaker property suggested in the original paper due to Bridgeland [Bri07] so that the stability condition can be deformed locally. After that we extend our construction to non rational numbers $(\beta, \omega) \in \mathbb{R} \times \mathbb{R}_{>0}$ by taking limits of rational ones.

Though we could not prove the support property of any stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} \in \mathcal{S}(\epsilon_1,\epsilon_2)$ directly, we will prove the support property via "specialization". More precisely, at the boundary of the family $\mathcal{S}(\epsilon_1,\epsilon_2)$, there is a stability condition $\Sigma_{1,0}^{\sigma_1,\sigma_2}$ satisfying the support property if σ_1 and σ_2 satisfy the support property (see also §7). Then the support property holds for any one in $\mathcal{S}(\epsilon_1,\epsilon_2)$ since the family is continuous.

Finally applying the semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ to $\mathbf{h}(\mathcal{C}^{\Delta^1}) = \langle \mathbf{h}(\mathcal{C}), \mathbf{h}(\mathcal{C}_{/0}) \rangle$ and $\mathbf{h}(\mathcal{C}^{\Delta^1}) = \langle \mathbf{h}(\mathcal{C}_{0/}), \mathbf{h}(\mathcal{C}) \rangle$, we obtain a path which connect $d_0^* \sigma$ and $d_1^* \sigma$. At the end of introduction, we would like to answer a question that readers may have. Problem 1.2 might be very simple for readers. However we hope that the construction of the answer, Theorem 8.5, could be interesting and that our construction will provide a new insight into deformation of gluing stability conditions in a future study.

2. Preliminaries

Let **D** be a triangulated category. We always assume that the rank of the Grothendieck group $K_0(\mathbf{D})$ is finite. The *t*-structure on **D** is denoted by $(\mathbf{D}^{\leq 0}, \mathbf{D}^{\geq 0})$. Recall that the subcategory $\mathbf{D}^{\leq 0} \cap \mathbf{D}^{\geq 0}$ is the heart \mathcal{A} of the *t*-structure. As usual, let us denote $\mathbf{D}^{\leq 0}[p]$ by $\mathbf{D}^{\leq -p}$ and denote $\mathbf{D}^{\geq 0}[p]$ by $\mathbf{D}^{\geq -p}$ for any integer p.

Through this article we frequently identify the complex plane \mathbb{C} with the 2-dimensional Euclidian space \mathbb{R}^2 via the canonical isomorphism. The upper half-plane is denoted by \mathcal{H} :

(2.1)
$$\mathcal{H} = \{ (\beta, \omega) \mid \beta \in \mathbb{R}, \omega \in \mathbb{R}_{>0} \}$$

The closure of \mathcal{H} in \mathbb{C} is denoted by $\overline{\mathcal{H}}$. For an arbitrary subset U of \mathcal{H} , the set of rational points in U is denoted by $U_{\mathbb{Q}}$:

$$U_{\mathbb{Q}} = \{ (\beta, \omega) \in U \mid \beta \in \mathbb{Q}, \omega \in \mathbb{Q} \}.$$

Given a complex number $z \in \mathbb{C}$, the real part of z is denoted by $\mathfrak{Re} z$ and the imaginary part of z is denote by $\mathfrak{Im} z$. We define the argument $\arg z$ of a non-zero complex number z so that

$$(2.2) \qquad \qquad \arg z \in (-\pi, \pi]$$

The following proposition is very elemental but is a key ingredient to prove the desired deformation property in Section 5.

Proposition 2.1. Let θ be in the open interval $(0, \pi)$. Suppose complex numbers z_1 and $z_2 \in \mathbb{C}$ satisfy $0 \leq \arg z_1 - \arg z_2 \leq \theta$. The following holds:

(2.3)
$$|z_1 + z_2| \ge \frac{\sqrt{2 + 2\cos\theta}}{2}(|z_1| + |z_2|)$$

Proof. Put $\phi = \arg z_1 - \arg z_2$. The following inequality holds for any $x \in \mathbb{R}$:

(2.4)
$$\sqrt{1+x^2+2x\cos\phi} \ge \frac{\sqrt{2+2\cos\phi}}{2}(x+1).$$

Indeed, a very elemental calculation below gives the proof:

$$x^{2} + 2x\cos\phi + 1 - \left(\frac{\sqrt{2+2\cos\phi}}{2}(x+1)\right)^{2} = \frac{1-\cos\phi}{2}(x-1)^{2} \ge 0.$$

On the other hand, we easily see $|z_1 + z_2| = \sqrt{|z_1|^2 + |z_2|^2 + 2|z_1| \cdot |z_2| \cdot \cos \phi}$. Then the inequality (2.4) implies

$$|z_1 + z_2| \ge |z_1| \cdot \frac{\sqrt{2 + 2\cos\phi}}{2} \left(\frac{|z_2|}{|z_1|} + 1\right) = \frac{\sqrt{2 + 2\cos\phi}}{2} (|z_2| + |z_1|)$$

Since $f(\phi) = \cos \phi$ is monotonically decreasing for $0 \le \phi < \pi$, this gives the proof.

Corollary 2.2. Let θ be in the open interval $(0, \pi)$. Then the supremum

$$\sup\left\{\frac{|z_2|}{|z_1+z_2|} \middle| z_1, z_2 \in \mathbb{C}, z_1+z_2 \neq 0, 0 \le |\arg z_1 - \arg z_2| \le \theta\right\}$$

is finite.

Proof. By Proposition 2.1, the following holds:

$$\frac{|z_2|}{|z_1+z_2|} \le \frac{2}{\sqrt{2+2\cos\theta}} \cdot \left(\frac{|z_2|}{|z_1|+|z_2|}\right) \le \frac{2}{\sqrt{2+2\cos\theta}}.$$

The right hand side is bounded above by the assumption for θ .

2.1. Stability conditions. This section is devoted to a review of basic notation and conventions for stability conditions. Let **D** be a triangulated category.

Definition 2.3 ([Bri07]). A stability condition $\sigma = (Z, \mathcal{P})$ on **D** is a collection of a group homomorphism $Z: K_0(\mathbf{D}) \to \mathbb{C}$ called the central charge and the collection of full subcategories $\mathcal{P}(\phi)$ of **D** for $\phi \in \mathbb{R}$ satisfying the following condition:

- (1) for any $E \in \mathcal{P}(\phi)$, then $Z(E) \in \mathbb{R}_{>0} \cdot \exp(\sqrt{-1\pi\phi})$,
- (2) E is in $\mathcal{P}(\phi)$ if and only if E[1] is in $\mathcal{P}(\phi+1)$,
- (3) if $\phi > \psi$ then $\operatorname{Hom}_{\mathbf{D}}(E, F) = 0$ for $E \in \mathcal{P}(\phi)$ and $F \in \mathcal{P}(\psi)$,
- (4) any object $E \in \mathbf{D}$ has a finite sequence of distinguished triangles

$$(2.5) \qquad 0 \xrightarrow{} E_1 \xrightarrow{} E_2 \xrightarrow{} \cdots \xrightarrow{} E_{n-1} \xrightarrow{} E_n = E$$

where $A_i \in \mathcal{P}(\phi) \setminus \{0\}$ with $\phi_i > \phi_{i+1}$.

A non-zero object $E \in \mathcal{P}(\phi)$ for some $\phi \in \mathbb{R}$ is said to be σ -semistable with the phase ϕ . The collection of full subcategories $\bigcup_{\phi \in \mathbb{R}} \mathcal{P}(\phi)$ satisfying the axiom (2), (3) and (4) is called the slicing of **D**.

Remark 2.4. By the definition, the zero object $0 \in \mathbf{D}$ is not σ -semistable. We note that $\mathcal{P}(\phi)$ is abelian. A σ -semistable object $A \in \mathcal{P}(\phi)$ is said to be σ -stable if there is no non-trivial subobject of A in $\mathcal{P}(\phi)$.

For any $E \in \mathbf{D}$, the filtration given by (2.5) is called a *Harder-Narasimhan filtration* of E. The filtration is unique up to isomorphism. The first term E_1 of the filtration is said to be the maximal destabilizing subobject of E, and the mapping cone E_n/E_{n-1} of $E_{n-1} \to E_n$ is called the maximal destabilizing quotient of E. Moreover the phase of the maximal destabilizing subobject (resp. quotient) of E is denoted by $\phi_{\sigma}^+(E)$ (resp. $\phi_{\sigma}^-(E)$).

Now we recall convention which is necessary in this article.

Definition 2.5. Let σ be a stability condition on **D**.

- (1) A stability condition σ is said to be *discrete* if the image of $Z: K_0(\mathbf{D}) \to \mathbb{C}$ is discrete in \mathbb{C} . A stability condition σ is said to be *rational* if the image of $Z: K_0(\mathbf{D}) \to \mathbb{C}$ is contained in rational coefficients complex numbers.
- (2) For an interval $I \subset \mathbb{R}$, the extension closure of $\{\mathcal{P}(\phi) \mid \phi \in I\}$ is denoted by $\mathcal{P}(I)$.
- (3) A stability condition $\sigma = (Z, \mathcal{P})$ is said to be *locally finite* if for any ϕ there exists an $\epsilon > 0$ such that the quasi-abelian category $\mathcal{P}(\phi \epsilon, \phi + \epsilon)$ is finite length.
- (4) For any $E \in \mathbf{D}$, the mass of E which will be denoted by $m_{\sigma}(E)$ is the sum $\sum_{A_i} |Z(A_i)|$ where A_i runs semistable factors of the Harder-Narasimhan filtration of E.
- (5) Let $\tau = (W, Q)$ be a stability condition on **D**. Set

$$\mathsf{d}(\mathcal{P},\mathcal{Q}) = \sup\{|\phi_{\sigma}^{-}(E) - \phi_{\tau}^{-}(E)|, |\phi_{\sigma}^{+}(E) - \phi_{\tau}^{+}(E)| \mid E \in \mathbf{D} \setminus \{0\}\},\$$

and a norm on $\operatorname{Hom}(K_0(\mathbf{D}), \mathbb{C})$ by

$$||U||_{\sigma} = \sup\left\{\frac{|U(E)|}{|Z(E)|}\middle| E \in \bigcup \mathcal{P}(\phi), E \neq 0\right\}.$$

We note that $||U||_{\sigma}$ might be infinity.

- (6) The stability condition σ is said to be *full* if the norm $||U||_{\sigma}$ is finite for any $U \in \text{Hom}(K_0(\mathbf{D}), \mathbb{C})$.
- (7) The stability condition σ is said to be *reasonable* if the infimum

$$\inf\left\{ |Z(E)| \left| E \in \bigcup_{\phi \in \mathbb{R}} \mathcal{P}(\phi) \right. \right\}$$

is positive.

- (8) The set of locally finite stability condition on \mathbf{D} is denoted by $\mathsf{Stab} \mathbf{D}$. In addition, the set of full (resp. reasonable) stability conditions is denoted by $\mathsf{Stab}^{\mathsf{fl}} \mathbf{D}$ (resp. $\mathsf{Stab}^{\mathsf{r}} \mathbf{D}$).
- (9) σ satisfies the support property if there exists a quadratic form q on $K_0(\mathbf{D}) \otimes \mathbb{R}$ such that
 - any σ -semistable object $E \in \mathbf{D}$, $q([E]) \ge 0$, and
 - q is negative definite on Ker $Z \subset K_0(\mathbf{D}) \otimes \mathbb{R}$.

The support property is important since it is necessary to discuss the continuousness of the family of stability conditions, or well-behaved wall crossing phenomena. The following gives some equivalent formulation of the support property.

Proposition 2.6 ([BM11], [KS08], [BMS16]). Let σ be a locally finite stability condition on **D**. The following are equivalent:

- (1) σ is full.
- (2) σ satisfies the support property.
- (3) Fix a norm on $K_0(\mathbf{D}) \otimes \mathbb{R}$. There exists a positive constant C > 0 such that the inequality

$$\frac{\|[E]\|}{|Z(E)|} \le C$$

holds for any σ -semistable object $E \in \mathbf{D}$.

By Proposition 2.6, one can deduce

Proposition 2.7. Let $\sigma = (Z, \mathcal{P})$ be a stability condition on a triangulated category **D**. If σ is full, then σ is reasonable. In addition, a reasonable stability condition is locally finite.

Proof. Let us fix a norm $\| * \| : K_0(\mathbf{D}) \otimes \mathbb{R} \to \mathbb{R}_{>0}$. If σ is full, by Proposition 2.6, there exists a constant C' > 0 such that the inequality

holds for any σ -semistable object E. Since ||E|| is discrete, σ is reasonable. The last part follows from [CP10, Lemma 1.1].

Remark 2.8. Thus we have a sequence of inclusion

$\mathsf{Stab}^\mathsf{fl}\,\mathbf{D}\subset\mathsf{Stab}^\mathsf{r}\,\mathbf{D}\subset\mathsf{Stab}\,\mathbf{D}.$

One can also check that the support property is open and closed by the argument in [Bri07, Lemmma 6.2]. Hence $\mathsf{Stab}^{\mathsf{fl}} \mathbf{D}$ is open and closed in $\mathsf{Stab} \mathbf{D}$. This implies that any $\sigma \in \mathsf{C}_0$ is full if there exists a full stability condition σ_0 in a connected component C_0 of $\mathsf{Stab} \mathbf{D}$. The same assertion for reasonable stability conditions holds by [CP10, Proposition 1.2].

Remark 2.9. If a locally finite stability condition σ satisfies the support property, then σ is said to be a *Bridgeland stability condition*. The set of Bridgeland stability condition on a triangulated category **D** is the same as $\mathsf{Stab}^{\mathsf{fl}} \mathbf{D}$ by Proposition 2.6. Recently the set of Bridgeland stability conditions is often denoted by $\mathsf{Stab} \mathbf{D}$ instead of $\mathsf{Stab}^{\mathsf{fl}} \mathbf{D}$, but in this paper we denote it by $\mathsf{Stab}^{\mathsf{fl}} \mathbf{D}$.

The following is a summary of [Bri07].

Theorem 2.10 ([Bri07]). Let **D** be a triangulated category and $\sigma = (Z, \mathcal{P})$ a locally finite stability condition.

(1) Set a subset $B_{\epsilon}(\sigma)$ of Stab **D** by

 $B_{\epsilon}(\sigma) = \{\tau = (W, \mathcal{Q}) \in \mathsf{Stab}\,\mathbf{D} \mid \mathsf{d}\,(\mathcal{P}, \mathcal{Q}) < \epsilon, \|W - Z\|_{\sigma} < \sin(\pi\epsilon)\}.$

Then there exists a topology on $\operatorname{Stab} \mathbf{D}$ such that $\{B_{\epsilon}(\sigma) \mid \sigma \in \operatorname{Stab} \mathbf{D}, \epsilon \in (0, 1/4)\}$ forms a basis of the topology.

(2) For an $\epsilon \in (0, 1/8)$, if $W \colon K_0(\mathbf{D}) \to \mathbb{C}$ satisfies

$$||W - Z||_{\sigma} < \sin(\pi\epsilon)$$

then there exists a unique stability condition τ whose central charge is W and slicing Q satisfies $d(Q, P) < \epsilon$.

(3) Each connected components of $\operatorname{Stab} \mathbf{D}$ is a complex manifold. The tangent space T_{σ} at $\sigma \in \operatorname{Stab} \mathbf{D}$ is isomorphic to the linear subspace

$$V_{\sigma} = \{ U \in \operatorname{Hom}(K_0(\mathbf{D}), \mathbb{C}) \mid ||U||_{\sigma} < \infty \}.$$

(4) Stab **D** has the right action of the universal cover $\widetilde{\operatorname{GL}}_2^+(\mathbb{R})$ of $\operatorname{GL}_2^+(\mathbb{R})$ and the left action of the autoequivalence group Aut (**D**) of **D**.

Definition 2.11. Let $\sigma = (Z, \mathcal{P})$ be a locally finite stability condition on **D**. A stability condition $\tau = (W, \mathcal{Q})$ on **D** is said to be a *deformation of* σ if τ is in $B_{\epsilon}(\sigma)$ for an $\epsilon \in (0, 1/8)$.

Remark 2.12. For a locally finite stability condition $\sigma = (Z, \mathcal{P})$ and a group homomorphism $W: K_0(\mathbf{D}) \to \mathbb{C}$, if $||W - Z||_{\sigma}$ is sufficiently small then there exists a unique deformation of σ by Theorem 2.10.

Finally we recall another definition of stability conditions which is helpful to construct σ in an explicit way.

Definition 2.13. Let \mathcal{A} be the heart of a bounded *t*-structure $(\mathbf{D}^{\leq 0}, \mathbf{D}^{\geq 0})$ on \mathbf{D} and let Z be a group homomorphism $Z: K_0(\mathcal{A}) \to \mathbb{C}$.

- (1) Z is said to be a central charge on \mathcal{A} if Z satisfies $0 < \arg Z(E) \leq \pi$ for any non-zero object $E \in \mathcal{A}$.
- (2) Given a central charge Z, an object E is said to be Z-semistable if the inequality $\arg Z(F) \leq \arg Z(E)$ holds for any non-trivial subobject F of E.
- (3) Let Z be a central charge on \mathcal{A} . The pair (\mathcal{A}, Z) has the Harder-Narashimhan property if any non-zero object E has a finite filtration of subobjects

$$0 = E_0 \subset E_1 \subset E_2 \subset \cdots \subset E_{n-1} \subset E_n = E$$

such that

(2.7)

- for each $i \in \{1, 2, \dots, n\}$, $A_i = E_i / E_{i-1}$ is Z-semistable, and
- $\arg Z(A_i)$ is decreasing.

In addition, the pair (\mathcal{A}, Z) with the Harder-Narasimhan property is called a *stability* condition on the heart \mathcal{A} .

Proposition 2.14 ([Bri07, Proposition 5.3]). Let **D** be a triangulated category. Then the following are equivalent:

- (1) To give a stability condition $\sigma = (Z, \mathcal{P})$ on **D**.
- (2) To give a pair (\mathcal{A}, Z) consisting of the heart \mathcal{A} of a bounded t-structure on \mathbf{D} and a central charge Z on \mathcal{A} which has the Harder-Narashimhan property.

Remark 2.15. Let us briefly recall that the heart \mathcal{A} is given by $\mathcal{P}(0, 1]$. Throughout the article, we call $\mathcal{A} = \mathcal{P}(0, 1]$ the *heart of a stability condition* σ and fluently identify the pair (Z, \mathcal{P}) with (\mathcal{A}, Z) . Namely we denote by \mathcal{P} the slicing of the stability condition (\mathcal{A}, Z) and by \mathcal{A} the heart of the stability condition (Z, \mathcal{P}) .

Given a stability condition $\sigma = (\mathcal{A}, Z)$ on **D**, we obtain a torsion pair on \mathcal{A} which is useful in this article:

Definition 2.16. Let **D** be a triangulated category and $\sigma = (\mathcal{A}, Z)$ a stability condition on **D**. An object $E \in \mathcal{A}$ is said to be σ -torsion if $\Im \mathfrak{m} Z(E) = 0$. An object $E' \in \mathcal{A}$ is said to be σ -free if $E \in \mathcal{P}(0, 1)$

The full subcategory of σ -torsion objects in \mathcal{A} is denoted by \mathcal{T}^{σ} and the full subcategory of σ -torsion free objects in \mathcal{A} is denoted by \mathcal{F}^{σ} .

Lemma 2.17. Keep the notation as in Definition 2.16. Then the pair $(\mathcal{T}^{\sigma}, \mathcal{F}^{\sigma})$ is a torsion pair on \mathcal{A} .

Proof. The definition of stability conditions directly implies the assertion.

2.2. Semiorthogonal decompositions and stability conditions. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition of \mathbf{D} . Namely we have pairs of adjoint functors

$$\mathbf{D}_2 \xrightarrow[\tau_2^R]{i_2} \mathbf{D} \xrightarrow[\tau_1^L]{i_1} \mathbf{D}_1 ; i_2 \dashv \tau_2^R, \tau_1^L \dashv i_1,$$

both i_2 and i_1 are fully faithful and $\tau_1^L \circ i_2 = \tau_2^R \circ i_1 = 0$. If i_1 has the right adjoint $\tau_1^R : \mathbf{D} \to \mathbf{D}_1$, then the composite $\tau_1^R \circ i_2[1] : \mathbf{D}_2 \to \mathbf{D}_1$ is said to be the *gluing functor* of the semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ which will be denoted by $\Phi : \mathbf{D}_2 \to \mathbf{D}_1$.

We introduce some important conditions derived from the category of morphisms.

Definition 2.18. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition of \mathbf{D} , and let $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$ be the bounded *t*-structure on \mathbf{D}_i . A torsion pair on the heart $\mathbf{D}_i^{\leq 0} \cap \mathbf{D}_i^{\geq 0}$ is denoted by $(\mathcal{T}_i, \mathcal{F}_i)$.

- (Mor1): $i_1: \mathbf{D}_1 \to \mathbf{D}$ has the right adjoint $\tau_1^R: \mathbf{D} \to \mathbf{D}_1$.
- (Mor2): In addition to (Mor1), there exist equivalences $\mathbf{D}_i \to \mathbf{C}$ such that the right adjoint $\tau_1^R : \mathbf{D} \to \mathbf{D}_1$ gives the left adjoint τ_2^L of $i_2 : \mathbf{D}_2 \to \mathbf{D}$ via the equivalences $\mathbf{D}_k \to \mathbf{C}$.
- (Mor3): In addition to (Mor1), the gluing functor $\Phi = \tau_1^R \circ i_2[1] \colon \mathbf{D}_2 \to \mathbf{D}_1$ is *t*-exact with respect to these *t*-structures $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$.
- (Mor4): In addition to (Mor3), the gluing functor Φ respects to free part of the torsion pairs, that is, $\Phi(\mathcal{F}_2) \subset \mathcal{F}_1$.

Moreover, let $\sigma_i = (\mathcal{A}_i, Z_i)$ be in Stab \mathbf{D}_i . If the gluing functor is *t*-exact with respect to the *t*-structures of σ_i , (σ_1, σ_2) is said to satisfy the condition (Mor3). Similarly, (σ_1, σ_2) is said to satisfy the condition (Mor4), if the torsion pair $(\mathcal{T}^{\sigma_i}, \mathcal{F}^{\sigma_i})$ on the heart \mathcal{A}_i satisfies the condition (Mor4). Finally let us introduce the following condition for σ_i :

(Mor5): In addition to (Mor4) for σ_i , the equality $Z_2(E_2) = Z_1(\Phi E_2)$ holds for any $E_2 \in \mathcal{A}_2$.

Remark 2.19. Suppose a semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ satisfies the condition (Mor2). Then the gluing functor Φ is isomorphic to [1] on \mathbf{C} . If *t*-structures $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$ satisfies (Mor3), then the heart \mathcal{A}_2 on $\mathbf{D}_2 \cong \mathbf{C}$ is $\mathcal{A}_1[-1]$ on $\mathbf{D}_1 \cong \mathbf{C}$.

Lemma 2.20. Suppose that a semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ satisfies (Mor1). The gluing functor $\mathbf{D}_2 \to \mathbf{D}_1$ is denoted by Φ . There exists a distinguished triangle in \mathbf{D}_1

$$\Phi(\tau_2^R F)[-1] \longrightarrow \tau_1^R F \longrightarrow \tau_1^L F \longrightarrow \Phi(\tau_2^R F)$$

for any $F \in \mathbf{D}$.

Proof. Consider the semiorthogonal decomposition of $F \in \mathbf{D}$:

(2.8)
$$i_2 \tau_2^R(F) \longrightarrow F \longrightarrow i_1 \tau_1^L(F) \longrightarrow i_2 \tau_2^R(F)[1].$$

Taking τ_1^R to the sequence (2.8), we obtain the desired sequence.

Lemma 2.21. Suppose that a semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ satisfies (Mor1). The gluing functor is denoted by $\Phi: \mathbf{D}_2 \to \mathbf{D}_1$. Let $\varphi: E \to F$ be a morphism in \mathbf{D} with $\tau_1^L \varphi = 0$ and $\tau_2^R \varphi = 0$. If $\operatorname{Hom}_{\mathbf{D}_1}(\tau_1^L E, \Phi(\tau_2^R F)[-1]) = 0$, the morphism f is zero.

Proof. Taking the semiorthogonal decomposition of E and $F \in \mathbf{D}$, we obtain the following diagram of exact sequences:

$$(2.9) \qquad \operatorname{Hom}_{\mathbf{D}}(i_{1}\tau_{1}^{L}E, i_{2}\tau_{2}^{R}F) \xrightarrow{\overline{s}} \operatorname{Hom}_{\mathbf{D}}(E, F) \xrightarrow{t_{1}} \operatorname{Hom}_{\mathbf{D}}(E, i_{1}\tau_{1}^{L}F) \xrightarrow{\overline{s}} \operatorname{Hom}_{\mathbf{D}}(E, F) \xrightarrow{t_{1}} \operatorname{Hom}_{\mathbf{D}}(E, i_{1}\tau_{1}^{L}F) \xrightarrow{\overline{t}_{2}} \xrightarrow{t_{2}} \xrightarrow{t_{2}} \xrightarrow{t_{2}} \operatorname{Hom}_{\mathbf{D}}(i_{2}\tau_{2}^{R}E, i_{2}\tau_{2}^{R}F) \xrightarrow{s} \operatorname{Hom}_{\mathbf{D}}(i_{2}\tau_{2}^{R}E, F) \xrightarrow{} \operatorname{Hom}(i_{2}\tau_{2}^{R}E, i_{1}\tau_{1}^{L}F).$$

Note that $\operatorname{Hom}_{\mathbf{D}}(i_1\tau_1^L E, i_2\tau_2^R F) \cong \operatorname{Hom}_{\mathbf{D}_1}(\tau_1^L E, \Phi(\tau_2^R F)[-1])$. Hence the morphism \overline{t}_2 is injective. The assumption $\tau_1^L \varphi = \tau_2^R \varphi = 0$ imply $t_1(\varphi) = t_2(\varphi) = 0$, and we see that there exists a $\psi \in \operatorname{Hom}_{\mathbf{D}}(E, i_2\tau_2^R F)$ such that $\overline{s}(\psi) = \varphi$. By the semiorthogonality, the morphism *s* above is an isomorphism. Since the composite $s \circ \overline{t}_2$ is injective, ψ has to be in the kernel ker \overline{t}_2 which is trivial. Thus ψ is zero and so is $\overline{s}(\psi) = \varphi$.

Corollary 2.22. Let \mathcal{C} be a stable infinity category, and put $\mathbf{D} = h(\mathcal{C}^{\Delta^1})$. For objects $[f: x \to y]$ and $[g: z \to w]$ in $h(\mathcal{C}^{\Delta^1})$, suppose a morphism $\tau: f \to g$ in $h(\mathcal{C}^{\Delta^1})$ satisfies $d_1\tau = d_0\tau = 0$. If $\operatorname{Hom}_{h(\mathcal{C})}(x, w[-1]) = 0$, then τ is a zero morphism.

Proof. Let $j_!: \mathcal{C} \to \mathcal{C}^{\Delta^1}$ be the right adjoint of $d_1: \mathcal{C}^{\Delta^1} \to \mathcal{C}$ and let $j_*: \mathcal{C} \to \mathcal{C}^{\Delta^1}$ be the left adjoint of $d_0: \mathcal{C}^{\Delta^1} \to \mathcal{C}$. Since the right adjoint τ_1^R of $j_!$ is taking the fiber of morphisms in \mathcal{C} , we have the diagram of the semiorthogonal decomposition with (Mor1):

$$\mathbf{h}(\mathcal{C}) \xrightarrow[d_0]{j_*} \mathbf{h}(\mathcal{C}^{\Delta^1}) \xrightarrow[\tau_1^L=d_1]{\tau_1^L=d_1} \mathbf{h}(\mathcal{C}).$$

Now clearly we see

$$\operatorname{Hom}_{\mathbf{h}(\mathcal{C}^{\Delta^{1}})}(\tau_{1}^{L}f, \Phi(\tau_{2}^{R}g)) = \operatorname{Hom}_{\mathbf{h}(\mathcal{C})}(x, w[-1]).$$

Then Lemma 2.21 implies the desired assertion.

The following lemma is a revision of an argument in [CP10, Lemma 2.1 and Theorem 3.6] in terms of the gluing functor.

Lemma 2.23. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor1) and let $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$ be a bounded t-structure on \mathbf{D}_i . Suppose that the t-structures $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$ satisfies (Mor3).

(1) Then the pair of following full subcategories determines a t-structure on \mathbf{D} :

$$\mathbf{D}^{\leq 0} = \{ E \in \mathbf{D} \mid \tau_2^R E \in \mathbf{D}_2^{\leq 0} \text{ and } \tau_1^L E \in \mathbf{D}_1^{\leq 0} \}, \text{ and} \\ \mathbf{D}^{\geq 0} = \{ E \in \mathbf{D} \mid \tau_2^R E \in \mathbf{D}_2^{\geq 0} \text{ and } \tau_1^L E \in \mathbf{D}_1^{\geq 0} \}.$$

(2) Put $\mathcal{A} = \mathbf{D}^{\leq 0} \cap \mathbf{D}^{\geq 0}$. Suppose a torsion pair $(\mathcal{T}_i, \mathcal{F}_i)$ on the heart $\mathcal{A}_i = \mathbf{D}_i^{\leq 0} \cap \mathbf{D}_i^{\geq 0}$ satisfies the condition (Mor4). Then the pair $(\mathcal{T}, \mathcal{F})$ (below) gives a torsion pair on \mathcal{A} .

$$\mathcal{T} = \{ E \in \mathcal{A} \mid \tau_2^R E \in \mathcal{T}_2 \text{ and } \tau_1^L E \in \mathcal{T}_1 \}, \text{ and} \\ \mathcal{F} = \{ E \in \mathcal{A} \mid \tau_2^R E \in \mathcal{F}_2 \text{ and } \tau_1^L E \in \mathcal{F}_1 \}.$$

Proof. Put $\mathcal{A}_i := \mathbf{D}_i^{\leq 0} \cap \mathbf{D}_i^{\geq 0}$. To prove the first assertion, by [CP10, Lemma 2.1], it is enough to show $\operatorname{Hom}_{\mathbf{D}}(i_1E_1, i_2E_2[p]) = 0$ for any $E_i \in \mathcal{A}_i$ (i = 1, 2) and $p \leq 0$. By the adjunction, we see

 $\operatorname{Hom}_{\mathbf{D}}(i_1E_1, i_2E_2[p]) \cong \operatorname{Hom}_{\mathbf{D}_1}(E_1, \tau_1^R i_2E_2[p]) = \operatorname{Hom}_{\mathbf{D}_1}(E_1, \Phi(E_2)[p-1]).$ Since the gluing functor is *t*-exact, we have $\Phi(E_2) \in \mathcal{A}_1$. Thus $\operatorname{Hom}_{\mathbf{D}_1}(i_1E_1, i_2E_2[p]) = 0.$

To prove the second assertion, put E_2 by $\tau_2^R E$ and E_1 by $\tau_1^L E$. let T_i (resp. F_i) be the torsion part of E_i (the free part of E_i). Now we claim Hom_D $(i_1T_1, i_2F_2[1]) = 0$. In fact, the adjunction and the condition (Mor4) imply the vanishing as follows:

 $\operatorname{Hom}_{\mathbf{D}}(i_1T_1, i_2F_2[1]) = \operatorname{Hom}_{\mathbf{D}_1}(T_1, \tau_1^R i_2F_2[1]) = \operatorname{Hom}_{\mathbf{D}_1}(T_1, \Phi(F_2)) = 0.$

Then the vanishing $\operatorname{Hom}_{\mathbf{D}}(i_1T_1, i_2F_2[1]) = 0$ implies a morphism $i_1T_1 \to i_2T_2[1]$ and $i_1F_1 \to i_2T_2[1]$ $i_2F_2[1]$ which commute the following diagram:

$$i_1T_1 \longrightarrow i_1E_1 \longrightarrow i_1F_1$$

$$\downarrow \qquad \qquad \downarrow$$

$$i_2T_2[1] \longrightarrow i_2E_2[1] \longrightarrow i_2F_2[1].$$

The middle vertical arrow represents the object E. Then the 3×3 lemma in triangulated categories gives the distinguished triangle

$$T \longrightarrow E \longrightarrow F$$


where T (resp. F) is the mapping cone of $i_1T_1 \rightarrow i_2T_2[1]$ (resp. $i_1F_1 \rightarrow i_2F_2[1]$). Finally one can show the vanishing Hom(T, F) = 0 by the vanishing $\text{Hom}(i_1T_1, i_2F_2) = 0$ and digram chasing. \square

Definition 2.24. Keep the notation as in Lemma 2.23. We denote by $g(A_1, A_2)$ the heart of the *t*-structure defined in Lemma 2.23.

Lemma 2.25. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Suppose that t-structures $(\mathbf{D}_i^{\leq 0}, \mathbf{D}_i^{\geq 0})$ on \mathbf{D}_i satisfies (Mor3). The heart of the t-structure is denoted by \mathcal{A}_i .

Then a subobject F of $\tau_1^L E$ of $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ in \mathcal{A}_1 determines a subobject \tilde{F} of E in $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2).$

Proof. Set $E_1 = \tau_1^L E$ and $E_2 = \tau_2^R E$. Let g_E be the morphism $E_1 \to \Phi(E_2)$ which represents the extension class $[E]: i_1 E_1 \to i_2 E_2[1]$ via the adjunction $\operatorname{Hom}_{\mathbf{D}}(i_1E_1, i_2E_2[1]) \cong \operatorname{Hom}_{\mathbf{D}_1}(E_1, \Phi(E_2))$. We denote by $f: F \to \Phi(E_2)$ the restriction of g_E to F. Under the equivalence $\mathbf{D}_1 \sim \mathbf{C}$, Then we have the following commutative diagram in **C**:



where im f is the image of f. Note that im $f \cong \Phi(\operatorname{im} f[-1])$. By the ismorphism $\operatorname{Hom}_{\mathbf{C}}(F, \operatorname{im} f) \cong$ $\operatorname{Hom}_{\mathbf{D}}(i_1F, i_2(\operatorname{im} f))$, we obtain the following diagram in **D**:

$$\begin{array}{c} i_1 E_1 \xrightarrow{[E]} (i_2 E_2)[1] \\ \downarrow \\ i_1 \ell & \uparrow \\ i_1 F \xrightarrow{} i_2(\operatorname{im} f). \end{array}$$

Then define $\tilde{F}[1]$ by the mapping cone of $i_1F \rightarrow i_2(\operatorname{im} f)$. Then there exists a morphism $j: F \to E$ such that $\tau_1^L j = \iota$. Since $\tau_2^R j$ is the inclusion, F gives a subobject of E by the 3×3 lemma in triangulated categories.

Remark 2.26. By the construction of \tilde{F} , $\tau_2^R \tilde{F}$ is isomorphic to (im f)[-1].

3. SLOPE ON THE GLUING HEART

The following construction of stability conditions on \mathbf{D} was motivated from [CP10, Theorem 3.6] and the previous work [Kaw19].

Proposition 3.1. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor1) and let $\sigma_i = (\mathcal{A}_i, Z_i)$ be a reasonable stability condition on \mathbf{D}_i . Suppose (σ_1, σ_2) satisfies (Mor5). Define $gl(Z_1, Z_2): K_0(\mathbf{D}) \to \mathbb{C}$ by

gl
$$(Z_1, Z_2)(E) := Z_1(\tau_1^L E) + Z_2(\tau_2^R E).$$

If $\Phi(E_2)$ is σ_1 -semistable for any σ_2 -semistable object $E_2 \in \mathcal{A}_2$, then the pair $(\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2), \mathfrak{gl}(Z_1, Z_2))$ which will be denoted by $\mathfrak{gl}(\sigma_1, \sigma_2)$ is a locally finite stability condition on \mathbf{D} .

Proof. Let \mathcal{P}_i be the slicing of σ_i for $i \in \{1, 2\}$. Following [CP10, Theorem 3.6], it is enough to show

- (a) Hom_{**D**} $(i_1E_1, i_2E_2[p]) = 0$ for any $E_i \in \mathcal{A}_i$ and any $p \leq 0$, and
- (b) there exists an a in the interval (0,1) such that $\operatorname{Hom}_{\mathbf{D}}(i_1F_1, i_2F_2[p]) = 0$ for any $F_i \in \mathcal{P}_i(a, a+1]$ and any $p \leq 0$.

Since $\Phi(\mathcal{A}_2) \subset \mathcal{A}_1$ by the condition (Mor5), we see

$$\operatorname{Hom}_{\mathbf{D}}(i_1 E_1, i_2 E_2[p]) \cong \operatorname{Hom}_{\mathbf{D}_1}(E_1, \Phi(E_2)[p-1]) = 0.$$

This gives the proof of (a).

Now for any $F_i \in \mathcal{P}_i(a, a+1]$, there exists a canonical triangle in \mathbf{D}_i

$$F_i^{\leq a}[1] \longrightarrow F_i \longrightarrow F_i^{>a} \longrightarrow F_i^{\leq a}[2],$$

where $F_i^{\leq a} \in \mathcal{P}_i(0, a]$ and $F_i^{>a} \in \mathcal{P}_i(a, 1]$. The assertion (a) implies $\operatorname{Hom}_{\mathbf{D}}(i_1 F_i^{\leq a}[1], F_2[p]) = 0$ and $\operatorname{Hom}_{\mathbf{D}}(i_1 F_1^{>a}, i_2 F_2^{>a}[p]) = 0$ for $p \leq 0$. Hence it is enough to show $\operatorname{Hom}_{\mathbf{D}}(i_1 F_1^{>a}, i_2 F_2^{\leq a}[p+1]) = 0$ for any $p \leq 0$.

By the assumption, we see $\Phi(\mathcal{P}_2(\phi)) \subset \mathcal{P}_1(\phi)$ for any $\phi \in \mathbb{R}$. Hence we have

$$\operatorname{Hom}_{\mathbf{D}}(i_1 F_1^{>a}, i_2 F_2^{\leq a}[p+1]) \cong \operatorname{Hom}_{\mathbf{D}_1}(F_1^{>a}, \Phi(F_2^{\leq a}[p])) = 0,$$

which gives the proof.

Remark 3.2. If the semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ satisfies (Mor2), then Φ is an equivalence. Hence $\Phi(E_2)$ is σ_1 -semistable for any σ_2 -semistable object E_2 .

Next we introduce a slope on the heart $gl(A_1, A_2)$ which is an analogy of the slope stability of coherent sheaves on projective surfaces. Similarly to the case of surfaces, a suitable "freeness" of objects in $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ is necessary.

Proposition-Definition 3.3. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition and let $\sigma_i = (\mathcal{A}_i, Z_i)$ be in Stab \mathbf{D}_i . Suppose that (σ_1, σ_2) satisfies the condition (Mor4).

Then the pair $(\mathcal{T}(\sigma_1, \sigma_2), \mathcal{F}(\sigma_1, \sigma_2))$

$$\mathcal{T}(\sigma_1, \sigma_2) = \{ E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2) \mid \tau_1^L E \in \mathcal{T}^{\sigma_1}, \tau_2^R E \in \mathcal{T}^{\sigma_2} \}, \text{ and} \\ \mathcal{F}(\sigma_1, \sigma_2) = \{ E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2) \mid \tau_1^L E \in \mathcal{F}^{\sigma_1}, \tau_2^R E \in \mathcal{F}^{\sigma_2} \}.$$

gives a torsion pair on the gluing t-structure $gl(A_1, A_2)$. We refer to the torsion pair as the gluing torsion pair by (σ_1, σ_2) .

An object $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ is said to be (σ_1, σ_2) -torsion (resp. (σ_1, σ_2) -free) if E is in $\mathcal{T}(\sigma_1, \sigma_2)$ (resp. $\mathcal{F}(\sigma_1, \sigma_2)$)

Proof. The assertion follows from Lemma 2.23.

Definition 3.4. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition of \mathbf{D} and let $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$.

Define $M^{Z_1,Z_2}_{\beta,\omega} \colon K_0(\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2)) \to \mathbb{C}$ by (3.1)
$$\begin{split} & M^{(1,1)}_{\beta,\omega}(E) = -\operatorname{\mathfrak{Im}} Z_1(\tau_1^L E) + \omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R E) - \beta \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E) + \sqrt{-1} (\operatorname{\mathfrak{Im}} Z_1(\tau_1^L E) + \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E)), \end{split}$$
where $\beta \in \mathbb{R}$ and $\omega \in \mathbb{R}_{>0}$.

Remark 3.5. If $\Im \mathfrak{m} M_{\beta,\omega}^{Z_1,Z_2}(E) = 0$, then $\Im \mathfrak{m} M_{\beta,\omega}^{Z_1,Z_2}(E)$ is non-negative and $\Re \mathfrak{m} M_{\beta,\omega}^{Z_1,Z_2}(E)$ is non-positive. Moreover $M_{\beta,\omega}^{Z_1,Z_2}(E) = 0$ if and only if $\tau_2^R E = 0$ and $\tau_1^L E$ is in the torsion part \mathcal{T}^{σ_1} of σ_1 .

Now we wish to define a stability condition which is analogous to the slope stability on algebraic surfaces. The stability is necessary for the construction of a torsion pair on the heart $\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2).$

Proposition 3.6. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition of \mathbf{D} with (Mor1). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor4). Define a full sub category \mathcal{B} of $gl(\mathcal{A}_1, \mathcal{A}_2)$ by

$$\mathcal{B} = \{ E \in \mathsf{gl}\left(\mathcal{A}_1, \mathcal{A}_2\right) \mid M^{Z_1, Z_2}_{\beta, \omega}(E) = 0 \}$$

- (1) \mathcal{B} is a Serre subcategory of $gl(\mathcal{A}_1, \mathcal{A}_2)$.
- (2) Let C be the Serre quotient category $gl(A_1, A_2)/B$. If both σ_1 and σ_2 are discrete, then the pair $\tau = (\mathcal{C}, M_{\beta,\omega}^{Z_1, Z_2})$ is a stability condition on the abelian category \mathcal{C} .
- (3) In addition to (2), suppose that E₁ satisfies 3m Z₁(E₁) > 0. Then i₁(E₁) is τ-semistable with arg M^{Z₁,Z₂}_{β,ω}(i₁E₁) = 3π/4.
 (4) The stability condition τ defined in (2) satisfies the support property.

Proof. Since $M_{\beta,\omega}^{Z_1,Z_2}$ is a group homomorphism, \mathcal{B} is closed under extensions and $M_{\beta,\omega}^{Z_1,Z_2}$ satisfies • $\mathfrak{Im} M_{\beta,\omega}^{Z_1,Z_2}(E) \ge 0$, and • If $\mathfrak{Im} M_{\beta,\omega}^{Z_1,Z_2}(E) = 0$ then $\mathfrak{Re}(E) \le 0$.

Hence \mathcal{B} is closed under subobjects and quotients. This gives the proof of the assertion (1).

Since σ_i is discrete, the hearts \mathcal{A}_i is a Noetherian abelian category by [CP10, Lemma 3.4]. Thus the heart $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)$ is also Noetherian and so is $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)/\mathcal{B}$. Since the set $\{\mathfrak{Im} Z(E) \mid \mathcal{I}(E) \mid \mathbb{C}\}$ $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ is discrete in $\mathbb{R}_{\geq 0}$, the pair $(\mathcal{C}, M_{\beta,\omega}^{Z_1, Z_2})$ has the Harder-Narasimhan property by [CP10, Lemma 3.4] (or [Bri08]). This gives the proof of the assertion (2).

To prove the assertion (3), let E_1 be in \mathbf{D}_1 . If necessary we may assume that E_1 is σ_1 -free, since there exists a σ_1 -free object $E' \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ such that E' is isomorphic to i_1E_1 in \mathcal{C} . Note that any subobject F of i_1E_1 in C gives a subobject $F' \subset i_1E_1$ in $g(A_1, A_2)$ such that F' is isomorphic to F in C. Since the essential image of \mathcal{A}_1 by i_1 is closed under subobject, we have $\arg M_{\beta,\omega}^{Z_1,Z_2}(i_1E_1) = \arg M_{\beta,\omega}^{Z_1,Z_2}(F)$ and this gives the proof of the assertion (3). To prove the last assertion, it is enough to construct a quadratic form

$$(3.2) q: K_0(\mathcal{C}) \otimes \mathbb{R} \to \mathbb{R}$$

satisfying

• the restriction of q to the subspace Ker $M_{\beta,\omega}^{Z_1,Z_2}$ of $K_0(\mathcal{C}) \otimes \mathbb{R}$ is negative definite, and • $q([E]) \geq 0$ if the class [E] is represented by a semistable object E.

We show that the quadratic form

(3.3)
$$q([E]) = \Im \mathfrak{m} Z_1(\tau_1^L E) \cdot \Im \mathfrak{m} Z_2(\tau_2^R E)$$

satisfies the desired property. Note that the first condition is trivial since Ker $M_{\beta,\omega}^{Z_1,Z_2}$ is trivial by the construction of \mathcal{C} .

Take a τ -semistable object $E \in \mathcal{C}$. Then we have

$$i_2 \tau_2^R E \longrightarrow E \longrightarrow i_1 \tau_1^L E \longrightarrow i_2 \tau_2^R E[1].$$

If $\mathfrak{Im} Z_1(i_1\tau_1^L E) = 0$, then the inequality $q([E]) \ge 0$ holds. If $\mathfrak{Im} Z_1(i_1\tau_1^L E) > 0$, then the assertion (3) implies

(3.4)
$$0 < \arg M_{\beta,\omega}^{Z_1,Z_2}(i_2\tau_2^R E) \le \arg M_{\beta,\omega}^{Z_1,Z_2}(E) \le \arg M_{\beta,\omega}^{Z_1,Z_2}(i_1\tau_1^L E) = \frac{3\pi}{4}.$$

Now let us canonically identify \mathbb{C} as \mathbb{R}^2 . Then the standard inner paring of $(1+\sqrt{-1})M_{\beta,\omega}^{Z_1,Z_2}(i_2\tau_2 E)$ and $M_{\beta,\omega}^{Z_1,Z_2}(i_1\tau_1^L E)$ on \mathbb{R}^2 is just

$$2 \operatorname{\mathfrak{Im}} Z_1(\tau_1^L E) \cdot \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E).$$

By (3.4) and the assertion (3) we have

(3.5)
$$|\arg(1+\sqrt{-1})M_{\beta,\omega}^{Z_1,Z_2}(i_2\tau_2 E) - \arg M_{\beta,\omega}^{Z_1,Z_2}(i_1\tau_1^L E)| < \pi/2,$$

and this gives the proof.

Remark 3.7. In the assertion (2), we assume that the stability conditions σ_i is discrete. If σ_i is rational then σ_i is discrete since $K_0(\mathbf{D})$ is finitely generated. However, the existence of rational stability conditions is subtle. Of course, if σ_i is full (equivalently satisfies the support property), then one can choose rational stability conditions by shrinking σ_i .

Definition 3.8. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor1). Let $\sigma_i = (\mathcal{A}_i, Z_i)$ be a discrete stability condition on \mathbf{D}_i . Suppose (σ_1, σ_2) satisfies (Mor4).

Let us define $\mu_{\beta,\omega}(E)$ for $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ by

$$\mu_{\beta,\omega}(E) = \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L E) - \omega \, \mathfrak{Re} \, Z_2(\tau_2^R E) + \beta \, \Im \mathfrak{m} \, Z_2(\tau_2^R E)}{\Im \mathfrak{m} \, Z_1(\tau_1^L E) + \Im \mathfrak{m} \, Z_2(\tau_2^R E)}$$

Similarly to the case of slope stability on projective surfaces, (σ_1, σ_2) -free object E is $M_{\beta,\omega}^{Z_1,Z_2}$ semistable if and only if for any nontrivial subobject $F \subset E$ so that $\tau_2^R F$ is a proper subobject
of $\tau_1^L E$, the inequality $\mu_{\beta,\omega}(F) \leq \mu_{\beta,\omega}(E)$ holds (cf. [HL97]).

Define the following subcategories of $gl(A_1, A_2)$ by

$$\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2} = \left\{ E \in \mathsf{gl}\left(\mathcal{A}_1, \mathcal{A}_2\right) \mid E \text{ is } (\sigma_1, \sigma_2) \text{-free with } \mu_{\beta,\omega}^+(E) \le 0 \right\}, \text{ and}$$
$$\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2} = \left\{ E \in \mathsf{gl}\left(\mathcal{A}_1, \mathcal{A}_2\right) \mid E \text{ is } (\sigma_1, \sigma_2) \text{-torsion or } (\sigma_1, \sigma_2) \text{-free part } E_{\mathrm{fr}} \text{ of } E \text{ has } \mu_{\beta,\omega}^-(E_{\mathrm{fr}}) > 0 \right\}$$

The tilting heart of $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ by the torsion pair $(\mathcal{T}^{\sigma_1, \sigma_2}_{\beta, \omega}, \mathcal{F}^{\sigma_1, \sigma_2}_{\beta, \omega})$ is denoted by $\mathcal{A}^{\sigma_1, \sigma_2}_{\beta, \omega}$.

Lemma 3.9. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ with the condition (Mor1). Suppose that rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ satisfy the condition (Mor4).

If a (σ_1, σ_2) -free object $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ satisfies $\mu_{\beta,\omega}^+(E) < 1$, then the canonical morphism $\tau_1^L E \to \Phi(\tau_2^R E)$ is a monomorphism in \mathcal{A}_1 .

Proof. By Lemma 2.20 we have the distinguished triangle in \mathbf{D}_1 :

$$\tau_1^R E \xrightarrow{f} \tau_1^L E \xrightarrow{g} \Phi(\tau_2^R E) \longrightarrow \tau_1^R E[1]$$

Since E is in gl $(\mathcal{A}_1, \mathcal{A}_2)$, both $\tau_1^L E$ and $\Phi(\tau_2^R E)$ is in \mathcal{A}_1 .

Let K be the kernel of the morphism g above. Note that the morphism $\iota: K \to \tau_1^L E$ lifts to $\tau_1^R E$, that is, there exists a morphism $\bar{\iota}: K \to \tau_1^R E$ such that $f \circ \bar{\iota} = \iota$. If K is non-zero, then we have a non-zero morphism in

$$\operatorname{Hom}_{\mathbf{D}_1}(K, \tau_1^R E) \cong \operatorname{Hom}_{\mathbf{D}}(i_1 K, E).$$

By Proposition 3.6 (3), i_1K is $\mu_{\beta,\omega}$ -semistable with $\mu_{\beta,\omega}(i_1K) = 1$. Hence K has to be zero since both i_1K and E are (σ_1, σ_2) -free.

4. RATIONAL COEFFICIENTS STABILITY CONDITIONS

In the previous section, we introduced the tilting heart $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$. The aim of this section is to show that the pair $(\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}, Z_{\beta,\omega}^{\sigma_1,\sigma_2})$ gives a locally finite stability condition on $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ if the semiorthogonal decomposition satisfies (Mor2) and (σ_1, σ_2) satisfies (Mor5). These assumptions are necessary to make all the arguments work well.

Definition 4.1. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ with the condition (Mor2). Suppose that rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ satisfy the condition (Mor5). Define $Z^{\sigma_1, \sigma_2}_{\beta, \omega} \colon K_0(\mathbf{D}) \to \mathbb{C}$ by

(4.1)
$$Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) = Z_1(\tau_1^L E) + (\beta - \sqrt{-1}\omega)Z_2(\tau_2^R E)$$

Moreover the pair $(\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}, Z_{\beta,\omega}^{\sigma_1,\sigma_2})$ is denoted by $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$.

Proposition 4.2. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with the condition (Mor2). Suppose that rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ satisfy the condition (Mor5). Then $Z^{\sigma_1,\sigma_2}_{\beta,\omega}$ is a central charge on $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$ for any $\beta \in \mathbb{R}$ and $\omega \in \mathbb{R}_{>0}$.

Proof. If an object $E \in \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is a (σ_1,σ_2) -torsion object, then $\Im \mathbb{I} Z_1(\tau_1^L E) = \Im \mathbb{I} Z_2(\tau_2^R E) = 0$. Hence we have $\Im \mathbb{I} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) = -\omega \Re \mathfrak{e} Z_2(\tau_2^R E) \ge 0$ and the equality holds if and only if $\tau_2^R E = 0$. Thus E is isomorphic to a σ_1 -torsion object $E_1 \in \mathcal{T}^{\sigma_1}$ and we see $\Re \mathfrak{e} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) = \Re \mathfrak{e} Z_1(E_1) \le 0$ and the equality holds if and only if $E_1 = 0$. Moreover if the object $E \in \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is (σ_1,σ_2) -free, the inequality $\mu_{\beta,\omega}(E) > 0$ implies $\Im \mathbb{I} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) > 0$.

Now take $E \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Without loss of generality, we may assume that the (σ_1, σ_2) -free object E is $M_{\beta,\omega}^{Z_1,Z_2}$ -semistable. Since $\Im m Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) \leq 0$ holds by the assumption $E \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$, we have to show the following:

• If $\mathfrak{Im} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) = 0$, then $\mathfrak{Re} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E)$ is positive.

Let us suppose $\mathfrak{Im} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) = 0$ holds. By the inequality $\omega > 0$, we have

(4.2)
$$\mathfrak{Re} Z_2(\tau_2^R E) = \frac{1}{\omega} \cdot (\mathfrak{Im} Z_1(\tau_1^L E) + \beta \, \mathfrak{Im} Z_2(\tau_2^R E))$$

Since the morphism $\tau_1^L E \to \Phi(\tau_2^R E)$ is a monomorphism in \mathcal{A}_1 by Lemma 3.9, the condition (Mor5) implies

(4.3)
$$\Im \mathfrak{m} Z_1(\tau_1^L E) \le \Im \mathfrak{m} Z_1\left(\Phi(\tau_2^R E)\right) = \Im \mathfrak{m} Z_2(\tau_2^R E)$$

Moreover there exists a subobject \tilde{F} of E in gl $(\mathcal{A}_1, \mathcal{A}_2)$ such that $\tau_1^L \tilde{F} \cong \tau_1^L E$ and $\Phi(\tau_2^R \tilde{F}) \cong \tau_1^L E$ by Lemma 2.25 and Remark 2.26. The $M_{\beta,\omega}^{Z_1,Z_2}$ -semistability of E implies $\mu_{\beta,\omega}^+(\tilde{F}) \leq \mu_{\beta,\omega}(E) \leq 0$.

Now the condition (Mor5) implies

$$\begin{split} \mu_{\beta,\omega}(\tilde{F}) &= \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L F) - \omega \, \mathfrak{Re} \, Z_2(\tau_2^R F) + \beta \, \Im \mathfrak{m} \, Z_2(\tau_2^R F)}{\Im \mathfrak{m} \, Z_1(\tau_1^L \tilde{F}) + \Im \mathfrak{m} \, Z_2(\tau_2^R \tilde{F})} \\ &= \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L \tilde{F}) - \omega \, \mathfrak{Re} \, Z_1(\Phi(\tau_2^R \tilde{F})) + \beta \, \Im \mathfrak{m} \, Z_1(\Phi(\tau_2^R \tilde{F}))}{\Im \mathfrak{m} \, Z_1(\tau_1^L \tilde{F}) + \Im \mathfrak{m} \, Z_1(\Phi(\tau_2^R \tilde{F}))} \\ &= \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L E) - \omega \, \mathfrak{Re} \, Z_1(\tau_1^L E) + \beta \, \Im \mathfrak{m} \, Z_1(\tau_1^L E)}{2 \, \Im \mathfrak{m} \, Z_1(\tau_1^L E)} \leq 0. \end{split}$$

Since $\Im \mathfrak{m} Z_1(\tau_1^L E)$ is positive, we see

(4.4)
$$\frac{1}{\omega} \left(\Im \mathfrak{m} \, Z_1(\tau_1^L E) + \beta \, \Im \mathfrak{m} \, Z_1(\tau_1^L E) \right) \leq \mathfrak{Re} \, Z_1(\tau_1^L E).$$

Thus (4.2), (4.3) and (4.4) imply

$$\begin{split} \mathfrak{Re} \ Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) &= \mathfrak{Re} \ Z_1(\tau_1^L E) + \beta \ \mathfrak{Re} \ Z_2(\tau_2^R E) + \omega \ \mathfrak{Im} \ Z_2(\tau_2^R E) \\ &= \mathfrak{Re} \ Z_1(\tau_1^L E) + \frac{\beta}{\omega} (\mathfrak{Im} \ Z_1(\tau_1^L E) + \beta \ \mathfrak{Im} \ Z_2(\tau_2^R E)) + \omega \ \mathfrak{Im} \ Z_2(\tau_2^R E) \\ &= \mathfrak{Re} \ Z_1(\tau_1^L E) + \frac{\beta}{\omega} \ \mathfrak{Im} \ Z_1(\tau_1^L E) + \frac{\beta^2 + \omega^2}{\omega} \ \mathfrak{Im} \ Z_2(\tau_2^R E) \\ &\geq \mathfrak{Re} \ Z_1(\tau_1^L E) + \frac{\beta}{\omega} \ \mathfrak{Im} \ Z_1(\tau_1^L E) + \frac{\beta^2 + \omega^2}{\omega} \ \mathfrak{Im} \ Z_1(\tau_1^L E) \\ &\geq \frac{1}{\omega} \left(\mathfrak{Im} \ Z_1(\tau_1^L E) + \beta \ \mathfrak{Im} \ Z_1(\tau_1^L E) \right) + \frac{\beta}{\omega} \ \mathfrak{Im} \ Z_1(\tau_1^L E) + \frac{\beta^2 + \omega^2}{\omega} \ \mathfrak{Im} \ Z_1(\tau_1^L E) \\ &= \frac{(1 + \beta)^2 + \omega^2}{\omega} \cdot \ \mathfrak{Im} \ Z_1(\tau_1^L E) > 0. \end{split}$$

Remark 4.3. Suppose that a semiorthogonal decomposition $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ satisfies (Mor2). If stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ satisfy (Mor5), then σ_2 is $\sigma_1[-1]$.

Proposition 4.4. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Suppose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \, \mathbf{D}_i$ are rational and satisfy (Mor5). If σ_1 and σ_2 are rational and both ω and β are rational then the pair $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} = (\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}, Z_{\beta,\omega}^{\sigma_1,\sigma_2})$ is a locally finite stability condition.

Proof. Since σ_i is rational, σ_i is discrete. In particular the heart \mathcal{A}_i of σ_i is Noetherian and so is $gl(\mathcal{A}_1, \mathcal{A}_2)$.

Suppose $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ satisfies $\Im \mathbb{I} \mathbb{I} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) = 0$. Due to [CP10, Lemma 3.4], it is enough to show that any increasing filtration $E^1 \subset E^2 \subset \cdots \subset E^n \subset \cdots$ of subobject of $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ terminates. Note that E^n also satisfies $\Im \mathbb{I} \mathbb{I} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^n) = 0$.

Taking the cohomology with respect to $gl(A_1, A_2)$, we have a sequence of monomorphisms

$$H^{-1}(E^n) \subset H^{-1}(E^{n+1}) \subset H^{-1}(E).$$

We can assume $H^{-1}(E^n)$ does not depend on n since $gl(\mathcal{A}_1, \mathcal{A}_2)$ is Noetherian. Now we claim $H^0(E^n)$ satisfies

(4.6)
$$\tau_1^L H^0(E^n) \in \mathcal{T}^{\sigma_1}.$$

Indeed, by the vanishing $\Im \mathfrak{Im} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^n) = 0$, we have $\Im \mathfrak{Im} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(H^0(E^n)) = 0$ which implies that $H^0(E^n)$ is (σ_1,σ_2) -torsion. The vanishing $\Im \mathfrak{Im} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(H^0(E^n)) = 0$ and the definition of $Z^{\sigma_1,\sigma_2}_{\beta,\omega}$ imply $\Im \mathfrak{Im} Z_1(\tau_1^L H^0(E^n)) = 0$ and $\tau_2^R H^0(E^n) = 0$. This gives the proof of the claim.

imply $\Im \mathfrak{I} Z_1(\tau_1^L H^0(E^n)) = 0$ and $\tau_2^R H^0(E^n) = 0$. This gives the proof of the claim. Let F^n be the quotient E^n/E^{n-1} in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Since $H^{-1}(E^n)$ is constant for n, we have a monomorphism in gl $(\mathcal{A}_1, \mathcal{A}_2)$

$$H^{-1}(F^n) \longrightarrow H^0(E^{n-1}).$$

Thus $\mathfrak{Im} \operatorname{gl}(Z_1, Z_2)(H^{-1}(F^n)) = 0$ holds by (4.5) and (4.6). On the other hand, if $H^{-1}(F^n)$ is non-zero, then $\mathfrak{Im} \operatorname{gl}(Z_1, Z_2)(H^{-1}(F^n))$ is positive by the (σ_1, σ_2) -freeness of $H^{-1}(F)$. So $H^{-1}(F^n)$ has to be 0 and the canonical morphism $H^0(E^{n-1}) \to H^0(E^n)$ is a monomorphism in $\operatorname{gl}(\mathcal{A}_1, \mathcal{A}_2)$.

To complete the proof, it is enough to show that $H^0(E^n)$ is independent of n. By (4.5) and (4.6), we have $\Re \mathfrak{e} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(H^0(E^n)) = \Re \mathfrak{e} Z_1(\tau_1^R H^0(E^n)) < 0$, it is enough to show that $\Re \mathfrak{e} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(H^0(E^n))$ is bounded below.

Let G^n be the quotient E/E^n in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Then we have the following exact sequence in $gl(\mathcal{A}_1,\mathcal{A}_2)$:

$$0 \longrightarrow H^{-1}(E^n) \longrightarrow H^{-1}(E) \xrightarrow{\varphi_n} H^{-1}(G^n) \xrightarrow{\delta_n} H^0(E^n) \xrightarrow{\psi_n} H^0(E) \xrightarrow{\pi_n} H^0(G^n) \longrightarrow 0$$

Since $H^{-1}(E^n) = \ker \varphi_n$ is independent of n, so is $\operatorname{im} \varphi_n = \ker \delta_n$. Moreover we may assume that $\ker \pi_n = \operatorname{im} \psi_n$ is independent of n since $\operatorname{gl}(\mathcal{A}_1, \mathcal{A}_2)$ is Noetherian. Then it is enough to show that $\operatorname{\mathfrak{Re}} Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\ker \psi_n)$ is bounded below since $H^0(E^n)$ is an extension by $\operatorname{im} \psi_n$ and $\ker \psi_n$:

$$0 \longrightarrow \ker \psi_n \longrightarrow H^0(E^n) \longrightarrow \operatorname{im} \psi_n \longrightarrow 0.$$

Note that $\tau_2^R(\ker \psi_n) = 0$ since the functor τ_2^R is *t*-exact with respect to $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ and \mathcal{A}_2 . Consider the exact sequence in $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$:

$$0 \longrightarrow \ker \delta_n \longrightarrow H^{-1}(G^n) \longrightarrow \ker \psi_n \longrightarrow 0.$$

Then Lemma 2.20 implies the following diagram of distinguished triangle in \mathbf{D}_1 :

The second and third row give short exact sequences in \mathcal{A}_1 . Since ker δ_n and $H^{-1}(G^n)$ are in $\mathcal{F}^{\sigma_1,\sigma_2}_{\beta,\omega}$, the morphism ν and ϵ are mono morphisms in \mathcal{A}_1 by Lemma 3.9. Thus we see the following isomorphism

$$\tau_1^L(\ker\psi_n) \cong \frac{\tau_1^L(H^{-1}(G^n))}{\tau_1^L(\ker\delta_n)} \cong \ker\left(\frac{\Phi(\ker\delta_n)}{\tau_1^L(\ker\delta_n)} \to \frac{\Phi(H^{-1}(G^n))}{\tau_1^L(H^{-1}(G^n))}\right).$$

The *t*-exactness of τ_1^L with respect to $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ and \mathcal{A}_1 implies $\tau_1^L(\ker \psi_n) \subset \tau_1^L(H^0(E^n))$. Since $H^0(E^n)$ satisfies (4.5) and (4.6), $\tau_1^L(\ker \psi_n)$ is in \mathcal{T}^{σ_1} and $\tau_2^R(\ker \psi_n) = 0$. Moreover $\tau_1^L(\ker \psi_n)$ is a subobject of the torsion part T of $\Phi(\ker \delta_n)/\tau_1^L(\ker \delta_n)$ with respect to the torsion pair $(\mathcal{T}^{\sigma_1}, \mathcal{F}^{\sigma_1})$.

Thus we see $Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\ker\psi_n) = Z_1(\tau_1^L(\ker\psi_n))$ and

(4.8)
$$\mathfrak{Re} Z_1(T) \leq \mathfrak{Re} Z_1(\ker \psi_n) = \mathfrak{Re} Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\ker \psi_n)$$

Since ker δ_n is independent of n, so is $\Phi(\ker \delta_n)/\tau_1^L(\ker \delta_n)$. Hence $\Re c Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\ker \psi_n)$ is bonded below and the pair $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} = (\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}, Z_{\beta,\omega}^{\sigma_1,\sigma_2})$ is a stability condition on **D**.

Finally by the rationality of σ_i and (β, ω) , the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ is rational. Since $K_0(\mathbf{D})$ is finitely generated, a rational stability condition is reasonable and in particular is locally finite.

5. Deformation property

Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Fixing $\sigma_i \in \mathsf{Stab} \mathbf{D}_i$ with (Mor5), we have constructed a family $\{\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} \mid \beta \in \mathbb{Q}, \omega \in \mathbb{Q}_{>0}\}$ of locally finite stability conditions on **D**. Now we wish to extend the family for real numbers and to show that the family is continuous for β and ω .

Although one of standard solutions is the support property of $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$, we show that the central charge $Z_{\beta',\omega'}^{\sigma_1,\sigma_2}$ for (β',ω') satisfies a finiteness condition

(5.1)
$$\|Z^{\sigma_1,\sigma_2}_{\beta',\omega'}\|_{\Sigma^{\sigma_1,\sigma_2}_{\beta,\omega}} < \infty$$

which has been established in Bridgeland's deformation of stability conditions (see also Theorem 2.10). To show this, by the definition of $Z^{\sigma_1,\sigma_2}_{\beta',\omega'}$, it is enough to show that the supremum

(5.2)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\middle|E \text{ is } \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}\text{-semistable}\right\}$$

is finite. This will be proven in Proposition 5.15. In the proposition, we only prove (5.2) for points (β, ω) in a contractible open set $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ of the upper half-plane \mathcal{H} . The restriction to $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ is necessary for Proposition 5.15 by a technical reason.

A key ingredient of Proposition 5.15 is Corollary 2.2. If the supremum of $|\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E)|$ for any semistable object E was smaller than π , then the desired assertion followed from Corollary 2.2. However, we do not see whether the canonical decomposition

$$i_2 \tau_2^R E \longrightarrow E \longrightarrow i_1 \tau_1^L E \longrightarrow i_2 \tau_2^R E[1]$$

satisfies the desired property or not since the decomposition is too "rough".

Thus a finer decomposition of the semistable object E is necessary. Roughly "finer decomposition" reflect the following principle:

• For a (σ_1, σ_2) -free object $F \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$, if the difference $\mu_{\beta,\omega}^+(F) - \mu_{\beta,\omega}^-(F)$ is sufficiently small, then the difference $|\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L F) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R F)|$ satisfies the assumption in Corollary 2.2.

To complete Proposition 5.15, it is necessary to observe properties of $\mu_{\beta,\omega}$ -semistable object.

Lemma 5.1. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Set a full subcategory of $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ by (5.3)

$$\mathcal{T}_{\beta,\omega}^{>1} = \{ E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2) \mid the \ (\sigma_1, \sigma_2) \text{-free part } F \text{ of } E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2) \text{ satisfies } \mu_{\beta,\omega}^-(F) > 1 \}.$$

(1) For any $E \in \mathcal{T}_{\beta,\omega}^{>1}$, the inequality $\arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E) \ge \arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$ holds. (2) The supremum

(5.4)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\middle|E\in\mathcal{T}_{\beta,\omega}^{>1}\right\}$$

is finite.

Proof. Let T be the (σ_1, σ_2) -torsion part of E and F the quotient E/T in $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$. The definition of $Z^{\sigma_1, \sigma_2}_{\beta, \omega}$ implies

(5.5)
$$\arg(-\beta + \sqrt{-1}\omega) \le \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T) \le \pi.$$

The assumption $\mu_{\beta,\omega}^{-}(F) > 1$ implies $\tau_1^L F = 0$ by the assertion (3) in Proposition 3.6. Hence we have

$$\mu_{\beta,\omega}(F) = \frac{-\omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R F) + \beta Z_2(\tau_2^R F)}{\operatorname{\mathfrak{Im}} Z_2(\tau_2^R F)} > 1$$

which implies $\Im \mathfrak{m} \left((\beta - 1 - \sqrt{-1}\omega) Z_2(\tau_2^R F) \right) > 0$. Thus we have

(5.6)
$$\arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega) < \arg(\beta - \sqrt{-1}\omega)Z_2(\tau_2^R F) = \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F).$$

Since $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E)$ is nothing but $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T) + Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F)$, the inequalities (5.5) and (5.6) give the proof of the assertion (1).

Since F satisfies $\tau_1^L F = 0$, we have $\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E) = \pi$ unless $\tau_1^L E$ is zero. Hence there exists a $\theta \in (0,\pi)$ such that the inequality

(5.7)
$$0 \le |\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E)| \le \theta$$

for any $E \in \mathcal{T}_{\beta,\omega}^{>1}$. Then Corollary 2.2 gives the proof of (2).

Lemma 5.2. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take ϵ_1 and ϵ_2 so that $\epsilon_2 \leq \epsilon_1 < 1$. Suppose a (σ_1, σ_2) -free object E satisfies $\mu^+_{\beta,\omega}(E) < 1$.

- (1) Suppose $\tau_1^L E \neq 0$. If E satisfies $\mu_{\beta,\omega}^+(E) \leq \epsilon_1$, then $\arg Z_1(\tau_1^L E) \leq \arg(\beta + 1 2\epsilon_1 + \sqrt{-1}\omega)$ for any $(\beta,\omega) \in \mathbb{R} \times \mathbb{R}_{>0}$.
- (2) If E satisfies $\mu_{\beta,\omega}^+(E) \leq \epsilon_1$, then $\arg Z_2(\tau_2^R E) \leq \arg(\beta \epsilon_1 + \sqrt{-1}\omega)$ for any $(\beta,\omega) \in \mathbb{R} \times \mathbb{R}_{>0}$.
- (3) If E satisfies $\epsilon_2 \leq \mu_{\beta,\omega}(E)$ then $\arg(\beta + 1 2\epsilon_2 + \sqrt{-1}\omega) \leq \arg Z_2(\tau_2^R E)$.

Proof. We first note that $\tau_2^R E$ is non-zero and in particular $\Im \mathbb{I} \mathbb{I} \mathbb{I} \mathbb{I} \mathbb{I} \mathbb{I} \mathbb{I} = 0$. Otherwise, E is in the essential image of $i_1: \mathbb{D}_1 \to \mathbb{D}$ and hence E is $\mu_{\beta,\omega}$ -semistable with $\mu_{\beta,\omega}(E) = 1$ by Proposition 3.6. This gives a contradiction.

Since $\mu_{\beta,\omega}^+(E) < 1$, the morphism $\tau_1^L E \to \Phi(\tau_2^R E)$ is a monomorphism in \mathcal{A}_1 by Lemma 3.9. Moreover Lemma 2.25 (and the proof) implies that there exists a subobject $F \subset E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ such that $\tau_1^L F \cong \tau_1^L E$ and $\Phi(\tau_2^R F) \cong \tau_1^L E$. Since the subobject F also satisfies $\mu_{\beta,\omega}^+(F) \leq \epsilon_1$ by the assumption for E, we see:

$$\begin{split} \epsilon_1 &\geq \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L F) - \omega \, \mathfrak{Re} \, Z_2(\tau_2^R F) + \beta \, \Im \mathfrak{m} \, Z_2(\tau_2^R F)}{\Im \mathfrak{m} \, Z_1(\tau_1^L F) + \Im \mathfrak{m} \, Z_2(\tau_2^R F)} \\ &= \frac{\Im \mathfrak{m} \, Z_1(\tau_1^L F) - \omega \, \mathfrak{Re} \, Z_1(\Phi(\tau_2^R F)) + \beta \, \Im \mathfrak{m} \, Z_1(\Phi(\tau_2^R F))}{\Im \mathfrak{m} \, Z_1(\tau_1^L F) + \Im \mathfrak{m} \, Z_1(\Phi(\tau_2^R F))} \\ &= \frac{-\omega \, \mathfrak{Re} \, Z_1(\tau_1^L E) + (\beta + 1) \, \Im \mathfrak{m} \, Z_1(\tau_1^L E)}{2 \, \Im \mathfrak{m} \, Z_1(\tau_1^L E)}. \end{split}$$

The last inequality implies

(5.8)
$$-\omega \mathfrak{Re} Z_1(\tau_1^L E) + (\beta + 1 - 2\epsilon_1) \mathfrak{Im} Z_1(\tau_1^L E) \le 0.$$

Since the left hand side of (5.8) is just the imaginary part of $(\beta + 1 - 2\epsilon_1 - \sqrt{-1}\omega)Z_1(\tau_1^L E)$, this gives the proof of the assertion (1).

Note that $i_2 \tau_2^R E$ is a subobject of E in gl $(\mathcal{A}_1, \mathcal{A}_2)$. Hence $i_2 \tau_2^R E$ also satisfies $\mu_{\beta,\omega}^+(i_2 \tau_2^R E) \leq \epsilon_1$. So we see

$$\epsilon_1 \ge \mu_{\beta,\omega}(i_2\tau_2^R E) = \frac{-\omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R E) + \beta \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E)}{\operatorname{\mathfrak{Im}} Z_2(\tau_2^R E)}.$$

Thus we have

(5.9)

$$-\,\omega\,\mathfrak{Re}\,Z_2(\tau_2^R E) + (\beta - \epsilon_1)\,\mathfrak{Im}\,Z_2(\tau_2^R E) \le 0$$

which gives the proof of the assertion (2).

We show the assertion (3). By the assumption $\epsilon_2 \leq \mu_{\beta,\omega}(E)$, we see

$$\begin{aligned} \epsilon_2(\Im\mathfrak{Im}\,Z_1(\tau_1^L E) + \Im\mathfrak{Im}\,Z_2(\tau_2^R E)) &\leq \Im\mathfrak{Im}\,Z_1(\tau_1^L E) - \omega\,\mathfrak{Re}\,Z_2(\tau_2^R E) + \beta\,\mathfrak{Im}\,Z_2(\tau_2^R E) \\ \iff (\epsilon_2 - 1)\,\mathfrak{Im}\,Z_1(\tau_1^L E) &\leq -\omega\,\mathfrak{Re}\,Z_2(\tau_2^R E) + (\beta - \epsilon_2)\,\mathfrak{Im}\,Z_2(\tau_2^R E). \end{aligned}$$

Lemma 3.9 and the condition (Mor5) implies $\Im \mathfrak{m} Z_1(\tau_1^L E) \leq \Im \mathfrak{m} Z_1(\Phi(\tau_2^R E)) = \Im \mathfrak{m} Z_2(\tau_2^R E)$. Hence we obtain

(5.10)
$$(\epsilon_2 - 1) \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E) \le -\omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R E) + (\beta - \epsilon_2) \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E)$$

(5.11)
$$\iff 0 \le -\omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R E) + (\beta - 2\epsilon_2 + 1) \operatorname{\mathfrak{Im}} Z_2(\tau_2^R E).$$

By the same argument for (1), we see $\arg(\beta - 2\epsilon_2 + 1 + \sqrt{-1}\omega) \leq \arg Z_2(\tau_2^R E)$.

Lemma 5.3. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \, \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take ϵ_1 and ϵ_2 which satisfy

$$0 \leq \epsilon_1, 0 < 1 - 2\epsilon_2, \epsilon_2 \leq \epsilon_1 < 1, and$$

(5.12)
$$0 < \omega^2 + (\beta + 1 - 2\epsilon_2)^2 + 2(1 - 2\epsilon_2)(\epsilon_2 - \epsilon_1).$$

Suppose that a (σ_1, σ_2) -free object $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ satisfies

(5.13)
$$\epsilon_2 \le \mu_{\beta,\omega}(E) \le \mu_{\beta,\omega}^+(E) \le \epsilon_1.$$

(1) There exists a $\theta \in (0,\pi)$ such that the argument of $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E)$ satisfies

$$0 < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E) - \arg(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega) \le \theta_0$$

for any E with the condition (5.13).

(2) The supremum

$$\sup\left\{\frac{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E)|}{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E)|}\right|E \text{ is } (\sigma_1,\sigma_2)\text{-free with } \epsilon_2 \leq \mu_{\beta,\omega}(E) \leq \mu^+_{\beta,\omega}(E) \leq \epsilon_1\right\}.$$

is finite.

Proof. Set $\theta_1 = \arg(\beta + 1 - 2\epsilon_1 + \sqrt{-1}\omega)$, $\theta_2 = \arg(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$, and $\theta_3 = \arg(\beta - \epsilon_1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$. Then the conditions $\epsilon_1 > 0$ and $1 - 2\epsilon_2 > 0$ imply that θ_1 is in the open interval $(0, \pi)$, θ_2 is non-positive, and θ_3 is in $[0, \pi)$.

Lemma 5.2 implies the following:

(5.14)
$$0 < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\tau_1^L E) \le \theta_1, \text{ and}$$

(5.15)
$$\theta_2 < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\tau_2^R E) \le \theta_3.$$

We claim that the assumption for (ϵ_1, ϵ_2) implies

$$(5.16) 0 < \theta_1 - \theta_2 < \pi.$$

Note that the inequalities (5.16) is equivalent to

(5.17)
$$\Im \mathfrak{m} \, \frac{\beta + 1 - 2\epsilon_1 + \sqrt{-1}\omega}{(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)} > 0.$$

Since the left hand side of (5.17) is $\omega \cdot (\omega^2 + (\beta + 1 - 2\epsilon_2)^2 + 2(1 - 2\epsilon_2)(\epsilon_2 - \epsilon_1))$ up to the positive constant $|(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)|^2$, the inequalities (5.16) hold.

Now the inequalities

$$(5.18) 0 < \theta_3 - \theta_2 < \pi$$

also hold as follows. Since $\Im \mathfrak{m}(\beta - \epsilon_1 + \sqrt{-1\omega})/(\beta + 1 + 2\epsilon_2 + \sqrt{-1\omega})$ is $\omega(1 + \epsilon_1 - 2\epsilon_2)$ up to positive constant, the assumption for ϵ_1 and ϵ_2 imply

$$1 + \epsilon_1 - 2\epsilon_2 \ge 1 + \epsilon_1 - 2\epsilon_1 = 1 - \epsilon_1 > 0.$$

The inequalities above imply (5.18).

Set θ_0 by max{ $\theta_1 - \theta_2, \theta_3 - \theta_2$ }. Since θ_2 is negative, the arguments arg $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E)$ and arg $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E)$ are in the interval $(\theta_2, \theta_0 + \theta_2]$. By inequalities (5.16) and (5.18), θ_0 is in $(0, \pi)$. Moreover (5.14) and (5.15) implies

$$\theta_2 < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E) \le \theta_0 + \theta_2.$$

Then Corollary 2.2 implies the assertion (2).

Remark 5.4. If $\epsilon_1 = 0$, then the inequality (5.12) is nothing but

(5.19)
$$\omega^2 + \beta^2 + (2\beta + 1)(2\epsilon_2 + 1) \ge 0.$$

Thus (β, ω) satisfies the (5.19) if $2\beta + 1 \ge 0$.

Proposition 5.5. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \, \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

For an $\epsilon_1 \in (0, 1/2)$, set an open subset $\mathcal{H}^+(\epsilon_1)$ by

(5.20)
$$\mathcal{H}^{+}(\epsilon_{1}) = \left\{ (\beta, \omega) \in \mathbb{R} \times \mathbb{R}_{>0} \middle| \begin{array}{l} 0 < \omega^{2} + (\beta + 1)^{2} - 2\epsilon_{1}, \text{ and} \\ 0 < \omega^{2} + (\beta + 1 - 2\epsilon_{1})^{2} + 2(1 - 2\epsilon_{1})(\epsilon_{1} - 1) \end{array} \right\}.$$

Set a full subcategory $\mathcal{T}_{\beta,\omega}^{<1}$ of gl $(\mathcal{A}_1, \mathcal{A}_2)$ by

(5.21)
$$\mathcal{T}_{\beta,\omega}^{<1} = \{ E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2) \mid E \text{ is } (\sigma_1, \sigma_2) \text{-free with } 0 < \mu_{\beta,\omega}^-(E) \le \mu_{\beta,\omega}^+(E) < 1 \}.$$

If $(\beta, \omega) \in \mathcal{H}^+(\epsilon_1)$,

(1) then there exists a $\theta \in (0, \pi)$ such that the following holds:

 $\sup\{\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E)\mid E\in \mathcal{T}^{<1}_{\beta,\omega}\}\leq \theta.$

(2) The supremum

(5.22)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\middle|E\in\mathcal{T}_{\beta,\omega}^{<1}\right\}$$

is finite.

Proof. Taking the Harder-Narasimhan filtration with respect to $\mu_{\beta,\omega}$ -semistability, we obtain a short exact sequence in gl $(\mathcal{A}_1, \mathcal{A}_2)$

$$(5.23) 0 \longrightarrow E^+ \longrightarrow E \longrightarrow E^- \longrightarrow 0,$$

where E^+ and E^- satisfy the following:

- E^+ is (σ_1, σ_2) -free with $\epsilon_1 < \mu^-_{\beta,\omega}(E^+) \le \mu^+_{\beta,\omega}(E^+) < 1$ and
- E^- is (σ_1, σ_2) -free with $0 < \mu_{\beta,\omega}^-(E^-) \le \mu_{\beta,\omega}^+(E^-) \le \epsilon_1$.

Now we note that two inequalities in (5.20) come from the inequality (5.12) in Lemma 5.3. Applying Lemma 5.3, there exist θ_+ and θ_- in $(0, \pi)$ such that $Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E^i) \leq \theta_i$ where $i \in \{+, -\}$. If set $\max\{\theta_+, \theta_-\}$ by θ , then the argument of $Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E) = Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E^+) + Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E^-)$ is in the interval $(0, \theta]$ and this gives the proof of the assertion (1) since θ is independent of the choice of $E \in \mathcal{T}^{<1}_{\beta, \omega}$.

Both arg $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^+)$ and arg $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-)$ are positive since $\mu_{\beta,\omega}(E^+)$ and $\mu_{\beta,\omega}(E^-)$ are positive. Moreover the inequality $|\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^+) - Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-)| \leq \theta$ holds. Then Proposition 2.1 implies that there exists a positive constant C > 0 such that the following holds:

(5.24)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)| \ge C|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^+)| + C|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)|.$$

By Lemma 5.2, there exist θ_0 and θ'_0 in the interval $(0, \pi)$ such that the following hold:

(5.25)
$$0 < |\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E^+) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E^+)| \le \theta_0, \text{ and}$$

(5.26)
$$0 < |\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L E^-) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E^-)| \le \theta'_0.$$

Hence we see

$$\begin{aligned} \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|} &\leq \frac{1}{C} \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^+) + Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)|} \\ &\leq \frac{1}{C} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)|} \right) \\ &\leq \frac{1}{C} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^+)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)|} \right) \end{aligned}$$

Corollary 2.2 implies that both $|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E^+)|/|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^+)|$ and $|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E^-)|/|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-)|$ are bounded above by (5.25) and (5.26). Hence the supremum is bounded above.

Remark 5.6. In the final section, we wish to deform the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ along a path from $(1,0) \in \overline{\mathcal{H}}$ to $(\cos 2\pi/3, \sin 2\pi/3) \in \mathcal{H}$. If set ϵ_1 by 1/3, then the set $\mathcal{H}^+(1/3)$ includes the desired path.

Corollary 5.7. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \, \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take $\epsilon_2 \leq 0$ and $(\beta, \omega) \in \mathbb{Q} \times \mathbb{Q}_{>0}$ so that

 $2\beta + 1 - 2\epsilon_2 > 0.$

(5.27)

Set $\mathcal{F}_{\beta,\omega}^{\leq \epsilon_2}$ by

$$\mathcal{F}_{\beta,\omega}^{\leq\epsilon_2} := \{ F \in \mathsf{gl}\left(\mathcal{A}_1, \mathcal{A}_2\right) \mid F \text{ is } (\sigma_1, \sigma_2) \text{-free with } \mu_{\beta,\omega}^+(F) \leq \epsilon_2 \}$$

and take $F \in \mathcal{F}_{\beta,\omega}^{\leq \epsilon_2}$.

(1) The argument of $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F)$ is greater than $\arg(\beta - \sqrt{-1}\omega)$. In particular the following holds for any $F \in \mathcal{F}^{\leq \epsilon_2}_{\beta,\omega}$:

(5.28)
$$\arg(\beta - \sqrt{-1}\omega) < Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F) \le 0$$

(2) The supremum

$$\sup\left\{\frac{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R F)|}{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F)|}\middle|F\in\mathcal{F}^{\leq\epsilon_2}_{\beta,\omega}\right\}$$

is finite.

Proof. The imaginary part $\Im \mathfrak{m} \frac{\beta+1-2\epsilon_2+\sqrt{-1}\omega}{\beta-\sqrt{-1}\omega}$ is $\omega(2\beta+1-2\epsilon_2)$ up to positive constant. Then The assumption $2\beta+1-2\epsilon_2 > 0$ directly implies

(5.29)
$$0 < \arg \frac{\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega}{\beta - \sqrt{-1}\omega} < \pi.$$

By the definition of $Z_{\beta,\omega}^{\sigma_1,\sigma_2}$, we have $\arg(-\beta + \sqrt{-1}\omega) < \arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)$. Lemma 5.2 implies

(5.30)
$$0 < \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L F) \le \arg(\beta + 1 - 2\epsilon_2 + \sqrt{-1\omega}), \text{ and}$$

(5.31)
$$\arg(\beta - \sqrt{-1}\omega) < \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R F) \le \arg(\beta - \epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega).$$

Since $\arg(\beta - \epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$ is negative, the inequalities (5.29), (5.30), and (5.31) imply $\arg(-\beta + \sqrt{-1}\omega) < Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F)$. In addition the inequalities $\mu^+_{\beta,\omega}(F) \leq \epsilon_2 < 0$ imply $\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F) \leq 0$, and thus we have

(5.32)
$$\arg(\beta - \sqrt{-1}\omega) < Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F) \le 0.$$

Finally, by (5.30) and (5.31), Corollary 2.2 implies the assertion (2).

Finally, by (5.30) and (5.31), Corollary 2.2 implies the assertion (2).

The following lemma might be technical for readers, but is necessary for the proof of Proposition 5.9

Lemma 5.8. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take $(\beta, \omega) \in \mathbb{Q} \times \mathbb{Q}_{>0}.$

- Suppose that $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ satisfies the following:
 - $\tau_1^L E$ is σ_1 -free, and
 - the (σ_1, σ_2) -free part $E_{\rm fr}$ of E has $\mu_{\beta,\omega}^-(E_{\rm fr}) \ge 1$.

The the following holds:

(1) The σ_2 -free part F of $\tau_2^R E$ has the property $\mu_{\beta,\omega}^-(i_2 F) \ge 1$.

(2)
$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R E) \ge \arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$$

Proof. Let E_{tor} be the (σ_1, σ_2) -torsion part of E. By the assumption we see $\tau_1^L E_{\text{tor}} = 0$ and (5.33) $\arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(E_{\text{tor}}) = \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\tau_2^R E_{\text{tor}}) = \arg(-\beta + \sqrt{-1}\omega).$

Let A be an arbitrary subobject of $\tau_2^R E$ such that the quotient $\tau_2^R E/A$ is σ_2 -free. Then $i_2 A$ gives a subobject of E. Hence the quotient $Q = E/i_2 A$ is (σ_1, σ_2) -free with $\mu_{\beta,\omega}^-(Q) \ge 1$. Then the inequalities $1 \le \mu_{\beta,\omega}^-(Q) \le \mu_{\beta,\omega}(Q)$ imply

(5.34)
$$-\omega \mathfrak{Re} Z_2(\tau_2^R Q) + (\beta - 1) \mathfrak{Im} Z_2(\tau_2^R Q) \ge 0,$$

which means $\arg Z_2(\tau_2^R Q) \ge \arg(\beta - 1 + \sqrt{-1}\omega)$. The inequality (5.34) also implies

(5.35)
$$\frac{-\omega \operatorname{\mathfrak{Re}} Z_2(\tau_2^R Q) + \beta \operatorname{\mathfrak{Im}} Z_2(\tau_2^R Q)}{\operatorname{\mathfrak{Im}} Z_2(\tau_2^R Q)} \ge 1.$$

Since A is arbitrary, the inequality (5.35) implies the assertion (1).

Recall $Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^{\vec{R}}E) = Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^{\vec{R}}E_{\text{tor}}) + Z_{\beta,\omega}^{\sigma_1,\sigma_2}(i_2F)$ and $Q = E/i_2$. Applying the inequality (5.34) A as σ_2 -torsion part of $\tau_2^R E$, we have

(5.36)
$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(i_2F) = \arg(\beta - \sqrt{-1}\omega)Z_2(F) \ge \arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega).$$

Then (5.33) and (5.36) imply the assertion (2).

Proposition 5.9. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take $(\beta, \omega) \in \mathcal{H}^+(\epsilon_1)_{\mathbb{Q}}$ for an $\epsilon_1 \in (0, 1/2)$ (cf. (5.20)).

Then the supremum

(5.37)
$$\sup\left\{\frac{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R T)|}{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T)|}\middle| T \in \mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega} \cap \mathcal{P}^{\sigma_1,\sigma_2}_{\beta,\omega}(\phi) \text{ where } \pi\phi \in (0,\arg(-\beta+\sqrt{-1}\omega)]\right\}$$

is finite.

Proof. We first note that the (σ_1, σ_2) -torsion part E_{tor} of any $E \in \mathcal{T}^{\sigma_1, \sigma_2}_{\beta, \omega}$ satisfies

$$\arg(-\beta + \sqrt{-1}\omega) \le \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E_{\mathrm{tor}}) \le \pi$$

by the definition of $Z^{\sigma_1,\sigma_2}_{\beta,\omega}$. Moreover the equality $\arg(-\beta + \sqrt{-1}\omega) = Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E_{\text{tor}})$ holds if and only if $\tau_1^L E_{\text{tor}} = 0$. Thus, for an object $T \in \mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega} \cap \mathcal{P}^{\sigma_1,\sigma_2}_{\beta,\omega}(\phi)$ with $\pi\phi \in (0, \arg(-\beta + \sqrt{-1}\omega)]$, we may assume that $\tau_1^L T$ is σ_1 -free since $\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T) \leq \arg(-\beta + \sqrt{-1}\omega)$.

Taking the Harder-Narasimhan filtration of T with respect to $\mu_{\beta,\omega}$ -semistability, we have the following short exact sequence in $gl(A_1, A_2)$

$$(5.38) 0 \longrightarrow T^+ \longrightarrow T \longrightarrow T^- \longrightarrow 0,$$

where T^+ and T^- satisfy

- the (σ_1, σ_2) -free part $T_{\rm fr}^+$ of T^+ has $\mu_{\beta,\omega}^-(T_{\rm fr}^+) \ge 1$, and
- T^- is (σ_1, σ_2) -free with $0 < \mu^-_{\beta,\omega}(T^-) \le \mu^+_{\beta,\omega}(T^-) < 1$.

Note that T^- also satisfies $\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T^-) > 0$. By Lemma 5.5, there exists a θ in $(0,\pi)$ such that the inequalities

(5.39)
$$0 < \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T^-) \le \theta$$

holds for any T.

Now the sequence (5.38) gives an exact sequence not only in $\mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega}$ but also in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$. Thus T^+ is a subobject of T in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$ and T^+ satisfies

(5.40)
$$0 < \phi^{-}(T^{+}) \le \phi^{+}(T^{+}) \le \phi^{+}(T) \le \arg(-\beta + \sqrt{-1}\omega)$$

with respect to the stability condition $\Sigma^{\sigma_{1},\sigma_{2}}_{\beta,\omega}$.

By Proposition 2.1, there exists a constant $C_1 > 0$ which is independent of T such that the inequality below holds:

(5.41)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T)| \ge C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)| + C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|.$$

Then, the inequality (5.41) implies the following:

(5.42)
$$\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T)|} \le \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|} \right)$$

(5.43)
$$\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|} \right)$$

(5.44)
$$\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|} \right)$$

By Proposition 5.5, there exists a positive constant $M_1 > 0$ such that the following holds:

(5.45)
$$\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^-)|} \le M_1$$

Taking semiorthogonal decomposition of T^+ , there exists a distinguished triangle below:

Since $\tau_1^L T_+$ is a subobject of $\tau_1^L T$ in \mathcal{A}_1 , $\tau_1^L T_+$ is σ_1 -free. Then, by Lemma 5.8, $\tau_2^R T$ is in $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and the inequality

(5.47)
$$\theta_1 \le \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\tau_2^R T^+)$$

holds where $\theta_1 = \arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$. In particular the distinguished triangle (5.46) gives a short exact sequence not only in gl $(\mathcal{A}_1, \mathcal{A}_2)$ but also in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Hence $\tau_2^R T^+$ is a subobject of T in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$, and the inequalities

(5.48)
$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R T^+) \le \pi \phi \le \theta_2 := \arg(-\beta + \sqrt{-1}\omega)$$

holds.

Now the closed interval $[\theta_1, \theta_2]$ is contained in $(0, \pi]$. Since $\arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(\tau_1^L T^+)$ is in the interval $(0,\pi]$, there exists a $\theta \in [0,\pi)$ such that the inequalities

$$0 \le |\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_1^L T^+) - \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R T^+)| \le \theta$$

hold for any T. By Corollary 2.2, there exists a positive constant M_2 which is independent of the choice of T and satisfies the following:

(5.49)
$$\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^+)|} = \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_1^L T^+) + Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^+)|} \le M_2.$$

Thus we have finished the proof.

Thus we have finished the proof.

Proposition 5.10. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). For an $\epsilon_2 < 0$ and set a subset $\mathcal{H}^-(\epsilon_2)$ of \mathcal{H} by

(5.50)
$$\mathcal{H}^{-}(\epsilon_{2}) := \left\{ (\beta, \omega) \in \mathcal{H} \middle| \begin{array}{l} 0 < 2\beta + 1 - 2\epsilon_{2}, \text{ and} \\ 0 < \omega^{2} + (\beta + 1 - 2\epsilon_{2})^{2} + 2\epsilon_{2}(1 - 2\epsilon_{2}) \end{array} \right\}.$$

Suppose that $(\beta, \omega) \in \mathcal{H}^{-}(\epsilon_2)$. Then the supremum

(5.51)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F)|}\middle|F\in\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}\right\}$$

is finite.

\

Proof. Choose $F \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Taking the Harder-Narasimhan filtration of F with respect to $\mu_{\beta,\omega}$ -semistability, we have the following short exact sequence in gl $(\mathcal{A}_1, \mathcal{A}_2)$

$$0 \longrightarrow F^+ \longrightarrow F \longrightarrow F^- \longrightarrow 0,$$

where F^+ and F^- satisfy

- F^+ is (σ_1, σ_2) -free with $\epsilon_2 < \mu^-_{\beta,\omega}(F^+) \le \mu^+_{\beta,\omega}(F^+) \le 0$, and
- F^- is (σ_1, σ_2) -free with $\mu^+_{\beta,\omega}(F^-) \leq \epsilon_2$.

Since $\mu_{\beta,\omega}^+(F^+) \leq 0$, applying Lemma 5.3 to F^+ , there exists a $\theta_+ \in (-\pi, 0)$ such that the inequalities hold:

(5.52)
$$\theta_+ < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F^+) \le 0.$$

Moreover, applying Corollary 5.7 to F^- , there exists a θ_- in the open interval $(-\pi, 0)$ such that the following holds:

(5.53)
$$\theta_{-} \leq Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F^-) \leq 0.$$

Thus one can use Proposition 2.1 by (5.52) and (5.53) and there exists a constant $C_1 > 0$ such that the following holds:

(5.54)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F)| \ge C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)| + C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|.$$

Similarly to the proof of Proposition 5.9, we see

$$\begin{aligned} \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F)|} &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|} \right) \\ &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|} \right) \\ &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^+)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^-)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|} \right) \end{aligned}$$

Now, by Lemma 5.3 and Corollary 5.7, there exist upper bounds for $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^+)|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^+)|$ and $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R F^-)|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^-)|$. This gives the proof.

Proposition 5.11. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take $(\beta, \omega) \in \mathcal{H}^+(\epsilon_1)$ for an $\epsilon_1 \in (0, 1/2)$.

Set $\mathcal{T}'_{\beta,\omega}$ by

(5.55)
$$\mathcal{T}_{\beta,\omega}' = \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2} \cap \mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2} \left[\frac{1}{\pi} \arg(-\beta + \sqrt{-1}\omega), 1 \right]$$

Then the supremum

(5.56)
$$\sup\left\{\frac{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(\tau_2^R T)|}{|Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T)|}\middle| T \in \mathcal{T}'_{\beta,\omega}\right\}$$

is finite.

Proof. Taking the Harder-Narasimhan filtration of $T \in \mathcal{T}'_{\beta,\omega}$ with respect to $\mu_{\beta,\omega}$ -semistability, there exists a short exact sequence in $gl(\mathcal{A}_1, \mathcal{A}_2)$

 $0 \longrightarrow T^{>1} \longrightarrow T \longrightarrow T^{\leq 1} \longrightarrow 0$

such that

• the (σ_1, σ_2) -free part $T_{\rm fr}^{>1}$ of $T^{>1}$ satisfies $\mu_{\beta,\omega}^-(T_{\rm fr}^{>1}) > 1$, and

•
$$T^{\leq 1}$$
 is (σ_1, σ_2) -free with $0 < \mu_{\beta,\omega}^-(T^{\leq 1}) \le \mu_{\beta,\omega}^+(T^{\leq 1}) \le 1$.

Then Lemma 5.1 implies

(5.57)

$$\arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega) \le \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(T^{>1}).$$

Moreover the canonical morphism $T \to T^{\leq 1}$ is an epimorphism not only in $\mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega}$ but also in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$. Hence the quotient $T^{\leq 1}$ in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$ also satisfies

(5.58)
$$\arg(-\beta + \sqrt{-1}\omega) \le Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T^{\le 1}).$$

Thus, by Proposition 2.1, there exists a constant C_1 such that the following holds:

(5.59)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T)| \ge C_1 \left(|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{>1})| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{\le 1})| \right)$$

Taking the Harder-Narasimhan filtration of $T^{\leq 1}$, we have the short exact sequence in gl $(\mathcal{A}_1, \mathcal{A}_2)$

$$0 \longrightarrow T^{=1} \longrightarrow T^{\leq 1} \longrightarrow T^{<1} \longrightarrow 0,$$

where $T^{=1}$ and $T^{<1}$ are

- $T^{=1}$ is (σ_1, σ_2) -free and is $\mu_{\beta,\omega}$ -semistable with $\mu_{\beta,\omega}(T^{=1}) = 1$, and
- $T^{<1}$ is (σ_1, σ_2) -free with $\mu^+_{\beta,\omega}(T^{<1}) < 1$.

Note that the inequality $\mu_{\beta,\omega}^{-}(T^{<1}) > 0$ also holds. Then the canonical morphism $T \to T^{<1}$ gives an epimorphism in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Thus $T^{<1}$ satisfies $\arg(-\beta + \sqrt{-1}\omega) < \arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{<1})$. By Proposition 5.5, there exists a $\theta_1 \in (0,\pi)$ such that the inequality $\arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{<1}) \leq \theta_1$ holds. Thus we obtain

(5.60)
$$\arg(-\beta + \sqrt{-1}\omega) < \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(T^{<1}) \le \theta_1$$

and we see that there exists a $\theta \in [0, \pi)$ such that $|\arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(T^{=1}) - \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(T^{<1})| < \theta$. By Proposition 2.1, there exists a positive constant C_2 such that the following holds:

(5.61)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{\leq 1})| \ge C_2 \left(|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{=1})| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{<1})| \right).$$

Thus we see

$$\begin{aligned} \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T)|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T)|} &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T)|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})| + |Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})| + |Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})| + |Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(\tau_{2}^{R}T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} + \frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|}{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{<1})|} \right) \\ &\leq \frac{1}{C_{1}} \left(\frac{|Z_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(T^{>1})|}{|Z_{\beta,\omega}^{$$

By Lemma 5.1, there exists an upper bound for $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^{>1})|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{>1})|$. The assumption $\mu_{\beta,\omega}(T^{=1}) = 1$ implies $\arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^{=1}) = \arg(\beta - 1 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$ and hence the difference $|\arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_1^L T^{=1}) - \arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^{=1})|$ is strictly smaller than π . Hence there exists an upper bound for $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^{=1})|/C_2|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{=1})|$ by Corollary 2.2. Finally, by Lemma

5.3, there exists an upper bound for $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R T^{<1})|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T^{<1})|$. Hence (5.62) is bounded above.

Lemma 5.12. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). If (β, ω) is in $\mathcal{H}^-(\epsilon_2)$ then the inequality

$$\arg(-\beta + \sqrt{-1}\omega) < Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F[1])$$

holds for any $F \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$.

Proof. Taking the Harder-Narashimhan filtration of $F \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$, we obtain the short exact sequence

 $0 \longrightarrow F^+ \longrightarrow F \longrightarrow F^- \longrightarrow 0,$

where F^+ and F^- satisfy

- F^+ is (σ_1, σ_2) -free with $\epsilon_2 < \mu^-_{\beta,\omega}(F^+) \le \mu^+_{\beta,\omega}(F^+) \le 0$, and
- F^- is (σ_1, σ_2) -free with $\mu^+_{\beta,\omega}(F^-) \leq \epsilon_2$.

In particular we have $\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F^{\pm}) \leq 0.$

Since ϵ_2 is negative, Lemma 5.3 implies

(5.63)
$$\arg(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega) \le \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F^+) \le 0.$$

In addition Corollary 5.7 implies

(5.64)
$$\arg(\beta - \sqrt{-1}\omega) < \arg Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F^-) \le 0.$$

Since the inequality $\arg(\beta - \sqrt{-1}\omega) < \arg(\beta + 1 - 2\epsilon_2 + \sqrt{-1}\omega)(\beta - \sqrt{-1}\omega)$ clearly holds, we obtain

$$\arg(\beta - \sqrt{-1}\omega) < Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F) \le 0$$

which gives the proof.

Proposition 5.13. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Suppose that (β, ω) is in $\mathcal{H}^-(\epsilon_2)$,

(Mor5). Suppose that (β, ω) is in $\mathcal{H}^{-}(\epsilon_{2})$, Let $E \in \mathcal{A}_{\beta,\omega}^{\sigma_{1},\sigma_{2}}$ be $\Sigma_{\beta,\omega}^{\sigma_{1},\sigma_{2}}$ -semistable with the phase ϕ . If $\pi\phi \leq \arg(-\beta + \sqrt{-1}\omega)$, then E is in $\mathcal{T}_{\beta,\omega}^{\sigma_{1},\sigma_{2}}$. In particular, $\mathcal{P}_{\beta,\omega}^{\sigma_{1},\sigma_{2}}(0,\arg(-\beta + \sqrt{-1}\omega)/\pi]$ is the full subcategory of $\mathcal{T}_{\beta,\omega}^{\sigma_{1},\sigma_{2}}$.

Proof. Since the pair $(\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2},\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1])$ is a torsion pair on $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$, there is an short exact sequence

 $0 \longrightarrow F[1] \longrightarrow E \longrightarrow T \longrightarrow 0,$

where $T \in \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and $F \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. By the $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ -semistability of E, we have

(5.65)
$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(F[1]) \le \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) \le \arg(-\beta + \sqrt{-1}\omega).$$

Then the inequalities (5.65) contradict Lemma 5.12. Hence F[1] has to be zero. The last assertion follows from the definition of $\mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(0, \arg(-\beta + \sqrt{-1}\omega)/\pi]$.

Proposition 5.14. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

Then the supremum

(5.66)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\middle|E\in\mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(\phi)\ where\ \pi\phi\in\left(\arg(-\beta+\sqrt{-1}\omega),\pi\right]\right\}$$

is finite if (β, ω) is in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$.

Proof. Let *E* be in the slicing $\mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(\phi)$ where $\pi\phi$ is in $\left(\arg(-\beta + \sqrt{-1}\omega),\pi\right]$. Since $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$, there exists a canonical distinguished triangle

 $E^{-1}[1] \longrightarrow E \longrightarrow E^0$

where $E^0 \in \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and $E^{-1} \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Then E^{-1} satisfies $\arg Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1]) > \arg(-\beta + \sqrt{-1}\omega)$ by Lemma 5.12. Since the morphism $E \to E^0$ is an epimorphism in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$, we see

(5.67)
$$\phi(E^0) \ge \phi^-(E^0) \ge \phi^-(E) = \phi(E) \ge \frac{1}{\pi} \arg(-\beta + \sqrt{-1}\omega).$$

Hence inequalities $0 \leq |\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^0) - Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^{-1}[1])| < \pi$ hold and there exists a positive constant C_1 such that the inequality

(5.68)
$$|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)| \ge C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)| + C_1 |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|$$

holds for any E. Then we see

$$\begin{aligned} \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|} &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|} \right) \\ &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^0)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R (E^{-1}[1]))|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|} \right) \\ &= \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(T_2^R E^0)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)| + |Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R (E^{-1}[1]))|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|} \right) \\ &\leq \frac{1}{C_1} \left(\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E^0)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)|} + \frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R (E^{-1}[1]))|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|} \right). \end{aligned}$$

Since E^{-1} is in $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$, Proposition 5.10 implies that $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R(E^{-1}[1]))|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^{-1}[1])|$ is bounded above. Moreover, by Proposition 5.11, $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R(E^0))|/|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^0)|$ is bounded above.

Proposition 5.15. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

If (β, ω) be in $(\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2))_{\mathbb{Q}}$ then the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ satisfies that the supremum

(5.69)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\right|E \text{ is } \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}\text{-semistable}\right\}$$

is finite.

Proof. Let E be a $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ -semistable object with the phase ϕ . Without loss of generality, we may assume that $\phi \in (0, 1]$.

If $\pi \phi \leq \arg(-\beta + \sqrt{-1}\omega)$, then *E* is in $\mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega}$ by Proposition 5.13. Then, by Proposition 5.9, there exists an $M_1 > 0$ such that

(5.70)
$$\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|} < M_1$$

Suppose $\pi \phi > \arg(-\beta + \sqrt{-1}\omega)$. By Proposition 5.14, we see the supremum

(5.71)
$$\sup\left\{\frac{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\right|E \text{ is } \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}\text{-semistable with } \pi\phi > \arg(-\beta + \sqrt{-1}\omega)\right\}$$

is bounded. Hence we have finished the desired assertion.

Corollary 5.16. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

If a rational points (β, ω) is in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$, then the supremum

(5.72)
$$\sup\left\{\frac{|Z_2(\tau_2^R E)|}{|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)|}\middle|E \text{ is } \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}\text{-semistable}\right\}$$

 $is\ finite.$

Proof. By the definition, we see $|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(\tau_2^R E)| = |\beta - \sqrt{-1}\omega| \cdot |Z_2(\tau_2^R E)|$ for any $E \in \mathbf{D}$. Thus Proposition 5.15 implies the desired assertion.

Now we will show that the following map

(5.73)
$$s_{\mathbb{Q}} \colon \mathcal{H}^{+}(\epsilon_{1})_{\mathbb{Q}} \cap \mathcal{H}^{-}(\epsilon_{2})_{\mathbb{Q}} \to \operatorname{Stab} \mathbf{D}, s_{\mathbb{Q}}(\beta, \omega) = \Sigma_{\beta, \omega}^{\sigma_{1}, \sigma_{2}}$$

is continuous.

Lemma 5.17. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Suppose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ satisfies (Mor5). Take $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and $E' \in \mathcal{A}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ where (β,ω) and $(\beta',\omega') \in \mathbb{R} \times \mathbb{R}_{>0}$. Then the vanishing $\operatorname{Hom}_{\mathbf{D}}(E[p], E') = 0$ holds for any $p \geq 2$.

Proof. Let \mathcal{P}^{σ_i} be the slicing of σ_i . By the construction of the heart $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$, if E is in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$ then $\tau_1^L E \in \mathcal{P}^{\sigma_1}(0,2]$ and $\tau_2^R E \in \mathcal{P}^{\sigma_2}(0,2]$. Since σ_i satisfies (Mor3), we see $\Phi(\tau_2^R E) \in \mathcal{P}^{\sigma_1}(0,2]$. Thus the vanishings

(5.74)
$$\begin{cases} \operatorname{Hom}_{\mathbf{D}_{1}}(\tau_{1}^{L}E[p],\tau_{1}^{L}E') = 0\\ \operatorname{Hom}_{\mathbf{D}_{2}}(\tau_{2}^{R}E[p],\tau_{2}^{R}E') = 0\\ \operatorname{Hom}_{\mathbf{D}_{1}}(\tau_{1}^{L}E[p],\Phi(\tau_{2}^{R}E')[-1]) = 0 \end{cases}$$

hold. Then Lemma 2.21 implies the desired assertion.

Proposition 5.18. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Take $a \ (\beta, \omega) \in (\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2))_{\mathbb{Q}}$ for an $\epsilon_1 \in (0, 1/2)$ and an $\epsilon_2 < 0$.

For an arbitrary $\varepsilon \in (0, \varepsilon_0)$, choose $(\beta', \omega') \in (\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2))_{\mathbb{Q}}$ so that the following holds:

(5.75)
$$\sup\{|\mu_{\beta',\omega'}^+(E) - \mu_{\beta,\omega}^+(E)|, |\mu_{\beta',\omega'}^-(E) - \mu_{\beta,\omega}^-(E)| \mid E \text{ is } (\sigma_1, \sigma_2)\text{-free}\} < \varepsilon.$$

(1) Let $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ be $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ -semistable with

$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) > \arg(\beta+1+\sqrt{-1}\omega)(\beta-\sqrt{-1}\omega) + \pi$$

Then the vanishing $\operatorname{Hom}_{\mathbf{D}}(E[1], E') = 0$ holds for any $E' \in \mathcal{A}_{\beta', \omega'}^{\sigma_1, \sigma_2}$.

(2) Let $E \in \mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ be $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ -semistable with

$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E[-1]) \le \arg(\beta+1+2\epsilon_0+\sqrt{-1}\omega)(\beta-\sqrt{-1}\omega).$$

Then the vanishing $\operatorname{Hom}_{\mathbf{D}}(E', E[-1]) = 0$ holds for any $E' \in \mathcal{A}_{\beta', \omega'}^{\sigma_1, \sigma_2}$.

Proof. Throughout the proof, let T (resp. T') be the free part of E (resp. E') and let F[1] (resp. F'[1]) be the torsion part of E (resp. E') with respect to the torsion pair $(\mathcal{F}^{\sigma_1,\sigma_2}_{\beta,\omega}[1],\mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega})$ on the heart $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta,\omega}$ (resp. on $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta',\omega'}$). Then one easily see $\operatorname{Hom}_{\mathbf{D}}(E[1],E') \cong \operatorname{Hom}_{\mathbf{D}}(T,F')$.

Basically the proof of the assertions are similar. Since F' is in $\mathcal{F}^{\sigma_1,\sigma_2}_{\beta',\omega'}$, we have $\mu^+_{\beta,\omega}(F') \leq \epsilon$ by (5.75). By truncating the Harder-Narasimhan filtration of F' with respect to the $\mu_{\beta,\omega}$ -semistability, there exists a subobject F'_+ of F' in gl $(\mathcal{A}_1, \mathcal{A}_2)$ such that

- F'_+ is (σ_1, σ_2) -free with $0 < \mu^-_{\beta,\omega}(F'_+) \le \mu^+_{\beta,\omega}(F'_+) \le \epsilon$,
- the quotient $F'_{-} := F'/F'_{+}$ in $\overrightarrow{\mathsf{gl}}(\mathcal{A}_1, \mathcal{A}_2)$ is (σ_1, σ_2) -free with $\mu^+_{\beta, \omega}(F'_{-}) \leq 0$.

Then any morphism $\varphi \colon T \to F'$ factors through F'_+ , that is, we have the following diagram:




Note that $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is not only the free part of $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ but also the torsion part of $\mathfrak{gl}(\mathcal{A}_1,\mathcal{A}_2)$. Since the morphism $\varphi': T \to F'_+$ above lives in $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ which is closed under subobjects in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$, the image im φ' of φ' in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ belongs to $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$.

Again by truncating the Harder-Narasimhan filtration of $\operatorname{im} \varphi'$ with respect to the $\mu_{\beta,\omega}$ semistability, we obtain a subobject G of $\operatorname{im} \varphi'$ in the abelian category $\operatorname{gl}(\mathcal{A}_1, \mathcal{A}_2)$ such that

- the (σ_1, σ_2) -free part $G_{\rm fr}$ of G_+ satisfies $\mu_{\beta,\omega}^-(G_{\rm fr}) > \epsilon$, and
- the quotient $G_{-} := \operatorname{im} \varphi'/G_{+}$ is (σ_{1}, σ_{2}) -free with $0 < \mu_{\beta,\omega}^{-}(G_{-}) \le \mu_{\beta,\omega}^{+}(G_{-}) \le \epsilon$.

Thus we have the diagram:



The vanishing $\operatorname{Hom}_{\mathbf{D}}(G_+, F'_+) = 0$ gives the lift of $\operatorname{im} \varphi' \to F'_+$ to G^- . The horizontal line gives a short exact sequence not only in $\operatorname{\mathsf{gl}}(\mathcal{A}_1, \mathcal{A}_2)$ but also in $\mathcal{A}^{\sigma_1, \sigma_2}_{\beta, \omega}$ since all objects are in $\mathcal{T}^{\sigma_1, \sigma_2}_{\beta, \omega}$. Hence the composite $E \to T \to \operatorname{im} \varphi' \to G_-$ gives an epi morphism in $\mathcal{A}^{\sigma_1, \sigma_2}_{\beta, \omega}$, in particular $G_$ is a quotient of E. So the argument $\operatorname{arg} Z^{\sigma_1, \sigma_2}_{\beta, \omega}(G^-)$ should satisfy

(5.76)
$$\arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(G^-) \ge \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E) \ge \arg(\beta+1+\sqrt{-1}\omega)(\beta-\sqrt{-1}\omega)+\pi.$$

Since G^- satisfies $0 < \mu^-_{\beta,\omega}(G_-) \le \mu^+_{\beta,\omega}(G_-) \le \epsilon$, the inequality (5.76) contradicts Lemma 5.3. Thus we have finished the proof of the assertion (1).

Similarly to the proof for (1), we easily see $\operatorname{Hom}_{\mathbf{D}}(E', E[-1]) \cong \operatorname{Hom}_{\mathbf{D}}(T', F)$. Since the (σ_1, σ_2) -free part of $T' \in \mathcal{T}^{\sigma_1, \sigma_2}_{\beta', \omega'}$ satisfies $\mu^-_{\beta, \omega}(T') > -\epsilon$, there exists a subobject T'_+ of T' in $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)$ such that

- the (σ_1, σ_2) -free part $T'_{\rm fr}$ of T'_+ satisfies $\mu^-_{\beta,\omega}(T'_{\rm fr}) > 0$, and
- the quotient $T'_{-} := T'/T'_{+}$ is (σ_1, σ_2) -free with $-\epsilon < \mu^-_{\beta,\omega}(T'_{-}) \le \mu^+_{\beta,\omega}(T'_{-}) \le 0$.


Then any morphism $\psi: T' \to F$ lifts to $\psi': T'_{-} \to F$ by $\mu^{+}_{\beta,\omega}(F) \leq 0$:



Since the morphism ψ' is in the torsion part $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$ of the abelian category $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}[-1]$, the image $\operatorname{im} \psi'$ of ψ' in $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}[-1]$ is also in $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. Truncating the Harder-Narasimhan filtration of $\operatorname{im} \psi'$ with respect to the $\mu_{\beta,\omega}$ -semistability, we find a subobject $H_+ \subset \operatorname{im} \psi'$ such that

- H_+ is (σ_1, σ_2) -free with $-\epsilon < \mu_{\beta,\omega}^- \le \mu_{\beta,\omega}^+(H_+) \le 0$, and
- the quotient $H_{-} := \operatorname{im} \psi'/H_{+}$ is (σ_{1}, σ_{2}) -free with $\mu_{\beta,\omega}^{+}(H_{-}) \leq -\epsilon$.

Thus we have the diagram:



By the assumption for T'_{-} , the morphism $T'_{-} \to \operatorname{im} \psi'$ factors through H_{+} . Moreover the composite $H_{+} \to \operatorname{im} \psi' \to F \to E[-1]$ is a mono morphism in $\mathcal{A}^{\sigma_{1},\sigma_{2}}_{\beta,\omega}$ since any object in the horizontal line lives in the torsion part $\mathcal{F}^{\sigma_{1},\sigma_{2}}_{\beta,\omega}$ of $\mathcal{A}^{\sigma_{1},\sigma_{2}}_{\beta,\omega}[-1]$. Hence H_{+} should satisfies

(5.77)
$$0 < \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(H_+) \le \arg Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E[-1]) \le \arg(\beta+1+2\epsilon_0+\sqrt{-1}\omega)(\beta-\sqrt{-1}\omega).$$

Then the inequality above gives a contradiction applying Lemma 5.3 as $\epsilon_2 = -\epsilon_0$ and $\epsilon_1 = 0$. \Box

Proposition 5.19. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Then the map $s_{\mathbb{Q}} \colon \mathcal{H}^+(\epsilon_1)_{\mathbb{Q}} \cap \mathcal{H}^-(\epsilon_2)_{\mathbb{Q}} \to \operatorname{Stab} \mathbf{D}$ defined by

(5.78)
$$s_{\mathbb{Q}}(\beta,\omega) = \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$$

Proof. We first show that $\mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1] \subset \mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(-1,2]$ for any (β,ω) and (β',ω') in $\mathbb{R} \times \mathbb{R}_{>0}$. To show this, Let E be in $\mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(0,1]$ and E' in $\mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1]$. We denote by \mathcal{A}_0 the heart $\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2)$ on \mathbf{D} . Then cohomologies of E and of E' with respect to \mathcal{A}_0 are concentrated in degree -1 and 0. Hence the vanishings $\operatorname{Hom}_{\mathbf{D}}(E[p], E') = \operatorname{Hom}_{\mathbf{D}}(E', E[-p]) = 0$ hold for any $p \geq 2$ by Lemma 5.17. Thus we see $\mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1] \subset \mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(-1,2]$.

Take $0 < \varepsilon < 1/8$ arbitrary. By Proposition 5.18, if (β', ω') is sufficiently close, then the inclusion $\mathcal{A}_{\beta',\omega'}^{\sigma_1,\sigma_2} \subset \mathcal{P}_{\beta,\omega}^{\sigma_1,\sigma_2}(-1+\varepsilon, 2-\varepsilon]$ holds. By Corollary 5.16, shrinking (β', ω') if necessary, we may assume that the following holds:

(5.79)
$$\sup\left\{\frac{\left|Z_{\beta',\omega'}^{\sigma_1,\sigma_2}(E) - Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)\right|}{\left|Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E)\right|}\right| E \text{ is } \Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}\text{-semistable}\right\} < \sin(\pi\varepsilon).$$

By [CP10, Proposition 4.2] we see that $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ is in the open neighborhood $B_{\varepsilon}(\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2})$ and this gives the proof.

6. Non-rational coefficients stability conditions

In the previous section, we have constructed stability conditions $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ for rational points (β,ω) in the open set $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$. These stability conditions $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ could be deformed for non-rational points in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ by Proposition 5.15 and Theorem 2.10. We wish to show that the extended stability condition is given by the same construction as rational stability conditions $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ and that the family $\{\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2} \mid (\beta,\omega) \in \mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)\}$ is continuous for (β,ω) .

Lemma 6.1. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Suppose $\tau = (Z_{\beta,\omega}^{\sigma_1,\sigma_2}, \mathcal{Q})$ is a stability condition obtained by a deformation of $\Sigma_{\beta_0,\omega_0}^{\sigma_1,\sigma_2}$ for a rational point $(\beta_0, \omega_0) \in \mathcal{H}^+(\epsilon_1)_{\mathbb{Q}} \cap \mathcal{H}^-(\epsilon_2)_{\mathbb{Q}}$ (see also Definition 2.11). Any (σ_1, σ_2) -torsion object is in $\mathcal{Q}(0, 1]$.

Proof. Let U be an open neighborhood of (β, ω) in \mathcal{H} . By the definition of $\Sigma_{\beta',\omega'}^{\sigma_1,\sigma_2}$ for any rational point $(\beta', \omega') \in U_{\mathbb{Q}}$, any (σ_1, σ_2) -torsion object T is in $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}$, in particular, in $\mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1]$. Taking limit (β', ω') to (β, ω) , we see that T is in $\mathcal{Q}[0,1]$.

Now we wish to show that T is in $\mathcal{Q}(0,1]$. To show this, let $\phi_{\beta',\omega'}(T)$ be the phase of the maximal destabilizing quotient of T with respect to $\Sigma_{\beta',\omega'}^{\sigma_1,\sigma_2}$ for $(\beta',\omega') \in U_{\mathbb{Q}}$. We claim

(6.1)
$$\phi_{\beta',\omega'}^{-}(T) \geq \frac{1}{\pi} \arg(-\beta' + \sqrt{-1}\omega').$$

In fact, otherwise, there exists a subobject K of T in $\mathcal{A}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ such that the quotient Q = T/K is $\Sigma_{\beta',\omega'}^{\sigma_1,\sigma_2}$ -semistable with $\arg Z_{\beta',\omega'}^{\sigma_1,\sigma_2}(Q) < \arg(-\beta' + \sqrt{-1}\omega')$. By Proposition 5.13, Q is in $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}$. Moreover, K is also in $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ since $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ is the free part of $\mathcal{A}_{\beta',\omega'}^{\sigma_1,\sigma_2}$. Hence the sequence

is exact not only in $\mathcal{A}^{\sigma_1,\sigma_2}_{\beta',\omega'}$ but also in $\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2)$. Thus Q has to be (σ_1,σ_2) -torsion. Then the definition of $Z^{\sigma_1,\sigma_2}_{\beta',\omega'}$ implies the inequality

$$\arg(-\beta' + \sqrt{-1}\omega') \le \arg Z^{\sigma_1,\sigma_2}_{\beta',\omega'}(Q)$$

which contradicts the assumption for Q.

Hence $\phi^{-}(T)$ has to satisfy (6.1). Since one can choose arbitrary close $(\beta', \omega') \in U_{\mathbb{Q}}$ to (β, ω) , T belongs to $\mathcal{Q}(0, 1]$.

Lemma 6.2. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Suppose $\tau = (Z_{\beta,\omega}^{\sigma_1,\sigma_2}, \mathcal{Q})$ is a stability condition obtained by a deformation of $\Sigma_{\beta_0,\omega_0}^{\sigma_1,\sigma_2}$ for a rational point $(\beta_0, \omega_0) \in \mathcal{H}^+(\epsilon_1)_{\mathbb{Q}} \cap \mathcal{H}^-(\epsilon_2)_{\mathbb{Q}}$. Then $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2} \subset \mathcal{Q}(0,1]$ and $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2} \subset \mathcal{Q}(-1,0]$.

Proof. We first show $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2} \subset \mathcal{Q}(0,1]$. Let E be in $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$. If E is (σ_1,σ_2) -torsion, then E is in $\mathcal{Q}(0,1]$ by Lemma 6.1. Now suppose that the object E is (σ_1,σ_2) -free and $\mu_{\beta,\omega}$ -semistable. Since $\mathcal{Q}(0,1]$ is closed under extension, it is enough to show that such an E belongs to $\mathcal{Q}(0,1]$.

Since the $\mu_{\beta,\omega}$ -stability satisfies the support property by Proposition 3.6, there exists an open neighborhood U of (β,ω) such that $\mu_{\beta',\omega'}^{-}(E) > 0$ for any $(\beta',\omega') \in U$. Hence E is in $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ which is a subcategory of $\mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1]$ for any $(\beta',\omega') \in U_{\mathbb{Q}}$. Taking the limit (β',ω') to (β,ω) , we see that E is in $\mathcal{Q}[0,1]$.

Now we claim the following:

Claim 6.3. Notations being as above, any object in $\mathcal{Q}(0,1]$ is quasi isomorphic to a 2-term complex concentrated in degree 0 and -1 with respect to the heart $\mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$.

Take $F \in \mathcal{Q}(\phi)$ where $\phi \in (0, 1]$. Since the deformation of a stability condition is locally unique, we can find a rational point $(\beta'', \omega'') \in \mathcal{H}^+(\epsilon_1)_{\mathbb{Q}} \cap \mathcal{H}^-(\epsilon_2)_{\mathbb{Q}}$ such that $\arg(-\beta + \sqrt{-1}\omega) < \arg(-\beta'' + \sqrt{-1}\omega'')$ and the stability condition $\Sigma_{\beta'',\omega''}^{\sigma_1,\sigma_2}$ satisfies $\tau \in B_{\theta}(\Sigma_{\beta'',\omega''}^{\sigma_1,\sigma_2})$ for some $\theta > 0$. Then F is in $\mathcal{P}_{\beta'',\omega''}^{\sigma_1,\sigma_2}(\phi - \theta, \phi + \theta)$. Shrinking (β'', ω'') if necessary, we may assume $\phi - \theta > 0$ and $\phi + \theta < \arg(-\beta + \sqrt{-1}\omega)/\pi + 1$. Then we have the sequence

$$[\phi - \theta, \phi + \theta] \subset (0, \arg(-\beta + \sqrt{-1}\omega)/\pi + 1) \subset (0, \arg(-\beta'' + \sqrt{-1}\omega'')/\pi + 1).$$

Hence the object F is given by an extension of objects in $\mathcal{P}_{\beta'',\omega''}^{\sigma_1,\sigma_2}(0,1]$ and $\mathcal{P}_{\beta'',\omega''}^{\sigma_1,\sigma_2}(1, \arg(-\beta'' + \sqrt{-1}\omega'')/\pi + 1]$. By Proposition 5.13, any object in $\mathcal{P}_{\beta'',\omega''}^{\sigma_1,\sigma_2}(1, \arg(-\beta'' + \sqrt{-1}\omega'')/\pi + 1]$ is in $\mathcal{T}_{\beta'',\omega''}^{\sigma_1,\sigma_2}[1]$, in particular, is concentrated in degree -1. Since any object in $\mathcal{P}_{\beta'',\omega''}^{\sigma_1,\sigma_2}(0,1]$ is concentrated in degree -1 and 0, so is F.

To show $\mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega} \subset \mathcal{Q}(0,1]$, consider the distinguished triangle

$$(6.3) E^+ \longrightarrow E^- \longrightarrow E^+[1]$$

where $E^+ \in \mathcal{Q}(0,1]$ and $E^- \in \mathcal{Q}(0)$. It is enough to show that E^- is zero. Taking the cohomology with respect to the heart $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)$, we see that both E^+ and E^- are concentrated in degree 0 part by Claim 6.3. Hence the triangle (6.3) gives a short exact sequence in $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)$.

Then the complex number $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-)$ is in the ray $\mathbb{R}_{>0}$. If E^- is (σ_1,σ_2) -torsion with $\Im \mathfrak{m} Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-) = 0$, then E^- satisfies $\tau_2^R E^- = 0$ and $\tau_1^L E^- \in \mathcal{P}_{\sigma_1}(1)$. This contradicts the fact $Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-) \in \mathbb{R}_{>0}$. Hence E^- is not (σ_1,σ_2) -torsion. Thus we see $\Im \mathbb{I} \mathbb{I} Z_1(\tau_1^L E^-) +$ $\mathfrak{Im} Z_2(\tau_2^R E^-) \neq 0$, and the exact sequence (6.3) implies

$$0 < \mu_{\beta,\omega}(E) \le \mu_{\beta,\omega}(E^-)$$

by the $\mu_{\beta,\omega}$ -semistability of E. Hence $\Im m Z_{\beta,\omega}^{\sigma_1,\sigma_2}(E^-)$ has to be positive and this contradicts $Z^{\sigma_1,\sigma_2}_{\beta,\omega}(E^-) \in \mathbb{R}_{>0}$. Thus E^- has to be zero.

We secondly show $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2} \subset \mathcal{Q}(-1,0]$. Let F be in $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and let U be an open neighborhood of (β,ω) . For any rational point $(\beta',\omega') \in U_{\mathbb{Q}}$, we have $\mathcal{F}_{\beta',\omega'}^{\sigma_1,\sigma_2} \subset \mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(-1,0]$ and $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2} \subset \mathcal{P}_{\beta',\omega'}^{\sigma_1,\sigma_2}(0,1]$. Since F is in the heart gl $(\mathcal{A}_1,\mathcal{A}_2)$ which is the extension closure of $\mathcal{F}_{\beta',\omega'}^{\sigma_1,\sigma_2}$ and $\mathcal{T}_{\beta',\omega'}^{\sigma_1,\sigma_2}(-1,1]$.

Taking the limit (β', ω') to (β, ω) , we see $F \in \mathcal{Q}[-1, 1]$. On the other hand any object in $\mathcal{Q}(-1) \subset \mathcal{Q}(-2,-1]$ is in the extension closure of $\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2)[-2]$ and $\mathsf{gl}(\mathcal{A}_1,\mathcal{A}_2)[-1]$ by Claim 6.3. Since F is in gl (A_1, A_2) , F is actually in $\mathcal{Q}(-1, 1]$. Thus we obtain the distinguished triangle

where $F^0 \in \mathcal{Q}(0,1]$ and $\mathcal{F}^1 \in \mathcal{Q}(-1,0]$. Again, by Claim 6.3, we see that both F^0 and F^1 are in gl $(\mathcal{A}_1, \mathcal{A}_2)$ by taking the cohomology of the sequence (6.4). Hence the sequence (6.4) gives a short exact sequence in gl $(\mathcal{A}_1, \mathcal{A}_2)$. Since F^0 is a subobject of F in gl $(\mathcal{A}_1, \mathcal{A}_2)$, F^0 is also in $\mathcal{F}^{\sigma_1, \sigma_2}_{\beta, \omega}$. Hence F^0 satisfies $\mu^+_{\beta, \omega}(F^0) \leq 0$ which implies

(6.5)
$$\Im \mathfrak{m} Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F^0) \le 0.$$

On the other hand, we have

(6.6)
$$\Im \mathfrak{m} \, Z^{\sigma_1, \sigma_2}_{\beta, \omega}(F^0) \ge 0$$

since $F \in \mathcal{Q}(0,1]$. Hence F must satisfy $\Im m Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^0) = 0$ and $\Re c Z_{\beta,\omega}^{\sigma_1,\sigma_2}(F^0) \leq 0$. This contradicts the proof of Proposition 4.2. Thus F^0 has to be zero and hence F is in $\mathcal{Q}(-1,0]$. \Box

Proposition 6.4. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5). Suppose $\tau = (Z_{\beta,\omega}^{\sigma_1,\sigma_2}, \mathcal{Q})$ is a stability condition is a deformation of $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ for a rational point $(\beta_0,\omega_0) \in \mathcal{H}_{\mathbb{Q}}$. Then the heart $\mathcal{Q}(0,1]$ is the tilting heart $\left\langle \mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}, \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1] \right\rangle$ defined in Definition 3.8.

Proof. Let us fix the pair (β, ω) through the proof. Both $\mathcal{Q}(0, 1]$ and $\mathcal{A}^{\sigma_1, \sigma_2}_{\beta, \omega} = \left\langle \mathcal{T}^{\sigma_1, \sigma_2}_{\beta, \omega}, \mathcal{F}^{\sigma_1, \sigma_2}_{\beta, \omega}[1] \right\rangle$ are hearts of bounded t-structures on **D**, it is enough to show

(6.7)
$$\left\langle \mathcal{T}^{\sigma_1,\sigma_2}_{\beta,\omega}, \mathcal{F}^{\sigma_1,\sigma_2}_{\beta,\omega}[1] \right\rangle \subset \mathcal{Q}(0,1].$$

By Proposition 6.2, we see both $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1]$ and $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ are contained in $\mathcal{Q}(0,1]$. Since $\mathcal{Q}(0,1]$ is a extension closure, (6.7) holds.

Theorem 6.5. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \operatorname{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor 5).

Then the map

(6.8)
$$s: \mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2) \to \operatorname{Stab} \mathbf{D}; (\beta, \omega) \mapsto \Sigma^{\sigma_1, \sigma_2}_{\beta, \omega}$$

is continuous, where $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ is the pair defined in Definition 4.1.

Proof. Recall that the map is continuous on rational points in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ by Proposition 5.19. By Corollary 5.16 and [Bri07, Theorem 1.2], there exists a stability condition τ given by a deformation of $\Sigma_{\beta_0,\omega_0}^{\sigma_1,\sigma_2}$ and whose central charge is $Z_{\beta,\omega}^{\sigma_1,\sigma_2}$ where $(\beta_0,\omega_0) \in (H^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2))_{\mathbb{Q}}$. Moreover the heart of τ is just $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ by Proposition 6.4. Hence τ is the pair $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ and the map s is continuous. \square

7. Support Property

We have obtained the continuous family $\mathcal{S}(\epsilon_1, \epsilon_2) = \{\Sigma_{\beta,\omega}^{\sigma_1, \sigma_2} \mid (\beta, \omega) \in \mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)\}$ of stability conditions on $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$. Unfortunately it was not possible to prove the support property for $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ directly. In this section, we show that any $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ satisfies the support property via "specialization". More precisely one can find a stability conditions $\Sigma_{1,0}^{\sigma_1,\sigma_2}$ which satisfies the support property in the boundary of the family.

The following proposition is a generalization of our previous work [Kaw19, Proposition 4.8].

Proposition 7.1. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

- (1) If $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ is $\mathsf{gl}(\sigma_1, \sigma_2)$ -semistable then $\tau_1^L(E)$ is σ_1 -semistable, and $\tau_2^R E$ is σ_2 semistable.
- (2) The equality $\arg Z_1(\tau_1^L E) = \arg Z_1(\Phi(\tau_2^R E))$ holds.

Proof. Any $E \in \mathsf{gl}(\mathcal{A}_1, \mathcal{A}_2)$ has the canonical decomposition:

$$0 \longrightarrow i_2 \tau_2^R E \longrightarrow E \longrightarrow i_1 \tau_1^L E \longrightarrow 0.$$

Suppose that E is gl (σ_1, σ_2) -semistable. Then, by the condition (Mor5), we obtain the following:

(7.1)
$$\arg Z_1(\Phi(\tau_2^R E)) = \arg Z_2(\tau_2^R E) \le \arg \operatorname{gl}(Z_1, Z_2)(E) \le Z_1(\tau_1^L E).$$

Suppose to the contrary that $\tau_1^L E$ is not σ_1 -semistable. Then there exists a subobject A of $\tau_1^L E$ which is σ_1 -semistable with

(7.2)
$$\arg Z_1(\tau_1^L E) < \arg Z_1(A).$$

By Lemma 2.25, there exists a subobject F of E such that $\tau_1^L F \cong A$ and $\tau_2^R F \cong \operatorname{im} g[-1]$ where g is the composite $A \to \tau_1^L E \to \Phi(\tau_2^R E)$. Since E is $\operatorname{\mathsf{gl}}(\sigma_1, \sigma_2)$ -semistable, we have

(7.3)
$$\arg \operatorname{\mathsf{gl}}(Z_1, Z_2)(F) \le \arg \operatorname{\mathsf{gl}}(Z_1, Z_2)(E).$$

The σ_1 -semistability of A and the condition (Mor2) imply

$$\arg Z_1(A) \le Z_1(\operatorname{im} g) = \arg Z_2(\operatorname{im} g[-1]).$$

By the construction of F, we have

(7.4)
$$\arg Z_1(A) \le \arg \operatorname{gl}(Z_1, Z_2)(F) \le \arg Z_2(\operatorname{im} g[-1]).$$

Then (7.1), (7.2) and (7.4) imply the following inequalities

(7.5)
$$\arg gl(Z_1, Z_2)(E) \le \arg Z_1(\tau_1^L E) < \arg Z_1(A) \le \arg gl(Z_1, Z_2)(F)$$


which contradict (7.3). Hence $\tau_1^L E$ is σ_1 -semistable. Now we wish to show that $\tau_2^R E$ is σ_2 -semistable. Suppose to the contrary that $\tau_2^R E$ is not σ_2 -semistable. Then there exists a quotient B of $\tau_2^R E$ in \mathcal{A}_2 which is σ_2 -semistable with

(7.6)
$$\arg Z_2(B) < \arg Z_2(\tau_2^R E).$$

Then (7.1) implies

$$\arg Z_2(B) = \arg Z_1(\Phi B) < \arg Z_1(\tau_1^L E).$$

Since $\tau_1^L E$ is σ_1 -semistable, we have $\operatorname{Hom}_{\mathbf{D}}(i_1\tau_1^L E[-1], i_2B) = \operatorname{Hom}_{\mathbf{D}_1}(\tau_1^L E, \Phi(B)) = 0$. Thus the morphism $q: i_2 \tau_2^R E \to i_2 B$ lifts to E, that is, we have the following commutative diagram in \mathbf{D} :



The morphism \bar{q} is an epi morphism in gl $(\mathcal{A}_1, \mathcal{A}_2)$ since $\tau_1^L q$ and $\tau_2^R q$ are respectively epi morphisms in \mathcal{A}_1 and in \mathcal{A}_2 . Since E is $\mathsf{gl}(\sigma_1, \sigma_2)$ -semistable, we have the inequalities

(7.7)
$$\arg gl(Z_1, Z_2)(E) \le \arg gl(Z_1, Z_2)(i_2B) = \arg Z_2(B)$$

which contradicts (7.1) and (7.6). Hence $\tau_2^R E$ is σ_2 -semistable.

For the proof of (2), suppose to contrary
$$\arg Z_2(\tau_2^R E) \neq \arg Z_1(\tau_1^L E)$$
. Then (7.1) implies

(7.8)
$$\arg Z_1(\Phi(\tau_2^R E)) = \arg Z_2(\tau_2^R E) < \arg Z_1(\tau_1^L E).$$

Since $\Phi(\tau_2^R E)$ is σ_1 -semistable, Then we have

$$0 = \operatorname{Hom}(\tau_1^L E, \Phi(\tau_2^R E)) \cong \operatorname{Hom}(i_1 \tau_1^L E, i_2 \tau_2^R E[1])$$

which implies $E \cong i_1 \tau_1^L E \oplus i_2 \tau_2^R E$. Since E is $\mathsf{gl}(\sigma_1, \sigma_2)$ -semistable, either $\tau_1^L E$ or $\tau_2^R E$ has to be zero.

Corollary 7.2. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \, \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

Then the supremum

(7.9)
$$\sup\left\{\frac{|Z_2(\tau_2^R E)|}{|\mathsf{gl}(Z_1, Z_2)(E)|}\middle|E \text{ is } \mathsf{gl}(\sigma_1, \sigma_2)\text{-semistable}\right\}$$

is smaller than or equal to 1.

Proof. Suppose that E is gl (σ_1, σ_2) -semistable. By Proposition 7.1, we have

$$gl(Z_1, Z_2)(E)| = |Z_1(\tau_1^L E)| + |Z_2(\tau_2^R E)|$$

which directly implies the desired assertion.

Proposition 7.3. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor5).

If $(\beta, \omega) \in \mathbb{R} \times \mathbb{R}_{>0}$ satisfies $|\beta - 1 + \sqrt{-1}\omega| < \sin(\pi\epsilon)$ for a sufficiently small $\epsilon > 0$, then the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ is in $B_{\epsilon}(\mathsf{gl}(\sigma_1,\sigma_2))$

Proof. By Corollary 7.2, we have

$$||Z_{\beta,\omega}^{\sigma_1,\sigma_2} - \mathsf{gl}(Z_1,Z_2)||_{\mathsf{gl}(\sigma_1,\sigma_2)} = ||(\beta - 1 + \sqrt{-1}\omega)Z \circ \tau_2^R||_{\mathsf{gl}(\sigma_1,\sigma_2)} < \sin(\pi\epsilon).$$

Let \mathcal{Q} be the slicing of the stability condition $\mathsf{gl}(\sigma_1, \sigma_2)$. To complete the proof, it is enough to

show the tilting heart $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is contained in $\mathcal{Q}(-1+\epsilon,2-\epsilon]$ by [CP10, Proposition 4.2]. Recall $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is the extension closure of $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ and $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1]$. The subcategory $\mathcal{T}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is clearly contained in gl $(\mathcal{A}_1,\mathcal{A}_2) = \mathcal{Q}(0,1]$. The inclusion $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2} \subset$ gl $(\mathcal{A}_1,\mathcal{A}_2)$ implies $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1] \subset \mathcal{Q}(1,2)$. $\mathcal{Q}(1,2]$. Hence we wish to show $\mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}[1] \subset \mathcal{Q}(1,2-\epsilon]$.

Now take $E \in \mathcal{F}_{\beta,\omega}^{\sigma_1,\sigma_2}$. let F be the maximal destabilizing quotient of E with respect to the stability condition $\mathfrak{gl}(\sigma_1, \sigma_2)$. Since $\mathcal{F}_{\beta, \omega}^{\sigma_1, \sigma_2}$ is the free part of $\mathfrak{gl}(\mathcal{A}_1, \mathcal{A}_2)$, the subobject F of Ein gl $(\mathcal{A}_1, \mathcal{A}_2)$ is also in $\mathcal{F}_{\beta, \omega}^{\sigma_1, \sigma_2}$. Lemma 5.2 implies the inequalities $0 < \arg Z_1(\tau_1^L F) \le \arg(\beta + 1 + \sqrt{-1}\omega)$ and $0 < \arg Z_2(\tau_2^R F) \le \arg(\beta + \sqrt{-1}\omega)$. Since gl $(Z_1, Z_2)(F) = Z_1(\tau_1^L F) + Z_2(\tau_2^R F)$, we see

(7.10)
$$\mathcal{F}^{\sigma_1,\sigma_2}_{\beta,\omega}[1] \subset \mathcal{Q}(1,1+\theta],$$

where $\theta = \arg(\beta + \sqrt{-1\omega})/\pi$. By the assumption $|\beta - 1 + \sqrt{-1\omega}| < \sin(\pi\epsilon)$, if ϵ is sufficiently small, then θ is smaller than 1/2. Thus we may assume the inequality $1 + \epsilon < 2 - \epsilon$ holds for a sufficiently small ϵ . Hence we have

$$\mathcal{F}^{\sigma_1,\sigma_2}_{\beta,\omega}[1] \subset \mathcal{Q}(1,1+\epsilon] \subset \mathcal{Q}(1,2-\epsilon].$$

Thus, if ϵ is sufficiently small, then the tilting heart $\mathcal{A}_{\beta,\omega}^{\sigma_1,\sigma_2}$ is contained in $\mathcal{Q}(0,2-\epsilon]$.

Remark 7.4. If (β, ω) is sufficiently close, the stability condition $\Sigma_{\beta,\omega}^{\sigma_1,\sigma_2}$ is close to $\mathsf{gl}(\sigma_1,\sigma_2)$. Hence $\mathsf{gl}(\sigma_1, \sigma_2)$ can be regarded as $\Sigma_{1,0}^{\sigma_1, \sigma_2}$.

Corollary 7.5. Let $\mathbf{D} = \langle \mathbf{D}_1, \mathbf{D}_2 \rangle$ be a semiorthogonal decomposition with (Mor2). Choose rational stability conditions $\sigma_i = (\mathcal{A}_i, Z_i) \in \mathsf{Stab} \mathbf{D}_i$ such that (σ_1, σ_2) satisfies the condition (Mor 5).

If σ_i satisfy the support property, then any stability condition in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ satisfies the support property.

Proof. Recall that the support property is open and closed by Remark 2.8. By Proposition 7.3, if (β, ω) is sufficiently closed to (1, 0), then $\Sigma_{\beta, \omega}^{\sigma_1, \sigma_2}$ satisfies the support property. Since the set $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ is connected, any $\Sigma_{\beta, \omega}^{\sigma_1, \sigma_2}$ for $(\beta, \omega) \in \mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$ satisfies the support property.

8. Construction of a path

The aim of this section is a construction of a path in the space of stability conditions on the category of morphisms. Before the construction, let us introduce notation. Let $(\mathbf{D}^{\leq 0}, \mathbf{D}^{\geq 0})$ be a bounded t-structure of a triangulated category **D**. To simplify notation, we denote by $\mathcal{A}_{[-q,-p]}$ the subcategory $\mathbf{D}^{\leq q} \cap \mathbf{D}^{\geq p}$ if $p \leq q$. The subcategory $\mathcal{A}_{[p,p]}$ is nothing but the shift $\mathcal{A}[p]$ of \mathcal{A} . Moreover the subcategory $\mathcal{A}_{[p,p+1]}$ is the extension closure of $\mathcal{A}[p]$ and $\mathcal{A}[p+1]$.

Now let us specialize **D** as the category of morphisms. Let \mathcal{C}^{Δ^1} be the infinity category of functors from Δ^1 to a stable infinity category \mathcal{C} . By [Lur17], \mathcal{C}^{Δ^1} is stable and hence the homotopy category $h(\mathcal{C}^{\Delta^1})$ is triangulated. We refer to $h(\mathcal{C}^{\Delta^1})$ as the category of morphisms in h(C).

Recall three functors between \mathfrak{C} and \mathfrak{C}^{Δ^1} introduced in (1.1). Then the adjoint pairs $d_0 \dashv s$ and $s \dashv d_1$ determine semiorthogonal decompositions on $h(\mathcal{C}^{\Delta^1}) = \langle \mathbf{D}_1^i, \mathbf{D}_2^i \rangle$ $(i \in \{0, 1\})$ respectively. Both subcategories \mathbf{D}_1^0 and \mathbf{D}_2^1 are given by full sub category of $h(\mathcal{C}^{\Delta^1})$ consisting of identity morphisms in h(\mathcal{C}). Moreover \mathbf{D}_1^2 is the full subcategory of morphisms to zero objects, and \mathbf{D}_1^1 is the full subcategory of morphisms from zero objects:

$$\mathbf{D}_1^0 = \mathbf{D}_2^1 = \{ [\mathrm{id} \colon x \to x] \mid x \in \mathrm{h}(\mathcal{C}) \},\$$

(8.1)
$$\mathbf{D}_{2}^{0} = \{[y \to 0] \mid y \in h(\mathcal{C})\} =: h(\mathcal{C}_{/0}), \text{ and}$$

(8.2)
$$\mathbf{D}_1^1 = \{ [0 \to z] \mid z \in \mathbf{h}(\mathcal{C}) \} =: \mathbf{h}(\mathcal{C}_{0/}).$$

These subcategories are canonically equivalent to $h(\mathcal{C})$. Under the equivalence, the inclusion functor $\mathbf{D}_2^0 \to \mathbf{h}(\mathcal{C}^{\Delta^1})$ (resp. $\mathbf{D}_1^1 \to \mathbf{h}(\mathcal{C}^{\Delta^1})$) is denoted by $j_!$ (resp. j_*). For a stability condition $\sigma = (\mathcal{A}, Z) \in \mathsf{Stab} \, \mathbf{D}_1^0$, set $(\sigma_1^0, \sigma_2^0) \in \mathsf{Stab} \, \mathbf{D}_1^0 \times \mathsf{Stab} \, \mathbf{D}_2^0$ by $(\sigma, \sigma[-1])$

and set $(\sigma_1^1, \sigma_2^1) \in \mathsf{Stab} \, \mathbf{D}_1^1 \times \mathsf{Stab} \, \mathbf{D}_2^1$ by $(\sigma[1], \sigma)$.

The gluing hearts derived from the semiorthogonal decompositions $h(\mathcal{C}^{\Delta^1}) = \langle \mathbf{D}_1^i, \mathbf{D}_2^i \rangle$ are denoted by $d_i^* \mathcal{A}$ respectively. Then we obtain the torsion pairs $(d_i^* \mathcal{T}_{\beta,\omega}, d_i^* \mathcal{F}_{\beta,\omega})$ on $d_i^* \mathcal{A}$ via Definition 3.8 and the stability conditions $\Sigma_{\beta,\omega}^{\sigma_1^i,\sigma_2^i}$ which will be denoted by $d_i^*\sigma_{\beta,\omega} = (d_i^*\mathcal{A}_{\beta,\omega}, d_i^*Z_{\beta,\omega}).$ By the construction of the continuous map d_i^* , the stability condition $d_i^*\sigma$ is nothing but $\sum_{1,0}^{\sigma_1^i,\sigma_2^i}$ (see also [Kaw19]).

Lemma 8.1. Let \mathcal{C} be a stable infinity category and let $\sigma = (\mathcal{A}, Z)$ be a discrete stability condition on h(\mathcal{C}). Take $f \in d_0^* \mathcal{T}_{\beta,\omega}$ and $g \in d_1^* \mathcal{F}_{\beta,\omega}$ arbitrary. For any integer $p \leq -1$, the vanishing $\operatorname{Hom}_{h(\mathcal{C}^{\Delta^1})}(f, g[p]) = 0$ holds.

Proof. Consider the semiorthogonal decomposition $h(\mathbb{C}^{\Delta^1}) = \langle \mathbf{D}_1^1, \mathbf{D}_2^1 \rangle$ associated with $s \dashv d_1$. For $[g: z \to w] \in d_1^* \mathcal{F}_{\beta,\omega}$, the gluing morphism is nothing but the canonical morphism $v: \operatorname{cof} g \to z[1]$. Since $g \in d_1^* \mathcal{F}_{\beta,\omega}$, Lemma 3.9 implies that v is a monomorphism in the abelian category $\mathcal{A}[1]$. Hence we see that both $d_1g = z$ and $d_0g = w$ are in \mathcal{A} and that the morphism $g: z \to w$ is an epi morphism in \mathcal{A} .

Moreover $f \in d_0^* \mathcal{T}_{\beta,\omega}$ satisfies $d_1 f \in \mathcal{A}_{[-1,0]}, d_0 f \in \mathcal{A}$ and $\operatorname{cof} f \in \mathcal{A}$. Hence we easily see $\operatorname{Hom}(d_1 f, d_1 g[p]) = \operatorname{Hom}(d_0 f, d_0 g[p]) = \operatorname{Hom}(d_1 f, d_0 g[p-1]) = 0$ for $p \leq -2$. Then Corollary 2.22 implies $\operatorname{Hom}(f, g[p]) = 0$ for $p \leq -2$.

To discuss the case p = -1, suppose that f is (σ_1, σ_2) -torsion. Then $d_0 f$ and $\operatorname{cof} f$ are both σ -torsion. Moreover the first cohomology $H^1(x)$ of $x = d_1 f$ is also σ -torsion since $H^1(x)$ is a quotient of $\operatorname{cof} f$ in \mathcal{A} . Since $d_1 g$ is σ -free, we see $\operatorname{Hom}(H^1(x)[-1], d_1 g[-1]) = 0$. The trivial vanishing $\operatorname{Hom}(H^0(x), d_1 g[-1]) = 0$ implies $\operatorname{Hom}(d_1 f, d_1 g[-1]) = 0$. Moreover the vanishings $\operatorname{Hom}(d_0 f, d_0 g[-1]) = \operatorname{Hom}(d_1 f, d_0 g[-2]) = 0$ hold by the cohomological degree reason. Again Corollary 2.22 implies $\operatorname{Hom}(f, g[-1]) = 0$.

Now let us suppose f is not (σ_1, σ_2) -torsion. Let u be the gluing morphism $u: d_0 f \to \operatorname{cof} f$ of f. Since $H^0(x)$ is the kernel ker u of u in \mathcal{A} , we obtain the following distinguished triangle $\operatorname{id}_{\ker u} \to f \to \overline{f}$ in $\operatorname{h}(\mathbb{C}^{\Delta^1})$ denoted by

$$\begin{array}{ccc} (8.3) & & & & & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & &$$

By the adjunction $s \dashv d_1$, we see

$$\operatorname{Hom}_{\mathbf{h}(\mathcal{C}\Delta^{1})}(\operatorname{id}_{\ker u}, g[-1]) \cong \operatorname{Hom}_{\mathbf{h}(\mathcal{C})}(\ker u, d_{1}g[-1]) = 0$$

by degree reason. Hence we may assume that $u: d_0 f \to \operatorname{cof} f$ is a mono morphism in \mathcal{A} without loss of generality. Then x is quasi-isomorphic to $(\operatorname{cof} f/d_0 f)[-1]$.


Take a morphism $\tau \in \text{Hom}_{h(\mathbb{C}^{\Delta^1})}(f, g[-1])$ arbitrary. We wish to show $\tau = 0$. Since $d_0g[-1]$ is in $\mathcal{A}[-1]$, it is easily see $\text{Hom}(d_0f, d_0g[-1]) = \text{Hom}(d_1f, d_0g[-2]) = 0$. Thus it is enough to show $d_1\tau = 0$.

Put the morphism $d_1\tau[1]: x[1] = \operatorname{cof} f/y \to z$ in \mathcal{A} by τ_1 . We obtain the following commutative diagram in h(\mathcal{C}):

(8.4)
$$x \xrightarrow{\tau_1[-1]} z[-1]$$

$$f \bigvee_{\substack{f \\ y \xrightarrow{d_0 \tau}}} w[-1].$$

The commutativity implies the composite $g \circ \tau_1$ is zero. Hence the morphism τ_1 factors through the kernel ker g, which means im $\tau_1 \subset \ker g$. Thus we obtain the following commutative diagram in $h(\mathcal{C})$:



By [Kaw19, Lemma 2.11], the above diagram gives a sequence $f \to j_!(\operatorname{im} \tau_1[-1]) \to g[-1]$. We claim that the morphism $\rho: f \to j_!(\operatorname{im} \tau_1[-1])$ is an epimorphism in $d_0^*\mathcal{A}$ and the morphism $\rho': j_!(\operatorname{im} \tau_1) \to g$ is a monomorphism in $d_1^*\mathcal{A}$.

Let h be the fiber of ρ in h(\mathcal{C}^{Δ^1}). Then clearly $d_0 h = y$, and cof h is the kernel of the morphism cof $f \to \operatorname{cof} f/y \to \operatorname{im} \delta$ in \mathcal{A} . Hence h is in $d_0^* \mathcal{A}$ which implies that ρ is an epimorphism in $d_0^* \mathcal{A}$.

To discuss ρ' , note that $j_!(\operatorname{im} \tau_1)$ is a subobject $j_!(\operatorname{ker} g)$ in $d_1^*\mathcal{A}$ by $\operatorname{im} \tau_1 \subset \operatorname{ker} g$. Since g is epi, there exists a morphism $\iota: j_!(\operatorname{ker} g) \to g$ in $\operatorname{h}(\mathbb{C}^{\Delta^1})$. Then one can easily see that the morphism ι is a monomorphism in $d_1^*\mathcal{A}$. Hence $j_!(\operatorname{im} \tau_1)$ is a subobject of g in $d_1^*\mathcal{A}$.

Since $j_!(\operatorname{im} \tau_1[-1])$ is a quotient of f in $d_0^* \mathcal{A}$, $j_!(\operatorname{im} \tau_1[-1])$ is in $d_0^* \mathcal{T}_{\beta,\omega}$. Hence $j_!(\operatorname{im} \tau_1[-1])$ satisfies $\mathfrak{Im} d_0^* Z_{\beta,\omega}(j_!(\operatorname{im} \tau_1[-1]))$ is positive which implies

(8.6)
$$\beta \operatorname{\mathfrak{Im}} Z(\operatorname{im} \tau_1) - \omega \operatorname{\mathfrak{Re}} Z(\operatorname{im} \tau_1) > 0.$$

On the other hand, $j_!(\operatorname{im} \tau_1)$ is in $d_1^* \mathcal{F}_{\beta,\omega}$ since it is a subobject of $g \in d_1^* \mathcal{F}_{\beta,\omega}$. Thus the non-positivity $d_1^* Z_{\beta,\omega}(j_!(\operatorname{im} \tau_1)) \leq 0$ implies

(8.7)
$$(\beta+1)\operatorname{\mathfrak{Im}} Z(\operatorname{\mathsf{im}} \tau_1) - \omega \operatorname{\mathfrak{Re}} Z(\operatorname{\mathsf{im}} \tau_1) \le 0.$$


Then (8.7) implies $\beta \Im \mathfrak{Im} Z(\operatorname{im} \tau_1) - \omega \Re \mathfrak{e} Z(\operatorname{im} \tau_1) \leq 0$ since $\operatorname{Im} Z(\operatorname{im} \tau_1)$ is non-negative. This clearly contradicts (8.6). Hence $\operatorname{im} \tau_1$ is zero which implies $d_1 \tau = 0$.

Lemma 8.2. Let \mathcal{C} be a stable infinity category and let $\sigma = (\mathcal{A}, Z)$ be a discrete stability condition on $h(\mathcal{C})$. Take $f \in d_0^* \mathcal{F}_{\beta,\omega}$ and $g \in d_1^* \mathcal{T}_{\beta,\omega}$ arbitrary. For any integer $p \leq 0$, the vanishing $\operatorname{Hom}_{h(\mathcal{C}\Delta^1)}(g, f[p]) = 0$ holds.

Proof. Since f is in $d_0^* \mathcal{F}_{\beta,\omega}$, the gluing morphism $d_0 f \to \operatorname{cof} f$ is mono by Lemma 2.21. Hence $d_1 f$ is in $\mathcal{A}[-1]$. Since g is in $d_1^* \mathcal{T}_{\beta,\omega}$, in particular in $d_1^* \mathcal{A}$, $d_1 g$ is in \mathcal{A} and $d_0 g$ is in $\mathcal{A}_{[0,1]}$. Hence Corollary 2.22 implies the vanishing $\operatorname{Hom}(g, f[p]) = 0$ for $p \leq -1$.

To discuss the case p = 0, suppose that g is (σ_1, σ_2) -torsion. Then d_1g and fib g are σ -torsion objects in \mathcal{A} . Moreover the 0-th cohomology $H^0(d_0g)$ of d_0g is σ -torsion since it is a quotient of the σ -torsion object d_1g . Then we see $\operatorname{Hom}(d_0g, d_0f) = 0$ by the σ -freeness of d_0f . In addition the vanishings $\operatorname{Hom}(d_1g, d_1f) = \operatorname{Hom}(d_1g, d_0f[-1]) = 0$ hold by the cohomological degree reason. Thus Corollary 2.22 implies $\operatorname{Hom}(g, f) = 0$.

Now suppose that g is not (σ_1, σ_2) -torsion. Let u be the universal morphism fib $g \to d_1 g$. Then there exists a diagram of the distinguished triangle in h(\mathcal{C}):



Put the right horizontal arrow by $[\bar{g}: z \to \operatorname{cok} u]$ which gives an object in $h(\mathcal{C}^{\Delta^1})$. By [Kaw19, Lemma 2.11], the above digram gives a distinguished triangle in $h(\mathcal{C}^{\Delta^1})$, in particular, a short exact sequence in $d_1^*\mathcal{A}$. Since the vanishings $\operatorname{Hom}(j_*(\ker u[1]), f) \cong \operatorname{Hom}(\ker u[1], d_0 f) = 0$ hold by the fact $d_0 f \in \mathcal{A}$, it is enough to show $\operatorname{Hom}(\bar{g}, f) = 0$.

Take $\bar{\varphi} \in \text{Hom}(\bar{g}, f)$. Since $\text{Hom}(d_1\bar{g}, d_1f) = \text{Hom}(d_1\bar{g}, d_0f[-1]) = 0$ hold by the cohomological degree reason, it is enough to show $d_0\bar{\varphi} = 0$. Since $(d_0\bar{\varphi})\circ\bar{g} = f\circ(d_1\bar{\varphi}) = 0$, \bar{g} factors through the kernel ker $d_0\bar{\varphi}$. Since \bar{g} is an epimorphism in \mathcal{A} , the canonical morphism ker $d_0\bar{\varphi} \to \operatorname{cok} u$ has to be an epimorphism. Thus $d_0\bar{\varphi}$ is zero morphism.

Proposition 8.3. Let C be a stable infinity category and let $\sigma = (A, Z)$ be a locally finite stability condition on h(C).

(1) $d_0^* \mathcal{T}_{\beta,\omega} \subset (d_1^* \mathcal{A}_{\beta,\omega})_{[-1,0]}.$ (2) $d_0^* \mathcal{F}_{\beta,\omega} \subset (d_1^* \mathcal{A}_{\beta,\omega})_{[-2,-1]}.$

 $(z) \quad a | b \quad p, \omega \subset (a | b \quad p, \omega) [-2, -1].$

Proof. Any object $g \in d_1^* \mathcal{A}_{\beta,\omega}$ has a decomposition

 $H^{-1}(g)[1] \longrightarrow g \longrightarrow H^0(g)$

with respect to the heart $d_1^* \mathcal{A}$ where $H^{-1}(g) \in d_1^* \mathcal{F}_{\beta,\omega}$ and $H^0(g) \in d_1^* \mathcal{T}_{\beta,\omega}$. Clearly $d_1(H^0(g))$ is in \mathcal{A} , and $d_0(H^0(g))$ is in $\mathcal{A}_{[0,1]}$. By Lemma 3.9, both $d_0(H^{-1}(g))$ and $d_1(H^{-1}(g))$ are in \mathcal{A} . Hence both d_0g and d_1g are in $\mathcal{A}_{[0,1]}$.

Take $f \in d_0^* \mathcal{T}_{\beta,\omega}$ and $g \in d_1^* \mathcal{A}_{\beta,\omega}$. To prove the assertion (1), it is enough to show

- (1a) $\operatorname{Hom}(g[p], f) = 0$ for $p \ge 1$ and
- (1b) Hom(f, g[q]) = 0 for $q \le -2$.

Since $d_1g[p]$ and $d_0g[p]$ are in $\mathcal{A}_{[p,p+1]}$, Hom $(d_1g[p], d_1f)$, Hom $(d_0g[p], d_0f)$, and Hom $(d_1g[p], d_0f[-1])$ vanish. Then Corollary 2.22 implies (1a).

Now suppose $g \in d_1^* \mathcal{T}_{\beta,\omega}$. Then $d_1 g[q] \in \mathcal{A}[q]$ and $d_0 g \in \mathcal{A}_{[q,q+1]}$. The assumption $q \leq -2$ implies the vanishings

$$Hom(d_1f, d_1g[q]) = Hom(d_0f, d_0g[q]) = Hom(d_1f, d_0g[q-1]) = 0.$$

Corollary 2.22 implies $\operatorname{Hom}(f, g[q]) = 0$ for $g \in d_1^* \mathcal{T}_{\beta,\omega}$. Now Lemma 8.1 implies $\operatorname{Hom}(f, g[q]) = 0$ for $g \in d_1^* \mathcal{T}_{\beta,\omega}[1]$. Thus we have proven (1b).

Take $f \in d_0^* \mathcal{F}_{\beta,\omega}$ and $g \in d_1^* \mathcal{A}_{\beta,\omega}$. To prove the assertion (2), it is enough to show

(2a) $\operatorname{Hom}(g[p], f) = 0$ for $p \ge 0$ and

(2b) Hom(f, g[q]) = 0 for $q \le -3$.

Lemma 8.2 implies $\operatorname{Hom}(g[p], f) = 0$ for $g \in d_1^* \mathcal{T}_{\beta,\omega}$. Suppose $g \in d_1^* \mathcal{F}_{\beta,\omega}[1]$. Then $d_1 g[p]$ and $d_0 g[p]$ are in $\mathcal{A}[p+1]$. Since f is in $d_0^* \mathcal{F}$, $d_1 f$ is in $\mathcal{A}[-1]$ and $d_0 f$ is in \mathcal{A} . Corollary 2.22 implies $\operatorname{Hom}(g[p], f) = 0$ for $g \in d_1^* \mathcal{F}_{\beta,\omega}[1]$. This gives the proof of claim (2a).

To prove (2b), recall g[q] satisfies that both $d_0^*g[q]$ and $d_1g[q]$ are in $\mathcal{A}_{[q,q+1]}$. Since f satisfies $d_0f \in \mathcal{A}$ and $d_1f \in \mathcal{A}[-1]$, Corollary 2.22 implies the claim (2b).

Corollary 8.4. Let \mathcal{C} be a stable infinity category and let $\sigma = (\mathcal{A}, Z)$ be a locally finite stability condition on $h(\mathcal{C})$. Then $d_0^* \mathcal{A}_{\beta,\omega} \subset (d_1^* \mathcal{A}_{\beta,\omega})_{[-1,0]}$.

Proof. Since the heart $d_0^* \mathcal{A}_{\beta,\omega}$ is the extension closure of $d_0^* \mathcal{T}_{\beta,\omega}$ and $d_0^* \mathcal{F}_{\beta,\omega}[1]$, the assertion follows from Proposition 8.3.

Theorem 8.5. Let \mathcal{C} be a stable infinity category and let $\sigma = (\mathcal{A}, Z)$ be a locally finite stability condition on $h(\mathcal{C})$. If σ is rational, then $d_0^*\sigma$ and $d_1^*\sigma$ are path connected in Stab $h(\mathcal{C}^{\Delta^1})$.

Proof. Let p be a path in $\overline{\mathcal{H}}$ given by

$$p: [0,1] \to \overline{\mathcal{H}}; p(t) = (\cos(2\pi t/3), \sin(2\pi t/3))$$

Set ϵ_1 as 1/3 and ϵ_2 as -1/2 in $\mathcal{H}^+(\epsilon_1) \cap \mathcal{H}^-(\epsilon_2)$. Then the domain $\mathcal{H}^+(1/3) \cap \mathcal{H}^-(-1/2)$ contains the image Im p of the path p. Hence the set $\{d_0^*\sigma_{\beta,\omega} \mid (\beta,\omega) \in \text{Im } p\}$ of stability conditions gives a continuous family.

Now, when $(\beta, \omega) \in \text{Im } p$, we denote by $d_i^* \sigma_{p(t)} = (d_i^* Z_{p(t)}, d_i^* \mathcal{A}_{p(t)})$ instead of $d_i^* \sigma_{\beta,\omega}$. By direct calculation, one can check that

$$d_0^* Z_{p(2\pi/3)}(f) = \exp(2\pi\sqrt{-1/3}) d_1^* Z_{p(2\pi/3)}(f)$$

for any $f \in h(\mathbb{C}^{\Delta^1})$. Corollary 8.4 implies that the heart of the stability condition $d_0^*\sigma_{p(2\pi/3)}$ is contained in $(d_1^*\mathcal{A}_{p(2\pi/3)})_{[-1,1]}$. Then [CP10, Proposition 4.1] implies the desired assertion. \Box

Remark 8.6. If σ is full then one can assume that σ is rational. Hence for any full stability σ , $d_0^*\sigma$ is path connected to $d_1^*\sigma$. Theorem 8.5 is a generalization of our previous result [Kaw19, Theorem 1.2]. Hence we obtain the following:

Corollary 8.7. Let C be a stable infinity category and let $\mathsf{Stab}^{\mathsf{fl}} h(C)$ be the space of full stability conditions. Then the images of maps

$$d_0^*, d_1^* \colon \mathsf{Stab}^{\mathsf{fl}} \operatorname{h}(\mathcal{C}) \rightrightarrows \mathsf{Stab}^{\mathsf{fl}} \operatorname{h}(\mathcal{C}^{\Delta^1})$$

are path connected to each other. In other word, the restricted map d_0^* and d_1^* gives the same map from the 0-th homotopy of Stab $h(\mathcal{C})$ to that of Stab $h(\mathcal{C}^{\Delta^1})$

Proof. Recall that the 0-th homotopy $\pi_0(X)$ of a topological space X is nothing but the set of connected components of X. Let [x] be the equivalence class of a point $x \in X$ with respect to path connectedness. By Theorem 8.5, $[d_0^*\sigma]$ is the same as $[d_1^*\sigma]$ for $\sigma \in \mathsf{Stab}^{\mathsf{fl}} h(\mathfrak{C})$. \Box

Acknoledgement. The author would like to thank his family for their great support and thank Atsushi Kanazawa for his encouragement. This work is partially supported by JSPS KAKENHI Grant Number JP21K03212.

References

- [AB13] Daniele Arcara and Aaron Bertram. Bridgeland-stable moduli spaces for K-trivial surfaces. J. Eur. Math. Soc. (JEMS), 15(1):1–38, 2013. With an appendix by Max Lieblich.
- [ABC⁺09] Paul S. Aspinwall, Tom Bridgeland, Alastair Craw, Michael R. Douglas, Mark Gross, Anton Kapustin, Gregory W. Moore, Graeme Segal, Balázs Szendroői, and P. M. H. Wilson. Dirichlet branes and mirror symmetry, volume 4 of Clay Mathematics Monographs. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2009.
- [BB17] Arend Bayer and Tom Bridgeland. Derived automorphism groups of K3 surfaces of Picard rank 1. Duke Math. J., 166(1):75–124, 2017.
- [BM11] Arend Bayer and Emanuele Macrì. The space of stability conditions on the local projective plane. Duke Math. J., 160(2):263–322, 2011.
- [BMS16] Arend Bayer, Emanuele Macri, and Paolo Stellari. The space of stability conditions on abelian threefolds, and on some Calabi-Yau threefolds. *Invent. Math.*, 206(3):869–933, 2016.
- [BMT14] Arend Bayer, Emanuele Macrì, and Yukinobu Toda. Bridgeland stability conditions on threefolds I: Bogomolov-Gieseker type inequalities. J. Algebraic Geom., 23(1):117–163, 2014.
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. Ann. of Math. (2), 166(2):317–345, 2007.
- [Bri08] Tom Bridgeland. Stability conditions on K3 surfaces. Duke Math. J., 141(2):241–291, 2008.
- [CP10] John Collins and Alexander Polishchuk. Gluing stability conditions. Adv. Theor. Math. Phys., 14(2):563–607, 2010.
- [HL97] Daniel Huybrechts and Manfred Lehn. The geometry of moduli spaces of sheaves. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [IUU10] Akira Ishii, Kazushi Ueda, and Hokuto Uehara. Stability conditions on A_n -singularities. J. Differential Geom., 84(1):87–126, 2010.
- [Kaw19] Kotaro Kawatani. Stability conditions on morphisms in a category. *arXiv e-prints*, pages arXiv:1905.05470, to appear in Kyoto Journal of Math., May 2019.
- [Kaw20] Kotaro Kawatani. Stability conditions on affine noetherian schemes, 2020.
- [KS08] Maxim Kontsevich and Yan Soibelman. Stability structures, motivic donaldson-thomas invariants and cluster transformations, 2008.
- [Lur17] Jacob Lurie. Higher Algebra. available on https://www.math.ias.edu/~lurie/, 2017.
- [Mac07] Emanuele Macrì. Stability conditions on curves. Math. Res. Lett., 14(4):657–672, 2007.

 $Email \ address: \tt kawatanikotaro@gmail.com$

Osaka Prefecture University, Gakuen-cho, Sakai-city, Osaka, Japan. Osaka /Yamato University, Katayama-cho, Suita-city, Osaka, Japan. Osaka