

CURVATURE ESTIMATES FOR p -CONVEX HYPERSURFACES OF PRESCRIBED CURVATURE

WEISONG DONG

ABSTRACT. In this paper, we establish the curvature estimates for p -convex hypersurfaces in \mathbb{R}^{n+1} of prescribed curvature with $p \geq \frac{n}{2}$. The existence of a star-shaped hypersurface of prescribed curvature is obtained. We also prove a type of interior C^2 estimates for solutions to the Dirichlet problem of the corresponding equation.

Mathematical Subject Classification (2010): 53C42; 53C21; 35J60.

Keywords: curvature estimates; p -convexity; interior estimates.

1. INTRODUCTION

Let $M \subset \mathbb{R}^{n+1}$ be a closed hypersurface and $\kappa(X) = (\kappa_1, \dots, \kappa_n)$ be the principle curvatures of M at X . Given $1 \leq p \leq n$, a C^2 regular hypersurface M is called p -convex if, at each $X \in M$, $\kappa(X)$ satisfies

$$\kappa_{i_1} + \dots + \kappa_{i_p} \geq 0, \forall 1 \leq i_1 < \dots < i_p \leq n.$$

In other words, the sum of the p smallest principle curvatures is nonnegative at each point in M . The notion of p -convexity goes back to Wu [35] and is studied extensively by Wu [35], Sha [29, 30] and Harvey-Lawson [18, 19].

In this paper, we are interested in finding a p -convex hypersurface $M \subset \mathbb{R}^{n+1}$ of prescribed curvature as below

$$(1.1) \quad \prod_{1 \leq i_1 < \dots < i_p \leq n} (\kappa_{i_1} + \dots + \kappa_{i_p}) = f(X, \nu(X)), \forall X \in M,$$

where $\nu(X)$ is the unit outer normal of M at X , the function $f(X, \nu) \in C^2(\Gamma)$ is positive and Γ is an open neighborhood of unit normal bundle of M in $\mathbb{R}^{n+1} \times \mathbb{S}^n$. The Gaussian curvature equation corresponds to $p = 1$ in (1.1) which was studied by Oliker [23]. The mean curvature equation corresponding to $p = n$ in (1.1) was studied by Bakelman-Kantor [2] and Treibergs-Wei [33]. For general curvature equations, see Caffarelli-Nirenberg-Spruck [4] and Gerhardt [11]. When $p = n - 1$, the equation was studied by Chu-Jiao [7] and, in complex settings, it is related to the Gauduchon conjecture which was solved by Székelyhidi-Tosatti-Weinkove [28]. For some previous work on this topic, see Tosatti-Weinkove [31, 32] and Fu-Wang-Wu [9, 10].

It is of great interest in geometry and PDE to derive a C^2 estimate for equation (1.1) for general $f(X, \nu(X))$. We have the following main result.

Theorem 1.1. *Suppose $M \subset \mathbb{R}^{n+1}$ is a closed star-shaped p -convex hypersurface with $p \geq \frac{n}{2}$ satisfying the curvature equation (1.1). Then, there is a positive constant C such that*

$$(1.2) \quad \sup_{X \in M, i=1, \dots, n} |\kappa_i(X)| \leq C,$$

where C depends on $n, p, |M|_{C^1}, \inf f$ and $|f|_{C^2}$.

We remark that in the theorem only a few conditions are assumed on f . Usually, to derive C^2 estimate for elliptic equations which are not strictly elliptic there should be some extra assumptions on f due to the dependency on $\nu(X)$. We refer the reader to Ivochkina [20, 21], Guan-Guan [13], Guan-Lin-Ma [16], Guan-Li-Li [15] and Guan-Jiao [14] for more details. Moreover, Guan-Ren-Wang in [17] showed that for the following curvature equation

$$\frac{\sigma_k}{\sigma_l}(\kappa(X)) = f(X, \nu(X)), \quad \forall X \in M,$$

where $0 < l < k \leq n$ and σ_k is the k -th elementary symmetric function, estimate (1.2) fails generally, though it may hold for special f as in Guan-Guan [13]. When $l = 0$, some results are known for general f . For instance, estimate (1.2) was proved for $k = n$ by Caffarelli-Nirenberg-Spruck [3] and, for $2 \leq k \leq n$, Guan-Ren-Wang [17] obtained the estimate for convex solutions. For $k = n - 1$ and $n - 2$, estimate (1.2) was established by Ren-Wang [24, 25]. They also conjectured that the estimate still holds for $k > \frac{n}{2}$ in [25, 26]. When f is independent of ν , Caffarelli-Nirenberg-Spruck [4] proved the C^2 estimate for a general class of fully nonlinear curvature equations. For hypersurfaces of prescribed curvature in Riemannian manifolds and Minkowski space, see [5] and [34, 27].

To obtain the existence of a p -convex hypersurface satisfying the prescribed curvature equation (1.1), we assume the following two conditions on f . The first one is that there exists two positive constants $r_1 < 1 < r_2$ such that

$$(1.3) \quad \begin{aligned} f\left(X, \frac{X}{|X|}\right) &\geq \frac{p^{C_n^p}}{r_1^{C_n^p}}, \quad \text{for } |X| = r_1; \\ f\left(X, \frac{X}{|X|}\right) &\leq \frac{p^{C_n^p}}{r_2^{C_n^p}}, \quad \text{for } |X| = r_2. \end{aligned}$$

This condition is used to derive C^0 estimate. The second one is that for any fixed unit vector ν ,

$$(1.4) \quad \frac{\partial}{\partial \rho} \left(\rho^{C_n^p} f(X, \nu) \right) \leq 0, \quad \text{where } \rho = |X|,$$

which will be used to derive C^1 estimate. Actually, with suitable assumptions of f , Li [22] proved that the interior gradient estimate holds.

By the continuity method argument as in [4], we can obtain the following result.

Theorem 1.2. *Let $f \in C^2(\overline{B_{r_2}} \setminus B_{r_1}) \times \mathbb{S}^n$ be a positive function satisfying (1.3) and (1.4). Then equation (1.1) has a unique $C^{3,\alpha}$ star-shaped p -convex solution M in $\{X \in \mathbb{R}^{n+1} : r_1 \leq |X| \leq r_2\}$ for any $\alpha \in (0, 1)$ as long as $p \geq \frac{n}{2}$.*

The method of proof of Theorem 1.1 can be applied to obtain an interior C^2 estimate for the Dirichlet problem of the corresponding equation in Euclidean space. Suppose that Ω is a bounded domain in \mathbb{R}^n . For a function $u \in C^2(\Omega)$, denote by $\lambda(D^2u) = (\lambda_1, \dots, \lambda_n)$ the eigenvalues of the Hessian D^2u . We call that $u \in C^2(\Omega)$ is p -plurisubharmonic if the eigenvalues of D^2u satisfy $\lambda_{i_1} + \dots + \lambda_{i_p} \geq 0, \forall 1 \leq i_1 < \dots < i_p \leq n$ (see [19]). Given a p -plurisubharmonic function v on $\bar{\Omega}$, suppose that there is a p -plurisubharmonic function $u \in C^4(\Omega) \cap C^{1,1}(\bar{\Omega})$ satisfying $u < v$

in Ω solves the equation

$$(1.5) \quad \prod_{1 \leq i_1 < \dots < i_p \leq n} (\lambda_{i_1} + \dots + \lambda_{i_p}) = f(x, u, Du), \text{ in } \Omega$$

with the Dirichlet boundary data

$$u = v, \text{ on } \partial\Omega,$$

where $f \in C^2(\Omega \times \mathbb{R} \times \mathbb{R}^n)$ is a positive function. By the same argument as Theorem 1.1, we can prove the following interior estimate.

Theorem 1.3. *For the Dirichlet problem of (1.5) with $p \geq \frac{n}{2}$, suppose that u and v are as above. Then there exists a constant C and β depending only on $n, p, |u|_{C^1}, \inf f, |f|_{C^2}$ and Ω such that*

$$(1.6) \quad \sup_{\Omega} (v - u)^{\beta} \Delta u \leq C.$$

Remark 1.4. As a byproduct of the proof of this theorem, one can conclude the following global C^2 estimate for equation (1.5)

$$\sup_{\Omega} |D^2 u| \leq C(1 + \sup_{\partial\Omega} |D^2 u|),$$

where C is a constant as in Theorem 1.3.

We shall only give an outline for the proof of Theorem 1.3, as it is almost the same as that of Theorem 1.1. The estimate (1.6) can also be seen in some sense as a generalization of Theorem 0.4 in [8], since there the right hand side function f doesn't depend on Du . Such estimate for $\sigma_k(\lambda)$ has been proved by Chou-Wang [6]. The function f depending on ν or Du creates substantial difficulties to derive C^2 estimate, as the bad term $-Ch_{11}$ or $-Cu_{11}$ appears when one applies the maximum principle to the test function. One way to overcome this is like in [17, 24, 25] to control the bad third order terms, which is very hard even for the σ_k equation. Another way is as in [7] to control $-Ch_{11}$ firstly by good terms (see Lemma 3.2), then the bad third order terms can be eliminated easily. Thank to Dinew [8], where many properties of the operator have been proved, we can follow the argument in [7] to prove the estimate.

The paper is organized as follows. In Section 2, we recall some properties of the operator from [8]. In Section 3, we prove the curvature estimate. In Section 4 we derive the gradient estimate and in Section 5 we apply the continuity method to prove Theorem 1.2. At last, we give an outline of the proof of Theorem 1.3 in Section 6.

Acknowledgements: The author is supported by the National Natural Science Foundation of China, No. 11801405.

2. PRELIMINARIES

Let \mathbb{S}^n be the unit sphere in \mathbb{R}^{n+1} and ∇ be the gradient on it. Assume that M is star-shaped with respect to the origin, i.e. the position vector X of M can be written as $X(x) = \rho(x)x$, where $x \in \mathbb{S}^n$. Then the unit outer normal of M is given by

$$\nu = \frac{\rho x - \nabla \rho}{\sqrt{\rho^2 + |\nabla \rho|^2}}.$$

Let $\{e_1, \dots, e_n\}$ be smooth local orthonormal frame on \mathbb{S}^n . Then the metric of M is given by $g_{ij} = \rho^2 \delta_{ij} + \rho_i \rho_j$, and the second fundamental form of M is

$$(2.1) \quad h_{ij} = \frac{\rho^2 \delta_{ij} + 2\rho_i \rho_j - \rho \rho_{ij}}{\sqrt{\rho^2 + |\nabla \rho|^2}}.$$

At a point X_0 in M , choose a local orthonormal frame $\{e_1, e_2, \dots, e_n\}$. The following geometric formulas are well known:

$$(2.2) \quad \begin{aligned} X_{ij} &= -h_{ij} \nu \quad (\text{Gauss formula}), \\ (\nu)_i &= h_{ij} e_j \quad (\text{Weingarten equation}), \\ h_{ijk} &= h_{ikj} \quad (\text{Codazzi formula}), \\ R_{ijkl} &= h_{ik} h_{jl} - h_{il} h_{jk} \quad (\text{Gauss equation}), \end{aligned}$$

where R_{ijkl} is the $(4, 0)$ -Riemannian curvature tensor and the formula

$$(2.3) \quad h_{ijkl} = h_{klij} + h_{mk}(h_{mj} h_{il} - h_{mi} h_{lj}) + h_{mi}(h_{mj} h_{kl} - h_{ml} h_{kj}).$$

We recall the p -convex cones introduced by Harvey and Lawson [19].

Definition 2.1. Let $p \in \{1, \dots, n\}$. The cone \mathcal{P}_p is defined by

$$\mathcal{P}_p = \{(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n \mid \forall 1 \leq i_1 < i_2 < \dots < i_p \leq n, \lambda_{i_1} + \dots + \lambda_{i_p} > 0\}.$$

Associated to \mathcal{P}_p is the cone of symmetric $n \times n$ matrices defined by

$$P_p = \{A \mid \forall 1 \leq i_1 < i_2 < \dots < i_p \leq n, \lambda_{i_1}(A) + \dots + \lambda_{i_p}(A) > 0\}.$$

We call A is p -positive if $A \in P_p$.

For convenience, we introduce the following notations

$$F(h_{ij}) := F(\kappa) = \prod_{1 \leq i_1 < \dots < i_p \leq n} (\kappa_{i_1} + \dots + \kappa_{i_p}) \quad \text{and} \quad \tilde{F} = F^{\frac{1}{C_n^p}}.$$

Equation (1.1) then can be written as

$$(2.4) \quad \tilde{F}(h_{ij}) := \tilde{F}(\kappa) = \tilde{f}(X, \nu(X)),$$

where $\kappa = (\kappa_1, \dots, \kappa_n)$ and $\tilde{f} = f^{\frac{1}{C_n^p}}$. Denote

$$F^{ij} = \frac{\partial F}{\partial h_{ij}}, \quad F^{ij,kl} = \frac{\partial^2 F}{\partial h_{ij} \partial h_{kl}}, \quad \text{and} \quad \mathcal{F} = \sum F^{ii}.$$

Direct calculations show that

$$\tilde{F}^{ij} = \frac{1}{C_n^p} F^{\frac{1}{C_n^p} - 1} F^{ij}$$

and

$$\tilde{F}^{ij,kl} = \frac{1}{C_n^p} F^{\frac{1}{C_n^p} - 1} F^{ij,kl} + \frac{1}{C_n^p} \left(\frac{1}{C_n^p} - 1 \right) F^{\frac{1}{C_n^p} - 2} F^{ij} F^{kl}.$$

We remark that \tilde{F} is concave with respect to h_{ij} by Lemma 1.13 and Corollary 1.14 in [8]. And the equation is elliptic as the matrix $\left\{ \frac{\partial \tilde{F}}{\partial h_{ij}} \right\}$ is positive definite for $\{h_{ij}\} \in P_p$.

Now we do some basic calculations which will be used in the next section. Our calculations are carried out at a point X_0 on the hypersurface M , and we use coordinates such that at this point $\{h_{ij}\}$ is diagonal and its eigenvalues with respect

to g_{ij} are ordered as $\kappa_1 \geq \kappa_2 \geq \cdots \geq \kappa_n$. Note that F^{ij} is also diagonal at X_0 and we have the following formulas

$$F^{kk} = \frac{\partial F}{\partial \kappa_k} = \sum_{k \in \{i_1, \dots, i_p\}} \frac{F(\kappa)}{\kappa_{i_1} + \cdots + \kappa_{i_p}},$$

which we refer to Lemma 1.10 in [8]. We also have formulas for the second order derivatives of F at X_0 as below

$$F^{kk, ll} = \frac{\partial^2 F}{\partial \kappa_k \partial \kappa_l} = \sum_{\substack{k \in \{i_1, \dots, i_p\} \\ l \in \{j_1, \dots, j_p\} \\ \{i_1, \dots, i_p\} \neq \{j_1, \dots, j_p\}}} \frac{F(\kappa)}{(\kappa_{i_1} + \cdots + \kappa_{i_p})(\kappa_{j_1} + \cdots + \kappa_{j_p})},$$

and, for $k \neq r$,

$$F^{kr, rk} = \frac{F^{kk} - F^{rr}}{\kappa_k - \kappa_r} = - \sum_{\substack{k \notin \{i_1, \dots, i_p\} \ni r \\ r \notin \{j_1, \dots, j_p\} \ni k}} \frac{F(\kappa)}{(\kappa_{i_1} + \cdots + \kappa_{i_p})(\kappa_{j_1} + \cdots + \kappa_{j_p})}.$$

Otherwise, we have $F^{ij, kl} = 0$. See Lemma 1.12 in [8] for the above formulas. These formulas can also be easily seen from Theorem 5.5 in [1]. The following properties of the function F which are very similar the the properties of σ_k are proved in Dinew [8].

Lemma 2.2 ([8]). *Suppose that the diagonal matrix $A = \text{diag}(\lambda_1, \dots, \lambda_n)$ belongs to P_p and $\lambda_1 \geq \cdots \geq \lambda_n$. Then,*

- (1) $\tilde{F}^{11}(A)\lambda_1 \geq \frac{1}{n}\tilde{F}(A)$;
- (2) $\sum_{k=1}^n \tilde{F}^{kk}(A) \geq C_{n-1}^{p-1}$;
- (3) $\sum_{k=1}^n F^{kk}(A)\lambda_k = C_n^p F(A)$;
- (4) *there is a constant $\theta = \theta(n, p)$ such that $F^{jj}(A) \geq \theta \sum F^{ii}$, $\forall j \geq n-p+1$;*
- (5) *given $\delta > 0$, there is an $\epsilon = \epsilon(p, \delta) > 0$ such that, if $\epsilon\lambda_1 \geq \lambda_{n-p+1}$, $-2F^{1i, i1}(A) \geq (1 + \delta)\frac{F^{ii}}{\lambda_1}$ holds for every $i = 2, 3, \dots, n$.*

3. CURVATURE ESTIMATES

Set $u = \langle X, \nu \rangle$, which is the support function of the hypersurface M . Clearly, we have

$$u = \frac{\rho^2}{\sqrt{\rho^2 + |\nabla \rho|^2}}.$$

There exists a positive constant C depending on $\inf_M \rho$ and $|\rho|_{C^1}$ such that

$$\frac{1}{C} \leq \inf_M u \leq u \leq \sup_M u \leq C.$$

In order to prove Theorem 1.1, we consider the following auxiliary function

$$G = \log \kappa_{max} - \log(u - a) + \frac{A}{2}|X|^2$$

where κ_{max} is the largest principle curvature, $a = \frac{1}{2} \inf_M u > 0$ and $A \geq 1$ is a large constant to be determined. Suppose the maximum of G is achieved at a point $X_0 \in M$. Choose a local coordinate frame $\{e_1, \dots, e_n\}$ around X_0 such that

$$h_{ij} = \delta_{ij} h_{ii} \text{ and } h_{11} \geq h_{22} \geq \cdots \geq h_{nn} \text{ at } X_0.$$

Since κ_{max} may not be differentiable, we define a new function \hat{G} near X_0 by

$$\hat{G} = \log h_{11} - \log(u - a) + \frac{A}{2}|X|^2.$$

It is easy to see \hat{G} achieves a maximum at X_0 . Now, differentiating \hat{G} at X_0 twice yields that

$$(3.1) \quad 0 = \frac{h_{11i}}{h_{11}} - \frac{u_i}{u - a} + A\langle X, e_i \rangle$$

and

$$(3.2) \quad 0 \geq \frac{h_{11ii}}{h_{11}} - \left(\frac{h_{11i}}{h_{11}}\right)^2 - \frac{u_{ii}}{u - a} + \left(\frac{u_i}{u - a}\right)^2 + A(1 + \langle X, X_{ii} \rangle).$$

Contracting (3.2) with \tilde{F}^{ii} , we get

$$(3.3) \quad 0 \geq \frac{\tilde{F}^{ii}h_{11ii}}{h_{11}} - \frac{\tilde{F}^{ii}h_{11i}^2}{h_{11}^2} - \frac{\tilde{F}^{ii}u_{ii}}{u - a} + \frac{\tilde{F}^{ii}u_i^2}{(u - a)^2} + A\tilde{F}^{ii}(1 + \langle X, X_{ii} \rangle).$$

Lemma 3.1. *We have*

$$(3.4) \quad \begin{aligned} 0 \geq & -\frac{2}{h_{11}} \sum_{i \geq 2} \tilde{F}^{1i,i1} h_{11i}^2 - \frac{\tilde{F}^{ii} h_{11i}^2}{h_{11}^2} - Ch_{11} \\ & + \frac{a\tilde{F}^{ii} h_{ii}^2}{u - a} + \frac{\tilde{F}^{ii} u_i^2}{(u - a)^2} + A \sum \tilde{F}^{ii} - CA. \end{aligned}$$

Proof. From the formula (2.3), we have

$$(3.5) \quad \tilde{F}^{ii} h_{11ii} = \tilde{F}^{ii} h_{ii} h_{11}^2 - \tilde{F}^{ii} h_{ii}^2 h_{11} + \tilde{F}^{ii} h_{ii11}.$$

Differentiating equation (1.1) twice at X_0 , we obtain

$$(3.6) \quad \tilde{F}^{ii} h_{iik} = (d_X \tilde{f})(e_k) + h_{kk}(d_\nu \tilde{f})(e_k)$$

and

$$(3.7) \quad \tilde{F}^{ii} h_{iikk} + \tilde{F}^{pq,rs} h_{pqk} h_{rsk} \geq -C - Ch_{11}^2 + h_{kk}(d_\nu \tilde{f})(e_i).$$

By the concavity of \tilde{F} and Codazzi formula, we have

$$(3.8) \quad -\tilde{F}^{pq,rs} h_{pq1} h_{rs1} \geq -2 \sum_{i \geq 2} \tilde{F}^{1i,i1} h_{11i}^2$$

Note that by Lemma 2.2 (3) we have $\tilde{F}^{ii} h_{ii} = \tilde{f}$. Hence, we see

$$(3.9) \quad \tilde{F}^{ii} h_{11ii} \geq -2 \sum_{i \geq 2} \tilde{F}^{1i,i1} h_{11i}^2 - \tilde{F}^{ii} h_{ii}^2 h_{11} + h_{11}(d_\nu \tilde{f})(e_i) - C - Ch_{11}^2$$

We now compute the term $\tilde{F}^{ii} u_{ii}$. By (2.2), we have

$$u_i = h_{ii} \langle X, e_i \rangle \text{ and } u_{ii} = \sum_k h_{iik} \langle X, e_k \rangle - u h_{ii}^2 + h_{ii}.$$

Hence, we obtain

$$(3.10) \quad \begin{aligned} \tilde{F}^{ii} u_{ii} &= \sum_k \tilde{F}^{ii} h_{iik} \langle X, e_k \rangle - u \tilde{F}^{ii} h_{ii}^2 + \tilde{f} \\ &\leq \sum_k h_{kk}(d_\nu \tilde{f})(e_k) \langle X, e_k \rangle - u \tilde{F}^{ii} h_{ii}^2 + C \end{aligned}$$

By Gauss formula, we have

$$(3.11) \quad \langle X, X_{ii} \rangle = -h_{ii} \langle X, \nu \rangle = -h_{ii} u$$

Substituting (3.9), (3.10) and (3.11) in (3.3), we obtain that

$$(3.12) \quad \begin{aligned} 0 \geq & -\frac{2}{h_{11}} \sum_{i \geq 2} \tilde{F}^{1i, i1} h_{11i}^2 - \tilde{F}^{ii} h_{ii}^2 + \frac{1}{h_{11}} \sum_l h_{l11} (d_\nu \tilde{f})(e_l) - Ch_{11} \\ & - \frac{\tilde{F}^{ii} h_{11i}^2}{h_{11}^2} - \frac{\sum_k h_{kk} (d_\nu \tilde{f})(e_k) \langle X, e_k \rangle}{u-a} + \frac{u \tilde{F}^{ii} h_{ii}^2}{u-a} - \frac{\tilde{f}}{u-a} \\ & + \frac{\tilde{F}^{ii} u_i^2}{(u-a)^2} + A \sum \tilde{F}^{ii} - CA \end{aligned}$$

By Codazzi formula, $u_k = h_{kk} \langle X, e_k \rangle$ and (3.1), we have

$$\frac{1}{h_{11}} \sum_k h_{k11} (d_\nu \tilde{f})(e_k) - \frac{h_{kk} (d_\nu \tilde{f})(e_k) \langle X, e_k \rangle}{u-a} \geq -CA.$$

Therefore, we arrive at

$$(3.13) \quad \begin{aligned} 0 \geq & -\frac{2}{h_{11}} \sum_{i \geq 2} \tilde{F}^{1i, i1} h_{11i}^2 - \tilde{F}^{ii} h_{ii}^2 - Ch_{11} - \frac{C}{u-a} \\ & - \frac{\tilde{F}^{ii} h_{11i}^2}{h_{11}^2} + \frac{u \tilde{F}^{ii} h_{ii}^2}{u-a} + \frac{\tilde{F}^{ii} u_i^2}{(u-a)^2} + A \sum \tilde{F}^{ii} - CA, \end{aligned}$$

which is just the inequality (3.4). \square

Next, we deal with the bad term $-Ch_{11}$.

Lemma 3.2. *If h_{11} is large enough, we have*

$$(3.14) \quad Ch_{11} \leq \frac{a \tilde{F}^{ii} h_{ii}^2}{2(u-a)} + \frac{A}{2} \sum \tilde{F}^{ii}$$

for sufficiently large A .

Proof. Note that

$$\kappa_{n-p+1} + \kappa_{n-p+2} + \cdots + \kappa_n > 0.$$

We divide the proof in two cases.

Case 1. Suppose $\kappa_n \leq -\delta \kappa_1$, where $\delta > 0$ is a small constant to be determined later. By Lemma 2.2 we see $\tilde{F}^{nn} \geq \theta \sum \tilde{F}^{ii} \geq \theta C_{n-1}^{p-1}$. We then obtain that

$$\tilde{F}^{nn} h_{nn}^2 \geq \delta^2 \kappa_1^2 \tilde{F}^{nn} \geq \theta C_{n-1}^{p-1} \delta^2 \kappa_1^2.$$

Therefore, for sufficiently large κ_1 , we have

$$Ch_{11} \leq \frac{a \tilde{F}^{nn} h_{nn}^2}{2(u-a)}.$$

Case 2. Now $\kappa_n \geq -\delta \kappa_1$. We further divide this case in two cases.

Subcase 2.1. Suppose $\kappa_{n-p+1} + \kappa_{n-p+2} + \cdots + \kappa_n < \frac{\delta}{\kappa_1}$ where $\delta > 0$ is a small constant to be chosen. Since

$$F^{nn} \geq \frac{F(\kappa)}{\kappa_{n-p+1} + \kappa_{n-p+2} + \cdots + \kappa_n},$$

we see that

$$\tilde{F}^{nn} = \frac{1}{C_n^p} F^{\frac{1}{C_n^p}-1} F^{nn} \geq \frac{F^{\frac{1}{C_n^p}} \kappa_1}{C_n^p \delta} = \frac{\tilde{f}}{C_n^p} \frac{\kappa_1}{\delta}$$

Choosing δ sufficiently small, we obtain that

$$C\kappa_1 \leq \tilde{F}^{nn}.$$

Subcase 2.2. Suppose $\kappa_{n-p+1} + \kappa_{n-p+2} + \cdots + \kappa_n > \frac{\delta}{\kappa_1}$. For a fixed $(p-1)$ -tuple $2 \leq i_1 < \cdots < i_{p-1} \leq n$, we have

$$\kappa_1 + \kappa_{i_1} + \cdots + \kappa_{i_{p-1}} \geq (1 - (p-1)\delta)\kappa_1.$$

Hence, we have

$$\begin{aligned} F^{nn} &\geq \prod_{2 \leq i_1 < \cdots < i_{p-1} \leq n} (\kappa_1 + \kappa_{i_1} + \cdots + \kappa_{i_{p-1}}) \\ &\quad \times \prod_{\substack{2 \leq i_1 < \cdots < i_p \leq n \\ (i_1, \dots, i_p) \neq (n-p+1, \dots, n)}} (\kappa_{i_1} + \kappa_{i_2} + \cdots + \kappa_{i_p}) \\ &\geq [(1 - (p-1)\delta)\kappa_1]^{C_{n-1}^{p-1}} \left[\frac{\delta}{\kappa_1} \right]^{C_{n-1}^{p-1}}. \end{aligned}$$

For $p \geq \frac{n}{2}$, direct calculations show that

$$C_{n-1}^{p-1} - C_{n-1}^p = \frac{(n-1) \cdots (n-p+1)}{(p-1)!} \left(1 - \frac{n-p}{p}\right) \geq 0.$$

Therefore, we obtain

$$F^{nn} \geq c_\delta \kappa_1,$$

where $c_\delta = [(1 - (p-1)\delta)]^{C_{n-1}^{p-1}} \delta^{C_{n-1}^p - 1}$. It then follows that, for sufficiently large A ,

$$C\kappa_1 \leq \frac{A}{2} \tilde{F}^{nn}.$$

□

By the above Lemma, (3.4) becomes

$$\begin{aligned} (3.15) \quad 0 &\geq -\frac{2}{h_{11}} \sum_{i \geq 2} \tilde{F}^{1i, i1} h_{11i}^2 - \frac{\tilde{F}^{ii} h_{11i}^2}{h_{11}^2} + \frac{a \tilde{F}^{ii} h_{ii}^2}{2(u-a)} \\ &\quad + \frac{\tilde{F}^{ii} u_i^2}{(u-a)^2} + \frac{A}{2} \sum \tilde{F}^{ii} - CA. \end{aligned}$$

Lemma 3.3. For κ_1 sufficiently large, we have

$$|\kappa_{n-p+1}|, \dots, |\kappa_n| \leq CA.$$

Proof. By the critical equation (3.1) and the Cauchy-Schwarz inequality, we have

$$-\frac{\tilde{F}^{ii} h_{11i}^2}{h_{11}^2} \geq -(1+\varepsilon) \frac{\tilde{F}^{ii} u_i^2}{(u-a)^2} - \left(1 + \frac{1}{\varepsilon}\right) A^2 \tilde{F}^{ii} \langle X, e_i \rangle^2.$$

From (3.15) and $-\tilde{F}^{1i, i1} \geq 0$, we see

$$(3.16) \quad 0 \geq \frac{a \tilde{F}^{ii} h_{ii}^2}{2(u-a)} - \frac{\varepsilon \tilde{F}^{ii} u_i^2}{(u-a)^2} - \frac{CA^2}{\varepsilon} \sum \tilde{F}^{ii} - CA.$$

Using $u_i = h_{ii}\langle X, e_i \rangle$ and choosing ε sufficiently small, we obtain from (3.16) that

$$(3.17) \quad 0 \geq \frac{a\tilde{F}^{ii}h_{ii}^2}{4(u-a)} - \frac{CA^2}{\varepsilon} \sum \tilde{F}^{ii},$$

where we also used $\sum \tilde{F}^{ii} \geq C_{n-1}^{p-1}$. By Lemma 2.2, we now arrive at

$$(3.18) \quad 0 \geq \frac{a\theta}{4(u-a)} \left(\sum \tilde{F}^{ii} \right) \left(\sum_{i \geq n-p+1} h_{ii}^2 \right) - \frac{CA^2}{\varepsilon} \sum \tilde{F}^{ii},$$

which implies that

$$\sum_{i \geq n-p+1} h_{ii}^2 \leq CA^2.$$

□

Now we continue to prove Theorem 1.1. By Lemma 3.3, we can assume

$$\epsilon\kappa_1 \geq \kappa_{n-p+1},$$

where ϵ is as in Lemma 2.2 (5). Then, by Lemma 2.2 (5) and $F^{ij} = 0$ for $i \neq j$, we have for every $i = 2, 3, \dots, n$,

$$-2\tilde{F}^{1i,i1} \geq \frac{\tilde{F}^{ii}}{\kappa_1}.$$

By the critical equation (3.1) and Cauchy-Schwarz inequality, we have

$$\begin{aligned} -\frac{\tilde{F}^{11}h_{11}^2}{h_{11}^2} &\geq -(1+\varepsilon)\frac{\tilde{F}^{11}u_1^2}{(u-a)^2} - \left(1 + \frac{1}{\varepsilon}\right)A^2\tilde{F}^{11}\langle X, e_1 \rangle^2 \\ &\geq -\frac{\tilde{F}^{11}u_1^2}{(u-a)^2} - C\varepsilon\frac{\tilde{F}^{11}h_{11}^2}{(u-a)^2} - \frac{CA^2}{\varepsilon}\tilde{F}^{11}. \end{aligned}$$

By the above two inequalities, we obtain from (3.15) that

$$(3.19) \quad \begin{aligned} 0 &\geq \sum_{i \geq 2} \frac{\tilde{F}^{ii}}{h_{11}^2} h_{11i}^2 - \sum_{i \geq 2} \frac{\tilde{F}^{ii}h_{11i}^2}{h_{11}^2} + \frac{a\tilde{F}^{ii}h_{ii}^2}{2(u-a)} - \frac{\tilde{F}^{11}u_1^2}{(u-a)^2} \\ &\quad - C\varepsilon\frac{\tilde{F}^{11}h_{11}^2}{(u-a)^2} - \frac{CA^2}{\varepsilon}\tilde{F}^{11} + \sum \frac{\tilde{F}^{ii}u_i^2}{(u-a)^2} + \frac{A}{2} \sum \tilde{F}^{ii} - CA. \end{aligned}$$

We can choose ε sufficiently small and assume h_{11} sufficiently large so that (3.19) becomes

$$(3.20) \quad 0 \geq \frac{a\tilde{F}^{ii}h_{ii}^2}{4(u-a)} + \frac{A}{2} \sum \tilde{F}^{ii} - CA.$$

It then follows that

$$\sum \tilde{F}^{ii} \leq C.$$

Next we prove that under this condition, one have

$$\tilde{F}^{11} \geq \frac{1}{C}.$$

Since $\sum \tilde{F}^{ii} \leq C$, especially we have $\frac{F(\kappa)}{\kappa_{n-p+1} + \dots + \kappa_n} \leq C$. It then implies that

$$\kappa_{n-p+1} + \dots + \kappa_n \geq \frac{1}{C}.$$

This yields that

$$\tilde{F}^{ii} \geq \frac{1}{C}, \forall 1 \leq i \leq n.$$

Substituting the above inequality into (3.20), we obtain

$$0 \geq \frac{h_{11}^2}{C} - CA,$$

from which we can derive an upper bound for h_{11} . Theorem 1.1 is proved.

4. GRADIENT ESTIMATES

Before we apply the continuity method to obtain a solution to equation (1.1), we need to derive C^1 estimate for the equation. We show that there exists a positive constant C depending on $n, p, \inf \rho, \sup \rho, \inf f$ and $|f|_{C^1}$ such that

$$|\nabla \rho| \leq C,$$

where ∇ denotes the gradient on \mathbb{S}^n . Note that

$$u = \frac{\rho^2}{\sqrt{\rho^2 + |\nabla \rho|^2}}.$$

We only need to derive a positive lower bound of u . As in [16] we consider the following quantity

$$w = -\log u + \gamma(|X|^2),$$

where the function $\gamma(\cdot)$ will be determined later. Suppose the maximum of w is achieved at $X_0 \in M$. If at X_0 , X is parallel to ν , we have

$$u = \langle X, \nu \rangle = \rho \geq \inf_M \rho,$$

which gives a lower bound since ρ is assumed to have a positive lower bound. If at X_0 , X is not parallel to ν , we can choose a local orthonormal frame $\{e_1, e_2, \dots, e_n\}$ such that

$$\langle X, e_1 \rangle \neq 0 \text{ and } \langle X, e_i \rangle = 0 \text{ for } i \geq 2.$$

Differentiate w at X_0 to obtain that

$$(4.1) \quad 0 = w_i = -\frac{u_i}{u} + 2\gamma' \langle X, e_i \rangle = -\frac{h_{i1} \langle X, e_1 \rangle}{u} + 2\gamma' \langle X, e_i \rangle,$$

where in the last equality we used Weingarten equation. Hence, we have

$$h_{11} = 2\gamma' u \text{ and } h_{1i} = 0 \text{ for } i \geq 2.$$

Without loss of generality, we can assume $\{h_{ij}\}$ is diagonal at X_0 . Differentiating w at X_0 a second time and contracting with $\{\tilde{F}^{ij}\}$, we obtain that

$$(4.2) \quad 0 \geq \tilde{F}^{ii} \left(-\frac{u_{ii}}{u} + \frac{u_i^2}{u^2} + \gamma''(|X|^2)_i + \gamma'(|X|^2)_{ii} \right).$$

Combining (4.1) and the above inequality, we arrive at

$$(4.3) \quad 0 \geq -\frac{\tilde{F}^{ii} u_{ii}}{u} + 4(\gamma'^2 + \gamma'') F^{11} \langle X, e_1 \rangle^2 + \gamma' \tilde{F}^{ii} (|X|^2)_{ii}.$$

By (3.10), we have

$$(4.4) \quad \tilde{F}^{ii} u_{ii} = \langle X, e_1 \rangle ((d_X \tilde{f})(e_1) + h_{11} (d_\nu \tilde{f})(e_1)) - u \tilde{F}^{ii} h_{ii}^2 + \tilde{f}.$$

Also, we have

$$\tilde{F}^{ii} (|X|^2)_{ii} = 2 \sum \tilde{F}^{ii} - 2u \tilde{f}$$

where we used $\tilde{F}^{ij}h_{ij} = \tilde{f}$. Recall that $h_{11} = 2\gamma'u$. Substituting the above two equalities into (4.3) we get

$$(4.5) \quad 0 \geq -\frac{1}{u} \left(\langle X, e_1 \rangle (d_X \tilde{f})(e_1) + \tilde{f} \right) - 2 \langle X, e_1 \rangle \gamma' (d_\nu \tilde{f})(e_1) \\ + \tilde{F}^{ii} h_{ii}^2 + 4(\gamma'^2 + \gamma'') F^{11} \langle X, e_1 \rangle^2 + 2\gamma' \left(\sum \tilde{F}^{ii} - u\tilde{f} \right).$$

At X_0 , we see $X = \langle X, e_1 \rangle e_1 + \langle X, \nu \rangle \nu$. It then follows that

$$(d_X \tilde{f})(X) = \langle X, e_1 \rangle (d_X \tilde{f})(e_1) + \langle X, \nu \rangle (d_X \tilde{f})(\nu).$$

From (1.4), we see that

$$0 \geq \frac{\partial}{\partial \rho} \left(\rho^{C_n^p} f(X, \nu) \right) = \frac{\partial}{\partial \rho} \left(\rho^{C_n^p} \tilde{f}^{C_n^p}(X, \nu) \right) \\ = C_n^p (\rho \tilde{f})^{C_n^p - 1} \left(\tilde{f} + (d_X \tilde{f})(X) \right) \\ = C_n^p (\rho \tilde{f})^{C_n^p - 1} \left(\tilde{f} + \langle X, e_1 \rangle (d_X \tilde{f})(e_1) + \langle X, \nu \rangle (d_X \tilde{f})(\nu) \right).$$

We therefore obtain

$$-(\tilde{f} + \langle X, e_1 \rangle (d_X \tilde{f})(e_1)) \geq \langle X, \nu \rangle (d_X \tilde{f})(\nu) = u (d_X \tilde{f})(\nu).$$

Substituting this into (4.5) we obtain

$$(4.6) \quad 0 \geq (d_X \tilde{f})(\nu) - 2 \langle X, e_1 \rangle \gamma' (d_\nu \tilde{f})(e_1) + \tilde{F}^{ii} h_{ii}^2 \\ + 4(\gamma'^2 + \gamma'') F^{11} \langle X, e_1 \rangle^2 + 2\gamma' \left(\sum \tilde{F}^{ii} - u\tilde{f} \right).$$

Now we choose $\gamma(t) = \frac{\alpha}{t}$, where α is a large constant to be determined. Recall that $h_{11} = 2\gamma'u$ at X_0 , which implies that $h_{11}(X_0) < 0$. This means that $h_{11} \in \{\kappa_{n-p+2}, \kappa_{n-p+3}, \dots, \kappa_n\}$ and therefore by Lemma 2.2

$$F^{11} \geq \theta \sum F^{ii}.$$

Similar to [7], we can assume $\langle X, e_1 \rangle^2 \geq \frac{1}{2} \inf_M \rho^2$. Now we arrive at

$$(4.7) \quad 0 \geq \left(\frac{\alpha^2}{C} - C\alpha \right) \sum \tilde{F}^{ii} - C\alpha.$$

Choosing α sufficiently large, we obtain a contradiction.

5. EXISTENCE OF A SOLUTION

We use the continuity method as in [4] to prove Theorem 1.2. Consider the following family of functions

$$f^t(X, \nu) = t f(X, \nu) + (1-t) p^{C_n^p} \left[\frac{1}{|X|^{C_n^p}} + \varepsilon \left(\frac{1}{|X|^{C_n^p}} - 1 \right) \right]$$

where ε is a small positive constant such that

$$\min_{r_1 \leq \rho \leq r_2} \left[\frac{1}{\rho^{C_n^p}} + \varepsilon \left(\frac{1}{\rho^{C_n^p}} - 1 \right) \right] \geq c_0 > 0,$$

for some positive constant c_0 . It is easy to see that $f^t(X, \nu)$ satisfies (1.3) and (1.4) with strict inequalities for $0 \leq t < 1$.

Let M_t be the solution of the equation

$$F(\kappa) = f^t(X_t, \nu_t),$$

where X_t and ν_t are position vector and unit outer normal of M_t respectively. Clearly, when $t = 0$, we have $M_0 = \mathbb{S}^n$ and $X_0 = x$. For $t \in (0, 1)$, suppose $\rho_t = |X_t|$ attains its maximum at the point x_0 . At this point, we have

$$g_{ij} = \rho_t^2 \delta_{ij} \text{ and } h_{ij} = -(\rho_t)_{ij} + \rho_t \delta_{ij} \geq \rho_t \delta_{ij}$$

Then, we have

$$F(\kappa) \geq F\left(\frac{1}{\rho_t}(1, \dots, 1)\right) = \frac{p^{C_n^p}}{\rho_t^{C_n^p}}.$$

On the other hand, at x_0 , the unit outer normal ν_t is parallel to X_t . If $\rho_t(x_0) = r_2$, we obtain

$$\frac{p^{C_n^p}}{r_2^{C_n^p}} \leq F(\kappa) = f^t(X_t, \nu_t) < \frac{p^{C_n^p}}{r_2^{C_n^p}},$$

which is a contradiction. So we have $\sup_{M_t} \rho_t \leq r_2$. Similarly argument at the minimum point of ρ_t gives that $\inf_{M_t} \rho_t \geq r_1$ on M_t . Hence, C^0 estimate follows. Combining our C^1 estimate, C^2 estimate, Evans-Krylov theorem and the argument in [4] we get the existence and uniqueness of solution to equation (1.1). Theorem 1.2 is proved.

6. PROOF OF THEOREM 1.3

By lemma 3.1, Lemma 3.2 and Lemma 3.3, one can prove Theorem 1.3. For completeness, we include an outline here.

Proof. We consider the following function

$$G(x, \xi) = \log u_{\xi\xi} + \frac{a}{2} |\nabla u|^2 + \frac{A}{2} |x|^2 + \beta \log(v - u),$$

where a, A, β are constants to be determined later. Suppose that G achieves its maximum at (x_0, ξ_0) . Around x_0 , we choose coordinate system such that $\xi_0 = e_1$ and $u_{ij}(x_0)$ is diagonal such that

$$u_{11} \geq u_{22} \geq \dots \geq u_{nn} \text{ at } x_0.$$

This can be done as in [12]. Thus the new function defined by

$$\hat{G}(x) = \log u_{11} + \frac{a}{2} |\nabla u|^2 + \frac{A}{2} |x|^2 + \beta \log(v - u)$$

also attains its maximum at x_0 . Differentiate it once to obtain

$$(6.1) \quad 0 = \frac{u_{11i}}{u_{11}} + a u_i u_{ii} + A x_i + \frac{\beta(v - u)_i}{v - u}.$$

Differentiating it twice and by similar computations as Lemma 3.1 and Lemma 3.2 we can arrive at

$$(6.2) \quad \begin{aligned} 0 \geq & -\frac{2}{u_{11}} \sum_{i \geq 2} \tilde{F}^{1i, i1} u_{11i}^2 - \frac{\tilde{F}^{ii} u_{11i}^2}{u_{11}^2} + \frac{a \tilde{F}^{ii} u_{ii}^2}{2} \\ & + \frac{A}{2} \sum \tilde{F}^{ii} - CA - \frac{\beta \tilde{F}^{ii} (v - u)_i^2}{(v - u)^2} - \frac{C\beta}{v - u}. \end{aligned}$$

Using the same argument as in Lemma 3.3, we obtain

$$u_{ii} \leq \frac{C}{v - u} \text{ for } i \geq n - p + 1,$$

where C depends on a, A and β . Now by Lemma 2.2 (5) we can assume without loss of generality that

$$\epsilon u_{11} \geq u_{n-p+1n-p+1}.$$

and

$$-2\tilde{F}^{1i,i1} \geq (1+\delta)\frac{\tilde{F}^{ii}}{u_{11}},$$

where ϵ and δ are as in (5).

On the other hand, by (6.1), we have

$$-\frac{\tilde{F}^{11}u_{11}^2}{u_{11}^2} \geq -Ca^2\tilde{F}^{11}u_{11}^2 - CA^2\tilde{F}^{11} - \frac{C\beta^2\tilde{F}^{11}}{(v-u)^2}$$

and

$$-\sum_{i \geq 2} \frac{\beta\tilde{F}^{ii}(v-u)_i^2}{(v-u)^2} \geq -\frac{3}{\beta} \sum_{i \geq 2} \frac{\tilde{F}^{ii}u_{11i}^2}{u_{11}^2} - \frac{Ca^2}{\beta} \sum_{i \geq 2} \tilde{F}^{ii}u_{ii}^2 - \frac{CA^2}{\beta} \sum_{i \geq 2} \tilde{F}^{ii}.$$

Substituting the above two inequalities in (6.2) we have

$$\begin{aligned} 0 &\geq -\frac{2}{u_{11}} \sum_{i \geq 2} \tilde{F}^{1i,i1}u_{11i}^2 - \left(1 + \frac{3}{\beta}\right) \sum_{i \geq 2} \frac{\tilde{F}^{ii}u_{11i}^2}{u_{11}^2} + \left(\frac{a}{2} - \frac{Ca^2}{\beta}\right) \tilde{F}^{ii}u_{ii}^2 \\ (6.3) \quad &+ \left(\frac{A}{2} - \frac{CA^2}{\beta}\right) \sum \tilde{F}^{ii} - Ca^2\tilde{F}^{11}u_{11}^2 - CA^2\tilde{F}^{11} - \frac{C\beta^2\tilde{F}^{11}}{(v-u)^2} \\ &- CA - \frac{2\beta\tilde{F}^{11}u_1^2}{(v-u)^2} - \frac{C\beta}{v-u}. \end{aligned}$$

Choose β sufficiently large and a small to see that

$$\begin{aligned} 0 &\geq -\frac{2}{u_{11}} \sum_{i \geq 2} \tilde{F}^{1i,i1}u_{11i}^2 - \left(1 + \frac{3}{\beta}\right) \sum_{i \geq 2} \frac{\tilde{F}^{ii}u_{11i}^2}{u_{11}^2} + \frac{a}{4}\tilde{F}^{ii}u_{ii}^2 \\ (6.4) \quad &+ \frac{A}{4} \sum \tilde{F}^{ii} - \left(CA^2 + \frac{C\beta^2}{(v-u)^2}\right) \tilde{F}^{11} - CA - \frac{C\beta}{v-u}. \end{aligned}$$

Now we choose β sufficiently large such that $\delta \geq \frac{3}{\beta}$. We then arrive at

$$\begin{aligned} 0 &\geq \frac{a}{4}\tilde{F}^{11}u_{11}^2 - \frac{C}{(v-u)^2}\tilde{F}^{11} - CA - \frac{C}{v-u} \\ (6.5) \quad &\geq \frac{a}{8}\tilde{F}^{11}u_{11}^2 - \frac{C}{v-u}, \end{aligned}$$

where in the second inequality we assumed u_{11} is large enough. From Lemma 2.2 (1) we have that $\tilde{F}^{11}u_{11} \geq c_0$, where $c_0 > 0$ depends on $\inf \tilde{f}$. We then derive that

$$(v-u)u_{11} \leq C$$

which implies the estimate (1.6). \square

REFERENCES

- [1] J.M. Ball, Differentiability properties of symmetric and isotropic functions, *Duke Math. J.* 51 (1984), 699-728.
- [2] I. Bakelman, B. Kantor, Existence of a hypersurface homeomorphic to the sphere in Euclidean space with a given mean curvature. (Russian) *Geometry and topology*, No. 1 (Russian), pp. 3-10. Leningrad. Gos. Ped. Inst. im. Gercena, Leningrad, 1974.
- [3] L. Caffarelli, L. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second order elliptic equations I. Monge-Ampère equations, *Comm. Pure Appl. Math.* 37, (1984), 369-402.
- [4] L. A. Caffarelli, L. Nirenberg and J. Spruck, Nonlinear second order elliptic equations IV: Starshaped compact Weingarten hypersurfaces, *Current topics in partial differential equations*, Y. Ohya, K. Kasahara and N. Shimakura (eds), Kinokunize, Tokyo, 1985, 1-26.
- [5] D. Chen, H. Li and Z. Wang, Starshaped compact hypersurfaces with prescribed Weingarten curvature in warped product manifolds, *Calc. Var. Partial Differential Equations* 57 (2018), no. 2, Paper No. 42, 26 pp.
- [6] K.-S. Chou and X.-J. Wang A variational theory of the Hessian equation, *Comm. Pure Appl. Math.* 54 (2001), no. 9, 1029-1064.
- [7] J. Chu and H. Jiao, Curvature estimates for a class of Hessian type equations, *Calc. Var. Partial Differential Equations* 60 (2021), no. 3, Paper No. 90, 18 pp.
- [8] S. Dinew, Interior estimates for p -plurisubharmonic functions, arXiv:2006.12979.
- [9] J. Fu, Z. Wang and D. Wu, Form-type Calabi-Yau equations, *Math. Res. Lett.* 17 (2010), no. 5, 887-903.
- [10] J. Fu, Z. Wang and D. Wu, Form-type equations on Kähler manifolds of nonnegative orthogonal bisectional curvature, *Calc. Var. Partial Differential Equations* 52 (2015), no. 1-2, 327-344.
- [11] C. Gerhardt, Hypersurfaces of prescribed Weingarten curvature, *Math. Z.* 224 (1997), 167-194.
- [12] B. Guan, Second-order estimates and regularity for fully nonlinear elliptic equations on Riemannian manifolds, *Duke Math. J.* 163 (2014), no. 8, 1491-1524.
- [13] B. Guan and P. Guan, Convex Hypersurfaces of Prescribed Curvature, *Annals of Mathematics*, 156 (2002), 655-674.
- [14] B. Guan and H. Jiao, Second order estimates for Hessian type fully nonlinear elliptic equations on Riemannian manifolds, *Calc. Var. Partial Differential Equations* 54 (2015), 2693-2712.
- [15] P. Guan, J. Li and Y. Li, Hypersurfaces of prescribed curvature measure, *Duke Math. J.* 161 (2012), no. 10, 1927-1942.
- [16] P. Guan, C. Lin and X.-N. Ma, The existence of convex body with prescribed curvature measures, *Int. Math. Res. Not. IMRN* 2009, no. 11, 1947-1975.
- [17] P.F. Guan, C. Ren and Z. Wang, Global C^2 -estimates for convex solutions of curvature equations, *Comm. Pure Appl. Math.* 68 (2015), 1287-1325.
- [18] F. R. Harvey and H. B. Lawson, Geometric plurisubharmonicity and convexity: an introduction, *Adv. Math.* 230 (2012), no. 4-6, 2428-2456.
- [19] F. R. Harvey and H. B. Lawson, p -convexity, p -plurisubharmonicity and the Levi problem, *Indiana Univ. Math. J.* 62 (2013), no. 1, 149-169.
- [20] N. M. Ivochkina, Solution of the Dirichlet problem for equations of m th order curvature. (Russian), *Mat. Sb.* 180 (1989), no. 7, 867-887, 991; translation in *Math. USSR-Sb.* 67 (1990), no. 2, 317-339.
- [21] N. M. Ivochkina, The Dirichlet problem for the curvature equation of order m , *Algebra i Analiz* 2 (1990), no. 3, 192-217; translation in *Leningrad Math. J.* 2 (1991), no. 3, 631-654.
- [22] Y.Y. Li, Interior gradient estimates for solutions of certain fully nonlinear elliptic equations, *J. Differential Equations* 90 (1991) 172-185.
- [23] V. Oliker, Hypersurfaces in \mathbf{R}^{n+1} with prescribed Gaussian curvature and related equations of Monge-Ampère type. *Commun. Partial Diff. Equations* 9 (1984), 807-838.
- [24] C. Ren and Z. Wang, On the curvature estimates for Hessian equations, *Amer. J. Math.* 141 (2019), no. 5, 1281-1315.
- [25] C. Ren and Z. Wang, The global curvature estimate for the $n - 2$ Hessian equation, arXiv:2002.08702.
- [26] C. Ren and Z. Wang, Notes on the curvature estimates for Hessian equations, arXiv:2003.14234v1.

- [27] C. Ren, Z. Wang and L. Xiao, The prescribed curvature problem for entire hypersurfaces in Minkowski space, arXiv:2007.04493v1.
- [28] G. Székelyhidi, V. Tosatti and B. Weinkove, Gauduchon metrics with prescribed volume form, Acta Math. 219 (2017), 181-211.
- [29] J. Sha, p -convex Riemannian manifolds, Invent. Math. 83 (1986), no. 3, 437-447.
- [30] J. Sha, Handlebodies and p -convexity. J. Differential Geom. 25 (1987), no. 3, 353-361.
- [31] V. Tosatti and B. Weinkove, The Monge-Ampère equation for $(n-1)$ -plurisubharmonic functions on a compact Kähler manifold, J. Amer. Math. Soc. 30 (2017), no. 2, 311-346.
- [32] V. Tosatti and B. Weinkove, Hermitian metrics, $(n-1, n-1)$ forms and Monge-Ampère equations, J. Reine Angew. Math. 755 (2019), 67-101.
- [33] A.E. Treibergs and S.W. Wei, Embedded hypersurfaces with prescribed mean curvature, J. Differential Geom. 18 (1983), 513-521.
- [34] Z. Wang and L. Xiao, Entire spacelike hypersurfaces with constant σ_k curvature in Minkowski space, arXiv:2007.01495v1.
- [35] H. Wu, Manifolds of partially positive curvature, Indiana Univ. Math. J. 36 (1987), no. 3, 525-548.

SCHOOL OF MATHEMATICS, TIANJIN UNIVERSITY, 135 YAGUAN ROAD, JINNAN, TIANJIN, CHINA, 300354

Email address: dr.dong@tju.edu.cn