

Singular solutions of some elliptic equations involving mixed absorption-reaction

Marie-Françoise Bidaut-Véron*,
Marta Garcia-Huidobro †
Laurent Véron ‡

Abstract

We study properties of nonnegative functions satisfying (E) $-\Delta u + u^p - M|\nabla u|^q = 0$ in a domain of \mathbb{R}^N when $p > 1$, $M > 0$ and $1 < q < p$. We concentrate our analysis on the solutions of (E) with an isolated singularity, or in an exterior domain, or in the whole space. The existence of such solutions and their behaviours depend strongly on the values of the exponents p and q and in particular according to the sign of $q - \frac{2p}{p+1}$, and when $q = \frac{2p}{p+1}$, also on the value of the parameter M which becomes a key element. The description of the different behaviours is made possible by a sharp analysis of the radial solutions of (E).

2010 Mathematics Subject Classification: 35J60-35J62-35J75-34A34-34C37

Keywords: Elliptic equations, isolated singularities, dynamical systems, linearization, super and sub solutions, Lyapounov functions, central manifold.

Contents

1	Introduction	3
2	The case $q = \frac{2p}{p+1}$	11
2.1	The equation on the sphere	11
2.1.1	Proof of Theorem 1.1: constant positive solutions	11
2.1.2	Proof of Theorem 1.1: positive solutions	12
2.2	Radial solutions	15
2.2.1	Linearisation at $(0,0)$	16
2.2.2	Linearisation at the non-trivial equilibrium points	17

*Laboratoire de Mathématiques et Physique Théorique, Université de Tours, 37200 Tours, France. E-mail: veronmf@univ-tours.fr

†Departamento de Matemáticas, Pontificia Universidad Católica de Chile Casilla 307, Correo 2, Santiago de Chile. E-mail: mgarcia@mat.puc.cl

‡Laboratoire de Mathématiques et Physique Théorique, Université de Tours, 37200 Tours, France. E-mail: veronl@univ-tours.fr

2.2.3	The vanishing curves of the vector field	18
2.3	Description of the radial solutions defined near 0	22
2.3.1	The case $1 < p < \frac{N}{N-2}$ and $M \geq 0$	22
2.3.2	The case $p = \frac{N}{N-2}$, $q = \frac{2p}{p+1} = \frac{N}{N-1}$ and $M > 0$	25
2.3.3	The case $p > \frac{N}{N-2}$ and $M > 0$	26
3	The radial case for $q \neq \frac{2p}{p+1}$	29
3.1	Non-autonomous systems associated to the equation	29
3.1.1	System describing the behaviour of Emden-Fowler equation	29
3.1.2	System describing the behaviour of the Riccati equation	30
3.1.3	System describing the behaviour of the eikonal equation	30
3.1.4	Lyapounov and slope functions	31
3.2	Asymptotic estimates for the Riccati equation	33
3.3	Estimates near 0	35
3.3.1	The case $\frac{2p}{p+1} < q < p$	35
3.3.2	The case $1 < q < \frac{2p}{p+1}$	39
3.4	Estimates at infinity	40
3.4.1	The case $q > \frac{2p}{p+1}$	40
3.4.2	The case $1 < q < \frac{2p}{p+1}$	42
3.5	Solutions of eikonal type	43
3.6	Local or global existence results	46
3.6.1	The systems of order 3	46
3.6.2	Singular solutions of eikonal type	48
3.6.3	Riccati type singular solutions	49
3.6.4	Emden-Fowler type singular solutions	52
3.6.5	Solutions behaving like the Newtonian potential	53
4	Isolated singularities of non-radial solutions	57
4.1	Existence and uniqueness of singular solutions	57
4.2	Characterization of singular solutions	61
4.3	The case $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{N}{N-1}$	62
4.4	The case $p \geq \frac{N}{N-2}$ and $q = \frac{2p}{p+1}$	68
5	Behaviour at infinity of non-radial solutions	73
5.1	The case $q = \frac{2p}{p+1}$	74
5.2	The case $q \neq \frac{2p}{p+1}$	78
A	Appendix	80
A.1	The a priori estimates	80
A.2	Equilibrium with a simple eigenvalue	81

1 Introduction

The aim of this article is to study existence and properties of nonnegative singular solutions of the following equation

$$\mathcal{L}_{p,q}^M u := -\Delta u + u^p - M|\nabla u|^q = 0 \quad (1.1)$$

in a domain Ω of \mathbb{R}^N or in \mathbb{R}^N ($N \geq 2$), where M is a real number and $p > q > 1$.

In the case $M < 0$ many results dealing with isolated singularities have been obtained in [17]. Therefore we will mainly concentrate on the case $M > 0$ where the two nonlinear terms act in a opposite direction: one is an absorption and the other is a source. Furthermore they are not of the same type, one involves the function and the other its gradient.

First we consider the case $q = \frac{2p}{p+1}$. Then (1.1) becomes

$$\mathcal{L}_{p, \frac{2p}{p+1}}^M u := -\Delta u + u^p - M|\nabla u|^{\frac{2p}{p+1}} = 0, \quad (1.2)$$

and this equation is invariant under the scaling transformation T_ℓ , $\ell > 0$, defined by

$$T_\ell[u](x) = \ell^\alpha u(\ell x). \quad (1.3)$$

In that case there may exist self-similar solutions, necessarily under the form $u(x) = u(r, s) = r^{-\alpha}\omega(s)$, where $(r, s) \in \mathbb{R}_+ \times S^{N-1}$ are the spherical coordinates in \mathbb{R}^N . The function ω is a solution of the following equation on S^{N-1}

$$-\Delta' \omega + \ell_{N,p} \omega + \omega^p - M(\alpha^2 \omega^2 + |\nabla' \omega|^2)^{\frac{p}{p+1}} = 0, \quad (1.4)$$

where Δ' and ∇' denote respectively the Laplace-Beltrami operator and the tangential gradient on S^{N-1} , identified with the covariant gradient on S^{N-1} for the metric induced by the standard one in \mathbb{R}^N , and where

$$\alpha = \frac{2}{p-1} \quad \text{and} \quad \ell_{N,p} = \alpha K \quad (1.5)$$

with

$$K = N - 2 - \alpha = \frac{(N-2)p - N}{N-2}. \quad (1.6)$$

The nonzero constant solutions of (1.4) are the positive zeros of the function

$$\mathcal{P}_M(x) = x^{p-1} - M \alpha^{\frac{2p}{p+1}} x^{\frac{p-1}{p+1}} + \ell_{N,p}. \quad (1.7)$$

The following value of the parameter M , which exists only if $N \geq 3$ and $p \geq \frac{N}{N-2}$, plays an important role in the study of (1.4):

$$m^* := (p+1) \left(\frac{(N-2)p - N}{2p} \right)^{\frac{p}{p+1}}. \quad (1.8)$$

The separable solutions obtained in the next theorem are at the core of the process of describing the behaviour of positive solutions of (1.1) near an isolated singularity or in an exterior domain of \mathbb{R}^N .

Theorem 1.1 *Let $p > 1$, then*

1- *If $M \leq 0$ equation (1.4) admits a positive solution if and only if $N = 2$ or $N \geq 3$ and $p < \frac{N}{N-2}$. Furthermore this solution is constant, unique and denoted by x_M .*

2- *If $M > 0$ and $p \leq \frac{N}{N-2}$ if $N \geq 3$, or any $p > 1$ if $N = 2$, equation (1.4) admits a unique positive solution. It is constant and denoted by x_M .*

3- *If $N \geq 3$, $p > \frac{N}{N-2}$ and $M = m^*$ there exists one positive solution to (1.4). It is constant and denoted by x_{m^*} .*

4- *If $N \geq 3$, $p > \frac{N}{N-2}$ and $0 < M < m^*$ there exists no positive solution to (1.4).*

5- *If $N \geq 3$, $p > \frac{N}{N-2}$ and $M > m^*$ there exist two constant positive solutions $x_{1,M} < x_{2,M}$ to (1.4) and any positive solution ω satisfies*

$$0 < \min_{S^{N-1}} \omega \leq x_{1,M} \leq \max_{S^{N-1}} \omega \leq x_{2,M}. \quad (1.9)$$

Furthermore, if

$$m^* < M < \tilde{m} := \frac{(p+1)^2}{2} \left(\frac{(N-2)p^2 - (N+2)}{4p^2} \right)^{\frac{p}{p+1}}, \quad (1.10)$$

then $x_{1,M}$ and $x_{2,M}$ are the only positive solutions, and

$$\frac{\tilde{m}}{m^*} > \left(\frac{p+1}{2p} \right)^{\frac{p}{p+1}} \frac{p+1}{2} > \frac{N-1}{N-2} \left(\frac{N-1}{N} \right)^{\frac{N}{2(N-1)}} > 1. \quad (1.11)$$

.

Not all the singular positive solutions of (1.2) are self-similar since there exist solutions with a *weak singularity*, which means

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow 0} |x|^{N-2} u(x) = k && \text{if } N \geq 3, \\ (ii) \quad & \lim_{x \rightarrow 0} |\ln |x||^{-1} u(x) = k && \text{if } N = 2. \end{aligned} \quad (1.12)$$

Thanks to the existence of positive radial singular solutions in $\mathbb{R}^N \setminus \{0\}$ we are able to prove the existence of non-radial positive solution in a punctured bounded domain with prescribed boundary value. This is a very general tool which is developed in Section 4 for obtaining singular solutions, and as an example we prove the following result.

Theorem 1.2 *Let Ω be a bounded smooth domain of \mathbb{R}^N ($N \geq 3$) containing 0 and $\phi \in W^{1,\infty}(\partial\Omega)$. If $1 < p < \frac{N}{N-2}$ then for any real $M > 0$ and $k > 0$ there exists a minimal positive solution u_k of (1.2) in $\Omega \setminus \{0\}$ satisfying (1.12) and such that $u = \phi$ on $\partial\Omega$. Furthermore, $k \mapsto u_k$ is increasing and $u_k \uparrow u_\infty$ where u_∞ is the minimal solution of (1.2) in $\Omega \setminus \{0\}$ satisfying (1.12), such that $u = \phi$ on $\partial\Omega$ and satisfying*

$$\lim_{x \rightarrow 0} |x|^{\frac{2}{p-1}} u_\infty(x) = x_M. \quad (1.13)$$

If $p \geq \frac{N}{N-2}$ it is proved in [7] that there exists no positive solution of (1.2) with weak singularity at 0 and that any positive solution in $\Omega \setminus \{0\}$ can be extended as a weak solution in whole Ω . However weak solutions may be unbounded. The different kinds of singular solutions play a key role for describing the behaviour near 0 of any positive solution of (1.2) in $\Omega \setminus \{0\}$. If Ω is replaced by \mathbb{R}^N there holds:

Theorem 1.3 *Let $N \geq 3$ and $1 < p < \frac{N}{N-2}$. Then for any real $M > 0$ and $k > 0$ there exists a unique positive solution u_k of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.12) and*

$$\lim_{|x| \rightarrow \infty} |x|^{\frac{2}{p-1}} u_k(x) = x_M. \quad (1.14)$$

Furthermore u_k is radial and $u_k \uparrow u_{x_M}$ as $k \rightarrow \infty$, where $u_{x_M}(x) = x_M |x|^{-\alpha}$.

When $p \geq \frac{N}{N-2}$ new phenomena appear.

Theorem 1.4 *Let $N \geq 3$, $p = \frac{N}{N-2}$ and $M > 0$. Then the function*

$$u_{x_M}(x) = x_M |x|^{2-N} = \left((N-2) M^{\frac{N-1}{N}} \right)^{N-2} |x|^{2-N}$$

is the unique radial positive solution of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying $\lim_{|x| \rightarrow 0} |x|^{N-2} u(x) = x_M$. Moreover there exists a positive solution u_s of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\begin{aligned} (i) \quad & \lim_{|x| \rightarrow 0} |x|^{N-2} |\ln |x||^{N-1} u_s(x) = \frac{1}{(N-2) ((N-1)M)^{N-1}} \\ (ii) \quad & \lim_{|x| \rightarrow \infty} |x|^{N-2} u_s(x) = \left((N-2) M^{\frac{N-1}{N}} \right)^{N-2}. \end{aligned} \quad (1.15)$$

Furthermore u_s is the unique positive solution (not only radial) satisfying (1.15).

The proof of existence is based upon a dynamical system formulation of the equation, see (2.16). Such a formulation, as well as similar ones, will be much used in the sequel.

Theorem 1.5 *Let $N \geq 3$ and $p > \frac{N}{N-2}$.*

1- *If $M > m^*$, besides the two self-similar solutions $u_{x_{j,M}}$ ($j = 1, 2$), there exists a radial positive solution u_s of (1.2) in $\mathbb{R}^N \setminus \{0\}$, unique among the radial ones up to the scaling transformation T_ℓ , satisfying*

$$\lim_{|x| \rightarrow 0} |x|^\alpha u_s(x) = x_{1,M} \quad \text{and} \quad \lim_{|x| \rightarrow \infty} |x|^\alpha u_s(x) = x_{2,M}. \quad (1.16)$$

For any $k > 0$ there exists also a radial positive solution u of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{|x| \rightarrow 0} |x|^\alpha u(x) = x_{1,M} \quad \text{and} \quad \lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k > 0. \quad (1.17)$$

It is unique among the radial positive solutions satisfying (1.17). Furthermore $T_\ell[u_c] = u_{c\ell^{\alpha+2-N}}$.

2- If $M = m^*$, the self-similar solution $u_{m^*}(x) = x_{m^*}|x|^{-\alpha}$ is the unique among the radial positive solutions of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{|x| \rightarrow 0} |x|^\alpha u(x) = x_{m^*} \quad \text{and} \quad \lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{m^*}. \quad (1.18)$$

Furthermore, for any $k > 0$ there exists also a radial positive solution u_k of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.17) with $x_{1,M}$ replaced by x_{m^*} . It is unique among the positive radial solutions satisfying (1.17) and it satisfies the same scaling invariance as (i).

The previous results allow to describe the behaviour at infinity of radial positive solutions of (1.2) in the complement of a ball. The next result will be partially extended to non-radial solutions in Section 5.

Proposition 1.6 *Let $N \geq 1$, $p > 1$, $M > 0$ and u be a positive radial solution of (1.2) in $\mathbb{R}^N \setminus B_R$ for some $R > 0$.*

1- *If $N = 2$, or $N \geq 3$ and $1 < p < \frac{N}{N-2}$, then $\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_M$.*

2- *If $N \geq 3$ and $p = \frac{N}{N-2}$, then $\lim_{|x| \rightarrow \infty} |x|^{N-2} (\ln |x|)^{\frac{N-2}{2}} u(x) = \left(\frac{N-2}{\sqrt{2}}\right)^{N-2}$ or $\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = x_M$.*

3- *If $N \geq 3$ and $p > \frac{N}{N-2}$,*

3-a- *if $0 < M < m^*$, then $\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k$ for some $k > 0$.*

3-b- *if $M = m^*$, then either $\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{m^*}$, or $\lim_{r \rightarrow \infty} r^{N-2} u(r) = k$ for some $k > 0$.*

2-c- *if $M > m^*$, then either $\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{1,M}$, or $\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{2,M}$ or*

$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k$ for some $k > 0$.

Next, we consider equation (1.1) when $q \neq \frac{2p}{p+1}$. In that case, the asymptotics of the solutions are governed either by the Emden-Fowler operator

$$u \mapsto \mathcal{L}_p u := -\Delta u + u^p, \quad (1.19)$$

or by the Riccati operator

$$u \mapsto \mathcal{R}_q^M u := -\Delta u - M|\nabla u|^q, \quad (1.20)$$

or by the eikonal operator

$$u \mapsto \mathcal{E}_{p,q}^M u := u^p - M|\nabla u|^q. \quad (1.21)$$

When $1 < q < \frac{2p}{p+1}$ the governing equation is the Emden-Fowler equation $\mathcal{L}_p u = 0$ near a singularity and the Riccati equation $\mathcal{R}_q^M u = 0$ at infinity. When $\frac{2p}{p+1} < q < p$, the situation is reversed. The following exponents play a crucial role

$$\alpha = \frac{2}{p-1}, \quad \beta = \frac{2-q}{q-1} \quad \text{and} \quad \gamma = \frac{q}{p-q} \quad \text{if } q \neq p, \quad (1.22)$$

and

$$\sigma = (p+1)q - 2p. \quad (1.23)$$

We also define

$$\kappa = \frac{(N-1)q - N}{q-1} \quad \text{if } q > 1, \quad (1.24)$$

and

$$\theta = \frac{(N-1)q - (N-2)p}{p-q} = \gamma + 2 - N \quad \text{if } q \neq p. \quad (1.25)$$

Theorem 1.7 *Let $N \geq 1$, $M > 0$ and $\frac{2p}{p+1} < q < p$. If there exists a radial positive solution u of (1.1) in $B_R \setminus \{0\}$ which is unbounded near 0, then*

1- either

$$\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M \quad \text{where } X_M = (M\gamma^q)^{\frac{1}{p-q}}, \quad (1.26)$$

2- or (1.26) does not hold. In that case $q \leq 2$, $N \geq 2$ and the following situation occurs:

2-a- if $\frac{N}{N-1} < q < 2$, then

$$\lim_{x \rightarrow 0} |x|^\beta u(x) = \xi_M \quad \text{where } \xi_M = \frac{1}{\beta} \left(\frac{\kappa}{M} \right) (M\gamma^q)^{\frac{1}{q-1}}, \quad (1.27)$$

2-b- if $q = 2$, then

$$\lim_{x \rightarrow 0} |\ln |x||^{-1} u(x) = \frac{N-2}{M} \quad \text{if } N \geq 3, \quad \text{or} \quad \lim_{x \rightarrow 0} (\ln |\ln |x||) u(x) = \frac{1}{M} \quad \text{if } N = 2, \quad (1.28)$$

2-c- if $q < \frac{N}{N-1}$, then there exists $k > 0$ such that

$$\lim_{x \rightarrow 0} |x|^{N-2} u(x) = k \quad \text{if } N \geq 3 \quad \text{and} \quad \lim_{x \rightarrow 0} |\ln |x||^{-1} u(x) = k \quad \text{if } N = 2, \quad (1.29)$$

2-d- if $q = \frac{N}{N-1}$, then

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow 0} ||x| \ln |x||^{N-1} u(x) = \frac{1}{N-1} \left(\frac{N-1}{M} \right)^{N-1} \quad \text{if } N \geq 3 \\ (ii) \quad & \lim_{x \rightarrow 0} |\ln |x||^{-1} u(x) = k > 0 \quad \text{if } N = 2 \end{aligned} \quad (1.30)$$

In the case $1 < q < \frac{2p}{p+1}$ the description of isolated singularities is simpler and it is similar to the one of the positive solutions of (1.19).

Theorem 1.8 *Let $M > 0$, $1 < p < \frac{N}{N-2}$ if $N \geq 3$, any $p > 1$ if $N = 1, 2$, and $1 < q < \frac{2p}{p+1}$. Assume that there exists a radial positive solution u of (1.1) in $B_R \setminus \{0\}$ which is unbounded near 0. Then the following alternative holds:*

1- either

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_0 := (\alpha |K|)^{\frac{1}{p-1}}, \quad (1.31)$$

2- or $N \geq 2$ and

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow 0} |x|^{N-2} u(x) = k > 0 \quad \text{if } N \geq 3 \\ (ii) \quad & \lim_{x \rightarrow 0} |\ln |x||^{-1} u(x) = k > 0 \quad \text{if } N = 2. \end{aligned} \quad (1.32)$$

It is noticeable that all the behaviours described in the previous two theorems occur. The behaviour at infinity of positive solutions of (1.1) in B_R^c inherits this complexity due to the value of q with respect to $\frac{2p}{p+1}$, and the situation is less intricated in the case $\frac{2p}{p+1} < q < p$ than in the case $1 < q < \frac{2p}{p+1}$.

Theorem 1.9 *Let $N \geq 1$, $M > 0$ and $\frac{2p}{p+1} < q < p$. Assume that there exists a radial positive solution u of (1.1) in B_R^c . Then*

1- If $1 < p < \frac{N}{N-2}$ (any $p > 1$ if $N = 1, 2$), there holds

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_0. \quad (1.33)$$

2- If $N \geq 3$ and $p > \frac{N}{N-2}$, there holds

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k > 0. \quad (1.34)$$

3- If $N \geq 3$ and $p = \frac{N}{N-2}$, there holds

$$\lim_{|x| \rightarrow \infty} (\ln |x|)^{\frac{N-2}{2}} |x|^{N-2} u(x) = \left(\frac{N-2}{\sqrt{2}} \right)^{N-2}. \quad (1.35)$$

Theorem 1.10 *Let $N \geq 2$, $M > 0$ and $1 < q < \frac{2p}{p+1}$. If u is a radial positive solution of (1.1) in B_R^c , there holds.*

1- If $q > \frac{N}{N-1}$, one of the three following situations occurs:

1-a- either

$$\lim_{|x| \rightarrow \infty} |x|^\gamma u(x) = X_M, \quad (1.36)$$

1-b- or

$$\lim_{|x| \rightarrow \infty} |x|^\beta u(x) = \xi_M, \quad (1.37)$$

1-c- or

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k > 0. \quad (1.38)$$

2- If $N = 1$ or $1 < q \leq \frac{N}{N-1}$, then only (1.36) can occur.

The existence of local or global singular solutions or asymptotic solutions with behaviour like $|x|^{-\gamma}$ (eikonal type) or like $|x|^{-\beta}$ (Riccati type) near 0 or ∞ will be proved in Section 3.6. For example we prove the following result by the method of sub and super solutions.

Theorem 1.11 *Let $N \geq 1$, $p > 1$ and $M > 0$.*

1- *If $\frac{2p}{p+1} < q < p$, then there exists a unique global positive solution u of (1.1) such that $\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M$, and its behaviour at infinity is given by Theorem 1.9.*

Moreover this solution is radial, and it is explicit if $N \geq 3$ and $q = \frac{(N-2)p}{N-1}$. Furthermore, for any bounded smooth domain Ω containing 0 there exists a positive solution of (1.1) in $\Omega \setminus \{0\}$ vanishing on $\partial\Omega$.

2- *If $\max\left\{1, \frac{(N-2)p}{N-1}\right\} < q < \frac{2p}{p+1}$, then for any $R > 0$ there exists a positive solution u in B_R^c satisfying (1.36).*

Introducing a new powerful autonomous system of order 3, we can construct local solutions behaving like $|x|^{-\beta}$ near 0.

Theorem 1.12 *Let $N \geq 2$, $p > 1$ and $M > 0$.*

1- *If $\max\left\{\frac{2p}{p+1}, \frac{N}{N-1}\right\} < q < \min\{2, p\}$. Then there exists at least one radial positive solution u of (1.1) in a neighborhood of 0 such that $\lim_{x \rightarrow 0} |x|^\beta u(x) = \xi_M$.*

2- *If $\frac{N}{N-1} < q < \frac{2p}{p+1}$ there exists a unique positive radial solution defined in a neighborhood of infinity satisfying such that $\lim_{|x| \rightarrow \infty} |x|^\beta u(x) = \xi_M$. There exists no radial positive solution in $\mathbb{R}^N \setminus \{0\}$ with such a behaviour at infinity.*

By a delicate method of super and sub solutions, we also prove the existence of radial positive singular solutions u of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.37) under more restrictive assumptions on the exponents p and q .

When $p < \frac{N}{N-2}$ we show the existence of the solutions of (1.1) in $\mathbb{R}^N \setminus \{0\}$, or in a neighborhood of 0, or at infinity having the behaviour described in Theorem 1.8 and Theorem 1.9. Such solutions are associated to the Emden-Fowler operator.

Theorem 1.13 *Let $M > 0$, $N \geq 3$ and $1 < p < \frac{N}{N-2}$, or $N = 1, 2$ and $p > 1$.*

1- *If $1 < q < \frac{2p}{p+1}$ there exists a unique positive solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying*

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow 0} |x|^\alpha u(x) = x_0 \\ (ii) \quad & \lim_{|x| \rightarrow \infty} |x|^\gamma u(x) = X_M. \end{aligned} \tag{1.39}$$

Furthermore this solution is radial and $|x|^\alpha u(x) \geq x_0$ in $\mathbb{R}^N \setminus \{0\}$. If Ω is a bounded domain containing 0 there exists a positive solution u of (1.1) in $\Omega \setminus \{0\}$ satisfying

(1.39)-(i) and vanishing on $\partial\Omega$.

2- If $\frac{2p}{p+1} < q < p$ there exists a positive radial solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\begin{aligned} (i) \quad & \lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M \\ (ii) \quad & \lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_0. \end{aligned} \tag{1.40}$$

Moreover this solution is unique among all the positive solutions satisfying (1.40).

We also give conditions on p and q for the existence of a positive radial solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$, necessarily singular at 0, with a behaviour at 0 given by (1.26), (1.31) or (1.27), and an asymptotic behaviour at infinity given by (1.38).

The last section of the article is devoted to non radial results. We first give a general existence statement which allows to construct positive singular solutions of (1.1) in a punctured bounded domain with prescribed boundary value, provided there exists a radial singular solution in $\mathbb{R}^N \setminus \{0\}$. This singular solution has been obtained by the phase plane analysis of Section 2 in the case $q = \frac{2p}{p+1}$, and by the radial analysis of section 3 in the other cases.

Theorem 1.14 *Let $\Omega \subset B_R \subset \mathbb{R}^N$ be a bounded smooth domain containing 0, M a real number, $p > 1$ and $1 \leq q \leq 2$. If there exists a radial positive and decreasing function v defined in $B_R \setminus \{0\}$ and satisfying $\mathcal{L}_{p,q}^M v = 0$ in $\Omega \setminus \{0\}$ and*

$$\lim_{x \rightarrow 0} v(x) = \infty,$$

then for any nonnegative function $\phi \in W^{1,\infty}(\Omega)$, there exists a solution u of $\mathcal{L}_{p,q}^M u = 0$ in $\Omega \setminus \{0\}$ satisfying $u = \phi$ on $\partial\Omega$ and

$$\lim_{x \rightarrow 0} u(x) = \infty.$$

Furthermore there holds

$$(v(x) - \max_{z \in \partial\Omega} \phi(z))_+ \leq u(x) \leq v(x) + \max_{z \in \partial\Omega} (\phi(z) - v(z))_+ \quad \text{for all } x \in \Omega \setminus \{0\}. \tag{1.41}$$

A second key result deals with the uniqueness of positive solutions in $\mathbb{R}^N \setminus \{0\}$ or in a punctured bounded domain $\Omega \setminus \{0\}$ starshaped with respect to 0. Using a general scaling method we prove the following

Theorem 1.15 *Assume $N \geq 2$, $p > 1$, $1 < q < 2$ and $M > 0$. Let a such that*

$$\begin{aligned} (i) \quad & 0 \leq a < \beta \quad \text{if } q \leq \frac{2p}{p+1} \\ (ii) \quad & \beta < a \quad \text{if } q > \frac{2p}{p+1}. \end{aligned} \tag{1.42}$$

There exists at most one positive solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{x \rightarrow 0} |x|^\alpha |\ln |x||^{\tilde{a}} u(x) = \Lambda \tag{1.43}$$

where Λ is some positive constant and \tilde{a} is a real number. If Ω is a bounded domain containing 0 and starshaped with respect to 0 and $\phi \in C(\partial\Omega)$ is nonnegative, there is at most one positive solution u of (1.1) in $\Omega \setminus \{0\}$ satisfying (1.43) with value ϕ on $\partial\Omega$.

This result admits various extensions valid when the exponent a above is equal to β .

With the help of these results we characterize *all the local positive solutions* of (1.1), *not necessarily radial*, either near 0 or near ∞ . An important tool is the intensive use of the tangency property of graphs of global solutions which has been introduced in [15] for the studying of isolated singularities of p -harmonic functions.

Acknowledgements. This article has been prepared with the support of the FONDECYT grants 1210241 and 1190102 for the three authors.

2 The case $q = \frac{2p}{p+1}$

2.1 The equation on the sphere

The existence of particular solutions of (1.4), and eventually their uniqueness, plays a key role in the description of the behaviour of all the solutions. Due to the invariance of the equation under the transformations T_ℓ these natural particular solutions are the ones which are self-similar, i.e. invariant by these transformations. In spherical coordinates $(r, s) \in \mathbb{R}_+ \times S^{N-1}$, they endow the form $(r, s) \mapsto u(r, s) = r^{-\alpha}\omega(s)$, and ω is a solution of (1.4). Since we are dealing with nonnegative solutions, by the strong maximum principle they are either positive or identically zero. This fact does not depend on the sign of M .

2.1.1 Proof of Theorem 1.1: constant positive solutions

Assume $M \leq 0$ and ω is a nonnegative solution of (1.4). Multiplying the equation by ω and integrating over S^{N-1} yields

$$\int_{S^{N-1}} \left(|\nabla' \omega|^2 + \ell_{N,p} \omega^2 + \omega^{p+1} - M (\alpha^2 \omega^2 + |\nabla' \omega|^2)^{\frac{p}{p+1}} \omega \right) dS = 0.$$

Since $\ell_{N,p} \geq 0$ if and only if $p \geq \frac{N}{N-2}$, we infer the non-existence statement 1.

For any M , constant positive solutions are the positive roots of $\mathcal{P}_M(x) = 0$. If we set $z = x^{\frac{p-1}{p+1}}$, $\mathcal{P}_M(x) = 0$ is equivalent to $\tilde{\mathcal{P}}_M(z) = 0$ where

$$\tilde{\mathcal{P}}_M(z) = z^{p+1} - M \alpha^{\frac{2p}{p+1}} z + \ell_{N,p}.$$

Since $\tilde{\mathcal{P}}_M'(z) = (p+1)z^p - M \alpha^{\frac{2p}{p+1}}$ the minimum of $\tilde{\mathcal{P}}_M$ on \mathbb{R}_+ is achieved at $z = 0$ if $M \leq 0$, or at $z_0 = \left(\frac{M}{p+1}\right)^{\frac{1}{p}} \alpha^{\frac{2}{p+1}}$ if $M > 0$. In the first case the function $\tilde{\mathcal{P}}_M$ is increasing on $(0, \infty)$. It vanishes therein if and only if $\tilde{\mathcal{P}}_M(0) = \ell_{N,p} < 0$, or

equivalently $1 < p < \frac{N}{N-2}$. In the second case, $\tilde{\mathcal{P}}_M$ is decreasing on $(0, z_0)$ and increasing on (z_0, ∞) . Its minimal value is

$$\tilde{\mathcal{P}}_M(z_0) = \ell_{N,p} - p \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \alpha^2 = \frac{4p}{(p-1)^2} \left(\left(\frac{m^*}{p+1} \right)^{\frac{p+1}{p}} - \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \right). \quad (2.1)$$

If $p \leq \frac{N}{N-2}$, then $\tilde{\mathcal{P}}_M(z_0) < \tilde{\mathcal{P}}_M(0) = \ell_{N,p} \leq 0$, hence $\tilde{\mathcal{P}}_M$ admits a unique positive zero and so does \mathcal{P}_M . This gives the existence of x_M in case 2.

If $p > \frac{N}{N-2}$, then $\tilde{\mathcal{P}}_M(0) = \ell_{N,p} > 0$. We obtain the existence of constant solutions in 3, 4 and 5 according $M > m^*$, $M = m^*$ and $0 < M < m^*$. \square

2.1.2 Proof of Theorem 1.1: positive solutions

Let ω be a nonnegative solution of (1.4). By regularity it is C^2 and either positive or identically 0. If it is not constant, we denote by $\bar{\omega}$ and $\underline{\omega}$ respectively the maximum and the minimum of ω on S^{N-1} . There holds $\mathcal{P}_M(\bar{\omega}) \leq 0$ and $\mathcal{P}_M(\underline{\omega}) \geq 0$, and if we set $\bar{\omega}^{\frac{p-1}{p+1}} = \bar{z}$ and $\underline{\omega}^{\frac{p-1}{p+1}} = \underline{z}$, we have that $\tilde{\mathcal{P}}_M(\bar{z}) \leq 0$ and $\tilde{\mathcal{P}}_M(\underline{z}) \geq 0$.

1- First we consider the case where $M < 0$ and $1 < p < \frac{N}{N-2}$. Since $\tilde{\mathcal{P}}_M$ is increasing on \mathbb{R}_+ we deduce $\bar{\omega} = \underline{\omega} = \omega = x_M$.

2- Next we assume $M > 0$ and $p \leq \frac{N}{N-2}$, then $\tilde{\mathcal{P}}_M$ is increasing on (z_0, ∞) . Hence it is negative on $[0, x_M^{\frac{p-1}{p+1}})$ and positive on $(x_M^{\frac{p-1}{p+1}}, \infty)$. This implies $\bar{\omega} \leq x_M \leq \underline{\omega}$ and finally $\omega = x_M$.

3- If $p > \frac{N}{N-2}$ and $M = m^*$, \mathcal{P}_{m^*} is positive on $\mathbb{R}^+ \setminus \{x_{m^*}\}$. This implies $\bar{\omega} \leq x_{m^*} \leq \underline{\omega}$ and finally $\omega = x_{m^*}$.

4- If $p > \frac{N}{N-2}$ and $M < m^*$, \mathcal{P}_{m^*} is positive on \mathbb{R}^+ , hence there exists no positive solution.

5- Finally, if $p > \frac{N}{N-2}$ and $M > m^*$, \mathcal{P}_M is positive on $(0, x_{1,M}) \cup (x_{2,M}, \infty)$ and negative on $(x_{1,M}, x_{2,M})$. This implies (1.9). The proof of the second assertion is more involved. Set $z = |\nabla' \omega|^2$ and $y = \alpha^2 \omega^2 + z^2$. Then

$$\Delta' \omega = \alpha K \omega + \omega^p - M y^{\frac{p}{p+1}}.$$

By Weitzenböck's formula

$$\frac{1}{2} \Delta' z = |\text{Hess}(\omega)|^2 + \langle \nabla' \Delta' \omega, \nabla' \omega \rangle + \text{Ric}_g(\nabla' \omega, \nabla' \omega), \quad (2.2)$$

where $\text{Hess}(\omega)$ is the Hessian and Ric_g is the curvature 2-tensor on S^{N-1} . In that case we have that $\text{Ric}_g = (N-2)g$. By Schwarz inequality

$$|\text{Hess}(\omega)|^2 \geq \frac{1}{N-1} (\Delta' \omega)^2,$$

therefore, replacing $\Delta' \omega$ by its value, we obtain the inequality

$$-\frac{1}{2} \Delta' z + (N-2)z + \frac{1}{N-1} (\Delta' \omega)^2 + (\alpha K + p \omega^{p-1})z - \frac{Mp}{p+1} y^{-\frac{1}{p+1}} \langle \nabla' y, \nabla' \omega \rangle \leq 0.$$

Since $\nabla'z = \nabla'y - 2\alpha^2\omega\nabla'\omega$ we infer

$$-\frac{1}{2}\Delta'z + \left(N - 2 + \alpha K + p\omega^{p-1} - \frac{2Mp}{p+1}y^{-\frac{1}{p+1}}\omega\right)z + \frac{1}{N-1}(\Delta'\omega)^2 - \frac{Mp}{p+1}y^{-\frac{1}{p+1}}\langle\nabla'z, \nabla'\omega\rangle \leq 0.$$

Let $s_0 \in S^{N-1}$ where z is maximal. Then $\nabla'z(s_0) = 0$ and $\Delta'z(s_0) \leq 0$. Hence at $s = s_0$ there holds

$$\left(\alpha K + N - 2 + p\omega^{p-1} - \frac{2Mp\alpha^2\omega}{(p+1)(\alpha^2\omega^2 + z)^{\frac{1}{p+1}}}\right)z + \frac{1}{N-1}\left(\alpha K\omega + \omega^p - My^{\frac{p}{p+1}}\right)^2 \leq 0.$$

Therefore

$$\left(\alpha K + N - 2 + p\omega^{p-1} - \frac{2Mp\alpha^{\frac{2p}{p+1}}\omega^{\frac{p-1}{p+1}}}{p+1}\right)z + \frac{1}{N-1}\left(\alpha K\omega + \omega^p - M(\alpha^2\omega^2 + z)^{\frac{p}{p+1}}\right)^2 \leq 0. \quad (2.3)$$

Set

$$F(t) = pt^{p+1} - \frac{2Mp\alpha^{\frac{2p}{p+1}}}{p+1}t + N - 2 + \alpha K$$

and $t_0 = \omega^{\frac{p-1}{p+1}}(s_0)$. If ω is non-constant, $z(s_0) > 0$, hence $F(t_0) \leq 0$. . If $t_i = x_{i,M}^{\frac{p-1}{p+1}}$, for $i = 1, 2$, there holds

$$t_i^{p+1} - M\alpha^{\frac{2p}{p+1}}t_i + \alpha K = 0,$$

hence

$$F(t_i) = pt_i^{p+1} - \frac{2p}{p+1}\left(\alpha K + t_i^{p+1}\right) + N - 2 + \alpha K = \frac{p(p-1)}{p+1}t_i^{p+1} + N - 2 - \frac{2K}{p+1}.$$

Since

$$F'(t) = p(p+1)t^p - \frac{2Mp\alpha^{\frac{2p}{p+1}}}{p+1},$$

F is minimal for $t = t_* = \left(\frac{2M}{(p+1)^2}\right)^{\frac{1}{p}}\alpha^{\frac{2}{p+1}}$ and

$$F(t_*) = N - 2 + \alpha K - p^2\left(\frac{2M}{(p+1)^2}\right)^{\frac{p+1}{p}}\alpha^2 \leq F(t_0) \leq 0. \quad (2.4)$$

This implies

$$\frac{2M}{(p+1)^2}\alpha^{\frac{2p}{p+1}} \geq \left(\frac{N-2+\alpha K}{p^2}\right)^{\frac{p}{p+1}}, \quad (2.5)$$

and equivalently $M \geq \tilde{m}$ where \tilde{m} is defined in (1.10). Therefore, if $M < \tilde{m}$ there cannot exist non-constant positive solution. In order to prove (1.11) we first notice that if $p > \frac{N}{N-2}$, then

$$\frac{(N-2)p^2 - (N+2)}{(N-2)p - N} > p + 1.$$

Therefore

$$\frac{\tilde{m}}{m^*} > \frac{p+1}{2} \left(\frac{p+1}{2p} \right)^{\frac{p}{p+1}}.$$

By taking the logarithm it is easy to check that the function $p \mapsto \frac{p+1}{2} \left(\frac{p+1}{2p} \right)^{\frac{p}{p+1}}$ is increasing, hence the right-hand side of the previous inequality is minorized by

$$\theta_N := \frac{N-1}{N-2} \left(\frac{N-1}{N} \right)^{\frac{N}{2(N-1)}}, \quad (2.6)$$

which is the desired estimate. Notice that $\theta_3 \sim 1.47$ and θ_N decreases to 1 when $N \rightarrow \infty$. \square

Remark. The following monotonicity properties of the points P_M are straightforward: in cases (i) and (ii) x_M is increasing with M . In case (iii) $M \mapsto x_{1,M}$ is decreasing while $M \mapsto x_{2,M}$ is increasing. Furthermore, if $M' > M > m^*$,

$$x_{1,M'} < x_{1,M} < x_{m^*} < x_{2,M} < x_{2,M'}. \quad (2.7)$$

The value of x_{m^*} is explicit

$$x_{m^*} = \left(\frac{2K}{p(p-1)} \right)^{\frac{1}{p-1}} = \left(\frac{\alpha K}{p} \right)^{\frac{1}{p-1}}. \quad (2.8)$$

We end this section by proving a result dealing with bifurcation from constant solutions.

Theorem 2.1 *When $M > m^*$ the solution $x_{j,M}$, $j = 1, 2$, is never a bifurcation point in the sense that the linearized equation at this point is singular.*

Proof. If we look for solutions of (1.4) under the form $\omega = x_{j,M} + \epsilon \phi_k$ where ϕ_k is an eigenfunction of $-\Delta'$ in $H^1(S^{N-1})$ associated to the eigenvalue $\lambda_k = k(N-2+k)$, we obtain that

$$\lambda_k + \alpha K + p x_{j,M}^{p-1} - \frac{2p}{p+1} \alpha^{\frac{2p}{p+1}} M x_{j,M}^{\frac{p-1}{p+1}} = 0 \quad (2.9)$$

We recall that \mathcal{P}_M is defined in (1.7). If $\mathcal{Q}_M(x) = x \mathcal{P}_M(x)$, then (2.9) is equivalent to

$$\lambda_k + x_{j,M} \mathcal{P}'_M(x_{j,M}) = 0. \quad (2.10)$$

We know that $\mathcal{P}'_M(x_{2,M}) > 0$, then for any $k \in \mathbb{N}^*$ identity (2.9) is impossible with $j = 2$. Concerning the case $j = 1$, (2.9) combined with $\mathcal{P}_M(x_{1,M}) = 0$ and the value of α yields

$$\lambda_k - 2K + \frac{p(p-1)}{p+1} \alpha^{\frac{2p}{p+1}} M x_{1,M}^{\frac{p-1}{p+1}} = 0, \quad (2.11)$$

which never occurs when $k \geq 2$ because of the values of λ_k and K . When $k = 1$, since $\mathcal{P}_M(x_{1,M}) = 0$ there holds

$$M\alpha^{\frac{2p}{p+1}}x_{1,M}^{\frac{p-1}{p+1}} > \alpha K$$

thus (2.11) yields

$$\lambda_1 - 2K + \frac{p(p-1)}{p+1}\alpha K < 0.$$

Because $\lambda_1 = N - 1$ and $\alpha(p-1) = 2$ we obtain $N - 1 - 2K + \frac{2p}{p+1}K < 0$, which is equivalent to $(p-1)(N-1) + 2 + 2\alpha < 0$, a contradiction. \square

2.2 Radial solutions

In this section we study in detail the nonnegative solutions of the ordinary differential equation

$$-u'' - \frac{N-1}{r}u' + |u|^{p-1}u - M|u'|^{\frac{2p}{p+1}} = 0, \quad (2.12)$$

when $p > 1$. Because of the scaling invariance (1.3) the equation can be transformed into an autonomous equation by setting

$$u(r) = r^{-\alpha}x(t), \quad t = \ln r. \quad (2.13)$$

Then $x(t)$ satisfies

$$x_{tt} + Lx_t - \alpha Kx - |x|^{p-1}x + M|\alpha x - x_t|^{\frac{2p}{p+1}} = 0, \quad (2.14)$$

where we recall that $K = \frac{(N-2)p-N}{p-1} = N - 2 - \alpha$ and where we set

$$L = \frac{(N-2)p - (N+2)}{p-1} = K - \alpha. \quad (2.15)$$

If we set $u'(r) = -r^{-(\alpha+1)}y(t)$, then (2.12) is equivalent to

$$\begin{aligned} x_t &= \alpha x - y \\ y_t &= -Ky - |x|^{p-1}x + M|y|^{\frac{2p}{p+1}}. \end{aligned} \quad (2.16)$$

Since we are interested in positive u we restrict to solutions of (2.16) in the half-space $\mathbb{R}_+^2 = \{(x, y) \in \mathbb{R}^2 : x > 0\} = Q_1 \cup Q_4$ where

$$Q_1 = \{(x, y) \in \mathbb{R}_+^2 : y > 0\}$$

is the first quadrant and

$$Q_4 = \{(x, y) \in \mathbb{R}_+^2 : y < 0\}$$

is the fourth quadrant. The regular solutions of (2.12) (with $u(0) = u_0 > 0$ and $u'(0) = 0$) are increasing near $r = 0$, so their trajectory $\mathcal{T}_1 := \{(x(t), y(t))\}$ lies in

Q_4 as $t \rightarrow -\infty$. The solutions defined in a neighborhood of $r = 0$ and unbounded near 0 are decreasing, so their trajectory lie in Q_1 as $t \rightarrow -\infty$. The solutions defined near $r = \infty$ are decreasing, so their trajectory remain in Q_1 as $t \rightarrow \infty$.

Theorem 1.1 can be reformulated in the following way:

- 1- If $M \leq 0$ and $1 < p < \frac{N}{N-2}$, the only non-trivial equilibrium in $\overline{\mathbb{R}}_+^2$, is $P_M = (x_M, \alpha x_M)$. If $p \geq \frac{N}{N-2}$ there exists no non-trivial equilibrium in this region.
- 2- If $1 < p \leq \frac{N}{N-2}$ and $M > 0$, the only non-trivial equilibrium in $\overline{\mathbb{R}}_+^2$ is $P_M = (x_M, \alpha x_M)$.
- 3- If $p \geq \frac{N}{N-2}$ and $M > m^*$, there exist two non-trivial equilibria in $\overline{\mathbb{R}}_+^2$, $P_{1,M} = (x_{1,M}, \alpha x_{1,M})$ and $P_{2,M} = (x_{2,M}, \alpha x_{2,M})$.
- 4- If $p \geq \frac{N}{N-2}$ and $M = m^*$, there exists one non-trivial equilibria in $\overline{\mathbb{R}}_+^2$, $P_{m^*} = (x_{m^*}, \alpha x_{m^*})$.
- 5- If $p \geq \frac{N}{N-2}$ and $0 < M < m^*$ there exists no non-trivial equilibrium in $\overline{\mathbb{R}}_+^2$.

We also recall the classical result concerning regular solutions, not only in the case $q = \frac{2p}{p+1}$.

Proposition 2.2 *Let $N \geq 1$, $1 < q < p$ and $M \geq 0$. Then for any $a > 0$ there exists a unique radial maximal positive solution u of (1.1) satisfying $u(0) = a$, $u'(0) = 0$. This solution denoted by $u_{[a]}$ is defined in B_R , where $R = R_a > 0$, and it satisfies $\lim_{|x| \uparrow R} u_{[a]} = \infty$.*

Proof. In the case $M = 0$ the result is a standard combination of Cauchy-Lipschitz theorem with the Keller-Osserman estimate. In the case $M > 0$ the proof can be obtained in a slightly similar way using also Proposition A.1. See also [2], [24] and [1] for many extensions concerning these regular (or large) solutions. \square

2.2.1 Linearisation at $(0, 0)$

The linearization at $(0, 0)$ is given by the system

$$\begin{aligned} x_t &= \alpha x - y \\ y_t &= -Ky \end{aligned} \tag{2.17}$$

with eigenvalues $\lambda_1 = -K$, $\lambda_2 = \alpha$ and corresponding eigenvectors $\xi_1 = (1, N-2)$ and $\xi_2 = (1, 0)$ if $K \neq -\alpha$ or equivalently $N \neq 2$. If $N = 2$, $\lambda_1 = \lambda_2 = \alpha$, the only eigenspace is $\text{span}(\xi_2)$. In any cases $\lambda_2 - \lambda_1 = N - 2$. There exists one trajectory located in Q_4 of the linearized system converging to 0 when $t \rightarrow -\infty$. To this trajectory is associated a trajectory \mathcal{T}_r of (2.16) such that $(x(t), y(t)) \approx c(e^{\alpha t}, -\frac{1}{N}e^{(\alpha+1)t})$ when $t \rightarrow -\infty$. These solutions are associated to the one parameter family of regular solutions mentioned above with $u(0) = u_0$ and $u'(0) = 0$.

(i) Assume first that $N \geq 3$.

If $p < \frac{N}{N-2}$ then $K < 0$ and $(0, 0)$ is a source. Then all trajectories of (2.17) defined

in a neighborhood of $(0, 0)$ converge to this point when $t \rightarrow -\infty$. Besides the trajectory \mathcal{T}_r , all the other trajectories converging to zero when $t \rightarrow -\infty$ start in Q_1 with initial slope $N - 2$. They satisfy $x(t) \sim ae^{-Kt}$ for some $a > 0$ by Lemma A.4. This means that $r^{N-2}u(r) \rightarrow a$ when $r \rightarrow 0$.

If $p > \frac{N}{N-2}$, then $K > 0$ and $(0, 0)$ is a saddle point. The trajectory \mathcal{T}_r converges to $(0, 0)$ at $-\infty$. There is also the unique trajectory \mathcal{T}_s which converges to $(0, 0)$ when $t \rightarrow \infty$. Their slope at $(0, 0)$ is $N - 2$ and they correspond to solutions $u(r) \sim cr^{2-N}$ when $r \rightarrow \infty$.

If $p = \frac{N}{N-2}$, then $K = 0$. Besides the regular trajectory which always exists, there exists an invariant trajectory passing through $(0, 0)$, with slope $N - 2$, by the theorem of the central manifold. We will see later on that it converges to $(0, 0)$ as $t \rightarrow -\infty$.

(ii) Assume now that $N = 1$ or 2 there still exists the regular trajectory \mathcal{T}_r .

If $N = 1$, then $\lambda_1 = \frac{p+1}{p-1} > \lambda_2$. There exist infinitely many trajectories different from \mathcal{T}_r , converging to $(0, 0)$ at $-\infty$, in Q_1 or Q_4 , corresponding to solutions such that $u(0) = u_0 > 0$ and $u'(0) = a \in \mathbb{R} \setminus \{0\}$. There exists one trajectory converging to $(0, 0)$ at $-\infty$ with slope -1 and located in Q_4 . It corresponds to solutions such that $u(0) = 0$ and $u'(0) > 0$.

If $N = 2$, then $\lambda_1 = \lambda_2 = \alpha$. The point $(0, 0)$ is a degenerate node. All the trajectories in a neighborhood of $(0, 0)$ tend to $(0, 0)$ when $t \rightarrow -\infty$ and are tangent to ξ_1 . However they behave like $c(-t)^{-1}$ for any $c > 0$. They correspond to solutions u such that $\lim_{r \rightarrow 0} (-\ln r)^{-1}u(r) = a > 0$.

2.2.2 Linearisation at the non-trivial equilibrium points

Lemma 2.3 1- If $1 < p \leq \frac{N}{N-2}$ and $M > 0$, or $1 < p < \frac{N}{N-2}$ and $M = 0$, P_M is a saddle point.

2- If $p > \frac{N}{N-2}$ and $M > m^*$, $P_{1,M}$ is a node point and a source and $P_{2,M}$ is a saddle point.

3- If $p > \frac{N}{N-2}$ and $M = m^*$, P_{m^*} is not hyperbolic. One eigenvalue is $N - 2$ and the other is 0 .

Proof. Set $y_M = \alpha x_M$. In view of (1.7), y_M satisfies

$$\alpha^{-p}y_M^{p-1} - My_M^{\frac{p-1}{p+1}} + K = 0. \quad (2.18)$$

Setting $x = x_M + \bar{x}$, $y = y_M + \bar{y}$, the linearized equation at (x_M, y_M) is

$$\begin{aligned} \bar{x}_t &= \alpha\bar{x} - \bar{y} \\ \bar{y}_t &= -px_M^{p-1}\bar{x} + \left(\frac{2Mp}{p+1}y_M^{\frac{p-1}{p+1}} - K \right)\bar{y}. \end{aligned} \quad (2.19)$$

The characteristic polynomial of the corresponding matrix is

$$T_{y_M}(X) = X^2 - \left(\frac{2Mp}{p+1}y_M^{\frac{p-1}{p+1}} - L \right)X + 2K - \frac{2Mp}{p+1}y_M^{\frac{p-1}{p+1}}. \quad (2.20)$$

1- If $p \leq \frac{N}{N-2}$ and $M > 0$, or $p < \frac{N}{N-2}$ and $M = 0$, P_M is unique. Since either $K \leq 0$ and $M > 0$ or $K < 0$ and $M = 0$, the product of the roots is negative. Hence P_M is a saddle point.

2-3- Next we assume $N \geq 3$ and $M \geq m^*$. The sum of the roots of $T_{y_M}(X)$ is equal to

$$\frac{2Mp}{p+1} y_M^{\frac{p-1}{p+1}} - L = \frac{2p}{p+1} (\alpha^{-p} y_M^{p-1} + K) - L > \frac{p-1}{p+1} K + \alpha > 0.$$

Concerning the product $\Pi(y_M)$ of the roots, we deduce from (2.3) that $T_{y_{m^*}}(0) = 0$ hence $\Pi(y_{m^*}) = 0$. Since by (2.18)

$$\Pi(y_M) = 2K - \frac{2Mp}{p+1} y_M^{\frac{p-1}{p+1}} = \frac{2K}{p+1} - \frac{2p}{p+1} \alpha^{-p} y_M^{p-1},$$

we infer that for $M > m^*$,

$$\Pi(y_{2,M}) < \Pi(y_{m^*}) = 0 < \Pi(y_{1,M}). \quad (2.21)$$

Hence $P_{2,M}$ is a saddle point and $P_{1,M}$ is a source. In order to characterize the nature of this source we denote by $D(T_{y_M})$ the discriminant of T_{y_M} . Then

$$\begin{aligned} D(T_{y_{1,M}}) &= \left(\frac{2Mp}{p+1} y_{1,M}^{\frac{p-1}{p+1}} - L \right)^2 + 4 \left(\frac{2Mp}{p+1} y_{1,M}^{\frac{p-1}{p+1}} - 2K \right) \\ &= \left(\frac{2Mp}{p+1} y_{1,M}^{\frac{p-1}{p+1}} - L + 2 + 2\sqrt{N-1} \right) \left(\frac{2Mp}{p+1} y_{1,M}^{\frac{p-1}{p+1}} - L + 2 - 2\sqrt{N-1} \right). \end{aligned}$$

By (2.18), $M y_{1,M}^{\frac{p-1}{p+1}} > K$, hence

$$\begin{aligned} \frac{2Mp}{p+1} y_{1,M}^{\frac{p-1}{p+1}} - L + 2 - 2\sqrt{N-1} &> \frac{p-1}{p+1} K + \frac{2}{p-1} + 2 - 2\sqrt{N-1} \\ &= \frac{N - 2\sqrt{N-1}}{p^2 - 1} \left(p - \frac{N-2}{N - 2\sqrt{N-1}} \right)^2. \end{aligned}$$

Hence $D(T_{y_{1,M}}) > 0$ which implies that the roots are real and $P_{1,M}$ is a node.

If $M = m^*$ the product of the roots is 0, hence one root is 0. Since their sum is equal to $\frac{2p}{p+1} m^* y_{m^*}^{\frac{p-1}{p+1}} - L = N - 2$, the nonzero root is equal to $N - 2$. \square

2.2.3 The vanishing curves of the vector field

The vector field associated to (2.16) is defined by

$$(x, y) \mapsto H(x, y) = (H_1(x, y), H_2(x, y)) := \left(\alpha x - y, -Ky - x^p + M|y|^{\frac{2p}{p+1}} \right). \quad (2.22)$$

We call vanishing curves of H in \mathbb{R}_+^2 the set of points where H_1 or H_2 vanishes.

$$\mathcal{L} = \{ (x, y) \in \mathbb{R}_+^2 : y = \alpha x \},$$

and

$$\mathcal{C} = \{(x, y) \in \mathbb{R}_+^2 : H_2(x, y) = 0\} = \mathcal{C}_1 \cup \mathcal{C}_4,$$

where

$$\mathcal{C}_1 = \left\{ (x, y) \in Q_1 : x^p = My^{\frac{2p}{p+1}} - Ky := \Phi(y) \right\},$$

and

$$\mathcal{C}_4 = \left\{ (x, y) \in Q_4 : x^p = M|y|^{\frac{2p}{p+1}} - Ky := \Psi(y) \right\}.$$

Those vanishing curves are the boundary of some semi-invariant regions in \mathbb{R}_+^2 . Their configuration depends on the intersection of these curves.

I- If $K \leq 0$ and $M > 0$ we denote by

(A) is the set of points $(x, y) \in Q_1$ such that $y > \max\{\alpha x, \Phi^{-1}(x)\}$.

(B) is the set of points $(x, y) \in Q_1$ such that $x \geq x_M$ and $\alpha x < y < \Phi^{-1}(x)$.

(C) is the union of the set of points $(x, y) \in Q_1$ such that $y < \min\{\alpha x, \Phi^{-1}(x)\}$ and the set of points $(x, y) \in Q_4$ such that $x > \Psi(y)$.

(D) is the set of points $(x, y) \in Q_1$ such that $\Phi^{-1}(x) < y < \alpha x$.

(E) is the set of points $(x, y) \in Q_4$ such that $x < \Psi(y)$.

II- If $K > 0$ and $M > m^*$ we denote by

(A) is the set of points $(x, y) \in Q_1$ such that $y > \max\{\alpha x, \Phi^{-1}(x)\}$.

(B) is the set of points $(x, y) \in Q_1$ such that $x \geq x_{2,M}$ and $\alpha x < y < \Phi^{-1}(x)$.

(C) is the union of the set of points $(x, y) \in Q_1$ such that $x > \Phi(y)$ and the set of points $(x, y) \in Q_4$ such that $x > \Psi(y)$.

(D) is the set of points $(x, y) \in Q_1$ such that $\Phi^{-1}(x) < y < \alpha x$.

(E) is the set of points $(x, y) \in Q_4$ such that $x < \Psi(y)$.

(F) is the set of points $(x, y) \in Q_1$ such that $x \leq x_{1,M}$ and $\alpha x < y < \Phi^{-1}(x)$.

III- If $K > 0$ and $M = m^*$, (D) is empty.

IV- If $K > 0$ and $0 < M < m^*$, (D) is empty and (B) and (F) are replaced by the set $(\tilde{B}) = \{(x, y) \in Q_1 \text{ such that } \alpha x < y < \Phi^{-1}(x)\}$ (note that \tilde{B} is connected).

We present below some graphics of the vector field H associated to system (2.16). We show the vanishing curves of the vector field H as well as the direction of the vector field along these curves.




Figure 1: $M > 0, K < 0 \iff p < \frac{N}{N-2}$.




Figure 2: $M > 0, K = 0 \iff p = \frac{N}{N-2}$.




Figure 3: $M > m^*, K > 0 \iff p > \frac{N}{N-2}$.




Figure 4: $M = m^*, K > 0 \iff p > \frac{N}{N-2}$.




Figure 5: $0 < M < m^*$, $K > 0 \iff p > \frac{N}{N-2}$.

2.3 Description of the radial solutions defined near 0

In this section we use the dynamical system (2.19) to describe all the positive solutions of (2.12) defined on a maximal interval $(0, R)$, $R \leq \infty$. The case $M = 0$ which is well-known will be used as a comparison model.

2.3.1 The case $1 < p < \frac{N}{N-2}$ and $M \geq 0$

In this range of exponents the fixed point P_M is unique, the problem is more rigid and some of our existence and uniqueness results hold without the assumption of radiality as shown in Theorem 4.6.

Theorem 2.4 *Let $N = 1, 2$ and $p > 1$ or $N \geq 3$ and $p < \frac{N}{N-2}$, and $M > 0$.*

1- *The function u_{x_M} is the unique positive solution of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying*

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_M. \quad (2.23)$$

2- *For any $k > 0$ there exists a unique positive solution $u = u_k$ of (1.2) in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.12). Furthermore u_k is radial and*

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u_k(x) = x_M. \quad (2.24)$$

To this set of solutions u_k is associated a unique heteroclinic orbit \mathcal{T}_1 of the system (2.16) connecting the origin when $t \rightarrow -\infty$ to P_M when $t \rightarrow \infty$.

3- *For any $R > 0$ there exists a unique positive solution of (1.2) in $\mathbb{R}^N \setminus \overline{B_R}$ satisfying (2.24) and $\lim_{|x| \rightarrow R} u(x) = \infty$. This solution is radial.*

4- For any $R > 0$ there exists a unique positive solution of (1.2) in $B_R \setminus \{0\}$ satisfying

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_M \quad \text{and} \quad \lim_{|x| \rightarrow R} u(x) = \infty, \quad (2.25)$$

and a unique positive solution satisfying

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_M \quad \text{and} \quad \lim_{|x| \rightarrow R} u(x) = 0. \quad (2.26)$$

Moreover these solutions are radial.

5- Assume $N > 2$. For any $k > 0$ there exists $R_k > 0$ and a unique radial positive solution of (1.2) in $B_{R_k} \setminus \{0\}$ satisfying (1.12) and vanishing on $\partial B_{B_{R_k}}$ or such that

$$\lim_{|x| \rightarrow R_k} u(x) = \infty. \quad (2.27)$$

Furthermore the mapping $k \rightarrow R_k$ is decreasing from $(0, \infty)$ onto $(0, \infty)$.

Proof. 1- All the uniqueness results, which are valid not only for radial solutions, are proved in Section 4, in particular in Theorem 4.6.

2- In the phase plane (x, y) , recall that $(0, 0)$ is a source equilibrium and there exist infinitely many trajectories different from the regular one \mathcal{T}_r , converging to $(0, 0)$ when $t \rightarrow -\infty$, with the initial slope $N - 2 < \alpha$ since $p < \frac{N}{N-2}$, so they start from $(0, 0)$ in Region (D) of Figure 1. The point P_M is a saddle point with eigenvalues $\lambda < 0 < \tilde{\lambda}$ and associated eigenvectors

$$\xi_1 = (1, \alpha + |\lambda|) \quad \text{and} \quad \xi_2 = (1, \alpha - \tilde{\lambda}).$$

We denote by \mathcal{T}_1 the trajectory such that $x(t)$ increases and converges to x_M when $t \rightarrow \infty$, and by \mathcal{T}_2 the trajectory such that $x(t)$ decreases and converges to x_M when $t \rightarrow \infty$. Their common slope is larger than α , then \mathcal{T}_1 lies in the region (D) and \mathcal{T}_2 lies in the region (B) when $t \rightarrow \infty$. Because (D) is negatively invariant and bounded, \mathcal{T}_1 is contained in (D), a region in which $x(t)$ and $y(t)$ are monotone. Hence \mathcal{T}_1 converges to a fixed point in (\overline{D}) which is necessarily $(0, 0y)$ as $t \rightarrow -\infty$. Therefore \mathcal{T}_1 is an heteroclinic orbit joining $(0, 0)$ to P_M , and it is necessarily unique since P_M is a saddle point. Its slope at $(0, 0)$ is $N - 2$. It corresponds to a solution u_k of (2.12) satisfying (1.12)-(i). This solution u_k is unique by Theorem 4.6. Furthermore if $k < k'$ then $u_k \leq u_{k'}$. We also notice that for any $\ell > 0$ and $x \in \mathbb{R}^N \setminus \{0\}$,

$$T_\ell[u_k](r) := \ell^\alpha u_k(\ell r) = u_{k\ell^{\alpha+2-N}}(r).$$

If we denote by u_∞ the limit of the increasing sequence $\{u_k\}$, then

$$T_\ell[u_\infty](r) = u_\infty(r)$$

This implies that u_∞ is a self-similar solution of (2.12), hence $u_\infty = u_{x_M}$ and (2.24) holds.

3- The trajectory \mathcal{T}_2 converges to P_M when $t \rightarrow \infty$ and remains in the region (B) which is negatively invariant. If \mathcal{T}_2 were defined on whole \mathbb{R} , it would imply that it remains bounded because of the a priori estimate Proposition A.1. But in the region (B) the two functions $x(t)$ and $y(t)$ are decreasing. Hence the trajectory would converge to an equilibrium in the closure of (B) different from P_M , which is impossible. Therefore the two functions $x(t)$ and $y(t)$ with image \mathcal{T}_2 are defined on some maximal interval (T, ∞) and if we set $R = e^T$, the corresponding solution u of (2.12) satisfies $\lim_{r \downarrow R} u(r) = \infty$. Uniqueness of a solution defined on (R, ∞) and blowing-up at $r = R$ follows from Theorem 4.5. By the scaling T_ℓ , the function u is transformed in a solution of (2.12) which blows-up at $r = \ell^{-1}R$ and which is associated with the same trajectory \mathcal{T}_2 . Hence R can be any positive real number.

4- There exist two unstable trajectories \mathcal{T}_3 and \mathcal{T}_4 converging to P_M when $t \rightarrow -\infty$. They are associated to the eigenvalue $\tilde{\lambda}$, and their slope at P_M is $\alpha - \tilde{\lambda}$. We denote by \mathcal{T}_4 the trajectory which enters in the region (C) . Since this region is positively invariant, \mathcal{T}_4 remains in it. Then either its components are defined on some maximal interval $(-\infty, T)$ and the corresponding solution u of (2.12) tends to ∞ when $r \uparrow R := e^T$, or they are defined on whole \mathbb{R} . In that case u would coincide with u_{x_M} by 1- which is contradictory. Hence u is defined on the maximal interval $(0, R)$. Notice also that u is decreasing on some interval $(0, r_0)$ and increasing on (r_0, R) by the phase plane analysis. Thanks to the scaling T_ℓ , R can be taken arbitrarily. Since this solution is uniquely determined by \mathcal{T}_4 , it is unique. This corresponds to a uniqueness result for solutions of (1.1) in the class of radial solutions. This proves (2.25)

Consider now the trajectory \mathcal{T}_3 . It belongs to region (A) when $t \rightarrow -\infty$, and in this region $x_t < 0$ and $y_t > 0$. Since P_M is the only equilibrium in the quadrant Q_1 the trajectory intersects the straight line $x = 0, y > 0$ at some $y_0 > y_M$ for some $t = T$. Hence the corresponding solution u vanishes for $r = R = e^T$. Furthermore R can be taken arbitrarily. This proves (2.26).

5- Since $(0, 0)$ is a source, there exists $\epsilon > 0$ such that any backward trajectory issued from $(x_0, y_0) \in B_\epsilon(0)$ converges to $(0, 0)$ when $t \rightarrow -\infty$. All these trajectories in the first quadrant Q_1 have initial slope $N - 2$. If $(x_0, y_0) \in B_\epsilon(0) \cap (D)$ is above the heteroclinic orbit \mathcal{T}_1 , it cannot converge to P_M , then it crosses \mathcal{L} , enters in (A) and crosses the axis $\{x = 0, y > 0\}$ for some T . By Appendix A2 the associated solution u of (2.12) satisfies (1.12) for some $k > 0$ and $u(e^T) = 0$. If $(x_0, y_0) \in B_\epsilon(0) \cap (D)$ is below \mathcal{T}_1 , it enters the region (C) which is positively invariant and for the same reasons as in Step 3 it blows-up for some $t = T$. The corresponding solution u satisfies (1.12) for some $k > 0$ and blows up for $r = e^T := R$. Uniqueness of this type of solutions in B_R follows either from the general result Theorem 4.6 or from the uniqueness of the trajectories of the system (W8). The correspondance $k \mapsto R$ is decreasing and onto from $(0, \infty)$ to $(0, \infty)$ by uniqueness and using the scaling transformation T_ℓ . \square

Remark. It follows from the analysis of the phase plane that all the positive radial solutions of (1.2) defined in a neighborhood of $x = 0$ or in the complement of a ball

have their behaviour described by 1 or 2.

Proof of Theorem 1.3. It is a direct consequence of Theorem 2.4 and Theorem 4.6.

2.3.2 The case $p = \frac{N}{N-2}$, $q = \frac{2p}{p+1} = \frac{N}{N-1}$ and $M > 0$

When $p = \frac{N}{N-2}$ and $M = 0$ the isolated singularities of solutions of (1.1) are removable and the behaviour at infinity of these solutions is described in [21]. When $M > 0$ it is no longer the case and the interaction of the two reaction terms yields new phenomena. The next result covers Theorem 1.4, up to uniqueness which will follow from Theorem 4.6.

Theorem 2.5 *Let $N \geq 3$, $p = \frac{N}{N-2}$ and $M > 0$.*

1- *If $M = 0$ any isolated singularity of a solution, not necessarily radial neither nonnegative, of $-\Delta u + |u|^{p-1}u = 0$ is removable. If u is any solution of this equation in B_R^c , there exists λ such that*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} (\ln |x|)^{\frac{N-2}{2}} u(x) = \lambda, \quad (2.28)$$

and λ can only take the three values $\left(\frac{N-2}{\sqrt{2}}\right)^{N-2}$, $-\left(\frac{N-2}{\sqrt{2}}\right)^{N-2}$ and 0.

2- *If $M > 0$, the function $u_s(x) = \left((N-2)M^{\frac{N}{N-1}}\right)^{N-2} |x|^{2-N}$ is the unique positive separable solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$. There exists a positive radial solution u , unique up to the scaling transformations T_ℓ , satisfying*

$$\begin{aligned} (i) \quad & \lim_{r \rightarrow 0} r^{N-2} |\ln r|^{N-1} u(r) = \frac{((N-1)M)^{1-N}}{N-2} \\ (ii) \quad & \lim_{r \rightarrow \infty} r^{N-2} u(r) = \left((N-2)M^{\frac{N}{N-1}}\right)^{N-2}. \end{aligned} \quad (2.29)$$

Furthermore, for any $R > 0$ there exists a positive and radial solution $u := u_R$ of (1.1) in $B_R \setminus \{0\}$ satisfying

$$\lim_{r \rightarrow 0} r^{N-2} u(r) = \left((N-2)M^{\frac{N}{N-1}}\right)^{N-2}, \quad (2.30)$$

and

$$\lim_{|x| \rightarrow R} u(x) = \infty. \quad (2.31)$$

Finally, there exists also a unique positive solution $u := \tilde{u}_R$ of (1.1) in $B_R \setminus \{0\}$ satisfying (2.29)-(i) and (2.31) or (2.29)-(i) and $u = 0$ on ∂B_R . In both cases the solution is radial.

Proof. The results of assertion 1 is proved in [21], [2].

Assertion 2- Since $p = \frac{N}{N-2}$, $K = 0$. As a consequence, the vanishing curve \mathcal{C}_4 goes through $(0, 0)$. Lemma 2.3 is still valid. The point $P_M = (x_M, y_M)$ is a saddle

point and with $x_M = (N-2)^{N-2} M^{\frac{N(N-2)}{N-1}}$ and $y_M = (N-2)^{N-1} M^{\frac{N(N-2)}{N-1}}$. The stable curve \mathcal{T}_1 is an heteroclinic orbit staying in the region (D) and connecting $(0,0)$ to P_M . The point $(0,0)$ is no longer hyperbolic since the characteristic values are $\lambda_1 = 0$ and $\lambda_2 = N-2$, and the behaviour of the solutions in its neighbourhood is more delicate. The vector $\xi_2 = (1,0)$ is the eigenvector associated to the nonzero eigenvalue $N-2$ and the unstable curve corresponds to the regular solutions $u_{[a]}$. By the central manifold theorem, the curve \mathcal{T}_1 is the central manifold of $(0,0)$ and is tangent at this point to the eigenvector $\xi_1 = (1, N-2)$. Therefore $\lim_{t \rightarrow -\infty} \frac{y(t)}{x(t)} = N-2$ on \mathcal{T}_1 . As a consequence

$$\frac{(u(r))^{\frac{N}{N-2}}}{|u'(r)|^{\frac{N}{N-1}}} = \frac{(x(t))^{\frac{N}{N-2}}}{(y(t))^{\frac{N}{N-1}}} \leq c(y(t))^{\frac{N}{(N-1)(N-2)}} \quad \text{for } r \leq \epsilon_0.$$

Consequently $u^{\frac{N}{N-2}} = o\left(|u'|^{\frac{N}{N-1}}\right)$ in a neighborhood of $r = 0$. Therefore, for any $\epsilon > 0$ there exists $r_\epsilon > 0$ such that

$$(1-\epsilon)Mr^{N-1}|u'|^{\frac{N}{N-1}} \leq -(r^{N-1}u)' \leq Mr^{N-1}|u'|^{\frac{N}{N-1}} \quad \text{on } (0, r_\epsilon]. \quad (2.32)$$

Putting $W = r^{N-1}|u'|$ these inequalities become completely integrable and we derive

$$(N-1)(W(r))^{-\frac{1}{N-1}} = M|\ln r|^{-1}(1+o(1)) \quad \text{as } r \rightarrow 0. \quad (2.33)$$

Integrating (2.33) implies (2.29)-(i). Uniqueness among the radial solutions follows from the uniqueness of the stable heteroclinic orbit. As in Theorem 2.4 the unstable trajectory of P_M entering (C) intersect the axis $y = 0$ at some $P_0 = (x_0, 0)$ and any corresponding solution u is defined in some $B_R, R_i 0$, and it blows-up when $|x| \uparrow R$. Using the transformation T_ℓ , $T_\ell[u]$ is a solution defined in the ball $B_{\frac{R}{\ell}}$ which blows-up for $|x| = \frac{R}{\ell}$ and still satisfies (2.30). The backward trajectory $\mathcal{T}_{[P]}^-$ of any point $P = (P, 0)$ on the segment $(0, P_0)$ converges to $(0, 0)$ when $t \rightarrow -\infty$. For the same reason as for \mathcal{T}_1 the corresponding solution u satisfies (2.29)-(i) and it blows-up ifor $r = R$ for some $R > 0$. Since the scaling transformation T_ℓ leaves (2.29)-(i) unchanged, $\frac{R}{\ell}$ can take any value, this ends the proof. \square

Remark.

2.3.3 The case $p > \frac{N}{N-2}$ and $M > 0$

In the range of exponent $p > \frac{N}{N-2}$, the positive parameter m^* defined by (1.8) plays a fundamental role. The following result covers Theorem 1.5 and describes all the positive solutions of (2.12) either defined near ∞ or near 0.

Theorem 2.6 *Let $N > 2$ and $p > \frac{N}{N-2}$.*

1- *If $M > m^*$, then $u_{x_{1,M}}$ and $u_{x_{2,M}}$ are the two self-similar solutions. Moreover (i) there exists a unique, up to the transformation T_ℓ , positive radial solution $u = u_{1,2}$ defined in $\mathbb{R}^N \setminus \{0\}$ satisfying*

$$\lim_{r \rightarrow 0} r^\alpha u(r) = x_{1,M} \quad \text{and} \quad \lim_{r \rightarrow \infty} r^\alpha u(r) = x_{2,M}. \quad (2.34)$$

(ii) For any $k > 0$ there exists a unique positive radial $u = u_{1,k}$ defined in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{r \rightarrow 0} r^\alpha u(r) = x_{1,M} \quad \text{and} \quad \lim_{r \rightarrow \infty} r^{N-2} u(r) = k. \quad (2.35)$$

(iii) For any $R > 0$ there exists a positive radial solution $u = u_{j,R}$ in $B_R \setminus \{0\}$ with $j=1,2$, satisfying

$$\lim_{r \rightarrow 0} r^\alpha u(r) = x_{j,M} \quad \text{and} \quad \lim_{r \uparrow R} u(r) = \infty. \quad (2.36)$$

This solution is unique if $j = 2$. There exists also a unique radial positive solution $\tilde{u} = \tilde{u}_{2,R}$ in $\mathbb{R}^N \setminus B_R$ satisfying

$$\lim_{r \rightarrow \infty} r^\alpha \tilde{u}(r) = x_{2,M} \quad \text{and} \quad \lim_{r \downarrow R} \tilde{u}(r) = \infty. \quad (2.37)$$

or

$$\lim_{r \rightarrow \infty} r^\alpha \tilde{u}(r) = x_{2,M} \quad \text{and} \quad \lim_{r \downarrow R} \tilde{u}(r) = 0, \quad (2.38)$$

2- If $M = m^*$, $u_{x_{m^*}}$ is the unique self-similar solution, and statement 1-(ii) still holds with $x_{1,M}$ replaced by x_{m^*} in (2.35). There exist infinitely many radial positive solutions in $B_R \setminus \{0\}$ satisfying (2.36) with $x_{j,M}$ replaced by x_{m^*} , and at least one in $\mathbb{R}^N \setminus B_R$ satisfying (2.37) or (2.38) with $x_{2,M}$ replaced by x_{m^*} .

3- If $0 \leq M < m^*$, there exists no singular solution. For any $R > 0$ there exist $k > 0$ and a unique positive radial solution in $\mathbb{R}^N \setminus B_R$ satisfying

$$\begin{aligned} (i) \quad & \lim_{r \rightarrow \infty} r^{N-2} u(r) = k \\ (ii) \quad & \lim_{r \downarrow R} u(r) = \infty. \end{aligned} \quad (2.39)$$

Any positive radial solution defined in $\mathbb{R}^N \setminus B_R$ has the same asymptotic behaviour as in (2.39)-(i).

Proof. Case 1: $M > m^$.* (i) From Lemma 2.3-2, $P_{2,M}$ is a saddle point with stable trajectories \mathcal{T}_1 , \mathcal{T}_2 , and unstable ones \mathcal{T}_3 and \mathcal{T}_4 defined as in the proof of Theorem 2.4. The trajectory \mathcal{T}_1 lies in (D) as $t \rightarrow \infty$ and remains in (D) for all t because (D) is negatively invariant. Hence it converges to $P_{1,M}$ when $t \rightarrow -\infty$. Therefore \mathcal{T}_1 is an heteroclinic orbit connecting $P_{1,M}$ to $P_{2,M}$. It is unique and it corresponds to a solution u satisfying (2.34), thus u is unique up to the scaling transformations T_ℓ for $\ell > 0$.

(ii) The point $(0, 0)$ is a saddle point with unstable trajectory \mathcal{T}_r and stable trajectory \mathcal{T}_s which converges to $(0, 0)$ as $t \rightarrow \infty$ with initial slope $N - 2$. To \mathcal{T}_s are associated the solutions u of (2.12) satisfying $\lim_{r \rightarrow \infty} r^{N-2} u(r) = k$, this solution is unique for fixed k and denoted by u_k . Since $N - 2 > \alpha$, this stable trajectory lies in the region (F) at infinity. Since (F) is negatively invariant, the two functions $x(t)$ and $y(t)$ are decreasing and thus \mathcal{T}_s converges to $P_{1,M}$ when $t \rightarrow -\infty$. Hence \mathcal{T}_s is a heteroclinic orbit connecting $P_{1,M}$ to $(0, 0)$ and it is unique. To this trajectory is associated a solution u of (2.12) satisfying (2.35) and unique up to the transformations T_ℓ .

(iii) The unstable trajectory \mathcal{T}_4 of $P_{2,M}$ enters the region (C), crosses the axis $0x$ and blows-up in finite time as in Theorem 2.4. Since $P_{1,M}$ is a source and a node, there exist trajectories different from \mathcal{T}_1 converging to $P_{1,M}$ when $t \rightarrow -\infty$ and with a slope at this point smaller than α . Consider one of them below \mathcal{T}_1 near $P_{1,M}$; either it enters the region (C), then intersects the axis $0x$ and finally blows-up, or it enters the region (D), but since it cannot converge to $P_{2,M}$, it leaves (D) and finally blows up as in the first case. In any case such a trajectory corresponds to a solution which satisfies (2.36) with $j = 1$. Because of the scaling invariance of the condition, R can take any positive value. Notice that since there may exist several trajectories converging to $P_{1,M}$ at $-\infty$ with the same slope at this point, the corresponding solution $u_{1,R}$ is not unique for R fixed.

As in the proof of Theorem 2.4 \mathcal{T}_3 corresponds to a solution satisfying (2.38). The stable trajectory \mathcal{T}_2 of $P_{2,M}$ lies in the region (B) near ∞ . Since this region is negatively invariant the trajectory remains in it, hence $x(t)$ and $y(t)$ are decreasing. If they were defined on \mathbb{R} , they would remain bounded by Proposition A.1 and the trajectory would converge to a fixed point in (B), different from $P_{2,M}$. Since such a point does not exist the functions $x(t)$ and $y(t)$ are defined on a maximal interval (T, ∞) and they blow-up when $t \downarrow T$. To this trajectory is associated a solution \tilde{u} of (2.12) satisfying (2.37) with $R = e^T$. The trajectory \mathcal{T}_2 is unique thus R can be fixed arbitrarily by using the scaling transformation T_ℓ .

Case 2: $M = m^$.* There exists a unique nontrivial equilibrium P_{m^*} . To the eigenvalue 0 is associated the eigenvector $(1, \alpha)$, while to the eigenvalue $N-2$ is associated the eigenvector $(1, -K)$. There exist two trajectories \mathcal{T}_3 and \mathcal{T}_4 converging to P_{m^*} when $t \rightarrow -\infty$. The trajectory \mathcal{T}_4 with slope $\alpha + 2 - N$ at P_{m^*} enters the region (C), crosses the axis $0x$ and blows-up in finite time. It corresponds to a solution u with $\lim_{r \rightarrow 0} r^\alpha u(r) = x_{m^*}$ when $r \rightarrow 0$ and which blows-up at $r = R$.

The point $(0, 0)$ is a saddle point. The stable manifold \mathcal{T}_1 has initial slope $N - 2$. It corresponds to a trajectory which converges to $(0, 0)$ when $t \rightarrow \infty$. Since the region (F) is negatively invariant this trajectory converges to P_{m^*} when $t \rightarrow -\infty$, and its slope at this point is α . Hence \mathcal{T}_1 is the central manifold at P_{m^*} . As in case (1) this trajectory corresponds to a positive solution u in $\mathbb{R}^N \setminus \{0\}$ which satisfies

$$\lim_{r \rightarrow 0} r^\alpha u(r) = x_{m^*} \quad \text{and} \quad \lim_{r \rightarrow \infty} r^{N-2} u(r) = k, \quad (2.40)$$

for some $k > 0$.

Moreover, any trajectory which has one point in the bounded negatively invariant region delimited by \mathcal{T}_s , \mathcal{T}_4 and the axis $0x$, converges to P_{m^*} when $t \rightarrow -\infty$, tangentially to the line \mathcal{L} . Since it cannot converge to $(0, 0)$, it crosses the axis $0x$ in finite time and it blows up for $t = T = \ln R$. This corresponds to a positive solution u of (2.12) in $B_R \setminus \{0\}$ which satisfies (2.36) with $x_{j,M}$ replaced by x_{m^*} .

We claim that there exists at least one trajectory belonging to the central manifold at P_{m^} which converges to P_{m^*} when $t \rightarrow \infty$ and blows up in finite time:* the backward trajectory \mathcal{T}_P of any $P \in \mathcal{C}_1 \cap \{(x, y) : x > x_{m^*}\}$, belongs locally to (B) for $t < 0$ since this region is negatively invariant. Furthermore its coordinates sat-

isfy $x(t) > x_{m^*}$ and $y(t) > y_{m^*}$ for $t < 0$. Next, the backward trajectory \mathcal{T}_P of any $P \in \mathcal{L} \cap \{(x, y) : x > x_{m^*}\}$ belongs to (B) and its coordinates satisfy also $x(t) > x_{m^*}$ and $y(t) > y_{m^*}$ for $t < 0$ and $x(t) > x_{m^*}$ for $t > 0$. Let \mathcal{U} be the set of points $P \in (B)$ such that \mathcal{T}_P crosses $\mathcal{C}_1 \cap \{(x, y) : x > x_{m^*}\}$ for some $t > 0$ and \mathcal{V} the set of points $P \in (B)$ such that \mathcal{T}_P crosses $\mathcal{L} \cap \{(x, y) : x > x_{m^*}\}$ for some $t > 0$. By standard transversality arguments \mathcal{U} and \mathcal{V} are open and disjoint. Since (B) is connected, it cannot be the union of the two sets \mathcal{U} and \mathcal{V} . Hence there exists P_0 in $(B) \setminus \{\mathcal{U} \cup \mathcal{V}\}$. By monotonicity, \mathcal{T}_{P_0} converges to p_{m^*} when $t \rightarrow \infty$. Clearly this trajectory cannot be defined on whole \mathbb{R} by Proposition A.1, hence it blows-up for $t = T = e^R$. This proves the existence of a solution u which satisfies

$$\lim_{r \rightarrow \infty} r^\alpha u(r) = x_{m^*} \quad \text{and} \quad \lim_{r \downarrow R} u(r) = 0. \quad (2.41)$$

Case 3: $0 < M < m^*$. There exists no equilibrium besides $(0, 0)$ which is a saddle point with unstable trajectory \mathcal{T}_r and stable trajectory \mathcal{T}_s with initial slope $N - 2 > \alpha$. The region (\tilde{B}) between \mathcal{C}_1 and \mathcal{L} is negatively invariant, hence \mathcal{T}_s remains in it and its two coordinate functions are decreasing and necessarily unbounded. The corresponding solution u of (2.12) cannot be defined for all $r > 0$ because of Proposition A.1, hence it blows-up for $r \downarrow R$. This proves (2.39). \square

Remark. It is noticeable that in the case $m = m^*$, the equilibrium P_{m^*} is not hyperbolic and the central manifold there consist in curves with the same slope at P_{m^*} but one is converging to this point when $t \rightarrow -\infty$ while the other (may be there are many) converges when $t \rightarrow \infty$.

3 The radial case for $q \neq \frac{2p}{p+1}$

In this section we study the nonnegative solutions of

$$-u'' - \frac{N-1}{r}u' + |u|^{p-1}u - M|u|^q = 0, \quad (3.1)$$

when $q \neq \frac{2p}{p+1}$.

3.1 Non-autonomous systems associated to the equation

Since $q \neq \frac{2p}{p+1}$ there exists no autonomous 2-dimensional system in which equation (2.12) can be transformed. The systems that we introduce below are suitable for specific range of singular phenomena characteristic of one of the following equations $\mathcal{L}_p u = 0$, $\mathcal{R}_q^M u = 0$ and $\mathcal{E}_{p,q}^M u = 0$.

3.1.1 System describing the behaviour of Emden-Fowler equation

We set

$$u(r) = r^{-\frac{2}{p-1}}x(t) = r^{-\alpha}x(t), \quad u'(r) = -r^{-\frac{p+1}{p-1}}y(t) = -r^{-\alpha-1}y(t), \quad t = \ln r. \quad (3.2)$$

If u is a positive solution of (2.12) there holds

$$\begin{aligned} x_t &= \alpha x - y \\ y_t &= -Ky - x^p + Me^{-\frac{\sigma t}{p-1}}|y|^q, \end{aligned} \quad (3.3)$$

where, we recall it, σ is defined in (1.23). Equivalently

$$x_{tt} + Lx_t - \alpha Kx - x^p + Me^{-\frac{\sigma t}{p-1}}|\alpha x - x_t|^q = 0, \quad (3.4)$$

where $K = N - 2 - \alpha$ and $L = K - \alpha$. If $M = 0$ this is the system which describes the radial solutions of $\mathcal{L}_p u = 0$.

3.1.2 System describing the behaviour of the Riccati equation

We set

$$u(r) = r^{-\frac{2-q}{q-1}}\xi(t) = r^{-\beta}\xi(t), \quad u'(r) = -r^{-\frac{1}{q-1}}\eta(t) = -r^{-\beta-1}\eta(t), \quad t = \ln r. \quad (3.5)$$

If u is a positive solution of (2.12), (ξ, η) satisfies the system

$$\begin{aligned} \xi_t &= \beta\xi - \eta \\ \eta_t &= -\kappa\eta - e^{\frac{\sigma t}{q-1}}\xi^p + M|\eta|^q, \end{aligned} \quad (3.6)$$

where $\kappa = N - \beta$ is defined at (1.24). The system admits a unique nontrivial equilibrium with $\xi \geq 0$ if and only if $\frac{N}{N-1} < q < 2$: it is

$$(\xi_M, \eta_M) = (\xi_M, \beta\xi_M) \quad \text{with} \quad \xi_M = \frac{1}{\beta} \left(\frac{\kappa}{M} \right)^{\beta+1}. \quad (3.7)$$

The system (3.6) is equivalent to

$$\xi_{tt} + (N - 2 - 2\beta)\xi_t - \beta\kappa\xi - e^{\frac{\sigma t}{q-1}}\xi^p + M|\beta\xi - \xi_t|^q = 0. \quad (3.8)$$

According to the sign of σ this system is a perturbation at $-\infty$ or at ∞ of

$$\begin{aligned} \xi_t &= \beta\xi - \eta \\ \eta_t &= -\kappa\eta + M|\eta|^q, \end{aligned} \quad (3.9)$$

which describes the radial positive solutions of $\mathcal{R}_q^M u = 0$.

3.1.3 System describing the behaviour of the eikonal equation

Assuming $p \neq q$, we set

$$u(r) = r^{-\frac{q}{p-q}}X(t) = r^{-\gamma}X(t), \quad u'(r) = -r^{-\frac{p}{p-q}}Y(t) = -r^{-\gamma-1}Y(t), \quad t = \ln r. \quad (3.10)$$

Then if u is a positive solution of (2.12), there holds

$$\begin{aligned} X_t &= \gamma X - Y \\ Y_t &= \theta Y + e^{-\frac{\sigma t}{p-q}} (M|Y|^q - X^p), \end{aligned} \quad (3.11)$$

where $\theta = \gamma + 2 - N$ is defined at (1.24). Equivalently

$$X_{tt} + (N - 2 - 2\gamma)X_t + \theta\gamma X + e^{-\frac{\sigma t}{p-q}} (M|\gamma X - X_t|^q - X^p) = 0. \quad (3.12)$$

According to the sign of σ this equation is a perturbation at $-\infty$ or at ∞ of

$$M|\gamma X - X_t|^q - X^p = 0$$

which corresponds to the eikonal equation $\mathcal{E}_{p,q}^M u = 0$. We note that in the case $q = \frac{N-2}{N-1}p$, then $\gamma = N - 2$, there exists an explicit radial solution of (1.1) which is

$$u_{M,p}^*(r) = C_p^* r^{2-N} \quad \text{with} \quad C_p^* = \left(M(N-2)^{\frac{N-2}{(N-1)p}} \right)^{\frac{N-1}{p}}. \quad (3.13)$$

The function $u_{M,p}^*$ is harmonic and satisfies $\mathcal{E}_{p,q}^M u_{M,p}^* = 0$. This solution which has already been noticed in [20] will be useful in the sequel.

Remark. The following relations between the solutions of the systems (3.3), (3.6) and (3.11) hold,

$$\begin{aligned} (i) \quad & u(r) = r^{-\alpha} x(t) = r^{-\beta} \xi(t) = r^{-\gamma} X(t) \\ (ii) \quad & u'(r) = -r^{\frac{p+1}{p-1}} y(t) = -r^{\frac{1}{q-1}} \eta(t) = -r^{\frac{p}{p-q}} Y(t), \end{aligned} \quad (3.14)$$

which implies

$$\begin{aligned} (i) \quad & \xi(t) = e^{-\frac{\sigma}{(q-1)(p-q)}t} X(t) = e^{-\frac{\sigma}{(q-1)(p-1)}t} x(t) \\ (ii) \quad & \eta(t) = e^{-\frac{\sigma}{(q-1)(p-q)}t} Y(t) = e^{-\frac{\sigma}{(q-1)(p-1)}t} y(t). \end{aligned} \quad (3.15)$$

This yields the following relations

$$x^{p-1} = X^{p-q} \xi^{q-1} \quad \text{and} \quad y^{p-1} = Y^{p-q} \eta^{q-1}. \quad (3.16)$$

3.1.4 Lyapounov and slope functions

There are several functions the variation of which along trajectories will be analyzed in the sequel. They are specific to the change of variable we use. The most surprising one is the function E described below.

Lemma 3.1 *Let $N \geq 1$, $p, q > 1$, $p \neq q$. We define E on $\mathbb{R}_+ \times \mathbb{R} \times \mathbb{R}$ by*

$$E(X, Y, t) = \frac{X^{p+1}}{p+1} - M\gamma^q \frac{X^{q+1}}{q+1} - e^{\frac{\sigma t}{p-q}} \left(\frac{(\gamma X - Y)^2}{2} + \frac{\gamma\theta X^2}{2} \right). \quad (3.17)$$

If u is a positive solution of (2.12) and X and Y are defined from (3.10), set

$$\mathcal{E}(t) = E(X(t), Y(t), t). \quad (3.18)$$

Then

$$\begin{aligned} \mathcal{E}_t(t) = & -M(\gamma X - Y)(\gamma^q X^q - |Y|^q) \\ & - e^{\frac{\sigma t}{p-q}} \left(\left(\frac{\alpha}{2} + \gamma + \theta \right) (\gamma X - Y)^2 + \frac{\sigma \gamma \theta}{2(p-q)} X^2 \right), \end{aligned} \quad (3.19)$$

where $(X, Y) = (X(t), Y(t))$.

Proof. There holds $Y = \gamma X - X_t$ and

$$X_{tt} = -\theta \gamma X - (N - 2 - 2\gamma)X_t - e^{-\frac{\sigma t}{p-q}}(M|Y|^q - X^p).$$

Multiplying by $e^{\frac{\sigma t}{p-q}} X_t$ we get

$$e^{\frac{\sigma t}{p-q}} \frac{d}{dt} \left(\frac{X_t^2}{2} + \theta \gamma \frac{X^2}{2} \right) - \frac{d}{dt} \left(\frac{X^{p+1}}{p+1} \right) = (\gamma + \theta) e^{\frac{\sigma t}{p-q}} X_t^2 - M|Y|^q X_t.$$

Putting

$$\mathcal{F}(t) = \frac{X^{p+1}}{p+1} - e^{\frac{\sigma t}{p-q}} \left(\frac{X_t^2}{2} + \theta \gamma \frac{X^2}{2} \right),$$

we obtain

$$\mathcal{F}_t(t) = -e^{\frac{\sigma t}{p-q}} \left(\frac{\sigma}{p-q} \left(\frac{X_t^2}{2} + \theta \gamma \frac{X^2}{2} \right) + (\gamma + \theta) X_t^2 \right) + M|Y|^q X_t.$$

Since

$$\mathcal{E}(t) = \mathcal{F}(t) - M \gamma^q \frac{X^{q+1}}{q+1},$$

and $X_t = \gamma X - Y$, we obtain

$$\mathcal{E}_t(t) = -M(\gamma X - Y)(\gamma^q X^q - |Y|^q) - e^{\frac{\sigma t}{p-q}} \left(\left(\frac{\sigma}{2(p-q)} + \gamma + \theta \right) X_t^2 + \frac{\sigma \gamma \theta}{2(p-q)} X^2 \right).$$

and (3.19) follows. \square

The slope of a trajectory has shown its importance in the previous section when studying solutions of (2.12) near an equilibrium. We introduce it as a Lyapounov type function the variations of which will be of particular interest for studying solutions of eikonal type.

Definition 3.2 *The slope of a solution u is*

$$S(t) = -\frac{ru'(r)}{u(r)} = \frac{y(t)}{x(t)} = \frac{\eta(t)}{\xi(t)} = \frac{Y(t)}{X(t)}. \quad (3.20)$$

Since $\frac{S_t}{S} = \frac{\eta_t}{\eta} - \frac{\xi_t}{\xi}$, there holds

$$\begin{aligned} S_t &= S(S + 2 - N) + \xi^{q-1} M |S|^q - x^{p-1} \\ &= S(S + 2 - N) + \xi^{q-1} (M |S|^q - X^{p-q}) \quad \text{if } q \neq p. \end{aligned} \quad (3.21)$$

Note that $S > 0$ if $Y > 0$.

3.2 Asymptotic estimates for the Riccati equation

The next lemma deals with estimates near 0 (resp. ∞) of radial subsolutions (resp. radial supersolutions) of the equation $\mathcal{R}_q^M u = 0$, which reduces to

$$\mathcal{R}_q^M u = -u'' - \frac{N-1}{r}u' - M|u'|^q = 0 \quad (3.22)$$

in the radial case.

Lemma 3.3 *Assume $N \geq 1$, $q > 1$ and $M > 0$.*

1- *Let u be any C^2 radial decreasing function satisfying $\mathcal{R}_q^M u \leq 0$ near 0.*

(i) *If $q > \frac{N}{N-1}$, then*

$$\liminf_{r \rightarrow 0} r^{\frac{1}{q-1}} |u'(r)| \geq \left(\frac{\kappa}{M}\right)^{\frac{1}{q-1}}. \quad (3.23)$$

Therefore

$$\liminf_{r \rightarrow 0} r^\beta u(r) \geq \frac{1}{\beta} \left(\frac{\kappa}{M}\right)^{\beta+1} \quad \text{if } q < 2, \quad (3.24)$$

$$\liminf_{r \rightarrow 0} |\ln r|^{-1} u(r) \geq \frac{N-2}{M} \quad \text{if } q = 2 \text{ and } N > 2. \quad (3.25)$$

(ii) *If $q = \frac{N}{N-1}$, then*

$$\liminf_{r \rightarrow 0} r^{N-1} |\ln r|^{N-1} |u'(r)| \geq ((N-1)M)^{1-N}. \quad (3.26)$$

Therefore

$$\liminf_{r \rightarrow 0} r^{N-2} |\ln r|^{N-1} u(r) \geq \frac{((N-1)M)^{1-N}}{N-2} \quad \text{if } N \geq 3, \quad (3.27)$$

$$\liminf_{r \rightarrow 0} (\ln |\ln r|)^{-1} u(r) \geq \frac{1}{M} \quad \text{if } N = 2. \quad (3.28)$$

(iii) *If $N = 1$, or if $N \geq 2$ and $1 < q < \frac{N}{N-1}$, then $r^{N-1}|u'(r)|$ admits a limit belonging to $(0, \infty]$. Therefore if $N \geq 3$, $r^{N-2}u(r)$ admits a limit c belonging to $(0, \infty]$. If $N = 2$, $r^{N-2}u(r)$ has to be replaced by $|\ln r|^{-1}u(r)$ and if $N = 1$ by $u(r)$ in the previous expression.*

2- *Let u be any C^2 radial decreasing function satisfying $\mathcal{R}_q^M u \geq 0$ near 0. Then all the previous statements (3.23)–(3.28) are valid, provided \geq is replaced by \leq , \liminf by \limsup and, in case (iii), the limit c belongs to $[0, \infty)$. Furthermore if $q > 2$ the function u is bounded.*

3- *Let u be any C^2 radial decreasing function satisfying $\mathcal{R}_q^M u \leq 0$ in B_R^c and tending to 0 at infinity, and assume $q > \frac{N}{N-1}$. Then*

(iv) *either $q < 2$ and*

$$r^{\frac{1}{q-1}} |u'(r)| \geq \left(\frac{\kappa}{M}\right)^{\frac{1}{q-1}} \quad \text{and} \quad r^\beta u(r) \geq \frac{1}{\beta} \left(\frac{\kappa}{M}\right)^{\frac{1}{q-1}}, \quad (3.29)$$

for r large enough,

(v) or $N > 2$ and

$$\lim_{r \rightarrow \infty} r^{N-1}|u'(r)| = c \geq 0 \quad \text{and} \quad \lim_{r \rightarrow \infty} r^{N-2}u(r) = C \geq 0. \quad (3.30)$$

4- Let u be any C^2 radial decreasing function satisfying $\mathcal{R}_q^M u \geq 0$ in B_R^c and tending to 0 at infinity. Then $q > \frac{N}{N-1}$, and either $N \geq 3$ and $\lim_{r \rightarrow \infty} r^{N-1}|u'(r)| = c > 0$, or

$$r^{\frac{1}{q-1}}|u'(r)| \leq \left(\frac{\kappa}{M}\right)^{\frac{1}{q-1}}; \quad \text{and if } q < 2, \text{ then } r^\beta u(r) \leq \frac{1}{\beta} \left(\frac{\kappa}{M}\right)^{\beta+1}, \quad (3.31)$$

for r large enough.

Proof. If u is a radial decreasing subsolution (resp. supersolution), there holds

$$(r^{N-1}u')' + Mr^{N-1}|u'|^q \geq 0 \quad (\text{resp. } \leq 0).$$

Set $W(r) = -r^{N-1}u'(r) = r^{N-1}|u'(r)|$, then

$$Mr^{-(N-1)(q-1)} - W^{-q}W' \geq 0 \quad (\text{resp. } \leq 0).$$

Hence the function

$$r \mapsto \phi(r) = \begin{cases} W^{1-q}(r) - \frac{M}{\kappa} r^{N-(N-1)q} & \text{if } \kappa = \frac{(N-1)q-N}{q-1} \neq 0 \\ W^{1-q}(r) + \frac{M}{N-1} \ln r & \text{if } \kappa = 0, \end{cases} \quad (3.32)$$

is nondecreasing (resp. nonincreasing). Notice in particular that if u is a decreasing radial solution, there holds

$$|u'(r)| = \begin{cases} r^{1-N} \left(C + \frac{M}{\kappa} r^{N-(N-1)q}\right)^{-\frac{1}{q-1}} & \text{if } \kappa \neq 0 \\ r^{1-N} \left(C - \frac{M}{N-1} \ln r\right)^{-\frac{1}{q-1}} & \text{if } \kappa = 0, \end{cases} \quad (3.33)$$

and the estimate on u follows by integration since $\beta = \frac{2-q}{q-1}$.

1- If u is a decreasing subsolution, ϕ is nondecreasing.

(i)- If $\kappa > 0$, then

$$W^{1-q}(r) \leq \frac{M}{\kappa} r^{N-(N-1)q} + c_0 \quad \text{for } 0 < r \leq r_0,$$

where $c_0 = W^{1-q}(r_0) - \frac{M}{\kappa} r_0^{N-(N-1)q}$. Since $N - (N-1)q < 0$, (3.23) follows.

(ii)- If $\kappa = 0$, then for any $\epsilon > 0$, there exists $r_\epsilon > 0$ such that

$$W^{1-q}(r) \leq \left(\frac{M}{N-1} + \epsilon\right) |\ln r| \quad \text{for } 0 < r \leq r_\epsilon,$$

and (3.26) follows.

(iii)- If $\kappa < 0$, then $r^{N-(N-1)q} \rightarrow 0$ as $r \rightarrow 0$. Therefore $W(r)$ admits a limit belonging to $(0, \infty]$ when $r \rightarrow 0$. We derive the estimates on u by integration.

2- If u is a decreasing supersolution the results follow in the same way. If $q > 2$, the estimate

$$\limsup_{r \rightarrow 0} r^{\frac{1}{q-1}} |u'(r)| \leq \left(\frac{M}{\kappa} \right)^{\frac{1}{q-1}},$$

implies that u is bounded near 0.

3- If u is a decreasing subsolution in an exterior domain, the function ϕ defined in (3.32) is nondecreasing, hence it admits a limit ν in $(-\infty, \infty]$.

(iv)- If $\kappa > 0$, then $r^{N-(N-1)q} \rightarrow 0$ as $r \rightarrow \infty$, hence $W^{1-q}(r) \rightarrow \nu \in [0, \infty]$, therefore $r^{N-1}|u'(r)| \rightarrow c \in [0, \infty]$. If $\nu \in (0, \infty]$, then $c \in [0, \infty)$. Since u tends to 0 at infinity, we obtain $r^{N-2}u(r) \rightarrow \frac{c}{N-2}$ when $r \rightarrow \infty$.

If $\nu = 0$, then $W^{1-q}(r) \leq \frac{M}{\kappa} r^{N-(N-1)q}$. This yields the estimate from below (3.29) of $|u'(r)|$, and therefore for $u(r)$ if $q < 2$.

If $q \geq 2$, we obtain $|u'(r)| \geq cr^{-\frac{1}{q-1}}$, and we derive a contradiction since $r^{-\frac{1}{q-1}}$ is not integrable at infinity.

4- If u is a decreasing supersolution in an exterior domain, then $r^{N-1}|u'(r)|$ is nondecreasing. Hence there exists $c > a$ such that $r^{N-1}|u'(r)| \geq a$, which implies $u(r) \geq cr^{2-N}$ for some $c > 0$. Since the function ϕ is nonincreasing, it admits a limit ν belonging to $[-\infty, \infty)$. If $\kappa > 0$ and because $r^{N-(N-1)q} \rightarrow 0$, it follows that $\nu \in [0, \infty)$. If $\nu > 0$, then $r^{N-1}|u'(r)|$ has a limit in $(0, \infty)$, and this implies that $r^{N-2}u(r)$ admits a limit in $(0, \infty)$ at infinity. If $\nu = 0$, then $W^{1-q}(r) - \frac{M}{\kappa} r^{N-(N-1)q} \geq 0$ and we obtain (3.31). If $\kappa \leq 0$, then $\phi(r) \rightarrow \infty$ as $r \rightarrow \infty$, contradiction. \square

3.3 Estimates near 0

In this paragraph we prove Theorem 1.7 and Theorem 1.8.

3.3.1 The case $\frac{2p}{p+1} < q < p$.

Here we prove Theorem 1.7. If u is a positive solution of (2.12) unbounded near 0, then $u' < 0$, hence the variable X and Y defined in (3.10) satisfy

$$\begin{aligned} X_t &= \gamma X - Y \\ Y_t &= \theta Y + e^{-\frac{\sigma t}{p-q}} (MY^q - X^p), \end{aligned} \tag{3.34}$$

where, we recall it, $\sigma = (p+1)q - 2p > 0$ and $\theta = \frac{(N-1)q - (N-2)p}{p-q}$. Since $q < p$, $X(t)$ remains bounded when $t \rightarrow -\infty$. The difficulty comes from the fact that the term $e^{-\frac{\sigma t}{p-q}}$ tends to infinity when $t \rightarrow -\infty$.

Lemma 3.4 *Assume $\frac{2p}{p+1} < q < p$. If u is a positive solution of (2.12) in $B_R \setminus \{0\}$ such that $u' < 0$, then $r^\gamma u(r)$ admits a limit when $r \rightarrow 0$ which can take only the values X_M or 0.*

Proof. We use the function \mathcal{E} introduced in (3.18). Because of Proposition A.1 and Proposition A.3, X and Y are bounded. By assumption σ is nonnegative, hence $\mathcal{E}(t)$ is bounded when $t \rightarrow -\infty$. Using (3.19) we have that

$$\mathcal{E}_t(t) + M(\gamma X - Y)(\gamma^q X^q - Y^q) = -e^{\frac{\sigma t}{p-q}} \left(\left(\frac{\alpha}{2} + \gamma + \theta \right) (\gamma X - Y)^2 + \frac{\sigma \gamma \theta}{2} X^2 \right),$$

which implies that

$$-\frac{C_2 \sigma}{p-q} e^{\frac{\sigma t}{p-q}} \leq \mathcal{E}_t(t) + M(\gamma X - Y)(\gamma^q X^q - Y^q) \leq \frac{C_2 \sigma}{p-q} e^{\frac{\sigma t}{p-q}}, \quad (3.35)$$

for some $C_2 > 0$. Because $(\gamma X - Y)(\gamma^q X^q - Y^q) \geq 0$, we deduce that the function $t \mapsto \mathcal{E}(t) - C_2 e^{\frac{\sigma t}{p-q}}$ is decreasing, therefore it admits a finite limit Λ when $t \rightarrow -\infty$, and Λ is also the limit of $\mathcal{E}(t)$. Hence

$$\lim_{t \rightarrow -\infty} \left(\frac{X^{p+1}(t)}{p+1} - M \gamma^q \frac{X^{q+1}(t)}{q+1} \right) = \Lambda. \quad (3.36)$$

Therefore $X(t)$ converges to some λ satisfying $\frac{\lambda^{p+1}}{p+1} - M \gamma^q \frac{\lambda^{q+1}}{q+1} = \Lambda$. The omega-limit set at $-\infty$ of the trajectory $\{(X(t), Y(t))\}_{t \in \mathbb{R}_-}$ is the set Γ of couples (X_0, Y_0) such that there exists a sequence $\{t_n\}$ decreasing to $-\infty$ such that $(X(t_n), Y(t_n)) \rightarrow (X_0, Y_0)$. It is non-empty since the trajectory is bounded, connected and compact. By La Salle's theorem, the function $\mathcal{E}(t) - C_2 e^{\frac{\sigma t}{p-q}}$ which is monotone decreasing is constant on Γ . This implies $M(\gamma X_0 - Y_0)(\gamma^q X_0^q - Y_0^q) = 0$, hence $Y_0 = \gamma X_0$. Because $X(t) \rightarrow \lambda$ then $X_0 = \lambda$, hence $Y_0 = \gamma \lambda$ and $Y(t) \rightarrow \gamma \lambda$ when $t \rightarrow -\infty$. If $M \gamma^q \lambda^q \neq \lambda^p$, it implies that

$$Y_t(t) = \theta \gamma \lambda + e^{-\frac{\sigma t}{p-q}} (M \gamma^q \lambda^q - \lambda^p + \epsilon(t)) \quad \text{where } \epsilon(t) \rightarrow 0 \text{ as } t \rightarrow -\infty.$$

Hence $Y_t(t) = c e^{-\frac{\sigma t}{p-q}} (1 + o(1))$ where $c \neq 0$. Clearly this implies that $Y(t)$ cannot be bounded, contradiction. Therefore $M \gamma^q \lambda^q = \lambda^p$. This implies that

$$\lambda \in \left\{ 0, M^{\frac{1}{p-q}} \gamma^\gamma \right\}, \quad (3.37)$$

which ends the proof. \square

Lemma 3.5 *Assume $N \geq 2$, and let u be a positive solution of (2.12) in B_R unbounded near 0.*

1- *If $q > \frac{N}{N-1}$ and $u(r) = o\left(r^{-\frac{q}{p(q-1)}}\right)$ near $r = 0$, then necessarily $q \leq 2$ and*

$$\lim_{r \rightarrow 0} r^{\frac{1}{q-1}} |u'(r)| = \eta_M := \left(\frac{\kappa}{M} \right)^{\frac{1}{q-1}}. \quad (3.38)$$

Therefore

$$\lim_{r \rightarrow 0} r^\beta u(r) = \xi_M := \frac{1}{\beta} \left(\frac{\kappa}{M} \right)^{\frac{1}{q-1}} \quad \text{if } q < 2, \quad (3.39)$$

$$\lim_{r \rightarrow 0} |\ln r|^{-1} u(r) = \frac{N-2}{M} \quad \text{if } q = 2. \quad (3.40)$$

2- If $\frac{2p}{p+1} < q = \frac{N}{N-1}$ and $r^{\beta+\epsilon} u(r) = r^{N-2+\epsilon} u(r)$ is bounded for any $\epsilon > 0$, then

$$\lim_{r \rightarrow 0} r^{N-2} |\ln r|^{N-1} u(r) = \frac{1}{N-2} \left(\frac{N-1}{M} \right)^{N-1} \quad \text{if } N \geq 3. \quad (3.41)$$

$$\lim_{r \rightarrow 0} r |\ln r| |u'(r)| = \lim_{r \rightarrow 0} \ln(|\ln r|) u(r) = \frac{1}{M} \quad \text{if } q = N = 2. \quad (3.42)$$

3- If $\frac{2p}{p+1} \leq q < \frac{N}{N-1}$ and $r^{N-2} u(r)$ is bounded if $N \geq 3$ or $r^\epsilon u(r)$ is bounded for any $\epsilon > 0$ if $N = 2$, then there exists $k > 0$ such that

$$\lim_{r \rightarrow 0} r^{N-2} u(r) = k \quad \text{if } N \geq 3, \quad (3.43)$$

and

$$\lim_{r \rightarrow 0} |\ln r|^{-1} u(r) = k \quad \text{if } N = 2. \quad (3.44)$$

Proof. We first notice that if u is unbounded near 0, then $u' < 0$ in a neighborhood of 0 and we can apply the results of Lemma 3.3 concerning subsolutions. Moreover, if $u^p(r) = o(|u'(r)|^q)$ when $r \rightarrow 0$, then for any $\delta > 0$ there exists $r_\delta > 0$ such that

$$M(1-\delta)|u'|^q \leq -\Delta u \leq M|u'|^q \quad \text{in } B_{r_\delta} \setminus \{0\}, \quad (3.45)$$

and we can also use the results of Lemma 3.3 dealing with supersolutions.

1- Since u is a decreasing subsolution of $\mathcal{R}_q^M u = 0$, $|u'(r)| \geq cr^{-\beta-1}$ near 0, hence $|u'(r)|^q \geq cr^{-\frac{q}{\beta+1}}$. By assumption $u^p(r) = o\left(r^{-\frac{q}{\beta+1}}\right)$. Then $u^p(r) = o(|u'(r)|^q)$ near 0, hence (3.45) applies. It follows by Lemma 3.3-(1)-(2) that

$$\left(\frac{\kappa}{M} \right)^{\beta+1} \leq \liminf_{r \rightarrow 0} r^{\beta+1} |u'(r)|^q \leq \limsup_{r \rightarrow 0} r^{\beta+1} |u'(r)|^q \leq \left(\frac{\kappa}{M(1-\delta)} \right)^{\beta+1}. \quad (3.46)$$

Since $\delta > 0$ is arbitrary, this implies (3.38). The other estimates (3.39) and (3.40) are obtained by integration.

2- By (3.26), $|u'(r)|^q \geq cr^{q(1-N)} |\ln r|^{q(1-N)} = cr^{-N} |\ln r|^{-N}$. From the assumptions, for any $\epsilon > 0$, $u^p(r) \leq c_\epsilon r^{p(2-N-\epsilon)}$, then

$$\frac{u^p(r)}{|u'(r)|^q} \leq c'_\epsilon r^{N-(N-2+\epsilon)p} |\ln r|^N.$$

Next $\frac{2p}{p+1} < \frac{N}{N-1}$ implies that $N > p(N-2)$. Therefore, we can take $\epsilon > 0$ small enough such that $N - (N-2+\epsilon)p > 0$. This implies that (3.45) holds in $B_{r_\delta} \setminus \{0\}$. Hence we get (3.41) and (3.42) by integration.

3- Suppose $q < \frac{N}{N-1}$ then $p < \frac{N}{N-2}$ if $N \geq 3$. By Lemma 3.3-(1), $r^{N-1} |u'(r)| \geq c > 0$ near 0, hence

$$\frac{u^p(r)}{|u'(r)|^q} \leq \tilde{c}^{-q} r^{(N-1)q - (N-2)p}.$$

Since $(N-1)q - (N-2)p \geq (N-1)\frac{2p}{p+1} - (N-2)p = \frac{p}{p+1}(N - (N-2)p) > 0$, we deduce that for any $\delta > 0$, (3.45) holds in $B_{r_\delta} \setminus \{0\}$. Then we use Lemma 3.3 and obtain (3.43) and (3.44) by integration. In the case $N = 2$ there holds

$$\frac{u^p}{|u'|^q} \leq cr^{q-\epsilon p}$$

for any $\epsilon > 0$. Choosing $\epsilon < \frac{q}{p}$, we find again (3.46). \square

For obtaining the next result, the key is the introduction of the slope function S which allows to make precise the behaviour of solutions such that $r^\gamma u(r) \rightarrow 0$.

Lemma 3.6 *Assume $N \geq 2$, $\frac{2p}{p+1} < q < p$ and $M > 0$. If u is a positive solution of (2.12) unbounded near 0 and such that $r^\gamma u(r) \rightarrow 0$ when $r \rightarrow 0$, then $q \leq 2$ and the following trichotomy holds.*

- 1- If $q > \frac{N}{N-1}$, then (3.39) or (3.40) is satisfied.
- 2- If $q = \frac{N}{N-1}$, then (3.41) or (3.42) is satisfied.
- 3- If $q < \frac{N}{N-1}$, then (3.43) or (3.44) is satisfied.

Proof. By assumption $X(t) \rightarrow 0$ as $t \rightarrow -\infty$. We recall that $S(t) = \frac{Y(t)}{X(t)}$ satisfies (3.21) hence

$$\begin{aligned} X_t &= X(\gamma - S) \\ S_t &= S(S + 2 - N) + \xi^{q-1}(MS^q - X^{p-q}). \end{aligned} \quad (3.47)$$

1- We first assume that $S(t) \rightarrow 0$ as $t \rightarrow -\infty$. Then for any $\epsilon > 0$, there exists $r_\epsilon > 0$ such that $0 < -\frac{ru'(r)}{u(r)} \leq \epsilon$ on $(0, r_\epsilon]$. Hence $r \mapsto r^\epsilon u(r)$ is increasing. This implies that $r^\epsilon u(r)$ is bounded near 0 and thus $q \leq 2$ by Lemma 3.5. If $\frac{N}{N-1} < q < 2$ it would follow from Lemma 3.5 that (3.39) holds, which is not possible. Hence $N \geq 3$, $q = 2$ and (3.40) holds. If $q = \frac{N}{N-1}$ and $N \geq 3$, (3.41) cannot hold; hence $N = q = 2$ and (3.42) holds. If $q < \frac{N}{N-1}$, (3.43) cannot be satisfied, hence $N = 2$ and (3.44) holds.

2- Now we assume that $\liminf_{t \rightarrow -\infty} S(t) = m > 0$. Then there exist $t_0 > -\infty$ and $m_0 \in (0, \infty)$ such that $S(t) \geq m_0$ for $t \leq t_0$. Hence $Y^q(t) \geq m_0 X^q(t)$ therefore $X^p(t) = X^{p-q}(t)X^q(t) = o(Y^q(t))$ as $t \rightarrow -\infty$. This implies $u^p(r) = o(|u'(r)|^q)$ as $r \rightarrow 0$. Then (3.45) holds. Using Lemma 3.3-(1)-(2), we have (3.39) or (3.40) if $q > \frac{N}{N-1}$, (3.41) or (3.42) if $q = \frac{N}{N-1}$ and (3.43) or (3.44) if $q < \frac{N}{N-1}$.

3- Next we assume that $0 = \liminf_{t \rightarrow -\infty} S(t) < \limsup_{t \rightarrow -\infty} S(t) = \Sigma \in (0, \infty]$. Then there exists a decreasing sequence $\{t_n\}$ converging to $-\infty$ such that $S_t(t_n) = 0$ and $S_n := S(t_n)$, which is a local maximum of $S(t)$, tends to Σ . Put $X_n = X(t_n)$ and $\xi_n = \xi(t_n)$, then

$$\xi_n^{q-1} = \frac{S_n(N-2-S_n)}{MS_n^q - X_n^{p-q}} = \frac{S_n(N-2-S_n)}{MS_n^q(1-\epsilon_n)} = \frac{N-2-S_n}{MS_n^{q-1}(1-\epsilon_n)}, \quad (3.48)$$

with $\epsilon_n \rightarrow 0$. This implies in particular $N > 2$ and $S_n < N - 2$. Since it holds for all local maximum of S_n we deduce $S < N - 2$, which implies $u(r) \leq Cr^{2-N}$. If $q < \frac{N}{N-1}$ (resp. $q = \frac{N}{N-1}$) we obtain (3.43) from Lemma 3.5-(3) (resp. (3.41) from Lemma 3.5-(2)). If $q > \frac{N}{N-1}$ we write (3.48) under the form

$$\xi_n^{q-1} M S_n^{q-1} (1 - \epsilon_n) = \eta_n^{q-1} M (1 - \epsilon_n) = N - 2 - S_n. \quad (3.49)$$

From (3.23), $\eta_n^{q-1} \geq \frac{\kappa}{M} (1 - \epsilon'_n)$ for n large enough and $\epsilon'_n \rightarrow 0$ when $n \rightarrow \infty$, hence

$$N - 2 - S_n \geq (1 - \epsilon_n)(1 - \epsilon'_n)\kappa \implies S_n \leq N - 2 - \kappa + \epsilon''_n = \beta + \epsilon''_n. \quad (3.50)$$

This implies that for any $\epsilon > 0$ there exists n_ϵ such that $S(t) \leq S_{n_\epsilon} \leq \beta + \frac{\epsilon}{2}$ for $t \leq t_{n_\epsilon}$. Hence $r^{\beta+\epsilon}u(r) \rightarrow 0$ as $r \rightarrow 0$. Since $\frac{q}{p(q-1)} = \beta + \frac{q(p+1)-2p}{p(q-1)}$, it implies that $u(r) = o(r^{-\frac{q}{p(q-1)}})$ as $r \rightarrow 0$. Therefore (3.39) and (3.40) hold. \square

Proof of Theorem 1.7. It follows from Lemma 3.4, Lemma 3.5 and Lemma 3.6. \square

3.3.2 The case $1 < q < \frac{2p}{p+1}$.

Proof of Theorem 1.8. If $1 < q < \frac{2p}{p+1}$ and $p \geq \frac{N}{N-2}$ it is proved in [7] that positive solutions of (1.1) in $B_R \setminus \{0\}$ can be extended as a C^2 solution in B_R . Next we suppose that $p < \frac{N}{N-2}$, or $N = 1, 2$, hence $q < \frac{N}{N-1}$. We use the change of variable (3.2) and (x, y) satisfies (3.3). It is important to notice that $\sigma = (p+1)q - 2p$ is negative, therefore the system satisfied by (x, y) is a perturbation at $-\infty$ of the system

$$\begin{aligned} x_t &= \alpha x - y \\ y_t &= -Ky - x^p \end{aligned} \quad (3.51)$$

where $K = N - 2 - \alpha$, associated to the Emden-Fowler equation $\mathcal{L}_p u = 0$ by the same change of variable. Since $(x(t), y(t))$ is bounded, the omega-limit set at $-\infty$ of the trajectory $\{(x(t), y(t))\}_{t \in \mathbb{R}_-}$ is a non-empty compact connected subset of the set of stationary solutions of (3.4). Therefore

$$\lim_{t \rightarrow -\infty} (x(t), y(t)) = (\ell, \alpha\ell) \quad \text{where } \ell \in \{0, x_0\}. \quad (3.52)$$

If $\ell = x_0$ the result is proved, thus let us assume that $\ell = 0$. By Lemma 3.3-1-(iii) $r^{N-1}u'(r)$ admits a limit $c \in (0, \infty]$ when $r \rightarrow 0$. If $c < \infty$, (1.32) follows by integration. Thus we are left with the case $c = \infty$. Hence $\lim_{r \rightarrow 0} r^{N-2}u(r) = \infty$ if $N \geq 3$, or $\lim_{r \rightarrow 0} |\ln r|^{-1}u(r) = \infty$ if $N = 2$. Therefore, for any $k > 0$, u is bounded from below in $B_R \setminus \{0\}$ by the function v_k which satisfies $\mathcal{L}_p v_k = 0$ in $B_R \setminus \{0\}$, $v_k = 0$ on ∂B_R and $\lim_{r \rightarrow 0} r^{N-2}v_k(r) = k$ if $N \geq 3$, or $\lim_{r \rightarrow 0} |\ln r|^{-1}v_k(r) = k$ if $N = 2$. Letting $k \rightarrow \infty$, $v_k \uparrow v_\infty$, and $\lim_{r \rightarrow 0} r^\alpha v_\infty(r) = x_0$ by [22]. This is a contradiction. \square

3.4 Estimates at infinity

3.4.1 The case $q > \frac{2p}{p+1}$.

Proof of Theorem 1.9. We recall that by Proposition A.1 and Proposition A.3 in the Appendix all the positive solutions of (1.1) in B_R^c satisfy

$$u(r) + r|u'(r)| \leq cr^{-\alpha} \quad \text{in } B_{R+1}^c. \quad (3.53)$$

where $c = c(N, p, q) > 0$, and by the maximum principle they are decreasing. Since u is continuous in B_R^c , $v = \min\{u(r) : r = R\}$ is well defined and positive. By the maximum principle, for any $n > R$, u is bounded from below in $B_n \setminus B_R$ by the solution \tilde{v}_n of

$$\mathcal{L}_p \tilde{v}_n = 0 \quad \text{in } B_n \setminus \overline{B}_R, \quad \tilde{v}_n = v \quad \text{on } \partial B_R, \quad \tilde{v}_n = 0 \quad \text{on } \partial B_n. \quad (3.54)$$

When $n \rightarrow \infty$, $\tilde{v}_n \uparrow \tilde{v}_\infty$ which satisfies $\mathcal{L}_p \tilde{v}_\infty = 0$ in B_R^c and $\tilde{v}_\infty = v$ on ∂B_R . Then $u \geq \tilde{v}_\infty$ and by [21], \tilde{v}_∞ satisfies

$$\begin{aligned} (i) \quad & \lim_{r \rightarrow \infty} r^\alpha \tilde{v}_\infty(r) = x_0 && \text{if } 1 < p < \frac{N}{N-2} \\ (ii) \quad & \lim_{r \rightarrow \infty} r^{N-2} (\ln r)^{\frac{N-2}{2}} \tilde{v}_\infty(r) = \left(\frac{N-2}{\sqrt{2}} \right)^{N-2} && \text{if } N \geq 3 \text{ and } p = \frac{N}{N-2} \\ (ii) \quad & \lim_{r \rightarrow \infty} r^{N-2} \tilde{v}_\infty(r) = c > 0 && \text{if } N \geq 3 \text{ and } p > \frac{N}{N-2}. \end{aligned} \quad (3.55)$$

We make the change of variable (3.2) and obtain the system (3.3) satisfied by the functions $t \mapsto (x(t), y(t))$. Since $q > \frac{2p}{p+1}$, σ is positive. Hence the omega-limit set of the trajectory of $\{(x(t), y(t))\}_{t \geq 0}$ as $t \rightarrow \infty$ is a non-empty compact connected set of the set of solutions of stationary solutions of (3.51), therefore

$$\lim_{t \rightarrow \infty} (x(t), y(t)) = (\ell, \alpha \ell) \quad \text{where } \ell \in \{0, x_0\}. \quad (3.56)$$

Therefore if $1 < p < \frac{N}{N-2}$ we obtain (1.33), and if $p \geq \frac{N}{N-2}$ we have that $\ell = 0$. If $p > \frac{N}{N-2}$, then $q > \frac{N}{N-1}$. From Lemma 3.3-(3), we have that either $q < 2$ and (3.29) holds, or $N > 2$ and (3.30) holds. However, since $q > \frac{2p}{p+1}$, one has $r^{-\beta} = o(r^{-\alpha})$ when $r \rightarrow \infty$, hence (3.29) does not hold and we deduce that (3.30) is verified. Finally we consider the case $p = \frac{N}{N-2}$. Then $\ell = 0$ and x satisfies

$$x_{tt} - (N-2)x_t - x^{\frac{N}{N-2}} + M e^{-((N-1)q-N)t} ((N-2)x - x_t)^q = 0 \quad \text{on } (\ln R, \infty), \quad (3.57)$$

and $q > \frac{2p}{p+1} = \frac{N}{N-1}$. Since u is bounded from below by v_∞ , we have that $x(t) \geq ct^{-\frac{N-2}{2}}$, with $c > 0$, for t large enough. Hence for any $\epsilon > 0$ there exists $t_\epsilon > \ln R$ such that

$$x_{tt} - (N-2)x_t - x^{\frac{N}{N-2}} \leq 0 \leq x_{tt} - (N-2)x_t - (1-\epsilon)x^{\frac{N}{N-2}} \quad \text{on } (\ln R, \infty). \quad (3.58)$$

Therefore $\theta_{2,\epsilon}(t) \leq x(t) \leq \theta_{1,\epsilon}(t)$ where

$$\begin{aligned} \frac{d^2}{dt^2}\theta_{j,\epsilon} - (N-2)\frac{d}{dt}\theta_{j,\epsilon} - (1+(-1)^j\epsilon)\theta_{j,\epsilon}^{\frac{N}{N-2}} &= 0 \quad \text{on } (t_\epsilon, \infty) \\ \theta_{j,\epsilon}(t_\epsilon) &= x(t_\epsilon). \end{aligned} \quad (3.59)$$

The asymptotic expansion of $\theta_{j,\epsilon}$ is obtained in [21, Lemme 3.2] using an old result due to Hardy. We give below a simpler proof.

$$\theta_{j,\epsilon}(t) = \left(\frac{N-2}{\sqrt{1+(-1)^j\epsilon}} \right)^{N-2} \left(\frac{1}{2t} \right)^{\frac{N-2}{2}} (1+o(1)). \quad (3.60)$$

This implies that for any $\epsilon > 0$ there holds

$$\left(\frac{N-2}{\sqrt{2(1+\epsilon)}} \right)^{N-2} \leq \liminf_{t \rightarrow -\infty} t^{\frac{N-2}{2}} x(t) \leq \limsup_{t \rightarrow -\infty} t^{\frac{N-2}{2}} x(t) \leq \left(\frac{N-2}{\sqrt{2(1-\epsilon)}} \right)^{N-2}, \quad (3.61)$$

which implies (1.35). \square

Remark. The proof of Hardy's theorem quoted in [3] is not easy to find. An alternative proof is to consider the following equation, to which (3.59) reduces by a suitable scaling transformation,

$$\theta'' - \theta' - \theta^n = 0 \quad \text{on } [0, \infty), \quad (3.62)$$

where $n > 1$ and $\theta > 0$. Since $\theta(t) \rightarrow 0$ as $t \rightarrow \infty$, it is easy to see that for any $t > 1$, $\theta(t) \leq Ct^{-\frac{1}{n-1}}$ by considering supersolutions under the form

$$\psi(t) = at^{-\frac{1}{n-1}} + bt^{-\frac{2}{n-1}}.$$

Since $\phi(t) = \left(\frac{1}{(n-1)(t+t_0)} \right)^{\frac{1}{n-1}}$ is a subsolution for some $t_0 > 0$, it is smaller than $\theta(t)$. Furthermore, for any $\epsilon > 0$, there exists $t_\epsilon > 0$ such that $t \mapsto \left(\frac{1+\epsilon}{(n-1)t} \right)^{\frac{1}{n-1}}$ is a supersolution on $[t_\epsilon, \infty)$ and is larger than θ . From that we infer

$$\lim_{t \rightarrow \infty} t^{\frac{1}{n-1}} \theta(t) = \left(\frac{1}{n-1} \right)^{\frac{1}{n-1}}. \quad (3.63)$$

An alternative proof of the convergence is to set $x(t) = t^{\frac{2-N}{2}} \mathcal{Z}(t)$. We get

$$\mathcal{Z}_{tt} - (N-2) \left(1 + \frac{1}{t} \right) \mathcal{Z}_t + \frac{1}{t} \left(\left(\frac{(N-2)^2}{2} + \frac{N(N-2)}{4t} \right) \mathcal{Z} - \mathcal{Z}^{\frac{N}{N-2}} \right) + \Phi = 0$$

where $\Phi(t) = O\left(t^{\frac{N}{N-2}} e^{(N-(N-1)q)t}\right)$. Applying [10, Corollary 4.2] we deduce that $\mathcal{Z}(t)$ converges to a limit ℓ which satisfies $\ell \left(\frac{(N-2)^2}{2} - \ell^{\frac{2}{N-2}} \right) = 0$. From the lower bound $u \geq \tilde{v}_\infty$ and (3.55) we infer that $\ell = 0$ is impossible.

3.4.2 The case $1 < q < \frac{2p}{p+1}$.

Proof of Theorem 1.10. If $1 < q < \frac{2p}{p+1}$, then $q < p$. Therefore $\gamma < \alpha < \beta$. Hence if u a nonnegative solution of (2.12) on $[R, \infty)$, $r^\gamma u(r)$ is bounded for $r \geq R > 0$. Therefore the natural system for describing the solution is the system (3.11) with bounded $X(t)$ and $Y(t)$ and we use an argument similar to the poof of Lemma 3.4.

Lemma 3.7 *Assume $p > 1$, $1 < q < \frac{2p}{p+1}$ and $M > 0$. If u is a positive radial solution of (1.1) in B_R^c , there holds*

$$\lim_{r \rightarrow \infty} r^\gamma u(r) = \ell \in \{0, X_M\}. \quad (3.64)$$

Proof. By Proposition A.3 Y is bounded, hence (3.35) still holds on $[T, \infty)$. We consider now the function \mathcal{E} defined in (3.18), then $t \mapsto \mathcal{E}(t) - \frac{C_2|\sigma|}{p-q} e^{\frac{\sigma t}{p-1}}$ is decreasing and bounded at infinity since $\sigma < 0$. Therefore $\mathcal{E}(t)$ converges to some real number Λ when $t \rightarrow \infty$. This implies that identity (3.36) is still valid provided $\lim_{t \rightarrow -\infty}$ is replaced by $\lim_{t \rightarrow \infty}$. Mutatis mutandis, the remaining of the proof of Lemma 3.4 still holds and we get (3.64). \square

Lemma 3.8 *Let the structural assumptions of Lemma 3.7 be satisfied. If u is a positive radial solution of (1.1) in B_R^c , such that $r^\gamma u(r) \rightarrow 0$ when $r \rightarrow \infty$, then necessarily $q > \frac{N}{N-1}$ and the following alternative holds:*

1- either $q < 2$ and

$$\lim_{r \rightarrow \infty} r^\beta u(r) = \xi_M, \quad (3.65)$$

where we recall that ξ_M is defined in (1.27),

2- or $N > 2$ and

$$\lim_{r \rightarrow \infty} r^{N-2} u(r) = k > 0. \quad (3.66)$$

Proof. Since $\mathcal{R}_q^M u \leq 0$, we can apply Lemma 3.3-(3) provided $q > \frac{N}{N-1}$. If this holds the following estimate from below of u holds:

either $q < 2$ and $\liminf_{r \rightarrow \infty} r^\beta u(r) \geq \xi_M$,

or $N > 2$ and $\lim_{r \rightarrow \infty} r^{N-2} u(r) = k \geq 0$.

1- We first prove that $r^\beta u(r)$ is bounded and we recall that $S(t) = \frac{Y(t)}{X(t)}$ denotes the slope function.

1-(i) If $S(t) \rightarrow 0$ as $t \rightarrow \infty$, then for any $\epsilon > 0$, $r \mapsto r^\epsilon u(r)$ is nondecreasing. Hence $u(r) \geq c_\epsilon r^{-\epsilon}$ for $r \geq R$, for some $c_\epsilon > 0$. This contradicts Proposition A.1

1-(ii) If $\liminf_{t \rightarrow \infty} S(t) = m > 0$. Then there exists $t_0 > \ln R$ and $m_0 > 0$ such that $S(t) \geq m_0$ on $[t_0, \infty)$. Hence $Y^q(t) \geq m_0^q X^q(t)$ for $t \geq t_0$ and $u^p = o(|u'|^q)$ as $r \rightarrow \infty$. Using Lemma 3.3-(3)-(4) we infer that $q > \frac{N}{N-1}$ and (3.65) or (3.66) holds, and in both cases $r^\beta u(r)$ is bounded.

1-(iii) If S satisfies $0 = \liminf_{t \rightarrow \infty} S(t) < \limsup_{t \rightarrow \infty} S(t) = \Sigma \in (0, \infty]$. There exists an increasing sequence $\{t_n\}$ tending to infinity of local maximum of $S(t)$. As in the

proof of Lemma 3.6-(3) we obtain that $N > 2$ and $u(r) \leq Cr^{2-N}$.

If $q \geq \frac{N}{N-1}$, equivalently $\beta \leq N - 2$, then $u(r) \leq Cr^{2-N} \leq Cr^{-\beta}$ for $r \geq 1$.

If $q < \frac{N}{N-1}$, we have from (3.6) and $\eta = \xi S$ that $\xi_t = \xi(\beta - S) > \xi(\beta - N + 2) \geq 0$. Hence $\xi(t)$ is increasing with limit $\xi_\infty \leq \infty$. Since at the points t_n of local maximum of $S(t)$, we also have (3.48), we obtain the implication

$$\xi_n^{q-1} S_n^{q-1} = \frac{N-2-S_n}{M(1-\epsilon'_n)} \implies \xi_\infty^{q-1} \Sigma^{q-1} = \frac{N-2-\Sigma}{M}. \quad (3.67)$$

Hence ξ_∞ is finite, which implies again that $r^\beta u(r)$ is bounded.

2- Convergence. Since $r^\beta u(r)$ is bounded, the trajectory $\{(\xi(t), \eta(t))_{t \geq \ln R}\}$ endows this property, and since $\sigma < 0$, its omega-limit set at infinity is non-empty, compact, connected and it is a subset of the nonnegative stationary solutions of (3.6).

If $q \leq \frac{N}{N-1}$ the set is reduced to $(0, 0)$. Since $\kappa \leq 0$, we deduce from (3.8) that $\xi(t)$ is monotone decreasing. It follows from (3.6) that $S(t) > \beta$, hence $u^p = o(|u'|^q)$ as in 1-(ii) and by Lemma 3.3-(3)-(4) necessarily $q > \frac{N}{N-1}$, contradiction.

If $q > \frac{N}{N-1}$, then either $(\xi(t), \eta(t))$ converges to (ξ_M, η_M) or it converges to $(0, 0)$, in which case $\lim_{r \rightarrow \infty} r^{N-2} u(r) = k \geq 0$ by Lemma 3.3. The function u is bounded from below in B_R^c by the solution v of

$$\begin{aligned} -\Delta v + v^p &= 0 && \text{in } B_R^c \\ v &= \min_{|x|=R} u(x) && \text{on } \partial B_R \end{aligned}$$

Since $\lim_{r \rightarrow \infty} r^{N-2} v(r) = k' > 0$ and $k \geq k' > 0$, this ends the proof. \square

Proof of Theorem 1.10. In all the cases, the basic convergence (1.36) holds true from Lemma 3.7. If the limit of $r^\gamma u(r)$ is zero, then necessarily $\frac{N}{N-1} < q < 2$ and we have (1.37) or (1.38). \square

3.5 Solutions of eikonal type

In order to study the properties of solutions of eikonal type we first give some asymptotic expansion results.

Lemma 3.9 *Let $M > 0$, $\frac{2p}{p+1} < q < p$ (resp. $1 < q < \frac{2p}{p+1}$) and $\theta \neq 0$ (see (1.25) for the definition of θ). If (X, Y) is a solution of (3.11) which converges to (X_M, Y_M) when $t \rightarrow -\infty$ (resp. $t \rightarrow \infty$), then $t \mapsto X(t) - X_M$ has a constant sign for $|t|$ large enough. Furthermore*

$$X(t) = X_M + \frac{\theta \gamma^{1-q} X_M^{2-q}}{p(q-1)M} e^{\frac{\sigma t}{p-q}} + O\left(e^{\frac{2\sigma t}{p-q}}\right) \text{ as } t \rightarrow -\infty \text{ (resp. } t \rightarrow \infty). \quad (3.68)$$

Equivalently, with $u(r) = r^{-\frac{q}{p-q}} X(t)$,

$$u(r) = X_M r^{-\frac{q}{p-q}} + \frac{\theta \gamma^{1-q} X_M^{2-q}}{p(q-1)M} r^{-\frac{p(2-q)}{p-q}} + O\left(r^{-\frac{(2p+1)q-2p}{p-q}}\right) \text{ as } r \rightarrow 0 \text{ (resp. } r \rightarrow \infty). \quad (3.69)$$

Proof. (i) *Expansion of $MY^q - X^p$.* Set

$$\Psi(t) = Y_t - \theta Y = e^{-\frac{\sigma t}{p-q}}(MY^q - X^p).$$

Then

$$\Psi_t = e^{-\frac{\sigma t}{p-q}} \left(-\frac{\sigma}{p-q}(MY^q - X^p) + MqY^{q-1}(\theta Y + \Psi) - pX^{p-1}(\gamma X - Y) \right).$$

If Ψ is not monotone, one has at the local extremum t_n of Ψ , denoting $\Psi_n = \Psi(t_n)$, $X_n = X(t_n)$ and $Y_n = Y(t_n)$,

$$MqY_n^{q-1}\Psi_n = -Mq\theta Y_n^q + pX_n^{p-1}(\gamma X_n - Y_n) + \frac{\sigma}{p-q}(MY_n^q - X_n^p).$$

But $\lim_{t \rightarrow -\infty} (X(t), Y(t)) = (X_M, Y_M)$, then $pX_n^{p-1}(\gamma X_n - Y_n) + \frac{\sigma}{p-q}(MY_n^q - X_n^p) \rightarrow 0$ when $t_n \rightarrow -\infty$. Therefore $\Psi_n \rightarrow -\theta\gamma X_M$. Since the limit is valid for local minima or maxima it follows that $\lim_{t \rightarrow -\infty} \Psi(t) = -\theta\gamma X_M$.

If Ψ is monotone, then $Y_t(t) - \theta Y(t)$ admits a limit when $t \rightarrow -\infty$. Since $Y(t) \rightarrow Y_M$, it follows that $Y_t(t)$ has a also a limit at $-\infty$ and the only possible one is 0. Hence $\Psi(t) \rightarrow -\theta\gamma X_M$. In both case it yields, since $\theta \neq 0$,

$$MY(t)^q - X(t)^p = -\theta\gamma X_M e^{\frac{\sigma t}{p-q}}(1 + o(1)) \quad \text{as } t \rightarrow -\infty. \quad (3.70)$$

(ii) *We claim that $X - X_M$ has a constant sign.* If X is nondecreasing (resp. nonincreasing) then $X(t) \geq X_M$ (resp. $X(t) \leq X_M$) for $t \leq 0$. Actually the inequality is strict, otherwise, if there is some t_0 such that $X(t_0) = X_M$, we would have $X_t(t) = 0$ for $t \leq t_0$ and $X(t) = X_M$ for $t \leq t_0$. If $\theta \neq 0$ this contradicts the fact that $M\gamma^q X_M^q - X_M^p = 0$. If $\theta > 0$ we deduce from (3.12) that if $X(t_n)$ is a local minimum we have

$$e^{-\frac{\sigma t_n}{p-q}}(M\gamma^q X^q(t_n) - X^p(t_n)) = -\theta X(t_n) - X_{tt}(t_n) < 0 \implies X(t_n) > X_M.$$

This implies that $X(t) > X_M$. Similarly, if $\theta < 0$ we get $X(t) < X_M$.

(iii) *Asymptotic expansion.* We write $X = X_M(1 + w)$ and $Y = \gamma X_M(1 + z)$. Then

$$\begin{aligned} w_t &= \gamma(w - z) \\ z_t &= \theta(1 + z) + M\gamma^{q-1}X_M^{q-1}e^{-\frac{\sigma t}{p-q}}((1 + z)^q - (1 + w)^p). \end{aligned} \quad (3.71)$$

There holds

$$(1 + z)^q - (1 + w)^p = qz - pw + \phi(z) - \psi(w),$$

where $\phi(z) = (1 + z)^q - 1 - qz$ and $\psi(w) = (1 + w)^p - 1 - pw$, therefore $0 \leq \phi(z) \leq cz^2$ and $0 \leq \psi(w) \leq cw^2$ for $t \leq t^*$. Next, from (3.71),

$$qz + \phi(z) = pw + \psi(w) + a_M e^{\frac{\sigma t}{p-q}}(z_t - \theta(1 + z)),$$

where $a_M = M^{-1}(\gamma X_M)^{1-q}$, and $z(t) \rightarrow 0$ and $z_t(t) \rightarrow 0$ when $t \rightarrow -\infty$. Therefore the previous identity becomes

$$qz + \phi(z) = pw + \psi(w) - \theta a_M e^{\frac{\sigma t}{p-q}}(1 + h(t)), \quad (3.72)$$

where $h(t) \rightarrow 0$ when $t \rightarrow -\infty$. Next

$$\begin{aligned} qz \leq qz + \phi(z) &\leq \begin{cases} qz(1 + \epsilon) & \text{if } z > 0 \\ qz(1 - \epsilon) & \text{if } z \leq 0 \end{cases} := qz(1 + \epsilon_z) \\ pw \leq pw + \psi(w) &\leq pw(1 + \epsilon_w), \end{aligned}$$

where $\epsilon_z = \epsilon \text{sign}(z)$ and $\epsilon_w = w \text{sign}(w)$. It follows from (3.72) that

$$\begin{aligned} qz - pw(1 + \epsilon_w) + \theta a_M e^{\frac{\sigma t}{p-q}}(1 + h(t)) &\leq 0 \\ qz(1 + \epsilon_z) - pw + \theta a_M e^{\frac{\sigma t}{p-q}}(1 + h(t)) &\geq 0. \end{aligned} \quad (3.73)$$

This leads to the following two inequalities verified by w_t

$$\begin{aligned} w_t = \gamma(w - z) &= \frac{1}{p-q}(qw - qz) \geq \frac{1}{p-q} \left(qw - pw(1 + \epsilon_w) + \theta a_M e^{\frac{\sigma t}{p-q}}(1 + h) \right) \\ &\geq -w \left(1 + \frac{p}{p-q} \epsilon_w \right) + \frac{\theta a_M}{p-q} e^{\frac{\sigma t}{p-q}}(1 + h), \end{aligned}$$

and

$$\begin{aligned} w_t &\leq \frac{q}{p-q} \left(w + \frac{1}{q(1 + \epsilon_z)}(-pw + \theta a_M e^{\frac{\sigma t}{p-q}}(1 + h)) \right) \\ &\leq w \left(-1 + \frac{p}{p-q} \epsilon_z \right) + \frac{\theta a_M}{(p-q)(1 + \epsilon_z)} e^{\frac{\sigma t}{p-q}}(1 + h), \end{aligned}$$

and we know from (i) that $w(t)$ keeps a constant sign when $t \rightarrow -\infty$. We deduce from the above inequalities that if $\theta < 0$ the function $t \mapsto e^{(1-\epsilon)t}w(t)$ is decreasing for some $\epsilon > 0$ and tends to 0, hence it is negative, while, if $\theta > 0$, $t \mapsto e^{(1+\epsilon)t}w(t)$ is increasing for another $\epsilon > 0$ and tends to 0, hence it is positive. Then, we can summarize as follows, with a new $\epsilon > 0$

$$(1 - \epsilon) \left(\theta^{-1}w - \frac{a_M}{p-q} e^{\frac{\sigma t}{p-q}} \right) \leq -(\theta^{-1}w)_t \leq (1 + \epsilon) \left(\theta^{-1}w - \frac{a_M}{p-q} e^{\frac{\sigma t}{p-q}} \right). \quad (3.74)$$

As we have $1 \pm \epsilon + \frac{\sigma}{p-q} = \frac{p(q-1)}{p-q} \pm \epsilon$, the function $t \mapsto e^{(1+\epsilon)t} \left(\theta^{-1}w - \frac{a_M}{p(q-1)+\epsilon} e^{\frac{\sigma t}{p-q}} \right)$ is increasing and tend to 0 as $t \rightarrow -\infty$. Hence it is positive. In the same way, the function $t \mapsto e^{(1-\epsilon)t} \left(\theta^{-1}w - \frac{a_M}{p(q-1)-\epsilon} e^{\frac{\sigma t}{p-q}} \right)$ is decreasing, tends to 0 hence it is negative. Therefore we infer that

$$w(t) = \frac{\theta a_M}{p(q-1)} e^{\frac{\sigma t}{p-q}}(1 + o(1)). \quad (3.75)$$

This implies $\psi(w) = O\left(e^{\frac{2\sigma t}{p-q}}\right)$. From (3.72), $z = O\left(e^{\frac{\sigma t}{p-q}}\right)$, then $\psi(z) = O\left(e^{\frac{2\sigma t}{p-q}}\right)$ and $\epsilon_z = O\left(e^{\frac{\sigma t}{p-q}}\right)$. Since $X - X_M = X_M w = \gamma Y_M w$, we deduce (3.68). Notice also that from (3.72) there holds

$$z = \frac{(2-q)\theta a_M}{q(q-1)} e^{\frac{\sigma t}{p-q}} + o\left(e^{\frac{\sigma t}{p-q}}\right),$$

hence

$$w_t = \gamma(w(t) - z(t)) = \frac{\sigma\theta a_M}{p(p-q)(q-1)} e^{\frac{\sigma t}{p-q}} (1 + o(1)). \quad (3.76)$$

In particular X_t has the sign of θ , and therefore X is monotone. \square

Remark. If $q = 2$ we obtain

$$X(t) = (M\gamma^2)^{\frac{1}{p-2}} + \frac{2(N-1) - (N-2)p}{2pM} e^{\frac{2t}{p-2}} + O\left(e^{\frac{4t}{p-2}}\right) \quad \text{as } t \rightarrow -\infty, \quad (3.77)$$

so we recover the result of [20].

3.6 Local or global existence results

3.6.1 The systems of order 3

Since $q \neq \frac{2p}{p+1}$, we can perform the transformation T_ℓ and assume that $M = 1$. For proving the existence of solutions to (1.1) there are essentially three methods: the methods of sub and super solutions which has already been developed in Section 2.3, the method of fixed points, and the use of a specific autonomous system of order 3. This last method appears to be entirely new and we explain it below. This system uses the variables $(X, \xi, S) \in \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+$,

$$\begin{aligned} X_t &= X(\gamma - S) \\ \xi_t &= \xi(\beta - S) \\ S_t &= S(S + 2 - N) + \xi^{q-1}(MS^q - X^{p-q}). \end{aligned} \quad (3.78)$$

Lemma 3.10 *Let $1 < q < p$ with $q \neq \frac{2p}{p+1}$ and $M > 0$. If u is a decreasing positive solution of (2.12), then*

$$(X(t), \xi(t), S(t)) = \left(r^\gamma u(r), r^\beta u(r), r \frac{|u'(r)|}{u(r)} \right) \quad \text{with } t = \ln r, \quad (3.79)$$

satisfies (3.78). Conversely, to each trajectory of (3.78) in $\mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+$ corresponds a unique solution of (2.12).

Proof. Let u be a decreasing solution of (2.12). We recall that (X, Y) are solutions of (3.11), $S = \frac{Y}{X}$ and $\xi(t) = r^\beta u(r)$ with $t = \ln r$. Then (X, S) satisfies the following system which is equivalent to (3.11),

$$\begin{aligned} X_t &= X(\gamma - S) \\ S_t &= S(S + 2 - N) + e^{-\frac{\sigma t}{p-q}} X^{q-1}(MS^q - X^{p-q}). \end{aligned} \quad (3.80)$$

Using (3.14) we have that $\xi^{q-1} = e^{-\frac{\sigma t}{p-q}} X^{q-1}$. Since by computation $\xi_t = \xi(\beta - S)$, we deduce that (X, ξ, S) satisfies (3.78).

Conversely, let $(X, \xi, S) \in \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+$ be a solution of (3.78), then

$$\frac{\xi_t}{\xi} - \frac{X_t}{X} = \beta - \gamma = -\frac{\sigma}{(p-q)(q-1)}.$$

Hence $\xi(t) = be^{-\frac{\sigma t}{(p-q)(q-1)}} X(t)$ for some $b > 0$. If we set $a = -\frac{(p-q)(q-1)}{\sigma} \ln b$, we see that

$$\xi(t) = e^{-\frac{\sigma(t+a)}{(p-q)(q-1)}} X(t).$$

Hence

$$\begin{aligned} X_t &= X(\gamma - S) \\ S_t &= S(S + 2 - N) + e^{-\frac{\sigma(t+a)}{p-q}} X^{q-1} (MS^q - X^{p-q}). \end{aligned} \quad (3.81)$$

Setting $\tau = t + a$, $X^{(a)}(\tau), S^{(a)}(\tau) = (X(t), S(t)) = (X(\tau - a), S(\tau - a))$, then

$$\begin{aligned} X_\tau^{(a)} &= X^{(a)}(\gamma - S^{(a)}) \\ S_\tau^{(a)} &= S^{(a)}(S^{(a)} + 2 - N) + e^{-\frac{\sigma\tau}{p-q}} (X^{(a)})^{q-1} (M(S^{(a)})^q - (X^{(a)})^{p-q}). \end{aligned} \quad (3.82)$$

Then the function $\rho \mapsto u^{(a)}(\rho) = \rho^{-\gamma} X^{(a)}(\ln \rho)$ satisfies (2.12). Let (X, ξ, S) and $(\tilde{X}, \tilde{\xi}, \tilde{S})$ be two solutions of (3.78). Then there exist a, \tilde{a} such that

$$\xi(t) = e^{-\frac{\sigma(t+a)}{(p-q)(q-1)}} X(t) \quad \text{and} \quad \tilde{\xi}(t) = e^{-\frac{\sigma(t+\tilde{a})}{(p-q)(q-1)}} \tilde{X}(t),$$

and

$$\begin{aligned} u^{(a)}(\rho) &= \rho^{-\gamma} X^{(a)}(\ln \rho) = \rho^{-\gamma} X(\ln \rho - a) \\ u^{(\tilde{a})}(\rho) &= \rho^{-\gamma} \tilde{X}(\ln \rho - \tilde{a}). \end{aligned}$$

If (X, ξ, S) and $(\tilde{X}, \tilde{\xi}, \tilde{S})$ correspond to the same trajectory, there exists $h \in \mathbb{R}$ such that $(\tilde{X}, \tilde{\xi}, \tilde{S})(t) = (X, \xi, S)(t + h)$ for all t , thus

$$\xi(t + h) = e^{-\frac{\sigma(t+a)}{(p-q)(q-1)}} X(t + h) = e^{-\frac{\sigma(t+a+h)}{(p-q)(q-1)}} X(t + h).$$

Therefore $\tilde{a} = a + h$. Hence

$$u^{(\tilde{a})}(\rho) = \rho^{-\gamma} \tilde{X}(\ln \rho - \tilde{a}) = \rho^{-\gamma} \tilde{X}(\ln \rho - a - h) = \rho^{-\gamma} X(\ln \rho - a) = u^{(a)}(\rho).$$

In conclusion, there is a one to one correspondence between the trajectories of (3.78) and the solutions of (2.12). \square

Remark. Using the relation (3.16) one can see that (3.78) is equivalent to the following system in the variables (x, ξ, S) ,

$$\begin{aligned} x_t &= x(\alpha - S) \\ \xi_t &= \xi(\beta - S) \\ S_t &= S(S + 2 - N) + \xi^{q-1} MS^q - x^{p-1}. \end{aligned} \quad (3.83)$$

This system is particularly suitable for constructing local solutions in $r^{-\alpha}$ or r^{2-N} , in particular when $r \rightarrow \infty$, in the case $q > \frac{2p}{p+1}$.

3.6.2 Singular solutions of eikonal type

Proof of Theorem 1.11. We recall that these solutions of eikonal type are the solutions which behave like $r^{-\gamma}$ near 0 or ∞ . For $c > 0$ and $A \geq 0$ we set $u_{c,A}(r) = cr^{-\gamma} + A$ and $u_c = u_{c,0}$. Then there exist $a, b > 0$ depending on p such that

$$c^p r^{-\gamma p} + aA^p + apc^{p-1} Ar^{-\gamma(p-1)} \leq u_{c,A}^p(r) \leq c^p r^{-\gamma p} + bA^p + bpc^{p-1} Ar^{-\gamma(p-1)}.$$

(i) *Subsolutions.* If $u_c(r) = cr^{-\gamma}$, then

$$\begin{aligned} \frac{1}{c} \mathcal{L}_{p,q}^M u_c &= -\gamma(\gamma + 2 - N)r^{-\gamma-2} + c^{p-1}r^{-\gamma p} - Mc^{q-1}\gamma^q r^{-(\gamma+1)q} \\ &= r^{-\frac{pq}{p-q}} \left(c^{p-1} - c^{q-1} X_M^{p-q} - \gamma\theta r^{\frac{\sigma}{p-q}} \right). \end{aligned} \quad (3.84)$$

Set $\Phi(c) = c^{p-1} - c^{q-1} X_M^{p-q}$. Then $\Phi(X_M) = 0$ and Φ achieves its minimum at $c_m = \left(\frac{q-1}{p-1} X_M^{p-q} \right)^{\frac{1}{p-q}}$ with minimal value $\Phi(c_m) = -\frac{p-q}{q-1} \left(\frac{q-1}{p-1} X_M^{p-q} \right)^{\frac{p-1}{p-q}}$. Notice that if $\theta = 0$ i.e. $q = \frac{(N-2)p}{N-1}$, then we find the explicit solution $u_{M,p}^*$ defined in (3.13).

(i-a) If $N = 1$ or $N = 2$, or if $N \geq 3$ and $q > \frac{N-2}{N-1}p$, then $\theta > 0$ and u_c is a subsolution in $\mathbb{R}^N \setminus \{0\}$ provided $c \leq X_M$.

(i-b) If $q > \frac{2p}{p+1}$ and $\theta < 0$ there exists $r_1 > 0$ such that u_{c_m} is a subsolution in $B_{r_1} \setminus \{0\}$. Hence $\tilde{u}_{c_m} = c_m(r^{-\gamma} - r_1^{-\gamma})_+$ is a subsolution in $\mathbb{R}^N \setminus \{0\}$.

(ii) *Supersolutions.* We have

$$\frac{1}{c} \mathcal{L}_{p,q}^M u_{c,A} \geq r^{-\frac{pq}{p-q}} \left(\Phi(c) - \gamma\theta r^{\frac{\sigma}{p-q}} \right) + \frac{aA^p}{c} + apc^{p-2} Ar^{-\gamma(p-1)} \quad (3.85)$$

(ii-a) If $\theta < 0$, then for $c \geq X_M$ and any $A \geq 0$, $u_{c,A}$ is a supersolution in $\mathbb{R}^N \setminus \{0\}$.

(ii-b) If $q > \frac{2p}{p+1}$ and $\theta > 0$, then for any $R > 0$ we take $c > X_M$ such that $\Phi(c) \geq \gamma\theta R^{\frac{\sigma}{p-q}}$, hence $\mathcal{L}_{p,q}^M u_{c,A} \geq 0$ in $B_R \setminus \{0\}$. Since $-\frac{pq}{p-q} + \frac{\sigma}{p-q} = \frac{q-2p}{p-q} < 0$, we take $A > 0$ such that $aA^p \geq \gamma\theta c R^{-\gamma-2}$, hence $\mathcal{L}_{p,q}^M u_{c,A} \geq 0$ in B_R^c . Consequently $u_{c,A}$ is a supersolution in $\mathbb{R}^N \setminus \{0\}$.

(ii-c) If $q < \frac{2p}{p+1}$ and $\theta > 0$, then we can take c such that $\Phi(c) \geq \gamma\theta R^{\frac{\sigma}{p-q}}$ and obtain that $u_{c,A}$ is a supersolution in B_R^c .

(iii) *Proof of statements 1 and 2.*

If $q > \frac{2p}{p+1}$ and whatever is the sign of θ there exist $c_m \leq c < X_M < c'$ and $A > 0$ such that $u_{c',A}$ is a supersolution in $\mathbb{R}^N \setminus \{0\}$ larger than the subsolution u_c . By [25, Theorem 1.4.5] there exists a radial solution u in $\mathbb{R}^N \setminus \{0\}$ satisfying $u_c \leq u \leq u_{c',A}$. Its behaviour at infinity is given by Theorem 1.9. This solution is decreasing by the maximum principle and it is unique by Theorem 4.6-(3).

The existence of a solution in a bounded domain Ω containing 0 and vanishing on $\partial\Omega$ satisfying (1.26) follows by Theorem 1.14 which is proved in Section 4. So we

deduce statement 1.

If $1 < q < \frac{2p}{p+1}$ and $\theta > 0$, one has a supersolution $u_{c',A}$ in B_R^c and a subsolution u_c in $\mathbb{R}^N \setminus \{0\}$. Up to increasing the value of A one has again a supersolution $u_{c',A}$ larger than the subsolution u_c . Hence there exists a solution u in between satisfying (1.36) which proves statement 2. \square

3.6.3 Riccati type singular solutions

Proof of Theorem 1.12. We recall that the Riccati equation (1.20) admits the radial solution $\xi_M |x|^{-\beta}$ if and only if $\frac{N}{N-1} < q < 2$. This function is a supersolution of (1.1) in $\mathbb{R}^N \setminus \{0\}$.

1- *Local existence in a neighborhood of 0.* Since $q > \frac{N}{N-1}$ the system (3.78) in variables (X, ξ, S) admits the equilibria $(0, 0, 0)$, $(0, 0, N-2)$ and $(0, \xi_M, \beta)$. Our aim is to construct local radial solutions of (2.12) satisfying $\lim_{r \rightarrow 0} r^\beta u(r) = \xi_M$ and $\lim_{r \rightarrow 0} r^{\beta+1} |u'(r)| = \eta_M = \beta \xi_M$, equivalently

$$\lim_{t \rightarrow -\infty} (X(t), \xi(t), S(t)) = (0, \xi_M, \beta). \quad (3.86)$$

Conversely, any solution (X, ξ, S) satisfying (3.86) corresponds to a solution u satisfying $\lim_{r \rightarrow 0} (r^\beta u(r), r^{\beta+1} |u'(r)|) = (\xi_M, \beta \xi_M)$. The system (3.78) may be singular at $\xi_M = 0$ or at $X = 0$; hence we desingularize it by setting $\hat{X} = X^{p-q}$ and $\hat{\xi} = \xi^{q-1}$. Then $(\hat{X}, \hat{\xi}, S)$ satisfies

$$\begin{aligned} \hat{X}_t &= (p-q)\hat{X}(\gamma - S) \\ \hat{\xi}_t &= (q-1)\hat{\xi}(\beta - S) \\ S_t &= S(S+2-N) + \hat{\xi}(MS^q - \hat{X}). \end{aligned} \quad (3.87)$$

So we are led to study solutions in a neighborhood of the equilibrium $(0, \hat{\xi}_M, \beta)$ where $\hat{\xi}_M = \xi_M^{q-1} = \frac{\kappa}{M\beta^{q-1}}$. We set $\hat{\xi} = \hat{\xi}_M + \bar{\xi}$, $\hat{X} = \bar{X}$ and $S = \beta + \bar{S}$ in order to reduce the study at $(0, 0, 0)$, and $(\bar{\xi}, \bar{X}, \bar{S})$ satisfies the following linearized system

$$\begin{aligned} \bar{X}_t &= \frac{\sigma}{q-1} \bar{X} \\ \bar{\xi}_t &= -(q-1)\hat{\xi}_M \bar{S} \\ \bar{S}_t &= -\hat{\xi}_M \bar{X} + M\beta^q \bar{\xi} + (\beta + \kappa(q-1))\bar{S}. \end{aligned} \quad (3.88)$$

If we denote by \mathcal{A} the matrix of this system, then its characteristic values are the roots of the polynomial

$$\det(\mathcal{A} - \mu I) = (\mu - \mu_1)(\mu - \mu_2)(\mu - \mu_3), \quad (3.89)$$

with $\mu_1 = \frac{\sigma}{q-1}$, $\mu_2 = \beta$ and $\mu_3 = (q-1)\kappa = (N-1)q - N$. Since $q > \max\left\{\frac{2p}{p+1}, \frac{N}{N-1}\right\}$ all the eigenvalues are positive. We find that

$$\mathbf{u}_2 = \left(0, 1, -\frac{M\beta^q}{\kappa(q-1)}\right) \text{ and } \mathbf{u}_3 = (0, 1, -M\beta^{q-1})$$

are eigenvectors corresponding to μ_2 and μ_3 respectively. If $\mu_1 \neq \mu_2$ and $\mu_1 \neq \mu_3$, we can take for eigenvector corresponding to μ_1 the vector $\mathbf{u}_1 = (1, b, c)$ for some real numbers b and c . Actually

$$b = -\frac{\mu_3}{M\mu_2^{q-1}\mu_1} \quad \text{where} \quad c = -\frac{\kappa\mu_1}{M\mu_2^{q-1}(\mu_1 - \mu_2)(\mu_1 - \mu_3)}.$$

Then there exists one trajectory of (3.87) with $X(t) > 0$ when $t \rightarrow -\infty$ such that $\xi(t) = \xi_M + O(e^{\frac{\sigma t}{q-1}})$ when $t \rightarrow -\infty$. Hence there exists at least one solution u of (2.12) such that $u(r) = r^{-\beta}\xi_M + Cr^{-\beta+(N-1)q-N}(1 + o(1))$ when $r \rightarrow 0$.

2- Local existence at infinity. Here we assume $\frac{N}{N-1} < q < \frac{2p}{p+1}$. Then $\mu_1 < 0$, $\mu_2 = \beta > 0$ and $\mu_3 = \kappa(q-1) > 0$. Then there exists a unique local trajectory which converges to $(0, \hat{\xi}, \beta)$ when $t \rightarrow \infty$, it corresponds to the stable manifold of this point. If there exists a positive solution in $\mathbb{R}^N \setminus \{0\}$, the solution can be extended as a solution in \mathbb{R}^N by [7, Theorem 1.1] since in this range of values of q one has $p > \frac{N}{N-2}$. By Proposition A.1 such a solution is identically 0. \square

Remark. Note that we have many types of trajectories converging to the origin and their geometry depends in their sign and their relative order. In this respect we denote

$$f(q) := (N-1)q - N + \frac{N - (N-2)q}{2-q}, \quad (3.90)$$

and we have

$$\begin{aligned} (i) \quad & \mu_3 = \kappa(q-1) > \mu_2 = \beta \iff q > 1 + \frac{1}{\sqrt{N-1}} \\ (ii) \quad & \mu_1 > \mu_2 \iff p < \frac{2(q-1)}{2-q} \iff q > \frac{2(p+1)}{p+2} \\ (iii) \quad & \mu_1 > \mu_3 \iff p < f(q). \end{aligned} \quad (3.91)$$

We have that $\mu_1 = \mu_2 = \mu_3$ only if $p = \frac{2}{\sqrt{N-1}-1}$ and $q = 1 + \frac{1}{\sqrt{N-1}}$, a condition which is compatible with $p > 1$ only if $2 \leq N \leq 9$.

Global (necessarily singular) solutions in $r^{-\beta}$ are difficult to construct. We give below a range of exponents in which there exists at least one.

Theorem 3.11 *Let $M > 0$, $p > 1$ and $1 < q < 2$, $q \neq 1 + \frac{1}{\sqrt{N-1}}$. If there holds*

$$p < \max \left\{ \frac{2(q-1)}{2-q}, f(q) \right\}, \quad (3.92)$$

in particular if $p < \frac{N}{N-2}$ and $q > \frac{N}{N-1}$, then there exists a positive radial solution of (1.1) defined in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{x \rightarrow 0} |x|^\beta u(x) = \xi_M. \quad (3.93)$$

Proof. The function $U(x) = \xi_M |x|^{-\beta}$ is a supersolution of (1.1) in $\mathbb{R}^N \setminus \{0\}$. We look for a subsolution under the form $\tilde{\xi}(t) = \xi_M(1 - Ae^{dt})_+$ for some $d, A > 0$. Set

$$H[\tilde{\xi}](t) = \tilde{\xi}_{tt} + D\tilde{\xi}_t - \kappa\beta\tilde{\xi} - e^{\mu_1 t}\tilde{\xi}^p + M|\beta\tilde{\xi} - \tilde{\xi}_t|^q,$$

where

$$D = \frac{Nq - N - 2}{q - 1} = \kappa - \beta.$$

Then on the interval $I_A := (-\infty, -\frac{\ln A}{d})$ one has

$$\beta\tilde{\xi} - \tilde{\xi}_t = \xi_M \left(\beta - A(\beta - d)e^{dt} \right).$$

In order $H[\tilde{\xi}] \geq 0$, one needs

$$\begin{aligned} -A(d^2 + Dd - \kappa\beta)e^{dt} + M\xi_M^{q-1}\beta^q \left(1 - A\frac{\beta-d}{\beta}e^{dt} \right)^q - \kappa\beta \\ - \xi_M^{p-1}e^{\mu_1 t} (1 - Ae^{dt})^p \geq 0. \end{aligned}$$

Since $M\xi_M^{q-1}\beta^q = \kappa\beta$, if we set $Z = Ae^{dt}$, then $0 < Z \leq 1$ on I_A and the previous inequality to be verified becomes

$$\frac{\xi_M^{p-1}}{A}e^{(\mu_1-d)t}Z(1-Z)^p \leq -(d^2 + Dd - \kappa\beta)Z + \kappa\beta \left(1 - \frac{\beta-d}{\beta}Z \right)^q - \kappa\beta.$$

We first impose $d \leq \mu_1$, then $\frac{e^{(\mu_1-d)t}}{A} \leq A^{-\frac{\mu_1}{d}}$ on I_A . We set

$$Q(Z) = \kappa\beta \left(1 - \frac{\beta-d}{\beta}Z \right)^q - \left(d^2 + Dd - \kappa\beta + \xi_M^{p-1}A^{-\frac{\mu_1}{d}} \right) Z - \kappa\beta. \quad (3.94)$$

Then

$$Q'(Z) = -q\kappa(\beta-d) \left(1 - \frac{\beta-d}{\beta}Z \right)^{q-1} - \left(d^2 + Dd - \kappa\beta + \xi_M^{p-1}A^{-\frac{\mu_1}{d}} \right),$$

and

$$Q''(Z) = \frac{\kappa(\beta-d)^2q(q-1)}{\beta} \left(1 - \frac{\beta-d}{\beta}Z \right)^{q-2}.$$

Since $\kappa > 0$, Q is convex on $[0, 1]$. Furthermore $Q(0) = 0$. Hence $H(\tilde{\xi}) \geq 0$ if $Q'(0) \geq 0$.

$$Q'(0) = -q\kappa(\beta-d) - \left(d^2 + Dd - \kappa\beta + \xi_M^{p-1}A^{-\frac{\mu_1}{d}} \right) \quad (3.95)$$

Replacing D by its value, (3.95) will be achieved, provided A is large enough, if

$$-d^2 + ((q-1)\kappa + \beta)d - (q-1)\kappa\beta = -(d - \mu_2)(d - \mu_3) > 0. \quad (3.96)$$

The condition is that $\mu_1 \geq d$ with d satisfying (3.96). It necessitates $\mu_2 \neq \mu_3$, equivalently $q \neq 1 + \frac{1}{\sqrt{N-1}}$, and

(i) either $\mu_2 < \inf\{\mu_1, \mu_3\}$, then we can choose any $d \in (\mu_2, \inf\{\mu_1, \mu_3\})$,
(ii) or $\mu_3 < \inf\{\mu_1, \mu_2\}$, then we can choose any $d \in (\mu_2, \inf\{\mu_1, \mu_2\})$.
These conditions are satisfied if $\mu_1 > \inf\{\mu_2, \mu_3\}$ which is equivalent to (3.92). If one of the above conditions is satisfied, it follows by [25, Corollary 1.4.5] that there exists a radial positive solution u of (1.1) in $\mathbb{R}^N \setminus \{0\}$ which satisfies

$$\xi_M (1 - A|x|^d)_+ |x|^{-\beta} \leq u(x) \leq \xi_M |x|^{-\beta} \quad \text{for all } x \in \mathbb{R}^N \setminus \{0\}. \quad (3.97)$$

□

Remark. Condition (3.92) is equivalent to

$$\begin{aligned} (i) \quad 1 < p < f(q) & \quad \text{if } 1 < q < 1 + \frac{1}{\sqrt{N-1}} \\ (ii) \quad 1 < p < \frac{2(q-1)}{2-q} & \quad \text{if } 1 + \frac{1}{\sqrt{N-1}} < q < 2. \end{aligned} \quad (3.98)$$

Condition (ii) is equivalent to

$$\frac{2(p+1)}{p+2} < q < 2. \quad (3.99)$$

Note that the nature of the variations of the function $p = f(q)$ differs according to the value of N .

If $N = 3$ or 4 , f is increasing and onto from $(\frac{3}{2}, 2)$ to $(3, \infty)$ when $N = 3$ and from $(\frac{4}{3}, 2)$ to $(2, \infty)$ when $N = 4$.

If $N \geq 5$, f achieves a maximal value \tilde{p} for $q = \tilde{q}$ with

$$\tilde{q} = 2 - \sqrt{\frac{N-4}{N-1}} \quad \text{and} \quad \tilde{p} = 2 \left(N - 2 - \sqrt{(N-4)(N-1)} \right). \quad (3.100)$$

In particular one has

$$f\left(\frac{N}{N-1}\right) = f\left(\frac{N}{N-2}\right) = \frac{N}{N-2}.$$

3.6.4 Emden-Fowler type singular solutions

Proof of Theorem 1.13-(1). Since $1 < p < \frac{N}{N-2}$ the function $x \mapsto U_{x_0}(x) = x_0|x|^{-\alpha}$ is a subsolution of (1.1) in $\mathbb{R}^N \setminus \{0\}$. In order $x \mapsto U_C(x) := C|x|^{-\alpha}$ to be a supersolution, one needs

$$C^{p-1} \geq x_0^{p-1} + \alpha^q C^{q-1} M |x|^{-\frac{\sigma}{p-1}}. \quad (3.101)$$

The function $C \mapsto C^{p-q} - x_0^{p-1} C^{1-q}$ is increasing and onto from $[x_0, \infty)$ to $[x_0, \infty)$. Hence there exists $C > x_0$ such that $\alpha^q C^{q-1} M = C^{p-1} - x_0^{p-1}$. For such a value we have that

$$\mathcal{L}_{p,q}^M U_C = M \alpha^q (1 - r^{-\frac{\sigma}{p-1}}) r^{-\frac{2p}{p-1}}.$$

Since $\sigma < 0$ the function U_C is a supersolution of (1.1) in $B_1 \setminus \{0\}$. For $A > 0$ we set $U_{C,A} = U_C + A$. Then

$$\mathcal{L}_{p,q}^M U_{C,A} = \mathcal{L}_{p,q}^M U_C + U_{C,A}^p - U_C^p \geq A^p + M\alpha^q(1 - |x|^{-\frac{\sigma}{p-1}})|x|^{-\frac{2p}{p-1}}. \quad (3.102)$$

Clearly $\mathcal{L}_{p,q}^M U_{C,A} \geq 0$ in $B_1 \setminus \{0\}$, and for $|x| > 1$, one has

$$\mathcal{L}_{p,q}^M U_{C,A} \geq A^p - M\alpha^q|x|^{-\frac{(p+1)q}{p-1}} \geq A^p - M\alpha^q.$$

Therefore, if $A = M\alpha^{\frac{q}{p}}$, the function $U_{C,A}$ is a supersolution in $\mathbb{R}^N \setminus \{0\}$. Since $U_{x_0} \leq U_{C,A}$, it follows by [25, Theorem 1.4.5] that there exists a solution u of (1.1) in $\mathbb{R}^N \setminus \{0\}$ such that $U_{x_0} \leq u \leq U_{C,A}$. Then by Theorem 1.8-(1), u satisfies (1.39)-(i), and by Theorem 1.10-(2), (1.39)-(ii) holds. Furthermore $r^\alpha u(r) \geq x_0$ for any $r > 0$. Uniqueness (not only for radial solutions) is a consequence of Theorem 4.6-(2). Obviously $|x| \mapsto u(x)$ is decreasing. Existence of a positive solution in a bounded domain Ω containing 0 is a consequence of Theorem 1.14, see Section 4. \square

Proof of Theorem 1.13-(2). It is a consequence of Theorem 1.11 and Theorem 1.9. \square

3.6.5 Solutions behaving like the Newtonian potential

There exist also solutions which behave like the Newtonian kernel at 0. They are described in the next result.

Theorem 3.12 *Let $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{N}{N-1}$. Then for any $M \geq 0$ and $k > 0$ there exists a minimal positive solution u_k of (1.1) in $\mathbb{R}^N \setminus \{0\}$ such that (1.12) holds. Furthermore it is radial and nonincreasing. If we assume $1 < q \leq \frac{2p}{p+1}$, this solution is unique among all the positive solutions.*

Proof. Proof of existence. If $M = 0$ the result is classical and for $k > 0$ we denote by v_k the solution of $\mathcal{L}_p v = 0$ in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.15). This is a natural subsolution of (1.1).

The construction of the supersolution is more involved.

(i) We first assume that $N \geq 3$ and prove that for any $k > 0$ there exists $M_k > 0$ such that for any $0 < M \leq M_k$ there exists a supersolution of (1.1) satisfying (1.12). Let $a > 0$ set

$$w_k(x) = k|x|^{2-N} + k^q|x|^{2-(N-1)q} + a$$

Then there exist $c_5, c_6 > 0$ depending on N and q such that.

$$\begin{aligned} \mathcal{L}_{p,q}^M w_k &= k^q((N-1)q-2)(N-(N-1)q)|x|^{(1-N)q} + (k|x|^{2-N} + k^q|x|^{2-(N-1)q} + a)^p \\ &\quad - M((N-2)k|x|^{1-N} + ((N-1)q-2)k^q|x|^{1-(N-1)q})^q \\ &\geq c_5 k^q|x|^{(1-N)q} + a^p - c_6 M \left(k^q|x|^{(1-N)q} + k^q|x|^{q-(N-1)q^2} \right) \\ &\geq k^q \left(c_5 - c_6 M(1 + k^{q^2-q}) \right) |x|^{(1-N)q} + a^p - k^q \left(c_6 M + k^{q^2-q} \right). \end{aligned} \quad (3.103)$$

Note that we have only used inequalities $2 \leq (N-1)q \leq N$. Set $M_k = \frac{c_5}{c_6 M(1+k^{q^2-q})}$. Then, for $M \leq M_k$ we take $a^p = k^q (c_6 M + k^{q^2-q})$ and we derive that $\mathcal{L}_{p,q}^M w_k \geq 0$. The supersolution w_k satisfies $v_k \leq k|x|^{2-N} \leq w_k$.
(ii) If $N = 2$ and for $b > 0$ we denote by ψ_k the solution of

$$-\Delta\psi + \psi^p = |x|^{-q} + 2\pi k\delta_0, \quad (3.104)$$

and we set $w_k = \psi_k + b$. Since $1 < q < 2$, $w_k = -k \ln|x|(1 + o(1))$ and $\nabla w_k = -k|x|^{-1}(1 + o(1))$ as $x \rightarrow 0$. Hence

$$0 \leq \psi_k(x) \leq c_7(-k \ln|x| + 1) \text{ and } |\nabla\psi_k(x)| \leq c_7(k+1)|x|^{-1} \text{ for } 0 < |x| \leq 1. \quad (3.105)$$

Furthermore, by Keller-Osserman technique combined with scaling method, there holds in $\mathbb{R}^2 \setminus B_1$,

$$\begin{aligned} (i) \quad 0 \leq \psi_k(x) &\leq c_8 \max \left\{ |x|^{-\alpha}, |x|^{-\frac{q}{p}} \right\}, \\ (ii) \quad |\nabla\psi_k(x)| &\leq c_8 \max \left\{ |x|^{-\frac{p+1}{p-1}}, |x|^{-\frac{p+q}{p}} \right\}. \end{aligned} \quad (3.106)$$

In the above inequalities, c_7 and c_8 are positive constants depending on p and q . Hence

$$\mathcal{L}_{p,q}^M w_k = |x|^{-q} + (\psi_k + b)^p - \psi_k^p - M|\nabla\psi_k|^q. \quad (3.107)$$

We infer

$$\mathcal{L}_{p,q}^M w_k \geq |x|^{-q} + b^p - M c_7^q k^q |x|^{-q} \text{ if } 0 < |x| \leq 1, \quad (3.108)$$

and

$$\mathcal{L}_{p,q}^M w_k \geq b^p - M c_8^q k^q \text{ if } |x| \geq 1. \quad (3.109)$$

If k is fixed, $M \leq M_k := k^{-q} c_7^{-q}$ and $b \geq M c_8^q k^q$ we conclude that w_k is a supersolution in $\mathbb{R}^2 \setminus \{0\}$ larger than v_k .

We deduce from (i) and (ii) that for any $k > 0$ there exists $M_k > 0$ such that for any $0 < M \leq M_k$ there exists a positive radial solution u_k of (1.1) satisfying (1.12). Furthermore u_k satisfies (A.1). Therefore u_k is necessarily decreasing.

End of the proof of existence. Let $q < \frac{N}{N-1}$ and q_1 such that $q < q_1 < \frac{N}{N-1}$. For $k > 0$ let $\epsilon > 0$ such that for any $0 < M' \leq \epsilon$ there exists a positive radial solution w_k to $\mathcal{L}_{q_1, M'} w = 0$ satisfying (1.12). If $M > M'$ there holds

$$M|X|^q \leq M'|X|^{q_1} + C \quad \text{for all } X \in \mathbb{R}^N,$$

where $C = \left(\frac{qM}{q_1 M'}\right)^{\frac{q}{q_1 - q}} \left(M - \frac{q}{q_1} M'\right) > 0$. Then

$$\mathcal{L}_{p,q}^M w_k = \mathcal{L}_{q_1, M'} w_k + (M' - M)|\nabla w_k^q| \geq -C$$

which implies that $w_k + C^{\frac{1}{p}}$ is a supersolution of (1.1) and $v_k \leq w_k + C^{\frac{1}{p}}$. We conclude as in the first step.

Uniqueness. It is proved in Theorem 4.6, this ends the proof.

When we do not assume $q \leq \frac{2p}{p+1}$ we have only the existence of a minimal positive solution. This is due to the fact that for two solutions u and u' as above, $\min\{u, u'\}$ is a supersolution larger than v_k . The conclusion follows easily. \square

In the next statements we prove the existence of radial solutions defined in the complement of a ball of \mathbb{R}^N , $N \geq 3$ which behaves like the Newtonian potential at infinity. We start with the following lemma dealing with the positive radial solutions of $\mathcal{L}_p v = 0$ in the complement of a ball.

Lemma 3.13 *Assume $N \geq 3$ and $p > \frac{N}{N-2}$. Then for any $c > 0$ there exists $k_c > 0$ such that the unique solution v_c of $\mathcal{L}_p v = 0$ in B_1^c verifying $v|_{\partial B_1} = c$ satisfies*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} v_c(x) = k_c. \quad (3.110)$$

Furthermore the mapping $c \mapsto k_c$ is continuous and increasing from $(0, \infty)$ onto $(0, k_\infty)$ for some $k_\infty < \infty$.

Proof. The existence and uniqueness of a solution v_c in an exterior domain and the fact that (3.110) holds is classical (see e.g. [21]). However the fact that $k_c > 0$ and the continuity of $c \mapsto k_c$ is not proved there. By the maximum principle $c \mapsto k_c$ is nondecreasing. Next we set $s = \frac{r^{N-2}}{N-2}$ and $v_c(r) = r^{2-N} \rho(s)$. Then $\rho_c := \rho$ satisfies

$$s^2 \rho_{ss} = c_{N,p} s^{\frac{N}{N-2}-p} \rho^p \quad \text{on } ((N-2)^{-1}, \infty) \quad \text{and} \quad \rho((N-2)^{-1}) = c, \quad (3.111)$$

where $c_{N,p} = (N-2)^{\frac{4-N}{N-2}-p}$. By the maximum principle $v_c(r) \leq cr^{2-N}$ (v_c is the positive harmonic function in B_1^c with value c on ∂B_1), hence $\rho(s)$ is bounded. Since ρ is convex and bounded, it is decreasing and $\rho'(s) \rightarrow 0$ as $s \rightarrow \infty$. Hence

$$-\rho'(s) = c_{N,p} \int_s^\infty \tau^{\frac{N}{N-2}-p-2} \rho^p(\tau) d\tau \leq c'_{N,p} s^{\frac{N}{N-2}-p-1} \rho^p(s). \quad (3.112)$$

Hence by integration the function $\rho \mapsto \Phi(\rho) = \rho^{1-p} - c'' \rho^{\frac{N}{N-2}-p}$ is increasing and bounded. Then it has a finite limit ℓ when $\rho \rightarrow \infty$ and $\Phi^{1-p}(\rho)$ has the same limit ℓ . Thus $\ell \neq 0$ and consequently $k_c > 0$. Let $\{c_n\}$ be a decreasing sequence in \mathbb{R}_+ converging to $c^* > 0$. Then the sequence of corresponding solutions $\{v_{c_n}\}$ is decreasing to v_{c^*} the sequence $\{k_{c_n}\}$ is nonincreasing with limit $k^* \geq k_{c^*}$. From (3.112) one get

$$c_n - k_{c_n} = c_{N,p} \int_{(N-2)^{-1}}^\infty \int_s^\infty \tau^{\frac{N}{N-2}-p-2} \rho_{c_n}^p(\tau) d\tau ds, \quad (3.113)$$

and the same identity holds in c_n is replaced by c^* . By the dominated convergence theorem, one has that

$$c^* - k^* = c_{N,p} \int_{(N-2)^{-1}}^\infty \int_s^\infty \tau^{\frac{N}{N-2}-p-2} \rho_{c^*}^p(\tau) d\tau ds, \quad (3.114)$$

which implies that $k^* = k_{c^*}$. A similar result holds if $\{c_n\}$ is an increasing sequence in \mathbb{R}_+ converging to $c_* > 0$. Hence $c \mapsto k_c$ is increasing and continuous. When $c \uparrow \infty$ v_c increases and converges to the unique positive solution v_∞ of $\mathcal{L}_p v = 0$ in B_1^c such that $\lim_{r \rightarrow 1} v(r) = \infty$. Hence $k_c \uparrow k_\infty$ and $k_\infty < \infty$. \square

Remark. Since the equation $\mathcal{L}_p v = 0$ is invariant by the transformation T_ℓ defined in (1.3), the ball B_1 can be replaced by B_R for any $R > 0$. The range of k_c , that we call $k_{c,R}$ is modified accordingly and $\lim_{c \rightarrow \infty} k_{c,R} = k_{\infty,R}$. Then one has

$$k_{\infty,R} = R^{N-2-\alpha} k_{\infty,1}. \quad (3.115)$$

Theorem 3.14 *Let $N \geq 3$, $M > 0$, $p > \frac{N}{N-2}$ and $\frac{N}{N-1} < q < p$.*

1- *For any $k > 0$ there exist $R := R_k > 0$ and a positive radial solution u of (1.1) in B_R^c satisfying*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k. \quad (3.116)$$

2- *If $\frac{2p}{p+1} < q < p$ there exist $\tilde{k} > 0$ and a positive radial solution, unique among all the positive solutions, u of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying $\lim_{r \rightarrow 0} r^\gamma u(r) = X_M$ and (3.116)*

with $k = \tilde{k}$. In the particular case $q = \frac{(N-2)p}{N-1}$ we have $u = u_{M,p}^$ (see (3.13)).*

3- *If $\frac{2p}{p+1} < q < 2$ and the assumption (3.92) of Theorem 3.11 is satisfied, there exist $k > 0$ and a radial positive solution u of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying (3.116) and $\lim_{x \rightarrow 0} |x|^\beta u(x) = \xi_M$. Furthermore u is unique among all the positive solutions satisfying (3.116).*

Proof. 1- If w is a positive radial and decreasing function such that $\mathcal{R}_q^M w = 0$ it satisfies (see (3.33))

$$-w'(r) = r^{1-N} \left(C + \frac{M}{\kappa} r^{N-(N-1)q} \right)^{-\frac{1}{q-1}},$$

where, $\kappa = \frac{(N-1)q-N}{q-1}$ and $C \in \mathbb{R}$. If $C > 0$, w is defined on $(0, \infty)$. Hence if $w(r) \rightarrow 0$ as $r \rightarrow \infty$, one has

$$w(r) = \int_r^\infty s^{1-N} \left(C + \frac{M}{\kappa} s^{N-(N-1)q} \right)^{-\frac{1}{q-1}} ds. \quad (3.117)$$

Then $w(r) = \frac{1}{(N-2)C^{\frac{1}{q-1}}} r^{2-N} (1+o(1))$ as $r \rightarrow \infty$. Hence, if $k > 0$ is given, we choose

$C > 0$ such that $\frac{1}{(N-2)C^{\frac{1}{q-1}}} = k$. In order that k is in the range of the application

$c \mapsto k_{c,R}$, one takes $R > 0$ such that $k < R^{N-2-\alpha} k_{\infty,1}$. For such an R , there exists $c > 0$ such that the solution v_c of $\mathcal{L}_p v = 0$ in B_R^c verifying $v = c$ on ∂B_R satisfies

(3.110). We then set $C = \left(\frac{1}{(N-2)k} \right)^{q-1}$. The function $w := w_C$ defined by (3.117) is

a supersolution of (1.1) in B_R^c , larger than the subsolution v_c and both v_c and w_C satisfy (3.116). Then by [25, Theorem 1.4.5] there exists a radial positive solution u of (1.1) in B_R^c such that $v_c \leq u \leq w_C$, hence (3.116) follows.

2- The existence of a unique positive and radial solution in $\mathbb{R}^N \setminus \{0\}$ satisfying (1.26) follows from Theorem 1.11. The asymptotic behaviour is a consequence of Theorem 1.9-(2).

3- Under the condition (3.92) of Theorem 3.11 there exists a unique positive solution in $\mathbb{R}^N \setminus \{0\}$ satisfying (3.93). From Theorem 1.9, and since $p > \frac{N}{N-2}$, its behaviour at infinity is given by (1.34) for some specific $k^* > 0$. Uniqueness follows from Theorem 4.6. \square

4 Isolated singularities of non-radial solutions

4.1 Existence and uniqueness of singular solutions

The results of this paragraph are independent of the description of the radial singular solutions performed in the previous sections and they provide a general tool for constructing singular solutions. The existence of singular solutions is based upon the next variant of [11, Theorem 2.1] proved in [25, Corollary 1.4.5].

Theorem 4.1 *Let G be a bounded domain in \mathbb{R}^N , $B \in C(G \times \mathbb{R} \times \mathbb{R}^N)$ a real valued function, $\Gamma \in C(\mathbb{R}_+, \mathbb{R}_+)$ an increasing function such that*

$$|B(x, r, \xi)| \leq \Gamma(|r|)(1 + |\xi|^2) \quad \text{for all } (x, r, \xi) \in G \times \mathbb{R} \times \mathbb{R}^N. \quad (4.1)$$

Let \mathcal{Q} be the operator defined by

$$\mathcal{Q}(u) = -\Delta u + B(x, u, \nabla u). \quad (4.2)$$

If there exist a supersolution $\phi \in W^{1,\infty}(G)$ and a subsolution $\psi \in W^{1,\infty}(G)$ such that $\psi \leq \phi$, then for any $\chi \in W^{1,\infty}(G)$ satisfying $\psi \leq \chi \leq \phi$ there exists a function $u \in W^{1,2}(G)$ verifying $\psi \leq u \leq \phi$, solution of $\mathcal{Q}(u) = 0$ and such that $u - \chi \in W_0^{1,2}(G)$.

One of the main application of this result is Theorem 1.14 which is proved below

Proof of Theorem 1.14. Let $\{\epsilon_n\}$ be a sequence decreasing to 0 and such that $\epsilon_1 < \text{dist}(0, \partial\Omega)$ and set $m = \max_{z \in \partial\Omega} v(z) + \max_{z \in \partial\Omega} (\phi(z) - v(z))_+$. Then $m \geq \phi$ on $\partial\Omega$ and the function $\bar{v} = v + m$ satisfies $\mathcal{L}_{p,q}^M \bar{v} \geq 0$ in $\Omega \setminus \{0\}$. The function $\underline{v} = (v - \max_{z \in \partial\Omega} \phi(z))_+$ satisfies $\mathcal{L}_{p,q}^M \underline{v} \leq 0$. Put $\chi = \sup\{\underline{v}, \inf\{\bar{v}, \phi\}\}$. Then $\chi \in W_{loc}^{1,\infty}(\Omega \setminus \{0\})$, $\underline{v} \leq \chi \leq \bar{v}$ and $\chi = \phi$ on $\partial\Omega$. By Theorem 4.1 for any $n \in \mathbb{N}^*$ there exists a function $u_n \in W^{1,2}(\Omega \setminus \bar{B}_{\epsilon_n})$ such that $(u_n - \chi)|_{\Omega \setminus \bar{B}_{\epsilon_n}} \in W_0^{1,2}(\Omega \setminus \bar{B}_{\epsilon_n})$ satisfying $\mathcal{L}_{p,q}^M u_n = 0$ in $\Omega \setminus \bar{B}_{\epsilon_n}$. Furthermore u_n is unique by the maximum principle. Since $u_n = \bar{v}$ on ∂B_{ϵ_n} , v , and therefore \bar{v} , is radially decreasing and $u_n = \chi$ on $\partial\Omega$ we infer that $u_n \leq u_{n'}$ in $\Omega \setminus B_{\epsilon_n}$ if $n' \geq n$. Hence the sequence $\{u_n\}$ is increasing and it satisfies

$$(v(x) - \max_{z \in \partial\Omega} \phi(z))_+ \leq u_n(x) \leq v(x) + \max_{z \in \partial\Omega} (\phi(z) - v(z))_+ \quad \text{for all } x \in \Omega \setminus B_{\epsilon_n}. \quad (4.3)$$

By standard regularity estimates, u_n is relatively compact in $C_{loc}^1(\Omega \setminus \{0\})$. Hence it converges to a solution u of $\mathcal{L}_{p,q}^M u = 0$ in $\Omega \setminus \{0\}$ which coincides with ϕ on $\partial\Omega$ and satisfies (1.41). \square

As a first application we have the following:

Corollary 4.2 *Let Ω be any bounded smooth domain containing 0 and $\phi \in W^{1,\infty}(\Omega)$ be nonnegative. There exists a positive solution u of $\mathcal{L}_{p,\frac{2p}{p+1}}^M u = 0$ in $\Omega \setminus \{0\}$ with value ϕ on $\partial\Omega$ such that $u(x) - a|x|^{-\alpha}$ remains bounded in Ω where a is equal to x_M or $x_{j,M}$ ($j=1,2$) or x_{m^*} according to we are in the cases (1)-(2) or (3) or (4) of Theorem 1.1.*

The existence of singular solutions is not restricted to the case $q = \frac{2p}{p+1}$ where they are explicit. The following easy to prove corollary shows that existence, and sometimes uniqueness, holds when $1 < q < \frac{N}{N-1}$. This range of exponents is analysed in [7] in connection with problems with Dirac measure data.

Corollary 4.3 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 1$, be any bounded smooth domain containing 0. Assume $1 < p < \frac{N}{N-2}$ if $N \geq 3$ or any $p > 1$ if $N = 1, 2$, $1 < q < \min\left\{p, \frac{N}{N-1}\right\}$ if $N \geq 2$ or any $q > 1$ if $N = 1$, $M > 0$ and $k > 0$. Then for any $\phi \in W^{1,\infty}(\Omega)$, $\phi \geq 0$, there exists a positive solution u of $\mathcal{L}_{p,q}^M u = 0$ in $\Omega \setminus \{0\}$ with value ϕ on $\partial\Omega$ satisfying (1.12).*

Proof of Theorem 1.2 and Theorem 1.3. It is a direct consequence of the above results. \square

More general uniqueness results valid for any positive solution, not necessarily radial, are obtained below. Furthermore the problems involved are either considered in $\mathbb{R}^N \setminus \{0\}$ or in a punctured bounded domain. If b is a positive parameter we define a continuous group of transformations acting on functions u defined in an open set G , $u \mapsto u_\ell^{(b)}$, for $\ell > 0$ by the formula

$$u_\ell^{(b)}(x) = \ell^b u(\ell x) \quad \text{for all } \ell > 0 \text{ and } x \in \ell^{-1}G. \quad (4.4)$$

If u satisfies (1.1) in G , then $u_\ell^{(b)}$ satisfies

$$-\Delta u_\ell^{(b)} + \ell^{2-b(p-1)}(u_\ell^{(b)})^p - M\ell^{2-q-b(q-1)}|\nabla u_\ell^{(b)}|^q = 0 \quad \text{in } \ell^{-1}G. \quad (4.5)$$

If $\ell > 1$, $u_\ell^{(b)}$ is a supersolution of (1.1) if and only if

$$\begin{aligned} (i) \quad & 2 - b(p-1) \leq 0 \quad \iff \alpha \leq b \\ (ii) \quad & 2 - q - b(q-1) \geq 0 \quad \iff \beta \geq b. \end{aligned} \quad (4.6)$$

This conditions are compatible if and only if $1 < q \leq \frac{2p}{p+1}$. Similarly, if $\ell < 1$, $u_\ell^{(b)}$ is a supersolution of (1.1) if and only if

$$\begin{aligned} (i) \quad & 2 - b(p-1) \geq 0 \quad \iff \alpha \geq b \\ (ii) \quad & 2 - q - b(q-1) \leq 0 \quad \iff \beta \leq b. \end{aligned} \quad (4.7)$$

This conditions are compatible if and only if $\frac{2p}{p+1} \leq q < 2$.

Proof of Theorem 1.15 First we note that two terms on the right hand-side of (1.42) in the statement of the theorem coincide only if $q = \frac{2p}{p+1}$ since $\alpha \leq \beta$ is equivalent to $q \leq \frac{2p}{p+1}$. We first study the problem in $\mathbb{R}^N \setminus \{0\}$. We have to consider two cases:
1- Suppose $\alpha \leq \beta$. We choose b such that

$$b \in (a, \infty) \cap [\alpha, \beta]. \quad (4.8)$$

Let u and \tilde{u} be two positive solutions satisfying (1.43). For $\ell > 1$, $u_\ell^{(b)}$ is a supersolution. Since

$$u_\ell^{(b)}(x) = \Lambda \ell^{b-a} |x|^{-a} (-\ln |x|)^{\tilde{a}} (1 + o(1)) \quad \text{as } x \rightarrow 0$$

and $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$, for any $\epsilon > 0$ the function $x \mapsto u_\ell^{(b)}(x) + \epsilon$ which is a supersolution is larger than \tilde{u} near 0 and at infinity. Then $u_\ell^{(b)} + \epsilon \geq \tilde{u}$ in $\mathbb{R}^N \setminus \{0\}$. Letting $\epsilon \downarrow 0$ and $\ell \downarrow 1$, yields $u \geq \tilde{u}$. Similarly $\tilde{u} \geq u$.

2- Suppose $\alpha > \beta$. We choose b such that

$$b \in (0, a) \cap [\beta, \alpha]. \quad (4.9)$$

Then for $\ell < 1$, $u_\ell^{(b)} + \epsilon$ is a supersolution in $\mathbb{R}^N \setminus \{0\}$ which is larger than \tilde{u} at 0 and at ∞ . Hence $\tilde{u} \leq u_\ell^{(b)} + \epsilon$ and we conclude as in the first case.

Next we consider the problem in Ω . Since the solutions are continuous in $\overline{\Omega} \setminus \{0\}$, for $\epsilon > 0$ we have that for $\ell > 1$ $u_\ell^{(b)} + \epsilon > \tilde{u}$ near $\partial(\ell^{-1}\Omega)$ provided $\ell - 1$ is small enough. Hence $u_\ell^{(b)}(x) + \epsilon \geq \tilde{u}$ in $\ell^{-1}\Omega \setminus \{0\}$. This implies that $u \geq \tilde{u}$ by letting $\ell \uparrow 1$ and then $\epsilon \rightarrow 0$. If $\ell < 1$ then $\Omega \subset \ell^{-1}\Omega$, and we compare $u_\ell^{(b)} + \epsilon$ and \tilde{u} in Ω . The proof follows. \square

The previous result necessitates to find some b satisfying either (4.8) or (4.9) which is not always possible in practice. We give below a variant of the result which necessitates a slightly sharper blow-up estimate.

Theorem 4.4 *Assume $N \geq 1$, $p > 1$, $1 < q \leq \frac{2p}{p+1}$ and $M > 0$. Let a such that*

$$0 \leq a \leq \beta \quad (4.10)$$

There exists at most one positive solution of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$u(x) = \Lambda |x|^{-a} + \Lambda' |x|^{-a'} (1 + o(1)) \quad \text{as } x \rightarrow 0, \quad (4.11)$$

or

$$u(x) = \Lambda |x|^{-a} (-\ln |x|)^{-a''} (1 + o(1)) \quad \text{as } x \rightarrow 0, \quad (4.12)$$

where Λ, Λ' are some positive constants and $a > a'$ and $a'' > 0$.

Proof. The principle of the proof is to replace (4.8) by

$$b \in [a, \infty) \cap [\alpha, \beta]. \quad (4.13)$$

when $\alpha \leq \beta$. Then, for $\ell > 1$, $u_\ell^{(b)}$ is a supersolution. If u satisfies (4.11) then, as $x \rightarrow 0$,

$$u_\ell^{(b)}(x) = \Lambda \ell^{b-a} |x|^{-a} + \Lambda' \ell^{b-a'} |x|^{-a} |x|^{-a'} (1 + o(1))$$

Since $b \geq a > a'$, $u_\ell^{(b)}$ is larger than another solution \tilde{u} near 0. Thus $u_\ell^{(b)} + \epsilon \geq \tilde{u}$ for any $\epsilon > 0$, which implies the claim.

If u satisfies (4.12), then

$$\begin{aligned} u_\ell^{(b)}(x) &= \Lambda \ell^{b-a} |x|^{-a} (-\ln |x| - \ln \ell)^{-a''} (1 + o(1)) \\ &= \Lambda \ell^{b-a} |x|^{-a} (-\ln |x|)^{-a''} \left(1 + a'' \frac{\ln \ell}{-\ln |x|} \right) (1 + o(1)). \end{aligned}$$

Again $u_\ell^{(b)}$ is larger than another solution \tilde{u} in a neighborhood of 0 and we end the proof as in the first case. \square

Remark. The method developed above allows to give uniqueness result for large solutions under some starshapedness assumption. Let $G \subset \mathbb{R}^N$ be a domain with compact boundary and $\rho_G(x) = \text{dist}(x, \partial G)$, we consider the problem

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^q &= 0 \quad \text{in } G \\ \lim_{\rho_G(x) \rightarrow 0} u(x) &= \infty. \end{aligned} \quad (4.14)$$

Such a solution, if it exists is called a *large solution*.

Theorem 4.5 *Assume $N \geq 1$, $M > 0$ and $p, q > 1$ and Ω is a bounded domain starshaped with respect to 0. There exists at most one positive function satisfying (4.14) in one of the following case:*

1- $\frac{2p}{p+1} \leq q < 2$ and $G = \Omega$.

2- $1 < q \leq \frac{2p}{p+1}$ and $G = \overline{\Omega}^c$.

Proof. Let u and \tilde{u} be two positive solutions of (4.14). In the first case with $G = \Omega$. Then for $\alpha \leq b \leq \beta$ and $\ell > 1$, u_ℓ^b is a supersolution of (4.14) in $\Omega_\ell := \ell^{-1}\Omega$. Since $\overline{\Omega}_\ell \subset \Omega$, $u_\ell^b \geq \tilde{u}$, it follows that $u \geq \tilde{u}$.

In the second case with $G = \overline{\Omega}^c$, then for $0 < \ell < 1$ and $\beta \leq b \leq \alpha$. Then u_ℓ^b is a supersolution in $\ell^{-1}\overline{\Omega}^c \subset \overline{\Omega}^c$. Then for $\epsilon > 0$, $u_\ell^b + \epsilon \geq \tilde{u}$ in $\ell^{-1}\overline{\Omega}^c$. Letting $\epsilon \rightarrow 0$ and $\ell \uparrow 1$ yields $u \geq \tilde{u}$. This ends the proof. \square

If we combine the results of existence of radial singular solutions in $\mathbb{R}^N \setminus \{0\}$ with the uniqueness results of Theorem 1.15 and Theorem 4.4 we have the following:

Theorem 4.6 Assume $N \geq 3$, $p, q > 1$ and $M > 0$. There exists one and only one positive solution u of (1.1) in $\mathbb{R}^N \setminus \{0\}$, if one of the following conditions holds:

1- $1 < p < \frac{N}{N-2}$, $q = \frac{2p}{p+1}$, $M > 0$ and u satisfies (1.12)-(i) for some $k > 0$.

2- $\frac{2p}{p+1} < q < p$ and $\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M$.

3- $1 < p < \frac{N}{N-2}$, $1 < q < \frac{2p}{p+1}$, $M > 0$ and either $\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_0$, or u satisfies (1.12)-(i) for some $k > 0$.

4- $p = \frac{N}{N-2}$, $q = \frac{2p}{p+1}$, $M > 0$ and u satisfies (1.15)-(i).

Furthermore, existence and uniqueness of a solution holds if the equation (1.1) is considered in $\Omega \setminus \{0\}$ where Ω is a bounded smooth domain starshaped with respect to 0 and is the function u is equal to some ϕ on $\partial\Omega$ where $\phi \in C^1(\partial\Omega)$ is nonnegative.

Proof. By applying Theorem 1.14 and Theorem 1.15 the proof is reduced to use results of existence of radial positive singular solutions in $\mathbb{R}^N \setminus \{0\}$ and to check that the parameters fulfill the conditions of Theorem 1.15.

Case 1- If $q = \frac{2p}{p+1}$, $\alpha = \beta$ and $N - 2 < \beta$. Existence of radial positive solutions satisfying (1.12)-(i) is proved in Theorem 1.3.

Case 2- Then $\gamma > \beta$. Existence of a radial positive solution satisfying $\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M$ is proved in Theorem 1.11-1.

Case 3- If $1 < q < \frac{2p}{p+1}$, then $\alpha < \beta$, and since $p < \frac{N}{N-2}$, $N - 2 < \alpha$. Existence of a radial positive solution satisfying $\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_0$ is proved in Theorem 1.13-1.

Furthermore the assumptions on p and q imply that $q < \frac{N}{N-1}$. The existence of a positive solution satisfying (1.12)-(i) for any $k > 0$ is proved in Theorem 3.12.

Case 4- When $p = \frac{N}{N-2}$, $q = \frac{2p}{p+1}$ there exists a radial global solution satisfying (1.15)-(i) by Theorem 1.4. We apply estimate (4.12) in Theorem 4.4 with $a = N - 2 = \alpha = \beta$ and $a'' = N - 1$. The result follows. \square

Remark. In the case $p = \frac{N}{N-2}$, $q = \frac{N}{N-1}$ we conjecture that the function u_{x_M} is the only positive solution of (1.1) defined in $\mathbb{R}^N \setminus \{0\}$ satisfying $\lim_{x \rightarrow 0} |x|^{N-2} u(x) = x_M$.

4.2 Characterization of singular solutions

In this section we give some results showing how the characterization of singularities of radial solutions can be extended to nonradial solutions. An important tool for studying positive isolated singularities is Harnack inequality.

Proposition 4.7 Assume $M > 0$, $p > 1$ and $1 < q \leq \frac{2p}{p+1}$. If u is a positive solution of (1.1) in $B_{R_0} \setminus \{0\}$, there exists $c_9 = c_9(N, p, q, R_0, M) > 0$ such that for any $R \in (0, \frac{R_0}{2}]$ there holds

$$\max_{|x|=R} u(x) \leq c_9 \min_{|x|=R} u(x). \quad (4.15)$$

Proof. We write (1.1) under the form

$$-\Delta u + C(x)u + V(x)|\nabla u| = 0, \quad (4.16)$$

where $C(x) = |u(x)|^{p-1}$ and $V(x) = M|\nabla u(x)|^{q-1}$. By Proposition A.1,

$$C(x) \leq c_1^{p-1} \max \left\{ M^{\frac{p-1}{p-q}} |x|^{-\frac{q(p-1)}{p-q}}, |x|^{-2} \right\}$$

and

$$V(x) \leq c_4^{q-1} \max \left\{ M^{\frac{q-1}{p-q}} |x|^{-\frac{p(q-1)}{p-q}}, |x|^{-\frac{(q-1)(p+1)}{(p-1)}} \right\}$$

Under the assumptions $1 < q \leq \frac{2p}{p+1}$, the terms $|x|^2 C(x)$ and $|x|V(x)$ are uniformly bounded in $B_{R_0} \setminus \{0\}$. The result follows by [13, Chapter 8]. \square

4.3 The case $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{N}{N-1}$

In this section, the results are obtained by a combination of Theorem 3.12 for existence of solutions and Theorem 4.6 for their uniqueness.

Theorem 4.8 *Let $N \geq 3$, $M > 0$, $1 < p < \frac{N}{N-2}$, $1 < q < \frac{N}{N-1}$ and Ω be a bounded domain containing 0. For any $k > 0$ there exists a unique positive function $u := u_k$ solution of (1.1) in $\Omega \setminus \{0\}$, vanishing on $\partial\Omega$ and satisfying*

$$\lim_{x \rightarrow 0} |x|^{N-2} u(x) = k. \quad (4.17)$$

Furthermore $k \mapsto u_k$ is increasing by the maximum principle and converges to a solution u_∞ of (1.1) in $\Omega \setminus \{0\}$, vanishing on $\partial\Omega$ and satisfying

$$\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M \quad \text{if } \frac{2p}{p+1} < q < \frac{N}{N-1}, \quad (4.18)$$

where X_M is defined at (1.26), or

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = \begin{cases} x_0 & \text{if } 1 < q < \frac{2p}{p+1} \\ x_M & \text{if } q = \frac{2p}{p+1}, \end{cases} \quad (4.19)$$

where x_0, x_M are the unique positive root of equation (1.4) with $M = 0$ and $M > 0$ respectively.

Proof. Let $0 < R_1 < R_2$ be such that $\overline{B}_{R_1} \subset \Omega \subset \overline{\Omega} \subset B_{R_2}$. By Theorem 1.13-1 and Theorem 4.6, for $k > 0$ there exists a unique solution $u_{1,k}$ (resp. $u_{2,k}$) of (1.1) in $B_{R_1} \setminus \{0\}$ (resp. $B_{R_2} \setminus \{0\}$) satisfying (1.12)-(i) and vanishing on ∂B_{R_1} (resp. ∂B_{R_2}). If we extend $u_{1,k}$ by 0 in $B_{R_1}^c$, we have,

$$u_{1,k}(|x|) \leq u_k(x) \leq u_{2,k}(|x|) \quad \text{in } \Omega \setminus \{0\}.$$

All the above functions are locally bounded in $\overline{\Omega} \setminus \{0\}$ and $\overline{B}_{R_2} \setminus \{0\}$ by Proposition A.1. Since the mappings $k \mapsto u_k$ and $k \mapsto u_{j,k}$ are increasing, we have, by letting $k \rightarrow \infty$,

$$u_{1,\infty}(|x|) \leq u_\infty(x) \leq u_{2,\infty}(|x|) \quad \text{in } \Omega \setminus \{0\}.$$

Then we obtain (4.18) by Theorem 1.7-(1) and (4.19) by Theorem 1.8-(1) and Theorem 1.3. \square

The main characterization of isolated singularities is the next result.

Theorem 4.9 *Let $N \geq 3$, Ω be an open subset containing 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $1 < q \leq \frac{2p}{p+1}$. If u is a positive solution of (1.1) in $\Omega \setminus \{0\}$, then either its behaviour at $x = 0$ is given by (4.18) or (4.19), or there exists $k \geq 0$ such that (4.17) holds. If $k = 0$ the singularity at 0 is removable.*

The proof needs a few intermediate steps.

Lemma 4.10 *Let $N \geq 3$, Ω be an open subset containing 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{N}{N-1}$. If u is a positive solution of (1.1) in $\Omega \setminus \{0\}$ vanishing on $\partial\Omega$ and such that*

$$\limsup_{x \rightarrow 0} |x|^{N-2} u(x) < \infty.$$

Then there exists $k \geq 0$ such that (4.17) holds. If $k = 0$, then u coincides in $\Omega \setminus \{0\}$ with a $C^2(\Omega)$ solution of (1.1) in Ω .

Proof. By assumption $u(x) \leq c|x|^{2-N}$ and by [4, Lemma 3.10] we have the following: if u is a solution of (1.1) in $\Omega \setminus \{0\}$ (not necessarily positive) such that $|x|^m |u(x)|$ is bounded near $x = 0$ for some $m < \inf\{\alpha, \frac{2-q}{q-1}\}$, then $|x|^{m+1} |\nabla u(x)|$ is also bounded near $x = 0$. Actually, in the reference the result is proved for a more general operator, without the absorption u^p , but the adaptation is straightforward. The result applies there with $m = N - 2$ and in particular $|\nabla u(x)| \leq c'|x|^{1-N}$. We write (1.1) under the form (4.16). Since $p < \frac{N}{N-2}$ and $q < \frac{N}{N-1}$ we have

$$|C(x)| \leq c|x|^{(2-N)(p-1)} \leq c|x|^{-2+\epsilon_1} \quad \text{and} \quad |V(x)| \leq c|x|^{(1-N)(q-1)} \leq c|x|^{-1+\epsilon_2},$$

for some $\epsilon_1, \epsilon_2 > 0$. It follows by Serrin's result that either the singularity at 0 is removable, or there exist $c_1 > c_2 > 0$ such that

$$c_2|x|^{2-N} \leq u(x) \leq c_1|x|^{2-N} \quad \text{for all } 0 < |x| \leq 1.$$

In order to make the convergence precise, we denote by u_2 the solution of

$$\begin{aligned} -\Delta u_2 &= M|\nabla u|^q - u^p && \text{in } B_1 \\ u_2 &= 0 && \text{on } \partial B_1. \end{aligned}$$

Then $-v_2 \leq u_2 \leq v_2'$ where

$$\begin{aligned} -\Delta v_2' &= M c'^q |x|^{(1-N)q} && \text{in } B_1 \\ v_2' &= 0 && \text{on } \partial B_1, \end{aligned}$$

and

$$\begin{aligned} -\Delta v_2 &= c^p |x|^{(2-N)p} && \text{in } B_1 \\ v_2 &= 0 && \text{on } \partial B_1, \end{aligned}$$

Because $(N-1)q < N$ and $(N-2)p < N$, v_2 and v_2' satisfy

$$0 \leq v_2(r) \leq c_{10} r^{2-N+\delta} \quad \text{and} \quad 0 \leq v_2'(r) \leq c_{10} r^{2-N+\delta'},$$

for some $c_{10} > 0$ and $\delta = N - (N - 2)p > 0$, $\delta' = N - (N - 1)q > 0$. Then u_2 satisfies

$$\lim_{x \rightarrow 0} |x|^{N-2} u_2(x) = 0.$$

The function $u_1 = u - u_2$ is harmonic in $B_1 \setminus \{0\}$ and is bounded from below by $-v_2'$ which satisfies $\lim_{x \rightarrow 0} |x|^{N-2} v_2'(x) = 0$. Hence by standard result on singularities of harmonic functions, $|x|^{N-2} u_1(x)$ admits a limit $k \geq 0$ when $x \rightarrow 0$. Combined with Serrin's estimates it follows that either $k = 0$ and the singularity is removable, or $k > 0$. Note that if $k = 0$, then u is a C^2 solution in Ω .

Another proof based on a perturbation is the following: let $u(x) = u(r, s) = |x|^{2-N} \phi(t, s)$ with $r = |x|$ and $t = \ln r$. Then

$$\begin{aligned} \phi_{tt} + (N - 2)\phi_t - e^{(N-p(N-2))t} \phi^p + \Delta' \phi \\ + M e^{(N-q(N-1))t} \left(((N - 2)\phi - \phi_t)^2 + |\nabla' \phi|^2 \right)^{\frac{q}{2}} = 0 \end{aligned}$$

Since $u = O(|x|^{2-N})$, we can write

$$\phi_{tt} + (N - 2)\phi_t + \Delta' \phi = -e^{at} \psi,$$

where $a = \min\{(N - p(N - 2)), (N - q(N - 1))\} > 0$ and ψ is bounded. Then the result follows by [10, Proposition 4.1]. \square

We give below another application of the perturbation method and specific to the case $q < \frac{2p}{p+1}$.

Proposition 4.11 *Assume Ω is an open subset containig 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{2p}{p+1}$. If u is a solution of*

$$-\Delta u + |u|^{p-1} u - M |\nabla u|^q = 0 \quad (4.20)$$

not necessarily nonnegative in $\Omega \setminus \{0\}$, then $r^\alpha u(r, s)$ converges uniformly with respect to $s \in S^{N-1}$ when $r \rightarrow 0$ to a non-empty compact and connected subset of the set of solutions ω of

$$-\Delta' \omega + \alpha(N - 2 - \alpha)\omega + |\omega|^{p-1} \omega = 0 \quad \text{on } S^{N-1}. \quad (4.21)$$

If $u \geq 0$, ω is either x_0 or 0.

Proof. We can assume that $\overline{B_1} \subset \Omega$ and set $\phi(t, s) = r^\alpha u(r, s)$ with $t = \ln r$. Then ϕ satisfies

$$\phi_{tt} + L\phi_t + \alpha K\phi + \Delta' \phi - |\phi|^{p-1} \phi + M e^{\frac{\sigma t}{p-1}} \left((\alpha\phi - \phi_t)^2 + |\nabla' \phi|^2 \right)^{\frac{q}{2}} = 0. \quad (4.22)$$

in $\mathbb{R}_- \times S^{N-1}$ where $K = N - 2 - \alpha$ and $L = K - \alpha$. By assumption $\sigma < 0$, hence (4.22) is an exponentially small perturbation of the autonomous equation associated to the Emden-Fowler equation by the same change of variables. The result follows

from [10, Theorem 4.1] but for the sake of comprehension, we recall its proof. By Proposition A.2 the function ϕ is uniformly bounded, and by (A.6) ϕ_t and $\nabla'\phi$ are also uniformly bounded. By standard local regularity theory for elliptic equations, there holds

$$\|\partial_{t^i}\nabla'^j\phi\|_{C[T-1, T+1]\times S^{N-1}} \leq c_{11} \quad \text{for all } (i, j) \in \mathbb{N} \times \mathbb{N}, i + j \leq 3 \quad \text{and } T \leq -2,$$

where ∇'^j is the covariant derivative of order j on S^{N-1} . Then the omega-limit at $-\infty$ of the trajectory $\{\phi(t, \cdot)\}_{t \in \mathbb{R}_-}$ in $C^2(S^{N-1})$ is a non-empty compact connected denoted by $\Gamma_\phi \subset C^2(S^{N-1})$. From (4.22) we have that

$$\frac{d}{dt}E = L \int_{S^{N-1}} \phi_t^2 dS(s) + M e^{\frac{\sigma t}{p-1}} \int_{S^{N-1}} ((\alpha\phi - \phi_t)^2 + |\nabla'\phi|^2)^{\frac{q}{2}} \phi_t dS(s),$$

where

$$J(t) = \frac{1}{2} \int_{S^{N-1}} \left(|\nabla'\phi|^2 + \frac{2}{p+1} |\phi|^{p+1} - \phi_t^2 \right) dS(s).$$

Because $L \neq 0$ and J is uniformly bounded, there holds

$$\int_{-\infty}^1 \int_{S^{N-1}} \phi_t^2 dS(s) dt < \infty.$$

Multiplying (4.22) by w_{tt} and using the previous estimate, we obtain

$$\int_{-\infty}^1 \int_{S^{N-1}} \phi_{tt}^2 dS(s) dt < \infty.$$

As ϕ_t and ϕ_{tt} are uniformly continuous in $(-\infty, 1] \times S^{N-1}$, this implies that

$$\lim_{t \rightarrow -\infty} \int_{S^{N-1}} (\phi_t^2(t, \cdot) + \phi_{tt}^2(t, \cdot)) dS(s) = 0.$$

Therefore $\Gamma_\phi \subset \{\omega \in C^2(S^{N-1}) : \alpha K\omega + \Delta'\omega - |\omega|^{p-1}\omega = 0 \text{ in } S^{N-1}\}$. If $u \geq 0$, then $\omega \geq 0$ for any $\omega \in \Gamma_\phi$ and the result follows by the maximum principle. \square

Lemma 4.12 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded open subset containing 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $1 < q < \frac{2p}{p+1}$. If u is a nonnegative solution of (1.1) in $\Omega \setminus \{0\}$ such that*

$$\limsup_{x \rightarrow 0} |x|^{N-2} u(x) = \infty, \tag{4.23}$$

then

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_0. \tag{4.24}$$

Proof. Without loss of generality we can assume that $\overline{B_1} \subset \Omega$. It follows from Proposition 4.7 that $\liminf_{x \rightarrow 0} |x|^{N-2} u(x) = \infty$, Hence there exists a decreasing sequence $\{r_n\}$ converging to 0 such that

$$\alpha_n := \inf_{|x|=r_n} |x|^{N-2} u(x) \uparrow \infty \quad \text{as } n \rightarrow \infty.$$

Let v_n be the solution of

$$\begin{aligned} -\Delta v + v^p &= c_N \alpha_n \delta_0 && \text{in } \mathcal{D}'(B_1) \\ v &= 0 && \text{on } \partial B_1, \end{aligned} \quad (4.25)$$

where c_N is the explicit constant such that $\mathbb{G}[c_N \delta_0](x) = |x|^{2-N}$, where \mathbb{G} is the Newtonian potential in \mathbb{R}^N ; note that v_n is radial because of uniqueness. Then the sequence $\{v_n\}$ is increasing and converges to the function v_∞ which satisfies (see [22])

$$\begin{aligned} -\Delta v + v^p &= 0 && \text{in } B_1 \setminus \{0\} \\ v &= 0 && \text{on } \partial B_1 \\ \lim_{x \rightarrow 0} |x|^\alpha v(x) &= x_0. \end{aligned} \quad (4.26)$$

Moreover, $v_n(x) \leq \alpha_n |x|^{2-N}$ and since v_n is a subsolution of (1.1), we have that $v_n \leq u$ in $\overline{B_1} \cap B_{r_n}^c$. Letting $n \rightarrow \infty$ implies that $v_\infty(x) \leq u(x)$ in $B_1 \setminus \{0\}$. Therefore

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) \geq x_0.$$

Combined with Proposition 4.11, this inequality implies (4.22). \square

This result admits an extension to the case $q = \frac{2p}{p+1}$.

Lemma 4.13 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded open subset containing 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $q = \frac{2p}{p+1}$. If u is a nonnegative solution of (1.1) in $\Omega \setminus \{0\}$ such that*

$$\limsup_{x \rightarrow 0} |x|^{N-2} u(x) = \infty, \quad (4.27)$$

then

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_M. \quad (4.28)$$

Proof. Assuming for simplicity that $\overline{B_1} \subset \Omega$ and using Proposition 4.7 we obtain that for some decreasing sequence $\{r_n\}$ converging to 0 we have

$$\alpha_n := \inf_{|x|=r_n} |x|^{N-2} u(x) \uparrow \infty \quad \text{as } n \rightarrow \infty.$$

Therefore, u is bounded from below in $B_1 \setminus B_{r_n}$ by the (radial) solution u_n of

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^q &= 0 && \text{in } B_1 \setminus \overline{B_{r_n}} \\ v &= 0 && \text{on } \partial B_1 \\ u &= \alpha_n r_n^{2-N} && \text{on } \partial B_{r_n}. \end{aligned} \quad (4.29)$$

The sequence $\{u_n\}$ may not be monotone, but $u_n \geq v_n$ where v_n has been defined in (4.25). Since $\{u_n\}$ is eventually locally bounded in $\overline{B_1} \setminus \{0\}$ by Proposition A.1 and standard regularity results (see e.g. [13]), up to a subsequence, it converges locally uniformly in $\overline{B_1} \setminus \{0\}$ to a radial function u_∞ which satisfies

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^q &= 0 && \text{in } B_1 \setminus \{0\} \\ v &= 0 && \text{on } \partial B_1, \end{aligned} \quad (4.30)$$

and

$$\liminf_{x \rightarrow 0} |x|^\alpha u_\infty(x) \geq \liminf_{x \rightarrow 0} |x|^\alpha v_\infty(x) = x_0. \quad (4.31)$$

By Theorem 2.4-(4) we have that

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) \geq \lim_{x \rightarrow 0} |x|^\alpha u_\infty(x) = x_M. \quad (4.32)$$

The upper estimate is obtained as follows. By Proposition 4.7 the function u is bounded from above in $B_1 \setminus B_{r_n}$ by $u'_n + \max_{|y|=1} u(y)$ where u'_n is the solution of

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^q &= 0 && \text{in } B_1 \setminus \overline{B_{r_n}} \\ v &= 0 && \text{on } \partial B_1 \\ u &= c_9 \alpha_n r_n^{2-N} && \text{on } \partial B_{r_n}. \end{aligned} \quad (4.33)$$

For the same reason as above there exists a subsequence, $\{u'_{n_\ell}\}$ which converges locally uniformly in $\overline{B_1} \setminus \{0\}$ to a radial solution u'_∞ of (4.30). By Theorem 2.4-(4) we have that $u'_\infty = u_\infty$. Then $\limsup_{x \rightarrow 0} |x|^\alpha u(x) \leq \lim_{x \rightarrow 0} |x|^\alpha u'_\infty(x) = x_M$, which ends the proof. \square

Proof of Theorem 4.9 It is a direct consequence of Lemma 4.10, Lemma 4.12 and Lemma 4.13. \square

When $q > \frac{2p}{p+1}$ and $|x|^{N-2}u(x)$ is not bounded, Harnack inequality may not hold. However we still have a dichotomy for the possible behaviour which extends Theorem 4.9.

Theorem 4.14 *Let $\Omega \subset \mathbb{R}^N$ ($N \geq 3$) be an open set containing 0, $M > 0$, $1 < p < \frac{N}{N-2}$ and $\frac{2p}{p+1} < q < \frac{N}{N-1}$. If u is a nonnegative solution of (1.1) in $\Omega \setminus \{0\}$, then the following dichotomy holds:*

- 1- either there exists $k \geq 0$ such that $r^{N-2}u(r, \cdot)$ converges to k in measure on S^{N-1} as $r \rightarrow 0$,
- 2- or $\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M$.

Proof. We recall Richard-Véron's isotropy theorem [20, Theorem 1.1]:

Let $g : \mathbb{R}_+ \mapsto \mathbb{R}_+$ be a continuous nondecreasing function satisfying

$$\int_0^1 g(r^{2-N}) r^{N-1} dr < \infty. \quad (4.34)$$

If $u \in C^1(\Omega \setminus \{0\})$ is a nonnegative function satisfying

$$\Delta u \leq g(u) + f \quad \text{in } \Omega \setminus \{0\}, \quad (4.35)$$

where $f \in L^1_{loc}(\Omega \setminus \{0\})$ is a nonnegative radial function, then we have the following:

- (i) either $r^{N-2}u(r, \cdot)$ converges in measure on S^{N-1} to some $k \geq 0$ as $r \rightarrow 0$,
- (ii) or

$$\lim_{x \rightarrow 0} |x|^{N-2}u(x) = \infty. \quad (4.36)$$

Since $p < \frac{N}{N-2}$, assumption (4.34) is satisfied with $g(r) = r^p$ and equation (4.35) with $f = 0$. Then either the statement 1 holds, or (4.36) holds. If it is the case, then for any $k > 0$, u is bounded from below in $B_1 \setminus \{0\}$ by the solution u_k of (1.1) in $B_1 \setminus \{0\}$ vanishing on ∂B_1 and satisfying (4.17). Such a solution exists and is unique by Theorem 4.8. Letting $k \rightarrow \infty$, this implies

$$\liminf_{x \rightarrow 0} |x|^\gamma u(x) \geq X_M. \quad (4.37)$$

Next, we denote by v the solution of (1.1) on $(0, \infty)$ (hence v is 1-dimensional) satisfying

$$\lim_{r \rightarrow 0} r^\gamma v(r) = X_M \quad \text{and} \quad \lim_{r \rightarrow \infty} r^\alpha v(r) = x_0.$$

Its existence is proved in Theorem 1.10, Theorem 1.11. It is decreasing. For $\epsilon > 0$ the function $r \mapsto v_\epsilon(x) = v(|x| - \epsilon)$ satisfies $\mathcal{L}_{p,q}^M(v_\epsilon) \geq 0$ in \overline{B}_ϵ^c and $v_\epsilon(x) \rightarrow \infty$ when $|x| \downarrow \epsilon$. If $c = \max u|_{\partial B_\delta}$ for some $\delta > \epsilon > 0$ such that $\overline{B}_\delta \subset \Omega$, then $v_\epsilon + c$ is a supersolution of (1.1) in $B_\delta \setminus \overline{B}_\epsilon$ which is larger than u for $|x| = \epsilon$ and $|x| = \delta$. Hence $u \leq gv_\epsilon + c$ in $B_\delta \setminus \overline{B}_\epsilon$. Letting $\epsilon \rightarrow 0$ yields $u(x) \leq v(x) + c$ for $0 < |x| \leq \delta$ and finally

$$\limsup_{x \rightarrow 0} |x|^\gamma u(x) \leq X_M. \quad (4.38)$$

Combining (4.37) and (4.38) we obtain $\lim_{x \rightarrow 0} |x|^\gamma u(x) = X_M$. \square

Remark. We conjecture that the convergence in Theorem 4.14-1 holds in the strong sense.

Remark. Most of the results of this section can be extended to the case $N = 2$. The subcritical case corresponds then to $p > 1$ and $1 < q < 2$. The main change is that $|x|^{2-N}$ has to be replaced by $-\ln|x|$.

4.4 The case $p \geq \frac{N}{N-2}$ and $q = \frac{2p}{p+1}$

The cases that we consider are $q = \frac{2p}{p+1}$, $p \geq \frac{N}{N-2}$ and $M > 0$. We recall that the stationary equation (1.4) admits two positive constant solutions $x_{1,M} < x_{2,M}$ if $p > \frac{N}{N-2}$ and $M > m^*$, and only one denoted by x_M if $p = \frac{N}{N-2}$ and $M > 0$ or if $p > \frac{N}{N-2}$ and $M = m^*$. The following result is an improvement of Proposition A.1

Lemma 4.15 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded domain containing 0 such that $\overline{B}_R \subset \Omega$ and $p \geq \frac{N}{N-2}$.*

1- *If $p > \frac{N}{N-2}$ and $M > m^*$, then any positive solution u of (1.2) in $\Omega \setminus \{0\}$ satisfies*

$$u(x) \leq x_{2,M} |x|^{-\alpha} + \sup_{|z|=R} u(z), \quad (4.39)$$

2- *If $p > \frac{N}{N-2}$ and $M = m^*$, or $p = \frac{N}{N-2}$ and $M > 0$, the same inequality holds with $x_{2,M}$ replaced by x_{m^*} and x_M respectively.*

Proof. We assume first that $M > m^*$.

1- *Construction of the maximal solution.* We claim that $x \mapsto x_{2,M}|x|^{-\alpha}$ is the maximal solution of (1.2) in $\mathbb{R}^N \setminus \{0\}$. For $a > 0$ we set $\phi_a(s) = as^\alpha$. Then

$$\tilde{L}\phi_a(s) := -\phi_a'' + \phi_a^p - M|\phi_a'|^{\frac{2p}{p+1}} = as^{-\alpha p} (a^{p-1} - \alpha^q M a^{q-1} - \alpha(\alpha+1)).$$

Taking a large enough we obtain that ϕ_a is a supersolution in $(0, \infty)$. We set $\Phi_{a,\epsilon}(x) = \phi_a(x_1 - \epsilon)$ for $x_1 > \epsilon > 0$ and as in the proof of Theorem 4.14 we deduce that the function $\tilde{\Phi}_{a,\epsilon}$ defined by

$$\tilde{\Phi}_{a,\epsilon} = \inf \{ \mathcal{R}[\Phi_{a,\epsilon}] : \mathcal{R} \in SO(N) \} \quad (SO(N) \text{ is the group of rotations in } \mathbb{R}^N),$$

is a positive and radial supersolution of (1.2) in $\mathbb{R}^N \setminus \overline{B_\epsilon}$ which tends to infinity on ∂B_ϵ . It dominates any positive solution of (1.2) in $\mathbb{R}^N \setminus \overline{B_\epsilon}$. Next we set

$$\tilde{\Psi}_\epsilon(x) = \sup \{ x_{2,M}|x-z|^{-\alpha} : |z| < \epsilon \}.$$

It is a subsolution of (1.2) in $\mathbb{R}^N \setminus \overline{B_\epsilon}$ dominated by $\tilde{\Phi}_{a,\epsilon}$. Since the supremum is achieved for $z = \epsilon \frac{x}{|x|}$, the function $\tilde{\Psi}_\epsilon$ is radial and positive. By [25, Theorem 1.4.5] there exists a solution U_ϵ of (1.2) in $\mathbb{R}^N \setminus \overline{B_\epsilon}$ such that

$$\tilde{\Psi}_\epsilon \leq U_\epsilon \leq \tilde{\Phi}_{a,\epsilon} \quad \text{in } \mathbb{R}^N \setminus \overline{B_\epsilon}.$$

The function U_ϵ is positive and radial. Since any positive solution u in $\mathbb{R}^N \setminus \{0\}$ is dominated by $\tilde{\Psi}_\epsilon$, the function U_ϵ is larger than u in $\mathbb{R}^N \setminus \overline{B_\epsilon}$. This implies the relation, valid for any $\ell > 0$,

$$T_\ell[U_\epsilon](x) := \ell^\alpha U_\epsilon(\ell x) = U_{\ell^{-1}\epsilon}. \quad (4.40)$$

When $\epsilon \downarrow 0$ the sequence $\{U_\epsilon\}$ decreases and converges to a positive radial solution U_0 of (1.2) in $\mathbb{R}^N \setminus \{0\}$ which dominates any other positive solution. Hence U_0 is the maximal solution in $\mathbb{R}^N \setminus \{0\}$. Letting $\epsilon \downarrow 0$ in (4.40) we infer that $T_\ell[U_0] = U_0$ for any $\ell > 0$. Hence U_0 is self-similar. Since it is radial and larger than any other positive solution, we deduce that

$$U_0(x) = x_{2,M}|x|^{-\alpha} \quad \text{for all } |x| > 0. \quad (4.41)$$

2- *End of the proof.* If u is any positive solution in $\Omega \setminus \{0\}$, then $U_\epsilon + \sup_{|z|=R} u(z)$ is a supersolution larger than u in $B_R \setminus B_\epsilon$. Letting $\epsilon \downarrow 0$ yields the result. The proof in the other case is similar. \square

Theorem 4.16 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded domain containing 0, $p > \frac{N}{N-2}$ and $M > m^*$. If u is a positive solution of (1.2) in $\Omega \setminus \{0\}$, there holds*

1- *If*

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) = 0, \quad (4.42)$$

then u can be extended as a C^2 solution of (1.2) in Ω .

2- If

$$\limsup_{x \rightarrow 0} |x|^\alpha u(x) = x_{2,M}, \quad (4.43)$$

then

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_{2,M}, \quad (4.44)$$

3- If

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) = x_{1,M} \quad \text{or} \quad \limsup_{x \rightarrow 0} |x|^\alpha u(x) = x_{1,M}, \quad (4.45)$$

then there exists a sequence $\{r_n\} \subset \mathbb{R}_*^+$ converging to 0 such that

$$\lim_{r_n \rightarrow 0} r_n^\alpha u(r_n, s) = x_{1,M} \quad \text{uniformly on } S^{N-1}. \quad (4.46)$$

4- Any one of the two following situations never occur

$$\begin{aligned} (i) \quad & 0 < \limsup_{x \rightarrow 0} |x|^\alpha u(x) < x_{1,M} \\ (ii) \quad & x_{1,M} < \liminf_{x \rightarrow 0} |x|^\alpha u(x) < x_{2,M}. \end{aligned} \quad (4.47)$$

Proof. 1- If relation (4.42) holds, it follows by Harnack inequality proved in Proposition 4.7, that there exists a sequence $\{r_n\}$ converging to 0 as $n \rightarrow \infty$ such that

$$\lim_{r_n \rightarrow 0} r_n^\alpha u(r_n, s) = 0 \quad \text{uniformly on } S^{N-1}. \quad (4.48)$$

For any $\epsilon > 0$ and $m = \sup\{u(z) : |z| = R\}$, the function $x \mapsto \epsilon|x|^\alpha + m$ is a supersolution of (1.2) in $B_R \setminus \{0\}$ which is larger than u near $x = 0$ and on ∂B_R . Hence $u(x) \leq \epsilon|x|^\alpha + m$. Letting $\epsilon \rightarrow 0$ implies $u \leq m$, and the result follows by standard regularity.

2- If (4.43) holds there exists a sequence $\{r_n\}$ converging to 0 such that

$$r_n^\alpha \max \{u(r_n, s) : s \in S^{N-1}\} = r_n^\alpha u(r_n, s_n) \rightarrow x_{2,M} \quad \text{as } r_n \rightarrow 0.$$

Furthermore, we can assume that $s_n \rightarrow s^*$ when $n \rightarrow \infty$. Using Lemma 4.15 there exist a nondecreasing sequence $\{w_{1,n}\}$ converging to $x_{2,M}$ and a bounded sequence $\{w_{2,n}\}$ such that $r_n^\alpha u(r_n, s_n) = w_{1,n} + r_n^\alpha w_{2,n}$. We set $w(t, s) = r^\alpha u(r, s)$ with $t = \ln r$, then there holds,

$$w_{tt} + Lw_t - \alpha K w + \Delta' w - w^p + M \left((w_t - \alpha w)^2 + |\nabla' w|^2 \right)^{\frac{p}{p+1}} = 0, \quad (4.49)$$

on $\mathbb{R}_- \times S^{N-1}$. By standard regularity estimates and Ascoli-Arzelà theorem there exist a subsequence $\{t_{n_j}\}$ of $\{t_n\} = \{\ln r_n\}$ and a nonnegative C^2 function W such that $w(t_{n_j} + t, s)$ converges to W in the C^2 topology of $[-a, a] \times S^{N-1}$, for any $a > 0$, and W is a solution of (4.49) in $\mathbb{R} \times S^{N-1}$. Furthermore $W(0, s^*) = x_{2,M}$. By Lemma 4.15, $x_{2,M}$ is the maximal solution of (4.49) in $\mathbb{R} \times S^{N-1}$, it then follows from

the strong maximum principle that $W = x_{2,M}$ and $w(t_n + t, s) \rightarrow x_{2,M}$ uniformly in $[-a, a] \times S^{N-1}$. Let $\epsilon > 0$, then there exists $n_\epsilon \in \mathbb{N}$ such that for any $n \geq n_\epsilon$, we have that $u(r_n, s) \geq r_n^{-\alpha}(x_{2,M} - \epsilon)$ for any $s \in S^{N-1}$. Since $r \mapsto r^{-\alpha}(x_{2,M} - \epsilon)$ is a subsolution of (1.2), it follows that for $m > n \geq n_\epsilon$, one has

$$u(r, s) \geq r^{-\alpha}(x_{2,M} - \epsilon) \quad \text{for all } (r, s) \in [r_m, r_n] \times S^{N-1}.$$

Letting $r_m \rightarrow 0$ yields

$$u(x) \geq |x|^{-\alpha}(x_{2,M} - \epsilon) \quad \text{for all } x \in B_{r_{n_\epsilon}} \setminus \{0\}. \quad (4.50)$$

Since ϵ is arbitrary we infer that

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) \geq x_{2,M}. \quad (4.51)$$

By assumption $\limsup_{x \rightarrow 0} |x|^\alpha u(x) \geq x_{2,M}$; then (4.44) holds.

3- Let us assume that the first condition in (4.45) holds. If the function

$$\tau \mapsto w(\tau, s_\tau) := \min_{s \in S^{N-1}} w(\tau, s) \quad (4.52)$$

is asymptotically monotone, nonincreasing or nondecreasing, then either $w(\tau, s_\tau) \uparrow x_{1,M}$ in the first case, or $w(\tau, s_\tau) \downarrow x_{1,M}$ in the second case. Using again the uniform $C^{2,\alpha}$ estimate and Ascoli-Arzelà theorem we have that there exists a sequence $\{\tau_n\}$ converging to $-\infty$ such that $w(t + \tau_n, s)$ converges in the C^2 -topology of $[-a, a] \times S^{N-1}$, for any $a > 0$, to a positive solution W of (4.49) in $\mathbb{R} \times S^{N-1}$ such that $W(t, s) \geq x_{1,M}$ and $\min_{s \in S^{N-1}} W(0, s) = W(0, s^*) = x_{1,M}$ for some $s^* \in S^{N-1}$ in the first case. By the strong maximum principle to w and to $x_{1,M}$ which are ordered solutions of (4.49) in $\mathbb{R} \times S^{N-1}$ we infer that $W \equiv x_{1,M}$, hence $w(t, s)$ converges to $x_{1,M}$ uniformly on S^{N-1} when $t \rightarrow -\infty$. In the second case we obtain that the limit function W satisfies $W(t, s) \geq x_{1,M}$ and $\min_{s \in S^{N-1}} W(0, s) = W(0, s^*) = x_{1,M}$. This implies again $W \equiv x_{1,M}$ by the strong maximum principle. Finally we do not suppose that the function $w(\tau, s_\tau)$ defined in (4.52) is monotone. By the definition of the liminf, there exist sequences $\{t_n\}$ tending to $-\infty$ and $\{s_n\} \subset S^{N-1}$ such that

$$w(t_n, s_n) = \inf\{w(t, s) : t \leq t_n, s \in S^{N-1}\} \uparrow x_{1,M} \quad \text{as } n \rightarrow \infty.$$

Using again Ascoli-Arzelà theorem we deduce that, up to a subsequence $\{t_{n_k}\}$, $w(t + t_{n_k}, s)$ converges in the C^2 -topology of $[-a, a] \times S^{N-1}$, for any $a > 0$, to a positive solution W of (4.49) in $\mathbb{R} \times S^{N-1}$ and $W \geq x_{1,M}$ and $W(0, s^*) = x_{1,M}$. Hence $W \equiv x_{1,M}$.

The proof of (4.46) under the second condition of (4.45) is similar.

4- Let (4.47)-(i) be satisfied and w be defined as in the previous steps. Then as in Step 3, $w(t_n, s)$ converges locally uniformly to a positive solution W of (4.49) defined on $\mathbb{R}_- \times S^{N-1}$, $w(t_n, s_n) \rightarrow W(0, s^*)$, $W(0, s^*)$ is a local maximum of W

and it is smaller than $x_{1,M}$. Hence $W_t(0, s^*) = |\nabla'W(0, s^*)| = 0$ and $W_{tt}(0, s^*) \leq 0$ and $\Delta'W(0, s^*) \leq 0$. Then

$$W^p(0, s^*) - M\alpha^q W^{\frac{2p}{p+1}}(0, s^*) + \alpha KW(0, s^*) \leq 0.$$

This contradicts the fact that \mathcal{P}_M defined in (1.7) is positive on the interval $(0, x_{1,M})$. Similarly, if (4.47)-(ii), we obtain that the limit function W and the limit point s^* where $W(0, s^*)$ is a local minimum of W satisfies $x_{1,M} < W(0, s^*) < x_{2,M}$ and

$$W^p(0, s^*) - M\alpha^q W^{\frac{2p}{p+1}}(0, s^*) + \alpha KW(0, s^*) \geq 0.$$

which is not compatible with the fact that \mathcal{P}_M is negative on $(x_{1,M}, x_{2,M})$. \square

Remark. We conjecture that the following stronger form of Theorem 4.16 holds:

1- Either u can be extended as a C^2 solution of (1.2) in Ω ,

2- or $\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_{2,M}$,

3- or $\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_{1,M}$.

In the case $M = m^*$, we prove the following.

Theorem 4.17 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded domain containing 0, $p > \frac{N}{N-2}$ and $M = m^*$. If u is a positive solution of (1.2) in $\Omega \setminus \{0\}$, there we have the following:*

1- either u can be extended as a C^2 solution in Ω ,

2- or there exists a sequence $\{r_n\}$ converging to 0 such that $r_n^\alpha u(r_n, s)$ converges to x_{m^*} uniformly on S^{N-1} .

Proof. If u satisfies (4.42), then the singularity of u at zero is removable since the function P_M is positive on $(0, m^*)$ and on (m^*, ∞) , see the argument in the proof of Theorem 4.16-(1). Thus we are left with the case

$$\liminf_{x \rightarrow 0} |x|^\alpha u(x) > 0. \quad (4.53)$$

If

$$\limsup_{x \rightarrow 0} |x|^\alpha u(x) = x_{m^*}, \quad (4.54)$$

then, as in the proof of Theorem 4.16-(3) we deduce that there exists a sequence $\{r_n\}$ converging to 0 such that $r_n^\alpha u(r_n, s)$ converges to x_{m^*} uniformly on S^{N-1} . If there exists $m \neq m^*$ such that

$$\limsup_{x \rightarrow 0} |x|^\alpha u(x) = m, \quad (4.55)$$

then there exists a sequence $\{t_n\}$ converging to $-\infty$ and $\{s_n\} \subset S^{N-1}$ such that $w(t+s_n, \cdot)$ converges in the C^2 topology of $[-a, a] \times S^{N-1}$ for any $a > 0$ to a solution W of

(4.49). Furthermore $w_t(t_n, s_n) \rightarrow 0$ and $\liminf_{t_n \rightarrow -\infty} w_{tt}(t_n, s_n) \leq 0$. Since $\nabla' w(t_n, s_n) = 0$ and $\Delta' w(t_n, s_n) \leq 0$, one has that

$$-\alpha K m - m^p + M(\alpha m)^{\frac{2p}{p+1}} = -m\tilde{P}_{m^*}(m^{\frac{p-1}{p+1}}) = -\Delta' W(0, s^*) - w_{tt}(0, s^*) \geq 0. \quad (4.56)$$

Since $\tilde{P}_{m^*} \geq 0$ and vanishes only at m^* , it implies $m = m^*$, contradiction. The proof of the uniform convergence of $w(t_n, \cdot)$ to m^* follows from the strong maximum principle since W is a positive solution (4.49) as in Theorem 4.16. \square

Remark. We conjecture that assertion (2) holds under the form

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_{m^*}.$$

Finally we have the following result dealing with the case $p = \frac{N}{N-2}$ and $M > 0$ where there exists a unique and explicit positive constant solution x_M to (1.4).

Theorem 4.18 *Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$, be a bounded domain containing 0, $p = \frac{N}{N-2}$ and $M > 0$. If u is a positive solution of (1.2) in $\Omega \setminus \{0\}$ which satisfies*

$$\limsup_{x \rightarrow 0} |x|^\alpha u(x) = x_M, \quad (4.57)$$

then

$$\lim_{x \rightarrow 0} |x|^\alpha u(x) = x_M, \quad (4.58)$$

Proof. Since the function \mathcal{P}_M is negative on $(0, x_M)$ and positive on (x_M, ∞) , for any $\epsilon > 0$ the function $x \mapsto (x_M - \epsilon)|x|^{-\alpha}$ is a subsolution of (1.2). The proof follows as in the proof of Theorem 4.16-(2). \square

Remark. We conjecture that the following dichotomy occurs: if u is a positive solution of (1.2) in $\Omega \setminus \{0\}$ unbounded near 0, then,

- 1- either (4.58) holds,
- 2- or (1.15)-(i) holds.

5 Behaviour at infinity of non-radial solutions

In this section we present some results dealing with the asymptotic behaviour of solutions which extend to the non-radial case what has already been proved in the radial one. The results are more complete if there exists only one possible behaviour for radial positive solutions; they have to be compared with what was obtained in [21] when $M = 0$.

5.1 The case $q = \frac{2p}{p+1}$

Theorem 5.1 *Let $p > 1$, $M > 0$ and u be a positive solution of (1.2) in $\mathbb{R}^N \setminus B_R$ with $N \geq 1$.*

1- *If $N = 1, 2$ and $p > 1$, or $N \geq 3$ and $1 < p < \frac{N}{N-2}$, then*

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_M. \quad (5.1)$$

2- *If $N \geq 3$, $p = \frac{N}{N-2}$, then*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} (\ln |x|)^{\frac{N-2}{2}} u(x) = \left(\frac{N-2}{\sqrt{2}} \right)^{N-2}. \quad (5.2)$$

3- *If $N \geq 3$, $p > \frac{N}{N-2}$ and $M < m^*$, then there exists $k > 0$ such that*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k. \quad (5.3)$$

Proof. The method of the proof is firstly to construct two positive radial solutions u_j , $j = 1, 2$ of (1.2) in B_R^c such that $u_1 \leq u \leq u_2$, and to use Proposition 1.6. The solution v of $\mathcal{L}_p v = 0$ in B_R^c with value $\min u|_{\partial B_R}$ for $|x| = R$ is a subsolution smaller than u . For cases 1 and 2, we can take for supersolution the function $u_{X_M} + \max u|_{\partial B_R}$. Therefore there exist two positive radial solutions u_1 and u_2 of (1.2) in B_R^c with respective value $\min u|_{\partial B_R}$ and $\max u|_{\partial B_R}$ on ∂B_R and such that

$$v(x) \leq u_1(x) \leq u(x) \leq u_2(x) \leq u_{X_M}(x) + \max u|_{\partial B_R} \quad \text{for } |x| \geq R, \quad (5.4)$$

by Theorem 2.4 and Theorem 2.5. Since u_1 and u_2 satisfy either (5.1) or (5.2) in cases 1 and 2 respectively, u shares this behaviour.

In case 3 with $p > \frac{N}{N-2}$ the function v satisfies the same behaviour (5.3) up to the constant $c > 0$ which is not fixed. By Theorem 2.6-(3), u_1 and u_2 satisfy (5.3) with two different constants $0 < c_1 \leq c_2$. In order to prove that (5.3) holds for some $c \in [c_1, c_2]$ we use the method introduced in [21]. We set $u(r, s) = r^{-\nu} w(t, s)$ with $\nu = N - 2$ and $t = \ln r$, then w satisfies

$$w_{tt} - \nu w_t + \Delta' w + e^{(N-p(N-2))t} \left(\nu^q M ((w_t - \nu w)^2 + |\nabla' w|^2)^{\frac{q}{2}} - w^p \right) = 0 \quad (5.5)$$

in $[0, \infty) \times S^{N-1}$. Since w and $\left(\nu^q M ((w_t - \nu w)^2 + |\nabla' w|^2)^{\frac{q}{2}} - w^p \right)$ are bounded, it follows from [10, Proposition 4.1] that there exists $c \geq 0$ such that $w(t, \cdot) \rightarrow c$ uniformly on S^{N-1} when $t \rightarrow \infty$. This ends the proof. \square

Theorem 5.2 *Let $N \geq 3$, $p > \frac{N}{N-2}$ and $M = m^*$. If u is a positive solution of (1.2) in $\mathbb{R}^N \setminus B_R$, we have the following alternative,*

1- *either there exists a sequence $\{r_n\}$ converging to infinity such that*

$$\lim_{r_n \rightarrow \infty} r_n^\alpha u(r_n, s) = x_{m^*}, \quad (5.6)$$

uniformly on S^{N-1} .

2- or there exists $k > 0$ such that

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k. \quad (5.7)$$

Proof. We can assume that u is continuous in B_R^c . By Theorem 2.6-(2), u is bounded from above by the function u_R where u_R is a radial solution of (1.2) in $\overline{B_R^c}$ which tends to ∞ when $r \downarrow R$ and satisfies $r^\alpha u_R(r) \downarrow x_{m^*}$ when $r \rightarrow \infty$. Hence

$$m := \limsup_{|x| \rightarrow \infty} |x|^\alpha u(x) \leq \sup_{|x| \geq R} |x|^\alpha u(x) \leq x_{m^*}. \quad (5.8)$$

We claim that either $m = 0$ or $m = x_{m^*}$. As in the proof of Theorem 4.17, there exists a sequence $\{t_n\}$ tending to ∞ and $\{s_n\} \subset S^{N-1}$ converging to s^* such that $w_n(t, \cdot) := w(t + t_n, \cdot)$ converges in the C^2 -topology of $[-a, a] \times S^{N-1}$ for any $a > 0$ to a solution W of (4.49) in $\mathbb{R} \times S^{N-1}$. Furthermore W achieves its maximal value m at $(0, s^*)$, hence $W_t(0, s^*) = 0$, $\nabla' W(0, s^*) = 0$, $\Delta' W(0, s^*) \leq 0$ and $W_{tt}(0, s^*) \leq 0$. Therefore

$$-m \tilde{\mathcal{P}}_{m^*}(m^{\frac{p-1}{p+1}}) = -\alpha K m - m^p + M(\alpha m)^{\frac{2p}{p+1}} \geq 0.$$

Since (4.56) holds this implies that either $m = 0$ or $m = x_{m^*}$. If $m = x_{m^*}$ it follows by the strong maximum principle, as in the proof of Theorem 4.16, that $w(t_n, s) \rightarrow x_{m^*}$ as $n \rightarrow \infty$, uniformly on S^{N-1} .

If $m = 0$, then

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = 0.$$

For any $a < x_{m^*}$ and $\rho > R$ such that $u|_{\partial B_\rho} \leq a$, we consider the problem

$$\begin{aligned} -v_{rr} - \frac{N-1}{r} v_r + v^p - M|v_r|^{\frac{2p}{p+1}} &= 0 \quad \text{in } (\rho, \infty) \\ v(\rho) &= \rho^{-\alpha} a \end{aligned} \quad (5.9)$$

Since the solution of

$$\begin{aligned} -v_{rr} - \frac{N-1}{r} v_r + v^p &= 0 \quad \text{in } (\rho, \infty) \\ v(\rho) &= \rho^{-\alpha} a \end{aligned} \quad (5.10)$$

is a subsolution and $x_{m^*} |x|^{-\alpha}$ a supersolution, the solution v exists and it is unique. By the phase plane analysis of Figure 4, the function $\tilde{v}(t) = e^{\alpha t} \tilde{u}(r)$ which initial value belongs to the region (F) converges to 0 when $t \rightarrow \infty$. Since $(0, 0)$ is a saddle point for the system (2.16) it follows that the corresponding trajectory is the unstable one of this point. The initial slope of this curve is $N - 2$. By Lemma A.4 it follows that there exists $\ell > 0$ such that

$$\lim_{r \rightarrow \infty} r^{N-2} v(r) = \ell. \quad (5.11)$$

Consequently $|x|^{N-2}u(x)$ is bounded, the proof of Theorem 5.1-3 applies and deduce from (5.8) there exists $c > 0$ such that $|x|^{N-2}u(x) \rightarrow c$ when $|x| \rightarrow \infty$. \square

In the case $M > m^*$ the situation is even more complicated and the results are still incomplete.

Theorem 5.3 *Let $N \geq 3$, $p > \frac{N}{N-2}$ and $M > m^*$. If u is a positive solution of (1.2) in $\mathbb{R}^N \setminus B_R$, we have the following,*

1-

$$\limsup_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{2,M} \implies \lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{2,M}, \quad (5.12)$$

2- *If $\liminf_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_{1,M}$, there exists a sequence $\{r_n\}$ tending to ∞ such that*

$$\lim_{r_n \rightarrow \infty} r_n^\alpha u(r_n, s) = x_{1,M}, \quad (5.13)$$

uniformly on S^{N-1} .

3- *If $\liminf_{|x| \rightarrow \infty} |x|^\alpha u(x) = 0$, there exists $k > 0$ such that*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2}u(x) = k. \quad (5.14)$$

Proof. By Theorem 2.6-(1) u is bounded from above by the solution u_R of (1.2) in $\mathbb{R}^N \setminus B_R$ which tends to infinity as $|x| \downarrow R$ and satisfies

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u_R(x) = x_{2,M}.$$

Hence

$$\tilde{m} := \liminf_{|x| \rightarrow \infty} |x|^\alpha u(x) \leq m := \limsup_{|x| \rightarrow \infty} |x|^\alpha u(x) \leq x_{2,M}$$

1- If $m = x_{2,M}$ there exists a sequence $\{t_n\}$ tending to ∞ and $\{s_n\} \subset S^{N-1}$ converging to s^* such that $w_n(t, \cdot) := w(t + t_n, \cdot)$ converges in the C^2 -topology of $[-a, a] \times S^{N-1}$ for any $a > 0$ to a solution W of (4.49) in $\mathbb{R} \times S^{N-1}$. Furthermore W achieves its maximal value m at $(0, s^*)$, hence $W_t(0, s^*) = 0$, $\nabla' W(0, s^*) = 0$, $\Delta' W(0, s^*) \leq 0$ and $W_{tt}(0, s^*) \leq 0$. Therefore

$$-m\tilde{\mathcal{P}}_M(m^{\frac{p-1}{p+1}}) = -\alpha K m - m^p + M(\alpha m)^{\frac{2p}{p+1}} \geq 0.$$

This implies that either $x_{1,M} \leq m \leq x_{2,M}$ or $m = 0$. For the liminf the same analysis yields that

$$-\tilde{m}\tilde{\mathcal{P}}_M(\tilde{m}^{\frac{p-1}{p+1}}) = -\alpha K \tilde{m} - \tilde{m}^p + M(\alpha \tilde{m})^{\frac{2p}{p+1}} \leq 0,$$

hence either $0 \leq \tilde{m} \leq x_{1,M}$ or $\tilde{m} \geq x_{2,M}$. Note that in the latter case (5.12) holds.

2- If $m = x_{2,M}$, then using the function W as in the proof of Theorem 5.2, we infer by the strong maximum principle that there exists a sequence $\{r_n\}$ tending to infinity such that $r_n^\alpha u(r_n, s)$ converges to $x_{2,M}$. For any $\epsilon > 0$ there exists $n_\epsilon > 0$ such that

for $n \geq n_\epsilon$ we have $r_n^\alpha u(r_n, s) \geq x_{2,M} - \epsilon$ for all $s \in S^{N-1}$. Since $\mathcal{P}_M(x_{2,M} - \epsilon) \leq 0$ the function $x \mapsto (x_{2,M} - \epsilon)|x|^{-\alpha}$ is a subsolution of (1.2). Then, for any $r_n > r_{n_\epsilon}$, $u(x) \geq (x_{2,M} - \epsilon)|x|^{-\alpha}$ in $\{x : r_{n_\epsilon} \leq |x| \leq r_n\}$. This implies

$$\liminf_{|x| \rightarrow \infty} |x|^\alpha u(x) \geq x_{2,M} - \epsilon.$$

Since ϵ is arbitrary, this yields (5.12).

If $\tilde{m} = x_{1,M}$ then we proceed as in the case above and deduce that there exist a function $W \geq 0$ satisfying (4.49) in $\mathbb{R} \times S^{N-1}$ and a sequence $\{t_n\}$ tending to infinity such that $w(t_n + t, s)$ converges in the C^2 -topology of $[-a, a] \times S^{N-1}$ to W for any $a > 0$. The function W is larger or equal to $x_{1,M}$ and coincides with $x_{1,M}$ at $(0, s^*)$ for some $s^* \in S^{N-1}$. By the strong maximum principle we have that $W \equiv x_{1,M}$. This implies assertion 2.

3- If $\liminf_{|x| \rightarrow \infty} |x|^\alpha u(x) = 0$, then we deduce by Harnack inequality that there exists a sequence $\{r_n\}$ tending to ∞ such that

$$\lim_{r_n \rightarrow \infty} r_n^\alpha u(r_n, s) = 0 \quad \text{uniformly on } S^{N-1}.$$

Then for any $\epsilon > 0$ the function there exists $n_\epsilon \in \mathbb{N}$ such that for any $n \geq n_\epsilon$, $u(r_n, s) \leq \epsilon |r_n|^{-\alpha}$ for all $s \in S^{N-1}$. The function $x \mapsto \epsilon |x|^{-\alpha}$ is a supersolution of (1.2). Since there exists a sequence $\{r_n\}$ tending to infinity such that $u(r_n, s) \leq \epsilon r_n^{-\alpha}$ for all $s \in S^{N-1}$ for $n \geq n_\epsilon$ it follows by the comparison principle applied to the sequence of annuli $\{x : r_{n_\epsilon} \leq |x| \leq r_n\}$, that $u(x) \leq \epsilon |x|^{-\alpha}$. Since the function $x \mapsto \epsilon |x|^{-\alpha}$ is a supersolution of (1.2), it follows by the comparison principle that $u(x) \leq \epsilon |x|^{-\alpha}$ in the annuli $\{x : r_{n_\epsilon} \leq |x| \leq r_n\}$. Letting $n \rightarrow \infty$ yields

$$u(x) \leq \epsilon |x|^{-\alpha} \quad \forall x \in B_{r_{n_\epsilon}} \implies \limsup_{|x| \rightarrow \infty} |x|^\alpha u(x) \leq \epsilon.$$

Since ϵ is arbitrary we infer that $|x|^\alpha u(x)$ converges to 0 when $|x| \rightarrow \infty$. By the phase plane analysis of section 2.4 (see Theorem 2.6), as in the proof of Theorem 5.2-2, we have that $|x|^{N-2} u(x)$ is bounded. Hence (5.14) follows as in the previous proof. \square

Remark. We conjecture that the results of Theorem 5.2 and Theorem 5.3 hold under the following forms:

For Theorem 5.2

1- either $\lim_{x \rightarrow \infty} |x|^\alpha u(x) = x_{m^*}$,

2- or there exists $k > 0$ such that $\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k$

For Theorem 5.3

1- either $\lim_{x \rightarrow \infty} |x|^\alpha u(x) = x_{2,M}$,

2- or $\lim_{x \rightarrow \infty} |x|^\alpha u(x) = x_{1,M}$,

3- or there exists $k > 0$ such that $\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k$.

5.2 The case $q \neq \frac{2p}{p+1}$

The next results extend the asymptotic behaviour described in Theorem 1.9 and Theorem 1.10 to non radial solutions. The following statement shows that equation (1.1) inherits the properties of the Emden-Fowler equation $\mathcal{L}_p u = 0$ if $\frac{2p}{p+1} < q < p$.

Theorem 5.4 *Let $N \geq 1$, $M > 0$, $\frac{2p}{p+1} < q < p$ and u be a positive solution of (1.1) in B_R^c . Then*

1- *If $N = 1, 2$ and $p > 1$, or $N \geq 3$ and $1 < p < \frac{N}{N-2}$, then*

$$\lim_{|x| \rightarrow \infty} |x|^\alpha u(x) = x_0. \quad (5.15)$$

2- *If $N \geq 3$ and $p > \frac{N}{N-2}$, then*

$$\lim_{|x| \rightarrow \infty} |x|^{N-2} u(x) = k > 0. \quad (5.16)$$

3- *If $N \geq 3$ and $p = \frac{N}{N-2}$, then*

$$\lim_{|x| \rightarrow \infty} (\ln |x|)^{\frac{N-2}{2}} |x|^{N-2} u(x) = \left(\frac{N-2}{\sqrt{2}} \right)^{N-2}. \quad (5.17)$$

Proof. In the first case, the solution v of $\mathcal{L}_p v = 0$ in B_R^c with value $\min u|_{\partial B_R}$ on ∂B_R is a subsolution smaller than u (it is obtained by minimization), and it has the behaviour expressed by (5.15). By Theorem 1.13 there exists a global positive solution \tilde{u} of (1.1) in $\mathbb{R}^N \setminus \{0\}$ satisfying (5.15). The difficulty is that this solution may not be larger than u for $|x| = R$. In such a case, for $a > \max u|_{\partial B_R}$, the function $\tilde{u}_a := \tilde{u} + a$ is a supersolution of (1.1) in B_R^c . The solution \tilde{v} which satisfies $\mathcal{L}_p \tilde{v} = 0$ in B_R^c with value a for $|x| = R$ is a subsolution smaller than \tilde{u}_a . Hence there exists a radial solution u_a of (1.1) in B_R^c such that $u_a(R) = a + \tilde{u}(R)$ and it dominates u in B_R^c . By Theorem 1.9-(1), the function u_a , and therefore u , satisfies (5.15).

In the second case, We proceed as in the proof of Theorem 5.1-(2), with the help of Theorem 1.9-(2). The function u satisfies $u_1 \leq u \leq u_2$ where u_1 and u_2 are radial solutions of (1.1) in B_R^c , hence $u(x) \leq C|x|^{2-N}$. If we set $w(t, s) = r^\nu u(r, s)$ with $t = \ln r$ and $\nu = N - 2$, then w satisfies

$$\begin{aligned} w_{tt} - \nu w_t + \Delta' w + \nu^q M e^{(N-q(N-1))t} ((w_t - \nu w)^2 + |\nabla' w|^2)^{\frac{q}{2}} \\ - e^{(N-p(N-2))t} w^p = 0 \quad \text{in } [0, \infty) \times S^{N-1}. \end{aligned} \quad (5.18)$$

Since w , w_t and $|\nabla' w|$ are bounded, it follows by the same argument [10, Proposition 4.1] that $w(t, \cdot) \rightarrow c \geq 0$ when $t \rightarrow \infty$ and $c > 0$ since u is bounded from below by

the solution of $\mathcal{L}^p v = 0$ in B_R^c which satisfies the same type of asymptotic behaviour with a positive limit of $r^{N-2}v(r)$ when $r \rightarrow \infty$.

In the third case, it is proved in [21, Théorème 3.1] that the solution v of $\mathcal{L}_p v = 0$ in B_R^c which coincide with $\min u|_{\partial B_R}$ for $|x| = R$ verifies the relation (5.17). In order to have the estimate from above, for $a > 0$ the function $h_a(x) = a|x|^{2-N}$ satisfies

$$\mathcal{L}_{p,q}^M h_a(x) = a^p|x|^{-p(N-2)} - Ma^q(N-2)^q|x|^{-q(N-1)} = a^p|x|^{-N} - Ma^q(N-2)^q|x|^{-q(N-1)}.$$

Since $q > \frac{2p}{p+1} = \frac{N}{N-1}$, we obtain for $|x| \geq R$

$$\begin{aligned} \mathcal{L}_{p,q}^M h_a(x) &= a^q|x|^{(1-N)q} (a^{p-a}|x|^{(N-1)q-N} - M(N-2)^q) \\ &\geq a^q|x|^{(1-N)q} (a^{p-a}R^{(N-1)q-N} - M(N-2)^q), \end{aligned}$$

Therefore, for $a > 0$ large enough $\mathcal{L}_{p,q}^M h_a \geq 0$ in B_R^c . Since the solution v_a of $\mathcal{L}_p v = 0$ in B_R^c with value aR^{2-N} for $|x| = R$ is a subsolution of (1.1) smaller than h_a it follows from [25, Theorem 1.4.5] that there exists a radial solution u_a of (1.1) in B_R^c such that $v_a \leq u_a \leq h_a$. If we choose a large enough so that $aR^{2-N} \geq \max u|_{\partial B_R}$, then u_a is larger than u in B_R^c . Finally, by Theorem 1.9-(3) the function u_a satisfies also (5.17), which ends the proof. \square

In the following result we extend Theorem 1.10-(2) to the non-radial case.

Theorem 5.5 *Let $N \geq 3$, $M > 0$, $p > 1$ and $1 < q < \frac{2p}{p+1}$ with $q \leq \frac{N}{N-1}$. If u is a positive solution of (1.1) in \overline{B}_R^c then there holds*

$$\lim_{|x| \rightarrow \infty} |x|^\gamma u(x) = X_M. \quad (5.19)$$

Proof. We can assume that u is continuous in B_ρ^c for any $\rho > R$.

For constructing a supersolution we proceed as in Theorem 4.14 using the solution v of (1.1) on $(0, \infty)$. The function $x \mapsto v(|x| - \rho) + \delta$ is a supersolution 1.1 in B_ρ^c which is larger than u for $|x| = \rho$ and at infinity. Hence it is larger than u in \overline{B}_ρ^c . Letting $\delta \rightarrow 0$ yields $u(x) \leq v(|x| - \rho)$ for all $x > \rho$.

Next, we construct a subsolution: we set $c = \min u|_{\partial B_\rho}$. For $n > \rho$ we denote by w_n the solution of

$$\begin{aligned} -\Delta w + w^p &= 0 \quad \text{in } \Gamma_{\rho,n} := B_n \setminus \overline{B}_\rho \\ w_n &= c \quad \text{on } \partial B_\rho, \quad w_n = 0 \quad \text{on } \partial B_n. \end{aligned}$$

The function w_n which is unique is a subsolution of 1.1 in $\Gamma_{\rho,n}$ where it satisfies $w_n \leq v(|x| - \rho)$. By Theorem 4.1 there exists a solution u_n of (1.1) in $\Gamma_{\rho,n}$ which coincides with w_n on $\partial\Gamma_{\rho,n}$ and is radial as w_n and $v(|\cdot| - \rho)$ are (or by uniqueness), and $u_n \leq u$ in $\Gamma_{\rho,n}$. When $n \rightarrow \infty$, $u_n \uparrow u_\infty$. The function u_∞ is a radial positive solution of (1.1) in \overline{B}_ρ^c and it satisfies

$$u_\infty(x) \leq u(x) \leq v(|x| - \rho) \quad \text{for all } |x| > \rho.$$

It follows from Theorem 1.10-(2) that u_∞ satisfies (5.19), as v does it too. This ends the proof. \square

A Appendix

A.1 The a priori estimates

The following a priori estimates proved in [7], [8] are fundamental throughout the paper. They do not depend on the sign of $q - \frac{2p}{p+1}$.

Proposition A.1 *Let $\Omega \subset \mathbb{R}^N$ be a domain containing 0, $1 < q < p$ and $M \in \mathbb{R}$. If $u \in C^1(\Omega \setminus \{0\})$ is a nonnegative solution of (1.1) in $\Omega \setminus \{0\}$, then there holds for any $0 < R \leq \frac{1}{2} \text{dist}(x, \partial\Omega)$:*

1- If $M > 0$,

$$u(x) \leq c_1 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\gamma}, |x|^{-\alpha} \right\} \quad \text{for all } x \in \overline{B}_R \setminus \{0\}, \quad (\text{A.1})$$

where $c_1 = c_1(N, p, q) > 0$.

2- If $M \leq 0$ and $q < 2$ if $M < 0$,

$$u(x) \leq \min \left\{ c_2 |x|^{-\alpha}, c_3 |M|^{-\frac{1}{q-1}} |x|^{-\beta} + \max_{|y|=R} u(y) \right\} \quad \text{for all } x \in \overline{B}_R \setminus \{0\}, \quad (\text{A.2})$$

where $c_2 = c_2(N, p) > 0$ and $c_3 = c_3(N, q) > 0$.

When u is a signed solution the following estimate holds [8, Corollary 2.2].

Proposition A.2 *Under the assumptions on Ω , p and q of Proposition A.1 and assuming that $M > 0$, any signed solution u of (4.20) in $\Omega \setminus \{0\}$ satisfies for any $0 < R \leq \frac{1}{2} \text{dist}(x, \partial\Omega)$,*

$$\begin{aligned} -\min \left\{ c_4 M^{\frac{1}{q-1}} |x|^{-\beta}, c_2 |x|^{-\alpha} \right\} &\leq -u_-(x) \leq 0 \\ &\leq u_+(x) \leq c_1 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{q}{p-q}}, |x|^{-\alpha} \right\}, \end{aligned} \quad (\text{A.3})$$

for all $x \in \overline{B}_R \setminus \{0\}$, where $c_1 = c_1(N, p, q) > 0$, $c_2 = c_2(N, p) > 0$ and $c_4 = c_4(N, p, q) > 0$.

Using scaling method when $1 < q \leq 2$ and the Bernstein method when $1 < q < p$, it is proved in [7, Proposition 2.3, Corollary 2.5] a gradient estimate that we recall.

Proposition A.3 *Let $\Omega \subset \mathbb{R}^N$ be a domain containing 0, $1 < q < p$ and $M > 0$. If $u \in C^1(\Omega \setminus \{0\})$ is a nonnegative solution of (1.1) in $\Omega \setminus \{0\}$, there holds for any $0 < R \leq R_0 := \frac{1}{2} \text{dist}(x, \partial\Omega)$ and some constants $c_4 = c_4(N, p, q, R_0) > 0$:*

1- When $1 < q \leq \frac{2p}{p+1}$

$$|\nabla u(x)| \leq c_4 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{p}{p-q}}, |x|^{-\frac{p+1}{p-1}} \right\}, \quad (\text{A.4})$$

for all $x \in \overline{B}_R \setminus \{0\}$. If $q = \frac{2p}{p+1}$, c_4 is independent of R_0 .

2- When $1 < q < p$

$$|\nabla u(x)| \leq c_5 \left(|x|^{-\frac{1}{q-1}} + \max \left\{ M^{\frac{p}{q(p-q)}} |x|^{-\frac{p}{p-q}}, |x|^{-\frac{2p}{q(p-1)}} \right\} \right), \quad (\text{A.5})$$

for all $x \in \overline{B}_R \setminus \{0\}$.

Remark. When $\frac{2p}{p+1} \leq q \leq 2$ any nonnegative solution of (1.1) in $\mathbb{R}^N \setminus B_{\frac{R}{2}}$ satisfies (A.4) in $\mathbb{R}^N \setminus B_R$.

Remark. If u is a signed solution, the Bernstein method that we developed in [7], [8] cannot be applied, however the scaling method can be used if $1 < q < p$. In the particular case $1 < q \leq \frac{2p}{p+1}$ there holds in a neighborhood of $x = 0$,

$$|\nabla u(x)| \leq c_5 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\gamma-1}, |x|^{-\alpha-1} \right\}. \quad (\text{A.6})$$

A.2 Equilibrium with a simple eigenvalue

Consider the system in \mathbb{R}^2

$$\begin{aligned} x_t &= ax + by + f(x, y) \\ y_t &= cx + dy + g(x, y) \end{aligned} \quad (\text{A.7})$$

where a, b, c, d are real numbers and f and g two C^1 real functions satisfying

$$|f(x, y) + g(x, y)| \leq c(|x|^s + |y|^s) \quad \text{for all } (x, y) \in B_1. \quad (\text{A.8})$$

Suppose that the matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of the system admits two eigenvalues $\mu_1 \neq \mu_2$ and $\mu_2 > 0$. By reduction to the diagonal form $\begin{pmatrix} x \\ y \end{pmatrix} = P \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix}$ the system becomes

$$\begin{aligned} \tilde{x}_t &= \mu_1 \tilde{x} + \tilde{f}(\tilde{x}, \tilde{y}) \\ \tilde{y}_t &= \mu_2 \tilde{y} + \tilde{g}(\tilde{x}, \tilde{y}). \end{aligned} \quad (\text{A.9})$$

Lemma A.4 *Under the above assumptions there exist at least two trajectories $\tilde{T}_1 = (\tilde{x}_1, \tilde{y}_1)$ and $\tilde{T}_2 = (\tilde{x}_2, \tilde{y}_2)$ tangent to the axis $0\tilde{y}$ converging to $(0, 0)$ when $t \rightarrow -\infty$, one with $\tilde{y}_1(t) > 0$, the other with $\tilde{y}_2(t) < 0$ for $t \leq -T$. Any trajectory $\tilde{T} = \{(\tilde{x}, \tilde{y})\}_{t \leq T}$ converging to $(0, 0)$ when $t \rightarrow -\infty$ and tangent at $(0, 0)$ to the axis $0\tilde{y}$ satisfies for some $\ell \neq 0$*

$$\lim_{t \rightarrow -\infty} e^{-\mu_2 t} \tilde{y}(t) = \ell. \quad (\text{A.10})$$

Proof. The existence of the solutions tangent to the axis $0\tilde{y}$ is classical. Consider a (\tilde{x}, \tilde{y}) converging to $(0, 0)$ tangentially to $0\tilde{y}$ and such that $\tilde{y}(t) > 0$ for $t \leq -T$. Then

$$|\tilde{g}(\tilde{x}(t), \tilde{y}(t))| \leq c(|\tilde{x}(t)|^s + |\tilde{y}(t)|^s) \leq c' |\tilde{y}(t)|^s$$

with $c' > c$, since $\frac{\tilde{x}(t)}{\tilde{y}(t)} \rightarrow 0$ when $t \rightarrow -\infty$. Put $v(t) = e^{-\mu_2 t} \tilde{y}(t)$. Then

$$|v_t(t)| = |\tilde{g}(\tilde{x}(t), \tilde{y}(t))| e^{-\mu_2 t} \leq c e^{-\mu_2 t + s \mu_2 t} v^s(t).$$

Therefore

$$\left| \left(\frac{v^{1-s}}{s-1} \right)_t \right| \leq c e^{\mu_2(s-1)t} \implies (v^{1-s})_t \in L^1(-\infty, T) \quad (\text{A.11})$$

Then $v^{1-s}(t)$ admits a nonnegative limit ℓ when $t \rightarrow -\infty$. If $\ell = 0$, it would follow from (A.11) that

$$(e^{\mu_2 t} v(t))^{s-1} \geq \tilde{c} \implies \tilde{y}(t) \geq C > 0,$$

which contradict the fact that $\tilde{y}(t) \rightarrow 0$. \square

Remark. This result is easily extendable to higher dimension, where A is a $N \times N$ matrix with a simple eigenvalue $\mu > 0$ and such that $\mathbb{R}^N = \ker(A - \mu I) \oplus E$, where E is A -invariant. Consider the system

$$X' = AX + F(X) \tag{A.12}$$

where $|F(X)| \leq c|X|^s$ in B_1 for some $s > 1$. If $X = X_1 + X'$ where $X_1 \in \ker(A - \mu I)$ and $X' \in E$, then there exist two trajectories $X_j(t)$ of (A.12) admitting a limit direction $\tau \in \ker(A - \mu I) \setminus \{0\}$ and , $j=1,2$, and they satisfy for some $a \neq 0$,

$$\lim_{t \rightarrow -\infty} e^{-\mu t} X(t) = a\tau. \tag{A.13}$$

References

- [1] C. Bandle and E. Giarrusso. Boundary blow-up for semilinear elliptic equation with nonlinear gradient term, *Adv. Diff. Equ.* **1**, 133-150 (1996).
- [2] C. Bandle and M. Marcus. Sur les solutions maximales de problèmes elliptiques non linéaires: bornes isopérimétriques et comportement asymptotique, *C. R. Acad. Sci. Paris* **311**, 91-93 (1990).
- [3] R. Bellman. *Stability theory of differential equations*, International Series in Pure and Appl. Math., Mac Graw-Hill (1953).
- [4] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron. Local and global properties of solutions of quasilinear Hamilton-Jacobi equations, *J. Funct. Anal.* **267**, 3294-3331 (2014).
- [5] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron. A priori estimates for elliptic equations with reaction terms involving the function and its gradient, *Math. Annalen* **378**, 13-56 (2020).
- [6] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron. Radial solutions of scaling invariant nonlinear elliptic equations with mixed reaction terms, *Discrete Cont. Dyn. Systems* **40**, 933-982 (2020).
- [7] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron. Measure data problems for a class of elliptic equations with mixed absorption-reaction, *Adv. Nonlinear Studies*, to appear <https://doi.org/10.1515/ans-2021-2124> (2021).
- [8] M.F. Bidaut-Véron, M. Garcia-Huidobro and L. Véron. Boundary singular solutions of a class of equations with mixed absorption-reaction, submitted, arXiv:2007.16097v2.
- [9] M.F. Bidaut-Véron and H. Giacomini. A new dynamical approach of Emden-Fowler equations and systems, *Adv. Diff. Equ.* **15**, 1033-1082 (2010).

- [10] M.F. Bidaut-Véron and Th. Raoux. Asymptotic of solutions of some nonlinear elliptic system, *Comm. Part. Diff. Equ.* **21**, 1035-1086 (1996).
- [11] L. Boccardo, F. Murat and J. P. Puel. Résultats d'existence pour certains problèmes quasilineaires, *Ann. Sc. Norm. Sup. Pisa, Classe di Scienze*, **Ser. 4, T. 11**, 213-255 (2015).
- [12] P. Felmer, A. Quass and B. Sirakov. Solvability of nonlinear elliptic equations with gradient terms, *J. Diff. Equ.*, **254**, 4327-4346 (2013).
- [13] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*, reprint of the 1998 edition, Classics in Mathematics, Springer-Verlag, New York (2001).
- [14] J. B. Keller. On the solutions of $\Delta u = f(u)$, *Comm. Pure Appl. Math.* **10**, 503-510 (1957).
- [15] S. Kichenassamy and L. Véron. Singular solutions of the p -Laplace equation, *Math. Ann* **275**, 599-615 (1986)
- [16] M. Marcus and L. Véron. Maximal solutions of semilinear elliptic equations with locally integrable forcing term, *Israel J. Math.* **152**, 333-348 (2006)
- [17] P. T. Nguyen. Isolated singularities of positive solutions with weighted gradient term, *Anal. P. D. E.* **9**, 1671-1692 (2016).
- [18] R. Osserman. On the inequality $\Delta u \geq f(u)$, *Pacific J. Math.* **7**, 1641-1647 (1957).
- [19] J. Serrin. Isolated singularities of solutions of quasilinear equations, *Acta Mathematica* **34**, 247-302 (1964).
- [20] Y. Richard and L. Véron. Isotropic singularities of solutions of nonlinear inequalities, *Ann. Inst. H. Poincaré* **6**, 37-72 (1989).
- [21] L. Véron. Comportement asymptotique des solutions d'équations elliptiques semi-linéaires dans \mathbb{R}^N , *Ann. Mat. Pura Appl.* **127**, 25-50 (1981).
- [22] L. Véron. Singular solutions of some nonlinear elliptic equations, *Nonlinear Anal. Theory, Methods Appl.* **5**, 225-242 (1981).
- [23] L. Véron. Singular solutions of some nonlinear elliptic equations, *Ann. Fac. Sci. Toulouse* **6**, 1-31 (1984).
- [24] L. Véron. Semilinear elliptic equations with uniform blow-up on the boundary, *J. Anal. Math.* **59**, 231-250 (1992).
- [25] L. Véron. *Local and Global Aspects of Quasilinear Degenerate Elliptic Equations*, World Scientific xvi+pp. 1-460 (2017).