ATIYAH SEQUENCE AND GAUGE TRANSFORMATIONS OF A PRINCIPAL 2-BUNDLE OVER A LIE GROUPOID

SAIKAT CHATTERJEE, ADITTYA CHAUDHURI, AND PRAPHULLA KOUSHIK

ABSTRACT. In this paper, a notion of a principal 2-bundle over a Lie groupoid has been introduced. For such principal 2-bundles, we produced a short exact sequence of VB-groupoids, namely, the Atiyah sequence. Two notions of connection structures viz. strict connections and semi-strict connections on a principal 2-bundle arising respectively, from a retraction of the Atiyah sequence and a retraction up to a natural isomorphism have been introduced. We constructed a class of principal $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundles and connections from a given principal G_0 -bundle $E_0 \rightarrow X_0$ over $[X_1 \rightrightarrows X_0]$ with connection. An existence criterion for the connections on a principal 2-bundle over a proper, étale Lie groupoid is proposed. The action of the 2-group of gauge transformations on the category of strict and semi-strict connections has been studied. Finally we noted an extended symmetry of the category of semi-strict connections.

CONTENTS

1.	Introduction	1
2.	Lie 2-groups, Lie crossed modules	4
3.	Principal 2-bundles over Lie groupoids	9
4.	Atiyah sequence for principal 2-bundles over Lie groupoids	18
5.	Connection on principal 2-bundles over Lie groupoids	21
6.	Gauge 2-group of a principal 2-bundle	33
Ac	knowledgements	37
Re	ferences	38

1. INTRODUCTION

The notion of (traditional) principal bundles and their connection structures over a manifold plays a central role at the interface of geometry and physics. In particular, a connection or, in the physics lexicon, a gauge field describes the dynamics of a particle. In the last decade or so, higher gauge theories were developed as frameworks to describe the dynamics of string-like extended 'higher dimensional objects'. A higher gauge theory typically involves suitable categorified versions of 'spaces'. For example, a manifold is replaced by a Lie groupoid or a category internal to smooth spaces, and a Lie group

²⁰¹⁰ Mathematics Subject Classification. Primary 53C08, Secondary 22A22, 58H05.

Key words and phrases. principal 2-bundles, Atiyah sequence, connections, gauge transformations, Lie groupoids.

is replaced by a Lie 2-group, a smooth map is replaced by a smooth functor, and so on, and an appropriate notion of a connection consistent with this categorification. In this context, intuitively, a principal 2-bundle would be understood as a smooth functor $\pi: \mathbb{E} \to \mathbb{B}$ between categories internal to smooth spaces with the structure group a suitably categorified version of a Lie group (weak or strict or coherent Lie 2-group). The precise descriptions of the 'categorical spaces' depend on the framework. Among the earliest works in this direction, we refer to the following papers [9, 6, 4, 3, 5], and the other papers cited therein. The list is, of course, far from complete.

More recently, Wockel in [45] introduced a notion of semi-strict principal 2-bundles over a discrete smooth 2-space and describes their classification up to Morita equivalence by non-abelian Cěch cohomology. A further generalization of Wockel's framework of principal 2-bundles can be found in [38]. Among other approaches to principal 2-bundles, one can mention non-abelian bundle gerbes of Aschieri, Cantini, and Jurčo [1], and *G*-gerbes [26] of Laurent-Gengoux, Stiénon, and Xu. [37] investigate the relation between non-abelian bundle gerbes of [1] and principal 2-bundles over a discrete smooth 2-space. Ginot and Stiénon introduced the notion of a principal 2-bundle over a Lie groupoid X with a strict structure Lie 2-group as a Hilsum & Skandalis generalized morphism of Lie 2-groupoids $X \to G$, where both X and G are treated as Lie 2-groupoids [21]. In particular, the paper shows that a *G*-gerbe is the same as (up to a Morita equivalence) a principal automorphism 2-group bundle.

The connections structure on the principal 2-bundles over a discrete smooth 2-space and its various aspects are also well studied. Some of the works which will be relevant in this context, are Breen-Messing [13], Baez-Schreiber [6, 39], Jurco-Samann-Wolf [23], Aschieri-Cantini-Jurco [1], Gengoux-Stienon-Xu [26]. In [43] Waldorf introduced the notion of a connection on Wockel's principal 2-bundle.

Some of the papers, which discuss higher gauge theory over path-space groupoids are [17, 19, 18, 31]. For higher gauge theories, which go beyond 2-spaces, we refer to [24, 7, 8] for a framework involving Kan simplicial manifolds, and more generally, for ∞ -topos theory framework, we refer to [20, 35, 36, 40].

In this paper, we introduce a notion of a principal 2-bundle $\mathbb{E} := [E_1 \Rightarrow E_0]$ over a Lie groupoid $\mathbb{X} := [X_1 \Rightarrow X_0]$, having a structure group a strict Lie 2-group $\mathbb{G} = [G_1 \Rightarrow G_0]$, as a smooth functor $\mathbb{E} \to \mathbb{X}$ with a functorial action $\mathbb{G} \times \mathbb{E} \to \mathbb{E}$ of \mathbb{G} on \mathbb{E} such that we have G_1 and G_0 -bundles $E_1 \to X_1$ and $E_0 \to X_0$ respectively over X_1, X_0 . The definition of 2-principal bundle we consider here similar to the one in [17] with suitable smoothness conditions, and the definition is slightly stronger than that of Ginot and Stiénon in [21].

The definition of a G-bundle over a Lie groupoid $[X_1 \rightrightarrows X_0]$, introduced in [27], can be viewed as a special case of ours. The Atiyah sequence associated to a principal G-bundle of [27] have been studied in [10, 11]. The main purpose of this paper is to explore the connections and gauge transformations on a strict Lie 2-group bundle over a Lie groupoid in terms of the Atiyah sequence. Recall that in the traditional set-up, with a principal G-bundle $P \rightarrow M$ one can associate the Atiyah sequence, a short exact sequence of vector bundles over M [2],

$$0 \to \operatorname{Ad}(P) \xrightarrow{j^{/G}} \operatorname{At}(P) \xrightarrow{\pi^{/G}_*} TM \to 0.$$

Here L(G) denote the Lie algebra of G, $Ad(P) := (P \times L(G))/G \to M$ and At(P) := $TP/G \rightarrow M$ are respectively the adjoint bundle and the Atiyah vector bundle of the principal G-bundle $\pi: P \to M$. Then a retraction of the Atiyah sequence is a connection on $P \to M$. We show that associated to a principal 2-bundle $\mathbb{E} := [E_1 \rightrightarrows E_0]$ over a Lie groupoid $\mathbb{X} := [X_1 \rightrightarrows X_0]$, we have a short exact sequence of VB-groupoids (for the discussion on VB-groupoids, we refer to [22, 29]). Then a 'strict connection' on the principal 2-bundle is defined as a retraction of the short exact sequence of the VBgroupoids. However, enrichment is more interesting when we impose an obstruction, given by a certain natural isomorphism, on the retraction. The corresponding structure thus obtained will be called a 'semi-strict connection'. Just as a side note on our terminology, we remark here that we reserve the word 'weak connection' for a more general set-up of a simplicial principal bundle obtained from a principal 2-bundle as above, which we will pursue in our subsequent work. The category of gauge transformations on our principal 2-bundles naturally defines a 2-group, which acts on the category of strict and semi-strict connections. We show that the category of semi-strict connections enjoys an 'extended gauge symmetry.' In particular, considering the category of semi-strict connections on a 2-bundle over a discrete Lie groupoid $[X \rightrightarrows X]$, this extended gauge symmetry offers a natural interpretation of the gauge transformation of the connection 1-forms in higher BF theories (for example, see [30, 33, 31]).

Along the course of this paper, we illustrate our constructions and results with several examples and highlight the relations with other relevant works. We believe that the framework provides a general treatment of connections on Lie 2-group bundles over a Lie groupoid. We show that given a Lie group *G*-bundle over a Lie groupoid $[X_1 \Rightarrow X_0]$ with connection, in the sense of [27], one can construct a Lie 2-group \mathbb{G} bundle with a connection, with $Obj(\mathbb{G}) = G$. Moreover, the construction gives us a criterion for the existence of strict and semi-strict connections on a principal 2-bundle over a proper, étale Lie groupoid (or an orbifold). As an outcome, we prove the existence of connections on a Lie 2-group bundle over a discrete Lie groupoid (compare, [43, Theorem 5.2.14]). We also emphasize that to the best of our knowledge, the approach of the Atiyah sequence has not been explored in the higher gauge theory literature, other than a casual reference in the lecture notes [42]. Particularly one interesting outcome of this approach is the notion of semi-strict connections on a Lie 2-group bundle. Having said that, we have not studied Chern-Weil theory or Morita invariance of our construction for this paper. In a sequel of this paper, we intend to study these topics in the set-up of simplicial principal bundles.

The outline and organization of the paper. Section 2 recalls the definition of a strict Lie 2-group, its correspondence with a Lie crossed module, and some properties and identities of the associated Lie 2-algebra.

In Section 3 we introduced the definition of a principal 2-bundle over a Lie groupoid and constructed a principal 2-bundle, namely a decorated bundle over a Lie groupoid from a Lie group bundle over the same. We introduce a notion of a categorical connection and characterize the decorated principal bundles. We also explored the relationship with a 'Lie groupoid G-extension'.

In Section 4 we show that a principal 2-bundle over a Lie groupoid produces a short exact sequence of VB-groupoids, namely the Atiyah sequence.

In Section 5 we introduce the strict and semi-strict connections on a principal 2-bundle, respectively, as a retraction of the associated Atiyah sequence and a retraction up to a certain natural isomorphism. We express the strict and semi-strict connections in terms of Lie 2-algebra valued differential forms. Further, we give an explicit construction of the connection structure on a decorated bundle and obtain a criterion for the existence of a connection on a principal 2-bundle over an orbifold.

In Section 6, we introduce the notion of the 2-group of the gauge transformations on a principal 2-bundle. The gauge 2-group naturally acts on the categories of strict and semi-strict connections. We observe an extended gauge symmetry of the category of semistrict connections, and we relate to the gauge transformations of the connection 1-forms in higher BF theories.

Notations and conventions: Here, we fix some conventions and notations which will be followed throughout this paper.

All manifolds assumed to be the smooth, second countable, Hausdorff. Let Man be category of such manifolds. For a smooth map $f: M \to N$, the differential at $m \in M$ will be denoted as $f_{*,m}: T_m M \to T_{f(m)}N$. A smooth right (resp. left) action of a Lie group Gon a smooth manifold P will be denoted as $(p,g) \mapsto pg$ (resp. $(g,p) \mapsto gp$), for $g \in G$ and $p \in P$. The corresponding differentials $T_pP \to T_{pg}P$ or $T_pP \to T_{gp}P$ will be denoted respectively as $v \mapsto v \cdot g$ or $v \mapsto g \cdot v$, for $v \in T_pP$. For a fixed $p \in P$, let $\delta_p: T_eG \to T_pP$ denote the differential of the map $G \to P$, $g \mapsto pg$, at the identity element. We will denote the fundamental vector field corresponding to an element $B \in L(G)$ evaluated at p by $\delta_p(B)$.

We assume our Lie groups to be matrix groups for computational simplicity. For a Lie group G, we denote its Lie algebra by L(G).

In any category, s, t will denote the source and target maps, and the composition of a pair of morphisms f_2, f_1 will be denoted as $f_2 \circ f_1$, where $t(f_1) = s(f_2)$.

By a Lie groupoid we mean a groupoid object in Man, whose source, target maps are surjective submersions. We use the blackboard bold notation to denote Lie groupoids; that is, \mathbb{E} for $[E_1 \Rightarrow E_0]$ and so on. Given a Lie groupoid $\mathbb{X} = [X_1 \Rightarrow X_0]$, we have the associated tangent Lie groupoid, denoted $T\mathbb{X} = [TX_1 \Rightarrow TX_0]$, structure maps of $T\mathbb{X}$ are given by the differentials of the respective structure maps of \mathbb{X} . In particular if $(\gamma_2, X_2), (\gamma_1, X_1)$ are composable morphisms in $T\mathbb{X}$, then we denote $(\gamma_2, X_2) \circ (\gamma_1, X_1) =$ $(m(\gamma_2, \gamma_1), m_{*(\gamma_2, \gamma_1)}(X_2, X_1))$ as $(\gamma_2 \circ \gamma_1, X_2 \circ X_1)$, where *m* is the composition map of the Lie groupoid \mathbb{X} .

2. LIE 2-GROUPS, LIE CROSSED MODULES

In this section, we briefly overview the notions of Lie 2-groups and Lie crossed modules. We then describe a correspondence between the two. This section does not contain any new material. For further reading on this topic, we refer to several papers by Baez, Schreiber, Lauda, Crans and others, [6, 5, 4, 3, 17, 19].

Definition 2.1 (strict 2-category [12]). A strict 2-category \mathcal{D} , consists of the following data,

(1) a collection of objects,

- (2) a small category $\mathcal{D}(x, y)$ for each pair of objects (x, y) in \mathcal{D} ,
- (3) a functor $\{1\} \to \mathcal{D}(x, x)$ for each object x in \mathcal{D} ,
- (4) a functor $\circ_{xyz} : \mathcal{D}(x,y) \times \mathcal{D}(y,z) \to \mathcal{D}(x,z)$ for each triple (x,y,z) of objects in \mathcal{D} ,

satisfying the usual strict associativity and unital compatibility conditions.

Definition 2.2. A strict Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$ is a group object in the category of Lie groupoids.

Though there are more general versions of Lie 2-groups, as we will be working with only strict Lie 2-groups, by a Lie 2-group we always mean a strict Lie 2-group in this paper. For a more general notion of Lie 2-groups, we refer the reader to [5].

Remark 2.1. One can show that a Lie 2-group is a Lie groupoid \mathbb{G} equipped with a morphism of Lie groupoids $\otimes : \mathbb{G} \times \mathbb{G} \to \mathbb{G}$ such that \otimes induce Lie group structures on both $\operatorname{Obj}(\mathbb{G})$ and $\operatorname{Mor}(\mathbb{G})$, and all the structure maps are Lie group homomorphisms. Let $k_2, k'_2 \in \operatorname{Mor}(\mathbb{G})$ and $k_1, k'_1 \in \operatorname{Mor}(\mathbb{G})$ be pairs of composable morphisms. Let $^{-1}$ and \circ respectively denote the group inverse functor and the composition. Then

$$(k_2 \otimes k_1) \circ (k'_2 \otimes k'_1) = (k_2 \circ k'_2) \otimes (k_1 \circ k'_1), (k_2 \circ k_1)^{-1} = k_2^{-1} \circ k_1^{-1}.$$

$$(2.1)$$

Another description of Lie 2-group is given by a Lie crossed module which we define below:

Definition 2.3. A Lie crossed module is a 4-tuple

$$(G, H, \tau, \alpha)$$

where, G, H are Lie groups, $\alpha : G \times H \to H$ is a smooth map giving an action of G on H, and $\tau : H \to G$ is a morphism of Lie groups such that the following conditions are satisfied:

$$\tau(\alpha(g,h)) = g\tau(h)g^{-1} \text{ for all } (g,h) \in G \times H,$$

$$\alpha(\tau(h),h') = hh'h^{-1} \text{ for all } h,h' \in H.$$
(2.2)

The conditions in Equation (2.2) are called *Peiffer identities*.

2.1. Lie 2-group associated to a Lie crossed module. Given a Lie crossed module (G, H, τ, α) , the associated Lie 2-group is given by the Lie groupoid $\mathbb{G} = [H \rtimes_{\alpha} G \rightrightarrows G]$, where $H \rtimes_{\alpha} G$ denotes the semidirect product of groups H and G with respect to the action α of G on H. The various maps and operations on this Lie 2-group are as listed below.

- the source map is given by s(h,g) = g,
- the target map is given by $t(h,g) = \tau(h)g$
- the composition map is given by $m((h_2, g_2), (h_1, g_1)) = (h_2 h_1, g_1),$
- the identity map is given by $1_g = (e_H, g)$
- the inverse map is given by $i(h,g) = (h^{-1}, \tau(h)g)$

• the group properties of $H \rtimes_{\alpha} G$ are those of the standard semidirect product of the groups, that is the bi-functor $\otimes : \mathbb{G} \times \mathbb{G} \to \mathbb{G}$ is defined as

$$\bigotimes_0 : (g_1, g_2) \mapsto g_1 g_2, \otimes_1 : ((h_2, g_2), (h_1, g_1)) \mapsto (h_2 \alpha(g_2, h_1), g_2 g_1),$$

$$(2.3)$$

whereas the group inverse and the identity elements are respectively given by $(h, g)^{-1} = (\alpha(g^{-1}, h^{-1}), g^{-1})$ and (e_H, e_G) . It is a routine verification that \mathbb{G} is a Lie 2-group. We call \mathbb{G} to be the Lie 2-group associated to the Lie crossed module (G, H, τ, α) .

2.2. Lie crossed module associated to a Lie 2-group. Let $\mathbb{G} = [G_1 \rightrightarrows G_0]$ be a Lie 2-group. Consider the Lie group ker $(s) = \{\gamma \in G_1 : s(\gamma) = 1_{G_0}\}$, and the morphism $\alpha : G_0 \times \ker(s) \rightarrow \ker(s)$ defined as $(a, \gamma) \mapsto 1_a \cdot \gamma \cdot 1_{a^{-1}}$. Then, the 4-tuple

 $(G_0, \ker(s), t|_{\ker(s)} : \ker(s) \to G_0, \alpha : G_0 \times \ker(s) \to \ker(s)),$

defines a Lie crossed module. We call the above Lie crossed module to be the *Lie crossed* module associated to the Lie 2-group \mathbb{G} .

Example 2.2. The Lie crossed module associated to the discrete Lie 2-group $[G \rightrightarrows G]$ is $(G, \{e\})$ with trivial τ and α .

Example 2.3. At the other extreme of the last example, one can consider the Lie 2-group associated to a crossed module ($\{e\}, H$) with trivial τ and $\alpha = \text{Id} \colon H \to H$. It is obvious that the Lie 2-group is the single object Lie 2-group $[H \rightrightarrows e]$

Example 2.4. The Lie 2-group associated to the Lie crossed module (G, G, τ, α) with $\tau: G \to G$ being the identity map, and $\alpha: G \times G \to G$ the action of G on itself by conjugation is the pair groupoid $[G \times G \rightrightarrows G]$.

Example 2.5. To any simply connected Lie group G one associates a Lie crossed module $(\operatorname{Aut}(G), G, \tau, \alpha)$, where $\tau: G \to \operatorname{Aut}(G)$ sends an element $g \in G$ to its innerautomorphism, and $\alpha: \operatorname{Aut}(G) \to \operatorname{Aut}(G)$ is the identity. The corresponding Lie 2-group is called the *automorphism* 2-group of G.

We will not spend much time on examples of Lie 2-groups and Lie crossed modules, which abound in higher gauge theory literature. An interested reader can go through the articles previously mentioned.

Given a Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$ we can associate the Lie groupoid $[L(G_1) \rightrightarrows L(G_0)]$ whose structure maps are given by taking differentials of the structure maps of \mathbb{G} at the identity.

Remark 2.6. The vector space structures on $L(G_1)$ and $L(G_0)$ are compatible with the composition of $[L(G_1) \rightrightarrows L(G_0)]$.

In particular, if the Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$ is obtained from a Lie crossed module $(G, H, \tau : H \rightarrow G, \alpha : G \times H \rightarrow H)$ then the groupoid $[L(G_1) \rightrightarrows L(G_0)]$ is described as $[L(H) \oplus L(G) \rightrightarrows L(G)]$ having structure maps

- the source map is given by s(A, B) = B,
- the target map is given by $t(A, B) = \tau(A) + B$,
- the composition map is given by $m((A_2, B_2), (A_1, B_1)) = (A_2 + A_1, B_1)$, where $s(A_2, B_2) = t(A_1, B_1)$,

- the identity map is given by $1_B = (0, B)$,
- the inverse map is given by $i(A, B) = (-A, \tau(A) + B)$.

Remark 2.7. It should be noted that $L(G_1)$ splits into the direct sum $L(H) \oplus L(G)$ only as a vector space. It is not a direct sum of Lie algebras.

The differential on the action α produces an action $\alpha_{*,(e,e)} \colon L(G) \times L(H) \to L(H)$ of L(G)on L(H) as a Lie algebra derivation, and the commutators on L(G) and $L(H \rtimes G)$ are respectively given as

$$(B_1, B_2) \mapsto [B_1, B_2],$$

$$((A_1, B_1), (A_2, B_2)) \mapsto [(A_1, B_1), (A_2, B_2)]$$

$$:= ([A_1, A_2] + \alpha_{*,(e,e)}(A_1, B_2) - \alpha_{*,(e,e)}(A_2, B_1), [B_1, B_2]),$$

$$(2.4)$$

for all $A_1, A_2 \in L(H), B_1, B_2 \in L(G)$. The Lie groupoid $[L(G_1) \rightrightarrows L(G_0)]$ is the usual strict Lie 2-algebra of the Lie 2-group $[G_1 \rightrightarrows G_0]$. We will not spell out the general definition of a Lie 2-algebra. For a detailed discussion on Lie 2-algebras and the associated Lie 2-algebra of a Lie 2-group we refer to [4].

Let $(G, H, \tau : H \to G, \alpha : G \times H \to H)$ be the Lie crossed module associated to the Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$. Consider the smooth map $\alpha : G \times H \to H$. For a fixed $g \in G$, we have the map $\alpha(g) : H \to H, h \mapsto \alpha(g, h)$ and the differential of this map at the identity element of H gives the Lie algebra homomorphism

$$\alpha(g)_{*,e_H} \colon L(H) \to L(H). \tag{2.5}$$

On the other hand for a fixed $h \in H$, we have the map $\bar{\alpha}(h) \colon G \to H, g \mapsto \alpha(g, h)$ and by taking the differential of this map at the identity element of G we obtain the linear map,

$$\bar{\alpha}(h)_{*,e_G} \colon L(G) \to T_h H.$$
 (2.6)

Remark 2.8. To simplify our notations and avoid notational cluttering, henceforth, we adopt the following conventions.

- τ_{*,e_H} will also be denoted as τ . Similarly $\alpha(g)_{*,h}$, $\alpha(g)_{*,e_H}$ and $\bar{\alpha}(h)_{*,e_G}$ will be denoted respectively as $\alpha(g)_h, \alpha(g)$ and $\bar{\alpha}(h)$.
- To avoid cluttering of parentheses we often write $\alpha(g)$ etc as α_g etc.
- Unless necessary, we do not notationally distinguish between identity elements of various groups. The common notation for the same will be e.

For future reference, we note the following identities. The identities are technical and would not be required anywhere else other than the proofs of Corollary 5.10 and Lemma 5.18. So a reader may postpone its reading till then.

Lemma 2.9. Let (G, H, τ, α) be a Lie crossed module.

(1) Then for any $h_2, h_1 \in H$ and $B \in L(G)$, we have

$$h_2 \cdot \left(\bar{\alpha}(h_2^{-1})(\mathrm{ad}_{\tau(h_1)}B)\right) + h_1 \cdot \left(\bar{\alpha}(h_1^{-1})(B)\right) = h_2 h_1 \cdot \left(\bar{\alpha}(h_1^{-1}h_2^{-1})(B)\right).$$

(2) Then for any $g \in G, h \in H, A \in L(H)$

$$\alpha_{g^{-1}}(h^{-1}) \cdot \left(\bar{\alpha}_{\alpha_{g^{-1}}(h)}(\mathrm{ad}_{g^{-1}}(\tau(A)) + \alpha_{g^{-1}}(A) = \alpha_{g^{-1}}(\mathrm{ad}_{h^{-1}}(A)).$$

Proof. (1) Note that the required identity follows directly as a infinitesimal version of the equation (1)

$$\left[h_2\left(\bar{\alpha}(h_2^{-1})(\mathrm{Ad}_{\tau(h_1)}(g))\right)\right]\left[h_1\left(\bar{\alpha}(h_1^{-1})(g)\right)\right] = h_2h_1\left(\bar{\alpha}(h_1^{-1}h_2^{-1})(g)\right)$$

for any $h_2, h_1 \in H$ and $g \in G$. In particular, using Equation (2.2) the expression

$$h_2\left[\left(\bar{\alpha}(h_2^{-1})(\operatorname{Ad}_{\tau(h_1)}(g))\right)\right]$$

reduces to $h_2 \left[h_1 \alpha(g) (h_1^{-1} h_2^{-1} h_1) h_1^{-1} \right]$. Then a straight forward calculation will prove the identity.

(2) A straightforward verification gives

$$\left[\alpha_{g^{-1}}(h^{-1}) \cdot \left(\bar{\alpha}_{\alpha_{g^{-1}}(h)}(\mathrm{Ad}_{g^{-1}}(\tau(h'))\right)\right] \left[\alpha_{g^{-1}}(h')\right] = \alpha_{g^{-1}}(\mathrm{Ad}_{h^{-1}}(h')),$$

for any $h, h' \in H, g \in G$. Then the identity follows as the infinitesimal version of the above equation.

2.3. Adjoint actions of a Lie 2-group. Let $\mathbb{G} = [G_1 \rightrightarrows G_0]$ be a Lie 2-group and (G, H, τ, α) the associated Lie crossed module. It would be convenient to write down the adjoint actions for the object and morphisms groups in terms of a crossed module. As we have seen $G_0 = G$ and $G_1 = H \rtimes_{\alpha} G$.

In order to simplify our calculation of the adjoint action for $H \rtimes_{\alpha} G$, we write $\operatorname{Ad}_{(h,g)}(h',g')$ as $\operatorname{Ad}_{(h,g)}(h',g') = [\operatorname{Ad}_{(h,g)}(h',e)][\operatorname{Ad}_{(h,g)}(e,g')]$ and then compute the bracketed terms on the right hand side separately. The group product in Equation (2.3) gives

$$Ad_{(h,g)}(h',e) = \left(Ad_{h}(\alpha_{g}(h')), e\right),$$

$$Ad_{(h,g)}(e,g') = \left(h\left(\alpha_{(Ad_{g}(g'))}(h^{-1})\right), Ad_{g}(g')\right) = \left(h\left(\bar{\alpha}_{h^{-1}}(Ad_{g}(g'))\right), Ad_{g}(g')\right), \quad (2.7)$$

$$Ad_{(h,g)}(h',g') = [Ad_{(h,g)}(h',e)][Ad_{(h,g)}(e,g')].$$

Now given an element $(A, B) \in L(H) \oplus L(G) = L(G_1)$, we write (A, B) = (A, 0) + (0, B). Then the adjoint actions of $H \rtimes G$ on $A \in L(H)$ and $B \in L(G)$ are respectively computed by the formulae $\operatorname{ad}_{(h,g)}(A) := \frac{\mathrm{d}}{\mathrm{d}t} \operatorname{Ad}_{(h,g)} \operatorname{e}^{tA}|_{t=0}$ and $\operatorname{ad}_{(h,g)}(B) := \frac{\mathrm{d}}{\mathrm{d}t} \operatorname{Ad}_{(h,g)} \operatorname{e}^{tB}|_{t=0}$,

$$ad_{(h,g)}(A,0) = \left(ad_{h}(\alpha_{g}(A)), 0\right),$$

$$ad_{(h,g)}(0,B) = \left(h \cdot (\bar{\alpha}_{h^{-1}}(ad_{g}(B))), ad_{g}(B)\right),$$

$$ad_{(h,g)}(A,B) = [ad_{(h,g)}(A,0)] + [ad_{(h,g)}(0,B)].$$

(2.8)

The notations $\alpha_g, \bar{\alpha}_{h^{-1}}$ in Equation (2.8) should be interpreted as per Remark 2.8 (third and fourth bullet points).

3. PRINCIPAL 2-BUNDLES OVER LIE GROUPOIDS

In this section, we introduce the notion of a principal 2-bundle over a Lie groupoid. The definition we introduce is a natural generalization of the classical definition.

Definition 3.1 (Action of a Lie 2-group on a Lie groupoid). An *action of a Lie 2-group* \mathbb{G} on a Lie groupoid \mathbb{X} is defined as a morphism of Lie groupoids $\rho : \mathbb{X} \times \mathbb{G} \to \mathbb{X}$ such that

- $\rho_0: X_0 \times G_0 \to X_0,$
- $\rho_1: X_1 \times G_1 \to X_1,$

are Lie group actions on manifolds X_0 and X_1 respectively.

Following proposition is evident.

Proposition 3.1. Let G be a Lie 2-group.

- (1) There is an action of Lie 2-group \mathbb{G} on Lie groupoid $L(\mathbb{G}) = [L(G_1) \rightrightarrows L(G_0)]$ by adjoint action.
- (2) Suppose \mathbb{G} acts on a Lie groupoid \mathbb{X} . Then there is an action of \mathbb{G} on Lie groupoids $T\mathbb{X} = [TX_1 \rightrightarrows TX_0]$, given by the differential of the given action.

A weaker definition compared to the one given in Definition 3.1 also exists in literature; for example, in [45], the identity and compatibility axioms of a group action at the object level hold up to an isomorphism. One can also generalize this action by replacing the category $\mathbb{X} \times \mathbb{G}$ by a 'twisted product category' $\mathbb{X} \rtimes_{\eta} \mathbb{G}$ of \mathbb{X} and \mathbb{G} with respect to certain map η , introduced in [18]. While we will stick to the definition of action given in Definition 3.1 for this paper and not pursue either of the other two generalizations very seriously, we recall the definition of a twisted product to relate with the notion of 'Lie groupoid *G*-extensions'.

Let \mathbb{G} and \mathbb{X} be as before. Consider a smooth map $\eta \colon \operatorname{Mor}(\mathbb{X}) \times \operatorname{Mor}(\mathbb{G}) \to \operatorname{Mor}(\mathbb{X})$ satisfying,

$$\eta(\gamma, k) \in \operatorname{Hom}_{\mathbb{X}}(x, y), \forall \gamma \in \operatorname{Hom}_{f} \mathbb{X}(x, y),$$

$$\eta(\gamma_{2} \circ \gamma_{1}, k) = \eta(\gamma_{2}, k) \circ \eta(\gamma_{1}, k),$$

$$\eta(1_{x}, k) = 1_{x}$$

$$\eta(\gamma, k_{2} \circ k_{1}) = \eta(\eta(\gamma, k_{2}), k_{1}),$$

$$\eta(\gamma, 1_{q}) = \gamma.$$

(3.1)

Then we have a smooth category $\mathbb{X} \rtimes_{\eta} \mathbb{G}$ with the following description [18, Proposition 5.1]:

$$\begin{aligned}
\operatorname{Obj}(\mathbb{X} \rtimes_{\eta} \mathbb{G}) &= \operatorname{Obj}(\mathbb{X}) \times \operatorname{Obj}(\mathbb{G}), \\
\operatorname{Mor}(\mathbb{X} \rtimes_{\eta} \mathbb{G}) &= \operatorname{Mor}(\mathbb{X}) \times \operatorname{Mor}(\mathbb{G}), \\
s(\gamma, k) &= (s(\gamma), s(k)), t(\gamma, k) = (t(\gamma), t(k)) \\
(\gamma_{2}, k_{2}) \circ_{\eta} (\gamma_{1}, k_{1}) &= (\gamma_{2} \circ \eta(\gamma_{1}, k_{2}), k_{2} \circ k_{1}).
\end{aligned}$$
(3.2)

The category $\mathbb{X} \rtimes_{\eta} \mathbb{G}$ will be called the η -twisted category.

Definition 3.2 (Twisted action of a Lie 2-group on a Lie groupoid). Let X be a Lie groupoid, \mathbb{G} a Lie 2-group and $\eta: \operatorname{Mor}(\mathbb{X}) \times \operatorname{Mor}(\mathbb{G}) \to \operatorname{Mor}(\mathbb{X})$ a smooth map satisfying conditions in Equation (3.1). An η -twisted action of \mathbb{G} on the Lie groupoid \mathbb{X} is defined as a smooth functor $\rho \colon \mathbb{X} \rtimes_{\eta} \mathbb{G} \to \mathbb{X}$ such that

- $\rho_0: X_0 \times G_0 \to X_0,$ $\rho_1: X_1 \times G_1 \to X_1,$

are Lie group actions on manifolds X_0 and X_1 respectively.

Remark 3.2. (1) Note that if we take $\eta = pr_1$ in Definition 3.2 then we recover the usual direct product of categories.

(2) Functoriality of $\rho : \mathbb{X} \rtimes_{\eta} \mathbb{G} \to \mathbb{X}$ implies $\rho(\gamma_2 \circ \eta(\gamma_1, k_2), k_2 \circ k_1) = \rho(\gamma_2, k_2) \circ \rho(\gamma_1, k_1)$ and following our convention for group actions we write,

$$(\gamma_2 \circ \eta(\gamma_1, k_2))(k_2 \circ k_1) = (\gamma_2 k_2) \circ (\gamma_1 k_1).$$

Definition 3.3. Let \mathbb{G} be a Lie 2-group. Suppose \mathbb{G} acts on a pair of Lie groupoids \mathbb{X} and \mathbb{Y} . Then a morphism of Lie groupoids $F := (F_1, F_0) : \mathbb{X} \to \mathbb{Y}$ is said to be \mathbb{G} -equivariant if F_0 is G_0 -equivariant and F_1 is G_1 -equivariant. A smooth natural transformation between two such functors $\eta: F \Longrightarrow F'$ is called a G-equivariant natural transformation if $\eta(xg) =$ $\eta(x)\mathbf{1}_{q}.$

Definition 3.4 (principal G-bundle over a Lie groupoid). Let G be a Lie 2-group. A principal G-bundle over a Lie groupoid X is given by a morphism of Lie groupoids π : $\mathbb{E} \to \mathbb{X}$ along with an action $\rho : \mathbb{E} \times \mathbb{G} \to \mathbb{E}$ of Lie 2-group \mathbb{G} on \mathbb{E} such that,


- $\pi_0: E_0 \to X_0$ is a principal G_0 -bundle over X_0 ,
- $\pi_1: E_1 \to X_1$ is a principal G_1 -bundle over X_1 .

We will call \mathbb{G} to be the structure 2-group of $\pi: \mathbb{E} \to \mathbb{X}$. We will denote the above principal \mathbb{G} -bundle over the Lie groupoid \mathbb{X} by $\mathbb{E}(\mathbb{X}, \mathbb{G})$.

If we replace the action functor $\rho \colon \mathbb{E} \times \mathbb{G} \to \mathbb{E}$ in Definition 3.4 by an η -twisted action $\rho \colon \mathbb{E} \rtimes_{\eta} \mathbb{G} \to \mathbb{E}$ for a smooth map $\eta \colon \operatorname{Mor}(\mathbb{E}) \times \operatorname{Mor}(\mathbb{G}) \to \operatorname{Mor}(\mathbb{E})$ satisfying conditions listed in Equation (3.1), then what we get will be called a *twisted principal* \mathbb{G} -bundle over the Lie groupoid X.

Remark 3.3. A slightly weaker version of a principal 2-bundle over a Lie groupoid has been introduced in [21, Definition 2.20]. Our main purpose in this paper is to study the Atiyah sequences and gauge transformations on the principal 2-bundles over Lie groupoids, so it is essential that we adopt a framework that provides us a pair of ordinary principal bundles (for objects and morphisms) related by the functoriality conditions.

Definition 3.5 (2-groupoid of principal 2-bundles). Let \mathbb{G} be a Lie 2-group. Let $\mathbb{E}(\mathbb{X}, \mathbb{G})$ and $\mathbb{E}'(\mathbb{X},\mathbb{G})$ be a pair of principal \mathbb{G} -bundles over a Lie groupoid \mathbb{X} . A morphism from $\mathbb{E}(\mathbb{X},\mathbb{G})$ to $\mathbb{E}'(\mathbb{X},\mathbb{G})$ is defined as a smooth \mathbb{G} -equivariant morphism $F:\mathbb{E}\to\mathbb{E}'$ such that the following diagram commutes up to nose



A 2-morphism from F to F' is defined as a smooth natural isomorphism $\eta: F \Longrightarrow F'$ such that $\eta(pg) = \eta(p)1_g$ and $\pi_1(\eta(p)) = 1_{\pi_0(p)}$ for all $p \in E_0$ and $g \in G_0$. We have a strict 2-groupoid, denoted Bun(X, G).

Definition 3.6 (Gauge 2-group). For an object \mathbb{E} in 2-groupoid $\operatorname{Bun}(\mathbb{X}, \mathbb{G})$, $\mathcal{G}(\mathbb{E})$ will denote the strict automorphism 2-group of \mathbb{E} , and will be called the *Gauge 2-group* of the principal \mathbb{G} -bundle $\mathbb{E} \to \mathbb{X}$.

Example 3.4. A traditional principal G-bundle $\pi: E \to X$ over a manifold X is same as a $[G \rightrightarrows G]$ -bundle $[E \rightrightarrows E]$ over the Lie groupoid $[X \rightrightarrows X]$.

Example 3.5. Let $\mathbb{X} = [X_1 \rightrightarrows X_0]$ be a Lie groupoid and $\mathbb{G} = [G_1 \rightrightarrows G_0]$ a Lie 2-group. Then we have the product principal 2-bundle $\mathbb{X} \times \mathbb{G} = [X_1 \times G_1 \rightrightarrows X_0 \times G_0]$ over \mathbb{X} .

Example 3.6. For any Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ and a (traditional) principal Gbundle $\pi: E_0 \to X_0$, we define a Lie groupoid $E_1 = \{(p, \gamma, q) | \gamma \in X_1, p \in \pi^{-1}(s(\gamma)), q \in \pi^{-1}(t(\gamma))\} \rightrightarrows E_0$ as follows. The source, target and composition maps are respectively given by $s(p, \gamma, q) = p, t(p, \gamma, q) = q$ and $(q, \gamma_2, r) \circ (p, \gamma_1, q) = (p, \gamma_2 \circ \gamma_1, r)$. The Lie 2-group $[G \times G \rightrightarrows G]$ naturally acts on $[E_1 \rightrightarrows E_0]$ by $(p, g) \mapsto pg$ and $((p, \gamma, q), (g_1, g_2)) = (pg_1, \gamma, qg_2)$. Then we have a $[G \times G \rightrightarrows G]$ -bundle $[E_1 \rightrightarrows E_0]$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$, with obvious projection functor.

We will provide several other examples of principal 2-bundles over Lie groupoids.

We recall the definition of a G-bundle over a Lie groupoid [27]. In the following subsection, we will describe a construction of a principal 2-bundle from a given G-bundle over a Lie groupoid. The structure is general enough to produce a large class of examples relevant in many contexts.

Definition 3.7 (principal *G*-bundle over Lie groupoid). For a Lie group *G*, a *principal G*-bundle over a Lie groupoid \mathbb{X} is a principal *G*-bundle $\pi: E_G \to X_0$ along with a smooth map $\mu: X_1 \times_{s,X_0,\pi} E_G \to E_G$ satisfying

- $\mu(1_{\pi(p)}, p) = p$ for all $p \in E_G$ where $1: X_0 \to X_1$ is the identity assigning map,
- for each $(\gamma, p) \in X_1 \times_{s, X_0, \pi} E_G$, we have $(\gamma, \mu(\gamma, p)) \in X_1 \times_{t, X_0, \pi} E_G$,
- if $\gamma_2, \gamma_1 \in X_1$ such that $t(\gamma_1) = s(\gamma_2)$ and $(\gamma_1, p) \in X_1 \times_{s, X_0, \pi} E_G$ then $\mu(\gamma_2 \circ \gamma_1, p) = \mu(\gamma_2, \mu(\gamma_1, p)),$
- for all $p \in E_G$, $g \in G$ and $\gamma \in X_1$ we have $\mu(\gamma, p)g = \mu(\gamma, pg)$.

We will denote the above principal G-bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ by the notation $(\pi \colon E_G \to X_0, \mu, [X_1 \rightrightarrows X_0])$. The map μ is called the *action map* of the principal G-bundle.

The collection of principal G-bundles over X forms a groupoid whose morphisms are defined naturally. We will denote this groupoid by Bun(X, G).

Example 3.7. Let $(\pi: E_G \to X_0, \mu, [X_1 \rightrightarrows X_0])$ be a principal *G*-bundle. The pull-back Lie groupoid $[X_1 \times_{s,X_0,\pi} E_G \rightrightarrows E_G]$, denoted $[s^*E_G \rightrightarrows E_G]$, gives the $[G \rightrightarrows G]$ -bundle over $[X_1 \rightrightarrows X_0]$. Conversely given a principal $[G \rightrightarrows G]$ -bundle $\pi: \mathbb{E} \to \mathbb{X}$ over the Lie groupoid \mathbb{X} , we take the underlying principal *G*-bundle to be $\pi_0: E_0 \to X_0$ and the action map $\eta: s^*E_0 \to E_0$ to be $(\gamma, p) \mapsto t(\tilde{\gamma})$, where $\tilde{\gamma} \in \pi_1^{-1}(\gamma) \subset s^*E_0$ is the unique element associated to $(\gamma, p) \in s^*E_0$ satisfying $s(\tilde{\gamma}) = p$. **Example 3.8.** As in Example 2.3 now we consider the single object Lie 2-group $[G \rightrightarrows e]$ of a Lie group G. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $[G \rightrightarrows e]$ -bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Then it is immediate from the definition of a principal 2-bundle that $E_0 = X_0$ and E_1 is a principal G-bundle over X_1 such that

$$\begin{array}{ccc} E_1 & \stackrel{\pi_1}{\longrightarrow} & X_1 \\ s & \downarrow t & s & \downarrow t \\ X_0 & \stackrel{\mathrm{Id}}{\longrightarrow} & X_0 \end{array}$$

commutes. Moreover the functoriality of the action of $[G \Rightarrow e]$ on $E_1 \Rightarrow X_0$ implies that the action $E_1 \times G \to E_1$ preserves the hom sets; that is, the restriction gives

$$\operatorname{Hom}_{\mathbb{E}}(x, y) \times G \to \operatorname{Hom}_{\mathbb{E}}(x, y) \tag{3.3}$$

for all $x, y \in X_0$ and for any pair of composable morphisms $\gamma_2, \gamma_1 \in E_1$, and $g, g' \in G$ we have

$$(\gamma_2 g) \circ (\gamma_1 g') = (\gamma_2 \circ \gamma_1) gg'. \tag{3.4}$$

On the other hand, suppose $[E \rightrightarrows X_0]$ is a Lie groupoid, and a compact Lie group G acts on E freely and smoothly. Assume that the source, target maps are G invariant and the action satisfies the conditions in Equations (3.3) and (3.4). Then we have a principal G-bundle $E \rightarrow E/G := X_1$, which defines a $[G \rightrightarrows e]$ -bundle $\mathbb{E} = [E \rightrightarrows X_0]$ over the Lie groupoid $\mathbb{X} = [E/G \rightrightarrows X_0]$.

The principal 2-bundle of Example 3.8 is related to *G*-extensions as we see below.

3.1. Lie groupoid G-extensions and twisted principal bundles. We recall the definition of a Lie groupoid G-extension [21].

Let G be a Lie group and M a smooth manifold. Consider the Lie groupoid $[M \times G \rightrightarrows M]$ with source, target maps both be projection at the first component $(m,g) \mapsto m$, composition $(m,g_2) \circ (m,g_1) = (m,g_2g_1)$, $1_m = (m,e)$ and $(m,g)^{-1} = (m,g^{-1})$. A Lie groupoid G-extension is a short exact sequence of Lie groupoids of the form

$$1 \to M \times G \xrightarrow{i} \Gamma_{2} \xrightarrow{\phi} \Gamma_{1} \to 1$$

$$\downarrow \downarrow \qquad \downarrow \downarrow \qquad \downarrow \downarrow \qquad ,$$

$$1 \longrightarrow M \xrightarrow{\mathrm{Id}} M \xrightarrow{\mathrm{Id}} M \xrightarrow{\mathrm{Id}} M \to 1$$

$$(3.5)$$

where i is an embedding and ϕ is a surjective submersion ([34, Chapter 4]).

Given a $[G \rightrightarrows e]$ -bundle $\mathbb{E} = [E_1 \rightrightarrows X_0]$ over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$, define the map $i: X_0 \times G \to E_1$ by

$$(x,g)\mapsto 1_x^{\mathbb{E}}g,$$

where $1^{\mathbb{E}}$ is the identity assigning map in the Lie groupoid $\mathbb{E} = [E_1 \rightrightarrows X_0]$. Since $1^{\mathbb{E}} \colon X_0 \to E_1$ is a diffeomorphism onto its image, by Equation (3.3), the map *i* is an embedding. As $E_1 \to X_1$ is a principal *G*-bundle, it follows that,

$$1 \longrightarrow X_0 \times G \xrightarrow{i} E_1 \xrightarrow{\pi_1} X_1 \longrightarrow 1$$
$$\downarrow \downarrow \qquad \qquad \downarrow \downarrow \qquad \qquad \downarrow \downarrow \qquad \qquad \downarrow \downarrow \qquad ,$$
$$1 \longrightarrow X_0 \xrightarrow{\mathrm{Id}} X_0 \xrightarrow{\mathrm{Id}} X_0 \xrightarrow{\mathrm{Id}} X_0 \longrightarrow 1$$

is a Lie groupoid G-extension.

Conversely, suppose we have a Lie groupoid G-extension as in Diagram 3.5. Observe that commutation of the second square from the left implies that

$$i(x,g) \in \operatorname{Hom}_{\Gamma_2}(x,x)$$

Lemma 3.9. (1) There exists a smooth free action $\Gamma_2 \times G \to \Gamma_2$ given by

$$(x \xrightarrow{\gamma} y, g) \mapsto \gamma \circ i(x, g). \tag{3.6}$$

- (2) The action is transitive on the fibers $\phi^{-1}(\lambda)$, for each $\lambda \in \Gamma_1$
- (3) ϕ is constant on the orbits of the action.
- (4) The action satisfies the condition in Equation (3.3);

$$\operatorname{Hom}_{\Gamma_2}(x, y) \times G \to \operatorname{Hom}_{\Gamma_2}(x, y) \tag{3.7}$$

for all $x, y \in M$.

Proof. (1) Functoriality of i implies that

$$\gamma(gg') = \gamma \circ i(x, gg') = \gamma \circ \left(i(x, g) \circ i(x, g')\right) = \left(\gamma \circ i((x, g))\right) \circ i(x, g') = (\gamma g)g'.$$

Injectivity of i ensures the action is free.

- (2) If $\phi(\gamma_2) = \phi(\gamma_1)$. Then $\phi(\gamma_2^{-1} \circ \gamma_1) = \mathbf{1}_x^{\Gamma_1}$. Thus by exactness of Diagram 3.5 we have $(x,g) \in M \times G$ such that $\gamma_2^{-1} \circ \gamma_1 = i(x,g)$. Hence $\gamma_1 = \gamma_2 g$.
- (3) $\phi(\gamma g) = \phi(\gamma \circ i(x,g)) = \phi(\gamma) \circ \phi(i(x,g))$. Since 3.5 is a short exact sequence $\phi(i(x,g)) = 1_x^{\Gamma_1}$. Therefore $\phi(\gamma g) = \phi(\gamma)$.
- (4) Readily follows from the observation that $i(x, g) \in \operatorname{Hom}_{\Gamma_2}(x, x)$.

In conclusion, since ϕ is a surjective submersion, the Lie groupoid *G*-extension in Diagram 3.5 produces a principal *G*-bundle $\phi: \Gamma_2 \to \Gamma_1$ such that the diagram

$$\begin{array}{ccc} \Gamma_2 & \stackrel{\phi}{\longrightarrow} & \Gamma_1 \\ s & \downarrow t & s & \downarrow t \\ M & \stackrel{\mathrm{Id}}{\longrightarrow} & M \end{array}$$

commutes, and the action satisfies Equation (3.7). The action defined in Lemma 3.9 in general does not satisfy the other functoriality condition in Equation (3.4)! Thus the principal G-bundle $\phi: \Gamma_2 \to \Gamma_1$ does not define a $[G \rightrightarrows e]$ -bundle $[\Gamma_2 \rightrightarrows M]$ over the Lie groupoid $[\Gamma_1 \rightrightarrows M]$. However we show that it is actually a twisted principal $[G \rightrightarrows e]$ bundle over the Lie groupoid $[\Gamma_1 \rightrightarrows M]$. For that we define

$$\eta \colon \operatorname{Mor}(\Gamma_2) \times G \to \operatorname{Mor}(\Gamma_2) (x \xrightarrow{\gamma} y, g) \mapsto i(y, g) \circ \gamma \circ i(x, g^{-1}).$$

$$(3.8)$$

It is readily verified that η satisfies all the required conditions of Equation (3.1). Thus we have the twisted product category $([\Gamma_2 \rightrightarrows M] \rtimes_{\eta} [G \rightrightarrows e])$. Then we verify,

$$(y \xrightarrow{\gamma_2} z g') \circ (x \xrightarrow{\gamma_1} y g) = (\gamma_2 \circ i(y, g')) \circ (\gamma_1 \circ i(x, g)) = (\gamma_2 \circ (i(y, g') \circ \gamma_1 \circ i(x, g'^{-1})) \circ i(x, g'g) = (\gamma_2 \circ \eta(\gamma_1, g')) \circ i(x, g'g) = (\gamma_2 \circ \eta(\gamma_1, g'))g'g,$$

$$(3.9)$$

which is precisely the condition, stated in Remark 3.2, required for the action in Equation (3.6)to be an η -twisted action

$$([\Gamma_2 \rightrightarrows M] \rtimes_\eta [G \rightrightarrows e]) \to [\Gamma_2 \rightrightarrows M].$$

Observe that in the other direction the construction proceeds for a twisted $[G \Rightarrow e]$ -bundle exactly same way as was for a *non-twisted* $[G \Rightarrow e]$ -bundle (using the third property in Equation (3.1)).

Proposition 3.10. Let G be a Lie group. A twisted principal $[G \rightrightarrows e]$ -bundle defines a Lie groupoid G-extension and vice-versa.

It is pertinent to note here that [21] also discussed the relation between Morita equivalent classes of Lie groupoid G-extensions and Morita equivalence classes of what the authors called $[G \rightrightarrows \operatorname{Aut}(G)]$ -bundles (see [21, Theorem 3.4]).

3.2. Decorated principal 2-bundles. In this subsection we are going to construct the main example for this paper. Given a Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$ and a principal G_0 bundle over a Lie groupoid X, we will construct a principal G-bundle over X. We will call them decorated principal 2-bundles. Let (G, H, τ, α) be the associated crossed module of G. Here, of course, $G_0 = G$.

An analogous notion for a principal 2-bundle over a path space groupoid already exists in literature, introduced in [17] by decorating the space of A-horizontal paths $\mathcal{P}_{\bar{A}}P$, for a connection \overline{A} on a principal G_0 -bundle $P \to M$.

Proposition 3.11. Let $(G, H, \tau \colon H \to G, \alpha \colon G \times H \to H)$ be a Lie crossed module and $\mathbb{G} = [H \rtimes G \rightrightarrows G]$ the associated Lie 2-group. Let $(\pi \colon E_G \to X_0, \mu, [X_1 \rightrightarrows X_0])$ a principal G-bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Let us denote $s^* E_G \times H =$ $(X_1 \times_{s,X_0,\pi} E_G) \times H$ by $(s^* E_G)^{\text{dec}}$.

- (1) The manifolds $(s^*E_G)^{\text{dec}}$ and E_G determines a Lie groupoid $[(s^*E_G)^{\text{dec}} \Rightarrow E_G]$ whose structure maps are given as
 - source map \tilde{s} : $(\gamma, p, h) \mapsto p$,

(

- target map \tilde{t} : $(\gamma, p, h) \mapsto \mu(\gamma, p)\tau(h^{-1})$,
- composition map \tilde{m} : $((\gamma_2, p_2, h_2), (\gamma_1, p_1, h_1)) \mapsto (\gamma_2 \circ \gamma_1, p_1, h_2 h_1),$
- unit map $\tilde{u}: p \mapsto (1_{\pi(p)}, p, e_H),$
- inverse map $\tilde{i} \colon (\gamma, p, h) \mapsto (\gamma^{-1}, \mu(\gamma, p)\tau(h^{-1}), h^{-1}).$

mdec

(2) The Lie groupoid $\mathbb{E}^{\text{dec}} := [(s^* E_G)^{\text{dec}} \Rightarrow E_G]$ forms a principal \mathbb{G} -bundle over $[X_1 \rightrightarrows X_0]$, whose action functor and the bundle functor respectively given by

mdec

$$\rho \colon \mathbb{E}^{\operatorname{dec}} \times \mathbb{G} \to \mathbb{E}^{\operatorname{dec}}$$

$$(p,g) \mapsto pg \qquad (3.10)$$

$$(\gamma, p, h), (h', g)) \mapsto (\gamma, pg, \alpha_{g^{-1}}(h'^{-1}h)),$$

and

$$\pi \colon \mathbb{E}^{\mathrm{dec}} \to \mathbb{X}$$

$$p \mapsto \pi(p) \tag{3.11}$$

$$(\gamma, p, h) \mapsto \gamma.$$

We call $\mathbb{E}^{dec} \to [X_1 \rightrightarrows X_0]$ the decorated \mathbb{G} -bundle associated to $(E_G \to X_0, [X_1 \rightrightarrows X_0])$.

Proof. (1) It is straightforward to verify that $[(s^*E_G)^{\text{dec}} \Rightarrow E_G]$ is a groupoid.

Since $E_G \to X_0$ is a surjective submersion, $(s^*E_G)^{\text{dec}}$ is a smooth manifold. Moreover, as \tilde{s} is the composition of surjective submersions, it (hence the target map as well) is a surjective submersion. The smoothness of other structure maps directly follows from the smoothness of μ and smoothness of structure maps of the Lie groupoid $[X_1 \Rightarrow X_0]$.

(2) Since the action of G on E_G is free and $\alpha_g \in \operatorname{Aut}(H)$ for each $g \in G$, the action of $H \rtimes_{\alpha} G$ on $(s^*E_G)^{\operatorname{dec}}$ in Equation (3.10) is free. Observe that for any $\gamma \in X_1$, $\pi^{-1}(\gamma) = \pi^{-1}(s(\gamma)) \times H$. Now it is straightforward to verify that the action is transitive on $\pi^{-1}(\gamma)$, and $\pi((\gamma, p, h)(h', g)) = \pi(\gamma, p, h)$.

The local trivialization on the principal G-bundle $\pi: E_G \to X_0$ induces a local trivialization on the map $\pi: (s^*E_G)^{\text{dec}} \to \mathbb{X}_1$ as follows. For any $\gamma \in X_1$, there exists a local trivialization $(U \subset X_0, \phi)$ such that $s(\gamma) \in U$ and $\phi: \pi^{-1}(U) \to U \times G$ a diffeomorphism. Then we have a local trivialization $(s^{-1}(U), \tilde{\phi})$ around γ , given by

$$\widetilde{\phi} \colon \pi^{-1}(s^{-1}(U)) \to s^{-1}(U) \times (H \rtimes_{\alpha} G)$$
$$(\gamma, p, h) \mapsto (\gamma, \alpha_g(h), g),$$

where $g = \text{pr}_2 \circ \phi(p) \in G$. This completes the proof.

Example 3.12. The product Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle $\mathbb{X} \times \mathbb{G}$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$ in Example 3.5 can be viewed as a decorated bundle. Note that $s^*(X_0 \times G_0) = X_1 \times G_0$. The action of X_1 on $X_0 \times G_0$ is given by $s^*(X_0 \times G_0) \rightarrow X_0 \times G_0$ by $(\gamma, g) \mapsto (t(\gamma), g)$. Thus the Lie groupoid $[X_1 \times G_0 \rightrightarrows X_0 \times G_0]$ is the pullback Lie groupoid, and by decoration we recover the $\mathbb{X} \times \mathbb{G}$.

Example 3.13. Let (G, H, τ, α) be a Lie crossed module associated to a Lie 2-group \mathbb{G} . Consider a traditional principal *G*-bundle $E \to X$. The Lie groupoid $[X \rightrightarrows X]$ trivially acts on *E* by $(x, p) \mapsto p$. Then the Lie groupoid $E^{\text{dec}} := [E \times H \rightrightarrows E]$ defines the associated decorated \mathbb{G} -bundle.

3.3. Categorical connections. We will, later on, discuss the differential geometric notion of a connection on a principal 2-bundle over a Lie groupoid. Apart from that, a principal 2-bundle can also be equipped with a lifting property for arrows, which we call a 'categorical connection'.

Definition 3.8. Let \mathbb{G} be a Lie 2-group and $\pi : \mathbb{E} \to \mathbb{X}$ a \mathbb{G} -bundle over \mathbb{X} . A categorical connection \mathcal{C} on $\mathbb{E}(\mathbb{X}, \mathbb{G})$ is an embedding $\mathcal{C} : s^* E_0 \to E_1$, and satisfy following conditions:

- $s(\mathcal{C}(\gamma, p)) = p$ for all $(\gamma, p) \in s^* E_0$,
- $\pi_1(\mathcal{C}(\gamma, p)) = \gamma$ for all $(\gamma, p) \in s^* E_0$,
- if $(\gamma_2, p_2), (\gamma_1, p_1) \in s^* E_0$ such that $s(\gamma_2) = t(\gamma_1)$ and $p_2 = t(\mathcal{C}(\gamma_1, p_1))$, then $\mathcal{C}(\gamma_2 \circ \gamma_1, p_1) = \mathcal{C}(\gamma_2, p_2) \circ \mathcal{C}(\gamma_1, p_1)$,
- $\mathcal{C}(1_x, p) = 1_p$ for any $x \in X_0$ and $p \in \pi^{-1}(x)$,
- $\mathcal{C}(\gamma, p.g) = \mathcal{C}(\gamma, p) \cdot 1_g$ for all $(\gamma, p) \in s^* E_0$ and $g \in G_0$.

We call $\mathcal{C}(\gamma, p)$ the (unique) \mathcal{C} -horizontal lift of $\gamma \in X_1$ through the point $p \in \pi_0^{-1}(s(\gamma))$.

16 SAIKAT CHATTERJEE, ADITTYA CHAUDHURI, AND PRAPHULLA KOUSHIK

The definition is motivated by the notion of horizontal lifting of a path by a connection in the traditional set-up of principal bundles. See [17] for an analogous notion of a categorical connection on the set-up of path space groupoid. We will observe that categorical connections characterize the decorated bundles.

Example 3.14. There exists a unique categorical connection on a $[G \rightrightarrows G]$ -bundle over a Lie groupoid.

Example 3.15. Let $[(s^*E_G)^{\text{dec}} \rightrightarrows E_G] \rightarrow [X_1 \rightrightarrows X_0]$ be the decorated G-bundle obtained in Proposition 3.11. Any smooth map $\beta \colon E_G \rightarrow H$ satisfying $\beta(pg) = \alpha_{g^{-1}}(\beta(p))$ for $p \in E_G, g \in G$, defines a categorical connection $\mathcal{C} \colon (\gamma, p) \mapsto (\gamma, p, \beta(p)\beta(\mu(\gamma, p))^{-1})$. For the constant map $\beta \colon p \mapsto e$, we have the categorical connection $(\gamma, p) \mapsto (\gamma, p, e)$. We call the categorical connection $(\gamma, p) \mapsto (\gamma, p, e)$, the *canonical categorical connection* on the decorated bundle.

Given a categorical connection \mathcal{C} on a \mathbb{G} -bundle $\pi \colon \mathbb{E} \to \mathbb{X}$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$, we can define an action of X_1 on E_0

$$\mu \colon s^* E_0 \to E_0, (\gamma, p) \mapsto t(\mathcal{C}(\gamma, p)),$$
(3.12)

which turns s^*E_0 into a Lie groupoid $[s^*E_0 \rightrightarrows E_0]$, and \mathcal{C} allows us to identify $[s^*E_0 \rightrightarrows E_0]$ as a Lie sub groupoid of $[E_1 \rightrightarrows E_0]$,

Note that μ satisfies $\mu(\gamma, p)g = t(\mathcal{C}(\gamma, p))g = t(\mathcal{C}(\gamma, p)1_g) = t(\mathcal{C}(\gamma, pg)) = \mu(\gamma, pg)$. Identifying G as a subgroup of $G_1 = H \rtimes G$, we observe that the fifth condition in Definition 3.8 implies $[s^*E_0 \Rightarrow E_0] \simeq [\mathcal{C}(s^*E_0) \Rightarrow E_0]$ is in fact a sub Lie groupoid bundle over $\mathbb{X} = [X_1 \Rightarrow X_0]$ with a reduced structure Lie 2-group $[G \Rightarrow G]$. Given any $\widetilde{\gamma} \in E_1$ it follows that we have a unique $h \in H$ such that $[\mathcal{C}(\pi(\widetilde{\gamma}), s(\widetilde{\gamma}))](h, e) = \widetilde{\gamma}$, since $\pi(\widetilde{\gamma}) = \pi(\mathcal{C}(\pi(\widetilde{\gamma}), s(\widetilde{\gamma})))$ and $s(\widetilde{\gamma}) = s(\mathcal{C}(\pi(\widetilde{\gamma}), s(\widetilde{\gamma}))$. In turn, we get an isomorphism of \mathbb{G} -bundles

$$\theta \colon s^* E_0 \times H \to E_1 ((\gamma, p), h) \to \mathcal{C}(\gamma, p)(h^{-1}, e).$$

$$(3.14)$$

Lemma 3.16. The map (θ, Id) is an isomorphism from $\mathbb{E}^{\mathrm{dec}}(\mathbb{X}, \mathbb{G})$ to $\mathbb{E}(\mathbb{X}, \mathbb{G})$.

Proof. The functoriality of θ follows from the composition law for the decorated Lie groupoid in Proposition 3.11 and the functoriality of the action of \mathbb{G} ,

$$\begin{aligned} \theta((\gamma_{2}, p_{2}), h_{2})) &\circ \theta((\gamma_{1}, p_{1}), h_{1}) \\ &= (\mathcal{C}(\gamma_{2}, p_{2})(h_{2}^{-1}, e)) \circ (\mathcal{C}(\gamma_{1}, p_{1})(h_{1}^{-1}, e)) \\ &= (\mathcal{C}(\gamma_{2}, p_{2})(h_{2}^{-1}, e)) \circ (\mathcal{C}(\gamma_{1}, p_{1}\tau(h_{1})^{-1})(e, \tau(h_{1}))(h_{1}^{-1}, e)) \\ &= (\mathcal{C}(\gamma_{2}, p_{2})(h_{2}^{-1}, e)) \circ (\mathcal{C}(\gamma_{1}, p_{1}\tau(h_{1})^{-1})(e, \tau(h_{1}))(h_{1}^{-1}, e)) \\ &= (\mathcal{C}(\gamma_{2}, p_{2})(h_{2}^{-1}, e)) \circ (\mathcal{C}(\gamma_{1}, p_{1}\tau(h_{1})^{-1})(e, \tau(h_{1}))(h_{1}^{-1}, e)) \\ &= (\mathcal{C}(\gamma_{2}, p_{2}) \circ \mathcal{C}(\gamma_{1}, p_{1}\tau(h_{1})^{-1}))((h_{2}^{-1}, e) \circ ((e, \tau(h_{1}))(h_{1}^{-1}, e))) \end{aligned}$$
(3.15)
$$&= \mathcal{C}(\gamma_{2} \circ \gamma_{1}, p_{1}\tau(h_{1})^{-1})(h_{2}^{-1}h_{1}^{-1}, \tau(h_{1})) \\ &= \mathcal{C}(\gamma_{2} \circ \gamma_{1}, p_{1}\tau(h_{1})^{-1})(h_{2}^{-1}h_{1}^{-1}, \tau(h_{1})) \\ &= \mathcal{C}(\gamma_{2} \circ \gamma_{1}, p_{1})(e, \tau(h_{1}^{-1}))(h_{2}^{-1}h_{1}^{-1}, \tau(h_{1})) \\ &= \mathcal{C}(\gamma_{2} \circ \gamma_{1}, p_{1})(h_{1}^{-1}h_{2}^{-1}, e) \\ &= \theta((\gamma_{2} \circ \gamma_{1}, p_{1}), h_{2}h_{1}). \end{aligned}$$

To see the equivarinacy, note that by the action defined in Proposition 3.11, $(\gamma, p, h)(h', g') = (\gamma, pg', \alpha_{g'^{-1}}(h'^{-1}h))$. Thus

$$\begin{aligned} \theta((\gamma, p, h)(h', g')) \\ &= \mathcal{C}(\gamma, pg') (\alpha_{g'^{-1}}(h^{-1}h'), e) \\ &= \mathcal{C}(\gamma, p)(e, g') (\alpha_{g'^{-1}}(h^{-1}h'), e) \\ &= \mathcal{C}(\gamma, p)(h^{-1}, e)(h, e)(e, g') (\alpha_{g'^{-1}}(h^{-1}h'), e) \\ &= \theta((\gamma, p), h)(h', g'). \end{aligned}$$
(3.16)

Obviously (θ, Id) is a morphism of principal 2-bundles.

In conclusion, we have the following characterization of principal 2-bundles over Lie groupoids in terms of the categorical connections.

Proposition 3.17. A principal 2-bundle over a Lie groupoid is a decorated bundle if and only if it admits a categorical connection.

Corollary 3.18. Let (G, H, τ, α) be a Lie crossed module associated to a Lie 2-group \mathbb{G} . Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a \mathbb{G} -bundle over $[X \rightrightarrows X]$. Then $\mathbb{E} = [E_1 \rightrightarrows E_0]$ is the decorated bundle defined in Example 3.13.

Proof. Any such principal 2-bundle admits a unique categorical connection $(1_x, p) \mapsto 1_p$ for $p \in E, x = \pi(p)$.

Example 3.19. The $[G \times G \rightrightarrows G]$ -bundle $[E_1 \rightrightarrows E_0]$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$ in Example 3.6 admits a categorical connection if and only if $E_0 \rightarrow X_0$ is a principal *G*-bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Suppose $E_0 \rightarrow X_0$ is a principal *G*-bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ with respect to the action $\mu \colon X_1 \times_s E_0 \rightarrow E_0$. Then the correspondence with the decorated bundle is given by $(p, \gamma, q) \mapsto ((p, \gamma), g)$, where $\mu(\gamma, p) = q g$.

4. ATIYAH SEQUENCE FOR PRINCIPAL 2-BUNDLES OVER LIE GROUPOIDS

Let G be a Lie group, and L(G) the Lie algebra of G. For a principal G-bundle $\pi \colon P \to M$ one can associate a short exact sequence

$$0 \to \operatorname{Ad}(P) \xrightarrow{j^{/G}} \operatorname{At}(P) \xrightarrow{\pi_*^{/G}} TM \to 0, \tag{4.1}$$

of vector bundles over the manifold M, known as Atyiah sequence [2]. Here Ad(P) := $(P \times L(G))/G \to M$ and $At(P) := TP/G \to M$ are respectively the adjoint bundle and the Atiyah bundle associated to the principal G-bundle $\pi: P \to M$, and the associated maps are respectively given by

$$j^{/G}: [(p,B)] \mapsto [(p,\delta_p(B))],$$

$$\pi_*^{/G}: [(p,v)] \mapsto (\pi(p), \pi_{*,p}(v)),$$

$$(4.2)$$

where $\delta(B)$ denotes the vertical vector field generated by the element $B \in L(G)$. The Atiyah bundle $\operatorname{At}(P) \to M$, with $\pi_*^{/G} : \operatorname{At}(P) \to TM$ being the anchor map, forms a Lie algebroid, known as the Atiyah Lie algebroid of the principal G-bundle $\pi: P \to M$.

A connection on a principal G-bundle P(M,G) is a splitting of the Atiyah sequence in Equation (4.1). Equivalently, a connection is same as a retract $R: \operatorname{At}(P) \to \operatorname{Ad}(P)$ of $j^{/G}$, or a section $\sigma: TM \to \operatorname{At}(P)$ of $\pi_*^{/G}$. The first equivalent condition for the splitting in turn gives an L(G)-valued differential 1-form $\omega: TP \to P \times L(G)$ satisfying,

- $\omega(pg, vg) = \operatorname{ad}_{q^{-1}}(\omega(p, v))$, for $g \in G, p \in P, v \in T_pP$ and
- $\omega(p, \delta_p(B)) = B$, for $p \in P$, $B \in L(G)$, where as before $\delta(B)$ denotes the vertical vector field generated by the element $B \in L(G)$,

while the second equivalent condition gives the horizontal sub-bundle $\mathcal{H} \subset TE$ satisfying,

- *H*_{pg} = (R_g)_{*,p}*H*_p for all p ∈ P, g ∈ G and
 giving a smooth splitting *H*_p ⊕ ker π_{*,p} = T_pP for each p ∈ P.

Here, $R_q: P \to P$ is the right translation map by the element $q \in G$.

We refer to [28, Appendix A] for a more detailed discussion on this association.

For a given $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle $\mathbb{E} = [E_1 \rightrightarrows E_0]$ over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows E_0]$ X_0 , we have a pair of Atiyah sequences,

for $i \in \{0, 1\}$. These pairs of short exact sequences are intertwined by the functoriality of various maps. We will see the right framework to describe the Ativah sequences of a principal 2-bundle is that of so-called 'a Vector bundle groupoid' or 'a VB-groupoid' in short. For more details about VB-groupoids, we refer the reader to [14]. Let us recall the definition given in [22].

Definition 4.1 (VB-groupoid [22, Definition 3.1.]). Let $[X_1 \rightrightarrows X_0]$ be a Lie groupoid. A *VB-groupoid over* $[X_1 \rightrightarrows X_0]$ is given by a morphism of Lie groupoids

$$V_1 \xrightarrow{\pi_1} X_1.$$

$$s_V \downarrow \downarrow t_V \qquad s_X \downarrow \downarrow t_X ,$$

$$V_0 \xrightarrow{\pi_0} X_0$$

satisfying the following conditions:

- (1) the maps $\pi_1: V_1 \to X_1$ and $\pi_0: V_0 \to X_0$ are vector bundles,
- (2) the maps $(s_V, s_X), (t_V, t_X)$ are morphisms of vector bundles,
- (3) for appropriate $\gamma_1, \gamma_2, \gamma_3, \gamma_4 \in E_1$, we have $(\gamma_3 \circ \gamma_1) + (\gamma_4 \circ \gamma_2) = (\gamma_3 + \gamma_4) \circ (\gamma_1 + \gamma_2)$.

Equivalently, a VB-groupoid is a Lie groupoid object in the category of smooth vector bundles.

A 1-morphism of VB-groupoids $[V_1 \rightrightarrows V_0] \rightarrow [V'_1 \rightrightarrows V'_0]$ over the base Lie groupoid $[X_1 \rightrightarrows X_0]$ is a morphism of Lie groupoids $\Phi = (\Phi_1.\Phi_0): [V_1 \rightrightarrows V_0] \rightarrow [V'_1 \rightrightarrows V'_0]$ such that $(\Phi_0, \operatorname{Id}_{X_0})$, and $(\Phi_1, \operatorname{Id}_{X_1})$ are morphisms of vector bundles. For a pair of such 1-morphisms $(\Phi, \Phi', a 2$ -morphism $\eta: \Phi \Longrightarrow \Phi'$ is a smooth natural transformation such that $(\eta, 1)$ is a morphism of vector bundles from $V_0 \rightarrow X_0$ to $V_1 \rightarrow X_1$.

Proposition 4.1. The collection of VB-groupoids, 1-morphisms and 2-morphisms forms a strict 2-category, and will be denoted as $2 - \text{VBGpd}(\mathbb{X})$.

4.1. Atiyah sequence of VB-groupoids. Let $\mathbb{E} = [E_1 \Rightarrow E_0]$ be a $\mathbb{G} = [G_1 \Rightarrow G_0]$ bundle over $\mathbb{X} = [X_1 \Rightarrow X_0]$. Then we show that the pair of short exact sequences in Diagram 4.3 give a short exact sequence of VB-groupoids. To be precise, each pair $\{TX_i\}_{i\in\{0,1\}}, \{\operatorname{Ad}(E_i)\}_{i\in\{0,1\}}$ and $\{\operatorname{At}(E_i)\}_{i\in\{0,1\}}$ forms a VB-groupoid over $\mathbb{X} = [X_1 \Rightarrow X_0]$, and the sequences define a short exact sequence of morphisms of VB-groupoids [29, Section 2.4].

VB-groupoid TX.

It is obvious that the tangent bundle maps $TX_1 \to X_1$ and $TX_0 \to X_0$ give a VBgroupoid $[TX_1 \rightrightarrows TX_0] \to [X_1 \rightrightarrows X_0]$.

VB-groupoid $Ad(\mathbb{E})$.

To see the groupoid structure of $[\operatorname{Ad}(E_1) \rightrightarrows \operatorname{Ad}(E_0)]$, we first define

• the source map $s^{/\sim}$: Ad $(E_1) \to$ Ad (E_0) as

$$[(\widetilde{\gamma}, K)] \mapsto [(s(\widetilde{\gamma}), s_{*,e}(K))],$$

• the target map $t^{\prime} : \operatorname{Ad}(E_1) \to \operatorname{Ad}(E_0)$ as

$$[(\widetilde{\gamma}, K)] \mapsto [(t(\widetilde{\gamma}), t_{*,e}(K))],$$

where $\tilde{\gamma} \in E_1$ and $K \in L(G_1)$. Next we make the following observation. Suppose

$$t^{\prime}([(\widetilde{\gamma}_{2}', K_{2}')]) = s^{\prime}([(\widetilde{\gamma}_{1}, K_{1})]).$$
(4.4)

Then there exists a $\theta \in G_0$ such that $s(\widetilde{\gamma}'_2)\theta = t(\widetilde{\gamma}_1)$ and $\mathrm{ad}_{\theta}(s_{*,e}(K'_2)) = t_{*,e}(K_1)$. That means $(\widetilde{\gamma}'_2, K'_2)\mathbf{1}_{\theta} = (\widetilde{\gamma}'_2 \mathbf{1}_{\theta}, \mathrm{ad}_{\mathbf{1}_{\theta}}(K'_2))$ is composable with $(\widetilde{\gamma}_1, K_1)$. Note that $(\widetilde{\gamma}'_2, K'_2)\mathbf{1}_{\theta} \in [(\widetilde{\gamma}_2, K_2)]$. In other words, whenever the condition in Equation (4.4) met, there exists a

composable pair respectively belonging to $[(\tilde{\gamma}_2, K_2)]$ and $[(\tilde{\gamma}_1, K_1)]$. We choose such a pair and define the composition as,

$$\left(\left[\left(\widetilde{\gamma}_{2}, K_{2}\right)\right]\right) \circ \left(\left[\left(\widetilde{\gamma}_{1}, K_{1}\right)\right]\right) = \left[\left(\widetilde{\gamma}_{2} \circ \widetilde{\gamma}_{1}, K_{2} \circ K_{1}\right)\right].$$
(4.5)

If $(\widetilde{\gamma}'_2, K'_2)$ and $(\widetilde{\gamma}'_1, K'_1)$ are another pair of such elements, then there exist composable $k_2, k_1 \in G_1$ such that $(\widetilde{\gamma}'_2, K'_2) = (\widetilde{\gamma}_2 k_2, \operatorname{ad}_{k_2}(K_2))$ and $(\widetilde{\gamma}'_1, K'_1) = (\widetilde{\gamma}_1 k_1, \operatorname{ad}_{k_1}(K_1))$. Then

$$\begin{aligned} &(\widetilde{\gamma}_{2}^{\prime} \circ \widetilde{\gamma}_{1}^{\prime}, K_{2}^{\prime} \circ K_{1}^{\prime}) \\ &= \left((\widetilde{\gamma}_{2}k_{2}) \circ (\widetilde{\gamma}_{1}k_{1}), (\operatorname{ad}_{k_{2}}(K_{2})) \circ (\operatorname{Ad}_{k_{1}}(K_{1})) \right) \\ &= \left((\widetilde{\gamma}_{2} \circ \widetilde{\gamma}_{1}) (k_{2}k_{1}), (\operatorname{ad}_{k_{2}k_{1}}(K_{2} \circ K_{1})) \right) \\ &= (\widetilde{\gamma}_{2} \circ \widetilde{\gamma}_{1}, K_{2} \circ K_{1}) (k_{2}k_{1}), \end{aligned}$$

$$(4.6)$$

where in the third step we have used the functoriality of the Lie 2-group action. Thus Equation (4.5) is well defined. The inverse and identity maps are the obvious ones.

Considering the following commutative diagram

we see that as $s_{\mathbb{E}}$ and $s_{\mathbb{G}_{*,e}}$ are surjective submersions, we have $s^{/\sim} \circ \delta_0$ a surjective submersion. Since δ_0 is a surjective submersion, it follows that $s^{/\sim}$ is also a surjective submersion. Similarly, $t^{/\sim}$ is also a surjective submersion. Thus $\operatorname{Ad}(\mathbb{E}) = [\operatorname{Ad}(E_1) \rightrightarrows$ $\operatorname{Ad}(E_0)]$ is a Lie groupoid.

The (quotient) vector bundles $\operatorname{Ad}(E_1) \to X_1$, and $\operatorname{Ad}(E_0) \to X_0$ form a VB-groupoid $[\operatorname{Ad}(E_1) \rightrightarrows \operatorname{Ad}(E_0)] \to [X_1 \rightrightarrows X_0].$

VB-groupoid $At(\mathbb{E})$.

We leave out the technical details of the construction of the Lie groupoid $[At(E_1) \rightrightarrows At(E_0)]$, which are almost identical to the construction of the Lie groupoid $Ad(\mathbb{E}) = [Ad(E_1) \rightrightarrows Ad(E_0)]$. The Lie groupoid structure on $[At(E_1) \rightrightarrows At(E_0)]$ has the following description. The source, target maps are respectively given by

•
$$s'^{\sim}_* : \operatorname{At}(E_1) \to \operatorname{At}(E_0),$$

$$[(\widetilde{\gamma}, \widetilde{X})] \mapsto [(s(\widetilde{\gamma}), s_{*,\widetilde{\gamma}}(\widetilde{X}))],$$
• $t'^{\sim}_* : \operatorname{At}(E_1) \to \operatorname{At}(E_0),$

$$[(\widetilde{\gamma}, \widetilde{X})] \mapsto [(t(\widetilde{\gamma}), t_{*,\widetilde{\gamma}}(\widetilde{X}))]$$

and composition is given by

$$[(\widetilde{\gamma}_2, \widetilde{X}_2)] \circ [(\widetilde{\gamma}_1, \widetilde{X}_1)] = [(\widetilde{\gamma}_2 \circ \widetilde{\gamma}_1, \widetilde{X}_2 \circ \widetilde{X}_1)]$$

$$(4.7)$$

for appropriate choice of representative elements of the equivalence classes. Then the vector bundles $\operatorname{At}(E_1) \to X_1$ and $\operatorname{At}(E_0) \to X_0$ give a VB-groupoid $[\operatorname{At}(E_1) \rightrightarrows \operatorname{At}(E_0)] \to [X_1 \rightrightarrows X_0]$.

Lemma 4.2. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. The vertical vector fields generating maps $\delta_p \colon L(G_0) \to T_p E_0$ and $\delta_{(p \xrightarrow{\tilde{\gamma}} q)} \colon L(G_1) \to T_{\tilde{\gamma}} E_1$ define a functor $\delta \colon \mathbb{E} \times L(\mathbb{G}) \to T(\mathbb{E})$. Moreover the functor δ is \mathbb{G} equivariant in the sense that, $\delta(pg, \mathrm{ad}_{g^{-1}}(B)) = \delta(p, B) \cdot g$ and $\delta(\tilde{\gamma}k, \mathrm{ad}_{k^{-1}}(K)) = \delta(\tilde{\gamma}, K) \cdot k$, for any $p \in E_0, g \in G_0, B \in L(G_0)$ and $\tilde{\gamma} \in E_1, k \in G_1, K \in L(G_1)$.

Proof. It is an immediate consequence of the functoriality of the Lie 2-group action. \Box

Now it is evident that the maps $\{j_i^{/G_i}\}_{i \in \{0,1\}}$ and $\{(\pi_i)_*^{/G_i}\}_{i \in \{0,1\}}$ in Diagram 4.3 respectively define the morphisms of VB-groupoids, $j^{/\mathbb{G}}$: Ad $\mathbb{E} \to \operatorname{At}(\mathbb{E})$ and $\pi_*^{/\mathbb{G}}$: At $(\mathbb{E}) \to T\mathbb{X}$. Hence we conclude the following.

Proposition 4.3. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Then we have a short exact sequence

$$0 \longrightarrow \operatorname{Ad}(\mathbb{E}) \xrightarrow{j^{/\mathbb{G}}} \operatorname{At}(\mathbb{E}) \xrightarrow{\pi^{/\mathbb{G}}_{*}} T\mathbb{X} \longrightarrow 0$$

$$(4.8)$$

of VB-groupoids over $\mathbb{X} = [X_1 \rightrightarrows X_0]$.

We call the short exact sequence above the *Atiyah sequence* associated to the $[G_1 \rightrightarrows G_0]$ bundle $[E_1 \rightrightarrows E_0] \rightarrow [X_1 \rightrightarrows X_0]$.

5. Connection on principal 2-bundles over Lie groupoids

We have observed that in a traditional principal bundle setup, a splitting of the associated Atiyah sequence gives a connection on the principal bundle. We can adapt the same idea for the framework of the Atiyah sequence of a principal 2-bundle (Diagram 4.8). We will see that the categorical structure involved in a principal 2-bundle offers a more enriched differential geometric connection structure.

5.1. Connection as a splitting of the Atiyah sequence.

Definition 5.1 (strict connection and semi strict connection). Let \mathbb{G} be a Lie 2-group. Let $\pi \colon \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} , and

$$0 \longrightarrow \operatorname{Ad}(\mathbb{E}) \xrightarrow{j^{/\mathbb{G}}} \operatorname{At}(\mathbb{E}) \xrightarrow{\pi^{/\mathbb{G}}} T\mathbb{X} \longrightarrow 0$$
(5.1)

its associated Atiyah sequence. A morphism $R : At(\mathbb{E}) \to Ad(\mathbb{E})$ of VB-groupoids is said to be a *strict connection* if

$$R \circ j^{/\mathbb{G}} = 1_{\mathrm{Ad}(\mathbb{E})}.$$

A morphism $R: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ of VB-groupoids is said to be a *semi-strict connection* if $R \circ j^{/\mathbb{G}}$ is 2-isomorphic to $1_{\operatorname{Ad}(\mathbb{E})}$,

$$R \circ j^{/\mathbb{G}} \simeq 1_{\mathrm{Ad}(\mathbb{E})}$$

in the category 2-VBGpd(X) (Proposition 4.1).

Definition 5.2 (category of strict and semi-strict connections). Let \mathbb{G} be a Lie 2-group, and $\pi: \mathbb{E} \to \mathbb{X}$ a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} . We define the category of strict (resp. semi strict) connections for $\mathbb{E}(\mathbb{X}, \mathbb{G})$ as a category whose objects are $R: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ such that R is a strict (resp. semi strict) connections on $\mathbb{E}(\mathbb{X}, \mathbb{G})$ and morphisms are 2-morphisms $\eta: R \Longrightarrow R'$ of 2-VBGpd(\mathbb{X}). We will denote the categories of strict and semi strict connections respectively by the notations $C_{\mathbb{E}}^{\operatorname{strict}}$ and $C_{\mathbb{E}}^{\operatorname{semi}}$.

It is evident that a strict connection gives a functorial section $T\mathbb{X} \xrightarrow{\Sigma} \operatorname{At}(\mathbb{E})$, with


respect to the VB-groupoid morphism $\operatorname{At}(\mathbb{E}) \xrightarrow{\pi_*^{\mathbb{G}}} T\mathbb{X}$, splitting the tangent bundles $TE_i \to E_i, i \in \{0, 1\}$ into horizontal and vertical subbundles. On the other hand for a semi-strict connection, the natural isomorphism $R \circ j^{/\mathbb{G}} \Longrightarrow 1_{\operatorname{Ad}(\mathbb{E})}$ gives an obstruction for the corresponding VB-groupoid morphism $T\mathbb{X} \xrightarrow{\Sigma} \operatorname{At}(\mathbb{E})$ to be a section of

At(\mathbb{E}) $\xrightarrow{\pi^{/\mathbb{G}}} T\mathbb{X}$. The issue is related to the parallel transport with respect to a connection, which we will pursue in a subsequent paper.

Before providing a description of the connections on a principal 2-bundle in terms of differential forms, we note an immediate consequence of our definition.

Lemma 5.1. Let π : $(\mathbb{E} = [E_1 \Rightarrow E_0]) \rightarrow (\mathbb{X} = [X_1 \Rightarrow X_0])$ be a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} . Let R_0 : At $(E_0) \rightarrow \operatorname{Ad}(E_0)$ and R_1 : At $(E_1) \rightarrow \operatorname{Ad}(E_1)$ be the splittings associated to be splittings of the respective Atiyah sequences. Then R_0, R_1 defines a strict connection on π : $\mathbb{E} \rightarrow \mathbb{X}$ if and only if R_0, R_1 define a morphism of Lie groupoids from At (\mathbb{E}) to Ad (\mathbb{E}) .

Remark 5.2. For a given $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle $\mathbb{E} = [E_1 \rightrightarrows E_0]$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$, we have a simplicial principal $\mathbb{G}_{\bullet} = \{G_i\}_{i\geq 0}$ bundles $\mathbb{E}_{\bullet} = \{E_i\}_{i\geq 0}$ over $\mathbb{X}_{\bullet} = \{X_i\}_{i\geq 0}$, where $G_{\bullet}, E_{\bullet}, X_{\bullet}$ are the respective simplicial manifolds of groupoid nerves associated to Lie groupoids $\mathbb{G} = [G_1 \rightrightarrows G_0], \mathbb{E} = [E_1 \rightrightarrows E_0]$ and $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Note each G_i has a natural Lie group structure compatible with the composition. Then one can further extend Diagram 4.3 to obtain a family of Atiyah sequences



for $i \ge 0$, where *i*-th level is related to i - 1-th level by the usual face and degeneracy maps. In our follow-up paper we will introduce a Chern-Weil theory by incorporating the Atiyah sequences as above into our framework.

5.2. Connections as differential forms.

Definition 5.3. Let $\mathbb{G} = [G_1 \rightrightarrows G_0]$ be a Lie 2-group, and $[E_1 \rightrightarrows E_0]$ a Lie groupoid. An $L(\mathbb{G})$ -valued 1-form on $[E_1 \rightrightarrows E_0]$ is a morphism of Lie groupoids $\omega := (\omega_1, \omega_0) : T\mathbb{E} \rightarrow L(\mathbb{G})$ such that ω_i is an $L(G_i)$ -valued differential 1-form on E_i , for $i \in \{0, 1\}$. If \mathbb{G} acts on \mathbb{E} and $\omega : T\mathbb{E} \rightarrow L(\mathbb{G})$ is \mathbb{G} -equivariant, then ω called a \mathbb{G} -equivariant 1-form.

The notion of a differential form on a Lie groupoid, called *multiplicative forms*, is already available in literature [15]. It would be illustrative to express an $L(\mathbb{G})$ valued differential 1-form in terms of the crossed module to see the relation between our definition and the notion of multiplicative forms. For that, suppose (G, H, τ, α) is the Lie crossed module associated to \mathbb{G} . Then $G_0 = G$ and $G_1 = H \rtimes_{\alpha} G$ and $L(G_0) = L(G)$, $L(G_1) =$ $L(H) \oplus L(G)$. Let $\omega := (\omega_1, \omega_0) : T\mathbb{E} \to L(\mathbb{G})$ is an $L(\mathbb{G})$ -valued 1-form. Let ω_{1G}, ω_{1H} respectively be the L(G) and L(H)-valued components of ω_1 . The functoriality condition on ω , can be expressed as:

$$\omega_{1G} = s^* \omega_0,$$

$$t^* \omega_0 = s^* \omega_0 + \tau(\omega_{1H}),$$

$$m^* \omega_{1H} = \operatorname{pr}_1^* \omega_{1H} + \operatorname{pr}_2^* \omega_{1H},$$

(5.2)

where $m: E_1 \times_{s,E_0,t} E_1 \to E_1$ is the composition map and $\operatorname{pr}_i: E_1 \times_{s,E_0,t} E_1 \to E_1, i \in \{1,2\}$ are the projection maps to respectively first and second components. Precisely the third condition implies ω_{1H} is an L(H)-valued multiplicative form on the groupoid $[E_1 \rightrightarrows E_0]$. In particular, if G = e in the crossed module set-up, we recover the definition of a multiplicative form given in [15].

We also refer to [43] for a different, but related, notion of an $L(\mathbb{G})$ -valued differential form on $\mathbb{E} = [E_1 \rightrightarrows E_0]$, where the author considers simplicial manifold \mathbb{E}_{\bullet} defined by the nerve of the Lie groupoid \mathbb{E} , and defines an $L(\mathbb{G})$ -valued differential form as a 'suitably chosen' subcomplex of the double complex $T^p := \bigoplus_{p=i+j+k} \Omega^i(E_j, \mathfrak{G}_k)$, where $\mathfrak{G}_{-1} = L(G), \mathfrak{G}_0 = L(H), \mathfrak{G}_i = 0 \forall i \neq -1 \text{ or } 0$. While the choice of the subcomplex eluded above is in fact motivated by the connection structure on a principal 2-bundle over a manifold, motivation for our definition of $L(\mathbb{G})$ -valued differential forms is to find an infinitesimal representation of the strict and semi-strict connection arising out of splitting of the Atiyah sequence. Note further if we attach an L(H)-valued differential 2-form on E_0 with our differential form ω in Definition 5.3 we obtain a differential 1-form as per the definition in [43].

Example 5.3. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a Lie groupoid. Suppose the Lie 2-group $\hat{H} := [H \rightrightarrows e]$, associated to a Lie group H, acts on \mathbb{E} . Then an $L(\hat{H})$ -valued 1-form on \mathbb{E} is same as an L(H)-valued multiplicative 1-form on \mathbb{E} . If the L(H)-valued multiplicative 1-form on \mathbb{E} is H-equivariant, then so is the corresponding $L(\hat{H})$ -valued 1-form on \mathbb{E} .

Example 5.4. If a Lie group G acts on a smooth manifold M, then an L(G)-valued 1-form is same as an $L(G \rightrightarrows G)$ -valued 1-form on the Lie groupoid $[M \rightrightarrows M]$. If the form on M is G-equivariant, then the corresponding $L(G \rightrightarrows G)$ -valued 1-form on the Lie groupoid $[M \rightrightarrows M]$ is $[G \rightrightarrows G]$ invariant.

Example 5.5. If the Lie 2-group $[G \rightrightarrows G]$ acts on a Lie groupoid $\mathbb{E} = [E_1 \rightrightarrows E_0]$. Then an $L(\mathbb{G})$ -valued 1-form on the Lie groupoid $[E_1 \rightrightarrows E_0]$ is an L(G)-valued 1-form on E_0 such that $t^*\omega = s^*\omega$. The equivariancy of one implies the equivariancy of the other.

Example 5.6. Let (G, H, τ, α) be a Lie crossed module associated to a Lie 2-group \mathbb{G} . Let the Lie 2-group $[G \rightrightarrows G]$ acts on a Lie groupoid $\mathbb{E} = [E_1 \rightrightarrows E_0]$. We can define another Lie groupoid $\mathbb{E} \times H := [E_1 \times H \rightrightarrows E_0]$ with source, target and composition maps are respectively given by $\tilde{s}(\gamma, h) = s(\gamma), \tilde{t}(\gamma, h) = t(\gamma)\tau(h^{-1})$ and $(\gamma_2, h_2) \circ (\gamma_1, h_1) = ((\gamma_2\tau(h_1)) \circ \gamma_1, h_2h_1)$. The unit and inverse maps respectively are $1_p = (1_p, e)$

and $(\gamma, h)^{-1} = (\gamma^{-1}\tau(h^{-1}), h^{-1})$. Moreover \mathbb{G} has an action on $\mathbb{E} \times H := [E_1 \times H \Rightarrow E_0]$ given by $(p, g) \mapsto pg$ for the objects and $(\gamma, h)(h', g') = (\gamma g, \alpha_{g^{-1}}(h'^{-1}h))$ for the morphisms. Let ω be an L(G)-valued G-equivariant 1-form on E_0 such that $s^*\omega = t^*\omega$. Then $\widetilde{\omega}$, defined as

$$\widetilde{\omega}_{\gamma,h}(X,\mathfrak{K}) := \mathrm{ad}_{(h,e)}\,\omega(s_{*\gamma}(X)) - \mathfrak{K} \cdot h^{-1},$$

and ω gives an $L(\mathbb{G})$ -valued \mathbb{G} -equivariant 1-form on Lie groupoid $\mathbb{E} \times H := [E_1 \times H \rightrightarrows E_0]$. In the next subsection we will see this example follows from a more general construction.

Our next goal is to express the connections (strict and semi-strict) on a principal 2bundle as differential forms. The following definition is natural.

Definition 5.4 (Category of \mathbb{G} -equivariant $L(\mathbb{G})$ -valued 1-forms). Let $\mathbb{G} = [G_1 \rightrightarrows G_0]$ be a Lie 2 group acting on a Lie groupoid $\mathbb{E} = [E_1 \rightrightarrows E_0]$. The *category of* \mathbb{G} -equivariant $L(\mathbb{G})$ -valued 1-forms on \mathbb{E} , denoted $\Omega_{\mathbb{E}^G}$, is defined as follows.

- objects are \mathbb{G} -equivariant 1-forms on \mathbb{E} ,
- morphisms are smooth natural isomorphisms $\eta: \omega \Longrightarrow \omega'$ such that $\eta((p, v) \cdot g) = \eta(p, v) \cdot 1_g$ for all $(p, v) \in TE_0, g \in G_0$, and $\eta: TE_0 \to L(G_1)$ is an $L(G_1)$ -valued 1-form on E_0 .

Observe that a morphism of VB-groupoids $S: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$

$$\begin{array}{ccc} \operatorname{At}(\mathbb{E}) & \stackrel{S}{\longrightarrow} & \operatorname{Ad}(\mathbb{E}) \\ & & \downarrow^{\pi} & & \downarrow^{\pi} \\ & & \mathbb{X} & \longrightarrow & \mathbb{X} \end{array}$$

is of the form

$$[(p,v)] \mapsto [(p,\omega(p,v))], \forall [p,v] \in \operatorname{At}(E_0), \\ [(\widetilde{\gamma},\widetilde{X})] \mapsto [(\widetilde{\gamma},\omega(\widetilde{\gamma},\widetilde{X}))], \forall [(\widetilde{\gamma},\widetilde{X})] \in \operatorname{At}(E_1)$$

and in turn defines a \mathbb{G} -equivariant $L(\mathbb{G})$ -valued 1-form $\omega \colon T\mathbb{E} \to L(\mathbb{G})$. Obviously converse also holds.

Proposition 5.7. Let $\pi: \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} . Let $R: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ be a morphism of VB-groupoids and $\omega: T\mathbb{E} \to L(\mathbb{G})$ the associated \mathbb{G} -equivariant $L(\mathbb{G})$ -valued differential 1-form on the Lie groupoid \mathbb{E} .

(1) $R: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ is a strict connection as in Definition 5.1 if and only if the following diagram of morphisms of Lie groupoids

commutes up to nose.

(2) $R: \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ is a semi-strict connection if and only if the Diagram 5.3 commutes up to a \mathbb{G} -equivariant, fiber-wise linear natural isomorphism.

Proof. (1) If R is a strict connection then $R \circ j^{/\mathbb{G}} = 1_{\mathrm{Ad}(\mathbb{E})}$. That means the corresponding \mathbb{G} -invariant $L(\mathbb{G})$ valued differential form $(\omega_1, \omega_0) = \omega \colon T\mathbb{E} \to L(\mathbb{G})$ define a pair of classical connections on G_0 -bundle $E_0 \to X_0$ and G_1 -bundle $E_1 \to X_1$ respectively. Hence ω_0, ω_1 satisfy

$$\omega_i(x_i, \delta_{x_i}(A_i)) = A_i, \qquad i \in \{0, 1\}$$

for all $x_i \in E_i, A_i \in L(G_i)$, giving the commutation relation in Diagram 5.3. The converse also easily follows.

(2) Now suppose R is a semi strict connection. Let $\epsilon \colon R \circ j^{/\mathbb{G}} \Longrightarrow 1_{\mathrm{Ad}(\mathbb{E})}$ be a 2 isomorphism in 2-VBGpd(X):

$$\epsilon_{([p,B])} \colon R([p, \delta_p(B)]) \longrightarrow [(p,B)].$$

We claim that, since $\pi(\epsilon([p, B]) = 1_{\pi(p)})$, we have a unique $(\omega(p, \delta_p(B)) \xrightarrow{\kappa_{(p,B)}} B) \in L(G_1)$ such that $\eta([p, B]) = [(1_p, \kappa_{(p,B)})]$, for a chosen $p \in E_0, B \in L(G_0)$. To see this note that if $\epsilon([p, B]) = [(\gamma_{p,B}, \alpha_{p,B})]$, for $\gamma_{p,B} \in E_1, \alpha_{p,B} \in L(G_1)$, then $[s(\gamma_{p,B})] = [p] \Rightarrow \exists ! g \in G_0$ such that $s(\gamma_{p,B})g = p$. That means $\epsilon([p, B]) = [(\gamma_{p,B}1_g, \operatorname{ad}_{1_g^{-1}}(\alpha_{p,B}))]$, with $\gamma_{p,B}1_g$ starts at p and projects down to $1_{\pi(p)} = \pi(1_p)$ in the G_1 -bundle $E_1 \to X_1$. Hence choosing the unique element of G_1 to translate $\gamma_{p,B}1_g$ to 1_p , we obtain the unique $\left(\omega(p, \delta_p(B)) \xrightarrow{\kappa_{(p,B)}} B\right) \in L(G_1)$ such that $\eta([p, B]) = [(1_p, \kappa_{(p,B)})]$. It is readily checked that $\kappa \colon \omega \circ \delta \Longrightarrow \operatorname{pr}_2$ is a natural isomorphism. Moreover smoothness of ϵ implies smoothness κ . If we represent the equivalence class [(p, B)] by some other element $(p', B') = (pg, \operatorname{ad}_{g^{-1}}(B))$, we end up with $\epsilon([p', B']) = [(1_{p'}, \kappa_{(p',B')})]$. Implying $(1_{p'}, \kappa_{(p',B')}) \sim (1_p, \kappa_{(p,B)})$. But that means $(1_{p'}, \kappa_{(p',B')}) = (1_p, \kappa_{(p,B)}) \cdot 1_g$ [this follows from the fact that p' = pg and freeness of action on the fiber of $E_1 \to X_1$]. Which gives us the equivalence of $\kappa :$

$$\kappa(pg, \mathrm{ad}_{g^{-1}}(B)) = \mathrm{ad}_{1_g^{-1}}(\kappa(p, B)).$$

Conversely given a G-equivariant, fiber-wise linear natural transformation $\kappa \colon \omega \circ \delta \implies \mathrm{pr}_2$, we define $\epsilon \colon R \circ j^{/\mathbb{G}} \implies 1_{\mathrm{Ad}(\mathbb{E})}$ by $\epsilon([p,B]) := [(1_p,\kappa_{(p,B)})]$. The map is well defined for the following reason. If $(p',B') \in [(p,A)]$, then $p' = pg, B' = \mathrm{ad}_g(B)$ for some $g \in G$. Then equivariancy of κ implies $(1_{p'},\kappa_{(p',B')}) = (1_p,\kappa_{(p,B)})1_g$. Thus $[(1_{p'},\kappa_{(p',B')})] = [(1_p,\kappa_{(p,B)})]$. ϵ satisfies $\pi(\epsilon_{[p,B]}) = 1_{\pi(p)}$. The map is obviously smooth. Verification that η is a natural transformation and a morphism in 2-VBGpd(X) is straightforward.

Let $\pi: \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} . A \mathbb{G} -equivariant $L(\mathbb{G})$ -valued 1-form $\omega: T\mathbb{E} \to L(\mathbb{G})$ satisfying the property (1) in Proposition 5.7 will be called a *strict* connection 1-form, whereas a \mathbb{G} -equivariant $L(\mathbb{G})$ -valued 1-form $\omega: T\mathbb{E} \to L(\mathbb{G})$ satisfying the property (2) in Proposition 5.7 will be called a *semi-strict connection* 1-form.

Definition 5.5 (Category of strict and semi-strict connection 1-forms). Let $\pi : \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over a Lie groupoid \mathbb{X} . The category of strict and semi-strict connections have objects respectively the strict connection 1-forms and semi-strict connection 1-forms and have morphisms the smooth natural isomorphisms $\eta : \omega \Longrightarrow \omega'$ such that $\eta((p, v) \cdot$

 $g) = \eta(p, v) \cdot 1_g$ for all $(p, v) \in TE_0$ and $g \in G_0$. The category of strict and semi-strict connections will be denoted respectively as $\Omega_{\mathbb{E}}^{\text{strict}}$ and $\Omega_{\mathbb{E}}^{\text{semi}}$.

Now it is obvious that we have a sequence of categories

$$\Omega^{\text{strict}}_{\mathbb{E}} \subset \Omega^{\text{semi}}_{\mathbb{E}} \subset \Omega_{\mathbb{E}^{\mathbb{G}}}$$

Let us further examine the natural isomorphism $\kappa \colon \omega \circ \delta \Longrightarrow \operatorname{pr}_2$ of the Diagram 5.3 for a semi-strict connection $\omega \colon T\mathbb{E} \to L(\mathbb{G})$. We express \mathbb{G} by its crossed module (G, H, τ, α) . Then for each $(p, B) \in E_0 \times L(G_0)$ we have a $\kappa(p, B)$ of the form $\kappa(p, B) = (\kappa_{\omega}(p, B), \omega(p, \delta_p(B)))$, where $\kappa_{\omega}(p, B) \in L(H)$ and

$$\omega(p,\delta_p(B)) - B = -\tau(\kappa_\omega(p,B))$$
(5.4)

Thus $\kappa_{\omega}(p, B)$ measures the deviation $\omega(p, \delta_p(B)) - B$. In turn given a κ we have a smooth fiber-wise linear map $\kappa_{\omega} \colon E_0 \times L(G) \to L(H)$. The equivariancy of κ implies

$$\kappa_{\omega}(pg, \operatorname{ad}_{g^{-1}} B) = \alpha_{g^{-1}}(\kappa_{\omega}(p, B)).$$
(5.5)

Since κ is a natural transformation, for $(A \xrightarrow{\kappa} B) \in L(G_1)$ and $(p \xrightarrow{\tilde{\gamma}} q) \in L(E_1)$ the following diagram commutes,

and we arrive at the condition

$$\omega_{1,\widetilde{\gamma}}(\delta_{\widetilde{\gamma}}(K)) - K = (\kappa_{\omega}(p,A) - \kappa_{\omega}(q,B), -\tau(\kappa_{\omega}(p,A))).$$
(5.6)

Consider the Lie 2-sub algebra $L(\mathbb{G})^{\tau} := [L(H) \rtimes \tau(L(H)) \rightrightarrows \tau(L(H))]$ of $L(\mathbb{G}) = [L(H) \rtimes L(G) \rightrightarrows L(G)]$. Observe that $L(\mathbb{G})^{\tau}$ is invariant under the adjoint action of \mathbb{G} .

Observe that Equation (5.6) gives us the following uniqueness property.

Lemma 5.8. Let ω be a semi strict connection 1-form on the $\mathbb{G} = (G, H, \tau, \alpha)$ -bundle $\mathbb{E} = [E_1 \rightrightarrows E_0]$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Then the natural transformation κ in Diagram 5.3 is unique.

Now we conclude the following.

Proposition 5.9. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $\mathbb{G} = (G, H, \tau, \alpha)$ -bundle over $\mathbb{X} = [X_1 \rightrightarrows X_0]$. A \mathbb{G} -equivariant 1-form $\omega \colon T\mathbb{E} \to L(\mathbb{G})$ is a semi-strict connection 1-form if and only if the functor $(\omega \circ \delta - \operatorname{pr}_2) := \widehat{\kappa}_{\omega}$ is of the form $\widehat{\kappa}_{\omega}(p, B) = \tau(\kappa_{\omega}(p, B))$ and $\widehat{\kappa}_{\omega}(\gamma, K) = (\kappa_{\omega}(p, A) - \kappa_{\omega}(q, B), -\tau(\kappa_{\omega}(p, A)))$ for a smooth, fiber wise linear map $\kappa_{\omega} \colon E_0 \times L(\mathbb{G}) \to L(H)$ satisfying Equation (5.5). In particular, if $\omega \circ \delta - \operatorname{pr}_2$ is the zero functor then we have a strict connection 1-form.

Corollary 5.10. Let $\omega = (\omega_0, \omega_1)$: $T\mathbb{E} \Rightarrow L(\mathbb{G})$ be a strict connection 1-form on the $\mathbb{G} = (G, H, \tau, \alpha)$ -bundle $\mathbb{E} = [E_1 \Rightarrow E_0]$ over $\mathbb{X} = [X_1 \Rightarrow X_0]$. Let $\lambda \colon TE_0 \to L(H)$ an L(H)-valued 1-form on E_0 satisfying the equivariance property $\lambda(pg, v \cdot g) = \alpha_{g^{-1}}(\lambda(p, v))$. Then $\widetilde{\omega}_0 = \omega_0 + \tau(\lambda)$ and $\widetilde{\omega}_1 = \omega_1 + \tau(s^*\lambda) + t^*\lambda - s^*\lambda$ define a semi strict connection $\widetilde{\omega} = (\widetilde{\omega}_0, \widetilde{\omega}_1) \colon T\mathbb{E} \to L(\mathbb{G})$.

Proof. The functoriality of $(\widetilde{\omega}_0, \widetilde{\omega}_1)$ is immediate. Since $\omega_{0p}(\delta_p(A)) = A$, for $A \in L(G_0)$, only thing remained to be verified is the \mathbb{G} -equivariancy of the functor

$$\Lambda: T\mathbb{E} \to L(\mathbb{G}),
(p,v) \mapsto \tau(\lambda(p,v)),
(\widetilde{\gamma}, \widetilde{X}) \mapsto \tau(\lambda(s(\widetilde{\gamma}), s_{*,\widetilde{\gamma}}(X))) + \lambda(t(\widetilde{\gamma}), t_{*,\widetilde{\gamma}}(X)) - \lambda(s(\widetilde{\gamma}), s_{*,\widetilde{\gamma}}(X)).$$
(5.7)

Then $\Lambda((p, v)g) = \tau(\lambda(pg, v \cdot g)) = \operatorname{ad}_{g^{-1}}(\Lambda(p, v))$. To see the equivariancy at the morphism level, consider $s(\widetilde{\gamma}, \widetilde{X}) = (p, v), t(\widetilde{\gamma}, \widetilde{X}) = (q, w)$. Then

$$\begin{split} \Lambda((\widetilde{\gamma},\widetilde{X})(h,g)) &= \tau(\lambda(pg,vg)) - \lambda(pg,vg) + \lambda(q\tau(h)g,w\tau(h)g) \\ &= \mathrm{ad}_{(e,g)^{-1}}\tau(\lambda(p,v)) - \alpha_{g^{-1}}(\lambda(p,v)) + \alpha_{g^{-1}\tau(h^{-1})}(\lambda(q,w)) \\ &= \mathrm{ad}_{(e,g)^{-1}}\tau(\lambda(p,v)) - \alpha_{g^{-1}}(\lambda(p,v)) + \underbrace{\alpha_{g^{-1}}(\mathrm{ad}_{h^{-1}}(\lambda(q,w)))}_{\mathrm{ad}_{(h,g)^{-1}}(\lambda(q,w))} \\ &= \underbrace{\mathrm{ad}_{(h,g)^{-1}}\tau(\lambda(p,v)) - \alpha_{g^{-1}}(h^{-1}) \cdot (\bar{\alpha}_{\alpha_{g^{-1}}(h)}(\mathrm{ad}_{g^{-1}}(\tau(\lambda(p,v)))))}_{\mathrm{ad}_{(e,g)^{-1}}\tau(\lambda(p,v))[using formulae in Equation (2.8)]} \\ &= \underbrace{\mathrm{ad}_{(h,g)^{-1}}((\lambda(p,v)) + (\alpha_{g^{-1}}(\mathrm{ad}_{h^{-1}}(\lambda(p,v))) - \alpha_{g^{-1}}(\lambda(p,v)))}_{-\alpha_{g^{-1}}(\lambda(p,v))[using formulae in Equation (2.8)]} \\ &= \operatorname{ad}_{(h,g)^{-1}}\tau(\lambda(p,v)) - \operatorname{ad}_{(h,g)^{-1}}((\lambda(p,v)) + \operatorname{ad}_{(h,g)^{-1}}(\lambda(q,w)) \\ &+ [-\alpha_{g^{-1}}(h^{-1}) \cdot (\bar{\alpha}_{\alpha_{g^{-1}}(h)}(\mathrm{ad}_{g^{-1}}(\tau(\lambda(p,v))))) + (\alpha_{g^{-1}}(\mathrm{ad}_{h^{-1}}(\lambda(p,v))) - \alpha_{g^{-1}}(\lambda(p,v)))] \\ &= \operatorname{ad}_{(h,g)^{-1}}\tau(\lambda(p,v)) - \operatorname{ad}_{(h,g)^{-1}}((\lambda(p,v)) + \operatorname{ad}_{(h,g)^{-1}}(\lambda(q,w)). \\ \\ &= \operatorname{ad}_{(h,g)^{-1}}\tau(\lambda(p,v)) - \operatorname{ad}_{(h,g)^{-1}}((\lambda(p,v)) + \operatorname{ad}_{(h,g)^{-1}}(\lambda(q,w)). \\ \end{array}$$

The Corollary 5.10 gives a very straightforward way of finding a semi-strict connection from a strict connection.

Example 5.11. Let $P \to M$ be a traditional principal *G*-bundle over the smooth manifold M. A connection ω on $P \to M$ defines a strict connection (ω, ω) on principal Lie 2-group $[G \rightrightarrows G]$ -bundle $[P \rightrightarrows P] \to [M \rightrightarrows M]$ over the Lie groupoid $[M \rightrightarrows M]$. A strict connection is same as a semi strict connection for such a Lie 2-group bundle.

Example 5.12. Let $[E_1 \Rightarrow E_0]$ be a principal $[G \Rightarrow G]$ -bundle over $[X_1 \Rightarrow X_0]$. Then as noted in Example 3.7, $E_0 \to X_0$ is a principal *G*-bundle over $[X_1 \Rightarrow X_0]$ and $[E_1 \Rightarrow E_0] =$ $[s^*E_0 \Rightarrow E_0]$. Then a strict connection ω on $[G \Rightarrow G]$ -bundle $[E_1 \Rightarrow E_0] \to [X_1 \Rightarrow X_0]$ is a connection on the principal *G*-bundle $E_0 \to X_0$ such that $s^*\omega = t^*\omega$. A strict connection is same as a semi strict connection for such a Lie 2-group bundle.

Remark 5.13. In Example 5.12 we recover the definition of a connection on a principal G-bundle over a Lie groupoid $[X_1 \rightrightarrows X_0]$ as in [27].

Example 5.14. Consider the principal $[H \Rightarrow e]$ -bundle $[E_1 \Rightarrow X_0]$ over $[X_1 \Rightarrow X_0]$ of Example 3.8. Then a strict connection is a classical connection ω on the principal Hbundle $[E_1 \rightrightarrows X_1]$ such that ω is a multiplicative 1-form on the Lie groupoid $[E_1 \rightrightarrows X_0]$. On the other hand any L(H)-valued H-equivariant multiplicative form on $[E_1 \rightrightarrows X_1]$ is a semi strict connection (see also Example 5.3).

Example 5.15. Given a connection ω_0 on a principal *G*-bundle $E_0 \to X_0$, we define a connection on the $[G \times G \rightrightarrows G]$ -bundle $[E_1 \rightrightarrows E_0]$ over $\mathbb{X} = [X_1 \rightrightarrows X_0]$ in Example 3.6 by

$$\omega_{1(p,\gamma,q)}(v_1, X, v_2) = \big(\omega_0(v_1), \omega_0(v_2)\big),\,$$

for all $(v_1, X, v_2) \in T_{(p,\gamma,q)}E_1$.

In Section 5.3, we will prescribe a construction of connections on the decorated bundle.

We have shown a bijective correspondence between strict (resp. semi strict) connections and strict (resp. semi strict) connection 1-forms for a principal 2-bundle. We show here this correspondence extends to the respective categories as well.

Theorem 5.16. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle over a Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0].$

- (1) The categories $C_{\mathbb{E}}^{\text{semi}}$ and $\Omega_{\mathbb{E}}^{\text{semi}}$ are isomorphic. (2) The categories $C_{\mathbb{E}}^{\text{strict}}$ and $\Omega_{\mathbb{E}}^{\text{strict}}$ are isomorphic.

(1) The object level association has already been established in Proposition 5.7. Proof. Let (G, H, τ, α) be the associated Lie cross module of \mathbb{G} . Let $R, R' \colon \operatorname{At}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ be a pair of strict connections and $\eta: R \implies R'$ a natural transformation such that $\pi(\eta([p,v])) = 1_{\pi(p)}$, for any $[(p,v)] \in \operatorname{At}(E_0) = TE_0/G_0$. Let $\omega, \omega' \colon T\mathbb{E} \to \mathbb{C}$ $L(\mathbb{G})$ be respective semi-strict connection 1-forms of R and R'. As noted earlier then $R[(p,v)] = [(p,\omega(p,v))], R'[(p,v)] = [(p,\omega'(p,v))].$ Let $\eta_{[p,v]} \colon [(p,\omega(p,v))] \to$ $[(p, \omega'(p, v))]$. We claim that there exists $\omega(p, v) \xrightarrow{\bar{\eta}(p, v)} \omega'(p, v) \in L(G_1)$ such that $[(1_p, \bar{\eta}(p, v))] = \eta_{[p,v]}$. To see this suppose $[(\gamma, K)] \in \eta_{[p,v]}$. That means there exist $g, g' \in G_0$ such that $s(\gamma)g = p, t(\gamma)g' = p, \operatorname{ad}_{g^{-1}}(s(K)) = \omega(p, v)$ and $\operatorname{ad}_{g'^{-1}}(t(K)) = \omega'(p, v).$ Then $(\gamma, K) \cdot 1_g = (\gamma 1_g, \operatorname{ad}_{1_g^{-1}}(K)) := (\gamma_p, K_\omega) \in \eta_{[p,v]}.$ Now as $\pi(\eta([p,v])) = 1_{\pi(p)}$, we have $\pi(\gamma_p) = \pi(1_p)$. Thus both $\gamma_p, 1_p$ are elements of the same fiber on the G_1 -bundle $E_1 \rightarrow X_1$ with the same source p. That means there exists a unique $h \in H$ such that $\gamma_p(h, e) = 1_p$. Comparing the targets $t(\gamma_p)\tau(h) = t(\gamma)g\tau(h) = p = t(\gamma)g'$. Which gives us g' = $g\tau(h) = g\tau(h)g^{-1}g = \tau(\alpha_g(h))g$. Thus $(\gamma_p, K_\omega) \cdot (h, e) = (1_p, \operatorname{ad}_{(h, e)^{-1}}(K_\omega)) \in \eta_{[p, v]}$. Now $\operatorname{ad}_{(h,e)^{-1}} K_{\omega} = \operatorname{ad}_{(h,e)^{-1}} \operatorname{ad}_{1g^{-1}}(K) = \operatorname{ad}_{(h,e)^{-1}} \operatorname{ad}_{(e,g)^{-1}}(K) = \operatorname{ad}_{(h^{-1},g^{-1})} K.$ Now using Equation (2.8) we see $s(\operatorname{ad}_{(h^{-1},g^{-1})}K) = \operatorname{ad}_{g^{-1}}(s(K)) = \omega(p,v)$ and $t(\mathrm{ad}_{(h^{-1},q^{-1})}K) = \mathrm{ad}_{\tau(h^{-1})q^{-1}}(t(K)) = \mathrm{ad}_{q'^{-1}}(t(K)) = \omega'(p,v).$ We define $\bar{\gamma}$

$$\bar{j}(p,v) := \operatorname{ad}_{(h,e)^{-1}} K_{\omega}$$

It is a straightforward verification that $\bar{\eta}$ defines a smooth natural transformation $\omega \Longrightarrow \omega'$. Moreover if $(p', v') = (p, v)\theta$, for some $\theta \in G_0$, then $(p', v') \in [(p, v)]$. Then above construction gives $(1_{p'}, \bar{\eta}(p', v')) \in [(1_p, \bar{\eta}(p, v))]$ and

$$(1_{p'}, \bar{\eta}(p', v')) = (1_p, \bar{\eta}(p, v)) 1_{\theta} = (1_{p\theta}, \operatorname{ad}_{1_{\theta}^{-1}} \bar{\eta}(p, v)).$$

Hence we obtain the \mathbb{G} equivariancy condition: $\operatorname{ad}_{1^{-1}}(\bar{\eta}(p,v)) = \bar{\eta}(p\theta,v\theta).$

In the other direction, if $\bar{\eta}: \omega \Longrightarrow \omega'$ is a smooth, \mathbb{G} -equivariant natural transformation between the semi-strict connection 1-forms, we define a natural transformation between the corresponding semi-strict connections $R, R': \operatorname{At}(\mathbb{E}) \longrightarrow \operatorname{Ad}\mathbb{E}$, by $\eta_{[p,v]} = [(1_p, \bar{\eta}(p, v))]$. The G-equivariancy implies the map is well defined. Now $\pi(\eta_{[(p,v)]}) = \pi(1_p) = 1_{\pi(p)}$. Thus we obtain a smooth natural transformation $\eta \colon R \Longrightarrow R'$ satisfying $\pi(\eta_{[(p,v)]}) = 1_{\pi(p)}$.

It is readily checked that both the maps are functorial and inverses of each other. (2) The proof is almost identical to that of the case of the semi-strict connections.

Let $\bar{C}_{\mathbb{E}}^{\text{strict}}, \bar{C}_{\mathbb{E}}^{\text{semi}}, \bar{\Omega}_{\mathbb{E}}^{\text{semi}}$, and $\bar{\Omega}_{\mathbb{E}}^{\text{strict}}$ be the respective collections of connected components of the categories $C_{\mathbb{E}}^{\text{strict}}, C_{\mathbb{E}}^{\text{semi}}, \Omega_{\mathbb{E}}^{\text{semi}}$, and $\Omega_{\mathbb{E}}^{\text{strict}}$. Then the Theorem 5.16 tells us the following.

Corollary 5.17. Let $\mathbb{E} = [E_1 \rightrightarrows E_0]$ be a $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle over a Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0].$

- (1) There exists a bijection between $\bar{C}_{\mathbb{E}}^{\text{strict}}$ and $\bar{\Omega}_{\mathbb{E}}^{\text{strict}}$. (2) There exists a bijection between $\bar{C}_{\mathbb{E}}^{\text{semi}}$ and $\bar{\Omega}_{\mathbb{E}}^{\text{semi}}$.

5.3. Connection on decorated principal 2-bundles. Let (G, H, τ, α) be a Lie crossed module and $\mathbb{G} = [G_1 \rightrightarrows G_0]$ the associated Lie 2-group. Let $\mathbb{E}^{\text{dec}} \to \mathbb{X}$ be the decorated bundle associated to a principal G-bundle $(\pi: E_0 \to X_0, \mu, [X_1 \rightrightarrows X_0])$ (Proposition 3.11). First we compute the differentials of the various structure maps and the action maps associated to the Lie groupoid $\mathbb{E}^{\text{dec}} = [(s^* E_0)^{\text{dec}} \rightrightarrows E_0].$

The source, target and composition maps of the tangent Lie groupoid $[T(s^*E_0)^{\text{dec}}]$ TE_0 are respectively given by

$$\widetilde{s}_{*((\gamma,p),h)}((X,v),\mathfrak{K}) = v,
\widetilde{t}_{*((\gamma,p),h)}((X,v),\mathfrak{K}) = \mu_{*(\gamma,p)}(X,v) \cdot \tau(h^{-1}) - \delta_{\mu(\gamma,p)\tau(h^{-1})}(\tau(\mathfrak{K}) \cdot \tau(h^{-1})),
((\gamma_{2}, p_{2}, h_{2}), (X_{2}, v_{2}, \mathfrak{K}_{2})) \circ ((\gamma_{1}, p_{1}, h_{1}), (X_{1}, v_{1}, \mathfrak{K}_{1}))
= ((\gamma_{2} \circ \gamma_{1}, p_{1}, h_{2}h_{1}), (X_{2} \circ X_{1}, v_{1}, h_{2} \cdot \mathfrak{K}_{1} + \mathfrak{K}_{2} \cdot h_{1}).$$
(5.8)

The differential of the target map \tilde{t} is computed by applying the chain-rule in the following composition of maps

$$s^*E_0 \times H \xrightarrow{(\mathrm{Id},^{-1})} s^*E_0 \times H \xrightarrow{(\mu,\tau)} E_0 \times G \to E_0.$$

Let us compute the vertical vector field generating functor $\delta \colon \mathbb{E}^{\mathrm{dec}} \times L(\mathbb{G}) \to T(\mathbb{E}^{\mathrm{dec}})$ with respect to the action defined in Equation (3.10). We have

$$\delta \colon \mathbb{E}^{\operatorname{dec}} \times L(\mathbb{G}) \to T(\mathbb{E}^{\operatorname{dec}})$$

$$(p, B) \mapsto \delta_p(B)$$

$$((\gamma, p, h)(A, B)) \mapsto \delta_p(B) - \bar{\alpha}_h(B) - A \cdot h.$$
(5.9)

Note that, for a fixed $((\gamma, p), h) \in s^* E_0 \times H$, the map $\delta_{((\gamma, p), h)} \colon H \rtimes G \to s^* E_0 \times H$ is given by $(h', g) \mapsto (\gamma, pg, \alpha_{q^{-1}}(h'^{-1}h))$. The first coordinate is the constant map, the second coordinate is the right translation map, whereas the third coordinate can be decomposed as

$$H \times G \xrightarrow{-1, -1} H \times G \xrightarrow{(-, -)} G \times H \xrightarrow{(\mathrm{Id}, R_h)} G \times H \xrightarrow{\alpha} H.$$

Then the formula in 5.9 is computed by applying the chain-rule.

The derivatives of the right action of \mathbb{G} on \mathbb{E} are given as, for fixed $g \in G, (h', g) \in G_1$,

$$(X, v, \mathfrak{K}) \mapsto (X, vg, \alpha_{g^{-1}}(h'^{-1} \cdot \mathfrak{K})),$$
(5.10)

for $v \in T_p E_0$, $((X, v), \mathfrak{K}) \in T_{((\gamma, p), h)}(s^* E_0^{\text{dec}})$. Here we have adopted the notations of Remark 2.8.

Suppose the principal G-bundle $(\pi: E_0 \to X_0, \mu, [X_1 \rightrightarrows X_0])$ over X admits a connection ω , as defined in [27]; that is ω is a connection on G-bundle $E_0 \to X_0$ satisfying $s^*\omega = t^*\omega$, where $s, t: [s^*E_0 \rightrightarrows E_0]$ respectively given by $s: (\gamma, p) \mapsto p, t: (\gamma, p) \mapsto \mu(\gamma, p)$. The condition explicitly reads

$$\omega_p(v) = \omega_{\mu(\gamma,p)} \big(\mu_{*,(\gamma,p)}(X,v) \big), \tag{5.11}$$

for any $(\gamma, p) \in s^* E_0, (X, v) \in T_{(\gamma, p)}(s^* E_0).$

Define an $L(H \rtimes G)$ -valued differential form on $s^* E_0 \times H$,

$$\omega_{(\gamma,p,h)}^{\mathrm{dec}}(X,v,\mathfrak{K}) = \mathrm{ad}_{(h,e)}(\omega_p(v)) - \mathfrak{K} \cdot h^{-1}.$$

We will prove here that $(\omega^{\text{dec}}, \omega)$ is a strict connection 1-form on \mathbb{G} -bundle \mathbb{E}^{dec} over \mathbb{X} . Note that

$$\omega_{(\gamma,p,h)}^{\text{dec}}(X,v,\mathfrak{K}) = \operatorname{ad}_{(h,e)}(\omega_p(v)) - \mathfrak{K} \cdot h^{-1}, \qquad (5.12)$$

is same as

$$\omega_{((\gamma, p), h)}^{\operatorname{dec}}(X, v, \mathfrak{K}) = \operatorname{ad}_{(h, e)} \big((s^* \omega)_{((\gamma, p), h)}((X, v), \mathfrak{K}) \big) - \Theta_h(\mathfrak{K}),$$

where Θ is the Maurer-Cartan form on H, $((\gamma, p), h) \in s^* E_0 \times H$ and $((X, v), \mathfrak{K}) \in T_{((\gamma, p), h)}(s^* E_0 \times H) = T_{(\gamma, p)}s^* E_0 \oplus T_h H.$

Let us verify that $(\omega^{\text{dec}}, \omega)$ is a functor.

Lemma 5.18. $(\omega^{\text{dec}}, \omega)$ define a functor $T\mathbb{E}^{\text{dec}} \to L(\mathbb{G})$.

Proof. Using Equation (2.8) we write

$$\begin{aligned} & \omega_{(\gamma,p,h)}^{\text{dec}}(X,v,\mathfrak{K}) \\ &= \operatorname{ad}_{(h,e)} \left(\omega_p(v) \right) - \mathfrak{K} \cdot h^{-1} \\ &= \underbrace{\omega_p(v)}_{\in L(G)} + \underbrace{h \cdot \left(\bar{\alpha}_{h^{-1}}(\omega_p(v)) \right) - \mathfrak{K} \cdot h^{-1}}_{\in L(H)}.
\end{aligned}$$

Let $((\gamma_2, p_2, h_2), (X_2, v_2, \mathfrak{K}_2))$ and $((\gamma_1, p_1, h_1), (X_1, v_1, \mathfrak{K}_1))$ be a pair of composable morphisms in $T\mathbb{E}^{dec}$, which means, by Equation (5.8), in particular

$$p_{2} = \mu(\gamma_{1}, p_{1})\tau(h_{1})^{-1}$$

$$v_{2} = \mu_{*(\gamma_{1}, p_{1})}(X_{1}, v_{1}) \cdot \tau(h_{1}^{-1}) - \delta_{\mu(\gamma_{1}, p_{1})\tau(h_{1}^{-1})}(\tau(\mathfrak{K}_{1}) \cdot \tau(h_{1}^{-1})).$$
(5.13)

Note

$$s\left(\omega_{(\gamma_{2},p_{2},h_{2})}^{\text{dec}}(X_{2},v_{2},\mathfrak{K}_{2})\right) = \omega_{p_{2}}(v_{2}),$$

$$t\left(\omega_{(\gamma_{1},p_{1},h_{1})}^{\text{dec}}(X_{1},v_{1},\mathfrak{K}_{1})\right) = \omega_{p_{1}}(v_{1}) + \tau(h_{1}^{-1} \cdot (\bar{\alpha}_{h_{1}}(\omega_{p_{1}}(v_{1}))) - \mathfrak{K}_{1} \cdot h_{1}^{-1}).$$
(5.14)

Now plugging the equations of Equation (5.13) into the equations above and noting that ω is a connection on *G*-bundle $E_0 \to X_0$ satisfying $\omega_{(\mu(\gamma_1, p_1))}(\mu_*(X_1, v_1)) = \omega_{p_1}(v_1)$ (see Equation (5.11)), we obtain the required source-target consistency.

To verify the composition we separately compute

$$\left(\omega_{(\gamma_2,p_2,h_2)}^{\operatorname{dec}}(X_2,v_2,\mathfrak{K}_2)\right)\circ\left(\omega_{(\gamma_1,p_1,h_1)}^{\operatorname{dec}}(X_1,v_1,\mathfrak{K}_1)\right)$$

and

$$\omega_{(\gamma_2,p_2,h_2)\circ(\gamma_1,p_1,h_1)}^{\operatorname{dec}}((X_2,v_2,\mathfrak{K}_2)\circ(X_1,v_1,\mathfrak{K}_1)),$$

using last of the equations in Equation (5.8). As before, we plug-in the relations in Equation (5.14) and use the properties of ω . After some calculation we derive,

$$\left(\omega_{(\gamma_2, p_2, h_2)}^{\text{dec}}(X_2, v_2, \mathfrak{K}_2) \right) \circ \left(\omega_{(\gamma_1, p_1, h_1)}^{\text{dec}}(X_1, v_1, \mathfrak{K}_1) \right)$$

= $\omega_{p_1}(v_1) - \mathfrak{K}_1 \cdot h_1^{-1} - \mathfrak{K}_2 \cdot h_2^{-1}$
+ $\left[h_2 \cdot \left(\bar{\alpha}(h_2^{-1})(\operatorname{ad}_{\tau(h_1)}(\omega_{p_1}(v_1))) \right) + h_1 \cdot (\bar{\alpha}(h_1^{-1})(\omega_{p_1}(v_1))) \right]$ (5.15)

and

$$\omega_{(\gamma_2,p_2,h_2)\circ(\gamma_1,p_1,h_1)}^{\text{dec}} \left((X_2, v_2, \mathfrak{K}_2) \circ (X_1, v_1, \mathfrak{K}_1) \right) \\ = \omega_{p_1}(v_1) - \mathfrak{K}_1 \cdot h_1^{-1} - \mathfrak{K}_2 \cdot h_2^{-1} + \left[h_2 h_1 \cdot \left(\bar{\alpha}(h_1^{-1} h_2^{-1})(\omega_{p_1}(v_1)) \right) \right].$$
(5.16)

Then Lemma 2.9 implies the terms inside [] on both equations are identical. Since ω was multiplicative, we conclude $(\omega^{\text{dec}}, \omega)$ is a functor $T\mathbb{E}^{\text{dec}} \to L(\mathbb{G})$.

Lemma 5.19. ω^{dec} is a connection on $H \rtimes_{\alpha} G = G_1$ -bundle $s^*E_0 \times H \to X_1$.

Proof. Using Equations (5.9) and (5.10) one directly verifies

$$\omega_{((\gamma,p),h)\cdot(h',g')}^{\operatorname{dec}}((X,v,\mathfrak{K}))\cdot(h',g')) = \operatorname{ad}_{(h'g')^{-1}}\left(\omega_{((\gamma,p),h)}^{\operatorname{dec}}((X,v),\mathfrak{K})\right)$$

and

$$\omega^{\mathrm{dec}}_{((\gamma,\,p),\,h)}\big(\delta^{(A+B)}_{((\gamma,\,p),\,h))}\big)=A+B$$

for $A \in L(H), B \in L(G)$.

Combining the last two lemmas we obtain the following result.

Proposition 5.20. Let $(\pi: E_0 \to X_0, \mu, [X_1 \rightrightarrows X_0])$ be a principal *G*-bundle over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ and ω a connection on it. Let $(G, H, \tau, \alpha) = \mathbb{G}$ be a Lie 2-group. Then $(\omega^{\text{dec}}, \omega)$ is a strict connection 1-form on \mathbb{G} -bundle \mathbb{E}^{dec} over \mathbb{X} , where ω^{dec} is as defined in Equation (5.12). Appropriately modifying the connection $(\omega^{\text{dec}}, \omega)$ using Corollary 5.10, we obtain a semistrict connection.

Corollary 5.21. Every principal 2-bundle over a discrete Lie groupoid admits a connection.

Proof. Let $[E_1 \rightrightarrows E_0]$ be a $[G_1 \rightrightarrows G_0]$ -bundle over a discrete Lie groupoid $[X \rightrightarrows X]$. Every such principal 2-bundle is a decorated bundle (Corollary 3.18). Then any choice of a connection on the underlying principal G_0 -bundle over X gives a connection on the principal 2-bundle (Proposition 5.20).

The pair of groups $(\mathbb{R}^n \rtimes G, G)$, where $G \subset GL(n, \mathbb{R})$ is of particular interest in Cartan geometry (See [25, 41, 16]). We make one passing observation here relating our construction with that of Cartan's connections (without going into the specifics).

Example 5.22. Suppose $E_0 \to X_0$ is a *G*-bundle. Consider the $\mathbb{G} = (G, H, \tau, \alpha)$ -bundle $[(E_0 \times H) \rightrightarrows E_0]$ over the Lie groupoid $[X_0 \rightrightarrows X_0]$ (see Example 3.13). Let θ be a *Cartan connection* for the pair $(G \subset \ker(\tau) \rtimes G, \ker(\tau) \rtimes G)$ on the principal *G*-bundle $E_0 \to X_0$ ([16, Definition 2.2]), that is a $L(\ker(\tau) \rtimes G)$ -valued 1-form θ on E_0 satisfying the properties,

$$\begin{aligned} \theta_p \colon T_p E_0 &\to L(\ker(\tau) \rtimes G) \text{ is a linear isomorphism for every } p \in E_0, \\ \theta_{pg}(v \cdot g) &= \operatorname{ad}_{(e,g)^{-1}}(\theta(v)), \qquad \forall p \in E_0, g \in G, \\ \theta_p(\delta_p(B)) &= B, \qquad \forall B \in L(G). \end{aligned}$$

Then one obtains a connection

$$\hat{\theta}_{(p,h)}(v, \mathfrak{K}) = \mathrm{ad}_{(h,e)}(\theta_p(v)) - \mathfrak{K} \cdot h^{-1}$$

on $H \rtimes_{\alpha} G$ -bundle $[(E_0 \times H) \rightrightarrows E_0]$. The source maps $E_0 \times H \to E_0, (p, h) \mapsto p$ and $H \rtimes_{\alpha} G \to G, (h, g) \mapsto g$ define a map from the principal $H \rtimes_{\alpha} G$ -bundle $E_0 \times H \to X_0$ to the principal G-bundle $E_0 \to X_0$. By push forwarding the connection $\tilde{\theta}$ on $H \rtimes_{\alpha} G$ -bundle $E_0 \times H \to X_0$ along this map of principal bundles we obtain a connection θ_0 on the G-bundle $E_0 \to X_0$. Then $(\tilde{\theta}, \theta_0)$ is a connection 1-form on \mathbb{G} -principal bundle $[E_0 \times H \rightrightarrows E_0]$ over $[X_0 \rightrightarrows X_0]$.

5.4. On the existence of connections on a principal 2-bundle. In this section, we comment on the existence of connections on principal 2-bundle over a Lie groupoid.

We call a Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ proper étale if the source map (and hence the target map as well) is a proper étale map. It is already known that a principal *G*-bundle $(E_0 \rightarrow X_0, \mu, \mathbb{X})$ over a proper étale Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$ admits a connection ([27, Theorem 3.16]), that is there exists a connection ω on the principal *G*-bundle $E_0 \rightarrow X_0$ satisfying


$$\omega_p = \omega_{\mu(\gamma, p)}.$$

Let $(G, H, \tau, \alpha) = \mathbb{G}$ be a Lie 2-group. We have seen in Section 5.3 that if a Lie group *G*-bundle $(E_0 \to X_0, \mu, \mathbb{X})$ admits a connection then the decorated bundle \mathbb{E}^{dec} over \mathbb{X} admits a connection. The Proposition 3.17 gives us a criteria for existence of a connection on a Lie 2-group bundle over a proper étale Lie groupoid.

Proposition 5.23. Let $(G, H, \tau, \alpha) = \mathbb{G}$ be a Lie 2-group, and $\mathbb{E} = [E_1 \rightrightarrows E_0]$ a \mathbb{G} -bundle over a proper étale Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. If the \mathbb{G} -bundle \mathbb{E} admits a categorical connection, then it also admits a connection.

6. Gauge 2-group of a principal 2-bundle

In Definition 3.6 we have defined Gauge 2-group $\mathcal{G}(\mathbb{E})$ of a \mathbb{G} -bundle $\mathbb{E} \to \mathbb{X}$. Consider the 2-group $C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$, whose objects are morphisms of Lie groupoids $\sigma \colon \mathbb{E} \to \mathbb{G}$ such that the diagram



commutes, and morphisms are natural transformations $\Phi: \sigma_1 \Longrightarrow \sigma_2$ satisfying $\Phi(pg) = 1_g^{-1}\Phi(p)1_g$. The strict 2-group structure in $C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$ is given by pointwise multiplication.

The 2-group isomorphism between $\mathcal{G}(\mathbb{E})$ and $C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$ is given by

$$F(x_i) = x_i \sigma(x_i), \forall x_i \in E_i, i \in \{0, 1\}$$

$$\Psi(p) = 1_p \Phi(p), \forall p \in E_0.$$
(6.1)

Let (G, H, τ, α) be the Lie crossed module associated to the Lie 2-group $\mathbb{G} = [G_1 \rightrightarrows G_0]$. Then a functor $\sigma \in C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$ is of the form $\sigma(p) \in G$ and $\sigma(\tilde{\gamma}) = (\sigma_{\tilde{\gamma}}, g_{\tilde{\gamma}}) \in H \rtimes_{\alpha} G$. The functoriality of σ imposes following conditions on $g_{\tilde{\gamma}} \in G, \sigma_{\tilde{\gamma}} \in H$,

$$g_{\widetilde{\gamma}} = \sigma_{s(\widetilde{\gamma})},$$

$$\sigma_{t(\widetilde{\gamma})} = \tau(\sigma_{\widetilde{\gamma}})\sigma_{s(\widetilde{\gamma})},$$

$$\sigma_{\widetilde{\gamma}_{2}}\sigma_{\widetilde{\gamma}_{1}} = \sigma_{\widetilde{\gamma}_{2}\circ\widetilde{\gamma}_{1}}.$$
(6.2)

A straightforward but tedious computation, using Equation (2.7), gives us the following \mathbb{G} -equivariancy condition on $(\sigma_{\tilde{\gamma}}, \sigma_p)$

$$\sigma_{pg} = \operatorname{Ad}_{g^{-1}}(\sigma_p)$$

$$\sigma_{\widetilde{\gamma}(h,g)} = \alpha_{g^{-1}} \left(\operatorname{Ad}_{h^{-1}}(\sigma_{\widetilde{\gamma}}) \right).$$
(6.3)

Now suppose $\sigma, \sigma' \colon \mathbb{E} \to \mathbb{G}$ are a pair of objects in $C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$ and $\tilde{\Phi} \colon \sigma \Longrightarrow \sigma'$ an arrow in $C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$. Then $\tilde{\Phi}(p) \colon \sigma(p) \to \sigma'(p)$ is of the form (Φ_p, σ_p) , where $\Phi(p) \in H$ satisfies the following conditions,

$$\sigma'(p) = \tau(\Phi(p)) \sigma(p),$$

$$\sigma'_{\tilde{\gamma}} = \Phi_q \, \sigma_{\tilde{\gamma}} \, \Phi_p^{-1},$$

$$\Phi_{pg} = \alpha_{g^{-1}}(\Phi_p),$$

(6.4)

for all $p \xrightarrow{\widetilde{\gamma}} q \in E_1, g \in G$.

Though our definition of gauge transformation is a straightforward categorification of the traditional definition, as we will see, it offers an enrichment that does not have a traditional counterpart.

Let $\pi : \mathbb{E} \to \mathbb{X}$ be a $\mathbb{G} = [G_1 \rightrightarrows G_0]$ -bundle over a Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$, which admits a categorical connection. Let $F : \mathbb{E} \to \mathbb{E}$ be a gauge transformation. Observe that F acts on the set of categorical connections by

$$(\mathcal{C} \cdot F)(\gamma, p) := F(\mathcal{C}(\gamma, F^{-1}(p))), \qquad (6.5)$$

where \mathcal{C} is a categorical connection and $(\gamma, p) \in s^* E_0$. The corresponding transformation of the action reads as $\mu(\gamma, p) \mapsto \mu_F(\gamma, p) := F(\mu(\gamma, F^{-1}(p)))$.

Choose a categorical connection \mathcal{C} to realize the \mathbb{G} -bundle as a decorated bundle $\mathbb{E}^{\text{dec}} = [s^*E_0 \times H \Rightarrow E_0] \Rightarrow \mathbb{X}$ over \mathbb{X} , as described in Section 3.3. Let $\overline{\mathbb{E}}^{\text{dec}}$ be the decorated bundle corresponding to the action μ_F . Note that $\overline{\mathbb{E}}^{\text{dec}}$ and \mathbb{E}^{dec} both have same set of objects and morphisms.

Proposition 6.1. Let $\theta_{\mathcal{C}\cdot F} \colon \overline{\mathbb{E}}^{\operatorname{dec}} \to \mathbb{E}$ be the map defined in Lemma 3.16. Let $\sigma \colon \mathbb{E} \to \mathbb{G}$ be a gauge transformation. Then $\sigma \circ \theta_{\mathcal{C}\cdot F} \colon \overline{\mathbb{E}}^{\operatorname{dec}} \to \mathbb{G}$ is a gauge transformation on $\overline{\mathbb{E}}^{\operatorname{dec}}$.

Proof. We verify $\sigma \circ \theta_{\mathcal{C}\cdot F}$ is a functor $\overline{\mathbb{E}}^{dec} \to \mathbb{G}$ satisfying conditions in Equation (6.3). \Box

We can further simplify the calculations for the gauge transformation on a decorated bundle as follows. Note that $((\gamma, p), h) = ((\gamma, p), e) (h^{-1}, e)$. Therefore if $F : \mathbb{E}^{dec} \to \mathbb{E}^{dec}$ is a gauge transformation, then $F(\gamma, p, h) = F(\gamma, p, e)(h^{-1}, e)$. In turn the gauge transformation $F : \mathbb{E}^{dec} \to \mathbb{E}^{dec}$ is completely determined by a smooth functor $F_0 : s^*E_0 \to s^*E_0 \times H$ satisfying $F_0(pg) = F_0(p)g, F_0(\gamma, pg) = F_0(\gamma, p) \mathbf{1}_g$. Consider the inclusion $1: [G \rightrightarrows G] \to [H \rtimes G \rightrightarrows G]$. Then, $(F_0, 1)$ is a map of principal 2-bundles $s^*E_0 \to \mathbb{E}^{dec}$ from the $[G \rightrightarrows G]$ -bundle $s^*E_0 \to \mathbb{X}$ to to the $[H \rtimes G \rightrightarrows G]$ -bundle $\mathbb{E}^{dec} \to \mathbb{X}$.

The association is given as $F(p) = F_0(p)$ and $F(\gamma, p, h) = F_0(\gamma, p)(h^{-1}, e)$. Equivalently if $\sigma \in C^{\infty}(\mathbb{E}^{dec}, \mathbb{G})^{\mathbb{G}}$ corresponds to $F \colon \mathbb{E}^{dec} \to \mathbb{E}^{dec}$, then σ is completely determined by a smooth functor $\sigma_0 \colon s^*E_0 \to \mathbb{G}$ satisfying $\sigma_0(pg) = g^{-1}\sigma_0(p)g, \sigma_0(\gamma, pg) = 1_g^{-1}\sigma_0(\gamma, p)1_g$, and then σ is given as $\sigma(p) = \sigma_0(p)$ and $\sigma((\gamma, p), h) = (h, e) \sigma_0(\gamma, p) (h^{-1}, e)$.

Proposition 6.2. Let $\mathbb{E}^{\text{dec}} = [s^* E_0 \times H \rightrightarrows E_0]$ be the decorated $(G, H, \tau, \alpha) = \mathbb{G}$ bundle associated to a *G*-bundle $(E_0 \to X_0 \mu, \mathbb{X})$ over the Lie groupoid $\mathbb{X} = [X_1 \rightrightarrows X_0]$. Then a gauge transformation $\sigma \colon \mathbb{E}^{\text{dec}} \to \mathbb{G}$ is completely determined by a smooth functor $\sigma_0 \colon s^* E_0 \to \mathbb{G}$ satisfying $\sigma_0(pg) = g^{-1}\sigma_0(p)g, \sigma_0(\gamma, pg) = 1_g^{-1}\sigma_0(\gamma, p)1_g$, and σ is given as

$$\sigma(p) = \sigma_0(p),$$

$$\sigma((\gamma, p), h) = (h, e) \sigma_0(\gamma, p) (h^{-1}, e).$$
(6.6)

Equivalently it is completely determined by a map of principal 2-bundles $(F_0, 1): s^*E_0 \to \mathbb{E}^{\text{dec}}$. The association is given as $F(p) = F_0(p)$ and $F((\gamma, p), h) = F_0(\gamma, p)(h^{-1}, e)$.

Example 6.3. Any gauge transformation on a principal *G*-bundle $E \to M$ defines a gauge transformation on the $[G \rightrightarrows G]$ -bundle $[E \rightrightarrows E]$ over $[X \rightrightarrows X]$ and vice versa.

Example 6.4. Recall that a principal $[G \rightrightarrows G]$ -bundle $[E_1 \rightrightarrows E_0] \rightarrow [X_1 \rightrightarrows X_0]$ is same as the pullback bundle $[s^*E_0 \rightrightarrows E_0] \rightarrow [X_1 \rightrightarrows X_0]$ defined with respect to an action $\mu: s^*E_0 \rightarrow E_0$ (Example 3.7). Then, any gauge transformation on a principal $[G \rightrightarrows G]$ bundle $[E_1 \rightrightarrows E_0] \rightarrow [X_1 \rightrightarrows X_0]$ is given by a gauge transformation $F: E_0 \rightarrow E_0$ on the principal G-bundle $E_0 \rightarrow X_0$ such that $F(\mu(\gamma, p)) = \mu(\gamma, F(p))$.

Example 6.5. Let $(G, H, \tau, \alpha) = \mathbb{G}$ be a Lie 2-group and $E \to X$ a principal *G*-bundle. Consider the decorated Lie 2-group (G, H, τ, α) -bundle $E^{\text{dec}} := [E \times H \rightrightarrows E]$ over $[X \rightrightarrows X]$ constructed in Example 3.13. Any gauge transformation $F \colon E \to E$ on the principal *G*-bundle $E \to X$ defines a gauge transformation on the (G, H, τ, α) -bundle $E^{\text{dec}} \to E^{\text{dec}}$ by $p \mapsto F(p)$ and $(p, h) \mapsto (F(p), h)$, for $p \in E, h \in H$. **Example 6.6.** Consider the $[H \rightrightarrows e]$ -bundle $[E_1 \rightrightarrows X_0]$ over $[X_1 \rightrightarrows X_0]$ of Example 3.8. Then a gauge transformation $F: E_1 \rightarrow E_1$ on the (traditional) principal *H*-bundle $E_1 \rightarrow X_1$ defines a gauge transformation on $[H \rightrightarrows e]$ -bundle $[E_1 \rightrightarrows X_0]$ over $[X_1 \rightrightarrows X_0]$ by $x \mapsto x$ and $\tilde{\gamma} \mapsto F(\tilde{\gamma})$, for $x \in X_0, \tilde{\gamma} \in E_1$.

6.1. Action of Gauge 2-group on the category of Connections. Let \mathbb{G} be a Lie 2-group, $\pi \colon \mathbb{E} \to \mathbb{X}$ a \mathbb{G} -bundle over \mathbb{X} and $\mathcal{G}(\mathbb{E})$ its gauge 2-group. Let $\Omega_{\mathbb{E}}^{\text{semi}}$ and $\Omega_{\mathbb{E}}^{\text{strict}}$ respectively be the categories of strict and semi-strict connection 1-forms. Let $F \colon \mathbb{E} \to \mathbb{E}$ be a gauge transformation. Then by taking the differentials of F we obtain the \mathbb{G} -equivariant Lie groupoid isomorphisms $F_* \colon T\mathbb{E} \to T\mathbb{E}$

$$(p,v) \mapsto F_*(p,v) := (p, F_{*,p}(v)),$$

$$(\widetilde{\gamma}, \widetilde{X}) \mapsto F_*(\widetilde{\gamma}, \widetilde{X}) := (\widetilde{\gamma}, F_{*,\widetilde{\gamma}}(\widetilde{X}))$$

for any $(p, v) \in TE_0, (\tilde{\gamma}, \tilde{X}) \in TE_1$. Similarly if $\Psi \colon F \Longrightarrow F'$ is an arrow in the category $\mathcal{G}(\mathbb{E})$, then the differential of Ψ gives us a \mathbb{G} -equivariant natural isomorphism $\Psi_* \colon F_* \Longrightarrow F'_*$:

$$(p,v) \mapsto \Psi_*(p,v) := \left(1_p, \Psi_{*,p}(v)\right) \colon F_*(p,v) \longrightarrow F'_*(p,v).$$

$$(6.7)$$

Now given an object $\omega \in \Omega_{\mathbb{E}}^{\text{strict}}$ or $\Omega_{\mathbb{E}}^{\text{semi}}$ and an arrow $(\zeta \colon \omega \Longrightarrow \omega') \in \Omega_{\mathbb{E}}^{\text{strict}}$ or $\Omega_{\mathbb{E}}^{\text{semi}}$ we respectively define

$$(F,\omega) \mapsto \omega \circ F_*^{-1} \colon T\mathbb{E} \to L(\mathbb{G}),$$

$$(\Psi \colon F \Longrightarrow F', \zeta \colon \omega \Longrightarrow \omega') \mapsto \zeta \circ_H \Psi_*^{-1} \colon \omega \circ F_*^{-1} \Longrightarrow \omega \circ F_*^{-1},$$
(6.8)

where \circ_H denote the horizontal composition of natural transformations. We claim that Equation (6.8) defines an action of the Lie 2-group $\mathcal{G}(\mathbb{E})$ on $\Omega_{\mathbb{E}}^{\text{strict}}$ and $\Omega_{\mathbb{E}}^{\text{semi}}$. If $\omega \in \Omega_{\mathbb{E}}^{\text{strict}}$ then it is obvious that $\omega \circ F_*^{-1} \in \Omega_{\mathbb{E}}^{\text{strict}}$. Moreover \mathbb{G} -equivariance of $\zeta : \omega \Longrightarrow \omega'$ and $\Psi: F \Longrightarrow F'$ give the \mathbb{G} -equivariance of $\zeta \circ_H \Psi_*^{-1}$ for a morphism Ψ either in $\Omega_{\mathbb{E}}^{\text{strict}}$ or $\Omega_{\mathbb{E}}^{\text{semi}}$. Perhaps the only non-trivial part is to show that $\omega \circ F_*^{-1} \in \Omega_{\mathbb{E}}^{\text{semi}}$ for $\omega \in \Omega_{\mathbb{E}}^{\text{semi}}$. Precisely we have to show that if there exists a \mathbb{G} -equivariant natural isomorphism $\kappa^* : (\omega \circ F_*^{-1}) \circ \delta \Longrightarrow \text{pr}_2$, where δ is the vertical vector field generating functor for the action of \mathbb{G} on \mathbb{E} . To see this we proceed as follows

For $(p, B) \in E_0 \times L(G_0)$ we define $\kappa_* : (\omega \circ F_*^{-1}) \circ \delta \Longrightarrow \operatorname{pr}_2$ as $\kappa_*(p, B) = \kappa(F^{-1}(p), B)$. Note that since a gauge transformation preserves the vertical vector field, $F_{*,p}^{-1}(\delta_p(B)) = \delta_{F^{-1}(p)}(B)$, we have $s(\kappa_*(p, B)) = s(\kappa(F^{-1}(p), B)) = \omega(\delta_{F^{-1}(p)}(B)) = (\omega \circ F_*^{-1}) \circ \delta(p, B)$ and $t(\kappa_*(p, B)) = t(\kappa(F^{-1}(p), B)) = B$. Thus

$$\kappa_*(p,B): (\omega \circ F_*^{-1}) \circ \delta(p,B) \to B.$$

Moreover since $\kappa \colon \omega \circ \delta \Longrightarrow \operatorname{pr}_2$ is a natural transformation, for any $(p, B) \xrightarrow{(\tilde{\gamma}, K)} (q, B') \in E_1 \times L(G_1)$, we have

$$K \circ \kappa(F^{-1}(p), B) = \kappa(F^{-1}(q), B') \circ \omega(F^{-1}(\gamma), \delta_{F^{-1}(\gamma)}(K)).$$
(6.9)

Substituting $\delta_{F^{-1}(\tilde{\gamma})}(K)$ by $F_{*,\tilde{\gamma}}^{-1}(\delta_{\tilde{\gamma}}(K))$ we confirm that κ_* is indeed a natural transformation. Now we verify G-equivariancy of κ_* ,

$$\kappa_*((p, B)g) = \kappa((F^{-1}(p), B)g) = \kappa(F^{-1}(p), B) \mathbf{1}_g = \kappa_*(p, B) \mathbf{1}_g$$

Proposition 6.7. Let $\pi \colon \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over \mathbb{X} and $\mathcal{G}(\mathbb{E})$ its gauge 2-group. Then Equation (6.8) defines an action of $\mathcal{G}(\mathbb{E})$ on $\Omega_{\mathbb{E}}^{\text{semi}}$. The action restricts to $\Omega_{\mathbb{E}}^{\text{strict}} \subset \Omega_{\mathbb{E}}^{\text{semi}}$.

If one represents F by $\sigma \in C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$, then the gauge transformations of a strict connection 1-form ω^{st} and a semi-strict connection 1-form ω^{se} are respectively expressed as

$$\begin{aligned} \omega^{\mathrm{st}} &\mapsto \mathrm{Ad}_{\sigma} \, \omega^{\mathrm{st}} - (d\sigma) \sigma^{-1}, \\ \omega^{\mathrm{se}} &\mapsto \mathrm{Ad}_{\sigma} \, \omega^{\mathrm{se}} - \widehat{\kappa}_{\omega^{\mathrm{se}}} \circ (d\sigma) \sigma^{-1} - (d\sigma) \sigma^{-1}, \end{aligned} \tag{6.10}$$

where $\widehat{\kappa}_{\omega^{se}} : \mathbb{E} \times L(\mathbb{G}) \longrightarrow L(\mathbb{G})^{\tau} = [L(H) \rtimes \tau(L(H))] \rightrightarrows \tau(L(H))]$ is as in Proposition 5.9. Note that here everything is defined point-wise.

Remark 6.8. Since $\Omega_{\mathbb{E}}^{\text{strict}}$ and $\Omega_{\mathbb{E}}^{\text{semi}}$ respectively are isomorphic to $C_{\mathbb{E}}^{\text{strict}}$ and $C_{\mathbb{E}}^{\text{semi}}$, the action in Proposition 6.7 induces actions of $\mathcal{G}(\mathbb{E})$ on $C_{\mathbb{E}}^{\text{strict}}$ and $C_{\mathbb{E}}^{\text{semi}}$. The induced action is given as $(R, F) \mapsto \overline{F} \circ R \circ \overline{TF^{-1}}$, where $\overline{F} : \operatorname{Ad}(\mathbb{E}) \to \operatorname{Ad}(\mathbb{E})$ sends $[x_i, A_i]$ to $[F(x_i), A_i]$ for any $[x_i, A_i] \in \operatorname{Ad}(E_i), i \in \{0, 1\}$ and $\overline{TF} : \operatorname{At}(\mathbb{E}) \to \operatorname{At}(\mathbb{E})$ sends an object $[x_i, v_i]$ to $[F(x_i), F_{*,x_i}(v_i)]$ for any $[x_i, v_i] \in \operatorname{At}(E_i), i \in \{0, 1\}$.

As with the connections, one can consider the set of categorical connected components $\overline{\mathcal{G}}(\mathbb{E})$ of the gauge 2-group $\mathcal{G}(\mathbb{E})$ for a \mathbb{G} -bundle $\pi \colon \mathbb{E} \to \mathbb{X}$. Then we have the following.

Corollary 6.9. Let $\pi \colon \mathbb{E} \to \mathbb{X}$ be a \mathbb{G} -bundle over \mathbb{X} and $\mathcal{G}(\mathbb{E})$ its connected components of the gauge 2-group. Then Equation (6.8) defines an action of $\overline{\mathcal{G}}(\mathbb{E})$ on $\overline{\Omega_{\mathbb{E}}}^{\text{semi}}$. The action restricts to $\overline{\Omega_{\mathbb{E}}}^{\text{strict}} \subset \overline{\Omega_{\mathbb{E}}}^{\text{semi}}$.

6.2. An Extended symmetry of $\Omega_{\mathbb{E}}^{\text{semi}}$. In fact, $\Omega_{\mathbb{E}}^{\text{semi}}$ enjoys a larger symmetry than given in Proposition 6.7. To see this, consider

$$\Omega^G(E_0, L(H)) = \{\lambda \colon TE_0 \to L(H) | \lambda_{p \cdot g}(v \cdot g) = \alpha_{g^{-1}}(\lambda_p(v))\},\$$

the *G*-equivariant L(H)-valued smooth 1-forms on E_0 . Observe that every such $\lambda \in \Omega^G(E_0, L(H))$ can be viewed as a \mathbb{G} -equivariant 1-form $\overline{\lambda} \colon T\mathbb{E} \to L(\mathbb{G})$ on \mathbb{E} taking values in $L(\mathbb{G})^{\tau} = [L(H) \rtimes \tau(L(H)) \rightrightarrows \tau(L(H))] \subset L(\mathbb{G}),$

$$\overline{\lambda} := \tau(\lambda) \colon TE_0 \to \tau(L(H)),$$

$$\overline{\lambda} := (t^*\lambda - s^*\lambda + (\tau(s^*\lambda))) \colon TE_1 \to L(H) \oplus \tau(L(H).$$
(6.11)

Let $\widehat{\Omega}^{\mathbb{G}}(E_0, L(H))$ be the category whose objects are functors of the form Equation (6.11) and morphisms are fiber-wise linear natural transformations.

Now, we intertwine the category $\widehat{\Omega}^{\mathbb{G}}(E_0, L(H))$ with the gauge 2-group $\mathcal{G}(\mathbb{E})$ to define a 2-group $\mathcal{G}^{\text{ext}} := \mathcal{G}(\mathbb{E}) \times \widehat{\Omega}^{\mathbb{G}}(E_0, L(H))$ with group products

$$(F,\lambda)(F',\lambda') = (FF',\lambda + (\lambda' \circ F_*^{-1})),$$

$$(\Psi,\theta)(\Psi',\theta') = (\Psi\Psi',\theta + (\theta' \circ_H \Psi_*^{-1})).$$
(6.12)

One has to apply Remark 2.6 to verify that \mathcal{G}^{ext} is a 2-group. Note that here we have made the identification $\text{Obj}(\mathcal{G}^{\text{ext}}) \simeq \text{Obj}(\mathcal{G}(\mathbb{E})) \times \Omega^G(E_0, L(H)).$

Then Proposition 5.9 allows us to define a left action of \mathcal{G}^{ext} on $\Omega_{\mathbb{R}}^{\text{semi}}$ by

$$((F,\lambda),\omega) \mapsto \omega \circ F_*^{-1} + \overline{\lambda} \colon T\mathbb{E} \to L(\mathbb{G}),$$

$$((\Psi,\theta),\zeta) \mapsto \zeta \circ_H \Psi_*^{-1} + \theta.$$

$$(6.13)$$

One can as well express the action above as

$$((\sigma,\lambda),\omega) \mapsto (\mathrm{Ad}_{\sigma}(\omega) - \kappa_{\omega} \circ (d\sigma)\sigma^{-1} - (d\sigma)\sigma^{-1} + \overline{\lambda}) \colon T\mathbb{E} \to L(\mathbb{G}), \tag{6.14}$$

where $\sigma \in C^{\infty}(\mathbb{E}, \mathbb{G})^{\mathbb{G}}$.

We call the action above an extended gauge transformation. It is obvious that this action does not restrict to $\Omega_{\mathbb{E}}^{\text{strict}}$ (see Corollary 5.10). Particularly, the action we derived in Equation (6.13) is of some importance in higher gauge theory and related areas of physics (for example 2-BF theories). We refer to the work of Martins and Miković [32], and works of Martins and Picken in this context [30, 33, 31]. It would be instructive to consider the following example.

Example 6.10. Consider the $\mathbb{G} = (G, H, \tau, \alpha)$ -bundle $E^{\text{dec}} := [E \times H \rightrightarrows E]$ over $[X \rightrightarrows X]$ defined by a traditional principal *G*-bundle $E \to X$ as in Example 3.13. A gauge transformation $\sigma \colon E^{\text{dec}} \to \mathbb{G}$ on $E^{\text{dec}} = [E \times H \rightrightarrows E]$ is given, by a gauge transformation $\sigma_0 \colon E \to G$ on the *G*-bundle $E \to X$, as $\sigma(p) = \sigma_0(p), \sigma(p, h) = (h, \sigma_0(p))$. Thus an extended gauge transformation (σ, λ) is given by a classical gauge transformation $\sigma_0 \colon E \to G$ on the *G*-bundle $E \to X$, as $\sigma(p) = \sigma_0(p), \sigma(p, h) = (h, \sigma_0(p))$. Thus an extended gauge transformation (σ, λ) is given by a classical gauge transformation $\sigma_0 \colon E \to G$ on the *G*-bundle $E \to X$ and a *G*-equivariant L(H)-valued 1-form $\lambda \in \Omega^G(E, L(H))$ on *E*. In Corollary 5.21 we have seen, any connection ω_0 on the principal *G*-bundle $E \to X$ gives a strict connection $\overline{\omega} := (\omega^{\text{dec}}, \omega_0)$. Under the action of (σ, λ) , the connection $\overline{\omega}$ transforms as:

$$\overline{\omega} \mapsto \operatorname{Ad}_{\sigma} \overline{\omega} - (d\sigma)\sigma^{-1} + \overline{\lambda}.$$

Which explicitly reads

$$\omega_{0}(p) \mapsto \operatorname{Ad}_{\sigma_{0}(p)} \omega_{0}(p) - (d\sigma_{0}(p))\sigma_{0}(p)^{-1} + \tau(\lambda(p)),$$

$$\omega^{\operatorname{dec}}(p, h) \mapsto \operatorname{Ad}_{(h, \sigma_{0}(p))} \omega^{\operatorname{dec}}(p, h) - \left(d(h, \sigma_{0}(p))\right)(h, \sigma_{0}(p))^{-1} + (\tau(\lambda(p)) - \lambda(p) + \operatorname{Ad}_{h} \lambda(p)).$$
(6.15)

The gauge transformation obtained here in Equation (6.15) is precisely the gauge transformation of a connection 1-form in the higher gauge theories (see [44] or [32]). Note that the extended gauge transformation is a direct consequence of obstruction to the splitting of the Atiyah sequence in Diagram 5.1, and global in nature. Of course, higher gauge theories also involve higher-order differential forms, which we have not dealt with here. In our follow-up paper, we will construct the Chern-Weil theory for the framework developed here, which is expected to make the relation with higher gauge theories more transparent.

ACKNOWLEDGEMENTS

The first named author acknowledges research support from SERB, DST, Government of India grant MTR/2018/000528.

References

- Paolo Aschieri, Luigi Cantini, and Branislav Jurčo. Nonabelian bundle gerbes, their differential geometry and gauge theory. Comm. Math. Phys., 254(2):367–400, 2005.
- [2] M. F. Atiyah. Complex analytic connections in fibre bundles. Trans. Amer. Math. Soc., 85:181–207, 1957.
- [3] John Baez and Urs Schreiber. Higher gauge theory: 2-connections on 2-bundles. arXiv:hepth/0412325, 2004.
- [4] John C. Baez and Alissa S. Crans. Higher-dimensional algebra. VI. Lie 2-algebras. Theory Appl. Categ., 12:492–538, 2004.
- [5] John C. Baez and Aaron D. Lauda. Higher-dimensional algebra. V. 2-groups. Theory Appl. Categ., 12:423–491, 2004.
- [6] John C. Baez and Urs Schreiber. Higher gauge theory. In Categories in algebra, geometry and mathematical physics, volume 431 of Contemp. Math., pages 7–30. Amer. Math. Soc., Providence, RI, 2007.
- [7] Igor Bakovic. The simplicial interpretation of bigroupoid 2-torsors. arXiv:0902.3436.
- [8] Igor Baković. Bigroupoid 2-torsors. PhD thesis, Ludwig Maximilian University of Munich, 2008.
- [9] Tobias Keith Bartels. Higher gauge theory: 2-bundles. ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)–University of California, Riverside.
- [10] Indranil Biswas, Saikat Chatterjee, Praphulla Koushik, and Frank Neumann. Chern-Weil theory for principal bundles over Lie groupoids. arXiv:2012.08447, 2020.
- [11] Indranil Biswas and Frank Neumann. Atiyah sequences, connections and characteristic forms for principal bundles over groupoids and stacks. C. R. Math. Acad. Sci. Paris, 352(1):59–64, 2014.
- [12] Francis Borceux. Handbook of categorical algebra. 1, volume 50 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 1994. Basic category theory.
- [13] Lawrence Breen and William Messing. Differential geometry of gerbes. Adv. Math., 198(2):732–846, 2005.
- [14] Henrique Bursztyn, Alejandro Cabrera, and Matias del Hoyo. Vector bundles over lie groupoids and algebroids. Advances in Mathematics, 290:163–207, 2016.
- [15] Henrique Bursztyn, Alejandro Cabrera, and Cristián Ortiz. Linear and multiplicative 2-forms. Letters in Mathematical Physics, 90(1-3):59, 2009.
- [16] Francesco Cattafi. Cartan geometries and multiplicative forms. Differential Geometry and its Applications, 75:101722, 2021.
- [17] Saikat Chatterjee, Amitabha Lahiri, and Ambar N. Sengupta. Path space connections and categorical geometry. J. Geom. Phys., 75:129–161, 2014.
- [18] Saikat Chatterjee, Amitabha Lahiri, and Ambar N. Sengupta. Twisted actions of categorical groups. Theory Appl. Categ., 29:No. 8, 215–255, 2014.
- [19] Saikat Chatterjee, Amitabha Lahiri, and Ambar N. Sengupta. Construction of categorical bundles from local data. *Theory Appl. Categ.*, 31:Paper No. 14, 388–417, 2016.
- [20] Domenico Fiorenza, Urs Schreiber, and Jim Stasheff. Cech cocycles for differential characteristic classes: an infinity-Lie theoretic construction. Adv. Theor. Math. Phys, 16(1):149–250, 2012.
- [21] Grégory Ginot and Mathieu Stiénon. G-gerbes, principal 2-group bundles and characteristic classes. J. Symplectic Geom., 13(4):1001–1047, 2015.
- [22] Alfonso Gracia-Saz and Rajan Amit Mehta. Vb-groupoids and representation theory of Lie groupoids. J. Symplectic Geom., 15(3):741–783, 2017.
- [23] Branislav Jurčo, Christian Sämann, and Martin Wolf. Semistrict higher gauge theory. J. High Energy Phys., (4):087, front matter+66, 2015.
- [24] Branislav Jurčo, Christian Sämann, and Martin Wolf. Higher groupoid bundles, higher spaces, and self-dual tensor field equations. *Fortschr. Phys.*, 64(8-9):674–717, 2016.
- [25] Shôshichi Kobayashi. On connections of cartan. Canadian Journal of Mathematics, 8:145–156, 1956.
- [26] Camille Laurent-Gengoux, Mathieu Stiénon, and Ping Xu. Non-abelian differentiable gerbes. Adv. Math., 220(5):1357–1427, 2009.
- [27] Camille Laurent-Gengoux, Jean-Louis Tu, and Ping Xu. Chern-Weil map for principal bundles over groupoids. Math. Z., 255(3):451–491, 2007.

- [28] Kirill C. H. Mackenzie. Lie groupoids and Lie algebroids in differential geometry, volume 124 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1987.
- [29] Kirill C. H. Mackenzie, Anatol Odzijewicz, and Aneta Sliżewska. Poisson geometry related to Atiyah sequences. SIGMA Symmetry Integrability Geom. Methods Appl., 14:Paper No. 005, 29, 2018.
- [30] João Faria Martins and Roger Picken. On two-dimensional holonomy. Trans. Amer. Math. Soc., 362(11):5657–5695, 2010.
- [31] João Faria Martins and Roger Picken. Surface holonomy for non-abelian 2-bundles via double groupoids. Adv. Math., 226(4):3309–3366, 2011.
- [32] João Martins, Aleksandar Miković, et al. Lie crossed modules and gauge-invariant actions for 2-bf theories. Advances in Theoretical and Mathematical Physics, 15(4):1059–1084, 2011.
- [33] Joao Faria Martins and Roger Picken. The fundamental gray 3-groupoid of a smooth manifold and local 3-dimensional holonomy based on a 2-crossed module. *Differential Geometry and its Applica*tions, 29(2):179–206, 2011.
- [34] Ieke Moerdijk. Introduction to the language of stacks and gerbes. arXiv:math/0212266, 2002.
- [35] Thomas Nikolaus, Urs Schreiber, and Danny Stevenson. Principal ∞-bundles: general theory. J. Homotopy Relat. Struct., 10(4):749–801, 2015.
- [36] Thomas Nikolaus, Urs Schreiber, and Danny Stevenson. Principal ∞-bundles: presentations. J. Homotopy Relat. Struct., 10(3):565–622, 2015.
- [37] Thomas Nikolaus and Konrad Waldorf. Four equivalent versions of nonabelian gerbes. Pacific J. Math., 264(2):355–419, 2013.
- [38] Christopher J. Schommer-Pries. Central extensions of smooth 2-groups and a finite-dimensional string 2-group. *Geom. Topol.*, 15(2):609–676, 2011.
- [39] Urs Schreiber. From loop space mechanics to nonabelian strings. arXiv:hep-th/0509163, 2005.
- [40] Urs Schreiber. Differential cohomology in a cohesive infinity-topos. arXiv:1310.7930, 2013.
- [41] Jan Slovák and Andreas Cap. Parabolic Geometries I, Background and General Theory, volume 1000. American Mathematical Society, 2009.
- [42] Danny Stevenson. Lie 2-algebras and the geometry of gerbes. available online as https://math.ucr.edu/home/baez/namboodiri/stevenson maclane.pdf.
- [43] Konrad Waldorf. A global perspective to connections on principal 2-bundles. Forum Math., 30(4):809–843, 2018.
- [44] Wei Wang. On the global 2-holonomy for a 2-connection on a 2-bundle. J. Geom. Phys., 117:151–178, 2017.
- [45] Christoph Wockel. Principal 2-bundles and their gauge 2-groups. Forum Math., 23(3):565–610, 2011.

School of Mathematics, Indian Institute of Science Education and Research Thiruvananthapuram, Maruthamala P.O., Vithura, Kerala 695551, India

Email address: saikat.chat01@gmail.com,adittyachaudhuri15@iisertvm.ac.in

School of Mathematics, IISER Pune, Dr. Homi Bhabha Road, Pashan, Pune, 411008, India

Email address: koushik16@iisertvm.ac.in, praphullakowshik@gmail.com