

Summability of the norms of differences of operator averages on a Hilbert space

Sakin Demir

Agri Ibrahim Cecen University

Faculty of Education

Department of Basic Education

04100 Ağrı, Turkey

E-mail: sakin.demir@gmail.com

July 30, 2021

Abstract

Let H be a Hilbert space and $U : H \rightarrow H$ be a unitary operator. Let

$$A_n f = \frac{1}{n} \sum_{j=1}^n U^j f$$

for $f \in H$. Suppose that (n_k) is a lacunary sequence, then there exists a constant $C > 0$ such that

$$\sum_{k=1}^{\infty} \|A_{n_{k+1}} f - A_{n_k} f\|_H \leq C \|f\|_H$$

for all $f \in H$.

Let T be a contraction on a Hilbert space H . Let (n_k) be lacunary sequence. Let

$$A_n(T)f = \frac{1}{n} \sum_{j=1}^n T^j f$$

2020 *Mathematics Subject Classification*: Primary 47B02; Secondary 37A30.

Key words and phrases: Hilbert Space, Unitary Operator, Contraction, Average.

for all $f \in H$. Then there exists a constant $C > 0$ such that

$$\sum_{k=1}^{\infty} \|A_{n_{k+1}}(T)f - A_{n_k}(T)f\|_H \leq C\|f\|_H$$

for all $f \in H$.

1 Introduction

Let H be a Hilbert space and $U : H \rightarrow H$ be a unitary operator. Define the average

$$A_n f = \frac{1}{n} \sum_{j=1}^n U^j f$$

for $f \in H$. When (n_k) is an increasing sequence it has been proved in [1] in 1996 that

$$\left(\sum_{k=1}^{\infty} \|A_{n_{k+1}} f - A_{n_k} f\|_H^2 \right)^{1/2} \leq 25\|f\|_H$$

for all $f \in H$ (see Theorem 1.2 in [1]).

Let T be a contraction on a Hilbert space H and (n_k) be an increasing sequence of positive integers. Let

$$A_n(T)f = \frac{1}{n} \sum_{j=1}^n T^j f$$

for all $f \in H$. Then it has also proven in [1] that

$$\left(\sum_{k=1}^{\infty} \|A_{n_{k+1}}(T)f - A_{n_k}(T)f\|_H^2 \right)^{1/2} \leq 25\|f\|_H$$

for all $f \in H$ (see Theorem 1.7 in [1]).

Since then it has remained as an open question if one can have the above inequalities when the square of the Hilbert space norms in the summation is removed. It is the main purpose of this work to answer this question. We prove that it is indeed the case with only an additional assumption of lacunarity on the sequence (n_k) .

Recall that a sequence (n_k) of real numbers is called lacunary if there exists a constant $\beta > 1$ such that

$$\frac{n_{k+1}}{n_k} \geq \beta$$

for all $k = 1, 2, 3, \dots$

2 Main Results

Our first result is the following:

Theorem 1. *Let (n_k) be a lacunary sequence and U be a unitary operator on a Hilbert space H . Then there exists a constant $C > 0$ such that*

$$\sum_{k=1}^{\infty} \|A_{n_{k+1}}f - A_{n_k}f\|_H \leq C\|f\|_H$$

for all $f \in H$.

Proof. It suffices to show that

$$\sum_{k=1}^N \|A_{n_{k+1}}f - A_{n_k}f\|_H \leq C\|f\|_H$$

for all $N \geq 1$. Let us recall the circle

$$\mathbb{T} = \{\gamma \in \mathbb{C} : |\gamma| = 1\}$$

and let

$$a_n(\gamma) = \frac{1}{n} \sum_{j=1}^n \gamma^j.$$

By the spectral theorem for unitary operators, it suffices to show the existence of a constant $C > 0$ such that

$$\sum_{k=1}^N |a_{n_{k+1}}(\gamma) - a_{n_k}(\gamma)| \leq C$$

for all $\gamma \in \mathbb{T}$.

Since $a_n(1) = 1$ for all $n \leq 1$, we can assume that $\gamma \neq 1$. Write $\gamma = e^{i\theta}$ where $\theta \in (0, 2\pi)$. If $\theta \in [\pi, 2\pi)$, then $\bar{\gamma} = e^{i\theta}$ where $\theta = 2\pi - \theta$ is in $(0, \pi]$. Since

$$|a_{n_{k+1}}(\gamma) - a_{n_k}(\gamma)| = |a_{n_{k+1}}(\bar{\gamma}) - a_{n_k}(\bar{\gamma})|$$

we may assume without loss of generality that $\theta \in (0, \pi]$.

Since $|e^{i\theta} - 1| \geq \frac{1}{4}\theta$ for all $\theta \in (0, \pi]$ we have

$$\begin{aligned} |a_{n_{k+1}}(\gamma) - a_{n_k}(\gamma)| &= \left| \frac{1}{n_{k+1}} \left(\frac{e^{in_{k+1}\theta} - 1}{e^{i\theta} - 1} \right) - \frac{1}{n_k} \left(\frac{e^{in_k\theta} - 1}{e^{i\theta} - 1} \right) \right| \\ &\leq 16 \left| \frac{e^{in_{k+1}\theta} - 1}{n_{k+1}\theta} - \frac{e^{in_k\theta} - 1}{n_k\theta} \right|. \end{aligned}$$

Let

$$I = \sum_{k=1}^{\infty} \left| \frac{e^{in_{k+1}\theta} - 1}{n_{k+1}\theta} - \frac{e^{in_k\theta} - 1}{n_k\theta} \right|.$$

First fix θ and let k_0 be the first k such that $\theta n_k \geq 1$, and let

$$\begin{aligned} I_1 &= \sum_{k:\theta n_k < 1} \left| \frac{e^{in_{k+1}\theta} - 1}{n_{k+1}\theta} - \frac{e^{in_k\theta} - 1}{n_k\theta} \right| \\ I_2 &= \sum_{k:\theta n_k \geq 1} \left| \frac{e^{in_{k+1}\theta} - 1}{n_{k+1}\theta} - \frac{e^{in_k\theta} - 1}{n_k\theta} \right|. \end{aligned}$$

We thus have $I = I_1 + I_2$, and to complete our theorem it suffices to control I_1 and I_2 .

To control I_1 first define a function $F: \mathbb{R} \rightarrow \mathbb{C}$ as

$$F(r) = \frac{e^{ir} - 1}{r}.$$

Then we have

$$F(n\theta) = \frac{e^{in\theta} - 1}{n\theta}.$$

By the Mean Value Theorem there exists a constant $\xi \in (n_k\theta, n_{k+1}\theta)$ such that

$$|F(n_{k+1}\theta) - F(n_k\theta)| = |F'(\xi)| |n_{k+1}\theta - n_k\theta|.$$

Also, it is easy to verify that

$$|F'(x)| \leq \frac{x+1}{x^2}$$

for $x > 0$.

Now we have

$$\begin{aligned} |F(n_{k+1}\theta) - F(n_k\theta)| &= |F'(\xi)| |n_{k+1}\theta - n_k\theta| \\ &\leq \frac{\xi+1}{\xi^2} \theta (n_{k+1} - n_k) \\ &\leq \frac{n_{k+1}\theta + 1}{\theta^2 n_k^2} \theta (n_{k+1} - n_k) \\ &= \frac{2n_{k+1}}{n_k^2} (n_{k+1} - n_k) \end{aligned}$$

Thus we have

$$\begin{aligned} I_1 &= \sum_{k:n_k\theta < 1} |F(n_{k+1}\theta) - F(n_k\theta)| \\ &\leq \sum_{k:n_k\theta < 1} \frac{2}{n_k\theta} \cdot \frac{2n_{k+1}}{n_k^2} (n_{k+1} - n_k) \\ &\leq \sum_{k:n_k\theta < 1} \frac{4n_{k+1}^2}{n_k^2\theta} \left(\frac{1}{n_k} - \frac{1}{n_{k+1}} \right) \\ &= \sum_{k:n_k\theta < 1} \frac{16}{\theta} \left(\frac{1}{n_k} - \frac{1}{n_{k+1}} \right) \\ &= \frac{16}{\theta} \left(\frac{1}{n_1} - \frac{1}{n_{k_0+1}} \right) \\ &\leq \frac{16}{n_{k_0+1}\theta} \\ &\leq 16. \end{aligned}$$

This shows that I_1 is bounded.

Since

$$\begin{aligned} I_2 &= \sum_{k:\theta n_k \geq 1} \left| \frac{e^{in_{k+1}\theta} - 1}{n_{k+1}\theta} - \frac{e^{in_k\theta} - 1}{n_k\theta} \right| \\ &\leq 2 \sum_{k:\theta n_k \geq 1} \frac{1}{n_{k+1}\theta} + 2 \sum_{k:\theta n_k \geq 1} \frac{1}{n_k\theta}. \end{aligned}$$

It suffices to control the second summation

$$2 \sum_{k:\theta n_k \geq 1} \frac{1}{n_k \theta}$$

since the first summation can be controlled similarly.

Since

$$\frac{n_{k_0}}{n_k} = \frac{n_{k_0}}{n_{k_0+1}} \cdot \frac{n_{k_0+1}}{n_{k_0+2}} \cdot \frac{n_{k_0+2}}{n_{k_0+3}} \cdots \frac{n_{k-1}}{n_k}$$

and (n_k) is lacunary we have

$$\frac{n_{k_0}}{n_k} \leq \frac{1}{\beta^{k-k_0}}.$$

Since $\frac{1}{\theta n_{k_0}} \leq 1$ we obtain

$$\begin{aligned} 2 \sum_{k:\theta n_k \geq 1} \frac{1}{n_k \theta} &= 2 \frac{1}{\theta n_{k_0}} \sum_{k:\theta n_k \geq 1} \frac{n_{k_0}}{n_k} \\ &\leq 2 \sum_{k:\theta n_k \geq 1} \frac{n_{k_0}}{n_k} \\ &\leq 2 \sum_{k:\theta n_k \geq 1} \frac{1}{\beta^{k-k_0}} \\ &\leq 2 \frac{1}{1-\beta} \end{aligned}$$

and this finishes our proof. □

The following is our second result:

Theorem 2. *Let T be a contraction on a Hilbert space H . Let (n_k) be lacunary sequence. Let*

$$A_n(T)f = \frac{1}{n} \sum_{j=1}^n T^j f$$

for all $f \in H$. Then there exists a constant $C > 0$ such that

$$\sum_{k=1}^{\infty} \|A_{n_{k+1}}(T)f - A_{n_k}(T)f\|_H \leq C \|f\|_H$$

for all $f \in H$.

Proof. We proceed with a classical argument as in the proof of Theorem 1.7 in [1]. By the dilation theorem (see [2]), there exists a Hilbert space L containing H as a closed subspace, an orthogonal projection $P : L \rightarrow H$, and a unitary operator $U : L \rightarrow L$ with $PU^j f = T^j f$ for all $j \geq 0$ and $f \in H$. But then for $f \in H$,

$$\begin{aligned} \sum_{k=1}^N \|A_{n_{k+1}}(T)f - A_{n_k}(T)f\|_H &= \sum_{k=1}^N \|P(A_{n_{k+1}}(U)f - A_{n_k}(U)f)\|_H \\ &= \|P\| \sum_{k=1}^N \|A_{n_{k+1}}(U)f - A_{n_k}(U)f\|_H \\ &\leq C \end{aligned}$$

by Theorem 1 and the fact that $\|P\| \leq 1$. □

References

- [1] R. L. Jones, I. V. Ostrovskii and J. M. Rosenblatt, *Square functions in ergodic theory*, Ergod. Th. & Dynam. Sys. 16 (1996) 267-305.
- [2] B. Sz-Nagy and C. Foias, *Analyse harmonique des operateurs de l'espace de Hilbert*, Akad. Kiado, Budapest. Mason, Paris, 1967.