-Lie-type maps on C^ -algebras

Ruth Nascimento Ferreira

Federal University of Technology, Avenida Professora Laura Pacheco Bastos, 800, 85053-510, Guarapuava, Brazil. e-mail: ruthferreira@utfpr.edu.br

Bruno Leonardo Macedo Ferreira

Federal University of Technology, Avenida Professora Laura Pacheco Bastos, 800, 85053-510, Guarapuava, Brazil. e-mail: brunolmfalg@gmail.com

Henrique Guzzo Junior

University of São Paulo, Rua do Matão, 1010, 05508-090, São Paulo, Brazil. e-mail: guzzo@ime.usp.br

Bruno Tadeu Costa

Federal University of Santa Catarina, Rua João Pessoa, 2750, 89036-256, Blumenau, Brazil. e-mail: b.t.costa@ufsc.br

Abstract

Let \mathfrak{A} and \mathfrak{A}' be two C^* -algebras with identities $I_{\mathfrak{A}}$ and $I_{\mathfrak{A}'}$, respectively, and P_1 and $P_2 = I_{\mathfrak{A}} - P_1$ nontrivial symmetric projections in \mathfrak{A} . In this paper we study the characterization of multiplicative *-Lie-type maps. In particular, if \mathcal{M} is a factor von Neumann algebra then every complex scalar multiplication bijective unital multiplicative *-Lie-type map is *-isomorphism.

AMS 2010 Subject Classification: 47B48, 46L05. Keywords: C*-algebra, factor von Neumann algebras, multiplicative *-Lietype maps, *-isomorphism

1 Introduction and Preliminaries

The study of additivity of maps strongly attracts attention of mathematicians, so that the first quite surprising result is due to Martindale who imposed conditions on a ring such that multiplicative bijective maps are all additive [17]. Thenceforth, diverse works have been published considering different types of associative and non-associative algebras. Among them we can mention [9, 11, 12, 8, 13, 10, 14, 3]. In order to further develop the study of additivity of maps, the researches incorporated two new product into the theory, presented by Brešar and Fošner in [2, 15], where the definition is as follows: for $X, Y \in \mathfrak{R}$, where \mathfrak{R} is a *-ring, we denote by $X \bullet Y = XY + YX^*$ and $[X, Y]_* = XY - YX^*$ the *-Jordan product and the *-Lie product, respectively. In [4], the authors proved that a map Φ between two factor von Newmann algebras is a *-ring isomorphism if and only if $\Phi([X,Y]_*) = [\Phi(X), \Phi(Y)]_*$. In [7], Ferreira and Costa extended these new products and defined two other types of applications, named multiplicative *-Lie n-map and multiplicative *-Jordan n-map and proved that, under certain conditions, an application between C^* -algebras that is multiplicative *-Lie n-map and multiplicative *-Jordan n-map is a *-ring isomorphism. In the second paper of this series [6], Ferreira and Costa prove when a multiplicative *-Jordan n-map is a *-ring isomorphism. As a consequence of their main result, they provide an application on von Neumann algebras, factor von Neumann algebras and prime algebras. Furthermore, they generalize the Main Theorem in [5]. With this picture in mind, in this article we will discuss when a multiplicative *-Lie n-map is a *-isomorphism and, just as it was done in [6], we provide an application on von Neumann algebras, factor von Neumann algebras and prime algebras.

Let us define the following sequence of polynomials, as presented in [7]:

$$p_{1_*}(x) = x$$
 and $p_{n_*}(x_1, x_2, \dots, x_n) = \left[p_{(n-1)_*}(x_1, x_2, \dots, x_{n-1}), x_n \right]_*$

for all integers $n \geq 2$. Thus, $p_{2_*}(x_1, x_2) = [x_1, x_2]_*$, $p_{3_*}(x_1, x_2, x_3) = [[x_1, x_2]_*, x_3]_*$, etc. Note that p_{2_*} is the product introduced by Brešar and Fošner [2, 15]. Then, using the nomenclature introduced in [7] we have a new class of maps (not necessarily additive): given two rings \mathfrak{R} and \mathfrak{R}' , $\varphi : \mathfrak{R} \longrightarrow \mathfrak{R}'$ is a *multiplicative* *-*Lie n*-*map* if

$$\varphi(p_{n_*}(x_1, x_2, ..., x_i, ..., x_n)) = p_{n_*}(\varphi(x_1), \varphi(x_2), ..., \varphi(x_i), ..., \varphi(x_n)),$$

where $n \ge 2$ is an integer. Multiplicative *-Lie 2-map, *-Lie 3-map and *-Lie *n*-map are collectively referred to as *multiplicative* *-Lie-type maps.

By a C^* -algebra we mean a complete normed complex algebra (say \mathfrak{A}) endowed with a conjugate-linear algebra involution *, satisfying $||a^*a|| =$ $||a||^2$ for all $a \in \mathfrak{A}$. Moreover, a C^* -algebra is a prime C^* -algebra if $A\mathfrak{A} B =$ 0 for $A, B \in \mathfrak{A}$ implies either A = 0 or B = 0.

We find it convenient to invoke the noted Gelfand-Naimark theorem that state a C^* -algebra \mathfrak{A} is *-isomorphic to a C^* -subalgebra $\mathfrak{D} \subset \mathcal{B}(\mathcal{H})$, where \mathcal{H} is a Hilbert space. So from now on we shall consider elements of a C^* -algebra as operators.

Let be P_1 a nontrivial projection in \mathfrak{A} and $P_2 = I_{\mathfrak{A}} - P_1$ where $I_{\mathfrak{A}}$ is the identity of \mathfrak{A} . Then \mathfrak{A} has a decomposition $\mathfrak{A} = \mathfrak{A}_{11} \oplus \mathfrak{A}_{12} \oplus \mathfrak{A}_{21} \oplus \mathfrak{A}_{22}$, where $\mathfrak{A}_{ij} = P_i \mathfrak{A} P_j$ (i, j = 1, 2).

2 Main theorem

In the following we shall prove a part of the main result of this paper.

Theorem 2.1. Let \mathfrak{A} and \mathfrak{A}' be two C^* -algebras with identities $I_{\mathfrak{A}}$ and $I_{\mathfrak{A}'}$, respectively, and P_1 and $P_2 = I_{\mathfrak{A}} - P_1$ nontrivial symmetric projections in \mathfrak{A} . Suppose that \mathfrak{A} satisfies

$$(\spadesuit) \qquad P_{i}\mathfrak{A} X = \{0\} \quad implies \quad X = 0.$$

Even more, suppose that $\varphi : \mathfrak{A} \to \mathfrak{A}'$ is a bijective unital map which satisfies

$$(\bullet) \varphi(p_{n_*}(A, B, \Xi, ..., \Xi)) = p_{n_*}(\varphi(A), \varphi(B), \varphi(\Xi), ..., \varphi(\Xi)),$$

for all $A, B \in \mathfrak{A}$ and $\Xi \in \{P_1, P_2, I_{\mathfrak{A}}\}$. Then φ is *-additive.

The following claims and lemmas have the same hypotheses as the Theorem 2.1 and we need them to prove the *-additivity of φ .

Claim 2.1. $*(\mathfrak{A}_{jk}) \subset \mathfrak{A}_{kj}$, for $j, k \in \{1, 2\}$.

Proof. It follows from the symmetry of P_1 and P_2 .

The next result can be found in [6] but for brevity we put the proof here

Claim 2.2. Let X, Y and H be in \mathfrak{A} such that $\varphi(H) = \varphi(X) + \varphi(Y)$. Then, given $Z \in \mathfrak{A}$,

$$\varphi(p_{n_*}(H, Z, \Xi, ..., \Xi)) = \varphi(p_{n_*}(X, Z, \Xi, ..., \Xi)) + \varphi(p_{n_*}(Y, Z, \Xi, ..., \Xi))$$

and

$$\varphi(p_{n_*}(Z,H,\Xi,...,\Xi)) = \varphi(p_{n_*}(Z,X,\Xi,...,\Xi)) + \varphi(p_{n_*}(Z,Y,\Xi,...\Xi))$$

for $\Xi \in \{P_1, P_2, I_{\mathfrak{A}}\}.$

Proof. Using the definition of φ and multilinearity of p_{n_*} we obtain

$$\begin{split} \varphi(p_{n_*}(H,Z,\Xi,...,\Xi)) &= p_{n_*}(\varphi(H),\varphi(Z),\varphi(\Xi),...,\varphi(\Xi)) \\ &= p_{n_*}(\varphi(X) + \varphi(Y),\varphi(Z),\varphi(\Xi),...,\varphi(\Xi)) \\ &= p_{n_*}(\varphi(X),\varphi(Z),\varphi(\Xi),...,\varphi(\Xi)) \\ &+ p_{n_*}(\varphi(Y),\varphi(Z),\varphi(\Xi),...,\varphi(\Xi)) \\ &= \varphi(p_{n_*}(X,Z,\Xi,...,\Xi)) \\ &+ \varphi(p_{n_*}(X,Z,\Xi,...,\Xi)). \end{split}$$

In a similar way we have

$$\varphi(p_{n_*}(Z,H,\Xi,...,\Xi))=\varphi(p_{n_*}(Z,X,\Xi,...,\Xi))+\varphi(p_{n_*}(Z,Y,\Xi,...,\Xi)). \quad \Box$$

Claim 2.3. $\varphi(0) = 0$.

Proof. Since φ is surjective, there exists $X \in \mathfrak{A}$ such that $\varphi(X) = 0$. Then, $\varphi(0) = \varphi(p_{n_*}(0, X, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}})) = p_{n_*}(\varphi(0), \varphi(X), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) = 0.$

We follow with a sequence of lemmas that show the additivity of φ .

Lemma 2.1. For any $A_{11} \in \mathfrak{A}_{11}$ and $B_{22} \in \mathfrak{A}_{22}$, we have

$$\varphi(A_{11} + B_{22}) = \varphi(A_{11}) + \varphi(B_{22}).$$

Proof. Since φ is surjective, given $\varphi(A_{11}) + \varphi(B_{22}) \in \mathfrak{A}'$ there exists $H \in \mathfrak{A}$ such that $\varphi(H) = \varphi(A_{11}) + \varphi(B_{22})$, with $H = H_{11} + H_{12} + H_{21} + H_{22}$. Besides, by claims 2.2 and 2.3

$$\varphi(p_{n_*}(P_1, H, P_1, ..., P_1)) = \varphi(p_{n_*}(P_1, A_{11}, P_1, ..., P_1)) + \varphi(p_{n_*}(P_1, B_{22}, P_1, ..., P_1)),$$

that is,

$$\varphi(-H_{21} + H_{21}^*) = \varphi(0) + \varphi(0) = 0$$

Then, by injectivity of φ , $-H_{21} + H_{21}^* = 0$. Thus $H_{21} = 0$. Moreover,

 $\varphi(p_{n_*}(P_2, H, P_2, ..., P_2)) = \varphi(p_{n_*}(P_2, A_{11}, P_2, ..., P_2)) + \varphi(p_{n_*}(P_2, B_{22}, P_2, ..., P_2)),$

that is,

$$\varphi(-H_{12} + H_{12}^*) = 0.$$

Again, by injectivity of φ we conclude that $H_{12} = 0$.

Furthermore, given $D_{21} \in \mathfrak{A}_{21}$,

$$\varphi(p_{n_*}(D_{21}, H, P_1, ..., P_1)) = \varphi(p_{n_*}(D_{21}, A_{11}, P_1, ..., P_1)) + \varphi(p_{n_*}(D_{21}, B_{22}, P_1, ..., P_1)),$$

that is,

$$\varphi(D_{21}H_{11} - (D_{21}H_{11})^*) = \varphi(D_{21}A_{11} - (D_{21}A_{11})^*).$$

Then we conclude, by injectivity of φ , that $D_{21}H_{11} - (D_{21}H_{11})^* = D_{21}A_{11} - (D_{21}A_{11})^*$, that is, $D_{21}(H_{11} - A_{11}) = 0$. Even more, $P_2\mathfrak{A}(H_{11} - A_{11}) = 0$, which implies that $H_{11} = A_{11}$ by (\blacklozenge).

Finally, given $D_{12} \in \mathfrak{A}_{12}$, a similar calculation gives us $H_{22} = B_{22}$. Therefore $H = A_{11} + B_{22}$.

Lemma 2.2. For any $A_{12} \in \mathfrak{A}_{12}$ and $B_{21} \in \mathfrak{A}_{21}$, we have $\varphi(A_{12} + B_{21}) = \varphi(A_{12}) + \varphi(B_{21})$.

Proof. Since φ is surjective, given $\varphi(A_{12}) + \varphi(B_{21}) \in \mathfrak{A}'$ there exists $H \in \mathfrak{A}$ such that $\varphi(H) = \varphi(A_{12}) + \varphi(B_{21})$, with $H = H_{11} + H_{12} + H_{21} + H_{22}$. Now, by Claims 2.2 and 2.3

$$\varphi(p_{n_*}(P_1, H, P_1, \dots, P_1)) = \varphi(p_{n_*}(P_1, A_{12}, P_1, \dots, P_1)) + \varphi(p_{n_*}(P_1, B_{21}, P_1, \dots, P_1)),$$

that is,

$$\varphi(-H_{21} + H_{21}^*) = \varphi(-B_{21} + B_{21}^*).$$

Then, by injectivity of φ , $-H_{21} + H_{21}^* = -B_{21} + B_{21}^*$. Thus $H_{21} = B_{21}$. Moreover,

$$\varphi(p_{n_*}(P_2, H, P_2, ..., P_2)) = \varphi(p_{n_*}(P_2, A_{12}, P_2, ..., P_2)) + \varphi(p_{n_*}(P_2, B_{21}, P_2, ..., P_2)),$$

that is,

$$\varphi(-H_{12} + H_{12}^*) = \varphi(-A_{12} + A_{12}^*)$$

Again, by injectivity of φ we conclude that $H_{12} = A_{12}$.

Furthermore, given $D_{21} \in \mathfrak{A}_{21}$,

$$\begin{aligned} \varphi(D_{21}H_{11} - (D_{21}H_{11})^*) &= \varphi(p_{n_*}(D_{21}, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(D_{21}, A_{12}, P_1, ..., P_1)) \\ &+ \varphi(p_{n_*}(D_{21}, B_{21}, P_1, ..., P_1)) = 0 \end{aligned}$$

Then we conclude, by injectivity of φ , that $D_{21}H_{11} - (D_{21}H_{11})^* = 0$, that is, $D_{21}H_{11} = 0$. Even more, $P_2\mathfrak{A}H_{11} = 0$, which implies that $H_{11} = 0$ by (\bigstar) .

Finally, given $D_{12} \in \mathfrak{A}_{12}$, a similar calculation gives us $H_{22} = 0$. Therefore, we conclude that $H = A_{12} + B_{21}$.

Lemma 2.3. For any $A_{11} \in \mathfrak{A}_{11}$, $B_{12} \in \mathfrak{A}_{12}$, $C_{21} \in \mathfrak{A}_{21}$ and $D_{22} \in \mathfrak{A}_{22}$ we have

$$\varphi(A_{11} + B_{12} + C_{21} + D_{22}) = \varphi(A_{11}) + \varphi(B_{12}) + \varphi(C_{21}) + \varphi(D_{22}).$$

Proof. Since φ is surjective, given $\varphi(A_{11}) + \varphi(B_{12}) + \varphi(C_{21}) + \varphi(D_{22}) \in \mathfrak{A}'$ there exists $H \in \mathfrak{A}$ such that $\varphi(H) = \varphi(A_{11}) + \varphi(B_{12}) + \varphi(C_{21}) + \varphi(D_{22})$, with $H = H_{11} + H_{12} + H_{21} + H_{22}$. Even more, by Lemmas 2.1 and 2.2

$$\varphi(H) = \varphi(A_{11}) + \varphi(B_{12}) + \varphi(C_{21}) + \varphi(D_{22}) = \varphi(A_{11} + D_{22}) + \varphi(B_{12} + C_{21}).$$

Now, observing that $p_{n_*}(P_1, A_{11} + D_{22}, ..., P_1) = 0 = p_{n_*}(P_1, B_{12}, P_1, ..., P_1)$ and by Claims 2.2 and 2.3 we obtain

$$\begin{split} \varphi(p_{n_*}(P_1, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(P_1, A_{11} + D_{22}, ..., P_1)) + \varphi(p_{n_*}(P_1, B_{12} + C_{21}, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(P_1, C_{21}, P_1, ..., P_1)), \end{split}$$

that is,

$$\varphi(-H_{21} + H_{21}^*) = \varphi(-C_{21} + C_{21}^*).$$

Then, by injectivity of φ , $-H_{21} + H_{21}^* = -C_{21} + C_{21}^*$. Thus $H_{21} = C_{21}$.

In a similar way, using P_2 rather than P_1 in the previous calculation, we conclude that $H_{12} = B_{12}$. Also, given $X_{21} \in \mathfrak{A}_{21}$,

$$\begin{split} &\varphi(p_{n_*}(X_{21}, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(X_{21}, A_{11} + D_{22}, ..., P_1)) + \varphi(p_{n_*}(X_{21}, B_{12} + C_{21}, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(X_{21}, A_{11}, P_1, ..., P_1)), \end{split}$$

since $p_{n_*}(X_{21}, B_{12} + C_{21}, P_1, ..., P_1) = 0 = p_{n_*}(X_{21}, D_{22}, ..., P_1)$. Again, by injectivity of φ we conclude, by following the same strategy as in the proof of Lemma 2.1, that $H_{11} = A_{11}$. Now, using P_2 rather than P_1 and X_{12} rather than X_{21} in the previous calculation we obtain $H_{22} = D_{22}$. Therefore, $H = A_{11} + B_{12} + C_{21} + D_{22}$.

Lemma 2.4. For all $A_{jk}, B_{jk} \in \mathfrak{A}_{jk}$ we have $\varphi(A_{jk} + B_{jk}) = \varphi(A_{jk}) + \varphi(B_{jk})$ for $j \neq k$.

Proof. We shall prove the case j = 1 and k = 2, since the other case is done in a similar way. Since φ is surjective, given $\varphi(A_{12}) + \varphi(B_{12}) \in \mathfrak{A}'$ and $\varphi(-A_{12}^*) + \varphi(-B_{12}^*)$ there exist $H \in \mathfrak{A}$ and $T \in \mathfrak{A}$ such that $\varphi(H) = \varphi(A_{12}) + \varphi(B_{12})$ and $\varphi(T) = \varphi(-A_{12}^*) + \varphi(-B_{12}^*)$, with $H = H_{11} + H_{12} + H_{21} + H_{22}$ and $T = T_{11} + T_{12} + T_{21} + T_{22}$.

Firstly, we show that $H \in \mathfrak{A}_{12}$: by Claim 2.2

$$\begin{aligned} \varphi(-H_{21} + H_{21}^*) &= \varphi(p_{n_*}(P_1, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(P_1, A_{12}, P_1, ..., P_1)) + \varphi(p_{n_*}(P_1, B_{12}, P_1, ..., P_1)) = 0 \end{aligned}$$

Then, by injectivity of φ we obtain $H_{21} = 0$. Also, given $D_{12} \in \mathfrak{A}_{12}$,

$$\begin{aligned} \varphi(D_{12}H_{22} - (D_{12}H_{22})^*) &= \varphi(p_{n_*}(D_{12}, H, P_2, ..., P_2)) \\ &= \varphi(p_{n_*}(D_{12}, A_{12}, P_2, ..., P_2)) \\ &+ \varphi(p_{n_*}(D_{12}, B_{12}, P_2, ..., P_2)) = 0, \end{aligned}$$

that is, $D_{12}H_{22} = 0$, which implies that $H_{22} = 0$ by (\bigstar). Now, using $D_{21} \in \mathfrak{A}_{21}$ rather than D_{12} in the previous calculation, we conclude that $H_{11} = 0$. Therefore, $H = H_{12} \in \mathfrak{A}_{12}$.

In a similar way, we obtain $T = T_{21} \in A_{21}$. Finally, by Lemma 2.3

$$\begin{split} \varphi(A_{12} + B_{12} - A_{12}^* - B_{12}^*) &= \varphi(p_{n*}(P_1 + A_{12}, P_2 + B_{12}, P_2, ..., P_2)) \\ &= p_{n*}(\varphi(P_1 + A_{12}), \varphi(P_2 + B_{12}), \varphi(P_2), ..., \varphi(P_2)) \\ &= p_{n*}(\varphi(P_1), \varphi(P_2), \varphi(P_2), ..., \varphi(P_2)) \\ &+ p_{n*}(\varphi(P_1), \varphi(B_{12}), \varphi(P_2), ..., \varphi(P_2)) \\ &+ p_{n*}(\varphi(A_{12}), \varphi(P_2), \varphi(P_2), ..., \varphi(P_2)) \\ &= \varphi(p_{n*}(P_1, P_2, P_2, ..., P_2)) \\ &+ \varphi(p_{n*}(A_{12}, P_2, P_2, ..., P_2)) \\ &+ \varphi(p_{n*}(A_{12}, B_{12}, P_2, ..., P_2)) \\ &= \varphi(A_{12} - A_{12}^*) + \varphi(B_{12} - B_{12}^*) \\ &= \varphi(A_{12}) + \varphi(B_{12}) + \varphi(-A_{12}^*) + \varphi(-B_{12}^*) \\ &= \varphi(H_{12}) + \varphi(T_{21}) = \varphi(H_{12} + T_{21}). \end{split}$$

Since φ is injective, we have $A_{12} + B_{12} - A_{12}^* - B_{12}^* = H_{12} + T_{21}$, i.e., $H = H_{12} = A_{12} + B_{12}$.

Lemma 2.5. For all $A_{jj}, B_{jj} \in \mathfrak{A}_{jj}$, we have $\varphi(A_{jj} + B_{jj}) = \varphi(A_{jj}) + \varphi(B_{jj})$ for $j \in \{1, 2\}$.

Proof. We shall prove the case j = 1, since the other case is done in a similar way. Since φ is surjective, given $\varphi(A_{11}) + \varphi(B_{11}) \in \mathfrak{A}'$ there exists $H \in \mathfrak{A}$ such that $\varphi(H) = \varphi(A_{11}) + \varphi(B_{11})$, with $H = H_{11} + H_{12} + H_{21} + H_{22}$. Now, by claim 2.2

$$\begin{aligned} \varphi(-H_{21} + H_{21}^*) &= \varphi(p_{n_*}(P_1, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(P_1, A_{11}, P_1, ..., P_1)) + \varphi(p_{n_*}(P_1, B_{11}, P_1, ..., P_1)) = 0 \end{aligned}$$

Then, by injectivity of φ we obtain $H_{21} = 0$. Also,

$$\varphi(-H_{12} + H_{12}^*) = \varphi(p_{n_*}(P_2, H, P_2, ..., P_2))$$

= $\varphi(p_{n_*}(P_2, A_{11}, P_2, ..., P_2)) + \varphi(p_{n_*}(P_2, B_{11}, P_2, ..., P_2)) = 0$

that is, $H_{12} = 0$ by injectivity of φ . Moreover, given $D_{12} \in \mathfrak{A}_{12}$,

$$\begin{aligned} \varphi(D_{12}H_{22} - (D_{12}H_{22})^*) &= \varphi(p_{n_*}(D_{12}, H, P_2, ..., P_2)) \\ &= \varphi(p_{n_*}(D_{12}, A_{11}, P_2, ..., P_2)) \\ &+ \varphi(p_{n_*}(D_{12}, B_{11}, P_2, ..., P_2)) = 0. \end{aligned}$$

Then, by injectivity of φ , $D_{12}H_{22} = 0$, which implies that $H_{22} = 0$ by (\blacklozenge). Finally, given $D_{21} \in \mathfrak{A}_{21}$, by Lemmas 2.3 and 2.4 we have

$$\begin{split} \varphi(D_{21}H_{11} - (D_{21}H_{11})^*) &= \varphi(p_{n_*}(D_{21}, H, P_1, ..., P_1)) \\ &= \varphi(p_{n_*}(D_{21}, A_{11}, P_1, ..., P_1)) \\ &+ \varphi(p_{n_*}(D_{21}, B_{11}, P_1, ..., P_1)) \\ &= \varphi(D_{21}A_{11} - (D_{21}A_{11})^*) \\ &+ \varphi(D_{21}B_{11} - (D_{21}B_{11})^*) \\ &= \varphi(D_{21}A_{11}) + \varphi(-(D_{21}A_{11})^*) \\ &+ \varphi(D_{21}B_{11}) + \varphi(-(D_{21}B_{11})^*) \\ &= \varphi(D_{21}A_{11} + D_{21}B_{11}) \\ &+ \varphi(-(D_{21}A_{11})^* - (D_{21}B_{11})^*) \\ &= \varphi(D_{21}(A_{11} + B_{11}) - (A_{11}^* + B_{11}^*)D_{21}^*), \end{split}$$

that is, $D_{21}H_{11} - (D_{21}H_{11})^* = D_{21}(A_{11} + B_{11}) - (A_{11}^* + B_{11}^*)D_{21}^*$, by injectivity of φ . Thus, $D_{21}(H_{11} - (A_{11} + B_{11})) = 0$, which implies that $H_{11} = A_{11} + B_{11}$ by (\blacklozenge).

Now we are able to show that φ is *-additive. Using Lemmas 2.3, 2.4 and 2.5 we have, for all $A, B \in \mathfrak{A}$,

$$\begin{split} \varphi(A+B) &= \varphi(A_{11}+A_{12}+A_{21}+A_{22}+B_{11}+B_{12}+B_{21}+B_{22}) \\ &= \varphi(A_{11}+B_{11}) + \varphi(A_{12}+B_{12}) + \varphi(A_{21}+B_{21}) + \varphi(A_{22}+B_{22}) \\ &= \varphi(A_{11}) + \varphi(B_{11}) + \varphi(A_{12}) + \varphi(B_{12}) \\ &+ \varphi(A_{21}) + \varphi(B_{21}) + \varphi(A_{22}) + \varphi(B_{22}) \\ &= \varphi(A_{11}+A_{12}+A_{21}+A_{22}) \\ &+ \varphi(B_{11}+B_{12}+B_{21}+B_{22}) = \varphi(A) + \varphi(B). \end{split}$$

Besides, given $A \in \mathfrak{A}$, by additivity of φ

$$2^{n-2}(\varphi(A) - \varphi(A)^*) = p_{n_*}(\varphi(A), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) = \varphi(p_{n_*}(A, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}))$$
$$= \varphi(2^{n-2}(A - A^*)) = 2^{n-2}\varphi(A - A^*) = 2^{n-2}(\varphi(A) - \varphi(A^*))$$

and then we conclude that $\varphi(A^*) = \varphi(A)^*$. This completes the proof of Theorem 2.1.

Theorem 2.2. Let \mathfrak{A} and \mathfrak{A}' be two C^* -algebras with identities $I_{\mathfrak{A}}$ and $I_{\mathfrak{A}'}$, respectively, and P_1 and $P_2 = I_{\mathfrak{A}} - P_1$ nontrivial symmetric projections in \mathfrak{A} . Suppose that \mathfrak{A} satisfies

$$(\spadesuit) \qquad P_{i}\mathfrak{A} X = \{0\} \quad implies \quad X = 0.$$

Even more, suppose that $\varphi : \mathfrak{A} \to \mathfrak{A}'$ is a complex scalar multiplication bijective unital map which satisfies

$$(\clubsuit) \qquad \varphi(P_j)\mathfrak{A}'Y = \{0\} \quad implies \quad Y = 0$$

and

$$(\bullet) \varphi(p_{n_*}(A, B, \Xi, ..., \Xi)) = p_{n_*}(\varphi(A), \varphi(B), \varphi(\Xi), ..., \varphi(\Xi)),$$

for all $A, B \in \mathfrak{A}$ and $\Xi \in \{P_1, P_2, I_{\mathfrak{A}}\}$. Then φ is *-isomorphism.

With this hypothesis, we have already proved that φ is *-additive. It remains for us to show that φ is multiplicative. In order to do that we will prove some more lemmas. Firstly, we observe that, for any $A \in \mathfrak{A}$,

Remark 2.1. Denoting by $\mathcal{R}(A)$ and $\mathcal{I}(A)$ the real part and imaginary part of $A \in \mathfrak{A}$, respectively, we have

$$\varphi(2^{n-1}\mathcal{R}(A)) = \varphi(2^{n-2}(A+A^*)) = \varphi(p_{n_*}(iA, -iI_{\mathfrak{A}}, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}))$$
$$= p_{n_*}(i\varphi(A), -i\varphi(I_{\mathfrak{A}}), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) = 2^{n-2}(\varphi(A) + \varphi(A)^*)$$
$$= 2^{n-1}\mathcal{R}(\varphi(A))$$

and

$$\varphi(2^{n-1}i\mathcal{I}(A)) = \varphi(2^{n-2}(A - A^*)) = \varphi(p_{n_*}(A, I_\mathfrak{A}, ..., I_\mathfrak{A}))$$

= $p_{n_*}(\varphi(A), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) = 2^{n-2}(\varphi(A) - \varphi(A)^*)$
= $2^{n-1}i\mathcal{I}(\varphi(A)).$

Even more,

Claim 2.4. $Q_j = \varphi(P_j)$ is a symmetric projection in \mathfrak{A}' , with $j \in \{1, 2\}$.

Proof. Since φ is a complex scalar multiplication, it follows that

$$2^{n-1}iQ_j = 2^{n-1}i\varphi(P_j) = \varphi(2^{n-1}iP_j) = \varphi(p_{n_*}(iP_j, P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}))$$

= $p_{n_*}(i\varphi(P_j), \varphi(P_j), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'})) = 2^{n-1}i\varphi(P_j)^2 = 2^{n-1}iQ_j^2$

Then we can conclude that $Q_j = Q_j^2$. Moreover, since P_j is a symmetric projection in \mathfrak{A} we have that $p_{n_*}(P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}) = 0$. Besides,

$$0 = \varphi(0) = \varphi(p_{n_*}(P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}})) = p_{n_*}(Q_j, I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}).$$

Thus, $Q_j - Q_j^* = 0$, that is, $Q_j = Q_j^*$.

Lemma 2.6. For all $A \in \mathfrak{A}$, $\varphi(P_jA) = \varphi(P_j)\varphi(A)$ and $\varphi(AP_j) = \varphi(A)\varphi(P_j)$.

Proof. Firstly, observe that

$$p_{n_*}(iA, P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}) = 2^{n-2}i(AP_j + P_jA^*)$$

and

$$p_{n_*}(A, P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}}) = 2^{n-2}(AP_j - P_jA^*)$$

Still, by (\bullet) and *-additivity of φ ,

$$\begin{split} \varphi(2^{n-2}i(AP_j + P_jA^*)) &= \varphi(p_{n_*}(iA, P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}})) \\ &= p_{n_*}(\varphi(iA), \varphi(P_j), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) \\ &= 2^{n-2}i(\varphi(A)\varphi(P_j) + \varphi(P_j)\varphi(A)^*) \end{split}$$

and

$$\begin{split} \varphi(2^{n-2}(AP_j - P_jA^*)) &= \varphi(p_{n_*}(A, P_j, I_{\mathfrak{A}}, ..., I_{\mathfrak{A}})) \\ &= p_{n_*}(\varphi(A), \varphi(P_j), I_{\mathfrak{A}'}, ..., I_{\mathfrak{A}'}) \\ &= 2^{n-2}(\varphi(A)\varphi(P_j) - \varphi(P_j)\varphi(A)^*). \end{split}$$

Now, since φ is *-additive, multiplying the second equality by *i* and adding these two equations we obtain $\varphi(AP_j) = \varphi(A)\varphi(P_j)$. The second statement is obtained in a similar way.

Lemma 2.7. $\varphi(\mathfrak{A}_{jk}) \subset \mathfrak{A}'_{jk}$.

Proof. Given $X \in \mathfrak{A}_{jk}$, we have $X = P_j X P_k$ and then, by Lemma 2.6, $\varphi(X) = \varphi(P_j)\varphi(XP_k) = \varphi(P_j)\varphi(X)\varphi(P_k) \in \mathfrak{A}'_{jk}$.

Lemma 2.8. To $j \neq k$:

- If $A_{jk} \in \mathfrak{A}_{jk}$ and $B_{kk} \in \mathfrak{A}_{kk}$ then $\varphi(A_{jk}B_{kk}) = \varphi(A_{jk})\varphi(B_{kk})$;
- If $A_{jj} \in \mathfrak{A}_{jj}$ and $B_{jk} \in \mathfrak{A}_{jk}$ then $\varphi(A_{jj}B_{jk}) = \varphi(A_{jj})\varphi(B_{jk})$;
- If $A_{jk} \in \mathfrak{A}_{jk}$ and $B_{kj} \in \mathfrak{A}_{kj}$ then $\varphi(A_{jk}B_{kj}) = \varphi(A_{jk})\varphi(B_{kj})$.

Proof. In order to prove the first statement, on the one hand, by Lemma 2.7

$$\varphi(A_{jk}B_{kk}) - \varphi(A_{jk}B_{kk})^* = \varphi(A_{jk}B_{kk} - (A_{jk}B_{kk})^*)$$
$$= \varphi(p_{n_*}(A_{jk}, B_{kk}, P_k, ..., P_k))$$
$$= p_{n_*}(\varphi(A_{jk}), \varphi(B_{kk}), Q_k, ..., Q_k)$$
$$= \varphi(A_{jk})\varphi(B_{kk}) - (\varphi(A_{jk})\varphi(B_{kk}))^*$$

and then $\mathcal{I}(\varphi(A_{jk}B_{kk})) = \mathcal{I}(\varphi(A_{jk})\varphi(B_{kk}))$. On the other hand, using iA_{jk} rather than A_{jk} we obtain $\mathcal{R}(\varphi(A_{jk}B_{kk})) = \mathcal{R}(\varphi(A_{jk})\varphi(B_{kk}))$. It concludes that $\varphi(A_{jk}B_{kk}) = \varphi(A_{jk})\varphi(B_{kk})$.

The others statements are proved in a similar way.

Lemma 2.9. If $A_{ij}, B_{ij} \in \mathfrak{A}_{ij}$ then $\varphi(A_{ij}B_{ij}) = \varphi(A_{ij})\varphi(B_{ij})$.

Proof. Let X_{kj} be an element of \mathfrak{A}_{kj} , with $j \neq k$. Using Lemma 2.8 we obtain

$$\varphi(X_{kj})\varphi(A_{jj}B_{jj}) = \varphi(X_{kj}A_{jj}B_{jj}) = \varphi(X_{kj}A_{jj})\varphi(B_{jj}) = \varphi(X_{kj})\varphi(A_{jj})\varphi(B_{jj})$$

that is,

$$\varphi(X_{kj})(\varphi(A_{jj}B_{jj}) - \varphi(A_{jj})\varphi(B_{jj})) = 0.$$

Now, by Lemma 2.7, $\varphi(X_{kj}) \in \mathfrak{A}'_{kj}$ as well as $\varphi(A_{jj}B_{jj})$ and $\varphi(A_{jj})\varphi(B_{jj}) \in \mathfrak{A}'_{jj}$. Then, $Q_k\mathfrak{A}'(\varphi(A_{jj}B_{jj}) - \varphi(A_{jj})\varphi(B_{jj})) = 0$, which implies that $\varphi(A_{jj}B_{jj}) = \varphi(A_{jj})\varphi(B_{jj})$ by (\clubsuit) .

By additivity of φ and Lemmas 2.8 and 2.9, it follows, for all $A, B \in \mathfrak{A}$, that $\varphi(AB) = \varphi(A)\varphi(B)$. It concludes the proof of Theorem 2.2.

3 Corollaries

Let us present some consequences of our main result. The first one provides the conjecture that appears in [7] to the case of multiplicative *-Lie-type maps:

Corollary 3.1. Let \mathfrak{A} and \mathfrak{A}' be two C^* -algebras with identities $I_{\mathfrak{A}}$ and $I_{\mathfrak{A}'}$, respectively, and P_1 and $P_2 = I_{\mathfrak{A}} - P_1$ nontrivial symmetric projections in \mathfrak{A} . Suppose that \mathfrak{A} satisfies

$$(\spadesuit) \qquad P_{j}\mathfrak{A} X = \{0\} \quad implies \quad X = 0.$$

Even more, suppose that $\varphi : \mathfrak{A} \to \mathfrak{A}'$ is a complex scalar multiplication bijective unital map which satisfies

$$(\clubsuit) \qquad \varphi(P_i)\mathfrak{A}'Y = \{0\} \quad implies \quad Y = 0.$$

Then $\varphi : \mathfrak{A} \to \mathfrak{A}'$ is a multiplicative *-Lie n-map if and only if φ is a *-isomorphism.

Observing that prime C^* -algebras satisfy $(\spadesuit), (\clubsuit)$ we have the following result:

Corollary 3.2. Let \mathfrak{A} and \mathfrak{A}' be prime C^* -algebras with identities $I_{\mathfrak{A}}$ and $I_{\mathfrak{A}'}$, respectively, and P_1 and $P_2 = I_{\mathfrak{A}} - P_1$ nontrivial projections in \mathfrak{A} . Then a complex scalar multiplication $\varphi : \mathfrak{A} \to \mathfrak{A}'$ is a bijective unital multiplicative *-Lie n-map if and only if φ is a *-isomorphism.

A von Neumann algebra \mathcal{M} is a weakly closed, self-adjoint algebra of operators on a Hilbert space \mathcal{H} containing the identity operator I. As an application on von Neumann algebras we have the following:

Corollary 3.3. Let \mathcal{M} be a von Neumann algebra without central summands of type I_1 . Then a complex scalar multiplication $\varphi : \mathcal{M} \to \mathcal{M}$ is a bijective unital multiplicative *-Lie n-map if and only if φ is a *-isomorphism.

Proof. Let \mathcal{M} be a von Neumann algebra. It is shown in [1] and [16] that if a von Neumann algebra has no central summands of type I_1 , then \mathcal{M} satisfies the following assumption:

• $P_i \mathcal{M} X = \{0\} \Rightarrow X = 0.$

Thus, by Theorem 2.2 the corollary is true.

To finish, \mathcal{M} is a factor von Neumann algebra if its center only contains the scalar operators. It is well known that a factor von Neumann algebra is prime and then we have the following:

Corollary 3.4. Let \mathcal{M} be a factor von Neumann algebra. Then a complex scalar multiplication $\varphi : \mathcal{M} \to \mathcal{M}$ is a bijective unital multiplicative *-Lie *n*-map if and only if φ is a *-isomorphism.

References

- Z. Bai and S. Du, Strong commutativity preserving maps on rings, Rocky Mountain J. Math., 44, (2014), 733-742.
- [2] M. Brešar and M. Fošner, On rings with involution equipped with some new product, Publicationes Math., 57 (2000).
- [3] Q. Chen and C. Li, Additivity of Lie multiplicative mappings on rings, Adv. in Math.(China), 46(1) (2017), 82-90.
- [4] J.Cui and C. Li, Maps preserving product XY YX* on factor von Neumann algebras, Linear Algebra and its Applications, 431, (2009), 833-842.
- [5] X. Fang, C. Li and F. Lu, Nonlinear mappings preserving product XY + YX* on factor von Neumann algebras, Linear Algebra and its Applications, 438, (2013), 2339-2345.
- [6] B. L. M. Ferreira and B. T. Costa, *-Jordan-type maps on C*-algebras, arXiv:2005.11430 [math.OA]
- [7] B. L. M. Ferreira and B. T. Costa, *-Lie-Jordan-type maps on C*algebras, arXiv:2003.11123 [math.OA]
- [8] B. L. M. Ferreira, Multiplicative maps on triangular n-matrix rings, International Journal of Mathematics, Game Theory and Algebra, 23, p. 1-14, 2014.
- [9] J. C. M. Ferreira and B. L. M. Ferreira, Additivity of n-multiplicative maps on alternative rings, Comm. in Algebra, 44 (2016), 1557-1568.
- [10] R. N. Ferreira and B. L. M. Ferreira, Jordan triple derivation on alternative rings, Proyectiones J. Math., 37 (2018), 171-180.
- [11] B. L. M. Ferreira, J. C. M. Ferreira, H. Guzzo Jr., Jordan maps on alternatives algebras, JP Journal of Algebra, Number Theory and Applications, **31**, p. 129-142, 2013.
- [12] B. L. M. Ferreira, J. C. M. Ferreira, H. Guzzo Jr., Jordan triple elementary maps on alternative rings, Extracta Mathematicae, 29, p. 1-18, 2014.

- [13] B. L. M. Ferreira, J. C. M. Ferreira, H. Guzzo Jr., Jordan triple maps of alternatives algebras, JP Journal of Algebra, Number Theory and Applications, 33, p. 25-33, 2014.
- [14] B. L. M. Ferreira and H. Guzzo Jr., *Lie maps on alternative rings*, Boll. Unione Mat. Ital., (2019), 1-12.
- [15] M. Fošner, Prime Rings with Involution Equipped with Some New Product, Southeast Asian Bull. Math., 26 (2002), 27-31.
- [16] W. S. Martindale III, Lie isomorphisms of operator algebras Pacific J. Math 38 (1971), 717-735.
- [17] W. S. Martindale III, When are multiplicative mappings additive? Proc. Amer. Math. Soc. 21 (1969), 695-698.