Stability theorems for multiplicities in graded S_n -modules

Marino Romero * Nolan Wallach

February 15, 2022

Abstract

In this paper, we prove several stability theorems for multiplicities of naturally defined representations of symmetric groups. The first such theorem states that if we consider the diagonal action of the symmetric group S_{m+r} on k sets of m + r variables, then the dimension of the invariants of degree m is the same as the dimension of the invariants of degree m for S_m acting on k sets of m variables. The second type of stability result is for Weyl modules. We prove that the dimension of the S_{n+r} invariants for a Weyl module, ${}_{m+r}F^{\lambda}$ (the Schur-Weyl dual of the $S_{|\lambda|}$ module V^{λ}) with $|\lambda| \leq m$ is of the same dimension as the space of S_m invariants for ${}_mF^{\lambda}$. Multigraded versions of the first type of result are given, as are multigraded generalizations to non-trivial modules of symmetric groups. Two related conjectures are given which if proved would give efficient calculations of multiplicities in the stable range.

1 Introduction

Consider the diagonal action of the symmetric group on n letters, S_n , on k sets of variables. A classical theorem of Hermann Weyl asserts that the polynomial invariants in k sets of variables are generated by the polarizations of the invariants in one set of variables. Thus if we take a standard

^{*}The author was partially supported by the University of California President's Post-doctoral Fellowship.

set, $f_1, ..., f_n$, of generators of the symmetric functions in n variables and do the repeated polarizations to a set of k variables, one gets a set of generators for the invariants for the diagonal action of the symmetric group on k sets of variables. A natural question is what are the relations among these polynomials (the so-called second fundamental theorem)? In this paper we show that the first relation is in degree n + 2, which has the following surprising (at least to us) implication: Looking upon the diagonal action of S_n on ksets of variables as the action of S_n on $\mathbb{C}^k \otimes \mathbb{C}^n$ given by sending s to $I \otimes s$, if $m \leq n$ then

$$\dim S^m(\mathbb{C}^k \otimes \mathbb{C}^m)^{S_m} = \dim S^m(\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n}.$$

Here if M is an S_n module, M^{S_n} is the space of invariants.

In this paper all of the results are true over arbitrary fields of characteristic 0. We state them over \mathbb{C} , but they can equally well be stated and proved in the same way over \mathbb{Q} .

This stability theorem is equivalent to another of a different sort. For this we need some notation. If $\lambda, \mu \in \mathbb{Z}^n$ then we will say that $\lambda \succ \mu$ if

$$\lambda_1 + ... + \lambda_i \ge \mu_1 + ... + \mu_i$$
, for $i = 1, ..., n$.

For $\mu \in \mathbb{Z}^n$ we define the character of $(\mathbb{C}^{\times})^n$, thought of as the subgroup of diagonal matrices in $GL(n, \mathbb{C})$, by

$$z\mapsto z^{\mu}=z_1^{\mu_1}z_2^{\mu_2}\cdots z_n^{\mu_n}.$$

If $\lambda = (\lambda_1, ..., \lambda_r) \in \mathbb{Z}^r$ with $\lambda_1 \geq \lambda_2 \geq ... \geq \lambda_r \geq 0$ (that is λ is dominant) and $r \leq n$, then the theorem of the highest weight implies that there is a unique up to equivalence irreducible representation of $GL(n, \mathbb{C})$, ${}_nF^{\lambda}$, such that if $\Lambda = (\lambda_1, ..., \lambda_r, 0, ..., 0)$ with n - r added 0's and if $\mu \in \mathbb{Z}^n$ is such that there exists $v \in {}_nF^{\lambda}$ such that $v \neq 0$ and $zv = z^{\mu}v$ for $z \in (\mathbb{C}^*)^n$, then $\Lambda \succ \mu$. Another characterization using the Weyl character formula is that the character of ${}_nF^{\lambda}$ is $s_{\lambda}(z_1, ..., z_n)$, the Schur symmetric function.

The so-called $GL(k, \mathbb{C}) - GL(n, \mathbb{C})$ duality implies that as a representation of $GL(k, \mathbb{C}) \times GL(n, \mathbb{C})$ the space $S^r(\mathbb{C}^k \otimes \mathbb{C}^n)$ is equivalent to

$$\bigoplus_{|\lambda|=r} {}_k F^\lambda \otimes {}_n F^\lambda$$

where the sum is over dominant λ with at most min(k, n) entries and $|\lambda| = \sum \lambda_i = r$. Using this formalism, our second main result is as follows. We

look upon S_n as the group of permutation matrices in $GL(n, \mathbb{C})$, and for $m \leq n$ we identify $GL(m, \mathbb{C})$ with the group of matrices

$$\left\{ \left[\begin{array}{cc} g & 0 \\ 0 & I \end{array} \right] \middle| g \in GL(m, \mathbb{C}) \right\}.$$

Then for $|\lambda| \leq m \leq n$, the following stability holds:

$$\dim \left({}_{m}F^{\lambda}\right)^{S_{m}} = \dim \left({}_{n}F^{\lambda}\right)^{S_{n}}$$

In the standard Young correspondence between irreducible representations of S_n and partitions of n in the guise of Young diagrams, the trivial representation of S_n corresponds to the partition (n). If μ is a partition of n let V^{μ} denote the corresponding Young representation of S_n . Then if $\mu = (m)$ our result says that

$$\dim \operatorname{Hom}_{S_m}(V^{\mu}, {}_mF^{\lambda}) = \dim \operatorname{Hom}_{S_{m+r}}(V^{\mu+re_1}, {}_{n+r}F^{\lambda})$$

with $e_1, ..., e_l$ the usual basis of \mathbb{Z}^l . We show that this stability can be extended to non-trivial representations of the symmetric group as follows: If $\mu = (\mu_1, \mu_2, ...)$ is a partition of m and if λ is dominant with

$$|\lambda| \le m - \mu_2,$$

then

$$\dim \operatorname{Hom}_{S_m}(V^{\mu}, {}_mF^{\lambda}) = \dim \operatorname{Hom}_{S_{m+r}}(V^{\mu+re_1}, {}_{m+r}F^{\lambda}).$$

(Here if $\mu = (\mu_1)$ then we take $\mu_2 = 0$.) One key aspect of the proof of this stabilization result is the following result of the first named author of this paper (see [7]). The total degree version found in Proposition 5.2 of [10] can also be used to provide an alternate proof. Considering $S(\mathbb{C}^k \otimes \mathbb{C}^n)$ as a kgraded representation of $GL(n, \mathbb{C})$, $S(\mathbb{C}^k \otimes \mathbb{C}^n)$ has a multigrade as follows. If we have $(\mathbb{C}^{\times})^k$ act on $\mathbb{C}^k \otimes \mathbb{C}^n$ by $z \mapsto z \otimes I$ then the $L = (l_1, ..., l_k)$ -level of the grade is the space of all $v \in S(\mathbb{C}^k \otimes \mathbb{C}^n)$ such that $zv = z_1^{l_1} \cdots z_k^{l_k} v$ for all $z \in$ $(\mathbb{C}^{\times})^k$. Denote the *L*-level of the grade by $S^L(\mathbb{C}^k \otimes \mathbb{C}^n)$. Clearly, $S^L(\mathbb{C}^k \otimes \mathbb{C}^n)$ is invariant under the action of $GL(n, \mathbb{C})$ and thus of S_n . Therefore for any partition μ ,

$$\operatorname{Hom}_{S_n}(V^{\mu}, S(\mathbb{C}^k \otimes \mathbb{C}^n)) = \bigoplus_L \operatorname{Hom}_{S_n}(V^{\mu}, S^L(\mathbb{C}^k \otimes \mathbb{C}^n))$$

is a grading and there is an associated Hilbert series

$$h_{k,n}^{\mu}(q_1,...,q_k) = \sum_{L} q_1^{l_1} q_2^{l_2} \cdots q_k^{l_k} \dim \operatorname{Hom}_{S_n}(V^{\mu}, S^{L}(\mathbb{C}^k \otimes \mathbb{C}^n)).$$

Fix a monomial order on the monomials in the $q_1, ..., q_k$ (e.g. lexicographic order). A semi-standard filling of the Young diagram of μ is a placement of monomials in the diagram such that they are in strictly increasing order in the columns and weakly increasing order in the rows. The weight of the filling is the product of the monomials in the filling. The result used is that the coefficient of $q^L = q_1^{l_1} q_2^{l_2} \cdots q_k^{l_k}$ in $h_{k,n}^{\mu}(q_1, ..., q_k)$ is the number of fillings if μ of weight q^L . This in plethystic notation says

$$h_{k,n}^{\mu}(q_1,...,q_k) = s_{\mu} \left[\frac{1}{(1-q_1)\cdots(1-q_k)} \right].$$

Contrast this with the corresponding Hilbert series

$$\sum_{L} q^{L} \dim \operatorname{Hom}_{GL(n,\mathbb{C})}({}_{n}F^{\lambda}, S^{L}(\mathbb{C}^{k} \otimes \mathbb{C}^{n})) = s_{\lambda}(q_{1}, ..., q_{k}).$$

So one has

$$s_{\mu}\left[\frac{1}{(1-q_1)\cdots(1-q_k)}\right] = \sum_{\lambda} s_{\lambda}(q_1,...,q_k) \dim \operatorname{Hom}_{S_n}(V^{\mu},{}_nF^{\lambda}).$$

The notion of stability of irreducible representations of S_n has appeared in several contexts. For one, the Kronecker coefficients have a stability condition of Murnaghan [4], [5], [2]; and there is stability through the character functions of Orellana and Zabrocki [9]. More connected to our work, Church, Ellenberg, and Farb have shown, in their theory of FI-modules [11], that Schur functors satisfy a notion of S_n -stability for sufficiently large n. Some of their stability assertions are made precise by our identities. Section 4 contains two conjectures that would imply a simple expression for the graded character of the S_n coinvariants up to degree n (see also [3]). This expression would also imply the stability condition for the space of coinvariants, as indicated in [11].

2 S_n invariants in $S(\mathbb{C}^k \otimes \mathbb{C}^n)$

We associate to $S(\mathbb{C}^k \otimes \mathbb{C}^n)$ the polynomial ring

$$\mathcal{R} = \mathbb{C}[ix_1, \dots, ix_n : i = 1, \dots, k]$$

in k sets of variables, viewing $_ix_j$ as the generator $e_i \otimes e_j \in \mathbb{C}^k \otimes \mathbb{C}^n$. The diagonal action of $s \in S_n$ on \mathcal{R} is given by sending $_ix_j$ to $_ix_{s(j)}$ for all i. The polynomials of total degree d will be denoted by \mathcal{R}_d , and we let $\mathcal{R}_{\leq d}$ denote the space of polynomials of total degree at most d.

A basis for the S_n invariants of \mathcal{R} can be found by first choosing a monomial

$$x^{\alpha} = \prod_{i=1}^{k} \prod_{j=1}^{n} {}_{i} x_{j}^{i\alpha_{j}}$$

and taking the sum of the orbit $S_n(x^{\alpha})$. Note that the orbit corresponds to taking the sequence of vectors $(P^1(\alpha), \ldots, P^n(\alpha))$, with

$$P^i(\alpha) = ({}_1\alpha_i, \ldots, {}_k\alpha_i),$$

and permuting the vectors in all possible ways, since S_n acts diagonally. This means that a representative of the orbit is the multiset of sequences $P^{\alpha} = [P^1(\alpha), \ldots, P^n(\alpha)]$. In other words, we look at the set created by $P^1(\alpha), \ldots, P^n(\alpha)$, including multiplicities.

The dimension of invariants of degree equal to d is equal to the cardinality of the set $\mathcal{P}_d^{n,k}$ of multisets $[P^1, \ldots, P^n]$ such that each $P^i = (P_1^i, \ldots, P_k^i)$ is a sequence of k nonnegative integers satisfying

$$|P^1| + \dots + |P^n| = d$$
 with $|P^i| = P_1^i + \dots + P_k^i$.

The union of all $\mathcal{P}_d^{n,k}$ with $d \leq r$ will be denoted by $\mathcal{P}_{\leq r}^{n,k}$. This implies, for the space of polynomials with degree less than or equal to d, that

$$\dim \mathcal{R}^{S_n}_{\leq d} = |\mathcal{P}^{n,k}_{\leq d}|.$$

In [6], Weyl showed that a set of generators for the invariants is given by the polarized power sums

$$p_a = p_{(a_1,\dots,a_k)} = \sum_{i=1}^n {}_1 x_i^{a_1} \cdots {}_k x_i^{a_k}$$

with $0 < |a| = a_1 + \cdots + a_k \leq n$. In other words, every invariant can be written as a polynomial in the polarized power sums of degree at most n.

The number of monomials $p_{a^1} \cdots p_{a^r}$ we can construct whose total degree is less than or equal to n is the same as picking a multiset of sequences

$$\left[a^1 = (a^1_1, \dots, a^1_k), \dots, a^r = (a^r_1, \dots, a^r_k)\right],$$

each with a nonzero entry, such that $|a^1| + \cdots + |a^r| \leq n$. Note however that since each a^i has a nonzero entry, there are at most n of them. This is in bijection with the set $\mathcal{P}_{\leq n}^{n,k}$ by adding a sufficient number of sequences whose only entries are zeroes. This proves that the set

$$\{p_{a^1} \cdots p_{a^r} : |a^1| + \dots + |a^r| \le n\}$$

is not only a spanning set for the space of invariants of degree at most n, but also a basis. Surprisingly, there is a stronger independence satisfied by these monomials.

Theorem 1 The collection

$$\{p_{a^1} \cdots p_{a^r} : |a^1| + \dots + |a^r| \le n+1\}$$

is a basis for $\mathcal{R}^{S_n}_{\leq n+1}$.

Proof. We have shown that the set of monomials $p_{a^1} \cdots p_{a^r}$ of degree at most n are a basis for the invariants of degree at most n. For monomials of degree n+1, we have to show that the number of possible monomials in the polarized power sums of degree n+1 is equal to the cardinality of $\mathcal{P}_{n+1}^{n,k}$.

An element $[P^1, \ldots, P^n] \in \mathcal{P}_{n+1}^{n,k}$ falls into one of two cases:

- 1. $|P^i| \leq n$ for all *i*, or
- 2. there is an *i* for which $|P^i| = n + 1$.

In the first case, we have the monomial

$$p_{P^1}\cdots p_{P^n}$$
.

The convention here gives $p_{P^i} = 1$ for $P^i = (0, ..., 0)$. This accounts for all monomials in the polarized power sums with at most n factors. In the second

case, we may assume that $|P^1| = n + 1$ and $|P^i| = 0$ for i > 1. The number of such choices is the number of ways of choosing $P^1 = (a_1, \ldots, a_k)$ with

$$a_1 + \dots + a_k = n + 1,$$

giving the binomial coefficient

$$\binom{n+1+k-1}{k-1} = \binom{n+k}{k-1}.$$

On the other hand, a monomial in the polarized power sums $p_{a^1} \cdots p_{a^{n+1}}$ is of degree n + 1 precisely when $|a^i| = 1$ for all *i*. Let a_j be the number of a^i whose component *j* is nonzero. Then picking such a monomial is equivalent to choosing $a_1 + \cdots + a_k = n + 1$, meaning the number of such choices is again $\binom{n+k}{k-1}$. This shows that the set of monomials in the polarized power sums whose degree is at most *n* is a basis for the invariants of degree at most n+1.

We put the standard symmetric bilinear form $\langle (z_1, ..., z_k), (w_1..., w_k) \rangle = \sum z_i w_i$ on \mathbb{C}^k . On $\mathbb{C}^k \otimes \mathbb{C}^n$ we use the tensor product of the forms on \mathbb{C}^k and \mathbb{C}^n . This form is invariant under the action of S_n given by $\sigma(v \otimes w) = v \otimes \sigma w$. Then \mathcal{R}_d is isomorphic with $S^d(\mathbb{C}^k \otimes \mathbb{C}^n)$, using the form to identify $\mathbb{C}^k \otimes \mathbb{C}^n$ with $(\mathbb{C}^k \otimes \mathbb{C}^n)^*$. Thus we have

Corollary 2 For $r \leq m \leq n$,

$$\dim S^r (\mathbb{C}^k \otimes \mathbb{C}^m)^{S_m} = \dim S^r (\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n}$$

and

$$\sum_{n=1}^{\infty} \dim S^n (\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n} q^n = \prod_{n=1}^{\infty} \frac{1}{(1-q^r)^{\binom{r+k-1}{k-1}}}$$

Proof. In the above notation, we have that $\mathcal{R}_r^{S_n} = S^r (\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n}$ has dimension equal to $|\mathcal{P}_r^{n,k}|$. It is therefore sufficient to show that $|\mathcal{P}_r^{m,k}| = |\mathcal{P}_r^{n,k}|$. Since $n \geq m$, any element $[P^1, \ldots, P^n] \in \mathcal{P}_r^{n,k}$ must have at most $r \leq m$ nonzero P^i . Therefore, the map that takes $P \in \mathcal{P}_r^{m,k}$ and adds n - mzero vectors is not only injective, but also surjective. The two sets then have the same cardinality, and the equality holds.

We will now prove the sum product formula. We have shown that

dim
$$S^{n}(\mathbb{C}^{k} \otimes \mathbb{C}^{n})^{S_{n}} = |\{p_{a^{1}} \cdots p_{a^{r}} : |a^{1}| + \dots + |a^{r}| = n\}|,$$

which is the coefficient of q^n in the generating series

$$\prod_{a} \frac{1}{1-q^{|a|}},$$

where the product runs over all nonzero $a \in \mathbb{Z}_{\geq 0}^k$. This product is independent of n, and the factor $1/1 - q^r$ appears with multiplicity the number of a with |a| = r. The number of $a = (a_1, ..., a_k)$ satisfying $\sum a_i = r$ is given by the binomial coefficient $\binom{r+k-1}{k-1}$.

Recall that if $k, n \in \mathbb{Z}_{>0}$ then $\otimes^k \mathbb{C}^n$ is a module for $S_k \times GL(n, \mathbb{C})$ with $s \in S_k$ permuting the tensor factors and $g \in GL(n, \mathbb{C})$ acting by $\otimes^k g$. As a representation of $S_k \times GL(n, \mathbb{C})$, the module $\otimes^k \mathbb{C}^n$ decomposes according to Schur-Weyl duality (c.f. [8],9.1.1) as follows:

$$\otimes^{k} \mathbb{C}^{n} \cong \bigoplus_{\substack{\lambda \ \neg \ k} \\ \ell(\lambda) \le \min\{n, k\}} V^{\lambda} \otimes_{n} F^{\lambda}.$$

Here, V^{λ} is the Young module for S_k corresponding to the partition $\lambda = (\lambda_1 \geq \cdots \geq \lambda_r > 0)$ of size $|\lambda| = \lambda_1 + \cdots + \lambda_r = k$ with length $\ell(\lambda) = r$; and ${}_n F^{\lambda}$ is the Weyl module for $GL(n, \mathbb{C})$ corresponding to λ . Let $\epsilon(\lambda) = \sum_{i=1}^r \lambda_i \varepsilon_i$. Here, ε_i is the linear functional on the space of diagonal matrices

$$h = \begin{bmatrix} h_1 & 0 & \cdots & 0 \\ 0 & h_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & h_n \end{bmatrix}$$

given by $\varepsilon_i(h) = h_i$. The space \mathfrak{h} of diagonal $n \times n$ matrices is a Cartan subalgebra of $M_n(\mathbb{C})$, the Lie algebra of $GL(n, \mathbb{C})$; and ${}_nF^{\lambda}$ is the irreducible representation of $GL(n, \mathbb{C})$ with highest weight $\epsilon(\lambda)$ relative to the choice of positive roots $\varepsilon_i - \varepsilon_j$, $1 \leq i < j \leq n$.

Consider the subgroup $GL(n-1,\mathbb{C})$ of $GL(n,\mathbb{C})$ consisting of the matrices

$$\left\{ \left[\begin{array}{cc} g & 0 \\ 0 & 1 \end{array} \right] \mid g \in GL(n-1,\mathbb{C}) \right\}.$$

Let $\mu = \sum_{i=1}^{n} \mu_i \varepsilon_i$ with $\mu_i \in \mathbb{Z}_{\geq 0}$ and $\mu_i \geq \mu_{i+1}$ for all $i = 1, \ldots, n-1$ (that is, μ is a dominant integral weight). The branching theorem (c.f. [8], Theorem

8.1.1) implies that if we consider ${}_{n}F^{\mu}$ to be a $GL(n-1,\mathbb{C})$ module, then

$${}_{n}F^{\mu}\Big|_{GL(n-1,\mathbb{C})} \cong \bigoplus_{\substack{\nu = \sum_{i=1}^{n-1} \nu_{i}\varepsilon_{i} \\ \mu_{1} \ge \nu_{1} \ge \mu_{2} \ge \dots \ge \mu_{n-1} \ge \nu_{n-1} \ge \mu_{n}} {}_{n-1}F^{\nu}$$

This implies that if $\ell(\mu) \leq m < n$ and if $GL(m, \mathbb{C})$ is imbedded in $GL(n, \mathbb{C})$ as

$$\left\{ \left[\begin{array}{cc} g & 0 \\ 0 & I \end{array} \right] \middle| g \in GL(m, \mathbb{C}) \right\},\$$

then ${}_{m}F^{\mu}$ occurs as a $GL(m, \mathbb{C})$ submodule of ${}_{n}F^{\mu}$.

Let $E_{i,j} \in M_n(\mathbb{C})$ be the matrix with all zero entries, except for a 1 in position (i, j). One has an action of $M_n(\mathbb{C})$ as the Lie algebra of $GL(n, \mathbb{C})$) on any polynomial representation of $GL(n, \mathbb{C})$. The theorem of the highest weight implies that the weight space ${}_nF^{\mu}_{\mu}$ is one dimensional, and if $E_{i,j}$ is the $n \times n$ matrix with all entries 0 except for a 1 in position i, j, then the ξ weight space of ${}_nF^{\mu}$ is spanned by the elements

$$E_{i_1+1,i_1}\cdots E_{i_r+1,i_r}v$$

with v a nonzero element of ${}_{n}F^{\mu}_{\mu}$ and $\varepsilon(\mu) - \sum_{j=1}^{r} (\varepsilon_{i_j} - \varepsilon_{i_{j+1}}) = \xi$. Using this observation we have

Lemma 3 If $|\mu| \leq m \leq n$ and if ξ is a dominant weight of ${}_{n}F^{\mu}$ then $({}_{n}F^{\mu})_{\xi} = ({}_{m}F^{\mu})_{\xi}$.

Proof. Since $\dim_{(n}F^{\mu})_{\mu} = 1$ and $\ell(\mu) \leq m \leq n$, the assertion is true for μ . Since $|\mu| \leq m$, $\varepsilon(\mu) = \sum_{i=1}^{m} \mu_i \varepsilon_i$, and since ξ is dominant, $\xi = \sum_{i=1}^{m} \xi_i \varepsilon_i$. Thus if $\xi = \varepsilon(\mu) - \sum_{j=1}^{r} (\varepsilon_{i_j} - \varepsilon_{i_{j+1}})$ then the maximum of the i_j occurring in the expression $E_{i_1+1,i_1} \cdots E_{i_r+1,i_r} v$ is less than or equal to m-1. Thus $E_{i_1+1,i_1} \cdots E_{i_r+1,i_r} v \in (mF^{\mu})_{\xi}$.

Lemma 4 If $|\mu| \le m \le n$, then dim $({}_mF^{\mu})^{S_m} \ge \dim ({}_nF^{\mu})^{S_n}$.

Proof. If ξ is a weight of ${}_{n}F^{\mu}$, then there exists an element s of S_{n} such that $s\xi$ is dominant. Thus if $({}_{n}F^{\mu})_{\xi}$ is the corresponding weight space, there exists $s \in S_{n}$ such that $s\xi$ is a weight of ${}_{m}F^{\mu}$ and $s ({}_{n}F^{\mu})_{\xi} = ({}_{n}F^{\mu})_{s\xi} = ({}_{m}F^{\mu})_{s\xi}$. This implies that the span of $\{s ({}_{m}F^{\mu}) | s \in S_{n}\}$ is ${}_{n}F^{\mu}$. We use the standard

notation $\mathbb{C}S_n$ for the group algebra of S_n . We have a surjective S_n module homomorphism

$$\operatorname{Ind}_{S_m}^{S_n}({}_mF^{\mu}) = \mathbb{C}S_n \otimes_{\mathbb{C}S_m} ({}_mF^{\mu}) \to {}_nF^{\mu} \to 0.$$

Now Frobenius reciprocity implies that dim $(\mathbb{C}S_n \otimes_{\mathbb{C}S_m} (_mF^{\mu}))^{S_n} = \dim(_mF^{\mu})^{S_m}$. This implies the inequality.

The $GL(k, \mathbb{C})$ - $GL(n, \mathbb{C})$ duality theorem says that as a representation of $GL(k, \mathbb{C}) \times GL(n, \mathbb{C})$,

$$S^{r}(\mathbb{C}^{k} \otimes \mathbb{C}^{n}) \cong \bigoplus_{\substack{\lambda \ \dashv \ r \\ \ell(\lambda) \le \min\{n, k\} \text{ parts}}} {}_{k}F^{\lambda} \otimes {}_{n}F^{\lambda}.$$

In particular, this implies that if we consider the S_n action on $S^r(\mathbb{C}^k \otimes \mathbb{C}^n)$ coming from the restriction of the action from $GL(n, \mathbb{C})$, then

$$\dim S^r (\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n} = \bigoplus_{\substack{\lambda \to r \\ \ell(\lambda) \le \min\{n, k\} \text{ parts}}} \dim_k F^\lambda \otimes \dim_k (nF^\lambda)^{S_n}.$$

Theorem 5 If λ is a dominant integral weight of $GL(m, \mathbb{C})$ and $|\lambda| \leq m \leq n$, then

$$\dim \left({}_{m}F^{\lambda}\right)^{S_{m}} = \dim \left({}_{n}F^{\lambda}\right)^{S_{n}}.$$

Proof. We have seen in Corollary 2 that if $r \leq m \leq n$ then

$$\dim S^r (\mathbb{C}^k \otimes \mathbb{C}^m)^{S_m} = \dim S^r (\mathbb{C}^k \otimes \mathbb{C}^n)^{S_n}.$$

This implies that

$$\sum_{\substack{\lambda \dashv r \\ \ell(\lambda) \leq k}} \dim_k F^{\lambda} \otimes \left(\dim \left({}_m F^{\lambda} \right)^{S_m} - \dim \left({}_n F^{\lambda} \right)^{S_n} \right) = 0.$$

If k is the number of parts of λ then $\dim_k F^{\lambda} > 0$. The previous lemma now implies the differences are nonnegative, and thus every term in the sum is nonnegative. For equality to hold, the differences must be 0, implying the result.

In general if G and H are finite groups and V is an H module then

$$\operatorname{Ind}_{H}^{G}V = \mathbb{C}G \otimes_{\mathbb{C}H} V.$$

As an aside we note

Lemma 6 As an S_n module

$${}_{n}F^{\lambda} \cong \bigoplus_{\mu \ a \ dominant \ weight \ of \ {}_{n}F^{\lambda}} \operatorname{Ind}_{S_{n,\mu}}^{S_{n}}({}_{n}F^{\lambda})_{\mu}$$

where $S_{n,\mu} = \{ s \in S_n | s\mu = \mu \}.$

Proof. If μ is a weight of ${}_{n}F^{\lambda}$ then there exists $s \in S_{n}$ such that $s\mu$ is dominant. Thus since

$${}_{n}F^{\lambda} = \bigoplus_{\mu \text{ a weight of } nF^{\lambda}} ({}_{n}F^{\lambda})_{\mu}$$

we see that

$$_{n}F^{\lambda} = \sum_{s \in S_{n}, \ \mu \text{ dominant}} s(_{n}F^{\lambda})_{\mu}.$$

We note that if μ, ν are dominant weights and if $s \in S_n$ is such that $s\mu = \nu$, then $\mu = \nu$. Indeed, the theorem of the highest weight implies that $s\mu = \mu - Q$ with Q a non-negative integral combination of elements of the form $e_i - e_{i+1}$ with i = 1, ..., m. Thus in particular, $\langle \nu, Q \rangle \geq 0$. Also, since $s\mu = \nu$, this implies that $\mu = \nu + Q$. Then

$$\langle \mu, \mu \rangle = \langle \nu, \nu \rangle + 2 \langle \nu, Q \rangle + \langle Q, Q \rangle \ge \langle \nu, \nu \rangle + \langle Q, Q \rangle.$$

Since

$$\langle \nu, \nu \rangle = \langle s\mu, s\mu \rangle = \langle \mu, \mu \rangle,$$

this implies that Q = 0. Thus $\nu = \mu$.

We therefore have

$${}_{n}F^{\lambda} = \bigoplus_{\mu \text{ dominant } s \in S_{n}} S({}_{n}F^{\lambda})_{\mu}.$$

We assert that the S_n module $\sum_{s \in S_n} s({}_nF^{\lambda})_{\mu}$ is equivalent with $\operatorname{Ind}_{S_{n,\mu}}^{S_n}({}_nF^{\lambda})_{\mu}$. Indeed, if $s \in S_n$ and

$$s({}_nF^{\lambda})_{\mu} = ({}_nF^{\lambda})_{\mu},$$

then $s\mu = \mu$ and thus $s \in S_{n,\mu}$. Let $s_1, ..., s_r$ be a set of representatives for $S_n/S_{n,\mu}$. Then the elements $s_1\mu, ..., s_r\mu$ are distinct and

$$\sum_{s \in S_n} s({}_n F^{\lambda})_{\mu} = \bigoplus_{i=1}^{\prime} ({}_n F^{\lambda})_{s_i \mu} = \bigoplus_{i=1}^{\prime} s_i ({}_n F^{\lambda})_{\mu},$$

 \mathbf{SO}

$$\dim \sum_{s \in S_n} s({}_n F^{\lambda})_{\mu} = r \dim({}_n F^{\lambda})_{\mu} = \dim \operatorname{Ind}_{S_{n,\mu}}^{S_n}({}_n F^{\lambda})_{\mu}.$$

Since the map

$$\mathbb{C}S_n \otimes_{\mathbb{C}S_{n,\mu}} ({}_nF^{\lambda})_{\mu} \to \sum_{s \in S_n} s({}_nF^{\lambda})_{\mu}$$

given by

 $s\otimes v\mapsto sv$

is surjective, we conclude that the map is an equivalence. \blacksquare

Let $|\lambda| \leq m \leq n$. It will be important to see that the action of $S_{n-\ell(\mu)}$ contained in $S_{n,\mu} = S_{m_1} \times \cdots \times S_{m_r} \times S_{n-\ell(\mu)}$ is trivial on $({}_nF^{\lambda})_{\mu} = ({}_{\ell(\mu)}F^{\lambda})_{\mu}$. Let $l = \ell(\lambda)$. Embedding $GL(l, \mathbb{C})$ in $GL(n, \mathbb{C})$ as before,

$$\left\{ \left[\begin{array}{cc} g & 0 \\ 0 & I \end{array} \right] \middle| g \in GL(l, \mathbb{C}) \right\},\$$

and embedding $GL(n-l,\mathbb{C})$ by

$$\left\{ \left[\begin{array}{cc} I & 0 \\ 0 & g \end{array} \right] \mid g \in GL(n-l, \mathbb{C}) \right\},\$$

gives an embedding of $GL(l, \mathbb{C}) \times GL(n-l, \mathbb{C})$ in $GL(n, \mathbb{C})$. The cyclic space of the $(\lambda_1, \ldots, \lambda_l, 0, \ldots, 0)$ weight space under $GL(l, \mathbb{C})$ is equivalent to ${}_lF^{\lambda}$. As a representation of $GL(l, \mathbb{C}) \times GL(n-l, \mathbb{C})$, the space ${}_nF^{\lambda}$ splits into a direct sum equivalent to

$$\bigoplus_{\xi_1,\xi_2} m_{\lambda}(\xi_1,\xi_2)_l F^{\xi_1} \otimes_{n-l} F^{\xi_2}.$$

for some multiplicities $m_{\lambda}(\xi_1, \xi_2)$. Consider $m_{\lambda}(\lambda, \xi)$. If it is nonzero, then (λ, ξ) is a weight for ${}_{n}F^{\lambda}$, and so $|\lambda| + |\xi| = |\lambda|$. So $\xi = 0$. This means that ${}_{n-l}F^{\xi}$ is one dimensional, and that the cyclic space under $GL(l, \mathbb{C}) \times GL(n-l, \mathbb{C})$ of the $(\lambda_1, \ldots, \lambda_l, 0, \ldots, 0)$ weight space is equivalent to

$${}_{l}F^{\lambda} \otimes {}_{n-l}F^{0} = {}_{l}F^{\lambda} \otimes \mathbb{C},$$

with $GL(n-l,\mathbb{C})$ acting trivially on \mathbb{C} .

Lemma 7 Let $\mu \neq 0$ be a dominant weight of ${}_{n}F^{\lambda}$ and let $\ell(\mu)$ be the last index of μ which is positive. Then

$$\ell(\mu) \ge \ell(\lambda).$$

Proof. The theorem of highest weight implies the weights of ${}_{n}F^{\lambda}$, other than $(\lambda_{1}, \ldots, \lambda_{l}, 0, \ldots, 0)$ are of the form $(\lambda_{1}, \ldots, \lambda_{\ell(\lambda)}, 0, \ldots, 0) - \xi$ with $(\xi_{1}, \ldots, \xi_{n}) \neq 0$ satisfying

$$\xi_1 \ge 0, \ \xi_1 + \xi_2 \ge 0, \dots, \ \xi_1 + \dots + \xi_{n-1} \ge 0, \ \text{and} \ \xi_1 + \dots + \xi_n = 0.$$

This implies that if m is the last nonzero index of ξ , then $\xi_m < 0$. If $\ell(\mu) < \ell(\lambda)$, we must have $\xi_{\ell(\lambda)} = \lambda_{\ell(\lambda)}$, with $\lambda_{\ell(\lambda)} > 0$, since $\lambda_{\ell(\lambda)} - \xi_{\ell(\lambda)} = \mu_{\ell(\lambda)} = 0$. This means we must have $m > \ell(\lambda)$, contradicting $\ell(\mu) < \ell(\lambda)$.

Proposition 8 If μ is a dominant weight of ${}_{n}F^{\lambda}$, then $({}_{\ell(\mu)}F^{\lambda})_{\mu} = ({}_{n}F^{\lambda})_{\mu}$. And if we imbed $GL(n - \ell(\mu), \mathbb{C})$ in $GL(n, \mathbb{C})$ as

$$\left\{ \left[\begin{array}{cc} I & 0 \\ 0 & g \end{array} \right] \middle| g \in GL(n - \ell(\mu), \mathbb{C}) \right\},\$$

then $GL(n - \ell(\mu), \mathbb{C})$ acts trivially on $(\ell(\mu) F^{\lambda})_{\mu}$.

Proof. As is mentioned in the proof of Lemma 3 we have that every element of $({}_nF^{\lambda})_{\mu}$ is a linear combination of elements of the form

$$E_{j_1,i_1}\cdots E_{j_k,i_k}v$$

with $v \in {}_{n}F^{\lambda}_{(\lambda_{1},...,\lambda_{\ell(\lambda)},0,...,0)}$; furthermore,

$$(\lambda_1,\ldots,\lambda_l,0,\ldots,0) + \sum_{r=1}^k \epsilon_{j_r} - \epsilon_{i_r} = \mu.$$

If j_u is the maximum of the j_r , then $j_u \leq \ell(\mu)$ so that all the E_{j_r,i_r} are in the Lie algebra of $GL(\ell(\mu), \mathbb{C})$. This implies the first assertion. The second statement follows from the above observations: The cyclic space for $GL(\ell(\mu), \mathbb{C}) \times GL(n - \ell(\mu), \mathbb{C})$ of ${}_nF^{\lambda}_{(\lambda_1, \dots, \lambda_{\ell(\lambda)}, 0, \dots, 0)}$ is ${}_{\ell(\mu)}F^{\lambda} \otimes \mathbb{C}$ with $GL(n - \ell(\mu), \mathbb{C})$ acting on \mathbb{C} trivially.

3 Stability for general partitions

Let \mathcal{R} be, as above, the polynomial ring in k sets of n variables. For a given $L = (l_1, \ldots, l_k) \in \mathbb{Z}_{\geq 0}^k$, let V_L denote the space of homogeneous polynomials of degree l_i in the variables ix_1, \ldots, ix_n . Then we have the grading

$$\mathcal{R} = \bigoplus_{L \in \mathbb{N}^k} V_L$$

Each homogeneous component decomposes into irreducible S_n -representations

$$V_L = \bigoplus_{\mu \vdash n} m_L^{\mu} V^{\mu}.$$

with multiplicities given by

$$m_L^{\mu} = \dim \operatorname{Hom}_{S_n}(V^{\mu}, S^L(\mathbb{C}^k \otimes \mathbb{C}^n)).$$

Now Corollary 2 can be stated as follows: For $n \leq l$, we have

$$\sum_{|L|=n} m_L^{(n)} = \sum_{|L|=n} m_L^{(l)}$$

In this section, we are going to prove the following extension. Let e_1 denote the vector (1, 0, ...). If μ is a partition of n thought of as an element of $\mathbb{Z}^{l(\mu)}$,

$$\mu + re_1 = (\mu_1 + r, \mu_2, \dots, \mu_{\ell(\mu)})$$

is the partition obtained from μ by adding r to the first component.

Theorem 9 Let μ be a partition of n, and let $d \leq n - \mu_2$. Then

$$\sum_{l_1+\dots+l_k=d} q_1^{l_1} \cdots q_k^{l_k} m_{(l_1,\dots,l_k)}^{\mu} = \sum_{l_1+\dots+l_k=d} q_1^{l_1} \cdots q_k^{l_k} m_{(l_1,\dots,l_k)}^{\mu+re_1}.$$

Note that Corollary 2 is a special case of this theorem by taking $\mu = (n)$ so that $\mu_2 = 0$, choosing d = n, and setting all $q_i = 1$.

Before beginning the proof, we recall that the multigraded character of S_n on \mathcal{R} is given by the Schur function evaluation

$$\sum_{l_1,\dots,l_k} q_1^{l_1} \cdots q_k^{l_k} m_{(l_1,\dots,l_k)}^{\mu} = s_{\mu} \left[\frac{1}{(1-q_1)\cdots(1-q_k)} \right]$$

The right-hand side is the Schur function s_{μ} evaluated at all monomials $q^a = q_1^{a_1} \cdots q_k^{a_k}$. This corresponds to taking the sum over semistandard tableaux whose entries are monomials in q_1, \ldots, q_k , written in some total order. We will use a monomial order such that, the monomial 1 is the minimal element. **Proof of Theorem 9.** Every semistandard tableau of shape μ with monomial entries in q_1, \ldots, q_k , and total degree $d \leq n - \mu_2$ has at least μ_2 cells filled by only 1's, meaning every cell in the first row and below the second part must contain a 1. Otherwise there would be more than $n - \mu_2$ cells with at least one q_i , meaning the degree would be larger than $n - \mu_2$. Therefore, the set of semistandard tableaux of shape μ with degree less than or equal to $n - \mu_2$. The bijection, adds r cells with entry 1 to the first row. Here is an example:



Viewing ${}_{n}F^{\lambda}$ as an S_{n} module, one has

$${}_{n}F^{\lambda} \cong \bigoplus_{\mu \vdash n} \operatorname{Hom}_{S_{n}}(V^{\mu}, {}_{n}F^{\lambda}) \otimes V^{\mu}.$$

We will use the notation

$$g_{\mu}^{\lambda} = \dim \operatorname{Hom}_{S_n}(V^{\mu}, {}_nF^{\lambda}).$$

Lemma 10 Let $\mu \vdash n$ and $\eta \vdash l < n$. Then for any $r \ge 0$, we have

$$\dim \operatorname{Hom}_{S_n}(\operatorname{Ind}_{S_l \times S_{n-l}}^{S_n}(V^{\eta} \otimes 1_{n-l}), V^{\mu}) \\\leq \dim \operatorname{Hom}_{S_n}(\operatorname{Ind}_{S_l \times S_{n+r-l}}^{S_{n+r}}(V^{\eta} \otimes 1_{n+r-l}), V^{\mu+re_1}).$$

Proof. We will use the following Pieri rule: Let 1_n represent the trivial representation for S_n . Then

$$\operatorname{Ind}_{S_l \times S_{n-l}}^{S_n}(V^{\eta} \otimes 1_{n-l}) \cong \bigoplus_{\substack{\mu_1 \ge \eta_1 \ge \mu_2 \ge \cdots \ge \eta_{\ell(\mu)} \ge \mu_{\ell(\mu)+1} \\ |\mu| = n}} V^{\mu}.$$

Note that if V^{μ} appears as an irreducible, the inequalities are then also satisfied by

$$\mu_1 + r \ge \eta_1 \ge \mu_2 \ge \cdots \ge \eta_{\ell(\mu)} \ge \mu_{\ell(\mu)+1}$$

meaning $V^{\mu+re_1}$ appears in $\operatorname{Ind}_{S_l \times S_{n+r-l}}^{S_n}(V^{\eta} \otimes 1_{n+r-l})$.

Theorem 11 For $\mu \vdash n$, $|\lambda| \leq n - \mu_2$ and $r \geq 0$, we have

$$g_{\mu}^{\lambda} = g_{\mu+re_1}^{\lambda}.$$

Proof. We start with writing, as an S_n module,

$${}_{n}F^{\lambda} = \bigoplus_{\xi \text{ dominant}} \operatorname{Ind}_{S_{n,\mu}n}^{S_{n}}(F^{\lambda})_{\xi}.$$

Each subgroup can be written as $S_{n,\xi} = S_{\ell(\xi),\xi} \times S_{n-\ell(\xi)}$, where now, by Proposition 8, $S_{n-\ell(\xi)}$ acts trivially on ${}_{n}(F^{\lambda})_{\xi}$. We then have

$$\operatorname{Ind}_{S_{n,\xi}n}^{S_n}(F^{\lambda})_{\xi} = \operatorname{Ind}_{S_{\ell(\xi)} \times S_{n-\ell(\xi)}}^{S_n}(\operatorname{Ind}_{S_{\ell(\xi),\xi}n}^{S_{\ell(\xi)}}(F^{\lambda})_{\xi} \otimes 1_{n-\ell(\xi)})_{\xi}$$

and from Lemma 3,

$$\operatorname{Ind}_{S_{n+r,\xi}n+r}^{S_n}(F^{\lambda})_{\xi} = \operatorname{Ind}_{S_{\ell(\xi)} \times S_{n+r-\ell(\xi)}}^{S_n}(\operatorname{Ind}_{S_{\ell(\xi),\xi}n}^{S_{\ell(\xi)}}(F^{\lambda})_{\xi} \otimes 1_{n+r-\ell(\xi)}).$$

It follows from the previous lemma that

 $\dim \operatorname{Hom}_{S_n}(\operatorname{Ind}_{S_{n,\xi}n}^{S_n}(F^{\lambda})_{\xi}, V^{\mu}) \leq \dim \operatorname{Hom}_{S_n}(\operatorname{Ind}_{S_{n+r,\xi}n+r}^{S_n}(F^{\lambda})_{\xi}, V^{\mu+re_1}),$

and therefore

$$g_{\mu}^{\lambda} \le g_{\mu+re_1}^{\lambda}$$

From Theorem 9, for $|\lambda| = m \le n - \mu_2$ we must have the equality

$$\sum_{\substack{\lambda \to n \\ \ell(\lambda) \le k}} \dim_k F^\lambda \otimes \left(g_\mu^\lambda - g_{\mu+re_1}^\lambda\right) = 0.$$

This implies, as in Theorem 5, that $g_{\mu}^{\lambda} = g_{\mu+re_1}^{\lambda}$.

4 Some conjectures related to stability

In this section we will describe two conjectures that sharpen the results in the previous sections. The first we call the quasi-freeness conjecture.

Let $\mathcal{R}_{k,n}$ and p_{α} be as in section 2. Let $d(\alpha)$ denote the degree of p_{α} . Let $\mathcal{H}_{k,n}$ be the graded space of S_n harmonics which form an orthogonal complement to the ideal, $\mathcal{I}_{k,n}$, generated by the p_{α} with $d(\alpha) > 0$.

Conjecture 12 If $m \leq n$ and for each α with $d(\alpha) \leq m$ we have homogeneous $h_{\alpha} \in \mathcal{H}_{k,n}$ such that $d(\alpha) + \deg(h_{\alpha}) = m$, and if

$$\sum_{d(\alpha) \le m} h_{\alpha} p_{\alpha} = 0,$$

then $h_{\alpha} = 0$ for all α with $d(\alpha) \leq m$.

We call the condition in the statement of the Lemma "quasi-freeness". We will make a few observations which imply the conjecture for all k, n and $m \leq 3$.

Write the sum that is 0 out as

$$h_0 + \sum_{d(\alpha)=1} h_{\alpha} p_{\alpha} + \dots + \sum_{d(\alpha)=m-1} h_{\alpha} p_{\alpha} + \sum_{d(\alpha)=m} h_{\alpha} p_{\alpha}.$$

Then by definition of $\mathcal{H}_{k,n}$, $h_0 = 0$. Observing that the only S_n -invariants in $\mathcal{H}_{k,n}$ are the constants and since h_{α} is constant if $d(\alpha) = m$, we see that $\sum_{d(\alpha)=m} h_{\alpha} p_{\alpha}$ is an S_n -invariant. If m = 1 then we have

$$\sum_{d(\alpha)=1} h_{\alpha} p_{\alpha} = 0$$

and if m > 1 then

$$-\sum_{d(\alpha)=m}h_{\alpha}p_{\alpha}=\sum_{d(\alpha)=1}h_{\alpha}p_{\alpha}+\cdots+\sum_{d(\alpha)=m-1}h_{\alpha}p_{\alpha}.$$

Noting that if $d(\alpha) < m$ then deg $h_{\alpha} \geq 1$, one sees that the projection onto the invariants of the right hand side of the equation is 0. Since the degrees of the h_{α} on the left hand side are 0, the left hand side is invariant. This implies that both sides of the equality are 0 and, hence, in both cases $\sum_{d(\alpha)=m} h_{\alpha} p_{\alpha} = 0$. Since $m \leq n$, the p_{α} are linearly independent so $h_{\alpha} = 0$ for $d(\alpha) = m$. Notice that our observations now proves the conjecture for m = 1, so assume that $m \geq 2$ and set

$$L_i = \sum_{j=1}^n \frac{\partial}{\partial_i x_j}$$

Then, since $L_i h = 0$ for $h \in \mathcal{H}_{k,n}$ and L_i is a derivation, we have for each i

$$-\sum_{d(\alpha)=1}h_{\alpha}L_{i}p_{\alpha}=\sum_{d(\alpha)=2}h_{\alpha}L_{i}p_{\alpha}+\cdots+\sum_{d(\alpha)=m-1}h_{\alpha}L_{i}p_{\alpha}.$$

The left hand side of the equation is $-nh_{\alpha_i}$ with $p_{\alpha_i} = \sum_{j=1}^n ix_j$ and the right hand side is in the ideal, thus $h_{\alpha} = 0$ for $d(\alpha) = 1$. This proves the conjecture for m = 2. For m = 3 we are left with

$$\sum_{d(\alpha)=2} h_{\alpha} p_{\alpha} = 0.$$

In this case using the fact that a basis of the space of elements of degree 1 in $\mathcal{H}_{k,n}$ is given by

$$\{i x_j - i x_{j+1} | i = 1, \dots, k, j = 1, \dots, n-1\}$$

one can prove that this implies that the vanishing of the sum implies that $h_{\alpha} = 0$ for all the pertinent α . The argument is complicated and will not generalize so we omit it.

If $\phi(q)$ is a polynomial in q equal to $\sum \phi_i q^i$, then we set $\phi(q)_{\leq m} = \sum_{i \leq m} a_i q^i$. The conjecture has the following corollary: If h_i , (q) is the Hilbert series of \mathcal{H}_i , then

If $h_{k,n}(q)$ is the Hilbert series of $\mathcal{H}_{k,n}$ then

$$h_{k,n}(q)_{\leq n} = \left(\prod_{r=1}^{n} \frac{(1-q^r)^{\binom{r+k-1}{k-1}}}{(1-q)^k}\right)_{\leq n}.$$

In fact the conjecture implies a more precise formula. Setting $\mathcal{H}_{k,n}^r$ and $\mathcal{R}_{k,l}^r$ equal to the subspace of elements of $\mathcal{H}_{k,n}(\text{resp. } \mathcal{R}_{k,l}^r)$ homogenous of degree r and for $\lambda \dashv n$,

$$\mu_{\lambda,k,n}(r) = \dim \operatorname{Hom}_{S_n}(V^{\lambda}, \mathcal{H}^r_{k,n}).$$

and

$$\tau_{\lambda,k,n}(r) = \dim \operatorname{Hom}_{S_n}(V^{\lambda}, \mathcal{R}^r_{k,n}).$$

Then

$$\sum_{r=0}^{m} \mu_{\lambda,k,n}(r)q^r = \left(\prod_{r=1}^{m} (1-q^r)^{\binom{r+k-1}{k-1}} \sum_{r=1}^{m} \tau_{\lambda,k,n}(r)q^r\right)_{\leq m}.$$

for all $m \leq n$. The arguments above show that this formula is proved for $m \leq 3$. We note that it is not even obvious that the right side of these equations has positive coefficients. In the case of k = 2 this conjecture gives a simple interpretation of the Frobenius characteristic of the diagonal Harmonics of Garsia and Haiman to degree n [1]. This conjecture has been checked for k = 2, $n \leq 6$ and for $n = 7, m \leq 4$.

The second conjecture involves lead monomials of Gröbner bases. Order the variables as

$$x_1 \le y_1 \le \dots \le x_n \le y_n.$$

Let $G_{k,n}$ be the Gröbner basis for $\mathcal{I}_{k,n}$. Let $LG_{k,n}$ be the set of leading monomials of $G_{k,n}$ and let $LG_{k,n}^{\leq m}$ be the set of leading monomials of total degree at most m.

Conjecture 13 $LG_{k,n}^{\leq m} = LG_{k,m}^{\leq m}$.

This conjecture has also been checked for k = 2, $n \le 6$ and for k = 2, n = 7 and $m \le 4$.

Using the theory of Gröbner Bases one sees that this conjecture implies: If $h_{k,n}$ is the Hilbert series of the coinvariants of degree k for the k-fold diagonal action of S_n and if m < n, then

$$(h_{k,n})_{\leq m} = \left(\frac{h_{k,m}}{(1-q)^{k(n-m)}}\right)_{\leq m}$$

References

- A. M. Garsia and M. Haiman , A remarkable q, t-Catalan sequence and q-Lagrange inversion, J. Algebraic Combin. 5 (1996), no. 3, 191–244. MR 1394305
- [2] E. Briand, R. Orellana, and M. Rosas, The stability of the Kronecker product of Schur functions, J. Algebra 331 (2011), no. 1, 11–27.

- F. Bergeron, Multivariate diagonal coinvariant spaces for complex reflection groups, Adv. Math. 239 (2013), 97–108. MR 3045143
- [4] F. D. Murnaghan, The Analysis of the Kronecker Product of Irreducible Representations of the Symmetric Group, Amer. J. Math. 60 (1938), no. 3, 761–784.
- [5] _____, On the analysis of the Kronecker product of irreducible representations of S_n , Proc. Nat. Acad. Sci. U.S.A. **41** (1955), 515–518.
- [6] Hermann Weyl, The Classical Groups: Their Invariants and Representations, Princeton University Press, 1939.
- [7] M. Romero, Kronecker powers of harmonics, polynomial rings, and generalized principal evaluations, arXiv:2103.14195v2 (2021), .
- [8] R. Goodman, N. Wallach, Symmetry, Representations, and Invariants, Springer, 2009.
- [9] R. Orellana and M. Zabrocki, A Combinatorial Model for the Decomposition of Multivariate Polynomial Rings as S_n -modules, The Electronic Journal of Combinatorics **27** (2020), no. 3.
- [10] ____, Howe duality of the symmetric group and a multiset partition algebra, arXiv:2007.07370 (2020).
- [11] T. Church, J. Ellenberg, and B. Farb, FI-modules and stability for representations of symmetric groups, Duke Mathematical Journal 164 (2015), no. 9, 5025–5030.

Marino Romero University of California, San Diego Department of Mathematics *E-mail*: mar007@ucsd.edu

Nolan Wallach University of California, San Diego Department of Mathematics *E-mail*: nwallach@ucsd.edu