Global existence in reaction-diffusion systems with mass control under relaxed assumptions merely referring to cross-absorptive effects

 ${\begin{tabular}{l} Johannes\ Lankeit*\\ Leibniz\ Universit"at\ Hannover,\ Institut\ f"ur\ Angewandte\ Mathematik,\\ Welfengarten\ 1,\ 30167\ Hannover,\ Germany\\ \end{tabular}}$

Michael Winkler[†] Institut für Mathematik, Universität Paderborn, Warburger Str. 100, 33098 Paderborn, Germany

Abstract

We introduce a generalized concept of solutions for reaction-diffusion systems and prove their global existence. The only restriction on the reaction function beyond regularity, quasipositivity and mass control is special in that it merely controls the growth of cross-absorptive terms. The result covers nonlinear diffusion and does not rely on an entropy estimate.

Key words: Reaction-diffusion equation; renormalized solution; global existence **MSC (2020)**: 35K57, 35D99, 35K59

^{*}lankeit@ifam.uni-hannover.de

[†]michael.winkler@math.uni-paderborn.de

1 Introduction

Reaction-diffusion equations arise in various applications in chemistry and biology (cf. [19, Ch. 2] and form an important class of model problems in the study of systems of parabolic equations (see [23, Ch. 33]). Already at the stage of basic theories of solvability, a major challenge for the analysis of such systems consists in the presence of commonly superlinear source terms. While the possibility of blow up then is apparrent as long as suitably destabilizing reaction mechanisms are admitted (cf. e.g. [12]), even the requirement of dissipation of mass – which is sufficient to yield global existence and boundedness in the corresponding ODE systems – cannot preclude its occurrence, as impressively demonstrated by the counterexamples in [21]. Global classical solutions have, accordingly, been searched for and found under certain restrictive conditions.

In the context of boundary value problems for systems of the general form

$$\partial_t u_i = d_i \Delta u_i^{m_i} + f_i(u_1, ..., u_N), \qquad i \in \{1, ..., N\},$$
(1.1)

in the linear diffusion case when $m_1 = ... = m_N = 1$ such results on global smooth solvability cover settings where boundedness of the first among two components is a priori known (from a sign of f_1 , cf. [17]), where the diffusion coefficients are close to each other ([5, 2]), or where sources exhibit subquadratic growth ([4]), and recently ideas of [11] have successfully been extended to show global solvability for quadratic or slightly superquadratic reaction functions ([25, 3, 8]).

Another line of investigations pursues solutions in a weaker sense. Weak solutions can be constructed if L^1 -bounds for the reaction terms $f_i(u)$ are known ([18]), or if the reaction functions are at most quadratic ([20]). For nonlinear diffusion of porous medium type, the existence of weak solutions to (1.4) with Dirichlet boundary data is shown in [14] under the assumptions that

$$\begin{cases}
 f_i \in C^1([0,\infty)^N) & \text{is such that} \\
 f_i(s_1,...s_{i-1},0,s_{i+1},...,s_N) \ge 0 & \text{for all } (s_1,...,s_N) \in [0,\infty)^N \text{ and } i \in \{1,...,N\},
\end{cases}$$
(1.2)

that with some $K \geq 0$ and $a \in (0, \infty)^N$ we have

$$\sum_{i=1}^{N} a_i f_i(s_1, ..., s_N) \le K \cdot \left(\sum_{i=1}^{N} a_i s_i + 1\right) \quad \text{for all } (s_1, ..., s_N) \in [0, \infty)^N \text{ and } i \in \{1, ..., N\}, \quad (1.3)$$

and that either a priori L^1 -bounds for the reaction terms are known, or $f_i(s_1,...,s_N) \ge -C\left(\sum_{j=1}^n s_j^{\beta_j} + 1\right)$ with $\beta_i < m_i + 1$ for all $i \in \{1,...,N\}$ An analogous result has been achieved for the corresponding Neumann problem in [13] with a different proof and less restrictive conditions on the initial data.

The concept of renormalized solutions, that is, the idea that not u itself, but a transformed quantity $\rho(u)$ solves (a weak form) of the equation, makes it possible to bypass even further restrictions on the form of the system. This concept has been successfully introduced for the Boltzmann equation by DiPerna and Lions [7] and was employed for reaction diffusion equations with quadratic reaction functions and linear diffusion in [6]. The apparently most far-reaching application of this idea to reaction-diffusion systems (with linear diffusion) can be found in [9], where essentially no growth restriction on the f_i is needed, but where the reaction function is supposed to obey a certain entropy

condition. The term in the definition of solutions for whose treatment this entropy condition is essential arises from the choice of renormalization functions $\xi \colon [0,\infty)^N \to \mathbb{R}$ with compactly supported $D\xi$, which in particular depend on all solution components simultaneously.

Main results. In the present manuscript, we intend to introduce an approach by which it becomes possible to avoid any requirement of the latter type, and it turns out that this can in fact be achieved by resorting to separate renormalization functions for each component u_i . Thereby, our main result, as stated in Theorem 1.1 below, partially answers the open problem [9, p.585] to find a similar notion of solution without requiring an entropy condition.

Specifically, we shall be concerned with the Neumann problem

$$\begin{cases}
\partial_t u_i = d_i \Delta u_i^{m_i} + f_i(u_1, ..., u_N), & x \in \Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
\partial_\nu u_i^{m_i} = 0, & x \in \partial\Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
u_i(x, 0) = u_{0i}(x), & x \in \Omega, \ i \in \{1, ..., N\},
\end{cases}$$
(1.4)

under the assumptions that (1.2) and (1.3) hold, and that

$$\begin{cases}
f_i(s_1, ..., s_N) \ge -\phi_i(s_i) \cdot \left(\sum_{\substack{j \in \{1, ..., N\} \\ j \ne i}} s_j^{\beta_j} + 1\right) & \text{for all } (s_1, ..., s_N) \in [0, \infty)^N \text{ and } i \in \{1, ..., N\}, \\
\text{with some nonnegative } \phi_i \in C^1([0, \infty)) \text{ such that } \phi_i' > 0 \text{ on } (0, \infty) \text{ for } i \in \{1, ..., N\}, \text{ and some} \\
\beta_i > 0 \text{ such that } \beta_i < m_i + 1 \text{ for all } i \in \{1, ..., N\}.
\end{cases}$$
(1.5)

Here we recall that the quasipositivity condition in (1.2) is important in order to avoid negative concentrations, and that (1.3) is a slightly generalized mass dissipation condition, and includes some stoichiometric coefficients a. In addition to this, (1.5) signifies a growth condition for the negative parts of the reaction functions, where in the special case of linear diffusion, subquadratic growth is admissible. It is important to note, however, that this restriction only applies to the cross-absorptive effects: For $(f_i)_-$, the possible growth with respect to the i-th argument remains unrestricted.

As for the initial data in (1.4), throughout this paper we shall suppose that

$$u_{0i}, i \in \{1, ..., N\}$$
, is a nonnegative function from $L^r(\Omega)$ with some
$$\begin{cases} r \geq 1 & \text{if } n = 1, \\ r > 1 & \text{if } n = 2, \\ r \geq \frac{2n}{n+2} & \text{if } n \geq 3. \end{cases}$$
 (1.6)

Postponing the precise description of the solution concept to be pursued here to Section 2, let us introduce our main result obtained in this framework, and give a few examples of its application.

Theorem 1.1 let $n \ge 1$ and $N \ge 1$ and $\Omega \subset \mathbb{R}^n$ a bounded domain with smooth boundary, and suppose that $d_1, ..., d_N$ and $m_1, ..., m_N$ are positive, and that $f_1, ..., f_N$ belong to $C^1([0, \infty))$ and satisfy (1.3), (1.2) and (1.5) with some positive constant K. Then given any $u_{01}, ..., u_{0N}$ fulfilling (1.6), one can find nonnegative functions $u_i \in L^{m_i+1}_{loc}(\overline{\Omega} \times [0, \infty))$ such that $(u_1, ..., u_N)$ is a generalized solution of (1.4) in the sense of Definition 2.2.

Application #1. A first application of Theorem 1.1 addresses the system

$$\begin{cases}
\partial_{t}u_{i} = d_{i}\Delta u_{i}^{m_{i}} + (p_{i} - q_{i}) \cdot \left(k_{2} \prod_{j=1}^{N} u_{j}^{q_{j}} - k_{1} \prod_{j=1}^{N} u_{j}^{p_{j}}\right), & x \in \Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
\partial_{\nu} u_{i}^{m_{i}} = 0, & x \in \partial\Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
u_{i}(x, 0) = u_{0i}(x), & x \in \Omega, \ i \in \{1, ..., N\},
\end{cases}$$

$$(1.7)$$

which describes a general reversible reaction of the form

$$p_1\mathcal{U}_1 + p_2\mathcal{U}_2 + \ldots + p_n\mathcal{U}_n \stackrel{k_1}{\rightleftharpoons} q_1\mathcal{U}_1 + q_2\mathcal{U}_2 + \ldots + q_n\mathcal{U}_n,$$

and for which we obtain the following.

Proposition 1.2 Let $N \ge 2$, and suppose that $k_1 > 0$ and $k_2 > 0$, and that for $i \in \{1, ..., N\}$, $d_i > 0$, $m_i > 0$, $p_i \ge 1$ and $q_i \ge 1$ are such that for some $a \in (0, \infty)^N$

$$\sum_{i=1}^{N} a_i p_i = \sum_{i=1}^{N} a_i q_i, \tag{1.8}$$

and that

$$\sum_{\substack{j \in \{1,...,N\} \\ j \neq i}} \frac{p_j}{m_j + 1} < 1 \quad \text{for all } i \in \{1,...,N\} \text{ such that } p_i > q_i$$
 (1.9)

as well as

$$\sum_{\substack{j \in \{1,...,N\} \\ j \neq i}} \frac{q_j}{m_j + 1} < 1 \quad \text{for all } i \in \{1,...,N\} \text{ such that } p_i < q_i.$$
 (1.10)

Then for any choice of $u_{01}, ..., u_{0n}$ complying with (1.6), the problem (1.7) admits a generalized solution in the sense of Definition 2.2.

PROOF of Proposition 1.2. Writing $f_i(s_1,...,s_N) := (p_i - q_i) \cdot \left(k_2 \prod_{j=1}^N s_j^{q_j} - k_1 \prod_{j=1}^N s_j^{p_j}\right)$ for $i \in \{1,...,N\}$ and $(s_1,...,s_N) \in [0,\infty)^N$, we see that (1.2) is fulfilled and (1.3) follows since $\sum_{j=1}^N a_i f_i \equiv 0$ due to (1.8). Moreover, if e.g. $i \in \{1,...,N\}$ is such that $p_i > q_i$, then (1.9) enables us to pick numbers $\theta_j > 1$, $j \in \{1,...,N\} \setminus \{i\}$, such that $p_j\theta_j < m_j + 1$ for all $j \in \{1,...,N\} \setminus \{i\}$ and $\sum_{j\neq i} \frac{1}{\theta_j} < 1$. An application of Young's inequality thus shows that for any such i,

$$f_{i}(s_{1},...,s_{N}) \geq -(p_{i}-q_{i})k_{1}s_{i}^{p_{i}} \cdot \prod_{j \neq i} s_{j}^{p_{j}}$$

$$\geq -(p_{i}-q_{i})k_{1}s_{i}^{p_{i}} \cdot \left(\sum_{j \neq i} s_{j}^{p_{j}\theta_{j}} + 1\right) \quad \text{for all } (s_{1},...,s_{N}) \in [0,\infty)^{N},$$

and complementing this by a similar reasoning for all $i \in \{1, ..., N\}$ for which $p_i < q_i$, we readily obtain that holds and Theorem 1.1 becomes applicable so as to yield the claim.

Proposition 1.2 corresponds to [14, Remark 2.10], where the existence of weak solutions is proved. The main difference is that there the summation in (1.9) and (1.10) extends over all $j \in \{1, ... N\}$.

For linear diffusion, weak solutions of (1.7) have been found in [22] if the reaction functions grow at most quadratically or if the diffusion coefficients are sufficiently close to each other. The same article also deals with their exponential convergence.

Application #2. A second application of our general theory is concerned with the variant of (1.7) given by

$$\begin{cases}
\partial_{t}u_{1} = d_{1}\Delta u_{1}^{m_{1}} + u_{1}^{\beta_{1}}g_{2}(u_{2}) - g_{1}(u_{1})u_{2}^{\beta_{2}}, & x \in \Omega, \ t > 0, \\
\partial_{t}u_{2} = d_{2}\Delta u_{2}^{m_{2}} - u_{1}^{\beta_{1}}g_{2}(u_{2}) + \lambda g_{1}(u_{1})u_{2}^{\beta_{2}}, & x \in \Omega, \ t > 0, \\
\partial_{\nu}u_{1}^{m_{1}} = \partial_{\nu}u_{2}^{m_{2}} = 0, & x \in \partial\Omega, \ t > 0, \\
u_{1}(x, 0) = u_{01}(x), \quad u_{2}(x, 0) = u_{02}(x), & x \in \Omega,
\end{cases}$$
(1.11)

and underlines the mildness of the assumptions in Theorem 1.1 by admitting widely arbitrary growth of the main ingredients g_1 and g_2 appearing herein:

Proposition 1.3 Let $d_1 > 0, d_2 > 0, m_1 > 0$, $m_2 > 0$ and $\lambda \in [0,1]$, let $\beta_1 \in [1, m_1 + 1)$ and $\beta_2 \in [1, m_2 + 1)$, and let $g_1 \in C^1([0, \infty))$ and $g_2 \in C^1([0, \infty))$ be such that $g_1(0) = g_2(0) = 0$ and that g_1 and g_2 are positive on $(0, \infty)$. Then for any pair (u_{01}, u_{02}) satisfying (1.6), there exists a generalized solution of (1.11) in the spirit of Definition 2.2.

PROOF of Proposition 1.3. Taking any nonnegative $\phi_i \in C^1([0,\infty))$ such that $\phi_i' > 0$ and $\phi_i \geq g_i$ on $(0,\infty)$ for $i \in \{1,2\}$, one can readily verify that for

$$f_1(s_1, s_2) := s_1^{\beta_1} g_2(s_2) - g_1(s_1) s_2^{\beta_2} \quad \text{and} \quad f_2(s_1, s_2) := -s_1^{\beta_1} g_2(s_2) + \lambda g_1(s_1) s_2^{\beta_2}, \qquad (s_1, s_2) \in [0, \infty)^2,$$

we have

$$f_1(s_1, s_2) + f_2(s_1, s_2) = -(1 - \lambda)g_1(s_1)s_2^{\beta_2} \le 0$$
 for all $(s_1, s_2) \in [0, \infty)^2$

as well as

$$f_1(s_1, s_2) > -q_1(s_1)s_2^{\beta_2} > -\phi_1(s_1)s_2^{\beta_2}$$
 for all $(s_1, s_2) \in [0, \infty)^2$

and, similarly,

$$f_s(s_1, s_2) \ge -\phi_2(s_2)s_1^{\beta_1}$$
 for all $(s_1, s_2) \in [0, \infty)^2$.

The assumptions $\beta_i < m_i + 1$, $i \in \{1,2\}$, therefore warrant applicability of Theorem 1.1 with the intended result.

Let us remark that since in Proposition 1.3 not only $f_1 + f_2 \leq 0$, but also $\lambda f_1(s_1, s_2) + f_2(s_1, s_2) = (\lambda - 1)s_1^{\beta_1}g_2(s_2) \leq 0$ for $(s_1, s_2) \in [0, \infty)^2$, [14, Cor. 2.11] could be applied to the variant of (1.11)

involving homogeneous Dirichlet boundary conditions (cf. [14, Remark 2.12]) so as to yield weak solutions for any L^1 -initial data; said corollary, however, requires that $m_1, m_2 < 2$.

Application #3. We shall next briefly address

$$\begin{cases}
\partial_{t}u_{1} = d_{1}\Delta u_{1}^{m_{1}} + k_{2}u_{1}^{q_{1}}u_{2}^{q_{2}} - k_{1}u_{1}^{p_{1}}u_{2}^{p_{2}}, & x \in \Omega, \ t > 0, \\
\partial_{t}u_{2} = d_{2}\Delta u_{2}^{m_{2}} - k_{2}u_{1}^{q_{1}}u_{2}^{q_{2}} + k_{1}u_{1}^{p_{1}}u_{2}^{p_{2}}, & x \in \Omega, \ t > 0, \\
\partial_{\nu}u_{1}^{m_{1}} = \partial_{\nu}u_{2}^{m_{2}} = 0, & x \in \partial\Omega, \ t > 0, \\
u_{1}(x, 0) = u_{01}(x), \quad u_{2}(x, 0) = u_{02}(x), & x \in \Omega,
\end{cases}$$
(1.12)

for which without imposing any smallness condition on q_2 nor p_1 we obtain the following.

Corollary 1.4 Let k_1, k_2, d_1, d_2, m_1 and m_2 be positive, and let $p_1 \ge 1, p_2 \ge 1, q_1 \ge 1$ and $q_2 \ge 1$ be such that

$$q_1 < m_1 + 1$$
 and $p_2 < m_2 + 1$.

Then for all (u_{01}, u_{02}) fulfilling (1.6), the problem (1.12) possesses a generalized solution in the sense of Definition 2.2.

PROOF of Corollary 1.4. With $g_1(s) = k_1 s^{p_1}$, $g_2(s) = k_2 s^{q_2}$, $\beta_1 = q_1$, $\beta_2 = p_2$, this immediately results from Proposition 1.3.

Application #4. As final example, let us consider the generalized Lotka-Volterra system

$$\begin{cases}
\partial_{t}u_{i} = d_{i}\Delta u_{i}^{m_{i}} + \gamma_{i}u_{i} + \sum_{j=1}^{N} a_{ij}u_{j}^{\beta_{ij}}u_{i}^{\beta_{ji}}, & x \in \Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
\partial_{\nu}u_{i}^{m_{i}} = 0, & x \in \partial\Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
u_{i}(x, 0) = u_{0i}(x), & x \in \Omega, \ i \in \{1, ..., N\},
\end{cases}$$
(1.13)

which does not obey the typical entropy condition (that is required for the renormalized solutions in [9] and for classical solvability e.g. in [25]). In [8], global classical solutions are shown to exist for the classical Lotka-Volterra system ($\beta_{ij} = 1$ for all i, j) with linear diffusion. If $m_i = 1$ for all i and $\beta_{ij} = \beta$ for all i, j, then the result of [8] covers $2\beta < 2 + \varepsilon$ (for sufficiently small $\varepsilon \in (0, 1)$, see [8, Step 1, (7)]).

Within the generalized solvability framework considered here, the following consequence of Theorem 1.1 shows that here actually the entire range $\beta < 2$ can be exhausted.

Proposition 1.5 Let $N \geq 2$, $d_i > 0$, $m_i > 0$ and $\gamma_i \in \mathbb{R}$ for $i \in \{1, ..., N\}$, and suppose that for $i, j \in \{1, ..., N\}$ the numbers $a_{ij} \in \mathbb{R}$ and $\beta_{ij} > 0$ are such that $a_{ij} + a_{ji} \leq 0$, and that

if
$$a_{ij} < 0$$
, then $\beta_{ij} < m_i + 1$. (1.14)

Then for all initial data $u_{01}, ..., u_{0n}$ as in (1.6), (1.13) has a generalized solution in the sense of Definition 2.2.

PROOF of Proposition 1.5. With $f_i(s) = \gamma_i s_i + \sum_{j=1}^N a_{ij} s_j^{\beta_{ij}} s_i^{\beta_{ji}}$, $i \in \{1, \dots, N\}$, (1.2) is clearly satisfied. As

$$\sum_{i=1}^{N} f_i(s) = \sum_{i=1}^{N} \gamma_i s_i + \sum_{i=1}^{N} \sum_{j=1}^{N} a_{ij} s_j^{\beta_{ij}} s_i^{\beta_{ji}} = \sum_{i=1}^{N} \gamma_i s_i + \sum_{i \neq j} (a_{ij} s_j^{\beta_{ij}} s_i^{\beta_{ji}} + a_{ji} s_i^{\beta_{ji}} s_j^{\beta_{ij}}) \le \max_i \gamma_i \sum_{i=1}^{N} s_i,$$

also (1.3) holds. Finally,

$$f_{i}(s) \geq \gamma_{i} s_{i} + \sum_{\substack{j \in \{1, \dots, N\} \\ a_{ij} < 0}} a_{ij} s_{j}^{\beta_{ij}} s_{i}^{\beta_{ji}} \geq -\left(|\gamma_{i}| + \sum_{\substack{j \in \{1, \dots, N\} \\ a_{ij} < 0}} |a_{ij}| s_{j}^{\beta_{ij}}\right) \left(s_{i} + \sum_{j=1}^{N} s_{i}^{\beta_{ji}}\right)$$

$$\geq -\phi_{i}(s) \left(1 + \sum_{\substack{j \in \{1, \dots, N\} \\ a_{ij} < 0}} s_{j}^{\beta_{ij}}\right)$$

for all $s = (s_1, ..., s_N) \in [0, \infty)^N$ if we set $\phi(s) = \max\{|\gamma_i|, |a_{ij}| \mid j \in \{1, ..., N\}\} \cdot \sum_{j=1}^N s_i^{\beta_{ji}}$ for any such s, so that, according to (1.14), (1.5) is fulfilled and Theorem 1.1 is applicable.

2 Solution concept

The first step toward the design of our solution concept is concerned with an appropriate supersolution feature required in each of the equations making up (1.4):

Definition 2.1 Suppose that for $i \in \{1,...,N\}$, $u_{0i}: \Omega \to \mathbb{R}$ and $u_i: \Omega \times (0,\infty) \to \mathbb{R}$ are measurable and nonnegative. Then $(u_1,...,u_N)$ will be called a renormalized supersolution of (1.4) if for every $\rho \in C^{\infty}([0,\infty))$ fulfilling $\rho' \in C^{\infty}_0([0,\infty))$, $\rho' \leq 0$ and $\rho'' \geq 0$, with

$$P_i^{(1)}(s) := \int_0^s \sigma^{\frac{m_i - 1}{2}} \sqrt{\rho''(\sigma)} d\sigma \quad and \quad P_i^{(2)}(s) := \int_0^s \sigma^{m_i - 1} \rho'(\sigma) d\sigma, \qquad s \ge 0, \ i \in \{1, ..., N\}, \quad (2.1)$$

we have

$$\rho'(u_i)f_i(u_1,...,u_N) \in L^1_{loc}(\overline{\Omega} \times [0,\infty)) \quad and \quad \nabla P_i^{(1)}(u_i) \in L^2_{loc}(\overline{\Omega} \times [0,\infty); \mathbb{R}^n) \quad for \ all \ i \in \{1,...,N\},$$

$$(2.2)$$

and if moreover

$$-\int_{0}^{\infty} \int_{\Omega} \rho(u_{i})\varphi_{t} - \int_{\Omega} \rho(u_{0i})\varphi(\cdot,0) \leq -d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} \varphi |\nabla P_{i}^{(1)}(u_{i})|^{2} + d_{1}m_{i} \int_{0}^{\infty} \int_{\Omega} P_{i}^{(2)}(u_{i})\Delta\varphi + \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i})f_{i}(u_{1},...,u_{N})\varphi$$

$$(2.3)$$

for all $i \in \{1, ..., N\}$ and each nonnegative $\varphi \in C_0^{\infty}(\overline{\Omega} \times [0, \infty))$ fulfilling $\partial_{\nu} \varphi = 0$ on $\partial \Omega \times (0, \infty)$.

Remark. i) In the above situation, both integrals on the left of (2.3) as well as the second integral on the right-hand side therein exist due to the readily verified fact that ρ and $P_i^{(2)}$, $i \in \{1, ..., N\}$, are bounded on $[0, \infty)$.

ii) The supersolution property in [14, Prop. 3.6] is obtained upon the choice of $\rho(x) = x$ (inadmissible in Definition 2.1), integration by parts in the integral involving $P_i^{(2)}$ (and addition of a corresponding integrability requirement) and, finally, exchange of \leq by \geq (resulting from the change of sign of ρ').

As discussed in several previous related approaches toward generalized solvability on the basis of supersolution features of the above flavor ([28], [27]), supplementing Definition 2.1 by a mere requirement on mass control is already sufficient to create a notion of solvability which within classes of suitably smooth functions indeed reduces to classical ones (see, e.g, [16] and [26] for detailed reasonings in this regard):

Definition 2.2 By a generalized solution of (1.4) we mean a vector $(u_1, ..., u_N)$ of nonnegative measurable functions on $\Omega \times (0, \infty)$ such that $(u_1, ..., u_N)$ is a renormalized supersolution of (1.4) in the sense of Definition 2.1, that with some $a \in (0, \infty)^N$

$$u_1, ..., u_N$$
 and
$$\sum_{i=1}^N a_i f_i(u_1, ..., u_N) \quad belong \text{ to } L^1_{loc}(\overline{\Omega} \times [0, \infty)), \tag{2.4}$$

and that

$$\int_{\Omega} \left(\sum_{i=1}^{N} a_i u_i(\cdot, t) \right) \le \int_{\Omega} \left(\sum_{i=1}^{N} a_i u_{0i} \right) + \int_{0}^{t} \int_{\Omega} \left(\sum_{i=1}^{N} a_i f_i(u_1, ..., u_N) \right) \quad \text{for a.e. } t > 0.$$
 (2.5)

3 Approximate systems

In order to construct such solutions through an essentially standard type of approximation, for $\varepsilon \in (0,1)$ we consider

$$\begin{cases}
\partial_t u_{i\varepsilon} = d_i \Delta (u_{i\varepsilon} + \varepsilon)^{m_i} + \frac{f_i(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^N f_i(u_{1\varepsilon}, ..., u_{N\varepsilon})}, & x \in \Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
\partial_{\nu} u_{i\varepsilon} = 0, & x \in \partial \Omega, \ t > 0, \ i \in \{1, ..., N\}, \\
u_{i\varepsilon}(x, 0) = u_{0i\varepsilon}(x), & x \in \Omega, \ i \in \{1, ..., N\},
\end{cases}$$
(3.1)

where

$$\begin{cases} (u_{0i\varepsilon})_{\varepsilon\in(0,1)} \subset C^1(\overline{\Omega}) & \text{is such that } u_{0i\varepsilon} \geq 0 \text{ for all } i \in \{1,...,N\}, \text{ that} \\ u_{0i\varepsilon} \to u_{0i} & \text{in } L^1(\Omega) \text{ and a.e. in } \Omega \text{ as } \varepsilon \searrow 0 \text{ for all } i \in \{1,...,N\}, \text{ and that} \\ \sup_{\varepsilon\in(0,1)} \|u_{0i\varepsilon}\|_{L^r(\Omega)} < \infty \end{cases}$$
(3.2)

with $r \ge 1$ taken from (1.6).

Due to boundedness of the reaction term therein and nondegeneracy of the diffusion, by [1, Theorems 14.4 and 14.6] and [15, Chapter V], for each fixed $\varepsilon \in (0,1)$ the problem (3.1) indeed admits a global classical solution.

General assumption. Throughout the sequel, we shall suppose that the assumptions of Theorem 1.1 and (3.2) are satisfied, and given $\varepsilon \in (0,1)$ we let $(u_{1\varepsilon},...,u_{N\varepsilon})$ denote the global classical solution of (3.1).

The following basic observation concerning L^1 -boundedness of these solutions is a fairly immediate consequence of (1.3).

Lemma 3.1 For all T > 0 there exists C(T) > 0 such that for all $i \in \{1, ..., N\}$ and any $\varepsilon \in (0, 1)$ we have

$$||u_{i\varepsilon}(\cdot,t)||_{L^1(\Omega)} \le C(T)$$
 for all $t \in (0,T)$. (3.3)

PROOF. By integrating in (3.1), we see that since $\partial_{\nu}u_{i\varepsilon}=0$ on $\partial\Omega\times(0,\infty)$, due to (1.3) we have

$$\frac{d}{dt} \sum_{i=1}^{N} a_{i} \int_{\Omega} u_{i\varepsilon} = \int_{\Omega} \frac{1}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \sum_{i=1}^{N} a_{i} f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})$$

$$\leq \int_{\Omega} \frac{1}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot K \left\{ \sum_{i=1}^{N} a_{i} u_{i\varepsilon} + 1 \right\}$$

$$\leq K \sum_{j=1}^{N} a_{i} \int_{\Omega} u_{i\varepsilon} + K |\Omega| \quad \text{for all } t > 0 \text{ and } \varepsilon \in (0, 1),$$

because $u_{i\varepsilon} \geq 0$ for all $i \in \{1,...,N\}$. By an ODE comparison, this shows that

$$\sum_{i=1}^{N} a_i \int_{\Omega} u_{i\varepsilon} \le \left\{ \sum_{i=1}^{N} a_i \int_{\Omega} u_{0i\varepsilon} + |\Omega| \right\} \cdot e^{Kt} \quad \text{for all } t > 0 \text{ and } \varepsilon \in (0,1)$$

and hence, again by nonnegativity of $u_{i\varepsilon}$ for $i \in \{1, ..., N\}$, establishes (3.3) due to (3.2).

The following estimate rests on a duality-based reasoning inspired by a corresponding argument from [14]. An important difference is given by the change of boundary conditions: Where [14] dealt with Dirichlet boundary data and thus could use the solution of Poisson's equation as test function, the non-invertibility of the Laplacian with homogeneous Neumann boundary data leads us to employ the solution of a Helmholtz equation instead. An alternative approach of working with the Neumann Laplacian after substracting the mean value has been followed in [13], but it seems unclear how far strategies of this type can be applied so as to successfully cover the present setting.

Lemma 3.2 For all T > 0 there exists C(T) > 0 such that for all $i \in \{1, ..., N\}$,

$$\int_{0}^{T} \int_{\Omega} u_{i\varepsilon}^{m_{i}+1} \le C(T) \quad \text{for all } \varepsilon \in (0,1).$$
 (3.4)

PROOF. Following [14, Proof of Theorem 2.7], we first observe that according to (3.1) and (1.3), writing

$$w_{\varepsilon}(\cdot,t) := e^{-Kt} \cdot \sum_{i=1}^{N} a_i u_{i\varepsilon}(\cdot,t)$$
 and $z_{\varepsilon}(\cdot,t) := \int_0^t e^{-Ks} \cdot \left\{ \sum_{i=1}^{N} a_i d_i (u_{i\varepsilon} + \varepsilon)^{m_i} \right\}$

for $\varepsilon \in (0,1)$ and $t \geq 0$, we have

$$\partial_{t} w_{\varepsilon} = e^{-Kt} \Delta \left\{ \sum_{i=1}^{N} a_{i} d_{i} (u_{i\varepsilon} + \varepsilon)^{m_{i}} \right\} + e^{-Kt} \sum_{i=1}^{N} \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} - Ke^{-Kt} \sum_{i=1}^{N} a_{i} u_{i\varepsilon}$$

$$\leq e^{-Kt} \Delta \left\{ \sum_{i=1}^{N} a_{i} d_{i} (u_{i\varepsilon} + \varepsilon)^{m_{i}} \right\} + Ke^{-Kt} \quad \text{in } \Omega \times (0, \infty) \quad \text{for all } \varepsilon \in (0, 1)$$

and hence

$$w_{\varepsilon}(\cdot, t) \le w_{\varepsilon}(\cdot, 0) + \Delta z_{\varepsilon}(\cdot, t) + 1 \quad \text{in } \Omega \quad \text{ for all } t > 0 \text{ and } \varepsilon \in (0, 1),$$
 (3.5)

because $\int_0^t Ke^{-Ks}ds = 1 - e^{-Ks} \le 1$ for all t > 0. Upon multiplication by $\partial_t z_{\varepsilon} \ge 0$ and integration over $\Omega \times (0,T)$ for T > 0, as in [14] we obtain that since $z_{\varepsilon}(\cdot,0) \equiv 0$,

$$\int_{0}^{T} \int_{\Omega} w_{\varepsilon} \partial_{t} z_{\varepsilon} \leq \int_{\Omega} w_{\varepsilon}(\cdot, 0) z_{\varepsilon}(\cdot, T) - \frac{1}{2} \int_{\Omega} |\nabla z_{\varepsilon}(\cdot, T)|^{2} + \int_{\Omega} z_{\varepsilon}(\cdot, T) \\
\leq \int_{\Omega} \left(w_{\varepsilon}(\cdot, 0) + 1 \right) z_{\varepsilon}(\cdot, T) \quad \text{for all } \varepsilon \in (0, 1), \tag{3.6}$$

where the right-hand side is now estimated in a way slightly deviating from that in [14] due to the different boundary conditions considered here. In fact, by nonnegativity of w_{ε} the inequality in (3.5) implies that for each fixed T > 0 and arbitrary $\varepsilon \in (0, 1)$,

$$\begin{cases} -\Delta z_{\varepsilon}(\cdot, T) + z_{\varepsilon}(\cdot, T) \leq w_{\varepsilon}(\cdot, 0) + 1 + z_{\varepsilon}(\cdot, T) & \text{in } \Omega \\ \partial_{\nu} z_{\varepsilon}(\cdot, T) = 0 & \text{on } \partial\Omega, \end{cases}$$
 and

so that since the Helmholtz operator $-\Delta + 1$ admits a comparison principle under homogeneous Neumann boundary conditions, we obtain that

$$z_{\varepsilon}(\cdot, T) \le \overline{z_{\varepsilon}} \quad \text{in } \Omega,$$
 (3.7)

where $\overline{z_{\varepsilon}}$ denotes the solution of

$$\begin{cases} -\Delta \overline{z_{\varepsilon}} + \overline{z_{\varepsilon}} = w_{\varepsilon}(\cdot, 0) + 1 + z_{\varepsilon}(\cdot T) & \text{in } \Omega, \\ \partial_{\nu} \overline{z_{\varepsilon}} = 0 & \text{on } \partial \Omega. \end{cases}$$

Now without loss of generality assuming that the number $r \geq 1$ in (1.6) satisfies r < 2, we take $r' \in (2, \infty]$ such that $\frac{1}{r} + \frac{1}{r'} = 1$, and employ a Sobolev embedding theorem and elliptic regularity theory ([10]) to find $c_1 > 0$ and $c_2 > 0$ such that for all $\varepsilon \in (0, 1)$,

$$\|\overline{z_{\varepsilon}}\|_{L^{r'}(\Omega)} \le c_1 \|\overline{z_{\varepsilon}}\|_{W^{2,r}(\Omega)} \le c_2 \|w_{\varepsilon}(\cdot,0) + 1 + z_{\varepsilon}(\cdot,T)\|_{L^r(\Omega)}$$

$$\le c_2 \|w_{\varepsilon}(\cdot,0) + 1\|_{L^r(\Omega)} + c_2 \|z_{\varepsilon}(\cdot,T)\|_{L^r(\Omega)}, \tag{3.8}$$

because the restrictions on r in (1.6) warrant that $2 - \frac{n}{r} \ge -\frac{n}{r'}$ if $n \ge 3$, and that $2 - \frac{n}{r} > 0$ if $n \le 2$. Here we note that according to Lemma 3.1 we know that if we let $\theta \in (0,1)$ small enough such that $\theta m_i \leq 1$ for all $i \in \{1,...,N\}$, then we can find $c_3(T) > 0$ fulfilling

$$\begin{split} \int_{\Omega} z_{\varepsilon}^{\theta}(\cdot,T) & \leq & T^{\theta} \sup_{t \in (0,T)} \int_{\Omega} \left\{ \sum_{i=1}^{N} a_{i} d_{i} (u_{i\varepsilon}(\cdot,t) + \varepsilon)^{m_{i}} \right\}^{\theta} \\ & \leq & \left\{ NT \max_{i \in \{1,\dots,N\}} a_{i} d_{i} \right\}^{\theta} \sup_{t \in (0,T)} \int_{\Omega} (u_{i\varepsilon}(\cdot,t) + \varepsilon)^{\theta m_{i}} \\ & \leq & \left\{ NT \max_{i \in \{1,\dots,N\}} a_{i} d_{i} \right\}^{\theta} \sup_{t \in (0,T)} \int_{\Omega} (u_{i\varepsilon}(\cdot,t) + 1) \\ & \leq & c_{3}(T) \quad \text{for all } \varepsilon \in (0,1), \end{split}$$

whence using that then $\theta < 1 < r < 2 < r'$ we may invoke Young's inequality to see that with some $c_4 > 0$ we have

$$c_{2}\|z_{\varepsilon}(\cdot,T)\|_{L^{r}(\Omega)} \leq \frac{1}{2}\|z_{\varepsilon}(\cdot,T)\|_{L^{r'}(\Omega)} + c_{4}\|z_{\varepsilon}(\cdot,T)\|_{L^{\theta}(\Omega)}$$

$$\leq \frac{1}{2}\|z_{\varepsilon}(\cdot,T)\|_{L^{r'}(\Omega)} + c_{3}^{\frac{1}{\theta}}(T)c_{4} \quad \text{for all } \varepsilon \in (0,1).$$

In view of (3.7) and the nonnegativity of z_{ε} , (3.8) thus implies that

$$||z_{\varepsilon}(\cdot,T)||_{L^{r'}(\Omega)} \leq ||\overline{z_{\varepsilon}}||_{L^{r'}(\Omega)} \leq c_2||w_{\varepsilon}(\cdot,0) + 1||_{L^r(\Omega)} + \frac{1}{2}||z_{\varepsilon}(\cdot,T)||_{L^{r'}(\Omega)} + c_3^{\frac{1}{\theta}}(T)c_4 \quad \text{for all } \varepsilon \in (0,1),$$

so that in (3.6) we can use the Hölder inequality to estimate

$$\int_{\Omega} \left(w_{\varepsilon}(\cdot,0) + 1 \right) z_{\varepsilon}(\cdot,T) \leq \|w_{\varepsilon}(\cdot,0) + 1\|_{L^{r}(\Omega)} \|z_{\varepsilon}(\cdot,T)\|_{L^{r'}(\Omega)} \\
\leq \|w_{\varepsilon}(\cdot,0) + 1\|_{L^{r}(\Omega)} \cdot \left\{ 2c_{2} \|w_{\varepsilon}(\cdot,0) + 1\|_{L^{r}(\Omega)} + 2c_{3}^{\frac{1}{\theta}}(T)c_{4} \right\}$$

for all $\varepsilon \in (0,1)$. As $\sup_{\varepsilon \in (0,1)} \|w_{\varepsilon}(\cdot,0)\|_{L^{r}(\Omega)}$ is finite according to the hypothesis (3.2), (3.6) thereby entails the existence of $c_5(T) > 0$ such that

$$\int_0^T \int_{\Omega} e^{-Kt} \cdot \left\{ \sum_{i=1}^N d_i (u_{i\varepsilon} + \varepsilon)^{m_i} \right\} \cdot \left\{ \sum_{i=1}^N u_{i\varepsilon} \right\} \le c_5(T) \quad \text{for all } \varepsilon \in (0,1),$$

from which (3.4) readily follows.

We next rely on (1.5) in deriving the following estimates for gradients and reaction terms. Testing (3.1) by $-\frac{1}{\phi_i(u_{i\varepsilon})}$, namely, enables us to successfully combine (1.5) with (3.4). In order to obtain a bound for, e.g., $|f_i(u_{1\varepsilon},...,u_{N\varepsilon})|$ instead of $|-\frac{1}{\phi_i(u_{i\varepsilon})}f_i(u_{1\varepsilon},...,u_{N\varepsilon})|$, we here restrict our attention to sets of the form $\{u_{i\varepsilon} \leq M\}$, where $|-\frac{1}{\phi_i(u_{i\varepsilon})}|$ can be estimated from below by a positive constant.

Lemma 3.3 Let M > 0 and T > 0. Then one can find C(M,T) > 0 such that

$$\int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} (u_{i\varepsilon} + \varepsilon)^{m_{i} - 1} |\nabla u_{i\varepsilon}|^{2} \leq C(M, T) \quad \text{for all } i \in \mathbb{N} \text{ and } \varepsilon \in (0, 1)$$
 (3.9)

and

$$\int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} \frac{|f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})|}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \leq C(M, T) \quad \text{for all } i \in \mathbb{N} \text{ and } \varepsilon \in (0, 1).$$
 (3.10)

PROOF. For fixed $i \in \{1, ..., N\}$, we take $\phi_i \in C^1([0, \infty))$ as in (1.5) and define

$$\Phi_i(s) := -\int_1^{s+1} \frac{d\sigma}{\phi_i(\sigma)}, \qquad s \ge 0.$$

Then since ϕ_i is nondecreasing, we have

$$0 \ge \Phi_i(s) \ge -c_{1i}s \qquad \text{for all } s \ge 0 \tag{3.11}$$

with $c_{1i} := \frac{1}{\phi_i(1)} > 0$, and moreover $\Phi'_i(s) = -\frac{1}{\phi_i(s+1)}, \ s \geq 0$, satisfies

$$0 \le -\Phi_i'(s) \le \frac{1}{\phi_i(s)} \quad \text{for all } s \ge 0.$$
 (3.12)

Now using (3.1), for $\varepsilon \in (0,1)$ and t>0 we compute

$$\frac{d}{dt} \int_{\Omega} \Phi_{i}(u_{i\varepsilon}) = \int_{\Omega} \Phi'_{i}(u_{i\varepsilon}) \cdot \left\{ d_{i}m_{i}\nabla \cdot \left((u_{i\varepsilon} + \varepsilon)^{m_{i}-1}\nabla u_{i\varepsilon} \right) + \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \right\} \\
= -d_{i}m_{i} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} \Phi''_{i}(u_{i\varepsilon}) |\nabla u_{i\varepsilon}|^{2} + \int_{\Omega} \Phi'_{i}(u_{i\varepsilon}) \cdot \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|},$$

so that splitting $f_i = |f_i| - 2(f_i)_-$, upon further integration we find that

$$d_{i}m_{i} \int_{0}^{T} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} \Phi_{i}''(u_{i\varepsilon}) |\nabla u_{i\varepsilon}|^{2} + \int_{0}^{T} \int_{\Omega} |\Phi_{i}'(u_{i\varepsilon})| \cdot \frac{|f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})|}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|}$$

$$= \int_{\Omega} \Phi_{i}(u_{0i\varepsilon}) - \int_{\Omega} \Phi_{i}(u_{i\varepsilon}(\cdot, T))$$

$$+2 \int_{0}^{T} \int_{\Omega} |\Phi_{i}'(u_{i\varepsilon})| \cdot \frac{(f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon}))_{-}}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \quad \text{for all } T > 0,$$

$$(3.13)$$

because Φ'_i is nonpositive. Now on the right-hand side of (3.13) we use (1.5) to see that thanks to (3.12),

$$2\int_{0}^{T} \int_{\Omega} |\Phi'_{i}(u_{i\varepsilon})| \cdot \frac{(f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon}))_{-}}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \leq 2\int_{0}^{T} \int_{\Omega} |\Phi'_{i}(u_{i\varepsilon})| \cdot \phi_{i}(u_{i\varepsilon}) \cdot \left\{ \sum_{j \neq i} u_{j\varepsilon}^{\beta_{j}} + 1 \right\}$$
$$\leq 2\sum_{j \neq i} \int_{0}^{T} \int_{\Omega} u_{j\varepsilon}^{\beta_{j}} + 2|\Omega|T \quad \text{for all } T > 0,$$

whence recalling Lemma 3.2 we obtain $c_{2i}(T) > 0$ such that for all $\varepsilon \in (0,1)$,

$$2\int_0^T \int_{\Omega} |\Phi_i'(u_{i\varepsilon})| \cdot \frac{(f_i(u_{1\varepsilon}, ..., u_{N\varepsilon}))_-}{1 + \varepsilon \sum_{i=1}^N |f_i(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \le c_{2i}(T).$$

$$(3.14)$$

As

$$\int_{\Omega} \Phi_i(u_{0i\varepsilon}) - \int_{\Omega} \Phi_i(u_{i\varepsilon}(\cdot, T)) \le c_{1i} \int_{\Omega} u_{i\varepsilon}(\cdot, T) \quad \text{for all } T > 0$$

by (3.11), in view of Lemma 3.1 we thus conclude from (3.13) that there exists $c_{3i}(T) > 0$ with the property that

$$d_{i}m_{i}\int_{0}^{T}\int_{\Omega}(u_{i\varepsilon}+\varepsilon)^{m_{i}-1}\Phi_{i}''(u_{i\varepsilon})|\nabla u_{i\varepsilon}|^{2}+\int_{0}^{T}\int_{\Omega}|\Phi_{i}'(u_{i\varepsilon})|\cdot\frac{|f_{i}(u_{1\varepsilon},...,u_{N\varepsilon})|}{1+\varepsilon\sum_{i=1}^{N}|f_{i}(u_{1\varepsilon},...,u_{N\varepsilon})|}\leq c_{3i}(T) \quad (3.15)$$

for all $\varepsilon \in (0,1)$. Noting that for fixed M>0 and any $i\in\{1,...,N\}$ the numbers

$$c_{4i} := \min_{s \in [0,M]} \Phi_i''(s) = \min_{s \in [0,M]} \frac{\phi_i'(s+1)}{\phi_i^2(s+1)} \quad \text{and} \quad c_{5i} := \min_{s \in [0,M]} |\Phi_i'(s)| = \frac{1}{\phi_i(M+1)}$$

are both positive according to our hypotheses on ϕ_i and ϕ_i' , from (3.15) we readily infer (3.9) and (3.10) if we let $C(M,T) := \max\left\{\max_{i \in \{1,\dots,N\}} \frac{c_{3i}(T)}{d_i m_i c_{4i}}, \max_{i \in \{1,\dots,N\}} \frac{c_{3i}(T)}{c_{5i}}\right\}$, for instance.

While the bound in Lemma 3.2 is sufficient for concluding relative compactness of $\{u_{i\varepsilon} \mid \varepsilon \in (0,1)\}$ in some weak topology, we are additionally interested in possible pointwise convergence of $u_{i\varepsilon_j}$ along some sequence $(\varepsilon_j)_{j\in\mathbb{N}} \setminus 0$. We thus strive to derive a suitable strong compactness property in $L^2(\Omega \times (0,T))$, at least of a power of $u_{i\varepsilon}$ which has been cut off at large values so as to ensure accessibility to the estimates of Lemma 3.3.

Lemma 3.4 Given $\zeta \in C_0^{\infty}([0,\infty))$, for $i \in \{1,...,N\}$ let $\rho_i(s) := s^{\kappa_i}\zeta(s)$, $s \geq 0$, where $\kappa_i := \max\{\frac{m_i+1}{2}, 2\}$. Then

$$\left(\rho_i(u_{i\varepsilon})\right)_{\varepsilon\in(0,1)}$$
 is relatively compact in $L^2(\Omega\times(0,T))$ for all $T>0$. (3.16)

PROOF. Let us first make make sure that for each T > 0,

$$\left(\rho_i(u_{i\varepsilon})\right)_{\varepsilon\in(0,1)}$$
 is bounded in $L^2((0,T);W^{1,2}(\Omega))$. (3.17)

To see this, we note that due to the compactness of supp ρ_i it is clear that $(\rho_i(u_{i\varepsilon}))_{\varepsilon\in(0,1)}$ is bounded in $L^{\infty}(\Omega\times(0,\infty))$. Thereore, (3.17) results upon the observation that if we fix M>0 such that $\zeta\equiv 0$ on (M,∞) , then by Young's inequality and (3.9) we see that for all T>0 there exists $c_1(T)>0$ such

that

$$\int_{0}^{T} \int_{\Omega} |\nabla \rho_{i}(u_{i\varepsilon})|^{2} = \int_{0}^{T} \int_{\Omega} (\rho'_{i}(u_{i\varepsilon}))^{2} |\nabla u_{i\varepsilon}|^{2}$$

$$= \int_{0}^{T} \int_{\Omega} \left(\kappa_{i} u_{i\varepsilon}^{\kappa_{i}-1} \zeta(u_{i\varepsilon}) + u_{i\varepsilon}^{\kappa_{i}} \zeta'(u_{i\varepsilon}) \right)^{2} |\nabla u_{i\varepsilon}|^{2}$$

$$\leq 2\kappa_{i}^{2} ||\zeta||_{L^{\infty}((0,\infty))}^{2} \int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} u_{i\varepsilon}^{2\kappa_{i}-2} |\nabla u_{i\varepsilon}|^{2}$$

$$+2||\zeta'||_{L^{\infty}((0,\infty))}^{2} \int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} u_{i\varepsilon}^{2\kappa_{i}} |\nabla u_{i\varepsilon}|^{2}$$

$$\leq 2\kappa_{i}^{2} ||\zeta||_{L^{\infty}((0,\infty))}^{2} M^{2\kappa_{i}-m_{i}-1} \int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} u_{i\varepsilon}^{m_{i}-1} |\nabla u_{i\varepsilon}|^{2}$$

$$+2||\zeta'||_{L^{\infty}((0,\infty))}^{2} M^{2\kappa_{i}-m_{i}+1} \int_{0}^{T} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} u_{i\varepsilon}^{m_{i}-1} |\nabla u_{i\varepsilon}|^{2}$$

$$\leq c_{1}(T) \quad \text{for all } \varepsilon \in (0,1),$$

where we have used that $2\kappa_i - m_i + 1 \ge 2\kappa_i - m_i - 1 \ge 0$ by hypothesis.

We next fix any integer $k \geq 1$ such that k > n, and claim that then for all T > 0,

$$\left(\partial_t \rho_i(u_{i\varepsilon})\right)_{\varepsilon \in (0,1)}$$
 is bounded in $L^1\left((0,T); (W^{k,2}(\Omega))^*\right)$. (3.18)

To verify this in quite a straightforward manner, we pick $\psi \in C^{\infty}(\overline{\Omega})$ and use (3.1) to see that for each t > 0 and any $\varepsilon \in (0, 1)$,

$$\left| \int_{\Omega} \partial_{t} \rho_{i}(u_{i\varepsilon}(\cdot,t)) \cdot \psi \right|$$

$$= \left| \int_{\Omega} \rho'_{i}(u_{i\varepsilon}) \psi \cdot \left\{ d_{i} m_{i} \nabla \cdot \left((u_{i\varepsilon} + \varepsilon)^{m_{i}-1} \nabla u_{i\varepsilon} \right) + \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \right\} \right|$$

$$= \left| - d_{i} m_{i} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} \rho''_{i}(u_{i\varepsilon}) |\nabla u_{i\varepsilon}|^{2} \psi - d_{i} m_{i} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} \rho'_{i}(u_{i\varepsilon}) \nabla u_{i\varepsilon} \cdot \nabla \psi \right|$$

$$+ \int_{\Omega} \rho'_{i}(u_{i\varepsilon}) \cdot \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \psi \right|$$

$$\leq d_{i} m_{i} \|\rho''_{i}\|_{L^{\infty}((0,\infty))} \cdot \left\{ \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} |\nabla u_{i\varepsilon}|^{2} \right\} \cdot \|\psi\|_{L^{\infty}(\Omega)}$$

$$+ d_{i} m_{i} \cdot \left\{ \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} |\nabla u_{i\varepsilon}|^{2} + |\Omega| \sup_{s \geq 0} (s + \varepsilon)^{m_{i}-1} |\rho'_{i}(s)|^{2} \right\} \|\nabla \psi\|_{L^{\infty}(\Omega)}$$

$$+ \|\rho'_{i}\|_{L^{\infty}((0,\infty))} \cdot \left\{ \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} \frac{|f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})|}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \right\} \cdot \|\psi\|_{L^{\infty}(\Omega)}, \tag{3.19}$$

where finiteness of both $\|\rho'\|_{L^{\infty}((0,\infty))}$ and $\|\rho''_i\|_{L^{\infty}((0,\infty))}$ is asserted by our restriction that $\kappa_i \geq 2$. Here we observe that in the case $m_i \geq 1$ we have

$$(s+\varepsilon)^{m_i-1}|\rho_i'(s)|^2 \le (M+1)^{m_i-1}||\rho_i'||_{L^{\infty}((0,\infty))}^2,$$

whereas if $m_i < 1$ then

$$(s+\varepsilon)^{m_{i}-1}|\rho'_{i}(s)|^{2} \leq s^{m_{i}-1}|\rho'_{i}(s)|^{2}$$

$$= s^{m_{i}-1} \cdot \left|\kappa_{i}s^{\kappa_{i}-1}\zeta(s) + s^{\kappa_{i}}\zeta'(s)\right|^{2}$$

$$\leq 2\kappa_{i}^{2}s^{2\kappa_{i}+m_{i}-3}\zeta^{2}(s) + 2s^{2\kappa_{i}+m_{i}-1}|\zeta'(s)|^{2}$$

$$\leq 2\kappa_{i}^{2}M^{2\kappa_{i}+m_{i}-3}\|\zeta\|_{L^{\infty}((0,\infty))}^{2} + 2M^{2\kappa_{i}+m_{i}-1}\|\zeta'\|_{L^{\infty}((0,\infty))}^{2},$$

because again by definition of κ_i , we have $2\kappa_i + m_i - 1 \ge 2\kappa_i + m_i - 3 \ge 0$. As furthermore $W^{k,2}(\Omega) \hookrightarrow W^{1,\infty}(\Omega)$ due to our restriction that k > n, from (3.19) we thus infer the existence of $c_2 > 0$ such that for all t > 0 and any $\varepsilon \in (0,1)$,

$$\left\| \partial_{t} \rho_{i}(u_{i\varepsilon}(\cdot,t)) \right\|_{(W^{k,2}(\Omega))^{\star}} \leq c_{2} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1} |\nabla u_{i\varepsilon}|^{2} + c_{2} + c_{2} \int_{\Omega} \chi_{\{u_{i\varepsilon} \leq M\}} \frac{|f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})|}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|},$$

which in light of (3.9) and (3.10) establishes (3.18) upon a time integration.

We finally only need to combine (3.17) with (3.18) to conclude that (3.16) is a consequence of an Aubin-Lions type lemma ([24, Cor. 4]).

Based on this compactness statement, we may conclude the existence of a limit object.

Lemma 3.5 There exist $(\varepsilon_j)_{j\in\mathbb{N}}\subset(0,1)$ and nonnegative functions $u_1,...,u_N$ defined on $\Omega\times(0,\infty)$ such that $\varepsilon_j\searrow 0$ as $j\to\infty$, and such that for all $i\in\{1,...,N\}$,

$$u_{i\varepsilon} \to u_i \qquad a.e. \ in \ \Omega \times (0, \infty)$$
 (3.20)

and

$$u_{i\varepsilon} \to u_i \quad in \ L^p_{loc}(\overline{\Omega} \times [0, \infty)) \quad for \ all \ p \in [1, m_i + 1)$$
 (3.21)

as well as

$$u_{i\varepsilon} \rightharpoonup u_i \qquad in \ L_{loc}^{m_i-1}(\overline{\Omega} \times [0,\infty))$$
 (3.22)

as $\varepsilon = \varepsilon_i \searrow 0$.

PROOF. In Lemma 3.4 choosing $\zeta = \zeta_l$, with $\zeta_l \in C_0^{\infty}([0,\infty))$ satisfying $\zeta_j \equiv 1$ in [0,l] for $l \in \mathbb{N}$, by means of a straightforward extraction procedure we obtain $(\varepsilon_j)_{j \in \mathbb{N}} \subset (0,1)$ and $u = (u_1, ..., u_N) : \Omega \times (0,\infty) \to \mathbb{R}^N$ such that for all $i \in \{1,...,N\}$ we have $u_0 \geq 0$ and $u_{i\varepsilon} \to u_i$ a.e. in $\Omega \times (0,\infty)$ as $\varepsilon = \varepsilon_j \searrow 0$. Since for each $i \in \{1,...,N\}$ and all T > 0 we know from Lemma 3.2 that $(u_{i\varepsilon})_{\varepsilon \in (0,1)}$ is bounded in $L^{m_i+1}(\Omega \times (0,T))$, and that hence $(u^p_{i\varepsilon})_{\varepsilon \in (0,1)}$ is uniformly integrable over $\Omega \times (0,T)$ for all $p \in [1, m_i + 1)$, by reflexivity of $L^{m_i+1}(\Omega \times (0,T))$ and the Vitali convergence theorem we readily

infer that on passing to a further subsequence if necessary we can also achieve simultaneous validity of (3.21) amd (3.22).

Our next goal is to show that the functions just constructed acutally form a solution. We begin by confirming that they enjoy a renormalized supersolution property in the style of Definition 2.1. The most crucial ingredient in our verification of this – and actually the reason for dealing with supersolutions – becomes apparent in (3.27), which is enlisted to control the integral involving the gradient (of $P_i^{(1)}(u)$) from above by means of lower semicontinuity.

Lemma 3.6 Let $u_1, ..., u_N$ be as given by Lemma 3.5. Then $u = (u_1, ..., u_N)$ forms a renormalized supersolution of (1.4) in the sense of Definition 2.1.

PROOF. We fix $i \in \{1,...,N\}$ and a nonincreasing concave $\rho \in C^{\infty}([0,\infty))$ such that $\rho' \in C^{\infty}_{0}([0,\infty))$, and use (3.1) to see that for all nonnegative $\varphi \in C^{\infty}_{0}(\overline{\Omega} \times [0,\infty))$ such that $\frac{\partial \varphi}{\partial \nu} = 0$ on $\partial \Omega \times (0,\infty)$, we have

$$-\int_{0}^{\infty} \int_{\Omega} \rho(u_{i\varepsilon})\varphi_{t} - \int_{\Omega} \rho(u_{0i\varepsilon})\varphi(\cdot,0) = + \int_{0}^{\infty} \int_{\Omega} \partial_{t}\rho(u_{i\varepsilon})\varphi$$

$$= \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon})\varphi \cdot \left\{ d_{i}m_{i}\nabla \cdot \left((u_{i\varepsilon} + \varepsilon)^{m_{i}-1}\nabla u_{i\varepsilon} \right) + \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \right\}$$

$$= -d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1}\rho''(u_{i\varepsilon})|\nabla u_{i\varepsilon}|^{2}\varphi - d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} (u_{i\varepsilon} + \varepsilon)^{m_{i}-1}\rho'(u_{i\varepsilon})\nabla u_{i\varepsilon} \cdot \nabla \varphi$$

$$+ \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \varphi$$

$$= -d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} |\nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon})|^{2}\varphi + d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} P_{i\varepsilon}^{(2)}(u_{i\varepsilon})\Delta\varphi$$

$$+ \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon})}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \varphi \quad \text{for all } \varepsilon \in (0, 1),$$
(3.23)

where we have set

$$P_{i\varepsilon}^{(1)}(s) := \int_0^s (\sigma + \varepsilon)^{\frac{m_i - 1}{2}} \sqrt{\rho''(\sigma)} d\sigma \quad \text{and} \quad P_{i\varepsilon}^{(2)}(s) := \int_0^s (\sigma + \varepsilon)^{m_i - 1} \rho'(\sigma) d\sigma, \qquad s \ge 0, \quad (3.24)$$

for $i \in \{1, ..., N\}$ and $\varepsilon \in (0, 1)$. Here we note that if we take M > 0 and T > 0 large enough fulfilling $\operatorname{supp} \rho' \subset [0, M]$ and $\operatorname{supp} \varphi \subset \overline{\Omega} \times [0, T]$, then for all $\varepsilon \in (0, 1)$,

$$\int_0^\infty \int_\Omega \left| \sqrt{\varphi} \nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon}) \right|^2 \le \|\rho''\|_{L^\infty((0,\infty))} \|\varphi\|_{L^\infty(\Omega\times(0,\infty))} \int_0^T \int_\Omega \chi_{\{u_{i\varepsilon} \le M\}}(u_{i\varepsilon} + \varepsilon)^{m_i - 1} |\nabla u_{i\varepsilon}|^2,$$

whence again employing Lemma 3.3 we infer that with $(\varepsilon_j)_{j\in\mathbb{N}}$ as provided by Lemma 3.5, we can find a subsequence $(\varepsilon_{j_k})_{k\in\mathbb{N}}$ such that

$$\sqrt{\varphi} \nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon}) \rightharpoonup z \quad \text{in } L^2(\Omega \times (0,\infty); \mathbb{R}^n) \qquad \text{as } \varepsilon = \varepsilon_{j_k} \searrow 0$$
 (3.25)

for some $z \in L^2(\Omega \times (0, \infty); \mathbb{R}^n)$. On the other hand, since (3.24) entails that $P_{i\varepsilon}^{(1)} \to P_i^{(1)}$ in $L_{loc}^{\infty}([0, \infty))$ as $\varepsilon \searrow 0$, with $P_i^{(1)}$ given by (2.1), and since moreover

$$|P_{i\varepsilon}^{(1)}(s)| \leq \|\rho''\|_{L^{\infty}((0,\infty))}^{\frac{1}{2}} \int_{0}^{M} (\sigma + \varepsilon)^{\frac{m_{i}-1}{2}} d\sigma$$

$$= \|\rho''\|_{L^{\infty}((0,\infty))}^{\frac{1}{2}} \cdot \frac{(M+\varepsilon)^{\frac{m_{i}+1}{2}} - \varepsilon^{\frac{m_{i}+1}{2}}}{\frac{m_{i}+1}{2}}$$

$$\leq \|\rho''\|_{L^{\infty}((0,\infty))}^{\frac{1}{2}} \cdot \frac{2(M+1)^{\frac{m_{i}+1}{2}}}{m_{i}+1} \quad \text{for all } s \geq 0 \text{ and } \varepsilon \in (0,1), \tag{3.26}$$

from (3.20) and the dominated convergence theorem it follows that

$$P_{i\varepsilon}^{(1)}(u_{i\varepsilon}) \to P_{i}^{(1)}(u_{i})$$
 in $L^{1}(\Omega \times (0,T))$ as $\varepsilon = \varepsilon_{j} \searrow 0$.

Therefore, a standard argument shows that in (3.25) we must have $z = \sqrt{\varphi} \nabla P_i^{(1)}(u_i)$ a.e. in $\{\varphi > \delta\}$ for all $\delta > 0$, and hence actually $\sqrt{\varphi} \nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon}) \rightharpoonup \sqrt{\varphi} \nabla P_i^{(1)}(u_i)$ in $L_{loc}^2(\Omega \times (0,\infty))$ as $\varepsilon = \varepsilon_{j_k} \searrow 0$, so that by lower semicontinuity of the norm in $L^2(\Omega \times (0,\infty))$ with respect to weak convergence,

$$d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} |\nabla P_{i}^{(1)}(u_{i})|^{2} \varphi \leq \liminf_{\varepsilon = \varepsilon_{j_{k}} \searrow 0} \left\{ d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} |\nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon})|^{2} \varphi \right\}. \tag{3.27}$$

Next addressing the integrals in (3.23) exclusively containing zero-order expressions with respect to $u_{i\varepsilon}$, we first observe that clearly

$$|\rho(s)| \le ||\rho'||_{L^{\infty}((0,\infty))} \cdot M + \rho(0)$$
 for all $s \ge 0$,

and that furthermore, by (3.24), similarly to (3.26) we can estimate

$$|P_{i\varepsilon}^{(2)}(s)| \leq \|\rho'\|_{L^{\infty}((0,\infty))} \int_{0}^{M} (\sigma+\varepsilon)^{m_{i}-1} d\sigma$$

$$\leq \|\rho'\|_{L^{\infty}((0,\infty))} \cdot \frac{(M+1)^{m_{i}}}{m_{i}} \quad \text{for all } s \geq 0 \text{ and } \varepsilon \in (0,1).$$

Therefore, three applications of the dominated convergence theorem on the basis of (3.20) and (3.2) show that if we take $P_i^{(2)}$ from (2.1) then

$$\int_0^\infty \int_\Omega \rho(u_{i\varepsilon})\varphi_t \to \int_0^\infty \int_\Omega \rho(u_i)\varphi_t \tag{3.28}$$

and

$$\int_{\Omega} \rho(u_{0i\varepsilon})\varphi(\cdot,0) \to \int_{\Omega} \rho(u_{0i})\varphi(\cdot,0) \tag{3.29}$$

as well as

$$d_i m_i \int_0^\infty \int_\Omega P_{i\varepsilon}^{(2)}(u_{i\varepsilon}) \Delta \varphi \to d_i m_i \int_0^\infty \int_\Omega P_i^{(2)}(u_i) \Delta \varphi \tag{3.30}$$

as $\varepsilon = \varepsilon_j \searrow 0$, the latter because in addition obviously $P_{i\varepsilon}^{(2)} \to P_i^{(2)}$ in $L_{loc}^{\infty}([0,\infty))$ as $\varepsilon \searrow 0$.

Finally, in the crucial rightmost summand in (3.23) containing the reactive contribution, we once more rewrite $f_i = |f_i| - 2(f_i)_-$ and note that fixing any $\delta > 0$ such that $(1 + \delta)\beta_j \leq m_j + 1$ for all $j \in \{1, ..., N\} \setminus \{i\}$, again relying on (1.5) we can estimate

$$\int_{0}^{T} \int_{\Omega} \left| \rho'(u_{i\varepsilon}) \cdot \frac{(f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon}))_{-}}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \varphi \right|^{1+\delta} \\
\leq \int_{0}^{T} \int_{\Omega} |\rho'(u_{i\varepsilon})|^{1+\delta} \cdot \left\{ \phi_{i}(u_{i\varepsilon}) \left(\sum_{j \neq i} u_{j\varepsilon}^{\beta_{j}} + 1 \right) \right\}^{1+\delta} \cdot \varphi^{1+\delta} \\
\leq \left\{ \|\rho'\|_{L^{\infty}((0,\infty))} \cdot \|\phi_{i}\|_{L^{\infty}((0,M))} \cdot \|\varphi\|_{L^{\infty}(\Omega \times (0,\infty))} \right\}^{1+\delta} \cdot N^{1+\delta} \cdot \left(\sum_{j \neq i} \int_{0}^{T} \int_{\Omega} u_{j\varepsilon}^{(1+\delta)\beta_{j}} + |\Omega|T \right)$$

for all $\varepsilon \in (0,1)$. In view of Lemma 3.2, by positivity of δ this implies uniform integrability of $\left(\rho'(u_{i\varepsilon}) \cdot \frac{(f_i(u_{1\varepsilon},...,u_{N\varepsilon}))_-}{1+\varepsilon \sum_{j=1}^N |f_j(u_{1\varepsilon},...,u_{N\varepsilon})|} \cdot \varphi\right)_{\varepsilon \in (0,1)}$ over $\Omega \times (0,T)$ and hence entails, when combined with (3.20), that

$$-2\int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{(f_{i}(u_{1\varepsilon}, ..., u_{N\varepsilon}))_{-}}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon}, ..., u_{N\varepsilon})|} \cdot \varphi \to -2\int_{0}^{\infty} \int_{\Omega} \rho'(u_{i})(f_{i}(u_{1}, ..., u_{N}))_{-}\varphi$$
 (3.31)

as $\varepsilon = \varepsilon_j \searrow 0$. Since apart from that, by nonnegativity of both $-\rho'$ and φ we can invoke Fatou's lemma to see that, again thanks to (3.20),

$$-\int_{0}^{\infty} \int_{\Omega} \rho'(u_{i})|f_{i}(u_{1},...,u_{N})|\varphi \leq \liminf_{\varepsilon=\varepsilon_{j} \searrow 0} \left\{-\int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{|f_{i}(u_{1\varepsilon},...,u_{N\varepsilon})|}{1+\varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon},...,u_{N\varepsilon})|} \cdot \varphi\right\}, \quad (3.32)$$

upon collecting (3.27)-(3.32) we altogether conclude from (3.23) that

$$\begin{split} d_{i}m_{i} & \int_{0}^{\infty} \int_{\Omega} |\nabla P_{i}^{(1)}(u_{i})|^{2} \varphi - \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i})|f_{i}(u_{1},...,u_{N})| \varphi \\ & \leq \lim\inf_{\varepsilon = \varepsilon_{j_{k}} \searrow 0} \left\{ d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} |\nabla P_{i\varepsilon}^{(1)}(u_{i\varepsilon})|^{2} \varphi - \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{|f_{i}(u_{1\varepsilon},...,u_{N\varepsilon})|}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon},...,u_{N\varepsilon})|} \cdot \varphi \right\} \\ & = \lim\inf_{\varepsilon = \varepsilon_{j_{k}} \searrow 0} \left\{ \int_{0}^{\infty} \int_{\Omega} \rho(u_{i\varepsilon}) \varphi_{t} + \int_{\Omega} \rho(u_{0i\varepsilon}) \varphi(\cdot,0) \right. \\ & \left. + d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} P_{i\varepsilon}^{(2)}(u_{i\varepsilon}) \Delta \varphi - 2 \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i\varepsilon}) \cdot \frac{(f_{i}(u_{1\varepsilon},...,u_{N\varepsilon}))_{-}}{1 + \varepsilon \sum_{j=1}^{N} |f_{j}(u_{1\varepsilon},...,u_{N\varepsilon})|} \cdot \varphi \right\} \\ & = \int_{0}^{\infty} \int_{\Omega} \rho(u_{i}) \varphi_{t} + \int_{\Omega} \rho(u_{0i}) \varphi(\cdot,0) \\ & \left. + d_{i}m_{i} \int_{0}^{\infty} \int_{\Omega} P_{i}^{(2)}(u_{i}) \Delta \varphi - 2 \int_{0}^{\infty} \int_{\Omega} \rho'(u_{i})(f_{i}(u_{1},...,u_{N}))_{-} \varphi, \end{split}$$

which is equivalent to the desired inequality (2.3). The integrability requirements in (2.2) are evident by-products of the above considerations.

But also the subsolution property encoded in (2.5) is fulfilled:

Lemma 3.7 The function $u = (u_1, ..., u_N)$ from Lemma 3.5 satisfies (2.4) and (2.5) of Definition 2.2.

PROOF. According to (3.21), we can pick a null set $N \subset (0, \infty)$ such that with $(\varepsilon_j)_{j \in \mathbb{N}}$ taken from Lemma 3.5, for each $t \in (0, \infty) \setminus N$ we have $u_{i\varepsilon}(\cdot, t) \to u_i(\cdot, t)$ in $L^1(\Omega)$ for all $i \in \{1, ..., N\}$ and hence

$$\int_{\Omega} \sum_{i=1}^{N} a_i u_{i\varepsilon}(\cdot, t) \to \int_{\Omega} \sum_{i=1}^{N} a_i u_i(\cdot, t)$$
(3.33)

as $\varepsilon = \varepsilon_j \searrow 0$, where $a \in (0, \infty)^N$ is taken from (1.3). We next let $F_{\varepsilon}(s_1, ..., s_N) := \sum_{i=1}^N \frac{a_i f_i(s_1, ..., s_N)}{1 + \varepsilon \sum_{j=1}^N |f_j(s_1, ..., s_N)|}$ and $F(s_1, ..., s_N) := \sum_{i=1}^N a_i f_i(s_1, ..., s_N)$ for $(s_1, ..., s_N) \in \mathbb{R}^N$ and $\varepsilon \in (0, 1)$, and then obtain from (1.3) that

$$\left(F_{\varepsilon}(u_{1\varepsilon},...,u_{N\varepsilon})\right)_{+} \leq K \sum_{i=1}^{N} a_{i} u_{i\varepsilon} + K \quad \text{in } \Omega \times (0,\infty) \qquad \text{for all } \varepsilon \in (0,1).$$

In view of (3.21) applied to p := 1, a version of the dominated convergence theorem thus ensures that

$$\int_{0}^{t} \int_{\Omega} \left(F_{\varepsilon}(u_{1\varepsilon}, ..., u_{N\varepsilon}) \right)_{+} \to \int_{0}^{t} \int_{\Omega} F_{+}(u_{1}, ..., u_{N}) \quad \text{for all } t > 0 \qquad \text{as } \varepsilon = \varepsilon_{j} \searrow 0, \tag{3.34}$$

because clearly $\left(F_{\varepsilon}(u_{1\varepsilon},...,u_{N\varepsilon})\right)_{+} \to F_{+}(u_{1},...,u_{N})$ a.e. in $\Omega \times (0,\infty)$ as $\varepsilon = \varepsilon_{j} \searrow 0$ by (3.20). Since from (3.1) we know that for all t > 0 and $\varepsilon \in (0,1)$ we have

$$\int_{\Omega} \sum_{i=1}^{N} a_i u_{i\varepsilon}(\cdot, t) + \int_{0}^{t} \int_{\Omega} \left(F_{\varepsilon}(u_{1\varepsilon}, ..., u_{N\varepsilon}) \right)_{-} = \int_{\Omega} \sum_{i=1}^{N} a_i u_{0i\varepsilon} + \int_{0}^{t} \int_{\Omega} \left(F_{\varepsilon}(u_{1\varepsilon}, ..., u_{N\varepsilon}) \right)_{+},$$

where by (3.20) also $\left(F_{\varepsilon}(u_{1\varepsilon},...,u_{N\varepsilon})\right)_{-} \to F_{-}(u_{1},...,u_{N})$ a.e. in $\Omega \times (0,\infty)$ as $\varepsilon = \varepsilon_{j} \searrow 0$, and where $\int_{\Omega} \sum_{i=1}^{N} a_{i}u_{0i\varepsilon} \to \int_{\Omega} \sum_{i=1}^{N} a_{i}u_{0i}$ as $\varepsilon \searrow 0$ due to (3.2), invoking Fatou's lemma we infer by means of (3.33) and (3.34) that

$$\int_{\Omega} \sum_{i=1}^{N} a_i u_i(\cdot, t) + \int_{0}^{t} \int_{\Omega} F_{-}(u_1, ..., u_N) \le \int_{\Omega} \sum_{i=1}^{N} a_i u_{0i} + \int_{0}^{t} \int_{\Omega} F_{+}(u_1, ..., u_N) \quad \text{for all } t \in (0, \infty) \setminus N.$$

For any such t, this firstly implies that $F(u_1, \ldots, u_N)$ belongs to $L^1(\Omega \times (0, t))$, and secondly entails that (2.5) holds.

The previous two lemmata already demonstrate that u is a generalized solution in the sense of Definition 2.2:

PROOF of Theorem 1.1. We take $u_1, ..., u_N$ as given by Lemma 3.5 and then only need to combine Lemma 3.6 with Lemma 3.7.

References

- [1] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992), volume 133 of Teubner-Texte Math., pages 9–126. Teubner, Stuttgart, 1993.
- [2] J. A. Cañizo, L. Desvillettes, and K. Fellner. Improved duality estimates and applications to reaction-diffusion equations. *Comm. Partial Differential Equations*, 39(6):1185–1204, 2014.
- [3] M. C. Caputo, T. Goudon, and A. F. Vasseur. Solutions of the 4-species quadratic reaction-diffusion system are bounded and C^{∞} -smooth, in any space dimension. *Anal. PDE*, 12(7):1773–1804, 2019.
- [4] M. C. Caputo and A. Vasseur. Global regularity of solutions to systems of reaction-diffusion with sub-quadratic growth in any dimension. *Comm. Partial Differential Equations*, 34(10-12):1228–1250, 2009.
- [5] B. P. Cupps, J. Morgan, and B. Q. Tang. Uniform boundedness for reaction-diffusion systems with mass dissipation. SIAM J. Math. Anal., 53(1):323–350, 2021.
- [6] L. Desvillettes, K. Fellner, M. Pierre, and J. Vovelle. Global existence for quadratic systems of reaction-diffusion. Adv. Nonlinear Stud., 7(3):491–511, 2007.
- [7] R. J. DiPerna and P.-L. Lions. On the Cauchy problem for Boltzmann equations: global existence and weak stability. *Ann. of Math.* (2), 130(2):321–366, 1989.
- [8] K. Fellner, J. Morgan, and B. Q. Tang. Global classical solutions to quadratic systems with mass control in arbitrary dimensions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 37(2):281–307, 2020.
- [9] J. Fischer. Global existence of renormalized solutions to entropy-dissipating reaction-diffusion systems. *Arch. Ration. Mech. Anal.*, 218(1):553–587, 2015.
- [10] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [11] Y. I. Kanel'. Solvability in the large of a system of reaction-diffusion equations with the balance condition. Differentsial' nye Uravneniya, 26(3):448–458, 549, 1990.
- [12] E. H. Laamri. Étude de l'existence de solutions globales d'un système de réaction-diffusion parabolique fortement non linéaire. Ann. Fac. Sci. Toulouse Math. (5), 12(3):373–390, 1991.
- [13] E.-H. Laamri and B. Perthame. Reaction-diffusion systems with initial data of low regularity. *J. Differential Equations*, 269(11):9310–9335, 2020.
- [14] E. H. Laamri and M. Pierre. Global existence for reaction-diffusion systems with nonlinear diffusion and control of mass. Ann. Inst. H. Poincaré Anal. Non Linéaire, 34(3):571–591, 2017.
- [15] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva. Linear and quasilinear equations of parabolic type. Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968.
- [16] J. Lankeit and M. Winkler. A generalized solution concept for the Keller-Segel system with logarithmic sensitivity: global solvability for large nonradial data. NoDEA Nonlinear Differential Equations Appl., 24(4):Paper No. 49, 33, 2017.
- [17] R. H. Martin and M. Pierre. Influence of mixed boundary conditions in some reaction-diffusion systems. Proc. Roy. Soc. Edinburgh Sect. A, 127(5):1053–1066, 1997.
- [18] M. Pierre. Weak solutions and supersolutions in L^1 for reaction-diffusion systems. volume 3, pages 153–168. 2003. Dedicated to Philippe Bénilan.
- [19] M. Pierre. Global existence in reaction-diffusion systems with control of mass: a survey. Milan J. Math., 78(2):417–455, 2010.
- [20] M. Pierre and G. Rolland. Global existence for a class of quadratic reaction-diffusion systems with nonlinear diffusions and L^1 initial data. Nonlinear Anal., 138:369–387, 2016.
- [21] M. Pierre and D. Schmitt. Blowup in reaction-diffusion systems with dissipation of mass. SIAM J. Math. Anal., 28(2):259–269, 1997.

- [22] M. Pierre, T. Suzuki, and H. Umakoshi. Global-in-time behavior of weak solutions to reaction-diffusion systems with inhomogeneous Dirichlet boundary condition. *Nonlinear Anal.*, 159:393–407, 2017.
- [23] P. Quittner and P. Souplet. Superlinear parabolic problems. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser/Springer, Cham, 2019. Blow-up, global existence and steady states, Second edition of [MR2346798].
- [24] J. Simon. Compact sets in the space $L^{p}(0,T;B)$. Ann. Mat. Pura Appl. (4), 146:65–96, 1987.
- [25] P. Souplet. Global existence for reaction-diffusion systems with dissipation of mass and quadratic growth. J. Evol. Equ., 18(4):1713–1720, 2018.
- [26] M. Winkler. Large-data global generalized solutions in a chemotaxis system with tensor-valued sensitivities. SIAM J. Math. Anal., 47(4):3092–3115, 2015.
- [27] M. Winkler. Global mass-preserving solutions in a two-dimensional chemotaxis-Stokes system with rotational flux components. J. Evol. Equ., 18(3):1267–1289, 2018.
- [28] A. Zhigun. Generalized global supersolutions with mass control for systems with taxis. SIAM J. Math. Anal., 51(3):2425–2443, 2019.