A FUNCTIONAL CHARACTERIZATION OF ALMOST-GREEDY AND PARTIALLY-BASES IN BANACH SPACES

P. M. BERNÁ AND D. MONDÉJAR

ABSTRACT. In 2003, S. J. Dilworth et al. ([8]) introduced the notion of almost-greedy (resp. partially-greedy) bases. These bases were characterized in terms of quasi-greediness and democracy (resp. conservativeness). In this paper we will show a new functional characterization of these type of bases in general Banach spaces following the spirit of the characterization of greediness proved in [5].

1. INTRODUCTION AND BACKGROUND

Assume that $(\mathbb{X}, \|\cdot\|)$ is a Banach space over the field $\mathbb{F} = \mathbb{R}$ or \mathbb{C} . Throughout the paper, we assume that $\mathcal{B} = (e_n)_{n=1}^{\infty}$ is a semi-normalized Markushevich basis, that is, there exists a unique sequence $(e_n^*)_{n=1}^{\infty} \subset \mathbb{X}^*$ such that

- $\overline{\operatorname{span}(e_n : n \in \mathbb{N})} = \mathbb{X},$
- $e_n^*(e_m) = \delta_{n,m}$, if $e_n^*(f) = 0$ for all $n \in \mathbb{N}$, then f = 0,
- there are $c_1, c_2 > 0$ such that

$$0 < c_1 := \inf_n \min\{\|e_n\|, \|e_n^*\|\} \le \sup_n \max\{\|e_n\|, \|e_n^*\|\} =: c_2 < \infty.$$

Hereinafter, by a *basis* for X, we mean a semi-normalized Markushevich basis. Under these conditions, for each $f \in \mathbb{X}$, we have that $f \sim \sum_{n=1}^{\infty} e_n^*(f)e_n$ where $(e_n^*(f))_n \in c_0$. The support of $f \in \mathbb{X}$ is denoted by $\operatorname{supp}(f)$, where $\operatorname{supp}(f) = \{n \in \mathbb{N} : |e_n^*(f)| \neq 0\}$. Finally, we will use the following notation: \mathbb{X}_{fin} is the subspace of \mathbb{X} with the elements with finite support, if $f, g \in \mathbb{X}$, $f \cdot g = 0$ means that $\operatorname{supp}(f) \cap \operatorname{supp}(g) = \emptyset$, $\tilde{f} = (e_n^*(f))_{n \in \mathbb{N}}$ and $\|\tilde{f}\|_{\infty} = \operatorname{sup}_{n \in \mathbb{N}} |e_n^*(f)|$. Moreover, if A and B are finite sets of natural numbers, A < B means that $\max_{n \in A} n < n$ $\min_{j\in B} j$, P_A is the projection operator, that is, $P_A(f) = \sum_{n\in A} e_n^*(f)e_n$, and S_k is the partial sum of order k, that is, $S_k(f) = P_{\{1,\dots,k\}}(f)$.

In 1999, S. V. Konyagin and V. N. Temlyakov ([11]) introduced one of the most studied algorithms in the field of non-linear approximation, the so called Thresholding Greedy Algorithm: for $f \in \mathbb{X}$ and $m \in \mathbb{N}$, we define a greedy sum of order m as

$$\mathcal{G}_m(f)[\mathbb{X},\mathcal{B}] = \mathcal{G}_m(f) := \sum_{n \in A_m(f)} e_n^*(f)e_n$$

²⁰⁰⁰ Mathematics Subject Classification. 46B15, 41A65.

Key words and phrases: greedy algorithm, non-linear approximation; bases; Banach spaces.

The first author was supported by Grants PID2019-105599GB-I00 (Agencia Estatal de Investigación, Spain) and 20906/PI/18 from Fundación Séneca (Región de Murcia, Spain). The second author was supported by the research project PGC2018-098321-B-I00 (Ministerio de Ciencia, Innovación y Universidades, Spain).

where $A_m(f)$ is a greedy set of order m, that is, $|A_m(f)| = m$ and

$$\min_{n \in A_m(f)} |e_n^*(f)| \ge \max_{n \notin A_m(f)} |e_n^*(f)|.$$

The collection $(\mathcal{G}_m)_{m\in\mathbb{N}}$ is the Thresholding Greedy Algorithm. As for every algorithm, one of the first question that we can ask to the audience is when the algorithm converges. To solve that question, S. V. Konyagin and V. N. Temlyakov introduced in [11] the notion of quasi-greediness.

Definition 1.1. We say that \mathcal{B} is quasi-greedy if there is a positive constant C such that

$$\|f - \mathcal{G}_m(f)\| \le C \|f\|, \ \forall m \in \mathbb{N}, \forall f \in \mathbb{X}.$$
(1)

The least constant verifying (1) is denoted by $C_q = C_q[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} is C_q -quasigreedy.

Although this definition only talks about the boundedness of the greedy sums, P. Wojtaszczyk showed in [12] that quasi-greediness is equivalent to the convergence of the algorithm.

Theorem 1.2 ([1, 12]). A basis \mathcal{B} in a Banach (or quasi-Banach) space is quasi-greedy if and only if

$$\lim_{m \to +\infty} \|f - \mathcal{G}_m(f)\| = 0, \ \forall f \in \mathbb{X}.$$

Then, quasi-greediness is the minimal condition in the convergence of the algorithm, but we are interested in others types of convergence. For instance, when does the algorithm produce the best possible approximation? To study this question, in [11], the authors introduced the notion of greediness: a basis \mathcal{B} is greedy if there is a positive constant C_g such that

$$\|f - \mathcal{G}_m(f)\| \le C_g \inf\{\|f - \sum_{n \in B} a_n e_n\| : a_n \in \mathbb{F}, |B| \le m\}, \ \forall m \in \mathbb{N}, \forall f \in \mathbb{X}.$$

There are several examples of greedy bases, for instance the canonical basis in the spaces ℓ_p with $1 \leq p < \infty$, the Haar system in $L_p((0, 1))$ with $1 or the trigonometric system in <math>L_2(\mathbb{T})$. To study greedy bases, S. V. Konyagin and V. N. Temlyakov gave a characterization in terms of unconditional and democratic bases, where a basis is unconditional if the projection operator is uniformly bounded, that is, there is K > 0 such that, for any finite set A,

$$||P_A(f)|| \le K||f||, \ \forall f \in \mathbb{X}.$$

Consider A a finite set and define the set of the collection of signs in A, $\mathcal{E}_A = \{\varepsilon = (\varepsilon_n)_{n \in A} : |\varepsilon_n| = 1\}$, and take the indicator sum

$$1_{\varepsilon A} = 1_{\varepsilon A}[\mathbb{X}, \mathcal{B}] := \sum_{n \in A} \varepsilon_n e_n.$$

If $\varepsilon \equiv 1$, we will use the notation 1_A .

Definition 1.3. [1, 4, 8, 9] We say that \mathcal{B} is symmetric for largest coefficients if there is a positive constant C such that

$$\|f + 1_{\varepsilon A}\| \le C \|f + 1_{\varepsilon' B}\|,\tag{2}$$

for any pair of sets $|A| \leq |B| < \infty$, $A \cap B = \emptyset$, for any $f \in \mathbb{X}$ such that $\operatorname{supp}(f) \cap (A \cup B) = \emptyset$, $|e_n^*(f)| \leq 1$ for all $n \in \mathbb{N}$ and for any choice of signs $\varepsilon \in \mathcal{E}_A$, $\varepsilon' \in \mathcal{E}_B$. The least constant verifying (2) is denoted by $\Delta = \Delta[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} is Δ -symmetric for largest coefficients. If (2) is satisfied with the extra condition that $A < \operatorname{supp}(f) \cup B$, then we say that \mathcal{B} is partially symmetric for largest coefficients with constant Δ_{pc} .

Definition 1.4. We say that \mathcal{B} is super-democratic if there is a positive constant C such that

$$\|1_{\varepsilon A}\| \le C \|1_{\eta B}\|,\tag{3}$$

for any pair of sets $A, B \subset \mathbb{N}$, $|A| \leq |B| < \infty$ and for any choice of signs $\varepsilon \in \mathcal{E}_A, \eta \in \mathcal{E}_B$. The least constant verifying (3) is denoted by $\Delta_s = \Delta_s[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} is Δ_s -superdemocratic.

If (3) is satisfied for A < B, we say that \mathcal{B} is Δ_{sc} -super-conservative.

If (3) is satisfied for $\varepsilon \equiv \eta \equiv 1$, we say that \mathcal{B} is Δ_d -democratic and if in addition, A < B, we say that \mathcal{B} is Δ_c -conservative.

With these definitions, we can find the following characterizations of greedy bases.

Theorem 1.5. Assume that \mathcal{B} is a basis in a Banach space \mathbb{X} .

• \mathcal{B} is greedy if and only if \mathcal{B} is democratic and unconditional (see [11]). Moreover,

 $\max\{K, \Delta_d\} \le C_g \le K + K^2 \Delta_d.$

• \mathcal{B} is greedy if and only if \mathcal{B} is super-democratic and unconditional (see [6]). Moreover,

$$\max\{K, \Delta_s\} \le C_g \le K + K\Delta_s.$$

• \mathcal{B} is greedy if and only if \mathcal{B} is symmetric for largest coefficients and unconditional (see [9]). Moreover,

$$\max\{K, \Delta_d\} \le C_g \le K\Delta.$$

The last two characterizations were studied with the objective to improve the boundedness constant of greedy bases. Moreover, all the characterizations were given under the assumption of unconditionality and one of the democracy-like properties but, in [5], we find a new and interesting property that is so useful to give a new characterization of greediness (see [5, Corollary 1.8]). This property is the so called Property (Q): there is a C > 0 such that

$$||f + 1_A|| \le C||f + g + 1_B||,$$

for any $|A| = |B| < \infty$, $A \cap B = \emptyset$ and $f, g \in \mathbb{X}_{fin}$ such that $\operatorname{supp}(f) \cap \operatorname{supp}(g) = \emptyset$, $\|\tilde{f}\|_{\infty} \leq 1$ and $\operatorname{supp}(f+g) \cap (A \cup B) = \emptyset$.

In that paper, we focus our attention in a closed inequality to characterize the so called almost-greedy and partially-greedy bases.

Definition 1.6 ([8]). We say that \mathcal{B} is almost-greedy if there is a positive constant C such that

$$\|f - \mathcal{G}_m(f)\| \le C \inf\{\|f - P_B(f)\| : |B| \le m\}, \quad \forall m \in \mathbb{N}, \forall f \in \mathbb{X}.$$
(4)

The least constant verifying (4) is denoted by $C_{al} = C_{al}[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} is C_{al} -almost-greedy.

Definition 1.7 ([4, 8]). We say that \mathcal{B} is partially-greedy if there is positive constant C such that

$$\|f - \mathcal{G}_m(f)\| \le C \inf_{k \le m} \|f - S_k(f)\|, \ \forall m \in \mathbb{N}, \forall f \in \mathbb{X}.$$
(5)

The least constant verifying (5) is denoted by $C_p = C_p[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} is C_p -partially-greedy.

Remark 1.8. In [8], the condition of partially-greediness was introduced as follows:

$$\|f - \mathcal{G}_m(f)\| \le C \|f - S_m(f)\|, \ \forall m \in \mathbb{N}, \forall f \in \mathbb{X}.$$
(6)

Under the condition of Schauder bases, (6) and (5) are equivalent notions and in [4], the authors proved that if (6) is satisfied with C = 1, then the basis is partially-greedy.

Of course, every greedy basis is almost-greedy and every almost-greedy basis is partiallygreedy. One example of an almost-greedy basis that is not greedy is the Lindestrauss basis in ℓ_1 ([10]). Recently, one basis that is partially-greedy and not almost-greedy is presented in [7, Proposition 6.10].

It is well known that a basis is almost-greedy if and only if the basis is quasi-greedy and democratic and a basis is partially-greedy if the basis is quasi-greedy and conservative ([8, 7]). Moreover, as for greedy bases, we have the following characterizations.

Theorem 1.9. Assume that \mathcal{B} is a basis in a Banach space.

• \mathcal{B} is almost-greedy if and only if \mathcal{B} is democratic and quasi-greedy ([8]). Moreover,

$$\max\{C_q, \Delta_d\} \le C_{al} \le 8C_q^4 \Delta_d + C_q + 1.$$

• \mathcal{B} is almost-greedy if and only if \mathcal{B} is super-democratic and quasi-greedy ([6]). Moreover,

$$\max\{C_q, \Delta_s\} \le C_{al} \le C_q + C_q \Delta_s.$$

• \mathcal{B} is almost-greedy if and only if \mathcal{B} is symmetric for largest coefficients and quasi-greedy ([6]). Moreover,

$$\max\{C_q, \Delta\} \le C_{al} \le C_q \Delta.$$

Theorem 1.10. [4, 8] Assume that \mathcal{B} is a basis in a Banach space.

• \mathcal{B} is partially-greedy if and only if \mathcal{B} is conservative and quasi-greedy. Moreover,

 $\max\{C_q, \Delta_c\} \le C_p \le C_q + C_q^2 (1 + C_q) \Delta_c.$

• \mathcal{B} is partially-greedy if and only if \mathcal{B} is super-conservative and quasi-greedy. Moreover,

 $\max\{C_q, \Delta_{sc}\} \le C_p \le C_q + C_q(1+C_q)\Delta_{sc}.$

• \mathcal{B} is partially-greedy if and only if \mathcal{B} is partially-symmetric for largest coefficients and quasi-greedy. Moreover,

$$\max\{C_q, \Delta_{pc}\} \le C_p \le C_q \Delta_{pc}.$$

The purpose of this paper is to get a new characterization of almost-greedy and partiallygreedy bases following the ideas of [5, Corollary 1.8] for greedy bases.

Definition 1.11. We say that \mathcal{B} has the Property (F) if there is a positive constant C such that

$$\|f + 1_A\| \le C \|f + g + 1_B\|,\tag{7}$$

for any A, B, f, g satisfying the following conditions:

i) $|A| \leq |B| < \infty$ and $A \cap B = \emptyset$,

ii) $f, g \in \mathbb{X}_{fin}, f \cdot g = 0$, $\operatorname{supp}(f+g) \cap (A \cup B) = \emptyset$, $\|\tilde{f}\|_{\infty} \le 1$ and $\|\tilde{f}\|_{\infty} \le \inf_{n \in \operatorname{supp}(g)} |e_n^*(g)|$.

The least constant verifying (7) is denoted by $\mathcal{F} = \mathcal{F}[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} has the Property (F) with constant \mathcal{F} .

Also, if (7) is satisfied with the extra condition that $A < \operatorname{supp}(g) \cup B$, we say that \mathcal{B} has the Property (F_p) with constant \mathcal{F}_p .

Definition 1.12. We say that \mathcal{B} has the Property (F^{*}) if there is a positive constant C such that

$$\|f + z\| \le C \|f + y\|,\tag{8}$$

for any $f, z, y \in \mathbb{X}_{fin}$ satisfying the following conditions:

- i) $f \cdot z = 0, f \cdot y = 0, z \cdot y = 0,$
- ii) $\max\{\|\tilde{f}\|_{\infty}, \|\tilde{z}\|_{\infty}\} \le 1$.
- iii) $|D| \ge |\operatorname{supp}(z)|$, where $D = \{n \in \operatorname{supp}(y) : |e_n^*(y)| = 1\}$.
- iv) $\inf_{n \in \operatorname{supp}(y)} |e_n^*(y)| \ge ||f||_{\infty}$.

The least constant verifying (8) is denoted by $\mathcal{F}^* = \mathcal{F}^*[\mathbb{X}, \mathcal{B}]$ and we say that \mathcal{B} has the Property (F^{*}) with constant \mathcal{F}^* .

Also, if (8) is satisfied with the extra condition that $\operatorname{supp}(z) < \operatorname{supp}(f + y)$, we say that \mathcal{B} has the Property (F_p^*) with constant \mathcal{F}_p^* .

The main theorems that we will prove are the following.

Theorem 1.13. Let \mathcal{B} be a basis in a Banach space \mathbb{X} .

- a) If \mathcal{B} is almost-greedy with constant C_{al} , then \mathcal{B} has the Property (F^*) with constant $\mathcal{F}^* \leq C_{al}(1+2C_{al})$.
- b) If \mathcal{B} has the Property (F^*) with constant \mathcal{F}^* , then the basis is almost-greedy with constant $C_{al} \leq (\mathcal{F}^*)^2$.

Theorem 1.14. Let \mathcal{B} be a basis in a Banach space X.

- a) If \mathcal{B} is partially-greedy with constant C_p , then \mathcal{B} has the Property (F_p^*) with constant $\mathcal{F}_p^* \leq C_p(1+2C_p)$.
- b) If \mathcal{B} has the Property (F_p^*) with constant \mathcal{F}_p^* , then the basis is partially-greedy with constant $C_p \leq (\mathcal{F}_p^*)^2$.

The structure of the paper is the following: in Section 2, we will show some basics about the Properties (F) and (F^{*}). In Section 3, we prove Theorem 1.13. In Section 4 we give a brief summary about the Properties (F_p) and (F^{*}_p), in Section 5 we prove Theorem 1.14 and, finally, in Section 6, we give some density results that we use in the paper.

2. Properties (F) and (F^*)

This section is focused in the study of the Properties (F) and (F^*) . In fact, we will show that these properties are equivalent. To show that we will need some auxiliary lemmas about convexity.

Lemma 2.1. [1, Corollary 2.3] Let X be a Banach space, let \mathcal{B} be a basis for X and J a finite set.

(i) For any scalars $(a_j)_{j \in J}$ with $0 \le a_j \le 1$ and any $g \in \mathbb{X}$,

$$||g + \sum_{j \in J} a_j e_j|| \le \sup\{||g + 1_A|| : A \subseteq J\}.$$

(ii) For any scalars $(a_j)_{j \in J}$ with $|a_j| \leq 1$ and any $g \in \mathbb{X}$,

$$\|g + \sum_{j \in J} a_j e_j\| \le \sup_{\varepsilon \in \mathcal{E}_J} \|g + 1_{\varepsilon J}\|$$

Lemma 2.2. Let \mathcal{B} be a basis of a Banach space X. Then,

$$\sup_{\varepsilon \in \mathcal{E}_A} \|f + 1_{\varepsilon A}\| \le 5 \sup_{B \subseteq A} \|f + 1_B\|.$$

Proof. If $\mathbb{F} = \mathbb{R}$, following the result [5, Lemma 2.3], we know that

$$\sup_{\varepsilon_n=\pm 1} \|f + 1_{\varepsilon A}\| \le 3 \sup_{B \subseteq A} \|f + 1_B\|$$

We prove now the result for the complex case. In that case,

$$\mathbf{1}_{\varepsilon A} = \sum_{n \in A} \operatorname{Re}(\varepsilon_n) e_n + i \sum_{n \in A} \operatorname{Im}(\varepsilon_n) e_n
= \sum_{n \in A_1} \operatorname{Re}^+(\varepsilon_n) e_n - \sum_{n \in A_2} \operatorname{Re}^-(\varepsilon_n) e_n
+ i \left(\sum_{n \in A_3} \operatorname{Im}^+(\varepsilon_n) e_n - \sum_{n \in A_4} \operatorname{Im}^-(\varepsilon_n) e_n \right),$$
(9)

where A_i are the corresponding subsets of A. Then,

$$\begin{aligned} \|f + 1_{\varepsilon A}\| &\leq \|f + \sum_{n \in A} \operatorname{Re}(\varepsilon_n) e_n\| + \|\sum_{n \in A} \operatorname{Im}(\varepsilon_n) e_n\| \\ &\leq \|f\| + \|f + \sum_{n \in A_1} \operatorname{Re}^+(\varepsilon_n) e_n\| + \|f + \sum_{n \in A_2} \operatorname{Re}^-(\varepsilon_n) e_n\| \\ &+ \|f + \sum_{n \in A_3} \operatorname{Im}^+(\varepsilon_n) e_n\| + \|f + \sum_{n \in A_4} \operatorname{Im}^-(\varepsilon_n) e_n\| \\ &\leq \sup_{B \subseteq A} \|f + 1_B\|. \end{aligned}$$

Theorem 2.3. Let \mathcal{B} be a basis in a Banach space \mathbb{X} . The basis is democratic (or symmetric for largest coefficients) and quasi-greedy if and only if the basis has the Property (F). Concretely:

(1) If \mathcal{B} has the Property (F) with constant \mathcal{F} , then the basis is C_q -quasi-greedy and Δ_d -democratic with

$$\max\{C_q, \Delta_d\} \le \mathcal{F}.$$

(2) If \mathcal{B} has the Property (F) with constant \mathcal{F} , then the basis is C_q -quasi-greedy and Δ -symmetric for largest coefficients with

$$C_q \leq \mathcal{F}, \ \Delta \leq 5(\mathcal{F} + 4\mathcal{F}^2 + 4\mathcal{F}^3).$$

(3) If \mathcal{B} is Δ_d -democratic and C_q -quasi-greedy, then the basis has the Property (F) with constant

$$\mathcal{F} \le C_q (1 + (1 + C_q)\Delta_d).$$

(4) If \mathcal{B} is Δ -symmetric for largest coefficients and C_q -quasi-greedy, then the basis has the Property (F) with constant

$$\mathcal{F} \leq 3\Delta C_q.$$

Proof. First of all, we show (1). Assume that the basis has the Property (F) with constant \mathcal{F} . To show that \mathcal{B} is quasi-greedy, we take $f \in \mathbb{X}_{fin}$ with $t = \|\tilde{f}\|_{\infty}$ and $m \in \mathbb{N}$. Then, if we take in the definition of the Property (F) $f' = \frac{f}{t} - \mathcal{G}_m(\frac{f}{t}), g' = \mathcal{G}_m(\frac{f}{t})$ and $A = B = \emptyset$, since $\|\tilde{f}'\|_{\infty} \leq \inf_{n \in \text{supp}(g')} |e_n^*(g')|$, we obtain that

$$|f - \mathcal{G}_m(f)|| = t||f'|| \le \mathcal{F} t||f' + g'|| = \mathcal{F}||f||,$$

so the basis is quasi-greedy with $C_q \leq \mathcal{F}$ for elements with finite support. To obtain that \mathcal{B} is quasi-greedy for any $f \in \mathbb{X}$, we use Corollary 6.2.

Prove now that the basis is democratic. For that, we take C and D two finite sets such that $|C| \leq |D|$. Now, we do the following decomposition:

$$D = (D \cap C) \cup D_1 \cup D_2,$$

where $|D_1| = |C \setminus D|$ and $D_1 \cap D_2 = \emptyset$. Hence, taking $f = 1_{D \cap C}$, $g = 1_{D_2}$, $A = C \setminus D$ and $B = D_1$,

$$||1_C|| = ||1_{C \cap D} + 1_{C \setminus D}|| \le \mathcal{F} ||1_{D \cap C} + 1_{D_2} + 1_{D_1}|| = \mathcal{F} ||1_D||.$$

Thus, \mathcal{B} is democratic with $\Delta_d \leq \mathcal{F}$.

Prove now (2). We only have to show that \mathcal{B} is symmetric for largest coefficients. For that, take $f \in \mathbb{X}_{fin}$, $\|\tilde{f}\|_{\infty} \leq 1$, $A \cap B = \emptyset$, $|A| \leq |B| < \infty$, $\operatorname{supp}(f) \cap (A \cup B) = \emptyset$, $\varepsilon \in \mathcal{E}_A$ and $\eta \in \mathcal{E}_B$. Using Lemmas 2.2 and 2.1, we only have to show that there is some absolute constant C such that

$$||f + 1_{A'}|| \le C ||f + 1_{\eta B}||, \ \forall A' \subseteq A.$$

Of course, since the Property (F) implies quasi-greediness with constant $C_q \leq \mathcal{F}$ by (1), if we take the element $h := f + 1_{\eta B}$ with $\|\tilde{f}\|_{\infty} \leq 1$, we have

$$||f|| = ||h - \mathcal{G}_{|B|}(h)|| \le \mathcal{F}||h|| = \mathcal{F}||f + 1_{\eta B}||.$$
(10)

Also, respect to the set A', we can have the following:

$$\|1_{A'}\| \le \mathcal{F}\|1_B\| \le 4\mathcal{F}^2\|1_{\eta B}\|,\tag{11}$$

where in the last inequality we have used [2, Proposition 2.1.11] or [1, Lemma 3.2].¹

Thus,

$$\begin{aligned} \|f + 1_{A'}\| &\leq \|f\| + \|1_{A'}\| \leq_{(10)+(11)} \mathcal{F}\|f + 1_{\eta B}\| + 4\mathcal{F}^2 \|1_{\eta B}\| \\ &\leq (\mathcal{F} + 4\mathcal{F}^2) \|f + 1_{\eta B}\| + 4\mathcal{F}^2 \|f\| \\ &\leq (\mathcal{F} + 4\mathcal{F}^2 + 4\mathcal{F}^3) \|f + 1_{\eta B}\|. \end{aligned}$$

Finally, applying convexity,

$$\|f + 1_{\varepsilon A}\| \leq 5 \sup_{A' \subseteq A} \|f + 1_{A'}\| \leq 5(\mathcal{F} + 4\mathcal{F}^2 + 4\mathcal{F}^3)\|f + 1_{\eta B}\|.$$

So, the basis is symmetry for largest coefficients for elements with finite support with constant

$$\Delta \le 5(\mathcal{F} + 4\mathcal{F}^2 + 4\mathcal{F}^3).$$

Applying Lemma 6.5, the result follows for any $f \in X$.

¹These results affirm that for quasi-greedy bases, $||1_{\varepsilon A}|| \leq 2\kappa C_q ||1_{\eta A}||$, for any $\eta, \varepsilon \in \mathcal{E}_A$ and any finite set A with $\kappa = 1$ if $\mathbb{F} = \mathbb{R}$ and $\kappa = 2$ if $\mathbb{F} = \mathbb{C}$.

(3) Assume now that \mathcal{B} is C_q -quasi-greedy and Δ_d -democratic and take $f, g \in \mathbb{X}_{fin}$ with $f \cdot g = 0$, $\inf_{n \in \operatorname{supp}(g)} |e_n^*(g)| \ge \|\tilde{f}\|_{\infty}, A \cap B = \emptyset, |A| \le |B| < \infty$ and $\operatorname{supp}(f+g) \cap (A \cup B) = \emptyset$. $\|f + 1_A\| \le \|f\| + \|1_A\| \le \|f\| + \Delta_d \|1_B\|.$ (12)

If we take $h := f + g + 1_B$, it is clear that $\operatorname{supp}(g + 1_B)$ is a greedy set of h. Then, if $|\operatorname{supp}(g + 1_B)| = n$,

$$||f|| = ||h - \mathcal{G}_n(h)|| \le C_q ||h|| = C_q ||f + g + 1_B||.$$
(13)

Since $\inf_{n \inf_{n \in \text{supp}(g)}} |e_n^*(g)| \ge \|\tilde{f}\|_{\infty}$ and $\|\tilde{f}\|_{\infty} \le 1$, we can decompose g as $g = g_1 + g_2$, where $\supp(g_1) = \{n \in \supp(g) : |e_n^*(g)| \ge 1\}$ and $\supp(g_2) = \{n \in \supp(g) : |e_n^*(g)| < 1\}$. Then, if we take $u := f + g_2 + 1_B$, B is a greedy set for u with of order k := |B|, and taking $v = u + g_1$, $\supp(g_1)$ is a greedy set of v of order $p := |\supp(g_1)|$. Thus,

$$\|1_B\| = \|\mathcal{G}_k(u)\| \le (1+C_q)\|u\| = (1+C_q)\|f+g_2+1_B\|$$

= $(1+C_q)\|v-\mathcal{G}_p(v)\| \le (1+C_q)C_q\|f+g+1_B\|.$ (14)

Adding up (13) and (14) in (12), we obtain the result, that is, the basis has the Property (F) with $\mathcal{F} \leq C_q(1 + (1 + C_q)\Delta_d)$.

(4) Finally, assume that \mathcal{B} is Δ -symmetric for largest coefficients and C_q -quasi-greedy. Take f, g, A and B as in the Property (F). Then,

$$\begin{aligned} \|f + 1_A\| &\leq \Delta \|f + 1_B\| \leq \Delta (\|f + g_1 + 1_B\| + \|g_1 + f\| + \|f\|) \\ &\leq 3C_q \Delta \|f + g + 1_B\|. \end{aligned}$$

Thus, the basis has the Property (F) with constant $\mathcal{F} \leq 3C_q\Delta$.

Theorem 2.4. Let \mathcal{B} be a basis in a Banach space \mathbb{X} . The basis has the Property (F) if and only if the basis has the Property (F^{*}). Moreover, if \mathcal{F} and \mathcal{F}^* are the constants of the corresponding properties, then

$$\mathcal{F} \le \mathcal{F}^* \le 5\mathcal{F}(1+2\mathcal{F}+8\mathcal{F}^2).$$

Proof. Assume that we have the Property (F^{*}) with constant \mathcal{F}^* and take f, g, A and B as in the Property (F), that is, $f \cdot g = 0$, $A \cap B = \emptyset$, $|A| \leq |B|$, $\operatorname{supp}(f+g) \cap (A \cup B) = \emptyset$, $\|\tilde{f}\|_{\infty} \leq 1$ and $\|\tilde{f}\|_{\infty} \leq \inf_{n \in \operatorname{supp}(g)} |e_n^*(g)|$. Taking $z = 1_A$ and $y = g + 1_B$ in the Property (F^{*}), f, z and y verify the conditions established in the Property (F^{*}). Then,

$$||f + 1_A|| = ||f + z|| \le \mathcal{F}^* ||f + y|| = \mathcal{F}^* ||f + g + 1_B||,$$

so the basis has the Property (F) with $\mathcal{F} \leq \mathcal{F}^*$.

Assume now that we have the Property (F) and take f, y and z in \mathbb{X}_{fin} as in the Property (F^{*}), that is, $f \cdot z = 0$, $f \cdot y = 0$, $z \cdot y = 0$, $\max\{\|\tilde{f}\|_{\infty}, \|\tilde{z}\|_{\infty}\} \leq 1$ and $|\operatorname{supp}(z)| \leq |D|$ where $D = \{n : |e_n^*(y)| = 1\}|$. Using now Lemmas 2.2 and 2.1, it is enough to prove that there exists $C_1 > 0$ such that

$$||f + 1_{A'}|| \le C_1 ||f + y||, \ \forall A' \subseteq A,$$

where A = supp(z). Using Property (F), we have that

$$\|h\| \le \mathcal{F}\|h + w\|,\tag{15}$$

for any h and w such that $h \cdot w = 0$ and $\inf_{n \in \text{supp}(w)} |e_n^*(w)| \ge ||h||_{\infty}$.

Taking $D = \{n : |e_n^*(y)| = 1\}$, observe that $y = P_{D^e}(y) + 1_{\eta D}$, where $\eta \equiv \{\operatorname{sign}(e_n^*(y))\}_n$. Then, if $A' \subseteq A$,

$$\|f + 1_{A'}\| \le \mathcal{F}\|f + P_{D^c}(y) + 1_D\| \le \mathcal{F}\left(\|f + y\| + \|1_{\eta D}\| + \|1_D\|\right).$$
(16)

A FUNCTIONAL CHARACTERIZATION OF ALMOST-GREEDY AND PARTIALLY-BASES IN BANACH SPACES

If we decompose $D^c = D_1 \cup D_2$ such that

$$D_1 = \{n : |e_n^*(y)| < 1\}, \ D_2 = \{n : |e_n^*(y)| > 1\},\$$

we obtain that

$$\|1_{\eta D}\| \le \|f + P_{D_1}(y) + 1_{\eta D}\| + \|f + P_{D_1}(y)\| \le 2\mathcal{F}\|f + y\|.$$
(17)

Following the idea of (11), we can obtain that

$$\|1_D\| \le 4\mathcal{F}\|1_{\eta D}\| \le 8\mathcal{F}^2 \|f + y\|.$$
(18)

Using (17) and (18) in (16), we have

$$||f + 1_{A'}|| \le \mathcal{F}(1 + 2\mathcal{F} + 8\mathcal{F}^2)||f + y||.$$

Using now Lemma 2.1, we obtain

$$\|f + z\| \le \sup_{\varepsilon \in \mathcal{E}_A} \|f + 1_{\varepsilon A}\| \le \sup_{\text{Lemma } 2.2} 5 \sup_{A' \subseteq A} \|f + 1_{A'}\| \le 5\mathcal{F}(1 + 2\mathcal{F} + 8\mathcal{F}^2)\|f + y\|.$$

So the basis has the Property (F*) with $\mathcal{F}^* \leq 5\mathcal{F}(1+2\mathcal{F}+8\mathcal{F}^2)$.

To finish this section, we give the following nice characterization of the Property (F^*) that will be useful to show our main theorem.

Proposition 2.5. Let \mathcal{B} be a basis in a Banach space \mathbb{X} . The following are equivalent:

i) There is a positive constant C such that

$$||f + 1_{\varepsilon A}|| \le C ||f + g + 1_{\eta B}||, \tag{19}$$

for any $f, g \in \mathbb{X}_{fin}$ such that $f \cdot g = 0$, $\|\tilde{f}\|_{\infty} \leq 1$ and $\inf_{n \in supp(g)} |e_n^*(g)| \geq \|\tilde{f}\|_{\infty}$, for any pair of finite sets A and B such that $A \cap B = \emptyset$, $|A| \leq |B|$, $supp(f+g) \cap (A \cup B) = \emptyset$, and for any $\varepsilon \in \mathcal{E}_A$, $\eta \in \mathcal{E}_B$.

- ii) The basis has the Property (F^*) with constant \mathcal{F}^* .
- iii) There is a positive constant C such that

$$||f|| \le C||f - P_A(f) + y||, \tag{20}$$

for any $f, y \in X_{fin}$ with $f \cdot y = 0$ and $A \subseteq supp(f)$ verifying

- a) $\|\tilde{f}\|_{\infty} \leq 1$.
- b) $\inf_{n \in supp(y)} |e_n^*(y)| \ge \|\tilde{f}\|_{\infty}$. c) $|D| \ge |A|$, where $D = \{n \in supp(y) : |e_n^*(y)| = 1\}$.

Moreover, if we denote by C_1 and C_2 the least constants verifying (19) and (20) respectively, we have

$$\mathcal{F}^* \le C_1, \ C_2 \le \mathcal{F}^*, \ C_1 \le C_2$$

Proof. First, we prove i) \Rightarrow ii). Take $f, z, y \in \mathbb{X}_{fin}$ as in the definition of the Property (F^{*}):

- $f \cdot y = 0, \ f \cdot z = 0, \ z \cdot y = 0,$
- $\max\{\|f\|_{\infty}, \|\tilde{z}\|_{\infty}\} \le 1$.
- $|D| \ge |\operatorname{supp}(z)|$, where $D = \{n : |e_n^*(y)| = 1\}.$
- $\inf_{n \in \operatorname{supp}(y)} |e_n^*(y)| \ge \|\tilde{f}\|_{\infty}$.

If z = 0, just take $A = B = \emptyset$ and the prove is over. Consider now that $z \neq 0$ and take $\operatorname{supp}(z) = A$. If we divide $y = 1_{\eta D} + P_{D^c}(y)$ with $\eta \equiv \{\operatorname{sign}(e_n^*(y))\}$, we have for all $\varepsilon \in \mathcal{E}_A$,

$$||f + 1_{\varepsilon A}|| \le C_1 ||f + P_{D^c(y)} + 1_{\eta D}|| = C_1 ||f + y||$$
(21)

Applying now Lemma 2.1, we obtain the result with $\mathcal{F}^* \leq C_1$.

Now, we show that ii) \Rightarrow iii). Of course, if $A = \emptyset$, the result is trivial. Take f, y and A as in iii) with $A \neq \emptyset$ and $A \subseteq \text{supp}(f)$. If in the Property (F*) we take $f' = f - P_A(f)$, $z' = 1_{\varepsilon A}$ with $\varepsilon \in \mathcal{E}_A$ and y' = y,

$$||f' + z'|| = ||f - P_A(f) + 1_{\varepsilon A}|| \le \mathcal{F}^* ||f' + y'|| = \mathcal{F}^* ||f - P_A(f) + y||,$$

so applying the item (ii) of Lemma 2.1, iii) is proved with $C_2 \leq \mathcal{F}^*$.

Finally, we make the proof to show that iii) \Rightarrow i). Take $f, g \in \mathbb{X}_{fin}$ such that $f \cdot g = 0$, $\|\tilde{f}\|_{\infty} \leq \inf_{n \in \text{supp}(g)} |e_n^*(g)|, |A| \leq |B| < \infty, A \cap B = \emptyset$, $\sup_{n \in \mathcal{E}_B} (f + g) \cap (A \cup B) = \emptyset$ and $\varepsilon \in \mathcal{E}_A$, $\eta \in \mathcal{E}_B$.

Taking $f' = f + 1_{\varepsilon A}$ and $y = g + 1_{\eta B}$,

$$||f + 1_{\varepsilon A}|| = ||f'|| \le C_2 ||f' - P_A(f') + y|| = C_2 ||f + g + 1_{\eta B}||,$$

so the proof is over and $C_1 \leq C_2$.

3. Proof of Theorem 1.13

To prove Theorem 1.13, we will use one of the most important tools in the world of quasigreedy bases: the truncation operator. To define this operator, we take $\alpha > 0$ and define, first of all, the α -truncation of $z \in \mathbb{C}$:

$$T_{\alpha}(z) = \alpha \operatorname{sign}(z), \quad \text{if } |z| \ge \alpha,$$

and

$$T_{\alpha}(z) = z, \text{ if } |z| \leq \alpha.$$

Now, it is possible to extend T_{α} to an operator in the space X by

$$T_{\alpha}(f) = \sum_{n \in \text{supp}(f)} T_{\alpha}(e_n^*(f))e_n = \sum_{n \in \Delta_{\alpha}} \alpha \frac{e_n^*(f)}{|e_n^*(f)|}e_n + \sum_{n \notin \Delta_{\alpha}} e_n^*(f)e_n$$

where the set $\Delta_{\alpha} = \{n \in \mathbb{N} : |e_n^*(f)| > \alpha\}$. Of course, since Δ_{α} is a finite set, T_{α} is well-defined for all $f \in \mathbb{X}$.

Lemma 3.1. [6, Lemma 2.5] Let \mathcal{B} be a C_q -quasi-greedy basis in a Banach space. Then, the truncation operator is uniformly bounded that is,

$$||T_{\alpha}(f)|| \le C_q ||f||, \ \forall \alpha > 0, \ \forall f \in \mathbb{X}.$$

Proof of Theorem 1.13. Assume that \mathcal{B} is almost-greedy with constant C_{al} and take f, z and y as in the Property (F^{*}) and decompose $y = P_{B_1}(y) + P_{B_2}(y) + 1_{\eta B}$, where $\eta \equiv \{ \text{sign}(e_n^*(y)) \}, B_1 \cup B_2 = B^c$ and

$$B_1 = \{n : |e_n^*(y)| < 1\}, \ B_2 = \{n : |e_n^*(y)| > 1\}.$$

10

Taking now $h := f + 1_{\varepsilon A} + P_{B_2}(y) + 1_{\eta B}$ with $A = \operatorname{supp}(z), \varepsilon \in \mathcal{E}_A$ and $n = |B_2| + |B|$, we obtain

$$\begin{aligned} \|f + 1_{\varepsilon A}\| &= \|h - \mathcal{G}_n(h)\| \le C_{al} \|h - P_A(h)\| = C_{al} \|f + P_{B_2}(y) + 1_{\eta B}\| \\ &\le C_{al}(\|f + y\| + \|P_{B_1}(y)\|) \le C_{al}(\|f + y\| + \|f + P_{B_1}(y)\| + \|f\|) \\ &\le C_{al}(\|f + y\| + 2C_{al}\|f + y\|) \\ &\le C_{al}(1 + 2C_{al})\|f + y\|. \end{aligned}$$

Thus, applying Lemma 2.1, the basis has the Property (F^{*}) with constant $\mathcal{F}^* \leq C_{al}(1+2C_{al})$.

Assume now that the basis has the Property (F^{*}). Take $f \in \mathbb{X}_{fin}$, $m \in \mathbb{N}$, $\mathcal{G}_m(f) = P_G(f)$ and $|A| \leq m$.

Consider now the elements $f' = \frac{1}{t}(f - \mathcal{G}_m(f))$ with $t = \min_{n \in G \setminus A} |e_n^*(f)|$, $B = A \setminus G$, $y = 1_{\eta(G \setminus A)}$ and $\eta \equiv \{ \operatorname{sign}(e_n^*(f)) \}$. Of course, $f' \cdot y = 0$, $\|\tilde{f}'\|_{\infty} \leq 1$ since $|e_n^*(f - \mathcal{G}_m(f))| \leq t$ for $n \in G^c$ and $|G \setminus A| \geq |B|$. Then, applying these elements in the item iii) of Proposition 2.5, we obtain the following:

$$\|f - \mathcal{G}_{m}(f)\| = t \|f'\| \le t \mathcal{F}^{*} \|f' - P_{B}(f') + y\|$$

$$= \mathcal{F}^{*} \|f - P_{G}(f) - P_{A \setminus G}(f) + t \mathbf{1}_{\eta(G \setminus A)} \|$$

$$= \mathcal{F}^{*} \|P_{(A \cup G)^{c}}(f - P_{A}(f)) + t \mathbf{1}_{\eta(G \setminus A)} \|.$$
(22)

Since the Property (F^{*}) implies that the basis is quasi-greedy with $C_q \leq \mathcal{F}^*$ (Theorems 2.3 and 2.4), applying Lemma 3.1,

$$\|P_{(A\cup G)^c}(f - P_A(f)) + t1_{\eta(G\setminus A)}\| = \|T_t(f - P_A(f))\| \le \mathcal{F}^* \|f - P_A(f)\|.$$
(23)

Thus, by (23) and (22), the basis is almost-greedy with constant $C_{al} \leq (\mathcal{F}^*)^2$ for elements $f \in \mathbb{X}_{fin}$. Now, applying Corollary 6.3, the results follows.

4. PROPERTIES (F_p) AND (F_n^*)

In all the results presented in Section 2 we can change democracy by conservativeness or super-conservativeness and Property (F) and (F^{*}) by Property (F_p) and (F^{*}_p) and obtain the same results. Here, we only present the fundamental theorem that is the version of Theorem 2.3 to study how the constants change.

Theorem 4.1. A basis \mathcal{B} in a Banach space \mathbb{X} has the Property (F_p) if and only if \mathcal{B} is quasi-greedy and conservative. Moreover,

 $\max\{\Delta_c, C_q\} \le \mathcal{F}_p \le 2 + C_q + 2C_q\Delta_c.$

Proof. Assume that \mathcal{B} has the Property (F_p) with constant \mathcal{F}_p . Taking $A = \emptyset$, we have that

$$||f|| \le C||f + 1_B + g||,\tag{24}$$

for any f, g and B as in the definition of the Property (F_p). Now, taking $B = \emptyset$ and considering $f' := f - \mathcal{G}_m(f)$ and $y = \mathcal{G}_m(f)$,

$$||f - \mathcal{G}_m(f)|| = ||f'|| \le \mathcal{F}_p ||f' + g|| = \mathcal{F}_p ||f||,$$
(25)

so the basis is quasi-greedy for elements with finite support. Applying now Corollary 6.2, the basis is quasi-greedy with $C_q \leq \mathcal{F}_p$. Now, on the other hand, taking f = g = 0, we obtain conservativeness with constant $\Delta_c \leq \mathcal{F}_p$.

Now, take f, g, A and B as in the definition of Property (F_p). If we have $g = g_1 + g_2$ where

$$supp(g_1) = \{n \in supp(g) : |e_n^*(g)| < 1\}$$

$$\begin{split} \|f + 1_A\| &\leq \|f + g + 1_B\| + \|g + 1_B\| + \|1_A\| \\ &\leq 2\|f + g + 1_B\| + \|f\| + \|1_A\| \\ &\leq (2 + C_q)\|f + g + 1_B\| + \Delta_c\|1_B\| \\ &\leq (2 + C_q)\|f + g + 1_B\| + \Delta_c\|f + g_1 + 1_B\| + \Delta_c\|f + g_1\| \\ &\leq (2 + C_q)\|f + g + 1_B\| + 2C_q\Delta_c\|f + g + 1_B\| \\ &= (2 + C_q + 2C_q\Delta_c)\|f + g + 1_B\|. \end{split}$$

5. Proof of Theorem 1.14

Proof of Theorem 1.14. Assume now that \mathcal{B} is C_p -partially-greedy and prove the Property (\mathbb{F}_p^*) . Take $f, z, y \in \mathbb{X}_{fin}$ satisfying from i) to iv) in the definition of Property (\mathbb{F}_p^*) . We write $y = 1_{\eta D} + y_1 + y_2$, where

$$\operatorname{supp}(y_1) = \{n \in \operatorname{supp}(y) : |e_n^*(y)| < 1\}, \ \operatorname{supp}(y_2) = \{n \in \operatorname{supp}(y) : |e_n^*(y)| > 1\},\$$

and $\eta \equiv {\text{sign}(e_n^*(y))}$. Consider A = supp(z).

If $A = \emptyset$, applying Theorem 1.10, the basis is quasi-greedy with $C_q \leq C_p$ and we can conclude that $||f|| \leq ||f + y||$.

Assume now that $A \neq \emptyset$ and consider $m = \max A$ and define $B = \{1, \ldots, m\} \setminus A$. It is clear that $m = |A \cup B| \leq |B \cup D|$. Now, for any choice $\varepsilon \in \mathcal{E}_A$, define $h := f + 1_{\varepsilon A} + y_2 + 1_{\eta D} + 1_B$.

Since partially-greediness implies quasi-greediness with constant C_p (see Theorem 1.10), we have

$$\|f + 1_{\varepsilon A}\| = \|h - \mathcal{G}_m(h)\| \le C_p \inf_{k \le m} \|h - S_k(h)\| \le C_p \|f + y_2 + 1_{\eta D}\|$$

$$\le C_p (\|f\| + \|y_2 + 1_{\eta D}\|).$$

For the first element of the sum, consider w := f + y and we have

$$\|f\| = \|w - \mathcal{G}_n(w)\|_{\mathcal{F}}$$

with $n = |\operatorname{supp}(y)|$. Then, applying quasi-greediness, we obtain $||f|| \leq C_p ||f + y||$. For the second one, we write $w = f + y_1 + y_2 + 1_{\eta D}$ and using quasi-greediness, we have

$$||y_2 + 1_{\eta D}|| = ||\mathcal{G}_m(w)|| \le (1 + C_p)||f + y||,$$

where $m = |\operatorname{supp}(y_2) \cup D|$.

Using both bounds, we obtain $||f + 1_{\varepsilon A}|| \leq C_p (1 + 2C_p) ||f + y||$. Because of Lemma2.1, we conclude that $||f + z|| \leq C_p (1 + 2C_p) ||f + y||$.

Prove now b). Without loss of generality we can assume that $f \in \mathbb{X}_{fin}$ using Corollary 6.4 and that $\|\tilde{f}\|_{\infty} \leq 1$. Start considering $A = \operatorname{supp}(\mathcal{G}_m(f)), k \leq m$ and $B = \{1, \ldots, k\}$. If A = B, then the result is trivial. If $A \neq B$, we can decompose

$$f - \mathcal{G}_m(f) = P_{(A \cup B)^c}(f - S_k(f)) + P_{B \setminus A}(f).$$

Let $f' = \frac{1}{t} P_{(A \cup B)^c}(f - S_k(f))$ and $z = \frac{1}{t} P_{B \setminus A}(f)$ with $t = \min_{n \in A} |e_n^*(f)|$ and $y = 1_{\varepsilon(A \setminus B)}$ with $\varepsilon \equiv \{ \operatorname{sign}(e_n^*(f)) \}$. Of course, $f' \cdot z = 0$, $f' \cdot y = 0$ and $y \cdot z = 0$, $\|\tilde{f}'\|_{\infty} \leq 1$ since $|e_n^*(P_{(A \cup B)^c}(f - S_k(f)))| \leq t$ for $n \in (A \cup B)^c$ and $|A \setminus B| \geq |B \setminus A|$. Then, f', z and y verify the items of the Property (\mathbf{F}_p^*) , so

$$||f - \mathcal{G}_m(f)|| = t||f' + z|| \le t\mathcal{F}_p^*||f' + 1_{\varepsilon(A \setminus B)}|| = \mathcal{F}_p^*||P_{(A \cup B)^c}(f - S_k(f)) + t1_{\varepsilon(A \setminus B)}||.$$

A FUNCTIONAL CHARACTERIZATION OF ALMOST-GREEDY AND PARTIALLY-BASES IN BANACH SPACES

It turns out that

$$P_{(A\cup B)^c}(f - S_k(f)) + t1_{\varepsilon A\setminus B} = T_t(f - S_k(f)),$$

where T_t is the *t*-truncation operator. Now, since the Property (\mathbf{F}_p^*) implies the Property (\mathbf{F}_p) with the same constant, because of Theorem 4.1 and Corollary 6.2, the basis is quasi-greedy with constant $C_q \leq \mathcal{F}_p^*$. Then, applying Lemma 3.1, we have that $\|T_t(f - S_k(f))\| \leq \mathcal{F}_p^* \|f - S_k(f)\|$. All together, we obtain $\|f - \mathcal{G}_m(f)\| \leq (\mathcal{F}_p^*)^2 \|f - S_k(f)\|$ for all $k \leq m$ and hence, \mathcal{B} is partially-greedy.

6. Annex

In this annex, we write the main lemmas about density that we use in the paper.

Lemma 6.1. [3, Lemma 7.2] Let \mathcal{B} be a basis for a Banach space \mathbb{X} . If A is a greedy set for $f \in \mathbb{X}$, for every $\varepsilon > 0$, there is $y \in \mathbb{X}_{fin}$ such that $||f - y|| \le \varepsilon$ and A is a greedy set for y.

Corollary 6.2. Assume that \mathcal{B} is a C_q -quasi-greedy basis of a Banach space \mathbb{X} for elements with finite support. Then, \mathcal{B} is quasi-greedy for every $f \in \mathbb{X}$.

Proof. Take $f \in \mathbb{X}$ and A a greedy set of f with order m. By Lemma 6.1, there is $y \in \mathbb{X}_{fin}$ such that $||f - y|| \leq \varepsilon$ for every $\varepsilon > 0$ with A a greedy set for y. Then

$$\begin{aligned} \|f - P_A(f)\| &= \|f - y - P_A(f) + y - P_A(y) + P_A(y)\| \\ &\leq \|f - y\| + \|y - P_A(y)\| + \|P_A(f - y)\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_q \|y\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_q \|f - y\| + C_q \|f\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_q \|f - y\| + C_q \|f\|. \end{aligned}$$

Taking $\varepsilon \to 0$, we obtain the result.

Corollary 6.3. Let \mathcal{B} be a basis for a Banach space \mathbb{X} . If \mathcal{B} is an almost-greedy basis for all $f \in \mathbb{X}_{fin}$ with constant C_{al} , then the basis is almost-greedy for every $f \in \mathbb{X}$ with the same constant.

Proof. Assume that \mathcal{B} is almost-greedy for elements with finite support. Take $f \in \mathbb{X}$ with A a greedy set of order m. Applying Lemma 6.1, for any $\varepsilon > 0$, there is $g \in \mathbb{X}_{fin}$ such that $||f - g|| \leq \varepsilon$ and A a greedy set for g. Consider the set B_1 the set such that

$$\inf_{|B| \le m} \|f - P_B(f)\| = \|f - P_{B_1}(f)\|.$$

<u>Case 1:</u> $B_1 = \emptyset$.

$$\begin{aligned} \|f - P_A(f)\| &= \|f - g + g - P_A(f) - P_A(g) + P_A(g)\| \\ &\leq \|f - g\| + \|g - P_A(g)\| + \|P_A(f - g)\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_{al} \inf_{|B| \le m} \|g - P_B(g)\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_{al} \|g\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_{al} \|f - g\| + C_{al} \|f\| \\ &\leq \varepsilon (1 + \|P_A\|) + C_{al} \|f - g\| + C_{al} \|f\| \\ &\leq \varepsilon (1 + C_{al} + \|P_A\|) + C_{al} \|f\|. \end{aligned}$$

Taking $\varepsilon \to 0$, we obtain the result.

<u>Case 2:</u> $B_1 \neq \emptyset$.

$$\begin{aligned} \|f - P_A(f)\| &= \|f - g + g - P_A(f) - P_A(g) + P_A(g)\| \\ &\leq \|f - g\| + \|g - P_A(g)\| + \|P_A(f - g)\| \\ &\leq \varepsilon(1 + \|P_A\|) + C_{al} \inf_{|B| \le m} \|g - P_B(g)\| \\ &\leq \varepsilon(1 + \|P_A\|) + C_{al} \|g - P_{B_1}(g)\| \\ &\leq \varepsilon(1 + \|P_A\|) + C_{al} \|g - f + f - P_{B_1}(g) + P_{B_1}(f) - P_{B_1}(f)\| \\ &\leq \varepsilon(1 + \|P_A\|) + C_{al} \|P_{B_1}(f - g)\| + \|f - P_{B_1}(f)\| \\ &\leq \varepsilon(1 + \|P_A\| + C_{al}) + C_{al} \|P_{B_1}(f - g)\| + \|f - P_{B_1}(f)\| \\ &\leq \varepsilon(1 + \|P_A\| + C_{al} + C_{al} \|P_{B_1}\|) + \|f - P_{B_1}(f)\| \end{aligned}$$

Taking $\varepsilon \to 0$, we obtain the result.

With the same arguments, it is straightforward to show the next result.

Corollary 6.4. Let \mathcal{B} be a basis for a Banach space \mathbb{X} . If \mathcal{B} is a partially-greedy basis for all $f \in \mathbb{X}_{fin}$ with constant C_p , then the basis is partially-greedy for every $f \in \mathbb{X}$ with the same constant.

Lemma 6.5. [7, Lemma 3.2] Let \mathbb{X} be a Banach space. Suppose D is a finite subset of \mathbb{N} , and $f \in \mathbb{X} \setminus \{0\}$ satisfies $supp(f) \cap D = \emptyset$. Then, for any $\varepsilon > 0$ there is $y \in \mathbb{X}_{fin}$ so that $||f - y|| < \varepsilon$, $supp(y) \cap D = \emptyset$ and $||\tilde{f}||_{\infty} = ||\tilde{y}||_{\infty}$.

References

- F. ALBIAC, J. L. ANSORENA, P. M. BERNÁ, P. WOJTASZCZYK, Greedy approximation for biorthogonal systems in quasi-Banach spaces, Dissertatione Math. 560 (2021), 1–88.
- [2] P. M. BERNÁ, Thresholding Greedy Algorithms in Banach spaces, PhD Thesis. https://repositorio.uam.es/handle/10486/688914
- [3] M. BERASATEGUI, P. M. BERNÁ, Quasi-greedy bases for sequence with gaps, Nonlinear Analysis 208 (2021), 112294.
- M. BERASATEGUI, P. M. BERNÁ, S. LASSALLE, Strong Partially Greedy Bases and Lebesgue-Type Inequalities, Constr. Approx. (2021) https://doi.org/10.1007/s00365-021-09531-8
- [5] P. M. BERNÁ, O. BLASCO, Characterization of greedy bases in Banach spaces, J. Approx. Theory, 217 (2017), 28-39.
- [6] P. M. BERNÁ, O. BLASCO, G. GARRIGÓS, Lebesgue inequalities for greedy algorithm in general bases, Rev. Mat. Complut. 30 (2017), 369–392.
- [7] P. M. BERNÁ, S. J. DILWORTH, D. KUTZAROVA, T. OIKHBERG, B. WALLIS, *The weighted property* (A) and the greedy algorithm, J. Approx. Theory **248** (2019), 105300.
- [8] S. J. DILWORTH, N. J. KALTON, D. KUTZAROVA, AND V. N. TEMLYAKOV, The Thresholding Greedy Algorithm, Greedy Bases, and Duality, Constr. Approx. 19 (2003), 575–597.
- [9] S.J. DILWORTH, D. KUTZAROVA, E. ODELL, T. SCHLUMPRECHT, A. ZSAK, *Renorming spaces with greedy bases*, J. Approx. Theory **188** (2014), 39-56.
- [10] G. GARRIGÓS, E. HERNÁNDEZ, T. OIKHBERG, Lebesgue-type inequalities for quasi-greedy bases, Constr. Approx. 38 (3) (2013), 447–470.
- [11] S.V.KONYAGIN, V.N.TEMLYAKOV, A remark on greedy approximation in Banach spaces, East J. Approx. 5 (1999), 365-379.
- P.WOJTASZCZYK, Greedy algorithm for general biorthogonal systems, J.Approx. Theory 107 (2000), no.2, 293-314.

A FUNCTIONAL CHARACTERIZATION OF ALMOST-GREEDY AND PARTIALLY-BASES IN BANACH SPACES

PABLO M. BERNÁ, DEPARTAMENTO DE MATEMÁTICA APLICADA Y ESTADÍSTICA, FACULTAD DE CIENCIAS ECONÓMICAS Y EMPRESARIALES, UNIVERSIDAD SAN PABLO-CEU, CEU UNIVERSITIES, MADRID, 28003 SPAIN.

Email address: pablo.bernalarrosa@ceu.es

DIEGO MONDÉJAR, DEPARTAMENTO DE MATEMÁTICA APLICADA Y ESTADÍSTICA, FACULTAD DE CIENCIAS ECONÓMICAS Y EMPRESARIALES, UNIVERSIDAD SAN PABLO-CEU, CEU UNIVERSITIES, MADRID, 28003 SPAIN.

Email address: diego.mondejarruiz@ceu.es