On the ν -zeros of the modified Bessel function $K_{i\nu}(x)$ of positive argument

R. B. PARIS

Division of Computing and Mathematics, Abertay University, Dundee DD1 1HG, UK

Abstract

The modified Bessel function of the second kind $K_{i\nu}(x)$ of imaginary order for fixed x > 0 possesses a countably infinite sequence of real zeros. Recently it has been shown that the *n*th zero behaves like $\nu_n \sim \pi n/\log n$ as $n \to \infty$. In this note we determine a more precise estimate for the bahaviour of these zeros for large *n* by making use of the known asymptotic expansion of $K_{i\nu}(x)$ for large ν . Numerical results are presented to illustrate the accuracy of the expansion obtained.

Mathematics subject classification (2020): 33C10, 34E05, 41A30, 41A60

Keywords: modified Bessel function, imaginary order, zeros, asymptotic expansion, method of steepest descents

1. Introduction

The modified Bessel function of the second kind (the Macdonald function) $K_{i\nu}(x)$ of purely imaginary order and argument x > 0 is given by

$$K_{1\nu}(x) = \int_0^\infty e^{-x\cosh t} \cos\nu t \, dt = \frac{1}{2} \int_{-\infty}^\infty e^{-x\cosh t + i\nu t} dt.$$
(1.1)

In a recent paper, Bagirova and Khanmamedov [1] have shown that $K_{i\nu}(x)$ has a countably infinite number of (simple) real zeros in ν when x > 0 is fixed. We label the zeros $\nu_n \equiv \nu_n(x)$ (n = 0, 1, 2...) and observe that it is sufficient to consider only the case $\nu > 0$ since $K_{i\nu}(x) = K_{-i\nu}(x)$. By transforming the differential equation satisfied by $K_{i\nu}(x)$ into a one-dimensional Schrödinger equation with an exponential potential, these authors employed the well-known quantisation rule to deduce the leading asymptotic behaviour of the *n*th zero given by

$$\nu_n \sim \frac{\pi n}{\log n} \qquad (n \to +\infty).$$
(1.2)

In this note, we consider the behaviour of the large- $n \nu$ -zeros of $K_{i\nu}(x)$ in more detail. To achieve this we make use of the known asymptotic expansion of $K_{i\nu}(x)$ for $\nu \to +\infty$ stated in Section 2. A typical plot of $e^{\pi\nu/2}K_{i\nu}(x)$ as a function of ν is shown in Fig. 1.

2. Derivation of the equation describing the large-n zeros

We start with the asymptotic expansion of $K_{i\nu}(x)$ for $\nu \to +\infty$ given by [5] (see also¹ [4, pp. 41–42])

$$e^{\pi\nu/2}K_{i\nu}(x) \sim \Re \sqrt{\frac{2\pi}{\nu\tanh\mu}} e^{i\Phi} \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k C_k(\mu)}{(\frac{1}{2}i\nu\tanh\mu)^k},$$
 (2.1)

¹There is a misprint in (1.4.8) of [4]: the argument of the Bessel function should be x.



Figure 1: Plot of $e^{\pi\nu/2}K_{i\nu}(x)$ as a function of ν when x = 1.

where

$$\Phi := \nu(\mu - \tanh \mu) - \frac{\pi}{4}, \quad \cosh \mu = \frac{\nu}{x}$$

and $(a)_k = \Gamma(a+k)/\Gamma(a)$ denotes the Pochhammer symbol. The first few coefficients $C_k(\mu)$ are

$$C_0(\mu) = 1, \quad C_1(\mu) = -\frac{1}{8} + \frac{5 \coth^2 \mu}{24}, \quad C_2(\mu) = \frac{3}{128} - \frac{77 \coth^2 \mu}{576} + \frac{385 \coth^4 \mu}{3456};$$

values of $C_k(\mu)$ for $k \leq 5$ are derived in the appendix. It is important to point out that the determination of the expansion (2.1) from the integral (1.1) involves (when $\nu > x$) an infinite number of contributing saddle points. The expansion (2.1) results from the two dominant saddles, the remaining saddles yielding an exponentially small contribution of $O(e^{-\pi\nu})$. Then, from (2.1), the zeros of $K_{i\nu}(x)$ are given asymptotically by

$$\cos \Phi \left\{ 1 - \frac{3C_2(\mu)}{(\nu \tanh \mu)^2} + \frac{105C_4(\mu)}{(\nu \tanh \mu)^4} + \cdots \right\} + \sin \Phi \left\{ \frac{C_1(\mu)}{\nu \tanh \mu} - \frac{15C_3(\mu)}{(\nu \tanh \mu)^3} + \frac{945C_5(\mu)}{(\nu \tanh \mu)^5} + \cdots \right\} = 0.$$
$$\Phi = (n + \frac{1}{2})\pi + \epsilon, \qquad (2.2)$$

Let

where n is a large positive integer and ϵ is a small quantity. Then

$$\begin{aligned} \tan \epsilon &= \frac{\frac{C_1(\mu)}{\nu \tanh \mu} - \frac{15C_3(\mu)}{(\nu \tanh \mu)^3} + \frac{945C_5(\mu)}{(\nu \tanh \mu)^5} + \cdots}{1 - \frac{3C_2(\mu)}{(\nu \tanh \mu)^2} + \frac{105C^4(\mu)}{(\nu \tanh \mu)^4} + \cdots} \\ &= \frac{1}{\nu \tanh \mu} \bigg\{ C_1(\mu) + \frac{3(C_1(\mu)C_2(\mu) - 5C_3(\mu))}{(\nu \tanh \mu)^2} \\ + \frac{9C_1(\mu)C_2^2(\mu) - 45C_2(\mu)C_3(\mu) - 105C_1(\mu)C_4(\mu) + 945C_5(\mu)}{(\nu \tanh \mu)^4} + \cdots \bigg\}, \end{aligned}$$

so that using $\arctan z = z - z^3/3 + z^5/5 - \dots (|z| < 1)$, we obtain

$$\epsilon = \frac{a_0(\mu)}{\nu} + \frac{a_1(\mu)}{\nu^3} + \frac{a_2(\mu)}{\nu^5} + \cdots,$$

where

$$a_0(\mu) = \frac{C_1(\mu)}{\tanh \mu}, \qquad a_1(\mu) = \frac{3(C_1(\mu)C_2(\mu) - 5C_3(\mu)) - \frac{1}{3}C_1^3(\mu)}{\tanh^3 \mu},$$

$$a_{2}(\mu) = \frac{1}{\tanh^{5}\mu} \left\{ 3(3C_{1}(\mu)C_{2}^{2}(\mu) - 15C_{2}(\mu)C_{3}(\mu) - 35(C_{1}(\mu)C_{4}(\mu) - 9C_{5}(\mu))) - 3C_{1}^{2}(\mu)(C_{1}(\mu)C_{2}(\mu) - 5C_{3}(\mu)) + \frac{1}{5}C_{1}^{5}(\mu) \right\}.$$

Since $\tanh \mu = \sqrt{1 - x^2/\nu^2}$, the coefficients $a_k(\mu)$ possess expansions in inverse powers of ν^2 . Some laborious algebra shows that

$$a_{0}(\mu) = \frac{1}{12} + \frac{x^{2}}{4\nu^{2}} + \frac{11x^{4}}{32\nu^{4}} + O(\nu^{-6})$$

$$a_{1}(\mu) = \frac{1}{360} - \frac{x^{2}}{4\nu^{2}} + O(\nu^{-4})$$

$$a_{2}(\mu) = \frac{1}{1260} + O(\nu^{-2}),$$

whence we obtain

$$\epsilon = \frac{1}{12\nu} + \frac{1}{\nu^3} \left(\frac{1}{360} + \frac{x^2}{4} \right) + \frac{1}{\nu^5} \left(\frac{1}{1260} - \frac{x^2}{4} + \frac{11x^4}{32} \right) + \cdots$$
(2.3)

Making use of the expansion

$$\mu - \tanh \mu = \log \,\lambda\nu + \frac{x^2}{4\nu^2} + \frac{x^4}{32\nu^4} + \frac{x^6}{96\nu^6} + O(\nu^{-8}), \qquad \lambda := \frac{2}{ex},$$

we then finally obtain from (2.2) and (2.3) the equation describing the large-*n* zeros of $K_{i\nu}(x)$ given by

$$\nu \log \lambda \nu = m + \frac{A_0}{\nu} + \frac{A_1}{\nu^3} + \frac{A_2}{\nu^5} + \cdots,$$
 (2.4)

where

$$A_0 = \frac{1}{12} - \frac{x^2}{4}, \quad A_1 = \frac{1}{360} + \frac{x^2}{4} - \frac{x^4}{32}, \quad A_2 = \frac{1}{1260} - \frac{x^2}{4} + \frac{11x^4}{32} - \frac{x^6}{96}$$

and, for convenience, we have put $m := (n + \frac{3}{4})\pi$.

3. Solution of the equation for the zeros

To solve (2.4) we expand ν as

$$\nu_n = \xi + \frac{c_0}{\xi} + \frac{c_1}{\xi^3} + \frac{c_2}{\xi^5} + \cdots,$$

where the c_k are constants to be determined and we suppose that ξ is large as $n \to \infty$. Substitution in (2.4) then produces

$$\xi \log \lambda \xi + \frac{c_0(1+\log \lambda \xi)}{\xi} + \frac{c_1(1+\log \lambda \xi) + \frac{1}{2}c_0^2}{\xi^3} + \frac{c_2(1+\log \lambda \xi) + c_0c_1 - \frac{1}{6}c_0^3}{\xi^5} + \cdots$$
$$= m + \frac{A_0}{\xi} + \frac{A_1 - A_0c_0}{\xi^3} + \frac{A_2 - 3A_1c_0 + A_0(c_0^2 - c_1)}{\xi^5} + \cdots$$

Equating coefficients of like powers of ξ , we obtain

$$\xi \log \lambda \xi = m, \tag{3.1}$$

and

$$c_0 = \frac{A_0}{1 + \log \lambda \xi}, \quad c_1 = \frac{A_1 - A_0 c_0 - \frac{1}{2} c_0^2}{1 + \log \lambda \xi},$$

$$c_2 = \frac{A_2 - 3A_1c_0 + A_0(c_0^2 - c_1) - c_0c_1 + \frac{1}{6}c_0^3}{1 + \log\lambda\xi} .$$

The solution of (3.1) for the lowest-order term ξ can be expressed in terms of the Lambert W function, which is the (positive) solution² of $W(z)e^{W(z)} = z$ for z > 0. Rearrangement of (3.1) shows that

$$\frac{m}{\xi} e^{m/\xi} = \lambda m,$$

$$\xi = \frac{m}{W(\lambda m)}.$$
(3.2)

whence

The asymptotic expansion of W(z) for $z \to +\infty$ is [2], [3, (4.13.10)]

$$W(z) \sim L_1 - L_2 + \frac{L_2}{L_1} + \frac{L_2^2 - 2L_2}{2L_1^2} + \frac{(6 - 9L_2 + 2L_2^2)L_2}{6L_1^3} + \frac{(-12 + 36L_2 - 22L_2^2 + 3L_2^3)L_2}{12L_1^4} + \cdots$$

where $L_1 = \log z$, $L_2 = \log \log z$, from which it follows that

$$\frac{1}{W(z)} \sim \frac{1}{L_1} \left\{ 1 + \frac{L_2}{L_1} + \frac{L_2(L_2 - 1)}{L_1^2} + \frac{(1 - 5L_2 + 2L_2^2)L_2}{2L_1^3} + \frac{(-6 + 21L_2 - 26L_2^2 + 6L_2^3)L_2}{6L_1^4} + \cdots \right\}.$$

Then we have the expansion for ξ as $\lambda m \to \infty$ given by

$$\xi \sim \frac{m}{\log \lambda m} \bigg\{ 1 + \frac{\log \log \lambda m}{\log \lambda m} + \frac{\log \log \lambda m (\log \log \lambda m - 1)}{(\log \lambda m)^2} + \cdots \bigg\},\tag{3.3}$$

where we recall that $m = (n + \frac{3}{4})\pi$.

If we define $\chi := \xi/m$, so that by (3.1) $1 + \log \lambda \xi = (1 + \chi)/\chi$ and introduce the coefficients

$$B_0 = \frac{A_0}{1+\chi}, \quad B_1 = \frac{A_1 - A_0c_0 - \frac{1}{2}c_0^2}{\chi^2(1+\chi)}, \quad B_2 = \frac{A_2 - 3A_1c_0 + A_0(c_0^2 - c_1) - c_0c_1 + \frac{1}{6}c_0^3}{\chi^4(1+\chi)}, \quad (3.4)$$

then we finally have the result:

Theorem 1. The expansion for the nth ν -zero of $K_{i\nu}(x)$ for fixed x > 0 is

$$\nu_n \sim \frac{m}{W(\lambda m)} + \frac{B_0}{m} + \frac{B_1}{m^3} + \frac{B_2}{m^5} + \dots \qquad (n \to \infty),$$
(3.5)

where $m = (n + \frac{3}{4})\pi$, $\lambda = 2/(ex)$ and the coefficients B_k are given in (3.4).

4. Numerical results

The leading form of ξ from (3.3) is, with $\lambda = 2/(ex)$,

$$\xi \sim \frac{(n+\frac{3}{4})\pi}{\log(n+\frac{3}{4})\pi + \log\lambda} \qquad (n \to \infty),$$

which yields the approximate estimate in (1.2). In numerical calculations it is found more expedient to use the expression for ξ in terms of the Lambert function in (3.2), rather than (3.3), since the asymptotic scale in this latter series is $\log \lambda m$ and so requires an extremely large value of n to attain reasonable accuracy.

We present numerical results in Table 1 showing the zeros of $K_{i\nu}(x)$ computed using the FindRoot command in *Mathematica* compared with the asymptotic values determined from the expansion (3.5) with coefficients B_k , $k \leq 2$, where ξ is evaluated from (3.2). The value of the zeroth-order approximation $\xi = m/W(\lambda m)$ is shown in the final column. It is seen that there is excellent agreement with the computed zeros, even for n = 1.

²In [3, p. 111] this is denoted by Wp(z).

n	$ u_n$	Asymptotic	ξ
1	4.5344907181	4.5345024086	4.550063
2	5.8798671997	5.8798689800	5.890918
4	8.2589364092	8.258936 5 588	8.265990
5	9.3550938258	9.3550938 8 60	9.361083
10	14.3318529171	14.33185291 9 8	14.335296
15	18.8230418511	18.8230418514	18.825473
20	23.0318794957	23.031879495 8	23.033764
30	30.9169674670	30.9169674670	30.918273

Table 1: Values of the zeros of $K_{i\nu}(x)$ and their asymptotic estimates when x = 1.

Appendix: The coefficients $C_k(\mu)$ in the expansion (2.1)

The integral in (1.1) can be cast in the form

$$K_{i\nu}(x) = \frac{1}{2} \int_{-\infty}^{\infty} e^{-x\psi(t)} dt, \qquad \psi(t) := \cosh t - it \cosh \mu,$$
 (A.1)

where $\cosh \mu = \nu/x$. In the right-half plane, the dominant saddle point (where $\psi'(t) = 0$) is situated at $t_0 = \mu + \frac{1}{2}\pi i$, with a similar saddle in the left-half plane at $-\mu + \frac{1}{2}\pi i$. When $\nu > x$, there are in addition two infinite strings of subdominant saddles at $\pm \mu + (2k + \frac{1}{2})\pi i$ (k = 1, 2, ...) parallel to the positive imaginary axis over which the integration path in (A.1) is deformed; see [5], [4, pp. 41–42] for details.

We introduce the new variable w by

$$w^{2} = \psi(t) - \psi(t_{0}) = \frac{1}{2}i\sinh\mu(t-t_{0})^{2} + \frac{1}{6}i\cosh\mu(t-t_{0})^{3} + \frac{1}{24}i\sinh\mu(t-t_{0})^{4} + \cdots$$

to find with the help of *Mathematica* using the InverseSeries command (essentially Lagrange inversion)

$$t - t_0 = e^{-\pi i/4} \sqrt{\frac{2}{\sinh \mu}} w + \frac{i \coth \mu}{2 \sinh \mu} w^2 - i e^{-\pi i/4} \sqrt{\frac{2}{\sinh \mu}} \frac{(-3 + 5 \coth^2 \mu)}{36 \sinh \mu} w^3 + \cdots$$

Differentiation then yields the expansion

$$\frac{dt}{dw} \stackrel{e}{=} e^{-\pi i/4} \sqrt{\frac{2}{\sinh \mu}} \sum_{k \ge 0} \frac{C_k(\mu)}{(\frac{1}{2}i \sinh \mu)^k} w^{2k}, \tag{A.2}$$

where $\stackrel{e}{=}$ signifies the inclusion of *only the even powers of* w, since odd powers will not enter into this calculation. The first few coefficients are found to be:

$$C_{0}(\mu) = 1, \qquad C_{1}(\mu) = -\frac{1}{8} + \frac{5 \coth^{2} \mu}{24},$$

$$C_{2}(\mu) = \frac{3}{128} - \frac{77 \coth^{2} \mu}{576} + \frac{385 \coth^{4} \mu}{3456},$$

$$C_{3}(\mu) = -\frac{5}{1024} + \frac{1521 \coth^{2} \mu}{25600} - \frac{17017 \coth^{4} \mu}{138240} + \frac{17017 \coth^{6} \mu}{248832},$$

$$C_{4}(\mu) = \frac{35}{32768} - \frac{96833 \coth^{2} \mu}{4300800} + \frac{144001 \coth^{4} \mu}{1720320} - \frac{1062347 \coth^{6} \mu}{9953280} + \frac{1062347 \coth^{8} \mu}{23887872},$$

$$C_{5}(\mu) = -\frac{63}{262144} + \frac{67608983 \coth^{2} \mu}{8670412800} - \frac{35840233 \coth^{4} \mu}{796262400} + \frac{3094663 \coth^{6} \mu}{31850496}$$

$$-\frac{154040315 \coth^8 \mu}{1719926784} + \frac{154040315 \coth^{10} \mu}{5159780352}.$$
 (A.3)

Then the integral (A.1) resulting from the saddle t_0 can be expressed as

$$\int_{-\infty}^{\infty} e^{-xw^2} \, \frac{dt}{dw} \, dw$$

which, after substitution of the expansion (A.2) followed by routine integration and taking into account the contribution from the saddle at $t = -\mu + \frac{1}{2}\pi i$, then yields the expansion for $K_{i\nu}(x)$ stated in (2.1).

References

- S.M. Bagirova and A. Kh. Khanmamedov, On zeros of the modified Bessel function of the second kind, Comput. Math. and Math. Phys. 60 (2020) 817–820.
- [2] R.M. Corless, G.H. Gonnet, D.E.G. Hare, D.J. Jeffrey and D.E. Knuth, On the Lambert W function, Adv. in Comp. Math. 5 (1996) 329–359.
- [3] F.W.J. Olver, D.W. Lozier, R.F. Boisvert and C.W. Clark (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press, Cambridge, 2010.
- [4] R.B. Paris, Hadamard Expansions and Hyperasymptotic Evaluation, Encyclopedia of Mathematics and its Applications Vol. 141, Cambridge University Press, Cambridge, 2011.
- [5] N.M. Temme, Steepest descent paths for integrals defining the modified Bessel functions of imaginary order, Methods Appl. Anal. 1 (1994) 14-24.