A new type of mirror symmetry in the set of protein amino acids

Miloje M. Rakočević

University of Niš, Faculty of Sciences and Mathematics, Department of Chemistry, Višegradska 33, 18000 Niš, Serbia

Abstract. In several previous works, I presented the mirror symmetry in the set of protein amino acids, expressed through the number of atoms. Here, however, the same thing is shown but over the number of nucleons and molecules mass.

The new type of the mirror symmetry, in the set of protein amino acids (AAs), which is the matter in this brief communication, was go in the form of a hint in a previous my work (MMR, 2004, Table 1, p. 223). But then it was not allowed to say something like that, because it would be declared as numerology. Now when that symmetry is the factual state, expressed though the number of atoms (Box 1), it makes sense to represent it also through the number of nucleons (Table 1 in relation to Figure 1).

We look now at Tables 1 and 2 together with Figure 1 and Survey 1. If we place a mirror at the middle row of Table 1, either from above or from below, we always see one and the same sequence-image; for the number of nucleons: 299, 289, 189 and for the molecule mass: 596, 586, 485, kao što je na Figure 1 naznačeno.⁴ This is directly obvious mirror symmetry, but it is existing a hidden one. The full meaning of balancing and nuancing, expressed through distinctions and differences between quantities, it appears only when the original and/or the mirror image of the double value of the third perfect

¹ The word is about a new type of mirror symmetry, in relation to enantiomer symmetry of amino acid molecules (MMR, 2018a, Solution 1, p. 42).

² It is the fact, however, that this new type of symmetry I have presented in some of the previous papers published only in preprint form; also once in the official journal, in a regular article (MMR, 2018a, at the end of page 38, in relation to Figure 4, p. 37.).

³ We are talking here about the set of standard protein amino acids that have the status of "canonical amino acids" because their incorporation into proteins is mediated by the genetic code.

⁴ In the middle row, in Table 1 and Figure 1, respectively, the number of nucleons 299 and the quantity of mass 596 irresistibly "invoke" the third perfect number: the quantity 596 because of the difference 596 - 496 is equal to 100, and the number 299 because that record is actually a mirror image of the double value of the third perfect number. That was the hint in my work from 17 years ago (MMR, 2004, Tab. 1), which, as I said, I was not allowed to refer to.

number is included: the image as in Table 2, and original as in Table 3; both Tables taken together as a single system).⁵

Box 1. Mirror symmetry through the number of atoms as a unity of numerical quantities and chemical qualities

Start: a unique sequence of Periodic System of Numbers (PSN) in front of a mirror, with superposition; then "crossing" with a unique 6-bit binary tree sequence (Box 2): 010/101 as 2/5 (3 x 2 and 2 x 5):

$$00-11-22/22-11-00 \rightarrow 00-11-22-11-00 \rightarrow 02, 13, 24, 16, 05$$

Then mirroring with the result as in Display 1 and Fig. 2 (MMR, 2019, Fig. 2, p. 17 and Tab. A3, p. 30; 2021, Fig. 2, p. 55 and Fig. A2, p. 88). PSN as in (MMR, 2019, Fig. A1, p. 28).

```
Display 1

G 01 + A 04 + N 08 + D 07 \rightarrow 20

V 10 + P 08 + S 05 + T 08 \rightarrow 31

I 13 + L 13 + C 05 + M 11 \rightarrow 42

K 15 + R 17 + F 14 + Y 15 \rightarrow 61

Q 11+ E 10 + W 18 + H 11 \rightarrow 50
```

Box 2. The determination of the series of the numbers 0-63 on the 6-bit binary tree

"The determination of the series of the numbers 0-63 [on the 6-bit binary tree]. When we look closely into the structure of the sequence 0-63 of the series of the natural numbers we come to the obvious and self-evident explanation of the reason why the genetic code must be six-bit code, no matter if it is the manifestation in the form of the Gray Code model (Swanson, 1984, p 188), or it is in the form of the Binary tree (Rakočević, 1994, p. 38) [Rakočević, 1998, Fig. 1, p. 284]. There must be 8 codon i.e. amino acid classes. The structure of the sequence 0-63 is strictly determined by third perfect number (496) and the sum chonsisted of the first pair of the friendly numbers (220 + 284 = 504) ..." (Rakočević, 1997b, Fig. 7, p. 60) www.rakocevcode.rs

⁵ That perfect and friendly numbers are determinants of the genetic code I have repeatedly explained, especially in a monograph from 25 years ago (MMR, 1997b, pp. 60-64). www.rakocevcode.rs

Remark 1. One can note in Table 2, in the area of the quantities of molecules mass, the difference **100** (bold) in the middle row, and in area of nucleons number the difference **000** (bold). The difference of difference is 100. On the other hand, in Survey 1 we find that the difference between the sums and the sums in Table 2 is also 100. As we see the principle of self-similarity is at work also here as well.

Remark 2. About "strange result" in Figure 1, where from differences of the quantities of molecules mass and the number of nucleons follows the number of atoms in the AAs molecules it is difficult to give any explanation. Instead that, we give only the facts. In the previous paper (MMR, Table 3a, p. 224) we showed that in the first two columns of GCT there are 196 + 100 = 296 atoms; the other two columns have 154 + 144 = 298 atoms. It is also indicated that in the first two columns there are 297 - 1 and in the other two columns there are 297 + 1 atoms.

Remark 3. The three equations we find reviewed in Survey 2 (very bottom, right) are also found in the standard GCT (MMR, 2004, Table 3a, p. 224).

*

Our expectations are that over time it will turn out that the presented results are strong support for two hypotheses and one prediction. The first hypothesis is the hypothesis of Vladimir Shcherbak, according that "The laws of additive-positions of numbers ... have analogies with quantum physics" (Shcherbak, 1994, p. 476, last passage). The second hypothesis is our hypothesis given in the previous paper, whose this brief communication is a specific supplement: "Hypothesis on a complete [prebiotic] genetic code" (MMR, 2004, Section 7.1, p. 231). The prediction we give here for the first time boils down to the expectation that future researches will show that the view of mirror symmetry, presented here, has a direct or indirect connection with the choice of only the left amino acid enantiomers in the act of the generating of life.

References

Rakočević, M.M. (1994) Logic of the Genetic Code, Naučna knjiga, Beograd. www.rakocevcode.rs)

Rakočević, M.M. (1997a) Two classes of the aminoacyl-tRNA synthetases in correspondence with the Codon path cube, Bull. Math. Biol. Vol. 59, No. 4, pp. 645-648.

Rakočević, M.M. (1997b) Genetic Code as a Unique System, Studentski kulturni centar, Niš. http://www.rakocevcode.rs

Rakočević, M.M. (1998) The genetic code as a Golden mean determined system, Biosystems, 46, 283–291.

Rakočević, M.M. (2004) A harmonic structure of the genetic code. J. Theor. Biol. 229, 221–234.

Rakočević, M.M. (2011) Genetic code as a coherent system, NeuroQuantology, 9 (4), 821–841. http://www.rakocevcode.rs

Rakočević, M.M. (2018a) The Cipher of the Genetic Code, BioSystems 171 (2018) 31–47.

Rakočević, M. M. (2018b) Analogies of Genetic and Chemical Code, Polyhedron, 153, 292–298.

Rakočević, M.M. (2019) Protein amino acids as a complete (periodic) system, Chemia Naissensis, Vol 2, Issue 1, pp. 1-43.

hhttps://www.pmf.ni.ac.rs/chemianaissensis/archives/volume-2-number-1-2019/.

Rakočević, M.M. (2021) System-directed pairing of protein amino acids. Part I, DOI 10.31219/osf.io/bpqzt

Shcherbak, V. I. (1994) Sixty-four Triplets and 20 Canonical Amino Acids of the Genetic Code: The Arithmetical Regularities. Part II. J. Theor. Biol. 166, 475-477.

Swanson, R. (1984) A unifying concept for the amino acid code. Bull. Math. Biol. 46, 187–207.

Tables

Table 1. "The harmonic structure with two 'acidic' and three 'basic' amino acid quartets"

	Amino	acids	n	Molecule mass	
D 59	N 58	A 15	L 57	189	485.49 ≈ 485
R 100	F 91	P 41	l 57	289	585.70 ≈ 586
K 72	Y 107	T 45	M 75	299	595.71 ≈ 596
H 81	W 130	S 31	C 47	289	585.64 ≈ 586
E 73	Q 72	G 01	V 43	189	485.50 ≈ 485
				2738.04	
				1255	2738 = 2 x (37 x 37)

Minimally reduced and technically adjusted Table from previous work (MMR, 2004, Table 1, p. 223). Designation (n): The number of nucleons within 20 AAs side chains, calculated from the first, the lightest nuclide (H-1, C-12, N-14, O-16, S-32); One can notice that molecule mass within five rows is realized through the same logic-patterns of notations as the first nuclide, i.e. isotope.

Table 2. The number of nucleons and the quantities of molecules mass in Table 1, in relation to the third perfect number and the mirror image of its double value

992 - <u>0</u> 11	189		(496 - 0 <u>1</u> 0)	485	
992 - 0 <u>1</u> 0	289	<u>1</u> 00	(496 + 0 <u>9</u> 0)	586	1 <u>0</u> 1
992 ± 000	299	0 <u>1</u> 0	496 + 100	596	0 <u>1</u> 0
992 - 0 <u>1</u> 0	289	0 <u>1</u> 0	(496 + 090)	586	0 <u>1</u> 0
992 - <u>0</u> 11	189	<u>1</u> 00	(496 - 0 <u>1</u> 0)	485	1 <u>0</u> 1
042		220	300		222

The sums of differences (the sums of absolute values) correspond to two numbers (220 and 300) in the "logical square" in Figure 2 [(0). 204, (1). 220, (2). 284, (3). 300]. The other two vertices off the logic square, the numbers 284 and 204, exist through the relation with the golden mean, as shown in Survey 1, below right, in relation to Survey 2.

Table 3. The number of nucleons and the quantities of molecules mass in Table 1, in relation to the third perfect number and its double value

496 - <u>307</u>	189		(992 - 507)	485	
		<u>1</u> 00			1 <u>0</u> 1
496 - 207	289		(992 - 406)	586	
		0 <u>1</u> 0			0 <u>1</u> 0
496 - 197	299	0.10	992 - 396	596	0.10
406 207	200	0 <u>1</u> 0	(002 406)	506	0 <u>1</u> 0
496 - 207	289	100	(992 - 406)	586	1 <u>0</u> 1
496 - <u>307</u>	189	<u>1</u> 00	(992 - 507)	485	1 <u>0</u> 1
170 <u>507</u>	10)		(222 301)	103	
1225		220	2222		222

$$(1225 + 220 = 1445) / (2222 + 222 = 2444) \rightarrow 999$$

Last result in Shcherbak's Table: 27 x 37 = 999 (Shcherbak, 1994, Table 1, p. 476)

The relatively small changes in the status of the third perfect number in this Table in relation to Table 2, lead to the large changes in the result. Here it does not follow from the logical square in Figure 2 but follows from two specific arithmetical system-arrangements, presented in Tables 3.1 and 3.2.

Table 3.1. *Specific Arithmetical System-Arrangement* (SASA I)

0	15	+	0	=	15
1	1 2 5	+	20	=	145
2	1225	+	220	=	1445
3	12225	+	2220	=	14445
0	15	+	9	=	24
1	1 4 5	+	99	=	244
2	1445	+	999	=	2444
3	14445 	+	9999	=	24444

This arithmetical system-arrangement, taken together with the results in Table 3, is a proof more that the key determinant of the Multiplication Table in the decimal number system (3 x 5 = 15) also the determinant of the genetic code (MMR, 2021a, Survey 8 and Tables C2 and C3). The numbers 2 and 4 are emphasized in both subsystems, upper and lower, to indicate the fact that the ratio of number 4 and its half, the number 2, is manual in the genetic code as we have shown in previous works, as also follows from the system-arrangement shown in Table 3.2 (MMR, 2018b, Eq. II in Solution 3, on page 293).

Table 3.2. Specific Arithmetical System-Arrangement (SASA II)

1	2	+	22	=	24
2	22	+	222	=	244
3	222	+	2222	=	2444
4	2222	+	22222	=	24444

For the explanation see Legend of Table 3.1.

Surveys

Survey 1. The relationships of quantities contained in Table 2

$$(042 + 300 = 342) (220 + 222 = 442) [442 - 342 = 100]$$

$$(220 + 284 = 504) (300 + 204 = 504)^{a}$$

$$(504 - 342 = 162) (504 - 442 = 062)^{b}$$

$$[162 + 062 = 8 \times 28]^{c} [162 = 666 - 504] [162 = 2 \times 81]^{d}$$

$$100: 162 = 0.617 \dots \approx 0.618 \dots (Golden mean)$$

$$342 + 442 = 784$$

$$784 = 28 \times 28$$

$$384 = 204 + 180$$

$$(300 + (2 \times 042) = 384)^{e}$$

$$(300 + (2 \times 042) = 384)^{e}$$

$$384 = 496 - (4 \times 28)$$

$$(222 = 284 - 062)^{b}$$

$$042 = 204 - 162$$

$$0.62 \approx 0.618 \dots$$

$$1.62 \approx 1.618 \dots$$

and other two are indirectly present through a relation with the golden mean. b Mediation by golden mean. c Mediation with the Golden mean and athe second perfect number, the number 28. d The quantities (81 versus 123) as the number of atoms in two classes of AAs, classified according to two classes of the enzymes aminoacyl-tRNA synthetases. [MMR, 2004, Fig. 1, p. 222: "Notice that '81' (as 9 x 9) is the first possible (zeroth) arithmetic square in module 9, and 0-1-2-3 is the first possible (zeroth) logical square (as 00-01-10-11)."] [About the said two classes of AAs one can see in (MMR, 1997a).] c The quantity 384 as the total number of atoms in 20 standard protein amino acids (204 in "bodies" and 180 in "heads") is also found in Plato's Timaeus as the main result of the "harmonization" of geometric progression (MMR, 2011, Table A.2., p. 839).

Survey 2. Golden mean, expressed in different number systems, in relation to the quantities in Tables 2 and 3

Sum of (I)	Golden mear	ı			
•••			504 - 162 = 342		
$(770)_{8}$	$(1.474)_8$		504 052 442		
(70.1)	(4.640)		504 - 062 = 442		
$(504)_{10}$	$(1.618)_{10}$				
(100)	(1.0E2		242 - 200 + 042		
$(1F8)_{16}$	$(1.9E3)_{16}$		342 = 300 + 042		
HTH FFN C' . C	. 11	1	440 220 222		
"I" \rightarrow The first f	riendly pair of num	ıb.	442 = 220 + 222		
		25	$256 = 16 \times 16$		
222 + 042	= 264	25	$256 = 8 \times 32$		
220 + 300	= 264 + 256				
		26	$264 = 8 \times 33$		
384 = 256 +	1/2 256	$297 = 9 \times 33$			
		33	$0 = 10 \times 33$		

For the three equations listed in Remark 3 (very bottom, right), the arithmetical template can be also found as in (MMR, 2018a, Survey 2, positions 8, 9 and 10) or in (MMR, 2021a, Table C10, with corrected cipher errors in line 3 and 27).

Figures

189 289 299 299 289 289 189 189	296 297 297 297 296	485 586 596 586 485	485 586 596 586 485
--	---------------------------------	--	--

Figure 1. The mirror symmetry of the quantities contained in Table 1. In the middle area there are the differences of the quantities of molecules mass and the number of nucleons per row, respectively. These differences simultaneously correspond to the quantities of the number of atoms in 61 amino acid molecules within the standard Genetic Code Table (GCT), from the aspect of pyrimidine-purine distinctions: on the left pyrimidine half 32 amino acid molecules have 296 atoms and on the right purine half 29 amino acid molecules have 298 atoms; which means that the mean value of the left and right sides is 297.

(0)	(1)	(2)	(3)	(4)	(5)	(6)		
00	02	04	06	08	10	12		
11	13	15	17	19	21	23		
22	24	26	28	30	32	34		
11	16	21	26	31	36	41		
00	05	10	15	20	25	30		
44	60	76	92	108	124	140		
	12	14	16	18	20	22		
	23	25	27	29	31	33		
	34	36	38	40	42	44		
	41	46	51	56	61	66		
	30	35	40	45	50	55		
	140	156	172	188	<u>204</u>	<u>220</u>		
	22	24	26	28	30	/32		
	33	35	37	39	41	43		
	44	46	48	50	52	54		
	66	71	76	81	86 /	91		
	55	60	65	70	75/	80		
	220	236	252	268	284	300		
(10) 204, (11) 220 / (15) 284, (16) 300								
(2	(220 + 284 = 504) (204 + 300 = 504)							

Figure 2. "The arrangement represents the Table of distinct 2-5 adding (TDA) with starting column 00- 11-22-11-00 which follows from PSN (Periodic system of numbers: Figure A1) in decimal number system by overlapping the real sequence of doubled the first possible triangle in Boolean space (0-1-2) with its mirror image through compression and superposition at the point '22'. In the 10th step we have a realization of the sequence (20-31-42-61-50), the same with the number of atoms in five AAs classes ... as it is here presented: all five results in the 10th step are mirror image of the first step." (MMR, 2011, Table A3, p. 30).