# Backward Stability of the Schur Decomposition under Small Perturbation

A. Minenkova, E. Nitch-Griffin, V. Olshevsky<sup>a</sup>

<sup>a</sup>University of Connecticut

# Abstract

In the present paper, we show the backward stability of the Schur decomposition for a given matrix under small perturbation.

*Keywords:* perturbation theory, Schur decomposition, unitary Hessenberg matrices, the Gohberg-Kaashoek numbers, invariant subspaces, gaps.

#### 1. Introduction

Lipschitz-Hölder stability was investigated for several canonical forms like Jordan, flipped-orthogonal, flipped-orthogonal conjugare symmetrical, real canonical forms [1, 3, 5, 6, 7, 8, 13, 14], but it was never addressed for the Schur decomposition.

We begin by recalling some classical results.

#### 1.1. Eigenvalues' Stability

The first question to consider is what happens to the eigenvalues of a given matrix under small perturbation. In general it might not be true but if we impose additional restrictions on eigenvalues of matrices then we have Lipschitz stability of the eigenvalues.

The following result can be found in [1].

**Proposition 1.1.** Let  $A_0$  be an  $n \times n$  matrix and  $\{\lambda_1, \ldots, \lambda_n\}$  be its eigenvalues, and A being its perturbation with  $||A - A_0|| < \varepsilon$  for sufficiently small  $\varepsilon$  depending on  $A_0$  and the eigenvalues  $\mu_j$ . If the number of eigenvalues of  $A_0$  is the same as of A, then there is a certain ordering of them such that for some positive  $K = K(A_0)$ 

$$|\mu_i - \lambda_i| \le K ||A - A_0||, \quad i = 1, 2, \dots, |\sigma(A_0)|.$$

For the general case of the eigenvalues stability we have the following result (see [11, Appendix K]).

Preprint submitted to TBD

October 29, 2021

**Proposition 1.2.** Let  $A_0$  be an  $n \times n$  matrix and  $\{\lambda_1, \ldots, \lambda_n\}$  be its eigenvalues. Then, there is an ordering of  $\lambda_j$ 's that for every A with  $||A - A_0|| < \varepsilon$  for sufficiently small  $\varepsilon$  depending on  $A_0$  there is an ordering of its eigenvalues  $\mu_j$ 's and a positive constant  $K = K(A_0)$  such that

$$|\mu_j - \lambda_j| \le K ||A - A_0||^{1/n}.$$
(1.1)

This type of bounds is called Hölder because of the power 1/n for the matrix norm.

The following example shows that the power 1/n in (1.1) cannot be relaxed and, in general, we can hope only for a Hölder type bound.

**Example 1.3.** Consider the following matrices  $A_0, A \in \mathbb{C}^{2 \times 2}$ .

$$A_0 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \text{ and } A = \begin{bmatrix} 0 & \epsilon \\ 1 & 0 \end{bmatrix}$$

Note that  $||A - A_0|| = \epsilon$ . Moreover,  $\sigma(A_0) = \{0\}$  and  $\sigma(A) = \{\pm \sqrt{\epsilon}\}$ . It is easy to see that in this case we have

$$|0 \mp \sqrt{\epsilon}| = \epsilon^{1/2} = ||A - A_0||^{1/2}.$$

This example can be easily modified for  $n \times n$ -matrices.

#### 1.2. Backward Stability of the Schur Decomposition

Every  $n \times n$ -matrix A is unitary similar to an upper triangular matrix T, i.e,  $A = UTU^*$  where U is unitary. This triangular matrix T is called a Schur Triangular form and the factorization is called the Schur Decomposition.

Note that diagonal entries of T are the eigenvalues of A. That is why the eigenvalues stability results give us the confidence to consider stability of the Schur decomposition.

But what kind of stability can we have for the Schur canonical form? We start by considering the following type of result.

**Conjecture 1.4** (Forward Stability). Let  $A_0 = U_0 T_0 U_0^* \in \mathbb{C}^{n \times n}$  where  $U_0$  is unitary and  $T_0$  is upper triangular. Then, there exist constants  $K, \epsilon > 0$  (depending on  $A_0$  only) such that for all A with  $||A - A_0|| < \epsilon$  there exists a factorization  $UTU^*$  of A such that

$$||U - U_0|| + ||T - T_0|| \le K ||A - A_0||^{1/n}$$

We call this property forward stability of the Schur form.

As the following example shows this conjecture is not valid in the form stated.

**Example 1.5.** Consider the following matrix and its perturbation,

$$A_0 = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \text{ and } A = \begin{bmatrix} 2 & 0 \\ \epsilon & 2 \end{bmatrix}.$$

Let us consider the following Schur factorization of  $A_0$ 

$$A_0 = U_0 T_0 U_0^* = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

and the Schur factorization of A

$$A = UTU^* = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & \epsilon \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Since the first column of U has to be an eigenvector of A, and the latter is essentially unique, the matrix U is essentially unique as well. Hence, the distance  $||U - U_0||$  is quite large. Since U was the only possible choice for triangulating A, we can conclude that our Conjecture 1.4 above is false in general. Although forward stability results have been obtained for other canonical forms (see [1, 3]), in case of the Schur canonical form we cannot have the stability mentioned in Conjecture 1.4. The next statement shows us why.

**Theorem 1.6** (Different Gohberg-Kaashoek Numbers). Let us fix matrix  $A_0$  and its fixed Schur decomposition  $A_0 = U_0 T_0 U_0^*$ . There exists K > 0 such that in any neighborhood of  $A_0$ , i.e.  $\{A : ||A - A_0|| < \varepsilon\}$  for any  $\varepsilon > 0$ ,

$$\sup_{A} \inf_{U,T} \|U - U_0\| + \|T - T_0\| > M > 0,$$
(1.2)

where the supremum is taken over all A in this neighborhood having different Gohberg-Kaashoek numbers from  $A_0$  and the infimum is taken over all their Schur factorizations  $A = UTU^*$ .

#### 1.2.1. Gohberg-Kaashoek Numbers

Theorem 1.6 uses Gohberg-Kaashoek (GK) numbers. Let us introduce these numbers now (see [4, 9, 10] for details).

Let  $A \in \mathbb{C}^{n \times n}$ ,  $\sigma(A)$  be the set of all its eigenvalues, and  $m_1(A, \lambda) \leq m_2(A, \lambda) \leq \cdots \leq m_t(A, \lambda)$  be the sizes of all blocks corresponding to  $\lambda \in \sigma(A)$  in the Jordan form of A. We set  $m_i(A, \lambda) = 0$   $(i = t + 1, \ldots, n)$  for convenience. The numbers

$$m_i(A) = \sum_{\lambda \in \sigma(A)} m_i(A, \lambda)$$

are called the Gohberg-Kaashoek numbers.

We can actually prove a more general result, for this we need to define the dual Gohberg-Kaashoek numbers.  $[m_1]$ 

Let 
$$m = \begin{bmatrix} m_1^2 \\ \vdots \\ m_n \end{bmatrix}$$
 be a vector with integer entries such that  $m_i \ge m_{i+1}$  for  
= 1, ...,  $n - 1$ . The vector  $k = \begin{bmatrix} k_1 \\ k_2 \\ \vdots \\ k_n \end{bmatrix}$  with  $k_i = \max_{1 \le l \le n} \{l : m_l \ge i\}$  is called

dual to m.

i

In terms of the Gohberg-Kaashoek numbers  $m_j$ 's it means that if we have

	$\lambda$	1	0	0					
	0	$\lambda$	1	0					
	0	0	$\lambda$	1					
	0	0	0	$\lambda$					
A =					λ	1	0		
					0	$\lambda$	1		
					0	0	$\lambda$		
								$\lambda$	1
								0	λ

then we can put the Jordan chains corresponding to  $\lambda$  in the following order.

						$m_j(A,\lambda)$
	e <sub>4</sub>	$\rightarrow e_3$ –	e <sub>2</sub>	$\rightarrow e_1 \rightarrow e_1$	→ <b>0</b>	4
	$e_7$ –	$ ightarrow \mathbf{e_6}$ —	$e_5$ –	→ <b>0</b>		3
	$e_9$ –	$ ightarrow \mathbf{e_8}$ –	ightarrow 0			2
$k_i$	3	3	2	1		

Therefore, for  $m(A) = (4, 3, 2, 0, 0, 0, 0, 0, 0)^{\top}$  the dual is going to be  $k(A) = (3, 3, 2, 1, 0, 0, 0, 0, 0)^{\top}$ . These numbers were introduced in [4], where the problem of complete description for the Jordan structure of a matrix, which is a small perturbation of a given matrix, was posed. This problem was solved independently in [2] and [8].

#### 1.2.2. Backward Stability


Although we are unable to obtain a general forward stability result, we can get the backward stability result.

**Theorem 1.7** (Backward Stability). Let  $A_0 \in \mathbb{C}^{n \times n}$  be given. There exist constants  $K, \epsilon > 0$  (depending on  $A_0$  only) such that for all A with  $||A - A_0|| < \epsilon$  and for any factorization  $UTU^*$  of A (U unitary and T is upper triangular) there exist  $U_0$  and  $T_0$  such that  $A_0 = U_0 T_0 U_0^*$  is a Schur factorization of  $A_0$  with

$$||U - U_0|| + ||T - T_0|| \le K ||A - A_0||^{1/n}.$$
(1.3)

# 1.2.3. Organization of the Paper

In Section 2 we consider some auxillary results about what happens to GK numbers after we apply a reduction step. In Section 3 we discuss facts related to unitary Hessenberg matrices. In Section 4 we give a short overview about theory of gaps and semigaps. In Section 5 and 6 we present the proofs of Theorem 1.7 and Theorem 1.6 respectively.



# 2. Auxiliary Results

Before proving Theorem 1.6 we need a couple of technical lemmas. Let us start by introducing the following fact.

**Lemma 2.1.** For every eigenvector x of A there is a Jordan basis of A including x.

*Proof.* Let us fix any Jordan basis of A, with the Jordan chains corresponding to  $\lambda_t$  ordered by length. Now, given another eigenvector x decompose it in that basis. Look for the last non-zero coefficient, say,  $\alpha$  that is corresponding to the eigenvector, say, y.

Then the chain for x has the same length as for y and we can replace the chain for y with the chain for x. All it remains to prove is the linear independence of the new set of vectors.

Let Y stands for the matrix whose columns are the Jordan basis we started with. Y is invertible. Denote by X the matrix where the chain for y is replaced by the chain for x. Then X = YR, where R is an upper triangular matrix that is invertible, since it has either 1 on its diagonal or  $\alpha$ . Note that for the generalized eigenvectors of the chain for x we have the same decompositions with the same coefficients as for x with the vectors from the corresponding chains for the original basis, so we can write down the matrix R.

The next result describes the recursion we will use. In particular, we want to figure out what happens to the GK numbers during each step of recursion. Here is the idea behind it:

$$m_1(A,\lambda_t) \ge m_2(A,\lambda_t) \ge m_3(A,\lambda_t) \ge \ldots \ge m_{l-1}(A,\lambda_t) \ge m_l(A,\lambda_t)$$
  
The corresponding Jordan chains stay the same.  
$$m_j(A,\lambda_t) < m_l(A,\lambda_t) \text{ for } j > l$$

So what happens when  $m_j(A, \lambda_t) = m_l(A, \lambda_t)$  for some j's greater than l? Let  $j^*$  be the maximal such index.

Recursion decreases this chain by one vector.

$$\overline{m_l(A,\lambda_t)} = m_{l+1}(A,\lambda_t) = \ldots = m_{j^*}(A,\lambda_t) > \overline{m_l(A,\lambda_t)} - 1$$

Now let us formalize it.

**Lemma 2.2.** Consider matrix B with the eigenvalues  $\{\lambda_j\}$ 's, having the GK numbers  $\{m_j(B,\lambda_i)\}$  and  $e_1$  as its eigenvector corresponding to the Jordan chain for  $\lambda_t$  and  $m_l(B,\lambda_t)$ , i.e.

$$B = \begin{bmatrix} \lambda_t & \star & \cdots & \star \\ 0 & & \\ \vdots & C & \\ 0 & & \end{bmatrix}$$

Then

- $m_j(C, \lambda_i) = m_j(B, \lambda_i)$  for all  $i \neq t$  or i = t and j > l + 1;
- $m_l(C,\lambda_t) = m_l(B,\lambda_t) 1, \ m_{l+1}(C,\lambda_t) = m_{l+1}(B,\lambda_t) \ \text{if } m_l(A,\lambda_t) > m_{l+1}(B,\lambda_t);$
- $m_{j^*}(C,\lambda_t) = m_l(B,\lambda_t) 1$ ,  $m_j(C,\lambda_t) = m_{j+1}(B,\lambda_t)$  for  $j = l, \ldots, j^* 1$  if  $m_l(B,\lambda_t) = m_{l+1}(B,\lambda_t) = \ldots = m_{j^*}(B,\lambda_t)$  and  $j^*$  is the maximal such index;
- $m_j(C,\lambda_t) = m_j(B,\lambda_t)$  if  $m_j(B,\lambda_1) < m_l(B,\lambda_t)$ .

*Proof.* Note that due to Lemma 2.1 there is a Jordan basis of B containing  $e_1$ . Let J be the canonical Jordan form of B where the first block corresponds to the Jordan chain for  $\lambda_t$  that we mentioned. Thus, there is a invertible matrix R containing the Jordan basis  $\{f_{i,j}^{(k)}\}_{i,j,k}$  (*i* is the place in the Jordan chain for  $f_{i,j}^{(k)}$  corresponding to  $m_j(B,\lambda_k)$ ) as its columns, where the first  $m_l(B,\lambda_t)$  vectors forms the chain of A, having  $f_{0,l}^{(t)} = e_1$ , i.e.  $R = \left[f_{0,l}^{(t)}|f_{1,l}^{(t)}|\dots|f_{m_l(B,\lambda_t)-1,l}^{(t)}|\dots\right]$ . That is R and  $R^{-1}$  are of the following form.

$$R = \begin{bmatrix} 1 & \bullet & \cdots & \bullet \\ 0 \\ \vdots \\ 0 \end{bmatrix} \quad \text{and} \quad R^{-1} = \begin{bmatrix} 1 & \bullet & \cdots & \bullet \\ 0 \\ \vdots \\ 0 \end{bmatrix} R_1^{-1}$$

This argument implies that

=

Note that  $J_1 = R_1^{-1}CR_1$  is the Jordan form of C.

So what is the difference between J and  $J_1$ ? The only Jordan chain that is affected is

$$0 \leftarrow f_{0,l}^{(t)} \leftarrow f_{1,l}^{(t)} \leftarrow \ldots \leftarrow f_{m_l(B,\lambda_t)-1,l}^{(t)}$$

We delete the eigenvector from this chain and truncate the rest of the vectors to get a Jordan chain of length  $m_l(B, \lambda_t) - 1$  of C. The length of the rest Jordan chains of C stay the same as they were in B. The conclusion of the lemma follows from this observation.

## 3. Unitary Hessenberg Matrices and Schur Canonical Forms

Throughout this paper we are going to use the special type of structured matrices that are called Hessenberg. So, let us introduce it to the reader first.

A matrix is called the *upper Hessenberg* if it has zero entries below the first subdiagonal. Similarly, it is called the *lower Hessenberg* if it has zeros above the first super diagonal.

The following is a well-known fact (for example see [12]).

**Proposition 3.1.** An  $n \times n$  lower unitary Hessenberg matrix can be repre-

sented in the following way

$$U = \begin{bmatrix} -\rho_1 & \mu_1 & 0 & \dots & 0\\ -\rho_2\mu_1 & -\rho_2\bar{\rho}_1 & \mu_2 & \dots & 0\\ \vdots & \vdots & \vdots & \ddots & \vdots\\ \vdots & \vdots & \ddots & 0\\ \vdots & \vdots & \vdots & \ddots & 0\\ \vdots & \vdots & \vdots & \ddots & \mu_{n-1}\\ -\rho_n\mu_{n-1}\dots\mu_1 & -\rho_n\mu_{n-1}\dots\mu_2\bar{\rho}_1 & -\rho_n\mu_{n-1}\dots\mu_3\bar{\rho}_2 & \dots & -\rho_n\bar{\rho}_{n-1} \end{bmatrix},$$

where  $\mu_j = \sqrt{1 - \rho_j^2}$  for all j's.

The following statement is an immediate consequence of the previous proposition.

**Corollary 3.2.** If  $\rho_j < 1$  for all j, the first column of a lower unitary Hessenberg matrix completely defines the whole matrix.

Let us consider the following properties of Hessenberg matrices first. Unitary Hessenberg matrices have a number of interesting properties, and are of particular importance in their relationship with the Schur form. In the classical proof of the construction of the Schur form, one builds an orthonormal set using eigenvectors of the matrix  $A_0$ , typically through the Gram-Schmidt process. With the above observation, we can derive the Schur form specifically through the use of unitary Hessenberg matrices.

**Lemma 3.3.** For any  $A \in \mathbb{C}^{n \times n}$  there exist U unitary and T upper triangular with the eigenvalues of A along the diagonal such that  $A = UTU^*$ .

*Proof.* Let  $\lambda_1, \ldots, \lambda_m$  be the eigenvalues of A and let x be a unit eigenvector of A corresponding to eigenvalue  $\lambda_1$ . Moreover, pick  $H_1$  to be a lower unitary Hessenberg matrix with x as its first column. By Proposition 3.1 this determines  $H_1$  completely.

Then, we have  $H_1^*AH_1e_1 = H_1^*Ax = H_1^*\lambda_1x = \lambda_1e_1$ . In other words,

$$H_1^* A H_1 = \begin{bmatrix} \lambda_1 & \star & \cdots & \star \\ 0 & & \\ \vdots & & A_2 \\ 0 & & \end{bmatrix}.$$
(3.5)

By repeating the process of reducing the matrix dimensions, i.e. for each matrix  $A_k$  constructing matrix  $H_k$  in a way we described, we get a string of matrices  $H_1, \ldots, H_n$  that are all unitary Hessenberg and

$$\widetilde{H}_{n-1}^* \dots \widetilde{H}_1^* A_0 \widetilde{H}_1 \dots \widetilde{H}_{n-1} = T_0, \qquad (3.6)$$

Where  $T_0$  is an upper triangular matrix,  $\widetilde{H}_1 = H_1$ , and  $\widetilde{H}_k = \begin{bmatrix} I_{k-1} & \mathbf{0} \\ \mathbf{0} & H_k \end{bmatrix}$ for  $k = 2, \ldots, n-1$  with  $I_j$  being the  $j \times j$  identity matrix. By taking  $U = \widetilde{H}_1 \cdot \ldots \cdot \widetilde{H}_{n-1}$  we get the result.

Observe that we constructed our unitary matrix U using only unitary Hessenberg matrices  $H_k$ , each of whose first column was an eigenvector of the corresponding matrix  $A_k$ .

Lemma 3.4. Every unitary matrix U admits a factorization

$$U = H_1 \cdot \ldots \cdot H_{n-1},$$

where matrices  $H_i$ 's are unitary Hessenberg.

*Proof.* Let  $H_1$  be the unitary Hessenberg matrix whose first column  $x_1$  is the same as U. Note that  $x_1^*x_j = \delta_{1,j}$ , since U is unitary. Then

$$H_1^* U = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & U_2 \\ 0 & & & \end{bmatrix},$$
(3.7)

where  $U_2$  is unitary. As before we repeat the process until we get

$$H_{n-1}^* \cdot \ldots \cdot H_1^* U = I.$$

The result follows from simply multiplying the both parts of this equality by  $H_1 \cdot \ldots \cdot H_{n-1}$ .

# 4. Gap and Semi-gap

In this section we discuss some topological properties of the set of subspaces in  $\mathbb{C}^n$ , since in order to prove our main result, we require some facts from the theory of gaps. We begin by stating some definitions.

A matrix  $P_{\mathcal{M}}$  is called an *orthogonal projector* onto a subspace  $\mathcal{M} \subset \mathbb{C}^n$  if

- $\operatorname{Im} P_{\mathcal{M}} = \mathcal{M};$
- $P^2_{\mathcal{M}} = P_{\mathcal{M}};$
- $P^*_{\mathcal{M}} = P_{\mathcal{M}}$ .

The following concept is the key definition.

Let  $\mathcal{M}, \mathcal{N}$  be subspaces of  $\mathbb{C}^n$ , and let  $P_{\mathcal{M}}, P_{\mathcal{N}}$  be the orthogonal projectors onto  $\mathcal{M}$  and  $\mathcal{N}$  respectively. We define the gap  $\theta(\mathcal{M}, \mathcal{N})$  between  $\mathcal{M}$ and  $\mathcal{N}$  as follows

$$\theta(\mathcal{M}, \mathcal{N}) = \|P_{\mathcal{M}} - P_{\mathcal{N}}\|$$

or, equivalently,

$$\theta(\mathcal{M}, \mathcal{N}) = \max \left\{ \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \inf_{\substack{y \in \mathcal{N} \\ \|y\|=1}} \|x - y\|, \sup_{\substack{y \in \mathcal{N} \\ \|y\|=1}} \inf_{\substack{x \in \mathcal{M} \\ \|y\|=1}} \|x - y\| \right\}.$$

It follows immediately from the definition that  $\theta(\mathcal{M}, \mathcal{N})$  is a metric on the set of all subspaces in  $\mathbb{C}^n$ . Moreover,  $\theta(\mathcal{M}, \mathcal{N}) \leq 1$ .

Note that the Hausdorff distance between sets Inv A and Inv B of all invariant subspaces matrices A and B can be defined as follows

dist (Inv A, Inv B) = max{  $\sup_{\mathcal{M} \in \text{Inv } A} \theta(\mathcal{M}, \text{Inv } B), \sup_{\mathcal{N} \in \text{Inv } B} \theta(\mathcal{N}, \text{Inv } A)$ }.

This distance is a metric as well.

We are going to use the following property of gaps between subspaces. It can be found in [5].

**Proposition 4.1.** For subspaces  $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$ , we have

$$\theta(\mathcal{M}, \mathcal{N}) = \theta(\mathcal{N}^{\perp}, \mathcal{M}^{\perp}).$$
(4.8)

The symmetry with respect to subspaces of the gap is actually a disadvantage.

**Proposition 4.2.** Let  $\mathcal{M}, \mathcal{N}$  be subspaces of  $\mathbb{C}^n$ .

- (i) If dim( $\mathcal{M}$ ) = dim( $\mathcal{N}$ ) then for any  $x \in \mathcal{M}$  there exists a  $y \in \mathcal{N}$  such that  $||x y|| \le \theta(\mathcal{M}, \mathcal{N})$ .
- (*ii*) If dim( $\mathcal{M}$ )  $\neq$  dim( $\mathcal{N}$ ) then  $\theta(\mathcal{M}, \mathcal{N}) = 1$ .

The above result shows us that the gap is often not useful to consider when  $\dim(\mathcal{M}) \neq \dim(\mathcal{N})$ . In our theorem, we wish to find bounds on the kernels of the matrices, however, the dimension of the kernels are, in general, not equal.

The gap provides many useful results in providing a variety of bounds but the usefulness is limited to when the dimensions are equal. The concept of a semi-gap can be helpful when the dimensions are not equal. This advantage is highly useful when considering matrix perturbations.

Let  $\mathcal{M}, \mathcal{N}$  be subspaces of  $\mathbb{C}^n$ . The quantity

$$\theta_0(\mathcal{M}, \mathcal{N}) = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \inf_{y \in \mathcal{N}} \|x - y\|$$

is called the *semigap* (or one-sided gap) from  $\mathcal{M}$  to  $\mathcal{N}$ .

We notice some immediate properties of the semi-gap.

**Lemma 4.3.** Let  $\mathcal{M}, \mathcal{N} \subset \mathbb{C}^n$  be two subspaces. Then the following statements hold.

(i)  $\theta(\mathcal{M}, \mathcal{N}) = \max\{\theta_0(\mathcal{M}, \mathcal{N}), \theta_0(\mathcal{N}, \mathcal{M})\}.$ 

(*ii*) 
$$\theta_0(\mathcal{M}, \mathcal{N}) = \sup_{\substack{x \in \mathcal{M} \\ \|x\|=1}} \|x - P_{\mathcal{N}}x\|.$$

(iii) If 
$$\mathcal{N}_1 \subset \mathcal{N}_2$$
, then  $\theta_0(\mathcal{M}, \mathcal{N}_2) \leq \theta_0(\mathcal{M}, \mathcal{N}_1)$ ,  $\theta_0(\mathcal{N}_1, \mathcal{M}) \leq \theta_0(\mathcal{N}_2, \mathcal{M})$ .

- (iv)  $\theta_0(\mathcal{M}, \mathcal{N}) \leq 1.$
- (v) If dim  $\mathcal{M} > \dim \mathcal{N}$ , then  $\theta_0(\mathcal{M}, \mathcal{N}) = 1$ .
- (vi)  $\theta_0(\mathcal{M}, \mathcal{N}) < 1$  if and only if  $\mathcal{M} \cap \mathcal{N}^{\perp} = \emptyset$ .

These facts are well-known and can be found e.g. in [5, 6].

To this end we will need some new results on gap and semigap. These results will be derived next.

**Proposition 4.4.** Let  $A_0$  be fixed. Then, there exist  $\epsilon, K > 0$  such that for all A with  $||A - A_0|| < \epsilon$ , we have

$$\theta_0(\ker(A), \ker(A_0)) \le K ||A - A_0||.$$
 (4.9)

Since  $\ker(A) \oplus \operatorname{Im}(A^{\top}) = \mathbb{C}^n$ , often times it is easier to prove a result for the image rather than for the kernel. Because of this, the above proposition will follow from the next results.

**Lemma 4.5.** Let  $A_0$  be fixed. Then, there exist  $\epsilon, K > 0$  such that for all A with  $||A - A_0|| < \epsilon$ , we have

$$\theta_0(\operatorname{Im}(A_0), \operatorname{Im}(A)) \le K \|A - A_0\|.$$

*Proof.* Let  $A_0$  be an  $n \times n$ -matrix with dim $(\text{Im}(A_0)) = k$ . Consider an orthonormal basis  $g_1, g_2, \ldots, g_k$  of  $\text{Im}(A_0)$ . That is there are  $f_1, f_2, \ldots, f_k$  such that  $g_i = A_0 f_i$  for  $i = 1, \ldots, k$ . Define  $h_i = A f_i$ , so that  $h_i \in \text{Im}(A)$  for all i. Then, we have that

$$||h_i - g_i|| = ||Af_i - A_0 f_i|| = ||(A - A_0)f_i|| \le ||f_i|| ||A - A_0||.$$

Now, let  $x \in \text{Im}(A_0)$  and ||x|| = 1. This means that  $x = \alpha_1 g_1 + \ldots \alpha_k g_k$  for some  $\alpha_i$ 's. Define  $y = \alpha_1 h_1 + \ldots + \alpha_k h_k$ . Clearly,  $y \in \text{Im}(A)$  and

$$\inf_{z \in \text{Im}A} \|x - z\| \le \|x - y\| \le \sum_{i=1}^{k} \alpha_i \|h_i - g_i\| \\
\le \max_j (|\alpha_j|) \sum_{i=1}^{k} \|f_i\| \|A - A_0\| \\
\le \max_j (|\alpha_j|) \max_i \|f_i\| \|A - A_0\|$$

Therefore, by taking the supremum over x we arrive at  $\theta_0(\operatorname{Im}(A_0), \operatorname{Im}(A)) \leq K ||A - A_0||$ , where  $K = k \max_j (|\alpha_j|) \max_i ||f_i||$ .

Next, in order to show that Lemma 4.5 implies Proposition 4.4, we need the following result.

**Lemma 4.6.** Let  $\mathcal{M}, \mathcal{N}$  be subspaces of  $\mathbb{C}^n$ . Then we have

$$\theta_0(\mathcal{M}, \mathcal{N}) = \theta_0(\mathcal{N}^\perp, \mathcal{M}^\perp). \tag{4.10}$$

Proof. First, notice that if  $\dim \mathcal{M} > \dim \mathcal{N}$  or  $\theta_0(M, N) = 1$ , the result follows immediately. Now, we consider the case where  $\dim \mathcal{M} \leq \dim \mathcal{N}$  and  $\theta_0(M, N) < 1$ . Define P to be the subspace of all the projection of vectors in  $\mathcal{M}$  to  $\mathcal{N}$ , i.e.  $P = \operatorname{proj}_{\mathcal{N}} \mathcal{M}$ . Since  $\theta_0(M, N) < 1$ , we have that  $\dim P =$  $\dim \mathcal{M}$ . Additionally, we have that  $\theta(M, P) = \theta_0(M, N)$ . Recall that  $P \subset \mathcal{N}$ implies that  $\mathcal{N}^{\perp} \subset P^{\perp}$ . By Proposition 4.1 we get  $\theta(\mathcal{M}, P) = \theta(P^{\perp}, \mathcal{M}^{\perp})$ . Then, using Lemma 4.3, we have  $\theta(P^{\perp}, \mathcal{M}^{\perp}) \geq \theta_0(P^{\perp}, \mathcal{M}^{\perp}) = \theta(\mathcal{N}^{\perp}, \mathcal{M}^{\perp})$ . That is,

$$\theta(\mathcal{N}^{\perp}, \mathcal{M}^{\perp}) \leq \theta(\mathcal{M}, \mathcal{N}).$$

Now, repeating the argument for  $\mathcal{N}^{\perp}$  and  $\mathcal{M}^{\perp}$  gives us the result.

Recall that  $\ker(A) = \operatorname{Im}(A^*)^{\perp}$  and  $\operatorname{Im}(A) = \ker(A^*)^{\perp}$ . So, by combining Lemma 4.5 and Lemma 4.6, we see that Proposition 4.4 holds true.

#### 5. Backward Stability

In this section we finalize the proof of Theorem 1.7.

Proof of Theorem 1.7. Let  $A_0$  be given. Then, according to Proposition 4.4 there exists  $K, \epsilon > 0$  such that (4.9) holds. Let A be a matrix such that  $||A - A_0|| < \epsilon$  and A has Schur decomposition  $A = UTU^*$ . Let  $\lambda_i$ 's denote the eigenvalues of  $A_0$  and  $\mu_j$ 's denote the eigenvalues of A. In the Schur decomposition for A, we have a sequence of unitary Hessenberg matrices  $\{V_k\}_1^n$  such that  $U = V_1 \cdot \ldots \cdot V_n$ . Since the Schur decomposition construction is an iteration of stepsas was shown in Lemma 3.3, it is enough to show that we can obtain a bound on the first step. Let  $\mu_1$  be an eigenvalue of A and let  $V_1$  be the unitary matrix such that

$$V_1^* A V_1 = \begin{bmatrix} \mu_1 & \star & \cdots & \star \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

where the first column  $v_1$  of  $V_1$  is a unit eigenvector of A corresponding to eigenvalue  $\mu_1$ , and the matrix forms an orthogonal basis for  $\mathbb{C}^n$ .

Thus, by Proposition 4.4 we can find a unit vector  $u_1 \in \ker(A_0 - \lambda_1 I)$ and constants  $K_i$ 's such that  $||v_1 - u_1|| \leq K_0 ||A - A_0 + (\lambda_1 - \mu_1)I|| \leq K_0 ||A - A_0|| + K_1 ||\lambda_1 - \mu_1| \leq K_0 ||A - A_0|| + \tilde{K}_1 ||A - A_0||^{\alpha} \leq K_2 ||A - A_0||^{\alpha}$ where  $\alpha$  is either 1 or 1/n.

Hence, we can find a corresponding orthonormal basis forming  $U_1$  such that  $||v_i - u_i|| \leq K_i ||A - A_0||^{\alpha}$ . Therefore, we have that  $||V_1 - U_1|| \leq M_1 ||A - A_0||^{\alpha}$  and by construction, we have that

$$U_1^* A_0 U_1 = \begin{bmatrix} \lambda_1 & \star & \cdots & \star \\ 0 \\ \vdots \\ 0 \end{bmatrix} A_{0,2}$$

Repeating the process the same way we did in the proof of Lemma 3.4, we acquire a unitary matrix  $U_0 = U_1 \dots U_n$  such that  $||V_i - U_i|| \le M_i ||A_i - A_{0,i}||^{\alpha}$  for all  $i = 1, \dots, n$ .

Note that by construction

$$\begin{split} \|A_{i+1} - A_{0,i+1}\| &\leq \|V_i^* A_i V_i - U_i^* A_{0,i} U_i\| \leq \|V_i^* A_i V_i - V_i^* A_{0,i} V_i + V_i^* A_{0,i} V_i - \\ -V_i^* A_{0,i} U_i + V_i^* A_{0,i} U_i - U_i^* A_{0,i} U_i\| \leq \|V_i^* A_i V_i - V_i^* A_{0,i} V_i\| + \|V_i^* A_{0,i} V_i - \\ -V_i^* A_{0,i} U_{0,i}\| + \|V_i^* A_{0,i} U_{0,i} - U_i^* A_{0,i} U_i\| \leq \|A_i - A_{0,i}\| + \|A_{0,i}\| \|V_i - \\ -U_{0,i}\| + \|V_i^* - U_i^*\| \|A_{0,i}\| \leq \|A_i - A_{0,i}\| + 2M_i \|A_{0,i}\| \|A_i - A_{0,i}\|^{\alpha} \leq \widetilde{M}_i \|A - A_0\|^{\alpha}. \end{split}$$
  
It follows by induction that  $\|U - U_0\| \leq M \|A - A_0\|^{1/n}$  and  $M := \sum_i \widetilde{M}_i$  with  $\widetilde{M}_i$  depending only on  $A_0$ .

Next, let us consider  $T - T_0$ . Using the argument similar to above, we conclude that

$$||T - T_0|| = ||U^*AU - U_0^*A_0U_0|| \le \widehat{M}||A - A_0||,$$

where  $\widehat{M}$  depends only on  $A_0$ .

Hence, we arrive at the conclusion of one of our main results, i.e. formula (1.3) holds true with  $K = M + \widehat{M}$ .

# 6. Different GK numbers and Failure of the Forward Stability of the Schur Decomposition

As it turns out the GK numbers of the original and perturbed matrices give us the information whether the forward stability of the Schur decomposition is impossible. The intuition behind the non-stable case comes from the following fact.

**Proposition 6.1** (see [9]). We have the inequality

$$\inf \operatorname{dist}(\operatorname{Inv} A, \operatorname{Inv} A_0) > 0$$

where the infimum is taken over all possible pairs of  $A, A_0$ , having different GK numbers.

That is why we got the backward stability result and could not get the general result for forward stability.

**Lemma 6.2.** Let  $A_0 \in \mathbb{C}^{n \times n}$  and  $A_0 = U_0 T_0 U_0^*$  its fixed Schur decomposition, where the (1,1)-entry of  $T_0$  is an eigenvalue  $\lambda$  with dim Ker $(A_0 -$ 

 $\lambda I \geq 2$ . There exists M > 0 such that in any neighborhood of  $A_0$ ,, i.e.  $\{A : ||A - A_0|| < \varepsilon\}$  for any  $\varepsilon > 0$ ,

$$\sup_{A} \inf_{\substack{U, T \\ Schur Form \\ of A}} \|U - U_0\| + \|T - T_0\| > M > 0,$$
(6.11)

where the supremum is taken over all  $A \in \mathcal{U}$  not having an eigenvector close to the first column  $u_1$  of  $U_0$ , i.e. we have  $||u_1 - v|| > M$  for all v, eigenvectors of A, and the infimum is taken over all their Schur factorizations.

*Proof.* Let  $u_1$  be the first column of  $U_0$  and  $\lambda$  is the corresponding eigenvalue of  $A_0$ . By Proposition 1.2 we know that there are eigenvalues of A that lie relatively close to  $\lambda$  and the difference is equivalent to  $||A - A_0||^{1/n}$ . We will list those eigenvalues as  $\mu_1, \ldots, \mu_l$ .

Let  $\{v_1, \ldots, v_{l_j}\}$  be a basis of  $\operatorname{Ker}(A - \mu_j)$  (for  $j = 1, \ldots, l$ ). Moreover, let us denote by  $\{u_1, \ldots, u_{k_1(A_0)}\}$  the basis of  $\operatorname{Ker}(A_0 - \lambda)$ , where  $u_1$  as before and for each  $v_j$  there is  $u_s$  ( $s \neq 1$ ) such that

$$||v_j - u_s|| \le C ||A - A_0||^{1/r}$$

for some positive number C. We can assume this by using the backward stability result proven above. Then,

$$||u_1 - v_j|| \ge ||u_1 - u_s|| - ||u_s - v_j|| \ge \min_{i \ne 1} ||u_1 - u_i|| - C||A - A_0||^{1/n}.$$

We can always choose A close to  $A_0$  so  $M := \frac{\min_{i \neq 1} \|u_1 - u_i\|}{2} > C \|A - A_0\|^{1/n}$ . Denote by  $j^*$  the index that minimize the left-hand side of the above inequalities. Therefore, when supremum and infimum is taken under the conditions of this lemma

$$\sup_{\widetilde{A}} \inf_{U,T} \|U - U_0\| + \|T - T_0\| \ge \inf_{U,T:A = UTU^*} \|U - U_0\| \ge \|u_1 - v_{j^*}\| > M.$$

**Remark 6.3.** The assumption that we can find such A in every neighborhood of  $A_0$  not having an eigenvector close to the first column of  $U_0$  is based on the following fact. Consider the Jordan form of  $A_0 = P_0 J_0 P_0^{-1}$  such that the first two blocks correspond to the eigenvalue  $\lambda$  with the second one having  $u_1$  as the eigenvector. For any  $\varepsilon > 0$  take  $A = P_0 (J_0 + J_{\varepsilon}) P_0^{-1}$ ,

where  $J_{\varepsilon}$  has the only non-zero entry equal to  $\frac{\varepsilon}{\|P_0\|\|P_0^{-1}\|}$  on the (j, j + 1)spot  $(j \times j)$  is the size of the first Jordan block in  $J_0$  and the second block corresponds to the eigenvector  $u_1$ ). Hence,  $u_1$  is not an eigenvector of A and  $\|A - A_0\| = \|P_0 J_{\varepsilon} P_0^{-1}\| \leq \varepsilon$ . So the set of A that we are taking supremum over in Lemma 6.2 is not empty.

Now let us show that the statement of Theorem 1.6 is valid.

Proof of Theorem 1.6. First, note that having different GK numbers for A and  $A_0$  implies that  $A_0$  is derogatory, i.e. there is an eigenvalue  $\lambda$  of  $A_0$  such that dim Ker $(A_0 - \lambda I) \geq 2$ . Moreover,  $A_0$  and  $T_0$  are similar so they have the same GK numbers and dim Ker $(T_0 - \lambda I) \geq 2$  as well. Therefore, we can show the equivalent fact instead, i.e.

$$\sup_{\widetilde{B}} \inf_{U,T} \|U - I\| + \|T - T_0\| \ge \inf_{U,T:B = UTU^*} \|U - I\| > M.$$

If  $\lambda$  is the (1,1) entry of  $T_0$  the we use Lemma 6.2 to get the desired result. If it is not then we are required to perform an extra step.

$$T_{0} = U_{0}^{*}A_{0}U_{0} = \begin{bmatrix} \star & \star & \cdots & \star & \star & \star & \star & \cdots & \star \\ 0 & \star & \cdots & \star & \star & \star & \star & \cdots & \star \\ 0 & 0 & \cdots & \star & \star & \star & \cdots & \star \\ \vdots & \ddots & \ddots & \ddots & \vdots & \star & \cdots & \star \\ 0 & 0 & \cdots & 0 & \star & \star & \cdots & \star \\ 0 & 0 & \cdots & 0 & \star & \star & \cdots & \star \\ \hline & 0 & & & T_{1} \end{bmatrix}$$

where the first say j rows do not have  $\lambda$  on its main diagonal and the next row of  $T_0$  is the first time we meet  $\lambda$ . In addition, the (1,1)-entry of  $T_1$  is  $\lambda$ . According to Lemma 2.2, it means that we have the same GK numbers related to  $\lambda$  for  $T_1$  as for  $A_0$ . Recall that the truncation was using eigenvectors not corresponding to  $\lambda$ , so  $T_1$  will have the same number and length of Jordan chains for  $\lambda$  as  $A_0$  has. Thus, dim Ker $(T_1 - \lambda I) \geq 2$ . Now, we use Lemma 6.2 to get the result for  $\tilde{A}_0$ ,  $V_0$ , and  $\tilde{T}_0$  by constructing  $B_1$ (note that B is not upper triangular, since  $e_1$  is not its eigenvector) for  $T_1$  as described by the lemma. Then define

where the first j rows marked with stars coincide with  $T_0$ . By the construction, we can see that (1.2) holds true and hence finishing the proof of Theorem 1.6.

To summarize, we have showed that the Schur decomposition is backward stable and why it fails to be forward stable.

## References

- T. Bella, V. Olshevsky, U. Prasad, Lipschitz stability of canonical Jordan bases of H-selfadjoint matrices under structurepreserving perturbations, Linear Algebra and its Applications, 428, 8–9, 2008, 2130–2176.
- [2] H. Den Boer, G.Ph. Thijsse, Semi-stability of sums of partial multiplicities under additive perturbations, Integral Equations and Operator Theory, 3, 1980, 23–42.
- [3] S. Dogruer Akgul, A. Minenkova, V. Olshevsky, Lipschitz stability of γ-FOCS and RC canonical Jordan bases of real Hselfadjoint matrices under small perturbations.
- [4] I. Gohberg, M. A. Kaashoek, Unsolved problems in matrix and operator theory, Integral Equations Operator Theory, 1, 1978, 278–283.
- [5] I. Gohberg, P. Lancaster, L. Rodman, *Invariant Subspaces of Matrices with Applications*, Canadian Mathematical Society Series of Monographs and Advanced Texts. A Wiley-Interscience Publication. John Wiley& Sons, Inc., New York, 1986, xviii+692 pp.

- [6] T. Kato, Perturbation theory for linear operators, Die Grundlehren der mathematischen Wissenschaften, Band 132 Springer-Verlag New York, Inc., New York, 1966, xix+592 pp.
- [7] M. Konstantinov, D. Gu, V. Mehrmann, P. Petkov, Perturbation Theory for Matrix Equations Studies in Computational Mathematics, 9, North Holland Publishing Co., Amsterdam, 2003, xii+429 pp.
- [8] A. Markus, E. Parilis, The change of the Jordan structure of a matrix under small perturbations, Mat. Issled., 54 (1980), 98– 109 (in Russian), English translation: Linear Algebra Appl., 54, 1983, 139–152.
- [9] V. Olshevsky, A condition for the nearness of sets of invariant subspaces of near matrices in terms of their Jordan structures., (in Russian), Siberian Math. Journal, **30**, 4, 1989, 102–110, English translation: Siberian Math. Journal, Plenum publishing corp., **30**, 4, 1989, 580–586.
- [10] V. Matsaev, V. Olshevsky, Cyclic dimensions, kernel multiplicities, and Gohberg-Kaashoek numbers, Linear Algebra Appl., 239, 1996, 161–174.
- [11] A.M. Ostrowski, Solution of equations in Euclidean and Banach spaces. Third edition of Solution of equations and systems of equations. Pure and Applied Mathematics, Vol. 9. Academic Press, New York-London, 1973. xx+412 pp.
- [12] B. Simon, Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory, AMS Colloquium Series, American Mathematical Society, Providence, RI, 2005.
- [13] G.W. Stewart, J.G. Sun, Matrix Perturbation Theory, Academic Press Inc., Boston, MA, 1990, xvi+365 pp.
- [14] J. H. Wilkinson, The algebraic eigenvalue problem, Clarendon Press, Oxford, 1965, xviii+662 pp.