

Analysis of nonconforming IFE methods and a new scheme for elliptic interface problems

Haifeng Ji^{*} Feng Wang[†] Jinru Chen[‡] Zhilin Li[§]

Abstract

In this paper, an important discovery has been found for nonconforming immersed finite element (IFE) methods using integral-value degrees of freedom for solving elliptic interface problems. We show that those IFE methods can only achieve suboptimal convergence rates (i.e., $O(h^{1/2})$ in the H^1 norm and $O(h)$ in the L^2 norm) if the tangential derivative of the exact solution and the jump of the coefficient are not zero on the interface. A nontrivial counter example is also provided to support our theoretical analysis. To recover the optimal convergence rates, we develop a new nonconforming IFE method with additional terms locally on interface edges. The unisolvence of IFE basis functions is proved on arbitrary triangles. Furthermore, we derive the optimal approximation capabilities of both the Crouzeix-Raviart and the rotated- Q_1 IFE spaces for interface problems with variable coefficients via a unified approach different from multipoint Taylor expansions. Finally, optimal error estimates in both H^1 - and L^2 - norms are proved and confirmed with numerical experiments.

keyword: interface problem, nonconforming, immersed finite element, unfitted mesh

AMS subject classification. 65N15, 65N30, 35J60

1 Introduction

Let $\Omega \subset \mathbb{R}^2$ be a convex polygonal domain and Γ be a C^2 -smooth interface immersed in Ω . Without loss of generality, we assume that Γ divides Ω into two disjoint sub-domains Ω^+ and Ω^- such that $\Gamma = \partial\Omega^-$, see Figure 1 for an illustration. We consider the following second-order elliptic interface problem

$$-\nabla \cdot (\beta(x) \nabla u(x)) = f(x) \quad \text{in } \Omega \setminus \Gamma, \quad (1.1)$$

$$[u]_\Gamma(x) = 0 \quad \text{on } \Gamma, \quad (1.2)$$

$$[\beta \nabla u \cdot \mathbf{n}]_\Gamma(x) = 0 \quad \text{on } \Gamma, \quad (1.3)$$

$$u(x) = 0 \quad \text{on } \partial\Omega, \quad (1.4)$$

where $\mathbf{n}(x)$ is the unit normal vector of the interface Γ at point $x \in \Gamma$ pointing toward Ω^+ , and the notation $[v]_\Gamma$ is defined as

$$[v]_\Gamma(x) := \lim_{p \rightarrow x, p \in \Omega^+} v(p) - \lim_{p \rightarrow x, p \in \Omega^-} v(p) \quad \forall x \in \Gamma, \quad (1.5)$$

^{*}School of Science, Nanjing University of Posts and Telecommunications, Nanjing 210023, China (hfji@njupt.edu.cn)

[†]Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, Nanjing 210023, China (fwang@njnu.edu.cn)

[‡]Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, Nanjing 210023, China (jrchen@njnu.edu.cn)

[§]Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA (zhilin@math.ncsu.edu)

for a piecewise continuous function v . The coefficient $\beta(x)$ can be discontinuous across the interface Γ and is assumed to be piecewise smooth

$$\beta(x) = \beta^+(x) \text{ if } x \in \Omega^+ \text{ and } \beta(x) = \beta^-(x) \text{ if } x \in \Omega^-, \quad (1.6)$$

with $\beta^s(x) \in C^1(\overline{\Omega^s})$, $s = +, -$. We also assume that there exist two positive constants β_{min} and β_{max} such that $\beta_{min} \leq \beta^s(x) \leq \beta_{max}$ for all $x \in \overline{\Omega^s}$, $s = +, -$.




Figure 1: A diagram of the geometries of an interface problem.

It is well-known that traditional finite element methods using an interface-fitted mesh can solve the interface problem with optimal convergence rates, see for example [3, 5, 24]. For complicated interfaces or moving interfaces, unfitted meshes, which are not necessarily aligned with interfaces, have some advantages over interface-fitted meshes. However, traditional finite element methods using unfitted meshes only achieve suboptimal convergence rates ($O(h^{1/2})$ in the H^1 norm and $O(h)$ in the L^2 norm) no matter how high degree of the polynomial is used, see [1, 7].

Immersed finite element (IFE) methods [19, 21, 9, 20, 13, 12, 17, 11, 23] are designed to recover the optimal convergence rates of traditional finite element methods on unfitted meshes while keeping the degrees of freedom and the structure unchanged. The basic idea of IFEs is to modify traditional shape functions on interface elements to satisfy interface conditions approximately. However, these modifications are done on each interface elements independently, which may cause discontinuities of IFE basis functions across interface edges. Even for the P_1 conforming IFE method, these discontinuities are not negligible [13, 22] and the optimal convergence rates cannot be achieved if the discontinuities are not treated appropriately. To overcome the difficulty, Lin et al. [22] proposed a partially penalized IFE method where extra penalty terms at interface edges were added to penalize the discontinuity. For nonconforming IFE methods (i.e., a modification to the traditional Crouzeix-Raviart or the rotated- Q_1 finite element method), we can choose midpoint values or integral-values on edges as degrees of freedom. If we choose midpoint values on edges as degrees of freedom, the discontinuities of IFE basis functions are also not negligible and the optimal convergence rates can be obtained by adding partially penalized terms (see [25]). However, in the case that integral-values on edges are used as degrees of freedom, the IFE basis functions have less severe discontinuity across interface edges since they now maintain the integral-value continuous, and the authors [17, 23] claimed the optimal convergence rates can be obtained without requiring any penalty terms.

In this paper, we show that those nonconforming IFE methods using integral-value degrees of freedom cannot achieve the optimal convergence rates without penalty terms unless the tangential derivative of the exact solution u or the jump of the coefficient β is zero on the interface (see Theorem 4.5 and Remark 4.4). Furthermore, a nontrivial counter example (see *Example 6.1*) is constructed to support our theoretical analysis. The numerical examples in the literature [17, 23] show the optimal convergence rates for the nonconforming IFE methods using integral-value degrees of freedom because the constructed exact solutions are constants along the interface (i.e., the tangential

derivative is zero).

To achieve the optimal convergence rates, we develop a new nonconforming IFE method using either the Crouzeix-Raviart or the rotated- Q_1 element for solving interface problems with variable coefficients. The bilinear form of the new method includes some additional terms locally on interface edges to take into account of the contributions from the discontinuity of IFE basis functions. The method is symmetric and the coercivity is ensured without requiring a sufficient large parameter by using a special designed lifting operator defined locally on interface edges. To avoid integrating on curved regions, we approximate the interface by line segments connecting the intersection points of the mesh and the interface. We derive the optimal error estimates that are verified by numerical experiments.

There are other contributions of this paper. First, we prove that Crouzeix-Raviart IFE basis functions are unisolvent on arbitrary triangles. Our recent study in [16] shows that, for the IFEs using nodal values as degrees of freedom, the maximum angle condition, $\alpha_{max} \leq \pi/2$ on interface triangles, is necessary to ensure the unisolvence of the basis functions. The unisolvence of basis functions on arbitrary triangles is a significant advantage of the nonconforming IFEs using integral-value degrees of freedom over the IFEs using nodal values as degrees of freedom. Another contribution is a unified proof of the optimal interpolation error estimates for both the Crouzeix-Raviart and the rotated- Q_1 nonconforming IFE spaces for interface problems with piecewise smooth coefficients. Different from multipoint Taylor expansions [11], our proof is based on auxiliary functions constructed on interface elements and some useful inequalities developed by Li et al. in [18] and by Bramble and King in [2] for estimating errors in the region near the interface. The other contribution is a new theoretical result that the interpolation polynomial on one side of the interface can approximate the extensions of the exact solution optimally on the whole element T no matter how small $T \cap \Omega^+$ or $T \cap \Omega^-$ might be (see Theorem 3.7 in Section 3.3). The result is useful for proving the optimal interpolation error estimates on interface edges (see (5.20) in the proof of Lemma 5.5).

The rest of the paper is organized as follows. In Section 2, we introduce some notations, assumptions, and basic lemmas that are needed for the analysis. In Section 3, we describe two nonconforming IFE spaces based on the Crouzeix-Raviart and the rotated- Q_1 elements with integral-value degrees of freedom and prove the unisolvence of IFE basis functions on arbitrary triangles. Furthermore, optimal approximation capabilities of these nonconforming IFE spaces are derived via a new approach. In Section 4, we show that the existing nonconforming IFE method using integral-value degrees of freedom only has suboptimal convergence rates. In Section 5, we develop a new nonconforming IFE method and derive optimal error estimates in H^1 - and L^2 - norms. Numerical examples are presented in Section 6 to validate our theoretical findings. We conclude in the last section.

2 Notations and preliminaries

Throughout the paper we adopt the standard notation $W_p^k(\Lambda)$ for Sobolev spaces on a domain Λ with the norm $\|\cdot\|_{W_p^k(\Lambda)}$ and the seminorm $|\cdot|_{W_p^k(\Lambda)}$. Specially, $W_2^k(\Lambda)$ is denoted by $H^k(\Lambda)$ with the norm $\|\cdot\|_{H^k(\Lambda)}$ and the seminorm $|\cdot|_{H^k(\Lambda)}$. As usual $H_0^1(\Lambda) = \{v \in H^1(\Lambda) : v = 0 \text{ on } \partial\Lambda\}$. For a domain Λ , we define $\Lambda^s := \Lambda \cap \Omega^s$, $s = +, -$ and a space

$$\tilde{H}^2(\Lambda) := \{v \in H^1(\Lambda) : v|_{\Lambda^s} \in H^2(\Lambda^s), s = +, -, [v]_{\Gamma \cap \Lambda} = 0, [\beta \nabla v \cdot \mathbf{n}]_{\Gamma \cap \Lambda} = 0\} \quad (2.1)$$

when $\Lambda^s \neq \emptyset$, $s = +, -$. The space $\tilde{H}^2(\Lambda)$ is equipped with the norm $\|\cdot\|_{H^2(\Lambda^+ \cup \Lambda^-)}$ and the semi-norm $|\cdot|_{H^2(\Lambda^+ \cup \Lambda^-)}$ satisfying

$$\|\cdot\|_{H^2(\Lambda^+ \cup \Lambda^-)}^2 = \|\cdot\|_{H^2(\Lambda^+)}^2 + \|\cdot\|_{H^2(\Lambda^-)}^2, \quad |\cdot|_{H^2(\Lambda^+ \cup \Lambda^-)}^2 = |\cdot|_{H^2(\Lambda^+)}^2 + |\cdot|_{H^2(\Lambda^-)}^2.$$

By integrating by parts, we can immediately derive the following weak formulation of the interface problem (1.1)-(1.4): find $u \in H_0^1(\Omega)$ such that

$$a(u, v) := \int_{\Omega} \beta(x) \nabla u \cdot \nabla v dx = \int_{\Omega} f v dx \quad \forall v \in H_0^1(\Omega). \quad (2.2)$$

We have the following regularity theorem for the weak solution (see [15] for problems with piecewise smooth coefficients and [14, 6] for problems with piecewise constant coefficients)

Theorem 2.1. *If $f \in L^2(\Omega)$, then the interface problem (1.1)-(1.4) has a unique solution $u \in \tilde{H}^2(\Omega)$ satisfying the following a priori estimate*

$$\|u\|_{H^2(\Omega^+ \cup \Omega^-)} \leq C \|f\|_{L^2(\Omega)}, \quad (2.3)$$

where C is a positive constant depending only on Ω , Γ and β .

Let $\{\mathcal{T}_h\}_{h>0}$ be a family of triangular or rectangular subdivisions of Ω such that no vertex of any element lies in the interior of an edge of another element. The diameter of $T \in \mathcal{T}_h$ is denoted by h_T . We define $h = \max_{T \in \mathcal{T}_h} h_T$ and assume that \mathcal{T}_h is shape regular, i.e., for every T , there exists a positive constant ϱ such that $h_T \leq \varrho r_T$ where r_T is the diameter of the largest circle inscribed in T . Denote \mathcal{E}_h as the set of edges of the subdivision, and let \mathcal{E}_h° and \mathcal{E}_h^b be the sets of interior edges and boundary edges. We adopt the convention that elements $T \in \mathcal{T}_h$ and edges $e \in \mathcal{E}_h$ are open sets. Then, the sets of interface elements and interface edges are defined as

$$\mathcal{T}_h^\Gamma := \{T \in \mathcal{T}_h : T \cap \Gamma \neq \emptyset\} \quad \text{and} \quad \mathcal{E}_h^\Gamma := \{e \in \mathcal{E}_h : e \cap \Gamma \neq \emptyset\},$$

and the sets of non-interface elements and non-interface edges are $\mathcal{T}_h^{non} = \mathcal{T}_h \setminus \mathcal{T}_h^\Gamma$ and $\mathcal{E}_h^{non} = \mathcal{E}_h \setminus \mathcal{E}_h^\Gamma$. We can always refine the mesh near the interface to satisfy the following assumption.

Assumption 2.2. *The interface Γ does not intersect the boundary of any interface element at more than two points. The interface Γ does not intersect the closure \bar{e} for any $e \in \mathcal{E}_h$ at more than one point.*

The interface Γ is approximated by Γ_h that is composed of all the line segments connecting the intersection points of the boundaries of interface elements and the interface. In addition, we assume that the approximated interface Γ_h divides Ω into two disjoint sub-domains Ω_h^+ and Ω_h^- such that $\Gamma_h = \partial\Omega_h^-$.

On each interface element $T \in \mathcal{T}_h^\Gamma$, we denote the intersection points of Γ and ∂T by D and E . The straight line DE divides T into $T_h^+ = T \cap \Omega_h^+$ and $T_h^- = T \cap \Omega_h^-$, see Figure 2 for an illustration.

Let $\mathbf{n}_h(x)$ be the unit normal vector of Γ_h pointing toward Ω_h^+ . The tangent vector of Γ_h can be defined as $\mathbf{t}_h(x) = R_{\pi/2} \mathbf{n}_h(x)$, where R_α is a rotation matrix

$$R_\alpha = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}.$$

Denote $\text{dist}(x, \Gamma)$ as the distance between a point x and the interface Γ , and $U(\Gamma, \delta) = \{x \in \mathbb{R}^2 : \text{dist}(x, \Gamma) < \delta\}$ as the neighborhood of Γ of thickness δ . Define the meshsize of \mathcal{T}_h^Γ by

$$h_\Gamma := \max_{T \in \mathcal{T}_h^\Gamma} h_T. \quad (2.4)$$

It is obvious that $h_\Gamma \leq h$ and $\bigcup_{T \in \mathcal{T}_h^\Gamma} T \subset U(\Gamma, h_\Gamma)$.

We also define a signed distance function near the interface as

$$\rho(x) = \begin{cases} \text{dist}(x, \Gamma) & \text{if } x \in \Omega^+ \cap U(\Gamma, \delta_0), \\ -\text{dist}(x, \Gamma) & \text{if } x \in \Omega^- \cap U(\Gamma, \delta_0). \end{cases}$$

Assumption 2.3. *There exists a constant $\delta_0 > 0$ such that the signed distance function $\rho(x)$ is well-defined in $U(\Gamma, \delta_0)$ and $\rho(x) \in C^2(U(\Gamma, \delta_0))$. We also assume that $h_\Gamma < \delta_0$ so that $T \subset U(\Gamma, \delta_0)$ for all interface elements $T \in \mathcal{T}_h^\Gamma$.*

The assumption is reasonable since the interface Γ is in C^2 . We extend the coefficients $\beta^s(x)$, $s = +, -$ smoothly to slight larger domains $\Omega_e^s := \Omega^s \cup U(\Gamma, \delta_0)$, $s = +, -$ such that

$$\beta^s(x) \in C^1(\overline{\Omega_e^s}) \quad \text{and} \quad \beta_{\min} \leq \beta^s(x) \leq \beta_{\max}, \quad s = +, -. \quad (2.5)$$

Thus, there exists a constant C_β such that

$$\|\nabla \beta^s\|_{L^\infty(\Omega_e^s)} \leq C_\beta, \quad s = +, -. \quad (2.6)$$

By using the signed distance function ρ , we can evaluate the unit normal and tangent vectors of the interface as

$$\mathbf{n}(x) = \nabla \rho, \quad \mathbf{t}(x) = \left(-\frac{\partial \rho}{\partial x_2}, \frac{\partial \rho}{\partial x_1} \right)^T, \quad (2.7)$$

which are well-defined in the region $U(\Gamma, \delta_0)$. We note that the functions $\mathbf{n}_h(x)$ and $\mathbf{t}_h(x)$ are also viewed as piecewise constant vectors defined on interface elements. On each interface element $T \in \mathcal{T}_h^\Gamma$, since Γ is in C^2 , by Rolle's Theorem, there exists at least one point $x^* \in \Gamma \cap T$, see Figure 2, such that

$$\mathbf{n}(x^*) = \mathbf{n}_h(x^*) \quad \text{and} \quad \mathbf{t}(x^*) = \mathbf{t}_h(x^*). \quad (2.8)$$

Since $\rho(x) \in C^2(U(\Gamma, \delta_0))$, we have

$$\mathbf{n}(x) \in (C^1(\overline{T}))^2 \quad \text{and} \quad \mathbf{t}(x) \in (C^1(\overline{T}))^2 \quad \forall T \in \mathcal{T}_h^\Gamma. \quad (2.9)$$

Using Taylor's expansion at x^* , we further have

$$\|\mathbf{n} - \mathbf{n}_h\|_{L^\infty(T)} \leq Ch_T \quad \text{and} \quad \|\mathbf{t} - \mathbf{t}_h\|_{L^\infty(T)} \leq Ch_T \quad \forall T \in \mathcal{T}_h^\Gamma. \quad (2.10)$$

The following lemma presents a δ -strip argument that will be used for the error estimate in the region near the interface (see Lemma 2.1 in [18]).

Lemma 2.4. *Let δ be sufficiently small. Then it holds for any $v \in H^1(\Omega)$ that*

$$\|v\|_{L^2(U(\Gamma, \delta))} \leq C\sqrt{\delta} \|v\|_{H^1(\Omega)}.$$

Furthermore, if $v|_\Gamma = 0$, then there holds

$$\|v\|_{L^2(U(\Gamma, \delta))} \leq C\delta \|\nabla v\|_{L^2(U(\Gamma, \delta))}.$$

Recalling $T^s = T \cap \Omega^s$, $T_h^s = T \cap \Omega_h^s$, $s = +, -$ for all $T \in \mathcal{T}_h^\Gamma$, we define

$$T^\Delta := (T^- \cap T_h^+) \cup (T^+ \cap T_h^-). \quad (2.11)$$

Since Γ is in C^2 , we have $|T^\Delta| \leq Ch_T^3$. We shall need the following estimate on the region T^Δ (see Lemma 2 in [2]).

Lemma 2.5. *Assume that $v \in H^1(T)$ and $T \in \mathcal{T}_h^\Gamma$. Then there is a constant C , independent of h and the interface location relative to the mesh, such that*

$$\|v\|_{L^2(T^\Delta)}^2 \leq C(h_T^2 \|v\|_{L^2(\Gamma \cap T)}^2 + h_T^4 \|\nabla v\|_{L^2(T^\Delta)}^2).$$

3 Nonconforming IFE spaces and their properties

In this section, we describe nonconforming IFE spaces based on the Crouzeix-Raviart element or the rotated- Q_1 element and present their properties. To begin with, we define IFE shape function spaces. On a non-interface element $T \in \mathcal{T}_h^{non}$, we use the traditional shape function space

$$V_h(T) = \begin{cases} \text{Span}\{1, x_1, x_2\}, & \text{for the Crouzeix-Raviart element } (T \text{ is a triangle}), \\ \text{Span}\{1, x_1, x_2, x_1^2 - (\kappa_T x_2)^2\}, & \text{for the rotated-}Q_1 \text{ element } (T \text{ is a rectangle}), \end{cases}$$

where $\kappa_T = |e_1|/|e_2|$, e_1 and e_2 are edges of the rectangle and parallel to the x_1 -axis and the x_2 -axis, respectively. On an interface element $T \in \mathcal{T}_h^\Gamma$, the IFE shape function space $S_h(T)$ is defined as the set of the following functions

$$\phi(x) = \begin{cases} \phi^+(x) \in V_h(T) & \text{if } x = (x_1, x_2)^T \in T_h^+, \\ \phi^-(x) \in V_h(T) & \text{if } x = (x_1, x_2)^T \in T_h^-, \end{cases} \quad (3.1)$$

satisfying

$$[\phi]_{\Gamma_h \cap T}(x) = \phi^+(x) - \phi^-(x) = 0 \quad \forall x \in \Gamma_h \cap T, \quad (3.2)$$

$$\beta_c^+(\nabla \phi^+ \cdot \mathbf{n}_h)(x_p) - \beta_c^-(\nabla \phi^- \cdot \mathbf{n}_h)(x_p) = 0, \quad (3.3)$$

where x_p is an arbitrary point on $\Gamma_h \cap T$ and the constants β_c^+ and β_c^- are chosen such that

$$\|\beta^s(x) - \beta_c^s\|_{L^\infty(T)} \leq Ch_T, \quad s = +, -. \quad (3.4)$$

Actually, we can choose $\beta_c^s = \beta^s(x_c^s)$ with arbitrary $x_c^s \in T$, $s = +, -$, to satisfy the condition (3.4) since we know that $\beta^s(x) \in C^1(\overline{T})$, $s = +, -$ from (2.5).

Remark 3.1. For the Crouzeix-Raviart element, the condition (3.2) is equivalent to

$$\phi^+(D) = \phi^-(D), \quad \phi^+(E) = \phi^-(E),$$

since $\phi^s(x)$, $s = +, -$, are linear functions. For the rotated- Q_1 element, we can write $\phi^s(x)$ as

$$\phi^s(x) = a^s + b^s x_1 + c^s x_2 + d^s(x_1^2 - (\kappa_T x_2)^2), \quad x = (x_1, x_2)^T, \quad s = +, -,$$

where a^s, b^s, c^s, d^s , $s = +, -$, are constants. If we define a functional $d : V_h(T) \rightarrow \mathbb{R}$ as

$$d(\phi^s) = \frac{1}{2} \frac{\partial^2 \phi^s}{\partial x_1^2} = \frac{1}{\sqrt{(4\kappa_T + 4)|T|}} |\phi^s|_{H^2(T)}, \quad (3.5)$$

then $d^s = d(\phi^s)$. Similar to Lemma 2.1 in [12], the condition (3.2) is equivalent to

$$\phi^+(D) = \phi^-(D), \quad \phi^+(E) = \phi^-(E), \quad d(\phi^+) = d(\phi^-).$$

Remark 3.2. For the Crouzeix-Raviart element, $\beta_c^s \nabla \phi^s \cdot \mathbf{n}_h$, $s = +, -$, are constants on the interface element. Thus, the condition (3.3) is equivalent to

$$\beta_c^+(\nabla \phi^+ \cdot \mathbf{n}_h)(x) = \beta_c^-(\nabla \phi^- \cdot \mathbf{n}_h)(x) \quad \forall x \in \Gamma_h \cap T. \quad (3.6)$$

However, for the rotated- Q_1 element, the relation (3.6) is no longer valid. In [23], the authors weakly enforce the continuity by using the following condition

$$\int_{\Gamma_h \cap T} \beta_c^+(\nabla \phi^+ \cdot \mathbf{n}_h) - \beta_c^-(\nabla \phi^- \cdot \mathbf{n}_h) ds = 0$$

which is a particular case of (3.3) since there exists a point $x_p \in \Gamma_h \cap T$ such that

$$\int_{\Gamma_h \cap T} (\beta_c^+ \nabla \phi^+ - \beta_c^- \nabla \phi^-) \cdot \mathbf{n}_h ds = |\Gamma_h \cap T| (\beta_c^+ \nabla \phi^+ - \beta_c^- \nabla \phi^-)(x_p) \cdot \mathbf{n}_h.$$

Let $\mathcal{I} = \{1, 2, 3\}$ for the Crouzeix-Raviart element and $\mathcal{I} = \{1, 2, 3, 4\}$ for the rotated- Q_1 element. The degrees of freedom are defined as the mean values over edges

$$N_i(\phi) := \frac{1}{|e_i|} \int_{e_i} \phi ds, \quad i \in \mathcal{I},$$

where $e_i, i \in \mathcal{I}$ are edges of the element T , and $|e_i|$ denotes the length of the edge e_i . On an interface element $T \in \mathcal{T}_h^\Gamma$, the immersed finite element is defined as $(T, S_h(T), \Sigma_T)$ with $\Sigma_T = \{N_i, i \in \mathcal{I}\}$.

The nonconforming IFE spaces. The nonconforming IFE space V_h^{IFE} is defined as the set of all functions satisfying

$$\begin{cases} \phi|_T \in S_h(T) & \forall T \in \mathcal{T}_h^\Gamma, \\ \phi|_T \in V_h(T) & \forall T \in \mathcal{T}_h^{non}, \\ \int_e [\phi]_e ds = 0 & \forall e \in \mathcal{E}_h^\circ. \end{cases}$$

We also need a space for homogeneous boundary conditions

$$V_{h,0}^{IFE} := \left\{ v \in V_h^{IFE} : \int_e v ds = 0 \quad \forall e \in \mathcal{E}_h^b \right\}.$$

3.1 The unsolvence of IFE basis functions

It was proved in [11] that the function $\phi \in S_h(T)$ is uniquely determined by $N_i(\phi)$, $i = 1, 2, 3, 4$ for the rotated- Q_1 element, and $i = 1, 2, 3$ for the Crouzeix-Raviart element when the interface element is a right triangle. Now we prove that the result is also valid for arbitrary triangles in the following lemma. Note that for the IFEs using nodal values as degrees of freedom, the maximum angle condition, $\alpha_{max} \leq \pi/2$ on interface triangles, is necessary to ensure the unsolvence (see [16]). This property of the unsolvence of basis functions is one of advantages of nonconforming IFEs compared with the IFEs using nodal values as degrees of freedom.




Figure 2: Typical interface elements. Left diagram: triangle element. Right diagram: rectangle element.

Lemma 3.3. *Let T be an arbitrary interface triangle. For the Crouzeix-Raviart element, the function $\phi \in S_h(T)$ is uniquely determined by $N_i(\phi)$, $i = 1, 2, 3$.*

Proof. We use the argument proposed in [11, 10]. Consider a triangle $\triangle A_1A_2A_3$ with edges $e_1 = \overline{A_2A_3}$, $e_2 = \overline{A_1A_3}$ and $e_3 = \overline{A_1A_2}$. The interface Γ cuts e_1 and e_2 at points D and E , see Figure 2 for an illustration. Without loss of generality, we assume $T_h^- = \triangle EDA_3$ since the case $T_h^+ = \triangle EDA_3$ can be treated by reversing β_c^+ and β_c^- . Let $\lambda_i(x)$, $i = 1, 2, 3$, be basis functions in $V_h(T)$ such that

$\frac{1}{|e_j|} \int_{e_j} \lambda_i(x) dx = \delta_{ij}$, $i, j = 1, 2, 3$, where δ_{ij} is the Kronecker function. By using the condition (3.2) and $|e_3|^{-1} \int_{e_3} \phi ds = N_3(\phi)$, we can write the IFE shape function $\phi(x)$ in (3.1) as

$$\phi(x) = \begin{cases} \phi^+(x) = c_1 \lambda_1(x) + c_2 \lambda_2(x) + N_3(\phi) \lambda_3(x) & \text{if } x = (x_1, x_2)^T \in T_h^+, \\ \phi^-(x) = \phi^+(x) + c_0 \mathbf{n}_h \cdot \vec{Dx} & \text{if } x = (x_1, x_2)^T \in T_h^-, \end{cases} \quad (3.7)$$

where c_0, c_1, c_2 are unknowns. Applying the condition (3.3), the unknown c_0 can be expressed as

$$c_0 = (\beta_c^+ / \beta_c^- - 1) \nabla \phi^+ \cdot \mathbf{n}_h = (\beta_c^+ / \beta_c^- - 1) (c_1 \nabla \lambda_1 + c_2 \nabla \lambda_2 + N_3(\phi) \nabla \lambda_3) \cdot \mathbf{n}_h. \quad (3.8)$$

Substituting (3.8) into (3.7) and using $N_i(\phi) = |e_i|^{-1} \int_{e_i} \phi ds$, $i = 1, 2$, we obtain the following linear system of equations for other coefficients (see [11, 10] for details),

$$(\mathbf{I} + (\beta_c^+ / \beta_c^- - 1) \delta \gamma^T) \mathbf{c} = \mathbf{b}, \quad (3.9)$$

where

$$\begin{aligned} \delta &= \left(|e_1|^{-1} \int_{A_3D} L(x) ds, |e_2|^{-1} \int_{A_3E} L(x) ds \right)^T, \quad L(x) = \mathbf{n}_h \cdot \vec{Dx}, \\ \mathbf{c} &= (c_1, c_2)^T, \quad \gamma = (\nabla \lambda_1 \cdot \mathbf{n}_h, \nabla \lambda_2 \cdot \mathbf{n}_h)^T, \\ \mathbf{b} &= \left(N_1(\phi) - \frac{(\beta_c^+ / \beta_c^- - 1) N_3(\phi) \nabla \lambda_3 \cdot \mathbf{n}_h}{|e_1|} \int_{A_3D} L(x) ds, \right. \\ &\quad \left. N_2(\phi) - \frac{(\beta_c^+ / \beta_c^- - 1) N_3(\phi) \nabla \lambda_3 \cdot \mathbf{n}_h}{|e_2|} \int_{A_3E} L(x) ds \right)^T. \end{aligned} \quad (3.10)$$

Set $k_1 = |A_3D| |e_1|^{-1}$ and $k_2 = |A_3E| |e_2|^{-1}$. Let M_i be the midpoint of the edge e_i , $i = 1, 2, 3$, and Q be the orthogonal projection of M_2 onto the line A_2A_3 . We can find out $\gamma(1)$ and $\delta(1)$ as below

$$\begin{aligned} \gamma(1) &= \nabla \lambda_1 \cdot \mathbf{n}_h = |M_2Q|^{-1} \overrightarrow{M_2Q} |M_2Q|^{-1} \cdot \mathbf{n}_h = |M_2Q|^{-1} R_{\pi/2} \left(\overrightarrow{M_2Q} |M_2Q|^{-1} \right) \cdot R_{\pi/2}(\mathbf{n}_h) \\ &= |M_2Q|^{-1} \overrightarrow{A_2A_3} |A_2A_3|^{-1} \cdot \mathbf{t}_h = \left(\frac{1}{2} |e_2| \sin \angle A_3 \right)^{-1} |e_1|^{-1} \overrightarrow{A_2A_3} \cdot \mathbf{t}_h, \end{aligned}$$

and

$$\delta(1) = |e_1|^{-1} \int_{A_3D} \mathbf{n}_h \cdot \vec{Dx} ds = |e_1|^{-1} |A_3D| \mathbf{n}_h \cdot \vec{DH} = -\frac{1}{2} k_1 |A_3A_{3,\perp}|,$$

where $\angle A_3 \in (0, \pi)$, H is the midpoint of the line segment $\overline{A_3D}$, and $A_{3,\perp}$ is the orthogonal projection of A_3 onto the line DE . Thus,

$$\begin{aligned} \gamma(1) \delta(1) &= \nabla \lambda_1 \cdot \mathbf{n}_h |e_1|^{-1} \int_{A_3D} \vec{Dx} ds = -k_1 \overrightarrow{A_2A_3} \cdot \mathbf{t}_h |A_3A_{3,\perp}| (|e_1| |e_2| \sin \angle A_3)^{-1} \\ &= -\overrightarrow{DA_3} \cdot \mathbf{t}_h |A_3A_{3,\perp}| (|e_1| |e_2| \sin \angle A_3)^{-1}. \end{aligned}$$

Analogously, we have

$$\gamma(2) \delta(2) = \nabla \lambda_2 \cdot \mathbf{n}_h |e_2|^{-1} \int_{A_3E} \vec{Dx} ds = -\overrightarrow{A_3E} \cdot \mathbf{t}_h |A_3A_{3,\perp}| (|e_1| |e_2| \sin \angle A_3)^{-1}.$$

Therefore,

$$\gamma^T \delta = \overrightarrow{ED} \cdot \mathbf{t}_h |A_3A_{3,\perp}| (|e_1| |e_2| \sin \angle A_3)^{-1} = |DE| |A_3A_{3,\perp}| (|e_1| |e_2| \sin \angle A_3)^{-1}.$$

As long as $\angle A_3ED \in (0, \pi)$, it holds $|A_3A_{3,\perp}| = k_2 |e_2| \sin \angle A_3ED$, which together with the relations $|DE| \sin^{-1} \angle A_3 = k_1 |e_1| \sin^{-1} \angle A_3ED$ yields

$$\gamma^T \delta = k_1 |e_1| (\sin \angle A_3ED)^{-1} k_2 |e_2| (\sin \angle A_3ED) (|e_1| |e_2|)^{-1} = k_1 k_2 \in [0, 1].$$

From the above inequality, it holds

$$1 + (\beta_c^+/\beta_c^- - 1)\boldsymbol{\gamma}^T\boldsymbol{\delta} \geq \min(1, \beta_c^+/\beta_c^-) \geq \beta_{\min}/\beta_{\max} > 0. \quad (3.11)$$

Hence, by the well-known Sherman-Morrison formula, the linear system (3.9) has a unique solution

$$\mathbf{c} = \mathbf{b} - \frac{(\beta_c^+/\beta_c^- - 1)(\boldsymbol{\gamma}^T\mathbf{b})\boldsymbol{\delta}}{1 + (\beta_c^+/\beta_c^- - 1)\boldsymbol{\gamma}^T\boldsymbol{\delta}}, \quad (3.12)$$

which implies the lemma. \square

3.2 Estimates of IFE basis functions

On an interface element $T \in \mathcal{T}_h^\Gamma$, define IFE basis functions as

$$\phi_i \in S_h(T), \quad N_j(\phi_i) = \delta_{ij} \quad \forall i, j \in \mathcal{I}. \quad (3.13)$$

Also define functions $\phi_i^s \in V_h(T)$, $s = +, -, i \in \mathcal{I}$ such that $\phi_i^s = \phi_i|_{T_h^s}$. Let λ_i be traditional basis functions, i.e.,

$$\lambda_i \in V_h(T), \quad N_j(\lambda_i) = \delta_{ij} \quad \forall i, j \in \mathcal{I}. \quad (3.14)$$

Note that these functions depend on elements. We omit this dependence in our notation for simplicity. It is well-known that the traditional basis functions satisfy

$$|\lambda_i|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad i \in \mathcal{I}, \quad m = 0, 1, 2, \quad (3.15)$$

where the constant C only depends on the shape regularity parameter ϱ . The following lemma shows that the IFE basis functions also have similar estimates, which is one of essential ingredients for the success of IFE methods.

Lemma 3.4. *There exists a constant C , depending only on β_c^+ , β_c^- and the shape regularity parameter ϱ , such that*

$$|\phi_i^+|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad |\phi_i^-|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad i \in \mathcal{I}, \quad m = 0, 1, 2, \quad (3.16)$$

$$|\phi_i|_{W_\infty^m(T_h^+ \cup T_h^-)} \leq Ch_T^{-m}, \quad i \in \mathcal{I}, \quad m = 0, 1, 2. \quad (3.17)$$

Proof. See Theorem 4.2 in [11] for the estimate (3.17) for the rotated- Q_1 element and the Crouzeix-Raviart element on right triangles. The estimate (3.16) can also be obtained easily from the proof of Theorem 5.6 in [10]. Next, we give a proof for general triangles without the constraint $\alpha_{\max} = \pi/2$. We just need to prove (3.16) since (3.17) is a direct consequence of (3.16). Using $|N_j(\phi_i)| \leq 1$, $\|\nabla \lambda_i\|_{L^\infty(T)} \leq Ch_T^{-1}$, $i, j \in \mathcal{I}$, we can estimate vectors in (3.10) as

$$\|\mathbf{b}\| \leq C, \quad \|\boldsymbol{\gamma}\| \leq Ch_T^{-1}, \quad \|\boldsymbol{\delta}\| \leq Ch_T,$$

which, together with (3.11) and (3.12) lead to $\|\mathbf{c}\| \leq C$, where the constant C is independent of h_T and the interface location relative to the mesh. Thus, from (3.7), it follows

$$|\phi_i^+|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad i \in \mathcal{I}, \quad m = 0, 1,$$

where we have used the fact $|\lambda_i|_{W_\infty^m(T)} \leq Ch_T^{-m}$. It follows from (3.8) that $|c_0| \leq Ch_T^{-1}$ which together with (3.7) yields

$$|\phi_i^-|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad i \in \mathcal{I}, \quad m = 0, 1. \quad \square$$

3.3 Optimal approximation capabilities of IFE spaces

On each element $T \in \mathcal{T}_h$, define a local interpolation operator $I_{h,T} : W(T) \rightarrow V_h(T)$ such that

$$N_i(I_{h,T}v) = N_i(v), \quad i \in \mathcal{I},$$

where $W(T) = \{v : N_i(v), i \in \mathcal{I} \text{ are well defined}\}$. Similarly, on each interface element $T \in \mathcal{T}_h^\Gamma$, define $I_{h,T}^{IFE} : W(T) \rightarrow S_h(T)$ such that

$$N_i(I_{h,T}^{IFE}v) = N_i(v), \quad i \in \mathcal{I}. \quad (3.18)$$

The global IFE interpolation operator is defined by $I_h^{IFE} : H^1(\Omega) \rightarrow V_h^{IFE}$ such that

$$(I_h^{IFE}v)|_T = \begin{cases} I_{h,T}^{IFE}v & \text{if } T \in \mathcal{T}_h^\Gamma, \\ I_{h,T}v & \text{if } T \in \mathcal{T}_h^{non}. \end{cases}$$

For simplicity, define $v^s := v|_{\Omega^s}$, $s = +, -$ for all $v \in L^2(\Omega)$. With a small ambiguity of notation, given a function $v_h \in S_h(T)$, we define $v_h^s \in V_h(T)$, $s = +, -$ such that

$$v_h^s = v_h|_{T_h^s}, \quad s = +, -. \quad (3.19)$$

To show that functions in V_h^{IFE} can approximate a function v in $\tilde{H}^2(\Omega)$ optimally, we need to interpolate extensions of v^s , $s = +, -$. It is well-known that (see [8]) for any $v \in \tilde{H}^2(\Omega)$ there exist extensions $v_E^s \in H^2(\Omega)$, $s = +, -$ such that

$$v_E^s|_{\Omega^s} = v^s \quad \text{and} \quad |v_E^s|_{H^i(\Omega)} \leq C|v^s|_{H^i(\Omega^s)}, \quad i = 0, 1, 2, \quad s = +, -. \quad (3.20)$$

On each interface element $T \in \mathcal{T}_h^\Gamma$, given two functions $v^+ \in L^2(T)$ and $v^- \in L^2(T)$, we define

$$\llbracket v^\pm \rrbracket(x) := v^+(x) - v^-(x) \quad \forall x \in T.$$

For example, for a function $v \in \tilde{H}^2(\Omega)$,

$$\begin{aligned} \llbracket v_E^\pm \rrbracket(x) &= v_E^+(x) - v_E^-(x) & \forall x \in T, \\ \llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket(x) &= \beta^+(x) \nabla v_E^+(x) \cdot \mathbf{n}(x) - \beta^-(x) \nabla v_E^-(x) \cdot \mathbf{n}(x) & \forall x \in T, \\ \llbracket I_h v_E^\pm \rrbracket(x) &= I_h v_E^+(x) - I_h v_E^-(x) & \forall x \in T, \\ \llbracket \beta^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket(x) &= \beta^+(x) \nabla(I_h v_E^+) \cdot \mathbf{n}_h(x) - \beta^-(x) \nabla(I_h v_E^-) \cdot \mathbf{n}_h(x) & \forall x \in T, \end{aligned} \quad (3.21)$$

and for a function $v_h \in V_h^{IFE}$,

$$\begin{aligned} \llbracket v_h^\pm \rrbracket(x) &= v_h^+(x) - v_h^-(x) & \forall x \in T, \\ \llbracket \beta_c^\pm \nabla v_h^\pm \cdot \mathbf{n}_h \rrbracket(x) &= \beta_c^+ \nabla v_h^+(x) \cdot \mathbf{n}_h(x) - \beta_c^- \nabla v_h^-(x) \cdot \mathbf{n}_h(x) & \forall x \in T. \end{aligned} \quad (3.22)$$

Note that the difference between $\llbracket \cdot \rrbracket(x)$ and $[\cdot]_\Gamma(x)$ is the range of x .

We introduce *auxiliary functions* on each interface element $T \in \mathcal{T}_h^\Gamma$. Recalling that D and E are intersection points of Γ and ∂T , define auxiliary functions $\Upsilon_1(x)$, $\Upsilon_2(x)$ and $\Psi(x)$ as

$$\Upsilon_i(x) := \begin{cases} \Upsilon_i^+(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^+, \\ \Upsilon_i^-(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^-, \end{cases} \quad i = 1, 2, \quad (3.23)$$

such that

$$\begin{aligned} N_j(\Upsilon_i) &= 0, \quad j = \mathcal{I}, \quad \llbracket \Upsilon_i^\pm \rrbracket(D) = 0, \quad i = 1, 2, \\ \llbracket \beta_c^\pm \nabla \Upsilon_1^\pm \cdot \mathbf{n}_h \rrbracket(x_p) &= 1, \quad \llbracket \nabla \Upsilon_1^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 0, \quad \llbracket d(\Upsilon_1^\pm) \rrbracket = 0, \\ \llbracket \beta_c^\pm \nabla \Upsilon_2^\pm \cdot \mathbf{n}_h \rrbracket(x_p) &= 0, \quad \llbracket \nabla \Upsilon_2^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 1, \quad \llbracket d(\Upsilon_2^\pm) \rrbracket = 0, \end{aligned} \quad (3.24)$$

and

$$\Psi(x) := \begin{cases} \Psi^+(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^+, \\ \Psi^-(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^-, \end{cases} \quad (3.25)$$

such that

$$\begin{aligned} N_j(\Psi) &= 0, \quad j = \mathcal{I}, \quad \llbracket \Psi^\pm \rrbracket(D) = 1, \\ \llbracket \beta_c^\pm \nabla \Psi^\pm \cdot \mathbf{n}_h \rrbracket(x_p) &= 0, \quad \llbracket \nabla \Psi^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 0, \quad \llbracket d(\Psi^\pm) \rrbracket = 0, \end{aligned} \quad (3.26)$$

where $d(\cdot)$ is defined in (3.5) and the point $x_p \in \Gamma_h \cap T$ is the same as that in (3.3). For the rotated- Q_1 element, another auxiliary function is needed

$$\Theta(x) := \begin{cases} \Theta^+(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^+, \\ \Theta^-(x) \in V_h(T) & \text{if } x = (x_1, x_2) \in T_h^-, \end{cases} \quad (3.27)$$

such that

$$\begin{aligned} N_j(\Theta) &= 0, \quad j = \mathcal{I}, \quad \llbracket \Theta^\pm \rrbracket(D) = 0, \\ \llbracket \beta_c^\pm \nabla \Theta^\pm \cdot \mathbf{n}_h \rrbracket(x_p) &= 0, \quad \llbracket \nabla \Theta^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 0, \quad \llbracket d(\Theta^\pm) \rrbracket = 1. \end{aligned} \quad (3.28)$$

In order to have a unified analysis, we also define $\Theta = 0$ for the Crouzeix-Raviart element. Note that these auxiliary functions depend on the element T . We omit the dependence for simplicity of notations.

Lemma 3.5. *On each interface element $T \in \mathcal{T}_h^\Gamma$, these functions $\Upsilon_1(x)$, $\Upsilon_2(x)$, $\Psi(x)$ and Θ defined in (3.23)-(3.28) exist and satisfy*

$$\begin{aligned} |\Upsilon_i^s|_{W_2^m(T)}^2 &\leq Ch_T^{4-2m}, \quad m = 0, 1, 2, \quad i = 1, 2, \quad s = +, -, \\ |\Psi^s|_{W_2^m(T)}^2 &\leq Ch_T^{2-2m}, \quad |\Theta^s|_{W_2^m(T)}^2 \leq Ch_T^{6-2m}, \quad m = 0, 1, 2, \quad s = +, -, \end{aligned} \quad (3.29)$$

where the constant C depends only on β_c^+ , β_c^- and the shape regularity parameter ϱ .

Proof. We construct $\Upsilon_i(x)$, $i = 1, 2$ as follows,

$$\Upsilon_i = z_i - I_{h,T}^{IFE} z_i, \quad z_i = \begin{cases} z_i^+ & \text{in } T_h^+, \\ 0 & \text{in } T_h^-, \end{cases}, \quad i = 1, 2, \quad (3.30)$$

where z_1^+ and z_2^+ are linear and satisfy

$$\begin{aligned} z_1^+(D) &= 0, \quad \beta_c^+ \nabla z_1^+ \cdot \mathbf{n}_h = 1, \quad \nabla z_1^+ \cdot \mathbf{t}_h = 0, \\ z_2^+(D) &= 0, \quad \beta_c^+ \nabla z_2^+ \cdot \mathbf{n}_h = 0, \quad \nabla z_2^+ \cdot \mathbf{t}_h = 1. \end{aligned} \quad (3.31)$$

It is easy to verify that z_1^+ and z_2^+ exist, and the constructed functions Υ_i , $i = 1, 2$ satisfy (3.23) and (3.24). From (3.31), we have

$$|z_i^+|_{L^\infty(T)} \leq Ch_T, \quad |\nabla z_i^+| \leq C, \quad |z_i^+|_{W_\infty^2(T)} = 0, \quad |z_i|_{L^\infty(T)} \leq Ch_T, \quad i = 1, 2. \quad (3.32)$$

Since $I_{h,T}^{IFE} z_i = \sum_j N_j(z_i) \phi_j$, it follows from (3.16) that, for $m = 0, 1, 2$, $i = 1, 2$, $s = +, -$,

$$\left| (I_{h,T}^{IFE} z_i)^s \right|_{W_\infty^m(T)} \leq \sum_{j \in \mathcal{I}} |N_j(z_i)| |\phi_j^s|_{W_\infty^m(T)} \leq Ch_T^{-m} \sum_{j \in \mathcal{I}} |N_j(z_i)|.$$

Noticing

$$|N_j(z_i)| \leq |e_j|^{-1} \int_{e_j} |z_i| ds \leq |z_i|_{L^\infty(T)} \leq Ch_T,$$

we have

$$\left| (I_{h,T}^{IFE} z_i)^s \right|_{W_\infty^m(T)} \leq Ch_T^{1-m}, \quad m = 0, 1, 2, \quad i = 1, 2, \quad s = +, -,$$

which together with (3.30) and (3.32) implies

$$|\Upsilon_i^s|_{W_\infty^m(T)} \leq Ch_T^{1-m}, \quad m = 0, 1, 2, \quad i = 1, 2, \quad s = +, -.$$

Finally, the first estimate in (3.29) is obtained by

$$|\Upsilon_i^s|_{W_2^m(T)}^2 \leq |\Upsilon_i^s|_{W_\infty^m(T)}^2 |T| \leq Ch_T^{4-2m}, \quad m = 0, 1, 2, \quad i = 1, 2, \quad s = +, -.$$

Other estimates in (3.29) can be proved similarly. We construct $\Psi(x)$ as

$$\Psi = z - I_{h,T}^{IFE} z, \quad z = \begin{cases} z^+ = 1 & \text{in } T_h^+, \\ 0 & \text{in } T_h^-. \end{cases}$$

It is easy to verify that the constructed function Ψ satisfies (3.25) and (3.26). Since

$$\begin{aligned} |z^+|_{L^\infty(T)} &= 1, \quad |z^+|_{W_\infty^m(T)} = 0, \quad |z|_{L^\infty(T)} = 1, \quad m = 1, 2, \\ |N_j(z)| &\leq |e_j|^{-1} \int_{e_j} |z| ds \leq |z|_{L^\infty(T)} \leq 1, \quad j \in \mathcal{I}, \end{aligned}$$

we get

$$\left| (I_{h,T}^{IFE} z)^s \right|_{W_\infty^m(T)} \leq \sum_{j \in \mathcal{I}} |N_j(z)| |\phi_j^s|_{W_\infty^m(T)} \leq Ch_T^{-m} \sum_{j \in \mathcal{I}} |N_j(z)| \leq Ch_T^{-m}$$

and

$$|\Psi^s|_{W_\infty^m(T)} \leq Ch_T^{-m}, \quad s = +, -, \quad m = 0, 1, 2,$$

which implies

$$|\Psi^s|_{W_2^m(T)}^2 \leq |\Psi^s|_{W_\infty^m(T)}^2 |T| \leq Ch_T^{2-2m}, \quad s = +, -, \quad m = 0, 1, 2.$$

For the rotated- Q_1 element, we construct $\Theta(x)$ as

$$\Theta = z - I_{h,T}^{IFE} z, \quad z = \begin{cases} z^+ = (x_1 - m_1)^2 - \kappa_T^2 (x_2 - m_2)^2 & \text{in } T_h^+, \\ 0 & \text{in } T_h^-, \end{cases}$$

where (m_1, m_2) is the center of the rectangle T . Using (3.5), we easily verify that the constructed function Θ satisfies (3.27) and (3.28). It follows that

$$\begin{aligned} |z^+|_{L^\infty(T)} &\leq Ch_T^2, \quad |z^+|_{L_\infty^1(T)} \leq Ch_T, \quad |z^+|_{W_\infty^2(T)} \leq C, \quad |z|_{L^\infty(T)} \leq Ch_T^2, \\ |N_j(z)| &\leq |e_j|^{-1} \int_{e_j} |z| ds \leq |z|_{L^\infty(T)} \leq Ch_T^2, \quad j \in \mathcal{I}, \end{aligned} \tag{3.33}$$

which implies

$$\left| (I_{h,T}^{IFE} z)^s \right|_{W_\infty^m(T)} \leq \sum_{j \in \mathcal{I}} |N_j(z)| |\phi_j^s|_{W_\infty^m(T)} \leq Ch_T^{-m} \sum_{j \in \mathcal{I}} |N_j(z)| \leq Ch_T^{2-m}$$

and

$$\begin{aligned} |\Theta^s|_{W_2^m(T)}^2 &\leq |\Theta^s|_{W_\infty^m(T)}^2 |T| \leq |\Theta^s|_{W_\infty^m(T)}^2 |T| \\ &\leq C \left(|(z)^s|_{W_\infty^m(T)}^2 + \left| (I_{h,T}^{IFE} z)^s \right|_{W_\infty^m(T)}^2 \right) |T| \\ &\leq Ch_T^{6-2m}, \quad s = +, -, \quad m = 0, 1, 2. \end{aligned} \tag{3.34}$$

For the Crouzeix-Raviart element, the above inequality (3.34) is also valid since we have defined $\Theta = 0$ if T is a triangle. \square

Lemma 3.6. *On each interface element $T \in \mathcal{T}_h^\Gamma$, for any $v \in \tilde{H}^2(\Omega)$, the following identity holds*

$$(I_h v_E^s)(x) - (I_h^{IFE} v)^s(x) = a\Psi^s(x) + \sum_{i=1,2} b_i \Upsilon_i^s(x) + \sum_{i \in \mathcal{I}} g_i \phi_i^s(x) + t\Theta^s(x) \quad \forall x \in T, \quad s = +, -, \quad (3.35)$$

with

$$\begin{aligned} a &= \llbracket I_h v_E^\pm \rrbracket(D), \quad b_1 = \llbracket \beta_c^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket(x_p), \quad b_2 = \llbracket \nabla(I_h v_E^\pm) \cdot \mathbf{t}_h \rrbracket(x_p), \\ g_i &= |e_i|^{-1} \left(\int_{e_i^+} (I_h v_E^+ - v_E^+) ds + \int_{e_i^-} (I_h v_E^- - v_E^-) ds \right), \quad i \in \mathcal{I}, \\ t &= \llbracket d(I_h v_E^\pm) \rrbracket, \end{aligned} \quad (3.36)$$

where $e_i^s = e_i \cap \Omega^s$, $s = +, -$, and $d(\cdot)$ is defined in (3.5). It is easy to see that $g_i = 0$ when $e_i^+ = \emptyset$ or $e_i^+ = e_i$.

Proof. For simplicity, define a function w_h such that

$$w_h|_{T_h^s} = w_h^s \quad \text{with} \quad w_h^s = I_h v_E^s - (I_h^{IFE} v)^s, \quad s = +, -. \quad (3.37)$$

It is obvious that $w_h^s \in V_h(T)$, $s = +, -$. Define another functions v_h by

$$\begin{aligned} v_h &:= \llbracket w_h^\pm \rrbracket(D) \Psi(x) + \llbracket \beta_c^\pm \nabla w_h^\pm \cdot \mathbf{n}_h \rrbracket(x_p) \Upsilon_1(x) \\ &\quad + \llbracket \nabla w_h^\pm \cdot \mathbf{t}_h \rrbracket(x_p) \Upsilon_2(x) + \sum_{i \in \mathcal{I}} N_i(w_h) \phi_i(x) + \llbracket d(w_h^\pm) \rrbracket \Theta(x). \end{aligned} \quad (3.38)$$

Next, we prove $w_h = v_h$. From (3.13), (3.1)-(3.3) and Remark 3.1, the IFE basis functions ϕ_i , $i \in \mathcal{I}$ satisfy the following identities

$$\begin{aligned} \llbracket \phi_i^\pm \rrbracket(D) &= 0, \quad \llbracket \beta_c^\pm \nabla \phi_i^\pm \cdot \mathbf{n}_h \rrbracket(x_p) = 0, \quad \llbracket \nabla \phi_i^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 0, \\ \llbracket d(\phi_i^\pm) \rrbracket &= 0, \quad N_j(\phi_i) = \delta_{i,j}, \quad i, j \in \mathcal{I}. \end{aligned} \quad (3.39)$$

Combining (3.38), (3.39) and (3.23)-(3.28), we find

$$\llbracket \beta_c^\pm \nabla v_h^\pm \cdot \mathbf{n}_h \rrbracket(x_p) = \llbracket \beta_c^\pm \nabla w_h^\pm \cdot \mathbf{n}_h \rrbracket(x_p), \quad N_i(v_h) = N_i(w_h), \quad i \in \mathcal{I} \quad (3.40)$$

and

$$\llbracket v_h^\pm \rrbracket(D) = \llbracket w_h^\pm \rrbracket(D), \quad \llbracket \nabla v_h^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = \llbracket \nabla w_h^\pm \cdot \mathbf{t}_h \rrbracket(x_p), \quad \llbracket d(v_h^\pm) \rrbracket = \llbracket d(w_h^\pm) \rrbracket. \quad (3.41)$$

It follows from (3.41) that $v_h - w_h$ is continuous across $\Gamma_h \cap T$, which together with (3.40) and (3.2)-(3.3), implies

$$v_h - w_h \in S_h(T) \quad \text{and} \quad N_i(v_h - w_h) = 0, \quad i \in \mathcal{I}.$$

Using the unisolvence of IFE shape functions (see Lemma 3.3), we know that the function $v_h - w_h$ is unique and $v_h - w_h = 0$ through a simple verification. Thus, from (3.38), we have

$$w_h = v_h = a\Psi(x) + b_1 \Upsilon_1(x) + b_2 \Upsilon_2(x) + \sum_{i \in \mathcal{I}} g_i \phi_i(x) + t\Theta(x), \quad (3.42)$$

where

$$a = \llbracket w_h^\pm \rrbracket(D), \quad b_1 = \llbracket \beta_c^\pm \nabla w_h^\pm \cdot \mathbf{n}_h \rrbracket(x_p), \quad b_2 = \llbracket \nabla w_h^\pm \cdot \mathbf{t}_h \rrbracket(x_p), \quad g_i = N_i(w_h), \quad t = \llbracket d(I_h v_E^\pm) \rrbracket.$$

Using the following facts from (3.2)-(3.3)

$$\llbracket I_h^{IFE} v \rrbracket(D) = 0, \quad \llbracket \beta_c^\pm \nabla(I_h^{IFE} v)^\pm \cdot \mathbf{n}_h \rrbracket(x_p) = 0, \quad \llbracket \nabla(I_h^{IFE} v)^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = 0,$$

we can further derive

$$\begin{aligned} a &= \llbracket w_h^\pm \rrbracket(D) = \llbracket I_h v_E^\pm - (I_h^{IFE} v)^\pm \rrbracket(D) = \llbracket I_h v_E^\pm \rrbracket(D) - \llbracket (I_h^{IFE} v)^\pm \rrbracket(D) = \llbracket I_h v_E^\pm \rrbracket(D), \\ b_1 &= \llbracket \beta_c^\pm \nabla w_h^\pm \cdot \mathbf{n}_h \rrbracket(x_p) = \llbracket \beta_c^\pm \nabla (I_h v_E^\pm - (I_h^{IFE} v)^\pm) \cdot \mathbf{n}_h \rrbracket(x_p) = \llbracket \beta_c^\pm \nabla (I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket(x_p), \\ b_2 &= \llbracket \nabla w_h^\pm \cdot \mathbf{t}_h \rrbracket(x_p) = \llbracket \nabla (I_h v_E^\pm - (I_h^{IFE} v)^\pm) \cdot \mathbf{t}_h \rrbracket(x_p) = \llbracket \nabla (I_h v_E^\pm) \cdot \mathbf{t}_h \rrbracket(x_p). \end{aligned}$$

It remains to consider g_i , $i \in \mathcal{I}$. Define

$$I_h^{BK} v := \begin{cases} I_h v_E^+ & \text{in } T_h^+, \\ I_h v_E^- & \text{in } T_h^-, \end{cases}$$

then we know from (3.37) that $w_h = I_h^{BK} v - I_h^{IFE} v$. Using the fact that $N_i(v - I_h^{IFE} v) = 0$ on interface elements from (3.18), we obtain

$$\begin{aligned} g_i &= N_i(w_h) = N_i(I_h^{BK} v - I_h^{IFE} v) = N_i(I_h^{BK} v - v + v - I_h^{IFE} v) \\ &= N_i(I_h^{BK} v - v) + N_i(v - I_h^{IFE} v) = N_i(I_h^{BK} v - v) \\ &= |e_i|^{-1} \left(\int_{e_i^+} (I_h v_E^+ - v_E^+) ds + \int_{e_i^-} (I_h v_E^- - v_E^-) ds \right). \end{aligned}$$

□

Theorem 3.7. *For any $v \in \tilde{H}^2(\Omega)$, there exists a constant C independent of h and the interface location relative to the mesh such that*

$$\sum_{T \in \mathcal{T}_h^\Gamma} |v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 \leq C h_\Gamma^{4-2m} \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2, \quad m = 0, 1, 2, \quad s = +, -,$$

where h_Γ is defined in (2.4).

Proof. On each interface element $T \in \mathcal{T}_h^\Gamma$, by the triangle inequality, we have

$$|v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)} \leq |v_E^s - I_h v_E^s|_{W_2^m(T)} + |I_h v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)}, \quad s = +, -. \quad (3.43)$$

The estimate of the first term is standard

$$|v_E^s - I_h v_E^s|_{W_2^m(T)}^2 \leq C h_T^{4-2m} |v_E^s|_{H^2(T)}^2, \quad m = 0, 1, 2, \quad s = +, -. \quad (3.44)$$

For the second term on the right-hand side of (3.43), from Lemma 3.6, Lemma 3.5 and (3.16), we have

$$\begin{aligned} &|I_h v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 \\ &\leq C \left(a^2 |\Psi^s|_{W_2^m(T)}^2 + \sum_{i=1,2} b_i^2 |\Upsilon_i^s|_{W_2^m(T)}^2 + \sum_{i \in \mathcal{I}} g_i^2 |\phi_i^s|_{W_2^m(T)}^2 + t^2 |\Theta^s|_{W_2^m(T)}^2 \right) \\ &\leq C h_T^{2-2m} \left(a^2 + \sum_{i \in \mathcal{I}} g_i^2 \right) + C h_T^{4-2m} \sum_{i=1,2} b_i^2 + C h_T^{6-2m} t^2, \quad m = 0, 1, 2. \end{aligned} \quad (3.45)$$

Here the constants t , a , b_1 , b_2 and g_i , $i \in \mathcal{I}$ are defined in (3.36). We now estimate these constants one by one. Firstly, (3.5) implies

$$\begin{aligned} t^2 &= |\llbracket d(I_h v_E^\pm) \rrbracket|^2 = \frac{1}{(4\kappa_T + 4)|T|} |\llbracket I_h v_E^\pm \rrbracket|_{H^2(T)}^2 \leq C h_T^{-2} \sum_{s=+,-} |I_h v_E^s|_{H^2(T)}^2 \\ &\leq C h_T^{-2} \sum_{s=+,-} \left(|v_E^s|_{H^2(T)}^2 + |v_E^s - I_h v_E^s|_{H^2(T)}^2 \right) \leq C h_T^{-2} \sum_{s=+,-} |v_E^s|_{H^2(T)}^2. \end{aligned} \quad (3.46)$$

Since $v \in \tilde{H}^2(\Omega) = 0$, we have $\llbracket v_E^\pm \rrbracket^2(D)$, which leads to

$$\begin{aligned} a^2 &= \llbracket I_h v_E^\pm \rrbracket^2(D) = \llbracket I_h v_E^\pm - v_E^\pm \rrbracket^2(D) \leq \|\llbracket I_h v_E^\pm - v_E^\pm \rrbracket\|_{L^\infty(T)}^2 \\ &\leq C \|I_h v_E^+ - v_E^+\|_{L^\infty(T)}^2 + C \|I_h v_E^- - v_E^-\|_{L^\infty(T)}^2 \leq Ch_T^2 (|v_E^+|_{H^2(T)}^2 + |v_E^-|_{H^2(T)}^2), \end{aligned} \quad (3.47)$$

where the standard interpolation error estimate is used in the last inequality, see Theorem 4.4.20 in [4].

For the constant b_1 , using the standard inverse inequality, we have

$$\begin{aligned} b_1^2 &= \|\beta_c^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h\|^2(x_p) \leq \|\llbracket \beta_c^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^\infty(T)}^2 \leq Ch_T^{-2} \|\llbracket \beta_c^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2 \\ &\leq Ch_T^{-2} \|\llbracket \beta^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2 + Ch_T^{-2} \|\llbracket (\beta_c^\pm - \beta^\pm) \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2. \end{aligned} \quad (3.48)$$

From (3.4), the second term can be estimate as

$$\begin{aligned} h_T^{-2} \|\llbracket (\beta_c^\pm - \beta^\pm) \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2 &\leq h_T^{-2} \sum_{s=+,-} \|\beta_c^s - \beta^s\|_{L^\infty(T)}^2 \|\nabla(I_h v_E^s) \cdot \mathbf{n}_h\|_{L^2(T)}^2 \\ &\leq C \sum_{s=+,-} \left(\|\nabla(I_h v_E^s) \cdot \mathbf{n}_h - \nabla v_E^s \cdot \mathbf{n}_h\|_{L^2(T)}^2 + \|\nabla v_E^s \cdot \mathbf{n}_h\|_{L^2(T)}^2 \right) \leq \sum_{s=+,-} |v_E^s|_{H^1(T)}^2. \end{aligned}$$

For the first term on the right-hand side of (3.48), using (2.10), we get

$$\begin{aligned} h_T^{-2} \|\llbracket \beta^\pm \nabla(I_h v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2 &= h_T^{-2} \|\llbracket \beta^\pm \nabla(I_h v_E^\pm - v_E^\pm) \cdot \mathbf{n}_h + \beta^\pm \nabla v_E^\pm \cdot (\mathbf{n}_h - \mathbf{n} + \mathbf{n}) \rrbracket\|_{L^2(T)}^2 \\ &\leq Ch_T^{-2} \left(\|\llbracket \beta^\pm \nabla(I_h v_E^\pm - v_E^\pm) \cdot \mathbf{n}_h \rrbracket\|_{L^2(T)}^2 + \|\mathbf{n} - \mathbf{n}_h\|_{L^\infty(T)}^2 \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(T)}^2 + \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(T)}^2 \right) \\ &\leq C \sum_{s=+,-} \left(|v_E^s|_{H^2(T)}^2 + |v_E^s|_{H^1(T)}^2 \right) + Ch_T^{-2} \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(T)}^2. \end{aligned}$$

Combining above three inequalities yields

$$b_1^2 \leq C \sum_{s=+,-} \left(|v_E^s|_{H^2(T)}^2 + |v_E^s|_{H^1(T)}^2 \right) + Ch_T^{-2} \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(T)}^2. \quad (3.49)$$

Analogously,

$$\begin{aligned} b_2^2 &= \|\nabla(I_h v_E^\pm) \cdot \mathbf{t}_h\|^2(x_p) \leq \|\llbracket \nabla(I_h v_E^\pm) \cdot \mathbf{t}_h \rrbracket\|_{L^\infty(T)}^2 \leq Ch_T^{-2} \|\llbracket \nabla(I_h v_E^\pm) \cdot \mathbf{t}_h \rrbracket\|_{L^2(T)}^2 \\ &\leq Ch_T^{-2} \|\llbracket \nabla(I_h v_E^\pm - v_E^\pm) \cdot \mathbf{t}_h + \nabla v_E^\pm \cdot (\mathbf{t}_h - \mathbf{t} + \mathbf{t}) \rrbracket\|_{L^2(T)}^2 \\ &\leq Ch_T^{-2} \left(\|\llbracket \nabla(I_h v_E^\pm - v_E^\pm) \cdot \mathbf{t}_h \rrbracket\|_{L^2(T)}^2 + \|\mathbf{t}_h - \mathbf{t}\|_{L^\infty(T)}^2 \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(T)}^2 + \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(T)}^2 \right) \\ &\leq C \sum_{s=+,-} \left(|v_E^s|_{H^2(T)}^2 + |v_E^s|_{H^1(T)}^2 \right) + Ch_T^{-2} \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(T)}^2. \end{aligned} \quad (3.50)$$

Finally, the constants g_i , $i \in \mathcal{I}$ in (3.36) can be estimated by the Cauchy-Schwarz inequality and the standard interpolation error estimate

$$\begin{aligned} g_i^2 &= |e_i|^{-2} \left| \int_{e_i^+} (I_h v_E^+ - v_E^+) ds + \int_{e_i^-} (I_h v_E^- - v_E^-) ds \right|^2 \leq Ch_T^{-1} \sum_{s=+,-} \|I_h v_E^s - v_E^s\|_{L^2(e_i)}^2 \\ &\leq C \sum_{s=+,-} \left(h_T^{-2} \|I_h v_E^s - v_E^s\|_{L^2(T)}^2 + \|\nabla(I_h v_E^s - v_E^s)\|_{L^2(T)}^2 \right) \leq Ch_T^2 \sum_{i=+,-} |v_E^s|_{H^2(T)}^2. \end{aligned} \quad (3.51)$$

We now combine (3.43)-(3.47), (3.49)-(3.51) to obtain the error estimate on interface elements

$$|v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 \leq Ch_T^{4-2m} \sum_{s=\pm} \|v_E^s\|_{H^2(T)}^2 + Ch_T^{2-2m} \left(\|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(T)}^2 + \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(T)}^2 \right).$$

Summing up the estimate over all interface elements $T \in \mathcal{T}_h^\Gamma$ and using Assumption (2.3), we get

$$\begin{aligned} \sum_{T \in \mathcal{T}_h^\Gamma} |v_E^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 &\leq Ch_\Gamma^{4-2m} \|v_E^s\|_{H^2(\Omega)}^2 + Ch_\Gamma^{2-2m} \left(\|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(U(\Gamma, h_\Gamma))}^2 \right. \\ &\quad \left. + \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(U(\Gamma, h_\Gamma))}^2 \right), \quad s = +, -, \quad m = 0, 1, 2. \end{aligned} \quad (3.52)$$

Since $v \in \tilde{H}^2(\Omega)$, we know from the definition (2.1) that $\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket(x) = 0$ and $\llbracket v_E^\pm \rrbracket(x) = 0$ for all $x \in \Gamma$, which also implies $\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket(x) = 0$ on Γ . Thus, by Lemma 2.4, and (2.5)-(2.6),

$$\begin{aligned} \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{L^2(U(\Gamma, h_\Gamma))}^2 &\leq Ch_\Gamma^2 \|\llbracket \beta^\pm \nabla v_E^\pm \cdot \mathbf{n} \rrbracket\|_{H^1(U(\Gamma, h_\Gamma))}^2 \leq Ch_\Gamma^2 \sum_{s=+,-} \|v_E^s\|_{H^2(\Omega)}^2, \\ \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{L^2(U(\Gamma, h_\Gamma))}^2 &\leq Ch_\Gamma^2 \|\llbracket \nabla v_E^\pm \cdot \mathbf{t} \rrbracket\|_{H^1(U(\Gamma, h_\Gamma))}^2 \leq Ch_\Gamma^2 \sum_{s=+,-} \|v_E^s\|_{H^2(\Omega)}^2. \end{aligned}$$

Substituting the above inequalities into (3.52) and using the extension result (3.20) we complete the proof. \square

Now we prove the optimal approximation capabilities of the nonconforming IFE spaces V_h^{IFE} , where the errors resulting from the mismatch of Γ and Γ_h are taken into account.

Theorem 3.8. *For any $v \in \tilde{H}^2(\Omega)$, there exists a constant C independent of h and the interface location relative to the mesh such that*

$$\sum_{T \in \mathcal{T}_h} |v - I_h^{IFE} v|_{W_2^m(T)}^2 \leq Ch^{4-2m} \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2, \quad m = 0, 1. \quad (3.53)$$

Proof. On each non-interface element $T \in \mathcal{T}_h^{non}$, the following estimate is standard

$$|v - I_h^{IFE} v|_{W_2^m(T)}^2 = |v - I_h v|_{W_2^m(T)}^2 \leq Ch_T^{4-2m} |v|_{H^2(T)}^2, \quad m = 0, 1. \quad (3.54)$$

On each interface element $T \in \mathcal{T}_h^\Gamma$, in view of the relations $T = T^+ \cup T^-$ and $T^s = (T^s \cap T_h^+) \cup (T^s \cap T_h^-)$, $s = +, -$, we have

$$\begin{aligned} |v - I_h^{IFE} v|_{W_2^m(T)}^2 &= \sum_{s=+,-} |v^s - (I_h^{IFE} v)^s|_{W_2^m(T^s \cap T_h^s)}^2 \\ &\quad + |v^- - (I_h^{IFE} v)^+|_{W_2^m(T^- \cap T_h^+)}^2 + |v^+ - (I_h^{IFE} v)^-|_{W_2^m(T^+ \cap T_h^-)}^2. \end{aligned} \quad (3.55)$$

By the triangle inequality,

$$\begin{aligned} |v^- - (I_h^{IFE} v)^+|_{W_2^m(T^- \cap T_h^+)}^2 &\leq 2|v^- - v_E^+|_{W_2^m(T^- \cap T_h^+)}^2 + 2|v_E^+ - (I_h^{IFE} v)^+|_{W_2^m(T^- \cap T_h^+)}^2, \\ |v^+ - (I_h^{IFE} v)^-|_{W_2^m(T^+ \cap T_h^-)}^2 &\leq 2|v^+ - v_E^-|_{W_2^m(T^+ \cap T_h^-)}^2 + 2|v_E^- - (I_h^{IFE} v)^-|_{W_2^m(T^+ \cap T_h^-)}^2. \end{aligned} \quad (3.56)$$

Substituting (3.56) into (3.55) and using the definition (2.11), we get

$$|v - I_h^{IFE} v|_{W_2^m(T)}^2 \leq C \sum_{s=+,-} |v^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 + C |v_E^+ - v_E^-|_{W_2^m(T^\Delta)}^2, \quad m = 0, 1. \quad (3.57)$$

It follows from Lemma (2.5) and the fact $\llbracket v_E^\pm \rrbracket = 0$ on $\Gamma \cap T$ that

$$\begin{aligned} |v_E^+ - v_E^-|_{L^2(T^\Delta)}^2 &\leq Ch_T^4 \|\llbracket v_E^\pm \rrbracket\|_{H^1(T^\Delta)}^2 \leq Ch_T^4 \sum_{s=+,-} |v_E^s|_{H^1(T)}^2, \\ \|\nabla(v_E^+ - v_E^-)\|_{L^2(T^\Delta)}^2 &\leq C \left(h_T^2 \|\nabla \llbracket v_E^\pm \rrbracket\|_{L^2(\Gamma \cap T)}^2 + h_T^4 \|\llbracket v_E^\pm \rrbracket\|_{H^2(T^\Delta)}^2 \right) \\ &\leq Ch_T^2 \sum_{s=+,-} \|\nabla v_E^s\|_{L^2(\Gamma \cap T)}^2 + Ch_T^4 \sum_{s=+,-} |v_E^s|_{H^2(T)}^2, \end{aligned}$$

which implies

$$|v_E^+ - v_E^-|_{W_2^m(T^\Delta)}^2 \leq Ch_T^{4-2m} \sum_{s=+,-} \left(|v_E^s|_{H^1(T)}^2 + \|\nabla v_E^s\|_{L^2(\Gamma \cap T)}^2 \right), \quad m = 0, 1. \quad (3.58)$$

Combining (3.54), (3.57) and (3.58), we arrive at

$$\begin{aligned} \sum_{T \in \mathcal{T}_h} |v - I_h^{IFE} v|_{W_2^m(T)}^2 &\leq Ch^{4-2m} \sum_{T \in \mathcal{T}_h^{non}} |v|_{H^2(T)}^2 + C \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} |v^s - (I_h^{IFE} v)^s|_{W_2^m(T)}^2 \\ &\quad + Ch^{4-2m} \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} |v_E^s|_{H^1(T)}^2 + Ch^{4-2m} \|\nabla v_E^s\|_{L^2(\Gamma)}^2 \quad m = 0, 1, \end{aligned}$$

which together with Theorem 3.7, the extension result (3.20) and the following global trace inequality

$$\sum_{s=+,-} \|\nabla v_E^s\|_{L^2(\Gamma)}^2 \leq C(\|v_E^+\|_{H^2(\Omega^+)}^2 + \|v_E^-\|_{H^2(\Omega^-)}^2) \quad (3.59)$$

yields the theorem. \square

4 Error estimates of the existing nonconforming IFE method

In this section, we show that the existing nonconforming IFE method [17, 23] cannot achieve the optimal convergence rates unless $[\beta]_\Gamma \nabla u \cdot \mathbf{t} = 0$ on Γ . The existing nonconforming IFE method is to find $u_h \in V_{h,0}^{IFE}$ such that

$$a_h(u_h, v_h) := \sum_{T \in \mathcal{T}_h} \int_T \beta(x) \nabla u_h \cdot \nabla v_h dx = \int_\Omega f v_h dx \quad \forall v_h \in V_{h,0}^{IFE}. \quad (4.1)$$

It is easy to see that $a_h(\cdot, \cdot) = a(\cdot, \cdot)$ on $H_0^1(\Omega)$ and it is positive-definite on $V_{h,0}^{IFE}$ because $a_h(v_h, v_h) = 0$, $v_h \in V_{h,0}^{IFE}$ implies $v_h = 0$. Thus, the discrete problem (4.1) has a unique solution. We define the energy norm

$$\|v\|_{a_h} := \sqrt{a_h(v, v)} \quad \forall v \in V_{h,0}^{IFE} + H_0^1(\Omega)$$

and quote the following well-known second Strang lemma (see Lemma 10.1.9 in [4]).

Lemma 4.1. *Let u and u_h be the solutions of the problems (2.2) and (4.1), respectively. Then*

$$\|u - u_h\|_{a_h} \leq C \left\{ \inf_{v_h \in V_{h,0}^{IFE}} \|u - v_h\|_{a_h} + \sup_{w_h \in V_{h,0}^{IFE} \setminus \{0\}} \frac{|a_h(u - u_h, w_h)|}{\|w_h\|_{a_h}} \right\}. \quad (4.2)$$

Since $u \in \tilde{H}^2(\Omega)$, Theorem 3.8 implies

$$\inf_{v_h \in V_{h,0}^{IFE}} \|u - v_h\|_{a_h} \leq \|u - I_h^{IFE} u\|_{a_h} \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)}. \quad (4.3)$$

For the second term on the right-hand side of (4.2), we have

$$a_h(u - u_h, w_h) = \sum_{T \in \mathcal{T}_h} \int_T \beta \nabla u \cdot \nabla w_h dx - \int_\Omega f w_h dx = \sum_{e \in \mathcal{E}_h} \int_e \beta \nabla u \cdot \mathbf{n}_e [w_h]_e ds,$$

where the jump $[w_h]_e \mathbf{n}_e$ across an edge e is defined as follows. Let e be an interior edge shared by two elements T_1^e and T_2^e , and \mathbf{n}_e the unit normal of e pointing towards the outside of T_1^e . Define

$$[w_h]_e \mathbf{n}_e := (w_h|_{T_1^e} - w_h|_{T_2^e}) \mathbf{n}_e \quad \text{on } e.$$

If e is an edge on the boundary of Ω , then define $[w_h]_e \mathbf{n}_e = w_h \mathbf{n}_e$, where \mathbf{n}_e is the unit normal of e pointing towards the outside of Ω . Given an edge e and an element T , define the standard L^2 projection operators P_0^e and P_0^T as

$$P_0^e f = |e|^{-1} \int_e f ds, \quad P_0^T f = |T|^{-1} \int_T f dx.$$

It follows from the fact $\int_e [w_h]_e ds = 0$ that

$$\begin{aligned} |a_h(u - u_h, w_h)| &= \left| \sum_{e \in \mathcal{E}_h} \int_e \beta \nabla u \cdot \mathbf{n}_e [w_h]_e ds \right| = \left| \sum_{e \in \mathcal{E}_h} \int_e (\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)) [w_h]_e ds \right| \\ &\leq \left(\sum_{e \in \mathcal{E}_h} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 \right)^{1/2} \left(\sum_{e \in \mathcal{E}_h} \|[w_h]_e\|_{L^2(e)}^2 \right)^{1/2}. \end{aligned} \quad (4.4)$$

Lemma 4.2. *There exists a constant C independent of h and the interface location relative to the mesh such that*

$$\|[w_h]_e\|_{L^2(e)}^2 \leq C|e| \left(|w_h|_{H^1(T_1^e)}^2 + |w_h|_{H^1(T_2^e)}^2 \right) \quad \forall e \in \mathcal{E}_h^\circ \quad \forall w_h \in V_h^{IFE}.$$

Proof. Using the fact that $P_0^e(w_h|_{T_1^e}) = P_0^e(w_h|_{T_2^e})$, we have

$$\begin{aligned} \|[w_h]_e\|_{L^2(e)}^2 &= \int_e (w_h|_{T_1^e} - P_0^e(w_h|_{T_1^e}) - w_h|_{T_2^e} + P_0^e(w_h|_{T_2^e}))^2 ds \\ &\leq 2 \sum_{i=1,2} \int_e (w_h|_{T_i^e} - P_0^e(w_h|_{T_i^e}))^2 ds. \end{aligned}$$

By the property of L^2 projection

$$\|f - P_0^e(f)\|_{L^2(e)} \leq \|f - c_e\|_{L^2(e)} \quad \forall c_e \in \mathbb{R}$$

and the standard trace inequality, we get

$$\begin{aligned} \|[w_h]_e\|_{L^2(e)}^2 &\leq 2 \sum_{i=1,2} \int_e (w_h|_{T_i^e} - P_0^e(w_h|_{T_i^e}))^2 ds \leq 2 \sum_{i=1,2} \int_e (w_h|_{T_i^e} - P_0^{T_i^e}(w_h|_{T_i^e}))^2 ds \\ &\leq C \sum_{i=1,2} \left(h_{T_i^e}^{-1} \left\| w_h|_{T_i^e} - P_0^{T_i^e}(w_h|_{T_i^e}) \right\|_{L^2(T_i^e)}^2 + h_{T_i^e} \left| w_h|_{T_i^e} - P_0^{T_i^e}(w_h|_{T_i^e}) \right|_{H^1(T_i^e)}^2 \right) \\ &\leq C \left(h_{T_1^e} |w_h|_{H^1(T_1^e)}^2 + h_{T_2^e} |w_h|_{H^1(T_2^e)}^2 \right), \end{aligned}$$

where the fact that $w_h|_T \in H^1(T)$ is used in the last inequality even though $T \in \mathcal{T}_h^\Gamma$. \square

If e is an edge on the boundary of Ω , we also have similar estimates. Thus,

$$\sum_{e \in \mathcal{E}_h} \|[w_h]_e\|_{L^2(e)}^2 \leq Ch \|w_h\|_{a_h}^2. \quad (4.5)$$

The next step is to estimate $\|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}$ in (4.4). Let T be an element such that $e \subset \partial T$. If $e \in \mathcal{E}_h^{non}$ and $T \in \mathcal{T}_h^{non}$, using the property of the L^2 projection operator P_0^e and the standard trace inequality, we have

$$\begin{aligned} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)} &\leq \|\beta \nabla u \cdot \mathbf{n}_e - P_0^T(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)} \\ &\leq Ch_T^{-1/2} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^T(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(T)} + Ch_T^{1/2} |\beta \nabla u \cdot \mathbf{n}_e|_{H^1(T)} \\ &\leq Ch_T^{1/2} |u|_{H^2(T)}. \end{aligned} \quad (4.6)$$

If $e \in \mathcal{E}_h^{non}$ and $T \in \mathcal{T}_h^\Gamma$, the term can be estimated by using the fact that $e \in \Omega^s$, $s = +$ or $-$, and the extension (3.20),

$$\begin{aligned} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)} &= \|\beta^s \nabla u_E^s \cdot \mathbf{n}_e - P_0^e(\beta^s \nabla u_E^s \cdot \mathbf{n}_e)\|_{L^2(e)} \\ &\leq Ch_T^{-1/2} \|\beta^s \nabla u_E^s \cdot \mathbf{n}_e - P_0^T(\beta^s \nabla u_E^s \cdot \mathbf{n}_e)\|_{L^2(T)} + Ch_T^{1/2} |\beta \nabla u \cdot \mathbf{n}_e|_{H^1(T)} \\ &\leq Ch_T^{1/2} |u_E^s|_{H^2(T)}. \end{aligned} \quad (4.7)$$

Hence, it follows from (4.6)-(4.7) and the extension result (3.20) that

$$\sum_{e \in \mathcal{E}_h^{non}} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 \leq Ch \sum_{i=+,-} |u_E^s|_{H^2(\Omega)}^2 \leq Ch |u|_{H^2(\Omega^+ \cup \Omega^-)}^2. \quad (4.8)$$

For interface edges $e \in \mathcal{E}_h^\Gamma$, we cannot conclude the optimal estimate since $(\beta \nabla u \cdot \mathbf{n}_e)|_e$ may have a jump across $e \cap \Gamma$. Noticing that $[u]_\Gamma = 0$ implies $[\nabla u \cdot \mathbf{t}]_\Gamma = 0$, the jump can be derived as

$$[\beta \nabla u \cdot \mathbf{n}_e]_\Gamma = [\beta \nabla u \cdot \mathbf{n}]_\Gamma \mathbf{n} \cdot \mathbf{n}_e + [\beta \nabla u \cdot \mathbf{t}]_\Gamma \mathbf{t} \cdot \mathbf{n}_e = [\beta]_\Gamma (\nabla u \cdot \mathbf{t})(\mathbf{t} \cdot \mathbf{n}_e), \quad (4.9)$$

where we have used

$$[\beta \nabla u \cdot \mathbf{n}]_\Gamma = 0 \quad \text{and} \quad [\beta \nabla u \cdot \mathbf{t}]_\Gamma = \frac{1}{2}(\beta^+ + \beta^-)[\nabla u \cdot \mathbf{t}]_\Gamma + [\beta]_\Gamma \nabla u \cdot \mathbf{t} = [\beta]_\Gamma \nabla u \cdot \mathbf{t}.$$

The following lemma gives an estimate on interface edges.

Lemma 4.3. *Let u be the solution of the problem (2.2). Assume the triangulation near the interface is quasi-uniform, i.e., $h_T^{-1} \leq Ch_\Gamma^{-1}$ for all $T \in \mathcal{T}_h^\Gamma$. Then it holds that*

$$\sum_{e \in \mathcal{E}_h^\Gamma} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 \leq Ch_\Gamma \sum_{T \in \mathcal{T}_h^\Gamma} |u|_{H^2(T^+ \cup T^-)}^2 + C \|[\beta]_\Gamma \nabla u \cdot \mathbf{t}\|_{H^{1/2}(\Gamma)}^2. \quad (4.10)$$

Proof. Define a function $z|_{\Omega^s} = z^s$, $s = +, -$, such that

$$\begin{aligned} -\Delta z^s + z^s &= 0 \quad \text{in } \Omega^s, \quad s = +, -, \\ z^s &= [\beta]_\Gamma \nabla u \cdot \mathbf{t} \quad \text{on } \Gamma, \quad \frac{\partial z^+}{\partial \nu} = 0 \quad \text{on } \partial\Omega, \quad s = +, -, \end{aligned}$$

where ν is the outward unit normal vector to $\partial\Omega$. Since $[\beta]_\Gamma \nabla u \cdot \mathbf{t} \in H^{1/2}(\Gamma)$, the function z exists and satisfies

$$z|_\Gamma = [\beta]_\Gamma \nabla u \cdot \mathbf{t} \quad \text{and} \quad \|z\|_{H^1(\Omega)} \leq \|[\beta]_\Gamma \nabla u \cdot \mathbf{t}\|_{H^{1/2}(\Gamma)}. \quad (4.11)$$

For an edge $e \in \mathcal{E}_h^\Gamma$, let $w = \beta \nabla u \cdot \mathbf{n}_e$ and T be an element that has e as one of its edges. Define

$$\tilde{w}(x) = z(x) \mathbf{t}(x) \cdot \mathbf{n}_e \quad \forall x \in T \quad \text{and} \quad \hat{w}(x) = \begin{cases} \tilde{w} & \text{if } x \in T^+, \\ 0 & \text{if } x \in T^-. \end{cases}$$

From (4.9) and (4.11), we have $[w - \hat{w}]_{\Gamma \cap T} = 0$. Thus, $w - \hat{w} \in H^1(T)$. By the property of the L^2 projection operator P_0^e and the standard trace inequality, we derive

$$\begin{aligned} \|w - P_0^e(w)\|_{L^2(e)}^2 &\leq \|w - P_0^T(w - \hat{w})\|_{L^2(e)}^2 = \|w - \hat{w} + \hat{w} - P_0^T(w - \hat{w})\|_{L^2(e)}^2 \\ &\leq 2\|w - \hat{w} - P_0^T(w - \hat{w})\|_{L^2(e)}^2 + 2\|\hat{w}\|_{L^2(e)}^2 \\ &\leq C(h_T^{-1}\|w - \hat{w} - P_0^T(w - \hat{w})\|_{L^2(T)}^2 + h_T|w - \hat{w}|_{H^1(T)}^2) + 2\|z\|_{L^2(e)}^2 \\ &\leq C(h_T|w - \hat{w}|_{H^1(T)}^2 + h_T^{-1}\|z\|_{L^2(T)}^2 + h_T|z|_{H^1(T)}^2) \\ &\leq C(h_T|w|_{H^1(T^+ \cup T^-)}^2 + h_T^{-1}\|z\|_{L^2(T)}^2 + h_T|z|_{H^1(T)}^2). \end{aligned}$$

Summing over all interface edges and using Lemma 2.4, we get

$$\begin{aligned}
\sum_{e \in \mathcal{E}_h^\Gamma} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 &\leq \sum_{T \in \mathcal{T}_h^\Gamma} C(h_T |w|_{H^1(T^+ \cup T^-)}^2 + h_T^{-1} \|z\|_{L^2(T)}^2 + h_T |z|_{H^1(T)}^2) \\
&\leq Ch_\Gamma \sum_{T \in \mathcal{T}_h^\Gamma} |u|_{H^2(T^+ \cup T^-)}^2 + Ch_\Gamma^{-1} \|z\|_{L^2(U(\Gamma, h_\Gamma))}^2 + Ch_\Gamma |z|_{H^1(\Omega)}^2 \\
&\leq Ch_\Gamma \sum_{T \in \mathcal{T}_h^\Gamma} |u|_{H^2(T^+ \cup T^-)}^2 + C \|z\|_{H^1(\Omega)}^2,
\end{aligned}$$

which together with (4.11) yields this lemma. \square




Figure 3: A triangulation of a region Λ contained in Ω ; red: interface edges.

Remark 4.4. The estimate (4.10) is sharp, i.e., we cannot find a better upper bound for the approximation error than $O(1)$ when $[\beta]_\Gamma \nabla u \cdot \mathbf{t} \neq 0$ on Γ . We explain it by a concrete example as illustrated in Figure 3. The domain Ω contains a region Λ such that $\Lambda = (0, 1)^2$, $\Lambda^+ = \{x = (x_1, x_2) \in \Lambda : x_1 > x_2\}$, $\Lambda^- = \{x = (x_1, x_2) \in \Lambda : x_1 < x_2\}$. The interface contained in the region is $\Gamma \cap \Lambda = \{x = (x_1, x_2) \in \Lambda : x_1 = x_2\}$. We use uniform triangulations constructed by connecting the midpoints of edges (see Figure 3) and only consider these interface edges which are contained in the region Λ . Obviously, $\mathbf{n}_e = \mathbf{t} = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})^T$ and $\mathbf{n} = (\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}})^T$. Let $\beta^+ = 2$, $\beta^- = 1$ and the exact solution $u(x_1, x_2)$ be a piecewise linear function on Λ^+ , Λ^- such that

$$\beta^+ \nabla u^+ \cdot \mathbf{n} = \beta^- \nabla u^- \cdot \mathbf{n} = 1 \quad \text{and} \quad u|_{\Gamma \cap \Lambda} = \frac{1}{\sqrt{2}}(x_1 + x_2).$$

Thus, $\nabla u \cdot \mathbf{t} = 1$ and $(\beta \nabla u \cdot \mathbf{n}_e)|_{e^+} = 2$, $(\beta \nabla u \cdot \mathbf{n}_e)|_{e^-} = 1$ for all $e \subset \Lambda$ and $e \in \mathcal{E}_h^\Gamma$. Therefore,

$$\begin{aligned}
\|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 &= \inf_{c_e \in \mathbb{R}} \|\beta \nabla u \cdot \mathbf{n}_e - c_e\|_{L^2(e)}^2 \\
&= \inf_{c_e \in \mathbb{R}} \left(\frac{|e|}{2} (2 - c_e)^2 + \frac{|e|}{2} (1 - c_e)^2 \right) \geq \frac{|e|}{4} \geq Ch.
\end{aligned}$$

Using the fact that the number of interface edges contained in Λ is $O(h^{-1})$, we observe that

$$\sum_{e \in \mathcal{E}_h^\Gamma} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 \geq \sum_{e \in \mathcal{E}_h^\Gamma, e \subset \Lambda} Ch \geq C.$$

With the above discussions, we have the following error estimate for the existing nonconforming IFE method.

Theorem 4.5. Let u and u_h be the solutions of the problems (2.2) and (4.1), respectively. Under the assumption of Lemma 4.3, the following discretization error estimate holds

$$\|u - u_h\|_{a_h} \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)} + Ch^{1/2} \|[\beta]_\Gamma \nabla u \cdot \mathbf{t}\|_{H^{1/2}(\Gamma)}. \quad (4.12)$$

Proof. It follows from (4.8) and (4.10) that

$$\sum_{e \in \mathcal{E}_h} \|\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)\|_{L^2(e)}^2 \leq Ch|u|_{H^2(\Omega^+ \cup \Omega^-)}^2 + C \|[\beta]_\Gamma \nabla u \cdot \mathbf{t}\|_{H^{1/2}(\Gamma)}^2, \quad (4.13)$$

which together with (4.4) and (4.5) yields

$$|a_h(u - u_h, w_h)| \leq Ch^{1/2} \|w_h\|_{a_h} \left(h^{1/2} |u|_{H^2(\Omega^+ \cup \Omega^-)} + \|[\beta]_\Gamma \nabla u \cdot \mathbf{t}\|_{H^{1/2}(\Gamma)} \right). \quad (4.14)$$

Combining (4.2), (4.3) and (4.14), we complete the proof. \square

Remark 4.6. The estimate (4.12) in Theorem 4.5 suggests that the solution of the nonconforming IFE method (4.1) only converges with a suboptimal convergence rate $O(h^{1/2})$ in the energy norm if $[\beta]_\Gamma \nabla u \cdot \mathbf{t} \neq 0$ on Γ . Applying global trace inequalities on Ω^+ and Ω^- to the second term on the right-hand side of (4.12), we get

$$\|u - u_h\|_{a_h} \leq Ch^{1/2} \|u\|_{H^2(\Omega^+ \cup \Omega^-)}.$$

Using the standard duality argument (see [4, p. 284]), we can also derive the following suboptimal L^2 error estimate

$$\|u - u_h\|_{L^2(\Omega)} \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)}. \quad (4.15)$$

5 A new nonconforming IFE method and error estimates

To recover the optimal convergence rates, we propose a new nonconforming IFE method. On each interface element $T \in \mathcal{T}_h^\Gamma$, define

$$W_h(T) := \{w_h \in (L^2(T))^2 : w_h = \nabla v_h \quad \forall v_h \in S_h(T)\}. \quad (5.1)$$

We also define a space associated with an edge $e \in \mathcal{E}_h^\Gamma$ as

$$W_e := \{w_h \in (L^2(\Omega))^2 : w_h|_{T_1^e} \in W_h(T_1^e), w_h|_{T_2^e} \in W_h(T_2^e), w_h|_{\Omega \setminus (T_1^e \cup T_2^e)} = 0\}. \quad (5.2)$$

For simplicity of the implementation, the coefficient $\beta(x)$ is approximated by

$$\beta_h(x) = \begin{cases} \beta^+(x) & \text{if } x \in \Omega_h^+, \\ \beta^-(x) & \text{if } x \in \Omega_h^-. \end{cases}$$

Define a local *lifting operator* $r_e : L^2(e) \rightarrow W_e$ such that

$$\int_\Omega \beta_h(x) r_e(\varphi) \cdot w_h dx = \int_e \{\beta_h w_h \cdot \mathbf{n}_e\}_e \varphi ds \quad \forall w_h \in W_e, \quad (5.3)$$

where $\{v\}_e = \frac{1}{2}(v|_{T_1^e} + v|_{T_2^e})$. Obviously, the support of $r_e(\varphi)$ is $T_1^e \cup T_2^e$.

The new nonconforming IFE method is to find $u_h \in V_{h,0}^{IFE}$ such that

$$A_h(u_h, v_h) := \tilde{a}_h(u_h, v_h) + b_h(u_h, v_h) + s_h(u_h, v_h) = \int_\Omega f v_h dx \quad \forall v_h \in V_{h,0}^{IFE}, \quad (5.4)$$

where

$$\begin{aligned} \tilde{a}_h(u_h, v_h) &= \sum_{T \in \mathcal{T}_h} \int_T \beta_h(x) \nabla u_h \cdot \nabla v_h dx, \\ b_h(u_h, v_h) &= - \sum_{e \in \mathcal{E}_h^\Gamma} \int_e (\{\beta_h \nabla u_h \cdot \mathbf{n}_e\}_e [v_h]_e + \{\beta_h \nabla v_h \cdot \mathbf{n}_e\}_e [u_h]_e) ds, \\ s_h(u_h, v_h) &= 4 \sum_{e \in \mathcal{E}_h^\Gamma} \int_{T_1^e \cup T_2^e} \beta_h(x) r_e([u_h]_e) \cdot r_e([v_h]_e) dx. \end{aligned} \quad (5.5)$$

Note that the new nonconforming IFE method is symmetric and does not require a manually chosen stabilization parameter. Now we analyze the new method. For all $v \in H_0^1(\Omega) \cap \tilde{H}^2(\Omega) + V_{h,0}^{IFE}$, define the following mesh-dependent norms

$$\|v\|_{\tilde{a}_h} := \sqrt{\tilde{a}_h(v, v)}$$

and

$$\| \| v \| \|_h := \left(\|v\|_{\tilde{a}_h}^2 + \sum_{e \in \mathcal{E}_h^\Gamma} |e| \|\{\beta_h \nabla v\}_e\|_{L^2(e)}^2 + \sum_{e \in \mathcal{E}_h^\Gamma} |e|^{-1} \|[v]_e\|_{L^2(e)}^2 + s_h(v, v) \right)^{1/2}. \quad (5.6)$$

The continuity of the bilinear form $A(\cdot, \cdot)$ is verified directly from the Cauchy-Schwarz inequality

$$|A_h(w, v)| \leq \| \| w \| \|_h \| \| v \| \|_h \quad \forall w, v \in H_0^1(\Omega) \cap \tilde{H}^2(\Omega) + V_{h,0}^{IFE}. \quad (5.7)$$

The following lemma demonstrates the coercivity of the bilinear form $A(\cdot, \cdot)$ on the IFE space $V_{h,0}^{IFE}$ with respect to the norm $\| \| \cdot \| \|_{\tilde{a}_h}$.

Lemma 5.1. *It holds that*

$$A_h(v_h, v_h) \geq \frac{1}{2} \|v_h\|_{\tilde{a}_h}^2 \quad \forall v_h \in V_{h,0}^{IFE}. \quad (5.8)$$

Proof. For all $v_h \in V_{h,0}^{IFE}$, choosing $w_h|_{T_1^e \cup T_2^e} = \nabla v_h$, $w_h|_{\Omega \setminus (T_1^e \cup T_2^e)} = 0$ in (5.3) and using the fact that the support of $r_e(\varphi)$ is $T_1^e \cup T_2^e$, we have

$$\int_{T_1^e \cup T_2^e} \beta_h(x) r_e(\varphi) \cdot \nabla v_h dx = \int_{\Omega} \beta_h(x) r_e(\varphi) \cdot \nabla v_h dx = \int_e \{\beta_h \nabla v_h \cdot \mathbf{n}_e\}_e \varphi ds.$$

It follows from the Cauchy-Schwarz inequality that

$$\begin{aligned} |b_h(v_h, v_h)| &= \left| 2 \sum_{e \in \mathcal{E}_h^\Gamma} \int_e \{\beta_h \nabla v_h \cdot \mathbf{n}_e\}_e [v_h]_e ds \right| \\ &\leq \left(4 \sum_{e \in \mathcal{E}_h^\Gamma} \int_{\Omega} \beta_h r_e([v_h]_e) \cdot r_e([v_h]_e) dx \right)^{1/2} \left(\sum_{e \in \mathcal{E}_h^\Gamma} \int_{T_1^e \cup T_2^e} \beta_h \nabla v_h \cdot \nabla v_h dx \right)^{1/2}. \end{aligned} \quad (5.9)$$

Since each interface element has at most two interface edges from Assumption 2.2, each element is calculated at most twice. Therefore,

$$\sum_{e \in \mathcal{E}_h^\Gamma} \int_{T_1^e \cup T_2^e} \beta_h \nabla v_h \cdot \nabla v_h dx \leq 2 \sum_{T \in \mathcal{T}_h} \int_T \beta_h(x) \nabla v_h \cdot \nabla v_h dx. \quad (5.10)$$

Combining (5.5), (5.9) and (5.10), we have

$$\begin{aligned} |b_h(v_h, v_h)| &\leq (s_h(v_h, v_h))^{1/2} \left(2 \sum_{T \in \mathcal{T}_h} \int_T \beta_h(x) \nabla v_h \cdot \nabla v_h dx \right)^{1/2} \\ &\leq s_h(v_h, v_h) + \frac{1}{2} \sum_{T \in \mathcal{T}_h} \int_T \beta_h(x) \nabla v_h \cdot \nabla v_h dx, \end{aligned}$$

which together with (5.5) yields the result

$$A(v_h, v_h) = \tilde{a}_h(v_h, v_h) + b_h(v_h, v_h) + s_h(v_h, v_h) \geq \frac{1}{2} \sum_{T \in \mathcal{T}_h} \int_T \beta_h(x) \nabla v_h \cdot \nabla v_h dx = \frac{1}{2} \|v_h\|_{\tilde{a}_h}^2.$$

□

Next, we show the equivalence of the two norms $\|\cdot\|_{\tilde{a}_h}$ and $|||\cdot|||_h$ on the IFE space $V_{h,0}^{IFE}$. To begin with, we need the following trace inequality for the IFE shape functions in $S_h(T)$ which can be verified via straightforward calculations. We also refer readers to [23, Theorem 2.7].

Lemma 5.2. *There exists a constant C independent of h and the interface location relative to the mesh such that*

$$\|\nabla v_h\|_{L^2(\partial T)} \leq Ch_T^{-1/2} \|\nabla v_h\|_{L^2(T)} \quad \forall v_h \in S_h(T) \quad \forall T \in \mathcal{T}_h^\Gamma. \quad (5.11)$$

We also need the following stability estimate for the local lifting operator r_e .

Lemma 5.3. *There exists a constant C independent of h and the interface location relative to the mesh such that*

$$\|r_e(\varphi)\|_{L^2(\Omega)} \leq C|e|^{-1/2} \|\varphi\|_{L^2(e)} \quad \forall \varphi \in L^2(e) \quad \forall e \in \mathcal{E}_h^\Gamma.$$

Proof. Since the support of $r_e(\varphi)$ is $T_1^e \cup T_2^e$, choosing $w_h = r_e(\varphi)$ in (5.3) yields

$$\begin{aligned} \|r_e(\varphi)\|_{L^2(\Omega)}^2 &\leq C \|\beta_h^{1/2} r_e(\varphi)\|_{L^2(T_1^e \cup T_2^e)}^2 = C \int_e \{\beta_h r_e(\varphi) \cdot \mathbf{n}_e\}_e \varphi ds \\ &\leq C \|\{\beta_h r_e(\varphi)\}_e\|_{L^2(e)} \|\varphi\|_{L^2(e)} \leq C \|\varphi\|_{L^2(e)} \sum_{i=1,2} \|r_e(\varphi)|_{T_i^e}\|_{L^2(e)}. \end{aligned} \quad (5.12)$$

Note $r_e(\varphi)|_{T_1^e} \in W_h(T_1^e)$, we know from (5.1) that there exists a function $v_h \in S_h(T_1^e)$ such that $r_e(\varphi)|_{T_1^e} = \nabla v_h$. By the trace inequality (5.11) for the IFE shape functions, we have

$$\|r_e(\varphi)|_{T_1^e}\|_{L^2(e)} = \|\nabla v_h\|_{L^2(e)} \leq Ch_T^{-1/2} \|\nabla v_h\|_{L^2(T_1^e)} = Ch_T^{-1/2} \|r_e(\varphi)\|_{L^2(T_1^e)},$$

which, together with (5.12) and a similar estimate on T_2^e , completes the proof of this lemma. \square

We now prove the norm-equivalence in the following lemma.

Lemma 5.4. *There exists a constant C independent of h and the interface location relative to the mesh such that*

$$\|v_h\|_{\tilde{a}_h} \leq |||v_h|||_h \leq C \|v_h\|_{\tilde{a}_h} \quad \forall v_h \in V_{h,0}^{IFE}. \quad (5.13)$$

Proof. We just need to prove the second inequality since the first inequality is obvious. By the trace inequality (5.11) for the IFE shape functions, it holds

$$\sum_{e \in \mathcal{E}_h^\Gamma} |e| \|\{\beta_h \nabla v_h\}_e\|_{L^2(e)}^2 \leq C \sum_{e \in \mathcal{E}_h^\Gamma} \sum_{i=1,2} \|\nabla v_h\|_{L^2(T_i^e)}^2 \leq C \|v_h\|_{\tilde{a}_h}^2. \quad (5.14)$$

From Lemma 4.2, we have

$$\sum_{e \in \mathcal{E}_h^\Gamma} |e|^{-1} \|[v_h]_e\|_{L^2(e)}^2 \leq C \sum_{e \in \mathcal{E}_h^\Gamma} \sum_{i=1,2} \|\nabla v_h\|_{L^2(T_i^e)}^2 \leq C \|v_h\|_{\tilde{a}_h}^2, \quad (5.15)$$

which, together with Lemma 5.3 for the local lifting operator, leads to

$$s_h(v_h, v_h) \leq C \sum_{e \in \mathcal{E}_h^\Gamma} \|r_e([v_h]_e)\|_{L^2(\Omega)}^2 \leq C \sum_{e \in \mathcal{E}_h^\Gamma} |e|^{-1} \|[v_h]_e\|_{L^2(e)}^2 \leq C \|v_h\|_{\tilde{a}_h}^2. \quad (5.16)$$

Combining (5.6), (5.14)-(5.16), we get the second inequality in (5.13). \square

The following lemma provides an optimal estimate for the interpolation error in terms of the norm $|||\cdot|||_h$.

Lemma 5.5. Suppose $v \in \tilde{H}^2(\Omega)$, then there exists a constant C independent of h and the interface location relative to the mesh such that

$$\|v - I_h^{IFE} v\|_h \leq Ch \|v\|_{H^2(\Omega^+ \cup \Omega^-)}.$$

Proof. The first term in the norm $\|\cdot\|$ can be bounded by Theorem 3.8

$$\|v - I_h^{IFE} v\|_{\tilde{a}_h} \leq Ch \|v\|_{H^2(\Omega^+ \cup \Omega^-)}. \quad (5.17)$$

Since $(v - I_h^{IFE} v)|_T \in H^1(T)$ for all $T \in \mathcal{T}_h^\Gamma$, by the standard trace inequality and Theorem 3.8, we have

$$\begin{aligned} \sum_{e \in \mathcal{E}_h^\Gamma} |e|^{-1} \|v - I_h^{IFE} v\|_{L^2(e)}^2 &\leq C \sum_{e \in \mathcal{E}_h^\Gamma} \sum_{i=1,2} \left(h_T^{-2} \|v - I_h^{IFE} v\|_{L^2(T_i^e)}^2 + |v - I_h^{IFE} v|_{H^1(T_i^e)}^2 \right) \\ &\leq C \sum_{T \in \mathcal{T}_h^\Gamma} \left(h_T^{-2} \|v - I_h^{IFE} v\|_{L^2(T)}^2 + |v - I_h^{IFE} v|_{H^1(T)}^2 \right) \leq Ch_\Gamma^2 \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2, \end{aligned} \quad (5.18)$$

which, together with Lemma 5.3, implies

$$s_h(v - I_h^{IFE} v, v - I_h^{IFE} v) \leq C \sum_{e \in \mathcal{E}_h^\Gamma} |e|^{-1} \|v - I_h^{IFE} v\|_{L^2(e)}^2 \leq Ch_\Gamma^2 \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2. \quad (5.19)$$

Let $e^s = e \cap \Omega^s$, $s = +, -$. Recalling the notations in (3.19)-(3.20), it holds

$$\begin{aligned} \|\{\beta_h \nabla(v - I_h^{IFE} v)\}_e\|_{L^2(e)}^2 &= \|\{\beta_h \nabla(v - I_h^{IFE} v)\}_e\|_{L^2(e^+)}^2 + \|\{\beta_h \nabla(v - I_h^{IFE} v)\}_e\|_{L^2(e^-)}^2 \\ &\leq C \|\{\nabla(v_E^+ - (I_h^{IFE} v)^+)\}_e\|_{L^2(e)}^2 + C \|\{\nabla(v_E^- - (I_h^{IFE} v)^-)\}_e\|_{L^2(e)}^2. \end{aligned}$$

Then using the standard trace inequality and Theorem 3.7, we can derive

$$\begin{aligned} \sum_{e \in \mathcal{E}_h^\Gamma} |e| \|\{\beta_h \nabla(v - I_h^{IFE} v)\}_e\|_{L^2(e)}^2 \\ \leq C \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} \left(|v_E^s - (I_h^{IFE} v)^s|_{H^1(T)}^2 + h_T^2 |v_E^s|_{H^2(T)}^2 \right) \leq Ch_\Gamma^2 \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2. \end{aligned} \quad (5.20)$$

The lemma follows from (5.6), (5.17)-(5.20). \square

The following lemma concerns the errors caused by replacing $\beta(x)$ by $\beta_h(x)$.

Lemma 5.6. Let $v \in \tilde{H}^2(\Omega)$ and $w \in V_h^{IFE} + H^1(\Omega)$. Then there exists a constant C independent of h and the interface location relative to the mesh such that

$$|a_h(v, w) - \tilde{a}_h(v, w)| \leq Ch_\Gamma \|v\|_{H^2(\Omega^+ \cup \Omega^-)} \left(\sum_{T \in \mathcal{T}_h^\Gamma} \|\nabla w\|_{L^2(T^\Delta)}^2 \right)^{1/2}. \quad (5.21)$$

Furthermore, if $w \in \tilde{H}^2(\Omega)$, there holds

$$|a_h(v, w) - \tilde{a}_h(v, w)| \leq Ch_\Gamma^2 \|v\|_{H^2(\Omega^+ \cup \Omega^-)} \|w\|_{H^2(\Omega^+ \cup \Omega^-)}. \quad (5.22)$$

Proof. The Cauchy-Schwarz inequality gives

$$\begin{aligned} |a_h(v, w) - \tilde{a}_h(v, w)| &= \left| \sum_{T \in \mathcal{T}_h^\Gamma} \int_{T^\Delta} (\beta - \beta_h) \nabla v \cdot \nabla w dx \right| \\ &\leq C \left(\sum_{T \in \mathcal{T}_h^\Gamma} \|\nabla v\|_{L^2(T^\Delta)}^2 \right)^{1/2} \left(\sum_{T \in \mathcal{T}_h^\Gamma} \|\nabla w\|_{L^2(T^\Delta)}^2 \right)^{1/2}. \end{aligned} \quad (5.23)$$

Using Lemma 2.5 and the global trace inequalities on Ω^+ and Ω^- , we derive

$$\begin{aligned}
\sum_{T \in \mathcal{T}_h^\Gamma} \|\nabla v\|_{L^2(T^\Delta)}^2 &= \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} \|\nabla v^s\|_{L^2(T^\Delta \cap T^s)}^2 \leq \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} \|\nabla v_E^s\|_{L^2(T^\Delta)}^2 \\
&\leq C \sum_{T \in \mathcal{T}_h^\Gamma} \sum_{s=+,-} \left(h_T^2 \|\nabla v^s\|_{L^2(T \cap \Gamma)}^2 + h_T^4 |v_E^s|_{H^2(T^\Delta)}^2 \right) \\
&\leq Ch_\Gamma^2 \sum_{s=+,-} \|\nabla v^s\|_{L^2(\Gamma)}^2 + Ch_\Gamma^4 \sum_{s=+,-} |v_E^s|_{H^2(\Omega)}^2 \\
&\leq Ch_\Gamma^2 \sum_{s=+,-} \|v\|_{H^2(\Omega^s)}^2 = Ch_\Gamma^2 \|v\|_{H^2(\Omega^+ \cup \Omega^-)}^2.
\end{aligned} \tag{5.24}$$

The estimate (5.21) follows from (5.23) and (5.24). If $w \in \tilde{H}^2(\Omega)$, then similar to (5.24),

$$\sum_{T \in \mathcal{T}_h^\Gamma} \|\nabla w\|_{L^2(T^\Delta)}^2 \leq Ch_\Gamma^2 \|w\|_{H^2(\Omega^+ \cup \Omega^-)}^2. \tag{5.25}$$

The estimate (5.22) then follows from (5.21) and (5.25). \square

With these preparations, we are ready to derive the H^1 error estimate for the new nonconforming IFE method.

Theorem 5.7. *Let u and u_h be the solutions of the problems (2.2) and (5.4), respectively. Then there exists a constant C independent of h and the interface location relative to the mesh such that*

$$\| \| u - u_h \| \|_h \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)}. \tag{5.26}$$

Proof. From Lemma 5.1 and Lemma 5.4, we know that the bilinear form $A_h(\cdot, \cdot)$ is also coercive on V_h^{IFE} with respect to the norm $\| \| \cdot \| \|_h$. Thus, the second Strang lemma implies

$$\| \| u - u_h \| \|_h \leq C \left\{ \inf_{v_h \in V_{h,0}^{IFE}} \| \| u - v_h \| \|_h + \sup_{w_h \in V_{h,0}^{IFE} \setminus \{0\}} \frac{|A_h(u - u_h, w_h)|}{\| \| w_h \| \|_h} \right\}. \tag{5.27}$$

The first term of the right-hand side of (5.27) can be bounded by Lemma 5.5

$$\inf_{v_h \in V_{h,0}^{IFE}} \| \| u - v_h \| \|_h \leq \| \| u - I_h^{IFE} u \| \|_h \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)}. \tag{5.28}$$

Multiplying (1.1) by $w_h \in V_{h,0}^{IFE}$ and integrating by parts, we obtain

$$\begin{aligned}
\int_{\Omega} f w_h dx &= a_h(u, w_h) + s_h(u, w_h) - \sum_{e \in \mathcal{E}_h} \int_e \{\beta \nabla u \cdot \mathbf{n}_e\}_e [w_h]_e + \{\beta \nabla w_h \cdot \mathbf{n}_e\}_e [u]_e ds \\
&= A_h(u, w_h) - \sum_{e \in \mathcal{E}_h^{non}} \int_e \beta \nabla u \cdot \mathbf{n}_e [w_h]_e ds + a_h(u, w_h) - \tilde{a}_h(u, w_h),
\end{aligned} \tag{5.29}$$

where $\beta_h(x) = \beta(x)$ on edges, $[u]_e = 0$, $[\beta \nabla u \cdot \mathbf{n}_e]_e = 0$ and $r_e([u]_e) = 0$ are used. It follows from (5.29) and (5.4) that

$$A_h(u - u_h, w_h) = \sum_{e \in \mathcal{E}_h^{non}} \int_e \beta \nabla u \cdot \mathbf{n}_e [w_h]_e ds + \tilde{a}_h(u, w_h) - a_h(u, w_h). \tag{5.30}$$

Hence, by (4.4), (4.5), (4.8), and Lemma (5.6), we have

$$|A_h(u - u_h, w_h)| \leq Ch \|u\|_{H^2(\Omega^+ \cup \Omega^-)} \| \| w_h \| \|,$$

which, together with (5.27) and (5.28), completes the proof of the theorem. \square

The optimal L^2 error estimate is also derived by using the standard duality argument below.

Theorem 5.8. *Let u and u_h be the solutions of the problems (2.2) and (5.4), respectively. Then there exists a constant C independent of h and the interface location relative to the mesh such that*

$$\|u - u_h\|_{L^2(\Omega)} \leq Ch^2 \|u\|_{H^2(\Omega \cup \Omega^-)}. \quad (5.31)$$

Proof. Let $z \in H_0^1(\Omega)$ be the solution of the following auxiliary problem

$$a(v, z) = \int_{\Omega} (u - u_h) v dx \quad \forall v \in H_0^1(\Omega). \quad (5.32)$$

Since $u - u_h \in L^2(\Omega)$, it follows from Theorem 2.1 that

$$z \in \tilde{H}^2(\Omega) \quad \text{and} \quad \|z\|_{H^2(\Omega \cup \Omega^-)} \leq C \|u - u_h\|_{L^2(\Omega)}. \quad (5.33)$$

Let $z_h \in V_{h,0}^{IFE}$ be the solution of the new nonconforming IFE method applied to the auxiliary problem (5.32), i.e.,

$$A_h(v_h, z_h) = \int_{\Omega} (u - u_h) v_h dx \quad \forall v_h \in V_{h,0}^{IFE}. \quad (5.34)$$

Recalling that $a_h(\cdot, \cdot) = a(\cdot, \cdot)$ on $H_0^1(\Omega)$, and applying (5.32) and (5.34), we have

$$\begin{aligned} \|u - u_h\|_{L^2(\Omega)}^2 &= a(u, z) - A_h(u_h, z_h) = A_h(u, z) - A_h(u_h, z_h) + \tilde{a}_h(u, z) - a_h(u, z) \\ &= A_h(u - u_h, z - z_h) + A_h(u - u_h, z_h) + A_h(u_h, z - z_h) + (\tilde{a}_h(u, z) - a_h(u, z)), \end{aligned} \quad (5.35)$$

where the relation $b_h(u, z) = s_h(u, z) = 0$ is used in the second identity since $[u]_e = [v]_e = 0$ for all edges. Lemma 5.6 provides the estimate for the last term

$$\tilde{a}_h(u, z) - a_h(u, z) \leq Ch^2 \|u\|_{H^2(\Omega \cup \Omega^-)} \|z\|_{H^2(\Omega \cup \Omega^-)}. \quad (5.36)$$

The first terms on the right-hand side of (5.35) can be estimated using Theorem 5.7,

$$A_h(u - u_h, z - z_h) \leq \|u - u_h\|_h \|z - z_h\|_h \leq Ch^2 \|u\|_{H^2(\Omega \cup \Omega^-)} \|z\|_{H^2(\Omega \cup \Omega^-)}. \quad (5.37)$$

We rewrite the second term on the right-hand side of (5.35) as

$$A_h(u - u_h, z_h) = A_h(u - u_h, z_h - I_h^{IFE} z) + A_h(u - u_h, I_h^{IFE} z_h). \quad (5.38)$$

It is easy to see that

$$A_h(u - u_h, z_h - I_h^{IFE} z) \leq \|u - u_h\|_h \|z_h - I_h^{IFE} z\|_h \leq Ch^2 \|u\|_{H^2(\Omega \cup \Omega^-)} \|z\|_{H^2(\Omega \cup \Omega^-)}. \quad (5.39)$$

From (5.30), we have

$$A_h(u - u_h, I_h^{IFE} z_h) = \sum_{e \in \mathcal{E}_h^{non}} \int_e \beta \nabla u \cdot \mathbf{n}_e [I_h^{IFE} z_h]_e ds + \tilde{a}_h(u, I_h^{IFE} z_h) - a_h(u, I_h^{IFE} z_h). \quad (5.40)$$

Since $[I_h^{IFE} z_h]_e = [z]_e = 0$, we can derive

$$\begin{aligned} \sum_{e \in \mathcal{E}_h^{non}} \int_e \beta \nabla u \cdot \mathbf{n}_e [I_h^{IFE} z_h]_e ds &= \sum_{e \in \mathcal{E}_h^{non}} \int_e (\beta \nabla u \cdot \mathbf{n}_e - P_0^e(\beta \nabla u \cdot \mathbf{n}_e)) [I_h^{IFE} z_h - z]_e ds \\ &\leq Ch^2 \|u\|_{H^2(\Omega \cup \Omega^-)} \|z\|_{H^2(\Omega \cup \Omega^-)}, \end{aligned} \quad (5.41)$$

where the Cauchy-Schwarz inequality, (4.8), the standard trace inequality and Theorem 3.8 are used. Applying Lemma 5.6 and Theorem 3.8 again we obtain

$$\begin{aligned} |\tilde{a}_h(u, I_h^{IFE} z_h) - a_h(u, I_h^{IFE} z_h)| &\leq |\tilde{a}_h(u, z) - a_h(u, z)| \\ &\quad + |\tilde{a}_h(u, I_h^{IFE} z_h - z) - a_h(u, I_h^{IFE} z_h - z)| \\ &\leq Ch^2 \|u\|_{H^2(\Omega^+ \cup \Omega^-)} \|z\|_{H^2(\Omega^+ \cup \Omega^-)}. \end{aligned} \quad (5.42)$$

Combining (5.38)-(5.42), we find

$$A_h(u - u_h, z_h) \leq Ch^2 \|u\|_{H^2(\Omega^+ \cup \Omega^-)} \|z\|_{H^2(\Omega^+ \cup \Omega^-)}, \quad (5.43)$$

and similarly,

$$A_h(u_h, z - z_h) \leq Ch^2 \|u\|_{H^2(\Omega^+ \cup \Omega^-)} \|z\|_{H^2(\Omega^+ \cup \Omega^-)}. \quad (5.44)$$

Applying (5.35)-(5.37), (5.43)-(5.44), we arrive at the estimate

$$\|u - u_h\|_{L^2(\Omega)}^2 \leq Ch^2 \|u\|_{H^2(\Omega^+ \cup \Omega^-)} \|z\|_{H^2(\Omega^+ \cup \Omega^-)},$$

which together with the regularity result (5.33) implies the estimate (5.31). \square

6 Numerical examples

In this section, we present some numerical examples to validate the theoretical analysis. To avoid redundancy, we only report numerical results of IFE methods based on the Crouzeix-Raviart element since the results of IFE methods based on the rotated- Q_1 element are almost the same.

For simplicity, we take $\Omega = (-1, 1) \times (-1, 1)$ as the computational domain and use uniform triangulations constructed as follows. We first partition the domain into $N \times N$ congruent rectangles, and then obtain the triangulation by cutting the rectangles along one of diagonals in the same direction. We examine the convergence rate of IFE solutions using the following norms

$$|e_h|_{H^1} := \left(\sum_{T \in \mathcal{T}_h} \|\sqrt{\beta_h} \nabla(u - u_h)\|_{L^2(T)}^2 \right)^{1/2} \quad \text{and} \quad \|e_h\|_{L^2} := \|u - u_h\|_{L^2(\Omega)}.$$

For comparison, we replace $\beta(x)$ by $\beta_h(x)$ in the existing nonconforming IFE method (4.1) in our computation. Thus, the difference of the existing nonconforming IFE method and our new nonconforming IFE method (5.4) is the terms $b_h(\cdot, \cdot)$ and $s_h(\cdot, \cdot)$. In view of the analysis for our new method, the error resulting from replacing $\beta(x)$ by $\beta_h(x)$ does not affect the error estimates in Theorem 4.5 for the existing nonconforming IFE method.

6.1 A counter example with $\nabla u \cdot \mathbf{t} \neq 0$ on Γ

We use this example to show that the existing nonconforming IFE method using integral values as degrees of freedom does not have the optimal convergence rates.

Example 6.1. The interface is $\Gamma = \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 = r_0^2\}$ such that $\Omega^- = \{(x_1, x_2) \in \mathbb{R}^2 : x_1^2 + x_2^2 < r_0^2\}$. Let (r, θ) be the polar coordinate of $x = (x_1, x_2)$. The exact solution is chosen as $u(x) = j(x)v(x)\omega(x)$, where $\omega(x) = \sin(\theta)$,

$$j(x) = \begin{cases} \exp\left(-\frac{1}{1 - (r - r_0)^2/\eta^2}\right) & \text{if } |r - r_0| < \eta, \\ 0 & \text{if } |r - r_0| \geq \eta, \end{cases}$$

and

$$v(x) = \begin{cases} 1 + (r^2 - r_0^2)/\beta^+(x) & \text{if } x \in \Omega^+, \\ 1 + (r^2 - r_0^2)/\beta^-(x) & \text{if } x \in \Omega^-. \end{cases}$$

Let $r_0 = 0.5$, $\eta = 0.45$, $\beta^+(x)$ and $\beta^-(x)$ be positive constants. It is easy to verify that the jump condition (1.2)-(1.3) is satisfied and $\nabla u \cdot \mathbf{t} \neq 0$ on Γ . We test two cases: $(\beta^+, \beta^-) = (10, 1000)$ and $(\beta^+, \beta^-) = (1000, 10)$. The exact solutions of these two cases are plotted in Figure 4.




Figure 4: Exact solutions of *Example 6.1*. Left: $(\beta^+, \beta^-) = (10, 1000)$; Right: $(\beta^+, \beta^-) = (1000, 10)$

We report numerical results in Tables 1-2 which clearly confirm our theoretical analysis. The second and third columns in Tables 1-2 indicate suboptimal convergence rates: $\|e_h\|_{L^2} \approx O(h)$, $|e_h|_{H^1} \approx O(h^{1/2})$ for the existing nonconforming IFE method (4.1). When the terms $b_h(\cdot, \cdot)$ and $s_h(\cdot, \cdot)$ are added to the scheme, i.e., the new nonconforming IFE method (5.4), we observe the optimal convergence rates (see last two columns in Tables 1-2).

Table 1: Numerical results of *Example 6.1* with $(\beta^+, \beta^-) = (10, 1000)$.

N	Existing nonconforming IFE method				New nonconforming IFE method			
	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate
8	2.221E-01		2.140E+01		1.617E-01		1.781E+01	
16	7.650E-02	1.54	1.037E+01	1.05	4.414E-02	1.87	6.889E+00	1.37
32	1.745E-02	2.13	5.970E+00	0.80	5.989E-03	2.88	3.851E+00	0.84
64	7.322E-03	1.25	3.597E+00	0.73	7.855E-04	2.93	1.784E+00	1.11
128	3.204E-03	1.19	2.309E+00	0.64	1.935E-04	2.02	8.932E-01	1.00
256	1.514E-03	1.08	1.548E+00	0.58	4.836E-05	2.00	4.461E-01	1.00
512	7.276E-04	1.06	1.056E+00	0.55	1.197E-05	2.01	2.229E-01	1.00
1024	3.603E-04	1.01	7.378E-01	0.52	2.992E-06	2.00	1.114E-01	1.00

Table 2: Numerical results of *Example 6.1* with $(\beta^+, \beta^-) = (1000, 10)$.

	Existing nonconforming IFE method				New nonconforming IFE method			
N	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate
8	1.736E-01		4.641E+01		1.540E-01		4.611E+01	
16	7.679E-02	1.18	1.686E+01	1.46	6.402E-02	1.27	1.579E+01	1.55
32	1.186E-02	2.69	8.927E+00	0.92	5.868E-03	3.45	8.055E+00	0.97
64	5.188E-03	1.19	4.822E+00	0.89	8.575E-04	2.77	3.888E+00	1.05
128	2.242E-03	1.21	2.802E+00	0.78	1.944E-04	2.14	1.949E+00	1.00
256	1.061E-03	1.08	1.746E+00	0.68	4.841E-05	2.01	9.738E-01	1.00
512	5.043E-04	1.07	1.133E+00	0.62	1.218E-05	1.99	4.868E-01	1.00
1024	2.483E-04	1.02	7.654E-01	0.57	2.982E-06	2.03	2.434E-01	1.00

6.2 An example with variable coefficients and a non-convex interface

Example 6.2. The interface is the zero level set of the function,

$$\varphi(x) = (3(x_1^2 + x_2^2) - x_1)^2 - x_1^2 - x_2^2 + 0.02.$$

The exact solution is chosen as $u(x) = \varphi(x)/\beta(x)$, where

$$\beta(x) = \begin{cases} \beta^+(x) = 300(2 + \sin(6x_1 + 6x_2)) & \text{if } \varphi(x) > 0, \\ \beta^-(x) = 2 + \cos(6x_1 + 6x_2) & \text{if } \varphi(x) < 0. \end{cases}$$

It is easy to verify that the jump condition (1.2)-(1.3) is satisfied and $\nabla u \cdot \mathbf{t} = 0$ on Γ . The exact solution and the interface are plotted in Figure 5.

For the problem with variable coefficients, we choose $\beta_c^+ = \beta^+(x_m)$, $\beta_c^- = \beta^-(x_m)$ on each interface element $T \in \mathcal{T}_h^\Gamma$, where x_m is the midpoint of $\Gamma_h \cap T$. Since $\nabla u \cdot \mathbf{t} = 0$ on Γ , our theoretical analysis suggests the optimal convergence rates for both the existing and the new IFE methods, which are confirmed by the results listed in Table 3.

In view of the requirement (3.4), we also test another choice: $\beta_c^\pm = \beta^\pm(A_i) + C^\pm h$, i.e., using the value at one vertex A_i of T and random constants C^\pm in $[0, 1]$. The numerical results are almost the same and thus are not presented to avoid redundancy.

Table 3: Numerical results of *Example 6.2*.

	Existing nonconforming IFE method				New nonconforming IFE method			
N	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate	$\ e_h\ _{L^2}$	rate	$ e_h _{H^1}$	rate
8	1.510E-02		1.744E+00		1.959E-02		1.747E+00	
16	5.245E-03	1.53	8.899E-01	0.97	6.609E-03	1.57	8.921E-01	0.97
32	1.540E-03	1.77	4.541E-01	0.97	2.151E-03	1.62	4.568E-01	0.97
64	2.785E-04	2.47	2.277E-01	1.00	4.067E-04	2.40	2.287E-01	1.00
128	5.968E-05	2.22	1.140E-01	1.00	7.657E-05	2.41	1.142E-01	1.00
256	1.356E-05	2.14	5.701E-02	1.00	1.601E-05	2.26	5.709E-02	1.00
512	3.335E-06	2.02	2.851E-02	1.00	3.773E-06	2.09	2.854E-02	1.00
1024	7.856E-07	2.09	1.425E-02	1.00	8.338E-07	2.18	1.426E-02	1.00




Figure 5: The exact solution and the interface of *Example 6.2*. Left: the plot of $-u$; Right: the plot of the interface

7 Conclusions

In this paper, we have shown that the nonconforming IFE methods using integral-value degrees of freedom cannot achieve optimal convergence rates although the continuity of IFE shape functions is weakly enforced through average values over edges. The suboptimal convergence rates have been confirmed by a counter numerical example where the tangential derivative of the exact solution is not zero on the interface. We think there is similar issue for nonconforming IFE methods using integral-value degrees of freedom for solving elasticity and Stokes interface problems, which is an interesting topic in our future research.

To recover the optimal convergence rates, we have developed a new nonconforming IFE method with additional terms at interface edges. The new nonconforming IFE method is symmetric and the coercivity is ensured by a local lifting operator without a sufficient large penalty parameter. We have also proved that IFE basis functions based on the Crouzeix-Raviart elements are unisolvent on arbitrary triangles which is one of advantages compared with the IFEs using nodal values as degrees of freedom. The optimal approximation capabilities of nonconforming IFE spaces based on the Crouzeix-Raviart and the rotated- Q_1 elements have been derived via a novel approach where the problem with variable coefficients are also considered. The optimal error estimates for the IFE solutions in the H^1 - and L^2 - norms have been derived and confirmed by some numerical examples.

Acknowledgments: H. Ji is partially supported by the National Natural Science Foundation of China (Grants Nos. 11701291 and 11801281) and the Natural Science Foundation of Jiangsu Province (Grant No. BK20200848); F. Wang is partially supported by the National Natural Science Foundation of China (Grant Nos. 12071227 and 11871281) and the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant No. 20KJA110001); J. Chen is partially supported by the National Natural Science Foundation of China (Grant Nos. 11871281, 11731007 and 12071227); Z. Li is partially supported by a Simons grant (No. 633724).

References

- [1] I. Babuška. The finite element method for elliptic equations with discontinuous coefficients. *Computing*, 5:207–213, 1970.

- [2] J. H. Bramble and J. T. King. A robust finite element method for nonhomogeneous Dirichlet problems in domains with curved boundaries. *Math. Comp.*, 63:1–17, 1994.
- [3] J. H. Bramble and J. T. King. A finite element method for interface problems in domains with smooth boundaries and interfaces. *Adv. Comput. Math.*, 6:109–138, 1996.
- [4] S. Brenner and L. Scott. *The mathematical theory of finite element methods*, volume 15. Springer, 2008.
- [5] Z. Chen and J. Zou. Finite element methods and their convergence for elliptic and parabolic interface problems. *Numer. Math.*, 79:175–202, 1998.
- [6] C.-C. Chu, I. G. Graham, and T.-Y. Hou. A new multiscale finite element method for high-contrast elliptic interface problems. *Math. Comp.*, 79:1915–1955, 2010.
- [7] S. Frei and T. Richter. A locally modified parametric finite element method for interface problems. *SIAM J. Numer. Anal.*, 52:2315–2334, 2014.
- [8] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [9] Y. Gong, B. Li, and Z. Li. Immersed-interface finite-element methods for elliptic interface problems with non-homogeneous jump conditions. *SIAM J. Numer. Anal.*, 46:472–495, 2008.
- [10] R. Guo and T. Lin. A group of immersed finite-element spaces for elliptic interface problems. *IMA J. Numer. Anal.*, 39:482–511, 2019.
- [11] R. Guo, T. Lin, and X. Zhang. Nonconforming immersed finite element spaces for elliptic interface problems. *Comput. Math. Appl.*, 75:2002–2016, 2018.
- [12] X. He, T. Lin, and Y. Lin. Approximation capability of a bilinear immersed finite element space. *Numer. Methods Partial Differential Equations*, 24:1265–1300, 2008.
- [13] X. He, T. Lin, and Y. Lin. The convergence of the bilinear and linear immersed finite element solutions to interface problems. *Numer. Methods Partial Differential Equations*, 28:312–330, 2012.
- [14] J. Huang and J. Zou. Some new a priori estimates for second-order elliptic and parabolic interface problems. *J. Differential Equations*, 184:570–586, 2002.
- [15] J. Huang and J. Zou. Uniform a priori estimates for elliptic and static maxwell interface problems. *Discrete and Continuous Dynamical Systems - Series B*, 7:145–170, 2007.
- [16] H. Ji, F. Wang, J. Chen, and Z. Li. A new parameter free partially penalized immersed finite element and the optimal convergence analysis. *arXiv:2103.10025*, 2021.
- [17] D. Y. Kwak, K. T. Wee, and K. S. Chang. An analysis of a broken P_1 -nonconforming finite element method for interface problems. *SIAM J. Numer. Anal.*, 48:2117–2134, 2010.
- [18] J. Li, J. Markus, B. Wohlmuth, and J. Zou. Optimal a priori estimates for higher order finite elements for elliptic interface problems. *Appl. Numer. Math.*, 60:19–37, 2010.
- [19] Z. Li. The immersed interface method using a finite element formulation. *Appl. Numer. Math.*, 27:253–267, 1998.
- [20] Z. Li, T. Lin, Y. Lin, and R. Rogers. An immersed finite element space and its approximation capability. *Numer. Methods Partial Differential Equations*, 20:338–367, 2004.

- [21] Z. Li, T. Lin, and X. Wu. New Cartesian grid methods for interface problems using the finite element formulation. *Numer. Math.*, 96:61–98, 2003.
- [22] T. Lin, Y. Lin, and X. Zhang. Partially penalized immersed finite element methods for elliptic interface problems. *SIAM J. Numer. Anal.*, 53:1121–1144, 2015.
- [23] T. Lin, D. Sheen, and X. Zhang. A nonconforming immersed finite element method for elliptic interface problems. *J. Sci. Comput.*, 79:442–463, 2019.
- [24] J. Xu. Error estimates of the finite element method for the 2nd order elliptic equations with discontinuous coefficients. *J. Xiangtan University*, 1:1–5, 1982.
- [25] X. Zhang. *Nonconforming Immersed Finite Element Methods for Interface Problems (Ph.D. Thesis)*. 2013.