

Non-self adjoint impedance in Generalized Optimized Schwarz Methods

X.Claeys¹

¹Sorbonne Université, Université Paris-Diderot SPC, CNRS, Inria, Laboratoire
Jacques-Louis Lions, équipe Alpines

Abstract

We present a convergence theory for Optimized Schwarz Methods that rely on a non-local exchange operator and covers the case of coercive possibly non-self-adjoint impedance operators. This analysis also naturally deals with the presence of cross-points in subdomain partitions of arbitrary shape. In the particular case of self-adjoint impedance, we recover the theory proposed in [Claeys & Parolin,2021].

Introduction

Although domain decomposition (DD) literature now offers a wide panel of well established techniques in the case of symmetric positive definite systems, many problems of high applicative relevance do not fit this case. Wave propagation problems in particular, on which the present article focuses, traditionally lead to indefinite systems and can only be addressed by a much smaller set of DD methods.

While there exist overlapping domain decomposition approaches adapted to the wave context see e.g. [38, Chap.11] and [28, 30] and references therein, we shall rather be interested in non-overlapping domain decomposition, also called substructuring. In the approach proposed by Després [14] also known, together with its variants, as Optimized Schwarz Method (OSM) [22, 26], the wave equation with outgoing Robin boundary condition is solved in each subdomain, and neighbouring subdomains are coupled by swapping Robin traces across each interface. Robin traces involve an impedance factor the choice of which has a strong impact on the speed of convergence of the global solution algorithm. While this impedance factor was the wave number in the thesis of Després, it can be chosen operator valued and exterior Dirichlet-to-Neumann (DtN) maps were spotted as an ideal choice for model configurations with no cross-point (i.e. points where three subdomains are adjacent), see [34]. Consequently, many contributions searched for the best tuning of this impedance parameter, trying to approximate exterior DtN maps typically by means of Pade approximants. In this direction we refer in particular to the work of Antoine, Geuzaine and their collaborators [5, 20, 33].

In the original work of Després, a convergence proof had been provided for general geometrical configurations, but it gave no estimate regarding the rate of convergence and was restricted to a continuous (i.e. non-discretized) setting. Later Collino & Joly [10] proposed a general theoretical framework for OSM that remained restricted to continuous settings and required no cross point in the subdomain partition. The question of cross-points remained a

thorny issue that hindered the development of a general theoretical framework for the analysis of OSM, despite a few scattered contributions that focused on the cross-point issue [2, 24, 21].

Until recently, all variants of OSM systematically enforced a coupling between subdomains by means of the same local exchange operator that swapped traces from both sides of each interface. Because this local swapping operator is not continuous when the subdomain partition involves cross-points, in [6] we proposed to replace it by a non-local counterpart which paved the way to a convergence analysis similar to [10] but in general geometrical configurations where cross-points are allowed. This idea was then extended to discrete settings in [8] where an explicit estimate of the convergence rate was provided for general geometrical partitioning, regardless of the presence of cross-points.

The theoretical framework of [6, 8] holds for a whole family of impedance operators. It relies on the hypothesis that the impedance is symmetric positive definite (SPD), which covers certain OSM approaches that pre-existed in the literature, including the original Després algorithm for which a novel convergence estimate was derived, see Example 11.4 in [8]. However many variants of OSM involve impedance operators that are not SPD and thus cannot be analyzed through the theory of [6, 8]. It then appears natural to try to extend the later theory to the case of impedance operators that are not SPD.

In the present contribution, we extend this theory to the case of impedance operators that only need to be coercive and can thus admit a non-self-adjoint part. In this context, the non-local exchange operator that maintains a coupling between subdomains is not orthogonal anymore. Besides this extension, the present contribution also contains several other novelties.

- We identify the spectral bounds of the impedance operator that played a pivotal role in the convergence estimate of [8] as the continuity modulus and inf-sup constant of the trace operator, see Section 7.
- In Theorem 8.2 we exhibit a factorized form for the inverse of the skeleton formulation involving an oblique projector onto the space of Cauchy traces parallel to the space of traces that comply with transmission conditions, see Proposition 8.1.
- We show that the local swapping operator considered elsewhere in the literature on OSM is simply the exchange operator associated to the identity matrix as impedance, and we study which other impedances lead to this same local exchange operator, see Section 9.
- We show that the OSM strategies presented in [11, 17] are particular cases of the non-self-adjoint theory presented here.

The present contribution aims at extending the theory of [6, 8]. To our knowledge this is the first contribution that establishes a convergence theory for Optimized Schwarz Methods for such a general class of impedance operators and geometric partitions including the possibility of cross-points. Of course, in the special case of symmetric positive definite impedance, we recover the framework of [8]. We believe that this unifying framework may help better understanding the convergence properties of the substructuring strategies belonging to the OSM family. Our goal is theoretical here, which is why we do not provide any numerical result. We refer the reader to Section 14 of [8] and to Chapter 11 of [35] for numerical results in 2D and 3D for acoustics and electromagnetics in both homogeneous and heterogeneous media.

1 Problem under study

In the present section we describe the problem to be solved, as well as basic notations regarding the discretization strategy. To present the theoretical novelties, we base our presentation on a problem and a geometric configuration very close to the ones of [8].

1.1 Wave propagation problem

We consider a boundary value problem modeling scalar wave propagation in an *a priori* heterogeneous medium in \mathbb{R}^d with $d = 1, 2$ or 3 . The computational domain $\Omega \subset \mathbb{R}^d$ is assumed bounded and polygonal ($d = 2$) or polyhedral ($d = 3$). The material characteristics of the propagation medium will be represented by two measurable functions satisfying the following

Assumption 1.

The functions $\kappa : \Omega \rightarrow \mathbb{C}$ and $\mu : \Omega \rightarrow (0, +\infty)$ satisfy

- (i) $\Im\{\kappa(\mathbf{x})\} \geq 0, \Re\{\kappa(\mathbf{x})\} \geq 0 \forall \mathbf{x} \in \Omega$
- (ii) $\kappa_* := \max(1, \sup_{\mathbf{x} \in \Omega} |\kappa(\mathbf{x})|) < +\infty$
- (iii) $\sup_{\mathbf{x} \in \Omega} |\mu(\mathbf{x})| + |\mu^{-1}(\mathbf{x})| < +\infty$.

These are both general and physically reasonable assumptions. Condition (i) above implies in particular that $\Im\{\kappa^2(\mathbf{x})\} \geq 0$ for all $\mathbf{x} \in \Omega$. It means that the medium can only absorb or propagate energy. In addition, we consider source terms $f \in L^2(\Omega)$ and $g \in L^2(\partial\Omega)$. The boundary value problem under consideration will be

$$\begin{cases} \text{Find } u \in H^1(\Omega) \text{ such that} \\ \operatorname{div}(\mu \nabla u) + \kappa^2 u = -f \quad \text{in } \Omega, \\ \mu \partial_{\mathbf{n}} u = g \quad \text{on } \partial\Omega. \end{cases} \quad (1)$$

where \mathbf{n} refers to the outward unit normal vector to $\partial\Omega$. As usual, for any domain $\omega \subset \mathbb{R}^d$, the space $H^1(\omega)$ refers to those $v \in L^2(\omega)$ such that $\nabla v \in L^2(\omega)$. It will be equipped with the frequency dependent norm

$$\|v\|_{H^1(\omega)}^2 := \|\nabla v\|_{L^2(\omega)}^2 + \kappa_*^2 \|v\|_{L^2(\omega)}^2.$$

As usual, Problem (1) can be put in variational form: Find $u \in H^1(\Omega)$ such that $a(u, v) = \ell(v) \forall v \in H^1(\Omega)$ where

$$\begin{aligned} a(u, v) &:= \int_{\Omega} \mu \nabla u \cdot \nabla \bar{v} - \kappa^2 u \bar{v} \, d\mathbf{x} \\ \ell(v) &:= \int_{\Omega} f \bar{v} \, d\mathbf{x} + \int_{\partial\Omega} g \bar{v} \, d\sigma \end{aligned} \quad (2)$$

1.2 Discrete formulation

We are interested in the numerical solution to this problem by means of a classical nodal finite element scheme. We consider a regular triangulation $\mathcal{T}_h(\Omega)$ of the computational domain $\bar{\Omega} = \cup_{\tau \in \mathcal{T}_h(\Omega)} \bar{\tau}$. Shape regularity of this mesh is *not* needed for the subsequent analysis. We denote $V_h(\Omega) \subset H^1(\Omega)$ a space of \mathbb{P}_k -Lagrange finite element functions constructed on $\mathcal{T}_h(\Omega)$,

$$\begin{aligned} V_h(\Omega) &:= \{v \in \mathcal{C}^0(\bar{\Omega}) : v|_{\tau} \in \mathbb{P}_k(\bar{\tau}) \forall \tau \in \mathcal{T}_h(\Omega)\} \\ \text{where } \mathbb{P}_k(\bar{\tau}) &:= \{\text{polynomials of order } \leq k \text{ on } \bar{\tau}\} \end{aligned}$$

If $\omega \subset \Omega$ is any open subset that is resolved by the triangulation i.e. $\bar{\omega} = \cup_{\tau \in \mathcal{T}_h(\omega)} \bar{\tau}$, where $\mathcal{T}_h(\omega) \subset \mathcal{T}_h(\Omega)$, then we denote $V_h(\omega) := \{\varphi|_{\omega}, \varphi \in V_h(\Omega)\}$ and also consider finite element spaces on boundaries $V_h(\partial\omega) := \{\varphi|_{\partial\omega}, \varphi \in V_h(\Omega)\}$. We will focus on the discrete variational formulation

$$\begin{aligned} &\text{Find } u_h \in V_h(\Omega) \text{ such that} \\ &a(u_h, v_h) = \ell(v_h) \quad \forall v_h \in V_h(\Omega). \end{aligned} \tag{3}$$

Devising domain decomposition algorithms to solve this discrete problem is the main goal of the present article. This is why we need to assume that it admits a unique solution, which is equivalent to assuming that the corresponding finite element matrix is invertible.

Assumption 2.

$$\alpha_h := \inf_{u \in V_h(\Omega) \setminus \{0\}} \sup_{v \in V_h(\Omega) \setminus \{0\}} \frac{|a(u, v)|}{\|u\|_{H^1(\Omega)} \|v\|_{H^1(\Omega)}} > 0. \tag{4}$$

Although in practice this constant is uniformly bounded from below $\liminf_{h \rightarrow 0} \alpha_h > 0$, this uniform bound is not required for our analysis.

1.3 Notation conventions

Here we fix a few notation conventions that will be used all through the manuscript. Since the analysis presented here is fully discrete, we will only deal with finite dimensional function spaces. If V is any vector space equipped with the norm $\|\cdot\|_V$, then V^* will refer to its topological dual i.e. the space of linear functionals $p : V \rightarrow \mathbb{C}$ continuous in the norm $\|\cdot\|_V$, equipped with the norm

$$\|p\|_{V^*} = \sup_{v \in V \setminus \{0\}} |\langle p, v \rangle| / \|v\|_V, \tag{5}$$

the duality pairing between two dual spaces being systematically denoted $\langle \cdot, \cdot \rangle$. When such pairing brackets are used, it shall be clear from the context which duality pair of spaces (V, V^*) is considered. For $v \in V$ and $p \in V^*$ we shall write indistinctly $\langle p, v \rangle = \langle v, p \rangle$. We emphasize that duality pairings involve no conjugation operation. The polar set of any subspace $W \subset V$ will be defined by

$$W^\circ := \{\varphi \in V^* : \langle \varphi, v \rangle = 0 \forall v \in W\}.$$

This polar space inherits the norm (5) from V^* . In addition if $L : V_1 \rightarrow V_2$ is any continuous linear map between two vector spaces V_1, V_2 , its adjoint will then be a map $L^* : V_2^* \rightarrow V_1^*$ defined by the identity

$$\langle L^*(u), \bar{v} \rangle = \langle u, \overline{L(v)} \rangle \quad \forall u \in V_1, \quad \forall v \in V_2.$$

We shall systematically consider finite dimensional vector spaces stemming from a finite element discretization of function spaces. The linear operators $L : V_1 \rightarrow V_2$ will simply be finite element matrices but, very much like [3, Chap.5], we will refer to them as "operators" so as to keep track of our abstract setting where the choice of appropriate norms matters.

2 Geometric partitioning

We shall study a substructuring domain decomposition strategy for the solution to Problem (3), which leads to introducing a non-overlapping subdomain partition of the computational domain.

$$\begin{aligned}\bar{\Omega} &= \cup_{j=1}^J \bar{\Omega}_j, \quad \text{with } \Omega_j \cap \Omega_k = \emptyset \quad \text{for } j \neq k \\ \Sigma &:= \cup_{j=1}^J \Gamma_j, \quad \text{where } \Gamma_j := \partial\Omega_j,\end{aligned}\tag{6}$$

where each $\Omega_j \subset \Omega$ is itself a polyhedral domain that is exactly resolved by the triangulation. We do not make any further assumption regarding the subdomain partitioning. As opposed to [36, §2.5.2] or [38, §4.2], we do not formulate any regularity assumption on the subdomains. Our geometrical framework covers in particular the possibility of so-called cross-points i.e. points where either three subdomains abut or an interface separating two subdomains touches the external boundary of the computational domain. The treatment of cross-points has been the subject of several recent contributions, see e.g. [33, 21, 17, 8, 6]. In accordance with the notations of the previous section, we have

$$V_h(\Sigma) := \{v|_{\Sigma} : v \in V_h(\Omega)\}$$

which is here a space of (single valued) finite element functions defined over the skeleton that is a surface with multiple branches i.e. the union of all interfaces. Next we introduce continuous and discrete function spaces naturally associated to the multi-domain setting

$$\begin{aligned}\mathbb{H}^1(\Omega) &:= \mathbb{H}^1(\Omega_1) \times \cdots \times \mathbb{H}^1(\Omega_J), \\ \mathbb{V}_h(\Omega) &:= V_h(\Omega_1) \times \cdots \times V_h(\Omega_J) \subset \mathbb{H}^1(\Omega), \\ \mathbb{X}_h(\Omega) &:= \{(v|_{\Omega_1}, \dots, v|_{\Omega_J}) \in \mathbb{V}_h(\Omega) : v \in V_h(\Omega)\}.\end{aligned}\tag{7}$$

Since they are cartesian products, these spaces are made of tuples of (volume based) functions. The "broken space" $\mathbb{V}_h(\Omega)$ is naturally identified with those functions that are piecewise \mathbb{P}_k -Lagrange in each subdomain whereas, due to the matching conditions, the space $\mathbb{X}_h(\Omega)$ is naturally identified with $V_h(\Omega)$ i.e. those functions that are globally \mathbb{P}_k -Lagrange in the whole computational domain, including through interfaces $\Gamma_j \cap \Gamma_k$. These spaces will be equipped with the norm

$$\|u\|_{\mathbb{H}^1(\Omega)}^2 := \|u_1\|_{\mathbb{H}^1(\Omega_1)}^2 + \cdots + \|u_J\|_{\mathbb{H}^1(\Omega_J)}^2\tag{8}$$

for $u = (u_1, \dots, u_J)$. Since we are interested in domain decomposition where behaviour of functions at interfaces play a crucial role, we need to consider spaces consisting in tuples of trace functions. These will be called Dirichlet multi-trace (resp. Dirichlet single-trace) space

$$\begin{aligned}\mathbb{V}_h(\Sigma) &:= V_h(\Gamma_1) \times \cdots \times V_h(\Gamma_J) \\ \mathbb{X}_h(\Sigma) &:= \{(p|_{\Gamma_1}, \dots, p|_{\Gamma_J}) \in \mathbb{V}_h(\Sigma) : p \in V_h(\Sigma)\}.\end{aligned}\tag{9}$$

The elements of $\mathbb{X}_h(\Sigma)$ can also be characterized as those tuples $\mathbf{p} = (p_1, \dots, p_J) \in \mathbb{V}_h(\Sigma)$ that match across interfaces $p_j = p_k$ on $\Gamma_j \cap \Gamma_k$. The spaces (9) can be obtained by taking interior traces of functions belonging to $V_h(\Omega)$ resp. $\mathbb{X}_h(\Omega)$. This motivates the introduction of the trace map B defined as follows

$$\begin{aligned}B : V_h(\Omega) &\rightarrow \mathbb{V}_h(\Sigma) \\ B(v) &:= (v_1|_{\Gamma_1}, \dots, v_J|_{\Gamma_J})\end{aligned}\tag{10}$$

for $v = (v_1, \dots, v_J) \in \mathbb{V}_h(\Omega)$. This trace operator (10) surjectively maps $\mathbb{V}_h(\Omega)$ onto $\mathbb{V}_h(\Sigma)$, and it is also surjective from $\mathbb{X}_h(\Omega)$ onto $\mathbb{X}_h(\Sigma)$. We emphasize that the boundary trace map (10) is subdomain-wise block-diagonal. Since we are in a finite dimensional context, according to e.g [37, Thm. 4.7 & 4.12], we have $\text{Range}(\mathbf{B}^*) = \text{Ker}(\mathbf{B})^\circ := \{\phi \in \mathbb{V}_h(\Omega)^* : \langle \phi, v \rangle = 0 \forall v \in \text{Ker}(\mathbf{B})\}$, which can be rephrased in more concrete terms as follows.

Lemma 2.1.

Consider $\phi \in \mathbb{V}_h(\Omega)^$ satisfying $\langle \phi, u \rangle = 0$ for all $u \in \mathbb{V}_h(\Omega)$ such that $\mathbf{B}(u) = 0$. Then there exists $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ satisfying $\langle \phi, v \rangle = \langle \mathbf{p}, \mathbf{B}(v) \rangle \forall v \in \mathbb{V}_h(\Omega)$.*

Observe that $\mathbb{X}_h(\Omega) = \{u \in \mathbb{V}_h(\Omega) : \mathbf{B}(u) \in \mathbb{X}_h(\Sigma)\}$. As a consequence, if $u \in \mathbb{V}_h(\Omega)$ and $\mathbf{B}(u) = 0$ then $u \in \mathbb{X}_h(\Omega)$. In other words $\text{Ker}(\mathbf{B}) \subset \mathbb{X}_h(\Omega)$. The single-trace space $\mathbb{X}_h(\Sigma)$ can be parameterized by means of the restriction operator

$$\begin{aligned} \mathbf{R} : \mathbb{V}_h(\Sigma) &\rightarrow \mathbb{X}_h(\Sigma) \subset \mathbb{V}_h(\Sigma) \\ \mathbf{R}(p) &:= (p|_{\Gamma_1}, \dots, p|_{\Gamma_J}) \quad \text{where} \\ \mathbb{V}_h(\Sigma) &:= \{v|_{\Sigma} : v \in \mathbb{V}_h(\Omega)\} \end{aligned} \tag{11}$$

The space $\mathbb{V}_h(\Sigma)^*$ shall be referred to as the Neumann multi-trace space, while the following polar set will be called Neumann single-trace space

$$\mathbb{X}_h(\Sigma)^\circ := \{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* : \langle \mathbf{p}, \mathbf{v} \rangle = 0 \forall \mathbf{v} \in \mathbb{X}_h(\Sigma)\}. \tag{12}$$

This later space yields a variational characterization of $\mathbb{X}_h(\Sigma)$ through a polarity identity i.e. for any $\mathbf{v} \in \mathbb{V}_h(\Sigma)$ we have $\mathbf{v} \in \mathbb{X}_h(\Sigma) \iff \langle \mathbf{p}, \mathbf{v} \rangle = 0 \forall \mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$, which is the discrete counterpart of a polarity result frequently used in Multi-Trace theory [7, Prop.2.1]. Since $\mathbb{X}_h(\Sigma)$ is the range of the operator \mathbf{R} , this polarity identity also rewrites

$$\begin{aligned} \mathbb{X}_h(\Sigma) &= \text{Range}(\mathbf{R}), \\ \mathbb{X}_h(\Sigma)^\circ &= \text{Ker}(\mathbf{R}^*). \end{aligned} \tag{13}$$

3 Reformulation of transmission conditions

Transmission conditions are a crucial ingredient of any domain decomposition strategy. As a consequence, we pay a special attention to the matching conditions at interfaces between subdomains. We need to introduce a so-called "impedance" operator \mathbf{T} (following the terminology of [5]) that shall play a central role in the subsequent analysis. All that needs to be assumed concerning this operator is the following.

Assumption 3.

The linear operator $\mathbf{T} : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^$ satisfies $\Re\{\langle \mathbf{T}(\mathbf{p}), \overline{\mathbf{p}} \rangle\} > 0 \forall \mathbf{p} \in \mathbb{V}_h(\Sigma) \setminus \{0\}$.*

We underline that, concerning the impedance operator \mathbf{T} , no other property than Assumption 3 will be needed in the subsequent analysis. This assumption covers the possibility that $\mathbf{T} \neq \mathbf{T}^*$, which is new compared to [8].

In the context of waves, many domain decomposition strategies belonging to the family of Optimized Schwarz Methods have considered particular instances of non-self-adjoint impedances. With Assumption 3, the present article extends the theory of [8], as it now covers the situations considered in [11, Eq.(2.13)], in [17, Prop.4] and in [4].

Assumption 3 can be rephrased by stating that the symmetric part $T_s := (T + T^*)/2$ induces a scalar product over $\mathbb{V}_h(\Sigma)$. As a consequence the operator $T_s^{-1} = 2(T + T^*)^{-1}$ induces a scalar product over the dual space $\mathbb{V}_h(\Sigma)^*$. We define

$$\begin{aligned}\|\mathbf{v}\|_{T_s}^2 &:= \langle T_s(\mathbf{v}), \overline{\mathbf{v}} \rangle \\ \|\mathbf{p}\|_{T_s^{-1}}^2 &:= \langle T_s^{-1}(\mathbf{p}), \overline{\mathbf{p}} \rangle \\ \text{where } T_s &:= (T + T^*)/2.\end{aligned}\tag{14}$$

With these definitions, for any $\mathbf{v} \in \mathbb{V}_h(\Sigma)$, $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$, we have $|\langle \mathbf{v}, \mathbf{p} \rangle| \leq \|\mathbf{v}\|_{T_s} \|\mathbf{p}\|_{T_s^{-1}}$ and $\|T_s(\mathbf{v})\|_{T_s^{-1}} = \|\mathbf{v}\|_{T_s}$ and $\|T_s^{-1}(\mathbf{p})\|_{T_s} = \|\mathbf{p}\|_{T_s^{-1}}$. The theory that we present here stems from a new treatment of transmission conditions that relies on a proper characterization of Dirichlet (resp. Neumann) single-trace space $\mathbb{X}_h(\Sigma)$ (resp. $\mathbb{X}_h(\Sigma)^\circ$). We need a first lemma.

Lemma 3.1.

For any impedance operator $T : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^$ satisfying Assumption 3, we have the following direct sums*

$$\begin{aligned}\mathbb{V}_h(\Sigma) &= \mathbb{X}_h(\Sigma) \oplus T^{-1}(\mathbb{X}_h(\Sigma)^\circ), \\ \mathbb{V}_h(\Sigma)^* &= \mathbb{X}_h(\Sigma)^\circ \oplus T(\mathbb{X}_h(\Sigma)).\end{aligned}$$

Moreover, if $T = T^ = T_s$ then these direct sums are respectively T_s -orthogonal and T_s^{-1} -orthogonal.*

Proof:

We prove the result for the second direct sum only, since the proof runs completely parallel for the first direct sum. First pick $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ \cap T(\mathbb{X}_h(\Sigma))$ so that there exists $\mathbf{u} \in \mathbb{X}_h(\Sigma)$ with $\mathbf{p} = T(\mathbf{u})$. Then we have

$$\begin{aligned}0 &= 2\Re\{\langle \mathbf{p}, \overline{\mathbf{u}} \rangle\} = 2\Re\{\langle T(\mathbf{u}), \overline{\mathbf{u}} \rangle\} = \langle T(\mathbf{u}), \overline{\mathbf{u}} \rangle + \langle \mathbf{u}, \overline{T(\mathbf{u})} \rangle \\ &= \langle (T + T^*)\mathbf{u}, \overline{\mathbf{u}} \rangle = 2\|\mathbf{u}\|_{T_s}^2 \implies \mathbf{u} = 0.\end{aligned}$$

From this we conclude that $\mathbf{p} = T(\mathbf{u}) = 0$ hence $\mathbb{X}_h(\Sigma)^\circ \cap T_s(\mathbb{X}_h(\Sigma)) = \{0\}$ since \mathbf{p} was chosen arbitrarily.

Next take an arbitrary $\mathbf{q} \in \mathbb{V}_h(\Sigma)^*$ and define \mathbf{v} as the only element of $\mathbb{X}_h(\Sigma)$ satisfying $\langle T(\mathbf{v}), \mathbf{w} \rangle = \langle \mathbf{q}, \mathbf{w} \rangle \forall \mathbf{w} \in \mathbb{X}_h(\Sigma)$. This variational problem admits a unique solution due to the coercivity of T from Assumption 3. By construction we have $\langle \mathbf{q} - T(\mathbf{v}), \mathbf{w} \rangle = 0 \forall \mathbf{w} \in \mathbb{X}_h(\Sigma)$ which means that $\mathbf{q}' = \mathbf{q} - T(\mathbf{v}) \in \mathbb{X}_h(\Sigma)^\circ$. We have just established that $\mathbf{q} = \mathbf{q}' + T(\mathbf{v}) \in \mathbb{X}_h(\Sigma)^\circ + T(\mathbb{X}_h(\Sigma))$, hence finally $\mathbb{V}_h(\Sigma)^* = \mathbb{X}_h(\Sigma)^\circ \oplus T(\mathbb{X}_h(\Sigma))$.

In the case where $T = T^* = T_s$, take arbitrary $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$, $\mathbf{q} \in T_s(\mathbb{X}_h(\Sigma))$. We have $\mathbf{q} = T_s(\mathbf{v})$ for some $\mathbf{v} \in \mathbb{X}_h(\Sigma)$, and thus $\langle T_s^{-1}(\mathbf{q}), \mathbf{p} \rangle = \langle \mathbf{v}, \mathbf{p} \rangle = 0$ by the very definition (12). We conclude that $\mathbb{X}_h(\Sigma)^\circ$ and $T_s(\mathbb{X}_h(\Sigma))$ are T_s^{-1} -orthogonal to each other. \square

The spaces $\mathbb{X}_h(\Sigma)$ and $\mathbb{X}_h(\Sigma)^\circ$ will play an important role in the sequel, and we need a convenient way to characterize them. This is our motivation for introducing an oblique (i.e. non-self-adjoint) counterpart of the exchange operator considered in [6, Cor.5.1] and [8, Lem.6.2].

Lemma 3.2.

Under Assumption 3, the operator $\Pi := (\mathsf{T} + \mathsf{T}^*)\mathsf{R}(\mathsf{R}^*\mathsf{T}^*\mathsf{R})^{-1}\mathsf{R}^* - \text{Id}$ with mapping property $\Pi : \mathbb{V}_h(\Sigma)^* \rightarrow \mathbb{V}_h(\Sigma)^*$ is an isometry in the norm (14) i.e. for all $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ we have

$$\|\Pi(\mathbf{p})\|_{\mathbb{T}_s^{-1}} = \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}.$$

Proof:

Pick an arbitrary $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ and set $\mathbf{u} := \mathsf{R}(\mathsf{R}^*\mathsf{T}^*\mathsf{R})^{-1}\mathsf{R}^*\mathbf{p}$. We have $\mathbf{u} \in \text{Range}(\mathsf{R}) = \mathbb{X}_h(\Sigma)$ according to (13) and $\mathsf{R}^*\mathsf{T}^*(\mathbf{u}) = \mathsf{R}^*\mathbf{p}$ hence, pairing with an arbitrary $w \in \mathbb{V}_h(\Sigma)$, we obtain $\langle \mathsf{T}^*(\mathbf{u}), \mathsf{R}(w) \rangle = \langle \mathbf{p}, \mathsf{R}(w) \rangle$. Since $\mathsf{R} : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{X}_h(\Sigma)$ is surjective and w is chosen arbitrarily in $\mathbb{V}_h(\Sigma)$, we conclude that \mathbf{u} is characterized as the unique solution to the following variational problem

$$\begin{aligned} \mathbf{u} \in \mathbb{X}_h(\Sigma) \quad \text{and} \\ \langle \mathsf{T}^*(\mathbf{u}), \mathbf{v} \rangle = \langle \mathbf{p}, \mathbf{v} \rangle \quad \forall \mathbf{v} \in \mathbb{X}_h(\Sigma). \end{aligned} \tag{15}$$

In particular, taking $\mathbf{v} = \overline{\mathbf{u}}$, we obtain

$$\begin{aligned} 2\Re\{\langle \mathbf{p}, \overline{\mathbf{u}} \rangle\} &= 2\Re\{\langle \mathsf{T}^*(\mathbf{u}), \overline{\mathbf{u}} \rangle\} = \langle \mathsf{T}^*(\mathbf{u}), \overline{\mathbf{u}} \rangle + \overline{\langle \overline{\mathbf{u}}, \mathsf{T}^*(\mathbf{u}) \rangle} \\ &= \langle \mathsf{T}^*(\mathbf{u}), \overline{\mathbf{u}} \rangle + \langle \mathbf{u}, \overline{\mathsf{T}^*(\mathbf{u})} \rangle = 2\langle \mathsf{T}_s(\mathbf{u}), \overline{\mathbf{u}} \rangle \\ &= 2\|\mathbf{u}\|_{\mathbb{T}_s}^2 = 2\|\mathsf{T}_s(\mathbf{u})\|_{\mathbb{T}_s^{-1}}^2. \end{aligned} \tag{16}$$

Next, observe that $\Pi(\mathbf{p}) = 2\mathsf{T}_s(\mathbf{u}) - \mathbf{p}$ by construction, which leads to

$$\begin{aligned} \|\Pi(\mathbf{p})\|_{\mathbb{T}_s^{-1}}^2 &= \|2\mathsf{T}_s(\mathbf{u}) - \mathbf{p}\|_{\mathbb{T}_s^{-1}}^2 \\ &= \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}^2 - 4\Re\{\langle \mathbf{p}, \overline{\mathbf{u}} \rangle\} + 4\|\mathsf{T}_s(\mathbf{u})\|_{\mathbb{T}_s^{-1}}^2 = \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}^2. \end{aligned}$$

□

Interestingly, since Π is a \mathbb{T}_s^{-1} -isometry and $\mathbb{V}_h(\Sigma)^*$ is finite dimensional, we deduce an expression of the inverse $\Pi^{-1} = (\mathsf{T} + \mathsf{T}^*)\mathsf{R}(\mathsf{R}^*\mathsf{T}\mathsf{R})^{-1}\mathsf{R}^* - \text{Id}$. This formula shall not be of much use in the present contribution though. In the special case of a self-adjoint impedance, this exchange operator becomes an orthogonal symmetry.

Lemma 3.3.

Let Assumption 3 hold and suppose further that $\mathsf{T} = \mathsf{T}^*$. Then $\Pi = 2\mathsf{T}\mathsf{R}(\mathsf{R}^*\mathsf{T}\mathsf{R})^{-1}\mathsf{R}^* - \text{Id}$ defined in Lemma 3.2 satisfies in addition the following properties

- i) $(\text{Id} - \Pi)/2$ is the \mathbb{T}^{-1} -orthogonal projection onto $\mathbb{X}_h(\Sigma)^\circ$,
- ii) $(\text{Id} + \Pi^*)/2$ is the \mathbb{T} -orthogonal projection onto $\mathbb{X}_h(\Sigma)$,
- iii) $\Pi^2 = \text{Id}$,
- iv) $\Pi^*\mathsf{R} = \mathsf{R}$.

Proof:

In the case $\mathsf{T} = \mathsf{T}^*$ we have $\mathsf{T} = \mathsf{T}_s$. The identities *iii*) and *iv*) follow from direct calculus. Then it is clear that $(\text{Id} \pm \Pi)/2$ and $(\text{Id} \pm \Pi^*)/2$ are projectors. The \mathbb{T}^{-1} -orthogonality of $(\text{Id} - \Pi)/2$ and the \mathbb{T} -orthogonality of $(\text{Id} + \Pi^*)/2$ are equivalent to the \mathbb{T}^{-1} -orthogonality of

$(\text{Id} + \Pi)/2 = \text{TR}(\text{R}^*\text{TR})^{-1}\text{R}^*$. The latter is proved by taking $\mathbf{p}, \mathbf{q} \in \mathbb{V}_h(\Sigma)^*$ arbitrary, and observing that

$$\begin{aligned} \langle \text{T}^{-1}(\text{Id} + \Pi)\mathbf{p}, \bar{\mathbf{q}} \rangle &= \langle (\text{R}^*\text{TR})^{-1}\text{R}^*\mathbf{p}, \text{R}^*\bar{\mathbf{q}} \rangle \\ &= \langle \mathbf{p}, \text{R}(\text{R}^*\text{TR})^{-1}\text{R}^*\bar{\mathbf{q}} \rangle = \langle \text{T}^{-1}\mathbf{p}, \overline{(\text{Id} + \Pi)\mathbf{q}} \rangle \end{aligned}$$

□

The previous lemma is consistent with the theory proposed in our previous contributions that only considered symmetric positive definite impedances, see Corollary 5.1 in [6] and Lemma 6.2 in [8]. Coming back to the general case of a priori non-self-adjoint impedance, let us show how this exchange operator can serve for the effective characterization of $\mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$.

Lemma 3.4.

Let Assumption 3 hold and define $\mathcal{X}_h(\Sigma) := \mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$ and $\Pi := 2\text{T}_s\text{R}(\text{R}^*\text{T}^*\text{R})^{-1}\text{R}^* - \text{Id}$ as in Lemma 3.2. Then for any pair $(\mathbf{u}, \mathbf{p}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$ we have the equivalence

$$(\mathbf{u}, \mathbf{p}) \in \mathcal{X}_h(\Sigma) \iff -\mathbf{p} + i\text{T}(\mathbf{u}) = \Pi(\mathbf{p} + i\text{T}^*(\mathbf{u})). \quad (17)$$

Proof:

Let us first investigate the action of Π on certain well chosen input arguments. If $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$ then $\text{R}^*(\mathbf{p}) = 0$ according to (13) hence $\Pi(\mathbf{p}) = -\mathbf{p}$. On the other hand, if $\mathbf{u} \in \mathbb{X}_h(\Sigma)$ then $\mathbf{u} = \text{R}(v)$ for some $v \in \mathbb{V}_h(\Sigma)$, and this yields

$$\begin{aligned} \Pi \cdot \text{T}^*(\mathbf{u}) &= (\text{T} + \text{T}^*)\text{R}(\text{R}^*\text{T}^*\text{R})^{-1}(\text{R}^*\text{T}^*\text{R})v - \text{T}^*(\mathbf{u}) \\ &= (\text{T} + \text{T}^*)\text{R}v - \text{T}^*(\mathbf{u}) = \text{T}(\mathbf{u}). \end{aligned} \quad (18)$$

The previous observations clearly imply that any pair $(\mathbf{u}, \mathbf{p}) \in \mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$ satisfies $-\mathbf{p} + i\text{T}(\mathbf{u}) = \Pi(\mathbf{p} + i\text{T}^*(\mathbf{u}))$. Reciprocally pick an arbitrary pair $(\mathbf{u}, \mathbf{p}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$ satisfying $-\mathbf{p} + i\text{T}(\mathbf{u}) = \Pi(\mathbf{p} + i\text{T}^*(\mathbf{u}))$. Plugging the expression of Π into this identity implies

$$\begin{aligned} -\mathbf{p} + i\text{T}(\mathbf{u}) &= (\text{T} + \text{T}^*)\text{R}(\text{R}^*\text{T}^*\text{R})^{-1}\text{R}^*(\mathbf{p} + i\text{T}^*(\mathbf{u})) - \mathbf{p} - i\text{T}^*(\mathbf{u}) \\ \iff i(\text{T} + \text{T}^*)\mathbf{u} &= (\text{T} + \text{T}^*)\text{R}(\text{R}^*\text{T}^*\text{R})^{-1}\text{R}^*(\mathbf{p} + i\text{T}^*(\mathbf{u})) \end{aligned} \quad (19)$$

Next multiply the last identity above on the left by $\text{R}^*\text{T}^*(\text{T} + \text{T}^*)^{-1}$ which leads to $i\text{R}^*\text{T}^*(\mathbf{u}) = i\text{R}^*\text{T}^*(\mathbf{u}) + \text{R}^*\mathbf{p}$ and finally $\text{R}^*\mathbf{p} = 0$ which is equivalent to $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$ according to (13). Coming back to the second identity of (19), and multiplying by $-i(\text{T} + \text{T}^*)^{-1}$ and taking account of (13), we obtain $\mathbf{u} = \text{R}(\text{R}^*\text{T}^*\text{R})^{-1}\text{R}^*\text{T}^*(\mathbf{u}) \in \text{Range}(\text{R}) = \mathbb{X}_h(\Sigma)$. This finishes the proof. □

4 Reformulation of the scattering problem

The equivalence (17) is a new way to impose transmission conditions across interfaces between subdomains. We now make use of this new ingredient to reformulate the wave propagation problem we are interested in. Let us introduce an operator $\mathbf{A} : \mathbb{V}_h(\Omega) \rightarrow \mathbb{V}_h(\Omega)^*$ and a source term $\mathbf{f} \in \mathbb{V}_h(\Omega)^*$ associated to the domain decomposed problem and defined by

$$\begin{aligned} \langle \mathbf{A}\mathbf{u}, \mathbf{v} \rangle &:= \sum_{j=1 \dots J} \int_{\Omega_j} \mu \nabla u_j \cdot \nabla v_j - \kappa^2 u_j v_j \, d\mathbf{x} \\ \langle \mathbf{f}, \mathbf{v} \rangle &:= \sum_{j=1 \dots J} \int_{\Omega_j} f v_j \, d\mathbf{x} + \int_{\partial\Omega_j \cap \partial\Omega} g v_j \, d\sigma \end{aligned} \quad (20)$$

for any $\mathbf{u} = (u_1, \dots, u_J), \mathbf{v} = (v_1, \dots, v_J)$ in $\mathbb{V}_h(\Omega)$. These are nothing but a finite element matrix and vector. A few remarks are in order concerning the operator A . First of all it admits a block diagonal form with respect to the subdomain decomposition. Taking account of Assumption 1, the imaginary part of A is signed

$$\Im\{\langle A\mathbf{u}, \bar{\mathbf{u}} \rangle\} \leq 0 \quad \forall \mathbf{u} \in \mathbb{V}_h(\Omega). \quad (21)$$

Besides, a comparison of (20) with (2)-(7) shows that, if $\mathbf{u} = (u|_{\Omega_1}, \dots, u|_{\Omega_J})$ and $\mathbf{v} = (v|_{\Omega_1}, \dots, v|_{\Omega_J})$ for some $u, v \in \mathbb{V}_h(\Omega)$, then $\mathbf{u}, \mathbf{v} \in \mathbb{X}_h(\Omega)$ according to (7) and, in this case, we have $\langle A\mathbf{u}, \bar{\mathbf{v}} \rangle = a(u, v)$ and $\langle \mathbf{f}, \bar{\mathbf{v}} \rangle = \ell(v)$. We introduce the continuity modulus $\|a\|$ of A , and also re-express the inf-sup constant α_h from Assumption 2 as follows

$$\begin{aligned} \alpha_h &= \inf_{\mathbf{u} \in \mathbb{X}_h(\Omega) \setminus \{0\}} \sup_{\mathbf{v} \in \mathbb{X}_h(\Omega) \setminus \{0\}} \frac{|\langle A(\mathbf{u}), \mathbf{v} \rangle|}{\|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbb{H}^1(\Omega)}} > 0 \\ \|a\| &:= \sup_{\mathbf{u} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|\langle A(\mathbf{u}), \mathbf{v} \rangle|}{\|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbb{H}^1(\Omega)}} \end{aligned} \quad (22)$$

The constant $\|a\|$ is simply the continuity modulus of the sesquilinear form $a(\cdot, \cdot)$ over the space of piecewise \mathbb{P}_k -Lagrange functions. We will now re-write Problem (3) in several equivalent forms more prone to domain decomposition. First we use operator A so as to put it in matrix form.

Lemma 4.1.

Assume that $u \in \mathbb{V}_h(\Omega)$ is solution to (3). Then, setting $\mathbf{u} = (u|_{\Omega_1}, \dots, u|_{\Omega_J}) \in \mathbb{V}_h(\Omega)$, there exists $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ such that

$$\begin{aligned} (\mathbf{u}, \mathbf{p}) &\in \mathbb{X}_h(\Omega) \times \mathbb{X}_h(\Sigma)^\circ \text{ and} \\ A\mathbf{u} - B^*\mathbf{p} &= \mathbf{f}. \end{aligned} \quad (23)$$

Reciprocally if the pair $(\mathbf{u}, \mathbf{p}) \in \mathbb{V}_h(\Omega) \times \mathbb{V}_h(\Sigma)^*$ solves (23), with $\mathbf{u} = (u_1, \dots, u_J)$, then the function defined by $u(\mathbf{x}) = u_1(\mathbf{x})1_{\Omega_1}(\mathbf{x}) + \dots + u_J(\mathbf{x})1_{\Omega_J}(\mathbf{x})$ belongs to $\mathbb{V}_h(\Omega)$ and solves (3).

Proof:

Assume that $u \in \mathbb{V}_h(\Omega)$ is solution to (3). By definition we have $\mathbf{u} \in \mathbb{X}_h(\Omega)$ where $\mathbf{u} = (u|_{\Omega_1}, \dots, u|_{\Omega_J})$ and (3) rewrites

$$\begin{aligned} \mathbf{u} &\in \mathbb{X}_h(\Omega) \text{ such that} \\ \langle A(\mathbf{u}), \mathbf{v} \rangle &= \langle \mathbf{f}, \mathbf{v} \rangle \quad \forall \mathbf{v} \in \mathbb{X}_h(\Omega) \end{aligned} \quad (24)$$

Since $\mathbf{v} \in \mathbb{X}_h(\Omega)$ whenever $\mathbf{v} \in \mathbb{V}_h(\Omega)$ satisfies $B(\mathbf{v}) = 0$, we can apply Lemma 2.1 to $A(\mathbf{u}) - \mathbf{f} \in \mathbb{V}_h(\Omega)^*$ which yields the existence of $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ such that $A\mathbf{u} - \mathbf{f} = B^*\mathbf{p}$ which is the second line of (23). Then the second line of (24) rewrites $0 = \langle B^*\mathbf{p}, \mathbf{v} \rangle = \langle \mathbf{p}, B(\mathbf{v}) \rangle \quad \forall \mathbf{v} \in \mathbb{X}_h(\Omega)$. Since B maps $\mathbb{X}_h(\Omega)$ onto $\mathbb{X}_h(\Sigma)$, we conclude that $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$. As a consequence (23) holds.

Now assume that $(\mathbf{u}, \mathbf{p}) \in \mathbb{V}_h(\Omega) \times \mathbb{V}_h(\Sigma)^*$ solves (23), where we denote $\mathbf{u} = (u_1, \dots, u_J)$. Besides, since $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$, we have $\langle B^*\mathbf{p}, \mathbf{v} \rangle = \langle \mathbf{p}, B(\mathbf{v}) \rangle = 0$ for all $\mathbf{v} \in \mathbb{X}_h(\Omega)$ which rewrites as (24). Since (24) implies (3) with the function $u(\mathbf{x}) = u_1(\mathbf{x})1_{\Omega_1}(\mathbf{x}) + \dots + u_J(\mathbf{x})1_{\Omega_J}(\mathbf{x})$ belonging to $\mathbb{V}_h(\Omega)$, this concludes the proof. \square

The tuple of unknowns \mathbf{p} in Formulation (23) should be understood as Neumann fluxes of the volume solution across boundaries of subdomains, and the condition $\mathbf{p} \in \mathbb{X}_h(\Sigma)^\circ$ should be understood as the Neumann part of classical transmission conditions across interfaces.

Note that $\mathbf{u} \in \mathbb{X}_h(\Omega)$ if and only if $\mathbf{u} \in \mathbb{V}_h(\Omega)$ and $\mathbf{B}(\mathbf{u}) \in \mathbb{X}_h(\Sigma)$. The property $\mathbf{B}(\mathbf{u}) \in \mathbb{X}_h(\Sigma)$ should be interpreted as the Dirichlet part of classical transmission conditions across interfaces. Thanks to the previous remarks, we can transform further Formulation (23) by taking account of the characterization of $\mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$ stemming from Lemma 3.4. This directly yields the following reformulation.

Lemma 4.2.

Under Assumption 3, the pair (\mathbf{u}, \mathbf{p}) solves (23) if and only if it satisfies

$$\begin{aligned} (\mathbf{u}, \mathbf{p}) &\in \mathbb{V}_h(\Omega) \times \mathbb{V}_h(\Sigma)^* \text{ and} \\ A\mathbf{u} - \mathbf{B}^*\mathbf{p} &= \mathbf{f}, \\ \mathbf{p} - iT\mathbf{B}(\mathbf{u}) &= -\Pi(\mathbf{p} + iT^*\mathbf{B}(\mathbf{u})). \end{aligned} \tag{25}$$

The formulation above only involves the multi-trace space and its dual where contributions from subdomains are decorrelated from each other. Transmission conditions across interfaces come only into play through the exchange operator Π . Although we do not write it as an iterative algorithm, Formulation (25) above is entirely analogous to e.g. [17, Eq.(8)], [8, Eq.(24)], [5, Eq.(3)-(4)] or [10, Eq.(59)].

5 Scattering operator

The next step of our analysis consists in eliminating the volume unknowns in (25). However the map $A : \mathbb{V}_h(\Omega) \rightarrow \mathbb{V}_h(\Omega)^*$ may be not invertible, so we avoid using A^{-1} and re-arrange local subproblems. The next lemma establishes that local subproblems become invertible if we add absorbing impedance conditions.

Lemma 5.1.

*Under Assumptions 1, 2 and 3, the operator $A - i\mathbf{B}^*T\mathbf{B} : \mathbb{V}_h(\Omega) \rightarrow \mathbb{V}_h(\Omega)^*$ is systematically invertible which rewrites in inf-sup condition form as*

$$\beta_h := \inf_{\mathbf{u} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|(A - i\mathbf{B}^*T\mathbf{B})\mathbf{u}, \mathbf{v}\rangle|}{\|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{v}\|_{\mathbb{H}^1(\Omega)}} > 0. \tag{26}$$

Proof:

Assume that $\beta_h = 0$ so that there exists $\mathbf{u} \in \mathbb{V}_h(\Omega)$ such that $(A - i\mathbf{B}^*T\mathbf{B})\mathbf{u} = 0$. From this and (21), we conclude that $0 = -\Im m\{\langle A(\mathbf{u}), \bar{\mathbf{u}} \rangle\} + \Im m\{i\langle T\mathbf{B}(\mathbf{u}), \mathbf{B}(\bar{\mathbf{u}}) \rangle\} \geq \Re e\{\langle T\mathbf{B}(\mathbf{u}), \mathbf{B}(\bar{\mathbf{u}}) \rangle\}$ which implies $\mathbf{B}(\mathbf{u}) = 0$ according to Assumption 3. Since we have $\text{Ker}(\mathbf{B}) \subset \mathbb{X}_h(\Omega)$ according to (13), this shows that $\mathbf{u} \in \mathbb{X}_h(\Omega)$ and $A(\mathbf{u}) = 0$, which implies $\mathbf{u} = 0$ according to (22). This establishes that $\text{Ker}(A - i\mathbf{B}^*T\mathbf{B}) = \{0\}$ and finishes the proof. \square

Now we can introduce a so called scattering operator. The following result is inspired by [16, Lemma 6].

Lemma 5.2.

Under Assumptions 1, 2 and 3, the operator $S := \text{Id} + 2iT_s B(A - iB^*TB)^{-1}B^*$ is a T_s^{-1} -contraction and, for all $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$, satisfies the identity

$$\|S(\mathbf{p})\|_{T_s^{-1}}^2 + 4|\Im\{\langle A\mathbf{u}, \bar{\mathbf{u}} \rangle\}| = \|\mathbf{p}\|_{T_s^{-1}}^2 \quad (27)$$

where $\mathbf{u} = (A - iB^*TB)^{-1}B^*\mathbf{p}$.

Proof:

We note that $S(\mathbf{p}) = \mathbf{p} + 2iT_s B(\mathbf{u})$, and simply expand the left hand side of (27), taking account of the sign property provided by (21). This yields the following calculus

$$\begin{aligned} \|S(\mathbf{p})\|_{T_s^{-1}}^2 &= \|\mathbf{p} + 2iT_s B(\mathbf{u})\|_{T_s^{-1}}^2 \\ &= \|\mathbf{p}\|_{T_s^{-1}}^2 - 4\Re\{i\langle \mathbf{p}, B(\bar{\mathbf{u}}) \rangle\} + 4\|T_s B(\mathbf{u})\|_{T_s^{-1}}^2 \\ &= \|\mathbf{p}\|_{T_s^{-1}}^2 - 4\Re\{i\langle (A - iB^*TB)\mathbf{u}, \bar{\mathbf{u}} \rangle\} + 4\langle T_s B(\mathbf{u}), B(\bar{\mathbf{u}}) \rangle \\ &= \|\mathbf{p}\|_{T_s^{-1}}^2 + 4\Im\{\langle A(\mathbf{u}), \bar{\mathbf{u}} \rangle\} = \|\mathbf{p}\|_{T_s^{-1}}^2 - 4|\Im\{\langle A(\mathbf{u}), \bar{\mathbf{u}} \rangle\}| \end{aligned} \quad (28)$$

□

The scattering operator S is subdomain-wise block diagonal under the additional assumption that T is subdomain-wise block-diagonal, which is a reasonable and easy property to fulfill in practice.

Identity (27) should be interpreted as energy conservation. It is similar to [14, Lemma 4.1], [10, Lemma 1] or [17, Lemma 11]. The scattering operator S takes a Neuman multi-trace as input, solves in each subdomain the associated ingoing impedance problem, and returns the outgoing impedance trace as output. This interpretation of the scattering operator is made explicit in the next result.

Proposition 5.3.

Define $\mathcal{C}_h(\Sigma) := \{(\mathbf{v}, \mathbf{p}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* : \exists \mathbf{u} \in \mathbb{V}_h(\Omega), A\mathbf{u} = B^*\mathbf{p}, B\mathbf{u} = \mathbf{v}\}$ which will be called the space of discrete Cauchy data. Let Assumptions 1, 2 and 3 hold. Then for any pair $(\mathbf{v}, \mathbf{p}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$, we have the equivalence

$$(\mathbf{v}, \mathbf{p}) \in \mathcal{C}_h(\Sigma) \iff \mathbf{p} + iT^*\mathbf{v} = S(\mathbf{p} - iT\mathbf{v}). \quad (29)$$

Proof:

Take an arbitrary pair $(\mathbf{v}, \mathbf{p}) \in \mathcal{C}_h(\Sigma)$ and set $\mathbf{q} = \mathbf{p} - iT\mathbf{v}$. By definition of the Cauchy data space, there exists $\mathbf{u} \in \mathbb{V}_h(\Omega)$ such that $A\mathbf{u} = B^*\mathbf{p}$ and $B\mathbf{u} = \mathbf{v}$ which implies in particular that $(A - iB^*TB)\mathbf{u} = B^*\mathbf{q}$ hence $\mathbf{u} = (A - iB^*TB)^{-1}B^*\mathbf{q}$. Next the definition of the scattering operator given in Lemma (5.2) yields $S(\mathbf{q}) = \mathbf{p} - iT\mathbf{v} + 2iT_s B(\mathbf{u}) = \mathbf{p} - iT\mathbf{v} + iT\mathbf{v} + iT^*\mathbf{v} = \mathbf{p} + iT^*\mathbf{v}$ which rewrites as (29).

Reciprocally take any $(\mathbf{v}, \mathbf{p}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$ satisfying $\mathbf{p} + iT^*\mathbf{v} = S(\mathbf{p} - iT\mathbf{v})$. Define $\mathbf{u} \in \mathbb{V}_h(\Omega)$ by $\mathbf{u} = (A - iB^*TB)^{-1}B^*(\mathbf{p} - iT\mathbf{v})$ so that, using the definition of S and the invertibility of $T_s := (T + T^*)/2$, we have $\mathbf{p} + iT^*\mathbf{v} = S(\mathbf{p} - iT\mathbf{v}) = \mathbf{p} - iT\mathbf{v} + 2iT_s B\mathbf{u} \iff 2iT_s \mathbf{v} = 2iT_s B\mathbf{u} \iff \mathbf{v} = B\mathbf{u}$. Next coming back to the definition of \mathbf{u} we conclude that $A\mathbf{u} - iB^*T\mathbf{v} = B^*(\mathbf{p} - iT\mathbf{v})$, and finally $A\mathbf{u} = B^*\mathbf{p}$. This proves that the pair (\mathbf{v}, \mathbf{p}) belongs to $\mathcal{C}_h(\Sigma)$. □

We draw the attention of the reader on the similarities between (29) and (17). Both characterizations are expressed in terms of ingoing traces (i.e. traces of the form $\mathbf{p} - iT\mathbf{v}$) and outgoing traces (i.e. traces of the form $\mathbf{p} + iT^*\mathbf{v}$). Let us also point an interesting property satisfied by the elements of $\mathcal{C}_h(\Sigma)$.

Lemma 5.4.

Under Assumptions 1, 2 and 3, if $(\mathbf{v}, \mathbf{p}) \in \mathcal{C}_h(\Sigma)$ then $\Im m\{\langle \mathbf{p}, \bar{\mathbf{v}} \rangle\} \leq 0$ and we have the energy identities

$$\begin{aligned} \|\mathbf{v}\|_{T_s}^2 + \|\mathbf{p}\|_{T_s^{-1}}^2 &= \|\mathbf{p} + iT_s\mathbf{v}\|_{T_s^{-1}}^2 + 2|\Im m\{\langle \mathbf{p}, \bar{\mathbf{v}} \rangle\}| \\ &= \|\mathbf{p} - iT_s\mathbf{v}\|_{T_s^{-1}}^2 - 2|\Im m\{\langle \mathbf{p}, \bar{\mathbf{v}} \rangle\}|. \end{aligned}$$

Proof:

By the very definition of $\mathcal{C}_h(\Sigma)$ given in Proposition 5.3, there exists $\mathbf{u} \in \mathbb{V}_h(\Omega)$ such that $A(\mathbf{u}) = B^*(\mathbf{p})$ and $B(\mathbf{u}) = \mathbf{v}$, hence $\langle \mathbf{p}, \bar{\mathbf{v}} \rangle = \langle \mathbf{p}, B(\bar{\mathbf{u}}) \rangle = \langle B^*(\mathbf{p}), \bar{\mathbf{u}} \rangle = \langle A(\mathbf{u}), \bar{\mathbf{u}} \rangle$. With (21) we obtain $\Im m\{\langle \mathbf{p}, \bar{\mathbf{v}} \rangle\} = \Im m\{\langle A(\mathbf{u}), \bar{\mathbf{u}} \rangle\} \leq 0$ which is the first desired result. The energy identities directly follow from $\|\mathbf{p} \pm iT_s\mathbf{v}\|_{T_s^{-1}}^2 = \|\mathbf{p}\|_{T_s^{-1}}^2 \pm 2\Im m\{\langle \mathbf{p}, \bar{\mathbf{v}} \rangle\} + \|\mathbf{v}\|_{T_s}^2$. \square

6 Skeleton formulation

We will now use the scattering operator introduced in the previous section to rewrite equivalently our wave propagation problem as an equation posed on the skeleton of the subdomain partition. In spite of a possible minor difference due to a sign convention, the formulation derived below is similar to [8, Eq.(27)], [6, Eq.(7.2)], [10, Eq.(45)&(51)], [14, §3.3],[31, Chap.6], [17].

Lemma 6.1.

*Let Assumptions 1, 2 and 3 hold. Set $\mathbf{g} := -2i\Pi T_s B(A - iB^*TB)^{-1}\mathbf{f} \in \mathbb{V}_h(\Sigma)^*$. If the pair $(\mathbf{u}, \mathbf{p}) \in \mathbb{V}_h(\Omega) \times \mathbb{V}_h(\Sigma)^*$ solves (25) then the tuple of traces $\mathbf{q} = \mathbf{p} - iTB\mathbf{u}$ solves*

$$\begin{aligned} \mathbf{q} &\in \mathbb{V}_h(\Sigma)^* \quad \text{and} \\ (\text{Id} + \Pi S)\mathbf{q} &= \mathbf{g}. \end{aligned} \tag{30}$$

*Reciprocally, if \mathbf{q} satisfies (30) then the pair (\mathbf{u}, \mathbf{p}) defined by $\mathbf{u} = (A - iB^*TB)^{-1}(B^*\mathbf{q} + \mathbf{f})$ and $\mathbf{p} = \mathbf{q} + iTB\mathbf{u}$ solves (25).*

Proof:

Assume that (\mathbf{u}, \mathbf{p}) is solution to (25) and consider $\mathbf{q} = \mathbf{p} - iTB\mathbf{u}$. We have $\mathbf{p} = \mathbf{q} + iTB\mathbf{u}$ and $\mathbf{p} + iT^*B\mathbf{u} = \mathbf{q} + 2iT_sB\mathbf{u}$. Then we can replace \mathbf{p} by $\mathbf{q} + iTB\mathbf{u}$ in (25) which leads to the equations

$$\begin{aligned} (A - iB^*TB)\mathbf{u} &= B^*\mathbf{q} + \mathbf{f}, \\ \mathbf{q} &= -\Pi(\mathbf{q} + 2iT_sB\mathbf{u}). \end{aligned}$$

Let us now decompose $\mathbf{u} = \tilde{\mathbf{u}} + \mathbf{u}_f$ where $\mathbf{u}_f = (A - iB^*TB)^{-1}\mathbf{f}$ and $\tilde{\mathbf{u}} = \mathbf{u} - \mathbf{u}_f$. This leads to $(A - iB^*TB)\tilde{\mathbf{u}} = B^*\mathbf{q}$ and $\mathbf{q} = -\Pi(\mathbf{q} + 2iT_sB\tilde{\mathbf{u}}) + \mathbf{g}$. There only remains to eliminate $\tilde{\mathbf{u}}$ in these equations, taking account of the definition of S from Lemma 5.2, which yields $(\text{Id} + \Pi S)\mathbf{q} = \mathbf{g}$.

Reciprocally assume that $\mathbf{q} \in \mathbb{V}_h(\Sigma)^*$ solves (30), and set $\mathbf{u} = (A - iB^*TB)^{-1}(B^*\mathbf{q} + \mathbf{f})$ and

$\mathbf{p} = \mathbf{q} + iT\mathbf{B}\mathbf{u}$. As a consequence, by construction, the first equation of (25) is satisfied, namely $\mathbf{A}\mathbf{u} = \mathbf{B}^*\mathbf{p} + \mathbf{f}$. There only remains to verify that the second equation of (25) is satisfied by (\mathbf{u}, \mathbf{p}) as well. Set $\mathbf{u}_f = (\mathbf{A} - i\mathbf{B}^*\mathbf{T}\mathbf{B})^{-1}\mathbf{f}$ so that $\mathbf{u} - \mathbf{u}_f = (\mathbf{A} - i\mathbf{B}^*\mathbf{T}\mathbf{B})^{-1}\mathbf{B}^*\mathbf{q}$ hence $\mathbf{S}\mathbf{q} = \mathbf{q} + 2iT_s\mathbf{B}(\mathbf{u} - \mathbf{u}_f)$. Plugging this in (30) yields

$$\begin{aligned}\mathbf{q} &= -\Pi\mathbf{S}(\mathbf{q}) + \mathbf{g} = -\Pi(\mathbf{q} + 2iT_s\mathbf{B}(\mathbf{u} - \mathbf{u}_f)) + \mathbf{g} \\ &= -\Pi(\mathbf{q} + 2iT_s\mathbf{B}\mathbf{u})\end{aligned}$$

Since $\mathbf{q} = \mathbf{p} - iT\mathbf{B}\mathbf{u}$ and $\mathbf{q} + 2iT_s\mathbf{B}\mathbf{u} = \mathbf{p} + iT^*\mathbf{B}\mathbf{u}$, the above equation rewrites $-\mathbf{p} + iT\mathbf{B}\mathbf{u} = \Pi(\mathbf{p} + iT^*\mathbf{B}\mathbf{u})$ which concludes the proof. \square

Previously we exhibited a chain of equivalent formulations that relates the initial discrete variational problem (3) to the skeleton equation (30). Well posedness of (3) thus immediately implies well posedness of (30) but there is more: the skeleton formulation provides a strongly coercive formulation of our Helmholtz problem. The following result is completely similar to Corollary 8.4 of [8].

Corollary 6.2.

Under Assumptions 1, 2 and 3, the operator $\text{Id} + \Pi\mathbf{S} : \mathbb{V}_h(\Sigma)^ \rightarrow \mathbb{V}_h(\Sigma)^*$ is an isomorphism that is T_s^{-1} -coercive. More precisely, for all $\mathbf{q} \in \mathbb{V}_h(\Sigma)^*$ we have*

$$\begin{aligned}\Re\{\langle (\text{Id} + \Pi\mathbf{S})\mathbf{q}, T_s^{-1}(\bar{\mathbf{q}}) \rangle\} &\geq \frac{\gamma_h^2}{2} \|\mathbf{q}\|_{T_s^{-1}}^2 \\ \text{where } \gamma_h &:= \inf_{\mathbf{q} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \|(\text{Id} + \Pi\mathbf{S})\mathbf{q}\|_{T_s^{-1}} / \|\mathbf{q}\|_{T_s^{-1}}\end{aligned}\tag{31}$$

Proof:

We first prove that $\text{Ker}(\text{Id} + \Pi\mathbf{S}) = \{0\}$ which will show that $\text{Id} + \Pi\mathbf{S}$ is an isomorphism (because $\dim(\mathbb{V}_h(\Sigma)^*) < +\infty$) and $\gamma_h > 0$. Assume that $(\text{Id} + \Pi\mathbf{S})\mathbf{q} = 0$ for some $\mathbf{q} \in \mathbb{V}_h(\Sigma)^*$. Set $\mathbf{g} = 0$, $\mathbf{f} = 0$, $\mathbf{u} = (\mathbf{A} - i\mathbf{B}^*\mathbf{T}\mathbf{B})^{-1}\mathbf{B}^*\mathbf{q}$ and $\mathbf{p} = \mathbf{q} + iT\mathbf{B}\mathbf{u}$. Applying Lemma 6.1 we see that the pair (\mathbf{u}, \mathbf{p}) solves (25) with $\mathbf{f} = 0$. Next applying the equivalences given by Lemma 4.2 and (4.1), and using Assumption 2 that yields existence and uniqueness of the solution to these boundary value problems, we conclude that $(\mathbf{u}, \mathbf{p}) = (0, 0)$, hence $\mathbf{q} = 0$.

Next, combining Lemma 3.2 and 5.2, we see that $\Pi\mathbf{S}$ is a contraction with respect to the norm (14) induced by T_s^{-1} . As a consequence, we obtain

$$\begin{aligned}\|\mathbf{q}\|_{T_s^{-1}}^2 &\geq \|\Pi\mathbf{S}\mathbf{q}\|_{T_s^{-1}}^2 = \|\mathbf{q} - (\text{Id} + \Pi\mathbf{S})\mathbf{q}\|_{T_s^{-1}}^2 \\ &= \|\mathbf{q}\|_{T_s^{-1}}^2 - 2\Re\{\langle (\text{Id} + \Pi\mathbf{S})\mathbf{q}, T_s^{-1}(\bar{\mathbf{q}}) \rangle\} + \|(\text{Id} + \Pi\mathbf{S})\mathbf{q}\|_{T_s^{-1}}^2 \\ \Re\{\langle (\text{Id} + \Pi\mathbf{S})\mathbf{q}, T_s^{-1}(\bar{\mathbf{q}}) \rangle\} / \|\mathbf{q}\|_{T_s^{-1}}^2 &\geq \frac{1}{2} \|(\text{Id} + \Pi\mathbf{S})\mathbf{q}\|_{T_s^{-1}}^2 / \|\mathbf{q}\|_{T_s^{-1}}^2 \geq \gamma_h^2/2.\end{aligned}$$

\square

According to Lemma 3.2 and 5.2, we have $\|\Pi\mathbf{S}(\mathbf{p})\|_{T_s^{-1}} \leq \|\mathbf{p}\|_{T_s^{-1}} \forall \mathbf{p} \in \mathbb{V}_h(\Sigma)^*$. Taking account of (31) in addition, we conclude that the field of values in the T_s^{-1} -scalar product $\{\langle (\text{Id} + \Pi\mathbf{S})\mathbf{p}, T_s^{-1}(\bar{\mathbf{p}}) \rangle : \|\mathbf{p}\|_{T_s^{-1}} = 1, \mathbf{p} \in \mathbb{V}_h(\Sigma)^*\}$ is contained in $\{\lambda \in \mathbb{C} : |\lambda - 1| \leq 1, \Re\{\lambda\} \geq \gamma_h^2/2\}$. Combined with e.g. Elman estimate [19], this readily yields an upper bound on the rate of convergence of GMRes [29, Prop. 10.35]. A similar remark holds for e.g. Richardson's algorithm see e.g. [29, §3.5].

7 Bounds on the trace operator

Before conducting a more detailed convergence analysis for the skeleton equation (30), we need to derive a few estimates related to the trace operator. The most natural constants related to this operator are its continuity modulus t_h^+ and its inf-sup constant t_h^- defined by

$$\begin{aligned} t_h^- &:= \inf_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|\langle \mathbf{B}(\mathbf{v}), \mathbf{p} \rangle|}{\|\mathbf{v}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}}, \\ t_h^+ &:= \sup_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|\langle \mathbf{B}(\mathbf{v}), \mathbf{p} \rangle|}{\|\mathbf{v}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}}. \end{aligned} \quad (32)$$

We have in particular $\|\mathbf{B}(\mathbf{v})\|_{\mathbb{T}_s} \leq t_h^+ \|\mathbf{v}\|_{\mathbb{H}^1(\Omega)}$. We provide an important alternative interpretation of t_h^\pm . Let us introduce a map $\mathbf{B}^\dagger : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Omega)$ defined as Moore-Penrose pseudo-inverse (see e.g. [27, §5.5.4] or [1, §2.6]) of the trace operator with respect to the volume norm

$$\begin{aligned} \mathbf{B}\mathbf{B}^\dagger &= \text{Id} \quad \text{and} \\ \|\mathbf{B}^\dagger(\mathbf{p})\|_{\mathbb{H}^1(\Omega)} &= \inf\{\|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} : \mathbf{u} \in \mathbb{V}_h(\Omega), \mathbf{B}(\mathbf{u}) = \mathbf{p}\}. \end{aligned} \quad (33)$$

By construction, for any $\mathbf{p} \in \mathbb{V}_h(\Sigma)$ and any $\mathbf{u} \in \mathbb{V}_h(\Omega)$ such that $\mathbf{B}(\mathbf{u}) = \mathbf{p}$ we have $\|\mathbf{B}^\dagger(\mathbf{p})\|_{\mathbb{H}^1(\Omega)} \leq \|\mathbf{u}\|_{\mathbb{H}^1(\Omega)}$. The map \mathbf{B}^\dagger is a classical object of domain decomposition literature that is sometimes referred to as discrete harmonic lifting, see [38, §4.4] and [36, Def. 1.55]. Because \mathbf{B} is subdomain-wise block-diagonal, \mathbf{B}^\dagger is itself block-diagonal. The operator $\mathbf{B}^\dagger\mathbf{B} : \mathbb{H}^1(\Omega) \rightarrow \mathbb{H}^1(\Omega)$ is a projector that is orthogonal with respect to the scalar product induced by (8).

Lemma 7.1.

Define $\Lambda : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ as the unique symmetric positive definite operator satisfying $\langle \Lambda(\mathbf{v}), \bar{\mathbf{v}} \rangle := \|\mathbf{B}^\dagger(\mathbf{v})\|_{\mathbb{H}^1(\Omega)}^2$ and set $\|\mathbf{v}\|_\Lambda^2 := \langle \Lambda(\mathbf{v}), \bar{\mathbf{v}} \rangle$. Then we have the identities

$$t_h^- = \inf_{\mathbf{v} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \frac{\|\mathbf{v}\|_{\mathbb{T}_s}}{\|\mathbf{v}\|_\Lambda}, \quad t_h^+ = \sup_{\mathbf{v} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \frac{\|\mathbf{v}\|_{\mathbb{T}_s}}{\|\mathbf{v}\|_\Lambda}. \quad (34)$$

Proof:

We only prove the identity related to t_h^- because the proof of the other identity follows a very similar path. First for any $\mathbf{v} \in \mathbb{V}_h(\Omega)$, setting $\tilde{\mathbf{v}} = \mathbf{B}^\dagger\mathbf{B}(\mathbf{v})$, we have $\mathbf{B}(\mathbf{v}) = \mathbf{B}(\tilde{\mathbf{v}})$ and $\|\tilde{\mathbf{v}}\|_{\mathbb{H}^1(\Omega)} \leq \|\mathbf{v}\|_{\mathbb{H}^1(\Omega)}$. Since the trace operator $\mathbf{B} : \mathbb{V}_h(\Omega) \rightarrow \mathbb{V}_h(\Sigma)$ is surjective, and $\mathbf{B}\mathbf{B}^\dagger\mathbf{B} = \mathbf{B}$ we conclude

$$\begin{aligned} \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|\langle \mathbf{B}(\mathbf{v}), \mathbf{p} \rangle|}{\|\mathbf{v}\|_{\mathbb{H}^1(\Omega)} \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}} &= \sup_{\mathbf{v} \in \mathbb{V}_h(\Omega) \setminus \{0\}} \frac{|\langle \mathbf{B}(\mathbf{v}), \mathbf{p} \rangle|}{\|\mathbf{B}^\dagger\mathbf{B}(\mathbf{v})\|_{\mathbb{H}^1(\Omega)} \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}} \\ &= \sup_{\mathbf{u} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \frac{|\langle \mathbf{u}, \mathbf{p} \rangle|}{\|\mathbf{u}\|_\Lambda \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}} = \frac{\|\mathbf{p}\|_{\Lambda^{-1}}}{\|\mathbf{p}\|_{\mathbb{T}_s^{-1}}} \end{aligned}$$

for any $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ with $\|\mathbf{p}\|_{\Lambda^{-1}}^2 := \langle \Lambda^{-1}\mathbf{p}, \bar{\mathbf{p}} \rangle$. The above calculus leads to the Rayleigh quotient expression $t_h^- = \inf_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \|\mathbf{p}\|_{\Lambda^{-1}} / \|\mathbf{p}\|_{\mathbb{T}_s^{-1}}$, which shows that $(t_h^-)^2$ can be characterized as an extremum of a generalized eigenvalue problem

$$\begin{aligned} (t_h^-)^2 &= \min\{\lambda \in \mathbb{R} : \text{Ker}(\Lambda^{-1} - \lambda\mathbb{T}_s^{-1}) \neq \{0\}\} \\ &= \min\{\lambda \in \mathbb{R} : \text{Ker}(\mathbb{T}_s - \lambda\Lambda) \neq \{0\}\} = \inf_{\mathbf{v} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \frac{\|\mathbf{v}\|_{\mathbb{T}_s}^2}{\|\mathbf{v}\|_\Lambda^2} \end{aligned}$$

□

From the previous result, we see that $\|\mathbf{B}^\dagger(\mathbf{v})\|_{\mathbb{H}^1(\Omega)} = \|\mathbf{v}\|_\Lambda \leq (1/t_h^-)\|\mathbf{v}\|_{\mathbb{T}_s}$ for all $\mathbf{v} \in \mathbb{V}_h(\Sigma)$. The constants t_h^\pm thus appear as key constants in the interplay between \mathbf{B} and \mathbb{T}_s .

Since \mathbf{B} and \mathbf{B}^\dagger are subdomain-wise block-diagonal, the operator $\Lambda : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ is itself subdomain-wise block-diagonal $\Lambda = \text{diag}_{j=1\dots J}(\Lambda_j)$, i.e. there are operators $\Lambda_j : \mathbb{V}_h(\Gamma_j) \rightarrow \mathbb{V}_h(\Gamma_j)^*, j = 1 \dots J$ such that

$$\langle \Lambda(\mathbf{u}), \mathbf{v} \rangle = \langle \Lambda_1(u_1), v_1 \rangle + \dots + \langle \Lambda_J(u_J), v_J \rangle$$

for $\mathbf{u}, \mathbf{v} \in \mathbb{V}_h(\Sigma)$ with $\mathbf{u} = (u_1, \dots, u_J)$ and $\mathbf{v} = (v_1, \dots, v_J)$. The operator Λ can actually serve as an effective choice of self-adjoint impedance, in which case $t_h^\pm = 1$. In any case, taking Λ as a reference scalar product over $\mathbb{V}_h(\Sigma)$, the constants t_h^\pm are then the extremal eigenvalues of the symmetrized impedance operator \mathbb{T}_s . In addition, if \mathbb{T} is also subdomain-wise block-diagonal $\mathbb{T} = \text{diag}_{j=1\dots J}(\mathbb{T}_j)$, then we have

$$\begin{aligned} (t_h^-)^2 &= \min_{j=1\dots J} \inf_{v \in \mathbb{V}_h(\Gamma_j) \setminus \{0\}} \Re\{\langle \mathbb{T}_j(v), \bar{v} \rangle / \|v\|_{\Lambda_j}^2\}, \\ (t_h^+)^2 &= \max_{j=1\dots J} \sup_{v \in \mathbb{V}_h(\Gamma_j) \setminus \{0\}} \Re\{\langle \mathbb{T}_j(v), \bar{v} \rangle / \|v\|_{\Lambda_j}^2\}. \end{aligned}$$

that is t_h^\pm can be interpreted as bounds on the (real part of the) field of values of the \mathbb{T}_j 's with respect to local reference norms induced by the Λ_j 's. To summarize and since only the case of subdomain-wise block-diagonal impedance can be regarded as computationally reasonable a situation, in practice, the constants t_h^\pm are determined by local (in each subdomain) behaviour of the impedance.

8 Coercivity estimate

The coercivity of the skeleton equation (30) is a valuable feature because it guarantees convergence of linear solvers. To properly estimate the speed of convergence though, we need to bound this coercivity estimate which is the focus of the present section. We first establish an intermediate estimation. We will consider

$$\begin{aligned} &\mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* \text{ equipped with} \\ &\|(\mathbf{v}, \mathbf{q})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}^2 := \|\mathbf{v}\|_{\mathbb{T}_s}^2 + \|\mathbf{q}\|_{\mathbb{T}_s^{-1}}^2 \end{aligned}$$

which is the natural cartesian product norm. For this space, we have previously considered two important subspaces: $\mathcal{X}_h(\Sigma) := \mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$ in Lemma 3.4, and the space of Cauchy data $\mathcal{C}_h(\Sigma)$ in Lemma 5.3. Besides t_h^\pm equivalently defined by (32) or (34), we shall also rely on the inf-sup constant α_h and the continuity modulus $\|a\|$ defined by (22).

Proposition 8.1.

Let Assumptions 1, 2 and 3 hold and set $\mathcal{X}_h(\Sigma) := \mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$. Then we have the following (a priori not orthogonal) direct sum

$$\mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* = \mathcal{X}_h(\Sigma) \oplus \mathcal{C}_h(\Sigma).$$

Moreover if $P : \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* \rightarrow \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$ refers to the projector with $\text{Ker}(P) = \mathcal{X}_h(\Sigma)$ and $\text{Range}(P) = \mathcal{C}_h(\Sigma)$, we have

$$\|P\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}} := \sup_{(\mathbf{v}, \mathbf{q}) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* \setminus \{(0,0)\}} \frac{\|P(\mathbf{v}, \mathbf{q})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}}{\|(\mathbf{v}, \mathbf{q})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}} \leq \frac{(t_h^+)^2 + (2\|a\|/t_h^-)^2}{\alpha_h}$$

Proof:

First assume that $(\mathbf{p}, \mathbf{q}) \in \mathcal{X}_h(\Sigma) \cap \mathcal{C}_h(\Sigma)$. By the very definition of $\mathcal{C}_h(\Sigma)$, there exists $\mathbf{u} \in \mathbb{V}_h(\Omega)$ such that $B\mathbf{u} = \mathbf{p}$ and $A\mathbf{u} = B^*\mathbf{q}$. Since $\mathbf{p} \in \mathbb{X}_h(\Sigma)$, we conclude that $\mathbf{u} \in \mathbb{X}_h(\Omega)$. Next for any $\mathbf{v} \in \mathbb{X}_h(\Omega)$ we have $B\mathbf{v} \in \mathbb{X}_h(\Sigma)$ and hence $\langle B^*\mathbf{q}, \mathbf{v} \rangle = \langle \mathbf{q}, B\mathbf{v} \rangle = 0$ as $\mathbf{q} \in \mathbb{X}_h(\Sigma)^\circ$. To sum up, we have $\mathbf{u} \in \mathbb{X}_h(\Omega)$ and $\langle A\mathbf{u}, \mathbf{v} \rangle = \langle B^*\mathbf{q}, \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathbb{X}_h(\Omega)$. Using (22), we deduce that $\mathbf{u} = 0$ hence $\mathbf{p} = B\mathbf{u} = 0$ and $B^*\mathbf{q} = A\mathbf{u} = 0 \Rightarrow \mathbf{q} = 0$ since B^* is one-to-one (because B is onto). We have just established that

$$\mathcal{X}_h(\Sigma) \cap \mathcal{C}_h(\Sigma) = \{0\}.$$

Now we show that $\mathcal{X}_h(\Sigma) + \mathcal{C}_h(\Sigma) = \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$. Pick an arbitrary pair $(\mathbf{p}_D, \mathbf{p}_N) \in \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$. Define $\tilde{\mathbf{u}}$ as the unique element of $\mathbb{X}_h(\Omega)$ such that $\langle A(\tilde{\mathbf{u}} + B^\dagger \mathbf{p}_D), \mathbf{v} \rangle = \langle \mathbf{p}_N, B(\mathbf{v}) \rangle$ for all $\mathbf{v} \in \mathbb{X}_h(\Omega)$. Taking account of (22) then yields

$$\alpha_h \|\tilde{\mathbf{u}}\|_{\mathbb{H}^1(\Omega)} \leq (\|a\|/t_h^-) \|\mathbf{p}_D\|_{\mathbb{T}_s} + t_h^+ \|\mathbf{p}_N\|_{\mathbb{T}_s^{-1}} \quad (35)$$

Next let us set $\mathbf{u} = \tilde{\mathbf{u}} + B^\dagger(\mathbf{p}_D)$ and $\mathbf{u}_D = B(\mathbf{u}) = B(\tilde{\mathbf{u}}) + \mathbf{p}_D$. For all $\mathbf{v} \in \mathbb{V}_h(\Omega)$ satisfying $B(\mathbf{v}) = 0$ we have $\langle A\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{p}_N, B(\mathbf{v}) \rangle = 0$ hence, applying Lemma 2.1 yields the existence of \mathbf{u}_N such that $\langle A\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}_N, B(\mathbf{v}) \rangle \forall \mathbf{v} \in \mathbb{V}_h(\Omega)$ which rewrites $A\mathbf{u} = B^*\mathbf{u}_N$. In particular we have $\langle \mathbf{u}_N, \mathbf{q} \rangle = \langle A\mathbf{u}, B^\dagger(\mathbf{q}) \rangle$ for all $\mathbf{q} \in \mathbb{V}_h(\Sigma)$. From this we deduce the estimates

$$\begin{aligned} \|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} &\leq \|\tilde{\mathbf{u}}\|_{\mathbb{H}^1(\Omega)} + (1/t_h^-) \|\mathbf{p}_D\|_{\mathbb{T}_s} \\ \|\mathbf{u}_D\|_{\mathbb{T}_s} &\leq t_h^+ \|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} \\ \|\mathbf{u}_N\|_{\mathbb{T}_s^{-1}} &\leq (\|a\|/t_h^-) \|\mathbf{u}\|_{\mathbb{H}^1(\Omega)} \end{aligned} \quad (36)$$

Now observe that, by construction, we have $(\mathbf{u}_D, \mathbf{u}_N) \in \mathcal{C}_h(\Sigma)$ see Proposition 5.3. Besides $\mathbf{p}_D - \mathbf{u}_D = -B(\tilde{\mathbf{u}}) \in \mathbb{X}_h(\Sigma)$ since $\tilde{\mathbf{u}} \in \mathbb{X}_h(\Omega)$. On the other hand $\langle \mathbf{p}_N - \mathbf{u}_N, B(\mathbf{v}) \rangle = \langle A(\mathbf{u}), \mathbf{v} \rangle - \langle A(\mathbf{u}), \mathbf{v} \rangle = 0$ for all $\mathbf{v} \in \mathbb{X}_h(\Omega)$ and, since $\mathbb{X}_h(\Sigma) = B(\mathbb{X}_h(\Omega))$, we conclude that $\mathbf{p}_N - \mathbf{u}_N \in \mathbb{X}_h(\Sigma)^\circ$. In conclusion we have proved that $(\mathbf{u}_D, \mathbf{u}_N) - (\mathbf{p}_D, \mathbf{p}_N) \in \mathcal{X}_h(\Sigma)$ hence $(\mathbf{u}_D, \mathbf{u}_N) = P(\mathbf{p}_D, \mathbf{p}_N)$. To conclude the proof, there only remains to combine (35) and (36). \square

The projection P introduced in the previous result is intimately connected to the inverse of the operator $\text{Id} + \text{IIS}$ of the skeleton formulation (30). This is made apparent by the factorized form provided by the next theorem.

Theorem 8.2.

Let Assumptions 1, 2 and 3 hold. Define $\mathcal{T} : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^*$ by $\mathcal{T}(\mathbf{v}) = (\mathbf{v}, -i\mathbb{T}^*\mathbf{v})$ and $\mathcal{T}' : \mathbb{V}_h(\Sigma) \times \mathbb{V}_h(\Sigma)^* \rightarrow \mathbb{V}_h(\Sigma)^*$ by $\mathcal{T}'(\mathbf{v}, \mathbf{p}) = \mathbf{p} - i\mathbb{T}(\mathbf{v})$. Then we have

$$(\text{Id} + \text{IIS})^{-1} = i\mathcal{T}' \cdot P \cdot \mathcal{T} \cdot \mathbb{T}_s^{-1}/2.$$

Proof:

Pick an arbitrary $\mathbf{f} \in \mathbb{V}_h(\Sigma)^*$ and define $\mathbf{p}_D = \mathbb{T}_s^{-1}(\mathbf{f})/2$ and $\mathbf{p}_N = -i\mathbb{T}^*\mathbb{T}_s^{-1}(\mathbf{f})/2$, which simply rewrites $(\mathbf{p}_D, \mathbf{p}_N) = \mathcal{T} \cdot \mathbb{T}_s^{-1}(\mathbf{f})/2$. Next define $(\mathbf{u}_D, \mathbf{u}_N) = \mathbb{P}(\mathbf{p}_D, \mathbf{p}_N)$ hence $(\mathbf{u}_D - \mathbf{p}_D, \mathbf{u}_N - \mathbf{p}_N) \in \mathcal{X}_h(\Sigma) = \mathbb{X}_h(\Sigma) \times \mathbb{X}_h(\Sigma)^\circ$. On the other hand $(\mathbf{u}_D, \mathbf{u}_N) \in \mathcal{C}_h(\Sigma)$, so applying Lemma 3.4 and Proposition 5.3 yields

$$\begin{aligned}
& -(\mathbf{u}_N - \mathbf{p}_N) + i\mathbb{T}(\mathbf{u}_D - \mathbf{p}_D) = \Pi((\mathbf{u}_N - \mathbf{p}_N) + i\mathbb{T}^*(\mathbf{u}_D - \mathbf{p}_D)) \\
\iff & \mathbf{u}_N - i\mathbb{T}\mathbf{u}_D + \Pi(\mathbf{u}_N + i\mathbb{T}^*\mathbf{u}_D) = (\text{Id} + \Pi)\mathbf{p}_N - i(\mathbb{T} - \Pi\mathbb{T}^*)\mathbf{p}_D \\
\iff & (\text{Id} + \Pi\mathbb{S})(\mathbf{u}_N - i\mathbb{T}\mathbf{u}_D) = (\text{Id} + \Pi)\mathbf{p}_N - i(\mathbb{T} - \Pi\mathbb{T}^*)\mathbf{p}_D \\
\iff & (\text{Id} + \Pi\mathbb{S})\mathcal{T}'\mathcal{P}\mathcal{T}\mathbb{T}_s^{-1}(\mathbf{f})/2 = (\text{Id} + \Pi)\mathbf{p}_N - i(\mathbb{T} - \Pi\mathbb{T}^*)\mathbf{p}_D
\end{aligned} \tag{37}$$

Coming back to the definition of the exchange operator Π given by Lemma 3.2, and setting $\mathbb{Q} = \mathbb{R}(\mathbb{R}^*\mathbb{T}^*\mathbb{R})^{-1}\mathbb{R}^*$, we have $\text{Id} + \Pi = 2\mathbb{T}_s\mathbb{Q}$ and $\mathbb{T} - \Pi\mathbb{T}^* = 2\mathbb{T}_s(\text{Id} - \mathbb{Q}\mathbb{T}^*)$. Combining these identities with the definition of $(\mathbf{p}_D, \mathbf{p}_N)$ leads to $(\text{Id} + \Pi)\mathbf{p}_N - i(\mathbb{T} - \Pi\mathbb{T}^*)\mathbf{p}_D = -i\mathbf{f}$. We have thus established the desired result. \square

Note that $\mathcal{T}' \cdot \mathcal{T} = 2\mathbb{T}_s/i$, so that $(i/2)\mathcal{T}\mathbb{T}_s^{-1}\mathcal{T}'$ is (a priori oblique) a projector, although this observation seems of no use in the present context. We will exploit Theorem 8.2 to establish a coercivity estimate for $\text{Id} + \Pi\mathbb{S}$. The non-self adjoint part of the impedance operator \mathbb{T} shall arise naturally in this analysis so we introduce a notation for bounding it

$$\begin{aligned}
t_h^* & := \sup_{\mathbf{v} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \left\| \left(\frac{\mathbb{T} - \mathbb{T}^*}{2} \right) \mathbf{v} \right\|_{\mathbb{T}_s^{-1}} / \|\mathbf{v}\|_{\mathbb{T}_s} \\
& = \sup_{\mathbf{v} \in \mathbb{V}_h(\Sigma) \setminus \{0\}} \left| \frac{\langle (\mathbb{T} - \mathbb{T}^*)\mathbf{v}, \bar{\mathbf{v}} \rangle}{\langle (\mathbb{T} + \mathbb{T}^*)\mathbf{v}, \bar{\mathbf{v}} \rangle} \right|.
\end{aligned} \tag{38}$$

Theorem 8.3.

Let Assumptions 1, 2 and 3 hold. Then the inf-sup constant γ_h from (31) admits the following lower bound

$$\gamma_h \geq \frac{2\alpha_h}{[1 + (1 + t_h^*)^2] [(t_h^+)^2 + (2\|a\|/t_h^-)^2]}$$

Proof:

To bound γ_h from below, it suffices to bound $1/\gamma_h$ from above. On the other hand, close inspection of the definition of γ_h from (31) shows that $1/\gamma_h$ is the continuity modulus of $(\text{Id} - \Pi\mathbb{S})^{-1}$ which writes

$$\frac{1}{\gamma_h} = \sup_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \frac{\|(\text{Id} + \Pi\mathbb{S})^{-1}\mathbf{p}\|_{\mathbb{T}_s^{-1}}}{\|\mathbf{p}\|_{\mathbb{T}_s^{-1}}}. \tag{39}$$

Next we need to obtain upper bounds for the continuity modulus of $\mathcal{T}, \mathcal{T}'$ and \mathbb{P} . A direct estimation using (38) yields

$$\begin{aligned}
\|\mathcal{T}'(\mathbf{v}, \mathbf{q})\|_{\mathbb{T}_s^{-1}} & \leq \|\mathbf{q}\|_{\mathbb{T}_s^{-1}} + (1 + t_h^*)\|\mathbf{v}\|_{\mathbb{T}_s} \\
& \leq (1 + (1 + t_h^*)^2)^{1/2} \|(\mathbf{v}, \mathbf{q})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}} \\
\|\mathcal{T}(\mathbf{v})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}} & \leq (\|\mathbf{v}\|_{\mathbb{T}_s}^2 + \|\mathbb{T}^*(\mathbf{v})\|_{\mathbb{T}_s^{-1}}^2)^{1/2} \\
& \leq (1 + (1 + t_h^*)^2)^{1/2} \|\mathbf{v}\|_{\mathbb{T}_s}
\end{aligned} \tag{40}$$

In the estimate above we used the elementary inequalities $\|\mathbf{T}(\mathbf{v})\|_{\mathbb{T}_s^{-1}} \leq (1 + t_h^*)\|\mathbf{v}\|_{\mathbb{T}_s}$ and similarly for \mathbb{T}^* , and $\|\mathbb{T}_s(\mathbf{v})\|_{\mathbb{T}_s^{-1}} = \|\mathbf{v}\|_{\mathbb{T}_s}$. Plugging the factorized form of Theorem 8.2, combined with (40) and Proposition 8.1, into (39) yields the desired estimate. \square

The next result, that agrees with the estimate provided by [8, Prop. 10.4], yields another variant of this coercivity bound that appears sharper in certain cases.

Proposition 8.4.

Let Assumptions 1, 2 and 3 hold, and suppose in addition that $\mathbb{T} = \mathbb{T}^ = \mathbb{T}_s$. Then we have the following estimate*

$$\gamma_h = \inf_{\mathbf{q} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \frac{\|(\text{Id} + \Pi\mathbb{S})\mathbf{q}\|_{\mathbb{T}_s^{-1}}}{\|\mathbf{q}\|_{\mathbb{T}_s^{-1}}} \geq \frac{1}{\|\mathbb{P}\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}}. \quad (41)$$

Proof:

Since $\mathbb{T} = \mathbb{T}_s$ we have $\|\mathcal{T}'(\mathbf{v}, \mathbf{p})\|_{\mathbb{T}_s^{-1}} \leq \sqrt{2}\|(\mathbf{v}, \mathbf{p})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}$ for all $\mathbf{v} \in \mathbb{V}_h(\Sigma)$, $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$. Moreover a direct calculation yields $\|\mathcal{T}\mathbb{T}_s^{-1}(\mathbf{p})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}^2 = 2\|\mathbf{p}\|_{\mathbb{T}_s^{-1}}^2$. Using the definition of the continuity modulus of \mathbb{P} , as well as Theorem 8.2 and the definition of γ_h , we obtain the inequality

$$\begin{aligned} \frac{1}{\gamma_h} &= \sup_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \frac{\|\mathcal{T}'\mathbb{P}\mathcal{T}\mathbb{T}_s^{-1}(\mathbf{p})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}}{2\|\mathbf{p}\|_{\mathbb{T}_s^{-1}}} \leq \sup_{\mathbf{p} \in \mathbb{V}_h(\Sigma)^* \setminus \{0\}} \frac{\|\mathbb{P}\mathcal{T}\mathbb{T}_s^{-1}(\mathbf{p})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}}{\|\mathcal{T}\mathbb{T}_s^{-1}(\mathbf{p})\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}} \\ &\leq \|\mathbb{P}\|_{\mathbb{T}_s \times \mathbb{T}_s^{-1}}. \end{aligned}$$

\square

9 Purely local exchange operator

In this section, we wish to show that the theory of the present contribution covers other pre-existing variants of OSM that involve non-self-adjoint impedance operators. We will in particular establish connections with the DDM strategies described in [11] and [17]. Additionally let us recall that, as explained in [8, Sect.12], our theoretical framework also covers the initial Després algorithm of [14, 13, 12, 15].

Applying the exchange operator introduced in Section 3 (i.e. the operation $\mathbf{p} \mapsto \Pi(\mathbf{p})$) is computationally non-trivial because the expression of Π involves the inverse operator $(\mathbf{R}^*\mathbf{T}^*\mathbf{R})^{-1}$ which is a priori not known explicitly. However in the particular case where the impedance is diagonal, calculations related to the exchange operator become much simpler. The present section will focus on the even simpler situation where the impedance is associated to the identity matrix which is already an interesting and instructive special case.

9.1 Derivation of the swapping operator

Let us denote $\text{dof}(\Omega)$ the set of degrees of freedom of the \mathbb{P}_k -Lagrange discretization over the whole computational domain, which will be considered a point cloud, and set $\text{dof}(\Gamma_j) =$

$\Gamma_j \cap \text{dof}(\Omega)$ and $\text{dof}(\Sigma) = \Sigma \cap \text{dof}(\Omega)$. In the present section we will assume that $\mathbf{T} = \mathbf{D} = \text{diag}_{j=1 \dots J}(\mathbf{D}_j)$ where $\mathbf{D} : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ is defined by

$$\begin{aligned} \langle \mathbf{D}(\mathbf{u}), \mathbf{v} \rangle &:= \langle \mathbf{D}_1(u_1), v_1 \rangle + \dots + \langle \mathbf{D}_J(u_J), v_J \rangle \\ \text{with } \langle \mathbf{D}_j(u_j), v_j \rangle &:= \sum_{\mathbf{x} \in \text{dof}(\Gamma_j)} u_j(\mathbf{x})v_j(\mathbf{x}) \end{aligned} \quad (42)$$

for any $\mathbf{u}, \mathbf{v} \in \mathbb{V}_h(\Sigma)$ with $\mathbf{u} = (u_1, \dots, u_J)$ and $\mathbf{v} = (v_1, \dots, v_J)$. Obviously this is a self-adjoint operator $\mathbf{D} = \mathbf{D}^*$. Algebraically, each \mathbf{D}_j simply corresponds to an identity matrix. We shall denote Π_{loc} the associated exchange operator which is defined by $\Pi_{\text{loc}} = 2\mathbf{D}(\mathbf{R}^*\mathbf{D}\mathbf{R})^{-1}\mathbf{R}^* - \text{Id}$ in accordance with Lemma 3.2.

To obtain a more explicit expression for the exchange operator Π_{loc} , we start from an arbitrary $\mathbf{p} = (p_1, \dots, p_J) \in \mathbb{V}_h(\Sigma)^*$ and we decompose each local component $p_j \in \mathbb{V}_h(\Gamma_j)^*, j = 1 \dots J$ according to its coordinates in the canonical dual basis which writes

$$\langle p_j, v \rangle = \sum_{\mathbf{x} \in \text{dof}(\Gamma_j)} p_j(\mathbf{x})v(\mathbf{x}) \quad \forall v \in \mathbb{V}_h(\Gamma_j).$$

In other words, defining $\delta_{\mathbf{x}} \in \mathbb{V}_h(\Gamma_j)^*$ by $\langle \delta_{\mathbf{x}}, v \rangle := v(\mathbf{x})$, we have $p_j = \sum_{\mathbf{x} \in \text{dof}(\Gamma_j)} p_j(\mathbf{x})\delta_{\mathbf{x}}$. We derive an expression of $\Pi(\mathbf{p})$ in terms of the coefficients $p_j(\mathbf{x})$. Examining the action of the operators $\mathbf{R}^* : \mathbb{V}_h(\Sigma)^* \rightarrow \mathbb{V}_h(\Sigma)^*$ and $\mathbf{R}^*\mathbf{D}\mathbf{R} : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ we obtain the following: for any $u, v \in \mathbb{V}_h(\Sigma)$

$$\begin{aligned} \langle \mathbf{R}^*(\mathbf{p}), v \rangle &= \langle \mathbf{p}, \mathbf{R}(v) \rangle = \sum_{\mathbf{x} \in \text{dof}(\Sigma)} v(\mathbf{x}) \sum_{j: \mathbf{x} \in \Gamma_j} p_j(\mathbf{x}) \\ \langle \mathbf{D}\mathbf{R}(u), \mathbf{R}(v) \rangle &= \sum_{\mathbf{x} \in \text{dof}(\Sigma)} \#\{j : \mathbf{x} \in \Gamma_j\} u(\mathbf{x})v(\mathbf{x}) \end{aligned} \quad (43)$$

In particular $\mathbf{R}^*\mathbf{D}\mathbf{R}$ appears diagonal. Now plugging the previous expressions into the definition $\Pi_{\text{loc}} := 2\mathbf{D}\mathbf{R}(\mathbf{R}^*\mathbf{D}\mathbf{R})^{-1}\mathbf{R}^* - \text{Id}$, for any $\mathbf{v} = (v_1, \dots, v_J) \in \mathbb{V}_h(\Sigma)$ we obtain

$$\begin{aligned} \langle \Pi_{\text{loc}}(\mathbf{p}), \mathbf{v} \rangle &= \\ \sum_{j=1 \dots J} \sum_{\mathbf{x} \in \text{dof}(\Gamma_j)} v_j(\mathbf{x}) &\left[-p_j(\mathbf{x}) + \frac{2}{\#\{n : \mathbf{x} \in \Gamma_n\}} \sum_{k: \mathbf{x} \in \Gamma_k} p_k(\mathbf{x}) \right] \end{aligned} \quad (44)$$

Let us inspect this expression, assuming for a moment that the subdomain partition does not contain any cross-point. In this case we have $\#\{n : \mathbf{x} \in \Gamma_n\} = 2 \forall \mathbf{x} \in \text{dof}(\Sigma)$ and the operator $-\Pi_{\text{loc}}$ simply consists in swapping the unknowns from both sides of each interface. This operator Π_{loc} should thus be understood as the exchange operator that is used in the rest of the literature on Optimized Schwarz Methods. This is indeed the operator considered in [4, 17, 16, 18, 10, 31, 9, 25, 32, 23, 15, 14, 12, 11, 5, 20, 33, 2] for enforcing coupling between neighbouring subdomains. Expression (44) can be re-arranged so as to take a more symmetric form

$$\langle \Pi_{\text{loc}}(\mathbf{p}), \mathbf{v} \rangle = -\langle \mathbf{p}, \mathbf{v} \rangle + 2 \sum_{\mathbf{x} \in \text{dof}(\Sigma)} \frac{\left[\sum_{j: \mathbf{x} \in \Gamma_j} p_j(\mathbf{x}) \right] \left[\sum_{k: \mathbf{x} \in \Gamma_k} v_k(\mathbf{x}) \right]}{\#\{n : \mathbf{x} \in \Gamma_n\}}. \quad (45)$$

Applying Lemma 3.3 with $\mathbf{T} = \mathbf{D}$ yields that $\Pi_{\text{loc}}^* \mathbf{R} = \mathbf{R}$, $\Pi_{\text{loc}}^2 = \text{Id}$ i.e. swapping traces twice leaves them unchanged, and $\Pi_{\text{loc}}(\mathbf{q}) = -\mathbf{q} \forall \mathbf{q} \in \mathbb{X}_h(\Sigma)^\circ$.

9.2 A criterion for locality

The expression (44) and (45) are fully explicit so that applying Π_{loc} is fast and straightforward. We proved that the exchange operator induced by D is $\Pi_{\text{loc}} = 2DR(R^*DR)^{-1}R^* - \text{Id}$. But other choices of impedance can induce the same exchange operator. A natural question arises then to determine those impedance operators that lead to Π_{loc} as exchange operator. The next lemma provides an explicit criterion for this.

Lemma 9.1.

For any linear map $T : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ satisfying Assumption 3 we have

$$\Pi_{\text{loc}}T^*R = \text{TR} \iff \Pi_{\text{loc}} = (T + T^*)R(R^*T^*R)^{-1}R^* - \text{Id}.$$

Proof:

Assume first that $\Pi_{\text{loc}} = (T + T^*)R(R^*T^*R)^{-1}R^* - \text{Id}$. Then a direct calculus shows that $\Pi_{\text{loc}}T^*R = 2T_sR(R^*T^*R)^{-1}(R^*T^*R) - T^*R = (T + T^*)R - T^*R = \text{TR}$.

Reciprocally assume that $\Pi_{\text{loc}}T^*R = \text{TR}$ holds. Pick an arbitrary $\mathbf{p} \in \mathbb{V}_h(\Sigma)^*$ and let $u \in \mathbb{V}_h(\Sigma)$ solve $\langle T^*R(u), R(v) \rangle = \langle \mathbf{p}, R(v) \rangle \forall v \in \mathbb{V}_h(\Sigma)$, which is uniquely solvable a variational problem thanks to the coercivity of T^* given by Assumption 3. Then we have $u = (R^*T^*R)^{-1}R^*\mathbf{p}$ and, by construction, $\mathbf{q} = \mathbf{p} - T^*R(u) \in \mathbb{X}_h(\Sigma)^\circ$. From this we conclude

$$\begin{aligned} \Pi_{\text{loc}}(\mathbf{p}) &= \Pi_{\text{loc}}(\mathbf{q}) + \Pi_{\text{loc}}T^*R(u) \\ &= -\mathbf{q} + \text{TR}(u) = -\mathbf{q} - T^*R(u) + (T + T^*)R(u) \\ &= -\mathbf{p} + (T + T^*)R(R^*T^*R)^{-1}R^*\mathbf{p} \end{aligned}$$

In the above calculus we have used the fact that $\Pi_{\text{loc}}(\mathbf{q}) = -\mathbf{q} \forall \mathbf{q} \in \mathbb{X}_h(\Sigma)^\circ$. Since the tuple of traces \mathbf{p} was chosen arbitrarily in $\mathbb{V}_h(\Sigma)^*$, we have proved the desired result. \square

As a corollary, the criterion exhibited in Lemma 9.1 can be simplified, taking the form of a commutation identity. This is not an equivalence anymore though.

Corollary 9.2.

For any linear map $T : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ satisfying Assumption 3 we have

$$\Pi_{\text{loc}}T = T^*\Pi_{\text{loc}}^* \implies \Pi_{\text{loc}} = (T + T^*)R(R^*T^*R)^{-1}R^* - \text{Id}.$$

Proof:

Observe that $(\Pi_{\text{loc}})^2 = \text{Id}$, hence multiplying $\Pi_{\text{loc}}T = T^*\Pi_{\text{loc}}^*$ on the left by Π_{loc} and on the right by Π_{loc}^* yields $T\Pi_{\text{loc}}^* = \Pi_{\text{loc}}T^*$. Since we have $\Pi_{\text{loc}}^*R = R$ systematically according to Lemma 3.3, we conclude that $\Pi_{\text{loc}}T^*R = T\Pi_{\text{loc}}^*R = \text{TR}$. There only remains to apply Lemma 9.1 which yields the desired result. \square

The criterion $\Pi_{\text{loc}}T = T^*\Pi_{\text{loc}}^*$ is precisely the one considered in [11, §2.1.2]. In this reference the authors write $T' = T^*$, and impose the same condition as Assumption 3 (see Eq.(2.13) in [11]) and only consider configuration with no cross-point. The criterion $\Pi_{\text{loc}}T = T^*\Pi_{\text{loc}}^*$ also directly corresponds to the compatibility assumption of [17, Def.12 & Lem.14] where a particular choice of impedance fulfilling Assumption 3 is considered, see [17, Prop.4].

The present discussion shows that our theory covers the strategies considered in [11] and [17] embedding them in a more general framework and, in passing, provides a refined convergence estimate for them through application of Theorem 8.3.

In conclusion, we would like to point that, in the special case of a self-adjoint impedance, the criterion provided by Corollary 9.2 does give rise to an equivalence.

Corollary 9.3.

For any linear map $T : \mathbb{V}_h(\Sigma) \rightarrow \mathbb{V}_h(\Sigma)^*$ satisfying Assumption 3, and that is in addition self-adjoint $T = T^*$, we have

$$\Pi_{\text{loc}} T = T \Pi_{\text{loc}}^* \iff \Pi_{\text{loc}} = 2\text{TR}(\text{R}^* \text{TR})^{-1} \text{R}^* - \text{Id}.$$

Proof:

Direct calculus indicates clearly that, if $\Pi_{\text{loc}} = 2\text{TR}(\text{R}^* \text{TR})^{-1} \text{R}^* - \text{Id}$, then $\Pi_{\text{loc}} T = T \Pi_{\text{loc}}^*$. Reciprocally, assume that $\Pi_{\text{loc}} T = T \Pi_{\text{loc}}^*$ and denote $Q := (\text{Id} + \Pi_{\text{loc}}^*)/2$. According to Lemma 3.3, Q is a projector onto $\mathbb{X}_h(\Sigma)$. By hypothesis we have $Q^* T = T Q$ so that Q is more precisely a T -orthogonal projector onto $\mathbb{X}_h(\Sigma)$. As a consequence, for any $\mathbf{w} \in \mathbb{V}_h(\Sigma)$ there exists $u \in \mathbb{V}_h(\Sigma)$ such that $Q(\mathbf{w}) = \text{R}(u)$ and $\langle \text{TR}(u), \text{R}(v) \rangle = \langle T(\mathbf{w}), \text{R}(v) \rangle \forall v \in \mathbb{V}_h(\Sigma)$, which rewrites $u = (\text{R}^* \text{TR})^{-1} \text{R}^* T(\mathbf{w})$ hence $Q = \text{R}(\text{R}^* \text{TR})^{-1} \text{R}^* T$. Finally, by hypothesis we have $\Pi_{\text{loc}} = T \Pi_{\text{loc}}^* T^{-1} = T(2Q - \text{Id})T^{-1} = 2\text{TR}(\text{R}^* \text{TR})^{-1} \text{R}^* - \text{Id}$. \square

References

- [1] G. Allaire and S.M. Kaber. *Numerical linear algebra*, volume 55. New York, NY: Springer, 2008.
- [2] A. Bendali and Y. Boubendir. Non-overlapping domain decomposition method for a nodal finite element method. *Numerische Mathematik*, 103(4):515–537, Jun 2006.
- [3] D. Boffi, F. Brezzi, and M. Fortin. *Mixed finite element methods and applications*, volume 44. Berlin: Springer, 2013.
- [4] Y. Boubendir. An analysis of the BEM-FEM non-overlapping domain decomposition method for a scattering problem. *J. Comput. Appl. Math.*, 204(2):282–291, 2007.
- [5] Y. Boubendir, X. Antoine, and C. Geuzaine. A Quasi-Optimal Non-Overlapping Domain Decomposition Algorithm for the Helmholtz Equation. *J. Comp. Phys.*, 213(2):262–280, 2012.
- [6] X. Claeys. Non-local variant of the Optimised Schwarz Method for arbitrary non-overlapping subdomain partitions. *ESAIM: M2AN*, 55(2):429–448, 2021.
- [7] X. Claeys and R. Hiptmair. Multi-trace boundary integral formulation for acoustic scattering by composite structures. *Comm. Pure Appl. Math.*, 66(8):1163–1201, 2013.
- [8] X. Claeys and E. Parolin. Robust treatment of cross points in optimized schwarz methods, 2020. preprint arXiv:2003.06657.
- [9] X. Claeys, B. Thierry, and F. Collino. Integral equation based optimized Schwarz method for electromagnetics. In *Domain decomposition methods in science and engineering XXIV*, volume 125 of *Lect. Notes Comput. Sci. Eng.*, pages 187–194. Springer, Cham, 2018.

- [10] F. Collino, S. Ghanemi, and P. Joly. Domain decomposition method for harmonic wave propagation: a general presentation. *Computer Methods in Applied Mechanics and Engineering*, 184(2):171 – 211, 2000.
- [11] F. Collino, P. Joly, and M. Lecouvez. Exponentially convergent non overlapping domain decomposition methods for the Helmholtz equation. *ESAIM, Math. Model. Numer. Anal.*, 54(3):775–810, 2020.
- [12] B. Després. Décomposition de domaine et problème de Helmholtz. *C. R. Acad. Sci. Paris Sér. I Math.*, 311(6):313–316, 1990.
- [13] B. Després. Domain decomposition method and the Helmholtz problem. In *Mathematical and numerical aspects of wave propagation phenomena (Strasbourg, 1991)*, pages 44–52. SIAM, Philadelphia, PA, 1991.
- [14] B. Després. *Méthodes de décomposition de domaine pour les problèmes de propagation d’ondes en régime harmonique. Le théorème de Borg pour l’équation de Hill vectorielle.* Institut National de Recherche en Informatique et en Automatique (INRIA), Rocquencourt, 1991. Thèse, Université de Paris IX (Dauphine), Paris, 1991.
- [15] B. Després. Domain decomposition method and the Helmholtz problem. II. In *Second International Conference on Mathematical and Numerical Aspects of Wave Propagation (Newark, DE, 1993)*, pages 197–206. SIAM, Philadelphia, PA, 1993.
- [16] B. Després, A. Nicolopoulos, and B. Thierry. Corners and stable optimized domain decomposition methods for the Helmholtz problem. preprint hal-02612368, May 2020.
- [17] B. Després, A. Nicolopoulos, and B. Thierry. On Domain Decomposition Methods with optimized transmission conditions and cross-points. preprint hal-03230250, May 2021.
- [18] V. Dolean, P. Jolivet, and F. Nataf. *An introduction to domain decomposition methods.* Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2015. Algorithms, theory, and parallel implementation.
- [19] S.C. Eisenstat, H.C. Elman, and M.H. Schultz. Variational iterative methods for non-symmetric systems of linear equations. *SIAM J. Numer. Anal.*, 20:345–357, 1983.
- [20] M. El Bouajaji, B. Thierry, X. Antoine, and C. Geuzaine. A quasi-optimal domain decomposition algorithm for the time-harmonic maxwell’s equations. *Journal of Computational Physics*, 294:38–57, 2015.
- [21] M. Gander and F. Kwok. On the applicability of Lions’ energy estimates in the analysis of discrete optimized schwarz methods with cross points. *Lecture Notes in Computational Science and Engineering*, 91, 01 2013.
- [22] M.J. Gander. Optimized Schwarz methods. *SIAM J. Numer. Anal.*, 44(2):699–731, 2006.
- [23] M.J. Gander, F. Magoulès, and F. Nataf. Optimized Schwarz methods without overlap for the Helmholtz equation. *SIAM J. Sci. Comput.*, 24(1):38–60, 2002.
- [24] M.J. Gander and K. Santugini. Cross-points in domain decomposition methods with a finite element discretization. *Electron. Trans. Numer. Anal.*, 45:219–240, 2016.

- [25] M.J. Gander and Y. Xu. Optimized Schwarz methods for model problems with continuously variable coefficients. *SIAM J. Sci. Comput.*, 38(5):A2964–A2986, 2016.
- [26] M.J. Gander and H. Zhang. A class of iterative solvers for the Helmholtz equation: factorizations, sweeping preconditioners, source transfer, single layer potentials, polarized traces, and optimized Schwarz methods. *SIAM Rev.*, 61(1):3–76, 2019.
- [27] G.H. Golub and C.F. Van Loan. *Matrix computations*. Baltimore, MD: The Johns Hopkins University Press, 2013.
- [28] I. G. Graham, E. A. Spence, and J. Zou. Domain decomposition with local impedance conditions for the Helmholtz equation with absorption. *SIAM J. Numer. Anal.*, 58(5):2515–2543, 2020.
- [29] W. Hackbusch. *Iterative solution of large sparse systems of equations*, volume 95. Cham: Springer, 2016.
- [30] J.-H. Kimn and M. Sarkis. Restricted overlapping balancing domain decomposition methods and restricted coarse problems for the Helmholtz problem. *Comput. Methods Appl. Mech. Eng.*, 196(8):1507–1514, 2007.
- [31] M. Lecouvez. *Iterative methods for domain decomposition without overlap with exponential convergence for the Helmholtz equation*. Theses, Ecole Polytechnique, July 2015.
- [32] F. Magoulès, P. Iványi, and B. H. V. Topping. Non-overlapping Schwarz methods with optimized transmission conditions for the Helmholtz equation. *Comput. Methods Appl. Mech. Engrg.*, 193(45–47):4797–4818, 2004.
- [33] A. Modave, A. Royer, X. Antoine, and C. Geuzaine. A non-overlapping domain decomposition method with high-order transmission conditions and cross-point treatment for Helmholtz problems. *Comput. Methods Appl. Mech. Eng.*, 368:23, 2020. Id/No 113162.
- [34] F. Nataf, F. Rogier, and E De Sturler. Optimal Interface Conditions for Domain Decomposition Methods. Technical Report 301, CMAP Ecole Polytechnique, 1994.
- [35] E. Parolin. *Non-overlapping domain decomposition methods with non-local transmission-operators for harmonic wave propagation problems*. Theses, Institut Polytechnique de Paris, December 2020.
- [36] C. Pechstein. *Finite and boundary element tearing and interconnecting solvers for multiscale problems*, volume 90 of *Lecture Notes in Computational Science and Engineering*. Springer, Heidelberg, 2013.
- [37] W. Rudin. *Functional analysis*. 2nd ed. New York, NY: McGraw-Hill, 2nd ed. edition, 1991.
- [38] A. Toselli and O. Widlund. *Domain decomposition methods—algorithms and theory*, volume 34 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 2005.