

# AUXILIARY SPACE PRECONDITIONERS FOR $C^0$ FINITE ELEMENT APPROXIMATION OF HAMILTON–JACOBI–BELLMAN EQUATIONS WITH CORDES COEFFICIENTS

GUANGWEI GAO    SHUONAN WU\*

**Abstract.** In the past decade, there are many works on the finite element methods for the fully nonlinear Hamilton–Jacobi–Bellman (HJB) equations with Cordes condition. The linearised systems have large condition numbers, which depend not only on the mesh size, but also on the parameters in the Cordes condition. This paper is concerned with the design and analysis of auxiliary space preconditioners for the linearised systems of  $C^0$  finite element discretization of HJB equations [Calcolo, 58, 2021]. Based on the stable decomposition on the auxiliary spaces, we propose both the additive and multiplicative preconditioners which converge uniformly in the sense that the resulting condition number is independent of both the number of degrees of freedom and the parameter  $\lambda$  in Cordes condition. Numerical experiments are carried out to illustrate the efficiency of the proposed preconditioners.

**Key words.** Non-divergence form, Hamilton–Jacobi–Bellman, Cordes condition,  $C^0$  finite element methods, auxiliary space precondition

**1. Introduction.** Let  $\Omega$  be a bounded, open, convex polytopal domain in  $\mathbb{R}^d$ , where  $d = 2, 3$  represent the dimension. In this paper, we are interested in the Hamilton–Jacobi–Bellman (HJB) equations of the following type:

$$(1.1) \quad \sup_{\alpha \in \Lambda} (L^\alpha u - f^\alpha) = 0 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where  $\Lambda$  is a compact metric space, and

$$L^\alpha v := A^\alpha : D^2 v + \mathbf{b}^\alpha \cdot \nabla v - c^\alpha v.$$

Here,  $D^2 u$  and  $\nabla u$  denote the Hessian and gradient of real-valued function  $u$ , respectively. The coefficient  $A^\alpha \in C(\bar{\Omega} \times \Lambda; \mathbb{R}^{d \times d})$  is assumed to be uniformly elliptic, i.e., there exist constants  $\bar{\nu}, \underline{\nu} > 0$  such that

$$(1.2) \quad \underline{\nu} |\boldsymbol{\xi}|^2 \leq \boldsymbol{\xi}^t A^\alpha(x) \boldsymbol{\xi} \leq \bar{\nu} |\boldsymbol{\xi}|^2 \quad \forall \boldsymbol{\xi} \in \mathbb{R}^d, \text{ a.e. in } \Omega, \forall \alpha \in \Lambda.$$

Further,  $\mathbf{b}^\alpha \in C(\bar{\Omega} \times \Lambda; \mathbb{R}^d)$  and  $c^\alpha \geq 0, f^\alpha \in C(\bar{\Omega} \times \Lambda; \mathbb{R})$ .

The HJB equations arise in many applications including stochastic optimal control, game theory, and mathematical finance [11]. In [20, 31], the HJB equations are shown to admit  $H^2$  strong solutions under the following Cordes condition.

**DEFINITION 1.1** (Cordes condition for (1.1)). *The coefficients satisfy that there exist  $\lambda > 0$  and  $\varepsilon \in (0, 1]$  such that*

$$(1.3) \quad \frac{|A^\alpha|^2 + |\mathbf{b}^\alpha|^2/2\lambda + (c^\alpha/\lambda)^2}{(\text{tr } A^\alpha + c^\alpha/\lambda)^2} \leq \frac{1}{d + \varepsilon} \quad \text{a.e. in } \Omega, \forall \alpha \in \Lambda.$$

In the past decade, several studies have been taken on the finite element approximation of  $H^2$  strong solutions of the HJB equations with Cordes coefficients (1.3). The first discontinuous Galerkin (DG) method was proposed in [31], which has been extended to the parabolic HJB equations in [32]. The  $C^0$ -interior penalty DG

---

\*Corresponding author. School of Mathematical Sciences, Peking University, Beijing 100871, China .

methods were developed in [22]. A mixed method based on the stable finite element Stokes spaces was proposed in [12]. Recently, the  $C^0$  (non-Lagrange) finite element method with no stabilization parameter was proposed in [34], where the element is required to be  $C^1$ -continuous at  $(d - 2)$ -dimensional subsimplex, e.g.,  $\mathcal{P}_k$ -Hermite family ( $k \geq 3$ ) in 2D and  $\mathcal{P}_k$ -Argyris family ( $k \geq 5$ ) [21, 9] in 3D. The above discretizations can be naturally applied to the linear elliptic equations in non-divergence form [30, 17, 22, 12, 34]. Other related topics include the unified analysis of DGFEM and  $C^0$ -IPDG [18], and the adaptivity of  $C^0$ -IPDG [7, 19].

For all these discretizations, the discrete well-posedness is analysed under the broken  $H^2$ -norm with possible jump terms across the boundary. This, after linearization, leads to the ill-conditioned systems with condition number  $\mathcal{O}(h^{-4})$  on quasi-uniform meshes, where  $h$  represents the mesh size. Due to the similar performance to the discrete system for fourth-order problems, it is conceivable that the linearised system from HJB equations can be effectively solved by the solvers for fourth-order problems, e.g., geometric multigrid [25, 5, 3, 33, 8] or domain decomposition [38, 6]. In [29], the nonoverlapping domain decomposition preconditioner was studied for the DGFEM discretization of HJB equations.

Traditional geometric multigrid methods depend crucially on the multilevel structures of underlying grids. On unstructured grids, the more user-friendly option is the algebraic multigrid method (AMG) that have been extensively studied for the second-order equations. In [24], the first biharmonic equation was converted to a Poisson system based on the boundary operator proposed in [13]. Under the framework of auxiliary space preconditioning [35], Zhang and Xu [37] proposed a class of optimal solvers based on the auxiliary discretization of mixed form for the fourth-order problems. As a generalization of [25, 24, 26], it works for a variety of conforming and nonconforming finite element discretizations on both convex and nonconvex domains with unstructured triangulation.

The propose of this work is to study the auxiliary space preconditioner to the  $C^0$  finite element discretization of HJB equations. More specifically, the numerical scheme for fully nonlinear HJB equations leads to a discrete nonlinear problem that can be solved iteratively by a semi-smooth Newton method [31, 22, 34]. The linear system obtained from the semi-smooth Newton linearization are generally non-symmetric but coercive. To handle the non-symmetry, the existing GMRES theory [10] will lead to a guaranteed minimum convergence rate with a symmetric FOV-equivalent preconditioner  $P_{\lambda,h}$  that satisfies (3.7). The construction of  $P_{\lambda,h}$  under the auxiliary preconditioning framework follows two steps:

1. Construct appropriate auxiliary spaces and corresponding transfer operators mapping functions from original space to the auxiliary spaces;
2. Devise solvers on auxiliary spaces so that the bounds in (3.7) are uniform with respect to both  $h$  and the parameter  $\lambda$  in the Cordes condition.

Based on the stable decomposition for auxiliary spaces, both additive and multiplicative preconditioners are shown to be efficient and  $\lambda$ -uniform for the linearised system. Further, the preconditioners only involve the Poisson-like solver which can be efficiently solved by AMG with nearly optimal complexity.

In general cases, the auxiliary space preconditioner is additive [35, 37, 14], which usually leads to a stable but relatively large condition number in practical applications. The first contribution of this work is the construction and analysis of a multiplicative preconditioner based on the specific structure of auxiliary spaces. Having a coarse subspace, the symmetrized two-level multiplicative precondition was shown to be positive definite provided that the smoother on the fine level has contraction property

[16]. The condition number estimate of multiplicative precondition at the matrix level can be found in [23]. In this work, we show that the contracted smoother together with the stable decomposition for auxiliary spaces leads to a robust multiplicative preconditioner, which is also numerical verified with better performance than the additive version.

The parameter  $\lambda$  in the Cordes condition balances the diffusion and the constant term. We emphasize that this parameter is not involved in the monotonicity constant (2.1), which makes it possible to consider the preconditioner with uniformity on  $\lambda$ . In this work, we carefully define the norm on the auxiliary space so that the induced preconditioner is uniform with respect to  $\lambda$ . Although the preconditioner is designed for the  $C^0$  finite element approximation, a similar idea can be applied to other discretizations.

The rest of the paper is organized as follows. In Section 2, we establish the notation and state some preliminaries results. In Section 3, we apply the FOV-equivalence preconditioner for the linear system, which can be used to solve non-symmetric systems appearing in applications to the HJB equations. In Section 4, we construct both the additive and multiplicative auxiliary space preconditioners. We also show that the condition numbers of the preconditioned systems are uniformly bounded with the stable decomposition assumption, which is verified in Section 5. Several numerical experiments are presented in Section 6 to illustrate the theoretical results.

For convenience, we use  $C$  to denote a generic positive constant which may depend on  $\Omega$ , share regularity of mesh and polynomial degree, but is independent of the mesh size  $h$ . The notation  $X \lesssim Y$  means  $X \leq CY$ .  $X \simeq Y$  means  $X \lesssim Y$  and  $Y \lesssim X$ .

**2. Preliminaries.** In this section, we first review the  $H^2$  strong solutions to the HJB equations (1.1) under the Cordes condition (1.3). Then we give a brief statement about the  $C^0$  finite element scheme in [34].

Given an integer  $k \geq 0$ , let  $H^k(\Omega)$  and  $H_0^k(\Omega)$  be the usual Sobolev spaces,  $\|\cdot\|_{H^k(\Omega)}$  and  $|\cdot|_{H^k(\Omega)}$  denote the Sobolev norm and semi-norm. We also denote  $V = H^2(\Omega) \cap H_0^1(\Omega)$ . For any Hilbert space  $X$ , we denote  $X'$  for the dual space of  $X$ , and  $\langle \cdot, \cdot \rangle$  for the corresponding dual pair. We also denote  $|\cdot|$  as the Euclidian norm for vectors and the Frobenius norm for matrices.

**2.1.  $H^2$  strong solutions to the HJB equations.** We now invoke the theory of  $H^2$  strong solutions of the HJB equations. In view of the Cordes condition (1.3), for each  $\alpha \in \Lambda$ , define

$$(2.1) \quad \gamma^\alpha := \frac{\text{tr } A^\alpha + c^\alpha/\lambda}{|A^\alpha|^2 + |\mathbf{b}^\alpha|^2/2\lambda + (c^\alpha/\lambda)^2}.$$

And for  $\lambda$  as in (1.3), define a linear operator  $L_\lambda : H^2(\Omega) \rightarrow L^2(\Omega)$  by

$$(2.2) \quad L_\lambda u := \Delta u - \lambda u \quad u \in H^2(\Omega).$$

Next, we define the operator  $F_\gamma : H^2(\Omega) \rightarrow L^2(\Omega)$  by

$$(2.3) \quad F_\gamma[u] := \sup_{\alpha \in \Lambda} \{\gamma^\alpha L^\alpha u - \gamma^\alpha f^\alpha\}.$$

Note that the continuity of data implies  $\gamma^\alpha \in C(\bar{\Omega} \times \Lambda; \mathbb{R})$ . As a consequence, it is readily seen that the HJB equation (1.1) is equivalent to the problem  $F_\gamma[u] = 0$  in  $\Omega$ , and  $u = 0$  on  $\partial\Omega$ . The Cordes condition leads to the following lemma; See [31, Lemma 1] for a proof.

LEMMA 2.1 (property of Cordes condition). *Under the Cordes condition (1.3), for any open set  $U \subset \Omega$  and  $w, v \in H^2(U)$ ,  $z = w - v$ , the following inequality holds a.e. in  $U$ :*

$$(2.4) \quad |F_\gamma[w] - F_\gamma[v] - L_\lambda z| \leq \sqrt{1 - \varepsilon} \sqrt{|D^2 z|^2 + 2\lambda |\nabla z|^2 + \lambda^2 z^2}.$$

Another key ingredient for the well-posedness of (1.1) is the Miranda-Talenti estimate stated as follows.

LEMMA 2.2 (Miranda-Talenti estimate, [15, 20]). *Suppose  $\Omega$  is a bounded convex domain in  $\mathbb{R}^d$ . Then, for any  $v \in V = H^2(\Omega) \cap H_0^1(\Omega)$ ,*

$$(2.5) \quad |v|_{H^2(\Omega)} \leq \|\Delta v\|_{L^2(\Omega)} \leq C|v|_{H^2(\Omega)},$$

where the constant  $C$  depends only on the dimension.

Let the operator  $M : V \rightarrow V'$  be

$$(2.6) \quad \langle M[w], v \rangle := \int_{\Omega} F_\gamma[w] L_\lambda v dx.$$

By using Miranda-Talenti estimate (2.5) and Cordes condition (1.3), one can show the strong monotonicity of  $M$ ,

$$\langle M[v], v \rangle \geq (1 - \sqrt{1 - \varepsilon}) \|v\|_\lambda^2 \quad \forall v \in V,$$

where  $\|v\|_\lambda^2 := \|D^2 v\|_{L^2(\Omega)}^2 + 2\lambda \|\nabla v\|_{L^2(\Omega)}^2 + \lambda^2 \|v\|_{L^2(\Omega)}^2$ . Together with the Lipschitz continuity of  $M$ , the compactness of  $\Lambda$  and the Browder-Minty Theorem [27, Theorem 10.49], one can show the existence and uniqueness of the following problem: Find  $u \in V$  such that

$$(2.7) \quad \langle M[u], v \rangle = 0 \quad \forall v \in V.$$

We refer to [31, Theorem 3] for a detailed proof.

**2.2.  $C^0$  finite element approximations of the HJB equations.** Let  $\mathcal{T}_h$  be a conforming shape regular simplicial triangulation of polytope  $\Omega$  and  $\mathcal{F}_h$  be the set of all faces of  $\mathcal{T}_h$ .  $\mathcal{F}_h^i := \mathcal{F}_h \setminus \partial\Omega$  and  $\mathcal{F}_h^\partial := \mathcal{F}_h \cap \partial\Omega$ . Let  $\mathcal{N}_h$  be the set of all the nodes of  $\mathcal{T}_h$ . Here  $h = \max_{T \in \mathcal{T}_h} h_T$ , where  $h_T$  is the diameter of  $T \in \mathcal{T}_h$ . We also denote  $h_F$  as the diameter of  $F \in \mathcal{F}_h$ . For  $F \in \mathcal{F}_h$  and  $T \in \mathcal{T}_h$ , we use  $(\cdot, \cdot)_T$ , respectively  $\langle \cdot, \cdot \rangle_F$ , to denote the  $L^2$ -inner product over  $T$ , respectively  $F$ .

Following [34], we adopt the  $\mathcal{P}_k$ -Hermite finite elements ( $k \geq 3$ ) in 2D and  $\mathcal{P}_k$ -Argyris finite elements in 3D to solve the HJB equations (1.1). Define the finite element spaces  $V_h$  as

1. For  $d = 2$ , with  $k \geq 3$  (cf. Fig. 1),

$$V_h := \{v \in H_0^1(\Omega) : v|_T \in \mathcal{P}_k(T), \forall T \in \mathcal{T}_h, v \text{ is } C^1 \text{ at all vertices}\},$$

2. For  $d = 3$ , with  $k \geq 5$  (cf. Fig. 2),

$$V_h := \{v \in H_0^1(\Omega) : v|_T \in \mathcal{P}_k(T), \forall T \in \mathcal{T}_h, v \text{ is } C^1 \text{ on all edges, } \\ v \text{ is } C^2 \text{ at all vertices}\},$$




Fig. 1: Degrees of freedom of 2D  $\mathcal{P}_k$  Hermite elements, in the case of  $k = 3$  and  $k = 4$




Fig. 2: Degrees of freedom of 3D  $\mathcal{P}_k$  Argyris elements, in the case of  $k = 5$  and  $k = 6$

where  $\mathcal{P}_k(T)$  denotes set of the polynomials of degree  $k$  on  $T$ .

For each  $F \in \mathcal{F}_h^i$ , we define the tangential Laplace operator  $\Delta_T : H^s(F) \rightarrow H^{s-2}(F)$  as follows, where  $s \geq 2$ . Let  $\{\mathbf{t}_i\}_{i=1}^{d-1}$  be a orthogonal coordinate system on  $F$ . Then, for  $w \in H^s(F)$  define

$$\Delta_T w = \sum_{i=1}^{d-1} \frac{\partial^2}{\partial \mathbf{t}_i^2} w.$$

Next, we define the jump of a vector function  $\mathbf{v}$  on an interior face  $F = \partial T^+ \cap \partial T^-$  as follows:

$$[[\mathbf{v}]]_F := \mathbf{v}^+ \cdot \mathbf{n}^+|_F + \mathbf{v}^- \cdot \mathbf{n}^-|_F,$$

where  $\mathbf{v}^\pm = \mathbf{v}|_{T^\pm}$  and  $\mathbf{n}^\pm$  is the unit outward normal vector of  $T^\pm$ , respectively. For scalar function  $w$  we define

$$[[w]] := w|_{T^+} - w|_{T^-}.$$

The following lemma is critical in the design and analysis of finite element approximation of HJB equations (1.1).

**LEMMA 2.3** (discrete Miranda-Talenti identity, [34]). *Let  $\Omega \subset \mathbb{R}^d$  be a convex polytopal domain and  $\mathcal{T}_h$  be a conforming triangulation. For each  $v_h \in V_h$ , it holds*

that

$$\sum_{T \in \mathcal{T}_h} \|\Delta v_h\|_{L^2(T)}^2 = \sum_{T \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(T)}^2 + 2 \sum_{F \in \mathcal{F}_h^i} \langle \llbracket \nabla v_h \rrbracket, \Delta_T v_h \rangle_F.$$

In light of (2.6), we define the operator  $M_h : V + V_h \rightarrow V'_h$  by

$$\begin{aligned} \langle M_h[w], v_h \rangle &:= \sum_{T \in \mathcal{T}_h} (F_\gamma[w], L_\lambda v_h)_T \\ &\quad - (2 - \sqrt{1 - \varepsilon}) \sum_{F \in \mathcal{F}_h^i} \langle \llbracket \nabla w \rrbracket, \Delta_T v_h - \lambda v_h \rangle_F. \end{aligned}$$

The following finite element scheme is proposed to approximate the solutions to the HJB equations (1.1): Find  $u_h \in V_h$  such that

$$(2.8) \quad \langle M_h u_h, v_h \rangle = 0 \quad \forall v_h \in V_h.$$

We refer to [34] for the well-posedness and approximation property of discrete systems (2.8).

**2.3. Semi-smooth Newton method.** It is shown in [31] that the discretized nonlinear system (2.8) can be solved by a semi-smooth Newton method, which leads to a sequence of discretized linear systems. We summarized the main ideas on semi-smooth Newton here and refer [31] for more details.

Following the discuss in [31], we define the admissible maximizers set for any  $v \in V_h + V$ ,

$$\Lambda[v] := \left\{ \alpha(\cdot) : \Omega \rightarrow \Lambda \mid \begin{array}{l} \alpha(x) \in \arg \max_{\alpha \in \Lambda} (A^\alpha : D_h^2 v + \mathbf{b}^\alpha \cdot \nabla v - c^\alpha v - f^\alpha) \\ \text{measurable} \quad \text{for almost every } x \in \Omega \end{array} \right\},$$

where  $D_h^2 v$  denotes the broken Hessian of  $v$ . As shown in [31, Lemma 9 & Theorem 10], the set  $\Lambda[v]$  is not empty for any  $v \in V + V_h$ .

The semi-smooth Newton method is now stated as follows. Start by choosing an initial iterate  $u_h^0 \in V_h$ . Then, for each nonnegative integer  $j$ , given the previous iterate  $u_h^j \in V_h$ , choose an  $\alpha_j \in \Lambda[u_h^j]$ . Next the function  $f^{\alpha_j} : \Omega \mapsto \mathbb{R}$  is defined by  $f^{\alpha_j} : x \rightarrow f^{\alpha_j(x)}(x)$ ; the functions  $A^{\alpha_j}$ ,  $\mathbf{b}^{\alpha_j}$ ,  $c^{\alpha_j}$  and  $\gamma^{\alpha_j}$  are defined in a similar way. Then find the solution  $u_h^{j+1} \in V_h$  of the linearised system

$$(2.9) \quad b_{\lambda,h}^j(u_h^{j+1}, v_h) = \sum_{T \in \mathcal{T}_h} (\gamma^{\alpha_j} f^{\alpha_j}, \Delta v_h)_T \quad \forall v_h \in V_h,$$

where the bilinear form  $b_{\lambda,h}^j : V_h \times V_h \rightarrow \mathbb{R}$  is defined by

$$\begin{aligned} b_{\lambda,h}^j(w_h, v_h) &:= \sum_{T \in \mathcal{T}_h} (\gamma^{\alpha_j} L^{\alpha_j} w_h, L_\lambda v_h)_T \\ &\quad - (2 - \sqrt{1 - \varepsilon}) \sum_{F \in \mathcal{F}_h^i} \langle \llbracket \nabla w_h \rrbracket, \Delta_T v_h - \lambda v_h \rangle_F. \end{aligned}$$

Following [34], we define inner product on  $V_h$  as

$$(w_h, v_h)_{\lambda,h} := \sum_{T \in \mathcal{T}_h} (D^2 w_h, D^2 v_h)_T + 2\lambda(\nabla w_h, \nabla v_h)_\Omega + \lambda^2(w_h, v_h)_\Omega,$$

and the norm  $\|v_h\|_{\lambda,h}^2 := (v_h, v_h)_{\lambda,h}$ . It is also shown in [34] that the bilinear forms  $b_{\lambda,h}^j$  are uniformly coercive and bounded on  $V_h$  with norm  $\|\cdot\|_{\lambda,h}$ , with constants independent of iterates. Since the preconditioners in this work take advantage on the coercivity and boundedness of  $b_{\lambda,h}^j$ , we summarize the relevant results in the following lemma (see [34, Lemmas 4.1 & 4.2] for a detailed proof).

LEMMA 2.4 (coercivity and boundedness of bilinear form). *For every  $w_h, v_h \in V_h$ , we have*

$$\begin{aligned} b_{\lambda,h}^j(v_h, w_h) &\leq C \|v_h\|_{\lambda,h} \|w_h\|_{\lambda,h}, \\ b_{\lambda,h}^j(v_h, v_h) &\geq (1 - \sqrt{1 - \varepsilon}) \|v_h\|_{\lambda,h}^2. \end{aligned}$$

Here, the constant  $C$  depends only on  $\Omega$ , shape regularity of the grid and polynomial degree  $k$ .

**3. FOV-equivalent preconditioners for GMRES methods.** The preconditioned GMRES (PGMRES) methods are among the most effective iterative methods for non-symmetric linear systems arising from discretizations of PDEs. Our study will start by discussing PGMRES methods in an operator form. Let  $G : \mathcal{X} \rightarrow \mathcal{X}'$  be a linear operator which may be non-symmetric or indefinite, defined on a finite dimensional space  $\mathcal{X}$ , and  $g$  be a given functional in its dual space  $\mathcal{X}'$ . The linear equation considered here is of the following form

$$(3.1) \quad Gx = g.$$

Let  $(\cdot, \cdot)_M$  be an inner product on  $\mathcal{X}$ , and  $P : \mathcal{X}' \rightarrow \mathcal{X}$  be the preconditioner. The PGMRES method for solving (3.1) is stated as follows: Begin with an initial guess  $x_0 \in \mathcal{X}$  and denote  $r_0 = g - Gx_0$  the initial residual, the  $k$ -th steps of PGMRES method seeks  $x_k$  such that

$$x_k = \underset{\tilde{x}_k \in \mathcal{K}_k(PG, Pr_0) + x_0}{\operatorname{argmin}} \|PG(x - \tilde{x}_k)\|_M,$$

where  $\mathcal{K}_k(PG, Pr_0)$  is the Krylov subspace of dimension  $k$  generated by  $PG$  and  $Pr_0$ .

In the semi-smooth Newton steps, the discrete linear equations (2.9) have a common form: Find  $u_h \in V_h$  such that

$$(3.2) \quad b_{\lambda,h}(u_h, v_h) = f_h(v_h) \quad \forall v_h \in V_h,$$

where we shall omit to denote independence of the bilinear form  $b_{\lambda,h}$  and of the right-hand side  $f_h$  on the iteration number of the semi-smooth Newton method. Define the operator  $B_{\lambda,h} : V_h \rightarrow V_h'$  by

$$(3.3) \quad \langle B_{\lambda,h} u_h, v_h \rangle := b_{\lambda,h}(u_h, v_h) \quad \forall u_h, v_h \in V_h,$$

then the discrete system (3.2) can be written in an operator form, namely

$$(3.4) \quad B_{\lambda,h} u_h = f_h.$$

Moreover, a general operator  $P_{\lambda,h} : V_h' \rightarrow V_h$  is used to denote the preconditioner. Given an inner product  $(\cdot, \cdot)_{M_{\lambda,h}}$ , we can estimate the convergence rate of the PGMRES method. It is proved in [10, 28] that if  $u_h^m$  is the  $m$ -iteration of PGMRES method and  $u_h$  is the exact solution of (3.4), then

$$\frac{\|P_{\lambda,h} B_{\lambda,h}(u_h - u_h^m)\|_{M_{\lambda,h}}}{\|P_{\lambda,h} B_{\lambda,h}(u_h - u_h^0)\|_{M_{\lambda,h}}} \leq \left(1 - \frac{\gamma^2}{\Gamma^2}\right)^{m/2},$$

where

$$(3.5) \quad \gamma \leq \frac{(v_h, P_{\lambda,h} B_{\lambda,h} v_h)_{M_{\lambda,h}}}{(v_h, v_h)_{M_{\lambda,h}}}, \quad \frac{\|P_{\lambda,h} B_{\lambda,h} v_h\|_{M_{\lambda,h}}}{\|v_h\|_{M_{\lambda,h}}} \leq \Gamma \quad \forall v_h \in V_h.$$

Therefore, we conclude that as long as we find an operator  $P_{\lambda,h}$  and a proper inner product  $(\cdot, \cdot)_{M_{\lambda,h}}$  such that condition (3.5) is satisfied with constants  $\gamma$  and  $\Gamma$  independent of the discretization parameter  $h$  and the Cordes condition parameter  $\lambda$ , then  $P_{\lambda,h}$  is a uniform preconditioner for GMRES method. Such preconditioners are usually referred to as *FOV-equivalent preconditioners*. In what follows, we always take

$$P_{\lambda,h} \text{ to be an SPD operator, and } M_{\lambda,h} = P_{\lambda,h}^{-1}.$$

Next, we give a general principle for constructing  $P_{\lambda,h}$ . Define an SPD operator  $A_{\lambda,h} : V_h \rightarrow V_h'$  by

$$(3.6) \quad \langle A_{\lambda,h} w_h, v_h \rangle := (w_h, v_h)_{\lambda,h} \quad \forall w_h, v_h \in V_h.$$

Recalling Lemma 2.4 (coercivity and boundedness of bilinear form),  $b_{\lambda,h}(\cdot, \cdot)$  is coercive and bounded on  $V_h$  with the inner product  $(\cdot, \cdot)_{\lambda,h}$ . It is therefore that an efficient preconditioner for  $A_{\lambda,h}$  can also be used as an FOV-preconditioner for the GMRES algorithm applied to  $B_{\lambda,h}$ , which is shown in the following lemma.

LEMMA 3.1 (FOV-equivalent preconditioner). *Let  $A_{\lambda,h}$  and  $B_{\lambda,h}$  be the operators defined in (3.6) and (3.3), respectively. If an SPD operator  $P_{\lambda,h} : V_h' \rightarrow V_h$  satisfies that*

$$(3.7) \quad \alpha \langle P_{\lambda,h}^{-1} v_h, v_h \rangle \leq \langle A_{\lambda,h} v_h, v_h \rangle \leq \beta \langle P_{\lambda,h}^{-1} v_h, v_h \rangle \quad \forall v_h \in V_h,$$

with constants  $\alpha, \beta$  independent of both  $\lambda$  and  $h$ , then  $P_{\lambda,h}$  is a uniform FOV-equivalent preconditioner of  $B_{\lambda,h}$ .

*Proof.* From (3.3), (3.6) and Lemma 2.4 (coercivity and boundedness of bilinear form), we see that for any  $u_h, v_h \in V_h$

$$(3.8a) \quad (1 - \sqrt{1 - \varepsilon}) \langle A_{\lambda,h} u_h, u_h \rangle \leq \langle B_{\lambda,h} u_h, u_h \rangle$$

$$(3.8b) \quad \langle B_{\lambda,h} u_h, v_h \rangle \leq C \langle A_{\lambda,h} u_h, u_h \rangle^{1/2} \langle A_{\lambda,h} v_h, v_h \rangle^{1/2},$$

where  $C$  is independent of both  $\varepsilon$  and  $\lambda$ .

Recalling  $M_{\lambda,h} := P_{\lambda,h}^{-1}$ , then for any  $u_h \in V_h$ , we have

$$\begin{aligned} \|P_{\lambda,h} B_{\lambda,h} u_h\|_{P_{\lambda,h}^{-1}} &= \sup_{v_h \in V_h, v_h \neq 0} \frac{(P_{\lambda,h} B_{\lambda,h} u_h, v_h)_{P_{\lambda,h}^{-1}}}{(v_h, v_h)_{P_{\lambda,h}^{-1}}^{1/2}} \\ &\leq \beta^{1/2} \sup_{v_h \in V_h, v_h \neq 0} \frac{\langle B_{\lambda,h} u_h, v_h \rangle}{\langle A_{\lambda,h} v_h, v_h \rangle^{1/2}} \quad (\text{by (3.7)}) \\ &\leq C \beta^{1/2} \langle A_{\lambda,h} u_h, u_h \rangle^{1/2} \quad (\text{by (3.8b)}) \\ &\leq C \beta \|u_h\|_{P_{\lambda,h}^{-1}}, \quad (\text{by (3.7)}) \end{aligned}$$



which yields the second inequality of (3.5) with  $\Gamma = C\beta$ . On the other side, we have

$$\begin{aligned} (u_h, P_{\lambda,h} B_{\lambda,h} u_h)_{P_{\lambda,h}^{-1}} &= \langle B_{\lambda,h} u_h, u_h \rangle \\ &\geq (1 - \sqrt{1 - \varepsilon}) \langle A_{\lambda,h} u_h, u_h \rangle \quad (\text{by (3.8a)}) \\ &\geq (1 - \sqrt{1 - \varepsilon}) \alpha (u_h, u_h)_{P_{\lambda,h}^{-1}}, \quad (\text{by (3.7)}) \end{aligned}$$

which yields the first inequality of (3.5) with  $\gamma = (1 - \sqrt{1 - \varepsilon})\alpha$ .  $\square$

**4. Fast auxiliary space preconditioners.** In this section, we construct both additive and multiplicative auxiliary space preconditioners for SPD operator  $A_{\lambda,h}$ . From Lemma 3.1 (FOV-equivalent preconditioner), those preconditioners can be applied to the discrete linearised systems (2.9) arising from each semi-smooth Newton step of solving the HJB equations.

**4.1. Space decomposition.** For the purpose of constructing auxiliary space preconditioners, we give the following space decomposition of  $V_h$  as

$$V_h = V_h + \Pi_0 V_0,$$

where the auxiliary space  $V_0 \subset H_0^1(\Omega)$  denotes the continuous piecewise linear element space on  $\mathcal{T}_h$  with homogeneous Dirichlet boundary condition, and  $\Pi_0 : V_0 \rightarrow V_h$  is a linear injective map which will be defined later. As a result, the induced operator  $A_0 := \Pi_0' A_{\lambda,h} \Pi_0 : V_0 \rightarrow V_0'$  is also SPD, and hence we define  $\|\cdot\|_{A_0}^2 := \langle A_0 \cdot, \cdot \rangle$  on  $V_0$ . We also introduce a projection  $P_0 : V_h \rightarrow V_0$  by

$$\langle A_0 P_0 v_h, w_h \rangle := (v_h, \Pi_0 w_h)_{\lambda,h} \quad \forall v_h \in V_h, w_h \in V_0.$$

A direct calculation shows the following identity

$$(4.1) \quad \Pi_0' A_{\lambda,h} = A_0 P_0.$$

*Smoother and norm on  $V_0$ .* Define the discrete Laplacian operator  $-\Delta_h : V_0 \rightarrow V_0$  by

$$(4.2) \quad (-\Delta_h w_h, v_h)_\Omega := (\nabla w_h, \nabla v_h)_\Omega \quad \forall w_h, v_h \in V_0.$$

Then, the smoother on  $V_0$ , denoted by  $R_0 : V_0' \rightarrow V_0$ , is defined by

$$(4.3) \quad \langle R_0^{-1} u_h, v_h \rangle = (\lambda u_h - \Delta_h u_h, \lambda v_h - \Delta_h v_h)_\Omega \quad \forall u_h, v_h \in V_0.$$

Note that for any given  $f_h \in V_0'$ ,  $u_h = R_0 f_h \in V_0$  can be obtained by solving the following two discrete Poisson-like equations

$$(4.4a) \quad \langle f_h, v_h \rangle = \lambda (z_h, v_h)_\Omega + (\nabla z_h, \nabla v_h)_\Omega \quad \forall v_h \in V_0,$$

$$(4.4b) \quad (z_h, w_h)_\Omega = \lambda (u_h, w_h)_\Omega + (\nabla u_h, \nabla w_h)_\Omega \quad \forall w_h \in V_0.$$

It can be shown that the above two equations can be solved within  $\mathcal{O}(N \log N)$  operations, where  $N$  denotes the number of degrees of freedom. We will give a detailed explanation in Remark 4.10 (computational complexity). The smoother  $R_0$  induces a norm on  $V_0$ , i.e.,  $\|\cdot\|_{R_0^{-1}}^2 := \langle R_0^{-1} \cdot, \cdot \rangle$ . By using (4.2) and (4.3), it is straightforward to show that

$$\|v_h\|_{R_0^{-1}}^2 = \|\Delta_h v_h\|_{L^2(\Omega)}^2 + 2\lambda \|\nabla v_h\|_{L^2(\Omega)}^2 + \lambda^2 \|v_h\|_{L^2(\Omega)}^2.$$

The relationship between  $\|\cdot\|_{R_0^{-1}}$  and  $\|\cdot\|_{A_0}$  is shown in the following lemma, whose proof is postponed to Section 5.

LEMMA 4.1 (spectral equivalence of  $R_0$ ). *Let  $R_0$  be the operator defined in (4.3) and  $A_0 = \Pi'_0 A_{\lambda,h} \Pi_0$ . Then,*

$$\|v_0\|_{R_0^{-1}} \simeq \|v_0\|_{A_0} \quad \forall v_0 \in V_0,$$

with hidden constants independent of both  $\lambda$  and  $h$ .

*Smoother on  $V_h$ .* Let  $R_h$  denote the Gauss-Seidel smoother for  $A_{\lambda,h}$  and  $\bar{R}_h$  be the symmetric Gauss-Seidel smoother, i.e.,

$$I - \bar{R}_h A_{\lambda,h} = (I - R'_h A_{\lambda,h})(I - R_h A_{\lambda,h}).$$

We also define  $\|\cdot\|_{\bar{R}_h^{-1}}^2 := \langle \bar{R}_h^{-1} \cdot, \cdot \rangle$  as a norm on  $V_h$ . Note that  $A_{\lambda,h}$  is an SPD operator, thus  $R_h$  has the following contraction property

$$(4.5) \quad \|I - R_h A_{\lambda,h}\|_{\lambda,h} < 1.$$

*Transfer operator.* We now give the definition of  $\Pi_0 : V_0 \rightarrow V_h$ . To this end, we first give some notation on the degrees of freedom of finite element space  $V_h$ . We denote the degrees of freedom as

$$\mathcal{N}_\alpha(\varphi) = \int_{D_\alpha} \nabla^{k_\alpha}(\varphi)(\mathbf{t}_1, \dots, \mathbf{t}_{k_\alpha})$$

where  $D_\alpha$  is the domain of the integral with respect to the degree of freedom,  $\int_{D_\alpha}$  denotes the integral average on  $D_\alpha$ . In general,  $D_\alpha$  is a subsimplex of the triangulation. When  $D_\alpha$  is a point, the average of the integral is reduced to the evaluation on the point.  $\mathbf{t}_1, \dots, \mathbf{t}_{k_\alpha}$  are  $k_\alpha$  identical or different unit vectors to denote the direction of the derivative where  $k_\alpha = 0, 1, 2$ . When  $k_\alpha = 1$  only one direction is involved for the derivative, and the direction is denoted by  $\mathbf{t}_\alpha$ . Let  $\varphi_\alpha$  be the nodal basis function corresponding to  $\mathcal{N}_\alpha$ . Define  $\omega_\alpha := \bigcup\{T : \overset{\circ}{T} \cap \text{supp}(\varphi_\alpha) \neq \emptyset, T \in \mathcal{T}_h\}$ ,  $\#\omega_\alpha := \#\{T : \overset{\circ}{T} \cap \text{supp}(\varphi_\alpha) \neq \emptyset, T \in \mathcal{T}_h\}$  and  $h_\alpha = \max_{T \subset \omega_\alpha} h_T$ . We are now ready to give the definition of  $\Pi_0$  as follows: For any  $p_h \in V_0$

$$(4.6) \quad \begin{aligned} \mathcal{N}_\alpha(\Pi_0 p_h) &= \frac{1}{\#\omega_\alpha} \sum_{T \subset \omega_\alpha} \int_{D_\alpha} \partial_{\mathbf{n}}(p_h|_T)(\mathbf{n} \cdot \mathbf{t}_\alpha) \\ &\quad \text{when } D_\alpha \subset \partial\Omega \text{ and } k_\alpha = 1, \\ \mathcal{N}_\alpha(\Pi_0 p_h) &= \frac{1}{\#\omega_\alpha} \sum_{T \subset \omega_\alpha} \mathcal{N}_\alpha(p_h|_T), \text{ else,} \end{aligned}$$

where  $\mathbf{n}$  is the unit outer normal vector of  $\partial\Omega$ . We note that the degrees of freedom corresponding to the second-order derivative vanish since  $p_h$  is piecewise linear.

The general theory of auxiliary space preconditioning simplifies the analysis of preconditioners to the verification of the following two key assumptions.

*Assumption 4.2 (stable decomposition).* There exists a uniform constant  $c_0$  independent of both  $\lambda$  and  $h$ , such that for any  $v \in V_h$ , there exist  $v_h \in V_h$  and  $v_0 \in V_0$  satisfy

$$(4.7a) \quad v = v_h + \Pi_0 v_0,$$

$$(4.7b) \quad \|v_h\|_{\bar{R}_h^{-1}}^2 + \|v_0\|_{R_0^{-1}}^2 \leq c_0^2 \|v\|_{\lambda,h}^2.$$

*Assumption 4.3* (boundedness). There exist uniform constants  $c_1$  and  $c_2$  independent of both  $\lambda$  and  $h$  such that

$$(4.8a) \quad \|v_0\|_{A_0} \leq c_1 \|v_0\|_{\bar{R}_0^{-1}} \quad \forall v_0 \in V_0,$$

$$(4.8b) \quad \|v_h\|_{\lambda,h} \leq c_2 \|v_h\|_{\bar{R}_h^{-1}} \quad \forall v_h \in V_h.$$

**4.2. Additive preconditioner.** Firstly, we introduce the additive preconditioner  $P_a : V'_h \rightarrow V_h$  as

$$(4.9) \quad P_a := \bar{R}_h + \Pi_0 R_0 \Pi'_0.$$

The following theorem plays a fundamental role in the theory of auxiliary space preconditioning [35].

**THEOREM 4.4** (spectral equivalence of additive preconditioner). *Let  $P_a : V'_h \rightarrow V_h$  be the preconditioner defined in (4.9). If Assumption 4.2 (stable decomposition) and Assumption 4.3 (boundedness) hold, then we have*

$$c_0^{-2}(v_h, v_h)_{\lambda,h} \leq (P_a A_{\lambda,h} v_h, v_h)_{\lambda,h} \leq (c_1^2 + c_2^2)(v_h, v_h)_{\lambda,h} \quad \forall v_h \in V_h,$$

*That is,  $P_a$  is a uniform spectral equivalence preconditioner of  $A_{\lambda,h}$ .*

In light of the above theorem and Lemma 3.1 (FOV-equivalent preconditioner), one can see that  $P_a$  is a uniform FOV-equivalent preconditioner of  $B_{\lambda,h}$  as long as Assumption 4.2 (stable decomposition) and Assumption 4.3 (boundedness) are verified. We postpone those verifications to Section 5.

*Remark 4.5* (additive preconditioner with Jacobi smoother). Let  $D_h^{-1}$  be the Jacobi smoother of  $A_{\lambda,h}$ . From the norm equivalence between  $\bar{R}_h$  and  $D_h^{-1}$  [36, Lemma 4.6], the additive preconditioner

$$\tilde{P}_a := D_h^{-1} + \Pi_0 R_0 \Pi'_0$$

is also a uniform spectral equivalence preconditioner of  $A_{\lambda,h}$ .

*Remark 4.6* (additive preconditioner with scaled parameter). When implementing the additive preconditioners, a positive parameter  $\omega$  is usually introduced to balance the two components, namely,

$$(4.10) \quad P_a := \bar{R}_h + \omega \Pi_0 R_0 \Pi'_0.$$

A proper choice of  $\omega$  may lead to a better preconditioning performance of  $P_a$  in practice.

**4.3. Multiplicative preconditioner.** We introduce the multiplicative preconditioner  $P_m : V'_h \rightarrow V_h$  by

$$(4.11) \quad I - P_m A_{\lambda,h} := (I - R_h A_{\lambda,h})(I - \Pi_0 R_0 \Pi'_0 A_{\lambda,h})(I - R'_h A_{\lambda,h}).$$

Let  $\hat{R}_0 = A_0^{-1}$  be the exact solver on  $V_0$ . To analyse the multiplicative preconditioner  $P_m$ , we introduce an auxiliary multiplicative preconditioner  $\hat{P}_m : V'_h \rightarrow V_h$  by

$$(4.12) \quad I - \hat{P}_m A_{\lambda,h} := (I - R_h A_{\lambda,h})(I - \Pi_0 \hat{R}_0 \Pi'_0 A_{\lambda,h})(I - R'_h A_{\lambda,h}).$$

We emphasize that  $\hat{P}_m$  is never computed but is useful for theoretical purposes in Theorem 4.8 (spectral equivalence of multiplicative preconditioner), which can be divided into two steps: (i) The spectral equivalence between  $P_m A_{\lambda,h}$  and  $\hat{P}_m A_{\lambda,h}$  by using Lemma 4.1 (spectral equivalence of  $R_0$ ); (ii) Estimate of  $\hat{P}_m A_{\lambda,h}$  by two-level convergence results. For the second step, let  $E := (I - R_h A_{\lambda,h})(I - \Pi_0 \hat{R}_0 \Pi'_0 A_{\lambda,h})$  be the error propagation operator the two-level method corresponding to  $\hat{P}_m$ . Since the solver on  $V_0$  is exact, the convergence rate can be obtained in the following theorem. We refer to [36, Theorem 5.3] for more details.

**THEOREM 4.7** (two-level convergence rate, [36]). *The following identity holds*

$$\|E\|_{\lambda,h}^2 = 1 - \frac{1}{K(V_0)},$$

where

$$K(V_0) := \max_{v \in V_h} \min_{v_0 \in V_0} \frac{\|v - \Pi_0 v_0\|_{R_h^{-1}}^2}{\|v\|_{\lambda,h}^2}.$$

Note that the identity (4.1) implies that  $\hat{R}_0 \Pi'_0 A_{\lambda,h} = P_0$ . For any  $v_h, w_h \in V_h$ , we have

$$\begin{aligned} (\Pi_0 P_0 v_h, w_h)_{\lambda,h} &= \langle P_0 v_h, \Pi'_0 A_{\lambda,h} w_h \rangle = \langle P_0 v_h, A_0 P_0 w_h \rangle = (v_h, \Pi_0 P_0 w_h)_{\lambda,h}, \\ (R'_h A_{\lambda,h} v_h, w_h)_{\lambda,h} &= \langle A_{\lambda,h} v_h, R_h A_{\lambda,h} w_h \rangle = (v_h, R_h A_{\lambda,h} w_h)_{\lambda,h}, \end{aligned}$$

which means that  $I - \Pi_0 P_0$  and  $I - R'_h A_{\lambda,h}$  are respectively the dual operators of  $I - \Pi_0 P_0$  and  $I - R_h A_{\lambda,h}$  under the inner product  $(\cdot, \cdot)_{\lambda,h}$ . As a consequence,

$$(4.13) \quad \|E\|_{\lambda,h}^2 = \|(I - R_h A_{\lambda,h})(I - \Pi_0 P_0)\|_{\lambda,h}^2 = \|(I - \Pi_0 P_0)(I - R'_h A_{\lambda,h})\|_{\lambda,h}^2.$$

Moreover, under the Assumption 4.2 (stable decomposition), we have  $K(V_0) \leq c_0^2$  and hence  $\|E\|_{\lambda,h}^2 \leq 1 - \frac{1}{c_0^2}$ , which leads to the following theorem.

**THEOREM 4.8** (spectral equivalence of multiplicative preconditioner). *Let  $R_h$  be the Gauss-Seidel smoother for  $A_{\lambda,h}$ . Under the Assumption 4.2 (stable decomposition), the multiplicative preconditioner  $P_m$  defined in (4.11) satisfies*

$$(P_m A_{\lambda,h} v_h, v_h)_{\lambda,h} \simeq (v_h, v_h)_{\lambda,h} \quad \forall v_h \in V_h,$$

with hidden constants independent of both  $\lambda$  and  $h$ . That is,  $P_m$  is a uniform spectral equivalence preconditioner of  $A_{\lambda,h}$ .

*Proof. Step (i):* For any  $v \in V_h$ , denote  $w = (I - R'_h A_{\lambda,h})v$ . By the definition of  $P_m$  (4.11), we have

$$\begin{aligned} (P_m A_{\lambda,h} v, v)_{\lambda,h} &= (v, v)_{\lambda,h} - ((I - \Pi_0 R_0 \Pi'_0 A_{\lambda,h})w, w)_{\lambda,h} \\ &= \|v\|_{\lambda,h}^2 - \|w\|_{\lambda,h}^2 + (\Pi_0 R_0 \Pi'_0 A_{\lambda,h} w, w)_{\lambda,h} \\ &= \|v\|_{\lambda,h}^2 - \|w\|_{\lambda,h}^2 + (R_0 A_0 P_0 w, P_0 w)_{A_0}, \end{aligned}$$

where we use the identity (4.1) in the last step. Similarly, for  $\hat{P}_m$  defined in (4.12), we have

$$(\hat{P}_m A_{\lambda,h} v, v)_{\lambda,h} = \|v\|_{\lambda,h}^2 - \|w\|_{\lambda,h}^2 + (P_0 w, P_0 w)_{A_0},$$

since  $\hat{R}_0 = A_0^{-1}$ . Invoking Lemma 4.1 (spectral equivalence of  $R_0$ ), we have

$$(4.14) \quad C_1(P_0w, P_0w)_{A_0} \leq (R_0A_0P_0w, P_0w)_{A_0} \leq C_2(P_0w, P_0w)_{A_0},$$

where  $C_1, C_2$  are constants independent of  $\lambda$  and  $h$ . Then,

$$(4.15) \quad \min\{1, C_1\}(\hat{P}_m A_{\lambda,h}v, v)_{\lambda,h} \leq (P_m A_{\lambda,h}v, v)_{\lambda,h} \leq \max\{1, C_2\}(\hat{P}_m A_{\lambda,h}v, v)_{\lambda,h}.$$

Here, we use the contraction property of  $R_h$ , namely  $\|v\|_{\lambda,h}^2 - \|w\|_{\lambda,h}^2 \geq (1 - \|I - R'_h A_{\lambda,h}\|_{\lambda,h}^2)\|v\|_{\lambda,h}^2 > 0$ .

*Step (ii):* In light of (4.15), we only need to show

$$(\hat{P}_m A_{\lambda,h}v, v)_{\lambda,h} \simeq (v, v)_{\lambda,h} \quad \forall v \in V_h,$$

with hidden constants independent of  $\lambda$  and  $h$ . From the identity (4.1), we see that  $(I - \Pi_0 P_0) = (I - \Pi_0 P_0)^2$  since  $P_0 \Pi_0 = A_0^{-1} \Pi'_0 A_{\lambda,h} \Pi_0 = I$ . Then, we obtain the upper bound of  $\hat{P}_m A_{\lambda,h}$ :

$$\begin{aligned} (\hat{P}_m A_{\lambda,h}v, v)_{\lambda,h} &= \|v\|_{\lambda,h}^2 - ((I - \Pi_0 P_0)w, w)_{\lambda,h} \\ &= \|v\|_{\lambda,h}^2 - ((I - \Pi_0 P_0)w, (I - \Pi_0 P_0)w)_{\lambda,h} \leq \|v\|_{\lambda,h}^2. \end{aligned}$$

Next, we estimate the lower bound of  $\hat{P}_m A_{\lambda,h}$  by Theorem 4.7 (two-level convergence rate) and Assumption 4.2 (stable decomposition),

$$\begin{aligned} (\hat{P}_m A_{\lambda,h}v, v)_{\lambda,h} &= \|v\|_{\lambda,h}^2 - \|(I - \Pi_0 P_0)(I - R'_h A_{\lambda,h})v\|_{\lambda,h}^2 \\ &\geq \left(1 - \|(I - \Pi_0 P_0)(I - R'_h A_{\lambda,h})\|_{\lambda,h}^2\right) \|v\|_{\lambda,h}^2 \\ &= \left(1 - \|(I - R_h A_{\lambda,h})(I - \Pi_0 P_0)\|_{\lambda,h}^2\right) \|v\|_{\lambda,h}^2 \quad (\text{by (4.13)}) \\ &= \left(1 - \|E\|_{\lambda,h}^2\right) \|v\|_{\lambda,h}^2 \\ &= \frac{1}{K(V_0)} \|v\|_{\lambda,h}^2 \geq c_0^{-2} \|v\|_{\lambda,h}^2. \end{aligned}$$

Combining *Step (i)* and *Step (ii)*, we obtain

$$\min\{1, C_1\}c_0^{-2} \|v\|_{\lambda,h}^2 \leq (P_m A_{\lambda,h}v, v)_{\lambda,h} \leq \max\{1, C_2\} \|v\|_{\lambda,h}^2.$$

The proof is thus complete.  $\square$

*Remark 4.9.* In the proof of Theorem 4.8 (spectral equivalence of multiplicative preconditioner), Assumption 4.3 (boundedness) is not directly used. That is because Lemma 4.1 (spectral equivalence of  $R_0$ ) leads to the boundedness on coarse space (4.8a), and the boundedness on fine space (4.8b) is a direct consequence of the contraction property (4.5) of Gauss–Seidel smoother  $R_h$ .

*Remark 4.10* (computational complexity). We now discuss the computational complexity of the action of preconditioner  $P_a$  (4.9) and  $P_m$  (4.11). Let  $N_h$  be the number of degrees of freedom,  $N_p$  be the number of interior points of the grids. Since the transfer operator  $\Pi_0$  (4.6) is local, the action except  $R_0$  can be done within  $\mathcal{O}(N_h)$  operations.

Invoking the definition of  $R_0$  in (4.3), we can see for any given  $f_h \in V'_0$ ,  $u_h = R_0 f_h$  can be obtained by solving two discrete Poisson-like equations (4.4). The computational complexity of  $(\lambda I - \Delta_h)^{-1}$  with classic geometric multigrid methods was shown

to be optimal for  $\lambda = \mathcal{O}(1)$  [2] and for arbitrary  $\lambda > 0$  [1]. For unstructured shape-regular grids with  $\lambda = \mathcal{O}(1)$ , the computational complexity turns to be  $\mathcal{O}(N_p \log N_p)$  by constructing of an auxiliary coarse grid hierarchy where the geometric multigrid can be applied [14]. Therefore, the nearly optimal computational complexity for arbitrary  $\lambda > 0$  on unstructured grids is to be expected by combining the techniques from [1] and [14].

*Remark 4.11* (Implement of action  $R_0$ ). Let  $\{\psi_i\}_{i=1}^{N_p}$  be the nodal basis functions of  $V_0$ . Denote  $\mathbf{A} = ((\nabla\psi_j, \nabla\psi_i)_\Omega) \in \mathbb{R}^{N_p \times N_p}$ ,  $\mathbf{M} = ((\psi_j, \psi_i)_\Omega) \in \mathbb{R}^{N_p \times N_p}$  as the stiffness and mass matrix, respectively. For any  $f_h \in V_0'$ , let  $\mathbf{f} \in \mathbb{R}^{N_p}$  be its vector representation, i.e.,  $(\mathbf{f})_i = \langle f_h, \psi_i \rangle$ . Then  $u_h = R_0 f_h$  can be obtain by solving the following two linear systems successively.

$$\begin{aligned} \mathbf{f} &= (\lambda \mathbf{M} + \mathbf{A}) \mathbf{z}, \\ \mathbf{M} \mathbf{z} &= (\lambda \mathbf{M} + \mathbf{A}) \mathbf{u}, \end{aligned}$$

where  $\mathbf{u} \in \mathbb{R}^{N_p}$  is the vector representation of  $u_h$ , i.e.,  $u_h = \sum_{i=1}^{N_p} (\mathbf{u})_i \psi_i$ .

**5. Analysis of the auxiliary space preconditioners.** In this section, we shall verify Assumption 4.2 (stable decomposition) and Assumption 4.3 (boundedness), then show the proof of Lemma 4.1 (spectral equivalence of  $R_0$ ). For this propose, we first show some properties about the space  $V_0$  introduced in Section 4. We refer to [4, 37] for details on those results.

LEMMA 5.1 (see [37], Lemma 3.6 and [4]). *Let  $\Pi_0 : V_0 \rightarrow V_h$  be the interpolation operator defined in (4.6). For any  $p_h \in V_0$ , it holds that*

$$(5.1) \quad \|\Pi_0 p_h\|_{L^2(\Omega)} \lesssim \|p_h\|_{L^2(\Omega)} \quad \text{and} \quad |\Pi_0 p_h|_{H^1(\Omega)} \lesssim |p_h|_{H^1(\Omega)}.$$

Moreover, we have

$$(5.2) \quad \sum_{T \in \mathcal{T}_h} h_T^{-4} \|p_h - \Pi_0 p_h\|_{L^2(T)}^2 \lesssim \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left\| \frac{\partial p_h}{\partial \mathbf{n}_F} \right\|^2 ds \quad \forall p_h \in V_0.$$

LEMMA 5.2. *Suppose  $\Omega \subset \mathbb{R}^d$  is a convex polytopal domain. Let  $\Delta_h$  be the discrete Laplacian operator defined in (4.2). Then it holds that*

$$(5.3) \quad \|\Delta_h p_h\|_{L^2(\Omega)}^2 \simeq \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left\| \frac{\partial p_h}{\partial \mathbf{n}_F} \right\|^2 ds \quad \forall p_h \in V_0.$$

*Proof.* (5.3) can be obtained by combining a similar technique in [37, Lemma 3.1] and the elliptic regularity in convex polytopal domain in  $\mathbb{R}^d$  [15, Chapter 3].  $\square$

For any  $v_h \in V_h$ , define nodal interpolation  $I_h : V_h \rightarrow V_0$  as

$$(5.4) \quad I_h v_h(\mathbf{x}) = v_h(\mathbf{x}) \quad \forall \mathbf{x} \in \mathcal{N}_h.$$

According to standard polynomial approximation theory [4], we have the following lemma.

LEMMA 5.3 (see [4]). *For any  $v_h \in V_h$ , it holds that*

$$(5.5) \quad \|I_h v_h\|_{L^2(\Omega)} \lesssim \|v_h\|_{L^2(\Omega)} \quad \text{and} \quad |I_h v_h|_{H^1(\Omega)} \lesssim |v_h|_{H^1(\Omega)}.$$

Moreover, for any  $T \in \mathcal{T}_h$  we have

$$(5.6) \quad \|v_h - I_h v_h\|_{L^2(T)} \lesssim h_T^2 |v_h|_{H^2(T)} \quad \text{and} \quad \left\| \frac{\partial(v_h - I_h v_h)}{\partial \mathbf{n}} \right\|_{L^2(\partial T)} \lesssim h_T^{1/2} |v_h|_{H^2(T)}.$$

LEMMA 5.4. *For any  $v \in V_h$ , it holds that*

$$(5.7) \quad \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v - \Pi_0 I_h v\|_{L^2(T)}^2 \lesssim \sum_{T \in \mathcal{T}_h} (1 + \lambda h_T^2)^2 \|D^2 v\|_{L^2(T)}^2.$$

*Proof.* It suffices to show:

$$\sum_{T \in \mathcal{T}_h} h_T^{-2\ell} \|v - \Pi_0 I_h v\|_{L^2(T)}^2 \lesssim \sum_{T \in \mathcal{T}_h} h_T^{-2\ell+4} \|D^2 v\|_{L^2(T)}^2,$$

for  $\ell = 0, 1, 2$ . The case when  $\ell = 2$  is proved in [37, Lemma 3.7]. When  $\ell = 0, 1$ , it can be proved by the same arguments in [37, Lemma 3.7].  $\square$

Next lemma gives a equivalence form of  $\|\cdot\|_{\bar{R}_h^{-1}}$  which will be used in the verification of Assumption 4.2 (stable decomposition) and Assumption 4.3 (boundedness).

LEMMA 5.5 (norm equivalence of  $\bar{R}_h$ ). *Let  $\mathcal{T}_h$  be a conforming shape regular triangulation of  $\Omega$ . Let  $D_h^{-1}$  and  $R_h$  be the Jacobi and Gauss-Seidel smoother for  $A_{\lambda,h}$ , respectively. Then, we have*

$$(5.8) \quad \|v_h\|_{\bar{R}_h^{-1}}^2 \simeq \|v_h\|_{D_h}^2 \simeq \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v_h\|_{L^2(T)}^2 \quad \forall v_h \in V_h.$$

*Proof.* By classical theory of iterative method [36, Lemma 4.6], the symmetric Gauss-Seidel smoother and the Jacobi smoother are spectral equivalent for sparse SPD operator, namely  $\|v_h\|_{\bar{R}_h^{-1}}^2 \simeq \|v_h\|_{D_h}^2$ . Standard scaling argument [4] gives that

$$(5.9a) \quad \|v_h\|_{L^2(T)}^2 \simeq \sum_{\omega_\alpha \supset T} h_T^{2k_\alpha+2} (\mathcal{N}_\alpha(v_h))^2 \quad \forall T \in \mathcal{T}_h, v_h \in V_h,$$

$$(5.9b) \quad \sum_{T \in \mathcal{T}_h} \int_T |D^j \varphi_\alpha|^2 dx \simeq h_\alpha^{2k_\alpha+2-2j} \quad j = 0, 1, 2,$$

where  $\varphi_\alpha$  is the nodal basis function corresponding to the degree of freedom  $\mathcal{N}_\alpha(\cdot)$ . Combining (5.9a) and (5.9b), we have

$$(5.10) \quad \begin{aligned} \sum_{T \in \mathcal{T}_h} h_T^{-4} \|v_h\|_{L^2(T)}^2 &\simeq \sum_{T \in \mathcal{T}_h} h_T^{-4} \sum_{\omega_\alpha \supset T} h_T^{2k_\alpha+2} (\mathcal{N}_\alpha(v_h))^2 \\ &\simeq \sum_{\alpha} (\#\omega_\alpha) h_\alpha^{2k_\alpha-2} (\mathcal{N}_\alpha(v_h))^2 \\ &\simeq \sum_{\alpha} \left( \sum_{T \in \mathcal{T}_h} \int_T |D^2 \varphi_\alpha|^2 dx \right) (\mathcal{N}_\alpha(v_h))^2. \end{aligned}$$

A similar argument leads to

$$(5.11) \quad \sum_{T \in \mathcal{T}_h} h_T^{-2} \|v_h\|_{L^2(T)}^2 \simeq \sum_{\alpha} \left( \|\nabla \varphi_\alpha\|_{L^2(\Omega)}^2 \right) (\mathcal{N}_\alpha(v_h))^2,$$

and

$$(5.12) \quad \sum_{T \in \mathcal{T}_h} \|v_h\|_{L^2(T)}^2 \simeq \sum_{\alpha} \left( \|\varphi_\alpha\|_{L^2(\Omega)}^2 \right) (\mathcal{N}_\alpha(v_h))^2.$$

Multiplying (5.10), (5.11) and (5.12) respectively by 1,  $2\lambda$  and  $\lambda^2$ , then summing these equations, we obtain

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v_h\|_{L^2(T)}^2 \\ & \simeq \sum_{\alpha} \left( \sum_{T \in \mathcal{T}_h} \int_T |D^2 \varphi_{\alpha}|^2 + 2\lambda |\nabla \varphi_{\alpha}|^2 + \lambda^2 \varphi_{\alpha}^2 dx \right) (\mathcal{N}_{\alpha}(v_h))^2 = \|v_h\|_{D_h}^2, \end{aligned}$$

which yields the norm equivalence (5.8).  $\square$

With the help of above lemmas, we are now ready to verify of Assumption 4.2 (stable decomposition) and Assumption 4.3 (boundedness).

**THEOREM 5.6** (verification of boundedness). *There exist constants  $c_1, c_2$  independent of  $\lambda$  and  $h$ , such that*

$$(5.13a) \quad \|v_0\|_{A_0} \leq c_1 \|v_0\|_{R_0^{-1}} \quad \forall v_0 \in V_0,$$

$$(5.13b) \quad \|v_h\|_{\lambda, h} \leq c_2 \|v_h\|_{\bar{R}_h^{-1}} \quad \forall v_h \in V_h.$$

*Proof.* (5.13b) follows from the standard inverse estimate and Lemma 5.5 (norm equivalence of  $\bar{R}_h$ ),

$$\begin{aligned} \|v_h\|_{\lambda, h}^2 &= \sum_{T \in \mathcal{T}_h} \|D^2 v_h\|_{L^2(T)}^2 + 2\lambda \|\nabla v_h\|_{L^2(\Omega)}^2 + \lambda^2 \|v_h\|_{L^2(\Omega)}^2 \\ &\lesssim \sum_{T \in \mathcal{T}_h} (h_T^{-4} + 2\lambda h_T^{-2} + \lambda^2) \|v_h\|_{L^2(T)}^2 \\ &= \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v_h\|_{L^2(T)}^2 \simeq \|v_h\|_{\bar{R}_h^{-1}}^2. \end{aligned}$$

Now turn to (5.13a), by combining Lemma 5.1, Lemma 5.2 and inverse estimate, we have

$$\begin{aligned} (5.14) \quad \sum_{T \in \mathcal{T}_h} \|D^2 \Pi_h v_0\|_{L^2(T)}^2 &= \sum_{T \in \mathcal{T}_h} \|D^2 (\Pi_h v_0 - v_0)\|_{L^2(T)}^2 \\ &\lesssim \sum_{T \in \mathcal{T}_h} h_T^{-4} \|\Pi_h v_0 - v_0\|_{L^2(T)}^2 \quad (\text{by inverse estimate}) \\ &\lesssim \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left[ \frac{\partial v_0}{\partial \mathbf{n}_F} \right]^2 ds \quad (\text{by Lemma 5.1}) \\ &\lesssim \|\Delta_h v_0\|_{L^2(\Omega)}^2. \quad (\text{by Lemma 5.2}) \end{aligned}$$

Then, (5.13a) follows from (5.14) and boundedness of  $\Pi_0$  in Lemma 5.1

$$\begin{aligned} (5.15) \quad \|v_0\|_{A_0}^2 &= \sum_{T \in \mathcal{T}_h} \|D^2 \Pi_0 v_0\|_{L^2(T)}^2 + 2\lambda \|\nabla \Pi_0 v_0\|_{L^2(\Omega)}^2 + \lambda^2 \|\Pi_0 v_0\|_{L^2(\Omega)}^2 \\ &\lesssim \|\Delta_h v_0\|_{L^2(\Omega)}^2 \quad (\text{by (5.14)}) \\ &\quad + 2\lambda \|\nabla v_0\|_{L^2(\Omega)}^2 + \lambda^2 \|v_0\|_{L^2(\Omega)}^2 \quad (\text{by Lemma 5.1}) \\ &= \|v_0\|_{R_0^{-1}}^2. \end{aligned}$$

This completes the proof.  $\square$



THEOREM 5.7 (verification of stable decomposition). *For any  $v \in V_h$ , there exist  $v_h \in V_h$  and  $v_0 \in V_0$  such that*

$$(5.16a) \quad v = v_h + \Pi_0 v_0,$$

$$(5.16b) \quad \|v_h\|_{\bar{R}_h}^2 + \|v_0\|_{R_0}^2 \leq c_0^2 \|v\|_{\lambda, h}^2,$$

where  $c_0$  is a constant independent with  $\lambda$  and  $h$ .

*Proof.* Recall the nodal interpolation  $I_h : V_h \rightarrow V_0$  defined in (5.4). For any  $v \in V_h$ , take  $v_0 = I_h v$  and  $v_h = v - \Pi_0 I_h v$  so that (5.16a) is satisfied. It follows from Lemma 5.4, Lemma 5.5 (norm equivalence of  $\bar{R}_h$ ) and inverse estimate that

$$\begin{aligned} \|v_h\|_{\bar{R}_h}^2 &\simeq \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v_h\|_{L^2(T)}^2 \\ &= \sum_{T \in \mathcal{T}_h} (h_T^{-2} + \lambda)^2 \|v - \Pi_0 I_h v\|_{L^2(T)}^2 \\ &\lesssim \sum_{T \in \mathcal{T}_h} (1 + \lambda h_T^2)^2 \|D^2 v\|_{L^2(T)}^2 && \text{(by Lemma 5.4)} \\ &\lesssim \sum_{T \in \mathcal{T}_h} \|D^2 v\|_{L^2(T)}^2 + 2\lambda \|\nabla v\|_{L^2(\Omega)}^2 + \lambda^2 \|v\|_{L^2(\Omega)}^2 && \text{(by inverse estimate)} \\ &= \|v\|_{\lambda, h}^2. \end{aligned}$$

On the other hand, combining Lemma 5.2 and the approximation property of  $I_h$  in Lemma 5.3, we have

$$(5.17) \quad \begin{aligned} \|\Delta_h I_h v\|_{L^2(\Omega)}^2 &\lesssim \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left[ \frac{\partial I_h v}{\partial \mathbf{n}_F} \right]^2 ds \\ &\lesssim \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left[ \frac{\partial (I_h v - v)}{\partial \mathbf{n}_F} \right]^2 ds + \sum_{F \in \mathcal{F}_h^i} h_F^{-1} \int_F \left[ \frac{\partial v}{\partial \mathbf{n}_F} \right]^2 ds \\ &\lesssim \sum_{T \in \mathcal{T}_h} \|D^2 v\|_{L^2(T)}^2. \end{aligned}$$

Here, in the last step, the standard scaling argument [4] gives that

$$h_F^{-1} \int_F \left[ \frac{\partial v}{\partial \mathbf{n}_F} \right]^2 ds \lesssim \sum_{T \in \{T^+, T^-\}} \|D^2 v\|_{L^2(T)}^2,$$

holds for any interior face  $F = \partial T^+ \cap \partial T^-$ , where the  $C^0$ -continuity at face and  $C^1$ -continuity at  $(d-2)$ -dimensional subsimplex guarantee that the piecewise linear function on  $\omega_F = T^+ \cup T^-$  has to be a linear function on the  $\omega_F$ .

Combining the boundedness of  $I_h$  in Lemma 5.2 and (5.17), we get

$$\begin{aligned} \|v_0\|_{R_0}^2 &= \|\Delta_h I_h v\|_{L^2(\Omega)}^2 + 2\lambda \|\nabla I_h v\|_{L^2(\Omega)}^2 + \lambda^2 \|I_h v\|_{L^2(\Omega)}^2 \\ &\lesssim \sum_{T \in \mathcal{T}_h} \|D^2 v\|_{L^2(T)}^2 && \text{(by (5.17))} \\ &\quad + 2\lambda \|\nabla v\|_{L^2(\Omega)}^2 + \lambda^2 \|v\|_{L^2(\Omega)}^2 && \text{(by (5.5))} \\ &= \|v\|_{\lambda, h}^2. \end{aligned}$$

The proof is thus complete.  $\square$

By the similar arguments in Theorem 5.6 (verification of boundedness) and Theorem 5.7 (verification of stable decomposition), we are now ready to give the proof of Lemma 4.1 (spectral equivalence of  $R_0$ ).

*Proof of Lemma 4.1 (spectral equivalence of  $R_0$ ).* Note that  $v_0 = I_h \Pi_h v_0$  for any  $v_0 \in V_0$ , then

$$\begin{aligned} \|v_0\|_{R_0^{-1}}^2 &= \|\Delta_h I_h \Pi_h v_0\|_{L^2(\Omega)}^2 + 2\lambda \|\nabla I_h \Pi_h v_0\|_{L^2(\Omega)}^2 + \lambda^2 \|I_h \Pi_h v_0\|_{L^2(\Omega)}^2 \\ &\lesssim \sum_{T \in \mathcal{T}_h} \|D^2 \Pi_h v_0\|_{L^2(T)}^2 && \text{(similar to (5.17))} \\ &\quad + 2\lambda \|\nabla \Pi_h v_0\|_{L^2(\Omega)}^2 + \lambda^2 \|\Pi_h v_0\|_{L^2(\Omega)}^2 && \text{(by (5.5))} \\ &= \|\Pi_h v_0\|_{\lambda, h}^2 = \|v_0\|_{A_0}^2. \end{aligned}$$

The other direction has been proved in (5.15), whence we obtain the spectral equivalence  $\|v_0\|_{A_0} \simeq \|v_0\|_{R_0^{-1}}$ .  $\square$

**6. Numerical experiments.** In this section, we present numerical experiments to illustrate the performance of PFMRES preconditioners for solving both linear and nonlinear problems.


Denote  $\kappa(\cdot)$  for the condition number of operator or matrix, and DOF for the number of degrees of freedom. On the fine level, we use Gauss-Seidel method for  $A_{\lambda, h}$  with three iterations as the smoother on  $V_h$ , which shares the similar properties as the Gauss-Seidel smoother  $R_h$ . For the actor of subspace smoother  $R_0$ , we apply algebraic multigrid method (AMG) with stop criterion  $\|\mathbf{b} - \mathbf{A}\mathbf{x}\|_2 / \|\mathbf{b}\|_2 \leq 10^{-8}$  for a linear system  $\mathbf{A}\mathbf{x} = \mathbf{b}$ .

**6.1. Preconditioning effect of  $P_a$  and  $P_m$  for  $A_{\lambda, h}$ .** We test the theoretical results in Theorem 4.4 (spectral equivalence of additive preconditioner) and Theorem 4.8 (spectral equivalence of multiplicative preconditioner) by examining the condition number of  $P_m A_{\lambda, h}$  and  $P_a A_{\lambda, h}$ . To showcase the flexibility of the preconditioner on non-uniform grids, we illustrate the performance of preconditioners on a sequence of graded bisection grids  $\{\mathcal{T}_\ell\}_{\ell \in \mathbb{N}_0}$  with grading factor 1/2 on  $\Omega = (-1, 1)^2$ , see Fig. 3. More specifically, we mark the elements which satisfy  $|T| > C(\|x_T\|_{\ell_2} - 1/2)^2 / \#\mathcal{T}_\ell$ , where  $x_T$  is barycenter of the element  $T$ ,  $\#\mathcal{T}_\ell$  is the number of elements in  $\mathcal{T}_\ell$ . In the experiment, we set  $C = 1000$ . Further, in the case of additive preconditioners, we apply the scaled form (4.10) in Remark 4.6 (additive preconditioner with scaled parameter) with  $\omega = 1/10$ .

The resulting condition numbers for additive and multiplicative preconditioners at different bisection levels are listed respectively in Table 1 and Table 2. We observe that both  $P_m$  and  $P_a$  are uniform preconditioners with respect to both  $\lambda$  and DOF, which is in agreement with the theoretical results in Theorem 4.8 (spectral equivalence of multiplicative preconditioner) and Theorem 4.4 (spectral equivalence of additive preconditioner). We also observe that the multiplicative preconditioners perform better than additive ones.

**6.2. Uniform preconditioning for the linearised problems.** In the second experiment, we consider the linearised problems in the semi-smooth Newton steps, i.e., elliptic equations in non-divergence form:

$$(6.1) \quad A : D^2 u + \mathbf{b}^\theta \cdot \nabla u - c^\theta u = f^\theta,$$



(a) Level 3: DOF = 2,387      (b) Level 5: DOF = 4,467      (c) Level 7: DOF = 10,027

 Fig. 3: Graded grids at different levels for Experiment 6.1, initial grid size  $h_0 = 1/8$ .

Table 1: Condition number of additive preconditioning for Experiment 6.1.

DOF	$\kappa(P_a A_{\lambda,h})$						
	$\lambda = 10^{-3}$	$\lambda = 10^{-2}$	$\lambda = 10^{-1}$	$\lambda = 1$	$\lambda = 10$	$\lambda = 10^2$	$\lambda = 10^3$
3,147	1.64e2	1.64e2	1.60e2	1.40e2	9.41e1	5.43e1	2.48e1
4,467	1.65e2	1.64e2	1.60e2	1.40e2	9.49e1	6.43e1	3.42e1
6,587	1.61e2	1.60e2	1.57e2	1.35e2	9.22e1	7.09e1	4.52e1
10,027	1.61e2	1.61e2	1.57e2	1.36e2	9.62e1	8.04e1	5.49e1
15,927	1.62e2	1.62e2	1.58e2	1.36e2	9.95e1	8.78e1	6.46e1

Table 2: Condition number of multiplicative preconditioning for Experiment 6.1.

DOF	$\kappa(P_m A_{\lambda,h})$						
	$\lambda = 10^{-3}$	$\lambda = 10^{-2}$	$\lambda = 10^{-1}$	$\lambda = 1$	$\lambda = 10$	$\lambda = 10^2$	$\lambda = 10^3$
3,147	5.76	5.76	5.76	5.75	5.65	5.21	4.16
4,467	5.76	5.76	5.75	5.75	5.68	5.42	4.77
6,587	5.63	5.63	5.63	5.62	5.56	5.39	4.99
10,027	6.03	6.03	6.03	6.03	6.02	5.94	5.59
15,927	6.07	6.07	6.07	6.07	6.06	6.00	5.75

on the domain  $\Omega = (-1, 1)^2$ . The coefficients are set to be

$$(6.2) \quad A = \begin{pmatrix} 2 & \frac{x_1 x_2}{|x_1 x_2|} \\ \frac{x_1 x_2}{|x_1 x_2|} & 2 \end{pmatrix} \quad \mathbf{b}^\theta = \sqrt{\theta}(x_1, x_2)^T \quad c^\theta = 3\theta,$$

where  $\theta$  is a parameter so that the  $\lambda$  in Cordes condition differs with varying  $\theta$ . Let exact solution be

$$(6.3) \quad u(x_1, x_2) = (x_1 e^{1-|x_1|} - x_1)(x_2 e^{1-|x_2|} - x_2),$$

the right hand side  $f^\theta$  is directly calculated from the equation (6.1). For any given  $\theta > 0$ , we may set  $\lambda = \theta$ , which yields

$$(6.4) \quad \frac{|A|^2 + |\mathbf{b}^\theta|^2/2\lambda + (c^\theta/\lambda)^2}{(\text{Tr } A + c^\theta/\lambda)^2} = \frac{10 + 1/2(x_1^2 + x_2^2) + 9}{(4 + 3)^2} \leq \frac{20}{49},$$

which means that the Cordes condition is satisfied for  $\varepsilon = 9/20$ . The PGMRES method is applied to solve the linear system arising from the discretization of (6.1) with varying parameter  $\theta > 0$ . We stop the iteration when the relative residual is smaller than  $10^{-6}$ . The iteration numbers for PGMRES are shown in Tables 3 and 4. Similarly, a better performance of multiplicative preconditioner is observed.

Table 3: The iteration steps of additive preconditioning for Experiment 6.2.

DOF	$\lambda = 10^{-3}$	$\lambda = 10^{-2}$	$\lambda = 10^{-1}$	$\lambda = 1$	$\lambda = 10^1$	$\lambda = 10^2$	$\lambda = 10^3$
	steps	steps	steps	steps	steps	steps	steps
5,055	120	119	116	117	99	71	36
20,351	133	132	132	130	114	95	59
81,663	137	137	136	134	118	109	72
32,7167	138	137	135	135	118	117	80

Table 4: The iteration steps of multiplicative preconditioning for Experiment 6.2.

DOF	$\lambda = 10^{-3}$	$\lambda = 10^{-2}$	$\lambda = 10^{-1}$	$\lambda = 1$	$\lambda = 10^1$	$\lambda = 10^2$	$\lambda = 10^3$
	steps	steps	steps	steps	steps	steps	steps
5,055	28	28	28	28	28	27	24
20,351	26	26	26	26	25	25	29
81,663	25	24	24	24	23	23	28
32,7167	23	23	23	22	21	21	26

**6.3. Application to the HJB equations.** In this experiment, we solve the nonlinear HJB equations (1.1) on the domain  $\Omega = (0, 1)^2$ . Following [31], we take  $\Lambda = [0, \pi/3] \times \text{SO}(2)$ , where  $\text{SO}(2)$  is the set of 2 rotation matrices. The coefficients are given by  $\mathbf{b}^\alpha = 0$ ,  $c^\alpha = \pi^2$ , and

$$(6.5) \quad A^\alpha = \frac{1}{2}\sigma^\alpha(\sigma^\alpha)^T, \quad \sigma^\alpha = R^T \begin{pmatrix} 1 & \sin \theta \\ 0 & \cos \theta \end{pmatrix}, \quad \alpha = (\theta, R) \in \Lambda.$$

We choose  $f^\alpha = \sqrt{3}\sin^2\theta/\pi^2 + g$ ,  $g$  independent of  $\alpha$  such that the exact solution of the HJB equations (1.1) is  $u(x_1, x_2) = \exp(x_1x_2)\sin(\pi x_1)\sin(\pi x_2)$ . That is

$$(6.6) \quad g = \sup_{\alpha \in \Lambda} \{A^\alpha : D^2u - c^\alpha u - \sqrt{3}\sin^2\theta/\pi^2\}.$$

In the semi-smooth Newton algorithm, the initial guess is  $u_h^0 = 0$ . For the discrete linearised systems arising from each Newton step, the multiplicative preconditioners is applied. We compute the average number of PGMERS iterations of per Newton step which require to reduce the residual norm below a relative tolerance of  $10^{-4}$ . Convergence of the Newton method was determined by requiring a step-increment  $L^2$ -norm below  $10^{-6}$ . These tolerances are chosen to balance the different sources of error originating from discretization.

The numbers of semi-smooth Newton iterations and average PGMRES iterations are listed in Table 5. As can be observed from [31, 34], the semi-smooth Newton algorithm convergences fast (within eight iterations in the numerical experiment). In

Table 5: Average PGMRES iterations (Newton steps)

DOF	$h$	Average PGMRES iterations (Newton steps)
71	1/4	14 (6)
303	1/8	18 (6)
1,247	1/16	18 (6)
5,055	1/32	18 (7)
20,351	1/64	18 (8)

each Newton step, we apply the PGMRES with multiplicative preconditioner due to its better performance than the additive one. Based on the results shown in the Table 5, we can conclude that our multiplicative preconditioner is also effective and robust in the application to the HJB equations.

**Acknowledgments.** The authors would like to express their gratitude to Prof. Jun Hu in Peking University for his helpful discussions.

## REFERENCES

- [1] J. BRAMBLE, J. PASCIAK, AND P. VASSILEVSKI, *Computational scales of sobolev norms with application to preconditioning*, Mathematics of Computation, 69 (2000), pp. 463–480.
- [2] J. BRAMBLE, J. PASCIAK, AND J. XU, *Parallel multilevel preconditioners*, Mathematics of Computation, 55 (1990), pp. 1–22.
- [3] S. BRENNER, *Convergence of nonconforming multigrid methods without full elliptic regularity*, Mathematics of computation, 68 (1999), pp. 25–53.
- [4] S. BRENNER AND R. SCOTT, *The Mathematical Theory of Finite Element Methods*, vol. 15, Springer Science & Business Media, 2007.
- [5] S. C. BRENNER, *An optimal-order nonconforming multigrid method for the biharmonic equation*, SIAM journal on numerical analysis, 26 (1989), pp. 1124–1138.
- [6] S. C. BRENNER, *A two-level additive Schwarz preconditioner for nonconforming plate elements*, Numerische Mathematik, 72 (1996), pp. 419–447.
- [7] S. C. BRENNER AND E. L. KAWECKI, *Adaptive  $C^0$  interior penalty methods for Hamilton–Jacobi–Bellman equations with Cordes coefficients*, Journal of Computational and Applied Mathematics, (2020), p. 113241.
- [8] C. CARSTENSEN AND J. HU, *Hierarchical argyris finite element method for adaptive and multigrid algorithms*, Computational Methods in Applied Mathematics, (2021).
- [9] S. H. CHRISTIANSEN, J. HU, AND K. HU, *Nodal finite element de Rham complexes*, Numerische Mathematik, 139 (2018), pp. 411–446.
- [10] H. C. ELMAN, *Iterative methods for large, sparse, nonsymmetric systems of linear equations*, PhD thesis, Yale University New Haven, Conn, 1982.
- [11] W. H. FLEMING AND H. M. SONER, *Controlled Markov processes and viscosity solutions*, vol. 25, Springer Science & Business Media, 2006.
- [12] D. GALLISTL AND E. SÜLI, *Mixed finite element approximation of the Hamilton–Jacobi–Bellman equation with Cordes coefficients*, SIAM Journal on Numerical Analysis, 57 (2019), pp. 592–614.
- [13] R. GLOWINSKI AND O. PIRONNEAU, *Numerical methods for the first biharmonic equation and for the two-dimensional Stokes problem*, SIAM review, 21 (1979), pp. 167–212.
- [14] L. GRASEDYCK, L. WANG, AND J. XU, *A nearly optimal multigrid method for general unstructured grids*, Numerische Mathematik, 134 (2016), pp. 637–666.
- [15] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, SIAM, 2011.
- [16] M. HOLST AND S. VANDEWALLE, *Schwarz methods: to symmetrize or not to symmetrize*, SIAM Journal on Numerical Analysis, 34 (1997), pp. 699–722.
- [17] E. L. KAWECKI, *A DGFEM for nondivergence form elliptic equations with Cordes coefficients on curved domains*, Numerical Methods for Partial Differential Equations, 35 (2019), pp. 1717–1744.
- [18] E. L. KAWECKI AND I. SMEARS, *Unified analysis of discontinuous Galerkin and  $C^0$ -interior*

- penalty finite element methods for Hamilton–Jacobi–Bellman and Isaacs equations*, arXiv preprint arXiv:2006.07202, (2020).
- [19] E. L. KAWECKI AND I. SMEARS, *Convergence of adaptive discontinuous Galerkin and  $C^0$ -interior penalty finite element methods for Hamilton–Jacobi–Bellman and Isaacs equations*, Foundations of Computational Mathematics, (2021).
  - [20] A. MAUGERI, D. K. PALAGACHEV, AND L. G. SOFTOVA, *Elliptic and Parabolic Equations with Discontinuous Coefficients*, vol. 109, WILEY-VCH Verlag GmbH & Co., 2000.
  - [21] M. NEILAN, *Discrete and conforming smooth de Rham complexes in three dimensions*, Mathematics of Computation, 84 (2015), pp. 2059–2081.
  - [22] M. NEILAN AND M. WU, *Discrete Miranda-Talenti estimates and applications to linear and nonlinear PDEs*, Journal of Computational and Applied Mathematics, 356 (2019), pp. 358–376.
  - [23] Y. NOTAY AND A. NAPOV, *Further comparison of additive and multiplicative coarse grid correction*, Applied Numerical Mathematics, 65 (2013), pp. 53–62.
  - [24] P. PEISKER, *On the numerical solution of the first biharmonic equation*, ESAIM: Mathematical Modelling and Numerical Analysis, 22 (1988), pp. 655–676.
  - [25] P. PEISKER AND D. BRAESS, *A conjugate gradient method and a multigrid algorithm for Morley’s finite element approximation of the biharmonic equation*, Numerische Mathematik, 50 (1987), pp. 567–586.
  - [26] P. PEISKER, W. RUST, AND E. STEIN, *Iterative solution methods for plate bending problems: Multigrid and preconditioned CG algorithm*, SIAM journal on numerical analysis, 27 (1990), pp. 1450–1465.
  - [27] M. RENARDY AND R. C. ROGERS, *An Introduction to Partial Differential Equations*, vol. 13, Springer Science & Business Media, 2006.
  - [28] Y. SAAD, *Iterative Methods for Sparse Linear Systems*, SIAM, 2003.
  - [29] I. SMEARS, *Nonoverlapping domain decomposition preconditioners for discontinuous Galerkin approximations of Hamilton–Jacobi–Bellman equations*, Journal of Scientific Computing, 74 (2018), pp. 145–174.
  - [30] I. SMEARS AND E. SÜLI, *Discontinuous Galerkin finite element approximation of nondivergence form elliptic equations with Cordes coefficients*, SIAM Journal on Numerical Analysis, 51 (2013), pp. 2088–2106.
  - [31] I. SMEARS AND E. SÜLI, *Discontinuous Galerkin finite element approximation of Hamilton–Jacobi–Bellman equations with Cordes coefficients*, SIAM Journal on Numerical Analysis, 52 (2014), pp. 993–1016.
  - [32] I. SMEARS AND E. SÜLI, *Discontinuous Galerkin finite element methods for time-dependent Hamilton–Jacobi–Bellman equations with Cordes coefficients*, Numerische Mathematik, 133 (2016), pp. 141–176.
  - [33] R. STEVENSON, *An analysis of nonconforming multigrid methods, leading to an improved method for the Morley element*, Mathematics of computation, 72 (2003), pp. 55–81.
  - [34] S. WU,  *$C^0$  finite element approximations of linear elliptic equations in non-divergence form and Hamilton–Jacobi–Bellman equations with Cordes coefficients*, Calcolo, 58 (2021).
  - [35] J. XU, *The auxiliary space method and optimal multigrid preconditioning techniques for unstructured grids*, Computing, 56 (1996), pp. 215–235.
  - [36] J. XU AND L. ZIKATANOV, *Algebraic multigrid methods*, Acta Numerica, 26 (2017), pp. 591–721.
  - [37] S. ZHANG AND J. XU, *Optimal solvers for fourth-order PDEs discretized on unstructured grids*, SIAM Journal on Numerical Analysis, 52 (2014), pp. 282–307.
  - [38] X. ZHANG, *Multilevel Schwarz methods for the biharmonic Dirichlet problem*, SIAM Journal on Scientific Computing, 15 (1994), pp. 621–644.