

**PERIODONTITIS AND PREECLAMPSIA IN PREGNANCY: A SYSTEMATIC  
REVIEW AND META-ANALYSIS**

**Quynh-Anh LE<sup>1</sup>, Rahena AKHTER<sup>2</sup>, Kimberly Mathieu COULTON<sup>2</sup>, Ngoc Truong  
Nhu VO<sup>3</sup>, Le Thi Yen DUONG<sup>4</sup>, Hoang Viet NONG<sup>4</sup>, Albert YAACOUB<sup>5</sup>, George  
CONDOUS<sup>6</sup>, Joerg EBERHARD<sup>7</sup>, Ralph NANAN<sup>8</sup>**

<sup>1</sup> PhD candidate, DDS, School of Dentistry and the Charles Perkins Center, Faculty of  
Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia.

<sup>2</sup> PhD, School of Dentistry and the Charles Perkins Center, Faculty of Medicine and Health,  
The University of Sydney, Sydney, New South Wales, Australia.

<sup>3</sup> Associate Professor, Department of Pediatric Dentistry, School of Odonto-Stomatology,  
Hanoi Medical University, Hanoi, Vietnam.

<sup>4</sup> DDS, School of Odonto-Stomatology, Hanoi Medical University, Hanoi, Vietnam.

<sup>5</sup> MPH, Nepean Centre for Oral Health, Nepean hospital, Kingswood, New South Wales,  
Australia.

<sup>6</sup> Associate Professor, Acute Gynaecology, Early Pregnancy, and Advanced Endoscopy  
Surgery Unit, Nepean Hospital, Sydney Medical School Nepean, University of Sydney,  
Sydney, Australia.

<sup>7</sup> Professor, School of Dentistry and the Charles Perkins Center, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia.

<sup>8</sup> Professor, Sydney Medical School Nepean and Charles Perkins Center Nepean, The University of Sydney, Sydney, New South Wales, Australia.

## CONFLICT OF INTEREST

The authors report no conflict of interest.

\* Corresponding author    Dr Quynh-Anh Le

Address                      School of Dentistry, Faculty of Medicine and Health, The  
University of Sydney, Sydney, New South Wales, Australia.

Telephone                  +61415076584

Email address              [Qule7436@uni.sydney.edu.au](mailto:Qule7436@uni.sydney.edu.au)

Total word count of abstract: 231

Total word count of main text: 6192

## 40 ABSTRACT

41 **Objectives:** A conflicting body of evidence suggests localized periodontal inflammation to  
 42 spread systemically during pregnancy inducing adverse pregnancy outcomes. This systematic  
 43 review and meta-analysis aims to specifically evaluate the relationship between periodontitis  
 44 and preeclampsia.

45 **Data sources:** Electronic searches were carried out in Medline, Pubmed, Embase, LiLacs,  
 46 Cochrane Controlled Clinical Trial Register, CINAHL, ClinicalTrials.gov and Google  
 47 Scholar with no restrictions on year of publication.

48 **Study eligibility criteria:** We identified and selected observational case-control and cohort  
 49 studies that analyzed the association between periodontal disease and preeclampsia.

50 **Study appraisal and synthesis methods:** This meta-analysis was conducted according to  
 51 PRISMA guidelines and MOOSE checklist. Pooled odds ratios, mean difference, and 95%  
 52 confidence intervals were calculated using the random effect model. Heterogeneity was  
 53 tested with Cochran's Q statistic.

54 **Results:** Thirty studies including six cohort and twenty-four case-control studies were  
 55 selected. Periodontitis was significantly associated with increased risk for preeclampsia (OR  
 56 3.18, 95% CI 2.26 – 4.48,  $p < 0.00001$ ), especially in a subgroup analysis including cohort  
 57 studies (OR 4.19, 95% CI 2.23 – 7.87,  $p < 0.00001$ ). The association was even stronger in a  
 58 subgroup analysis with lower middle-income countries (OR 6.70, 95% CI 2.61 – 17.19,  $p <$   
 59  $0.0001$ ).

60 **Conclusion:** Periodontitis appears as a significant risk factor for preeclampsia, which might  
 61 be even more pronounced in lower middle-income countries. Future studies to investigate if  
 62 maternal amelioration of periodontitis prevents preeclampsia might be warranted.

63 **Keywords:** Periodontitis, periodontal disease, preeclampsia, pre-eclampsia, hypertension,  
 64 pregnancy outcome.

65    **Abbreviations and Acronyms:**

66    PD:    pocket depth

67    CAL:   clinical attachment loss

68

## INTRODUCTION

Preeclampsia is the onset of pregnancy-related hypertensive disorder and proteinuria arising most commonly after 20 weeks of gestation, which could lead to eclampsia and induce maternal and perinatal morbidity and mortality. The prevalence of preeclampsia is between 2% to 8% of all pregnancies worldwide.<sup>1</sup> Preeclampsia affected pregnancies had a higher risk of poor maternal outcomes including cerebrovascular bleeding, HELLP syndrome, eclampsia, poorer outcomes of their offspring including premature birth, intrauterine growth restriction and the complications may manifest over years postpartum.<sup>2, 3</sup>

Contributing to USD 6.4 billion short-term estimated costs for preeclamptic pregnancies in US healthcare system, USD1.03 billion were spent on maternal healthcare and USD1.15 billion were expended for infants born to these mothers while the remaining expenses were for peripartum and postpartum care<sup>4</sup>.

Herein, it is of utmost importance to manage risk factors of preeclampsia to improve maternal and perinatal outcomes as well as lessen the burden on the health economic aspects.

Depending on geographical regions approximately 14.2% and 54.8% of pregnant women suffer from periodontal disease.<sup>5-7</sup> Especially, periodontitis, a more severe type of periodontal diseases affecting 11% of the pregnant women, can cause the destruction of periodontal tissue and cause systemic dissemination of bacteria and other inflammatory mediators.<sup>8, 9</sup> Systemic inflammatory processes triggered by focal periodontal infections have been attributed to cardiovascular, cerebrovascular diseases and respiratory diseases.<sup>10</sup> Periodontitis has independently been linked to several pregnancy complications such as preterm birth, low birth weight and gestational diabetes.<sup>11, 12</sup>

Socioeconomic status is a recognized factor associated with medical outcomes, including pregnancy outcomes.<sup>13</sup> Women with lower socioeconomic status are at a higher risk of pregnancy complications such as gestational diabetes, preterm delivery and preeclampsia.<sup>14-16</sup> Women with high socioeconomic status have a statistically significant reduced risk of preeclampsia with an odds ratio of 0.899 (95% CI, 0.862 – 0.937,  $p < 0.001$ ) compared to women with lower socioeconomic statuses.<sup>17</sup> At the same time, the proportion of periodontitis in pregnancy is linked to low socioeconomic status with 42.6%, compared to high socioeconomic status with 15.0%.

Two previous meta-analyses both published in 2013 reported positive associations between preeclampsia and periodontitis with OR 2.17, 95% CI 1.38-3.41,  $p = 0.008$  and OR of 2.79, 95% CI 2.01-3.01,  $p < 0.0001$ , but did not consider socioeconomic factors.<sup>18,19</sup> Since then further case-control and cohort studies have been published on this research topic.<sup>20,21,22</sup> Nonetheless, the causal relationship between periodontal disease and preeclampsia remains unclear.<sup>23,24</sup> In this review, we included all available new studies, to re-evaluate the potential association between periodontitis and preeclampsia and also to take socioeconomic factors into consideration.<sup>25</sup>

## **METHODS**

### **Eligibility criteria**

The studies were screened according to the following inclusion criteria: (1) case-control, prospective cohort study, (2) studies analysing the association between periodontal disease

and preeclampsia, (3) study population was pregnant women without systemic diseases, and (4) data was presented in such a way where Odds Ratio and 95% Confidence Interval could be calculated. Studies were excluded if they did not report adequate data or outcome of interest.

### **Information sources**

We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines with the checklist of 27 items to conduct our study.<sup>26</sup> We also adopted the MOOSE checklist for Meta-analysis of Observational studies.<sup>27</sup> A systematic search of the electronic database including Medline (from 1950), Pubmed (from 1946), Embase (from 1949), Lilacs, Cochrane Controlled Clinical Trial Register, CINAHL, ClinicalTrials.gov and Google Scholar (from 1990) to identify relevant articles .

### **Search strategy**

We used the following search terms: periodontitis, periodontal disease, preeclampsia, pre-eclampsia, pregnancy outcomes, pregnancy complications, hypertension. The combinations of search terms were used to explore above databases. The search strategy was peer reviewed by two independent reviewers (QA and LD). The reference lists of relevant articles were also scanned for appropriate studies. No language restrictions were adopted in either the search or study selection. No search for unpublished literature was carried out. Authors were contacted for translation and information.

### **Study selection**

Two independent reviewers (LD and HN) reviewed the titles, abstracts and methods of retrieved results to assess for the eligibility criteria. When there was a disagreement in a selection process between reviewers, consensus with the third reviewer (QA) was obtained.

## 144 **Data extraction**

145 Data extraction was carried out using a standardized extraction form, collecting information  
 146 on the first author's name, publication year, study design, number of cases, number of  
 147 controls, total sample size, country, national income (according to World Bank  
 148 classification<sup>28</sup>), mean age, the risk of estimates or data used to calculate the risk estimates,  
 149 Cis or data used to generate CI. The researchers cross-checked all extracted data and  
 150 discussed if there were disagreements.

151

## 152 **Assessment of risk of bias**

153 Risk of bias was executed using the Newcastle Ottawa Scale by two reviewers (QA and  
 154 LD)<sup>29</sup> with disagreements resolved by consensus attainment between reviewers. This Scale  
 155 has three components including Selection, Comparability and Outcome/Exposure assessment  
 156 with maximum overall score of nine. Studies were rated as low risk of bias if they received  
 157 nine score, moderate risk of bias if they received seven or eight score and high risk of bias if  
 158 they received less than seven scores.

159

## 160 **Data synthesis**

161 Data were imported in a statistical software (RevMan, Version 5, 2008, The Nordic Cochrane  
 162 Center, The Cochrane Collaboration, Copenhagen, Denmark). Pooled Odds Ratios, mean  
 163 difference, and 95% Confidence Intervals were calculated for the association between  
 164 periodontitis and preeclampsia using a random effects model. The pooled effect was  
 165 considered significant if p-value was less than 0.05. Forest plots for primary analysis and  
 166 subgroup analysis show the raw data, Odds Ratio and CIs, Means and SDs for the chosen  
 167 effect, heterogeneity statistic ( $I^2$ ), total number of participants per group, overall Odds Ratio  
 168 and Mean difference.



Subgroup analysis was carried out according to the study design (case-control or cohort), definition of periodontitis (defined by pocket depth (PD) and/or clinical attachment loss<sup>30</sup>), mean CAL, mean PD, national income (high-income or middle-income or low-income countries). Heterogeneity was tested with Cochran's Q statistic, with  $P < 0.10$  indicating heterogeneity, and quantified the degree of heterogeneity using the  $I^2$  statistic, which represents the percentage of the total variability across studies which is due to heterogeneity.  $I^2$  values of 25, 50 and 75% corresponded to low, moderate and high degrees of heterogeneity respectively<sup>31</sup>. We quantified publication bias using the Egger's regression model with the effect of bias assessed using the fail-safe number method.<sup>32</sup> The fail-safe number was the number of studies that we would need to have missed for our observed result to be nullified to statistical non-significance at the  $p < 0.05$  level. Publication bias is generally regarded as a concern if the fail-safe number is less than  $5n + 10$ , with  $n$  being the number of studies included in the meta-analysis.<sup>33</sup> Publication bias was assessed using Stata (16.1, StataCorp LLC, College Station, TX).

## RESULTS

### Study selection

A total of 3450 articles were found through the manual and electronic searches. After duplicates' removal, we screened 110 records for relevance. There were 67 papers excluded on a basis of evaluation of the title and abstract, leaving 43 articles to be assessed for eligibility. Of these, 30 articles were included in the quantitative analysis. A PRISMA flow diagram is provided in Figure 1. There were 9650 participants included in this systematic review and meta-analysis.

## Study characteristics

Table 1 depicts the characteristics of the included studies. There were six cohort studies<sup>21, 34-38</sup>, whilst the remaining studies were case - control<sup>20, 22, 30, 39-59</sup>. The definitions of periodontitis varied among these studies, while the definition of preeclampsia is presented comparatively consistent as blood pressure  $\geq 140/90$  mmHg and proteinuria during second trimester of gestation. The oral examination was conducted at different timepoints among studies, during pregnancy<sup>21, 34-39, 41, 47, 55, 56, 59</sup>, within 24 to 48 hours prior to delivery<sup>40, 43, 52, 54</sup> or after delivery<sup>22, 30, 37, 42, 44-46, 48-51, 53, 57, 58</sup>. The sample size ranged from 40 participants<sup>20</sup> to 1240 subjects<sup>53</sup>. There were seven studies with no evidence of an association between periodontitis and preeclampsia<sup>22, 30, 35, 39, 45, 46, 50, 55, 56</sup>, while the remaining studies reported a positive association.

## Risk of bias of included studies

Newcastle Ottawa Scale was used to evaluate the quality of evidence of these reports. Two reviewers marked the scores for each paper based on the tool provided by the Scale. Nine studies<sup>21, 37, 38, 48, 51, 54-57</sup> obtained the maximum score in Selection outcome while fourteen studies were marked with maximum score in the Comparability outcome and none of the studies could achieve ultimately 3 marks in the Exposure outcome. Table 2 describes the evaluation of risk of bias for this review.

## Synthesis of results

The results of the meta-analysis showed that periodontitis was associated with increased risk for preeclampsia (OR 3.18, 95% CI 2.26 – 4.48,  $p < 0.00001$ ; Figure 2). The heterogeneity was high ( $I^2 = 81\%$ ,  $p < 0.00001$ ) revealing a significant variation among studies.

## 219 Subgroup analysis.

220 According to the study type, the results revealed the increased risk of preeclampsia in  
 221 periodontitis patients in the cohort studies (OR 4.19, 95% CI 2.23 – 7.87,  $p < 0.00001$ ; Figure  
 222 3) and in case-control studies (OR 2.96, 95% CI 2.00 – 4.39,  $p < 0.00001$ ; Figure 4).  
 223 Heterogeneity was moderate for cohort ( $I^2 = 55\%$ ,  $p = 0.05$ ) but high for case-control ( $I^2 =$   
 224  $83\%$ ,  $p < 0.00001$ ).

225 When analyzing according to the national income, increased risk of preeclampsia were found  
 226 in periodontitis group in high-income countries (OR 2.67, 95% CI 1.59 – 4.49,  $p = 0.0002$ ;  
 227 Figure 5), upper middle-income countries (OR 2.40, 95% CI 1.73 – 3.31,  $p < 0.00001$ ; Figure  
 228 6) and especially lower-middle income countries (OR 6.7, 95% CI 2.61 – 17.19,  $p < 0.0001$ ;  
 229 Figure 7). Heterogeneity in the group of high-income, upper middle-income and lower  
 230 middle-income countries were moderate ( $I^2 = 59\%$ ,  $p = 0.006$ ;  $I^2 = 64\%$ ,  $p = 0.003$ ;  $I^2 = 86\%$ ,  
 231  $p < 0.00001$ , respectively).

232 When the results were analyzed according to the definition of periodontitis, an increased risk  
 233 of preeclampsia was observed in all subgroups, including PD only (OR 3.13, 95% CI 1.51 –  
 234 6.50,  $p = 0.002$ ), CAL and PD (OR 3.30, 95% CI 2.02 – 5.41,  $p < 0.00001$ ), CAL alone (OR  
 235 2.74, 95% CI 1.50 – 5.01,  $p = 0.001$ ). Heterogeneity was moderate for the subgroups in  
 236 which periodontitis was defined by PD alone ( $I^2 = 59\%$ ,  $p = 0.03$ ) and CAL alone ( $I^2 = 66\%$ ,  
 237  $p = 0.01$ ), while significantly high heterogeneity was found in the subgroup which  
 238 periodontitis was defined by CAL and PD ( $I^2 = 86\%$ ,  $p < 0.00001$ ).

239 When analyzing the periodontal condition between both groups, mean CAL was statistically  
 240 higher in the preeclamptic patients than in the healthy group (MD = 0.62, 95% CI 0.27 –  
 241 0.98,  $p = 0.0006$ ). Likewise, the preeclamptic group had a statistically higher mean PD  
 242 compared to healthy group (MD = 0.79, 95% CI -0.47 – 1.11,  $p < 0.00001$ ). The heterogeneity  
 243 was 98% in both subgroup analysis, ( $I^2 = 98\%$ ,  $p < 0.00001$ ).

## Publication bias

The funnel plot for the association between periodontitis and preeclampsia revealed the symmetry (Figure 13). There was no publication bias.

## DISCUSSION:

The aim of this meta-analysis was to re-evaluate the potential association between preeclampsia and periodontitis. The results confirm that periodontitis is a risk factor for preeclampsia, which was similar to the finding of a meta-analysis in 2013 by Sgolastra et al<sup>18</sup>. Our review has fifteen additional studies with three more cohort studies considerably increasing the sample size and hence generating more robust effect sizes and significance levels.

By stratifying according to study designs, periodontitis and preeclampsia showed significant associations in both case-control and cohort studies, whereas Sgolastra et al<sup>18</sup> could not report the statistical significance in the subgroup analysis of cohort studies (OR 2.2, 95% CI 0.66 – 7.36,  $p=0.2$ ). As a review of cohort studies provides higher level of evidence compared to case-control studies<sup>60</sup>, our data considerably strengthens the evidence of a positive association between preeclampsia and periodontitis. Moreover, the heterogeneity in an analysis of cohort studies in our study ( $I^2=55\%$ ,  $p=0.05$ ) was substantially lower than in a study by Sgolastra et al ( $I^2=89\%$ ,  $p=0.0001$ ), again strengthening the reliability of our results.

<sup>18</sup>

268 When analyzed according to the definition of periodontitis, three subgroup analysis with  
 269 studies defining periodontitis by PD alone, CAL and PD and CAL alone showed statistically  
 270 significant differences, whereas the previous meta-analysis showed only significance with a  
 271 subgroup analyzing periodontitis by CAL and PD. This could be explained by the number of  
 272 studies included in our review was more than in the previous analysis, thus, providing a more  
 273 comprehensive finding. However, according to the most recent case definition developed by  
 274 the Centre for Disease Control and Prevention in partnership with the American Academy of  
 275 Periodontology, the diagnostic criteria of periodontitis is at least 2 interproximal sites with  
 276 the minimum of attachment loss of 3 mm and at least 2 interproximal sites with the minimum  
 277 pocket depth of 4 mm (not on the same tooth) or one site with pocket depth  $\geq 5$  mm.<sup>61</sup>  
 278 Moreover, pregnant women who were preeclamptic had higher mean CAL and PD, however,  
 279 the heterogeneity in both subgroup analysis was high, indicating significant variations among  
 280 these studies in each subgroup. This could result from the difference in the periodontal probes  
 281 used in the dental examination in each study.  
 282 Several mechanisms have been proposed for the link between periodontitis and preeclampsia.  
 283 Higher levels of some periodontal pathogens such as *P.gingivalis* and *F. nucleatum* were  
 284 found in placenta of patients with preeclampsia.<sup>62</sup> Moreover, inflammatory responses  
 285 including the shifting of Th2 toward Th1, increasing oxidative stress, anti-angiogenic  
 286 proteins, vascular endothelial growth factor receptor 1 and complement C5a could potentially  
 287 enhance the development of preeclampsia.<sup>63</sup> Ananth et al. has reported the association  
 288 between intrauterine growth restriction and maternal periodontitis.<sup>64</sup> Since severe and early  
 289 onset preeclampsia were associated significantly with fetal growth restriction, this could  
 290 contribute to the mechanism underlying the association between preeclampsia and  
 291 periodontitis.<sup>65</sup> Furthermore, the mechanisms might be a reflection of dietary patterns.  
 292 Recently, some evidence has indicated that pathogenesis of preeclampsia involves maternal

gut microbiota, specifically, high-fiber diet which promote short chain fatty acid production and are associated with reduced risk of preeclampsia.<sup>66</sup> Similarly, high-fiber foods such as fruit and grains have been linked to the reduction of the progression of periodontal disease, suggesting the role of dietary intake in the potential relationship between preeclampsia and periodontal disease.<sup>66, 67</sup> However, future studies are required to elucidate these hypotheses.

When analyzed according to national income, this review revealed the significant difference in the subgroup analysis of high-income and upper middle-income countries (OR=2.67 and OR=2.40, respectively). Moreover, the subgroup analysis with lower middle-income countries, which generated Odds ratio of 6.70, indicated the considerable significance in the relationship between periodontitis and preeclampsia in this specific country group. Lower middle-income countries have poorer oral health condition.<sup>68</sup> Moreover, they have lower access to oral health care services (35%) to compare with upper middle-income (60-75%) and high-income countries (82%).<sup>69</sup> Therefore, it is notably important to improve the access to oral health services in pregnant women to lessen the risk of having preeclampsia.

We used Newcastle Ottawa Scale to evaluate the risk of bias and found twenty studies with moderate risk of bias and ten remaining studies with high risk of bias. Furthermore, there was no publication bias.

The strength of this systematic review and meta-analysis includes the large sample size of 9650 subjects. Six cohort studies comprising 2840 subjects were analyzed and revealed the statistically significant difference. Because cohort approach is the best methodical design in epidemiology, this finding provided the reliable evidence for the association between periodontitis and preeclampsia. Furthermore, by stratifying into subgroup analysis of national

income, our review has pointed out the association between these two diseases differed according to economical inequalities, thus, providing recommendation for health policy improvement. Pregnant women in low socioeconomic areas should be given access to oral healthcare services and encouraged to have their periodontal health checked and treated during pregnancy to potentially lower the risk of preeclampsia as well as other pregnancy complications. Jeffcoat et al reported non-surgical periodontal therapy could significantly reduce the medical costs for pregnant women by 73.7%.<sup>70</sup> There were also some limitations. Firstly, the heterogeneity of the overall analysis for the association between periodontitis and preeclampsia was high, pointing out the variations among studies included. This could be due to the synthesis of cohort and case-control studies in our review. Secondly, the general consensus in the definition and diagnosis of periodontitis was not clear enough which could influence the results of our meta-analysis. Therefore, future studies could take these into consideration and confirm our results.

## CONCLUSIONS AND IMPLICATIONS

This meta-analysis not only confirms previous findings of an association between periodontitis and preeclampsia but also shows larger effect size overall and specifically for lower-middle income countries in comparison to high and upper-middle income countries. Our results warrant future studies to investigate the mechanisms of this association and whether targeted interventions to prevent or treat periodontitis preconception or during pregnancy can lead to better pregnancy outcomes.

## FUNDING

This research did not receive any funding support from any parts.

## REFERENCE

1. Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol.* 2009;33(3):130-7.
2. Hung T-H, Hsieh Ts-Ta, Chen S-F. Risk of abnormal fetal growth in women with early- and late-onset preeclampsia. *Pregnancy hypertension.* 2018;12:201-6.
3. Turbeville HR, Sasser JM. Preeclampsia beyond pregnancy: long-term consequences for mother and child. *American journal of physiology Renal physiology.* 2020;318(6):F1315-F26.
4. Stevens W, Shih T, Incerti D, Ton TGN, Lee HC, Peneva D, et al. Short-term costs of preeclampsia to the United States health care system. *American Journal of Obstetrics & Gynecology.* 2017;217(3):237-48.e16.
5. Gesase N, Miranda-Rius J, Brunet-Llobet L, Lahor-Soler E, Mahande MJ, Masenga G. The association between periodontal disease and adverse pregnancy outcomes in Northern Tanzania: a cross-sectional study. *African health sciences.* 2018;18(3):601-11.
6. Govindasamy R, Dhanasekaran M, Varghese S, Balaji V, Karthikeyan B, Christopher A. Maternal risk factors and periodontal disease: A cross-sectional study among postpartum mothers in Tamil Nadu. *Journal of Pharmacy And Bioallied Sciences.* 2017;9(5):50-4.
7. Alchalabi HA, Al Habashneh R, Jabali OA, Khader YS. Association between periodontal disease and adverse pregnancy outcomes in a cohort of pregnant women in Jordan. *Clin Exp Obstet Gynecol.* 2013;40(3):399-402.
8. Piscoya MD, Ximenes RA, Silva GM, Jamelli SR, Coutinho SB. Periodontitis-associated risk factors in pregnant women. *Clinics (Sao Paulo, Brazil).* 2012;67(1):27-33.
9. Bui FQ, Almeida-da-Silva CLC, Huynh B, Trinh A, Liu J, Woodward J, et al. Association between periodontal pathogens and systemic disease. *Biomedical Journal.* 2019;42(1):27-35.
10. Winning L, Linden GJ. Periodontitis and systemic disease. *BDJ Team.* 2015;2(10):15163.
11. Corbella S, Taschieri S, Francetti L, De Siena F, Del Fabbro M. Periodontal disease as a risk factor for adverse pregnancy outcomes: a systematic review and meta-analysis of case-control studies. *Odontology.* 2012;100(2):232-40.
12. Abariga SA, Whitcomb BW. Periodontitis and gestational diabetes mellitus: a systematic review and meta-analysis of observational studies. *BMC Pregnancy Childbirth* 2016;16(344).
13. Kivimäki M, Batty GD, Pentti J, Shipley MJ, Sipilä PN, Nyberg ST, et al. Association between socioeconomic status and the development of mental and physical health conditions in adulthood: a multi-cohort study. *The Lancet Public Health.* 2020;5(3):e140-e9.
14. Bo S, Menato G, Bardelli C, Lezo A, Signorile A, Repetti E, et al. Low socioeconomic status as a risk factor for gestational diabetes. *Diabetes Metab.* 2002;28(2):139-40.
15. Silva LM, Coolman M, Steegers EAP, Jaddoe VWV, Moll HA, Hofman A, et al. Low socioeconomic status is a risk factor for preeclampsia: the Generation R Study. *Journal of Hypertension.* 2008;26(6).
16. Peacock JL, Bland JM, Anderson HR. Preterm delivery: effects of socioeconomic factors, psychological stress, smoking, alcohol, and caffeine. *Bmj.* 1995;311(7004):531-5.
17. Ross KM, Dunkel Schetter C, McLemore MR, Chambers BD, Paynter RA, Baer R, et al. Socioeconomic Status, Preeclampsia Risk and Gestational Length in Black and White Women. *J Racial Ethn Health Disparities.* 2019;6(6):1182-91.
18. Sgolastra F, Petrucci A, Severino M, Gatto R, Monaco A. Relationship between periodontitis and pre-eclampsia: a meta-analysis. *PloS one.* 2013;8(8):e71387.
19. Wei BJ, Chen YJ, Yu L, Wu B. Periodontal disease and risk of preeclampsia: a meta-analysis of observational studies. *PLoS One.* 2013;8(8):e70901.
20. Varshney S, Gautam A. Poor periodontal health as a risk factor for development of pre-eclampsia in pregnant women. *Journal of Indian Society of Periodontology.* 2014;18(3):321-5.
21. Soucy-Giguère L, Tétu A, Gauthier S, Morand M, Chandad F, Giguère Y, et al. Periodontal Disease and Adverse Pregnancy Outcomes: A Prospective Study in a Low-Risk Population. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC.* 2016;38(4):346-50.



22. Lafaurie GI, Gómez LA, Montenegro DA, De Avila J, Tamayo MC, Lancheros MC, et al. Periodontal condition is associated with adverse perinatal outcomes and premature rupture membranes in low income pregnant women in Bogota, Colombia: a case control study. *The journal of maternal-fetal & neonatal medicine : the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstetricians*. 2018;33(1):16-23.
23. Lavigne SE, Forrest JL. An umbrella review of systematic reviews of the evidence of a causal relationship between periodontal disease and adverse pregnancy outcomes: A position paper from the Canadian Dental Hygienists Association. *Can J Dent Hyg*. 2020;54(2):92-100.
24. Kunnen A, van Doormaal JJ, Abbas F, Aarnoudse JG, van Pampus MG, Faas MM. Periodontal disease and pre-eclampsia: a systematic review. *J Clin Periodontol*. 2010;37(12):1075-87.
25. Australian Institute of Health and Welfare 2010. Socioeconomic variation in periodontitis among Australian adults 2004–06. Research report series no. 50 . Cat. no. DEN 207. Canberra: AIHW.
26. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *J Clin Epidemiol*. 2009;62:1006-12.
27. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama*. 2000;283(15):2008-12.
28. Country Income Groups (World Bank Classification) [Internet]. 2011. Available from: <http://data.worldbank.org/about/country-classifications/country-and-lending-groups>.
29. Lo CK, Mertz D, Loeb M. Newcastle-Ottawa Scale: comparing reviewers' to authors' assessments. *BMC Med Res Methodol*. 2014;14(45).
30. Taghzouti N, Xiong X, Gornitsky M, Chandad F, Voyer R, Gagnon G, et al. Periodontal Disease is Not Associated With Preeclampsia in Canadian Pregnant Women. *Journal of periodontology*. 2012;83(7):871-7.
31. Higgins JP, Thompson SG. Quantifying heterogeneity in a metaanalysis. . *Stat Med*. 2002;21:1539-58.
32. Egger M, Davey SG, Schneider M, Minder C. Bias in metaanalysis detected by a simple, graphical test. *Br Med J*. 1997;315:629-34.
33. Orwin R. A fail-safe N for effect size in meta-analysis. . *J Educ Stat*. 1983;8:157-9.
34. Boggess KA, Lieff S, Murtha AP, Moss K, Beck J, Offenbacher S. Maternal periodontal disease is associated with an increased risk for preeclampsia. *Obstetrics and gynecology*. 2003;101(2):227-31.
35. Horton AL, Boggess KA, Moss KL, Beck J, Offenbacher S. Periodontal disease, oxidative stress, and risk for preeclampsia. *Journal of periodontology*. 2010;81(2):199-204.
36. Kumar A, Basra M, Begum N, Rani V, Prasad S, Lamba AK, et al. Association of maternal periodontal health with adverse pregnancy outcome. *The journal of obstetrics and gynaecology research*. 2013;39(1):40-5.
37. Ha JE, Jun JK, Ko HJ, Paik DI, Bae KH. Association between periodontitis and preeclampsia in never-smokers: a prospective study. *Journal of clinical periodontology*. 2014;41(9):869-74.
38. Lee HJ, Ha JE, Bae KH. Synergistic effect of maternal obesity and periodontitis on preterm birth in women with pre-eclampsia: a prospective study. *Journal of clinical periodontology*. 2016;43(8):646-51.
39. Chaparro A, Sanz A, Quintero A, Inostroza C, Ramirez V, Carrion F, et al. Increased inflammatory biomarkers in early pregnancy is associated with the development of pre-eclampsia in patients with periodontitis: a case control study. *Journal of periodontal research*. 2013;48(3):302-7.
40. Canakci V, Canakci CF, Canakci H, Canakci E, Cicek Y, Ingec M, et al. Periodontal disease as a risk factor for pre-eclampsia: a case control study. *The Australian & New Zealand journal of obstetrics & gynaecology*. 2004;44(6):568-73.

- 448 41. Contreras A, Herrera JA, Soto JE, Arce RM, Jaramillo A, Botero JE. Periodontitis Is  
449 Associated With Preeclampsia in Pregnant Women. *Journal of Periodontology*. 2006;77(2).
- 450 42. Cota LO, Guimarães AN, Costa JE, Lorentz TC, Costa FO. Association between maternal  
451 periodontitis and an increased risk of preeclampsia. *Journal of periodontology*.  
452 2006;77(12):2063-9.
- 453 43. Canakci V, Canakci CF, Yildirim A, Ingec M, Eltas A, Erturk A. Periodontal disease  
454 increases the risk of severe pre-eclampsia among pregnant women. *J Clin Periodontol*.  
455 2007;34:639-45.
- 456 44. Siqueira FM, Cota LO, Costa JE, Haddad JP, Lana AM, Costa FO. Maternal periodontitis as  
457 a potential risk variable for preeclampsia: a case-control study. *Journal of periodontology*.  
458 2008;79(2):207-15.
- 459 45. Lohsoonthorn V, Kungsadalpipob K, Chanchareonsook P, Limpongsanurak S,  
460 Vanichjakvong O, Sutdhibhisal S, et al. Maternal periodontal disease and risk of preeclampsia: a  
461 case-control study. *American journal of hypertension*. 2009;22(4):457-63.
- 462 46. Shetty M, Shetty PK, Ramesh A, Thomas B, Prabhu S, Rao A. Periodontal disease in  
463 pregnancy is a risk factor for preeclampsia. *Acta obstetrica et gynecologica Scandinavica*.  
464 2009;89(5):718-21.
- 465 47. Politano GT, Passini R, Nomura ML, Velloso L, Morari J, Couto E. Correlation between  
466 periodontal disease, inflammatory alterations and pre-eclampsia. *Journal of Dental Research*.  
467 2011;46:505-11.
- 468 48. Ha JE, Oh KJ, Yang HJ, Jun JK, Jin BH, Paik DI, et al. Oral Health Behaviors, Periodontal  
469 Disease, and Pathogens in Preeclampsia: A Case-Control Study in Korea. *Journal of*  
470 *periodontology*. 2011;82(12):1685-92.
- 471 49. Sayar F, Sadat Hoseini M, Abbaspour S. Effect of Periodontal Disease on Preeclampsia.  
472 *Iranian J Publ Health*. 2011;40(3):122-7.
- 473 50. Hirano E, Sugita N, Kikuchi A, Shimada Y, Sasahara J, Iwanaga R, et al. The association of  
474 *Aggregatibacter actinomycetemcomitans* with preeclampsia in a subset of Japanese pregnant  
475 women. *J Clin Periodontol*. 2012;39:229-38.
- 476 51. Moura da Silva G, Coutinho SB, Piscoya MD, Ximenes RA, Jamelli SR. Periodontitis as a  
477 Risk Factor for Preeclampsia. *Journal of periodontology*. 2012;83(11):1388-96.
- 478 52. Pralhad S, Thomas B, Kushtagi P. Periodontal Disease and Pregnancy Hypertension: A  
479 Clinical Correlation. *Journal of periodontology*. 2013;84(8):1118-25.
- 480 53. Desai K, Desai P, Duseja S, Kumar S, Mahendra J, Duseja S. Significance of maternal  
481 periodontal health in preeclampsia. *Journal of International Society of Preventive & Community*  
482 *Dentistry*. 2015;5(2):103-7.
- 483 54. Jaiman G, Nayak PA, Sharma S, Nagpal K. Maternal periodontal disease and preeclampsia  
484 in Jaipur population. *Journal of Indian Society of Periodontology*. 2018;22(1):50-4.
- 485 55. Pattanashetti JI, Nagathan VM, Rao SM. Evaluation of Periodontitis as a Risk for Preterm  
486 Birth among Preeclamptic and Non-Preeclamptic Pregnant Women - A Case Control Study. .  
487 *Journal of clinical and diagnostic research : JCDR*. 2013;7(8):1776-8.
- 488 56. Khalighinejad N, Aminoshariae A, Kulild JC, Mickel A. Apical Periodontitis, a Predictor  
489 Variable for Preeclampsia: A Case-control Study. *Journal of endodontics*. 2017;43(10):1611-4.
- 490 57. Khader YS, Jibreal M, Al-Omiri M, Amarin Z. Lack of association between periodontal  
491 parameters and preeclampsia. *Journal of periodontology*. 2006;77(10):1681-7.
- 492 58. Yaghini J, Mostajeran F, Afshari E, Naghsh N. Is periodontal disease related to  
493 preeclampsia? *Dental research journal*. 2012;9(6):770-3.
- 494 59. Kunnen A, Blaauw J, van Doormaal JJ, van Pampus MG, van der Schans CP, Aarnoudse JG,  
495 et al. Women with a recent history of early-onset pre-eclampsia have a worse periodontal  
496 condition. *Journal of clinical periodontology*. 2007;34(3):202-7.
- 497 60. Guyatt GH, Haynes RB, Jaeschke RZ, Cook DJ. Users' guides to the medical literature:  
498 XXV. evidence-based medicine: principles for applying the users' guides to patient care. *Jama*.  
499 2000;284:1290-6.
- 500 61. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions for  
501 population-based surveillance of periodontitis. *J Periodontol*. 2012;83(12):1449-54.

62. Barak S, Oettinger-Barak O, Machtei EE, Sprecher H, Ohel G. Evidence of periopathogenic microorganisms in placentas of women with preeclampsia. *J Periodontol.* 2007;78(4):670-6.
63. Nourollahpour Shiadeh M, Behboodi Moghadam Z, Adam I, Saber V, Bagheri M, Rostami A. Human infectious diseases and risk of preeclampsia: an updated review of the literature. *Infection.* 2017;45(5):589-600.
64. Ananth CV, Andrews HF, Papapanou PN, Ward AM, Bruzelius E, Conicella ML, et al. History of periodontal treatment and risk for intrauterine growth restriction (IUGR). *BMC Oral Health.* 2018;18(1):161.
65. Odegård RA, Vatten LJ, Nilsen ST, Salvesen KA, Austgulen R. Preeclampsia and fetal growth. *Obstet Gynecol.* 2000;96(6):950-5.
66. Hu M, Eviston D, Hsu P, Mariño E, Chidgey A, Santner-Nanan B, et al. Decreased maternal serum acetate and impaired fetal thymic and regulatory T cell development in preeclampsia. *Nature Communications.* 2019;10(1):3031.
67. Schwartz N, Kaye EK, Nunn ME, Spiro A, 3rd, Garcia RI. High-fiber foods reduce periodontal disease progression in men aged 65 and older: the Veterans Affairs normative aging study/Dental Longitudinal Study. *Journal of the American Geriatrics Society.* 2012;60(4):676-83.
68. Watt R, Sheiham A. Inequalities in oral health: a review of the evidence and recommendations for action. *Br Dent J* 1999;187:6-12.
69. Hosseinpour AR, Itani L, Petersen PE. Socio-economic inequality in oral healthcare coverage: results from the World Health Survey. *J Dent Res.* 2012;91(3):275-81.
70. Jeffcoat MK, Jeffcoat RL, Gladowski PA, Bramson JB, Blum JJ. Impact of Periodontal Therapy on General Health: Evidence from Insurance Data for Five Systemic Conditions. *American Journal of Preventive Medicine.* 2014;47(2):166-74.

**Table 1. Descriptions of included studies**

No	Reference	Country	Design	Participants and age	Definition of PE	Definition of PD	Examination Time	Finding (conclusion)	OR (95% CI)
1	Boggess et al. <sup>34</sup> (2003)	US	Cohort	39 cases  763 controls	BP $\geq$ 140/90 on 2 separate occasions, and $\geq$ 1+ proteinuria on catheterized urine specimen	PD $\geq$ 4 and CAL $\geq$ 3mm without BOP.  Mild: PD $\geq$ 4 mm or BOP on 1-15 teeth  Severe: PD $\geq$ 4 mm on > 15 teeth  Disease progression: $\geq$ 4 sites that increased $\geq$ 2mm in PD, resulting in $\geq$ 4 mm in PD	At the first or second prenatal visit and then repeated within 48 hours antepartum.  Enrolled at < 26 weeks 'gestation and followed until delivery	Women were at higher risk for preeclampsia if they had severe periodontal disease at delivery or if they had periodontal disease progression during pregnancy	Severe periodontal disease: 2.4 (1.1- 5.3)  Periodontal disease progression: 2.1 (1.0 – 4.4)

2	Canakci et al. <sup>40</sup> (2004)	Turkey	Case-control	41 cases ,41 controls	BP $\geq$ 140/90 mmHg and proteinuria $\geq$ 300mg/24h or 2+ proteinuria on dip sticks, on 2 occasions $\geq$ 6h apart if 24h urine specimen is unavailable	$\geq$ 4 teeth with $\geq$ 1 sites with PD $\geq$ 4mm and BOP+ and CAL $\geq$ 3mm at the same site	within 48 h prior to delivery	multiple logistic regression results showed that pre-eclamptic patients were 3.47 times more likely to have periodontal disease than normotensive patients	3.47 (1.07–11.95)
3	Contreras et al. <sup>41</sup> 2006	Colombia	Case-control	130 case, 243 controls	2+ proteinuria, confirmed by $\geq$ 0.3 g proteinuria/24 hours and hypertension ( $\geq$ 140/ 90 mmHg)	$\geq$ 4 sites showed PD $\geq$ 4 mm), CAL $\geq$ 4 mm, and bleeding on probing.  Incipient: CAL from 4 to 5  Moderate/severe: CAL $\geq$ 6 mm	between 26 to 36 weeks of pregnancy	Chronic periodontal disease was significantly associated with preeclampsia in pregnant women	3.0 (1.91 - 4.87)

4	Cota et al. <sup>42</sup> 2006	Brazil	Case-control	109 cases, 479 controls	BP >140/90 mm Hg and $\geq$ 1+proteinuria after 20 weeks of gestation	$\geq$ 4 teeth with $\geq$ 1 sites with a PD $\geq$ 4 mm and CAL $\geq$ 3mm at the same site	Within 48 hours of delivery	Maternal periodontitis was determined to be associated with an increased risk of preeclampsia	1.88 (1.1 - 3.0)
5	Khader et al. <sup>57</sup> 2006	Jordan	Case-control	115 cases 230 controls	Preeclampsia was defined as the development of blood pressure of $\geq$ 140/90 mmHg after 20 weeks of gestation, combined with proteinurea of at least 1+ on a midstream urine specimen or on a	Not mentioned	Within 24 hours after delivery	This study did not support the association between periodontal parameters and preeclampsia	

					catheter specimen, provided urinary tract infection was not the contributing factor to the proteinuria in women who were known to be normotensive and nonproteinuric before pregnancy or in early pregnancy				
6	Kunnen et al. <sup>59</sup> 2007	Netherlands	Case-control	17 cases, 35 controls	DBP $\geq$ 90mmHg on 2 occasions and proteinuria $\geq$	PD $\geq$ 4mm	before 34 weeks of pregnancy	These results indicate that Caucasian	7.9 (1.9–32.8)

					30mg/dl (or 1+ on a urine dip stick) on $\geq 2$ random specimens collected $\geq 4$ h apart.	Mild PD: BOP and PD $\geq 4$ mm on 1–15 sites  Severe: BOP and PD $\geq 4$ mm on $> 15$ sites		women with a recent history of early-onset pre-eclampsia have a worse periodontal condition, as compared with women with uncomplicated deliveries	
7	Canakci et al. <sup>43</sup> 2007	Turkey	Case-control	20 Mild PE, 18 Severe PE, 21 Controls	DBP $\geq 90$ mmHg and proteinuria(300 mg/24h urine sample) and the presence of edema	Mild: BOP and $\geq 4$ mm PD on 1–15 sites Severe: : BOP and $\geq 4$ mm PD on $\geq 15$ sites	within 48h preceding delivery	The results of multivariate logistic regression showed a highly significant association between mild to	Severe PD: 3.78 (1.77–12.74)



					<p>Mild: BP <math>\geq</math> 140/90mmHg on <math>\geq 2</math> occasions 6h apart, with or without proteinuria</p> <p>Severe: SBP <math>\geq</math> 160 or DBP <math>\geq</math> 110mmHg on 2 occasions <math>\geq</math>6h apart and proteinuria <math>\geq</math> 5g/24h urine sample or <math>\geq</math> 3l on dip stick in <math>\geq 2</math> random clean-catch samples <math>\geq</math> 4h apart</p>			severe pre-eclampsia and severe periodontal disease	
--	--	--	--	--	--	--	--	---	--

8	Siqueira et al. <sup>44</sup> 2008	Brazil	Case-control	164 cases, 1042 controls	BP > 90mmHg on 2 occasions after 20 GW and $\geq 1+$ proteinuria	$\geq 4$ mm and CAL $\geq 3$ mm at the same site in $\geq 4$ teeth	within 48 hours of delivery	Maternal periodontitis is a risk factor associated with preeclampsia	1.52 ( 1.01 - 2.29)
9	Lohsoonthorn et al. <sup>45</sup> 2009	Thailand	Case-control	150 cases, 150 controls	BP $\geq 140/90$ mmHg and proteinuria $\geq 30$ mg/dl (or $1+$ on a urine dip stick) on $\geq 2$ random specimens collected $\geq 4$ h apart.	Mild: $\geq 1$ teeth with interproximal sites showing $\geq 4$ mm CAL and $\geq 4$ mm PD  Moderate: $\geq 2$ nonadjacent teeth with interproximal sites showing $\geq 5$ mm CAL and $\geq 4$ mm PD  Severe: $\geq 2$ nonadjacent teeth with interproximal sites showing $\geq 6$ mm CAL and $\geq 4$ mm PD	within 48 h after delivery	This study provides no convincing evidence that periodontal disease is associated with preeclampsia risk among Thai women	Severe PD: 0.92 ( 0.26–3.28)

10	Horton et al. <sup>35</sup> 2010	US	Cohort	34 cases (pre-eclampsia) 757 controls (non pre-eclampsia)	BP >140/90 mmHg and ≥ 1+ proteinuria on a catheterized urine specimen	Mild: <15 sites with the presence of one or more pockets ≥4 mm or one or more pockets with bleeding.  Moderate/severe: ≥15 sites demonstrated a probing depth ≥4 mm	<26 weeks of gestation	Among women with periodontal disease, the presence of 8-isoprostane ≥ 75th percentile did not significantly increase the odds for the development of preeclampsia	2.08 (0.65 - 6.60)
11	Shetty et al. <sup>46</sup> 2009	India	Case-control	30 cases 100 controls	BP > 140/90 mmHg on more than 2 occasions 4 hours apart and 1+ or more proteinuria by Dipstick on a	CAL of ≥ 3 mm and a PD of ≥ 4 mm. (The teeth examined were 16, 22, 24, 36, 42, and 44)	within 48 hours of delivery	periodontitis both at enrolment as well as within 48 hours of delivery may be associated with	Enrolment: 5.78 (2.41–13.89) Delivery: 20.15 (4.55–89.29)

					random urine sample			an increased risk of preeclampsia.	
<b>12</b>	Politano et al. <sup>47</sup> 2011	Brazil	Case-control	58 cases 58 controls	increase in systolic arterial pressure ( $\geq$ 140 mmHg) and/or diastolic pressure ( $\geq$ 90mmHg) and proteinuria ( $\geq$ 300 mg/24 h), after 20 wk of gestation	two or more sites showed pocket formation ( $\geq$ 4 mm), clinical attachment level ( $\geq$ 4 mm) and bleeding on probing	after 20 wk of gestation	periodontal disease may increase the risk of pre-eclampsia	3.73 (1.32–10.58)
<b>13</b>	Ha et al. <sup>48</sup> 2011	Korea	Case-control	16 cases 48 controls	BP >140/90 mm Hg on two separate occasions and $\geq$ 1+ proteinuria on	Localized periodontitis: periodontal clinical attachment loss $\geq$ 3.5 mm on two or three sites not on the same tooth.	5 days after delivery	preeclampsia could be associated with the maternal periodontal condition and	Localized periodontitis: 4.79 (1.02 - 29.72)

					a random sample of urine	Generalized periodontitis: CAL $\geq 3.5$ mm on $\geq 4$ sites not on the same tooth		interdental cleaning	Generalized periodontitis: 6.60 (1.25 - 41.61)
14	Sayar et al. <sup>49</sup> 2011	Iran	Case- control	105 cases 105 controls	blood pressure $\geq 140/90$ mmHg and proteinuria +1  Mild: CAL $\leq 2$ mm  Moderate to Severe: CAL $\geq 3$ mm	No definition specified (results based on periodontal parameters)  Mild: CAL $\leq 2$ mm  Moderate to Severe: CAL $\geq 3$ mm	48 hours after child delivery.	Preeclamptic cases significantly had higher attachment loss and gingival recession than the control group	4.1 (1.5-11.5)
15	Taghzouti et al. <sup>30</sup> 2012	Canada	Case- control	92 cases 245 controls	BP $\geq 140/90$ mm Hg on two occasions $\geq 4$ hours apart after 20 weeks of	periodontitis is defined as $\geq 4$ sites exhibiting PD $\geq 5$ mm and CAL $\geq 3$ mm at the same sites.	within 48 hours after delivery	This study does not support the hypothesis of an association between	1.13 (0.59 - 2.17)

					gestation, and 0.3 g proteinuria on a 24-hour urine collection, or $\geq 1$ on a dipstick			periodontal disease and preeclampsia	
16	Chaparro et al. <sup>39</sup> 2013	Chile	Case-control	11 cases 43 controls	During the second and third trimester of pregnancy, BP > 140/90 and proteinuria, which was considered to be present when one 24-h urine collection showed a total protein excretion $\geq 300$ mg.	PD $\geq 4$ mm and CAL $\geq 3$ mm at the same site of $\geq 4$ teeth, inflammation and bleeding on probing (BOP)	Blood samples were collected at enrolment  Gingival crevicular fluid samples were collected between 11–14 wk	Preeclamptic women shows increased levels of IL-6 in GCF and CRP in plasma during early pregnancy. Periodontal disease could contribute to systemic inflammation in early pregnancy via a local	1.36 (0.252–7.372)

								increase of IL-6 and the systemic elevation of CRP. Therefore, both inflammatory markers could be involved in the relationship between periodontal disease and pre- eclampsia.	
17	Hirano et al. <sup>50</sup> 2012	Japan	Case- control	18 cases 109 controls	hypertension (systolic blood pressure > 140 mmHg and/or diastolic blood pressure > 90	having over 60% of sites with CAL $\geq$ 3 mm	5 days after labor	No statistically significant association between any of the periodontal clinical	1.7 (1.1 - 2.7)

					mmHg) with proteinuria ( 300 mg/day) occurring after the 20th week of gestation, but being resolved by the 12th postpartum week			parameters or the presence of periodontitis and preeclampsia.	
<b>18</b>	Kumar et al. <sup>36</sup> 2012	India	Cohort	35 cases 305 controls	systolic blood pressure 140 mm of mercury and diastolic blood pressure 90 mm of mercury at two occasions at least 4 h apart after 20 weeks of gestation in a	clinical attachment loss and probing depth $4 \geq$ mm in one or more sites were diagnosed as those with periodontitis	14–20 weeks period of gestation	Maternal periodontitis is associated with an increased risk of pre-eclampsia	7.48 (2.72–22.42)



					woman with previously normal blood pressure along with development of proteinuria				
19	Da Silva et al. <sup>15</sup> 2012	Brazil	Case-control	284 cases 290 controls	a systolic blood pressure $\geq 140$ mmHg or a diastolic pressure $\geq 90$ mmHg and proteinuria $\geq 300$ mg/24 hours or 2+ on dipsticks, developed after week 20 of gestation in previously	$\geq 4$ teeth with $\geq 1$ sites with a PD $\geq 4$ mm and AL $\geq 3$ mm in the same site	within 48 hours of childbirth	periodontitis was a risk factor for preeclampsia	8.60 ( 3.92 - 18.88)

					normotensive females				
20	Pralhad et al. <sup>52</sup> 2012	India	Case-control	100 cases 100 controls	The resting blood pressure was $\geq 140/90$ mmHg after 20 weeks of gestation with or without associated proteinuria	Any of the following is present: 1) OHI >3 69; 2) GI >1 (moderate-to-severe gingival inflammation); 3) mean PD >4 mm; and 4) CAL >3 mm	within 72 hours of their hospital admission for delivery	Periodontal disease is more prevalent in females with pregnancy hypertension	5.5 (2.7 -11.4)
21	Yaghini et al. <sup>58</sup> 2012	Iran	Case-control	26 cases 25 controls	blood pressure >140/90 mmHg and > or = 1+ proteinuria on a catheterized urine specimen.	Not mentioned	48 hours after delivery	Maternal periodontal disease during pregnancy is not associated with preeclampsia.	
22	Pattanashetti et al. <sup>55</sup> 2013	India	Case-control	100 cases 100 controls	Pregnancy induced hypertension	- Mild periodontal disease: One or more sites with probing	Sixth month of pregnancy and within 48	Pregnant women with preeclampsia are	Periodontitis in the case group was 72%, in

					<p>occurs after the 20th week of gestation and is characterized by:</p> <p>Hypertension – High blood pressure, usually higher than 140/90 mm Hg.</p> <p>The rise of blood pressure should be evident at least on two occasions, four or more hours apart.</p> <p>Edema – Demonstration of pitting oedema</p>	<p>depth <math>\geq 3</math> mm that bleed upon probing but less than 25 sites with probing depth <math>\geq 4</math>mm.</p> <ul style="list-style-type: none"> <li>- Moderate/Severe: 15 or more sites with periodontal probing <math>\geq 4</math>mm.</li> <li>- Worsening periodontal status was defined as four or more sites had increased by at least 2mm in pocket depth between the two oral health examinations</li> </ul>	hours post-partum	<p>at greater risk for preterm delivery if periodontal disease is present during pregnancy or progress during pregnancy and also the rate of preterm delivery is more in preeclamptic women having moderate to severe periodontal disease.</p>	control group was 62%, $p < 0.001$ .
--	--	--	--	--	--	---	-------------------	--	--------------------------------------

					<p>over the ankles after 12 hours of bed rest or rapid gain in weight of more than 1 lb per week or more than 5 lb a month in the later month of pregnancy may be the easiest evidence of preeclampsia.</p> <p>Proteinuria – Presence of protein in 24 hours urine with more than 1gm per litre in 2 or more midstream</p>				
--	--	--	--	--	--	--	--	--	--

					specimens obtained 6 hours apart in the absence of urinary tract infection is considered significant.				
23	Ha et al. <sup>37</sup> 2014	Korea	Cohort	13 cases  270 controls	BP > 140/90 mmHg on two separate occasions, and at least 1+ proteinuria on a random urine screen after the 20th week of pregnancy	CAL $\geq$ 4.0 mm on two or more sites on different teeth.	21–24 weeks of gestation	Periodontitis increased the risk of preeclampsia among never- smokers	4.51 (1.13– 17.96)

24	Varshney et al. <sup>20</sup> 2014	India	Case-control	20 cases 20 controls	BP $\geq$ 140/90 mm of Hg on two separate occasions after 20 week of gestation and $\geq$ 1+ proteinuria	PD $\geq$ 4 mm and CAL $\geq$ 3 mm at the same site on at least 4 different non-neighboring teeth	within the 48 h after delivery	Maternal clinical periodontal disease at delivery is associated with an increased risk for the development of pre-eclampsia, independent of the effects of maternal age, race, smoking, gestational age at delivery	4.33 (1.15- 16.32)
25	Desai et al. <sup>53</sup> 2015	India	Case-control	120 cases 1120 controls	Blood pressure $\geq$ 140/90 mm Hg on two separate occasions after	PD $\geq$ 4 mm and CAL $\geq$ 3 mm at the same site in at least four teeth.	Periodontal examination was	Maternal clinical periodontal disease at delivery is	19.898 (7.80-48.94)

					week 20 of gestation.		performed 48h after delivery	associated with an increased risk for the development of pre-eclampsia, independent of the effects of maternal age, race, smoking, gestational age at delivery	
26	Soucy-Giguere et al. <sup>21</sup> 2015	Canada	Cohort	11 cases 237 controls	Not mentioned	the presence of at least one site with probing depths $\geq 4$ mm and $\geq 10\%$ bleeding on probing	in the seven days following amniocentesis.	Pregnant women with periodontal disease were more likely to develop preeclampsia	RR 5.89 (1.24-28.05)

27	Lee et al. <sup>38</sup> 2016	Korea	Cohort	15 cases 313 controls	BP > 140/90 mmHg on two separate occasions with at least 1+ proteinuria on a random urine screen after the 20th week of pregnancy	two or more inter-proximal sites with CAL $\geq$ 4 mm that were not on the same tooth	at 21–24 weeks of gestation	The association was much stronger in women with both obesity and periodontitis	15.94 (3.31–76.71)
28	Khalighinejad et al. <sup>56</sup> 2017	USA	Case-control	50 cases 50 controls	A systolic blood pressure $\geq$ 140 mm HG or a diastolic pressure $\geq$ 90 mm HG and proteinuria > 300 mg/24h developed after	The presence of 4 or more teeth with 1 or more sites with PD $\geq$ 4 mm and with clinical attachment loss $\geq$ 3mm at the same site	Before 26 weeks of gestation	Apical periodontitis was significantly more prevalent in the experimental group.	2.23 (1.92-6.88)



					the 20 <sup>th</sup> week of gestation				
29	Lafaurie et al. <sup>22</sup> 2018	Colombia	Case-control	76 cases 304 controls	not mentioned	the patients were classified according to the presence of periodontal pockets (code 3: periodontal pockets of 4-5mm or code 4: periodontal pockets >5mm)	during the first week after birth	Periodontal pockets presence was not associated with preeclampsia.	5.46 (1.84-16.1)
30	Jaiman et al. <sup>54</sup> 2018	India	Case-control	15 cases 15 controls	preeclampsia as the appearance of a diastolic blood pressure $\geq 90$ mmHg mercury measured at two different occasions at least 4 h apart in combination with proteinuria ( $\geq 300$	According to the criteria of L��e and Silness.	24 hours before delivery	The preeclamptic women were associated with significantly higher periodontitis and lower fetal birth weight than normotensive women	Periodontitis in case group was 93.3% and in control group was 33.3% (p<0.05)

					mg/24 h or +1 dipstick) developing after a gestational age of 20 weeks in a previously normotensive woman				
BP: blood pressure; DBP: diastolic blood pressure; SBP: systolic blood pressure; GW: gestational week; PD: pocket depth; CAL: clinical attachment loss; OHI: oral hygiene index; GI: gingival index; BOP: bleeding on probing; GCF: gingival crevicular fluid									

**Table 2. Risk of bias in included studies based on Newcastle-Ottawa scale.**

Study, year	Selection (Max 4*)	Comparability (Max 2*)	Exposure (Max 3*)	Risk of bias
Bogges 2003	***	*	**	High
Canakci 2004	***	**	**	Moderate
Contreras 2006	***	*	**	High
Cota 2006	***	*	**	High
Khader 2006	****	**	**	Moderate
Kunnen 2007	***	**	**	Moderate
Canakci 2007	***	**	**	Moderate
Siqueira 2008	***	*	**	High
Lohsoonthorn 2009	***	**	**	Moderate
Horton 2010	***	*	**	High
Shetty 2010	***	**	**	Moderate
Politano 2011	***	**	**	Moderate
Ha 2011	****	**	**	Moderate
Sayar 2011	***	**	**	Moderate
Taghzouti 2012	***	**	**	Moderate
Chaparro 2012	***	*	**	High

Hirano 2012	***	*	**	High
Kumar 2012	***	*	**	High
da Silva 2012	****	**	**	Moderate
Pralhad 2012	***	**	**	Moderate
Yaghini 2012	***	**	**	Moderate
Pattanashetti 2013	****	**	**	Moderate
Ha 2014	****	**	**	Moderate
Varshney 2014	***	**	**	Moderate
Desai 2015	***	**	**	Moderate
Soucy-Giguere 2015	****	**	***	Low
Lee 2016	****	*	**	Moderate
Khalighinejad 2017	****	**	**	Moderate
Lafaurie 2018	***	*	**	High
Jaiman 2018	****	**	**	Moderate

## **FIGURE LEGENDS**

**Figure 1. Flow diagram of study selection**

**Figure 2. Forest plot for the association between periodontitis and preeclampsia.**

**Figure 3. Forest plot for the subgroup analysis according to the type of study design (cohort study).**

**Figure 4. Forest plot for the subgroup analysis according to the type of study design (case-control study)**

**Figure 5. Forest plot for the subgroup analysis according to the national income (high income countries)**

**Figure 6. Forest plot for the subgroup analysis according to the national income (upper middle-income countries)**

**Figure 7. Forest plot for the subgroup analysis according to the national income (lower middle- income countries)**

**Figure 8. Forest plot for the subgroup analysis according to the definition of periodontitis (PD alone)**


**Figure 9. Forest plot for the subgroup analysis according to the definition of periodontitis (PD and CAL)**

**Figure 10. Forest plot for the subgroup analysis according to the definition of periodontitis (CAL alone)**


**Figure 11. Forest plot for the subgroup analysis of mean CAL between preeclamptic and healthy groups.**

**Figure 12. Forest plot for the subgroup analysis of mean PD between preeclamptic and healthy groups.**


**Figure 13. Funnel plot for the association between periodontitis and preeclampsia.**



**Figure 1. Flow diagram of study selection**




**Figure 2. Forest plot for the association between periodontitis and preeclampsia.**




**Figure 3. Forest plot for the subgroup analysis according to the type of study design (cohort study)**







**Figure 4. Forest plot for the subgroup analysis according to the type of study design (case-control study)**




**Figure 5. Forest plot for the subgroup analysis according to the national income (high income countries)**




**Figure 6. Forest plot for the subgroup analysis according to the national income (upper middle-income countries)**




**Figure 7. Forest plot for the subgroup analysis according to the national income (lower middle- income countries)**



**Figure 8. Forest plot for the subgroup analysis according to the definition of periodontitis (PD alone)**



**Figure 9. Forest plot for the subgroup analysis according to the definition of periodontitis (PD and CAL)**



**Figure 10. Forest plot for the subgroup analysis according to the definition of periodontitis (CAL alone)**






Figure 11. Forest plot for the subgroup analysis of mean CAL between preeclamptic and healthy groups.





**Figure 12. Forest plot for the subgroup analysis of mean PD between preeclamptic and healthy groups.**

healthy groups.



**Figure 13. Funnel plot for the association between periodontitis and preeclampsia.**

