REVERSE ORDER LAW FOR GENERALIZED INVERSES WITH INDEFINITE HERMITIAN WEIGHTS

K. KAMARAJ 1 , P. SAM JOHNSON 2 AND ATHIRA SATHEESH K 3

ABSTRACT. In this paper, necessary and sufficient conditions are given for the existence of Moore-Penrose inverse of a product of two matrices in an indefinite inner product space (IIPS) in which reverse order law holds good. Rank equivalence formulas with respect to IIPS are provided and an open problem is given at the end.

1. Introduction

The reverse order law for generalized inverse plays an important role in the theoretic research and numerical computations in many areas, including the singular matrix problems, ill-posed problems, optimization problems, and statistics problems (see, for instance, [1, 4, 7, 10, 9, 13, 6]). A classical result of Greville [12] gives necessary and sufficient conditions for the two term reverse order law for the Moore-Penrose inverse in the Euclidean space. It is known that the reverse order law does not hold for various classes of generalized inverses [2, 5]. Hence, a significant number of papers treat the sufficient or equivalent conditions such that the reverse order law holds in some sense. Sun and Wei established some sufficient and necessary conditions for inverse order rule for weighted generalized inverses with positive definite weights [14, 15]. The concept of the Moore-Penrose inverse between indefinite inner product spaces has been introduced and mentioned in [8] that if the weights are positive definite, then the weighted generalized inverse and the Moore-Penrose inverse between indefinite inner product spaces are the same. In this paper, we give some necessary and sufficient conditions for the existence of Moore-Penrose inverse of a product of two matrices and to hold reverse order law in an IIPS. Also, we claim that our results are more general than the existing ones for weighted Moore-Penrose inverse.

2. Preliminaries

We consider matrices on the field \mathbb{C} of complex numbers and denote the space of complex matrices of order $m \times n$ by $\mathbb{C}^{m \times n}$. The range and the rank of $A \in \mathbb{C}^{m \times n}$ are denoted by R(A) and rank(A) respectively. The index of $A \in \mathbb{C}^{n \times n}$ is the least positive integer p such that $rank(A^p) = rank(A^{p+1})$ and it is denoted by ind(A).

For a complex square matrix A, we call it *Hermitian* if $A = A^*$, where A^* denotes the adjoint of A with respect to the Hermitian inner product $\langle \cdot, \cdot \rangle$ on \mathbb{C}^n (i.e., complex conjugate transpose). Let N be an invertible Hermitian matrix of order n. An *indefinite inner product* in \mathbb{C}^n is defined by an equation

$$[x, y] = \langle x, Ny \rangle$$

where $x, y \in \mathbb{C}^n$. Such a matrix N is called a weight. A space with an indefinite inner product is called an indefinite inner product space (IIPS). Let M and N be weights of order m and n, respectively. The MN-adjoint of an $m \times n$ matrix A denoted $A^{[*]}$ is defined by

$$A^{[*]} = N^{-1}A^*M.$$

Sun and Wei [14] used the terminology weighted conjugate transpose for MN-adjoint. In an IIPS, by considering the same weights M = N, a complex square matrix A is called N-Hermitian if $A^{[*]} = A$; it is called N-range Hermitian if $R(A) = R(A^{[*]})$. If the IIPS is understood from the context, then

Date: December 10, 2021.

 $Key\ words\ and\ phrases.$ Moore-Penrose inverse, reverse order law, indefinite inner product space, weighted generalized inverse.

2

instead of saying that A is N-Hermitian (N-range Hermitian), we may simply say that A is Hermitian (range Hermitian).

The MN-Moore-Penrose inverse $A^{[\dagger]}$ of $A \in \mathbb{C}^{m \times n}$ between IIPSs is defined to be the unique solution $X \in \mathbb{C}^{n \times m}$, if it exists, to the equations

$$AXA = A \tag{2.1}$$

$$XAX = X (2.2)$$

$$(AX)^{[*]} = AX \tag{2.3}$$

$$(XA)^{[*]} = XA.$$
 (2.4)

The reference to MN will be dropped when there is no ambiguity and $A^{[\dagger]}$ will be simply called the Moore-Penrose inverse of A. If A is invertible, then $A^{[\dagger]} = A^{-1}$. It is easy to observe that if M and N are the identity matrices, then $A^{[\dagger]} = A^{\dagger}$, where A^{\dagger} denotes the usual Moore-Penrose inverse in an Euclidean space. Sun and Wei used the notation A^{\dagger}_{MN} for $A^{[\dagger]}$ to emphasize on the weights of positive definite Hermite matrices M and N. In this case, A^{\dagger}_{MN} exists for all matrices A and $A^{\dagger}_{MN} = N^{-\frac{1}{2}}(M^{\frac{1}{2}}AN^{-\frac{1}{2}})^{\dagger}M^{\frac{1}{2}}$ [14]. Unlike the Euclidean case and weighted Moore-Penrose inverse, a matrix need not have a Moore-Penrose inverse between IIPSs [8]. The following result gives a necessary and sufficient condition for the existence of Moore-Penrose inverse of a matrix between IIPSs.

Theorem 2.1 ([8], Theorem 1). Let $A \in \mathbb{C}^{m \times n}$. Then $A^{[\dagger]}$ exists iff $rank(A) = rank(AA^{[*]}) = rank(A^{[*]}A)$.

For the sake of clarity as well as for easier reference we mention the following properties of Moore-Penrose inverse between IIPSs.

Theorem 2.2 ([8], Section 4). Let $A \in \mathbb{C}^{m \times n}$ be such that $A^{[\dagger]}$ exists. Then the following statements hold:

- (i) $A^{[*]} = A^{[*]}AA^{[\dagger]} = A^{[\dagger]}AA^{[*]}$.
- (ii) $(A^{[*]})^{[\dagger]} = (A^{[\dagger]})^{[*]}$.
- (iii) $(AA^{[*]})^{[\dagger]}$ and $(A^{[*]}A)^{[\dagger]}$ exist. In this case, $(AA^{[*]})^{[\dagger]} = (A^{[*]})^{[\dagger]}A^{[\dagger]}$ and $(A^{[*]}A)^{[\dagger]} = A^{[\dagger]}(A^{[*]})^{[\dagger]}$.
- (iv) $A^{[\dagger]} = A^{[*]} (AA^{[*]})^{[\dagger]} = (A^{[*]}A)^{[\dagger]} A^{[*]}$.
- (v) $(AA^{[*]})^{[\dagger]}(AA^{[*]})A = A = (AA^{[*]})(AA^{[*]})^{[\dagger]}A.$
- (vi) $(AA^{[*]})^{[\dagger]}(AA^{[*]}) = (AA^{[*]})(AA^{[*]})^{[\dagger]}$.

This section is ended with some known results which will be used in the sequel.

Lemma 2.3 ([1], p.173). Let A be a square matrix of order n with ind(A) = 1. Let $B \in \mathbb{C}^{n \times \ell}$ be a matrix such that $R(AB) \subseteq R(B)$. Then

$$R(AB) = R(A) \cap R(B).$$

Lemma 2.4 ([15], Lemma 2.1). Let A, B, C and D be matrices with suitable orders. Then

$$rank\left(\begin{array}{cc}A & AB\\ CA & D\end{array}\right) = rank(A) + rank(D-CAB).$$

Lemma 2.5 ([17], Theorem 2.7). Let P and Q are two idempotent matrices of suitable orders. Then

$$\begin{split} rank(PQ-QP) &= rank\left(\begin{array}{c} P \\ Q \end{array}\right) + rank\left(\begin{array}{c} P \\ Q \end{array}\right) + rank(PQ) \\ &+ rank(QP) - 2rank(P) - 2rank(Q). \end{split}$$

Lemma 2.6 ([3], Corollary). Let $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. Then rank(M) = rank(A) if and only if $D - CA^{[\dagger]}B = 0$, $N(A) \subseteq N(C)$ and $N((A)^*) \subseteq N(B)^*$.

Lemma 2.7 ([16], Theorem 1.2). Let A, B, C and D be matrices with suitable orders. Then

$$rank \left(\begin{array}{cc} A^*AA^* & A^*B \\ CA^* & D \end{array} \right) = rank(A) + rank(D - CA^{\dagger}B).$$

3. Reverse Order Law

We start the section with examples which illustrate that between IIPSs, $(AB)^{[\dagger]}$ may not exist although $A^{[\dagger]}$ and $B^{[\dagger]}$ exist, and even though Moore-Penrose inverses of A,B and AB exist, the reverse order law $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$ does not hold.

Example 3.1. Let
$$A = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$
, $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $M = N = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Clearly, $B^{[*]} = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}$ and $(AB)^{[*]} = \begin{pmatrix} 0 & 0 \\ -1 & 1 \end{pmatrix}$. Then A is non-singular and $rank(B) = rank(BB^{[*]}) = rank(B^{[*]}B)$, so both $A^{[\dagger]}$ and $B^{[\dagger]}$ exist. Also, $AB = \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$ and $rank(AB) \neq rank((AB)^{[*]}AB)$, hence $(AB)^{[\dagger]}$ does not exist.

$$\begin{array}{l} \textbf{Example 3.2. Let } A = \begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}, \ B = \begin{pmatrix} 2 & 1 \\ 0 & 0 \end{pmatrix} \ \text{and } M = N = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \ \text{Clearly, } A^{[*]} = \\ \begin{pmatrix} 1 & 0 \\ -2 & 0 \end{pmatrix} \ \text{and } B^{[*]} = \begin{pmatrix} 2 & 0 \\ -1 & 0 \end{pmatrix}. \ \text{Then } AA^{[*]} = \begin{pmatrix} -3 & 0 \\ 0 & 0 \end{pmatrix}, A^{[*]}A = \begin{pmatrix} 1 & 2 \\ -2 & -4 \end{pmatrix}, \\ BB^{[*]} = \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix} \ \text{and } B^{[*]}B = \begin{pmatrix} 4 & 2 \\ -2 & -1 \end{pmatrix}. \ \text{Hence } A^{[\dagger]} = -\frac{1}{3}\begin{pmatrix} 1 & 0 \\ -2 & 0 \end{pmatrix} \ \text{and } B^{[\dagger]} = \frac{1}{3}\begin{pmatrix} 2 & 0 \\ -1 & 0 \end{pmatrix}. \\ \text{Moreover, } (AB)^{[\dagger]} = \frac{1}{3}\begin{pmatrix} 2 & 0 \\ -1 & 0 \end{pmatrix} \ \text{and } B^{[\dagger]}A^{[\dagger]} = -\frac{1}{9}\begin{pmatrix} 2 & 0 \\ -1 & 0 \end{pmatrix}. \ \text{Thus } (AB)^{[\dagger]} \neq B^{[\dagger]}A^{[\dagger]}. \end{array}$$

Motivated by the above examples, we show when the reverse order law holds good in an indefinite inner product space. Before presenting the main results, we collect some basic results.

Lemma 3.3. Let A, B, C and D be matrices with suitable orders. If

$$rank \left(egin{array}{cc} A & B \\ C & D \end{array} \right) = rank(A) = rank(B) = rank(C),$$

then rank(A) = rank(D).

Proof. Let $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. It is given that rank(M) = rank(A). Then by Lemma 2.6, we have $D - CA^{\dagger}B = 0$, $R(C^*) \subseteq R(A^*)$ and $R(B) \subseteq R(A)$. This implies $D = CA^{\dagger}B \implies rank(D) \le rank(C) = rank(A)$.

Now to prove the reverse inequality, $rank(A) = rank(B) = rank(C) \implies R(A) = R(B)$ and $R(A^*) = R(C^*)$. Since $D = CA^{\dagger}B$ we get $C^{\dagger}D = C^{\dagger}CA^{\dagger}B = C^{\dagger}(C^{\dagger})^{\dagger}A^{\dagger}B$. Now, $R(A^*) = R(C^*) \implies R(A^{\dagger}) = R(C^{\dagger})$. Using the fact that if $R(E) \subseteq R(F)$ then $FF^{\dagger}E = E$, we get $C^{\dagger}(C^{\dagger})^{\dagger}A^{\dagger} = A^{\dagger}$. This implies $C^{\dagger}D = A^{\dagger}B \implies C^{\dagger}DB^{\dagger} = A^{\dagger}BB^{\dagger} = A^{\dagger}(B^{\dagger})^{\dagger}B^{\dagger}$. Now, $R(A) = R(B) \implies R((A^{\dagger})^*) = R((B^{\dagger})^*)$ and using the fact that if $R(E^*) \subseteq R(F^*)$ then $EF^{\dagger}F = E$ we get, $C^{\dagger}DB^{\dagger} = A^{\dagger} \implies rank(A^{\dagger}) \le rank(D) \implies rank(A) \le rank(D)$. This completes the proof. \square

Next we prove the indefinite version of Lemma 2.7.

Theorem 3.4. Let A, B, C and D be matrices with suitable orders. If $A^{[\dagger]}$ exists, then

$$rank \begin{pmatrix} A^{[*]}AA^{[*]} & A^{[*]}B \\ CA^{[*]} & D \end{pmatrix} = rank \begin{pmatrix} D & CA^{[*]} \\ A^{[*]}B & A^{[*]}AA^{[*]} \end{pmatrix}$$
$$= rank(A) + rank(D - CA^{[\dagger]}B).$$

Proof. By Theorem 2.2 we can easily verify the following relations.

$$\left(\begin{array}{cc} (A^{[\dagger]})^{[*]} & 0 \\ 0 & I \end{array} \right) \left(\begin{array}{cc} A^{[*]}AA^{[*]} & A^{[*]}B \\ CA^{[*]} & D \end{array} \right) \left(\begin{array}{cc} (A^{[\dagger]})^{[*]} & 0 \\ 0 & I \end{array} \right) = \left(\begin{array}{cc} A & AA^{[\dagger]}B \\ CA^{[\dagger]}A & D \end{array} \right)$$

and

$$\left(\begin{array}{cc}A^{[*]}&0\\0&I\end{array}\right)\left(\begin{array}{cc}A&AA^{[\dagger]}B\\CA^{[\dagger]}A&D\end{array}\right)\left(\begin{array}{cc}A^{[*]}&0\\0&I\end{array}\right)=\left(\begin{array}{cc}A^{[*]}AA^{[*]}&A^{[*]}B\\CA^{[*]}&D\end{array}\right).$$

Thus

$$\begin{aligned} rank \left(\begin{array}{cc} A^{[*]}AA^{[*]} & A^{[*]}B \\ CA^{[*]} & D \end{array} \right) & = rank \left(\begin{array}{cc} A & AA^{[\dagger]}B \\ CA^{[\dagger]}A & D \end{array} \right) \\ & = rank(A) + rank(D - CA^{[\dagger]}AA^{[\dagger]}B) \text{ (by Lemma 2.4)} \\ & = rank(A) + rank(D - CA^{[\dagger]}B). \end{aligned}$$

Similarly we can prove the other equality.

It is known in the Euclidean case that the single expression $R(A^*ABB^*) = R(BB^*A^*A)$ is a necessary and sufficient condition for the reverse order law to hold ([1], p.161). This condition was later shown ([11], p.231) to hold in a more general setting. The main result and its proof closely follow those of Greville [12].

Theorem 3.5. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$. If $A^{[\dagger]}$ and $B^{[\dagger]}$ exist, then the following are equivalent:

- (i) $A^{[*]}ABB^{[*]}$ is range Hermitian.
- (ii) $R(A^{[*]}AB) \subseteq R(B)$ and $R(BB^{[*]}A^{[*]}) \subseteq R(A^{[*]})$.
- (iii) $BB^{[\dagger]}A^{[\ast]}A$ and $A^{[\dagger]}ABB^{[\ast]}$ are range Hermitian.
- (iv) $BB^{[\dagger]}A^{[*]}AB = A^{[*]}AB$ and $A^{[\dagger]}ABB^{[*]}A^{[*]} = BB^{[*]}A^{[*]}$.

Proof. $\underline{(i) \Rightarrow (ii)}$: As $B = BB^{[\dagger]}B = BB^{[\ast]}(B^{[\dagger]})^{[\ast]}$, we have $R(A^{[\ast]}ABB^{[\ast]}) = R(A^{[\ast]}AB)$. Suppose that $A^{[\ast]}ABB^{[\ast]}$ is range Hermitian. Then

$$R(A^{[*]}AB) = R(A^{[*]}ABB^{[*]}) = R(BB^{[*]}A^{[*]}A) \subseteq R(B).$$

The second part follows similarly.

 $(ii) \Leftrightarrow (iv)$: Straight forward.

 $\overline{(ii) \Rightarrow (iii)}$: Suppose that $R(A^{[*]}AB) \subseteq R(B)$. As $R(A^{[*]}ABB^{[\dagger]}) \subseteq R(A^{[*]}AB) \subseteq R(B)$, we get $A^{[*]}ABB^{[\dagger]} = BB^{[\dagger]}A^{[*]}ABB^{[\dagger]}$ and hence it can be shown that

$$R((BB^{[\dagger]}A^{[*]}A)^{[*]}) = R(A^{[*]}ABB^{[\dagger]}) = R(BB^{[\dagger]}A^{[*]}A).$$

Thus $BB^{[\dagger]}A^{[*]}A$ is range Hermitian. In a similar way, using the inclusion relation

$$R(BB^{[*]}A^{[*]}) \subseteq R(A^{[*]}),$$

we can prove that $A^{[\dagger]}ABB^{[*]}$ is also range Hermitian.

 $\underbrace{(iii) \Rightarrow (ii):} \text{Suppose } BB^{[\dagger]}A^{[*]}A \text{ is range Hermitian. Then } R(BB^{[\dagger]}A^{[*]}A) = R(A^{[*]}ABB^{[\dagger]}). \text{ It clear } \text{that } R(A^{[*]}AB) = R(A^{[*]}ABB^{[\dagger]}B) \subseteq R(A^{[*]}ABB^{[\dagger]}) = R(BB^{[\dagger]}A^{[*]}A) \subseteq R(B). \text{ Thus } BB^{[\dagger]}A^{[*]}AB = A^{[*]}AB. \text{ Similarly, we can prove } A^{[\dagger]}ABB^{[*]}A^{[*]} = BB^{[*]}A^{[*]}.$

Theorem 3.6. Let $A \in \mathbb{C}^{m \times n}$, $B \in \mathbb{C}^{n \times \ell}$ and D = AB. If $A^{[\dagger]}$ and $B^{[\dagger]}$ exist, then the following are equivalent:

(i)
$$rank \begin{pmatrix} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{pmatrix} = rank(D), where D = AB.$$

(ii) $(AB)^{[\dagger]}$ exists and $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$.

Proof. $(i) \Rightarrow (ii)$: First we prove the existence of Moore-Penrose inverse of AB. For that, let $E = AA^{[*]}D$. It is easy to observe that $D = AB = (AA^{[*]})^{[\dagger]}(AA^{[*]})AB = (AA^{[*]})^{[\dagger]}E$. Thus $rank(D) = rank((AA^{[*]})^{[\dagger]}E) \leq rank(E) = rank(AA^{[*]}D) \leq rank(D)$. It shows that $rank(D) = rank(AA^{[*]}D)$. Similarly, we can prove that $rank(D) = rank(DB^{[*]}B)$.

Suppose $rank \begin{pmatrix} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{pmatrix} = rank(D)$. Then $rank(D) = rank(DD^{[*]}D)$ by Lemma 3.3. It concludes that $rank(D) = rank(DD^{[*]}) = rank(D^{[*]}D)$. Thus $(AB)^{[\dagger]}$ exists. By Theorem 3.4,

$$rank \begin{pmatrix} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{pmatrix} = rank(D^{[*]}) + rank(D - AA^{[*]}(D^{[*]})^{[\dagger]}B^{[*]}B)$$
$$= rank(D) + rank(D^{[*]} - B^{[*]}BD^{[\dagger]}AA^{[*]}).$$

Hence, by the assumption $rank(D^{[*]} - B^{[*]}BD^{[\dagger]}AA^{[*]}) = 0$. Thus $D^{[*]} = B^{[*]}BD^{[\dagger]}AA^{[*]}$. Premultiplying by $(B^{[*]}B)^{[\dagger]}$ and post-multiplying by $(AA^{[*]})^{[\dagger]}$ we get

$$(B^{[*]}B)^{[\dagger]}B^{[*]}A^{[*]}(AA^{[*]})^{[\dagger]} = (B^{[*]}B)^{[\dagger]}B^{[*]}BD^{[\dagger]}AA^{[*]}(AA^{[*]})^{[\dagger]}.$$

By Theorem 2.2 (v) and (vi),

$$\begin{split} B^{[\dagger]}A^{[\dagger]} &= (B^{[*]}B)^{[\dagger]}B^{[*]}BB^{[*]}A^{[*]}(DD^{[*]})^{[\dagger]}AA^{[*]}(AA^{[*]})^{[\dagger]} \\ &= B^{[*]}A^{[*]}(DD^{[*]})^{[\dagger]}AA^{[*]}(AA^{[*]})^{[\dagger]} \\ &= D^{[\dagger]}AA^{[*]}(AA^{[*]})^{[\dagger]} = (D^{[*]}D)^{[\dagger]}D^{[*]}AA^{[*]}(AA^{[*]})^{[\dagger]} \\ &= D^{[\dagger]} = (AB)^{[\dagger]}. \end{split}$$

 $(ii) \Rightarrow (i)$: By Theorem 3.4,

$$\begin{aligned} rank \left(\begin{array}{cc} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{array} \right) & = rank(D^{[*]}) + rank(D - AA^{[*]}(D^{[*]})^{[\dagger]}B^{[*]}B) \\ & = rank(D) + rank(D - AA^{[*]}(B^{[\dagger]}A^{[\dagger]})^{[*]}B^{[*]}B) \\ & = rank(D) + rank(D - AA^{[*]}(A^{[\dagger]})^{[*]}(B^{[\dagger]})^{[*]}B^{[*]}B) \\ & = rank(D) + rank(D - AB) \\ & = rank(D). \end{aligned}$$

Theorem 3.7. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$ such that $A^{[\dagger]}$ and $B^{[\dagger]}$ exist. If any one of the conditions listed in Theorem 3.5 holds, then

$$rank \left(\begin{array}{cc} D & A^{[*]}AD \\ DB^{[*]}B & DD^{[*]}D \end{array} \right) = rank(D).$$

Proof. Suppose that $BB^{[\dagger]}A^{[*]}AB = A^{[*]}AB$ and $A^{[\dagger]}ABB^{[*]}A^{[*]} = BB^{[*]}A^{[*]}$ hold. Then, we have

$$\begin{pmatrix} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{pmatrix} = \begin{pmatrix} AB & ABB^{[\dagger]}A^{[*]}AB \\ ABB^{[*]}A^{[\dagger]}AB & AB(AB)^{[*]}AB \end{pmatrix}$$

$$= \begin{pmatrix} D & DB^{[\dagger]}A^{[*]}D \\ DB^{[*]}A^{[\dagger]}D & DD^{[*]}D \end{pmatrix}.$$

By Theorem 3.4,

$$rank \left(\begin{array}{ccc} D & AA^{[*]}D \\ DB^{[*]}B & DD^{[*]}D \end{array} \right) & = rank \left(\begin{array}{ccc} D & DB^{[\dagger]}A^{[*]}D \\ DB^{[*]}A^{[\dagger]}D & DD^{[*]}D \end{array} \right) \\ & = rank(D) + rank(DD^{[*]}D - DB^{[*]}A^{[\dagger]}DB^{[\dagger]}A^{[*]}D) \\ & = rank(D) + rank(DD^{[*]}D - ABB^{[*]}A^{[\dagger]}ABB^{[\dagger]}A^{[*]}AB) \\ & = rank(D) + rank(DD^{[*]}D - ABB^{[*]}A^{[\dagger]}AA^{[*]}AB) \\ & = rank(D) + rank(DD^{[*]}D - ABB^{[*]}A^{[*]}AB) \\ & = rank(D) + rank(DD^{[*]}D - DD^{[*]}D) \\ & = rank(D). \\ \end{aligned}$$

Corollary 3.8. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$ such that $A^{[\dagger]}$ and $B^{[\dagger]}$ exist. If any one of the conditions listed in Theorem 3.5 holds, then $(AB)^{[\dagger]}$ exists and $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$.

Lemma 3.9. Let $A \in \mathbb{C}^{m \times n}$, $B \in \mathbb{C}^{m \times \ell}$ and $C \in \mathbb{C}^{\ell \times n}$. Then

(i)

$$rank \left(\begin{array}{cc} A & B \end{array} \right) = rank \left(\begin{array}{cc} A^{[*]} \\ B^{[*]} \end{array} \right)$$

(ii)

$$rank \left(\begin{array}{c} A \\ C \end{array} \right) = rank \left(\begin{array}{cc} A^{[*]} & C^{[*]} \end{array} \right).$$

Proof.

$$\begin{split} rank \left(\begin{array}{c} A^{[*]} \\ B^{[*]} \end{array} \right) &= rank \left(\begin{array}{c} N^{-1}A^*M \\ L^{-1}B^*M \end{array} \right) &= rank \left(\begin{array}{c} N^{-1}A^* \\ L^{-1}B^* \end{array} \right) \quad M \quad \\ &= rank \left(\begin{array}{c} N^{-1}A^* \\ L^{-1}B^* \end{array} \right) \\ &= rank \left(\begin{array}{c} AN^{-1} & BL^{-1} \\ 0 & L^{-1} \end{array} \right) \\ &= rank \left(\begin{array}{c} A & B \end{array} \right) \left(\begin{array}{c} N^{-1} & 0 \\ 0 & L^{-1} \end{array} \right) \quad \\ &= rank \left(\begin{array}{c} A & B \end{array} \right). \end{split}$$

Similarly we can prove (ii).

Lemma 3.10. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{g \times h}$. If $A^{[\dagger]}$ and $B^{[\dagger]}$ exist, then

(i)

$$\left(\begin{array}{cc}A&0\\0&B\end{array}\right)^{[*]}=\left(\begin{array}{cc}A^{[*]}&0\\0&B^{[*]}\end{array}\right)$$

(ii)

$$\left(\begin{array}{cc} A & 0 \\ 0 & B \end{array}\right)^{[\dagger]} = \left(\begin{array}{cc} A^{[\dagger]} & 0 \\ 0 & B^{[\dagger]} \end{array}\right)$$

(iii)

$$\left(\begin{array}{cc} 0 & A \\ B & 0 \end{array}\right)^{[\dagger]} = \left(\begin{array}{cc} 0 & B^{[\dagger]} \\ A^{[\dagger]} & 0 \end{array}\right).$$

Proof. (i) Let $K = \begin{pmatrix} M & 0 \\ 0 & G \end{pmatrix}$ and $L = \begin{pmatrix} N & 0 \\ 0 & H \end{pmatrix}$. Without loss of generality we may assume that

$$T^{[*]} = L^{-1}T^*K, \text{where } T = \left(\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right).$$

Then

$$\begin{split} T^{[*]} & &= \left(\begin{array}{cc} N^{-1} & 0 \\ 0 & H^{-1} \end{array} \right) \left(\begin{array}{cc} A^* & 0 \\ 0 & B^* \end{array} \right) \left(\begin{array}{cc} M & 0 \\ 0 & G \end{array} \right) \\ & &= \left(\begin{array}{cc} N^{-1}A^*M & 0 \\ 0 & H^{-1}B^*G \end{array} \right) \\ & &= \left(\begin{array}{cc} A^{[*]} & 0 \\ 0 & B^{[*]} \end{array} \right). \end{split}$$

(ii) Suppose $A^{[\dagger]}$ and $B^{[\dagger]}$ exist.

$$T^{[*]}T = \begin{pmatrix} A^{[*]}A & 0 \\ 0 & B^{[*]}B \end{pmatrix} \text{ and } TT^{[*]} = \begin{pmatrix} AA^{[*]} & 0 \\ 0 & BB^{[*]} \end{pmatrix}.$$

Thus $rank(T^{[*]}T) = rank(A) + rank(B) = rank(TT^{[*]}) = rank(T)$, which implies $T^{[\dagger]}$ exists. Also it is easy to verify that $T^{[\dagger]} = \begin{pmatrix} A^{[\dagger]} & 0 \\ 0 & B^{[\dagger]} \end{pmatrix}$ satisfies the Moore-Penrose equations.

(iii) is similiar to (ii).

Theorem 3.11. Let A, B, C, D, P and Q be matrices with suitable orders such that $P^{[\dagger]}$ and $Q^{[\dagger]}$ exist. Then

$$rank(D-CP^{[\dagger]}AQ^{[\dagger]}B) = rank \left(\begin{array}{ccc} P^{[*]}AQ^{[*]} & P^{[*]}PP^{[*]} & 0 \\ Q^{[*]}QQ^{[*]} & 0 & Q^{[*]}B \\ 0 & CP^{[*]} & -D \end{array} \right) - rank(P) - rank(Q).$$

Proof. It is observed that

$$\begin{pmatrix} A & AQ^{[\dagger]}B \\ CP^{[\dagger]}A & D \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} + \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix} \begin{pmatrix} 0 & Q^{[\dagger]} \\ P^{[\dagger]} & 0 \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$$

$$= \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} + \begin{pmatrix} A & 0 \\ 0 & C \end{pmatrix} \begin{pmatrix} 0 & P \\ Q & 0 \end{pmatrix}^{[\dagger]} \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}$$

(by Lemma 3.10 (iii)).

Thus by Theorem 3.4,

$$rank\left(\begin{array}{cc}A & AQ^{[\dagger]}B\\ CP^{[\dagger]}A & D\end{array}\right) \ = \ rank\left(\begin{array}{cc}M^{[*]}MM^{[*]} & M^{[*]}\begin{pmatrix}A & 0\\ 0 & B\end{pmatrix}\\ \begin{pmatrix}A & 0\\ 0 & C\end{pmatrix}M^{[*]} & \begin{pmatrix}-A & 0\\ 0 & -D\end{pmatrix}\right) - rank\left(\begin{array}{cc}0 & P\\ Q & 0\end{array}\right),$$

where $M=\left(\begin{array}{cc} 0 & P \\ Q & 0 \end{array}\right)$. By Lemma 3.10 (ii),

$$rank \left(\begin{array}{ccc} A & AQ^{[\dagger]}B \\ CP^{[\dagger]}A & D \end{array} \right) = rank \left(\begin{array}{cccc} 0 & Q^{[\ast]}QQ^{[\ast]} & 0 & Q^{[\ast]}B \\ P^{[\ast]}PP^{[\ast]} & 0 & P^{[\ast]}A & 0 \\ 0 & AQ^{[\ast]} & -A & 0 \\ CP^{[\ast]} & 0 & 0 & -D \end{array} \right)$$
$$-rank(P) - rank(Q).$$

Also

$$\begin{pmatrix} 0 & I & P^{[*]} & 0 \\ I & 0 & 0 & 0 \\ 0 & 0 & 0 & I \\ 0 & 0 & I & 0 \end{pmatrix} \begin{pmatrix} 0 & Q^{[*]}QQ^{[*]} & 0 & Q^{[*]}B \\ P^{[*]}PP^{[*]} & 0 & P^{[*]}A & 0 \\ 0 & AQ^{[*]} & -A & 0 \\ CP^{[*]} & 0 & 0 & -D \end{pmatrix} \begin{pmatrix} 0 & I & 0 & 0 \\ I & 0 & 0 & 0 \\ Q^{[*]} & 0 & 0 & I \\ 0 & 0 & I & 0 \end{pmatrix}$$

$$= \begin{pmatrix} P^{[*]}AQ^{[*]} & P^{[*]}PP^{[*]} & 0 & 0 \\ Q^{[*]}QQ^{[*]} & 0 & Q^{[*]}B & 0 \\ 0 & CP^{[*]} & -D & 0 \\ 0 & 0 & 0 & -A \end{pmatrix}.$$

Thus

$$rank \left(\begin{array}{c} A & AQ^{[\dagger]}B \\ CP^{[\dagger]}A & D \end{array} \right) = rank \left(\begin{array}{ccc} P^{[\ast]}AQ^{[\ast]} & P^{[\ast]}PP^{[\ast]} & 0 & 0 \\ Q^{[\ast]}QQ^{[\ast]} & 0 & Q^{[\ast]}B & 0 \\ 0 & CP^{[\ast]} & -D & 0 \\ 0 & 0 & 0 & -A \end{array} \right)$$

$$-rank(P) - rank(Q)$$

$$= rank \left(\begin{array}{ccc} P^{[\ast]}AQ^{[\ast]} & P^{[\ast]}PP^{[\ast]} & 0 \\ Q^{[\ast]}QQ^{[\ast]} & 0 & Q^{[\ast]}B \\ 0 & CP^{[\ast]} & -D \end{array} \right) + rank(A)$$

$$-rank(P) - rank(Q).$$

But by Lemma 2.4,

$$rank\left(\begin{array}{cc}A & AQ^{[\dagger]}B\\ CP^{[\dagger]}A & D\end{array}\right) \qquad = rank(A) + rank(D - CP^{[\dagger]}AQ^{[\dagger]}B).$$

Thus

$$rank(D - CP^{[\dagger]}AQ^{[\dagger]}B) = rank \begin{pmatrix} P^{[\ast]}AQ^{[\ast]} & P^{[\ast]}PP^{[\ast]} & 0 \\ Q^{[\ast]}QQ^{[\ast]} & 0 & Q^{[\ast]}B \\ 0 & CP^{[\ast]} & -D \end{pmatrix} - rank(P) - rank(Q).$$

Corollary 3.12. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$ such that $A^{[\dagger]}$ and $B^{[\dagger]}$ exist. Then

$$rank(AB-ABB^{[\dagger]}A^{[\dagger]}AB)=rank\left(\begin{array}{cc}B^{[\ast]}A^{[\ast]}&B^{[\ast]}B\\AA^{[\ast]}&AB\end{array}\right)+rank(AB)-rank(A)-rank(B).$$

Proof. Replace D by AB, C by AB, P by B, A by I, Q by A and B by AB in Theorem 3.11, we get

$$rank(AB - ABB^{[\dagger]}A^{[\dagger]}AB) = rank \begin{pmatrix} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} & 0 \\ A^{[*]}AA^{[*]} & 0 & A^{[*]}AB \\ 0 & ABB^{[*]} & -AB \end{pmatrix} - rank(A) - rank(B).$$

Also,

$$\begin{pmatrix} I & 0 & 0 \\ 0 & I & A^{[*]} \\ 0 & 0 & I \end{pmatrix} \begin{pmatrix} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} & 0 \\ A^{[*]}AA^{[*]} & 0 & A^{[*]}AB \\ 0 & ABB^{[*]} & -AB \end{pmatrix} \begin{pmatrix} I & 0 & 0 \\ 0 & I & 0 \\ 0 & B^{[*]} & I \end{pmatrix}$$

$$= \begin{pmatrix} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} & 0 \\ A^{[*]}AA^{[*]} & A^{[*]}ABB^{[*]} & 0 \\ 0 & 0 & -AB \end{pmatrix}$$

Therefore

$$rank(AB - ABB^{[\dagger]}A^{[\dagger]}AB) = rank \begin{pmatrix} B^{[\ast]}A^{[\ast]} & B^{[\ast]}BB^{[\ast]} \\ A^{[\ast]}AA^{[\ast]} & A^{[\ast]}ABB^{[\ast]} \end{pmatrix} + rank(AB) - rank(A) - rank(B).$$

Moreover, by observing the following facts

$$\left(\begin{array}{cc} I & 0 \\ 0 & A^{[\dagger]}^{[*]} \end{array} \right) \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} \\ A^{[*]}AA^{[*]} & A^{[*]}ABB^{[*]} \end{array} \right) \left(\begin{array}{cc} I & 0 \\ 0 & B^{[\dagger]}^{[*]} \end{array} \right) = \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right)$$

and

$$\left(\begin{array}{cc} I & 0 \\ 0 & A^{[*]} \end{array} \right) \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right) \left(\begin{array}{cc} I & 0 \\ 0 & B^{[*]} \end{array} \right) = \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} \\ A^{[*]}AA^{[*]} & A^{[*]}ABB^{[*]} \end{array} \right),$$

we have

$$rank \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right) = rank \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}BB^{[*]} \\ A^{[*]}AA^{[*]} & A^{[*]}ABB^{[*]} \end{array} \right).$$

Thus

$$rank(AB-ABB^{[\dagger]}A^{[\dagger]}AB)=rank\left(\begin{array}{cc}B^{[\ast]}A^{[\ast]}&B^{[\ast]}B\\AA^{[\ast]}&AB\end{array}\right)+rank(AB)-rank(A)-rank(B).$$

Lemma 3.13. Let P and Q be two N-Hermitian idempotent matrices of suitable orders. Then $rank(PQ - QP) = 2 \ rank\left(\begin{array}{cc} P & Q \end{array}\right) + 2 \ rank(PQ) - 2 \ rank(P) - 2 \ rank(Q).$

Proof. Since P and Q are two idempotent matrices, by Lemma 2.5,

$$\begin{aligned} rank(PQ-QP) &= rank\left(\begin{array}{c} P \\ Q \end{array}\right) + rank\left(\begin{array}{c} P \\ Q \end{array}\right) + rank(PQ) \\ &+ rank(QP) - 2 \ rank(P) - 2 \ rank(Q). \end{aligned}$$

By Lemma 3.9,

$$\begin{aligned} rank(PQ-QP) &= rank\left(\begin{array}{cc} P^{[*]} & Q^{[*]} \end{array}\right) + rank\left(\begin{array}{cc} P & Q \end{array}\right) + rank(PQ) \\ &+ rank(P^{[*]}Q^{[*]}) - 2 \ rank(P) - 2 \ rank(Q). \end{aligned}$$

Since P and Q are Hermitian we get,

$$rank(PQ-QP) = 2\; rank\left(\begin{array}{cc} P & Q \end{array}\right) + 2\; rank(PQ) - 2\; rank(P) - 2\; rank(Q).$$

 $\textbf{Lemma 3.14.} \ \textit{If} \ A^{[\dagger]} \ \textit{and} \ B^{[\dagger]} \ \textit{exist, then } rank \left(\begin{array}{cc} BB^{[\dagger]} \end{array} \right. A^{[\dagger]}A \ \left. \right) = rank \left(\begin{array}{cc} B \end{array} \right. A^{[\ast]} \ \left. \right).$

Proof. The conclusion may be arrived easily by using the following two equations

$$\begin{pmatrix} BB^{[\dagger]} & A^{[\dagger]}A \end{pmatrix} \begin{pmatrix} B & 0 \\ 0 & A^{[*]} \end{pmatrix} = \begin{pmatrix} B & A^{[*]} \end{pmatrix} \text{ and }$$

$$\begin{pmatrix} B & A^{[*]} \end{pmatrix} \begin{pmatrix} B^{[\dagger]} & 0 \\ 0 & (A^{[\dagger]})^{[*]} \end{pmatrix} = \begin{pmatrix} BB^{[\dagger]} & A^{[\dagger]}A \end{pmatrix}.$$

Theorem 3.15. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$. If $A^{[\dagger]}$ and $B^{[\dagger]}$ exist, then $rank(BB^{[\dagger]}A^{[\dagger]}A - A^{[\dagger]}ABB^{[\dagger]}) = 2 \ rank\left(\begin{array}{cc} A^{[\ast]} & B \end{array}\right) + 2 \ rank(AB) - 2 \ rank(A) - 2 \ rank(B).$

Proof. Clearly $A^{[\dagger]}A$ and $BB^{[\dagger]}$ are Hermitian and idempotent, then by Lemma 3.13,

$$\begin{split} rank(BB^{[\dagger]}A^{[\dagger]}A - A^{[\dagger]}ABB^{[\dagger]}) &= 2\; rank\left(\; BB^{[\dagger]} \quad A^{[\dagger]}A \;\right) + 2\; rank(BB^{[\dagger]}A^{[\dagger]}A) \\ &- 2\; rank(BB^{[\dagger]}) - 2\; rank(A^{[\dagger]}A) \\ &= 2\; rank\left(\; B \quad A^{[\ast]} \;\right) + 2\; rank(AB) \\ &- 2\; rank(B) - 2\; rank(A) \;\; \text{(by Lemma 3.14)}. \end{split}$$

Also,

$$\begin{aligned} rank(BB^{[\dagger]}A^{[\dagger]}A) & = rank(B(B^{[\ast]}B)^{[\dagger]}B^{[\ast]}A^{[\ast]}(AA^{[\ast]})^{[\dagger]}A) \\ & \leq rank(B^{[\ast]}A^{[\ast]}) = rank(AB) \\ \text{and } rank(AB) = rank(B^{[\ast]}A^{[\ast]}) & = rank(B^{[\ast]}BB^{[\dagger]}A^{[\dagger]}AA^{[\ast]}) \\ & \leq rank(BB^{[\dagger]}A^{[\dagger]}A). \end{aligned}$$

Thus $rank(AB) = rank(BB^{[\dagger]}A^{[\dagger]}A)$.

Theorem 3.16. Let $A \in \mathbb{C}^{m \times n}$ and $B \in \mathbb{C}^{n \times \ell}$ such that $A^{[\dagger]}$ and $B^{[\dagger]}$ exist. If

$$rank \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right) = rank \left(\begin{array}{cc} A^{[*]} & B \end{array} \right),$$

then $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$ is equivalent to any one of the conditions given in Theorem 3.5.

Proof. Since

$$rank \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right) = rank \left(\begin{array}{cc} A^{[*]} & B \end{array} \right),$$

by Theorem 3.15 and Corollary 3.12,

$$2 \operatorname{rank}(AB - ABB^{[\dagger]}A^{[\dagger]}AB) = \operatorname{rank}(BB^{[\dagger]}A^{[\dagger]}A - A^{[\dagger]}ABB^{[\dagger]}).$$

Thus if $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$, then $BB^{[\dagger]}A^{[\dagger]}A = A^{[\dagger]}ABB^{[\dagger]}$. Therefore

$$BB^{[*]}A^{[*]} = BB^{[\dagger]}A^{[\dagger]}ABB^{[*]}B^{[*]}A^{[*]} = A^{[\dagger]}ABB^{[\dagger]}BB^{[*]}A^{[*]} = A^{[\dagger]}ABB^{[*]}A^{[*]}.$$

Similarly, we prove $BB^{[\dagger]}A^{[*]}AB = A^{[*]}AB$. Thus we obtain condition (iv) of Theorem 3.5.

4. AN OPEN PROBLEM

We can observe from Theorem 3.16 that $(AB)^{[\dagger]} = B^{[\dagger]}A^{[\dagger]}$ is equivalent to any one of the conditions given in Theorem 3.5 by assuming the rank equality

$$rank \left(\begin{array}{cc} B^{[*]}A^{[*]} & B^{[*]}B \\ AA^{[*]} & AB \end{array} \right) = rank \left(\begin{array}{cc} A^{[*]} & B \end{array} \right).$$

But in the Euclidean case such a rank equality assumption is not required. Thus it is an open question for giving proof for Theorem 3.16 without assuming the rank equality, or finding a counter example.

References

- [1] Adi Ben-Israel, Thomas N.E. Greville, Generalized inverses: Theory and applications, Second Edition, Springer-Verlag, New York, 2003.
- [2] Chongguang Cao, Xian Zhang, Xiaomin Tang, Reverse order law of group inverses of products of two matrices, Appl. Math. Comput. 158 (2004) 489-495.
- [3] David Carlson, Emile Haynsworth and Thomas Markham, A Generalization of the Schur Complement by Means of the Moore-Penrose Inverse, SIAM Journal on Applied Mathematics Vol. 26, No. 1 (Jan., 1974), pp. 169-175.
- [4] Gene H. Golub, Charles F. Van Loan, Matrix Computations, The John Hopkins University Press, Baltimore, MD, 1983.
- [5] Guorong Wang, The reverse order law for the Drazin inverses of multiple matrix products, Lin. Alg. Appl. 348 (2002) 265–272.

- [6] Hans Joachim Werner, G-inverse of matrix products, in: S. Schach, G. Trenkler (Eds.), Data Analysis and Statistical Inference, Eul Verlag, Bergisch-Gladbach, 1992, pp.531-546.
- [7] Israel Gohberg, Peter Lancaster, Leiba Rodman, Indefinite linear algebra and applications, Birkhäuser Verlag, Basel, 2005.
- [8] Kamaraj K, Sivakumar K.C, Moore-Penrose inverse in an indefinite inner product space, J. Appl. Math. and Computing 19 (2005) 297-310.
- [9] Radhakrishna Rao C, Sujit Kumar Mitra, Generalized inverse of matrices and its applications, Wiley, New York,
- [10] Radojević, Ivana M. and Djordjević, Dragan S., Moore-Penrose inverse in indefinite inner product spaces, Filomat, 31:12 (2017), 3847-3857.
- [11] Robert E. Hartwig, Block generalized inverses, Arch. Rational Mech. Anal. 61 (1976) 197-251.
- [12] Thomas N.E. Greville, Note on the generalized inverse of a matrix product, SIAM Review, 8 (1966) 518-521.
- [13] Wenyu Sun, Ya-Xiang Yuan, Optimization Theory and Methods, Science Press, Beijing, 1996.
- [14] Wenyu Sun, Yimin Wei, Inverse order rule for weighted generalized inverse, SIAM. J. Matrix Anal. Appl. 19 (1998) 772-775
- [15] Wenyu Sun, Yimin Wei, Triple reverse-order law for weighted generalized inverse, Appl. Math. Comput. 125 (2002) 221-229.
- [16] Yongge Tian, How to characterize equalities for the Moore-Penrose Inverse of a matrix, Kyungpook Math. J. 41 (2001) 1-15.
- [17] Yongge Tian, George P.H. Styan, Rank equalities for idempotent and involutory matrices, Linear Algebra Appl. 335 (2001), 101-117.

¹Department of Mathematics, University College of Engineering Arni, Thatchur, Arni-632326. India. Email address: krajkj@yahoo.com

 $^2\mathrm{Department}$ of Mathematical and Computational Sciences, National Institute of Technology Karnataka, Surathkal, Karnataka - 575 025, India.

Email address: sam@nitk.edu.in

 $^3\mathrm{Department}$ of Mathematical and Computational Sciences, National Institute of Technology Karnataka, Surathkal, Karnataka - 575 025, India.

Email address: athirachandri@gmail.com