

NON-PARAMETRIC ESTIMATION OF CUMULATIVE (RESIDUAL) EXTROPY

Sudheesh K. Kattumannil^a and Sreedevi E. P.^b

^aIndian Statistical Institute, Chennai, India,

^bSNGS College, Pattambi, India.

ABSTRACT. Extropy and its properties are explored to quantify the uncertainty. In this paper, we obtain alternative expressions for cumulative residual extropy and negative cumulative extropy. We obtain simple estimators of cumulative (residual) extropy. Asymptotic properties of the proposed estimators are studied. We also present new estimators of cumulative (residual) extropy when the data is right censored. The finite sample performance of the estimators is evaluated through Monte Carlo simulation studies. We use the proposed estimators to analyse different real data sets. Finally, we obtain the relationship between different dynamic and weighted extropy measures and reliability concepts, which leads to several open problems associated with these measures.

Keywords: Entropy, Extropy; Right censoring; U-statistics.

1. Introduction

The uncertainty associated with a random variable can be evaluated using information measures. The widely used measure of information theory is entropy. For an absolutely continuous and non-negative random variable X , the differential entropy proposed by Shannon (1948) is given by

$$H(X) = -E(\log f(X)) = \int_0^\infty f(x) \log(f(x)) dx,$$

where $f(x)$ is the probability density function of x , provided above integral exists. Later, various measures of entropy are defined in the literature, each one suitable for specific situations. The widely used versions of entropy measures involved cumulative residual

[†]Corresponding E-mail: sreedeviep@gmail.com.

entropy (Rao et al., 2004), cumulative entropy (Di Crescenzo and Longobardi, 2009) and the corresponding weighed measures by Mirali et al. (2016) and Mirali and Baratpour (2017). Recently an alternative measure of uncertainty called extropy is proposed by Lad et al. (2015) as the complementary dual of entropy as following. For non-negative random variable X , extropy is defined as

$$J(X) = -\frac{1}{2} \int_0^{\infty} f^2(x) dx.$$

For more properties of $J(X)$ see Lad et al. (2015). Recently, several researchers studied various forms of extropy and proposed some new measures. Extropy of order statistics and record values is studied by Qiu (2017) and Jose and Sathar (2019) among others. Qiu and Jia (2018a) studied the residual extropy of order statistics. Qiu and Jia (2018b) developed goodness of fit of uniform distribution using extropy. Qiu et al. (2019) studied the properties of extropy for mixed systems. Jahanshahi et al. (2020) and Tahmasebi and Toomaj (2020) studied cumulative residual extropy and negative cumulative extropy in detail. Recently, Balakrishnan et al. (2020), Banal and Gupta (2021) and Sathar and Nair (2021a, 2021b, 2021c) studied different weighted versions of extropy.

The rest of the article is organised as follows. In Section 2, we provide an alternative expression for cumulative residual extropy and then obtained an estimator of the same. We also propose a new estimator of cumulative residual extropy when the data contain right censored observations. In Section 3, we obtain simple non-parametric estimators of the negative cumulative extropy for complete and right censored data. We prove that proposed estimators are consistent and asymptotically normally distributed. In section 4, we report the results of the Monte Carlo simulation study conducted to evaluate the finite sample performance of the proposed estimators. Analysis of several real data sets using the proposed estimators are given in Section 5. In Section 6, we obtain the relationship between different extropy measures and reliability concepts which leads to several open problems in this area.

2. Cumulative residual extropy

Let X be a non-negative continuous random variable with density and distribution function f and F , respectively. Define survival function of X at x as $\bar{F}(x) = 1 - F(x)$. Assume that X has finite mean μ . The cumulative residual extropy is given by (Jahanshahi et al., 2020)

$$\mathcal{CRE} = -\frac{1}{2} \int_0^\infty \bar{F}^2(x) dx. \quad (1)$$

Next we discuss the estimation of \mathcal{CRE} under complete and censored situations.

2.1. Uncensored case. Let X_1 and X_2 be two independent random variables having distribution function F . Then the survival function of $\min(X_1, X_2)$ is given by $\bar{F}^2(x)$. For a positive random variable X we have

$$E(X) = \int_0^\infty x f(x) dx = \int_0^\infty \bar{F}(x) dx.$$

Hence we can represent \mathcal{CRE} as

$$\mathcal{CRE} = -\frac{1}{2} E(\min(X_1, X_2)). \quad (2)$$

An estimator of \mathcal{CRE} based on U-statistics is given by

$$T_1 = \frac{-1}{n(n-1)} \sum_{i=1}^{n-1} \sum_{j=i+1}^n \min(X_i, X_j).$$

Clearly T_1 is an unbiased estimator of \mathcal{CRE} . Using order statistics we can express above estimator as

$$T_1 = \frac{-1}{n(n-1)} \sum_{i=1}^n (n-i) X_{(i)}, \quad (3)$$

where $X_{(i)}$ denote the i -th order statistics based on a random sample X_1, \dots, X_n from F .

Next we study the asymptotic properties of T_1 using the limit theorems of U-statistics.

Since T_1 is a U statistic, it is consistent estimator of \mathcal{CRE} (Lehmann, 1951).

Theorem 1. *As $n \rightarrow \infty$, $\sqrt{n}(T_1 - \mathcal{CRE})$ converges in distribution to Gaussian random variable with mean zero and variance $4\sigma_1^2$ where*

$$\sigma_1^2 = V\left(X\bar{F}(X) + \int_0^X y dF(y)\right). \quad (4)$$

Proof: By central limit theorem of U-statistics, we have the asymptotic normality. The asymptotic variance is $4\sigma_1^2$, where σ_1^2 is given by (Lee, 2019)

$$\sigma_1^2 = \text{Var} (E(\min(X_1, X_2)|X_1)) .$$

Consider

$$\begin{aligned} E(\min(X_1, X_2)|X_1 = x) &= E(xI(x < X_2) + X_2I(X_2 \leq x)) \\ &= x\bar{F}(x) + \int_0^x ydF(y). \end{aligned}$$

Hence we have the variance expression specified as in the theorem.

2.2. Right censored case. In this section, we obtain a simple estimator of \mathcal{CRE} for right censored data. Suppose we have randomly right-censored observations where the censoring times are independent of the lifetimes. Let C be the censoring random variable with survival function K . We are interested to find an estimator of the cumulative residual extropy based on n independent and identical observations $\{(Y_i, \delta_i), 1 \leq i \leq n\}$, where $Y_i = \min(X_i, C_i)$ and $\delta_i = I(X_i \leq C_i)$, is the censoring indicator. A U-statistic defined for censored data is given by (Datta et al., 2010)

$$T_{1c} = -\frac{1}{n(n-1)} \sum_{i=1}^n \sum_{j < i; j=1}^n \frac{\min(Y_i, Y_j)\delta_i\delta_j}{\widehat{K}(Y_i-)\widehat{K}(Y_j-)}, \quad (5)$$

where $\widehat{K}(\cdot)$ is the Kaplan-Meier estimator of $K(\cdot)$. Now we study the asymptotic properties of T_{1c} . In the next theorem, we state the consistency of T_{1c} . The following results can be proved in similar lines as Theorem 2 of Sudheesh et al. (2021).

Theorem 2. *As $n \rightarrow \infty$, T_{1c} converges in probability to \mathcal{CRE} .*

To derive the asymptotic distribution of T_{1c} , define $N_i^c(t) = I(Y_i \leq t, \delta_i = 0)$ as the counting process corresponding to censoring for the i -th subject and $R_i(u) = I(Y_i \geq u)$. Let $\lambda_c(t)$ be the hazard rate of the censoring variable C . The martingale associated

with the counting process $N_i^c(t)$ is given by

$$M_i^c(t) = N_i^c(t) - \int_0^t R_i(u) \lambda_c(u) du.$$

Let $H_c(x) = P(Y_1 \leq x, \delta = 1)$, $y(t) = P(Y_1 > t)$ and

$$w(t) = \frac{1}{y(t)} \int \frac{h_1(x)}{\widehat{K}(x-)} I(x > t) dH_c(x),$$

where $h_1(y) = E(h(Y_1, Y_2) | Y_1 = y)$. The proof of the next theorem follows from Datta et al. (2010) with choice of the kernel $h(Y_1, Y_2) = \min(Y_1, Y_2)$.

Theorem 3. Assume $E(\min^2(Y_1, Y_2)) < \infty$, $\int \frac{h_1(x)}{\widehat{K}^2(x-)} dH_c(x) < \infty$ and $\int_0^\infty w^2(t) \lambda_c(t) dt < \infty$. As $n \rightarrow \infty$, the distribution of $\sqrt{n}(T_{1c} - \mathcal{CRE})$ is Gaussian with mean zero and variance $4\sigma_{1c}^2$, where σ_{1c}^2 is given by

$$\sigma_{1c}^2 = \text{Var}\left(\frac{h_1(X)\delta_1}{\widehat{K}(Y-)} + \int w(t) dM_1^c(t)\right). \quad (6)$$

3. Cumulative extropy

In this section, we discuss the estimation of negative cumulative extropy in uncensored and censored situations. For a non-negative random variable X , negative cumulative extropy is defined as (Tahmasebi and Toomaj, 2020)

$$\mathcal{CE} = \frac{1}{2} \int_0^\infty (1 - F^2(x)) dx. \quad (7)$$

3.1. Uncensored case. Let X_1 and X_2 be two independent random variables with same distribution function F . Then the distribution function of $\max(X_1, X_2)$ is $F^2(x)$. Hence we can represents \mathcal{CE} as

$$\mathcal{CE} = \frac{1}{2} E(\max(X_1, X_2)). \quad (8)$$

A U-statistic given by

$$T_2 = \frac{1}{n(n-1)} \sum_{i=1}^{n-1} \sum_{j=i+1}^n \max(X_i, X_j),$$

is an unbiased and consistent estimator of \mathcal{CE} . The above estimator can be written as

$$T_2 = \frac{1}{n(n-1)} \sum_{i=1}^n (i-1)X_{(i)}.$$

Next we obtain the asymptotic distribution of T_2 and the proof the following result is similar to Theorem 1.

Theorem 4. *As $n \rightarrow \infty$, $\sqrt{n}(T_2 - \mathcal{CE})$ converges in distribution to Gaussian random variable with mean zero and variance $4\sigma_2^2$ where*

$$\sigma_2^2 = V\left(XF(X) + \int_X^\infty ydF(y)\right). \quad (9)$$

3.2. Right censored case. In this subsection, we find a simple estimator of \mathcal{CE} in the presence of right censored observations. Using the same notations used in Section 2.2, a U-statistic for \mathcal{CE} is given by

$$T_{2c} = \frac{1}{n(n-1)} \sum_{i=1}^n \sum_{j < i; j=1}^n \frac{\max(Y_i, Y_j)\delta_i\delta_j}{\widehat{K}(Y_{i-})\widehat{K}(Y_{j-})}. \quad (10)$$

Next results establish the consistency and asymptotic normality of T_{2c} .

Theorem 5. *As $n \rightarrow \infty$, T_{2c} converges in probability to \mathcal{CE} .*

Theorem 6. *Let $h_1(y) = E(\max(Y_1, Y_2)|Y_1 = y)$. Assume $E(\max^2(Y_1, Y_2)) < \infty$, $\int \frac{h_1(x)}{\widehat{K}^2(x-)} dH_c(x) < \infty$ and $\int_0^\infty w^2(t)\lambda_c(t)dt < \infty$. As $n \rightarrow \infty$, the distribution of $\sqrt{n}(T_{2c} - \mathcal{CE})$ is Gaussian with mean zero and variance $4\sigma_{2c}^2$, where σ_{2c}^2 is given by*

$$\sigma_{2c}^2 = Var\left(\frac{h_1(X)\delta_1}{\widehat{K}(Y-)} + \int w(t)dM_1^c(t)\right). \quad (11)$$

4. Empirical evidence

To evaluate the finite sample performance of the proposed estimators we conduct a Monte Carlo simulation studies using R software. The simulation is repeated ten thousand times.

4.1. Cumulative Extropy.

4.1.1. *Complete data.* We generate observations from exponential, gamma, Weibull and log normal distributions. We find the bias and MSE of the proposed estimator based on samples with size $n = 10, 20, 30, 40$ and 50 . We also compare the estimator with the estimator of \mathcal{CRE} proposed by Tahmasebi and Toomaj (2020) (TT_1). The results are given in Table 1.

TABLE 1. Bias and MSE of estimator of \mathcal{CRE}

n	Exponential (1)				Gamma (2,1)			
	T_1		TT_1		T_1		TT_1	
	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
10	0.00076	0.00879	0.02541	0.00776	0.00199	0.02647	0.19588	0.05776
20	0.00074	0.00416	0.01283	0.00398	0.00168	0.01270	0.13943	0.03153
30	0.00038	0.00274	0.00861	0.00263	0.00124	0.00851	0.11270	0.02151
40	0.00018	0.00211	0.01283	0.00202	0.00094	0.00638	0.09819	0.01632
50	0.00004	0.00164	0.02541	0.00160	0.00036	0.00507	0.08221	0.01348
n	Weibull (2,1)				Lognormal (0,1)			
	T_1		TT_1		T_1		TT_1	
	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
10	0.00078	0.00431	0.12641	0.01951	0.00115	0.01765	0.08175	0.02327
20	0.00058	0.00217	0.09280	0.01105	0.00098	0.00844	0.06710	0.01264
30	0.00054	0.00142	0.07621	0.00764	0.00049	0.00546	0.05867	0.00892
40	0.00017	0.00104	0.06703	0.00600	0.00030	0.00410	0.05367	0.00705
50	0.00001	0.00087	0.05987	0.00484	0.00024	0.00325	0.04867	0.00572

From Table 1, we can see that the bias and MSE of the proposed estimator is very low compared to that proposed by Tahmasebi and Toomaj (2020). Also bias and MSE decrease as sample size increases. This ensures the efficiency of the new estimator.

4.1.2. *Censored data.* For generating lifetimes we consider the same distributions used in uncensored case. In all the cases, the censored observations are generated from exponential distribution with parameter λ where $P(T > C) = 0.2$. The bias and MSE of the estimates when 20% of the lifetimes are censored are given in Table 2. From Table 2, we observe that bias and MSE of the proposed estimator for censored observations are small and decreases with sample size.

TABLE 2. Bias and MSE of estimator of \mathcal{CRE} when 20% of lifetimes are censored

n	Exponential (1)		Gamma(2,1)		Weibull(2,1)		Lognormal(0,1)	
	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
50	0.08418	0.00830	0.17328	0.04083	0.08358	0.00914	0.26264	0.07059
75	0.08287	0.00788	0.17321	0.03822	0.08128	0.00852	0.26235	0.06979
100	0.08222	0.00761	0.17318	0.03735	0.07956	0.00807	0.26228	0.06956
200	0.08199	0.00741	0.17309	0.03585	0.07942	0.00762	0.26189	0.06895

4.2. Cumulative Extropy. The same simulation set up is employed to evaluate the efficiency of the estimator of negative cumulative extropy. Here also we compare the estimator with the estimator of \mathcal{CE} proposed by Tahmasebi and Toomaj (2020) (TT_2). The bias and MSE of the estimators are given in Table 3. In this case also the proposed estimator has less bias and MSE compared to the estimator of \mathcal{CE} proposed by Tahmasebi and Toomaj (2020).

TABLE 3. Bias and MSE of estimator of \mathcal{CE}

n	Exponential (1)				Gamma (2,1)			
	T_2		TT_2		T_2		TT_2	
	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
10	0.00118	0.01473	0.03877	0.01455	0.00074	0.02564	0.13489	0.04161
20	0.00098	0.00722	0.01938	0.00719	0.00052	0.01243	0.08878	0.02026
30	0.00050	0.00491	0.00861	0.00488	0.00036	0.00827	0.06992	0.01344
40	0.00038	0.00356	0.01283	0.00357	0.00032	0.00649	0.05864	0.01025
50	0.00037	0.00297	0.02541	0.00295	0.00028	0.00493	0.05128	0.00783
n	Weibull (2,1)				Lognormal (0,1)			
	T_2		TT_2		T_2		TT_2	
	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
10	0.00098	0.00845	0.15349	0.03316	0.00764	0.35391	0.16508	0.34391
20	0.00092	0.00419	0.10558	0.01656	0.00430	0.18710	0.10203	0.18699
30	0.00031	0.00281	0.08592	0.01120	0.00385	0.12033	0.08764	0.12357
40	0.00013	0.00210	0.07303	0.00828	0.00064	0.08892	0.07814	0.09248
50	0.00001	0.00165	0.06441	0.00655	0.00044	0.07305	0.06620	0.07591

4.2.1. Censored data. The same parameter setup as in Section 4.1.2 is used in the simulation. The bias and MSE of the proposed estimator of \mathcal{CE} when 20% of lifetimes are censored are given in Table 4. From Table 4, we observe that the bias and MSE are little higher in case of exponential and gamma distributions. This may be due to the

fact that exponential and gamma distributions are positively skewed and the estimator is based on maxima.

TABLE 4. Bias and MSE of estimator of \mathcal{CE} when 20% of lifetimes are censored

	Exponential (1)		Gamma (2,2)		Weibull (2,1)		Lognormal (0,1)	
n	Bias	MSE	Bias	MSE	Bias	MSE	Bias	MSE
50	0.19183	0.04122	0.28382	0.09566	0.11409	0.02219	0.00712	0.04865
75	0.19056	0.03923	0.28137	0.08928	0.10925	0.01876	0.00639	0.03171
100	0.19048	0.03834	0.27948	0.08552	0.10690	0.01792	0.00447	0.02334
200	0.18831	0.03646	0.27659	0.08010	0.10540	0.01546	0.00227	0.01142

5. Data analysis

We apply the proposed estimation procedures to several real life data sets. The U-statistic based estimators for cumulative residual extropy and negative cumulative extropy are estimated in uncensored and censored cases.

5.1. Complete data. Three real life data sets are used for illustration.

Example 1: First, we consider the failure times of 84 mechanical components which is reported in Example 4.1 Tahmasebi and Toomaj (2020). We obtain $T_1 = -0.9636$ and the corresponding estimator of Tahmasebi and Toomaj (2020) as $TT_1 = -0.9494$. Cumulative extropy of this data is estimated as $T_2 = 1.9509$ and the estimate of the same by Tahmasebi and Toomaj (2020) as $TT_2 = 1.9251$.

Example 2: Next, we consider the data set on active repair time (in hours) for an airborne communication transceiver reported by Balakrishnan et al. (2009) and studied by Jahanshahi et al. (2020). The complete data set of 46 repair times is given in Jahanshahi et al. (2020) in Example 8. For this data estimator of cumulative residual extropy is calculated as $T_1 = -0.7454$ ($TT_1 = -0.6684$) and the estimators of negative cumulative extropy is obtained as $T_2 = 2.8610$ ($TT_2 = 2.7380$).

Example 3: Now we consider the data on the number of million revolutions before

failure for each of 23 ball bearings studied in Lawless (2011). The data is given in Example 3.3.1 of Lawless (2011). For this data, we find $T_1 = -25.6843$ ($TT_1 = -17.1976$) and $T_2 = 46.5365$ ($TT_2 = 37.1432$).

5.2. Censored data. Here also, we consider three real life data sets for demonstration.

Example 1: We consider the data on survival times (in months) of patients with Hodgkin's disease given in Lawless (2011) (Example 3.2, Page 139) for illustration. The data consists of 35 lifetimes out of which 9 are censored. Hence censoring percentage is around 25.7. Now we estimate cumulative residual extropy as $T_{1c} = -3.4154$ and negative cumulative extropy as $T_{2c} = 12.0849$ using the proposed procedures.

Example 2: We examine the data on lifetimes of disk break pads on 40 cars studied in Lawless (2011) (see Table 6.11, Page 337). Out of the 40 observed lifetimes 22.5% are censored observations. In this case, cumulative residual extropy is estimated as $T_{1c} = -17.3063$ and negative cumulative extropy as $T_{2c} = 25.3462$.

Example 3: We also analyse stanford heart transplant data available in R package 'stanford2'. The data consist of 184 lifetimes where 72 (38.5%) of them are censored. For this data, we obtain $T_{1c} = -711.5523$ and $T_{2c} = 2028.036$.

6. Further Discussions

We gave alternative expressions for cumulative residual extropy and negative cumulative extropy. Using the alternative expressions we obtained estimators of these measures for complete and censored cases. We studied the asymptotic properties of the proposed estimators. Numerical illustrations are given through Monte Carlo simulation study and real data analysis.

Next, we give simple alternative expressions for different extropy measures. These alternative expression enables us to connect extropy and reliability measures. Using these alternative expressions one can study the inference and other problems associated with these quantities.

Sathar and Nair (2021b) defined dynamic survival extropy as

$$J_t(X) = -\frac{1}{2\bar{F}^2(t)} \int_t^\infty \bar{F}^2(x) dx.$$

For various properties of $J_t(X)$ interest readers may refer Jahanshahi et. al. (2020).

The mean residual life function of X (denoted by $m(t)$) can be express as

$$m(t) = E(X - t | X > t) = \frac{1}{\bar{F}(t)} \int_t^\infty \bar{F}(x) dx.$$

Hence using the survival function of $\min(X_1, X_2)$, we can express $J_t(X)$ as

$$J_t(X) = -\frac{1}{2} E(\min(X_1, X_2) - t | \min(X_1, X_2) > t).$$

From the above expression, we can easily observe that $-2J_t(X)$ is the mean residual life function of a series system having two identical components.

Kundu (2020) defined dynamic cumulative extropy given by

$$H_t(X) = -\frac{1}{2F^2(t)} \int_0^t F^2(x) dx.$$

The mean past life function of X (denoted by $r(t)$) is given by

$$r(t) = E(t - X | X \leq t).$$

Using some algebraic manipulations we can rewrite $r(t)$ as

$$r(t) = \frac{1}{F(t)} \int_0^t F(x) dx.$$

Hence using the distribution function of function of $\max(X_1, X_2)$, we can express $H_t(X)$ as

$$H_t(X) = -\frac{1}{2} E(t - \max(X_1, X_2) | \max(X_1, X_2) \leq t).$$

Hence $-2H_t(X)$ is the mean past life function of a parallel system having two identical components.

Weighted version of the survival extropy is given by (see Sathar and Nair, 2021c)

$$J(X, w) = -\frac{1}{2} \int_0^\infty x \bar{F}^2(x) dx.$$

With simple algebraic manipulation we can express it as

$$J(X, w) = -\frac{1}{4} E(\min(X_1, X_2)^2).$$

The weighted version of the cumulative extropy is given by

$$H(X, w) = -\frac{1}{2} \int_0^\infty x(1 - F^2(x)) dx.$$

Again with simple algebraic manipulation we obtain

$$H(X, w) = -\frac{1}{4} E(\max(X_1, X_2)^2).$$

Sathar and Nair (2021c) also defined weighted dynamic survival extropy as

$$J_t(X, w) = -\frac{1}{2\bar{F}^2(t)} \int_t^\infty x \bar{F}^2(x) dx.$$

Consider

$$\begin{aligned} E(\min(X_1, X_2) - t^2 | \min(X_1, X_2) > t) &= \frac{1}{\bar{F}^2(t)} \int_t^\infty (y^2 - t^2) 2\bar{F}(y) dF(y) \\ &= \frac{1}{\bar{F}^2(t)} \int_t^\infty \left(\int_t^y 2x dx \right) 2\bar{F}(y) dF(y) \\ &= \frac{1}{\bar{F}^2(t)} \int_t^\infty \left(\int_x^\infty 2\bar{F}(y) dF(y) \right) 2x dx \\ &= \frac{2}{\bar{F}^2(t)} \int_t^\infty x \bar{F}^2(x) dx. \end{aligned}$$

Hence

$$4J_t(X, w) = -E(\min(X_1, X_2) - t^2 | \min(X_1, X_2) > t).$$

They also defined weighted dynamic cumulative extropy as

$$H_t(X, w) = -\frac{1}{2F^2(x)} \int_0^t x F^2(x) dx.$$

Similar to above we obtain

$$4H_t(X, w) = -E(t^2 - \max(X_1, X_2) | \max(X_1, X_2) \leq t).$$

The alternative representations given above can be exploited for studying more about different extropy measures.

References

- [1] Bansal, S. and Gupta, N. (2020). Weighted extropies and past extropy of order statistics and k-record values. *Communications in Statistics-Theory and Methods*, 1–24.
- [2] Balakrishnan, N., Leiva, V., Sanhueza, A. and Cabrera, E. (2009). Mixture inverse Gaussian distributions and its transformations, moments and applications. *Statistics*, 43, 91–104.
- [3] Balakrishnan, N., Buono, F. and Longobardi, M. (2020). On weighted extropies. *Communications in Statistics-Theory and Methods*, 1–31.
- [4] Datta, S., Bandyopadhyay, D. and Satten, G. A. (2010). Inverse probability of censoring weighted U-statistics for right-censored data with an application to testing hypotheses. *Scandinavian Journal of Statistics*, 37, 680–700.
- [5] Blischke, W. R., and Murthy, D. P. (2011). *Reliability: Modeling, Prediction and Optimization*. John Wiley and Sons, New York.
- [6] Di Crescenzo, A. and Longobardi, M. (2009). On cumulative entropies. *Journal of Statistical Planning and Inference*, 139, 4072–4087.
- [7] Jose, J. and Sathar, E. A. (2019). Residual extropy of k-record values. *Statistics and Probability Letters*, 146, 1–6.
- [8] Jahanshahi, S. M. A., Zarei, H. and Khammar, A. H. (2020). On cumulative residual extropy. *Probability in the Engineering and Informational Sciences*, 34, 605–625.
- [9] Kundu, C. (2020). On Cumulative residual (past) extropy of extreme order statistics. *arXiv preprint*, arXiv:2004.12787.
- [10] Lad, F., Sanfilippo, G. and Agro, G. (2015). Extropy: complementary dual of entropy. *Statistical Science*, 30, 40–58.
- [11] Lawless, J. F. (2011). *Statistical Models and Methods for Lifetime Data*, John Wiley & Sons, New Jersey.
- [12] Lee, A. J. (2019). *U-statistics: Theory and Practice*, CRC press, Boca Raton.

- [13] Lehmann, E. L. (1951). Consistency and unbiasedness of certain non-parametric tests. *Annals of Mathematical Statistics*, 22, 165–179.
- [14] Mirali, M., Baratpour, S. and Fakoor, V. (2016). On weighted cumulative residual entropy. *Communications in Statistics- Theory and Methods* 46, 2857–2869.
- [15] Mirali, M. and Baratpour, S. (2017). Some results on weighted cumulative entropy. *Journal of The Iran Statistical Society*, 16, 21–32.
- [16] Rao, M., Chen, Y., Vemuri, B. and Wang, F. (2004). Cumulative residual entropy: A new measure of information. *IEEE transactions on Information Theory*, 50, 1220–1228.
- [17] Sathar, E. A. and Nair, R. D. (2021a). On dynamic weighted extropy. *Journal of Computational and Applied Mathematics*, 393, 113507.
- [18] Sathar, E. A. and Nair R, D. (2021b). On dynamic survival extropy. *Communications in Statistics-Theory and Methods*, 50, 1295–1313.
- [19] Sathar, E. A. and Nair, R. D. (2021c). A study on weighted dynamic survival and failure entropies. *Communications in Statistics-Theory and Methods*, 1–20.
- [20] Shannon, C. E. (1948). A mathematical theory of communication. *The Bell system technical journal*, 27, 379–423.
- [21] Sudheesh, K. K., Dewan, I. and Sreelaksmi, N. (2021). Non-parametric estimation of Gini index with right censored observations. *Statistics and Probability Letters*, 175, 109113.
- [22] Qiu, G. (2017). The extropy of order statistics and record values. *Statistics and Probability Letters*, 120, 52–60.
- [23] Qiu, G. and Jia, K. (2018a). The residual extropy of order statistics. *Statistics and Probability Letters*, 133, 15–22.
- [24] Qiu, G. and Jia, K. (2018b). Extropy estimators with applications in testing uniformity. *Journal of Nonparametric Statistics*, 30, 182–196.
- [25] Qiu, G., Wang, L. and Wang, X. (2019). On extropy properties of mixed systems. *Probability in the Engineering and Informational Sciences*, 33, 471–486.
- [26] Tahmasebi, S. and Toomaj, A. (2020). On negative cumulative extropy with applications. *Communications in Statistics-Theory and Methods*, 1–23.