

# The Continuity Postulate in Economic Theory: A Deconstruction and an Integration\*

Metin Uyanik<sup>†</sup> (r) M. Ali Khan<sup>‡</sup>

August 26, 2021

**Abstract:** This paper presents six theorems and ten propositions that can be read as deconstructing and integrating the continuity postulate under the rubric of pioneering work of Eilenberg, Wold, von Neumann-Morgenstern, Herstein-Milnor and Debreu. Its point of departure is the fact that the adjective *continuous* applied to a function or a binary relation does not acknowledge the many meanings that can be given to the concept it names, and that under a variety of technical mathematical structures, its many meanings can be whittled down to novel and unexpected equivalences that have been missed in the theory of choice. Specifically, it provides a systematic investigation of the two-way relation between *restricted* and *full* continuity of a function and a binary relation that, under convex, monotonic and differentiable structures, draws out the behavioral implications of the postulate. (135 words)

*Journal of Economic Literature* Classification Numbers: C00, C02, D01, D11, D21, D51, D81

*2020 Mathematics Subject* Classification Numbers: 26B05, 91B02, 91B08, 91B42, 06A06

**Key Words:** Wold-continuous, weakly Wold-continuous, mixture-continuous, scalar-continuous, linear-continuous, arc-continuous, graph-continuous, section-continuous, separately-continuous, jointly-continuous, restriction-continuous, Archimedean

*Running Title:* The Continuity Postulate

---

\*This work is presented as a celebration of Tajliing Koopmans' 110<sup>th</sup>-birthday: its authors constitute the second and third generation of his student-followers, the first being represented by Beckman, Christ and Nerlove, the editors of the first volume of his collected papers. The authors thank Max Amarante, Elena Antoniadou, Aurelien Baillon, Juan Dubra, Hülya Eraslan, Yorgos Gerasimou, Aniruddha Ghosh, Alfio Giarlotta, Carlos Hervés-Beloso, Jeff Kline, David Levy, Paulo Monteiro, Efe Ok, Eddie Schlee and Eric Schliesser for encouragement and stimulating conversation and correspondence. The authors also thank seminar participants at the School of Economics at UTS and ANU, and conference participants at the 38th AETW held at the University of Adelaide, Australia. Some of the results reported here were presented by the second author at *Positivity X*, a conference held at the University of Pretoria July 8-12, 2019, and also announced without proof at the NSF/NBER/CEME *Conference on Mathematical Economics: In Honor of Robert M. Anderson* in Berkeley, October 25, 2019. Khan should like to thank the respective organizers for their hospitality.

<sup>†</sup>School of Economics, University of Queensland, Brisbane, QLD 4072. **E-mail** m.uyanik@uq.edu.au. We use the *certified random order* in order to list the authors; see Ray-Robson (2018, American Economic Review).

<sup>‡</sup>Department of Economics, Johns Hopkins University, Baltimore, MD 21218. **E-mail** akhan@jhu.edu

# Contents

<b>Contents</b>	<b>1</b>
<b>1 Introduction</b>	<b>1</b>
<b>2 Mathematical Antecedents</b>	<b>5</b>
2.1 Eilenberg's Paper: A Conspicuous Omission . . . . .	5
2.2 Rosenthal's Theorem: Suggested Directions . . . . .	7
<b>3 The Theory</b>	<b>8</b>
3.1 On the Continuity of Functions . . . . .	8
3.2 On the Continuity of Relations . . . . .	10
<b>4 The Theory Applied</b>	<b>17</b>
4.1 Representation of Preferences: Benchmark Results . . . . .	17
4.1.1 An Euclidean Space: Wold (1943–44) . . . . .	18
4.1.2 A Mixture Space: Herstein and Milnor (1953) . . . . .	20
4.1.3 A Finite State Space: Gul (1992) . . . . .	20
4.1.4 A Topological Vector Space: Neuefeind-Trockel (1995) . . . . .	21
4.2 Neoclassical Consumer Theory . . . . .	22
4.2.1 Debreu's (1959) Remark . . . . .	22
4.2.2 Schmeidler's (1969) Non-decisive Consumer . . . . .	23
4.2.3 Shafer's (1974) Non-transitive Consumer . . . . .	24
4.2.4 Hidden Assumptions in Walrasian Equilibrium Theory . . . . .	25
4.3 Consistency and Decisiveness: A Reconsideration . . . . .	25
4.4 Other Potential Applications . . . . .	26
<b>5 Concluding Remarks</b>	<b>28</b>
<b>A Appendix</b>	<b>30</b>
A.1 Proofs of the Results . . . . .	30
A.2 Seven Technical Examples . . . . .	50
<b>B Bibliography</b>	<b>54</b>

---

*The postulational structure of the mathematical tool parallels that of the substantive theory to be constructed, and the two are studied and apprehended simultaneously. The welcome result is that “mathematical” and “literary” economics are moving closer. They meet on the ground of a common requirement of good hard thought from explicit basic postulates, rather than for manipulative skills. If there is a difference, it is one of succinctness of expression rather than of content, concepts or objective.*<sup>2</sup> Tjalling Koopmans (1957)

## 1 Introduction

Tjalling Koopmans mentions the continuity postulate early on in the first of his 1957 magisterial essays “on the state of economic science.” After a footnote reference to a *continuous* function, an adjective he does not define but freely use, and then to a *continuously representable* preference ordering that he does define, he masks the continuity postulate under the assumption of *local non-satiation*, referring to the latter as “a rather weak continuity property of preferences.” Subsequently, and in the interest of precision, he sees the completeness postulate on preferences as “suggested by considerations of continuity and nonsaturation.”<sup>3</sup> It is only in the second half of the essay, well after he has developed the basic theorems that the essay is to present, that he uses the adjective *continuous* for a correspondence, but here again does not define the “appropriate generalization of the concept of continuity.”<sup>4</sup> The point is that the notion of continuity is one

---

<sup>1</sup>See Gottfried Wilhelm Leibniz *Nouveaux essais sur l’entendement humain*, 1704, p. 50, and his 1702 essay *On the Principles of Continuity* in Weiner (1951). Also see Alexander Baumgarten *Metaphysics: A Critical Translation with Kant’s Elucidations*, translated and edited by Courtney D. Fugate and John Hymers, Bloomsbury, 2013, “Preface of the Third Edition (1750)”, p. 79, n. d: “[Baumgarten] must also have in mind Leibniz’s “natura non facit saltus [nature does not make leaps]” (NE IV, 16).” It is worth stating that sentence is also an epigraph of Marshall’s *Principles*. We engage this epigraph in the concluding section of this essay by emphasizing the importance of *perception* to *theorization* as the methodological backbone of this essay. This is in response to a referee’s comment that “the paper should convey some awareness of the approximations in use, and that it intends continuity to convey something at the perceptible level.” We take this opportunity to thank the referee for his/her generous and erudite comments.

<sup>2</sup>See page 177 in Koopmans (1957); we skip some phrases. Koopmans offers as an illustration: “[I]n some intuitive sense the “distance” between Lerner’s *The Economics of Control* and the mathematical formulations of the propositions of welfare economics .. is, I believe, not large.” We return to Koopmans in Footnote 63, and the text it footnotes.

<sup>3</sup>See Footnote 5 on page 11 for the first reference; page 19 for the second; and Footnote 1 on page 20 for the third, all in Koopmans (1957). For his free use of the notion of a continuous function and a continuously representable binary relation, see respectively Footnote 2 on page 33, Footnote 3 on page 34, and page 55. It is of interest that the problem of representation of a binary relation by a function is introduced so early on the essay, right after the notion of profit maximization, and in the context of supply and consumption decisions. More to the point, Koopmans disregards the chronological order to introduce Wold before the 1934 paper of Hicks-Allen.

<sup>4</sup>See page 57 in Koopmans (1957) for the discussion of a “class of theorems known as fixed point theorems, the appropriate concepts and theorems [for which] are found in the relatively young mathematical field of *topology* –

of the signature notions in the *Essays*,<sup>5</sup> and his reliance on it continues on with more or less increasing force and emphasis throughout his work subsequent to them.<sup>6</sup>

We shall have more to say about Koopmans' fraught relationship to the continuity postulate: its exploitation of the interplay between its bearing on a function as opposed to that on a binary relation and/or a correspondence is an interplay that runs throughout his later *oeuvre*. However, this is not a paper on Koopmans' place in the history of economic thought, and we begin with him only because of the impact that the 1957 *Essays*, and his subsequent papers, have had on the direction of the literature: it is this direction and its consequence for ongoing research that is of primary concern of this paper's attempted examination of the postulate. In this connection, it is worth pointing out that in his 1960 and 1964 work, Koopmans assumes preferences to be parametrized by a continuous function, and investigates additional axioms under which such a function takes on a specific form. Later in 1972, in his two papers in honour of Jacob Marschak, he considers the representation of a binary (preference) relation by a (utility) function, the so-called representation problem, and of necessity, introduces continuous relations. Since the work reported here draws on the insights of the antecedent mathematical literature on the continuity of functions, specifically on the application of Rosenthal's 1955 theorem to continuous relations (see below), both aspects of his work are relevant for us.<sup>7</sup>

Remaining with the *Essays*, but now shifting from the first to the second, Koopmans returns in his masterful discussion of Herstein-Milnor (1953) and Savage (1954) to the continuity postulate in the context of a "single decision maker faced with uncertainty." In this discussion, he refers to the "surprisingly far-reaching implications of simple postulates of consistency and sharpness of preferences concerning objects capable of a particular type of continuous variation," but again masks the continuity property of preferences, herein *scalar continuity*, in a postulate that exerts the existence of a probability mixture equivalent to the one intermediate between three ranked objects of choice, referred to as *prospects*.<sup>8</sup> In his discussion of the representability problem, he writes:

---

[which] crudely put, [is] the study of ways in which a set can "hang together." Arrow-Hahn (1971) will refer to "deeper methods," see Footnote 10 below.

<sup>5</sup>A tracking of the word and its derivatives yields 84 instances, ranging its technical use in economic and statistical theory to its colloquial use in ordinary discourse, as for example, in the importance of the continuity of research support for the investigation of mathematical tools.

<sup>6</sup>In his 1957 lectures at Stanford Summer School, (Koopmans-Bausch in Koopmans (1970), the notion is precisely defined in the context of a function, correspondence and a binary relation. In the *Essays* more generally, there are 21 uses of the word and its derivatives, but in his 1964 paper "On flexibility of future preference," originally dating to a 1950 abstract in which he tackles "utility analysis of consumer choice," any reference to continuity is conspicuously missing.

<sup>7</sup>For Koopmans' work referred to here, see Footnote 66 below. The reference to Rosenthal's theorem follows in Section 2 below.

<sup>8</sup>See Section 10 on pages 155 to 161 in Koopmans (1957). The quote is taken from the last paragraph on page 159. Also see Footnote 2 on page 155 for the relevant literature concerning the Herstein-Milnor paper.

Herman Wold and Gerard Debreu have realized that the [assertion that] every complete ordering can be represented by an ordinal utility function needs proof, and have successively and with increasing generality formulated conditions, essentially of continuity, under which they prove that a complete ordering permits expression by a continuous utility function.<sup>9</sup>

However, in this entire discussion there is no reference to Eilenberg (1941), a pioneering paper only acknowledged as a precursor of Debreu fifteen years later in Koopmans (1972).<sup>10</sup> More generally, it is this omission of Eilenberg’s paper, as well as the masking of the continuity postulate, both in the theory of competitive equilibrium and in uncertainty theory as presented by von Neumann-Morgenstern, that serves as the point of departure for, and motivational center of, the results to follow. There is no bridging between the two subjects, and they are confined to separate essays, a separation and a consequent obscuring that is seen here as a consequence of not beginning with the primitive raw notion of continuity. In short, this essay is an angled contribution to the theory, its two arms are represented by the twin *omission* and *masking*, each hitched to Koopmans’ *Essays*.

After a section on mathematical underpinnings, we discuss the omission in Section 3 below and turn to what we mean by the term “masking.” This is really brought out in Koopmans’ presentation of continuity as a non-satiation or an existence property. The point is that there is no single and unambiguous rendering of the continuity postulate: different notions of continuity have been used in economic and decision theory depending on the different needs and requirements of a particular result, as indeed they ought. There is no standard continuity assumption even when differences in topologies are disregarded. In terms of a further unpacking, one can discern three types of continuity pertaining to a binary relation and labelled as *graph* continuity, *section* continuity, and *scalar* continuity. All impose topological assumptions on the relation itself – graph continuity on it as a subset of the product space and section continuity on its sections – but while the first two use the topological structure of the space on which the choice function or relation may be defined, the third, in its reference to the *mixture-continuity*, and *Archimedean* properties, works with an algebraic structure on the choice set, and limits itself only to the topological structure of the unit interval.<sup>11</sup> All properties are widely used in economic theory, but perhaps the last is more pervasive in decision theory, while the first two

---

<sup>9</sup>See p. 19 in Koopmans (1957). In Footnote 2 on the same page the author cites the 1943 paper of Wold and 1954 paper of Debreu. The readers should note that the authors have rearranged this sentence hopefully without any loss or gain in meaning. Koopmans’ emphasis on Wold and Debreu is repeated in Arrow-Hahn (1971).

<sup>10</sup>In his reproduction of Debreu’s theorem [Proposition 1] on the sufficiency of connectedness of a choice set in a finite-dimensional Euclidean space, Koopmans corrects his oversight by observing that “Debreu credits a paper of Eilenberg (1941) as containing the mathematical essence of [his] Proposition 1.” Arrow-Hahn (1971) cite Eilenberg, but in a dismissive way: “A very considerable generalization, based on a mathematical paper by Eilenberg (1941), was achieved with the deeper methods by Debreu (1954).” We return to this in Section 2.

<sup>11</sup>As we show in the sequel, *Wold-continuity* and its various versions also need to be added to the mix, and they overlap with questions of *solvability*; see Figure 3 below.

constitute more of a staple of Walrasian general equilibrium and game-theoretic analyses. The point that constantly guides this essay is that one has to go back to the most raw, primitive and undefined, meaning of the term if the analytical and substantive connections between the subsequent conceptual proliferations are not to be missed.

With this preamble to the antecedent literature on “decision theory,” we can turn to another dissonance and exclusion, that in the now-standard expositions of neoclassical theory of the consumer. In seminal work on the theory of individual choice, Wold (1943–44) and Wold-Jureen (1953), also touch on the problem of the representation of a preference relation by a real-valued (utility) function in the context of a finite-dimensional Euclidean space. However, unlike Eilenberg (1941), they work with binary relations that are not necessarily anti-symmetric, and whose indifference curves are therefore not singletons. In their discussion of Wold’s theorem, Arrow-Hahn (1971, p. 106) write:

Since the introduction of indifference surfaces by Pareto and Irving Fisher, it has been taken for granted that they could be represented by a utility function. Wold seems to have been the first to see the need of specifying assumptions under which the representation by a continuous utility function exists.

In terms of bringing additional mathematical (non-topological) registers to bear on the continuity postulate, Wold’s theorem for relations constitutes another motivating result for this paper, and especially when coupled with Rosenthal’s 1955 theorem for functions. Wold seems to be the first author to formulate and use *scalar continuity* in economics in the framework of the consumer theory, and not only do Herstein and Milnor not connect to him, but as stated above, Koopmans separates them in two different essays. But even more than the representation theorem itself, Wold’s continuity assumption serves as a link between the results of Rosenthal and Herstein-Milnor, as well as articulating of a viewpoint that sees Debreu’s generalization of Wold’s work in the context of the criteria of *inessentiality* in Kim-Richter (1986), and *hiddenness* and *redundancy* explicated in Khan-Uyanik (2019). In the reference to Wold in his own study of the representation problem, Debreu (1954) found his assumptions to be “restrictive,” and in particular, singled out the Euclidean assumption, noting that to “treat the problem in a more general frame involves no additional mathematical cost.” In his comment on the antecedent literature, he found only the “particular case of the set of *prospects* as having received a “rigorous and extensive investigation,”<sup>12</sup> and in his subsequent treatment of the problem a decade later, he dropped any reference to Wold.

With this introductory framing as the backdrop, we are now in a position to spell out succinctly the substantive contribution of the paper. In a nutshell, we introduce new continuity

---

<sup>12</sup>See paragraph 2 of Debreu (1954) where he singles out von Neumann-Morgenstern (1947), Marschak (1950) and Herstein-Milnor (1953). We return to a reading of Debreu’s work below.

concepts for relations by using the concepts developed for functions, and use them not only to unify and simplify a range of results, but also to bring to light novel equivalences hitherto unseen in the economic literature. And perhaps most importantly, to see the neoclassical consumer theory as including decision theory now associated with the names of von Neumann, Morgenstern, Savage, Anscombe and Aumann. To be sure, as brought out above, some of the the continuity concepts for functions are part of the historical record, linear continuity and Rosenthal’s restricted continuity readily come to mind, but the translation from functions to relations has also yielded a new result for functions themselves; see, for example, the very first theorem below.<sup>13</sup> In Section 2, we recall the theorems of Rosenthal and Eilenberg along with the Genocchi-Peano example. In Section 3, we present six theorems, three propositions and a corollary: rather than repeat and summarize them – they are already clean and clear enough in their statements – we can invite the reader to quickly see how they articulate of the basic motivations delineated above, and how they relate to previous work in economic theory. In Section 4 we present seven propositions and three corollaries, and further discuss the implications of the results to the antecedent literature: these applications of the theory constitute the deliverables of the theory, applied theory so to speak. Section 5 concludes the paper with a summary and some observations regarding future and open direction. We reserve Appendix A.1 to the proofs of the results and the technical lemmata they require and Appendix A.2 to a battery of seven examples that close lacunae that may suggest themselves to an interested reader.

## 2 Mathematical Antecedents

### 2.1 Eilenberg’s Paper: A Conspicuous Omission

We begin with the omission: rather than the issue of priority, the importance of the Eilenberg paper lies in its substantial consequences for economic theory. First, as delineated in some detail in Khan-Uyanik (2019; 2020a), Eilenberg is the first to emphasize the behavioral consequences of what subsequently came to be regarded as mere technical assumptions made to ensure tractability, and thereby a professional missing-out of considerations later seminally considered by Sonnenschein (1965, 1967) and Schmeidler (1971). To elaborate, Eilenberg (1941) showed that a decision-maker (DM) with a continuous preference relation defined over a topologically connected choice set, must of necessity be consistent if he or she is strictly decisive, consistency being formalized by *transitivity*, and strict decisiveness by *completeness* of a binary relation that is assumed in addition to be anti-symmetric. Sonnenschein delineated circumstances under which Eilenberg’s theorem can be generalized to binary relations that are not necessarily anti-

---

<sup>13</sup>We are surprised that could not find any reference to the fact that for a quasi-concave (or a quasi-convex) function, linear continuity is equivalent to its continuity!

symmetric. In a parallel result, Schmeidler showed that under topological circumstances identical to Eilenberg’s, which is to say continuity and connectedness, a DM must of necessity be decisive if he or she is consistent. Under the rubric of what is referred to as the Eilenberg-Sonnenschein research program, these results have been comprehensively generalized and integrated by the authors:<sup>14</sup> they underscore the behavioral consequences of technical topological properties, and most particularly, the behavioral consequences of the continuity postulate.

The second contribution of Eilenberg’s paper is his theorem that a continuous, anti-symmetric, complete and transitive relation on a connected space can be represented by a continuous real-valued function. Prompted by Halmos, Debreu (1954, 1964) reached back to Eilenberg in choosing as his general frame of reference a topological space, albeit a preordered rather than an ordered one, but in which the preorder was not restricted by any ancillary monotonicity assumptions particular to the Euclidean setting. He extended Eilenberg’s result to relations that are not necessarily anti-symmetric, and sharpened it to highlight connectedness, separability and second-countability assumptions on the set over which the given relation is defined, formally defining a notion of a *natural* topology as being one in which the weak sections of the order are closed, and focussing on the quotient topology for his generalization.<sup>15</sup> Debreu’s extension of Eilenberg’s result has a clear and evident parallel to Sonnenschein’s extension of the first result, and has since emerged as a central preoccupation of decision theory as pursued in economic theory literature. To sum up, Eilenberg puts his entire emphasis on the topological register,<sup>16</sup> – what Arrow-Hahn refer to as the “deeper methods”<sup>17</sup> – and unlike Koopmans, metric considerations are totally bypassed and rendered inconsequential.<sup>18</sup> Second, the fact that these, and other issues, are considered in a single work allows one to see how the meaning given to the continuity postulate migrates from issue to issue and problem to problem.

---

<sup>14</sup>In this, Khan-Uyanik (2019) and Galaabaatar-Khan-Uyanik (2019) go against the grain of the literature.

<sup>15</sup>It is worth pointing out that this generalization is straightforward observation, if not a trivial one: it is the second theorem, and the alternative proof through the “open-gap lemma” that is the contribution of Debreu (1954); see Debreu (1964) and the definitive analysis of Beardon-Mehta (1994) and Beardon’s expository essay in Bosi-Campión-Candéal-Indurain (2020).

<sup>16</sup>This topological aspect was zeroed in by Peleg (1970) and subsequently by Lee (1972), Mehta (1983) and Herden (1995). Indeed, the work has been integrated into mainstream mathematics through a series of equivalence theorems that trace Cantor’s fundamental papers as the source of rich stream. See Herden (1995) for an omnibus result that involves an equivalence of eleven theorems one being used to provide a proof of the other; also see the texts Bridges-Mehta (1995) and Aleskerov-Monjardet (2002).

<sup>17</sup>In their masterly overview of the literature, Arrow-Hahn refer to the “very considerable generalization, based on a mathematical paper by Eilenberg, [that] was achieved with deeper methods by Debreu” in his reliance only on the topological notions of the continuity of preferences and the connectedness of the choice space on which the preferences were defined.

<sup>18</sup>A tracking of the word “distance” in Koopmans (1957) yields rewards in this connection.



## 2.2 Rosenthal’s Theorem: Suggested Directions

A point worthy of appreciation is the fact that even the move from a binary relation to the more primitive setting of a real-valued function does not yield an unambiguously single continuity postulate. The distinction between joint and separate continuity in each individual variable is to be sure a staple of the first course in real analysis, but *linear* continuity is perhaps less a part of the vernacular, at least in so far it is current in economics curriculum. Defining a real valued function on the plane to be *linearly* continuous when its restriction to a line is continuous, Genocchi-Peano (1884, pp. 173–174) provide the following example of a discontinuous function which is linearly continuous.

**Example (Genocchi-Peano).** *The function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  defined below is linearly continuous but not jointly continuous.*

$$f(x) = \frac{2x_1x_2^2}{x_1^2 + x_2^4} \text{ for } x \neq 0 \text{ and } f(0) = 0.$$

This basic example underscores the fact that linear continuity of a function is not enough to obtain “full continuity” on the entire domain.<sup>19</sup> It led one to ask for a concrete characterization of family of subsets in the domain of a function that ensure full continuity from continuity on the restricted domain. The answer to this question was given by a remarkable result of Rosenthal (1955) whose special case for the plane reads as follows:

**Theorem (Rosenthal).** *Every real-valued function is jointly continuous iff its restrictedly continuous, the restriction being any subset of the domain that can be represented as the graph of a smooth function of one variable in terms of the other.*

In terms of a general Euclidean space, the theorem established that the joint continuity of a real-valued function follows from the continuity of the restriction of the function to the graphs of all smooth curves in its domain.<sup>20</sup>

---

<sup>19</sup>The example, and along with the other GP examples more generally, is an important benchmark of a rich trajectory dating to Cauchy in the early part of the 18<sup>th</sup>-century. The fact that joint continuity is stronger than separate continuity was, even then in the time of Cauchy, standard material in textbooks on multivariate calculus, but an investigation of the relationship between joint continuity and restricted continuity properties of a function constituted a rich development to which many mathematicians, including Heine, Baire and Lebesgue, contributed; see for example Glatzer (2013) and the recent survey Ciesielski-Miller (2016).

<sup>20</sup>Note that in the functional form neither of the variables have to be fixed and the dependent variable can be one or the other. Furthermore, we take this continuity idea of Rosenthal and present an analogous definition for binary relations; see Definition 2 below. This is still admittedly loose in that we defer a formal definition of smoothness to the subsequent section and to the references in Footnote 41 below. Rosenthal restricts himself, for the plane, to the graph of a function that is both smooth and convex, and for the relation of convexity, see Glatzer (2013, Theorem 1.3.4).

In any case, Rosenthal’s theorem is the second important benchmark of the trajectory under consideration, and it motivates the deconstruction of the continuity postulate pursued here. Taking our cue from Rosenthal’s invocation of a restricted domain to generate an equivalence theorem, we invoke algebraic, order-theoretic and analytical structures to investigate the relationships between the myriad variety of continuity assumptions. This is to say, to investigate how convexity, monotonicity and differentiability structures, separately and together, render seemingly unrelated continuity assumptions equivalent for functions *and* relations. While of intrinsic interest, it bears emphasis that our attempt at a comprehensive and unified treatment is motivated primarily for the sharpening and the generalizing of the behavioural consequences that are already scattered in the literature: to identify resemblances and basic patterns obscured in a non-holistic view of the subject of choice theory.

### 3 The Theory

In this section we present the pure theoretical part of the work in the form of six theorems and two propositions collected in two subsections. It provides results on the relationship between different continuity postulates on functions and binary relations. The three categorizations of the continuity of preferences we mention in the introduction are a first exploratory overview, bringing out the important that the boundaries between them are not clear-cut and precise, and merit precise delineation.

#### 3.1 On the Continuity of Functions

As emphasized in the introduction, Rosenthal (1955) is the first to face the full consequences of the GP examples and ask for a strengthening of the linear continuity postulate on a function that would guarantee its full continuity: it offers a remarkable and elegant solution in terms of continuity of an *arbitrary* function on every smooth curve in a Euclidean space. In this paper we offer two results inspired by Rosenthal’s theorem. The first goes in a direction opposite to his by asking for restrictions on the class of functions rather than the subsets of their domains, for which linear and full continuity are identical; we present this result in this subsection. The second proceeds in a direction identical to his but for binary relations rather than a function; we present this result in the next subsection. It is a little surprising that these questions have not been asked before even in the mathematical literature, but their interest for us lies in that they set the stage for the results to follow.

We shall need the following notation for our first result.

**(C)** *A set  $X \subseteq \mathbb{R}^n$  satisfies property **C** if it is either open or a polyhedron where a polyhedron is a subset of  $\mathbb{R}^n$  that is an intersection of a finite number of closed half-spaces.*

We can now present our first result.

**Theorem 1.** *Let  $X \subseteq \mathbb{R}^n$  be a convex set with property **C**. Then every real-valued quasi-concave (or quasi-convex) function defined on  $X$  is linearly continuous iff<sup>21</sup> it is jointly continuous.*

We note for emphasis that the theorem is false without the sufficient conditions it assumes:

- (i) the quasi-concavity assumption cannot be dropped by virtue of the GP example;
- (ii) the polyhedron assumption in property **C** cannot be replaced by an arbitrary convex set by virtue of Example 7 in Appendix A.2.

Property **C** covers a wide range of applications in economics and mathematics, but it is still restrictive. Hence, we provide the following weaker property and leave it to the reader to check that the results using property **C** in this paper are true under this weaker property.

(**C'**) *A set  $X \subseteq \mathbb{R}^n$  satisfies property **C'** if there exist finitely many hyperplanes  $\{h_j\}_{j=1}^m$  in  $\text{aff}X$  such that  $X \setminus \text{ri}X \subseteq \bigcup_{j=1}^m h_j$ , where  $\text{aff}X$  denotes the affine hull of  $X$  and  $\text{ri}X$  denotes its relative interior.*<sup>22</sup>

Property **C'** imposes restrictions only on those points which are both in  $X$  and on its boundary in the affine hull of  $X$ . For example, the following sets satisfy property **C'** but fail property **C**:  $X = \{x \in \mathbb{R}^3 | x_3 = 0, x_1^2 + x_2^2 < 1\}$  and  $X' = \{x \in \mathbb{R}_+^2 | x_1^2 + x_2^2 < 1\}$ . Note that  $X$  is not open in  $\mathbb{R}^3$  but open in the affine hull of  $X$ , and  $X'$  is neither open nor a polyhedron, but the points that are both in  $X$  and on its boundary are contained in the union of two hyperplanes in  $X'$ .

We now illustrate myriad relationships among the different continuity postulates on functions in Figure 1.




Figure 1: The Continuity Postulates on Functions

The equivalence of joint-continuity and arc-continuity is used in Rosenthal (1955); the first red implication is proved in Theorem 1 above, the second red implication is due to Young (1910) and Kruse-Deely (1969); see also Ciesielski-Miller (2016) and Ghosh-Khan-Uyanik (2020) for implications of the monotonicity assumption for the relationship between separate and full continuity of functions and binary relations.

<sup>21</sup>The expression “iff” denotes “if and only if” hereafter.

<sup>22</sup>See Appendix A.1 for the definitions of the affine hull and the relative interior of a set.

It is well-known that a concave (or a convex) function on a non-empty, open and convex subset of  $\mathbb{R}^n$  is continuous. We next show that Theorem 1 above can be used to provide an alternative proof of this result.<sup>23</sup> First, we show that

**Proposition 1.** *Every concave (or a convex) real-valued function on a non-empty, open and convex subset of  $\mathbb{R}^n$  is linearly continuous.*

Since every concave (convex) function is quasi-concave (quasi-convex), the following is a direct corollary of Theorem 1 and Proposition 1.

**Corollary 1.** *Every real-valued concave (or a convex) function on a non-empty, open and convex subset of  $\mathbb{R}^n$  is continuous.*

We end this subsection by a remark on mid-point convexity for functions and relations; see for example Herstein-Milnor (1953) and Jensen (1967) for preferences, and Behringer (1992) and Kuczma (2009) for functions. The first two references already testify to the fact that the continuity and the convexity postulates have a certain complementary. It is also well-known that mid-point convexity of a function is equivalent to full convexity under the continuity postulate. It is an interesting question whether, under various versions of continuity mid-point convexity (mid-point versions of concavity, quasi-convexity and quasi-concavity) and its generalizations is equivalent to convexity (concavity, quasi-convexity and quasi-concavity). Further complementarities between various versions of convexity and continuity postulates are interesting and worth investigating.

### 3.2 On the Continuity of Relations

It is by now well understood that graph continuity implies section continuity; for complete and transitive relations, they are equivalent, however, for incomplete and non-transitive relations, further technical or behavioral assumptions on preferences are required in order to obtain the equivalence.<sup>24</sup> In this paper our emphasis is not on the investigation on this relationship. Pioneering treatment on the continuity assumptions are introduced in Birkhoff (1948), Nachbin (1965) and Villegas (1964). Section continuity of preferences implies certain scalar continuity postulates; Inoue (2010) illustrates that the example of GP shows that mixture-continuity is weaker than section continuity.<sup>25</sup> The relationship between the two are recently picked up in

---

<sup>23</sup>We thank an anonymous referee for pointing out this direction.

<sup>24</sup>See Schmeidler (1969) and Bergstrom-Parks-Rader (1976) for equivalence results for strict preferences; Ward (1954), Deshpandé (1968), Shafer (1974, Lemma) and Gerasimou (2015) for weak preferences; and Gerasimou (2013) for a comprehensive treatment for both weak and strict preferences.

<sup>25</sup>Similarly, Neufeld-Trockel (1995) provide an example for infinite-dimensional spaces; see Footnote 2 in their Section 3 and the text it footnotes. Also, see Footnote 57 below for details.

decision theory. It has been shown that under independence, or betweenness, assumptions, scalar continuity of a transitive preference relation is equivalent to its section continuity.<sup>26</sup> We investigate the relationship between section and scalar continuity postulates and introduce new continuity notions that are motivated by the literature in mathematics on the continuity of functions.

Let  $X$  be a set. A subset  $\succsim$  of  $X \times X$  denote a *binary relation* on  $X$ . We denote an element  $(x, y) \in \succsim$  as  $x \succsim y$ . The *asymmetric part*  $\succ$  of  $\succsim$  is defined as  $x \succ y$  if  $x \succsim y$  and  $y \not\succsim x$ , and its *symmetric part*  $\sim$  is defined as  $x \sim y$  if  $x \succsim y$  and  $y \succsim x$ . We call  $x \bowtie y$  if  $x \not\succ y$  and  $y \not\succ x$ . The inverse of  $\succsim$  is defined as  $x \precsim y$  if  $y \succsim x$ . Its asymmetric part  $\prec$  is defined analogously and its symmetric part is  $\sim$ . We provide the descriptive adjectives pertaining to a relation in a tabular form for the reader's convenience in the table below.

<i>reflexive</i>	$x \succsim x \ \forall x \in X$
<i>complete</i>	$x \succsim y \text{ or } y \succsim x \ \forall x, y \in X$
<i>non-trivial</i>	$\exists x, y \in X$ such that $x \succ y$
<i>transitive</i>	$x \succ y \succ z \Rightarrow x \succ z \ \forall x, y, z \in X$
<i>negatively transitive</i>	$x \not\succ y \not\succ z \Rightarrow x \not\succ z \ \forall x, y, z \in X$
<i>semi-transitive</i>	$x \succ y \sim z \Rightarrow x \succ z$ and $x \sim y \succ z \Rightarrow x \sim z \ \forall x, y, z \in X$

Table 1: Properties of Binary Relations

We now define “section continuity,” “graph continuity” and “restriction continuity” of a binary relation. Let  $\succsim$  be a binary relation on a set  $X$  that is endowed with a topology. For any  $x \in X$ , let  $A_{\succsim}(x) = \{y \in X | y \succsim x\}$  denote the *upper section* of  $\succsim$  at  $x$ , and  $A_{\precsim}(x) = \{y \in X | y \precsim x\}$  its *lower section* at  $x$ . The relation  $\succsim$  has *closed (open) graph* if it is closed (open) in the product space  $X \times X$ ; has *closed (open) upper sections* if it has closed upper sections; has *closed (open) lower sections* if it has closed lower sections; and has closed (open) sections if it has closed (open) upper and lower sections. Moreover,  $\succsim$  is *graph continuous* if it has closed graph and its asymmetric part  $\succ$  has open graph, and it is *continuous* if it has closed sections and its asymmetric part  $\succ$  has open sections. For any subset  $S$  of  $X$ , let  $\succsim|_S$  denote the *restriction* of  $\succsim$  to  $S$  and defined as follows: for all  $x \in X$ ,

$$A_{\succsim|_S}(x) = A_{\succsim}(x) \cap S, A_{\precsim|_S}(x) = A_{\precsim}(x) \cap S, A_{\succ|_S}(x) = A_{\succ}(x) \cap S, A_{\prec|_S}(x) = A_{\prec}(x) \cap S.$$

Notice that the restricted relation excludes the complement of  $S$  from the upper and lower sections of the relation. This definition of restricted continuity of a relation is different than the alternative requirement that  $\succsim|_S = \succsim \cap (S \times S)$ , and is natural in our context since the usual

<sup>26</sup>See, for example, Dubra-Maccheroni-Ok (2004, Proposition 1), Gilboa-Maccheroni-Marinacci-Schmeidler (2010, Lemma 3), Dubra (2011) and Karni-Safra (2015). We return to these references below.

continuity of a binary relation is defined by using its upper and lower sections. We want to focus on the restriction of the better-than- and worse-than-sets on a straight line for *all* points, not only at those contained in the line.<sup>27</sup> Note also that  $\succsim|S$  is continuous if  $A_{\succsim}(x) \cap S$  and  $A_{\preccurlyeq}(x) \cap S$  are closed (in  $S$ ) and  $A_{\succ}(x) \cap S$  and  $A_{\prec}(x) \cap S$  are open (in  $S$ ) for all  $x \in X$ .

We now turn to “scalar continuity” properties of a relation  $\succsim$  on a convex subset  $X$  of a vector space which is based on the topology of the unit interval. For all  $x, y \in X$  and  $\lambda \in [0, 1]$ , let  $x\lambda y$  denote  $\lambda x + (1 - \lambda)y$ . The relation  $\succsim$  is *upper (lower) mixture-continuous* if  $x, y, z \in X$  implies the set  $\{\lambda | x\lambda y \succsim z\}$  ( $\{\lambda | x\lambda y \preccurlyeq z\}$ ) is closed; *upper (lower) Archimedean* if  $x, y, z \in X, x \succ y$  implies that there exists  $\lambda \in (0, 1)$  ( $\delta \in (0, 1)$ ) such that  $x\lambda z \succ y$  ( $x \succ y\delta z$ ); and *upper (lower) strict-Archimedean* if  $x, y, z \in X$  implies the set  $\{\lambda | x\lambda y \succ z\}$  ( $\{\lambda | x\lambda y \prec z\}$ ) is open. Moreover,  $\succsim$  is *mixture-continuous* if it is upper and lower mixture-continuous. The *Archimedean*<sup>28</sup> and *strict-Archimedean* properties are defined analogously.<sup>29</sup> The relation  $\succsim$  is *order dense* if  $x \succ y$  implies that there exists  $z$  such that  $x \succ y \succ z$ . The relation  $\succsim$  is *weakly Wold-continuous* if (i) it is order dense and (ii)  $x \succ z \succ y$  implies that the straight line<sup>30</sup> joining  $x$  to  $y$  meets the indifference class of  $z$ .<sup>31</sup> Now, assume the space containing  $X$  is a finite dimensional Euclidean space. The relation  $\succsim$  is *Wold-continuous* if (i) it is order dense and (ii)  $x \succ z \succ y$  implies that any curve<sup>32</sup> joining  $x$  to  $y$  meets the indifference class of  $z$ .

We next define a restriction continuity concept for preferences which is analogous to linear continuity of a function.

**Definition 1.** *A binary relation on a convex subset  $X$  of a Euclidean space is linearly continuous if its restriction to any straight line in  $X$  is continuous.*

The following result illustrates that linear continuity of a relation is not a mere mathematical curiosity by showing that it is equivalent to mixture-continuity and Archimedean postulates.

**Proposition 2.** *A binary relation on a convex subset of a Euclidean space is linearly continuous iff it is mixture-continuous and Archimedean.*

<sup>27</sup>Example 5 in Appendix A.2 shows the crucial aspect of our definition. Moreover, both versions of restricted relations are used in mathematics; see for example Rebenciu (2008, Section 3).

<sup>28</sup>The Archimedean property in this paper is slightly stronger than the standard Archimedean property used in the literature: for all  $x, y, z \in S$  with  $x \succ y \succ z$ , there exists  $\lambda, \delta \in (0, 1)$  such that  $x\lambda z \succ y$  ( $x \succ y\delta z$ ) (or there exists  $\lambda \in (0, 1)$  such that  $x\lambda z \sim y$  as presented in Yokoyama). However, for complete and transitive relations, any of independence or strong monotonicity is enough to establish the equivalence between these two conditions.

<sup>29</sup>Strict-Archimedean property is stronger than Archimedean property, and the two properties are equivalent under mixture-continuity; see Galaabaatar-Khan-Uyanik (2019, Proposition 1).

<sup>30</sup>A *straight line in  $X$*  is defined as the intersection of a one dimensional affine subspace of  $\mathbb{R}^n$  and  $X$ .

<sup>31</sup>Note that the second part of weak Wold-continuity is equivalent to the *solvability* assumption in measurement theory; see for example Marschak (1950) and Krantz-Luce-Suppes-Tversky (1971).

<sup>32</sup>A *curve* on  $X$  is the range of a continuous injective function  $m : [0, 1] \rightarrow X$ ; see page 14 for details.

Next we provide convexity properties of a binary relation. Let  $\succsim$  be a binary relation on a convex subset  $X$  of a vector space.<sup>33</sup> The relation  $\succsim$  is *convex* if it has convex upper sections, and it *demonstrates convex indifference* if for all  $x, y \in X$  and all  $\lambda \in (0, 1)$   $x \sim y$  implies  $x \sim x\lambda y$ . A subset  $Y$  of a topological vector space is *locally convex* if each point of the closure of  $Y$ , denoted by  $\text{cl}Y$ , has a convex open neighborhood whose intersection with  $Y$  is convex. It is clear that every convex set is locally convex. However, the converse is not true since every finite subset of the space is locally convex but it is not convex.<sup>34</sup> A relation  $\succsim$  on  $Y$  is *locally convex* if it has locally convex upper sections.

We now provide monotonicity properties of a binary relation. Let  $\succsim$  be a binary relation on a subset  $X$  of  $\mathbb{R}^n$ . The relation  $\succsim$  is *strongly monotonic* if  $x > y$  implies<sup>35</sup>  $x \succ y$  and *weakly monotonic* if for all  $x, y \in X$ ,  $x > y$  implies  $x \succsim y$ . We call a subset  $A$  of  $X$  is *bounded by  $\succsim$*  if for all  $x, y \in A$  there exists  $a, b \in X$  such that  $a \succsim x, y$  and  $x, y \succsim b$ .

**(B)** A set  $X \subseteq \mathbb{R}^n$  satisfies property **B** if it is bounded by the usual relation  $\geq$ .

We now present an equivalence result among the seven continuity postulates we introduced above. It is an analogue of Theorem 1 above for preferences.

**Theorem 2.** Let  $\succsim$  be a complete and transitive relation on a convex subset of a Euclidean space with (i) property **C** and  $\succsim$  is convex, or (ii) property **B** and  $\succsim$  is weakly monotonic. Then the following seven continuity postulates for  $\succsim$  are equivalent: graph continuous, continuous, linearly continuous, mixture-continuous, Archimedean, Wold-continuous, and weakly Wold-continuous.

We note for emphasis that the theorem is false if properties **B** and **C** fail to hold; see Examples 6 and 7 in Appendix A.2. Property **C** holds for a simplex and the non-negative orthant in  $\mathbb{R}^n$ , the former is pervasive in decision theory and the latter in consumer theory, with convexity of preferences is a common assumption in both literatures. As to property **B**, when monotonicity of preferences is imposed (with or without convexity), the lattice structure is also assumed; see Gul (1992) in decision theory and Schmeidler (1969) in Walrasian economies where property **B** holds. Since properties **B** and **C** are used as substitutes in the theorem, and are not required to hold simultaneously, and the theorem can cover a wide range of models in economics. This being said, we note that properties **B**, **C**, and **C'**, the weakening of the latter introduced in Section 3.1 above, are still restrictive. For example, a simplex is a common assumption in decision theory which satisfies property **C** but fails property **B**. If a version of the convexity assumption, such as independence and betweenness, is not imposed in these models, then the theorem above does

<sup>33</sup>Diewert-Avriel-Zang (1981) discuss different convexity assumptions on functions.

<sup>34</sup>See Example 4 and the paragraph previous to it in Appendix A.2 for a discussion of local convexity.

<sup>35</sup>For vectors  $x$  and  $y$ , “ $x \geq y$ ” means  $x_i \geq y_i$  in every component; “ $x > y$ ” means  $x \geq y$  and  $x \neq y$ ; and “ $x \gg y$ ” means  $x_i > y_i$  in every component.

not apply even if the preference relation is assumed to be monotone.<sup>36</sup> Hence, the generalization of the results by weakening the assumptions on the choice sets is worthwhile and interesting.

Theorem 2 is a strong integration of continuity postulates. We next provide a result that first completely deconstruct the postulates by dropping all assumptions on preferences, and then a partial integration of the postulates by adding completeness and transitivity assumptions. It shows that some of the relationships in Theorem 2 can be recovered.

**Proposition 3.** *For any relation on a convex subset of a Euclidean space, only the following relationships hold among the seven continuity postulates.*

- (a) *Graph continuity  $\Rightarrow$  continuity  $\Rightarrow$  linear continuity  $\Leftrightarrow$  mixture-continuity & Archimedean  $\Rightarrow$  strict-Archimedean  $\Rightarrow$  Archimedean.*
- (b) *Wold-continuity  $\Rightarrow$  weak Wold-continuity.*

*If the relation is complete and transitive, then only the following additional relationships hold.*

- (c) *Continuity  $\Rightarrow$  graph continuity & Wold-continuity.*
- (d) *Strict-Archimedean  $\Rightarrow$  mixture-continuity  $\Rightarrow$  Archimedean & weak Wold-continuity.*
- (e) *Weak Wold-continuity  $\Rightarrow$  Archimedean.*

Please note the use of the word “only” in Proposition 3.<sup>37</sup> We next present a result establishing the equivalence between continuity and linear continuity of a relation which is not necessarily complete or transitive under different convexity assumptions.

**Theorem 3.** *Let  $\succsim$  be a binary relation on a convex set in  $\mathbb{R}^n$  with property **C**. Then  $\succsim$  is linearly continuous iff it is continuous, provided that any of the following holds.*

- (a)  *$\succsim$  is complete, and  $\succsim$  and  $\succ$  have locally convex upper sections,*
- (b)  *$\succsim$  and  $\succ$  have locally convex sections,*
- (c)  *$\succsim$  is reflexive and demonstrates convex indifference, and  $\sim$  is transitive.*

Figure 2 below illustrates the relationship among different continuity postulates on binary relations. Panel (a) illustrates the partial relationships among the continuity postulates based on Proposition 3, and Panel (b) further relationships in the presence of convexity or monotonicity assumptions based on Theorems 2 and 3.

Although independence axiom has been commonly used in decision theory, additivity is also an important convexity assumption. For example de Finetti’s (1974) axiomatic decision

<sup>36</sup>This paragraph is in response to a comment of an anonymous referee.

<sup>37</sup>Note that the Archimedean postulate or the other assumptions do not imply the continuity postulate. To belabour the point, there are no other relationships in addition to those illustrated in Proposition 3 without imposing additional assumptions on the choice set or on the binary relation.







Figure 2: The Continuity Postulates on Relations

theory model assumes the decision-maker has additive preference relation; see also Blackwell-Girshick (1954), Wakker (1988a,b, 1989, 1993) and Neufeind-Trockel (1995). We next present a result which provides a further deconstruction of the continuity postulate by showing that upper and lower scalar continuity properties are equivalent for additive relations.<sup>38</sup> Before that we need the following concepts. A *convex cone* is a subset  $X$  of a vector space such that  $\alpha x + \beta y \in X$  for all scalars  $\alpha, \beta > 0$  and all  $x, y \in X$ . A binary relation  $\succsim$  on a convex cone  $X$  is *additive* if for all  $x, y, z \in X$  with  $x + z, y + z \in X$ ,  $x \succsim y$  iff<sup>39</sup>  $x + z \succsim y + z$ .

**Theorem 4.** *Let  $\succsim$  be an additive binary relation on a convex cone. Then*

<sup>38</sup>Theorem 4 below shows that lower scalar continuity is equivalent to upper scalar continuity under the additivity assumption on a convex cone. McCarthy-Mikkola (2018) provide analogous equivalence results for an independent, reflexive and transitive relation on a convex set. Theorem 4 generalizes their result by dropping the transitivity and reflexivity assumption and weaken the independence assumptions. The works of Neufeind-Trockel (1995, Proposition), Candeal-Haro-Induráin-Eraso (1995, Theorem) and Gerasimou (2015, Theorem 1) provide equivalence results between lower and upper section continuity of a relation. Theorem 4 provides an analogous result for scalar continuity. Example 3 in Appendix A.2 illustrates that neither our result implies the results in Neufeind-Trockel and Gerasimou, nor their results imply ours.

<sup>39</sup>When the set  $X$  is a vector space, then “iff” can be equivalently replaced by “if”.

- (a)  $\succsim$  is upper mixture-continuous iff it is lower mixture-continuous iff it is mixture-continuous,
- (b)  $\succ$  is upper strict-Archimedean iff it is lower strict-Archimedean iff it is strict-Archimedean.

The following result shows that some of the equivalence results pertaining to scalar continuity postulates in Theorem 2 do not require finite-dimensionality or a topological structure on the choice set.

**Theorem 5.** *For every convex, complete and transitive relation on a convex subset of a vector space, the following continuity postulates are equivalent: Archimedean, strict Archimedean, mixture-continuous and weakly Wold-continuous.*

It is easy to show that the result above is true for mixture sets.

Next, we proceed in a direction identical to Rosenthal's by asking for a condition on an *arbitrary* binary relation that guarantees its continuity. We show that continuity along every smooth curve is enough for full continuity. The question of course is the kind of continuity that is at issue. The GP example was recently picked up (without citation) by Inoue (2010, Example 1) in order to illustrate that mixture-continuity is weaker than the continuity of a complete and transitive relation by showing that the relation induced by the function in GP example is mixture-continuous but not continuous. The interesting question of course is what condition works? Before we develop this, it is well to put down some basic terminology.

We now define continuity of preferences restricted to continuously differentiable curves. An *arc* in  $\mathbb{R}^n$  is a continuous injective function  $m : [0, 1] \rightarrow \mathbb{R}^n$  where  $m(\lambda) = (m_1(\lambda), \dots, m_n(\lambda))$ . An arc is called *smooth* if  $m_i$  is continuously differentiable for all  $i$  and  $m'(\lambda) = (m'_1(\lambda), \dots, m'_n(\lambda)) \neq 0$  for all  $\lambda \in [0, 1]$ . A *curve* in  $\mathbb{R}^n$  is the range of an arc and a *smooth curve* is the range of a smooth arc. Since an arc  $m$  is continuous and injective, it is a bijection from  $[0, 1]$  to its image  $m([0, 1])$ . Since  $[0, 1]$  is compact and  $m([0, 1])$  is Hausdorff,  $m$  is an homeomorphism between  $[0, 1]$  and  $m([0, 1])$ ; see for example Dugundji (1966, Theorem 2.1, p. 226). Therefore, an arc and the curve induced by it are homeomorphic.<sup>40</sup>

**Definition 2.** *A binary relation on a subset  $X$  of a Euclidean space is arc-continuous if its restriction to any smooth curve in  $X$  is continuous.*

We next introduce stronger versions of mixture-continuity and Archimedean properties by using smooth arcs instead of straight lines. The following notation for subsets of  $[0, 1]$  is useful. Let  $\mathcal{M}$  be the set of all smooth arcs. For all  $m \in \mathcal{M}$ , when the values  $m(1) = x$  and  $m(0) = y$  are of interest, we use the notation  $m_{xy}(\lambda)$  for  $m(\lambda)$ . Define the set of all smooth curves induced

---

<sup>40</sup>Note that an arc induces a unique curve but a curve can be induced by distinct arcs. For example, any closed segment  $[x, y]$  of the diagonal in  $\mathbb{R}^2$  is induced by every arc with  $m_1(\lambda) = m_2(\lambda)$  for all  $\lambda \in [0, 1]$  where  $m_1(0) = x, m_1(1) = y$ .

by the set of smooth arcs as  $\mathcal{C}_{\mathcal{M}}$ . For any binary relation  $\succsim$  on a convex set  $X$ , any  $x \in X$  and any function  $f : [0, 1] \rightarrow X$ , define

$$I_{\succsim}(f, x) = \{\lambda \mid f(\lambda) \succsim x\} \quad \text{and} \quad I_{\preccurlyeq}(f, x) = \{\lambda \mid x \succsim f(\lambda)\}. \quad (1)$$

A binary relation  $\succsim$  on a convex subset  $X$  of  $\mathbb{R}^n$  is *strongly mixture-continuous* if for all  $x, y, z \in X$  and all  $m_{xy} \in \mathcal{M}$ , the sets  $I_{\succsim}(m_{xy}, z)$  and  $I_{\preccurlyeq}(m_{xy}, z)$  are closed, is *strongly Archimedean* if for all  $x, y, z \in X$  with  $x \succ y$  and all  $m_{xz}, m_{yz} \in \mathcal{M}$ , there exists  $\lambda, \delta \in (0, 1)$  such that  $m_{xz}(\lambda) \succ y$  and  $x \succ m_{yz}(\delta)$  and is *strongly strict-Archimedean* if for all  $x, y, z \in X$  and all  $m_{xy} \in \mathcal{M}$ , the sets  $I_{\succ}(m_{xy}, z)$  and  $I_{\prec}(m_{xy}, z)$  are open.

**Theorem 6.** *Let  $\succsim$  be a binary relation on a convex set in  $\mathbb{R}^n$  with property **C**. Then  $\succsim$  is arc-continuous iff it is continuous iff it is strongly mixture-continuous and strongly Archimedean.*

Rosenthal's theorem characterizes the family of subsets such that restriction continuity of an arbitrary function is equivalent to its continuity. Theorem 6 generalizes Rosenthal's result to binary relations by showing that continuity of an arbitrary binary relation is equivalent to its restriction continuity on smooth curves.<sup>41</sup>

## 4 The Theory Applied

This section takes to opportunity to connect and discuss the contributions of our results on the continuity of preferences and functions on the antecedent literature in economics. Theorem 2 of the previous section brought out the many different meanings – eight to be precise – that can be given to the continuity postulate, and that in so doing, give it various colorings that, in conjunction with the five other results, thereby also impinge on the postulate's form and content. We now ask for the deliverables of these results in so far as applications are concerned. Our discussion is sectioned into four parts: ranging from representability, to neoclassical consumer theory and its incorporation into Walrasian equilibrium theory, to the behavioral implications of the transitivity and continuity postulates and finally, all that still remains to be done.

### 4.1 Representation of Preferences: Benchmark Results

In applying the theorems presented in Section 3, we make three introductory observations: (i) whereas scalar continuity assumptions have been mainly used in choice theory under uncertainty

---

<sup>41</sup>Rosenthal's theorem uses simple arcs which is a subset of the class of smooth curves. It is easy to observe that our results hold for simple arcs. We present our result by using smooth curves for the exposition purposes which is by now standard in this literature. See Young-Young (1910), Kershner (1943), Rosenthal (1955) and Ciesielski-Miller (2016) for the deconstruction of the continuity postulate for functions; also Footnote 20 above.

regarding cardinal representation of preferences, Wold had already relied on them in his treatment of non-stochastic consumer theory, (ii) in Walrasian general equilibrium theory, scalar continuity is dropped and section continuity is used. The scalar continuity assumptions, by themselves, are weaker than section continuity. In decision theory, consumer theory and Walrasian economies, either convexity or monotonicity assumptions are imposed on preferences. We show in this section that the distinct use of continuity postulates is inessential – in the presence of the other assumptions of the model, scalar and section continuity assumptions are identical. It is interesting that this relationship is not asked in consumer theory and Walrasian economies. There are some scattered special partial equivalences in decision theory,<sup>42</sup> yet a comprehensive analysis, which connects different literatures and results, is missing. We wish to emphasize that the following categories are not mutually exclusive.

In this subsection, we present three corollaries and three propositions. The first corollary illustrates that any of the eight continuity postulates can be equivalently used in Wold’s representation theorem, the second shows that in the expected utility theorem of Herstein-Milnor (1953), mixture continuity can be replaced with weak Wold-continuity, and the third obtains the expected utility theorem of Neufeld-Trockel (1995) as a corollary of our results. The first proposition establishes the link between additivity and independence postulates, the second obtains an expected utility representation theorem under additivity postulate and the last shows that the section continuity assumption of Gul (1992) can be replaced by mixture-continuity.

#### 4.1.1 An Euclidean Space: Wold (1943–44)

Wold (1943–44) in his seminal paper proved a utility representation theorem. The following is a direct corollary of Wold’s theorem and Theorems 2 and 6.

**Corollary 2.** *Every complete, transitive, weakly monotonic preference relation on  $\mathbb{R}_+^n$  is representable by a continuous utility function iff the preference relation satisfies any of the following eight continuity postulates: graph continuous, continuous, arc-continuous, linearly continuous, mixture-continuous, Archimedean, Wold-continuous, and weakly Wold-continuous.*

Note that this result provides a novel form to a classical theorem by providing equivalences under eight different continuity postulates; hence in addition to a consolidation of the antecedent literature, it unmask new connections. In particular, this result shows that we can replace Wold-continuity with any of the continuity properties listed in Theorem 2. The precise continuity hypotheses assumed by Wold in his theorems are fundamentally germane to this paper. In Wold’s proof, monotonicity of the relation plays a crucial role. Wold’s representation theorem in

---

<sup>42</sup>See Gilboa-Maccheroni-Marinacci-Schmeidler (2010), Dubra (2011) and Karni-Safra (2015).

his seminal work Wold (1943–44) provides an equivalence relationship between Wold-continuity and section continuity of a relation on a Euclidean space under a weak monotonicity postulate. Almost ten years after his papers, Wold provides separate sufficient conditions for the existence of a continuous utility representation in Wold-Jureen (1953). The new theorem replaces weak monotonicity with strong monotonicity, and Wold-continuity with weak Wold-continuity.<sup>43</sup> There is a dissonance in the literature in the way that these two theorems were received, but this need not concern us here.<sup>44</sup> The point is that the two separate continuity assumptions of Wold can be considered as a link between the scalar continuity assumption of Rosenthal which is based on smooth curves and of Herstein-Milnor which is based on straight lines. We elaborate this link in this paper by showing that under a weak monotonicity assumption, commonly used scalar continuity assumptions on preferences are equivalent to each other, and also each is equivalent to the continuity of the relation.

As we mention in the Introduction, Eilenberg’s and Debreu’s representation result is more general than Wold’s theorem since they simply drop the monotonicity assumption.<sup>45</sup> And, even though Wold’s continuity assumption is by itself weaker than the section continuity property used in Eilenberg and Debreu, we showed that they are equivalent in the presence of other assumptions of Wold.<sup>46</sup> Moreover, Arrow-Hahn (1971) in their representation theorem do not impose convexity, or monotonicity, assumptions on preferences, hence our results are silent about the possibility of replacing the continuity assumption with mixture-continuity. In fact, as we already noted in the Introduction, the Genocchi-Peano example provides an example which satisfy the assumption of Arrow-Hahn except section continuity. Hence, it is indeed not possible to replace the continuity assumption with scalar continuity in Arrow-Hahn’s theorem.<sup>47</sup> Note,

---

<sup>43</sup>Even though Wold assumes only part (ii) of weak Wold-continuity, it is easy to show that its part (i) follows from strong monotonicity under the transitivity postulate; see Example 2 and the discussion following it in Section A.2.

<sup>44</sup>Yokoyama (1956), Arrow-Hahn (1971) and Mitra-Ozbek (2013) mainly picked the set of assumptions listed in Wold’s book. Yokoyama (1956) provides a direct proof of the relationship between weak Wold-continuity and continuity of a relation on  $\mathbb{R}_+^n$  under strong monotonicity postulate. On the other hand, Beardon-Mehta (1994) pick up the hypothesis of Wold’s theorem as stated in his papers and show that Wold’s method of proof can be used in order to obtain a simple alternative proof of Debreu’s theorem. Moreover, Lloyd-Rohr-Walker (1967) provide an explicit proof of Wold’s theorem he presented in his papers, but under strong monotonicity. See also Yokoyama (1954) for a discussion on Wold’s strong continuity assumption in consumer theory and Banerjee-Mitra (2018) for a recent reference to Wold’s work.

<sup>45</sup>Even though Eilenberg’s result is for anti-symmetric relations, it is trivial to observe that by using the tools of the quotient space, the anti-symmetry assumption can simply be dropped; see for example Debreu (1954). Of course this does not devalue Debreu’s or Wold’s contributions – Debreu’s celebrated open gap lemma and Wold’s simple method-of-proof have substantial effect in economic theory; see Beardon-Mehta (1994) for a discussion.

<sup>46</sup>For ordinal representation of preferences, see Hervés-Beloso-del Valle-Inclán Cruces (2019) and its reference to Hausdorff’s work.

<sup>47</sup>Beardon-Mehta (1994) has an excellent survey and extension on Arrow-Hanah’s method and its relation to other method’s including Debreu’s and Wold’s methods. In particular, they consider the unit interval endowed with an equivalence relation, and under the assumption of closed equivalence classes, show the existence of

however, that the fact that the utility representation results of AH and Debreu does not use convexity, or monotonicity assumptions in their utility representation theorems, in many of their applications they assume continuous and convex preferences, which have quasi-concave utility representations. In these results we can replace the continuity assumption with various continuity assumptions we listed in Theorem 2. This observation also connects us to our Theorem 1.

#### 4.1.2 A Mixture Space: Herstein and Milnor (1953)

The following is a corollary of Herstein-Milnor (1953), von Neumann-Morgenstern (1947), Jensen (1967) and Theorem 5.

**Corollary 3.** *A binary relation defined on a mixture set is representable by a mixture-linear<sup>49</sup> function iff it is complete, transitive, independent and satisfies any of the following three continuity postulates: mixture-continuous, Archimedean, and weakly Wold-continuous.*

As we mention above, this result provides a novel form to the classical expected utility theorem and connects it to Wold’s work. Note that the necessity and sufficiency of mixture continuity is proved by Herstein-Milnor (1953), of Archimedean by von Neumann-Morgenstern (1947) and Jensen (1967),<sup>50</sup> and of weak Wold-continuity follows from Theorem 5.

#### 4.1.3 A Finite State Space: Gul (1992)

Gul (1992) re-does Savage’s work for finite state space by imposing topological structure on the set of outcomes.<sup>51</sup> His framework is as follows. Let  $N = \{1, \dots, n\}$ ,  $\mathcal{N} = 2^N \setminus \emptyset$  and  $X = [0, 1]^n$ . Let  $\Delta = \{x \in X | x_i = x_j \text{ for all } i, j \in N\}$ . For  $x \in \Delta$ , we use  $x$  and  $x_i$  interchangeably for any  $i \in N$ . For  $I \in \mathcal{N}$  and  $x, y \in \Delta$ ,  $xIy$  denotes  $z$  such that  $z_i = x_i$  for  $i \in I$  and  $z_i = y_i$  for  $i \notin I$ . Let  $\succsim$  be a preference relation on  $X$ . A set  $I \in \mathcal{N}$  will be called *null* if  $x_i = y_i$  for all  $i \in I^c$  (i.e.,  $x = y$  on  $I^c$ ) implies  $x \sim y$ . For simplicity, we assume no  $i \in N$  is null. Consider the following axioms.

---

a a non-decreasing real-valued continuous function on the interval that is constant on each equivalence class. Ascribing this result of real analysis to Wold, they use it to prove Debreu’s representation theorems as well as his “open-gap lemma” which he used to prove his theorems.<sup>48</sup> Furthermore, they go beyond this equivalence and a corresponding revisionary re-framing of Wold’s work to prove a generalized version of Arrow-Hahn’s representation theorem. However, the point of departure for the investigation pursued and reported here in this paper is not this question of whether one theorem can be used to prove, and be proved by, another, but rather the conceptual move that Wold makes from the topology on the choice space to that of the topology on the operation under which these objects of choice can be combined. Finally, see also Bosi-Campi3n-Candeal-Indurain (2020) on recent results on representation of preferences and in particular Bosi-Zuanon (2020, Theorem 2.23) for equivalence results on section and graph continuity postulates.

<sup>49</sup>A function  $l : X \rightarrow \mathbb{R}$  defined on a convex subset of a vector space  $X$  is *mixture-linear* if  $l(x\lambda y) = \lambda l(x) + (1 - \lambda)l(y)$  for all  $x, y \in X$  and  $\lambda \in [0, 1]$ .

<sup>50</sup>See Fishburn (1982) for a detailed discussion on this.

<sup>51</sup>See also Wakker (1988a, 1989) and Gravel-Merchant-Sen (2012).

- (A1)  $\succsim$  is complete and transitive.
- (A2) For  $I \in \mathcal{N}$ ,  $x'_i \sim x_i I z_i, y'_i \sim y_i I z_i$  for all  $i \in N$  implies  $x \succ y$  if and only if  $x' \succ y'$ .
- (A3) For all  $x, y \in \Delta$ ,  $x \succ y$  implies  $x \succsim y$ . Furthermore, there exist  $I^* \subseteq N$  such that for all  $x, y \in \Delta$ ,  $x I^* y \sim y I^* x$ .
- (A4)  $\succsim$  is continuous.

The following result and Theorem 2 imply that A4 and mixture-continuity of  $\succsim$  are equivalent, hence A4 in Gul (1992, Theorem) can be replaced by mixture continuity.<sup>52</sup>

**Proposition 4.** *Under A1–A3, if  $\succsim$  is mixture-continuous, then it is strongly monotone.*

#### 4.1.4 A Topological Vector Space: Neufeld-Trockel (1995)

We now present the relationship between additivity and the following two convexity assumptions on binary relations. A relation  $\succsim$  on a convex cone  $X$  is *independent* if for all  $x, y, z \in X$  and all  $\lambda \in (0, 1)$ ,  $x \succsim y$  implies  $x\lambda z \succsim y\lambda z$ , and it is *homothetic* if for all  $x, y \in X$  and all  $\lambda \in \mathbb{R}_{++}$  with  $\lambda x, \lambda y \in X$ ,  $x \succsim y$  implies  $\lambda x \succsim \lambda y$ . Independence and additivity assumptions are used commonly in decision theory, homotheticity is used mainly in consumer theory. The following result provides an equivalence relationship between these three common convexity postulates in decision theory as well as in consumer theory: additivity, homotheticity and independence.

**Proposition 5.** *The following are true for any transitive relation  $\succsim$  defined on a vector space.*

- (a)  $\succsim$  is additive and homothetic iff it is independent.
- (b) If  $\succsim$  is complete and mixture-continuous, then it is additive iff it is independent.

We next present a result that provides an expected utility representation theorem under minimal (explicit) technical and behavioral assumptions on preferences.

**Proposition 6.** *The following are equivalent for any non-trivial relation on a vector space.*

- (a) The relation is non-trivial, semi-transitive, additive, upper mixture-continuous and upper Archimedean.
- (b) The relation is representable by a mixture-linear function.

It should be noted that this is a new representation theorem by virtue of the fact that it replaces section continuity assumptions of Neufeld-Trockel (1995, Proposition) by weaker scalar continuity assumptions. Note also that Proposition 6 does not explicitly assumes completeness,

---

<sup>52</sup>Similar results can be obtained in Chew-Karni (1994), which generalize Gul (1992) and Nakamura (1990); Dekel (1986), Ahn-Ergin (2010), and Maccheroni-Marinacci-Rustichini (2006).

transitivity, independence and full mixture-continuity of the relation. However, if a mixture-linear representation exists, then all of these properties hold by necessity. Therefore, these assumptions are *hidden* in the hypothesis of this result. Since Proposition 6 and Herstein-Milnor's theorem have *essentially* the same hypothesis, it is an *inessential* generalization of the theorem of Herstein-Milnor in the sense of Kim-Richter (1986).

The following is a corollary of Proposition 6 and the representation theorem of Debreu (1954, Theorem I).

**Corollary 4.** *Every non-trivial, semi-transitive and additive relation  $\succsim$  on a topological vector space with  $A_{\succsim}(0)$  is closed and  $A_{\succ}(0)$  is open, is complete, transitive, continuous and representable by a continuous linear utility function.*

This corollary is presented first in Neuefeind-Trockel (1995, Proposition). Proposition 6 replaces closed upper sections of the weak relation and open upper sections of the strict relation with the weaker upper mixture-continuity and upper Archimedean properties, respectively. Hence, we obtain a representation theorem under a weaker continuity assumption. Of course the utility function Proposition 6 yields may not be continuous. However, since the preference relation is complete, the preference relation has closed upper and lower sections. Then, Debreu's representation theorem imply that there exists a continuous utility representation which is also linear, due to additivity.

## 4.2 Neoclassical Consumer Theory

In this subsection, we present two observations and two propositions. The first observation illustrates an *inessential* remark of Debreu, and the second illustrates that and of the eight continuity postulates above can be equivalently used for the results in the Walrasian equilibrium theory and neoclassical consumer theory. The two propositions show that some of the equivalences survive even if completeness or transitivity assumptions are dropped.

### 4.2.1 Debreu's (1959) Remark

We have already related our results to the antecedent literature, but in a sharp way that may be missing the outlines of the forest in its focus on its individual trees. In this section, we step back and provide an overview of the results from the criterial perspective of *inessentiality* in Kim-Richter (1986), and its invocation in Epstein (1987), and those of *hiddenness* and *redundancy* explicated in Khan-Uyanik (2019).

We already referred to Debreu's generalization of Wold's representation theorem from Euclidean spaces by using the section continuity assumption to eliminate monotonicity. We have now already seen in Theorem 2 that under the hypothesis of Wold's theorem Debreu's



section continuity assumption is not only equivalent to Wold-continuity but also the commonly used mixture-continuity, Archimedean and strong Wold-continuity properties.<sup>53</sup> If we drop the monotonicity assumption, these scalar continuity properties are weaker than section continuity. However, as we presented above, monotonicity can be compensated by convexity in order to obtain the equivalences. Our results in Section 3 illustrate the fact that seemingly unrelated topological, order and convexity properties together have behavioral topological implications on preferences. Hence, a scalar continuity assumption on preferences is strictly weaker than the section continuity property, however, in the presence of additional assumptions on preferences they may be equivalent. We illustrate this point by providing an example from the classic work of Debreu. He writes

Certain theorems whose statements list (a) of 4.6 [closed upper sections] among their hypotheses can, in fact, be proved by using weaker continuity assumptions on preferences [upper mixture-continuity] inspired by I. N. Herstein and J. Milnor. Debreu (1959, p.73 )

He then notes that he can replace the closed upper sections assumption with the weaker upper mixture-continuity property in order to obtain a relationship between certain convexity assumptions on preferences and in the statement of the second welfare theorem. Lemma 6 implies that under upper mixture-continuity assumption, Debreu's convexity assumption (star-convexity of  $\succ$ ) is equivalent to convexity of  $\succ$ . And Lemma 3 implies that upper mixture-continuity and closed upper sections are equivalent properties under the hypothesis of Debreu's results.

**Observation 1.** *Debreu's remark about replacing closed sections assumption with the upper mixture-continuity property is inessential.*

Wold (1943), Wold (1943–44) provides a model of consumer theory which assumes the consumers have complete, transitive, weakly monotonic and Wold-continuous preference relations. Wold-Jureen (1953) replaces the last two assumptions with strong monotonicity and weak Wold-continuity assumptions. Our results show that we can replace his continuity assumption with any of the scalar, section or graph continuity assumptions listed in Theorem 2. We get back to this in Section 4.2.4 below.

#### 4.2.2 Schmeidler's (1969) Non-decisive Consumer

Schmeidler (1969) generalizes Aumann's (1966) work on the existence of a general equilibrium with continuum of players by dropping the completeness assumption. Schmeidler's consumer has a *strict preference relation defined on  $\mathbb{R}_+^n$  which is irreflexive, strongly monotonic, transitive*

---

<sup>53</sup>To be precise, the equivalence of parts (a) and (d) of Theorem 2 under completeness, transitivity and weak monotonicity assumptions.

with open sections. The following result shows that Schmeidler’s continuity assumption can be replaced with the Archimedean property.<sup>54</sup>

**Proposition 7.** *The preference relation of Schmeidler’s consumer is Archimedean iff it is strict-Archimedean iff it has open sections.*

Note that in the classical consumer theory, completeness and transitivity are assumed. We show that different continuity assumptions are equivalent under these assumptions. Our discussion of the work of Shafer-Schmeidler show that when one of these two behavioral assumptions, we can still have equivalence results.

In his result, Schmeidler does not assume full continuity, instead he assumes only that the strict relation has open sections. With the added full continuity and transitivity assumptions, however, Schmeidler’s result would be a straightforward corollary of Aumann’s theorem. Therefore, Schmeidler’s result is an *essential* generalization as a result of weaker continuity and transitivity assumptions – without such a weakening, completeness and full transitivity would already be hidden in his seemingly more general assumptions. The recent work on Walrasian general equilibrium and game theory with discontinuous and/or non-ordered preferences is in line with this observation; see Reny (2020) for a recent survey on this literature.

#### 4.2.3 Shafer’s (1974) Non-transitive Consumer

Shafer (1974) re-works the neoclassical theory of demand for a consumer with possibly non-transitive preferences defined on  $\mathbb{R}_+^n$ . Shafer’s consumers have *continuous, complete and convex preferences whose asymmetric part is also convex*. The following result shows that it is possible to replace the section continuity assumption of Shafer with and of the mixture-continuity or Archimedean properties.

**Proposition 8.** *The preference relation of Shafer’s consumer is Archimedean iff it is mixture-continuous iff it is continuous.*

Moreover, Khan-Uyanik (2019) noted that if the preferences of Shafer’s non-transitive consumer satisfy semi-transitivity, or transitive indifference, or transitive strict relation, then the preference relation is transitive. That is to say, a “little” bit of consistency implies “fully” consistency in Shafer’s model of non-transitive consumer, thereby destructive of “all” non-transitivity. Therefore, Shafer’s non-transitive consumer has to be, by necessity, a “fundamentally” non-transitive agent.

---

<sup>54</sup>Note that we define Archimedean property for an arbitrary relation  $\succsim$  where its asymmetric part is  $\succ$ . In Schmeidler’s model, the primitive is a transitive and irreflexive relation  $\succ$ . Then it is asymmetric. Hence, its asymmetric part is equal to the relation itself.

#### 4.2.4 Hidden Assumptions in Walrasian Equilibrium Theory

The classical Walrasian equilibrium theory and neoclassical consumer theory assume that the preferences of the consumers are complete, transitive and convex (and/or monotone); see for example Debreu (1959), Aumann (1966), Arrow-Hahn (1971) and McKenzie (2002). For suitable consumption sets, it follows from Theorem 2 above that

**Observation 2.** *The continuity assumptions in the Walrasian equilibrium theory and neoclassical consumer theory can be replaced by any of the eight continuity postulates above.*

In many results in consumer theory and Walrasian equilibrium, our results imply that completeness and full transitivity properties follow from the other assumptions; see Khan-Uyanik(2019) for an extended discussion on the Walrasian equilibrium theory and consumer theory in the above-mentioned work. We will return to the hiddenness of completeness and full transitivity properties in the subsequent subsection.

### 4.3 Consistency and Decisiveness: A Reconsideration

In this subsection, we present two propositions on the behavioral implications of continuity postulates. The first proposition uses the *Dubra-method* to show that convexity and scalar continuity imply completeness and full-transitivity of preferences, and the second shows that additivity, a stronger convexity property, allows weakening of the continuity postulates in the first proposition.

We start with

**Proposition 9.** *Let  $X$  be a convex set in  $\mathbb{R}^n$  with property **C**. Then every non-trivial, reflexive, mixture-continuous and Archimedean binary relation on  $X$  which demonstrates convex indifference and has a transitive symmetric part, is continuous, complete and transitive.*

Scalar continuity, such as Archimedean or mixture-continuity assumptions, is used extensively in decision theory. An equivalence of this type has useful implications. Dubra-Maccheroni-Ok (2004, Proposition 1) shows that for a reflexive, transitive and independent relation, a stronger version of mixture-continuity, which is due to Shapley-Baucells (1998), is equivalent to closed graph property on a finite simplex. Gilboa-Maccheroni-Marinacci-Schmeidler (2010, Lemma 3) shows that if a reflexive and transitive relation on a normed space satisfies independence and monotonicity properties, then its mixture-continuity is equivalent to the closed graph property.<sup>55</sup> Dubra (2011) and Karni-Safra (2015) show that if a reflexive and transitive relation on a simplex satisfies any of the independence, betweenness, and cone monotonicity properties, then mixture-continuity and Archimedean properties imply continuity if the relation on a finite simplex.

---

<sup>55</sup>See also Chateauneuf-Cohen-Jaffray (2013) and Abdellaoui-Wakker (2020).

Proposition 9 in generalizes these results by weakening, or dropping, the convexity, monotonicity, reflexivity and transitivity postulates, and by expanding the choice set in the context of Euclidean spaces.<sup>56</sup>

We next provide a result for arbitrary vector spaces illustrating the behavioral implications of scalar continuity in the presence of additivity.

**Proposition 10.** *Every non-trivial, semi-transitive, additive, upper mixture-continuous and upper Archimedean relation on a vector space, is complete and transitive.*

Neuefeind-Trockel (1995, Proposition) prove a special case of this result for topological vector spaces under stronger section continuity postulates. Proposition 10 provides a considerable generalization of Neuefeind-Trockel’s result by replacing their continuity assumption with the weaker linear continuity postulate and allowing the space to be an arbitrary vector space.<sup>57</sup> Lastly, this result illustrates the strength of the additivity assumption by showing that it allows us to obtain both completeness and transitivity of a relation under weak continuity and transitivity assumptions. Note that the transitivity of  $\sim$  plays a crucial role in the antecedent results in the Eilenberg-Sonnenschein research program. This result, by replacing the transitive indifference assumption with the semi-transitivity postulate under the additivity assumption complements this research program.

#### 4.4 Other Potential Applications

Our investigation has limited itself to finite-dimensional Euclidean spaces, as well as to a vector space whose dimension is not of any consequence. As such, it asks for a natural extension to an infinite-dimensional setting that brings topological and other algebraic structures to bear on

---

<sup>56</sup>Note that the result presented in Gilboa-Maccheroni-Marinacci-Schmeidler (2010, Lemma 3) is for infinite dimensional spaces, hence our result is not a full generalization of their result. Moreover, Karni-Safra (2015) replace the independence assumption in Dubra’s theorem with the betweenness, or cone-monotonicity, assumption. Proposition 9 generalizes their results by weakening, or dropping, the convexity, reflexivity and transitivity postulates, by expanding the choice set, and more importantly by obtaining *both* completeness and transitivity as necessary conditions. Note that Galaabaatar-Khan-Uyanik (2019, Theorem 1) provide more general results than Proposition 9. However, our method-of-proof is different than theirs – their method-of-proof is direct and also their choice set does not have a topological structure. We, on the other hand, first obtain the continuity of a linearly continuous relation, and then apply a result due to Khan-Uyanik (2019, Theorem 2) in order to obtain completeness and transitivity postulates. This method-of-proof is in line with the method of proof of Dubra which is then picked up by Karni-Safra.

<sup>57</sup>This generalization requires a proof technique which is based on a closure concept for subsets of a vector space by using the topological structure of the unit interval; see the proof in Appendix A.1 for details. Moreover, Neuefeind-Trockel in Footnote 2 in their Section 3 note the fact that scalar continuity assumption is weaker than section continuity by providing an example of a preference relation defined on an infinite-dimensional space which satisfies the assumptions of Herstein-Milnor’s theorem yet it is not continuous. Inoue’s illustration of GP’s example is on a simplest possible choice set: on  $\mathbb{R}^2$ . Our results above show that the infinite dimensionality in Neuefeind-Trockel’s example is essential since under the hypothesis of the Herstein-Milnor’s theorem, continuity and linear continuity postulates are equivalent in finite dimensional spaces.

the discussion. Generalization of our results that are pertaining to finite-dimensional spaces to infinite-dimension is of interest. Since the finite-dimension assumption is used crucially in some of our results, it is not a routine exercise to generalize them to infinite-dimensional spaces.<sup>58</sup> To this end, generalization of the results in the literature pertaining to locally convex sets are provided in Schoenberg (1942) and Klee (1951) and their followers. Similarly, Borwein-Lewis (1992) and Borwein-Goebel (2003) provide generalizations of the results pertaining to relative interior to infinite-dimensional spaces. These references may be of use in order to generalize the results we present in this paper to infinite-dimensions. To the best of our knowledge, there is no generalization of Rosenthal (1955) to infinite-dimensional spaces, hence generalization of our results related to arc-continuity to infinite-dimensions seems even harder.

The recent book Bosi-Campión-Candéal-Indurain (2020) contains representation theorems in abstract topological spaces, which contains finite and infinite dimensional spaces. The subtleties in the behavioral implications of continuity postulates reveal themselves. For example, Theorem 4.2 of Bosi-Zuanon (2020) is, as stated false, since it lacks the open sections assumption. The usual order  $\leq$  on  $\mathbb{R}^2$  is a counterexample. It is reflexive, transitive, non-trivial has closed graph and closed sections, and the space is Hausdorff. It does not have the open sections. And the relation is not complete. As another example, Theorem 2.2(a) of Herves-Beloso-Monteiro in the same volume is false. It lacks the non-triviality assumption. The following relation on  $[0,1]$  is a counterexample: each point is comparable itself and there are no other points that are comparable. This relation satisfies reflexivity, transitivity as well as closed and open sections (of the asymmetric part) assumptions. But it is clear that it is not representable.

There is a rich literature on the structure of the discontinuity of linearly continuous functions; see Young-Young (1910), Kershner (1943) and Ciesielski-Miller (2016). This direction has been neglected for preferences in both mathematics and in economics literature.<sup>59</sup> It is possible to study this problem not only for linear continuity but also for graph and section continuity properties. This will be also in line with the *essentiality* concept of Kim-Richter (1986): if the set of discontinuities is not very rich, then the weakening of the continuity assumption may be inessential.

Finally, in this paper we provide a deconstruction and an integration of the continuity

---

<sup>58</sup>Theorem 7 and Lemma 2 are not true for infinite-dimensional spaces; see Borwein-Lewis (1992, p. 20). The non-emptiness of the relative interior is a crucial property (for convex optimization and general equilibrium) which we also use in the construction of the proofs of our results. There are various extensions of the concept of relative interior for infinite-dimensional topological vector spaces; see Borwein-Lewis (1992) and Borwein-Goebel (2003) for a detailed reference. Moreover, Schaefer (1966, p. 38) proves that “*For any convex subset  $A$  of a t.v.s, if  $x$  is interior to  $A$  and  $y$  in the closure of  $A$ , then the open line segment joining  $x$  and  $y$  is interior to  $A$ .*” This result may be useful for the generalization of our results to infinite dimensional spaces, but a different method-of-proof is needed.

<sup>59</sup>Banerjee-Mitra (2018) provide some preliminary results on the structure of discontinuity of a utility function representing a scalarly continuous relation.

postulate based on restricted continuity. There is a rich literature taking a binary relation with some desirable properties on a space and studies the extension of it to a larger space by keeping the desirable properties of the relation; see for example Yi (1993), Duggan (1999), Kopylov (2016) and Evren-Hüseyin (2021). It is an interesting direction to investigate the relationship between the results in the current paper to those in this literature.

## 5 Concluding Remarks

We conclude this essay with two remarks, both of an epistemological nature, and ones that underscore the methodological preconceptions of the continuity postulate, and also in so far as it feeds into theorization. The first remark supplements the Leibnizian dictum that “nature does not make leaps” by noting with Arkreyd (2005) that he “also viewed it as a carrier of sets of points denser than the set of reals, sets including ideal infinitely small elements greater than zero and smaller than any positive real number.”<sup>60</sup> The basic issue hinges on *perception* and *barely* perceptible differences.<sup>61</sup> An extended quotation from Arrow (1966) goes to this.

The assumption of Continuity seems, I believe correctly, to be the harmless simplification almost inevitable in the formalization of any real-life problem. It is sometimes held that certain possible consequences, such as death, are incommensurably greater than others, such as receiving one cent. Let action  $a^1$  involve receiving one cent with no risk of life,  $a^2$  receiving nothing with no risk of life, and  $a^3$  receiving one cent with an exceedingly small probability of death. Clearly,  $a^1$  is preferred to  $a^2$ . Continuity would demand that  $a^3$  be preferred to  $a^2$  if the probability of death under  $a^3$  is sufficiently small. This may sound outrageous at first blush, but I think a little reflection will demonstrate the reasonableness of the result. The probability in question may be  $10^{-6}$  or  $10^{-10}$ , inconceivably small magnitudes. Also, if in the above example, one cent were replaced by one billion dollars, one would hardly raise the same argument, and yet to go from one cent to one billion dollars certainly involves no discontinuity, however big the difference in scale may be. “Every journey, no matter how long, begins with a single step.”<sup>62</sup>

So it all hinges on “inconceivably small magnitudes” and some “incommensurably greater than

---

<sup>60</sup>A discussion of Leibniz on the continuity postulate, and more generally the relevance of his thought to *nonstandard analysis* is beyond the scope of this essay. This is even more true of the rich philosophical issues stemming from his capacious *oeuvre*. In addition to Arkreyd (2005), we refer the reader to Kulstad (1991), Mancosu (1996), Reyes (2004), Jorgensen (2009) and their references.

<sup>61</sup>For the rich philosophical literature associated with *apperception* and these Leibnizian terms; see, for example, Kulstad (1974, 1991), Jorgensen (2009) and their references.

<sup>62</sup>Arrow continues, “Blaise Pascal, or perhaps one of his co-authors of the *Port-Royal Logic*, indeed suggested that the salvation of the soul or the avoidance of eternal damnation might be of infinitely greater value than any earthly regard; but the humble economist may be excused for regarding such choices as beyond the scope of his theories.”

others.” It leads to a requirement of asymptotic implementation of the theory, and leads Le Cam (1986) to write:

Indeed, limit theorems “as  $n$  tends to infinity” are logically devoid of content about what happens at any particular  $n$ . All they can do is suggest certain approaches whose performance must then be checked on the case at hand. Unfortunately the approximation bounds we could get were too often too crude and cumbersome to be of any practical use. Thus we have let  $n$  tend to infinity, but we would urge the reader to think of the material in approximation terms.<sup>63</sup>

Our second observation concerns our second epigraph: a reading of Koopmans’ 1957 magisterial overview of the state of economic science and in terms of a summarizing overview, one that brings out the following points into salience:

- (i) an *omission* of a pioneering paper, Eilenberg (1941), that can now be seen as inaugurating an important aspect of the modern neoclassical theory of individual choice, one relating to ‘nice’ preference relations and their representation as well as their behavioral underpinnings,
- (ii) a *masking* of the continuity postulate that led to the obscuring of the connection between axiomatic decision theory and the neoclassical theory of choice in both its deterministic and stochastic modes.

It is then a contention of this reading that both aspects have had profound consequences on the reception and direction of subsequent work, and even the most perfunctory examination of these origins allows one to obtain both a systematic and comprehensive reading of the current literature, and also to highlight connections that go into shaping it. It is also the contention of this work that the history of the problematic is relevant to a successful resolution of the problematic itself, and that the line between exposition and research may have been overplayed in mainstream decision theory: Abdellaoui-Wakker (2020) and Hartmann (2020) have furnished embarrassing instances of this disregard. A successful theory must keep in the foreground the economic phenomenon to be explained; a successful generalization of a theorem needs orientation to the direction of the generalization and it cannot be oblivious of the history of the theorem.<sup>64</sup> But beyond this, surely our investigation raises questions beyond the finite-dimensional case to

---

<sup>63</sup>Since this essay is dedicated to Tjalling Koopmans, it is worth pointing out that he was very strict in his requirement of asymptotic implementation of any result for an idealized model; see Brown-Robinson (1972) and Khan (1973). To return to the first epigraph, a referee writes, “There is now very strong evidence that this is not true at small enough scales of measurement, indeed, time itself may be discrete. This is an argument that nature is nothing but jumps, but that this happens at so small a scale that none of our human sensory apparatus can distinguish the differences, and this is the source of the intuition that the quote is correct.”

<sup>64</sup>This sentence is in response to an anonymous referee who asked for implications that stem from the claim in a sentence previous to it.

which it has been confined:<sup>65</sup> Koopman’s seminal characterizations of impatience, axiomatization of the overtaking criterion, axiomatization of the Suppes-Sen grading principle and Diamond’s impossibility theorem are all germane to our work and concern infinite-dimensional spaces.<sup>66</sup>

## A Appendix

In this Appendix, we present the proofs of the results and seven technical examples.

### A.1 Proofs of the Results

The proofs of the results we present in this paper use an important result in convex analysis due to Rockafellar (1970, p. 45) which, in Rockafellar’s words, “provides a fundamental relationship between the closure and relative interior of convex sets.” Let  $X$  be a subset of  $\mathbb{R}^n$ . Since any lower dimensional subset of  $\mathbb{R}^n$  has empty interior, it is more convenient to work with the concept of relative interior. A subset  $X$  of a (real) vector space is called *affine* if for all  $x, y \in X$  and  $\lambda \in \mathbb{R}$ ,  $\lambda x + (1 - \lambda)y \in X$ . It is clear that  $A$  is affine if and only if  $A - \{a\}$  is a subspace of  $X$  for all  $a \in A$ . The *affine hull* of  $X$ ,  $\text{aff}X$ , is the smallest affine set containing  $X$ . The *relative interior* of a subset  $X$  of  $\mathbb{R}^n$  is defined as

$$\text{ri}X = \{x \in \text{aff}X \mid \exists N_\varepsilon, \text{ an } \varepsilon \text{ neighborhood of } x, \text{ such that } N_\varepsilon \cap \text{aff}X \subseteq X\}.$$

That is, the relative interior of  $X$  is the interior of  $X$  with respect to the smallest affine subspace containing  $X$ .

**Theorem 7** (Rockafellar). *Let  $X$  be a non-empty and convex subset of  $\mathbb{R}^n$ . Then  $\text{ri}X$  is non-empty, and for all  $x \in \text{ri}X$ ,  $y \in \text{cl}X$  and all  $\lambda \in [0, 1)$ ,  $y\lambda x \in \text{ri}X$ .*

We are now ready to present the proof of Theorem 1.

*Proof of Theorem 1.* Let  $X$  be a non-empty and convex subset of  $\mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}$  a quasi-concave function. Note that  $f$  is continuous if and only if it is both upper and lower semi-continuous, i.e., for all  $\alpha \in \mathbb{R}$ , the set  $A_\alpha = \{x \in X \mid f(x) \geq \alpha\}$  is closed (in the subspace  $X$ ) and the set  $B_\alpha = \{x \in X \mid f(x) > \alpha\}$  is open (in the subspace  $X$ ). Quasi-concavity of  $f$  implies  $A_\alpha$  and  $B_\alpha$  are convex.

---

<sup>65</sup>Of course to the extent that our results rely on a Euclidean space structure and not only on a vector space one, as for example in Theorems 4 and 5.

<sup>66</sup>See Koopmans (1970, 387–428) and Koopmans (1985, Chapters 4 and 5), and his references. For extensions, see Kettering-Kochov (2020); for the overtaking criterion, see Brock’s (1970); for the extension of Diamond’s impossibility theorem and axiomatization of the Suppes-Sen grading principle, see Basu-Mitra (2003; 2007).



Pick  $\alpha \in \mathbb{R}$ . If  $A_\alpha$  is empty or a singleton, then it is closed. Otherwise, pick  $x \in \text{cl}A_\alpha$ . Since  $A_\alpha$  is convex, Theorem 7 implies that its relative interior is non-empty and for all  $y \in \text{ri}A_\alpha$  and all  $\lambda \in [0, 1)$ ,  $x\lambda y \in \text{ri}A_\alpha$ , hence for all  $\lambda^n \rightarrow 1$ ,  $f(x\lambda^n y) \in A_\alpha$  for all  $n$ . Pick  $y \in \text{ri}A_\alpha$  and let  $L$  denote the straight line in  $X$  passing through  $x$  and  $y$ . Linear continuity implies that  $f|L$  is continuous, hence  $x \in A_\alpha$ . Then  $A_\alpha$  is closed. Therefore,  $f$  is upper semi-continuous.

If  $B_\alpha$  is empty or is equal to  $X$ , then it is open. Assume  $B_\alpha$  is not open. Then there exists  $x$  in  $B_\alpha$  which lies on the boundary of  $B_\alpha$ , i.e.,  $x \in B_\alpha \cap \text{cl}B_\alpha^c \cap \text{cl}B_\alpha$ . Let  $\mathcal{H}_x$  denote the set of all supporting hyperplanes at  $x$ , and for all  $h \in \mathcal{H}_x$ , let  $H$  denote the closed half space determined by  $h$  which contains  $B_\alpha$ . First, assume there exists  $h \in \mathcal{H}_x$  such that  $C = H^c \cap X \neq \emptyset$ . Since  $H^c$  and  $X$  are convex, Theorem 7 implies that the relative interior of  $C$  is non-empty and for all  $y \in \text{ri}C$  and all  $\lambda \in [0, 1)$ ,  $x\lambda y \in \text{ri}C$ . Pick  $y \in \text{ri}C$  and let  $L$  denote the straight line in  $X$  passing through  $x$  and  $y$ . Note that  $f(z) \leq \alpha$  for all  $z \in C$ . Linear continuity implies that  $f|L$  is continuous, hence  $f(x) \leq \alpha$ , i.e.,  $x \in C$ . This furnishes us a contradiction with  $x \in B_\alpha$ . Second, assume for all  $h \in \mathcal{H}_x$ , the set  $C = H^c \cap X = \emptyset$ . This case happens only if  $x$  lies on one of the hyperplanes that determines the space  $X$ .<sup>67</sup> (Hence, when  $X$  is open (in  $\mathbb{R}^n$ ), the first case is exhaustive.) Let  $\mathcal{H}'_x$  be the set of all such hyperplanes. Recall that a hyperplane is an affine and closed subspace of  $\mathbb{R}^n$ . Moreover,  $B_\alpha \cap h$  is non-empty and convex for all  $h \in \mathcal{H}'_x$ . Furthermore, for all  $h \in \mathcal{H}'_x$ , let  $H'$  denote the half space in the affine subspace  $h$  that contains  $B_\alpha \cap h$ . It follows from  $x \in B_\alpha \cap \text{cl}B_\alpha^c \cap \text{cl}B_\alpha$  that there exists  $h \in \mathcal{H}'_x$  such that  $(H')^c \cap X \neq \emptyset$ . Then, an argument analogous to the first case furnishes a contradiction. Hence,  $B_\alpha$  is open. Therefore,  $f$  is lower semi-continuous. ■

*Proof of Proposition 1.* Let  $X$  be a non-empty, open and convex subset of  $\mathbb{R}^n$  and  $f : X \rightarrow \mathbb{R}$  a convex function on  $X$ . Pick a straight line  $L \subseteq X$  and  $x_0 \in L$ . Since  $X$  is open, there exist  $a, b \in L$ ,  $a \neq b$ , such that  $x_0 = (a + b)/2$ .

We next show that  $f$  is bounded on  $L_{ab} = \{a\lambda b \mid \lambda \in [0, 1]\}$ . By definition, for any  $x \in L_{ab}$ , there exists a unique  $\lambda_x \in [0, 1]$  such that  $x = a\lambda_x b$  (where  $\lambda_x = (b_i - x_i)/(b_i - a_i)$  for all  $i = 1, \dots, n$ ). Let  $M_0 = \max\{f(a), f(b)\}$ . Then, by the convexity of  $f$ ,

$$f(x) \leq f(a)\lambda_x f(b) \leq M_0\lambda_x M_0 \leq M_0.$$

Hence,  $f$  is bounded above on  $L_{ab}$ . For all  $y \in L_{\frac{a+b}{2}b}$ , there exists a unique  $\lambda_y \in [0.5, 1]$  such that  $(a + b)/2 = a\lambda_y y$  (where  $\lambda_y = (2y_i - a_i - b_i)/(2(y_i - a_i))$  for all  $i = 1, \dots, n$ ). Then, by the

---

<sup>67</sup>This step of the proof uses the polyhedron assumption. Example 7 below shows that the polyhedron assumption is not redundant. We thank two anonymous referees for questioning the need for the polyhedron assumption. Also see Section 3.1 for a discussion on property **C** and on a weaker property **C'**, and Footnote 77 on the role of the polyhedron assumption in another context.

convexity of  $f$ ,

$$f\left(\frac{a+b}{2}\right) \leq f(a)\lambda_y f(y).$$

Then,  $f(y) \geq f((a+b)/2)$  for  $y = (a+b)/2$ . It follows from  $\lambda_y = (2y_i - a_i - b_i)/(2(y_i - a_i))$  for all  $i = 1, \dots, n$  that for all  $y \in L_{\frac{a+b}{2}b}$ ,  $y \neq (a+b)/2$ ,

$$f(y) \geq \frac{1}{1-\lambda_y} \left( f\left(\frac{a+b}{2}\right) - \lambda_y f(a) \right) = \frac{2(y_i - a_i)}{b_i - a_i} f\left(\frac{a+b}{2}\right) - f(a) \text{ for all } i = 1, \dots, n.$$

It follows from  $(y_i - a_i)/(b_i - a_i) = (y_j - a_j)/(b_j - a_j) \in [0, 1]$  for all  $i, j = 1, \dots, n$  that for all  $y \in L_{\frac{a+b}{2}b}$ ,

$$f(y) \geq -2 \left| f\left(\frac{a+b}{2}\right) \right| - |f(a)|.$$

Hence,  $f$  is bounded below on  $L_{\frac{a+b}{2}b}$ . The proof of  $f$  is bounded below on  $L_{a\frac{a+b}{2}}$  is analogous. Hence,  $f$  is bounded on  $L_{ab}$ . Therefore, there exists  $M > 0$  such that  $|f(x)| \leq M$  for all  $x \in L_{ab}$ .

Let  $\delta = \|a - x_0\| = \|b - x_0\|$  and  $L_{ab}^\circ = L_{ab} \setminus \{a, b\}$ . Hence, for all  $x \in L_{ab}^\circ$ ,  $\|x - x_0\| < \delta$ . Pick  $x \in L_{ab}^\circ \setminus \{x_0\}$ . Let  $y \in L_{ab}^\circ$  and  $\lambda \in (0, 1)$  be such that

$$\lambda\delta \in (\|x - x_0\|, 2\|x - x_0\|) \text{ and } x = y\lambda x_0.$$

Pick  $z \in L_{ab}^\circ$  such that  $(y + z)/2 = x_0$ . Then,  $x_0 = x(1 + \lambda)^{-1}z$ . By convexity of  $f$ ,

$$f(x) \leq f(y)\lambda f(x_0) \text{ and } f(x_0) \leq f(x)(1 + \lambda)^{-1}f(z),$$

hence,  $\lambda(f(x_0) - f(z)) \leq f(x) - f(x_0) \leq \lambda(f(y) - f(x_0))$ . Then, by  $x, y, z \in L_{ab}^\circ$  and  $|f(x')| \leq M$  for all  $x' \in L_{ab}$ ,  $-2M\lambda \leq f(x) - f(x_0) \leq 2M\lambda$ . Since  $\lambda < (2\|x - x_0\|)/\delta$ ,

$$|f(x) - f(x_0)| < \frac{4M}{\delta} \|x - x_0\|.$$

Since  $L$  and  $x_0$  are arbitrarily chosen and  $|f(x_0) - f(x_0)| = \|x_0 - x_0\|$ ,  $f$  is linearly continuous.

The proof is analogous for a concave function. ■

We now show that the definition of linear continuity of a relation can be equivalently stated by using straight line segments instead of straight lines. Let  $X$  be a convex set and  $\succsim$  a binary relation on it. For any  $x, y \in X$ , the set  $L_{xy} = \{x\lambda y | \lambda \in [0, 1]\}$  denotes the *straight line segment in  $X$  connecting  $x$  and  $y$* .

**Lemma 1.** *A relation on a convex subset of a Euclidean space is linearly-continuous iff its restriction to any straight line segment in  $X$  is continuous.*

*Proof of Lemma 1.* Let  $X$  be a convex subset of  $\mathbb{R}^n$  and  $\succsim$  a binary relation on it. Assume  $\succsim$  is linearly continuous. Pick  $x, y \in X$ . Let  $L$  be the straight line containing  $x$  and  $y$ . Then  $L_{xy} \subseteq L$ . Hence the restriction of  $\succsim$  on  $L_{xy}$  is continuous.

Now assume the restriction of  $\succsim$  to any straight line segment in  $X$  is continuous. Assume there exists a straight line  $L$  in  $X$  such that  $\succsim|L$  is not continuous. First, assume  $A_{\succsim}(x) \cap L$  is not closed in  $L$  for some  $x \in X$ . Then there exists  $y \in \text{cl}A_{\succsim}(x)$  and  $y \not\succsim x$ . Since  $L$  is a straight line, therefore there exists  $z \in L$  and  $\lambda^t \rightarrow 0$  such that  $z\lambda^t y \in A_{\succsim}(x) \cap L$  for all  $t$ . This furnishes us a contradiction with the continuity of  $\succsim|L_{zy}$ . Second, assume  $A_{\succ}(x) \cap L$  is not open in  $L$  for some  $x \in X$ . Then there exists  $y \in A_{\succ}(x) \cap L$  which is not an interior point of  $A_{\succ}(x) \cap L$ . Pick  $z \in L$  such that  $z \neq y$ . Assume there exists  $z' \in L$  such that  $y = z\delta z'$  for some  $\delta \in (0, 1)$ . Then  $\succ|L_{zz'}$  does not have open sections (in  $L_{zz'}$ ). Now assume there does not exist  $z' \in L$  such that  $y = z\delta z'$  for some  $\delta \in (0, 1)$ . Then  $\succ|L_{zy}$  does not have open sections (in  $L_{zy}$ ). These contradict the continuity of  $\succsim|L_{zz'}$  and  $\succsim|L_{zy}$ , respectively. Analogous arguments yield contradictions if for some  $x \in X$ ,  $A_{\preccurlyeq}(x) \cap L$  is not closed in  $L$  or  $A_{\prec}(x) \cap L$  is not open in  $L$ . ■

*Proof of Proposition 2.* Assume  $\succsim$  is linearly continuous. It follows from Lemma 1 that the restriction of  $\succsim$  to any straight line segment is continuous. Pick  $x, y, z \in X$  and let  $L_{xy} \subseteq X$  be the straight line segment connecting  $x$  and  $y$ . Then linear continuity implies that  $\{x' \in L_{xy} | x' \succsim z\}$  is closed in  $L_{xy}$ . Let  $l_{xy}$  denote the mixture-linear function passing through  $x$  and  $y$ . Since  $l_{xy}$  is a homeomorphism between  $[0, 1]$  and  $L_{xy}$ , therefore the set  $l_{xy}^{-1}(\{x' \in L_{xy} | x' \succsim z\}) = I_{\succsim}(l_{xy}, z)$  is closed in  $[0, 1]$ . Hence  $\succsim$  is upper mixture-continuous. Similarly, it follows from linear continuity that  $\{x' \in L_{xy} | x' \succ z\}$  is open in  $L_{xy}$ . Then  $l_{xy}^{-1}(\{x' \in L_{xy} | x' \succ z\}) = I_{\succ}(l_{xy}, z)$  is open in  $[0, 1]$ . Hence,  $\succsim$  is upper strict-Archimedean which implies it is upper Archimedean. Analogous arguments imply that  $\succsim$  is lower mixture-continuous and lower Archimedean.

Now assume  $\succsim$  is mixture-continuous and Archimedean. Assume for some straight line  $L$  in  $X$  and some  $x \in X$ , the set  $\{x' \in L | x' \succsim x\}$  is not closed in  $L$ . Then there exist a sequence  $\{y^n\}$  and a point  $y$  in  $L$  such that  $y^n \succsim x$  for all  $n$ ,  $y^n \rightarrow y$  and  $y \not\succsim x$ . Pick  $k$  such that the straight line segment connecting  $y^k$  and  $y$  contains infinitely many members of the sequence. Then, since the mixture-linear function  $l_{y^k y}$  passing through  $y^k$  and  $y$  is an isomorphism between  $[0, 1]$  and the straight line segment  $L_{y^k y}$  connecting  $y^k$  and  $y$ , the set  $I_{\succsim}(l_{y^k y}, x)$  is not closed. This contradicts mixture-continuity. An analogous argument shows that for all straight line  $L$ , the lower sections of  $\succsim|L$  is closed in  $L$ .

It remains to prove that for all straight lines  $L$  in  $X$ ,  $\succ|L$  has open sections in  $L$ . It follows from of Galaabaatar-Khan-Uyanik (2019, Proposition 1) that mixture-continuity and Archimedean properties imply strict-Archimedean property. Pick a straight line  $L$  in  $X$ ,  $x \in X$  and  $y \in A_{\succ}(x) \cap L$ . Pick an  $\varepsilon$ -neighborhood  $N_{\varepsilon}(y)$  of  $y$  in the subspace  $L$ . Let  $\underline{y}, \bar{y}$  are the boundary points of  $N_{\varepsilon}(y)$  in  $L$ . Then strict-Archimedean property implies that  $I_{\succ}(l_{\underline{y}\bar{y}}, x)$  is open

in  $[0, 1]$ . Note that there exists  $\lambda_y \in [0, 1]$  such that  $y = l_{\underline{y}\bar{y}}(\lambda_y)$ . It is clear that  $\lambda_y \in I_{\succ}(l_{\underline{y}\bar{y}}, x)$ . For some small  $\delta > 0$ , the set  $V(y) = l_{\underline{y}\bar{y}}\left(I_{\succ}(l_{\underline{y}\bar{y}}, x) \cap N_{\delta}(\lambda_y)\right)$  is an open neighborhood of  $y$  in the subspace  $L$ . It is clear that  $V(y) \subseteq A_{\succ}(x) \cap L$ . Therefore,  $\succ|L$  has open upper sections in  $L$ . An analogous argument implies that the restricted relation has open lower sections. ■

*Proof of Proposition 3.* (a) The relationship between graph continuity and continuity follows their definitions. The relationship between continuity and linear continuity follows from taking the restriction of the sections on straight lines. The equivalence between linear continuity and mixture continuity & Archimedean follows from Proposition 2. Moreover, Proposition 1 in Galaabaatar-Khan-Uyanik (2019), and the discussion following it, imply the relationship among mixture continuity, Archimedean and strict-Archimedean.

(b) The relationship between Wold-continuity and weak Wold-continuity follows from their definitions.

Before moving to part (c), we show that there is no further relationship between these continuity postulates in the absence of additional assumptions. It follows from the examples in Galaabaatar-Khan-Uyanik (2019, Section 7) that there is no further relationship between mixture continuity, Archimedean and strict-Archimedean postulates. The fact that linear continuity does not imply continuity follows from Proposition 2 and Inoue (2010, Example 1). For the fact that graph continuity is stronger than continuity, see Bergstrom-Parks-Rader (1976). The following example shows that Wold continuity is stronger than weak Wold-continuity. Let  $\succsim$  be a binary relation of  $X = \mathbb{R}^2$  such that  $x \succsim y$  iff  $f(x) \geq f(y)$  for all  $x, y \in X$ , where  $f$  is defined as in Example (Genocchi-Peano). It is easy to show that  $\succsim$  is order-dense and weakly Wold-continuous. Pick  $x \in \mathbb{R}^2$  with  $x_2 = 1$  and  $x_1 \in (0, 1)$ . Then,  $f(x) \in (0, 1)$ . Hence  $(1, 1) \succ x \succ (0, 0)$ . However, for the unbroken curve connecting  $(0, 0)$  and  $(1, 1)$  and consisting of points  $\hat{x}$  such that  $\hat{x}_1^2 = \hat{x}_2$ , all points except  $(0, 0)$  is indifferent to  $(1, 1)$ . Hence,  $\succsim$  is not Wold-continuous. We finally illustrate that Wold continuity or weak Wold-continuity neither implies nor implied by any of the continuity postulates in Part (a). Define a relation  $\succsim$  on  $[0, 1]$  as  $x \succ y$  if  $x \in (0.5, 1]$ ,  $y \in [0, 0.5)$ , and  $x \sim y$  if  $x, y \in [0, 0.5)$  or  $x, y \in (0.5, 1]$  or  $x = 0.5, y \in [0, 1]$ . It is easy to check that  $\succsim$  has closed graph, but it is not weakly Wold-continuous since  $1 \succ 0$  but there is no  $z \in [0, 1]$  such that  $1 \succ z \succ 0$ .<sup>68</sup>

Now assume the ration is complete and transitive.

(c) When the relation is complete, graph continuity is equivalent to continuity, see Ward (1954, Lemma 1). We next show that mixture-continuity implies order-denseness. Let  $X$  be a convex subset of  $\mathbb{R}^n$  and  $\succsim$  a complete, transitive and mixture-continuous relation on it. Pick  $x, y \in X$

<sup>68</sup>Note that this counterexample hinges on the order-denseness property in the definition of Wold-continuity. It is not difficult to show that for order-dense preferences, continuity implies Wold-continuity and mixture continuity implies weak Wold-continuity.

such that  $x \succ y$ . Assume for all  $z \in X$ ,  $x \not\succ z$  or  $z \not\succ y$ . Then by completeness, either  $z \succcurlyeq x$  or  $y \succcurlyeq z$ . Then by transitivity,  $\{\lambda | x\lambda y \succcurlyeq x\}$  and  $\{\lambda | x\lambda y \preccurlyeq y\}$  constitute a partition of  $[0, 1]$ . This contradicts mixture-continuity of  $\succcurlyeq$  and connectedness of  $[0, 1]$ .

Now we show that continuity implies Wold-continuity. Let  $X$  be a convex subset of  $\mathbb{R}^n$  and  $\succcurlyeq$  a complete, transitive and continuous relation on it. Pick  $x, y \in X$  such that  $x \succ y$ . Then part (a) implies  $\succcurlyeq$  is mixture-continuous, hence it is order-dense. Therefore, there exists  $z$  such that  $x \succ z \succ y$ . By continuity, the sets  $R(z)$  and  $R^{-1}(z)$  are closed, hence their restrictions to any unbroken curve  $C_{xy}$  connecting  $x$  and  $y$  are closed in the subspace. By completeness  $C_{xy} \cap (R(z) \cup R^{-1}(z)) = C_{xy}$ . Then, connectedness of  $C_{xy}$  implies that  $C_{xy} \cap (R(z) \cap R^{-1}(z)) \neq \emptyset$ . Hence, there exists  $z' \in C_{xy}$  such that  $z' \sim z$ . Therefore,  $\succcurlyeq$  is Wold-continuous.

(d) The equivalence between strict-Archimedean and mixture-continuity follows from the completeness of the relation. The fact that mixture continuity implies Archimedean follows from Galaabaatar-Khan-Uyanik (2019, Proposition 1). We next show that mixture-continuity implies weak Wold-continuity. Let  $X$  be a convex subset of  $\mathbb{R}^n$  and  $\succcurlyeq$  a complete, transitive and continuous relation on it. Pick  $x, y \in X$  such that  $x \succ y$ . Mixture-continuity implies order-denseness, hence there exists  $z$  such that  $x \succ z \succ y$ . By mixture-continuity, the sets  $\{\lambda | x\lambda y \succcurlyeq z\}$  and  $\{\lambda | x\lambda y \preccurlyeq z\}$  are closed, and by completeness  $[0, 1] = \{\lambda | x\lambda y \succcurlyeq z\} \cup \{\lambda | x\lambda y \preccurlyeq z\}$ . Then, connectedness of  $[0, 1]$  implies that  $\{\lambda | x\lambda y \succcurlyeq z\} \cap \{\lambda | x\lambda y \preccurlyeq z\} \neq \emptyset$ . Hence,  $x\lambda y \sim z$  for some  $\lambda \in (0, 1)$ . Therefore  $\succcurlyeq$  is weakly Wold-continuous.

(e)<sup>69</sup> Assume there exists  $x, y, z \in X$  such that  $x \succ y$  but for all  $\lambda \in (0, 1)$ ,  $x\lambda z \not\succ y$  (the proof of the case where  $x \not\succ y\lambda z$  is analogous). By completeness,  $y \succcurlyeq x\lambda z$ . Pick  $\lambda \in (0, 1)$ . If  $y \succ x\lambda z$ , then  $x \succ y \succ x\lambda z$  and weak Wold-continuity imply there exist  $\delta \in (0, 1)$  such that  $y \sim x\delta z$ . If  $y \sim x\lambda z$ , then set  $\delta = \lambda$ . Then by transitivity,  $x \succ x\delta z$ . By weak Wold-continuity, there exists  $\gamma \in (0, 1)$  such that  $x \succ x\gamma z \succ x\delta z \sim y$ . Hence, by transitivity,  $x\gamma z \succ y$ . This furnishes us a contradiction with the assumption that for all  $\lambda' \in (0, 1)$ ,  $x\lambda' z \not\succ y$ .

The binary relation in the example we provide above for the relationship between linear continuity and continuity is complete and transitive, hence the fact that continuity is stronger than linear continuity still holds under completeness and transitivity assumptions. The fact that Archimedean property does not imply any other continuity postulates listed in the Proposition, see Ghosh-Khan-Uyanik (2020, Example 5, Appendix B).

Now consider the following example. Let  $X = \mathbb{R}_+$  and  $f(x) = \sin(1/x)$  if  $x > 0$  and  $f(0) = 1$ . Then, it is clear that  $f(x) \in [-1, 1]$  for all  $x \in X$ . Define a binary relation  $\succcurlyeq$  on  $X$  as  $x \succcurlyeq y$  is and only if  $f(x) \geq f(y)$ . Pick  $\bar{x}$  such that  $f(\bar{x}) \in (0, 1)$ . The set  $A_{\preccurlyeq}(\bar{x}) = \{x' \in X | \bar{x} \succcurlyeq x'\}$  is not closed since it contains a sequence  $x_n \rightarrow 0$  but  $0 \succ \bar{x}$ . Therefore,  $\succcurlyeq$  is not continuous.

<sup>69</sup>The proof of this claim is presented in (Ghosh, Khan, and Uyanik, 2020, Theorem 1). We present a proof here for completeness.

Moreover, since  $X$  is one dimensional,  $\succsim$  is not mixture-continuous.

We next show that  $\succsim$  satisfies Wold-continuity. Pick  $x, y \in X$  such that  $x \succ y$ . Assume without loss of generality  $x > y$ . Since  $[f(y), f(x)] \subseteq f([y, x])$ , therefore there exists  $z$  such that  $x \succ z \succ y$ . Analogously, for all  $z$  such that  $x \succ z \succ y$ , there exists a point  $w$  on the line connecting  $x$  and  $y$  such that  $w \sim z$ . Therefore,  $\succsim$  is weakly Wold-continuous. Since  $X$  is one dimensional,  $\succsim$  is Wold-continuous. Hence, Wold-continuity does not imply continuity.

We finally show that  $\succsim$  satisfies Archimedean property. Pick  $x, y, z \in X$  such that  $x \succ y$ . Assume without loss of generality  $x > y$ . Recall that  $[f(a), f(b)] \cup [f(b), f(a)] \subseteq f([a, b])$  for all  $a < b$ . Therefore, there exists  $\lambda, \delta \in (0, 1)$  such that  $f(x\lambda z) > f(y)$  and  $f(x) > f(y\delta z)$ . Hence, Archimedean and weak Wold-continuity postulates do not imply mixture-continuity. ■

We next turn to the proof of Theorem 3. Before that, we present a generalization of Rockafellar's theorem to locally convex sets, and three preliminary results on continuity and convexity of a binary relation. We start with the generalization of Rockafellar's theorem.

**Lemma 2.** *Let  $X$  be a nonempty locally convex subset of  $\mathbb{R}^n$ . Then for all  $y \in \text{cl}X$  there exists an open neighborhood  $U^y$  of  $y$  such that  $\text{ri}(X \cap U^y)$  is non-empty, and for all  $x \in \text{ri}(X \cap U^y)$  and all  $\lambda \in [0, 1)$ ,  $y\lambda x \in \text{ri}(X \cap U^y)$ .*

*Proof of Lemma 2.* Pick  $y \in \text{cl}X$  and let  $U(y)$  be the open neighborhood of  $y$  such that  $U(y) \cap X$  is convex. Note that  $y \in \text{cl}(X \cap U(y))$ . Otherwise there exists an open neighborhood  $V(y)$  of  $y$  such that  $(X \cap U(y)) \cap V(y) = \emptyset$ . However,  $U(y) \cap V(y)$  is an open neighborhood of  $y$ , hence  $(X \cap U(y)) \cap V(y) = X \cap (U(y) \cap V(y))$  furnishes us a contradiction with  $y \in \text{cl}X$ . Then,  $U(y) \cap X$  is non-empty as its closure is non-empty. The conclusion follows from Theorem 7. ■

The following two lemmata provide partial equivalences between scalar and section continuity postulates in the presence of a convexity assumption without the completeness and transitivity assumptions.

**Lemma 3.** *Let  $\succsim$  be a binary relation on a convex subset  $X$  of a Euclidean space.*

- (a) *If  $\succsim$  has locally convex upper sections, then it has closed upper sections iff it is upper mixture-continuous.*
- (b) *If  $\succsim$  has locally convex lower sections, then it has closed lower sections iff it is lower mixture-continuous.*

*Proof of Lemma 3.* Assume  $\succsim$  has closed upper sections. Pick  $x, y, z \in X$  and a sequence  $\{\lambda^t\}$  such that  $x\lambda^t y \succsim z$  for all  $t$  and  $\lambda^t \rightarrow \lambda$ . Define  $w^t = (x_1\lambda^t y_1, \dots, x_n\lambda^t y_n)$ . Then  $w^t \rightarrow w = x\lambda y$ . Since  $\succsim$  has closed upper sections,  $x\lambda y \succsim z$ . Hence  $\succsim$  is upper mixture-continuous. (Note that this direction does not require the local convexity assumption.)

In order to prove the forward direction, assume  $\succsim$  is upper mixture-continuous but it does not have closed upper sections. Then there exists  $x_0 \in X$  such that  $A_{\succsim}(x_0)$  is not closed, hence there exists  $x \in \text{cl}A_{\succsim}(x_0)$  such that  $x \not\succsim x_0$ . Since its closure is nonempty,  $A_{\succsim}(x_0)$  is non-empty. Then local convexity property and Lemma 2 implies that there exists  $y \succsim x_0$  such that  $x\lambda y \succsim x_0$  for all  $\lambda \in [0, 1)$ . Since  $x1y = x \not\succsim x_0$ , this furnishes us a contradiction with upper mixture-continuity. Therefore,  $\succsim$  has closed upper sections.

The proof of the equivalence between the closed lower sections and lower mixture-continuity properties is analogous. ■

**Lemma 4.** *Let  $\succsim$  be a binary relation on a set  $X$  with property C.<sup>70</sup>*

- (a) *If  $\succ$  has locally convex upper sections, then it has open upper sections iff  $\succsim$  is upper strict-Archimedean.*
- (b) *If  $\succ$  has locally convex lower sections, then it has open lower sections iff  $\succsim$  is lower strict-Archimedean.*

The construction in the proof of Lemma 4 is similar to that of Theorem 1.

*Proof of Lemma 4.* We provide a detailed proof of part (a), the proof of part (b) is analogous. Assume  $\succ$  has open upper sections. Pick  $x, y, z \in X$  and  $\lambda$  such that  $x\lambda y \succ z$ . Then open upper sections property implies that there exists  $\varepsilon > 0$  such that for all  $w$  in the  $\varepsilon$  neighborhood  $N_\varepsilon$  of  $x\lambda y$ ,  $w \succ z$ . Since there exists  $\epsilon > 0$  such that  $x\delta y \in N_\varepsilon$  for all  $\delta \in (\lambda - \epsilon, \lambda + \epsilon) \cap [0, 1]$ , the relation  $\succsim$  satisfies the upper strict-Archimedean property. (Note that this direction does not require the local convexity assumption.)

In order to prove the backward direction, assume  $\succsim$  is upper strict-Archimedean but it does not have open upper sections. Then there exists  $x_0 \in X$  such that  $A_{\succ}(x_0)$  is not open (in  $X$ ), hence there exists  $x \succ x_0$  such that  $x$  is not an interior point of  $A_{\succ}(x_0)$ . It follows from Lemma 2 that  $x$  has a neighborhood  $V \subseteq X$  which is open in  $X$  such that  $V \cap A_{\succ}(x_0)$  is convex and has non-empty relative interior. Let  $\mathcal{H}_x$  denote the set of all supporting hyperplanes of  $V \cap A_{\succ}(x_0)$  at  $x$ , and for all  $h \in \mathcal{H}_x$ , let  $H_h$  denote the closed half space determined by  $h$  which contains  $V \cap A_{\succ}(x_0)$ .

First, assume there exists  $h \in \mathcal{H}_x$  such that  $C = H_h^c \cap V \neq \emptyset$ . Since  $H_h^c \cap V$  is non-empty and convex, Theorem 7 implies that  $x\delta w \not\succ x_0$  for all  $\delta \in [0, 1)$  and all  $w$  in the relative interior of  $H_h^c \cap V$ . Since  $x \succ x_0$ , therefore  $I_{\succ}(l_{xw}, x_0) = \{1\}$ , which is not open in  $[0, 1]$ . This furnishes us a contradiction with the upper strict-Archimedean property.

Second, assume for all  $h \in \mathcal{H}_x$ , the set  $H_h^c \cap V = \emptyset$ . This case happens only if  $X$  is a polyhedron and  $x$  lies in one of the hyperplanes that determines the set  $X$ . (Hence, when  $X$

---

<sup>70</sup>This proposition is false for an arbitrary convex subset of a Euclidean space; see Example 1 in Appendix A.2.

is open, the first case is exhaustive.) Let  $\mathcal{H}'_x$  be the set of all such hyperplanes. It is easy to show that there exists  $h \in \mathcal{H}'_x$  and  $y \in h$  such that for all  $\delta \in [0, 1]$ ,  $x\delta y \notin V \cap A_{\succ}(x_0)$ . (Otherwise, there exists an open ball around  $x$  that is contained in  $V \cap A_{\succ}(x_0)$ .) This furnishes us a contradiction with the upper strict-Archimedean property. Therefore,  $\succ$  has open upper sections.  $\blacksquare$

**Lemma 5.** *Let  $\succsim$  be a reflexive, mixture-continuous and Archimedean binary relation on a convex subset  $X$  of  $\mathbb{R}^n$  which demonstrates convex indifference and has a transitive symmetric part. Then,  $\succsim$  and its asymmetric part  $\succ$  have convex sections.*

*Proof of Lemma 5.* The convexity of the sections of  $\succsim$  follows from Lemma 2 of Galaabaatar-Khan-Uyanik (2019). The following claim implies that in order to show  $\succ$  is convex, it is enough to prove that  $I_{\succ}(l_{xy}, z)$  and  $I_{\prec}(l_{xy}, z)$  are convex for all  $x, y, z \in X$ .

*Claim 1.*  $\succ$  is convex  $\iff I_{\succ}(l_{xy}, z)$  and  $I_{\prec}(l_{xy}, z)$  are convex for all  $x, y, z \in X$ .

Pick  $x, y, z \in X$ . We next show that  $I_{\succ}(l_{xy}, z)$  is convex. Note that Theorem 2 and Proposition 1 of Galaabaatar-Khan-Uyanik (2019) imply that  $\succsim$  is semi-transitive and strict-Archimedean. If  $I_{\succ}(l_{xy}, z)$  is empty, then it is convex. If  $I_{\sim}(l_{xy}, z)$  is empty, then  $I_{\succ}(l_{xy}, z) = I_{\succsim}(l_{xy}, z)$ , hence it is convex. Now assume  $I_{\sim}(l_{xy}, z)$  is non-empty. Then  $I_{\succsim}(l_{xy}, z)$  is non-empty. It follows from the fact that  $\succsim$  demonstrates convex indifference and is mixture-continuous that both  $I_{\sim}(l_{xy}, z)$  and  $I_{\succsim}(l_{xy}, z)$  are closed intervals in  $[0, 1]$ .

Now assume  $I_{\succ}(l_{xy}, z)$  is not convex. The convexity and strict-Archimedean property of  $\succsim$  imply that there exist two nonempty, disjoint, open intervals  $I_1, I_2$  such that  $I_{\succ}(l_{xy}, z) = I_1 \cup I_2$  where the intervals of type  $[0, \alpha)$  or  $(\alpha, 1]$  are allowed for  $\alpha \in (0, 1)$ . Note that  $I_{\succsim}(l_{xy}, z) = I_1 \cup I_{\sim}(l_{xy}, z) \cup I_2$  and  $I_{\sim}(l_{xy}, z) = [\lambda_1, \lambda_2]$  for some  $\lambda_1 \leq \lambda_2$ . Therefore,  $I_1 \cup I_2 = [0, \lambda_1) \cup (\lambda_2, 1]$ . Without loss of generality assume  $I_1 = [0, \lambda_1)$ .

Assume there exists  $\lambda \in I_1$  such that  $I_2 \not\subseteq I_{\prec}(l_{xy}, x\lambda y)$ , that is, there exist  $\lambda \in I_1, \delta \in I_2$  (hence,  $x\lambda y \succ z, x\delta y \succ z$ ) such that  $x\lambda y \not\succ x\delta y$ . It follows from  $x\lambda y \succ z$  and semi-transitivity of  $\succsim$  that  $[\lambda_1, \lambda_2] \subseteq I_{\prec}(l_{xy}, x\lambda y)$ . Then it follows from  $\succ$  has open sections that there exist  $\delta_1 \in I_1$  and  $\delta_2 \in I_2$  such that  $\delta_1, \delta_2 \in I_{\prec}(l_{xy}, x\lambda y)$ . Reflexivity of  $\succsim$  implies that  $\lambda \in I_{\sim}(l_{xy}, x\lambda y) \cap I_1$ . Let  $\gamma_1$  be the smallest scalar  $\lambda'$  such that  $x\lambda'y \sim x\lambda y$ . It is clear that  $\gamma_1 \leq \lambda$ , hence  $\gamma_1 \in I_1$ . We next show that there exists  $\gamma_2 \in I_2$  such that  $x\gamma_2 y \sim x\lambda y$ . Assume such  $\gamma_2$  does not exist. Then  $I_{\preccurlyeq}(l_{xy}, x\lambda y) \cap I_2 = I_{\prec}(l_{xy}, x\lambda y) \cap I_2$ . Convexity of the set on the left hand side of the equation, the openness and the non-emptiness of the set on the right hand side and  $I_2 \not\subseteq I_{\prec}(l_{xy}, x\lambda y)$  imply that those sets are equal to  $(\lambda_2, t)$  for some  $t < 1$ . Since  $I_{\preccurlyeq}(l_{xy}, x\lambda y)$  is convex and  $I_2 = (\lambda_2, 1]$ , therefore  $I_{\preccurlyeq}(l_{xy}, x\lambda y) = [\gamma_1, t)$ . This furnishes us a contradiction with mixture-continuity. Hence there exists  $\gamma_2 \in I_2$  such that  $x\gamma_2 y \sim x\lambda y$ . Then  $\gamma_1, \gamma_2 \in I_{\succsim}(l_{xy}, x\lambda y)$ . Since  $\succsim$  is convex, therefore  $[\gamma_1, \gamma_2] \subseteq I_{\succsim}(l_{xy}, x\lambda y)$ . This furnishes us a contradiction with  $[\lambda_1, \lambda_2] \subseteq I_{\prec}(l_{xy}, x\lambda y)$ .



Now assume  $I_2 \subseteq I_{\prec}(l_{xy}, x\lambda'y)$  for all  $\lambda' \in I_1$ . Pick  $\lambda \in I_2$ . An argument analogous to the one presented in the previous paragraph yields a contradiction.

Therefore,  $I_{\succ}(l_{xy}, z)$  is convex. The proof of the convexity of  $I_{\prec}(l_{xy}, z)$  is analogous. Then, Claim 1 implies that  $\succ$  is convex.

*Proof of Claim 1.* Assume  $\succ$  is convex. Pick  $x, y, z \in X$  and  $\lambda, \delta \in I_{\succ}(l_{xy}, z)$  and  $\beta \in [0, 1]$ . Define  $w_{\lambda} = x\lambda y$  and  $w_{\delta} = x\delta y$ . By construction,  $w_{\lambda} \succ z$  and  $w_{\delta} \succ z$ . It follows from  $\succ$  is convex that  $w_{\lambda}\beta w_{\delta} \succ z$ . A simple algebra implies  $w_{\lambda}\beta w_{\delta} = x(\beta\lambda + (1 - \beta)\delta)y$ . Therefore,  $\beta\lambda + (1 - \beta)\delta \in I_{\succ}(l_{xy}, z)$ . Hence,  $I_{\succ}(l_{xy}, z)$  is convex. An analogous argument proves the convexity of  $I_{\prec}(l_{xy}, z)$ . Now assume  $I_{\succ}(l_{xy}, z)$  is convex for all  $x, y, z \in X$ . Pick  $x, y, z \in X$  and  $\lambda \in [0, 1]$  such that  $x \succ z$  and  $y \succ z$ . Then  $0, 1 \in I_{\succ}(l_{xy}, z)$ . Since  $I_{\succ}(l_{xy}, z)$  is convex, therefore  $I_{\succ}(l_{xy}, z) = [0, 1]$ . Hence,  $x\lambda y \succ z$ . Therefore,  $\succ$  has convex upper sections. An analogous argument implies the convexity of lower sections of  $\succ$ . ■

The proof of Lemma 5 is complete. ■

We are now ready to prove Theorem 3.

*Proof of Theorem 3.* For each part, the backward direction is clear. It remains to prove that linear continuity implies continuity.

(a) Proposition 2 implies that linear continuity of  $\succcurlyeq$  is equivalent to its mixture-continuity and its Archimedean. It follows from the local convexity assumption and Lemmata 3 and 4 that the upper sections of  $\succcurlyeq$  are closed and of  $\succ$  are open. Then completeness of  $\succcurlyeq$  implies the upper sections of  $\succcurlyeq$  are closed and of  $\succ$  are open. Therefore,  $\succcurlyeq$  is continuous.

(b) The proof follows from Proposition 2 and Lemmata 3 and 4.

(c) Lemma 5 implies that the sections of  $\succcurlyeq$  and of  $\succ$  are convex. Then part (b) above completes the proof.

The proof of Theorem 3 is complete. ■

We now define additional -convexity postulates and provides a result on their relationship to those we defined in Section 3. The relation  $\succcurlyeq$  is *star-convex* if for all distinct  $x, y \in X$  and all  $\lambda \in (0, 1)$ ,  $x \succcurlyeq y$  implies  $x\lambda y \succcurlyeq y$ ; and *strictly star-convex* if for all distinct  $x, y \in X$  and all  $\lambda \in (0, 1)$ ,  $x \sim y$  implies  $x\lambda y \succ y$ .

**Lemma 6.** *Let  $\succcurlyeq$  be a complete and transitive relation on a convex subset  $X$  of a vector space. Then the following are equivalent: (C1)  $\succcurlyeq$  is convex, (C2)  $\succ$  is convex, (C3)  $\succcurlyeq$  is star-convex.*

If  $\succsim$  is lower Archimedean, then star-convexity of  $\succ$  implies the convexity properties (C1)–(C3). Moreover, if  $\succsim$  is weakly Wold-continuous, then its strict star-convexity implies  $\succ$  is star-convex, hence the convexity properties (C1)–(C3).<sup>71</sup>

*Proof of Lemma 6.* (C1)  $\Rightarrow$  (C2) Pick  $x \in X$  and  $y, z \in A_{\succ}(x)$  and  $\lambda \in (0, 1)$ . Assume without loss of generality that  $z \succsim y$ . Since  $A_{\succ}(y)$  is convex, therefore  $y\lambda z \succsim y$ . Therefore, transitivity of  $\succsim$  and  $y \succ x$  imply<sup>72</sup> that  $y\lambda z \succ x$ .

(C2)  $\Rightarrow$  (C3) Assume there exists  $x, y \in X$  and  $\lambda \in (0, 1)$  such that  $y \succsim x$  and  $y\lambda x \prec x$ . Transitivity of  $\succsim$  implies  $y\lambda x \prec y$ . Then convexity of  $\succ$  implies  $y\lambda x \succ y\lambda x$ . This furnishes us a contradiction.

(C3)  $\Rightarrow$  (C1) Pick  $x \in X$  and  $y, z \in A_{\succ}(x)$  and  $\lambda \in (0, 1)$ . Assume without loss of generality that  $z \succsim y$ . Then star-convexity of  $\succsim$  implies  $y\lambda z \succsim y$ . It follows from transitivity of  $\succsim$  that  $y\lambda z \succ x$ .

Now assume  $\succsim$  is lower Archimedean and  $\succ$  is star-convex. Assume there exist  $x, y \in X$  and  $\lambda \in (0, 1)$  such that  $y \succsim x$  and  $x \succ x\lambda y$ . Then lower Archimedean property and completeness of  $\succsim$  imply that there exists  $\delta \in (0, 1)$  such that  $x \succ (x\lambda y)\delta y = x(\lambda\delta)y$ . Then star-convexity of  $\succ$  implies that  $x\lambda y \succ x(\lambda\delta)y$ . It follows from transitivity and  $y \succsim x \succ x\lambda y$  that  $y \succ x\lambda y$ . Then star-convexity of  $\succ$  implies that  $x(\lambda\delta)y \succ x\lambda y$ . This furnishes us a contradiction. Therefore  $\succsim$  is star-convex.<sup>73</sup>

Lastly assume  $\succsim$  is weakly Wold-continuous. Now let  $\succsim$  be strictly star-convex. Assume there exist  $x, y \in X$  and  $\lambda \in (0, 1)$  such that  $y \succ x$  and  $x\lambda y \preccurlyeq x$ . Let  $z = x\lambda y$ . If  $z \prec x$ , then weak Wold-continuity implies that there exists  $\delta \in (0, 1)$  with  $x \succ x\delta z \succ z$ . If  $z \sim x$ , then strict star-convexity implies that for all  $\delta \in (0, 1)$ ,  $x\delta z \succ z$ . Pick  $\delta \in (0, 1)$  such that  $x\delta z \succ z$ . Then, it follows from weak Wold-continuity and  $y \succ x\delta z \succ z$  that there exists  $\gamma \in (0, 1)$  such that  $y\gamma z \sim x\delta z$ . Since  $z = (y\gamma z)\mu(x\delta z)$  for some  $\mu \in (0, 1)$ , strict star-convexity implies  $z \succ x\delta z$ . Then,  $x\delta z \succ z$  and transitivity of  $\succsim$  yield a contradiction. ■

*Proof of Theorem 2.* Proposition 3 provides partial relationships among the continuity postulates without the convexity or monotonicity postulates. We first show that the remaining relationships hold under convexity of the relation, and then we replace the convexity assumption with weak monotonicity.

<sup>71</sup>Some of the relationships can be found elsewhere; see for example Debreu (1959). We provide the proofs of them also for completeness.

<sup>72</sup>See Khan-Uyanik (2019, Proposition 2, Theorems 5 and 6 and Figure 3) for the relationship among different transitivity postulates.

<sup>73</sup>Note that, the lower Archimedean property is not redundant in this result. In order to see this, consider the following example:  $X = [0, 1]$ ,  $\succsim$  is a reflexive preference relation on  $X$  such that for all  $x, y \neq 0.5$ ,  $x \sim y$  and  $x \succ 0.5$ . It is clear that  $\succsim$  is complete and transitive, not lower Archimedean and  $\succ$  is star-convex. However,  $\succsim$  is not star-convex since  $0 \succsim 1$  and  $0\lambda 1 \prec 1$  for  $\lambda = 0.5$ .

In the first part of the proof, let  $\succsim$  be a convex, complete and transitive binary relation on a convex subset of  $\mathbb{R}^n$  which satisfies property **C**.

First, by Lemma 6, convexity of  $\succsim$  implies convexity of  $\succ$ . Therefore, by Theorem 3, linear continuity implies continuity.

Second, we show that Archimedean property implies mixture-continuity.<sup>74</sup> To this end pick  $x, y, z \in X$ . Assume there exists a convergent sequence  $\lambda^n$  in  $[0, 1]$  such that  $x\lambda^n y \succsim z$  for all  $n$  and  $z \succ x\lambda y$ . Since  $A_{\succsim}(z)$  is convex, if  $\lambda^k \leq \lambda \leq \lambda^m$  for some  $k, m$ , then  $x\lambda y \succsim z$ . Therefore, either  $\lambda^n > \lambda$  for all  $n$  or  $\lambda^n < \lambda$  for all  $n$ . Assume wlog  $\lambda^n > \lambda$  for all  $n$ . Then convexity of  $\succsim$  implies that  $x\lambda' y \succsim z$  for all  $\lambda' \in (\lambda, \lambda^1]$ . It follows from Archimedean property and  $z \succ x\lambda y$  that there exists  $\delta \in (0, 1)$  such that  $z \succ (x\lambda y)\delta(x\lambda^1 y)$ . Since  $(x\lambda y)\delta(x\lambda^1 y) = x(\delta\lambda + (1-\delta)\lambda^1)y$  and  $\delta\lambda + (1-\delta)\lambda^1 \in (\lambda, \lambda^1]$ , this furnishes us a contradiction. Therefore,  $\succsim$  is upper mixture-continuous.

Now assume there exists a convergent sequence  $\lambda^n$  in  $[0, 1]$  such that  $z \succ x\lambda^n y$  for all  $n$  and  $x\lambda y \succ z$ . It follows from Archimedean property that  $x\delta(x\lambda y) \succ z$  and  $(x\lambda y)\gamma y \succ z$  for some  $\delta, \gamma \in (0, 1)$ . Note that  $x\delta(x\lambda y) = x(\lambda + \delta(1-\lambda))$  and  $(x\lambda y)\gamma y = x(\lambda\gamma)y$ , and  $\lambda\gamma \leq \lambda \leq \lambda + \delta(1-\lambda)$ . Note that  $\lambda \in (0, 1)$  implies both inequalities are strict,  $\lambda = 1$  implies  $\lambda\gamma < \lambda$  and  $\lambda = 0$  implies that  $\lambda < \lambda + \delta(1-\lambda)$ . These furnish us a contradiction with the convexity assumption and  $x\lambda^n y \succsim z$  for all  $n$ . Therefore,  $\succsim$  is lower mixture-continuous.

Third, we prove that weak Wold-continuity implies mixture-continuity. Assume there exist  $x, y, z \in X$  and a convergent sequence  $\lambda^k \rightarrow \lambda$  such that  $x\lambda^k y \succsim z$  for all  $k$  and  $z \succ x\lambda y$ . If  $\lambda^k \leq \lambda \leq \lambda^m$  for some  $k, m$ , then convexity of  $\succsim$  implies that  $x\lambda y \succsim z$ . Therefore, either  $\lambda^n > \lambda$  for all  $n$  or  $\lambda^n < \lambda$  for all  $n$ . Assume without loss of generality that  $\lambda^n > \lambda$  for all  $n$ . Then convexity of  $\succsim$  implies that  $x\lambda' y \succsim z$  for all  $\lambda' \in (\lambda, \lambda^1]$ . We next show that there exists  $\bar{\lambda} \in (\lambda, \lambda^1]$  such that  $z \sim x\bar{\lambda} y$ . If  $x\lambda' y \succ z$  for some  $\lambda' \in (\lambda, \lambda^1]$ , then  $z \succ x\lambda y$  and weak Wold-continuity imply that there exists  $\delta \in (0, 1)$  such that  $z \sim (x\lambda' y)\delta(x\lambda y) = x(\delta\lambda' + (1-\delta)\lambda)y$ . Otherwise,  $x\lambda' y \sim z$  for all  $\lambda' \in (\lambda, \lambda^1]$ . Now, it follows from transitivity and weak Wold-continuity of  $\succsim$ , and  $x\bar{\lambda} y \sim z \succ x\lambda y$  that there exists  $\hat{\lambda} \in (\lambda, \bar{\lambda})$  such that  $x\bar{\lambda} y \succ x\hat{\lambda} y \succ x\lambda y$ . The transitivity of  $\succsim$  and  $z \sim x\bar{\lambda} y \succ x\hat{\lambda} y$  imply that  $z \succ x\hat{\lambda} y$ . Then,  $\hat{\lambda} \in (\lambda, \lambda^1]$  contradicts convexity of  $\succsim$ . Therefore,  $\succsim$  is upper mixture-continuous.

Now assume there exist  $x, y, z \in X$  and a convergent sequence  $\lambda^k \rightarrow \lambda$  such that  $z \succ x\lambda^k y$  for all  $k$  and  $x\lambda y \succ z$ . Assume without loss of generality that there exists a subsequence  $\lambda^{k_i}$  of  $\lambda^k$  such that  $\lambda^{k_i} > \lambda$  for all  $i = 1, 2, \dots$ . We next show that there exists  $\bar{\lambda} \in (\lambda, \lambda^{k_1}]$  such that  $z \sim x\bar{\lambda} y$ . If  $z \succ x\lambda^{k_1} y$ , then  $x\lambda y \succ z$  and weak Wold-continuity imply that there exists

<sup>74</sup>Note that the second and third parts of the proof, that is the proof of mixture continuity follows from each of Archimedean and weak Wold-continuity properties, are independent of the dimension of the space. We return to this in the subsequent results.

$\delta \in (0, 1)$  such that  $z \sim (x\lambda^{k_1}y)\delta(x\lambda y) = x(\delta\lambda^{k_1} + (1 - \delta)\lambda)y$ . Otherwise,  $z \sim x\lambda^{k_1}y$ . Then, it follows from transitivity and weak Wold-continuity of  $\succsim$ , and  $x\lambda y \succ z \sim x\bar{\lambda}y$  that there exists  $\hat{\lambda} \in (\lambda, \bar{\lambda})$  such that  $x\lambda y \succ x\hat{\lambda}y \succ x\bar{\lambda}y$ . The transitivity of  $\succsim$  and  $z \sim x\bar{\lambda}y \prec x\hat{\lambda}y$  imply that  $z \prec x\hat{\lambda}y$ . Moreover, convexity of  $\succsim$  implies that for all  $\lambda' \in [\lambda, \hat{\lambda}]$ ,  $x\lambda'y \succsim x\hat{\lambda}y$ . Since  $\lambda^{k_i} \rightarrow \lambda$ , there exists  $j$  such that  $\lambda^{k_j} \in (\lambda, \hat{\lambda})$ . Then,  $z \succsim x\lambda^{k_j}y \succ x\hat{\lambda}y \succ z$  furnishes us a contradiction. Therefore,  $\succsim$  is lower mixture-continuous.

In the second part of the proof, assume  $\succsim$  is a weakly monotonic, complete and transitive binary relation on a convex subset  $X$  of  $\mathbb{R}_+^n$  which satisfies property **B**. It remains to show that each of Archimedean and weak Wold-continuity postulates implies the continuity postulate. Towards this end, assume  $\succsim$  is Archimedean.

Assume there exists  $x \in X$  such that  $A_{\preccurlyeq}(x)$  is not closed, i.e., there exists  $y^k \rightarrow y$  such that  $y^k \in A_{\preccurlyeq}(x)$  for all  $k$  and  $y \succ x$ . By property **B**, there exists  $b \in X$  such that  $x, y \geq b$ . The Archimedean property implies that there exists  $\lambda \in (0, 1)$  such that  $y\lambda b \succ x$ . Define  $z = y\lambda b$ . Since  $y > b$ , therefore  $z > b$ . Define  $\varepsilon = \min_{z_i \neq b_i} (y_i - z_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' > z$ . Then weak monotonicity implies that  $y' \succ z$ . Since  $z \succ x$ , it follows from transitivity that  $y' \succ x$  for all  $y'$  in the  $\varepsilon$  neighborhood of  $y$ . This furnishes us a contradiction with  $y^k \rightarrow y$ .

Now assume there exists  $x \in X$  such that  $A_{\succ}(x)$  is not closed, i.e., there exists  $y^k \rightarrow y$  such that  $y^k \in A_{\succ}(x)$  for all  $k$  and  $x \succ y$ . By property **B**, there exists  $x' \in X$  such that  $x' \geq x, y$ . Then Archimedean property implies that there exists  $\lambda \in (0, 1)$  such that  $x \succ x'\lambda y$ . Note that  $x' > y$ . Define  $z = x'\lambda y$ . Since  $x' > y$ , therefore  $z > y$ . Define  $\varepsilon = \min_i (z_i - y_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' < z$ . Then weak monotonicity implies that  $z \succ y'$ . Since  $x \succ z$ , it follows from transitivity that  $x \succ y'$ . This furnishes us a contradiction with  $y^k \rightarrow y$ . Therefore,  $\succsim$  is continuous.

Finally, we show that continuity follows from weak Wold-continuity. To this end, assume  $\succsim$  is weakly Wold-continuous. Assume there exists  $x \in X$  such that  $A_{\preccurlyeq}(x)$  is not closed, i.e., there exists  $y^k \rightarrow y$  such that  $y^k \in A_{\preccurlyeq}(x)$  for all  $k$  and  $y \succ x$ . By property **B**, there exists  $b \in X$  such that  $x, y \geq b$ . Order denseness of  $\succsim$  implies that there exists  $z'$  such that  $y \succ z' \succ x$ . It follows from  $\succsim$  is transitive and weakly monotonic, and  $y, z' > b$  that  $y \succ z' \succ b$ . Then, weak Wold-continuity implies that  $z' \sim y\lambda b$  for some  $\lambda \in (0, 1)$ . Define  $z = y\lambda b$ . Since  $y > b$ , therefore  $z > b$ . Define  $\varepsilon = \min_{z_i \neq b_i} (y_i - z_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' > z$ . Then weak monotonicity implies that  $y' \succ z$ . Since  $z \sim z' \succ x$ , it follows from transitivity that  $y' \succ x$  for all  $y'$  in the  $\varepsilon$  neighborhood of  $y$ . This furnishes us a contradiction with  $y^k \rightarrow y$ .

Now assume there exists  $x \in X$  such that  $A_{\succ}(x)$  is not closed, i.e., there exists  $y^k \rightarrow y$  such that  $y^k \in A_{\succ}(x)$  for all  $k$  and  $x \succ y$ . Then order denseness of  $\succsim$  implies that there exists  $z'$  such that  $x \succ z' \succ y$ . By property **B**, there exists  $x' \in X$  such that  $x' \geq x, y$ . Therefore, weak monotonicity and transitivity properties imply that  $x' \succ z' \succ y$ . Then, weak Wold-continuity

implies that  $z' \sim x'\lambda y$  for some  $\lambda \in (0, 1)$ . Define  $z = x'\lambda y$ . Since  $x' > y$ , therefore  $z > y$ . Define  $\varepsilon = \min_i (z_i - y_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' < z$ . Then weak monotonicity implies that  $z \succ y'$ . Since  $x \succ z$ , it follows from transitivity that  $x \succ y'$ . This furnishes us a contradiction with  $y^k \rightarrow y$  and  $y^k \in A_{\succ}(x)$  for all  $k$ . Therefore,  $\succ$  is continuous. ■

*Proof of Theorem 4. (a)* Assume  $\succ$  is upper mixture-continuous. Pick  $x, y, z \in X$  and a sequence  $\lambda^n \in [0, 1]$  such that  $\lambda^n \rightarrow \lambda$  and  $z \succ x\lambda^n y$  for all  $n$ . Since  $X$  is a convex cone,  $z + y\lambda^n x$  and  $x + y$  are in  $X$ . Note that  $z + x + y - x\lambda^n y = z + y\lambda^n x = (z + y)\lambda^n(z + x)$  for all  $n$ . Additivity implies that  $(z + y)\lambda^n(z + x) = z + x + y - x\lambda^n y \succ x + y$ . Since  $\succ$  is upper mixture-continuous,  $(z + y)\lambda(z + x) = z + x + y - x\lambda y \succ x + y$ . Additivity implies that  $z + x + y \succ x + y + x\lambda y$ . Since  $z, x\lambda y, x + y \in X$ , therefore additivity implies that  $z \succ x\lambda y$ . Hence  $\succ$  is lower mixture-continuous. The proof of the converse statement is analogous. Then, by definition, mixture continuity follows from any of the upper and lower mixture-continuity.

*(b)* Assume  $\succ$  is upper strict-Archimedean. Pick  $x, y, z \in X$  and  $\lambda \in (0, 1)$  such that<sup>75</sup>  $z \succ x\lambda y$ . Additivity implies that  $z + x + y - x\lambda y = z + y\lambda x \succ x + y$ . Note that  $z + y\mu x = (z + y)\mu(z + x)$  for all  $\mu \in [0, 1]$ . Upper strict-Archimedean property implies that there exists  $\underline{\lambda} < \lambda < \bar{\lambda}$  such that  $z + y\delta x \succ x + y$  for all  $\delta \in (\underline{\lambda}, \bar{\lambda})$ . Then additivity and  $z, x\delta y \in X$  imply that  $z \succ x\delta y$  for all  $\delta \in (\underline{\lambda}, \bar{\lambda})$ . Hence  $\succ$  is lower strict-Archimedean. The proof of the converse statement is analogous. Then, by definition, strict-Archimedean property follows from any of the upper and lower strict-Archimedean property. ■

*Proof of Theorem 5.* Proposition 3 shows that mixture continuity is equivalent to the strict Archimedean postulate, and implies Archimedean and weak Wold-continuity postulates. The proof of mixture-continuity follows from each of Archimedean and weak Wold-continuity postulates is identical to that provided in the proof of Theorem 2 since the dimension of the space is not used in the proof; see Footnote 74 for details. ■

We next turn to the proof of Theorem 6. Before that we present an important result due to Rosenthal (1955, Theorem 3) which is crucial for the proof of his remarkable result we provided in the Introduction, and a lemma on the continuity of binary relations. We restate the theorem of Rosenthal with the notation of our paper.

**Theorem 8** (Rosenthal). *Any bounded infinite subset of set  $X$  in  $\mathbb{R}^n$  with property **C** contains an infinite subset through which a smooth curve can be laid.*<sup>76</sup>

<sup>75</sup>When  $\lambda = 0, 1$ , then the proof is analogous.

<sup>76</sup>Rosenthal states the theorem for  $X = \mathbb{R}^n$ , but it is easy to observe that his theorem is true for any subset  $X$  of  $\mathbb{R}^n$  with property **C**.

The following lemma provides a characterization of strong Archimedean property under strong mixture-continuity.

**Lemma 7.** *Let  $\succsim$  be a strongly mixture-continuous binary relation on a convex subset of  $\mathbb{R}^n$ . Then  $\succsim$  is strong Archimedean iff it is strongly strict-Archimedean.*

*Proof of Lemma 7.* Assume  $\succsim$  is strongly mixture-continuous and strongly Archimedean. Pick  $x, y, z \in X$  and  $m_{xy} \in \mathcal{M}$ . If  $I_{\succ}(m_{xy}, z)$  is empty, then it is open. Otherwise, pick  $\lambda \in I_{\succ}(m_{xy}, z)$ . It follows from strong mixture-continuity and  $\lambda \notin I_{\preccurlyeq}(m_{xy}, z)$  that there exists  $t > 0$  such that  $N_t(\lambda) = \{\beta \mid |\beta - \lambda| < t\}$  is contained in the complement of  $I_{\preccurlyeq}(m_{xy}, z)$ .

Assume there exists  $\beta \in N_t(\lambda) \cap I_{\bowtie}(m_{xy}, z)$ . It follows from strong mixture-continuity that  $I_{\bowtie}(m_{xy}, z)$  is open. Therefore, as an open set in  $[0, 1]$ ,  $I_{\bowtie}(m_{xy}, z)$  is union of at most countably many mutually disjoint open intervals such that the intervals of type  $[0, \alpha)$  or  $(\alpha, 1]$  are allowed for any  $\alpha \in (0, 1)$ . Among these open intervals, there exists open interval  $I$  such that  $\beta \in I$ . Assume  $\beta > \lambda$ . Set  $\delta = \inf I$ . Then  $\delta \in I_{\succ}(m_{xy}, z)$ . Note that there exists  $m_{m_{xy}(\delta)m_{xy}(\beta)} \in \mathcal{M}$  such that  $m_{m_{xy}(\delta)m_{xy}(\beta)}((0, 1)) = m_{xy}((\delta, \beta))$ . Since  $(\delta, \beta) \subseteq I$ , therefore  $(\delta, \beta) \subseteq I_{\bowtie}(m_{xy}, z)$ . This furnishes us a contradiction with  $\succsim$  is strongly Archimedean. Now assume  $\beta < \lambda$ . Then setting  $\delta = \sup I$  and repeating the step above by replacing  $(\delta, \beta)$  with  $(\beta, \delta)$  furnishes us a contradiction with strongly Archimedean property. Since  $\beta \neq \lambda$ , therefore,  $I_{\succ}(m_{xy}, z)$  is open. An analogous argument implies  $I_{\prec}(m_{xy}, z)$  is open. Therefore  $\succsim$  is strongly strict-Archimedean.

Now assume  $\succsim$  is strongly mixture-continuous and strongly strict-Archimedean. Pick  $x, y, z \in X, m_{xz}, m_{yz} \in \mathcal{M}$ . Assume  $x \succ y$ . Then  $m_{xz}(1) = x$  implies that  $1 \in I_{\succ}(m_{xz}, y)$ . Since  $I_{\succ}(m_{xz}, y)$  is open, therefore there exists  $\lambda < 1$  such that  $(\lambda, 1] \subseteq I_{\succ}(m_{xz}, y)$ . Similarly,  $1 \in I_{\prec}(m_{yz}, x)$ . Since  $A_{\prec}(m_{yz}, x)$  is open, therefore there exists  $\delta < 1$  such that  $(\delta, 1] \subseteq I_{\prec}(m_{yz}, x)$ . Hence,  $\succsim$  is strongly Archimedean.  $\blacksquare$

We are now ready to prove Theorem 6.

*Proof of Theorem 6.* Lemma 7 and an argument analogous to the one presented in the proof of Proposition 2 establish the equivalence between arc-continuity postulate and the postulates of strong mixture-continuity and strong Archimedean.

It remains to prove the equivalence between arc-continuity and continuity postulates. The backward direction is obvious. In order to prove the forward direction assume  $\succsim$  is arc-continuous and pick  $x_0 \in X$  and  $x \in \text{cl}A_{\succ}(x_0)$ . If  $A_{\succ}(x_0)$  is empty or singleton, then it is closed. Otherwise pick a sequence  $x_n$  in  $A_{\succ}(x_0)$  such that  $x_n \rightarrow x$ . It follows from Theorem 8 that there exists a smooth curve<sup>77</sup> containing  $x$  and a subsequence  $x_{n_k}$  of  $x_n$ . Then arc-continuity implies that  $x \in A_{\succ}(x_0)$ . The closedness of the lower sections of  $\succsim$  follows from an analogous argument.

<sup>77</sup>We note for the convenience of the reader that this step of the proof uses the polyhedron assumption. The existence of such a smooth curve is also guaranteed when the boundary of the domain of the binary relation is

Now assume  $A_{\succ}(x_0)$  is not open for some  $x_0 \in X$ . Then there exists  $x \in A_{\succ}(x_0) \cap \text{cl}(A_{\succ}(x_0))^c$ . Pick a sequence  $x_n$  in  $(A_{\succ}(x_0))^c$  such that  $x_n \rightarrow x$ . It follows from Theorem 8 that there exists a smooth curve containing  $x$  and a subsequence  $x_{n_k}$  of  $x_n$ . This implies that  $\succ$  is not strongly strict-Archimedean. Then the first equivalence we proved above and Lemma 7 furnish us a contradiction. The openness of the lower sections of  $\succ$  follows from an analogous argument.  $\blacksquare$

Next, we present the proof of Proposition 4 which uses the construction in Gul's (1992) proof.

*Proof of Proposition 4.* Let  $I^* \in \mathcal{N}$  satisfying A3(ii), i.e., for all  $x, y \in \Delta$ ,  $xI^*y \sim yI^*x$ .

*Claim 2.* For all  $x, y, z, z' \in \Delta$  with  $x > y$  and  $z \geq z'$ , (i)  $x \succ xI^*y \succ y$  and (ii)  $xI^*z \succ yI^*z'$ .

Note that Claim 2 and mixture continuity imply that for all  $x, y \in \Delta$ , there exists a unique  $z \in \Delta$  such that  $z \sim xI^*y$ .

Let  $S = (x^1, \dots, x^m)$  where  $x^t \in \Delta$  for all  $t$ . We say that  $y^0$  reaches  $x$  through  $S$  if  $y^t \sim y^{t-1}I^*x^t$  for  $t = 1, \dots, m$  and  $y^m = x$ .

*Claim 3.* (i) For any  $x, y^0 \in \Delta$ , define  $y^t \sim y^{t-1}I^*x$  where  $y^t \in \Delta$  for  $t \geq 1$ . Then for all  $t$  there exists a unique  $\lambda^t \in [0, 1]$  such that  $y^t = 1\lambda^t 0, \lambda^t \rightarrow \bar{\lambda}$  and  $1\bar{\lambda} 0 = x$ .

(ii) For  $y^0 \in \Delta$  and  $x \in (0, 1)^n$ , there exists some finite  $S$  such that  $y^0$  reaches  $x$  through  $S$ .

Let  $\bar{S} = (x^1, \dots, x^m)$  where  $x^t \in X$  for all  $t$ . We say that  $y^0 \in X$  reaches  $x \in X$  through  $\bar{S}$  if for all  $i \in N$ ,  $y_i^t \sim y_i^{t-1}I^*x_i^t$  for  $t = 1, \dots, m$  and  $y^m = x$ .

*Claim 4.* The following is true.

- (i)  $y^0 \in X$  and  $x \in (0, 1)^n$  imply that there exists a finite  $\bar{S}$  such that  $y^0$  reaches  $x$  through  $\bar{S}$ .
- (ii) If  $y^0 \in X$  reaches  $x \in X$  through  $\bar{S}$  and  $\hat{y}^0$  reaches  $\hat{x}$  through  $\bar{S}$ , then  $y^0 \succ \hat{y}^0$  iff  $x \succ \hat{x}$  and for all  $i \in N$ ,  $y_i^0 \succ \hat{y}_i^0$  iff  $x^t \succ \hat{x}^t$ .

Now pick  $x, y \in X$  such that  $x > y$ , i.e.,  $x_i \geq y_i$  for all  $i$  and  $x_j > y_j$  for some  $j$ . We next show that  $x \succ y$ . We establish the result for the case in which  $x_{-j} = y_{-j}$ . Then the transitivity of  $\succ$  yields the desired conclusion. It follows from Claim 4(i) that for  $z \in \Delta$ , there exists  $\bar{S}$  such that  $x$  reaches  $z$  through  $\bar{S}$ . Define  $\bar{y} \in X$  as  $\bar{y}_{-j} = z$  and  $\bar{y}_j = w$  where  $w \in \Delta$  with  $w < z$ . Then Claim 4(ii) implies that  $y$  reaches  $\bar{y}$  through  $\bar{S}$ . By Claim 3(i), we can choose  $w$  arbitrarily close to  $z$ , hence  $1 \succ \bar{y} \succ 0$ . Then mixture continuity implies there exists  $\bar{z}$  such that  $\bar{z} \sim \bar{y}$ . Then A2 implies that  $z \succ \bar{z} \sim \bar{y}$ . Therefore Claim 4(ii) implies that  $x \succ y$ .

It remains to prove Claims 2, 3 and 4.

---

locally smooth at every point. A natural question arises whether these requirements can be dispensed with, and we leave this question for further investigation. On this issue, also see Footnote 67 above.

*Proof of Claim 2.* Pick  $x, y, z, z' \in \Delta$  such that  $x > y$  and  $z \geq z'$ . Then A3(i) implies that  $x \succ y$  and  $z \succcurlyeq z'$ .

(i) Assume towards a contradiction that  $xI^*y \not\succcurlyeq x$ . Then  $y \prec x$ , completeness and mixture-continuity of  $\succcurlyeq$  imply that there exists  $\lambda \in (0, 1]$  such that  $(x\lambda y)I^*y \sim x$ . Moreover, A3(i) implies that  $(x\lambda y)I^*(x\lambda y) \succ yI^*y$ . Then  $x \sim yI^*(x\lambda y)$ ,  $(x\lambda y) \succ y$  and A2 imply that  $(x\lambda y) \succ x$ . This contradicts A3(i). Hence  $x \succ xI^*y$ . An analogous argument implies that  $xI^*y \succ y$ .

(ii) It follows from mixture continuity, A3(i) and part (i) above that there exists  $x', y' \in \Delta$  such that  $x' \sim xI^*z$  and  $y' \sim yI^*z$ . Then  $x \succ y$  and A2 imply that  $x' \succ y'$ , hence  $xI^*z \succ yI^*z$ . If  $z = z'$ , then the proof is complete. If  $z > z'$ , then A3(i) implies that  $z \succ z'$ . Then A3(ii) and the argument in part (i) above imply that there exists  $y'' \in \Delta$  such that  $y'' \sim yI^*z'$ . Hence A2 imply that  $xI^*z \succ yI^*z'$ . ■

*Proof of Claim 3.* (i) Assume wlog that  $x > y^0$ . Then A3(i) and Claim 2 imply that  $y^t$  is a strictly increasing sequence and  $y^t < x$  for all  $t$ . Since  $1 \geq x > y^t \geq 0$ , therefore for all  $t$  there exists a unique  $\lambda^t \in [0, 1]$  such that  $y^t = 1\lambda^t 0$ . Assume  $1\bar{\lambda}0 < x$  where  $\bar{\lambda} = \lim \lambda^t$ . Define  $\bar{y} = 1\bar{\lambda}0$ . Let  $\hat{y} \in \Delta$  be such that  $\hat{y} \sim \bar{y}I^*x$ . Define  $\hat{\lambda}$  such that  $1\hat{\lambda}0 = \hat{y}$ . Claim 2 implies that  $\bar{y} < \hat{y} < x$ . Hence  $\hat{y} > 0.5(\hat{y} + \bar{y}) > \bar{y} > y^{t+1} \sim y^t I^*x$  for all  $t$ . Then A3(i) implies that  $\hat{y} \succ 0.5(\hat{y} + \bar{y}) \succ y^t I^*x$  for all  $t$ . Hence  $\hat{y} \succ 0.5(\hat{y} + \bar{y}) \succ (1\lambda^t 0)I^*x$ . Note that  $(1\bar{\lambda}0)I^*x = \bar{y}I^*x \sim \hat{y} \succ 0.5(\hat{y} + \bar{y})$ . This contradicts mixture-continuity of  $\succcurlyeq$ , i.e., the closedness of  $\{\lambda \in [0, 1] \mid 0.5(\hat{y} + \bar{y}) \succcurlyeq (1I^*x)\lambda(0I^*x)\}$ .

(ii) Assume wlog  $x > y^0$ . Let  $y^t \sim y^{t-1}I^*1$ . Part (i) implies that  $y^t \rightarrow 1$ . Let  $\tau = \inf\{t \mid y^t > x\} - 1$ . Since  $x < M$  and  $y^t \rightarrow 1$ ,  $\tau$  is well-defined. Hence  $y^{\tau-1} \leq x < y^\tau \sim y^{\tau-1}I^*1$ . By A3(i),  $y^{\tau-1}I^*y^{\tau-1} \preccurlyeq x \prec y^{\tau-1}I^*1$ . Then mixture continuity implies that there exists  $\lambda \in [0, 1]$  such that  $x \sim (y^{\tau-1}I^*1)\lambda(y^{\tau-1}I^*y^{\tau-1}) = y^{\tau-1}I^*(1\lambda y^{\tau-1})$ . Define  $z = 1\lambda y^{\tau-1}$ . Setting  $x^i = M$  for  $i = 1, \dots, n-1$  and  $x^n = z$  establishes the desired set  $S$ . ■

*Proof of Claim 4.* (i) follows from a repeated application of Claim 3.

(ii) The first statement in (ii) follows from a repeated application of A2; the second statement follows from Claim 2 and A3. ■

We next turn to the proof of Proposition 5.

*Proof of Proposition 5.* (a) Note that independence property,  $x \succcurlyeq y$  and  $\lambda \in (0, 1)$  imply that  $\lambda x = \lambda x + (1 - \lambda)0 \succcurlyeq \lambda y + (1 - \lambda)0 = \lambda y$ . Then  $\succcurlyeq$  is lower-homothetic.<sup>78</sup> The following result of

<sup>78</sup>A binary relation is *lower-homothetic* if for all  $x, y \in X$  and all  $\lambda \in (0, 1)$  with  $\lambda x, \lambda y \in X$ ,  $x \succcurlyeq y$  implies  $\lambda x \succcurlyeq \lambda y$ .



Gerasimou (2015, Claim 2) finishes the proof except that independence implies homotheticity.

*Claim 5. A lower-homothetic relation on a vector space is additive iff it is independent.*

Assume  $\succsim$  is independent. The discussion above and Claim 5 imply that the relation is additive and lower-homothetic. Additivity and transitivity imply that if  $x \succsim y$ , then  $kx = \sum_{i=1}^k x \succsim \sum_{i=1}^k y = ky$  for all natural number  $k$ . In order to see this,  $x \succsim y$  implies that  $x + x \succsim x + y \succsim y + y$ . It follows from transitivity that  $2x \succsim 2y$ . Repeating this step  $k$ -many times provides the desired result. Now pick  $x \succsim y$  and  $\lambda > 1$ . Let  $k$  be the largest integer smaller than or equal to  $\lambda$  and set  $\lambda' = \lambda - k$ . Note that  $\lambda x = kx + \lambda'x$ . It follows from additivity that  $kx \succsim ky$  and from lower-homotheticity that  $\lambda'x \succsim \lambda'y$ . Then additivity implies that  $\lambda x = kx + \lambda'x \succsim ky + \lambda'y = \lambda y$ . Hence  $\succsim$  is homothetic.

(b) Let  $\succsim$  be a complete, transitive, mixture-continuous and additive relation on  $X$ . If we show that  $\succsim$  is lower homothetic, then Claim 5 above shows that  $\succsim$  is independent. To this end assume it is not lower homothetic, i.e., there exist  $x, y \in X$  with  $x \succsim y$  and  $\alpha \in (0, 1)$  such that  $\alpha x \prec \alpha y$ . Additivity implies that  $z = x - y \succsim 0$  and  $\alpha z \prec 0$ . Additivity and transitivity imply that  $kz \succsim 0$  for any positive integer  $k$ . Completeness and mixture-continuity imply that  $\succsim$  is strict-Archimedean. Therefore  $I_{\prec}(l_{z0}, 0)$  is open. Then there exists an interval  $(\alpha_1, \alpha_2)$  which contains  $\alpha$  such that  $\alpha'z \prec 0$  for all  $\alpha' \in (\alpha_1, \alpha_2)$ . Pick a rational  $\lambda \in (\alpha_1, \alpha_2)$ . Then there exist positive integers  $m, n$  such that  $\lambda = m/n$ . Then, transitivity and additivity of  $\succsim$  imply that  $mz = \sum_{i=1}^n (m/n)z \prec 0$ . This furnishes us a contradiction. Therefore,  $\succsim$  is lower homothetic.

The backward direction follows from part (a). Therefore the proof of Proposition 5 is complete. ■

*Proof of Proposition 6.* It is clear that part (b) implies part (a) is true. Assume part (a). It follows from Proposition 10 that the relation is complete and transitive, from Theorem 4 that it is mixture-continuous, and from Proposition 5 that it is independent. The celebrated representation theorem of Herstein-Milnor completes the proof. ■

*Proof of Proposition 7.* It is easy to show that open sections property implies that the relation has the strict Archimedean property which implies the relation has the Archimedean property. Therefore, it remains to prove that Archimedean property implies open sections property. To this end, assume  $\succsim$  is Archimedean. Pick  $x \in \mathbb{R}_+^n$  and  $y \in A_{\succ}(x)$ . Then Archimedean property implies that there exists  $\lambda \in (0, 1)$  such that  $y\lambda 0 \succ x$ . Define  $z = y\lambda 0$ . Since  $y > 0$ , therefore  $z > 0$ . Define  $\varepsilon = \min_{z_i \neq 0} (y_i - z_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' > z$ . Then strong monotonicity implies that  $y' \succ z$ . Since  $z \succ x$ , it follows from transitivity of  $\succ$  that  $y' \succ x$  for all  $y'$  in the  $\varepsilon$  neighborhood of  $y$ . Therefore,  $\succ$  has open upper sections.

Now pick  $x \in \mathbb{R}_+^n, y \in A_{\prec}(x)$  and  $x' \geq x, y$ . Then Archimedean property implies that there exists  $\lambda \in (0, 1)$  such that  $x \succ x'\lambda y$ . Note that  $x' > y$ . Define  $z = x'\lambda y$ . Since  $x' > y$ , therefore  $z > y$ . Define  $\varepsilon = \min_i (z_i - y_i)$ . For any  $y'$  in the  $\varepsilon$  neighborhood of  $y$ ,  $y' < z$ . Then strong monotonicity implies that  $z \succ y'$ . Since  $x \succ z$ , it follows from transitivity of  $\succ$  that  $x \succ y'$ . This furnishes us a contradiction with  $y^k \rightarrow y$ . Therefore,  $\succ$  has open lower sections. ■

*Proof of Proposition 8.* Continuity implies mixture-continuity, and the converse implication follows from completeness and convexity assumptions and Theorem 3. It follows from completeness of the relation that mixture-continuity implies Archimedean property. In order to show the converse implication pick  $x, y, z \in \mathbb{R}_+^n$  and  $\lambda \in (0, 1)$  such that  $x\lambda y \succ z$ . It follows from Archimedean property that there exists  $\delta, \gamma \in (0, 1)$  such that  $x(\lambda\delta)y \succ z$  and  $y((1-\lambda)\gamma)x \succ z$ . Note that  $y((1-\lambda)\gamma)x = x(1 - (1-\lambda)\gamma)y$  and  $\underline{\lambda} = \lambda\delta < \lambda < 1 - (1-\lambda)\gamma = \bar{\lambda}$ . Then the convexity assumption implies that for all  $\lambda' \in (\underline{\lambda}, \bar{\lambda})$ ,  $x\lambda'y \succ z$ . Hence,  $\succ$  is upper strict-Archimedean. Therefore, it is lower mixture-continuous.

Now assume that there exists  $x, y, z \in \mathbb{R}_+^n$  and  $\lambda^k \in [0, 1]$  such that  $x\lambda^k y \not\succ z$  for all  $k$  with  $\lambda^k \rightarrow \lambda$  such that  $x\lambda y \prec z$ . If there exists  $k, l$  such that  $\lambda^k < \lambda < \lambda^l$ , then the convexity assumption implies that  $x\lambda y \succ z$ . Then assume wlog that  $\lambda^k > \lambda$  for all  $k$ . It follows from the convexity assumption that  $x\lambda'y \succ z$  for all  $\lambda' \in (\lambda^1, \lambda)$ . Archimedean property and  $z \succ x\lambda y$  implies that there exists  $\delta \in (0, 1)$  such that  $z \succ (x\lambda y)\delta(x\lambda^1 y) = x(\delta\lambda + (1-\delta)\lambda^1)y$ . Then  $\lambda < \delta\lambda + (1-\delta)\lambda^1 < \lambda^1$  furnishes us a contradiction. Hence,  $\succ$  is upper mixture-continuous. ■

*Proof of Proposition 9.* It follows from Proposition 2 that the relation is linearly continuous, from Theorem 3 that it is continuous and from Khan-Uyanik (2019, Theorem 2) that it is complete and transitive. ■

We now turn to the proof of Proposition 10. The construction of our proof has some similarity to the proof of Neufeind-Trockel (1995, Proposition), however their method-of-proof does not work since the topological structure of the space has a central role in their proof. We, instead, define a closure concept for subsets of vector spaces by using the topological structure of the unit interval and then obtain our result by using this closure concept. Let  $A$  be a subset of a vector space  $X$ . The *linear closure* of  $A$  is defined as  $\bar{A} = \{z \in X | x, y \in X, \lambda^n \rightarrow \lambda, x\lambda^n y \in A, z = x\lambda y\}$ . In other words, for all  $x, y \in X$  and  $\lambda^n \rightarrow \lambda$ , if  $x\lambda^n y \in A$  for all  $n$ , then  $x\lambda y \in \bar{A}$ .<sup>79</sup>

*Proof of Proposition 10.* Theorem 4 implies that  $\succ$  is mixture-continuous. It is clear from the argument in the proof of Proposition 1 of Galaabaatar-Khan-Uyanik (2019) that mixture-continuity and upper Archimedean properties imply upper strict-Archimedean property. Then,

---

<sup>79</sup>Note that linear closure of a set contains its “algebraic closure.”

Theorem 4 implies that  $\succsim$  is strict-Archimedean. Therefore, if we show that  $\succsim$  is reflexive and  $\sim$  is transitive, then the proof follows from Galaabaatar-Khan-Uyanik (2019, Theorem 1).

The following claim implies that the restriction of  $\succ$  on the line segment connecting any two points is negatively transitive.

*Claim 6.* For all  $x, y \in X$  with  $x \succ y$ ,  $[x, y] \subseteq A_{\succ}(y) \cup A_{\prec}(x)$ , where  $[x, y] = \{x\lambda y \mid \lambda \in [0, 1]\}$ .

It is easy to see that transitivity of  $\sim$  of an additive relation on a vector space is equivalent to the condition  $A_{\sim}(0) + A_{\sim}(0) \subseteq A_{\sim}(0)$ . Similarly, semi-transitivity of  $\succsim$  is equivalent to the condition  $A_{\succ}(0) + A_{\sim}(0) \subseteq A_{\succ}(0)$ .

*Claim 7.*  $A_{\succsim}(0) = \overline{A_{\succ}(0)}$  and  $\overline{A_{\succ}(0)} + A_{\sim}(0) \subseteq \overline{A_{\succ}(0) + A_{\sim}(0)}$ .

Semi-transitivity and Claim 7 imply that  $A_{\succsim}(0) + A_{\sim}(0) \subseteq \overline{A_{\succ}(0)} = A_{\succ}(0)$ . Therefore,  $A_{\sim}(0) + A_{\sim}(0) \subseteq A_{\succ}(0)$ . An analogous argument implies that  $A_{\sim}(0) + A_{\sim}(0) \subseteq A_{\prec}(0)$ , hence  $\sim$  is transitive.

*Claim 8.*  $\succsim$  is reflexive.

It follows from Galaabaatar-Khan-Uyanik (2019, Theorem 1) that  $\succsim$  is complete and transitive. It remains to prove Claims 6, 7 and 8 in order to complete the proof.

*Proof of Claim 6.* Pick  $x, y \in X$  with  $x \succ y$  and  $\lambda \in [0, 1]$ . Assume  $\lambda \in I_{\succ}(l_{xy}, y)$  and  $\lambda \notin I_{\succ}(l_{xy}, y)$ . Then  $x\lambda y \sim y$ . Semi-transitivity implies that  $x \succ x\lambda y$ , hence  $\lambda \in I_{\prec}(l_{xy}, x)$ . Now assume  $\lambda \in I_{\prec}(l_{xy}, x)$  and  $\lambda \notin I_{\prec}(l_{xy}, y)$ . Then  $x\lambda y \sim x$ . Semi-transitivity implies that  $x\lambda y \succ y$ , hence  $\lambda \in I_{\succ}(l_{xy}, y)$ . Therefore, it follows from  $x \succ y$  and  $[0, 1]$  is connected that

$$[0, 1] = I_{\succ}(l_{xy}, y) \cup I_{\prec}(l_{xy}, x) = I_{\succ}(l_{xy}, y) \cup I_{\prec}(l_{xy}, x).$$

Since  $A_{\succ}(y) \cap [x, y] = \{x\lambda y \mid \lambda \in I_{\succ}(l_{xy}, y)\}$ ,  $A_{\prec}(x) \cap [x, y] = \{x\lambda y \mid \lambda \in I_{\prec}(l_{xy}, x)\}$  and  $l_{xy}([0, 1]) = [x, y]$ , therefore  $[x, y] \subseteq A_{\succ}(y) \cup A_{\prec}(x)$ . ■

*Proof of Claim 7.* We start by showing  $A_{\succsim}(0) \subseteq \overline{A_{\succ}(0)}$ . Note that non-triviality and additivity imply that  $\bar{x} \succ 0$  for some  $\bar{x} \in X$ . Hence  $A_{\succ}(0) \neq \emptyset$ . Since  $A_{\succsim}(0) = A_{\succ}(0) \cup A_{\sim}(0)$  and , therefore showing that  $A_{\sim}(0) \subseteq \overline{A_{\succ}(0)}$  is enough. To this end pick  $x \sim 0$  and set  $\lambda = 0.5$ . It follows from Claim 6 and semi-transitivity that  $\bar{x}\lambda x \in A_{\succ}(0) \cup A_{\prec}(\bar{x})$ . We now show that  $\bar{x}\lambda x \in A_{\succ}(0)$ . If  $\bar{x}\lambda x \succ 0$ , then we are done. If  $\bar{x} \succ \bar{x}\lambda 0$ , then it follows from additivity that  $\bar{x}\lambda x \succ x$ . And semi-transitivity implies that  $\bar{x}\lambda x \succ 0$ . Repeating this argument by using Claim 6, semi-transitivity and additivity imply that  $\bar{x}\lambda^n x \in A_{\succ}(0)$  for all natural number  $n$ . Since  $\bar{x}\lambda^n x \rightarrow x$ , therefore  $x \in \overline{A_{\succ}(0)}$ .

For the inverse inclusion, pick  $y\lambda^n z \in \overline{A_{\succ}(0)}$  such that  $\lambda^n \rightarrow \lambda$ . Then mixture-continuity implies that  $y\lambda z \in A_{\succ}(0)$ . Hence  $\overline{A_{\succ}(0)} \subseteq A_{\succ}(0)$ .

We next prove  $\overline{A_{\succ}(0)} + A_{\sim}(0) \subseteq \overline{A_{\succ}(0) + A_{\sim}(0)}$ . Pick  $x^0 \in \overline{A_{\succ}(0)}$  and  $z \in A_{\sim}(0)$ . Then there exists  $x\lambda^n y \in A_{\succ}(0)$  such that  $\lambda^n \rightarrow \lambda$  and  $x^0 = x\lambda y$ . Observing that  $x\lambda^n y + z = (x + z)\lambda^n(y + z) \rightarrow (x + z)\lambda(y + z) = x\lambda y + z \in \overline{A_{\succ}(0) + A_{\sim}(0)}$  finishes the proof. ■

*Proof of Claim 8.* Reflexivity of an additive relation is equivalent to the condition  $0 \in A_{\sim}(0)$ . Claim 7 implies that showing  $0 \in \overline{A_{\succ}(0)}$  is enough. Set  $\lambda = 0.5$ . It follows from Claim 6 that  $\bar{x}\lambda 0 \in A_{\succ}(0) \cup A_{\prec}(\bar{x})$ . We first show that  $\bar{x}\lambda 0 \in A_{\succ}(0)$ . If  $\bar{x}\lambda 0 \succ 0$ , then we are done. Then assume  $\bar{x} \succ \bar{x}\lambda 0$ . It follows from additivity that  $\bar{x}\lambda 0 \succ 0$ . Repeating this argument by using Claim 6 and additivity imply that  $\bar{x}\lambda^n 0 \in A_{\succ}(0)$  for all natural number  $n$ . Since  $\bar{x}\lambda^n 0 \rightarrow 0$ , therefore  $0 \in \overline{A_{\succ}(0)}$ . ■

The proof of Proposition 10 is complete. ■

## A.2 Seven Technical Examples

Section 3 established equivalence between different continuity postulates under convexity or monotonicity assumptions, and Section 4 applied the results to the antecedent economic literature. Through illustrative examples, this section shows that some of the assumptions in the various theorems are not redundant.

The first example illustrates that linear continuity of a relation is not equivalent to its continuity on arbitrary convex subsets of a Euclidean space. Hence, the restrictive assumptions on the choice set we impose in this paper are not redundant.

**Example 1.** Let  $X = \{x \in \mathbb{R}^2 | x_1^2 + x_2^2 \leq 1\}$  be the unit sphere,  $A = \{x \in X | x_1 < 0, x_1^2 + x_2^2 \neq 1\} \cup \{(-1, 0)\}$  and  $B = \{x \in X | x_1 > 0, x_1^2 + x_2^2 \neq 1\} \cup \{(1, 0)\}$ . Define a binary relation  $\succsim$  on  $X$  as follows. For all  $x \in A$  and all  $y \in B$ ,  $x \succ y$ . Assume there are no other comparable points. It is easy to see that  $\succsim$  has convex sections and satisfies strict-Archimedean property. However, it does not have open sections since the sections of  $\succsim$  at any  $x \in X$  is either  $A$  or  $B$  which are not open.

The following example illustrates that if we drop the order-denseness property from the (weak) Wold-continuity postulate, then the binary relation can be discontinuous under weak monotonicity.<sup>80</sup>

**Example 2.** Define a relation  $\succsim$  on  $\mathbb{R}_+^2$  as follows:  $0 \sim 0$ ,  $x \succ 0$  for all  $x \neq 0$  and  $x \sim y$  for all  $x, y > 0$ . It is clear that  $\succsim$  is complete, transitive and weakly monotonic. Moreover, it trivially

<sup>80</sup>This example is originally due to Mitra-Ozbek (2013). They show that the relation in this example violates their scalar continuity property.

satisfies the second part of Wold-continuity (hence of weak Wold-continuity) since there does not exist  $x, y, z$  such that  $x \succ y \succ z$ . Since  $A_{\prec}(x) = \{0\}$  is not open for all  $x \neq 0$ ,  $\succ$  is not continuous.

Note that, under strong monotonicity, however, the second part of (weak) Wold-continuity implies the order-denseness property for a transitive relation. In order to see this pick  $x \succ y$ . If  $y = 0$ , then  $x \succ x\lambda 0 \succ 0$  for all  $\lambda \in (0, 1)$ . Otherwise pick  $z$  on the diagonal such that  $z \gg x, y$ . Then  $z \succ x \succ y \succ 0$ . Then there exists  $0 < \lambda < \delta < 1$  such that  $x \sim z\delta 0$  and  $y \sim z\lambda 0$ . For any  $\mu \in (\lambda, \delta)$ ,  $x \sim z\delta 0 \succ z\mu 0 \succ z\lambda 0 \sim y$ . Then transitivity completes the proof.

The following example shows that convexity and additivity properties of a relation do not imply each other.

**Example 3.** It is clear that a relation with convex (upper) sections may not be additive. In order to see that an additive relation may not have convex sections, consider the following preorder on  $\mathbb{R}$ :  $x \succ y$  if and only if  $x - y \in \mathbb{Z}$ . It is clear that  $\succ$  is additive. Since  $A_{\succ}(x) = A_{\prec}(x) = \{x + k | k \in \mathbb{Z}\}$  for all  $x \in \mathbb{R}$ , therefore the sections of  $\succ$  is not convex.

This implies that Theorem 4 above is neither a corollary of the other results we present in this paper, nor of those presented in Neuefeind-Trockel (1995) or in Gerasimou (2015), and that our results do not imply theirs.

The concept of local convexity of a set we define is slightly stronger than the ordinary local convexity concept in the literature which does not impose any restriction on those points that are not in the set; see Klee (1951, p. 448). However, in terms of the main result in this literature, *for closed, connected, and locally convex subset of  $\mathbb{R}^n$ , the two properties are equivalent*,<sup>81</sup> this definition is equivalent to the local convexity in the ordinary sense since the relevant set is closed in their result. The following example illustrates that our version of local convexity property is stronger.

**Example 4.** Let  $X = \bigcup_{n=1}^{\infty} (\frac{1}{n+1}, \frac{1}{n})$ . It is clear that  $X$  is locally convex in the usual sense since it is open. However,  $0 \in \text{cl}X$  and for all neighborhood  $N_{\varepsilon}(0)$  of 0,  $N_{\varepsilon}(0) \cap X$  contains infinitely many disjoint intervals, hence the intersection is not convex. Therefore  $X$  is not locally convex as we define.

The following example highlights the importance of the definition of the restricted relation given in the text above.

**Example 5.** Let  $X = [0, 1]^2$  and  $Y = \{y \in X \mid y_1 + y_2 = 1, y_1 > 0, y_2 > 0\}$ . Define  $\succ$  on  $X$  as follows: for all  $y \in Y$ ,  $0 \sim y$  and there are no other comparable points. Then,

---

<sup>81</sup>Unlike Rockafellar's theorem, this main result, which is due to 1928 work of Tietze and Nakajima, is true for arbitrary topological vector spaces; see for example Klee (1951).

$A_{\succ}(x) = A_{\prec}(x) = \emptyset$  for all  $x \in X$ , hence they are open and convex. Moreover,  $A_{\succneq}(x)$  and  $A_{\preceqneq}(x)$  are convex for all  $x \in X$ . However,  $A_{\succneq}(0)$  is not closed. Hence,  $\succneq$  is convex but not continuous. This example satisfies all assumptions of Theorem 3(b):  $X$  is a polyhedron,  $\succ$  and  $\succneq$  have (locally) convex sections. We next show that if we define the restricted relation on a set  $S$  as  $\succneq \cap (S \times S)$ , as mentioned in the text, then  $\succneq$  satisfies linear continuity. Note that the restriction of  $A_{\succneq}(x)$  and  $A_{\preceqneq}(x)$  on any straight line in  $X$  for all  $x \in X$  is either a singleton or empty, hence closed. It is trivial that the restriction of  $A_{\succ}(x)$  and  $A_{\prec}(x)$  on any straight line in  $X$  for all  $x \in X$  is empty, hence open. Therefore,  $\succneq$  is restricted continuous. Since  $\succneq$  is not continuous, Theorem 3 fails under this modified version of restricted relation.

The following example illustrates a preference relation that is linearly continuous, Wold-continuous and (trivially) weakly monotone but not continuous whose domain is a convex set that fails property **B**.

**Example 6.** Let  $X = \{x \in [-1, 0] \times [0, 1] \mid x_1 = -x_2\}$  and  $f(x) = \sin(1/x_1)$  if  $x_1 > 0$  and  $f(0, 0) = 1$ . Then, it is clear that  $f(x) \in [-1, 1]$  for all  $x \in X$ . Define a binary relation  $\succneq$  on  $X$  as  $x \succneq y$  is and only if  $f(x) \geq f(y)$ . Note that  $X$  does not satisfy property **B** since for all  $x, y \in X$ , there is no  $a, b \in X$  such that  $x, y \geq a$  and  $b \geq x, y$ . Therefore,  $\succneq$  is trivially monotone. By the arguments in the proof of Proposition 3,  $\succneq$  is Archimedean and Wold-continuous, but not mixture continuous or continuous.

The following example illustrates a function that is quasi-convex, linearly continuous but not jointly continuous whose domain is a convex and closed set that is not a polyhedron, hence property **C** fails.

**Example 7.** Let  $X = \{x \in \mathbb{R}_+^2 \mid x_1^2 \leq x_2\}$ ,  $Y = \{x \in [0, 1]^2 \mid x_1^2 \leq x_2 \leq ax_1\}$ ,  $a > 1$ , and  $f : X \rightarrow \mathbb{R}$  defined as

$$f(x) = \frac{2x_1^2x_2}{x_1^4 + x_2^2} \text{ for } x \neq 0 \text{ and } f(0) = 0.$$

It is clear that  $X, Y$  are non-empty, convex and closed, and  $Y \subseteq X$ . Note that, as  $f$  is the function in the Genocchi-Peano example, both  $f$  and its restriction on  $Y$  are linearly continuous but they are not jointly continuous.

We now show that  $f$  is quasi-convex on  $Y$ . First, we show that  $f$  is quasi-convex on  $\text{int}X$  (the interior of  $X$ ). Note that  $f$  is twice differentiable on  $\text{int}X$ . Hence if the determinant of the first and second order bordered Hessian of  $f$  is negative for all  $x$  in  $\text{int}X$ , then  $f$  is quasi-convex

on  $\text{int}X$ . Pick  $x \in \text{int}X$ . Note that  $x_1, x_2 > 0$ . The first order bordered Hessian of  $f$  at  $x$  is

$$BH_1 = \begin{bmatrix} 0 & \frac{4x_1x_2}{x_1^4+x_2^2} - \frac{8x_1^5x_2}{(x_1^4+x_2^2)^2} \\ \frac{4x_1x_2}{x_1^4+x_2^2} - \frac{8x_1^5x_2}{(x_1^4+x_2^2)^2} & -\frac{56x_1^4x_2}{(x_1^4+x_2^2)^2} + \frac{4x_2}{x_1^4+x_2^2} + \frac{64x_1^8x_2}{(x_1^4+x_2^2)^3} \end{bmatrix}$$

The determinant of  $BH_1(x)$  is

$$\det(BH_1) = -\frac{16x_1^2x_2^2(x_1^2+x_2)^2(x_1^2-x_2)^2}{(x_1^4+x_2^2)^4}$$

Note that  $x_1, x_2 > 0$  and  $x_2 > x_1^2$ . Therefore,  $\det(BH_1) < 0$ . The second order bordered Hessian of  $f$  at  $x$  is

$$BH_2 = \begin{bmatrix} 0 & \frac{4x_1x_2}{x_1^4+x_2^2} - \frac{8x_1^5x_2}{(x_1^4+x_2^2)^2} & \frac{2x_1^2}{x_1^4+x_2^2} - \frac{4x_1^2x_2^2}{(x_1^4+x_2^2)^2} \\ \frac{4x_1x_2}{x_1^4+x_2^2} - \frac{8x_1^5x_2}{(x_1^4+x_2^2)^2} & -\frac{56x_1^4x_2}{(x_1^4+x_2^2)^2} + \frac{4x_2}{x_1^4+x_2^2} + \frac{64x_1^8x_2}{(x_1^4+x_2^2)^3} & \frac{4x_1}{x_1^4+x_2^2} - \frac{8x_1x_2^2+8x_1^5}{(x_1^4+x_2^2)^2} + \frac{32x_1^5x_2^2}{(x_1^4+x_2^2)^3} \\ \frac{2x_1^2}{x_1^4+x_2^2} - \frac{4x_1^2x_2^2}{(x_1^4+x_2^2)^2} & \frac{4x_1}{x_1^4+x_2^2} - \frac{8x_1x_2^2+8x_1^5}{(x_1^4+x_2^2)^2} + \frac{32x_1^5x_2^2}{(x_1^4+x_2^2)^3} & \frac{16x_1^2x_2^3}{(x_1^4+x_2^2)^3} - \frac{12x_1^2x_2}{(x_1^4+x_2^2)^2} \end{bmatrix}$$

The determinant of  $BH_2(x)$  is

$$\det(BH_2) = -\frac{16x_1^4x_2(x_2^2-x_1^4)^3}{(x_1^4+x_2^2)^6}$$

It follows from  $x_1, x_2 > 0$  and  $x_2 > x_1^2$  that  $\det(BH_2) < 0$ . Since  $x$  is an arbitrary point in  $\text{int}X$ ,  $f$  is quasi-convex on  $\text{int}X$ .

We next show that  $f$  is quasi-convex on  $Y$ . Recall that  $f$  is quasi-convex if for all  $x, y \in X$ , all  $\lambda \in (0, 1)$ ,  $f(\lambda x + (1 - \lambda)y) \leq \max\{f(x), f(y)\}$ . Pick  $x, y \in Y$  and  $\lambda \in (0, 1)$ . If  $x, y \in \text{int}X$ , then by the argument above, quasi-convexity holds. Hence assume at least one of  $x$  and  $y$  is on the boundary of  $Y$ . Note that the part of the boundary of  $Y$  that is not contained in  $\text{int}X$  consists of  $\{z \in Y \mid z = 0 \text{ or } z_2 = z_1^2\}$ . It is easy to observe that  $f(x) \leq 1$  for all  $x \in Y$ . Hence, if  $x_2 = x_1^2$  or  $y_2 = y_1^2$ , then  $f(\lambda x + (1 - \lambda)y) \leq \max\{f(x), f(y)\}$  is trivially true. Finally, let  $y = 0$  and  $x \in \text{int}X$ . Then,

$$f(\lambda x + (1 - \lambda)y) = \frac{2\lambda^3x_1^2x_2}{\lambda^2(\lambda^2x_1^4+x_2^2)} = \frac{2\lambda x_1^2x_2}{\lambda^2x_1^4+x_2^2} \leq \frac{2x_1^2x_2}{x_1^4+x_2^2} = f(x) = \max\{f(x), f(y)\}.$$

Therefore,  $f$  is quasi-convex.

Note that even though this example is presented for functions, the binary relation  $\succsim$  on  $X$  induced by the function  $-f$  is complete, transitive, convex and linearly continuous but not continuous.

## B Bibliography

- ABDELLAOUI, M., AND P. WAKKER (2020): “Savage for dummies and experts,” *Journal of Economic Theory*, 186, 104991, 1–20.
- AHN, D. S., AND H. ERGIN (2010): “Framing contingencies,” *Econometrica*, 78(2), 655–695.
- ALESKEROV, F., AND B. MONJARDET (2002): *Utility Maximization, Choice and Preference*. Berlin: Springer-Verlag.
- ARKREYD, L. (2005): “Nonstandard analysis,” *The American Mathematical Monthly*, 112(10), 926–928.
- ARROW, K. J. (1966): “Exposition of the theory of choice under uncertainty,” *Synthese*, pp. 253–269.
- ARROW, K. J., AND F. H. HAHN (1971): *General Competitive Analysis*. Amsterdam: North-Holland.
- AUMANN, R. J. (1966): “Existence of competitive equilibria in markets with a continuum of traders,” *Econometrica*, pp. 1–17.
- BANERJEE, K., AND T. MITRA (2018): “On Wold’s approach to representation of preferences,” *Journal of Mathematical Economics*, 79, 65–74.
- BASU, K., AND T. MITRA (2003): “Aggregating infinite utility streams with intergenerational equity: the impossibility of being Paretian,” *Econometrica*, 71(5), 1557–1563.
- (2007): “Utilitarianism for infinite utility streams: A new welfare criterion and its axiomatic characterization,” *Journal of Economic Theory*, 133(1), 350–373.
- BEARDON, A. F., AND G. B. MEHTA (1994): “The utility theorems of Wold, Debreu, and Arrow-Hahn,” *Econometrica*, 62(1), 181–186.
- BEHRINGER, F. A. (1992): “Convexity is equivalent to midpoint convexity combined with strict quasiconvexity,” *Optimization*, 24(3-4), 219–228.
- BERGSTROM, T. C., R. P. PARKS, AND T. RADER (1976): “Preferences which have open graphs,” *Journal of Mathematical Economics*, 3(3), 265–268.
- BIRKHOFF, G. (1948): *Lattice Theory (Revised Edition)*, vol. XXV. New York: American Mathematical Society.
- BLACKWELL, D., AND M. GIRSHICK (1954): *Theory of Games and Statistical Decisions*. New York: John Wiley and Sons.
- BORWEIN, J. M., AND R. GOEBEL (2003): “Notions of relative interior in Banach spaces,” *Journal of Mathematical Sciences*, 115(4), 2542–2553.
- BORWEIN, J. M., AND A. S. LEWIS (1992): “Partially finite convex programming, part I: Quasi relative interiors and duality theory,” *Mathematical Programming*, 57(1), 15–48.



- BOSI, G., M. J. CAMPIÓN, J. C. CANDEAL, AND E. INDURAIN (2020): *Mathematical Topics on Representations of Ordered Structures and Utility Theory*. Springer: Switzerland.
- BOSI, G., AND M. ZUANON (2020): “Continuity and continuous multi-utility representations of nontotal preorders: some considerations concerning restrictiveness,” in *Mathematical Topics on Representations of Ordered Structures and Utility Theory: Essays in Honor of Professor Ghanshyam B. Mehta*, ed. by G. Bosi, M. J. Campión, J. C. Candeal, and E. Indurain, pp. 213–236. Springer: Switzerland.
- BRIDGES, D. S., AND G. B. MEHTA (1995): *Representations of Preference Orderings*. Berlin: Springer-Verlag.
- BROCK, W. A. (1970): “An Axiomatic Basis for the Ramsey-Weizsäcker Overtaking Criterion,” *Econometrica*, 38(6), 927–929.
- BROWN, D. J., AND A. ROBINSON (1972): “A limit theorem on the cores of large standard exchange economies,” *Proceedings of the National Academy of Sciences, USA*, 69(5), 1258–1260 (A Correction is published in 1972, volume 69, no 10, page 3068).
- CANDEAL-HARO, J. C., AND E. INDURÁIN-ERASO (1995): “A note on linear utility,” *Economic Theory*, 6(3), 519–522.
- CHATEAUNEUF, A., M. COHEN, AND J.-Y. JAFFRAY (2013): “Decision under uncertainty: the classical models,” in *Decision Making Process: Concepts and Methods*, ed. by D. Bouyssou, D. Dubois, H. Prade, and M. Pirlot, pp. 385–400. New Jersey: John Wiley & Sons.
- CHEW, S. H., AND E. KARNI (1994): “Choquet expected utility with a finite state space: Commutativity and act-independence,” *Journal of Economic Theory*, 62(2), 469–479.
- CHIPMAN, J. (1971): “Consumption theory without transitive indifference,” in *Preferences, Utility and Demand: A Minnesota Symposium*, ed. by J. Chipman, L. Hurwicz, M. Richter, and H. Sonenschein, pp. 224–253. New York: Harcourt Brace Jovanovich.
- CIESIELSKI, K. C., AND D. MILLER (2016): “A continuous tale on continuous and separately continuous functions,” *Real Analysis Exchange*, 41(1), 19–54.
- DE FINETTI, B. (1974): *Theory of Probability*, vol. 1. New York: Wiley.
- DEBREU, G. (1954): “Valuation equilibrium and Pareto optimum,” *Proceedings of the National Academy of Sciences*, 40(8), 588–592.
- (1959): *Theory of Value: An Axiomatic Analysis of Economic Equilibrium*, no. 17. Connecticut: Yale University Press.
- (1964): “Continuity properties of Paretian utility,” *International Economic Review*, 5(3), 285–293.
- DEKEL, E. (1986): “An axiomatic characterization of preferences under uncertainty: weakening the independence axiom,” *Journal of Economic Theory*, 40(2), 304–318.
- DESHPANDÉ, J. V. (1968): “On continuity of a partial order,” *Proceedings of the American Mathematical Society*, 19(2), 383–386.
- DIEWERT, W. E., M. AVRIEL, AND I. ZANG (1981): “Nine kinds of quasiconcavity and concavity,” *Journal of Economic Theory*, 25(3), 397–420.
- DUBRA, J. (2011): “Continuity and completeness under risk,” *Mathematical Social Sciences*,

61(1), 80–81.

- DUBRA, J., F. MACCHERONI, AND E. A. OK (2004): “Expected utility theory without the completeness axiom,” *Journal of Economic Theory*, 115(1), 118–133.
- DUGGAN, J. (1999): “A general extension theorem for binary relations,” *Journal of Economic Theory*, 86(1), 1–16.
- DUGUNDJI, J. (1966): *Topology*. Boston: Allyn and Bacon.
- EILENBERG, S. (1941): “Ordered topological spaces,” *American Journal of Mathematics*, 63(1), 39–45.
- EPSTEIN, L. G. (1987): “The unimportance of the intransitivity of separable preferences,” *International Economic Review*, pp. 315–322.
- EVREN, Ö., AND F. HÜSSEINOV (2021): “Extension of monotonic functions and representation of preferences,” *Mathematics of Operations Research*.
- FISHBURN, P. C. (1982): *The Foundations of Expected Utility*. Boston: D. Riedel Publishing Company.
- GALAABAATAR, T., M. A. KHAN, AND M. UYANIK (2019): “Completeness and transitivity of preferences on mixture sets,” *Mathematical Social Sciences*, 99, 49–62.
- GENOCCHI, A., AND G. PEANO (1884): *Calcolo Differenziale e Principii di Calcolo*. Torino: Fratelli Bocca.
- GERASIMOU, G. (2013): “On continuity of incomplete preferences,” *Social Choice and Welfare*, 41(1), 157–167.
- (2015): “(Hemi)continuity of additive preference preorders,” *Journal of Mathematical Economics*, 58, 79–81.
- GHOSH, A., M. A. KHAN, AND M. UYANIK (2020): “Solvability axioms in mathematical psychology and continuity postulates in economic theory: A consolidation in the theory of individual choice,” *Johns Hopkins University, mimeo*.
- GIARLOTTA, A., AND S. GRECO (2013): “Necessary and possible preference structures,” *Journal of Mathematical Economics*, 49(2), 163–172.
- GIARLOTTA, A., AND S. WATSON (2019): “A bi-preference interplay between transitivity and completeness: Reformulating and extending Schmeidler’s theorem,” *Journal of Mathematical Psychology*, (forthcoming).
- GILBOA, I., F. MACCHERONI, M. MARINACCI, AND D. SCHMEIDLER (2010): “Objective and subjective rationality in a multiple prior model,” *Econometrica*, 78(2), 755–770.
- GLATZER, T. J. (2013): “Continuities on Subspaces,” Ph.D. thesis, West Virginia University.
- GORMAN, W. M. (1971): “Preference, revealed preference, and indifference,” in *Preferences, Utility and Demand: A Minnesota Symposium*, ed. by J. Chipman, L. Hurwicz, M. Richter, and H. Sonenschein, pp. 81–113. New York: Harcourt Brace Jovanovich.
- GRANT, S., A. KAJII, AND B. POLAK (1992): “Many good risks: an interpretation of multi-variate risk and risk aversion without the independence axiom,” *Journal of Economic Theory*, 56(2), 338–351.

- GRAVEL, N., T. MARCHANT, AND A. SEN (2012): “Uniform expected utility criteria for decision making under ignorance or objective ambiguity,” *Journal of Mathematical Psychology*, 56(5), 297–315.
- GUL, F. (1992): “Savage’s theorem with a finite number of states,” *Journal of Economic Theory*, 57(1), 99–110.
- HARPER, J. (2016): “Defining continuity of real functions of real variables,” *BSHM Bulletin: Journal of the British Society for the History of Mathematics*, 31(3), 189–204.
- HARTMANN, L. (2020): “Savage’s P3 is redundant,” *Econometrica*, 88(1), 203–205.
- HERDEN, G. (1995): “On some equivalent approaches to mathematical utility theory,” *Mathematical Social Sciences*, 29(1), 19–31.
- HERSTEIN, I. N., AND J. MILNOR (1953): “An axiomatic approach to measurable utility,” *Econometrica*, 21(2), 291–297.
- HERVÉS-BELOSÓ, C., AND H. DEL VALLE-INCLÁN CRUCES (2019): “Continuous preference orderings representable by utility functions,” *Journal of Economic Surveys*, 33(1), 179–194.
- INOUE, T. (2010): “A utility representation theorem with weaker continuity condition,” *Journal of Mathematical Economics*, 46(1), 122–127.
- JENSEN, N. E. (1967): “An introduction to Bernoullian utility theory: I. Utility functions,” *The Swedish journal of Economics*, pp. 163–183.
- JORGENSEN, L. M. (2009): “The principle of continuity and Leibniz’s theory of consciousness,” *Journal of the History of Philosophy*, 47(2), 223–248.
- KARNI, E., AND Z. SAFRA (2015): “Continuity, completeness, betweenness and cone-monotonicity,” *Mathematical Social Sciences*, 74, 68–72.
- KERSHNER, R. (1943): “The continuity of functions of many variables,” *Transactions of the American Mathematical Society*, 53(1), 83–100.
- KETTERING, J., AND A. KOCHOV (2020): “Consumption Smoothing and Discounting in Infinite-Horizon, Discrete-Choice Problems,” *working paper*.
- KHAN, M. A. (1973): “Large exchange economies,” Ph.D. thesis, Yale University.
- KHAN, M. A., AND M. UYANIK (2019): “Topological connectedness and behavioral assumptions on preferences: a two-way relationship,” *Economic Theory*, (published online, doi: 10.1007/s00199-019-01206-7).
- (2020a): “Binary relations in mathematical economics: on the continuity, additivity and monotonicity postulates in Eilenberg, Villegas and DeGroot,” *arXiv preprint arXiv:2007.01952*.
- (2020b): “On an extension of a theorem of Eilenberg and a characterization of topological connectedness,” *Topology and its Applications*, 273, 107–117.
- KIM, T. (1987): “Intransitive indifference and revealed preference,” *Econometrica*, pp. 163–167.
- KIM, T., AND M. K. RICHTER (1986): “Nontransitive-nontotal consumer theory,” *Journal of Economic Theory*, 38(2), 324–363.
- KLEE, V. (1951): “Convex sets in linear spaces,” *Duke Mathematical Journal*, 18(2), 443–466.

- KOOPMANS, T. C. (1957): *Three Essays on the State of Economic Science*. New York: McGraw-Hill.
- (1970): *Scientific Papers of Tjalling C. Koopmans*, vol. I, Martin Beckmann, Carl F. Christ and Marc Nerlove (eds.). Berlin: Springer-Verlag.
- (1972): “Representation of preference orderings with independent components of consumption,” in *Decision and Organization*, ed. by C. B. McGuire, and R. Radner, pp. Chapter 4, 79–100. Minneapolis: University of Minnesota Press.
- (1985): *Scientific Papers of Tjalling C. Koopmans*, vol. II. Massachusetts: MIT Press.
- KOPYLOV, I. (2016): “Canonical utility functions and continuous preference extensions,” *Journal of Mathematical Economics*, 67, 32–37.
- KRANTZ, D., D. LUCE, P. SUPPES, AND A. TVERSKY (1971): *Foundations of Measurement, Volume I: Additive and Polynomial Representations*. New York: Academic Press.
- KRUSE, R., AND J. DEELY (1969): “Joint continuity of monotonic functions,” *The American Mathematical Monthly*, 76(1), 74–76.
- KUCZMA, M. (2009): *An Introduction to the Theory of Functional Equations and Inequalities: Cauchy’s Equation and Jensen’s Inequality*. Berlin: Birkhäuser, 2 edn.
- KULSTAD, M. (1974): “Two arguments on *petites perceptions*,” *Rice University Studies*, 63(1), 57–68.
- (1991): *Leibniz on Apperception, Consciousness, and Reflection*. Munich: Philosophia.
- LECAM, L. (1986): *Asymptotic Methods in Statistical Decision Theory*. Berlin: Springer-Verlag.
- LEE, L.-F. (1972): “The theorems of Debreu and Peleg for ordered topological spaces,” *Econometrica*, 40(6), 1151–1153.
- LLOYD, C., R. ROHR, AND M. WALKER (1967): “A calculus proof of the existence of a continuous utility function,” *Metroeconomica*, 19(2), 103–112.
- MACCHERONI, F., M. MARINACCI, AND A. RUSTICHINI (2006): “Ambiguity aversion, robustness, and the variational representation of preferences,” *Econometrica*, 74(6), 1447–1498.
- MANCOSU, P. (1996): *Philosophy of Mathematics and Mathematical Practice in the Seventeenth Century*. Oxford: Oxford University Press.
- MARSCHAK, J. (1950): “Rational behavior, uncertain prospects, and measurable utility,” *Econometrica*, 18(2), 111–141.
- MCCARTHY, D., AND K. MIKKOLA (2018): “Continuity and completeness of strongly independent preorders,” *Mathematical Social Sciences*, 93, 141–145.
- McKENZIE, L. W. (2002): *Classical General Equilibrium Theory*, vol. 1. Cambridge: The MIT Press.
- MEHTA, G. (1983): “Ordered topological spaces and the theorems of Debreu and Peleg,” *Indian Journal of Pure and Applied Mathematics*, 14(9), 1174–1182.
- MITRA, T., AND M. K. OZBEK (2013): “On representation of monotone preference orders in a sequence space,” *Social Choice and Welfare*, 41, 1–15.
- NACHBIN, L. (1965): *Topology and Order*, vol. 4. New Jersey: Krieger Pub Co.

- NAKAMURA, Y. (1990): “Subjective expected utility with non-additive probabilities on finite state spaces,” *Journal of Economic Theory*, 51(2), 346–366.
- NEUEFEIND, W., AND W. TROCKEL (1995): “Continuous linear representability of binary relations,” *Economic Theory*, 6(2), 351–356.
- PELEG, B. (1970): “Utility functions for partially ordered topological spaces,” *Econometrica*, 38(1), 93–96.
- REBENCIUC, M. (2008): “Binary relations – addenda 1: Kernel, restrictions and inducing, relational morphisms,” *UPB, Sci. Bull. Series A*, 70(3), 11–22.
- RENY, P. J. (2020): “Nash equilibrium in discontinuous games,” *Annual Review of Economics*, 12, 439–470.
- REYES, M. (2004): “The rhetoric in mathematics: Newton, Leibniz, the calculus, and the rhetorical force of the infinitesimal,” *Quarterly Journal of Speech*, 90(2), 163–188.
- ROCKAFELLAR, R. T. (1970): *Convex Analysis*. New York: Princeton University Press.
- ROSENTHAL, A. (1955): “On the continuity of functions of several variables,” *Mathematische Zeitschrift*, 63(1), 31–38.
- SAVAGE, L. J. (1954): *The Foundations of Statistics*. Wiley, New York (Second edition 1972, Dover, New York).
- SCHAEFER, H. H. (1966): *Topological Vector Spaces*. New York: Springer-Verlag.
- SCHERVISHA, M. J., T. SEIDENFELD, R. GONG, J. B. KADANE, AND R. B. STERN (2020): “When no price is right,” *working paper*.
- SCHMEIDLER, D. (1969): “Competitive equilibria in markets with a continuum of traders and incomplete preferences,” *Econometrica*, 37(4), 578–585.
- (1971): “A condition for the completeness of partial preference relations,” *Econometrica*, 39(2), 403–404.
- SCHOENBERG, I. J. (1942): “On local convexity in Hilbert space,” *Bulletin of the American Mathematical Society*, 48(6), 432–436.
- SHAFFER, W. (1974): “The nontransitive consumer,” *Econometrica*, 42(5), 913–919.
- SHAPLEY, L. S., AND M. BAUCCELLS (1998): “Multiperson utility,” Discussion paper, UCLA Economics Working Paper 779, UCLA Department of Economics.
- SONNENSCHN, H. (1965): “The relationship between transitive preference and the structure of the choice space,” *Econometrica*, 33(3), 624–634.
- (1967): “Reply to “a note on orderings”,” *Econometrica*, 35(3, 4), 540.
- STANCA, L. (2020): “A simplified approach to subjective expected utility,” *Journal of Mathematical Economics*, 87, 151–160.
- UYANIK, M., AND M. A. KHAN (2019): “On the consistency and the decisiveness of the double-minded decision-maker,” *Economics Letters*, 185, 108657, 1–5.
- VILLEGAS, C. (1964): “On qualitative probability  $\sigma$ -algebras,” *The Annals of Mathematical Statistics*, 35(4), 1787–1796. Mathematical Reviews (R. V. Chacon): MR167588.

- VON NEUMANN, J., AND O. MORGENSTERN (1947): *Theory of Games and Economic Behavior*. New Jersey: Princeton University Press, 2nd edn.
- WAKKER, P. (1988a): "The algebraic versus the topological approach to additive representations," *Journal of Mathematical Psychology*, 32(4), 421–435.
- (1988b): "Continuity of preference relations for separable topologies," *International Economic Review*, 29(1), 105–110.
- (1989): *Additive Representations of Preferences: A New Foundation of Decision Analysis*. Boston: Kluwer Academic Publishers.
- (1993): "Additive representations on rank-ordered sets: II. The topological approach," *Journal of Mathematical Economics*, 22(1), 1–26.
- WARD, L. (1954): "Partially ordered topological spaces," *Proceedings of the American Mathematical Society*, 5(1), 144–161.
- WEINER, P. P. E. (1951): *Leibniz Selections*. New York: Charles Scribner's Sons.
- WOLD, H. (1943–44): "A synthesis of pure demand analysis, I–III," *Scandinavian Actuarial Journal*, 26, 85–118, 220–263; 27, 69–120.
- WOLD, H., AND L. JUREEN (1953): *Demand Analysis*. New York: John Wiley and Sons, Inc.
- YAMAZAKI, A. (1983): "On open preferences," *Hitotsubashi Journal of Economics*, 24, 149–152.
- YI, G. (1993): "Continuous extension of preferences," *Journal of Mathematical Economics*, 22(6), 547–555.
- YOKOYAMA, T. (1954): "On uniformity and continuity condition in the theory of consumer's choice," *Osaka Economic Papers*, 3, 29–35.
- (1956): "Continuity conditions of preference ordering," *Osaka Economic Papers*, 4, 39–45.
- YOUNG, W. (1910): "A note on monotone functions," *The Quarterly Journal of Pure and Applied Mathematics (Oxford Ser.)*, 41, 79–87.
- YOUNG, W. H., AND G. C. YOUNG (1910): "Discontinuous functions continuous with respect to every straight line," *Quarterly Journal of Pure and Applied Mathematics*, 41, 87–93.