

MOTIVIC COHOMOLOGY OF FAT POINTS IN MILNOR RANGE VIA FORMAL AND RIGID GEOMETRIES

JINHYUN PARK

ABSTRACT. We present a formal scheme based cycle model for the motivic cohomology of the fat points defined by the truncated polynomial rings $k[t]/(t^m)$ with $m \geq 2$, in one variable over a field k . We compute their Milnor range cycle class groups when the field has sufficiently many elements.

With some aids from rigid analytic geometry and the Gersten conjecture for the Milnor K -theory resolved by M. Kerz, we prove that the resulting cycle class groups are isomorphic to the Milnor K -groups of the truncated polynomial rings, generalizing a theorem of Nesterenko-Suslin and Totaro.

CONTENTS

1. Introduction	1
2. Some definitions and recollections	6
3. On the generator cycles	14
4. The graph homomorphisms and the regulators	21
5. The proofs of the main theorems	33
References	37

1. INTRODUCTION

In this article, we propose a new cycle model through formal geometry for the motivic cohomology of the singular non-reduced scheme $Y := \text{Spec}(k_m)$, where $k_m := k[t]/(t^m)$, and we compute its Milnor range for all base fields k except a finite number of finite fields, using some ideas from rigid analytic geometry. The main theorem is summarized in Theorem 1.2.1 below.

1.1. Background and motivation. The motivic cohomology of smooth k -schemes is given by higher Chow groups of S. Bloch [5], but the higher Chow complexes fail to give correct models of motivic cohomology when the schemes have singularities.

For such non-smooth schemes, approaches via additive higher Chow groups ([6], [38], [42]) and cycles with modulus ([4]) were proposed as part of potential extensions. Various theoretical advances were made (e.g. [25], [26], [27], [32]) in this direction. All of them use the idea of “modulus conditions” imposed on cycles.

Recently in [41], the author and S. Ünver had tried a very different approach for $Y = \text{Spec}(k_m)$. Embedding $\text{Spec}(k_m)$ into the henselian scheme $\text{Spec}(k[[t]])$, the article *ibid.* considered certain cycles on $\text{Spec}(k[[t]]) \times \square_k^n$ subject to a few requirements, where $\square_k := \mathbb{P}^1 \setminus \{1\}$. On the cycles, the *ibid.* puts a “mod t^m equivalence

2020 *Mathematics Subject Classification.* Primary 14C25; Secondary 19D45, 13F25, 14B20, 16W60.

Key words and phrases. algebraic cycle, Chow group, singular scheme, formal scheme, Tate algebra, rigid geometry, motivic cohomology, Milnor K -theory, algebraic de Rham cohomology, de Rham-Witt form.

relation”, and the homology group of the resulting complex in the Milnor range was identified with the Milnor K -group $K_n^M(k_m)$ when k is of characteristic 0.

The author had received a few feedbacks since then. One of them from Fumiharu Kato at a social meeting in Tokyo, Japan in 2019 was that, for future generalizations to general singular schemes, it may be better to use formal schemes. For instance, instead of the scheme $\text{Spec}(k[[t]])$, the formal scheme $\widehat{X} = \text{Spf}(k[[t]])$ is the formal neighborhood of the immersion $Y = \text{Spec}(k_m) \hookrightarrow \mathbb{A}^1$, so Kato’s suggestion seems to be naturally consistent with the philosophy of the construction of the algebraic de Rham cohomology in A. Grothendieck [18] and R. Hartshorne [20].

Once one accepts to try this perspective via formal schemes, it becomes necessary to ask what the right notions of (higher Chow) cycles for formal schemes are. To have a higher Chow cycle-stye theory, for instance, one should take the fiber product $\text{Spf}(k[[t]]) \times_k \square_k^n$ in the category of formal schemes, instead of the fibre product $\text{Spec}(k[[t]]) \times_k \square_k^n$ of schemes. Here, one needs to answer what requirements one should impose on the closed formal subschemes to define the right theory.

The first nontrivial test for a potential theory is to see whether its homology in the Milnor range for Y is the Milnor K -group $K_n^M(k_m)$. One far harder test is to show that it extends to a functorial theory on the category of schemes of finite type over k , where on the smooth schemes, it coincides with the classical higher Chow theory.

The objective of this article is to prove that the theory we build here passes the first test in the Milnor range. The above harder test of constructing a functorial theory is studied in the sequel [39]. A related discussion on algebraic K -theory of singular schemes is studied in [40].

1.2. The central result. More specifically, we do the following in this article. Consider the formal scheme $\widehat{X} = \text{Spf}(k[[t]])$, with the closed immersion $Y := \text{Spec}(k_m) \hookrightarrow \widehat{X}$. Consider the formal scheme $\widehat{X} \times_k \square_k^n$.

We will construct (Definition 2.4.5) a complex of abelian groups

$$(1.2.1) \quad \cdots \rightarrow z^q(\widehat{X} \bmod Y, n+1) \xrightarrow{\partial} z^q(\widehat{X} \bmod Y, n) \xrightarrow{\partial} z^q(\widehat{X} \bmod Y, n-1) \rightarrow \cdots,$$

where $z^q(\widehat{X} \bmod Y, n)$ is the quotient

$$(1.2.2) \quad z^q(\widehat{X} \bmod Y, n) = \frac{z^q(\widehat{X}, n)}{\mathcal{M}^q(\widehat{X}, Y, n)}$$

of the group $z^q(\widehat{X}, n)$ consisting of certain “admissible cycles” of codimension q on the formal scheme $\widehat{X} \times_k \square_k^n$ modulo a subgroup $\mathcal{M}^q(\widehat{X}, Y, n)$ defined using the embedding $Y \hookrightarrow \widehat{X}$. The constructions of the groups $z^q(\widehat{X}, n)$ and $\mathcal{M}^q(\widehat{X}, Y, n)$ are rapidly sketched in §1.3 below. They are defined in full in Definitions 2.2.2 and 2.4.1. The homology $H_n(z^q(\widehat{X} \bmod Y, \bullet))$ of the complex (1.2.1) is denoted by any one of

$$\mathbf{CH}^q(k_m, n) = \mathbf{CH}^q(Y, n) = \mathbf{CH}^q(\widehat{X} \bmod Y, n) = H_n(z^q(\widehat{X} \bmod Y, \bullet)),$$

where the first two are written in boldface to distinguish them from the usual higher Chow group [5]. The central result of the article is to show that, under a mild assumption on the field k , we have an isomorphism $K_n^M(k_m) \simeq \mathbf{CH}^q(k_m, n)$. The following (from Proposition 2.4.6 and Theorem 5.2.6) gives more details. This includes a generalization to $\text{Spec}(k_m)$ of theorems of Nesterenko-Suslin [37] and B. Totaro [47] proven for $\text{Spec}(k)$:

Theorem 1.2.1. *Let $m \geq 2, n \geq 1$ be integers, k be a field, and $k_m = k[t]/(t^m)$.*

- (1) If $q > n$, then we have $\mathbf{CH}^q(\mathrm{Spec}(k_m), n) = 0$.
(2) If $q = n$, there is a homomorphism, called the graph map

$$(1.2.3) \quad gr_{k_m} : K_n^M(k_m) \rightarrow \mathbf{CH}^n(\mathrm{Spec}(k_m), n),$$

and it is an isomorphism of abelian groups when the cardinality $|k|$ is sufficiently large.

The phrase ‘‘sufficiently large’’ in Theorem 1.2.1 means that the cardinality of the field k is larger than a positive integer M_n given by M. Kerz [30, Proposition 10-(5), p.181] so that the Milnor K -theory $K_n^M(k[[t]])$ and the improved Milnor K -theory $\widehat{K}_n^M(k[[t]])$ of Gabber-Kerz coincide. For instance, when $n = 1$, this holds for an arbitrary field k .

1.3. The cycles. Recall $\widehat{X} = \mathrm{Spf}(k[[t]])$ and $Y = \mathrm{Spec}(k_m)$. We sketch the construction of $z^q(\widehat{X} \bmod Y, n)$ of (1.2.2). The formal scheme \widehat{X} has the largest ideal of definition \mathcal{I}_0 given by (t) . The reduction by \mathcal{I}_0 is $\widehat{X}_{\mathrm{red}} = \mathrm{Spec}(k)$.

1.3.1. The two requirements. The group $z^q(\widehat{X}, n)$ is (Definition 2.2.2) the free abelian group generated by the integral closed formal subschemes $\mathfrak{Z} \subset \widehat{X} \times_k \square_k^n$ of codimension q , satisfying two conditions **(GP)** and **(SF)**.

The general position property **(GP)** requires that the intersections of \mathfrak{Z} with all faces $\widehat{X} \times F$ are proper, where $F \subset \square_k^n$ are given by a finite set of equations of the form $\{y_i = \epsilon_i\}$ with $\epsilon_i \in \{0, \infty\}$. The special fiber property **(SF)** requires that the intersections of \mathfrak{Z} with all special fiber faces $\widehat{X}_{\mathrm{red}} \times F = \mathrm{Spec}(k) \times F$ are proper.

1.3.2. The mod Y -equivalence. We discuss the subgroup $\mathcal{M}^q(\widehat{X}, Y, n) \subset z^q(\widehat{X}, n)$ of (1.2.2), which defines the mod Y -equivalence on the cycles.

This subgroup is generated by the cycles of the form

$$[\mathcal{A}_1] - [\mathcal{A}_2],$$

where $(\mathcal{A}_1, \mathcal{A}_2)$ runs over all possible pairs of coherent $\mathcal{O}_{\widehat{X} \times \square^n}$ -algebras whose associated cycles belong to $z^n(\widehat{X}, n)$, such that there is an isomorphism

$$(1.3.1) \quad \mathcal{A}_1|_{Y \times \square^n} \simeq \mathcal{A}_2|_{Y \times \square^n}$$

as $\mathcal{O}_{Y \times \square^n}$ -algebras (see Definition 2.4.1).

1.4. Benefits of the formal geometry model. One point of using the formal scheme model in studying the motivic cohomology of $\mathrm{Spec}(k[t]/(t^m))$ is its natural generalizability to more general schemes Y of finite type in \mathbf{Sch}_k . This aspect is studied in [39] in terms of some ideas of the derived algebraic geometry of J. Lurie.

Yet, there is another aspect that we mention, compared to the scheme model in [41]. In *ibid.*, the generating cycles in the Milnor range were integral closed subschemes in $\mathrm{Spec}(k[[t]]) \times \square_k^n$ subject to the three requirements: (i) the proper intersection with the faces, (ii) the proper intersection with the special fiber times faces, and (iii) the extra requirement that each integral cycle is *proper* over $k[[t]]$, thus finite over $k[[t]]$.

The first two conditions (i) and (ii) correspond to our requirements **(GP)** and **(SF)**, respectively. However, to exclude potentially undesirable cycles, the extra condition (iii) was necessary there. One can ask what happens to this condition in our formal scheme model.

One interesting aspect we observe in the formal scheme model is that, in fact, in the Milnor range we *automatically* have the finiteness of the generating integral cycles over $k[[t]]$. This surprising conclusion in the formal scheme model is deduced

in §3.2, especially in Corollary 3.2.3. This argument does not work in the scheme model, as we really exploit the situation in the formal geometry.

In connection to this, in Example 2.2.6, we consider an integral closed cycle $\mathfrak{Z}_1 \subset \text{Spec}(k[[t]]) \times \square_k^2$ given by two concrete equations, such that it satisfies the two conditions (i), (ii) but not (iii); \mathfrak{Z}_1 is not finite over $k[[t]]$. It is forcibly removed in the scheme model by (iii).

In contrast, the corresponding formal scheme $\mathfrak{Z}_2 \subset \text{Spf}(k[[t]]) \times \square_k^2$ given by the same equations is empty in the formal scheme model, thanks to the convergence of a sequence in the (t) -adic topology.

This shows the formal scheme model of this article has various stronger points over the scheme model, e.g. of [41].

1.5. Connections to rigid geometry. We explain how and where the rigid analytic geometry enters into the landscape of algebraic cycles.

Using the automorphism $y \mapsto y/(y-1)$ of \mathbb{P}^1 , identify \square^1 with \mathbb{A}^1 . So for $y'_i := y_i/(y_i-1)$ where $(y_1, \dots, y_n) \in \square^n$, the ambient space $\widehat{X} \times \square_k^n$ can be seen as the formal spectrum of the ring $k[[t]]\{y'_1, \dots, y'_n\}$ of the *restricted formal power series*, i.e. the formal power series $\sum_I a_I y'^I$ in y'_1, \dots, y'_n such that the coefficients $a_I \in k[[t]]$ converge to 0 in the (t) -adic norm as the degree $|I| \rightarrow \infty$ (see Example 2.2.1).

Here, not all prime ideals of $k[[t]]\{y'_1, \dots, y'_n\}$ are suitable for our cycles. In (SF) in §1.3.1 or Definition 2.2.2, we ruled out those behave badly with respect to the ideals of definition of $\widehat{X} \times_k \square_k^n$. Those ideals give the “special fibers” $\text{Spec}(k) \times_k F$ for faces $F \subset \square_k^n$, and the condition (SF) requires the proper intersection of the cycles with such special fibers.

In particular, via localization to the “generic fibers”, our admissible prime ideals of height n in the Milnor range define prime ideals of height n of the Tate algebra $T_n = k((t))\{y'_1, \dots, y'_n\}$ over the non-archimedean complete discrete valued field $k((t))$.

The Tate algebras were introduced by J. Tate in [46]. Since T_n has the Krull dimension n , the height n primes are its maximal ideals. They define finite dimensional Banach algebras over $k((t))$. One can show that in T_n , all such maximal ideals are complete intersections given by n *polynomials* in y'_1, \dots, y'_n . Thus, taking closures, we deduce algebraic generators for the integral admissible cycles in $z^n(\widehat{X}, n)$. This is covered in §3.1. This allows us to see in §3.2 that each integral cycle \mathfrak{Z} can be written as $\text{Spf}(R)$, where $R = k[[t]]\{y_1, \dots, y_n\}/P$ for some prime ideal, and it is finite over $k[[t]]$.

Another place, where the rigid geometry plays a role, is in the middle of the proof that the graph map (1.2.3) is an isomorphism.

As a first (eventually wrong) attempt, one might try to “construct” its inverse

$$(1.5.1) \quad \mathbf{CH}^n(\text{Spec}(k_m), n) \rightarrow K_n^M(k_m),$$

as follows: let \bar{y}_i be the image of y_i in R . Then one may try to define

$$\rho : z^n(\text{Spf}(k[[t]]), n) \rightarrow K_n^M(k[[t]])$$

by sending an integral cycle \mathfrak{Z} to the norm $N_{R/k[[t]]}\{\bar{y}_1, \dots, \bar{y}_n\} \in K_n^M(k[[t]])$ of the Milnor symbol $\{\bar{y}_1, \dots, \bar{y}_n\} \in K_n^M(R)$. Then one may try to show that its composition with the natural surjection $K_n^M(k[[t]]) \rightarrow K_n^M(k[t]/(t^m))$ kills the boundary $\partial z^n(\text{Spf}(k[[t]]), n+1)$ as well as the subgroup $\mathcal{M}^n(\text{Spf}(k[[t]]), Y, n)$.

Unfortunately, the above proposed story does not work well. The problem is that we *do not* know whether there is the norm map $N_{R/k[[t]]} : K_n^M(R) \rightarrow K_n^M(k[[t]])$ unless $k[[t]] \rightarrow R$ is finite étale (see M. Kerz [30, Propositions 3, 10]). In general, its existence is unknown at present as far as the author is aware of.

To get around this difficulty, we take a detour through the rigid geometry via the generic fiber of the cycles, and use the Gersten conjecture for Milnor K -theory proven by M. Kerz in *ibid.*

We first (see §4.2) construct the composite

$$\widehat{\Upsilon} : z^n(\mathrm{Spf}(k[[t]]), n) \xrightarrow{\eta} \mathrm{CH}^n(\mathrm{Sp}(k((t))), n) \xrightarrow{\tilde{\Upsilon}} K_n^M(k((t))),$$

where the middle term is a rigid analytic analogue of the higher Chow group (see Definition 4.2.1), η is the localization map, and $\tilde{\Upsilon}$ is constructed using the norm map of Bass-Tate [3] and Kato [28] on the Milnor K -theory for *fields*. We check that the image of $\widehat{\Upsilon}$ belongs to the *subgroup* $K_n^M(k[[t]])$ of $K_n^M(k((t)))$, which requires the Gersten conjecture for K_n^M of M. Kerz [30]. The rest of the paper is, roughly speaking, about proving that this map descends mod t^m , and that gr_{k_m} in (1.2.3) is an isomorphism.

These offer an interesting new observation: the motivic cohomology of the truncated polynomials can be partially understood through the rigid analytic geometry. Already in [41], the topology given by the non-archimedean (t) -adic norm was very convenient. The author hopes that the new door opened here could be useful for the researchers in the field.

1.6. The relative parts. Taking the relative parts of the groups of Theorem 1.2.1, we deduce the following result, stated as Theorem 5.2.7:

Theorem 1.6.1. *Let $m \geq 1, n \geq 1$ be integers. Let k be a field of characteristic 0.*

Then we have an isomorphism from the group of the big de Rham-Witt forms

$$(1.6.1) \quad \mathbb{W}_m \Omega_k^{n-1} \xrightarrow{\sim} \mathbf{CH}^n((\mathrm{Spec}(k_{m+1}), (t)), n)$$

to the relative group of the new higher Chow group.

While we guess that the statement of Theorem 1.6.1 may stay valid for any field k , at this moment the author was unable to give a proof of it without imposing the assumption on the characteristic. It should be removed in the future. The group of the big de Rham-Witt forms on the left of (1.6.1) is defined by Hesselholt-Madsen [24], while the relative group on the right of (1.6.1) is defined to be (see (5.2.4))

$$\ker(\mathbf{CH}^n(\mathrm{Spec}(k_{m+1}), n) \xrightarrow{ev_0} \mathrm{CH}^n(k, n)),$$

where $\mathrm{CH}^n(k, n)$ is the usual higher Chow group of S. Bloch [5].

This Theorem 1.6.1 is an analogue of the theorem of K. Rülling [42] proven in terms of additive higher Chow groups (see also [32] for its higher dimensional generalization). An interesting point is that our theory in this article provides a single unified platform for all of Nesterenko-Suslin [37], Totaro [47] and Rülling [42], and it is also relatively conceptual compared to the approaches through “cycles with modulus” in the literature.

Conventions. In this paper k is a given arbitrary field, unless said otherwise. For the set inclusion symbol \subset , this allows the equality $=$ as well. The strict inclusion is denoted by \subsetneq , while we won’t use the symbol \subseteq .

The notations k_m, Y, X, \widehat{X} in this article exclusively mean the following: for an integer $m \geq 1$, we let $k_m := k[t]/(t^m) = k[[t]]/(t^m)$, $Y = \mathrm{Spec}(k_m)$, $Y \hookrightarrow X = \mathbb{A}^1$ is the closed immersion given by $k[t] \rightarrow k_m$, and \widehat{X} is the completion of X along Y , so that $\widehat{X} = \mathrm{Spf}(k[[t]])$. \square

2. SOME DEFINITIONS AND RECOLLECTIONS

In §2, we discuss the notion of higher Chow cycles on some noetherian affine formal k -schemes of finite Krull dimension. Based on these, in §2.4 we define the main objects of studies in the article, the groups $\mathbf{CH}^q(k_m, n)$. In §2.5, we also discuss basic definitions and results on Tate algebras, that will be used later to compute the Milnor range $\mathbf{CH}^n(k_m, n)$, when $q = n$.

2.1. Noetherian affine formal schemes and cycles. Let $\mathfrak{X} = \mathrm{Spf}(A)$ be a noetherian affine formal scheme of finite Krull dimension.

Definition 2.1.1. An *integral closed formal subscheme* $\mathfrak{Y} \subset \mathfrak{X}$ is of the form $\mathrm{Spf}(A/P)$ for a prime ideal $P \subset A$.

The *naive group* of algebraic cycles $\underline{z}_*(\mathfrak{X})$ on \mathfrak{X} is defined to be the free abelian group on such integral closed formal subschemes. This is equivalent to the group of algebraic cycles on the *scheme* $\mathrm{Spec}(A)$, that we call the associated scheme of \mathfrak{X} . Consequently, we have the same notion of intersections of cycles, and proper intersections of them.

Its subgroup of d -dimensional cycles is denoted by $\underline{z}_d(\mathfrak{X})$. In case \mathfrak{X} is equidimensional (i.e. A is equidimensional), we denote the group of the codimension q -cycles by $\underline{z}^q(\mathfrak{X})$.

Not all of the cycles in $\underline{z}_*(\mathfrak{X}) = z_*(\mathrm{Spec}(A))$ are suitable for our discussions. A better one is the subgroup $z_*(\mathfrak{X}) \subset \underline{z}_*(\mathfrak{X})$ of cycles that intersect properly with $\mathfrak{X}_{\mathrm{red}}$. This $\mathfrak{X}_{\mathrm{red}}$ is the closed subscheme of \mathfrak{X} given by the largest ideal of definition, which exists by EGA I [16, Proposition (10.5.4), p.187]. \square

When \mathcal{F} is a coherent $\mathcal{O}_{\mathfrak{X}}$ -module, we have the associated cycle $[\mathcal{F}] \in \underline{z}_*(\mathfrak{X})$ via minimal associated primes (see e.g. [44, Section 02QV]). This is possible because $\mathcal{F} = M^\Delta$ for a finitely generated A -module M (see EGA I [16, Proposition (10.10.2), p.201]), and the associated cycle of M on $\mathrm{Spec}(A)$ defines $[\mathcal{F}]$.

2.2. Higher Chow cycles over affine formal schemes. Let $\square_k := \mathbb{P}_k^1 \setminus \{1\}$. Let $\square_k^0 := \mathrm{Spec}(k)$. For $n \geq 1$, let \square_k^n be the n -fold fiber product of \square_k over k . Let $(y_1, \dots, y_n) \in \square_k^n$ be the coordinates.

Recall that in the cubical version of higher Chow theory, using the k -rational points $\{0, \infty\} \subset \square_k$, we define a face $F \subset \square_k^n$ to be a closed subscheme given by a system of equations of the form $\{y_{i_1} = \epsilon_1, \dots, y_{i_s} = \epsilon_s\}$ for an increasing sequence $1 \leq i_1 < \dots < i_s \leq n$ of indices and $\epsilon_j \in \{0, \infty\}$. When the set of equations is empty, we let $F = \square_k^n$ by convention. For $s = 1$, we have the codimension 1 faces given by $\{y_i = \epsilon\}$ for some $1 \leq i \leq n$ and $\epsilon \in \{0, \infty\}$.

For an equidimensional noetherian affine formal k -scheme \mathfrak{X} of finite Krull dimension, consider the fiber product $\mathfrak{X} \times_k \square_k^n$, which we often write $\square_{\mathfrak{X}}^n$. The fiber product exists in the category of formal schemes; see EGA I [16, Proposition (10.7.3), p.193]. A face $F_{\mathfrak{X}}$ of $\square_{\mathfrak{X}}^n$ is given by $\mathfrak{X} \times_k F$ for a face $F \subset \square_k^n$.

Example 2.2.1. Suppose $\mathfrak{X} = \mathrm{Spf}(A)$. Using the automorphisms of \mathbb{P}^1 given by $y_i \mapsto y'_i := y_i/(y_i - 1)$ for $1 \leq i \leq n$, we may identify $\square_{\mathfrak{X}}^n$ with the integral noetherian affine formal k -scheme $\mathfrak{X} \times_k \mathbb{A}_k^n$ given by the ring $A\{y'_1, \dots, y'_n\}$ of restricted formal power series.

In case $A = k[[t]]$ with the (t) -adic topology, this ring is

$$k[[t]]\{y'_1, \dots, y'_n\} = \varprojlim_m k_m[y'_1, \dots, y'_n].$$

This ring can also be regarded as the subring of $k[[t]][[y'_1, \dots, y'_n]]$, whose members are formal power series

$$p(y'_1, \dots, y'_n) = \sum_I \alpha_I y'^I$$

with $\alpha_I \in k[[t]]$ for multi-indices $I = (i_1, \dots, i_n)$ with $i_j \geq 0$, such that when $|I| = i_1 + \dots + i_n \rightarrow \infty$, we have $\alpha_I \rightarrow 0$ in the (t) -adic topology. \square

We define cubical higher Chow cycles over the formal scheme \mathfrak{X} (cf. [5]) as follows.

Definition 2.2.2. Let \mathfrak{X} be an equidimensional noetherian affine formal k -scheme of finite Krull dimension. Let $q, n \geq 0$ be integers. Let $\mathfrak{X}_{\text{red}}$ be the closed subscheme given by the largest ideal \mathcal{I}_0 of definition of \mathfrak{X} .

Let $\underline{z}^q(\mathfrak{X}, n) \subset \underline{z}^q(\square_{\mathfrak{X}}^n)$ be the subgroup generated by the codimension q integral closed formal subschemes $\mathfrak{Z} \subset \square_{\mathfrak{X}}^n$ subject to the following conditions:

- (**GP**) (General Position) For each face $F \subset \square_k^n$, \mathfrak{Z} intersects $\mathfrak{X} \times F$ properly.
- (**SF**) (Special Fiber) For each face $F \subset \square_k^n$, we have

$$\text{codim}_{\mathfrak{X}_{\text{red}} \times F} (\mathfrak{Z} \cap (\mathfrak{X}_{\text{red}} \times F)) \geq q.$$

A cycle in $\underline{z}^q(\mathfrak{X}, n)$ will be called *admissible*. \square

We have several remarks on the conditions we introduced in Definition 2.2.2.

Remark 2.2.3. One notes that in addition to the usual general position condition (the condition (**GP**)) with respect to the faces for higher Chow cycles, we require the additional condition (**SF**), which is related to the “special fiber” given by the largest ideal of definition of \mathfrak{X} . In case \mathfrak{X} is a *scheme*, regarded as a formal scheme, the requirement (**SF**) holds automatically, so that Definition 2.2.2 is compatible with the definition of the cubical version of the higher Chow cycles of [5], e.g. in [47]. \square

Remark 2.2.4. Note that by the special fiber condition (**SF**), the cycles given by open prime ideals of $A\{y'_1, \dots, y'_n\}$ are all excluded from $\underline{z}^q(\mathfrak{X}, n)$, where $\mathfrak{X} = \text{Spf}(A)$ and $y'_i := y_i/(y_i - 1)$ for $1 \leq i \leq n$ as before. \square

We make one elementary observation in Lemma 2.2.5 below. In an earlier version of the article, an analogue of this property was considered as part of the requirements in Definition 2.2.2, called the topological support condition (**TS**). Joseph Ayoub as well as the referee pointed out that, in fact, each integral cycle satisfies it automatically, so that it is redundant.

Lemma 2.2.5. *Let \mathfrak{X} be a noetherian formal scheme and let $\mathfrak{Z} \subset \mathfrak{X}$ be an integral closed formal subscheme. Let \mathcal{I} be an ideal of definition of \mathfrak{X} and let Z be the reduction of \mathfrak{Z} modulo \mathcal{I} . Then we have the equality of the topological supports $|\mathfrak{Z}| = |Z|$.*

Proof. Let $\mathcal{J} \subset \mathcal{O}_{\mathfrak{X}}$ be the ideal sheaf of \mathfrak{Z} . The ideal sheaf of Z is given by $\mathcal{I} + \mathcal{J}$. However an ideal of definition of \mathfrak{Z} is also given by $\mathcal{I} + \mathcal{J}$. \square

Example 2.2.6. When $\mathfrak{X} = \text{Spf}(A)$ for an adic ring A , one may ask whether we can consider the scheme $\text{Spec}(A) \times \square^n$ (that uses polynomials) instead of the formal scheme as we do here (that uses restricted formal power series).

Here is an example that shows why our formal scheme based model via $\text{Spf}(k[[t]]) \times \square_k^n$ considered in this article is better than the scheme based model via $\text{Spec}(k[[t]]) \times \square_k^n$, e.g. considered in [41]. The author thanks Joseph Ayoub and the referee for sharing their insights and encouraging him to try to write it down.

In the scheme model, we may have some undesirable cycles such as (here, $n = 2$ and $(y_1, y_2) \in \square^2$)

$$\mathfrak{Z}_1 : \{y_1 = 1 + t, \quad y_2 = t\},$$

which is not finite over $k[[t]]$. This is part of the reason why in [41], the extra finiteness requirement was imposed to exclude forcibly this kind of cycles.

On the other hand, in the formal scheme model of ours, the corresponding closed formal subscheme $\mathfrak{Z}_2 \subset \mathrm{Spf}(k[[t]]) \times \square_k^2$ given by the same defining equations,

$$\mathfrak{Z}_2 : \{y_1 = 1 + t, \quad y_2 = t\},$$

is empty, thus it vanishes itself.

To see this, applying the automorphism $y \mapsto y/(y-1)$ of \mathbb{P}^1 that sends $(\square, \{0, \infty\})$ to $(\mathbb{A}^1, \{0, 1\})$, we have a closed formal subscheme of $\mathrm{Spf}(k[[t]]\{y_1, y_2\})$ defined by

$$\left\{ \frac{y_1}{y_1 - 1} = 1 + t, \quad \frac{y_2}{y_2 - 1} = t \right\},$$

which gives

$$(2.2.1) \quad \{1 - (y_1 - 1)t = 0, \quad y_2 = (y_2 - 1)t\}.$$

However, the left hand side $1 - (y_1 - 1)t$ of (2.2.1) is a unit in $k[[t]]\{y_1, y_2\}$ because the power series

$$1 + (y_1 - 1)t + (y_1 - 1)^2 t^2 + \cdots + (y_1 - 1)^i t^i + \cdots$$

is convergent in $k[[t]]\{y_1, y_2\}$ and it is the inverse of $1 - (y_1 - 1)t$. In particular, the first equation of (2.2.1) cannot hold. Hence \mathfrak{Z}_2 in $\mathrm{Spf}(k[[t]]) \times \square_k^2$ is empty, and such \mathfrak{Z}_2 is automatically eliminated in the formal scheme model, without forcibly imposing an extra condition as in the scheme model of [41].

Furthermore, later we will see that each integral admissible cycle in $\mathrm{Spf}(k[[t]]) \times \square_k^n$ in the Milnor range is always finite over $k[[t]]$ in Corollary 3.2.3. \square

Coming back to our definition, we want to form a cycle complex:

Definition 2.2.7. Suppose $n \geq 1$. For each $1 \leq i \leq n$ and each $\epsilon \in \{0, \infty\}$, let $\iota_i^\epsilon : \square_{\mathfrak{X}}^{n-1} \hookrightarrow \square_{\mathfrak{X}}^n$ be the closed immersion given by the single equation $\{y_i = \epsilon\}$. For each integral cycle $\mathfrak{Z} \in \underline{z}^q(\mathfrak{X}, n)$, define the face $\partial_i^\epsilon(\mathfrak{Z})$ to be the cycle associated to the intersection $\mathfrak{Z} \cap \{y_i = \epsilon\} = [(\iota_i^\epsilon)^* \mathcal{O}_{\mathfrak{Z}}]$. One defines $\partial := \sum_{i=1}^n (-1)^i (\partial_i^\infty - \partial_i^0)$ and by the usual cubical formalism, we check that $\partial^2 = 0$.

This gives a complex $(\underline{z}^q(\mathfrak{X}, \bullet), \partial)$ of abelian groups. The complex $(z^q(\mathfrak{X}, \bullet), \partial)$ is defined to be the quotient of $(\underline{z}^q(\mathfrak{X}, \bullet), \partial)$ by the subcomplex $\underline{z}^q(\mathfrak{X}, \bullet)_{\mathrm{degn}}$ of degenerate cycles generated by pull-backs by the projections $\square_{\mathfrak{X}}^n \rightarrow \square_{\mathfrak{X}}^{n-1}$ that drop one of the coordinates. We define $\mathrm{CH}^q(\mathfrak{X}, n)$ to be the n -th homology of the complex $(z^q(\mathfrak{X}, \bullet), \partial)$. \square

The group $\mathrm{CH}^q(\mathfrak{X}, n)$ is *not* the primary object of studies of the article. In this article where we mostly consider $\mathfrak{X} = \widehat{X} = \mathrm{Spf}(k[[t]])$, we still need to take a quotient of $z^q(\widehat{X}, \bullet)$ by a subcomplex that gives the “mod Y -equivalence”. It will be given in Definition 2.4.1.

We mention that in the case of $\widehat{X} = \mathrm{Spf}(k[[t]])$ the following Tor independence property holds (see also Corollary 2.2.10):

Lemma 2.2.8. *Let $Y = \mathrm{Spec}(k_m)$ and $\widehat{X} = \mathrm{Spf}(k[[t]])$. Let \mathcal{F} be a coherent sheaf on $\square_{\widehat{X}}^n$, whose associated cycle $[\mathcal{F}]$ belongs to $z^*(\widehat{X}, n)$.*

Then we have $\mathrm{Tor}_i^{\mathcal{O}_{\widehat{X}}^n}(\mathcal{F}, \mathcal{O}_{\square_Y^n}) = 0$ for $i > 0$.

Proof. As before, for $y'_i := y_i/(y_i - 1)$, note that we have the short exact sequence

$$0 \rightarrow k[[t]]\{y'_1, \dots, y'_n\} \xrightarrow{\times t^m} k[[t]]\{y'_1, \dots, y'_n\} \rightarrow k_m[y'_1, \dots, y'_n] \rightarrow 0,$$

which gives a free $\mathcal{O}_{\square_{\widehat{X}}^n}$ -resolution of $\mathcal{O}_{\square_Y^n}$:

$$0 \rightarrow \mathcal{O}_{\square_{\widehat{X}}^n} \xrightarrow{\times t^m} \mathcal{O}_{\square_{\widehat{X}}^n} \rightarrow \mathcal{O}_{\square_Y^n} \rightarrow 0.$$

Hence

$$\mathcal{F} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}}^{\mathbf{L}} \mathcal{O}_{\square_Y^n} = \mathcal{F} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \left(\mathcal{O}_{\square_{\widehat{X}}^n} \xrightarrow{\times t^m} \mathcal{O}_{\square_{\widehat{X}}^n} \right) = \left(\mathcal{F} \xrightarrow{\times t^m} \mathcal{F} \right),$$

where the objects in the above complexes are placed in the homological degrees +1 and 0, respectively. This shows that $\mathrm{Tor}_i^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{F}, \mathcal{O}_{\square_Y^n}) = 0$ for $i > 1$, while $\mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{F}, \mathcal{O}_{\square_Y^n}) = \ker(\times t^m : \mathcal{F} \rightarrow \mathcal{F})$.

It remains to prove the vanishing of Tor_1 . There is a finite decreasing filtration of $\mathcal{O}_{\square_{\widehat{X}}^n}$ -submodules

$$0 = \mathcal{F}_r \subsetneq \mathcal{F}_{r-1} \subsetneq \dots \subsetneq \mathcal{F}_0 = \mathcal{F}$$

such that for each $0 \leq j \leq r-1$, we have $\mathcal{F}_j/\mathcal{F}_{j+1} \simeq \mathcal{O}_{\mathfrak{Z}_j}$ as $\mathcal{O}_{\square_{\widehat{X}}^n}$ -modules, for some integral closed formal subscheme $\mathfrak{Z}_j \subset \square_{\widehat{X}}^n$ (see [44, Lemma 01YF]).

We prove the vanishing of Tor_1 by induction on r .

First consider the case when $r = 1$, namely $\mathcal{F} = \mathcal{O}_{\mathfrak{Z}}$ for an integral cycle $\mathfrak{Z} \in z^*(\widehat{X}, n)$. Since \mathfrak{Z} is integral and it intersects the closed subscheme \square_Y^n properly by the condition **(SF)** of Definition 2.2.2, the element t is not a zero-divisor on $\mathcal{O}_{\mathfrak{Z}}$. In particular, we have $\ker(\times t^m : \mathcal{O}_{\mathfrak{Z}} \rightarrow \mathcal{O}_{\mathfrak{Z}}) = 0$, thus,

$$(2.2.2) \quad \mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{O}_{\mathfrak{Z}}, \mathcal{O}_{\square_Y^n}) = 0.$$

When $r > 1$, suppose we have the vanishing $\mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{F}_j, \mathcal{O}_{\square_Y^n}) = 0$ for all $j = 1, \dots, r$. We want to prove its vanishing for $j = 0$. Indeed, we have the short exact sequence of $\mathcal{O}_{\square_{\widehat{X}}^n}$ -modules

$$0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_0 \rightarrow \mathcal{O}_{\mathfrak{Z}_0} \rightarrow 0$$

from which we deduce part of the Tor long exact sequence

$$\dots \rightarrow \mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{F}_1, \mathcal{O}_{\square_Y^n}) \rightarrow \mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{F}_0, \mathcal{O}_{\square_Y^n}) \rightarrow \mathrm{Tor}_1^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{O}_{\mathfrak{Z}_0}, \mathcal{O}_{\square_Y^n}) \rightarrow \dots$$

Here the first term vanishes by the induction hypothesis, and the third term vanishes by (2.2.2). Hence the middle term also vanishes. Since $\mathcal{F}_0 = \mathcal{F}$, this proves the desired assertion of the lemma. \square

Remark 2.2.9. As was also remarked by the referee, a bit more general statement than the one given in Lemma 2.2.8 may hold for various regular adic rings A . Since we do not need it in this article, we resist the temptation to generalize it. \square

Corollary 2.2.10. *Let $Y = \mathrm{Spec}(k_m)$ and $\widehat{X} = \mathrm{Spf}(k[[t]])$. Let \mathcal{A} be a coherent $\mathcal{O}_{\square_{\widehat{X}}^n}$ -algebra such that $[\mathcal{A}] \in z^*(\widehat{X}, n)$.*

Then we have an isomorphism of the graded $\mathcal{O}_{\square_Y^n}$ -algebras

$$\mathrm{Tor}_*^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{A}, \mathcal{O}_{\square_Y^n}) \simeq \mathrm{Tor}_0^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{A}, \mathcal{O}_{\square_Y^n}).$$

Proof. The graded $\mathcal{O}_{\square_Y^n}$ -algebra structure on $\mathrm{Tor}_*^{\mathcal{O}_{\square_{\widehat{X}}^n}}(\mathcal{A}, \mathcal{O}_{\square_Y^n})$ is defined in [44, Section 068G]. Since all higher Tor's vanish by Lemma 2.2.8, we deduce the corollary. \square

2.3. Some finite push-forwards. The primary case we are interested in is when \mathfrak{X} is $\widehat{X} = \mathrm{Spf}(k[[t]])$, but in the middle of our arguments later (§4.4) we need a bit of push-forwards for cycles on some formal schemes that are finite over \widehat{X} . In §2.3, we minimize our discussions only to those needed in this article.

Recall (Fujiwara-Kato [14, Definition 4.7.1, p.341]) a morphism $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ of noetherian formal schemes is *proper* if it is separated of finite type ([14, Definition 4.6.2, p.336]) and universally closed ([14, Definition 4.5.5, p.334]). When \mathfrak{X} and \mathfrak{Y} are both affine, a proper morphism f between them is necessarily finite; recall ([14, Proposition 4.2.1, Definition 4.2.2, pp.324–325]) that a morphism $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ of noetherian formal schemes is *finite*, if for any affine open subset $V = \mathrm{Spf}(A)$ of \mathfrak{Y} , $f^{-1}(V)$ is affine of the form $\mathrm{Spf}(B)$ for a ring B that is a finitely generated A -module.

Definition 2.3.1. For a finite morphism $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ of noetherian affine formal k -schemes, define the finite push-forward on the naive groups (§2.1)

$$f_* : z_d(\mathfrak{X}) \rightarrow z_d(\mathfrak{Y})$$

by sending an integral closed formal subscheme $\mathfrak{Z} \subset \mathfrak{X}$ of dimension d to the d -dimensional cycle $[f_*\mathcal{O}_{\mathfrak{Z}}]_d$ associated to the coherent sheaf $f_*\mathcal{O}_{\mathfrak{Z}}$. Since f is proper, the sheaf push-forward $f_*\mathcal{O}_{\mathfrak{Z}}$ is a coherent $\mathcal{O}_{\mathfrak{Y}}$ -algebra. \square

Write $\mathfrak{X} = \mathrm{Spf}(A)$ and $\mathfrak{Y} = \mathrm{Spf}(B)$. By definition $z_d(\mathfrak{X}) = z_d(\mathrm{Spec}(A))$ and $z_d(\mathfrak{Y}) = z_d(\mathrm{Spec}(B))$ and f is finite. Hence f_* is identical to the usual finite push-forward on the cycles on the scheme $\mathrm{Spec}(A)$ via the associated finite morphism $f : \mathrm{Spec}(A) \rightarrow \mathrm{Spec}(B)$.

We deduce that the cycle push-forward and the sheaf push-forward are compatible:

Lemma 2.3.2. *Let $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a finite morphism of noetherian affine formal k -schemes. Let \mathcal{F} be a coherent $\mathcal{O}_{\mathfrak{X}}$ -module. Then we have the equality of cycles*

$$(2.3.1) \quad f_*[\mathcal{F}] = [f_*\mathcal{F}] \in z_*(\mathfrak{Y}),$$

where the left hand side is the push-forward of the cycle $[\mathcal{F}]$, while the right hand side is the cycle $[f_*\mathcal{F}]$ associated to the push-forward sheaf $f_*\mathcal{F}$.

Proof. Let $\mathfrak{X} = \mathrm{Spf}(A)$ and $\mathfrak{Y} = \mathrm{Spf}(B)$ so that the natural homomorphism $B \rightarrow A$ is finite. By construction we have $z_*(\mathfrak{X}) = z_*(\mathrm{Spec}(A))$ and $z_*(\mathfrak{Y}) = z_*(\mathrm{Spec}(B))$, and we are given $[\mathcal{F}] \in z_*(\mathrm{Spec}(A))$. The push-forward f_* on cycles is the push-forward of cycles on the scheme $\mathrm{Spec}(A)$. Hence the statement follows from the classical case of schemes, which is in [44, Lemma 0EP3], for instance. \square

The finite push-forwards as in Definition 2.3.1 induce push-forwards on higher Chow cycles on noetherian affine formal k -schemes under an additional mild assumption:

Lemma 2.3.3. *Let $f : \mathfrak{X} \rightarrow \mathfrak{Y}$ be a finite surjective morphism of equidimensional noetherian affine formal k -schemes.*

Then we have the induced push-forward morphism

$$f_* : z^q(\mathfrak{X}, \bullet) \rightarrow z^q(\mathfrak{Y}, \bullet)$$

of complexes of abelian groups.

Proof. Since f is finite surjective, so is the morphism $f \times_k 1 : \mathfrak{X} \times_k \square_k^n \rightarrow \mathfrak{Y} \times_k \square_k^n$, which we also denote by f . Furthermore, $\dim \mathfrak{X} = \dim \mathfrak{Y}$ and $\dim \mathfrak{X}_{\mathrm{red}} = \dim \mathfrak{Y}_{\mathrm{red}}$.

For an integral cycle $\mathfrak{Z} \in z^q(\mathfrak{X}, n)$, the cycle push-forward f_* is defined to be $f_*(\mathfrak{Z}) := [f_*\mathcal{O}_{\mathfrak{Z}}]_d$ as in Definition 2.3.1, where $d = \dim \mathfrak{Z}$. We need to check that

$f_*(\mathfrak{Z})$ satisfies the conditions of Definition 2.2.2 over \mathfrak{Y} . Let $Z = \mathfrak{Z} \cap (\mathfrak{X}_{\text{red}} \times \square_k^n)$, the reduction by the largest ideal of definition of $\mathfrak{X} \times \square_k^n$.

For a face $F \subset \square_k^n$, we have $f_*(\mathfrak{Z}) \cap (\mathfrak{Y} \times F) = f_*(\mathfrak{Z} \cap (\mathfrak{X} \times F))$. Thus $\dim(f_*(\mathfrak{Z}) \cap (\mathfrak{Y} \times F)) \leq \dim(\mathfrak{Z} \cap (\mathfrak{X} \times F))$.

The general position condition **(GP)** of Definition 2.2.2 for \mathfrak{Z} says

$$\dim(\mathfrak{Z} \cap (\mathfrak{X} \times F)) \leq \dim \mathfrak{X} \times F - q.$$

So, we have

$$(2.3.2) \quad \dim(f_*(\mathfrak{Z}) \cap (\mathfrak{Y} \times F)) \leq \dim \mathfrak{X} \times F - q = \dim \mathfrak{Y} \times F - q,$$

which is the condition **(GP)** for $f_*(\mathfrak{Z})$.

For the special fiber condition **(SF)** for $f_*(\mathfrak{Z})$, first note that

$$(2.3.3) \quad \dim f_*(\mathfrak{Z}) \cap (\mathfrak{Y}_{\text{red}} \times F) \leq \dim \mathfrak{Z} \cap (\mathfrak{X}_{f^{-1}(\mathcal{I}_0)} \times F),$$

where $\mathfrak{X}_{f^{-1}(\mathcal{I}_0)}$ is the scheme associated to the ideal $f^{-1}(\mathcal{I}_0)\mathcal{O}_{\mathfrak{X}}$. Since $\mathfrak{Z} \cap (\mathfrak{X}_{\text{red}} \times F)$ and $\mathfrak{Z} \cap (\mathfrak{X}_{f^{-1}(\mathcal{I}_0)} \times F)$ are both noetherian schemes, their dimensions as schemes are equal to the dimensions of their respective underlying noetherian topological spaces. But, their underlying topological spaces are both equal to $|\mathfrak{Z} \cap (\mathfrak{X} \times F)|$ (Lemma 2.2.5) so that we have

$$(2.3.4) \quad \dim \mathfrak{Z} \cap (\mathfrak{X}_{f^{-1}(\mathcal{I}_0)} \times F) = \dim \mathfrak{Z} \cap (\mathfrak{X}_{\text{red}} \times F).$$

On the other hand, by the condition **(SF)** for \mathfrak{Z} , we have

$$(2.3.5) \quad \dim \mathfrak{Z} \cap (\mathfrak{X}_{\text{red}} \times F) \leq \dim \mathfrak{X}_{\text{red}} \times F - q = \dim \mathfrak{Y}_{\text{red}} \times F - q.$$

Combining (2.3.3), (2.3.4), and (2.3.5), we deduce that

$$\dim f_*(\mathfrak{Z}) \cap (\mathfrak{Y}_{\text{red}} \times F) \leq \dim \mathfrak{Y}_{\text{red}} \times F - q,$$

which is the condition **(SF)** for $f_*(\mathfrak{Z})$.

Thus we have checked that $f_*(\mathfrak{Z}) \in z^q(\mathfrak{Y}, n)$, so that f_* maps $z^q(\mathfrak{X}, n)$ to $z^q(\mathfrak{Y}, n)$.

For its compatibility with the codimension 1 faces, take the face $F = F_i^\epsilon$ for some $1 \leq i \leq n$ and $\epsilon \in \{0, \infty\}$ in the proof of the condition **(GP)** in the above. By (2.3.2), we obtain the square

$$\begin{array}{ccc} z^q(\mathfrak{X}, n) & \xrightarrow{\partial_i^\epsilon} & z^q(\mathfrak{X}, n-1) \\ \downarrow f_* & & \downarrow f_* \\ z^q(\mathfrak{Y}, n) & \xrightarrow{\partial_i^\epsilon} & z^q(\mathfrak{Y}, n-1). \end{array}$$

That this diagram commutes is checked exactly as done in [5, Proposition (1.3)] using [15, Theorem 6.2(a), p.98]. We omit details. This implies that f_* commutes with ∂ . This proves the lemma. \square

2.4. The mod Y -equivalence on cycles. Returning back to $Y := \text{Spec}(k_m)$ and $\widehat{X} = \text{Spf}(k[[t]])$, we define the mod Y -equivalence on $z^q(\widehat{X}, n)$ as follows.

Definition 2.4.1. Under the above notations:

- (1) Let $\mathcal{R}^q(\widehat{X}, n)$ be the set of all coherent $\mathcal{O}_{\square_{\widehat{X}}^n}$ -algebras \mathcal{A} such that the associated cycle $[\mathcal{A}] \in z^q(\widehat{X}, n)$.

- (2) Let $\mathcal{L}^q(\widehat{X}, Y, n)$ be the set of all pairs $(\mathcal{A}_1, \mathcal{A}_2)$ of coherent $\mathcal{O}_{\widehat{X}}^n$ -algebras with $\mathcal{A}_j \in \mathcal{R}^q(\widehat{X}, n)$ such that there is an isomorphism

$$(2.4.1) \quad \mathcal{A}_1 \otimes_{\mathcal{O}_{\widehat{X}}^n} \mathcal{O}_{\square_Y^q} \simeq \mathcal{A}_2 \otimes_{\mathcal{O}_{\widehat{X}}^n} \mathcal{O}_{\square_Y^q}$$

of $\mathcal{O}_{\square_Y^q}$ -algebras.

- (3) Let $\mathcal{M}^q(\widehat{X}, Y, n) \subset z^q(\widehat{X}, n)$ be the subgroup generated by the cycles of the form $[\mathcal{A}_1] - [\mathcal{A}_2]$ over all pairs $(\mathcal{A}_1, \mathcal{A}_2) \in \mathcal{L}^q(\widehat{X}, Y, n)$.

We say that two cycles \mathfrak{Z}_1 and $\mathfrak{Z}_2 \in z^q(\widehat{X}, n)$ are mod Y -equivalent and write $\mathfrak{Z}_1 \sim_Y \mathfrak{Z}_2$, if $\mathfrak{Z}_1 - \mathfrak{Z}_2 \in \mathcal{M}^q(\widehat{X}, Y, n)$. \square

We have two remarks regarding Definition 2.4.1.

Remark 2.4.2. Corollary 2.2.10 offers a hint on how one may generalize the mod Y -equivalence of Definition 2.4.1 to more general k -schemes of finite type to be discussed in [39].

For a general formal neighborhood, in [39] the author uses some ideas from derived algebraic geometry. The isomorphism (2.4.1) is replaced by an isomorphism of a suitable pair of “derived rings”. While it is of a similar form, it is finer than the derived Milnor patching for perfect complexes studied by S. Landsburg [33], which generalizes the classical Milnor patching of J. Milnor [36].

It seems it is worth studying whether one can avoid using this heavy machine of derived algebraic geometry, by improving Lemma 2.2.8 to the fullest possible extent in the future. \square

Remark 2.4.3. Let $I := (t^m) \subset k[[t]]$. Instead of the above relation \sim_Y on the cycles, one may ask whether it might be better to use the relation \sim_I defined as follows: first declare $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$ for a pair of *integral* cycles \mathfrak{Z}_1 and $\mathfrak{Z}_2 \in z^q(\widehat{X}, n)$ when their structure sheaves $(\mathcal{O}_{\mathfrak{Z}_1}, \mathcal{O}_{\mathfrak{Z}_2})$ satisfy (2.4.1) of Definition 2.4.1, namely

$$\mathcal{O}_{\mathfrak{Z}_1} \otimes_{\mathcal{O}_{\widehat{X}}^n} \mathcal{O}_{\square_Y^q} \simeq \mathcal{O}_{\mathfrak{Z}_2} \otimes_{\mathcal{O}_{\widehat{X}}^n} \mathcal{O}_{\square_Y^q}.$$

Define the subgroup $\mathcal{N}^q(\widehat{X}, Y, n) \subset z^q(\widehat{X}, n)$ generated by $[\mathfrak{Z}_1] - [\mathfrak{Z}_2]$ for such pairs $(\mathcal{O}_{\mathfrak{Z}_1}, \mathcal{O}_{\mathfrak{Z}_2})$. We define \sim_I on $z^q(\widehat{X}, n)$ using this subgroup $\mathcal{N}^q(\widehat{X}, Y, n)$. It is similar to the version taken in [41], called the “mod t^m -equivalence” there.

A benefit of \sim_I is that it is relatively easier to work with; when one wishes to check whether certain operations on cycles respect the equivalence, one can test them just on pairs of integral cycles. A draw-back of this is that, the property of a scheme being integral is not preserved well under base change in general (see, e.g. EGA IV₂ [19, Définition (4.6.2), p.68] for the definition of geometric integrality, which does behave well under base change), so that this approach counting on putting relations on pairs integral cycles is unlikely to be generalizable in a functorial way, in general.

However, at least in the Milnor range (i.e. $q = n$), we will show that our definition of \sim_Y via $\mathcal{M}^n(\widehat{X}, Y, n)$ in Definition 2.4.1 and \sim_I in the above paragraph via $\mathcal{N}^n(\widehat{X}, Y, n)$ are equivalent. The proof requires a few properties of the generating cycles in $z^n(\widehat{X}, n)$, so that it is postponed until Lemma 4.3.1. This result will be used in §4.4 and §5. \square

Coming back to our construction, we observe:

Lemma 2.4.4. *Let $Y = \text{Spec}(k_m)$ and $\widehat{X} = \text{Spf}(k[[t]])$. Let $n \geq 1$. For $1 \leq i \leq n$ and $\epsilon \in \{0, \infty\}$, we have*

$$\partial_i^\epsilon \mathcal{M}^q(\widehat{X}, Y, n) \subset \mathcal{M}^q(\widehat{X}, Y, n-1).$$

In particular, $\mathcal{M}^q(\widehat{X}, Y, \bullet) \subset z^q(\widehat{X}, \bullet)$ is a subcomplex.

Proof. Let $\iota_i^\epsilon : \square_{\widehat{X}}^{n-1} \hookrightarrow \square_{\widehat{X}}^n$ be the closed immersion given by $\{y_i = \epsilon\}$.

When $\mathcal{A} \in \mathcal{R}^q(\widehat{X}, n)$, we have $[\mathcal{A}] \in z^q(\widehat{X}, n)$. Here $[(\iota_i^\epsilon)^*(\mathcal{A})] = \partial_i^\epsilon[\mathcal{A}] \in z^q(\widehat{X}, n-1)$, so that $(\iota_i^\epsilon)^*\mathcal{A} \in \mathcal{R}^q(\widehat{X}, n-1)$.

When $(\mathcal{A}_1, \mathcal{A}_2) \in \mathcal{L}^q(\widehat{X}, Y, n)$, we have $\mathcal{A}_j \in \mathcal{R}^q(\widehat{X}, n)$ and there is an isomorphism

$$(2.4.2) \quad \mathcal{A}_1 \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n} \simeq \mathcal{A}_2 \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n}$$

of $\mathcal{O}_{\square_Y^n}$ -algebras. We already saw that $(\iota_i^\epsilon)^*\mathcal{A}_j \in \mathcal{R}^q(\widehat{X}, n-1)$. Applying $(\iota_i^\epsilon)^*$ to (2.4.2), we deduce an isomorphism

$$(\iota_i^\epsilon)^*\mathcal{A}_1 \otimes_{\mathcal{O}_{\square_{\widehat{X}}^{n-1}}} \mathcal{O}_{\square_Y^{n-1}} \simeq (\iota_i^\epsilon)^*\mathcal{A}_2 \otimes_{\mathcal{O}_{\square_{\widehat{X}}^{n-1}}} \mathcal{O}_{\square_Y^{n-1}}$$

of $\mathcal{O}_{\square_Y^{n-1}}$ -algebras. This means $((\iota_i^\epsilon)^*\mathcal{A}_1, (\iota_i^\epsilon)^*\mathcal{A}_2) \in \mathcal{L}^q(\widehat{X}, Y, n-1)$. Thus

$$\partial_i^\epsilon([\mathcal{A}_1] - [\mathcal{A}_2]) = [(\iota_i^\epsilon)^*\mathcal{A}_1] - [(\iota_i^\epsilon)^*\mathcal{A}_2] \in \mathcal{M}^q(\widehat{X}, Y, n-1).$$

Since the cycles of the form $[\mathcal{A}_1] - [\mathcal{A}_2]$ generate $\mathcal{M}^q(\widehat{X}, Y, n)$, this implies that $\partial_i^\epsilon \mathcal{M}^q(\widehat{X}, Y, n) \subset \mathcal{M}^q(\widehat{X}, Y, n-1)$ as desired. \square

Definition 2.4.5. Let $m \geq 2$. Let $Y = \text{Spec}(k_m)$ and $\widehat{X} = \text{Spf}(k[[t]])$. Define

$$z^q(\widehat{X} \bmod Y, \bullet) := \frac{z^q(\widehat{X}, \bullet)}{\mathcal{M}^q(\widehat{X}, Y, \bullet)},$$

and define $\text{CH}^q(\widehat{X} \bmod Y, n) := \text{H}_n(z^q(\widehat{X} \bmod Y, \bullet))$. We denote this group also by $\mathbf{CH}^q(k_m, n) = \mathbf{CH}^q(\text{Spec}(k_m), n)$ in the boldface letters, to distinguish it from the higher Chow group of S. Bloch [5]. \square

The following special range is easy to compute. It answers the part (1) of the first main theorem, Theorem 1.2.1:

Proposition 2.4.6. For $q > n$, $z^q(\widehat{X} \bmod Y, n) = 0$. In particular, we have

$$\text{CH}^q(\widehat{X} \bmod Y, n) = \mathbf{CH}^q(k_m, n) = 0.$$

Proof. Note that $\dim \widehat{X} \times \square^n = n+1$. Thus if $q \geq n+2$, due to the dimension reason, we have $z^q(\widehat{X}, n) = 0$. Thus $z^q(\widehat{X} \bmod Y, n) = 0$ as well.

Suppose $q = n+1$. Let $\mathfrak{Z} \in z^{n+1}(\widehat{X}, n)$ be an integral cycle. Note that $\widehat{X}_{\text{red}} = \text{Spec}(k)$. By the special fiber condition (**SF**) of Definition 2.2.2, the intersection $\mathfrak{Z} \cap (\text{Spec}(k) \times \square^n)$ has the codimension $\geq n+1$ in the space $\text{Spec}(k) \times \square^n$ of dimension n . This shows the intersection $\mathfrak{Z} \cap (\text{Spec}(k) \times \square^n)$ is empty. On the other hand, we have

$$|\mathfrak{Z} \cap (\text{Spec}(k) \times \square_k^n)| \subset |\mathfrak{Z} \cap (\text{Spec}(k_m) \times \square_k^n)| \subset |\mathfrak{Z} \cap (\widehat{X} \times \square_k^n)|,$$

and the first and the third topological spaces are equal (Lemma 2.2.5). Thus $\mathfrak{Z} \cap (\text{Spec}(k_m) \times \square^n) = \emptyset$ as well. It means

$$\mathcal{O}_{\mathfrak{Z}} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n} = 0 = \mathcal{O}_{\emptyset} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n},$$

thus the cycle \mathfrak{Z} is mod Y -equivalent to the empty scheme \emptyset . Thus $[\mathfrak{Z}] \in \mathcal{M}^{n+1}(\widehat{X}, Y, n)$ and the class of \mathfrak{Z} in $z^{n+1}(\widehat{X} \bmod Y, n)$ is 0. This shows $z^{n+1}(\widehat{X} \bmod Y, n) = 0$. \square

Remark 2.4.7. If we specialize to the Milnor range $q = n$ in Definition 2.4.5, the group $\text{CH}^n(\widehat{X} \bmod Y, n)$ can be expressed as

$$(2.4.3) \quad \text{CH}^n(\widehat{X} \bmod Y, n) = \frac{z^n(\widehat{X} \bmod Y, n)}{\partial(z^n(\widehat{X} \bmod Y, n+1))}$$

because $z^n(\widehat{X} \bmod Y, n-1) = 0$ by Proposition 2.4.6 so that

$$\ker(\partial : z^n(\widehat{X} \bmod Y, n) \rightarrow z^n(\widehat{X} \bmod Y, n-1)) = z^n(\widehat{X} \bmod Y, n).$$

Thus we may rewrite (2.4.3) also as

$$(2.4.4) \quad \text{CH}^n(\widehat{X} \bmod Y, n) := \frac{z^n(\widehat{X}, n)}{\partial(z^n(\widehat{X}, n+1)) + \mathcal{M}^n(\widehat{X}, Y, n)}.$$

One may wonder whether the numerator in (2.4.4) is also equal to $\ker(\partial : z^n(\widehat{X}, n) \rightarrow z^n(\widehat{X}, n-1))$. The answer is yes, and we will see it in Lemma 3.2.4. \square

2.5. Tate algebras. In §2.5, we recall some basic definitions and results related to the Tate algebras of J. Tate [46]. We will use them to deal with the “generic fibers” of the cycles in this article. For general references on the Tate algebras, there are a few books in the literature, e.g. Bosch-Güntzer-Remmert [7], Bosch [8], Fresnel-van der Put [13], Fujiwara-Kato [14]. At some places, Tate algebras are also called the standard affinoid algebras or the free affinoid algebras, as well.

For a complete discrete valued field K with a nontrivial non-archimedean norm, such as $k((t))$, the Tate algebra T_n over K with n -variables z_1, \dots, z_n is defined to be the subring $K\{z_1, \dots, z_n\} \subset K[[z_1, \dots, z_n]]$ consisting of the restricted formal power series, i.e.

$$p = \sum_{I=(i_1, \dots, i_n), i_j \geq 0} a_I z^I,$$

with $z^I := z_1^{i_1} \cdots z_n^{i_n}$ such that $|a_I| \rightarrow 0$ as $|I| = i_1 + \cdots + i_n \rightarrow \infty$. The norm on K extends to the Gauß norm on T_n by taking $|p| := \max |a_I|$, making T_n a Banach algebra over K of at most countable type. See e.g. any one of [7, §5.1.1, p.192], [8, §2.2, Definition 2.2, p.13], [13, §3.1, p.46], or [14, §9.3(a), p.227].

The following is a compilation of a few facts on Tate algebras, that we need:

Lemma 2.5.1. *Let K be a complete discrete valued field with a nontrivial non-archimedean norm, and let T_n be the Tate algebra over K in n -variables. Then:*

- (1) T_n is a noetherian regular UFD of Krull dimension n .
- (2) T_1 is a Euclidean domain, in particular a PID.
- (3) For each maximal ideal $\mathfrak{m} \subset T_n$, the injective ring homomorphism $K \hookrightarrow T_n/\mathfrak{m}$ gives a finite extension of fields.
- (4) For any finite extension $K \subset L$ of fields, there exists a unique extension of the discrete valuation on K to L .
- (5) Every $f \in T_1$ can be written uniquely as $f = ug$, where $u \in T_1^\times$ and $g \in K^\circ[z_1]$ for the valuation ring $K^\circ = \{x \in K \mid |x| \leq 1\}$.

Proof. (1) is scattered at a few places. Collect them from [7, §5.2.6, Theorem 1, p.207], [8, §2.2, Proposition 14, 15, p.20, Proposition 17, p.22], [13, Theorem 3.2.1-(1),(2), p.48], and [14, Exercise 0.9.4, p.23, Proposition 9.3.9, p.229].

(2) : see [8, §2.2, Corollary 10, p.19].

(3) : see [8, Corollary 12, p.19] or [13, Theorem 3.2.1-(5), p.48].

(4) : see e.g. H. Hasse [23, Ch.12, Prolongation theorem, p.183].

(5) : see [13, Exercises 3.2.2-(1), p.50], as a consequence of the Weierstraß division theorem [13, Theorem 3.1.1, p.46], or e.g. [7, §5.2.1, Theorem 2, p. 200, §5.2.2, Theorem 1, p.201], [8, §2.2, Theorem 8, p.17, Corollary 9, p.18], [14, Exercise 0.A.3, p.236]. \square

3. ON THE GENERATOR CYCLES

We continue to use the notations $Y = \text{Spec}(k_m)$ and $\widehat{X} = \text{Spf}(k[[t]])$.

In §3, we study a few general aspects of the cycles in the group $z^n(\widehat{X}, n)$. These cycles generate the groups $\mathrm{CH}^n(\widehat{X}, n)$ and $\mathbf{CH}^n(\mathrm{Spec}(k_m), n)$.

The arguments here are a hybrid of basic commutative algebra, basic rigid analytic geometry, and algebraic geometry of cycles on formal schemes. An essential observation is that the “generic fiber” of an integral cycle in $z^n(\widehat{X}, n)$ induces a maximal ideal of a Tate algebra over the Laurent field $k((t))$. Some results of rudimentary rigid analytic geometry then allow us to obtain “triangular polynomial generators” of the ideal that define the given integral cycle over $\mathrm{Spf}(k[[t]])$ in §3.3.

3.1. Algebraicity and generic fibers. For a complete discrete valuation field K with a nontrivial non-archimedean norm, consider $T_n = K\{z_1, \dots, z_n\}$, the Tate algebra in n variables z_1, \dots, z_n . Let’s start with the standard fact that each maximal ideal $\mathfrak{m} \subset T_n$ of the Tate algebra is generated by n elements (in particular it is a complete intersection); see S. Bosch [8, §2.2, Proposition 17, p.22]. Here, these n generators can be chosen from the subring $K[z_1, \dots, z_n] \subset T_n$ of polynomials in z_1, \dots, z_n ; see Fresnel-van der Put [13, Exercises 3.2.2-(2), p.50] or Bosch-Güntzer-Remmert [7, §7.1.1, Proposition 3, p.261].

We improve this fact a bit further as follows. It might be of independent interest:

Lemma 3.1.1. *Let K be a complete discrete valuation field with a nontrivial non-archimedean norm, and K° be its valuation ring. Let $T_n = K\{z_1, \dots, z_n\}$ be the Tate algebra over K , and let $\mathfrak{m} \subset T_n$ be a maximal ideal.*

*Then we have a sequence of **polynomials** in z_1, \dots, z_n*

$$\begin{cases} p_1(z_1) \in K^\circ[z_1], \\ p_2(z_1, z_2) \in K[z_1, z_2], \\ \vdots \\ p_n(z_1, \dots, z_n) \in K[z_1, \dots, z_n], \end{cases}$$

where

- (1) each p_i is monic in z_i for $1 \leq i \leq n$,
- (2) for $2 \leq i \leq n$, when p_i is regarded as a polynomial in z_i , its coefficients in $K[z_1, \dots, z_{i-1}]$ have their z_j -degrees strictly less than $\deg_{z_j}(p_j)$ for all $1 \leq j < i$, and
- (3) $(p_1, \dots, p_n) = (p_1, \dots, p_n)T_n = \mathfrak{m}$.

Proof. The proof is in part based on the argument of S. Bosch [8, §2.2, Proposition 17, p.22], with some improvements to deduce our stronger statements.

For the sequence of inclusions $T_1 \subset T_2 \subset \dots \subset T_n$, consider the prime ideals $\mathfrak{m}_i := T_i \cap \mathfrak{m}$ for $1 \leq i \leq n$. Here $\mathfrak{m}_n = \mathfrak{m}$. These induce injections

$$K \hookrightarrow T_1/\mathfrak{m}_1 \hookrightarrow \dots \hookrightarrow T_n/\mathfrak{m}_n,$$

where T_n/\mathfrak{m}_n is a finite extension of K of fields (Lemma 2.5.1-(3)). Hence each intermediate T_i/\mathfrak{m}_i is also a field, thus $\mathfrak{m}_i \subset T_i$ is a maximal ideal for $1 \leq i \leq n$.

When $n = 1$, the ideal $\mathfrak{m}_1 \subset T_1$ is maximal, while T_1 is a PID by Lemma 2.5.1-(2). Hence we have $(f) = \mathfrak{m}_1$ for some nonzero $f \in T_1$. By Lemma 2.5.1-(5), we have $f = ug$ for some $u \in T_1^\times$ and a monic polynomial $g \in K^\circ[z_1]$. Thus $(f) = (g) = \mathfrak{m}_1$. Taking $p_1 := g$ answers the lemma in this case.

Now suppose $n \geq 2$, and suppose the lemma holds for all positive integers $< n$, so that we have

$$p_1(z_1) \in K^\circ[z_1], p_2(z_1, z_2) \in K[z_1, z_2], \dots, p_{n-1}(z_1, \dots, z_{n-1}) \in K[z_1, \dots, z_{n-1}]$$

satisfying the conclusions of the lemma for the maximal ideal $\mathfrak{m}_{n-1} \subset T_{n-1}$.

Note that (e.g. from *loc.cit*) we have the following commutative diagram of continuous K -algebra homomorphisms

$$(3.1.1) \quad \begin{array}{ccc} T_{n-1} \otimes_{T_{n-1}} T_n = T_{n-1}\{z_n\} & \xlongequal{\quad\quad\quad} & T_n \\ \text{can} \otimes \text{Id}_{T_n} =: \phi' \downarrow & & \downarrow \phi \\ (T_{n-1}/\mathfrak{m}_{n-1}) \otimes_{T_{n-1}} T_n = (T_{n-1}/\mathfrak{m}_{n-1})\{z_n\} & \xrightarrow{\quad \pi \quad} & T_n/\mathfrak{m}, \end{array}$$

where the vertical maps ϕ, ϕ' are given by the reduction mod \mathfrak{m} and \mathfrak{m}_{n-1} , respectively, and π maps z_n to its residue class in T_n/\mathfrak{m} . Note that both of the maps ϕ' and π are surjective.

Take $\ker(\pi)$. Since T_n/\mathfrak{m} is a field and π is surjective, by the first isomorphism theorem $\ker(\pi) \subset (T_{n-1}/\mathfrak{m}_{n-1})\{z_n\}$ is a maximal ideal. On the other hand, by Lemma 2.5.1-(4), the field $T_{n-1}/\mathfrak{m}_{n-1}$ is also a complete discrete valued field with respect to the unique norm extending the norm on K . Hence by the case $n = 1$ we proved already, there exists a polynomial $\bar{p}_n \in (T_{n-1}/\mathfrak{m}_{n-1})[z_n]$ monic in z_n , such that $(\bar{p}_n) = \ker(\pi)$.

Since ϕ' is surjective, we can choose a lifting $p_n(z_1, \dots, z_n) \in T_{n-1}[z_n] \subset T_{n-1}\{z_n\}$ of \bar{p}_n which is still a monic polynomial in z_n .

By the commutativity of the diagram (3.1.1), we have $(p_1, \dots, p_n) = \mathfrak{m}$, but one issue yet to resolve is that, while $p_n(z_1, \dots, z_n)$ is in $T_{n-1}[z_n]$, it may not be yet in $K[z_1, \dots, z_{n-1}][z_n]$.

To improve this, when $N := \deg_{z_n}(p_n)$, write

$$p_n = \sum_{i=0}^N \alpha_i z_n^i \in T_{n-1}[z_n],$$

for $\alpha_i \in T_{n-1}$, $0 \leq i \leq N$ with $\alpha_N = 1$. By the induction hypothesis p_{n-1} is a monic polynomial in z_{n-1} . By the Weierstraß division theorem (Fresnel-van der Put [13, Theorem 3.1.1, p.46]), for $0 \leq i < N$, we can find $q_i \in T_{n-1}$ and $r_i \in T_{n-2}[z_{n-1}]$ such that

$$\alpha_i = q_i p_{n-1} + r_i,$$

where r_i is a polynomial in z_{n-1} with $\deg_{z_{n-1}}(r_i) < \deg_{z_{n-1}}(p_{n-1})$. We let $r_N = 1$. Then for

$$p'_n := \sum_{i=1}^N r_i z_n^i \in T_{n-2}[z_{n-1}][z_n]$$

we have $\mathfrak{m} = (p_1, \dots, p_{n-1}, p_n) = (p_1, \dots, p_{n-1}, p'_n)$. Hence we may replace p_n by p'_n . We can inductively apply the Weierstraß division theorem with the divisions by $p_{n-2}, p_{n-3}, \dots, p_1$, and after these backward inductive replacements, in finite steps we obtain $p_n \in K[z_1, \dots, z_n]$. This proves the lemma. \square

As an immediate application, we deduce that our cycles in the Milnor range can be seen as separated $k[[t]]$ -schemes of finite presentation. It will be further improved in the next subsections.

Corollary 3.1.2. *Let $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$ be an integral cycle. Let $y'_i := y_i/(y_i - 1)$ using the automorphism $y_i \mapsto y'_i$ of \mathbb{P}^1 .*

*Then there are **polynomials** in $k[[t]][y'_1, \dots, y'_n]$ of the form*

$$(3.1.2) \quad \begin{cases} P_1(y'_1) \in k[[t]][y'_1], \\ \vdots \\ P_n(y'_1, \dots, y'_n) \in k[[t]][y'_1, \dots, y'_n], \end{cases}$$

such that the ideal $(P_1(y'_1), \dots, P_n(y'_1, \dots, y'_n))$ defines \mathfrak{Z} .

In particular, \mathfrak{Z} can be also regarded as a separated $k[[t]]$ -scheme of finite presentation.

Proof. Let $K := k((t)) = \text{Frac}(k[[t]])$. Write $z_i := y'_i$ for notational simplicity as well as to be consistent with the notation of Lemma 3.1.1.

The cycle \mathfrak{Z} is given by a height n prime ideal $I(\mathfrak{Z})$ of the Tate algebra $T_n = k[[t]]\{z_1, \dots, z_n\}$. So, we have the canonical homomorphisms

$$k[[t]] \hookrightarrow k[[t]]\{z_1, \dots, z_n\} \twoheadrightarrow k[[t]]\{z_1, \dots, z_n\}/I(\mathfrak{Z}),$$

and, we may regard \mathfrak{Z} as a scheme over $\text{Spec}(k[[t]])$ as well.

Let $\eta \in \text{Spec}(k[[t]])$ be the generic point. The generic fiber \mathfrak{Z}_η is obtained by the flat base change via $\eta \rightarrow \text{Spec}(k[[t]])$, and it gives a height n prime ideal of the Tate algebra T_n . Since the Krull dimension of T_n is n (Lemma 2.5.1-(1)), this height n prime ideal is maximal. Hence the generic fiber \mathfrak{Z}_η is defined by a maximal ideal $\mathfrak{m} \subset T_n$.

By Lemma 3.1.1, we then have a set of generators $p_1(z_1) \in K^\circ[z_1]$, $p_2(z_1, z_2) \in K[z_1, z_2]$, \dots , $p_n(z_1, \dots, z_n) \in K[z_1, \dots, z_n]$ with the properties there.

To obtain the generators for its closure \mathfrak{Z} , we clear the denominators of the coefficients of the terms of p_i by the l.c.m. in $k[[t]]$. Since $k[[t]]$ is a UFD, this is possible. So obtained elements are denoted by $P_1(z_1) \in k[[t]][z_1]$, \dots , $P_n(z_1, \dots, z_n) \in k[[t]][z_1, \dots, z_n]$. \square

In §3.2, after proving the finiteness, we will show that the above can be also stated for a sequence of polynomials in $k[[t]]\{y_1, \dots, y_n\}$ (not just for $k[[t]]\{y'_1, \dots, y'_n\}$), where the polynomials P_i can satisfy more requirements.

3.2. Finiteness. In §3.2, we study the generators of $z^n(\text{Spf}(k[[t]]), n)$ from the algebraic perspective to obtain additional properties.

Lemma 3.2.1. *Let $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$ be an integral cycle.*

Then \mathfrak{Z} regarded as a scheme over $k[[t]]$, is quasi-finite over $k[[t]]$.

Proof. By Corollary 3.1.2, we may regard \mathfrak{Z} as a scheme over $k[[t]]$.

We check that for each $\mathfrak{p} \in \text{Spec}(k[[t]])$, the fiber $\mathfrak{Z}_\mathfrak{p}$ is a finite set.

As before, via a suitable automorphism of \mathbb{P}^1 , use $y'_i = y_i/(y_i - 1)$ for $1 \leq i \leq n$. The generic fiber \mathfrak{Z}_η is given by a maximal ideal of the Tate algebra $T_n = k((t))\{y'_1, \dots, y'_n\}$. Hence $|\mathfrak{Z}_\eta| = 1 < \infty$.

For the closed point $\mathfrak{m} \in \text{Spec}(k[[t]])$, by the special fiber condition (**SF**) of Definition 2.2.2, this cycle intersects $\mathfrak{m} \times \square_k^n$ properly. Hence $\mathfrak{Z}_\mathfrak{m}$ gives a closed subscheme of codimension $\geq n$ in $\square_{\kappa(\mathfrak{m})}^n = \square_k^n$. Since $\dim \square_k^n = n$, we have $\dim(\mathfrak{Z}_\mathfrak{m}) \leq 0$ so that $|\mathfrak{Z}_\mathfrak{m}| < \infty$ again.

Thus we proved that $\mathfrak{Z} \rightarrow \text{Spec}(k[[t]])$ is quasi-finite, as desired. \square

We will refine Lemma 3.2.1 even further. To do so, recall the following (See Conrad-Stein [10, Lemma 8.3]. See also EGA IV₄ [19, Théorème (18.5.11)-(c), p.130]):

Lemma 3.2.2. *Let A be a henselian local ring, and let G be a quasi-finite separated A -scheme of finite presentation.*

Then there is a unique decomposition $G = G_f \coprod G'$ into disjoint clopen pieces such that $G_f \rightarrow \text{Spec}(A)$ is finite, and G' has the empty special fiber.

An interesting and important point is that the nonzero integral cycles in $z^n(\widehat{X}, n)$ are all finite in the Milnor range:

Corollary 3.2.3. *Let $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$ be a nonzero integral cycle. Then \mathfrak{Z} is finite over $k[[t]]$.*

Proof. The given integral cycle \mathfrak{Z} is quasi-finite over the henselian local ring $k[[t]]$ by Lemma 3.2.1. Furthermore, it can be regarded as a separated $k[[t]]$ -scheme of finite presentation by Corollary 3.1.2.

Hence we may apply Lemma 3.2.2. By this, \mathfrak{Z} can be written as the disjoint union of the finite part \mathfrak{Z}_f , and the part \mathfrak{Z}' that has the empty special fiber. But, \mathfrak{Z} is integral, so we have either (a) $\mathfrak{Z} = \mathfrak{Z}_f$, in which case $\mathfrak{Z} \rightarrow \text{Spec}(k[[t]])$ is finite, or (b) $\mathfrak{Z} = \mathfrak{Z}'$, in which case $\mathfrak{Z} \rightarrow \text{Spec}(k[[t]])$ is not finite, and the special fiber is empty.

Since the cycle \mathfrak{Z} *does have* the nonempty special fiber (Lemma 2.2.5), the above case (b) cannot happen. Hence the case (a) is the only possibility and $\mathfrak{Z} \rightarrow \text{Spec}(k[[t]])$ is finite. \square

Being finite over $k[[t]]$ have a few important consequences. The first is on intersection with the faces:

Lemma 3.2.4. *Let $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$ be an integral cycle. Then \mathfrak{Z} does not intersect any proper face of $\text{Spf}(k[[t]]) \times \square^n$. In particular, $\partial(\mathfrak{Z}) = 0$.*

Proof. We imitate the argument of Krishna-Park [31, Lemma 2.21, p.1005]. Let $\widehat{X} = \text{Spf}(k[[t]])$. Let $F \subsetneq \square_k^n$ be a proper face. We show that $\mathfrak{Z} \cap (\widehat{X} \times F) = \emptyset$.

Suppose not. As before, we regard \mathfrak{Z} as a scheme over $\text{Spec}(k[[t]])$. By Corollary 3.2.3, the morphism $\mathfrak{Z} \rightarrow \text{Spec}(k[[t]])$ is finite. Consider the composite of finite morphisms

$$\mathfrak{Z} \cap (\widehat{X} \times F) \hookrightarrow \mathfrak{Z} \rightarrow \text{Spec}(k[[t]]).$$

Its image in $\text{Spec}(k[[t]])$ is closed. Since we supposed that $\mathfrak{Z} \cap (\widehat{X} \times F)$ is nonempty, the image must intersect the unique closed point \mathfrak{m} of $\text{Spec}(k[[t]])$. Hence $\mathfrak{Z} \cap (\mathfrak{m} \times F) \neq \emptyset$.

On the other hand, by the special fiber condition (**SF**) of Definition 2.2.2 for \mathfrak{Z} , the codimension of $\mathfrak{Z} \cap (\mathfrak{m} \times F)$ in $\mathfrak{m} \times F$ is $\geq n$. This is equivalent to saying that

$$\dim(\mathfrak{Z} \cap (\mathfrak{m} \times F)) \leq \dim(\mathfrak{m} \times F) - n = \dim F - n < \dagger 0,$$

where \dagger holds because F is a proper face. But this is a contradiction for a nonempty set. Hence we must have $\mathfrak{Z} \cap (\widehat{X} \times F) = \emptyset$. This proves the lemma. \square

We discuss additional algebraic properties of the cycles in $z^n(\text{Spf}(k[[t]]), n)$. Recall:

Lemma 3.2.5. *Let (R, \mathfrak{m}) be a henselian (complete, resp.) local ring and let $R \hookrightarrow B$ be a finite extension of rings.*

Then B is a finite direct product of henselian (complete, resp.) local rings. Furthermore, there is a bijection between the factors and the maximal ideals of B .

In particular, if B is an integral domain in addition to the above assumptions, then B is a henselian (complete, resp.) local domain with a unique maximal ideal.

Proof. See EGA IV₄ [19, Propositions (18.5.9), (18.5.10), p.130] for the henselian case, and D. Eisenbud [11, Corollary 7.6, p.190] for the complete case. \square

Corollary 3.2.6. *Let $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$ be an integral cycle, and let B be the integral domain such that $\mathfrak{Z} = \text{Spf}(B)$.*

Then the canonical homomorphism $k[[t]] \rightarrow B$ of rings is finite, and B is a complete local integral domain, which is a free $k[[t]]$ -module of finite rank.

Proof. By Corollary 3.2.3, the ring homomorphism $k[[t]] \rightarrow B$ is finite. Here t is a non-zero-divisor on B by the special fiber condition (**SF**) of Definition 2.2.2 for \mathfrak{Z} . Since B is a finitely generated $k[[t]]$ -module, where t is not a zero-divisor in B , it is torsion free. Thus, by the fundamental theorem of finitely generated modules over a PID, B is a free $k[[t]]$ -module of finite rank.

That B is a complete local integral domain follows from Lemma 3.2.5. \square

Definition 3.2.7. For an integral cycle $\mathfrak{Z} \in z^n(\mathrm{Spf}(k[[t]]), n)$, let $k[[t]] \rightarrow B$ define \mathfrak{Z} as the above. We can define the norm map $N : B^\times \rightarrow k[[t]]^\times$ as follows.

By Corollary 3.2.6, B is a free $k[[t]]$ -module of finite rank. Hence for each $b \in B$, the left multiplication by b

$$L_b : B \rightarrow B, \quad x \mapsto bx$$

is a $k[[t]]$ -linear endomorphism of the free $k[[t]]$ -module B of finite rank. When $b \in B^\times$, the map L_b is an automorphism. We define $N(b)$ to be the determinant $\det(L_b) \in k[[t]]^\times$. It is independent of the choice of a $k[[t]]$ -basis of B , and the map N is multiplicative, by standard linear algebra. \square

The finiteness property of \mathfrak{Z} over $k[[t]]$ has another set of consequences. The following is an analogue in the formal setting of the finiteness criterion in [31, Lemma 2.9] :

Lemma 3.2.8. *Let $\mathfrak{Z} \subset \widehat{X} \times \square^n$ be an integral closed formal subscheme. Consider the open immersion $\widehat{X} \times \square^n \subset \widehat{X} \times (\mathbb{P}^1)^n$.*

Then the following are equivalent:

- (1) *The morphism $\mathfrak{Z} \rightarrow \widehat{X}$ is finite.*
- (2) *The formal scheme \mathfrak{Z} is closed in $\widehat{X} \times (\mathbb{P}^1)^n$.*

Proof. The proof is almost identical to that of [31, Lemma 2.9], except that we use the corresponding properties for formal schemes.

(1) \Rightarrow (2) : Consider the factorization $\mathfrak{Z} \hookrightarrow \widehat{X} \times (\mathbb{P}^1)^n \rightarrow \widehat{X}$ of the finite morphism. The second morphism is separated of finite type. Thus by Fujiwara-Kato [14, Corollary 4.6.15, p.340], the first morphism is also finite. An immersion is finite only when it is a closed immersion. This implies (2).

(2) \Rightarrow (1): Since \mathfrak{Z} is closed in $\widehat{X} \times (\mathbb{P}^1)^n$, in particular this inclusion morphism is proper. Thus the composite $\mathfrak{Z} \hookrightarrow \widehat{X} \times (\mathbb{P}^1)^n \rightarrow \widehat{X}$ is also proper. This implies that the reduction by an ideal of definition of \widehat{X} is also proper ([14, Proposition 4.7.3, p.341]). But both \mathfrak{Z} and \widehat{X} are affine formal schemes, so their reductions are affine schemes. A morphism between affine schemes is proper if and only if it is finite ([21, Exercise II-4.6, p.106]). But this means the morphism $\mathfrak{Z} \rightarrow \widehat{X}$ of formal schemes is also finite ([14, Proposition 4.2.3, p.325]). This implies (1). \square

Corollary 3.2.9. *Let $\mathfrak{Z} \subset z^n(\widehat{X}, n)$ be an integral cycle. Consider $(\mathbb{P}^1)^n \supset \square^n$, where y_1, \dots, y_n are the coordinates.*

Then it is closed in $\widehat{X} \times (\mathbb{P}^1)^n$ and it does not intersect any divisor of the form $\{y_i = \epsilon\}$, where $1 \leq i \leq n$ and $\epsilon = 0, \infty, 1$.

In particular, for $\mathbb{A}^n \subset (\mathbb{P}^1)^n$ with the coordinates y_1, \dots, y_n , we can regard \mathfrak{Z} as an integral closed formal subscheme of $\widehat{X} \times \mathbb{A}^n$, and there exists a prime ideal $I \subset k[[t]]\{y_1, \dots, y_n\}$ such that

$$(3.2.1) \quad \mathfrak{Z} = \mathrm{Spf}(k[[t]]\{y_1, \dots, y_n\}/I).$$

Proof. From Corollary 3.2.3, we know that \mathfrak{Z} is finite over $k[[t]]$, and by Lemma 3.2.8, we know that \mathfrak{Z} is closed in $\widehat{X} \times (\mathbb{P}^1)^n$. Because $\mathfrak{Z} \subset \widehat{X} \times \square^n$ does not intersect the divisors $\{y_i = 1\} \subset (\widehat{X} \times (\mathbb{P}^1)^n) \setminus (\widehat{X} \times \square^n)$, in fact $\mathfrak{Z} \subset \widehat{X} \times_k \mathbb{A}_k^n$, and it is closed. This shows (3.2.1).

The remaining statement that $\overline{\mathfrak{Z}} \cap \{y_i = \epsilon\}$ with $\epsilon = 0, \infty$ follows from Lemma 3.2.4. \square

Hence, we no longer need to use the ugly coordinate changes $y_i \mapsto y'_i = y_i/(y_i - 1)$. The generic fibers also enjoy the same property. Namely, Corollary 3.2.9 immediately implies:

Corollary 3.2.10. *Let $\mathfrak{Z} \in z^n(\widehat{X}, n)$ be an integral cycle. Let \mathfrak{Z}_η be its generic fiber. Then there exists a maximal ideal $\mathfrak{m} \subset k((t))\{y_1, \dots, y_n\}$ such that \mathfrak{Z}_η is given by the affinoid algebra $k((t))\{y_1, \dots, y_n\}/\mathfrak{m}$.*

3.3. Triangular generators. For a given integral cycle $\mathfrak{Z} \in z^n(\widehat{X}, n)$, in Corollary 3.1.2 we found n polynomial generators in the variables y'_1, \dots, y'_n , where $y'_i = y_i/(y_i - 1)$. While it was enough for our purposes of proving finiteness at that time, we now have obtained the finiteness and some of its consequences. This time, we want to improve it further.

The goal of §3.3 is to find a nice “triangular” generating set for each integral cycle $\mathfrak{Z} \in z^n(\mathrm{Spf}(k[[t]]), n)$, this time as polynomials in y_1, \dots, y_n . We will use Corollaries 3.2.9 and 3.2.10 we obtained in §3.2 as consequences of the finiteness of \mathfrak{Z} over $k[[t]]$.

Proposition 3.3.1. *Let $\mathfrak{Z} \in z^n(\mathrm{Spf}(k[[t]]), n)$ be an integral cycle.*

*Then there are **polynomials** in $k[[t]][y_1, \dots, y_n]$ of the form*

$$(3.3.1) \quad \begin{cases} P_1(y_1) \in k[[t]][y_1], \\ \vdots \\ P_n(y_1, \dots, y_n) \in k[[t]][y_1, \dots, y_n], \end{cases}$$

such that

- (1) *the ideal $(P_1(y_1), \dots, P_n(y_1, \dots, y_n))$ in $k[[t]][y_1, \dots, y_n]$ defines \mathfrak{Z} ,*
- (2) *for each i , the polynomial P_i is monic in y_i . When $i = 1$, the constant term of P_1 is in $k[[t]]^\times$, while for $2 \leq i \leq n$ and for the image $\bar{P}_i(y_i)$ of P_i in the quotient ring $\left(\frac{k[[t]][y_1, \dots, y_{i-1}]}{(P_1, \dots, P_{i-1})}\right)[y_i]$, its constant term is a unit in $\left(\frac{k[[t]][y_1, \dots, y_{i-1}]}{(P_1, \dots, P_{i-1})}\right)^\times$, and*
- (3) *for $2 \leq i \leq n$, for P_i regarded as a polynomial in y_i , its coefficients in $k[[t]][y_1, \dots, y_{i-1}]$ have their y_j -degrees strictly less than $\deg_{y_j} P_j$ for all $1 \leq j < i$.*

Proof. Let $\eta \in \mathrm{Spec}(k[[t]])$ be the generic point. By Corollary 3.2.10, the generic fiber \mathfrak{Z}_η is given by a maximal ideal $\mathfrak{m} \subset T_n = k((t))\{y_1, \dots, y_n\}$.

Applying Lemma 3.1.1 to \mathfrak{m} with $z_i = y_i$ for $1 \leq i \leq n$, we obtain a triangular shaped set of polynomials $p_1(y_1) \in k[[t]][y_1]$, $p_2(y_1, y_2) \in k((t))[y_1, y_2]$, \dots , $p_n(y_1, \dots, y_n) \in k((t))[y_1, \dots, y_n]$ that generate the maximal ideal $\mathfrak{m} \subset T_n$, and satisfy the properties there in the lemma.

To obtain the generators for its closure \mathfrak{Z} , for each $1 \leq i \leq n$ we clear the denominators of the coefficients in $k((t))$ of the terms of p_i by the l.c.m. of them in $k[[t]]$. We can do it because $k[[t]]$ is a UFD. So obtained elements are denoted by $P_1(y_1) \in k[[t]][y_1]$, \dots , $P_n(y_1, \dots, y_n) \in k[[t]][y_1, \dots, y_n]$, and they satisfy the properties that

- (a) *the ideal (P_1, \dots, P_n) generated in $k[[t]][y_1, \dots, y_n]$ defines \mathfrak{Z} ,*
- (b) *for each i , the highest y_i -degree term of P_i involves no other variable, and*
- (c) *for $2 \leq i \leq n$, when P_i is regarded as a polynomial in y_i , each of its coefficients in $k[[t]][y_1, \dots, y_{i-1}]$ has its y_j -degree strictly less than $\deg_{y_j} P_j$ for all $1 \leq j < i$.*

The conditions (1) and (3) of the proposition are satisfied by the above properties (a) and (c), but the condition (2) is not yet achieved. We show that (2) can be achieved for \mathfrak{Z} .

For $1 \leq i \leq n$, write $S_i := k[[t]]\{y_1, \dots, y_i\}$. Let $\mathfrak{p}_n \subset S_n$ be the prime ideal that defines \mathfrak{Z} (by Corollary 3.2.9). For the sequence of injective ring homomorphisms

$$k[[t]] \hookrightarrow S_1 \hookrightarrow \dots \hookrightarrow S_{n-1} \hookrightarrow S_n,$$

consider the prime ideals $\mathfrak{p}_i := \mathfrak{p}_n \cap S_i$ for $1 \leq i \leq n-1$. By construction, $\mathfrak{p}_i = (P_1, \dots, P_i)$, and

$$k[[t]] \hookrightarrow S_1/\mathfrak{p}_1 \hookrightarrow \dots \hookrightarrow S_{n-1}/\mathfrak{p}_{n-1} \hookrightarrow S_n/\mathfrak{p}_n.$$

Since $k[[t]] \rightarrow S_n/\mathfrak{p}_n$ is finite and free (Corollary 3.2.6), and $k[[t]]$ is a PID, all homomorphism $k[[t]] \rightarrow S_i/\mathfrak{p}_i$ for $1 \leq i \leq n$ are also finite and free because a submodule of a free module of finite rank over a PID is again free by the fundamental theorem on finitely generated modules over a PID.

We let $\mathfrak{Z}^{(i)} \subset \mathrm{Spf}(k[[t]]) \times \square^i$ be the integral closed formal subscheme given by $\mathrm{Spf}(S_i/\mathfrak{p}_i)$. By construction, its dimension is equal to that of \mathfrak{Z} , so that its codimension is i . This $\mathfrak{Z}^{(i)}$ is equal to the image of \mathfrak{Z} under the projection $pr_i : \mathrm{Spf}(k[[t]]) \times \square^n \rightarrow \mathrm{Spf}(k[[t]]) \times \square^i$, that sends $(y_1, \dots, y_n) \mapsto (y_1, \dots, y_i)$.

Claim: $\mathfrak{Z}^{(i)} \in z^i(\mathrm{Spf}(k[[t]]), i)$ for $1 \leq i \leq n$.

For each codimension 1 face $F = F_j^\varepsilon \subset \square_k^n$ with $j \leq i$, the intersection $\mathfrak{Z}^{(i)} \cap (\widehat{X} \times F)$ is the image of $\mathfrak{Z} \cap (\widehat{X} \times F)$ under pr_i . But the latter is empty by Lemma 3.2.4, thus so is the former. This means $\mathfrak{Z}^{(i)}$ has the empty intersection with all proper faces. In particular, the general position condition (**GP**) of Definition 2.2.2 for $\mathfrak{Z}^{(i)}$ holds trivially.

Since $\mathfrak{Z}^{(i)}$ has the empty intersection with all proper faces, to check the condition (**SF**) for $\mathfrak{Z}^{(i)}$, it remains to take $F = \square_k^n$, and we need to look at its special fiber. Since $\mathfrak{Z}^{(i)}$ is finite over $k[[t]]$ and the special fiber is given by its unique closed point by Lemma 3.2.5, the condition (**SF**) holds for $\mathfrak{Z}^{(i)}$. Thus $\mathfrak{Z}^{(i)} \in z^i(\mathrm{Spf}(k[[t]]), i)$.

We prove the remaining part (2) of the proposition.

Suppose $i = 1$. Then the intersection of $\mathfrak{Z}^{(1)}$ with the faces $\{y_1 = \infty\}$ and $\{y_1 = 0\}$ are empty by Lemma 3.2.4 and Claim. Thus the leading coefficient and the constant term of $P_1(y_1)$ in y_1 are both units in $k[[t]]^\times$. After scaling by the leading coefficient of $P_1(y_1)$ in y_1 , the leading coefficient becomes 1, and the constant term is still a unit in $k[[t]]^\times$, proving (2) for $i = 1$.

Let $i \geq 2$. Among the defining polynomials P_1, \dots, P_i of $\mathfrak{Z}^{(i)}$, the only one that involves the variable y_i is P_i . Furthermore, the highest y_i -degree term of P_i does not involve any other variables. Hence the empty intersection of $\mathfrak{Z}^{(i)}$ with the codimension 1 face $\{y_i = \infty\}$ (Lemma 3.2.4 and the above Claim) means that the leading coefficient of P_i in y_i is a unit in $k[[t]]^\times$. Hence after scaling by the leading coefficient of P_i in y_i , the leading coefficient becomes 1, so that we may assume P_i is monic in y_i .

Note that scaling by units of $k[[t]]^\times$ does not disturb the pre-existing properties (1), (3).

On the other hand, that the intersection of $\mathfrak{Z}^{(i)}$ with $\{y_i = 0\}$ is empty means that the image \bar{P}_i of P_i in the quotient ring $\left(\frac{k[[t]]\{y_1, \dots, y_{i-1}\}}{(P_1, \dots, P_{i-1})}\right)[y_i]$ has a unit constant term, when regarded as a polynomial in y_i .

Thus we have achieved all of (1), (2), (3). \square

4. THE GRAPH HOMOMORPHISMS AND THE REGULATORS

4.1. Milnor K -theory and the graph homomorphism. We discuss the graph homomorphisms from the Milnor K -groups. Recall from Elbaz-Vincent–Müller-Stach [12, Lemma 2.1] that for an equidimensional k -algebra R essentially of finite

type, there is the graph homomorphism

$$gr : K_n^M(R) \rightarrow \mathrm{CH}^n(\mathrm{Spec}(R), n).$$

We have a similar homomorphism for the higher Chow groups of some affine formal schemes. To avoid unnecessary complexities, we consider the relevant case only:

Lemma 4.1.1. *Consider $\widehat{X} = \mathrm{Spf}(k[[t]])$. For each Milnor symbol $\{a_1, \dots, a_n\} \in K_n^M(k[[t]])$, with $a_i \in k[[t]]^\times$, consider the integral closed formal subscheme $\Gamma_{(a_1, \dots, a_n)} \subset \square_{\widehat{X}}^n$ defined by the system of linear polynomials*

$$y_1 - a_1, \quad \dots, \quad y_n - a_n.$$

Then sending $\{a_1, \dots, a_n\}$ to the graph cycle $\Gamma_{(a_1, \dots, a_n)}$ defines the graph homomorphism

$$(4.1.1) \quad gr : K_n^M(k[[t]]) \rightarrow \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n).$$

Proof. One checks immediately that $\Gamma_{(a_1, \dots, a_n)} \in z^n(\mathrm{Spf}(k[[t]]), n)$. By Lemma 3.2.4, this represents a class in $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$.

The rest of the argument is essentially a repetition of [12, Lemma 2.1]. We remark that *loc.cit.* makes a general assumption that their ring is essentially of finite type, but this is not used there, so we can still follow a large part of the argument.

Firstly, for $a, a_i \in k[[t]]^\times$ such that $1 - a \in k[[t]]^\times$ for $3 \leq i \leq n$, consider the parametrized cycle given by the closure of the graph of the rational map

$$\mathfrak{W}_1 : \square_{\widehat{X}}^1 \dashrightarrow \square_{\widehat{X}}^{n+1}, \quad x \mapsto \left(x, 1 - x, \frac{a - x}{1 - x}, a_3, \dots, a_n \right).$$

One checks $\mathfrak{W}_1 \in z^n(\mathrm{Spf}(k[[t]]), n + 1)$. By straightforward calculations, one sees that the only nonzero face of \mathfrak{W}_1 is

$$\partial_3^0 \mathfrak{W}_1 = \Gamma_{(a, 1 - a, a_3, \dots, a_n)}.$$

In particular, the Steinberg symbol $\{a, 1 - a, a_3, \dots, a_n\} \in K_n^M(k[[t]])$ maps to 0 in $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$.

Secondly, for $a, b, a_i \in k[[t]]^\times$ for $2 \leq i \leq n$, consider the parametrized cycle

$$\mathfrak{W}_2 : \square_{\widehat{X}}^1 \dashrightarrow \square_{\widehat{X}}^{n+1}, \quad x \mapsto \left(x, \frac{ax - ab}{x - ab}, a_2, \dots, a_n \right).$$

One checks $\mathfrak{W}_2 \in z^n(\mathrm{Spf}(k[[t]]), n + 1)$, and that its only nonzero faces are

$$\begin{cases} \partial_1^\infty \mathfrak{W}_2 = \Gamma_{(a, a_2, \dots, a_n)}, \\ \partial_2^0 \mathfrak{W}_2 = \Gamma_{(b, a_2, \dots, a_n)}, \\ \partial_2^\infty \mathfrak{W}_2 = \Gamma_{(ab, a_2, \dots, a_n)}. \end{cases}$$

Hence killing $\partial \mathfrak{W}_2$ in $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$, we deduce

$$\Gamma_{(ab, a_2, \dots, a_n)} \equiv \Gamma_{(a, a_2, \dots, a_n)} + \Gamma_{(b, a_2, \dots, a_n)} \quad \text{in } \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n).$$

Considering various permutations of the cycle \mathfrak{W}_2 , we thus deduce

$$\bigotimes_{i=1}^n k[[t]]^\times \rightarrow \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n),$$

and then using the permutations of \mathfrak{W}_1 , it further descends to (4.1.1). \square

Lemma 4.1.2. *Let $m \geq 2$. The graph map (4.1.1) induces the homomorphism*

$$(4.1.2) \quad gr_{k_m} : K_n^M(k_m) \rightarrow \mathbf{CH}^n(\mathrm{Spec}(k_m), n),$$

where the codomain is the new higher Chow group defined in Definition 2.4.5.

Proof. Since $k[[t]]$ is a local ring, the natural homomorphism $k[[t]]^\times \rightarrow k_m^\times$ is surjective (see e.g. Hartshorne-Polini [22, Lemma 5.2]). Thus $K_n^M(k[[t]]) \rightarrow K_n^M(k_m)$ is also surjective. Hence to see that (4.1.1) descends to (4.1.2), it remains to check that $\ker(K_n^M(k[[t]]) \rightarrow K_n^M(k_m))$ is mapped to 0 under the graph map (4.1.1) composed with the natural map $\mathbf{CH}^n(\mathrm{Spf}(k[[t]]), n) \rightarrow \mathbf{CH}^n(\mathrm{Spec}(k_m), n)$.

Since the above kernel is generated by elements of the form $\{1 + t^m c, c_2, \dots, c_n\}$, where $c \in k[[t]]$ and $c_i \in k[[t]]^\times$ for $2 \leq i \leq n$ (see Kato-Saito [29, Lemma 1.3.1, p.261]), it is enough to show that

$$(4.1.3) \quad \Gamma_{(1+t^m c, c_2, \dots, c_n)} \equiv 0 \quad \text{in } \mathbf{CH}^n(\mathrm{Spec}(k_m), n),$$

where $\Gamma_{(1+t^m c, c_2, \dots, c_n)}$ is the graph cycle in $z^n(\widehat{X}, n)$ of the above Milnor symbol, defined in Lemma 4.1.1.

Before we do so, we first prove:

Claim: *Let $a_i, b_i \in k[[t]]^\times$ such that $a_i - b_i \in (t^m)k[[t]]$ for $1 \leq i \leq n$. Then have*

$$\Gamma_a - \Gamma_b \in \mathcal{M}^n(\widehat{X}, Y, n),$$

where $\Gamma_a := \Gamma_{(a_1, \dots, a_n)}$ and $\Gamma_b := \Gamma_{(b_1, \dots, b_n)}$ denote the corresponding integral graph cycles in $z^n(\widehat{X}, n)$. In particular, $\Gamma_a \equiv \Gamma_b$ in $\mathbf{CH}^n(\mathrm{Spec}(k_m), n)$.

This claim is immediate because we have an isomorphism

$$\mathcal{O}_{\Gamma_a} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n} \simeq \mathcal{O}_{\Gamma_b} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n}$$

of $\mathcal{O}_{\square_Y^n}$ -algebras.

Going back to the proof of (4.1.3), let $d_i := \bar{c}_i \in k^\times$ for $2 \leq i \leq n$. Via the obvious injective homomorphism $k^\times \hookrightarrow k[[t]]^\times$, regard $d_i \in k[[t]]^\times$. Then we have

$$1 + t^m c \equiv 1, \quad c_2 \equiv d_2, \quad \dots, \quad c_n \equiv d_n \quad \text{mod } (t^m)k[[t]].$$

Thus by the Claim, we have

$$(4.1.4) \quad \Gamma_{(1+t^m c, c_2, \dots, c_n)} \equiv \Gamma_{(1, d_2, \dots, d_n)} \quad \text{in } \mathbf{CH}^n(\mathrm{Spec}(k_m), n).$$

However, $\square = \mathbb{P}^1 \setminus \{1\}$ so that $\Gamma_{(1, d_2, \dots, d_n)} = \emptyset$ in $\square_{\widehat{X}}^n$. Thus (4.1.4) proves (4.1.3). This induces (4.1.2). \square

The goal of this article, Theorem 1.2.1, is to prove that (4.1.2) is an isomorphism when $|k| \gg 0$. It will be completed in Theorem 5.2.6.

4.2. Regulators via rigid analytic spaces. In §4.2, we construct a homomorphism

$$(4.2.1) \quad \widehat{\Upsilon} : \mathbf{CH}^n(\mathrm{Spf}(k[[t]]), n) \rightarrow K_n^M(k((t))).$$

Under the additional assumption that $|k| \gg 0$, the homomorphism $K_n^M(k[[t]]) \rightarrow K_n^M(k((t)))$ is injective (Gersten conjecture for K_n^M proven by M. Kerz [30, Proposition 10, p.181]). We will see that the image of $\widehat{\Upsilon}$ lies in $K_n^M(k[[t]])$.

To facilitate our discussions, we introduce a rigid analytic analogue of higher Chow groups in the Milnor range (see also J. Ayoub [2, Notations 2.4.14, p.297 and beyond]):

Definition 4.2.1. Let K be a complete discrete valued field with a nontrivial non-archimedean norm. Let K° be its valuation ring and suppose $k \subset K^\circ$. We define the rigid analytic analogue of higher Chow groups for $\mathrm{Sp}(K)$ for the Milnor range, but its generalization to the off-Milnor ranges is similar.

We consider the rigid analytic space $\mathrm{Sp}(K) \times \square_k^n$. (The fiber product exists by Bosch-Güntzer-Remmert [7, §7.1.4, Proposition 4, p.268].) We define $z^n(\mathrm{Sp}(K), n)$

to be the free abelian group on the codimension n points not lying on $\{y_i = 0, 1, \infty\}$ for $1 \leq i \leq n$, and define $\underline{z}^n(\mathrm{Sp}(K), n+1)$ be the free abelian group on the codimension n integral closed subsets that intersect all faces of $\mathrm{Sp}(K) \times \square_k^{n+1}$ properly, where the faces are similarly defined as in §2.2. We define $z^n(\mathrm{Sp}(K), *)$ for $* = n, n+1$ by modding out $\underline{z}^n(\mathrm{Sp}(K), *)$ by the degenerate cycles. Via the closed immersions $\iota_i^\epsilon : \mathrm{Sp}(K) \times \square_k^n \hookrightarrow \mathrm{Sp}(K) \times \square_k^{n+1}$ for $1 \leq i \leq n+1$ and $\epsilon = 0, \infty$, we have the natural boundary map $\partial : z^n(\mathrm{Sp}(K), n+1) \rightarrow z^n(\mathrm{Sp}(K), n)$ defined by $\partial = \sum_{i=1}^{n+1} (-1)^i (\partial_i^\infty - \partial_i^0)$, where $\partial_i^\epsilon = (\iota_i^\epsilon)^*$. Its cokernel is denoted by $\mathrm{CH}^n(\mathrm{Sp}(K), n)$. \square

Example 4.2.2. If we take $K = k((t))$ and an integral cycle $\mathfrak{Z} \in \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$, then the generic fiber \mathfrak{Z}_η gives a member of $\mathrm{CH}^n(\mathrm{Sp}(k((t))), n)$. With Corollaries 3.2.9 and 3.2.10 in mind, note that the localization homomorphisms $k[[t]]\{y_1, \dots, y_n\} \rightarrow k((t))\{y_1, \dots, y_n\}$ over $n \geq 1$ induce the commutative diagram

$$\begin{array}{ccc} z^n(\mathrm{Spf}(k[[t]]), n+1) & \longrightarrow & z^n(\mathrm{Sp}(k((t))), n+1) \\ \downarrow \partial & & \downarrow \partial \\ z^n(\mathrm{Spf}(k[[t]]), n) & \longrightarrow & z^n(\mathrm{Sp}(k((t))), n), \end{array}$$

from which we deduce $\eta : \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \rightarrow \mathrm{CH}^n(\mathrm{Sp}(k((t))), n)$. \square

What follows below is essentially an argument in B. Totaro [47, §3], with some suitable modifications to fit it into the rigid analytic situation being considered.

Definition 4.2.3. Let $\mathfrak{p} \in z^n(\mathrm{Sp}(k((t))), n)$ be a point. It gives a maximal ideal \mathfrak{m} of $T_n = k((t))\{y_1, \dots, y_n\}$ (cf. Corollary 3.2.10). By Lemma 2.5.1-(3), the injection $k((t)) \hookrightarrow L := T_n/\mathfrak{m}$ is a finite extension of fields. This point \mathfrak{p} defines an n -tuple (x_1, \dots, x_n) whose values lie in $L \setminus \{0, 1\}$, namely $x_i = \bar{y}_i$ in $T_n/\mathfrak{m} = L$. We consider the Milnor symbol $\{x_1, \dots, x_n\} \in K_n^M(L)$.

Define

$$(4.2.2) \quad \tilde{\Upsilon}(\mathfrak{p}) := N_{L/k((t))}\{x_1, \dots, x_n\} \in K_n^M(k((t))),$$

via the norm $N_{L/k((t))} : K_n^M(L) \rightarrow K_n^M(k((t)))$ of Bass-Tate [3] and K. Kato [28]. \square

Lemma 4.2.4. *The composite*

$$z^n(\mathrm{Sp}(k((t))), n+1) \xrightarrow{\partial} z^n(\mathrm{Sp}(k((t))), n) \xrightarrow{\tilde{\Upsilon}} K_n^M(k((t)))$$

is zero. In particular, we have the induced homomorphism

$$(4.2.3) \quad \tilde{\Upsilon} : \mathrm{CH}^n(\mathrm{Sp}(k((t))), n) \rightarrow K_n^M(k((t))).$$

Proof. We continue to follow B. Totaro [47, §3] with suitable modifications.

Let $C \in z^n(\mathrm{Sp}(k((t))), n+1)$ be an integral cycle. Take the normalization $D \rightarrow C$; since C is defined by a 1-dimensional noetherian integral domain, it has a unique normalization (see B. Conrad [9] for normalizations of more general rigid analytic spaces). The normalization map is finite because every affinoid $k((t))$ -algebra that is an integral domain is a Japanese ring (see Bosch-Güntzer-Remmert [7, §6.1.2, Proposition 4, p.228] or EGA IV₂ [17, Scholie (7.8.3)-(vi), p.215]). As in [47, §3], using W. Fulton [15, Example 1.2.3, p.9], it is enough to show that the boundary (defined by pull-back of the faces and push-forward) of D maps to 0 in $K_n^M(k((t)))$.

Let \bar{D} be the unique regular compactification of D , which exists as $\dim D = 1$. By the rigid analytic GAGA (see e.g. [13, Theorem 4.10.5, p.113]), this \bar{D} can be regarded as a regular projective algebraic curve over $k((t))$. Let \mathbb{K} be the rational function field of \bar{D} , which is equal to that of C and D . Since $\mathbb{K} \supset k((t))$, it is an infinite field.

The map $D \rightarrow \mathrm{Sp}(k((t))) \times \square_k^{n+1}$ is given by $(n+1)$ rational functions $g_1, \dots, g_{n+1} \in \mathbb{K}$ on D . By the given proper intersection conditions with the faces, no g_i is identically equal to 0 or ∞ , and any closed point $w \in D$ with $g_i(w) = 0$ or ∞ has $g_j(w) \notin \{0, \infty\}$ for all $j \neq i$.

For each closed point $w \in \overline{D}$, there is its associated discrete valuation on \mathbb{K} with the associated boundary map $\partial_w : K_{n+1}^M(\mathbb{K}) \rightarrow K_n^M(\kappa(w))$ (see J. Milnor [35, Lemma 2.1]), where $\kappa(w)$ is the residue field of w .

Since \mathbb{K} is infinite, by Suslin reciprocity (see [45]), for the Milnor symbol $\{g_1, \dots, g_{n+1}\} \in K_{n+1}^M(\mathbb{K})$, we have

$$\sum_{w \in \overline{D}} N_{\kappa(w)/k((t))} \partial_w \{g_1, \dots, g_{n+1}\} = 0 \quad \text{in } K_n^M(k((t))).$$

If $w \in \overline{D} \setminus D$, then one of g_i has $g_i(w) = 1$ so that $\partial_w \{g_1, \dots, g_{n+1}\} = 0$. Hence the above sum can be written without such points, and

$$(4.2.4) \quad \sum_{w \in D} N_{\kappa(w)/k((t))} \partial_w \{g_1, \dots, g_{n+1}\} = 0 \quad \text{in } K_n^M(k((t))).$$

On the other hand, by following the definition of ∂_w , we have the commutative diagram

$$\begin{array}{ccc} z^n(\mathrm{Sp}(k((t))), n+1) & \xrightarrow{\partial} & z^n(\mathrm{Sp}(k((t))), n) \\ \downarrow & & \downarrow \tilde{\Upsilon} \\ K_{n+1}^M(\mathbb{K}) & \xrightarrow{\sum_w N_{\kappa(w)/k((t))} \partial_w} & K_n^M(k((t))), \end{array}$$

where the left vertical arrow associates C to the Milnor symbol $\{g_1, \dots, g_{n+1}\}$ in $K_{n+1}^M(\mathbb{K})$. Thus $\tilde{\Upsilon} \circ \partial C$ is equal to (4.2.4), which is 0. Thus, we deduce the map (4.2.3) as desired. \square

Definition 4.2.5. We define the map $\hat{\Upsilon}$ as the composite

$$\hat{\Upsilon} : \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \xrightarrow{\eta} \mathrm{CH}^n(\mathrm{Sp}(k((t))), n) \xrightarrow{\tilde{\Upsilon}} K_n^M(k((t))),$$

where η is the localization map (Example 4.2.2) and $\tilde{\Upsilon}$ is from Definition 4.2.3 and Lemma 4.2.4. \square

Lemma 4.2.6. *Let k be a field with $|k| \gg 0$ so that $K_n^M(k[[t]]) = \hat{K}_n^M(k[[t]])$.*

Then the image of the composite

$$K_n^M(k[[t]]) \xrightarrow{gr} \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \xrightarrow{\hat{\Upsilon}} K_n^M(k((t))),$$

lands into the subgroup $K_n^M(k[[t]]) \subset K_n^M(k((t)))$, and the composite is the identity on $K_n^M(k[[t]])$.

In particular, the map gr is injective and the image of $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$ under $\hat{\Upsilon}$ is in the subgroup $K_n^M(k[[t]])$ of $K_n^M(k((t)))$.

Before the proof, we remark that the condition $K_n^M(k[[t]]) = \hat{K}_n^M(k[[t]])$ implies that the natural homomorphism $K_n^M(k[[t]]) \rightarrow K_n^M(k((t)))$ is injective (see M. Kerz [30, Proposition 10, p.181]), thus we can regard $K_n^M(k[[t]]) \subset K_n^M(k((t)))$.

When $n = 1$, this always holds for any field k .

Proof. Consider the Milnor symbol $\{a_1, \dots, a_n\} \in K_n^M(k[[t]])$ with $a_i \in k[[t]]^\times$. Its image under the first map gr is given by the integral graph cycle \mathfrak{Z} defined by the system of polynomials

$$y_1 - a_1, \quad \dots, \quad y_n - a_n.$$

Its coordinates are given by $a_1, \dots, a_n \in k[[t]]^\times \subset k((t))^\times$, and they are of degree 1 over $k((t))$. Hence the generic fiber \mathfrak{Z}_η gives $L = k((t))$ in the notations of Definition

4.2.3, and the norm map in (4.2.2) is the identity map. Hence $\tilde{\Upsilon}(\mathfrak{Z}) = \{a_1, \dots, a_n\}$ and it belongs to the subgroup $K_n^M(k[[t]])$. This shows that the composite is the identity map of $K_n^M(k[[t]])$. In particular, gr is injective.

Since the image of the composite lies in $K_n^M(k[[t]])$, so does the image of the group $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$. \square

The following is a rigid analytic analogue of [47]:

Corollary 4.2.7. *For any field k , the composite*

$$K_n^M(k((t))) \xrightarrow{gr} \mathrm{CH}^n(\mathrm{Sp}(k((t))), n) \xrightarrow{\tilde{\Upsilon}} K_n^M(k((t)))$$

is the identity map, and all arrows are isomorphisms.

Proof. For the injectivity of gr , we simply note that the composite $\tilde{\Upsilon} \circ gr$ evaluated at the Milnor symbols $\{a_1, \dots, a_n\} \in K_n^M(k((t)))$ for some $a_i \in (k((t)))^\times$ is again $\{a_1, \dots, a_n\}$. Thus $\tilde{\Upsilon} \circ gr = \mathrm{Id}$. In particular, gr is injective.

For the surjectivity of gr , we may follow B. Totaro [47, §4]. When F is a field, to prove that the graph homomorphism $K_n^M(F) \rightarrow \mathrm{CH}^n(\mathrm{Spec}(F), n)$ to the ordinary cubical higher Chow group, is surjective, Totaro used the norm $N : L^\times \rightarrow F^\times$ to relate the class of a given closed point to a cycle given by F -rational points, where L is the residue field of the point.

Specializing to the case $F = k((t))$ (which is infinite), we can repeat the argument of Totaro *mutatis mutandis*. As this corollary does not play an important role for the rest of the article, we omit details. \square

4.3. A comparison of equivalence relations. We continue to use the notations $Y = \mathrm{Spec}(k_m)$ and $\widehat{X} = \mathrm{Spf}(k[[t]])$. The closed immersion $Y \hookrightarrow \widehat{X}$ is given by the ideal $I := (t^m) \subset k[[t]]$ so that $k[[t]]/I = k_m$.

Recall from Remark 2.4.3 that, we have the mod Y -equivalence \sim_Y on $z^n(\widehat{X}, n)$ given by the subgroup $\mathcal{M}^n(\widehat{X}, Y, n)$, and we have another equivalence \sim_I given by the subgroup $\mathcal{N}^n(\widehat{X}, Y, n)$ generated by the differences of a pair of integral cycles that are equal after taking the mod I -reductions. We prove in Lemma 4.3.1 below that these two are equivalent in the Milnor range. It will simplify our discussions from §4.4.

Lemma 4.3.1. *We have the equality*

$$(4.3.1) \quad \mathcal{M}^n(\widehat{X}, Y, n) = \mathcal{N}^n(\widehat{X}, Y, n)$$

of the subgroups in $z^n(\widehat{X}, n)$.

Proof. For each generator $[\mathfrak{Z}_1] - [\mathfrak{Z}_2] \in \mathcal{N}^n(\widehat{X}, Y, n)$ associated to a pair of integral cycles \mathfrak{Z}_1 and $\mathfrak{Z}_2 \in z^n(\widehat{X}, n)$, by definition we have

$$\mathcal{O}_{\mathfrak{Z}_1} \otimes_{\mathcal{O}_{\widehat{X}}} \mathcal{O}_{\square_Y^n} \simeq \mathcal{O}_{\mathfrak{Z}_2} \otimes_{\mathcal{O}_{\widehat{X}}} \mathcal{O}_{\square_Y^n}$$

as $\mathcal{O}_{\square_Y^n}$ -algebras. Hence $[\mathfrak{Z}_1] - [\mathfrak{Z}_2] \in \mathcal{M}^n(\widehat{X}, Y, n)$, and this shows $\mathcal{M}^n(\widehat{X}, Y, n) \supset \mathcal{N}^n(\widehat{X}, Y, n)$. It remains to prove the opposite inclusion.

Let $(\mathcal{A}_1, \mathcal{A}_2) \in \mathcal{L}^n(\widehat{X}, Y, n)$. We have an isomorphism of $\mathcal{O}_{\square_Y^n}$ -algebras

$$(4.3.2) \quad \mathcal{A}_1 \otimes_{\mathcal{O}_{\widehat{X}}} \mathcal{O}_{\square_Y^n} \simeq \mathcal{A}_2 \otimes_{\mathcal{O}_{\widehat{X}}} \mathcal{O}_{\square_Y^n}.$$

Let B_j be the ring of the global sections of \mathcal{A}_j . Since $[\mathcal{A}_j] \in z^n(\widehat{X}, n)$, by Corollary 3.2.3 we see that B_j is a $k[[t]]$ -algebra which is a finite $k[[t]]$ -module. In particular, it is a semi-local $k[[t]]$ -algebra.

Having Corollary 3.2.9 in mind, the $k_m[y_1, \dots, y_n]$ -algebra isomorphism (4.3.2) in particular implies that we have an isomorphism of Artin rings

$$(4.3.3) \quad B_1/(t^m)B_1 \simeq B_2/(t^m)B_2.$$

By the structure theorem for Artin rings (Atiyah-MacDonald [1, Theorem 8.7, p.90] or D. Eisenbud [11, Corollary 2.16, p.76]), each Artin ring decomposes uniquely into a product of Artin local rings, up to isomorphism. So, both sides of (4.3.3) decompose uniquely into products of Artin local rings, and under the isomorphism, there is a 1-1 correspondence between the Artin local factors of both sides.

On the other hand, since $k[[t]]$ is henselian and complete, by Lemma 3.2.5, each B_j decomposes into a direct product $\prod_{i=1}^{r_j} B_{ji}$ of complete local domains, one for each maximal ideal, so that each Artin local factor of $B_j/(t^m)B_j$ corresponds to a local domain factor of B_j .

The above two paragraphs summarize into the following 1-1 correspondences:

$$\begin{array}{ccc} \left\{ \begin{array}{c} \text{Artin local} \\ \text{factors of } B_1/(t^m)B_1 \end{array} \right\} & \xleftrightarrow{1-1} & \left\{ \begin{array}{c} \text{Artin local} \\ \text{factors of } B_2/(t^m)B_2 \end{array} \right\} \\ \uparrow \text{1-1} & & \uparrow \text{1-1} \\ \left\{ \begin{array}{c} \text{local factors} \\ \text{of } B_1 \end{array} \right\} & & \left\{ \begin{array}{c} \text{local factors} \\ \text{of } B_2 \end{array} \right\} \end{array}$$

In particular, we have the equality of the number of the local factors $r_1 = r_2 =: r$.

For each fixed j , after relabeling B_{ji} over i if necessary, the above discussion also shows that we have an isomorphism of $k_m[y_1, \dots, y_n]$ -algebras

$$(4.3.4) \quad \phi_m : B_{1i}/(t^m)B_{1i} \simeq B_{2i}/(t^m)B_{2i}$$

for all $1 \leq i \leq r$.

Let \mathcal{A}_{ji} be the coherent $\mathcal{O}_{\square_{\bar{x}}^n}$ -algebra corresponding to B_{ji} . Let $f_{ji} : \text{Spec}(\mathcal{A}_{ji}) = \text{Spec}(B_{ji}) \rightarrow \text{Spec}(k[[t]]\{y_1, \dots, y_n\})$ be the corresponding finite morphism (R. Hartshorne [21, Exercise II-5.17-(c),(d), p.128] or more generally Malgoire-Voisin [34, 1.4.2, 1.4.3]). Here we have $\mathcal{A}_{ji} \simeq f_{ji*}\mathcal{O}_{\text{Spec}(B_{ji})}$ so that $[\mathcal{A}_{ji}] = f_{ji*}[\text{Spec}(B_{ji})] = d_{ji}[f_{ji}(\text{Spec}(B_{ji}))]$, where d_{ji} is the degree of f_{ji} .

Note that $\mathfrak{Z}_{ji} := f_{ji}(\text{Spec}(B_{ji}))$ is integral and closed in $\text{Spec}(k[[t]]\{y_1, \dots, y_n\})$, and its defining ideal is the kernel of the ring homomorphism $f_{ji}^\# : k[[t]]\{y_1, \dots, y_n\} \rightarrow B_{ji}$. Let $C_{ji} = \text{Im}(f_{ji}^\#)$. By the first isomorphism theorem, this C_{ji} is the coordinate ring of \mathfrak{Z}_{ji} . We have the natural factorization

$$f_{ji}^\# : k[[t]]\{y_1, \dots, y_n\} \twoheadrightarrow C_{ji} \hookrightarrow B_{ji}.$$

Going mod $I = (t^m)$, we have a commutative diagram of the induced ring homomorphisms, where ϕ_m is the isomorphism of (4.3.4) and $\bar{f}_{ji}^\#$ is the induced homomorphism for $f_{ji}^\#$

$$\begin{array}{ccc} k_m[y_1, \dots, y_n] & \xrightarrow{\bar{f}_{1i}^\#} & B_{1i}/(t^m)B_{1i} \\ & \searrow \bar{f}_{2i}^\# & \simeq \downarrow \phi_m \\ & & B_{2i}/(t^m)B_{2i}. \end{array}$$

As C_{ji} is the image of $f_{ji}^\#$, $C_{ji}/(t^m)C_{ji}$ is also the image of $\bar{f}_{ji}^\#$. In particular, under the identification ϕ_m , we have an isomorphism of $k_m[y_1, \dots, y_n]$ -algebras

$$(4.3.5) \quad C_{1i}/(t^m)C_{1i} \simeq C_{2i}/(t^m)C_{2i}.$$

This is equivalent to having an $\mathcal{O}_{\square_Y^n}$ -algebra isomorphism

$$\mathcal{O}_{\mathfrak{Z}_{1i}} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n} \simeq \mathcal{O}_{\mathfrak{Z}_{2i}} \otimes_{\mathcal{O}_{\square_{\widehat{X}}^n}} \mathcal{O}_{\square_Y^n}.$$

In particular, $[\mathfrak{Z}_{1i}] - [\mathfrak{Z}_{2i}] \in \mathcal{N}^n(\widehat{X}, Y, n)$.

On the other hand, B_{ji} is a flat $k[[t]]$ -algebra (as there is no t -torsion, and $k[[t]]$ is a PID), so that the degree d_{ji} can be computed by specializing $f_{ji}^\# \bmod t$. We have the field extensions $k \hookrightarrow C_{ji}/(t) \hookrightarrow B_{ji}/(t)$, and $d_{ji} = [B_{ji}/(t) : C_{ji}/(t)]$. Since $B_{1j}/(t) \simeq B_{2j}/(t)$ and $C_{1j}/(t) \simeq C_{2j}/(t)$ by (4.3.4) and (4.3.5), plugging in 1 into m , we deduce that $d_{1i} = d_{2i} =: d_i$.

Thus combining the above discussions, we have

$$[\mathcal{A}_1] - [\mathcal{A}_2] = \sum_{i=1}^r ([\mathcal{A}_{1i}] - [\mathcal{A}_{2i}]) = \sum_{i=1}^r d_i ([\mathfrak{Z}_{1i}] - [\mathfrak{Z}_{2i}]) \in \mathcal{N}^n(\widehat{X}, Y, n),$$

thus $\mathcal{M}^n(\widehat{X}, Y, n) \subset \mathcal{N}^n(\widehat{X}, Y, n)$. This proves the equality (4.3.1). \square

So, for the cycles in the Milnor range over $\mathrm{Spf}(k[[t]])$, we may use the simpler equivalence \sim_I , if needed. We do so in §4.4 and beyond.

4.4. An argument for pairs of cycles. The goal of §4.4 is to prove the following:

Proposition 4.4.1. *Let $m \geq 2$. Let $\widehat{X} = \mathrm{Spf}(k[[t]])$ and $I = (t^m) \subset k[[t]]$. Let \mathfrak{Z}_1 and $\mathfrak{Z}_2 \in z^n(\widehat{X}, n)$ be integral cycles such that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$.*

Then there exist cycles $E_{\mathfrak{Z}_1}$ and $E_{\mathfrak{Z}_2} \in z^n(\widehat{X}, n+1)$ such that

$$\partial E_{\mathfrak{Z}_\ell} = \mathfrak{Z}_\ell - \mathfrak{Z}'_\ell, \quad \ell = 1, 2$$

for some positive integer multiples of integral graph cycles, \mathfrak{Z}'_1 and $\mathfrak{Z}'_2 \in z^n(\widehat{X}, n)$, satisfying $\mathfrak{Z}'_1 \sim_I \mathfrak{Z}'_2$.

Once proven, we will have two applications. The first is the surjectivity of the graph homomorphism (4.1.1), to be discussed in §5.1. The second pertains to showing that gr_{k_m} is an isomorphism, discussed in §5.2.

Coming back to the proof of Proposition 4.4.1, let $\mathfrak{Z} \in z^n(\widehat{X}, n)$ be an integral cycle. By Corollary 3.2.9, this is given by a prime ideal $\mathfrak{p} \subset k[[t]]\{y_1, \dots, y_n\}$ of height n . For $1 \leq i \leq n$, we let \mathfrak{p}_i be the prime ideal $\mathfrak{p} \cap k[[t]]\{y_1, \dots, y_i\}$. This corresponds to the image $\mathfrak{Z}^{(i)}$ of \mathfrak{Z} under the projection $\mathrm{Spf}(k[[t]]) \times \square_k^n \rightarrow \mathrm{Spf}(k[[t]]) \times \square_k^i$ that sends $(y_1, \dots, y_n) \mapsto (y_1, \dots, y_i)$. For $i = 0$, we let $\mathfrak{Z}^{(0)} = \mathrm{Spf}(k[[t]])$ for our notational conveniences. We have $\mathfrak{Z}^{(i)} \in z^i(\widehat{X}, i)$ for each $1 \leq i \leq n$ as seen before, in the Claim in the middle of the proof of Proposition 3.3.1.

Definition 4.4.2. For an integral $\mathfrak{Z} \in z^n(\widehat{X}, n)$, for each pair $0 \leq i < j \leq n$ of indices, the induced morphism $f^{j/i} : \mathfrak{Z}^{(j)} \rightarrow \mathfrak{Z}^{(i)}$ is a finite surjective morphism of integral formal schemes. Define $d^{j/i}(\mathfrak{Z}) := \deg(f^{j/i} : \mathfrak{Z}^{(j)} \rightarrow \mathfrak{Z}^{(i)})$.

Observe the following, whose proof is obvious by definition:

Lemma 4.4.3. *The following are equivalent:*

- (1) \mathfrak{Z} is a graph cycle.
- (2) $d^{n/0}(\mathfrak{Z}) = 1$
- (3) For each $1 \leq i \leq n$, we have $d^{i/(i-1)}(\mathfrak{Z}) = 1$.

We will show that any integral cycle $\mathfrak{Z} \in z^n(\widehat{X}, n)$ can be turned into an integer multiple of an integral graph cycle modulo the boundary of a cycle in $z^n(\widehat{X}, n+1)$, by eventually achieving Lemma 4.4.3-(3). In the process, we will show that each

step of this construction respects the mod I -equivalence. This requires the following discussions for pairs of cycles.

Let $\mathfrak{Z}_\ell \in z^n(\widehat{X}, n)$ be two integral cycles for $\ell = 1, 2$ such that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$. For each ℓ , by Proposition 3.3.1, we have a system of polynomials in y_1, \dots, y_n for \mathfrak{Z}_ℓ

$$(4.4.1) \quad \begin{cases} P_\ell^{(1)}(y_1) \in k[[t]][y_1], \\ P_\ell^{(2)}(y_1, y_2) \in k[[t]][y_1, y_2], \\ \vdots \\ P_\ell^{(n)}(y_1, \dots, y_n) \in k[[t]][y_1, \dots, y_n], \end{cases}$$

such that all the properties in Proposition 3.3.1 hold. Note that $\mathfrak{Z}_1^{(i)} \sim_I \mathfrak{Z}_2^{(i)}$ for each $1 \leq i \leq n$ as well. Furthermore we have:

Lemma 4.4.4. *Under the above notations and assumptions, let $\bar{P}_\ell^{(1)} := P_\ell^{(1)}$ and $R_\ell^{(0)} := k[[t]]$, while for $2 \leq i \leq n$, let $\bar{P}_\ell^{(i)}$ be the image of $P_\ell^{(i)}(y_1, \dots, y_i)$ in $R_\ell^{(i-1)}[y_i]$, where*

$$(4.4.2) \quad R_\ell^{(i-1)} := k[[t]]\{y_1, \dots, y_{i-1}\} / (P_\ell^{(1)}, \dots, P_\ell^{(i-1)}).$$

For each $1 \leq i \leq n$ and each $\ell = 1, 2$, let $d_\ell^i := \deg_{y_i} \bar{P}_\ell^{(i)}$. Let $(-1)^{d_\ell^i} c_\ell^{(i)} \in R_\ell^{(i-1)}$ be the constant term of $\bar{P}_\ell^{(i)}$. Let $\bar{R}_\ell^{(i-1)} = R_\ell^{(i-1)} / IR_\ell^{(i-1)}$.

Then we have:

- (1) $d_1^i = d_2^i$.
- (2) $\bar{R}_1^{(i-1)} = \bar{R}_2^{(i-1)}$.
- (3) In the common ring $\bar{R}^{(i-1)}$ of the above, we have $c_1^{(i)} \equiv c_2^{(i)}$.

Note that by Proposition 3.3.1-(2), we have $c_\ell^{(i)} \in (R_\ell^{(i-1)})^\times$.

Proof. We prove (1), (2), (3) at the same time. By the properties of Proposition 3.3.1 for $P_\ell^{(i)}$, we know that $d_\ell^i = \deg_{y_i} \bar{P}_\ell^{(i)} \geq 1$. Going further down mod I , let $\bar{\bar{P}}_\ell^{(i)}$ be the image of $\bar{P}_\ell^{(i)}(y_i)$ in $\bar{\bar{R}}_\ell^{(i-1)}[y_i]$.

Note that since $\mathfrak{Z}_1^{(i-1)} \sim_I \mathfrak{Z}_2^{(i-1)}$, we have $\bar{\bar{R}}_1^{(i-1)} = \bar{\bar{R}}_2^{(i-1)}$, proving (2).

Since $\bar{P}_\ell^{(i)}(y_i)$ is monic in y_i in the ring $R_\ell^{(i-1)}[y_i]$ by Proposition 3.3.1, its further image $\bar{\bar{P}}_\ell^{(i)}(y_i)$ in $\bar{\bar{R}}_\ell^{(i-1)}[y_i]$ is also monic in y_i with the same y_i -degree. Thus

$$(4.4.3) \quad \deg_{y_i} \bar{P}_\ell^{(i)} = \deg_{y_i} \bar{\bar{P}}_\ell^{(i)}.$$

Furthermore, $\mathfrak{Z}_1^{(i)} \bmod I = \mathfrak{Z}_2^{(i)} \bmod I$ as closed subschemes of $\text{Spec}(\bar{\bar{R}}^{(i-1)}[y_i])$, and they are respectively given by the monic polynomials $\bar{\bar{P}}_\ell^{(i)}[y_i] \in \bar{\bar{R}}^{(i-1)}[y_i]$. Hence we have the equality

$$(4.4.4) \quad \bar{\bar{P}}_1^{(i)} = \bar{\bar{P}}_2^{(i)} \text{ in } \bar{\bar{R}}^{(i-1)}[y_i].$$

In particular,

$$(4.4.5) \quad \deg_{y_i} \bar{\bar{P}}_1^{(i)} = \deg_{y_i} \bar{\bar{P}}_2^{(i)}.$$

Hence, combining (4.4.3) and (4.4.5), we deduce (1).

On the other hand, from the equality (4.4.4), we have the equality of the constant terms $(-1)^{d_1^i} c_1^{(i)} \equiv (-1)^{d_2^i} c_2^{(i)}$ in $\bar{\bar{R}}^{(i-1)}$. Since $d_1^i = d_2^i$ by (1), we now have $c_1^{(i)} \equiv c_2^{(i)}$ in $\bar{\bar{R}}^{(i-1)}$, proving (3). \square

Definition 4.4.5. By Lemma 4.4.4-(1), $d_1^i = \deg_{y_i} \bar{\bar{P}}_1^{(i)} = \deg_{y_i} \bar{\bar{P}}_2^{(i)} = d_2^i$. We now simply call it d_i .

We remark that by definition, we have $d_i = d^{i/(i-1)}(\mathfrak{Z}_\ell)$ for $\ell = 1, 2$. \square

The following procedure will be used repeatedly.

Lemma 4.4.6. *Suppose $1 \leq i \leq n$. Let $\ell = 1, 2$. Let \mathfrak{Z}_1 and $\mathfrak{Z}_2 \in z^n(\widehat{X}, n)$ be two integral cycles such that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$, and*

(\star) *they have the generators as in (4.4.1) satisfying the properties of Proposition 3.3.1.*

Suppose $d^{j/(j-1)}(\mathfrak{Z}_\ell) = 1$ for $j > i$. (We allow $i = n$ as well, in which case there is no condition.)

Then there exist cycles $C_{\mathfrak{Z}_\ell} \in z^n(\widehat{X}, n+1)$ and positive integer multiples of integral cycles, \mathfrak{Z}'_1 and $\mathfrak{Z}'_2 \in z^n(\widehat{X}, n)$, such that for $\ell = 1, 2$, we have $\mathfrak{Z}_\ell - \mathfrak{Z}'_\ell = \partial(C_{\mathfrak{Z}_\ell})$, $\mathfrak{Z}'_1 \sim_I \mathfrak{Z}'_2$, the above (\star) holds for \mathfrak{Z}'_ℓ , and

$$d^{j/(j-1)}(\mathfrak{Z}'_\ell) = \begin{cases} \text{not necessarily } 1 & \text{for } j \geq i+1, \\ 1 & \text{for } j = i, \\ d^{j/(j-1)}(\mathfrak{Z}_\ell) & \text{for } j < i \end{cases}$$

Remark 4.4.7. Before we prove Lemma 4.4.6, let us give an illustration on how one can use it repeatedly to transform an integral cycle \mathfrak{Z} into a positive integer multiple of an integral graph cycle, \mathfrak{Z}' . For instance, take $n = 3$, and suppose that for an integral cycle \mathfrak{Z} , we had the “degree vector” $(d^{3/2}, d^{2/1}, d^{1/0})^t = (3, 2, 2)^t$, where the superscript t denotes the transpose of the matrix.

The following shows how the “degree vectors” change when we repeatedly apply Lemma 4.4.6, and also how to apply it.

The numbers over the arrows are the indices i for which we apply Lemma 4.4.6, and the entries $*$ are the numbers not known with certainty:

$$\begin{array}{l} \text{3rd} \rightarrow \begin{bmatrix} \mathfrak{Z} \\ 2 \\ 2 \end{bmatrix} \xrightarrow{i=3} \begin{bmatrix} \mathbf{1} \\ 2 \\ 2 \end{bmatrix} \xrightarrow{i=2} \begin{bmatrix} * \\ \mathbf{1} \\ 2 \end{bmatrix} \xrightarrow{i=3} \begin{bmatrix} \mathbf{1} \\ 1 \\ 2 \end{bmatrix} \xrightarrow{i=1} \begin{bmatrix} * \\ * \\ \mathbf{1} \end{bmatrix} \xrightarrow{i=3} \begin{bmatrix} \mathbf{1} \\ * \\ \mathbf{1} \end{bmatrix} \xrightarrow{i=2} \begin{bmatrix} * \\ \mathbf{1} \\ \mathbf{1} \end{bmatrix} \xrightarrow{i=3} \begin{bmatrix} \mathbf{1} \\ \mathbf{1} \\ \mathbf{1} \end{bmatrix}. \\ \text{2nd} \rightarrow \\ \text{1st} \rightarrow \end{array}$$

In general, for $n \geq 1$, if we are given $(d^{n/(n-1)}, \dots, d^{1/0})^t = (a_n, \dots, a_1)^t$ with $a_i \in \mathbb{N}$, then after a finite number of applications of Lemma 4.4.6, we will get to $(1, \dots, 1)^t$. The upper bound for the number of times this operation is needed to get to $(1, \dots, 1)^t$ may be computed by solving a recurrence relation. We guess we need at most $1 + 2 + \dots + 2^{n-1} = 2^n - 1$ times. \square

Proof of Lemma 4.4.6. This is a bit technical, although its basic philosophical ideas go back to part of the arguments of [47].

Since i is fixed here, let $R_\ell := k[\mathfrak{Z}_\ell^{(i-1)}]$ for $\ell = 1, 2$. They were previously called $R_\ell^{(i-1)}$ in Lemma 4.4.4. These are complete local domains finite and free over $k[[t]]$ (Lemma 3.2.5). Similarly, let $\tilde{R}_\ell := k[\mathfrak{Z}_\ell^{(i)}]$. They were previously called $R_\ell^{(i)}$. Here, $R_\ell \hookrightarrow \tilde{R}_\ell$ is the finite extension of rings corresponding to the projection $f_\ell^{i/(i-1)} : \mathfrak{Z}_\ell^{(i)} \rightarrow \mathfrak{Z}_\ell^{(i-1)}$.

By the given assumption that $d^{j/(j-1)}(\mathfrak{Z}_\ell) = 1$ for $j > i$, the images $\bar{P}_\ell^{(j)}$ of the polynomials $P_\ell^{(j)}(y_1, \dots, y_j)$ in $\tilde{R}_\ell[y_{i+1}, \dots, y_n]$ are given by

$$(4.4.6) \quad \bar{P}_\ell^{(j)}(y_{i+1}, \dots, y_n) = y_j - c_\ell^{(j)} \quad \text{for } i < j \leq n,$$

where we recall that for $j > i$, we have $d_j = 1$ and $(-1)^{d_j} c_\ell^{(j)} = -c_\ell^{(j)}$ is the constant term of $\bar{P}_\ell^{(j)}$.

On the other hand, recall that $(-1)^{d_i} c_\ell^{(i)}$ is the constant term of the polynomial $\bar{P}_\ell^{(i)} \in R_\ell[y_i]$, and we have $c_\ell^{(i)} \in R_\ell^\times$ by Proposition 3.3.1-(2). Let (y_i, y'_i) be the coordinates of \square^2 in $\text{Spf}(R_\ell) \times \square^2$, and consider the closed formal subscheme

$C_\ell \subset \mathrm{Spf}(R_\ell) \times \square^2$ given by the polynomial (cf. [47, Lemma 2])

$$Q_{\mathfrak{Z}_\ell}(y_i, y'_i) := \bar{P}_\ell^{(i)}(y_i) - (y_i - 1)^{d_i - 1} (y_i - c_\ell^{(i)}) y'_i \in R_\ell[y_i, y'_i].$$

Its codimension 1 faces are

$$(4.4.7) \quad \begin{cases} C_\ell \cap \{y_i = 0\} = \{(-1)^{d_i} c_\ell^{(i)} (1 - y'_i) = 0\} = \{y'_i = 1\} = \emptyset, \\ C_\ell \cap \{y_i = \infty\} = \dagger \{1 - y'_i = 0\} = \emptyset, \\ C_\ell \cap \{y'_i = 0\} = \{\bar{P}_\ell^{(i)}(y_i) = 0\}, \\ C_\ell \cap \{y'_i = \infty\} = \{(y_i - 1)^{d_i - 1} (y_i - c_\ell^{(i)}) = 0\} = \{y_i = c_\ell^{(i)}\}, \end{cases}$$

where \dagger uses that the polynomial $\bar{P}_\ell^{(i)}(y_i) \in R_\ell[y_i]$ is monic in y_i . Since these faces in (4.4.7) are of codimension ≥ 1 in $\mathrm{Spf}(R_\ell) \times \square^1$, where they intersect no additional face (cf. Lemma 3.2.4), we see that C_ℓ intersects no codimension ≥ 2 face. Thus we have the condition **(GP)** of Definition 2.2.2 for C_ℓ . The condition **(SF)** of Definition 2.2.2 is immediate. Thus we have $C_\ell \in z^1(\mathrm{Spf}(R_\ell), 2)$. Furthermore, from the calculations in (4.4.7), we deduce that

$$(4.4.8) \quad \partial C_\ell = -\{\bar{P}_\ell^{(i)}(y_i) = 0\} + [c_\ell^{(i)}] \in z^1(\mathrm{Spf}(R_\ell), 1).$$

For $1 \leq j \leq n$, let $\alpha_\ell^{(j)}$ be the image of y_j in $k[[t]]\{y_1, \dots, y_n\}/I(\mathfrak{Z}_\ell)$. For $j > i$, we are given that $d^{j/(j-1)}(\mathfrak{Z}_\ell) = 1$ so that we have

$$(4.4.9) \quad k[[t]]\{y_1, \dots, y_n\}/I(\mathfrak{Z}_\ell) = \tilde{R}_\ell.$$

Under this identification, for $j > i$, $\alpha_\ell^{(j)}$ coincides with the previously defined $c_\ell^{(j)}$ as used in (4.4.6). However these may differ for $j \leq i$.

Let $(Y_1, \dots, Y_{i-1}, Y_i, Y'_i, Y_{i+1}, \dots, Y_n)$ be the coordinates of \square^{n+1} in $\mathrm{Spf}(R_\ell) \times \square^{n+1}$. Define the cycle $\tilde{C}_{\mathfrak{Z}_\ell} \in z^n(\mathrm{Spf}(R_\ell), n+1)$ given by the system of polynomials

$$\tilde{C}_{\mathfrak{Z}_\ell} : \begin{cases} P_\ell^{(1)}(Y_1), \\ \vdots \\ P_\ell^{(i-1)}(Y_1, \dots, Y_{i-1}), \\ Q_{\mathfrak{Z}_\ell}(Y_i, Y'_i), \\ P_\ell^{(i+1)}(Y_1, \dots, Y_{i+1}), \\ \vdots \\ P_\ell^{(n)}(Y_1, \dots, Y_n) \end{cases}$$

in $R_\ell\{Y_1, \dots, Y_{i-1}, Y_i, Y'_i, Y_{i+1}, \dots, Y_n\}$. Taking (4.4.6) into account, we use the concatenation notations as in

$$\tilde{C}_{\mathfrak{Z}_\ell} = \mathfrak{Z}_\ell^{(i-1)} \boxtimes C_\ell \boxtimes [c_\ell^{(i+1)}] \boxtimes \dots \boxtimes [c_\ell^{(n)}].$$

Define $C_{\mathfrak{Z}_\ell} \in z^n(\widehat{X}, n+1)$ to be

$$C_{\mathfrak{Z}_\ell} := (-1)^i \pi_{\ell,*} \left(\tilde{C}_{\mathfrak{Z}_\ell} \right),$$

where $\pi_\ell := f_\ell^{(i-1)/0} : \mathrm{Spf}(R_\ell) \rightarrow \widehat{X}$ is the finite surjective morphism, and

$$\pi_{\ell,*} : z^n(\mathrm{Spf}(R_\ell), *) \rightarrow z^n(\widehat{X}, *)$$

is the finite push-forward as in Lemma 2.3.3.

We can compute the boundary of $C_{\mathfrak{Z}_\ell}$ by looking at the boundary of $\tilde{C}_{\mathfrak{Z}_\ell}$. There, note that both $\mathfrak{Z}_\ell^{(i-1)}$ and $[c_\ell^{(j)}]$ for $j > i$ have no proper face (cf. Lemma 3.2.4).

Consequently, combined with (4.4.8), we can write

$$\begin{aligned}
(4.4.10) \quad \partial \tilde{C}_{\mathfrak{Z}_\ell} &= (-1)^{i-1} \mathfrak{Z}_\ell^{(i-1)} \boxtimes (\partial C_\ell) \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}] \\
&= (-1)^i \mathfrak{Z}_\ell^{(i-1)} \boxtimes \{\bar{P}_\ell^{(i)} = 0\} \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}] \\
&\quad - (-1)^i \mathfrak{Z}_\ell^{(i-1)} \boxtimes [c_\ell^{(i)}] \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}] \in z^n(\mathrm{Spf}(R_\ell), n).
\end{aligned}$$

We analyze the two terms of (4.4.10). Consider the closed formal subscheme $\tilde{\mathfrak{Z}}_\ell \subset \mathrm{Spf}(\tilde{R}_\ell) \times \square^n$ given by the system of polynomials in $\tilde{R}_\ell\{Y_1, \dots, Y_n\}$

$$\tilde{\mathfrak{Z}}_\ell : \{Y_1 - \alpha_\ell^{(1)}, \dots, Y_n - \alpha_\ell^{(n)}\}.$$

Since \tilde{R}_ℓ is an integral k -domain and all $\alpha_\ell^{(j)} \in \tilde{R}_\ell$ for $1 \leq j \leq n$ under the identification (4.4.9), the formal scheme $\tilde{\mathfrak{Z}}_\ell$ is integral. Then by definition, the first term of (4.4.10) without the sign

$$\mathfrak{Z}_\ell^{(i-1)} \boxtimes \{\bar{P}_\ell^{(i)} = 0\} \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}]$$

is just $f_{\ell,*}^{i/(i-1)}[\tilde{\mathfrak{Z}}_\ell]$ because $\alpha_\ell^{(j)} = c_\ell^{(j)}$ for $j > i$. By definition, we have $f_{\ell,*}^{i/0}(\tilde{\mathfrak{Z}}_\ell) = \mathfrak{Z}_\ell$ as well. Since $f_\ell^{(i-1)/0} \circ f_\ell^{i/(i-1)} = f_\ell^{i/0}$ and $f_\ell^{(i-1)/0} = \pi_\ell$, we thus have $\pi_{\ell,*}(\mathfrak{Z}_\ell^{(i-1)} \boxtimes \{\bar{P}_\ell^{(i)} = 0\} \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}]) = \mathfrak{Z}_\ell$.

This time, consider the closed formal subscheme $\tilde{\mathfrak{Z}}'_\ell \subset \mathrm{Spf}(\tilde{R}_\ell) \times \square^n$ given by the polynomials in $\tilde{R}_\ell\{Y_1, \dots, Y_n\}$

$$\tilde{\mathfrak{Z}}'_\ell : \{Y_1 - \alpha_\ell^{(1)}, \dots, Y_{i-1} - \alpha_\ell^{(i-1)}, Y_i - c_\ell^{(i)}, Y_{i+1} - c_\ell^{(i+1)}, \dots, Y_n - c_\ell^{(n)}\}.$$

Here all $\alpha_\ell^{(j)}$ for $j < i$, and $c_\ell^{(j)}$ for $j \geq i$ are in \tilde{R}_ℓ so that $\tilde{\mathfrak{Z}}'_\ell$ is an integral formal scheme. Note that $c_\ell^{(i)} \in R_\ell \subset \tilde{R}_\ell$, and it may not be equal to $\alpha_\ell^{(i)}$ in \tilde{R}_ℓ . By definition

$$f_{\ell,*}^{i/(i-1)}(\tilde{\mathfrak{Z}}'_\ell) = \mathfrak{Z}_\ell^{(i-1)} \boxtimes [c_\ell^{(i)}] \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}],$$

which is the second term of (4.4.10) without the sign, so this is an integral cycle. Hence applying $\pi_{\ell,*} = f_{\ell,*}^{(i-1)/0}$, we obtain a cycle $\pi_{\ell,*}(\mathfrak{Z}_\ell^{(i-1)} \boxtimes [c_\ell^{(i)}] \boxtimes [c_\ell^{(i+1)}] \boxtimes \cdots \boxtimes [c_\ell^{(n)}])$, which is a positive integer multiple of an integral cycle.

Since the boundary operators commute with push-forwards (Lemma 2.3.3), we have

$$\partial C_{\mathfrak{Z}_\ell} = (-1)^i \pi_{\ell,*}(\partial \tilde{C}_{\mathfrak{Z}_\ell}) = \mathfrak{Z}_\ell - \pi_{\ell,*}(f_{\ell,*}^{i/(i-1)}(\tilde{\mathfrak{Z}}'_\ell)) = \mathfrak{Z}_\ell - f_{\ell,*}^{i/0}(\tilde{\mathfrak{Z}}'_\ell).$$

So, letting $\mathfrak{Z}'_\ell := f_{\ell,*}^{i/0}(\tilde{\mathfrak{Z}}'_\ell)$, we have $\partial C_{\mathfrak{Z}_\ell} = \mathfrak{Z}_\ell - \mathfrak{Z}'_\ell$.

We now claim that this cycle \mathfrak{Z}'_ℓ satisfies the desired properties. This is a positive integer multiple of an integral cycle. We can still define $d^{j/(j-1)}(\mathfrak{Z}'_\ell)$, ignoring the integer multiplicity. By definition, for $j < i$, we have $(\mathfrak{Z}'_\ell)^{(j)} = \mathfrak{Z}_\ell^{(j)}$, thus $d^{j/(j-1)}(\mathfrak{Z}'_\ell) = d^{j/(j-1)}(\mathfrak{Z}_\ell)$.

Since $c_\ell^{(i)} \in R_\ell$ and $(\mathfrak{Z}'_\ell)^{(i)} = \mathfrak{Z}_\ell^{(i-1)} \boxtimes [c_\ell^{(i)}]$, we have $k[(\mathfrak{Z}'_\ell)^{(i)}] = k[(\mathfrak{Z}'_\ell)^{(i-1)}] = R_\ell$. So, the projection $(\mathfrak{Z}'_\ell)^{(i)} \rightarrow (\mathfrak{Z}'_\ell)^{(i-1)} = \mathfrak{Z}_\ell^{(i-1)}$ is of degree 1, i.e. $d^{i/(i-1)}(\mathfrak{Z}'_\ell) = 1$.

On the other hand, we have $c_\ell^{(j)} \in \tilde{R}_\ell = k[\mathfrak{Z}'_\ell]$ for $j > i$, so that $d^{(i+1)/i}(\mathfrak{Z}'_\ell)$ may not be equal to 1.

Finally, from that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$, by Lemma 4.4.4, we know that $\mathfrak{Z}_1^{(i-1)} \sim_I \mathfrak{Z}_2^{(i-1)}$ so that $c_1^{(i)} \equiv c_2^{(i)}$ in $R_1/IR_1 = R_2/IR_2$, while we also know that $\mathfrak{Z}_1^{(i)} \sim_I \mathfrak{Z}_2^{(i)}$ so that $c_1^{(j)} \equiv c_2^{(j)}$ in $\tilde{R}_1/I\tilde{R}_1 = \tilde{R}_2/I\tilde{R}_2$. Hence $\mathfrak{Z}'_1 \sim_I \mathfrak{Z}'_2$. This finishes the proof of Lemma 4.4.6. \square

Proof of Proposition 4.4.1. We begin to apply Lemma 4.4.6 to a pair of integral cycles \mathfrak{Z}_ℓ in $z^n(\widehat{X}, n)$ such that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$. By Proposition 3.3.1, they satisfy the assumptions of Lemma 4.4.6. So we can apply it repeatedly.

As seen in Remark 4.4.7, in finite number of applications of Lemma 4.4.6, we eventually get to a pair of integral cycles with the equal multiplicities, whose degree vectors are $(d^{n/(n-1)}, \dots, d^{1/0})^t = (1, \dots, 1)^t$. By Lemma 4.4.3, they are positive integer multiples of integral graph cycles in $z^n(\widehat{X}, n)$.

Take the sums, say C_ℓ , of the cycles in $z^n(\widehat{X}, n+1)$ appearing in repeated applications of Lemma 4.4.6. Taking their boundaries, we have the cancellations of all the intermediate \mathfrak{Z}'_ℓ 's, and we have $\partial(C_\ell) = \mathfrak{Z}_\ell - \mathfrak{Z}'_\ell$, where \mathfrak{Z}_ℓ is the given cycle in $z^n(\widehat{X}, n)$, and the resulting \mathfrak{Z}'_ℓ is a positive integer multiple of an integral graph cycle, such that $\mathfrak{Z}'_1 \sim_I \mathfrak{Z}'_2$. This proves Proposition 4.4.1. \square

5. THE PROOFS OF THE MAIN THEOREMS

5.1. The isomorphism over $k[[t]]$. The first application of Proposition 4.4.1 is the following:

Lemma 5.1.1. *Let k be a field. Then the graph homomorphism $gr : K_n^M(k[[t]]) \rightarrow \text{CH}^n(\text{Spf}(k[[t]]), n)$ of (4.1.1) is surjective.*

Proof. Proposition 4.4.1 is stated for pairs of integral cycles that are mod I -equivalent. We apply it to the identical pair $(\mathfrak{Z}, \mathfrak{Z})$ for each integral cycle $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$. In particular, for a cycle $E_\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n+1)$ and a positive integer multiple of a graph cycle, call it Γ , we have $\partial E_\mathfrak{Z} = \mathfrak{Z} - \Gamma$. Since Γ belongs to the image of the graph map gr and $\mathfrak{Z} \equiv \Gamma$ in $\text{CH}^n(\text{Spf}(k[[t]]), n)$, this shows that gr is surjective. \square

Remark 5.1.2. One may give another proof of Lemma 5.1.1 based on arguments of B. Totaro [47, §4]. We outline what extra modifications are needed in addition to the arguments of *loc.cit.*

In *loc.cit.*, when F is a field, to prove that the graph homomorphism $gr : K_n^M(F) \rightarrow \text{CH}^n(\text{Spec}(F), n)$ is an isomorphism, Totaro used the norm $N : L^\times \rightarrow F^\times$, where L is the residue field of a given point and $F \subset L$ is a finite extension of fields.

Specializing to the case $F = k((t))$, what we need is that this norm operation for fields is compatible with the finiteness of our cycles over $k[[t]]$. Indeed if a finite ring extension $k[[t]] \rightarrow B$ describes an integral cycle $\mathfrak{Z} \in z^n(\text{Spf}(k[[t]]), n)$, then B is a free $k[[t]]$ -module and there is the norm map $N : B^\times \rightarrow k[[t]]^\times$ by Corollary 3.2.6 and Definition 3.2.7. This is compatible with the norm $N : L^\times \rightarrow k((t))^\times$, when $L = \text{Frac}(B)$, in that the following diagram commutes:

$$(5.1.1) \quad \begin{array}{ccc} B^\times & \hookrightarrow & L^\times \\ \downarrow N & & \downarrow N \\ k[[t]]^\times & \hookrightarrow & k((t))^\times. \end{array}$$

So, the norms of elements of B^\times belong to $k[[t]]^\times$. Following along the argument of [47, §4], together with the above, we may prove that the graph map of Lemma 5.1.1 is surjective. We do not attempt to give details as this is redundant. \square

Corollary 5.1.3. *Let k be a field with $|k| \gg 0$ so that $K_n^M(k[[t]]) = \widehat{K}_n^M(k[[t]])$.*

Then the graph map

$$gr : K_n^M(k[[t]]) \rightarrow \text{CH}^n(\text{Spf}(k[[t]]), n)$$

is an isomorphism.

Proof. The map gr is surjective by Lemma 5.1.1 and injectivity by Lemma 4.2.6. \square

We deduce the following formal-rigid geometric analogue of the Gersten conjecture for higher Chow groups (cf. S. Bloch [5, Theorem (10.1)]):

Corollary 5.1.4. *Let k be a field with $|k| \gg 0$. Then the localization map*

$$\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \rightarrow \mathrm{CH}^n(\mathrm{Sp}(k((t))), n)$$

is injective.

Proof. This follows from Corollaries 4.2.7 and 5.1.3, and the Gersten conjecture for K_n^M of $k[[t]]$ (see M. Kerz [30, Proposition 10, p.181]). \square

Lemma 5.1.1 for $k[[t]]$ also implies the surjectivity for k_m :

Corollary 5.1.5. *Let $m \geq 2$. Let k be a field, $k_m := k[t]/(t^m)$, and $Y := \mathrm{Spec}(k_m)$.*

Then the graph homomorphism $gr_{k_m} : K_n^M(k_m) \rightarrow \mathbf{CH}^n(k_m, n)$ of (4.1.2) is surjective.

Proof. Recall from (2.4.4) that

$$\mathbf{CH}^n(k_m, n) = \frac{z^n(\mathrm{Spf}(k[[t]]), n)}{\partial(z^n(\mathrm{Spf}(k[[t]]), n+1) + \mathcal{M}^n(\mathrm{Spf}(k[[t]]), Y, n))}.$$

Thus together with Lemma 3.2.4, we have the natural surjective map

$$\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \rightarrow \mathbf{CH}^n(k_m, n),$$

and we deduce the commutative diagram

$$\begin{array}{ccc} K_n^M(k[[t]]) & \xrightarrow{gr} & \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n) \\ \downarrow & & \downarrow \\ K_n^M(k_m) & \xrightarrow{gr_{k_m}} & \mathbf{CH}^n(k_m, n). \end{array}$$

Since the top horizontal arrow is surjective by Lemma 5.1.1, so is the bottom arrow. \square

5.2. The isomorphism over k_m . In Corollary 5.1.5, we saw that gr_{k_m} is surjective. We want to show that it is an isomorphism when $|k| \gg 0$. For this, we want to construct its inverse:

Proposition 5.2.1. *Let $m \geq 2$. Let k be a field with $|k| \gg 0$ so that $K_n^M(k[[t]]) = \widehat{K}_n^M(k[[t]])$.*

Then there is a homomorphism of groups

$$(5.2.1) \quad \Upsilon_m : \mathbf{CH}^n(k_m, n) \rightarrow K_n^M(k_m)$$

such that the composite $\Upsilon_m \circ gr_{k_m} : K_n^M(k_m) \rightarrow \mathbf{CH}^n(k_m, n) \rightarrow K_n^M(k_m)$ is the identity map.

To construct the homomorphism (5.2.1), first consider the composite

$$(5.2.2) \quad \Upsilon_m : z^n(\mathrm{Spf}(k[[t]]), n) \rightarrow \mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$$

$$\xrightarrow[\sim]{gr^{-1}} K_n^M(k[[t]]) \rightarrow K_n^M(k_m).$$

Recall that by Lemma 3.2.4 the first arrow of (5.2.2) is the quotient map, while the map gr is an isomorphism by Corollary 5.1.3, so that the inverse gr^{-1} exists. For the graph cycles, we have the following:

Lemma 5.2.2. *Let $m \geq 2$. For $\Gamma_{(a_1, \dots, a_n)} \in z^n(\mathrm{Spf}(k[[t]]), n)$ with $a_i \in k[[t]]^\times$ for $1 \leq i \leq n$, we have*

$$\Upsilon_m(\Gamma_{(a_1, \dots, a_n)}) = \{\bar{a}_1, \dots, \bar{a}_n\} \text{ in } K_n^M(k_m),$$

where \bar{a}_i is the image of a_i under $k[[t]]^\times \rightarrow k_m^\times$.

Proof. Since $gr(\{a_1, \dots, a_n\})$ is the class $[\Gamma_{(a_1, \dots, a_n)}]$ in $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$, we have $gr^{-1}([\Gamma_{(a_1, \dots, a_n)}]) = \{a_1, \dots, a_n\}$ in $K_n^M(k[[t]])$. The lemma follows immediately. \square

To show that (5.2.2) induces (5.2.1), we need to check that Υ_m respects the equivalence relations coming from the boundary $\partial(z^n(\mathrm{Spf}(k[[t]]), n+1))$ and the subgroup $\mathcal{M}^n(\mathrm{Spf}(k[[t]]), Y, n)$. We check these two reciprocity results in Lemma 5.2.3 and Corollary 5.2.5, respectively:

Lemma 5.2.3. *The composite*

$$z^n(\mathrm{Spf}(k[[t]]), n+1) \xrightarrow{\partial} z^n(\mathrm{Spf}(k[[t]]), n) \xrightarrow{\Upsilon_m} K_n^M(k_m)$$

is 0.

Proof. By definition, the map Υ_m of (5.2.2) already factors through the group $\mathrm{CH}^n(\mathrm{Spf}(k[[t]]), n)$. From this we deduce that $\Upsilon_m \circ \partial = 0$. \square

Note that we have:

Lemma 5.2.4. *Let $m \geq 2$. For two integral graph cycles $\Gamma_1 = \Gamma_{(a_1, \dots, a_n)}$ and $\Gamma_2 = \Gamma_{(a'_1, \dots, a'_n)}$ in $z^n(\mathrm{Spf}(k[[t]]), n)$ with $a_i, a'_i \in k[[t]]^\times$, we have $\Gamma_1 \sim_I \Gamma_2$ if and only if $a_i \equiv a'_i \pmod{t^m}$ for all $1 \leq i \leq n$.*

Corollary 5.2.5. *Let $m \geq 2$. For two cycles \mathfrak{Z}_1 and \mathfrak{Z}_2 in $z^n(\mathrm{Spf}(k[[t]]), n)$ such that $\mathfrak{Z}_1 \sim_Y \mathfrak{Z}_2$, we have $\Upsilon_m(\mathfrak{Z}_1) = \Upsilon_m(\mathfrak{Z}_2)$.*

Proof. By Lemma 4.3.1, it is enough to show the statement for two integral cycles \mathfrak{Z}_1 and \mathfrak{Z}_2 such that $\mathfrak{Z}_1 \sim_I \mathfrak{Z}_2$, i.e. when their mod I -reductions are equal.

By Proposition 4.4.1, there is a pair $\mathfrak{Z}'_1, \mathfrak{Z}'_2$ of positive integer multiples of integral graph cycles such that $\mathfrak{Z}'_1 \sim_I \mathfrak{Z}'_2$, and $\mathfrak{Z}_\ell - \mathfrak{Z}'_\ell = \partial E_{\mathfrak{Z}_\ell}$ for some cycles $E_{\mathfrak{Z}_\ell} \in z^n(\mathrm{Spf}(k[[t]]), n+1)$ for $\ell = 1, 2$.

Since $\Upsilon_m \circ \partial = 0$ by Lemma 5.2.3, we deduce $\Upsilon_m(\mathfrak{Z}_\ell) - \Upsilon_m(\mathfrak{Z}'_\ell) = \Upsilon_m(\partial E_{\mathfrak{Z}_\ell}) = 0$ for $\ell = 1, 2$. However $\Upsilon_m(\mathfrak{Z}'_1) = \Upsilon_m(\mathfrak{Z}'_2)$ by Lemmas 5.2.2 and 5.2.4. Hence

$$\Upsilon_m(\mathfrak{Z}_1) = \Upsilon_m(\mathfrak{Z}'_1) = \Upsilon_m(\mathfrak{Z}'_2) = \Upsilon_m(\mathfrak{Z}_2),$$

as desired. \square

Proof of Proposition 5.2.1. By Lemma 5.2.3 and Corollary 5.2.5, the regulator Υ_m kills the boundaries of cycles and respects the mod Y -equivalence. Hence Υ_m of (5.2.2) is well-defined on $\mathbf{CH}^n(k_m, n)$, inducing the homomorphism (5.2.1).

On the other hand, for an integral graph cycle $\Gamma_{(a_1, \dots, a_n)} \in z^n(\mathrm{Spf}(k[[t]]), n)$ for some $a_1, \dots, a_n \in k[[t]]^\times$, by Lemma 5.2.2, we have $\Upsilon_m(\Gamma_{(a_1, \dots, a_n)}) = \{\bar{a}_1, \dots, \bar{a}_n\} \in K_n^M(k_m)$, where \bar{a}_i is the image of a_i in k_m^\times . This implies that $\Upsilon_m \circ gr_{k_m}$ is the identity on $K_n^M(k_m)$. \square

The above discussions now summarize into the following. This proves the part (2) of the first main theorem, Theorem 1.2.1:

Theorem 5.2.6. *Let $m \geq 2$. Let k be a field with $|k| \gg 0$ so that $K_n^M(k[[t]]) = \widehat{K}_n^M(k[[t]])$.*

Then the graph homomorphism

$$gr_{k_m} : K_n^M(k_m) \rightarrow \mathbf{CH}^n(k_m, n)$$

is an isomorphism.

Proof. By Corollary 5.1.5, the map gr_{k_m} is surjective. On the other hand, under our assumption on the cardinality $|k|$, by Proposition 5.2.1, there is a homomorphism Υ_m such that the composite $\Upsilon_m \circ gr_{k_m}$ is the identity of $K_n^M(k_m)$. In particular, the map gr_{k_m} is injective. This proves the theorem. \square

Recall that by the condition **(SF)** of Definition 2.2.2, we have the Gysin map $ev_{\mathbf{CH}} : \mathbf{CH}^n(k_{m+1}, n) \rightarrow \mathbf{CH}^n(k, n)$ given by the evaluation $t = 0$. This is surjective because $gr_k : K_n^M(k) \rightarrow \mathbf{CH}^n(k, n)$ is an isomorphism by [37] and [47], and the map ev_0 in the following commutative diagram is surjective:

$$(5.2.3) \quad \begin{array}{ccc} K_n^M(k_{m+1}) & \xrightarrow{ev_0} & K_n^M(k) \\ \simeq \downarrow gr_{k_{m+1}} & & \simeq \downarrow gr_k \\ \mathbf{CH}^n(k_{m+1}, n) & \xrightarrow{ev_{\mathbf{CH}}} & \mathbf{CH}^n(k, n). \end{array}$$

Taking the kernels of the horizontal maps, we define the relative parts

$$(5.2.4) \quad \begin{cases} K_n^M(k_{m+1}, (t)) := \ker(ev_0) \text{ and} \\ \mathbf{CH}^n((k_{m+1}, (t)), n) := \ker(ev_{\mathbf{CH}}). \end{cases}$$

We deduce the second main theorem, Theorem 1.6.1:

Theorem 5.2.7. *Let $m \geq 1$. Let k be a field of characteristic 0.*

Then we have an isomorphism of the big de Rham-Witt forms of k of Hesselholt-Madsen [24] with the relative higher Chow groups

$$\mathbb{W}_m \Omega_k^{n-1} \simeq \mathbf{CH}^n((k_{m+1}, (t)), n).$$

Proof. The diagram (5.2.3) with the isomorphic vertical maps induces the isomorphism of the relative parts in (5.2.4),

$$K_n^M(k_{m+1}, (t)) \xrightarrow{\cong} \mathbf{CH}^n((k_{m+1}, (t)), n).$$

Here $K_n^M(k_{m+1}, (t)) \simeq \bigoplus_{i=1}^m \Omega_{k/\mathbb{Z}}^{n-1}$ (Park-Ünver [41, Proposition 5.4.2]), while $\mathbb{W}_m \Omega_k^{n-1} \simeq \bigoplus_{i=1}^m \Omega_{k/\mathbb{Z}}^{n-1}$ (Rülling [42, Remark 1.12]). This proves the theorem. \square

Theorem 5.2.7 is the counterpart to a result known for additive higher Chow groups of a field by K. Rülling [42]. In an earlier version, it was mistakenly claimed for all fields k using [43], but Matthew Morrow pointed out that the reasoning had a jump. The characteristic $p > 0$ case remains to be checked. The referee has suggested a good way to argue for this remaining case, and it will be discussed in a separate work.

Acknowledgments. JP thanks Fumiharu Kato and Shane Kelly for some remarks during JP's visit to the Tokyo Institute of Technology in Spring 2019, that influenced the direction of the project. JP thanks Joseph Ayoub, Chang-Yeon Chough, Youngsu Kim, Wansu Kim, Matthew Morrow and the referee for numerous comments that helped improving the article.

JP thanks Spencer Bloch for his teaching long ago that the Laurent fields should be related to the topic. JP thanks Damy for peace of mind at home and Seungmok Yi for helping him in enduring the long time he works on the project.

During this work, JP was supported by the National Research Foundation of Korea (NRF) grant (2018R1A2B6002287) funded by the Korean government (Ministry of Science and ICT).

REFERENCES

- [1] M. F. Atiyah and I. G. MacDONald, *Introduction to commutative algebra*, Addison-Wesley, 1969. ix+128 pp.
- [2] J. Ayoub, *Motifs des variétés analytiques rigides*, Mémoire. Soc. Math. France, (2015), Num. 140-141, Soc. Math. France.
- [3] H. Bass and J. Tate, *The Milnor ring of a global field*, in Algebraic K -theory II (Battelle Memorial Inst., Seattle, Washington, 1972), Lect. Notes in Math., **342**, Springer-Verlag, New York, 1973.
- [4] F. Binda and S. Saito, *Relative cycles with moduli and regulator maps*, J. Inst. Math. Jussieu, **18**, (2019), no. 6, 1233–1293.
- [5] S. Bloch, *Algebraic cycles and higher K -theory*, Adv. Math., **61**, (1986), no. 3, 267–304.
- [6] S. Bloch and H. Esnault, *The additive dilogarithm*, Documenta Math. **Extra Vol.** for Kazuya Kato’s fiftieth birthday conference, (2003), 131–155.
- [7] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, Grundlehren der Math. Wissen., **261**, Springer, 1984.
- [8] S. Bosch, *Lectures on formal and rigid geometry*, Lect. Notes in Math., **2105**, Springer, Cham, 2014, viii+254pp.
- [9] B. Conrad, *Irreducible components of rigid spaces*, Ann. Inst. Fourier, Grenoble, **49**, (1999), no. 2, 473–541.
- [10] B. Conrad and W. Stein, *Component groups of purely toric quotients*, Math. Res. Lett., **8**, (2001), 745–766.
- [11] D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Grad. Text in Math., **150**, Springer-Verlag, 1994.
- [12] P. Elbaz-Vincent and S. Müller-Stach, *Milnor K -theory of rings, higher Chow groups and applications*, Invent. Math., **148**, (2002), no. 1, 177–206.
- [13] J. Fresnel and M. van der Put, *Rigid analytic geometry and its applications*, Progress in Math., **218**, Birkhäuser, 2004.
- [14] K. Fujiwara and F. Kato, *Foundations of Rigid Geometry I*, EMS Monographs in Mathematics, European Math. Soc., Zürich, 2018.
- [15] W. Fulton, *Intersection theory, Second Edition*, Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer-Verlag, Berlin, 1998. xiv+470 pp.
- [16] A. Grothendieck, *Éléments de Géométrie Algébrique I, Le langage des schémas*. Publ. Math. de l’Inst. Hautes Études Sci., **4**, (1960), 228pp.
- [17] A. Grothendieck, *Éléments de Géométrie Algébrique IV, Étude locale des schémas et des morphismes de schémas, Seconde partie*, Publ. Math. de l’Inst. Hautes Études Sci., **24**, (1965), 5–231.
- [18] A. Grothendieck, *On the de Rham cohomology of algebraic varieties*, Publ. Math. de l’Inst. Hautes Études Sci., **29**, (1966), 95–103.
- [19] A. Grothendieck, *Éléments de Géométrie Algébrique IV, Étude locale des schémas et des morphismes de schémas, Quatrième partie.*, Publ. Math. de l’Inst. Hautes Études Sci., **32**, (1967), 5–361.
- [20] R. Hartshorne, *On the de Rham cohomology of algebraic varieties*, Publ. Math. de l’Inst. Hautes Études Sci., **45**, (1975), 5–99.
- [21] R. Hartshorne, *Algebraic Geometry*, Grad. Texts in Math., **52**, Springer-Verlag, New York-Heidelberg, 1977. xvi+496 pp.
- [22] R. Hartshorne and C. Polini, *Divisor class groups of singular surfaces*, Trans. Amer. Math. Soc., **367**, (2015), no. 9, 6357–6385.
- [23] H. Hasse, *Number Theory*, English translation, Edited and prepared by Horst Günter Zimmer, Classics in Math., Springer-Verlag, 2002.
- [24] L. Hesselholt and I. Madsen, *On the K -theory of nilpotent endomorphisms*, in Homotopy methods in Algebraic Topology (Boulder, CO, 1999), edited by Greenlees, J. P. C. et al., 127–140. Contempo. Math. **271**, Providence, RI, American Math. Soc. 2001.
- [25] B. Kahn, H. Miyazaki, S. Saito and T. Yamazaki, *Motives with modulus, I: modulus sheaves with transfers for non-proper modulus pairs*, Épijournal Géom. Algébr. **5**, (2021), Article Nr. 1.
- [26] B. Kahn, H. Miyazaki, S. Saito and T. Yamazaki, *Motives with modulus, II: modulus sheaves with transfers for proper modulus pairs*, Épijournal Géom. Algébr. **5**, (2021), Article Nr. 2.

- [27] B. Kahn, H. Miyazaki, S. Saito and T. Yamazaki, *Motives with modulus, III, The categories of motives*, Ann. K-theory, **7**, (2022), no. 1, 119–178.
- [28] K. Kato, *A generalization of local class field theory by using K-groups. II*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27**, (1980), 603–683.
- [29] K. Kato and S. Saito, *Global class field theory of arithmetic schemes*, in Applications of Algebraic K-theory to Algebraic Geometry and Number Theory, Proceedings of a Summer Research Conference, June 12–18, 1983, Eds. S. J. Bloch et. al., Contemporary Math., **55**, part I, 255–331.
- [30] M. Kerz, *Milnor K-theory of local rings with finite residue fields*, J. Algebr. Geom., **19**, (2010), 173–191.
- [31] A. Krishna and J. Park, *A moving lemma for relative 0-cycles*, Algebra & Number Theory, **14**, (2020), no. 4, 991–1054.
- [32] A. Krishna and J. Park, *De Rham-Witt sheaves via algebraic cycles* (with an appendix by Kay Rülling), Compositio Math., **158**, (2021), no. 10, 2089–2132.
- [33] S. E. Landsburg, *K-theory and patching for categories of complexes*, Duke Math. J., **62**, (1991), no. 2, 359–384.
- [34] J. Malgoire and C. Voisin, *Factorisation de Stein topologique découpe*, Topology, **20**, (1981), 191–207.
- [35] J. Milnor, *Algebraic K-theory and quadratic forms*, Invent. Math., **9**, (1970), 318–344.
- [36] J. Milnor, *Introduction to algebraic K-theory*, Annals Math. Studies **72**, Princeton Univ. Press, Princeton, NJ, 1971.
- [37] Yu. P. Nesterenko and A. A. Suslin, *Homology of the general linear group over a local ring, and Milnor’s K-theory*, (Russian) Izv. Akad. Nauk SSSR Ser. Mat., **53**, (1989), no. 1, 121–146; English translation: Math. USSR-Izv., **34**, (1990), no. 1, 121–145.
- [38] J. Park, *Regulators on additive higher Chow groups*, Amer. J. Math., **131**, (2009), no. 1, 257–276.
- [39] J. Park, *On extension of the motivic cohomology beyond smooth schemes*, preprint, arXiv:2108.13594, 2021.
- [40] J. Park and P. Pelaez, *On the coniveau filtration on algebraic K-theory of singular schemes*, preprint, 2021.
- [41] J. Park and S. Ünver, *Motivic cohomology of fat points in Milnor range*, Documenta Math., **23**, (2018), 759–798.
- [42] K. Rülling, *The generalized de Rham-Witt complex over a field is a complex of zero-cycles*, J. Algebr. Geom., **16**, (2007), no. 1, 109–169.
- [43] K. Rülling and S. Saito, *Higher Chow groups with modulus and relative Milnor K-theory*, Trans. Amer. Math. Soc., **370**, (2018), no. 2, 987–1043.
- [44] <https://stacks.math.columbia.edu>, The Stacks Project.
- [45] A. A. Suslin, *Reciprocity laws and the stable rank of rings of polynomials* (Russian), Izv. Akad. nauk SSSR Ser. Mat., **43**, (1979), no. 6, 1394–1429. English translation: Math. USSR Izv., **15**, (1980), no. 3, 589–623.
- [46] J. Tate, *Rigid analytic spaces*, Invent. Math., **12**, (1971), 257–289.
- [47] B. Totaro, *Milnor K-theory is the simplest part of algebraic K-theory*, K-Theory, **6**, (1992), no. 2, 177–189.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 291 DAEHAK-RO YUSEONG-GU, DAEJEON, 34141, REPUBLIC OF KOREA (SOUTH)

Email address: jinhyun@mathsci.kaist.ac.kr; jinhyun@kaist.edu