A short geometric derivation of the dual Steenrod algebra

Kiran Luecke

October 24, 2021

$$A \mapsto (X \mapsto EA^*X := E^*X \otimes_{E^*} A)$$

$$\eta_E : \operatorname{Spec} E_* E \to \operatorname{Hom}_{MT}(E, E-)$$

$$(E_*E \xrightarrow{f} A) \mapsto \eta_E(f) := (E^*X \xrightarrow{1 \land \mathrm{id}} (E \land E)^*X \simeq E(E_*E)^*X \xrightarrow{F} EA^*X)$$

Dotober 24, 2021 **Dotober 24**, 2021 **Abstrat**This two-page note gives a non-computational derivation of the dual Steenrod algebra as the automorphisms of the formal additive group. Instead of relying on computational tools like spectral sequences and Steenrod operations, the argument uses a few simple universal properties of certain cohomology theories.
All algebras are graded and commutative. C₂ is the cyclic group of order 2. **Preamble:** Let MT be the category of pairs (F, s) where F is a contravariant functor from the homotopy category of finite spectra to the category of graded abelian groups that takes sums to products and s is a natural transformation F → F' that takes s to s'. Let E be a homotopy ring spectrum and E^{*} = π_-*E. Consider the "Conner-Floyd" functor Alg_{E*} → MT
A → (X → EA*X := E*X ⊗_{E*} A).
Suppose that E*E is flat over E*, so that E(E*E) is represented by E ∧ E. Consider the natural transformation
M = SpecE*E → Hom_{MT}(E, E−)
(E*E ^f ∧ 1) → η_E(f) := (E*X ^{1.Δid} (E ∧ E)*X ≃ E(E*E)*X ^f → EA*X).
The coproduct on E*E induced by the map E ∧ E ^[MalAid] E ∧ E ∧ E in the increase of η_E. If EA is a cohomology theory (e.g. if A is flat over E*), then it corresponds to a (homotopy ring) spectrum which is an E-module, harmlessly denoted by E ∧ A, and there is an inverse to η given by sending a transformation T : E → EA to the map on homotopy induced by E ∧ E. ^[MalAid] E ∧ EA → EA.
Lemma 0.0.1. Let E be a real oriented homotopy ring spectrum such that E*E is flat over F*. Write G_E for the formal group with coordinate SpE*BC2 and write Au&E for the group-valued greshead on Alg_{E*} sending R to the group of strict argoup with coordinate SpE*BC2 and write Au&E for the some-valued greshead on Alg_E sending R to the group of Sint argoup with coordinate SpE*BC2 and there is a morphism Spec(E*E) ^(C*). The due and comparison of G_E([X]). Then there is a morphism Spec(E*E) ^(C*) an and oriented from Mar(E

to the multiplication map $BC_2 \times BC_2 \to BC_2$, the power series f(e) must be an endomorphism of the formal group law of \mathbb{G}_E . The linear coefficient f_0 is forced to be 1 by considering the pullback of e along $S^1 \hookrightarrow BC_2$ and invoking stability of T. Hence f(e) defines a strict automorphism of \mathbb{G}_E . Precomposition of the assignment $T \mapsto f(e)$ with η_E defines the desired map ev_E : Spec $E_*E(R) \rightarrow Aut \mathbb{G}_E(R)$ which is clearly natural in R and commutes with the monoidal structure on both sides by inspection.

Lemma 0.0.2. Suppose that there is a homotopy commutative ring spectrum M such that

- 1. There is a map $M \to H$ inducing an isomorphism on π_0 .
- 2. M is real oriented with real orientation $e_M \in M^1BC_2$ and formal group \mathbb{G}_M with formal group law F_M .
- 3. M_*M is flat over M^* , and the map $ev_M(R)$ afforded by the previous lemma is injective when MR is a cohomology theory.
- 4. For an \mathbb{F}_2 -algebra R let $Aut \mathbb{G}_M(R)$ be the groupoid whose objects are \mathbb{F}_2 -algebra maps $M^* \xrightarrow{f} R$ and whose morphisms from f to g are strict isomorphisms of formal group laws ϕ from f_*F_M to g_*F_M . Write R_f for the induced M^* -algebra

structure whose underlying ring is R. Note that MR_f is canonically real oriented by the class $e_{MR_f} = [e_M \otimes 1] \in MR_f^*BC_2 = M^*BC_2 \otimes_{M^*} R_f$. Then there is a functor

$$\gamma: Aut \mathbb{G}_M(R) \to MT$$

which on objects sends f to MR_f and on morphisms sends ϕ to the transformation $\gamma(\phi) : MR_f \to MR_g$ such that $\gamma(\phi)(e_{MR_f}) = \phi(e_{MR_g}) \in MR_g^*BC_2 \simeq R_g[[e_{MR_g}]]$ as a power series.

Then the evaluation map ev_H from the previous lemma is an isomorphism. In particular if \mathcal{A}_2 denotes a Hopf algebra corepresenting $Aut \mathbb{G}_H$ then there is an isomorphism of Hopf algebras $H_*H \simeq \mathcal{A}_2$.

Proof. By item 1 \mathbb{F}_2 is an algebra over M^* . By item 2 the formal group law of M^*BC_2 has vanishing 2-series $[2](e_M) = 0$ and so it is isomorphic to the additive one. Transporting such an isomorphism ϕ with γ produces a natural isomorphism

$$M^*X \simeq M^*X \otimes_{M^*} M^* \xrightarrow{\gamma(\phi)} M^*X \otimes_{M^*} \mathbb{F}_2 \otimes_{\mathbb{F}_2} M^* = M\mathbb{F}_2 \otimes_{\mathbb{F}_2} M^*$$

Thus $M\mathbb{F}_2$ is a summand of a cohomology theory and hence a cohomology theory. The map in item 1 induces a map $M\mathbb{F}_2 \to H$ which by Eilenberg-Steenrod uniqueness must be an isomorphism. Thus, combining item 3 with the pullback along the surjection $M \to M\mathbb{F}_2 \simeq H$, the map ev_H must be injective. So to produce an inverse it suffices to produce a section. Now since \mathbb{F}_2 is a field we have $MR \simeq HR$ for all \mathbb{F}_2 -algebras R. Note that there is an inclusion $B\operatorname{Aut}\mathbb{G}_H \hookrightarrow \operatorname{Aut}\mathbb{F}_H$ since F_M is sent to the additive formal group law by the map $M^* \to \mathbb{F}_2$ in item 1. Then the desired section of ev_H is given as follows. Write $\mathbb{F}_2 \xrightarrow{u} R$ for the unit map of an \mathbb{F}_2 -algebra R. Start with an automorphism of $\operatorname{Aut}\mathbb{G}_H(R)$ and view this as an automorphism of the object $M^* \to \mathbb{F}_2 \to R$ in $\operatorname{Aut}\mathbb{G}_M(R)$. Using γ that produces an automorphism of $MR \simeq HR$, which one then precomposes with the morphism $H \to HR$ induced by u.

Lemma 0.0.3. A cohomology theory M satisfying the conditions of the previous lemma exists. It is MO.

Proof. Let e_{MO} denote the homotopy class of the inclusion $\mathbb{RP}^{\infty} \hookrightarrow \Sigma MO$. This exhibits (MO, e_{MO}) as the universal real oriented multiplicative cohomology theory. Hence there is a map $MO \to H$ sending e_{MO} to $e \in H^*BC_2$. Thus items 1 and 2 are satisfied. Since MO is real oriented, an easy calculation¹ shows that $MO_*MO \simeq MO_*[a_1, a_2, ...]$ which is free and hence flat over MO^* . Let A be an MO^* -algebra and $T: MO \to MOA$ a multiplicative natural transformation. Let $\Theta_n \in MO^*MO(n)$ be the universal Thom class. By the splitting principle $T(\Theta_1)$ determines $T(\Theta_n)$, which determines all of T by universality. Therefore if MOA is a cohomology theory then $ev_{MO}(A)$ is the composition of an isomorphism $(\eta_{MO}(A))$ and an injection $(T \mapsto T(\Theta_1))$ so item 3 is satisfied. Let A be an \mathbb{F}_2 -algebra receiving two maps $f, g: MO^* \to A$ and let $\phi \in A[[x]]$ be an isomorphism from f_*F_{MO} to g_*F_{MO} . Note that MOA_f and MOA_g are canonically oriented by e_{MOA_f} and e_{MOA_g} as defined in item 4. It remains to construct (functorially) an automorphism $\gamma(\phi): MOA_f \to MOA_g$ such that $T(e_{MOA_f}) = \phi(e_{MOA_g})$. By the calculation $MO_*MO \simeq MO_* \otimes_{\mathbb{F}_2} \mathbb{F}_2[a_1, a_2, \ldots]$, the pair (g, ϕ) corresponds² to an \mathbb{F}_2 -algebra map $\Phi: MO_*MO \to A$ and so using the natural transformation η_{MO} from the preamble one gets multiplicative transformation $\eta_{MO}(\Phi) : MO \to MOA_q$ with the property that $\eta_{MO}(\Phi)(e_{MO}) = f(e_{MOA_q})$. Indeed the image of e_{MO} under the map $MO^*BC_1 \xrightarrow{1 \wedge \mathrm{id}} (MO \wedge MO)^*BC_2 \simeq MO^*BC_2 \otimes_{MO^*} MO_*MO$ is $[e_{MO} \otimes 1] + [e_{MO}^2 \otimes a_1] + \dots$ by a simple calculation using the pairing between homology and cohomology³. Note that the map $\eta_{MO}(\Phi)(\text{pt}): MO^* \to A$ is not equal to g. Now $\eta_{MO}(\Phi)$ induces an A-linear map $MOA_f^*X \to MOA_q^*X$ and produces the desired map $\gamma(\phi): MOA_f \to MOA_g$ which is what we are really after. Functoriality⁴ is proved by noting that $\gamma(\phi)$ is uniquely characterized by the properties of being A-linear, multiplicative, and sending e_{MOA_f} to $\phi(e_{MOA_g})$.

Remark 0.0.4. There is a similar but slightly messier derivation of the dual Steenrod algebra at odd primes which uses a mod p analog of MO constructed by Shaun Bullett [2] using manifolds with singularities. I decided to leave the odd primes to forthcoming work⁵ since the added technical difficulties (which in this business are usual at odd primes) obscures the simplicity and brevity of the exposition, which is a cardinal virtue of this note.

Acknowledgements: Thanks to Tim Campion for extensive comments on previous drafts, and to Robert Burklund for catching a gap in a proof.

¹The real orientation e_{MO} implies that all the differentials in the relevant Atiyah-Hirzebruch sequences are trivial.

²By strictness $f(e_{MOA_g}) = e_{MOA_g} + f_1 e_{MOA_g}^2 + \dots$ and the corresponding map $\Phi : MO_*[a_1, \dots] \to A$ sends a_i to f_{i+1} .

 $^{^3 \}mathrm{See}$ e.g. Adams [1] Lemma 6.3 page 60 for the complex version.

⁴This argument for functoriality is due to Quillen in [4].

⁵This work will also discuss derivations of algebras of cohomology operations of other spectra as well as a derivation of the Dyer-Lashof algebra.

1 References

[1] Adams, F. Stable homotopy and generalised homology. Chicago Lectures in Mathematics. The University of Chicago Press, 1995.

[2] Bullett, S. A Z/p Analogue for Unoriented Bordism. University of Warwick PhD Thesis (1973)

[3] Peterson, E. Formal geometry and bordism operations. Cambridge Studies in Advanced Mathematics, v17 (2019)

[4] Quillen, D. Elementary Proofs of Rome Results of Cobordism Theory Using Steenrod Operations. Advances in Mathematics, 7 29-56 (1971)