

December 1, 2021

Cross-correlation Power Spectra and Cosmic Birefringence of the CMB via Photon-neutrino Interaction

Roohollah Mohammadi*

*Iranian National Museum of Science and Technology (INMOST), PO Box 11369-14611, Tehran, Iran.
School of Astronomy, Institute for Research in Fundamental Sciences (IPM), P. O. Box 19395-5531, Tehran, Iran.*

Jafar Khodagholizadeh†

Farhangian University, P.O. Box 11876-13311, Tehran, Iran.

Mahdi Sadegh‡

*School of Particles and Accelerators, Institute for Research in
Fundamental Sciences (IPM), P. O. Box 19395-5531, Tehran, Iran.*

Ali Vahedi§

*Department of Astronomy and High Energy Physics,
Faculty of Physics, Kharazmi University, P. O. Box 15614, Tehran, Iran.*

It is shown that in the context of the standard model of particles, weak interaction of cosmic microwave background (CMB) and cosmic neutrino background (CvB), in the order of one loop forward scattering, can generate non-vanishing TB and EB power spectra in the presence of scalar perturbation which is in contrast with the standard scenario cosmology. By comparing our results with the current experimental data, significant information can be extracted about the nature of CvB: the contribution of CMB-CvB forward scattering for TB, TE, and EB power spectra in the case of CvB as Majorana particles is twice larger than that in the case of Dirac one. Finally, we have discussed the evidence of cosmic neutrino background in the Planck 2018 polarization data reported on a new measurement of the cosmic birefringence (CB) angle β . It is shown that the mean opacity due to cosmic neutrino background can behave as an anisotropic birefringent medium changing the linear polarization rotation angle. CB angle from the average value of $\beta|_v \simeq 1/2\bar{\kappa}$ is about four (eight) times larger than what reported in [40], in the case of CvB as Dirac (Majorana) particles. This point will be discussed more precisely.

PACS numbers: 13.15.+g,34.50.Rk,13.88.+e

* rmohammadi@ipm.ir

† gholizadeh@ipm.ir

‡ m.sadegh@ipm.ir

§ vahedi@ipm.ir

I. Introduction

Cosmic neutrinos can decouple deep earlier from matter about 1 sec after the Big Bang (BB) at the temperature of $\cong 1\text{MeV} \cong 10^{10}\text{K}$. Therefore, along with the cosmic microwave background (CMB) that is coming from 380000 years after BB, cosmic neutrino background (CvB) contains valuable information about the history of the universe [1]. Today, due to very weak interactions and low energy ($1.95\text{K} \cong 1.68 \times 10^{-4}\text{eV}$) of CvB, it is very hard to detect them.

Moreover, relativistic neutrino beam can not help make difference, at least in observation, between the Dirac and Majorana neutrino in the case of electromagnetic interaction. This fact directly relates to the left-handed interacting neutrino in the SM context, which is called "Practical Dirac-Majorana confusion theorem" [2]. But the very low energy of CvB that makes these particles is in the non-relativistic regime: so, the Dirac or Majorana nature of neutrino has different electromagnetic effects [3] that can be detected by the upgrading experiments. Recently, some experiments are prepared to detect CvB, e.g., KATRIN equipped to see CvB by the most probable interaction channel which is capturing of the electron neutrino from the background by Tritium ($\nu_e(1.95\text{Kelvin}) + {}^3\text{H} \rightarrow {}^3\text{He} + e^-$) [4]. Nevertheless, many theoretical attempts have been proposed in the literature to search for these relic neutrinos [5–12]. The detection method of CvB is divided into three categories: (1) direct detections via momentum transfer in neutrino elastic scattering off nuclei target, (2) neutrino capture in nuclei, or (3) indirect method by spectral distortion generated by CvB and ultra-high energy neutrinos or protons originating from unknown sources [13]. But the common denominator of all these categories is that no search has been done in any of them for the nature of Dirac or Majorana neutrinos. Another indirect detection method of CvB has been recently proposed by Mohammadi et al. [14–17]. In this method, photon-neutrino forward scattering is considered as a tool probing CvB which distinguishes Majorana and Dirac nature of CvB.

As is well-known, the CMB photons can interact electromagnetically with the matter: so, this can be used to test very vast theoretical models of particle physics and cosmology [18–23]. Intensity and polarization of CMB are impressed by the interactions which are ways to extract the information.

The most commonly used observables which can be investigated theoretically and experimentally are the CMB polarization power spectra. B-mode (C_ℓ^{BB}) power spectrum can be used to study primordial gravitational waves [24], because in the standard scenario of cosmology, the CMB B-mode polarization due to the Thomson scattering can only generate in the presence of tensor metric perturbations [25–27]. Thereupon, the B-mode power spectrum can calculate one of the most important cosmological parameters as tensor-to-scalar (r-parameter) [28, 29]. E-mode polarization is generated by both scalar and tensor metric fluctuations. Nowadays, E-mode power spectrum (C_ℓ^{EE}) measurements are almost consistent with the ΛCDM cosmological model [30–32]. It should be noted that due to the lower levels of polarization foreground measurements, temperature-polarization power spectrum (C_ℓ^{TE}) measurements can surpass our information about CMB temperature anisotropy (C_ℓ^{TT}) and put some stronger constraints on cosmological models [32, 33]. This point motivates us to investigate some of the main cross-correlation power spectra such as TE , TB , and EB .

In the standard scenario of cosmology, cross-correlations of EB and TB are due to the parity invariant vanish [24, 34]. So, we can use these cross-correlations to check some parity-violating interactions and go beyond the standard models [35]. In [36, 37], the authors have theoretically investigated that the quintessence and Axion-like dark matter particle coupled to the electromagnetic strength tensor and its dual ($\mathcal{L} = \phi(t)F_{\mu\nu}\tilde{F}^{\mu\nu}$) can generate non-zero EB and TB power spectra. In these theoretical models, the plane of CMB linear polarization (E-mode and B-mode polarizations) can rotate by an angle proportional to $\int_{t_{\text{LSS}}}^{t_0} dt \dot{\phi}$ which is known as cosmic birefringence (CB) [38]. In conventional physics, the cosmological birefringent medium is caused by some radiation anisotropy of the background along the cosmological path of CMB since the last scattering surface time. For example, finite displacement of electrons in the ionized plasma, photon-photon scattering due to the virtual electron-positron pairs in vacuum, and also, lack of hydrogen atom potential as a simple harmonic oscillator [39].

Moreover, we show that in the presence of cosmic neutrino background, their interaction with cosmic microwave background via parity violating forward scattering makes an opaque medium: this opacity acts as a birefringent medium. Using the phenomenological value of this opacity, we can calculate the contribution of CMB-CvB interaction at the measured CB angle. Recently, by analyzing the 2018 cosmological foreground polarization the data of Planck, this CB has been measured ($\beta = 0.35^\circ \pm 0.14^\circ$) for small l at 68% confidence level [40]. Also, some other experiments have measured CB that rely on the polarization angle calibration based on hardware calibrator and astrophysical sources [41–43]. Contrary to [40] their results were consistent to zero.

In [14], we studied the generation of CMB B-mode polarization via forward scattering of photon-neutrino in the case of tensor metric perturbation and showed that the effect of Majorana neutrinos on the r -parameter was different from that of the Dirac neutrinos.

Distinguishing Dirac from Majorana neutrinos can be used in the microwave cavity [44], $SU(6)$ grand unified theories [45], weak interaction of aggregate matter [46], neutrino decay[47], pure leptonic decays at the LHC [48], neutrino-electron elastic scattering in the presence of all possible Lorentz-invariant interactions [49], and non-cyclic geometric phase for neutrinos [50–52].

In this work, we will solve the quantum Boltzmann equation for the time-evolution of matrix density (Stokes parameters) of an ensemble of CMB photons. The CMB-CvB scattering is involved here as a collision term. The CMB-CvB weak interaction not only can generate non-zero EB and TB power spectra but also can identify the Dirac and Majorana nature of CvB. Also, the generated neutrino opacity via CMB-CvB interaction behaves as a birefringent medium that can change the CMB linear polarization angle. Therefore the observed cross power spectra between E-mode, B-mode, and temperature in the background of scalar perturbations has a significant contribution to measured CB and this scattering should be regarded as one of the most effective phenomena. As one can see in Fig.2, EB cross power spectrum has a good consistency with data. To give an effective sense, we show that the CB angle β can be roughly estimated by the average value of opacity due to CvB medium $\beta|_v \simeq 1/2\bar{\kappa}$ where $\kappa(\eta) = \int_{\eta}^{\eta_0} d\eta \frac{\sqrt{2}}{6\pi} \alpha G^F \frac{n_\nu}{2}$, η is the conformal time, n_ν is neutrino number density, G^F and α are Fermi coupling constant and electromagnetic fine structure constant. Estimated CB angle from the average value of $\beta|_v \simeq 1/2\bar{\kappa}$ is about two (four) times larger than what reported in [40], in the case of CvB as Dirac (Majorana) particles. This point will be discussed more precisely in Sec.III.

II. Cross power spectrum in the presence of Thomson scattering and CMB-CvB

To calculate cross power spectra of CMB radiation, we start with CMB radiation as an ensemble of photons described by

$$\begin{aligned} \hat{\rho}_{ij} &= \frac{1}{tr(\hat{\rho})} \int \frac{d^3k}{(2\pi)^3} \rho_{ij}(k) D_{ij}(k), \\ \hat{\rho}_{ij} &\equiv \frac{1}{2} \begin{pmatrix} I+Q & U-iV \\ U+iV & I-Q \end{pmatrix} \end{aligned} \quad (1)$$

where I , Q (U) and V are the Stokes parameters indicating the total intensities, linear polarizations intensities and the difference between clockwise and anti-clockwise circular polarization intensities, respectively [53]. $D_{ij}(\vec{k}) \equiv a_i^\dagger(\vec{k}) a_j(\vec{k})$ is the photon number operator. Then, to obtain the time evolution of Stokes parameters, we solved the following quantum Boltzmann equation

$$\begin{aligned} (2\pi)^3 2k^0 \frac{d}{dt} \rho_{ij}(k) &= i \langle [H_I^0, D_{ij}(k)] \rangle \\ &- \frac{1}{2} \int_{-\infty}^{+\infty} dt \langle [H_I^0(t), [H_I^0(t), D_{ij}(k)]] \rangle, \end{aligned} \quad (2)$$




Figure 1: Two Feynman diagrams describe the interaction between photons and Dirac neutrinos at the one-loop level. Along the loop, there are two QED-vertexes and two weak interacting vertexes. Note that the neutral channel diagram with a Z^0 -boson exchange does not contribute to the interacting Hamiltonian H_I^0 . In the case of neutrino as Majorana particles, the “conjugated” Feynman diagrams should be considered which can be yielded by replacing all the internal lines in the above diagrams with their conjugated lines.

where H_I^0 indicates the leading order term of the interacting Hamiltonian. The first term on RHS of the above equation is called forward scattering (interacting without momentum exchange) with linear dependence on H_I^0 and the second one is the high-order collision terms (representing ordinary cross-section term) with square dependence on H_I^0 (see [23] for more detail). As shown in [16], in the case of CvB-CMB interaction given by Fig.(1), the contribution of forward scattering is non-zero which is proportional to ($\propto \alpha G_F$). Note, the usual cross-section of photon- neutrino scattering is proportional to ($\propto \alpha^2 G_F^2$), which is very smaller than forward scattering term. The Boltzmann equation for the linear polarization and temperature anisotropy parameters of CMB radiation in the presence of compton scattering and photon-neutrino forward scattering could be written as [54]

$$\begin{aligned} & \frac{d}{d\eta} \Delta_T^{(S)} + iK\mu \Delta_T^{(S)} + 4[\psi - iK\mu\phi] \\ & = \hat{\tau}_{e\gamma} \left[-\Delta_T^{(S)} + \Delta_T^{0(S)} + i\mu v_b + \frac{1}{2} P_2(\mu) \Pi \right], \end{aligned} \quad (3)$$

$$\begin{aligned} & \frac{d}{d\eta} \Delta_P^{\pm(S)} + iK\mu \Delta_P^{\pm(S)} \\ & = \hat{\tau}_{e\gamma} \left[-\Delta_P^{\pm(S)} - \frac{1}{2} [1 - P_2(\mu)] \Pi \right] \mp a(\eta) \dot{\kappa}_{\pm} \Delta_P^{\pm(S)}, \end{aligned} \quad (4)$$

where $\Delta_P^{\pm(S)} = Q^{(S)} \pm iU^{(S)}$, $\Pi \equiv \Delta_I^{2(S)} + \Delta_P^{2(S)} + \Delta_P^{0(S)}$, ψ , and ϕ are scalar functions defining scalar perturbation of metric in longitudinal gauge, v_b is neutrino bulk velocity, $\hat{\tau}_{e\gamma} = an_e x_e \sigma_T$ is differential optical depth for Thomson scattering which $a(\eta)$ is the expansion factor normalized to unity, x_e is the ionization fraction and σ_T is Thomson cross-section. $\dot{\kappa}_{\pm} = d\kappa_{\pm}(\eta)/d\eta$ in the case of a Majorana neutrino is twice larger than that in the Dirac neutrino case (for more details, please see [55]).

The CMB polarization $\Delta_P^{\pm(S)}(\eta_0, k, \mu)$ is separated into the divergence-free part (B-mode) and curl-free (E-mode) as follows:

$$\Delta_E^{(S)}(\eta_0, K, \mu) = -\frac{3}{4} \int_0^{\eta_0} d\eta g(\eta) \Pi(\eta, K) \{ \partial_{\mu}^2 [(1 - \mu^2)^2 e^{ix\mu} \cos \kappa(\eta)] \}, \quad (5)$$

$$\Delta_B^{(S)}(\eta_0, K, \mu) = \frac{3}{4} \int_0^{\eta_0} d\eta g(\eta) \Pi(\eta, K) \{ \partial_{\mu}^2 [(1 - \mu^2)^2 e^{ix\mu} \sin \kappa(\eta)] \}, \quad (6)$$

where $g(\eta) = \hat{\tau}_{e\gamma} e^{-\tau_{e\gamma}}$, $\partial_{\mu} = \partial/\partial\mu$, and $\mu = \hat{\mathbf{n}} \cdot \hat{\mathbf{K}}$ is the cosine angle of the CMB photon direction $\hat{\mathbf{n}} = \mathbf{k}/|\mathbf{k}|$ and the wave vectors \mathbf{K} , $x = \mathbf{K}(\eta_0 - \eta)$ and $\kappa(\eta) = \int_{\eta}^{\eta_0} d\eta a(\eta) \dot{\kappa}_{\pm}(\eta)$.

Eq. (6) shows that the photon-neutrino scattering ($\kappa \neq 0$) results in the non-trivial B-mode $\Delta_B^{(S)}$ and




Figure 2: EB power spectrum via CvB-CMB forward scattering is plotted in terms of l by considering CvB as Dirac and Majorana neutrino (left panel). Data of EB power spectrum are added by considering constraints on cosmological parameters using Planck, WMAP, and New BICEP2/Keck observations in 95 and 150 GHz signals (right panel).




Figure 3: $\Delta C_{TEI}^{(S)} = C_{TEI}^{(S)}|_{CvB} - C_{TEI}^{(S)}$ via CMB-CvB forward scattering is plotted in terms of l by considering CvB as Dirac and Majorana neutrino. We can estimate the contribution of CMB-CvB interaction on TE cross power spectrum by $\Delta C_{TEI}^{(S)}/C_{TEI}^{(S)} \simeq 1 - \cos(\bar{\kappa}) = 0.00026$ in the case of neutrino as Dirac particles which may be detected by increasing the precision of future experiments.




Figure 4: TB power spectrum via CMB-CvB forward scattering is plotted in terms of l by considering CvB as Dirac and Majorana neutrino (left panel). Data of TB power spectrum are added by considering constraints on cosmological parameters and using Planck, WMAP, and New BICEP2/Keck observations in 95 and 150 GHz signals (right panel). Since C_{TB}^l is zero in the standard scenario of cosmology, this result is full contribution of neutrino-photon scattering.

modifies the E-mode $\Delta_E^{(S)}$. This agrees that the Compton scattering by itself can not generate B-mode without taking into account the tensor type of metric perturbations [27, 56, 57]. Therefore, there are

three combined power spectra among B-mode, E-mode and T as in

$$C_{EB}^{\ell(S)} = (4\pi)^2 \frac{9}{16} \frac{(\ell+2)!}{(\ell-2)!} \int K^2 dKP_S(K) \Delta_E^{(S)} \Delta_B^{(S)}, \quad (7)$$

$$C_{TE}^{\ell(S)} = (4\pi)^2 \sqrt{\frac{9}{16} \frac{(\ell+2)!}{(\ell-2)!}} \int K^2 dKP_S(K) \Delta_T^{(S)} \Delta_E^{(S)}, \quad (8)$$

$$C_{TB}^{\ell(S)} = (4\pi)^2 \sqrt{\frac{9}{16} \frac{(\ell+2)!}{(\ell-2)!}} \int K^2 dKP_S(K) \Delta_T^{(S)} \Delta_B^{(S)}, \quad (9)$$

where

$$\begin{aligned} \Delta_T^{(S)}(K, \eta_0) &= \int_0^{\eta_0} d\eta g(\eta) S_T^{(S)}(\eta, K) j_\ell(x), \\ \Delta_E^{(S)}(K, \eta_0) &= \int_0^{\eta_0} d\eta g(\eta) \Pi(\eta, K) \frac{j_\ell(x)}{x^2} \cos \kappa(\eta), \\ \Delta_B^{(S)}(K, \eta_0) &= \int_0^{\eta_0} d\eta g(\eta) \Pi(\eta, K) \frac{j_\ell(x)}{x^2} \sin \kappa(\eta), \end{aligned} \quad (10)$$

and

$$\begin{aligned} S_T^{(S)}(K, \eta) &= g(\Delta_{T0} + 2\dot{\alpha} + \frac{\dot{v}_b}{K} + \frac{\Pi}{4} + \frac{3\ddot{\Pi}}{4K^2}) \\ &+ e^{-\tau_{e\nu}(\dot{\eta} + \ddot{\alpha})} + \dot{g}(\alpha + \frac{v_b}{K} + \frac{3\dot{\Pi}}{4K^2}) + \frac{3\ddot{g}\Pi}{4K^2}, \end{aligned} \quad (11)$$

in which $j_\ell(x)$ is a spherical Bessel function of rank ℓ , $\dot{\alpha} = (\dot{h} + 6\dot{\rho})/2K^2$, h , and ρ are additional sources of temperature anisotropy. The contribution of photon-neutrino (CvB) scattering to the cross power spectrum is negligible for small ℓ ; however, it is significantly large for $50 < \ell < 200$ (see Fig.(2-4)). As we already know, the non-zero $C_{TB}^{\ell(S)}$ may also appear in exotic theories due to the presence of helical flows in the primordial plasma at the time of recombination [58]. Therefore the CMB-CvB scattering is a new source of TB correlation as well.

The mean value κ in Eq.(10) is the effective average opacity of neutrino from the last scattering time (redshift $z_l = 1100$) to the present time ($z_0 = 0$). Using the Friedmann equation in the matter dominated era, $H^2/H_0^2 = \Omega_M^0(1+z)^3 + \Omega_\Lambda^0$, $H_0 \approx 74 \text{ km/s/Mpc}$, $\Omega_M^0 \approx 0.3 \pm 0.007$, $\Omega_\Lambda^0 \approx 0.7$ [59], and $ad\eta = -dz/H(1+z)$, as well as the conservation of total neutrino number $n_\nu = n_\nu^0(1+z)^3$, we obtain

$$\begin{aligned} \kappa(z) &= \int_\eta^{\eta_0} ad\eta \dot{\kappa}_\pm = \frac{\sqrt{2}}{12\pi} \alpha G^F n_\nu^0 \int_0^z dz' \frac{(1+z')^2}{H(z')} \\ &= \frac{\sqrt{2}}{12\pi} \alpha G^F n_\nu^0 \frac{2H(z')}{3\Omega_M^0 H_0^2} \Big|_0^z \end{aligned} \quad (12)$$

$$\beta|_{\nu D} = \frac{1}{2} \beta|_{\nu M} = \frac{1}{2} \bar{\kappa} \equiv \frac{1}{z_1 - z_0} \int_{z_1}^{z_0} dz \kappa(z) \approx 0.088 \left(\frac{n_\nu^0}{340 \text{ cm}^3} \right) \text{ rad} \quad (13)$$

where the present number-density of all flavor neutrinos and anti-neutrinos $n_\nu^0 = \sum(n_\nu^0 + n_{\bar{\nu}}^0) \approx 340 \text{ cm}^{-3}$. Actually, κ is the anisotropic opacity of CMB photons against the photon-neutrino (CvB) scattering. According to Fig.(5), β has strong dependence on the redshift. So, this dependency can not be ignored in the integration of Eq.(10). But if we want to have an estimation of our results, we approximately write $C_{EB}^{\ell(S)}$, $C_{TE}^{\ell(S)}$ and $C_{TB}^{\ell(S)}$ as follows




Figure 5: Variation of β in radian in terms of redshift.

$$\begin{aligned}
 C_{EB}^{l(S)} &\cong \frac{1}{2} \sin(2\bar{\kappa}) \bar{C}_{EE}^{l(S)} \\
 C_{TE}^{l(S)} &\cong \cos(\bar{\kappa}) \bar{C}_{TE}^{l(S)} \\
 C_{TB}^{l(S)} &\cong \sin(\bar{\kappa}) \bar{C}_{TE}^{l(S)}
 \end{aligned} \tag{14}$$

where $\bar{C}_{EE}^{l(S)}$ and $\bar{C}_{TE}^{l(S)}$ are power spectra of E-mode polarization and its correlation with temperature anisotropy contributed from the Compton scattering alone in the case of scalar perturbation.

III. Cosmic birefringence calculation

In the presence of CB, E-mode can be rotated to B-mode and vice versa, so:

$$\begin{aligned}
 \Delta_E^{(S)} &= \bar{\Delta}_E^{(S)} \cos(2\beta) - \bar{\Delta}_B^{(S)} \sin(2\beta), \\
 \Delta_B^{(S)} &= \bar{\Delta}_B^{(S)} \cos(2\beta) + \bar{\Delta}_E^{(S)} \sin(2\beta),
 \end{aligned} \tag{15}$$

where $\bar{\Delta}_E^{(S)}$ and $\bar{\Delta}_B^{(S)}$ are the value of E-mode and B-mode in standard scenario of cosmology without considering CB effects [60]. Note that in the standard scenario of cosmology, $\bar{\Delta}_B^{(S)} = 0$; so, we can rewrite the above equations as

$$\begin{aligned}
 \Delta_E^{(S)} &= \bar{\Delta}_E^{(S)} \cos(2\beta), \\
 \Delta_B^{(S)} &= \bar{\Delta}_E^{(S)} \sin(2\beta),
 \end{aligned} \tag{16}$$

As shown in Eqs.(10), actually we are faced with an anisotropic birefringence medium. In the other hand, the κ value is strongly dependent on η as well as red-shift (see Fig.(5)). The presence of Bessel functions (dependent on η too) in Eqs.(10) highlights the importance of κ dependence on red-shift. In such case, the comparing the theoretical results of $C_l^{(EB)}$ to the reported measurements to extract information is more logical and the red-shift dependency should be considered in the integration of Eqs.(10). But to give an effective sense, the CB angle β can be roughly estimated by the average value of opacity $\kappa(\eta)$. If we want to have more reliable estimation, we can use κ value around of the last scattering redshift ($z = 1100$), because, visibility function ($g(\eta)$) in the integrand of Eq.(10) has strong peak in that time. By substituting $\bar{\kappa}$ given in Eq.(13) instead of κ in Eq.(10) and comparing the results by Eqs.(16), we

have $\beta \simeq \frac{1}{2}\bar{\kappa}$. Finally we have

$$C_l^{(EE)} = \bar{C}_l^{(EE)} \cos^2 \bar{\kappa}, \quad (17)$$

$$C_l^{(BB)} = \bar{C}_l^{(EE)} \sin^2 \bar{\kappa}, \quad (18)$$

$$C_l^{(EB)} = \frac{1}{2}\bar{C}_l^{(EE)} \sin 2\bar{\kappa}. \quad (19)$$

Note that from the above equations, CB angle β is:

$$\beta|_{vD} = \frac{1}{2}\beta|_{vM} \cong \frac{1}{2}\bar{\kappa} = 2.54^\circ, \quad (20)$$

where $\beta|_{vD}$ and $\beta|_{vM}$ refer to the contribution of CvB-CMB forward scattering for CB angle in the case of Dirac (D) and Majorana (M), respectively. As mention above, κ has strong dependency on redshift, so averaging didn't seem to applicable. This point becomes more important, when the properties of the Bessel functions $j_\ell(x)$ present in Eqs.(10) takes into account.

Next point is that, here we only calculate the contribution of photon-neutrino scattering. There may be some other mechanisms generating EB power spectrum that can suppress the contribution of photon-neutrino interaction as well as CB angle. Also, the role of other cosmological parameters such as neutrino density, effective flavor number of neutrinos, and, Hubble parameter for the exact value of CB angle can not be ignored.

IV. Summary

In this study, by considering CMB-CvB forward scattering in the context of the standard model of particles as the collision term, we rewrote quantum Boltzmann equations in the presence of scalar perturbation of metric for the temperature fluctuations and linear polarization of CMB radiation by using CMB Stokes parameters (see Eq.(4)). Then, the explicit expressions for TE, TB, and EB cross power spectra were derived in the form of the line-of-sight integral solutions. We showed that the parity violating photon-neutrino forward scattering generated non-zero TB and EB power spectra in the presence of scalar perturbation. TE, TB, and EB power spectra via Compton scattering with the effect of CMB-CvB forward scattering were plotted in terms of l for CvB as Dirac and Majorana particles. By considering constraints on cosmological parameters using Planck, WMAP, and recent BICEP2/Keck observations in 95 and 150 GHz signals, cross power spectra data were added to our plots (see Figs.(2-4)). As a result, TB and EB power spectra in terms of l behaved like TE and EE power spectra, respectively. One of the main significant pieces of information that could be inferred from our study was about the nature of CvB. By analyzing the all cross power spectra, the contribution of CMB-CvB interaction could be completely separable by factor 2 for the Majorana neutrino with respect to Dirac neutrino.

Also, it was proved that CvB-CMB forward scattering can make an opaque medium; this opacity could rotate the linear polarization angle of CMB. As shown in Eqs.(10), this medium actually is an anisotropic birefringence medium and κ value is strongly dependent on red-shift Fig.(5). The presence of Bessel functions (and also dependence on η) in Eqs.(10) highlights the importance of κ dependence on redshift. In such case, the comparing $C_l^{(EB)}$ to its reported measurements is the most reasonable decision. As a result, our results for $C_l^{(EB)}$ is shown in Fig.(2).

Despite the extreme dependence of κ on red-shift, to give an effective sense, we show that the CB angle β can be roughly estimated by the average value of opacity $\beta|_v \simeq 1/2\bar{\kappa}$. As a result, we obtained $\beta|_{vD} \simeq 2.54^\circ$ and $\beta|_{vM} \simeq 5.08^\circ$, which referred to the contribution of CvB-CMB forward scattering for CB angle in the case of Dirac and Majorana respectively. It seems this result is not in agreement with reported measurement of [40]. As mentioned in the previous section, κ and the Bessel functions $j_\ell(x)$ present in Eqs.(10) have a strong dependency on redshift, so averaging didn't seem to be reliable.

Next point is that, here we only calculate the contribution of photon-neutrino scattering. There may be some other mechanisms generating EB power spectrum that can suppress the contribution of photon-neutrino interaction as well as CB angle. Also, the role of other cosmological parameters such as neutrino density and effective flavor number of neutrinos for the exact value of CB angle can not be ignored. We hope that by upgrading data of EB-cross power spectrum and theoretical and studies in the future, we could obtain more information about the origin of the reported CB angle and the nature of CvB.

Acknowledgments

We would like to thank E. Komatsu for his valuable comments on the calculation which provided the basis for the successful result. Further thanks go to Arthur Kosowsky for his helpful suggestions on this topic.

-
- [1] S. Weinberg, "Cosmology", Oxford University Press 2008.
 - [2] L.F. Li and F. Wilczek, Phys. Rev. D **25** (1982) 143; B. Kayser and R.E. Shrock, Phys. Lett. B **112** (1982) 137; B. Kayser, Phys. Rev. D **26** (1982) 1662
 - [3] R. N. Mohapatra and P. B. Pal, "Massive Neutrinos in Physics and Astrophysics", 3rd Edition, World Scientific (London, 2004).
 - [4] A. Faessler, R. Hodak, S. Kovalenko and F. Simkovic, Int. J. Mod. Phys. E **26**, no.01n02, 1740008 (2017) [[arXiv:1602.03347](#)].
 - [5] A. Faessler, R. Hodak, S. Kovalenko and F. Simkovic, [[arXiv:1102.1799](#)].
 - [6] R. Lazauskas, P. Vogel and C. Volpe, J. Phys. G **35**, 025001 (2008) [[arXiv:0710.5312](#)].
 - [7] A. Ringwald and Y. Y. Y. Wong, JCAP **12**, 005 (2004) [[arXiv:hep-ph/0408241](#) [hep-ph]].
 - [8] R. Wigmans, Astropart. Phys. **19**, 379-392 (2003) [[arXiv:astro-ph/0205360](#)].
 - [9] A. Ringwald, Strong and Electroweak Matter **2002**, pp. 497-503 (2003) [[arXiv:hep-ph/0301157](#) [hep-ph]].
 - [10] T. Weiler, Phys. Rev. Lett. **49**, 234 (1982).
 - [11] F. Munyaneza and P. L. Biermann, Astron. Astrophys. **458**, L9-L12 (2006) [[arXiv:astro-ph/0609388](#)].
 - [12] H. Pas and T. J. Weiler, Phys. Rev. D **63**, 113015 (2001) [[arXiv:hep-ph/0101091](#)].
 - [13] C. Yanagisawa, Frontiers in Physics, **2**, 30 (2014).
 - [14] R. Mohammadi, J. Khodagholizadeh, M. Sadegh and S. S. Xue, Phys. Rev. D **93**, no.12, 125029 (2016) [[arXiv:1602.00237](#)].
 - [15] J. Khodagholizadeh, R. Mohammadi and S. S. Xue, Phys. Rev. D **90**, no.9, 091301 (2014) [[arXiv:1406.6213](#) [astro-ph.CO]].
 - [16] R. Mohammadi and S. S. Xue, Phys. Lett. B **731**, 272-278 (2014) [[arXiv:1312.3862](#) [hep-ph]].
 - [17] R. Mohammadi, Eur. Phys. J. C **74**, no.10, 3102 (2014) [[arXiv:1312.2199](#) [astro-ph.CO]].
 - [18] M. Kamionkowski and E. D. Kovetz, Ann. Rev. Astron. Astrophys. **54**, 227-269 (2016) [[arXiv:1510.06042](#)].
 - [19] D. Jeong and M. Kamionkowski, Phys. Rev. Lett. **124**, no.4, 041301 (2020) doi:10.1103/PhysRevLett.124.041301 [[arXiv:1908.06100](#)].
 - [20] A. Zucca, Y. Li and L. Pogosian, Phys. Rev. D **95**, no.6, 063506 (2017) doi:10.1103/PhysRevD.95.063506 [[arXiv:1611.00757](#)].
 - [21] K. N. Abazajian *et al.* [Topical Conveners: K.N. Abazajian, J.E. Carlstrom, A.T. Lee], Astropart. Phys. **63**, 66-80 (2015) doi:10.1016/j.astropartphys.2014.05.014 [[arXiv:1309.5383](#)].
 - [22] K. N. Abazajian *et al.* [CMB-S4], [[arXiv:1610.02743](#)].
 - [23] A. Kosowsky, Annals Phys. **246**, 49-85 (1996) [[arXiv:astro-ph/9501045](#)].
 - [24] U. Seljak and M. Zaldarriaga, Phys. Rev. Lett. **78**, 2054-2057 (1997) [[arXiv:astro-ph/9609169](#) [astro-ph]].
 - [25] M. Zaldarriaga and U. Seljak, Phys. Rev. D **55**, 1830-1840 (1997) [[arXiv:astro-ph/9609170](#)].
 - [26] M. Zaldarriaga, D. N. Spergel and U. Seljak, Astrophys. J. **488**, 1-13 (1997) [[arXiv:astro-ph/9702157](#)].
 - [27] W. Hu and M. J. White, New Astron. **2**, 323 (1997) [[arXiv:astro-ph/9706147](#)].
 - [28] M. Betoule, E. Pierpaoli, J. Delabrouille, M. Le Jeune, J. F. Cardoso, Astron. Astrophys., **503**, 691-706 (2009). [[arXiv:0901.1056](#)].
 - [29] J. A. Vazquez, M. Bridges, Y. Z. Ma and M. P. Hobson, JCAP **08**, 001 (2013) [[arXiv:1303.4014](#) [astro-

- ph.CO]].
- [30] J. W. Henning *et al.* [SPT], *Astrophys. J.* **852**, no.2, 97 (2018) [[arXiv:1707.09353](#)].
 - [31] N. Aghanim *et al.* [Planck], *Astron. Astrophys.* **641**, A5 (2020) [[arXiv:1907.12875](#)].
 - [32] T. Louis *et al.* [ACTPol], *JCAP* **06**, 031 (2017) [[arXiv:1610.02360](#)].
 - [33] S. Galli, K. Benabed, F. Bouchet, J. F. Cardoso, F. Elsner, E. Hivon, A. Mangilli, S. Prunet and B. Wandelt, *Phys. Rev. D* **90**, no.6, 063504 (2014) [[arXiv:1403.5271](#)].
 - [34] M. Kamionkowski, A. Kosowsky and A. Stebbins, *Phys. Rev. Lett.* **78**, 2058-2061 (1997) [[arXiv:astro-ph/9609132](#) [astro-ph]].
 - [35] A. Lue, L. M. Wang and M. Kamionkowski, *Phys. Rev. Lett.* **83**, 1506-1509 (1999) [[arXiv:astro-ph/9812088](#)].
 - [36] G. C. Liu, S. Lee and K. W. Ng, *Phys. Rev. Lett.* **97**, 161303 (2006) [[arXiv:astro-ph/0606248](#)].
 - [37] S. M. Carroll, G. B. Field and R. Jackiw, *Phys. Rev. D* **41**, 1231 (1990)
 - [38] V. Gluscevic, M. Kamionkowski and A. Cooray, *Phys. Rev. D* **80**, 023510 (2009) [[arXiv:0905.1687](#)].
 - [39] P. Montero-Camacho and C. M. Hirata, *JCAP* **08**, 040 (2018) [[arXiv:1803.04505](#) [astro-ph.CO]].
 - [40] Y. Minami and E. Komatsu, *Phys. Rev. Lett.* **125**, no.22, 221301 (2020) [[arXiv:2011.11254](#)].
 - [41] T. Namikawa *et al.* [Atacama Cosmology Telescope] *Phys. Rev. D* **101** (2020) no.8, 083527 [[arXiv:2001.10465](#) [astro-ph.CO]].
 - [42] P. A. R. Ade *et al.* [POLARBEAR], *Astrophys. J.* **794** (2014) no.2, 171 [[arXiv:1403.2369](#) [astro-ph.CO]].
 - [43] F. Bianchini *et al.* [SPT], *Phys. Rev. D* **102** (2020) no.8, 083504 [[arXiv:2006.08061](#) [astro-ph.CO]].
 - [44] M. Abdi, R. Mohammadi, S. S. Xue and M. Zarei, [[arXiv:1909.01536](#)].
 - [45] Z. Chacko, P. S. Bhupal Dev, R. N. Mohapatra and A. Thapa, *Phys. Rev. D* **102**, no.3, 035020 (2020) [[arXiv:2005.05413](#)].
 - [46] A. Segarra and J. Bernabéu, *Phys. Rev. D* **101**, no.9, 093004 (2020) [[arXiv:2001.05900](#)].
 - [47] A. B. Balantekin, A. de Gouvêa and B. Kayser, *Phys. Lett. B* **789**, 488-495 (2019) [[arXiv:1808.10518](#)].
 - [48] C. Arbelaéz, C. Dib, I. Schmidt and J. C. Vasquez, *Phys. Rev. D* **97**, no.5, 055011 (2018) [[arXiv:1712.08704](#)].
 - [49] W. Rodejohann, X. J. Xu and C. E. Yaguna, *JHEP* **05**, 024 (2017) [[arXiv:1702.05721](#)].
 - [50] A. Capolupo, S. M. Giampaolo, B. C. Hiesmayr and G. Vitiello, *Phys. Lett. B* **780**, 216-220 (2018) [[arXiv:1610.08679](#)].
 - [51] J. Lu, *Phys. Lett. B* **818**, 136376 (2021) [[arXiv:2105.07658](#) [hep-ph]].
 - [52] L. Johns, [[arXiv:2107.11434](#) [hep-ph]].
 - [53] S. Chandrasekhar, *Radiative Transfer*, Dover, New York, 1960.
 - [54] J. Khodagholizadeh, R. Mohammadi and S. S. Xue, *Phys. Rev. D* **90**, no. 9, 091301 (2014). [[arXiv:1406.6213](#)].
 - [55] R. Mohammadi and S. S. Xue, *Phys. Lett. B* **731**, 272-278 (2014) [[arXiv:1312.3862](#)]; R. Mohammadi, *Eur. Phys. J. C* **74**, no.10, 3102 (2014) [[arXiv:1312.2199](#)].
 - [56] M. Zaldarriaga and D. D. Harari, *Phys. Rev. D* **52**, 3276-3287 (1995) [[arXiv:astro-ph/9504085](#)].
 - [57] U. Seljak, *Astrophys. J.* **482**, 6 (1997) [[arXiv:astro-ph/9608131](#)].
 - [58] L. Pogosian, T. Vachaspati and S. Winitzki, *Phys. Rev. D* **65**, 083502 (2002) [[arXiv:astro-ph/0112536](#)].
 - [59] N. Aghanim *et al.* [Planck], *Astron. Astrophys.* **641** (2020), A6 [[arXiv:1807.06209](#) [astro-ph.CO]].
 - [60] Y. Minami, H. Ochi, K. Ichiki, N. Katayama, E. Komatsu and T. Matsumura, *PTEP* **2019**, no.8, 083E02 (2019) [[arXiv:1904.12440](#) [astro-ph.CO]].