

INEQUALITIES AND LIMITS OF WEIGHTED SPECTRAL GEOMETRIC MEAN

LUYINING GAN AND TIN-YAU TAM

In memory of Professor Boying Wang who passed away on August 20, 2021.

ABSTRACT. We establish some new properties of spectral geometric mean. In particular, we prove a log majorization relation between $(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}$ and the t -spectral mean $A\sharp_t B := (A^{-1}\sharp B)^t A (A^{-1}\sharp B)^t$ of two positive semidefinite matrices A and B , where $A\sharp B$ is the geometric mean, and the t -spectral mean is the dominant one. The limit involving t -spectral mean is also studied. We then extend all the results in the context of symmetric spaces of negative curvature.

1. INTRODUCTION

Let $\mathbb{C}_{n \times n}$ be the space of all $n \times n$ complex matrices, \mathbb{H}_n the real space of $n \times n$ Hermitian matrices, \mathbb{P}_n the set of $n \times n$ positive definite matrices in $\mathbb{C}_{n \times n}$ and $U(n)$ the group of $n \times n$ unitary matrices. For any $X \in \mathbb{C}_{n \times n}$, both e^X and $\exp X$ denote the exponential of X . Given $A \in \mathbb{C}_{n \times n}$, we use $A \geq 0$ to denote that A is positive semidefinite. Given $A, B \in \mathbb{H}_n$, denote by $A \leq B$ the Löwner order, that is, $B - A \geq 0$. Given $X \in \mathbb{C}_{n \times n}$, denote by $\|X\|$ the spectral norm of X , that is, the largest singular value of X , and denote by $\sigma(X)$ the spectrum of X . If the eigenvalues of X are all real, we write $\lambda(X) = (\lambda_1, \dots, \lambda_n)$, where $\lambda_1(X) \geq \dots \geq \lambda_n(X)$ are the eigenvalues of X .

The metric geometric mean (geometric mean, for short) of $A, B \in \mathbb{P}_n$

$$A\sharp B := A^{1/2}(A^{-1/2}BA^{-1/2})^{1/2}A^{1/2}, \quad (1.1)$$

was first introduced by Pusz and Woronowicz [28] in 1975 and further studied by Kubo and Ando [20] in the 1970s. Since then, it has been extensively studied. Though the definition (1.1) looks awkward, it is indeed a natural generalization of the classical geometric mean \sqrt{ab} of two positive numbers a, b [7]. Besides the algebraic formulation and properties, geometric mean has a rich geometric flavor which is due to the fact that \mathbb{P}_n is a Riemannian manifold [12] and that the geometric mean $A\sharp B$ is the mid-point of the unique geodesic joining A and B [7, 24].

The spectral geometric mean (spectral mean, for short) of $A, B \in \mathbb{P}_n$ was introduced by Fiedler and Pták [9] in 1997 and one of the formulations is

$$A\sharp_t B := (A^{-1}\sharp B)^{1/2} A (A^{-1}\sharp B)^{1/2}. \quad (1.2)$$

2020 *Mathematics Subject Classification.* 15A16, 15A45, 15B48, 22E46.

Key words and phrases. Geometric mean, weighted spectral mean, log majorization, Kostant's pre-order.

Professor Boying Wang, a faculty member of Beijing Normal University during 1960–2002, was a pioneer of developing and promoting Multilinear Algebra in China.

They named it as spectral geometric mean because the square of $A\sharp B$ is similar to AB , which means that the eigenvalues of their spectral mean are the positive square roots of the corresponding eigenvalues of AB [9, Theorem 3.2 and Remark 3.4]. As the spectral mean comes from the geometric mean, it possesses some important properties of the geometric mean and has been studied [17, 21, 23, 25]. However, unlike the geometric mean, not many results have been obtained on the spectral mean. Thus, this paper aims to obtain new results on the spectral mean and its extension, namely, the t -spectral mean. Some of the results are analogous to the geometric mean.

For each $t \in [0, 1]$, the t -metric geometric mean (t -geometric mean, for short) and t -spectral geometric mean (t -spectral mean, for short) of A and B are naturally defined by

$$A\sharp_t B := A^{1/2}(A^{-1/2}BA^{-1/2})^t A^{1/2}, \quad t \in [0, 1], \quad (1.3)$$

$$A\sharp_t B := (A^{-1}\sharp B)^t A (A^{-1}\sharp B)^t, \quad t \in [0, 1]. \quad (1.4)$$

Lee and Lim [22] first introduced the t -spectral mean in 2007. In the same year, it was also studied by Ahn, Kim and Lim [1, p.191] (also see [15, p.446]). Its further algebraic and geometric meaning has been recently studied by Kim [16]. When $t = 1/2$, they are abbreviated as $A\sharp_{1/2} B = A\sharp B$ and $A\sharp_{1/2} B = A\sharp B$.

Both (1.3) and (1.4) are paths joining A (when $t = 0$) and B (when $t = 1$) in \mathbb{P}_n . Here is a good way to interpret (1.3). Move the points A and B to I and $A^{-1/2}BA^{-1/2}$ respectively, via the congruence action by $A^{-1/2}$:

$$X \mapsto A^{-1/2} X A^{-1/2}, \quad X \in \mathbb{P}_n.$$

The t -geometric mean of the commuting I and $A^{-1/2}BA^{-1/2}$ is

$$I\sharp_t(A^{-1/2}BA^{-1/2}) = (A^{-1/2}BA^{-1/2})^t.$$

Then apply the inverse action on $(A^{-1/2}BA^{-1/2})^t$

$$Y \mapsto A^{1/2} Y A^{1/2}, \quad Y \in \mathbb{P}_n,$$

to have $A\sharp_t B = A^{1/2}(A^{-1/2}BA^{-1/2})^t A^{1/2}$. This nice feature follows from the fact that $S\mathbb{P}_n := \{A \in \mathbb{P}_n : \det A = 1\}$ is a symmetric space [12, p.208-209], when it is identified with $SL_n(\mathbb{C})/SU(n)$ via the polar decomposition, where $SL_n(\mathbb{C})$ is the special linear group over \mathbb{C} and $SU(n)$ is the special unitary group.

In this paper, we focus our study on t -spectral mean and organize the sections as follows. In Section 2 we provide a similarity property. In Section 3, we review a log majorization result of t -geometric mean and show that a similar result for the t -spectral mean. In particular, we prove that $(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}$ is log majorized by $(A^s\sharp_t B^s)^{1/s}$ for $s > 0$, and also log majorized by the t -spectral mean $A\sharp_t B$ for a range of s with respect to t . In Section 4, we provide the limit of t -spectral mean when p tends to 0. The results in Sections 2, 3 and 4 are then extended in the context of symmetric spaces associated with a noncompact semisimple Lie group in Section 5. In Section 6, some remarks are given.

2. PROPERTIES

In this section, we establish a similarity property of t -spectral mean. Let $A, B \in \mathbb{P}_n$. We say that A is *positively similar* to B if there exists $C \in \mathbb{P}_n$ such that $A = CBC^{-1}$. Let us first recall some basic properties of the t -spectral mean [7, 10, 22] in the following theorem.

Theorem 2.1. Let $A, B \in \mathbb{P}_n$ and $t \in [0, 1]$. Then

- (1) $(A\sharp_t B)^{-1} = A^{-1}\sharp_t B^{-1}$ and $A\sharp_t B = B\sharp_{1-t} A$.
- (2) $A^{-1}\sharp(A\sharp_t B) = (B\sharp_t A)^{-1}\sharp B = (A^{-1}\sharp B)^t$.
- (3) If $G_t = A^{-1}\sharp(A\sharp_t B)$, then $A\sharp_t B = G_t A G_t$ and $B\sharp_t A = G_t^{-1} B G_t^{-1}$.
- (4) $(A\sharp_r B)\sharp_t(A\sharp_s B) = A\sharp_{(1-t)r+ts} B$ for all $t, r, s \in [0, 1]$.

Fiedler and Pták [9, Theorem 5.5(5)] obtained a positively similarity relation between the geometric mean and the spectral mean.

Theorem 2.2. (Fiedler and Pták 1997) Given $A, B \in \mathbb{P}_n$, the geometric mean $A\sharp B$ is positively similar to $(A\sharp B)^{1/2} U (A\sharp B)^{1/2}$ for some $U \in U(n)$.

Motivated by the above theorem, we prove that the geometric mean and t -spectral mean are positively similar, which is reduced to the result of Fiedler and Pták when $t = 1/2$.

Theorem 2.3. Let $A, B \in \mathbb{P}_n$ and $t \in [0, 1]$. Then $A\sharp B$ is positively similar to

$$(A\sharp_{1-t} B)^{1/2} U (A\sharp_t B)^{1/2}$$

for some $U \in U(n)$.

Proof. By Theorem 2.1(3),

$$A\sharp_t B = G_t A G_t, \quad B\sharp_t A = G_t^{-1} B G_t^{-1},$$

where $G_t = A^{-1}\sharp(A\sharp_t B) \in \mathbb{P}_n$, that is,

$$A = G_t^{-1} (A\sharp_t B) G_t^{-1}, \quad B = G_t (B\sharp_t A) G_t.$$

Set

$$W := G_t (B\sharp_t A)^{1/2}, \quad V := (A\sharp_t B)^{-1/2} G_t.$$

Then

$$W^* V^{-1} = (B\sharp_t A)^{1/2} (A\sharp_t B)^{1/2}. \tag{2.1}$$

Now

$$G_t (B\sharp_t A) G_t = B = (A\sharp B) A^{-1} (A\sharp B) = (A\sharp B) G_t (A\sharp_t B)^{-1} G_t (A\sharp B),$$

where the second equality holds because $A^{-1}\sharp_t B = (A\sharp B)^t A^{-1} (A\sharp B)^t$, $0 \leq t \leq 1$, is a curve joining A^{-1} (when $t = 0$) and B (when $t = 1$) in view of (1.4). Then

$$\begin{aligned} & (A\sharp_t B)^{-1/2} G_t^2 (B\sharp_t A) G_t^2 (A\sharp_t B)^{-1/2} \\ &= (A\sharp_t B)^{-1/2} G_t (A\sharp B) G_t (A\sharp_t B)^{-1} G_t (A\sharp B) G_t (A\sharp_t B)^{-1/2} \\ &= [(A\sharp_t B)^{-1/2} G_t (A\sharp B) G_t (A\sharp_t B)^{-1/2}]^2. \end{aligned}$$

Thus we have

$$\begin{aligned} A\sharp B &= [(A\natural_t B)^{-1/2} G_t]^{-1} [(A\natural_t B)^{-1/2} G_t^2 (B\natural_t A) G_t^2 (A\natural_t B)^{-1/2}]^{1/2} [G_t (A\natural_t B)^{-1/2}]^{-1} \\ &= V^{-1} (V W W^* V^*)^{1/2} (V^*)^{-1}. \end{aligned}$$

Set $R := VW$ and from (2.1) $R = VV^*(A\natural_t B)^{1/2}(B\natural_t A)^{1/2}$. The matrix $U := R^{-1}(RR^*)^{1/2}$ is unitary since

$$UU^* = R^{-1}(RR^*)^{1/2}(RR^*)^{1/2}(R^*)^{-1} = R^{-1}(RR^*)(R^*)^{-1} = I.$$

Then we have

$$\begin{aligned} (A\natural_{1-t} B)^{1/2} U (A\natural_t B)^{1/2} &= (B\natural_t A)^{1/2} U (A\natural_t B)^{1/2} \\ &= (B\natural_t A)^{1/2} R^{-1} (RR^*)^{1/2} (A\natural_t B)^{1/2} \\ &= (B\natural_t A)^{1/2} [VV^*(A\natural_t B)^{1/2}(B\natural_t A)^{1/2}]^{-1} (VWW^*V^*)^{1/2} (A\natural_t B)^{1/2} \\ &= (A\natural_t B)^{-1/2} (V^*)^{-1} V^{-1} (VWW^*V^*)^{1/2} (V^*)^{-1} V^* (A\natural_t B)^{1/2} \\ &= [V^*(A\natural_t B)^{1/2}]^{-1} V^{-1} (VWW^*V^*)^{1/2} (V^*)^{-1} [V^*(A\natural_t B)^{1/2}] \\ &= [V^*(A\natural_t B)^{1/2}]^{-1} (A\sharp B) [V^*(A\natural_t B)^{1/2}], \end{aligned}$$

where $V^*(A\natural_t B)^{1/2} = G_t^* \in \mathbb{P}_n$ since $V = (A\natural_t B)^{-1/2} G_t$. Thus we complete the proof. \square

3. LOG MAJORIZATION

Let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ be in \mathbb{R}^n . Let $x^\downarrow = (x_{[1]}, x_{[2]}, \dots, x_{[n]})$ denote the rearrangement of the components of x such that $x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n]}$. We say that x is *majorized* by y [27], denoted by $x \prec y$, if

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}, \quad k = 1, 2, \dots, n-1, \quad \text{and} \quad \sum_{i=1}^n x_{[i]} = \sum_{i=1}^n y_{[i]}.$$

Among many equivalent conditions for majorization, the following is geometric in nature, noted by Rado [29] and A. Horn [14]:

$$x \prec y \quad \Leftrightarrow \quad \text{conv } S_n \cdot x \subset \text{conv } S_n \cdot y,$$

where $\text{conv } S_n \cdot x$ denotes the convex hull of the orbit of x under the action of the symmetric group S_n . See a good summary in Marshall, Olkin, and Arnold [27, p.10-14, p.34]. When x and y are nonnegative vectors, we say that x is *log majorized* by y , denoted by $x \prec_{\log} y$ if

$$\prod_{i=1}^k x_{[i]} \leq \prod_{i=1}^k y_{[i]}, \quad k = 1, 2, \dots, n-1 \quad \text{and} \quad \prod_{i=1}^n x_{[i]} = \prod_{i=1}^n y_{[i]}.$$

When x and y are positive vectors, $x \prec_{\log} y$ if and only if $\log x \prec \log y$, where $\log x := (\log x_1, \log x_2, \dots, \log x_n)$.

A natural way to extend the notion of log majorization from nonnegative real vectors to positive semidefinite matrices is via their eigenvalues, that is, given $X, Y \geq 0$, we write $X \prec_{\log} Y$ when $\lambda(X) \prec_{\log} \lambda(Y)$. As a relation, log majorization is transitive, reflexive but not anti-symmetric, so it is not a partial order. Needless to say, it is different from

the Löwner order \leq which is a partial order. We would like to point out that neither one implies the other.

Remark 3.1. We would like to point out that $X \prec_{\log} Y$ can be extended to X, Y which are diagonalizable with nonnegative eigenvalues. For example if A, B are positive semidefinite, then AB is diagonalizable with nonnegative eigenvalues, though AB is not Hermitian in general.

The t -geometric mean has been studied extensively and a lot of nice properties have been discovered. For example, the following result of Ando and Hiai [3, Theorem 2.1] gives a log majorization relation between the t -geometric mean of r -powers of positive semidefinite A and B and the r -power of the t -geometric mean of A and B .

Theorem 3.2. (Ando and Hiai 1994) For every $A, B \geq 0$ and $0 \leq t \leq 1$,

$$A^r \sharp_t B^r \prec_{\log} (A \sharp_t B)^r, \quad r \geq 1, \quad (3.1)$$

$$(A \sharp_t B)^r \prec_{\log} A^r \sharp_t B^r, \quad 0 < r \leq 1, \quad (3.2)$$

$$(A^p \sharp_t B^p)^{1/p} \prec_{\log} (A^q \sharp_t B^q)^{1/q}, \quad 0 < q \leq p. \quad (3.3)$$

Theorem 3.2 was extended to symmetric spaces of negative curvature by Liao, Liu and Tam [24, Theorem 3.7]. See Remark 5.5 for the geometry associated with (3.1), (3.2) and (3.3). Motivated by Theorem 3.2, we would like to know if analogous relation holds for the t -spectral mean. The following theorem shows that such relation does exist, but in reverse order.

Theorem 3.3. For every $A, B \geq 0$ and $0 \leq t \leq 1$,

$$(A \natural_t B)^r \prec_{\log} A^r \natural_t B^r, \quad r \geq 1, \quad (3.4)$$

$$A^r \natural_t B^r \prec_{\log} (A \natural_t B)^r, \quad 0 < r \leq 1, \quad (3.5)$$

$$(A^q \natural_t B^q)^{1/q} \prec_{\log} (A^p \natural_t B^p)^{1/p}, \quad 0 < q \leq p, \quad (3.6)$$

that is, $p \rightarrow (A^p \natural_t B^p)^{1/p}$ is a log majorization increasing function on $(0, \infty)$.

Proof. We first prove (3.4). We may consider $A, B > 0$ by continuity argument. It is easy to see

$$\det((A \natural_t B)^r) = (\det A)^{(1-t)r} (\det B)^{tr} = \det(A^r \natural_t B^r).$$

Recall [27, p.776-777] that

$$\prod_{i=1}^k \lambda_i(A) = \lambda_1(C_k(A)), \quad k = 1, \dots, n, \quad (3.7)$$

where $C_k(A)$ denotes the k th compound of $A \geq 0$. Thus, we need to show

$$\lambda_1(C_k((A \natural_t B)^r)) \leq \lambda_1(C_k(A^r \natural_t B^r)), \quad k = 1, \dots, n-1.$$

Note that [3, Lemma 1.2], [8, p.781] $C_k(A \sharp_t B) = C_k(A) \sharp_t C_k(B)$ for $k = 1, \dots, n$, and $C_k : \text{GL}_n(\mathbb{C}) \rightarrow \text{GL}_{\binom{n}{k}}(\mathbb{C})$ is a group representation of the general linear group $\text{GL}_n(\mathbb{C})$.

So from (1.4), we have for $r \in \mathbb{R}$ and $t \in [0, 1]$,

$$C_k(A^r \natural_t B^r) = C_k(A)^r \natural_t C_k(B)^r \quad (3.8)$$

and

$$C_k((A\sharp_t B)^r) = (C_k(A)\sharp_t C_k(B))^r.$$

Hence it suffices to show that

$$\lambda_1(A\sharp_t B)^r \leq \lambda_1(A^r\sharp_t B^r). \quad (3.9)$$

By joint homogeneity of t -spectral mean, we have for $\alpha, \beta > 0$

$$((\alpha A)\sharp_t(\beta B))^r = \alpha^{r(1-t)}\beta^{rt}(A\sharp_t B)^r \quad (3.10)$$

and

$$((\alpha A)^r\sharp_t(\beta B)^r) = (\alpha^r A^r)\sharp_t(\beta^r B^r) = \alpha^{r(1-t)}\beta^{rt}(A^r\sharp_t B^r). \quad (3.11)$$

In other words, both sides of (3.9) have the same order of homogeneity for A, B . Thus to prove (3.9), we may show that

$$\lambda_1(A^r\sharp_t B^r) \leq 1 \Rightarrow \lambda_1(A\sharp_t B)^r \leq 1. \quad (3.12)$$

Suppose that $\lambda_1(A^r\sharp_t B^r) \leq 1$, that is, $\lambda_1((A^{-r}\sharp_t B^r)^t A^r (A^{-r}\sharp_t B^r)^t) \leq 1$, we have

$$(A^{-r}\sharp_t B^r)^t A^r (A^{-r}\sharp_t B^r)^t \leq I$$

and thus

$$A^r \leq (A^{-r}\sharp_t B^r)^{-2t}.$$

For $r \geq 1$, that is, $0 < 1/r \leq 1$, we have

$$A \leq (A^{-r}\sharp_t B^r)^{-2t/r}. \quad (3.13)$$

When $r \geq 1$, we have from (3.1)

$$A^r \sharp_t B^r \prec_{\log} (A\sharp_t B)^r$$

so that

$$A^{-r}\sharp_t B^r \prec_{\log} (A^{-1}\sharp_t B)^r.$$

As $r \geq 1$, from (3.13) and (3.1) we have

$$\lambda_1(A) \leq \lambda_1((A^{-r}\sharp_t B^r)^{-2t/r}) \leq \lambda_1((A^{-1}\sharp_t B)^{-2t}),$$

which is equivalent to

$$\lambda_1((A^{-1}\sharp_t B)^{2t})\lambda_1(A) \leq 1.$$

Since $A, (A^{-1}\sharp_t B)^{2t} \geq 0$, we have

$$\lambda_1((A^{-1}\sharp_t B)^{2t})\lambda_1(A) \geq \lambda_1((A^{-1}\sharp_t B)^{2t}A) = \lambda_1((A^{-1}\sharp_t B)^t A (A^{-1}\sharp_t B)^t) = \lambda_1(A\sharp_t B).$$

Thus we get $\lambda_1(A\sharp_t B) \leq 1$, that is, (3.12) is established. Thus we complete the proof of (3.4). We omit the proofs of (3.5) and (3.6) due to the similar idea as that of (3.4). \square

Let us recall some interesting results in the following theorem.

Theorem 3.4. (Ando and Hiai 1994, Araki 1990) Let $A, B \in \mathbb{P}_n$. For any $t \in [0, 1]$ and $s > 0$,

$$A\sharp_t B \prec_{\log} e^{(1-t)\log A + t\log B} \quad (3.14)$$

$$\prec_{\log} (B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}. \quad (3.15)$$

The first inequality (3.14) is a result of Ando and Hiai [3, Corollary 2.3] as the complementary counterpart of the famous Golden-Thompson inequality for Hermitian matrices A and B :

$$\operatorname{tr} e^{A+B} \leq \operatorname{tr} (e^A e^B).$$

We remark that the complementary Golden-Thompson inequality

$$\operatorname{tr} (e^{pA} \sharp_t e^{pB})^{1/p} \leq \operatorname{tr} e^{(1-t)A+tB}, \quad p > 0, \quad 0 \leq t \leq 1,$$

was first proved by Hiai and Petz [13] and then extended to log majorization by Ando and Hiai [3]. The second inequality (3.15) follows from a result of Araki [4].

Very recently Gan, Liu and Tam [10] have proved the following result which asserts that the t -geometric mean of two positive definite matrices is log majorized by their t -spectral mean.

Theorem 3.5. (Gan, Liu, and Tam 2021) For all $A, B \in \mathbb{P}_n$ and $t \in [0, 1]$,

$$A \sharp_t B \prec_{\log} A \natural_t B. \quad (3.16)$$

Motivated by Theorem 3.4 and Theorem 3.5, it is natural to ask whether an analogous log majorization relation exists between $(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}$ and $A \natural_t B$, or between $e^{(1-t)\log A + t\log B}$ and $A \natural_t B$. The former would be a stronger result than the latter. We state this stronger result in Theorem 3.7 in which the range of s is specified. Before stating Theorem 3.7, we prove another interesting inequality between $(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}$ and $(A^s \natural_t B^s)^{1/s}$ for all positive s .

Theorem 3.6. Let $A, B \in \mathbb{P}_n$. For $t \in [0, 1]$ and $s > 0$. We have

$$(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s} \prec_{\log} (A^s \natural_t B^s)^{1/s}. \quad (3.17)$$

In particular, setting $s = 1$ yields

$$A \sharp_t B \prec_{\log} e^{(1-t)\log A + t\log B} \prec_{\log} B^{t/2} A^{1-t} B^{t/2} \prec_{\log} A \natural_t B, \quad 0 \leq t \leq 1. \quad (3.18)$$

Proof. It is easy to see that for $s > 0$, we have

$$\det(A^s \natural_t B^s)^{1/s} = (\det A)^{1-t} (\det B)^t = \det(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}.$$

Indeed, it is true for $s \in \mathbb{R}$. By (3.7), we need to show that for $s > 0$,

$$\lambda_1(C_k((B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s})) \leq \lambda_1(C_k((A^s \natural_t B^s)^{1/s})), \quad k = 1, \dots, n-1.$$

As the compound $C_k : \operatorname{GL}_n(\mathbb{C}) \rightarrow \operatorname{GL}_{\binom{n}{k}}(\mathbb{C})$ is a group representation of the general linear group $\operatorname{GL}_n(\mathbb{C})$, we have

$$C_k((B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}) = (C_k(B)^{ts/2} C_k(A)^{(1-t)s} C_k(B)^{ts/2})^{1/s},$$

for all $k = 1, \dots, n$. From (3.8), we have

$$C_k((A^s \natural_t B^s)^{1/s}) = (C_k(A)^s \natural_t C_k(B)^s)^{1/s}.$$

Hence it suffices to show that for all $A, B \in \mathbb{P}_n$ and $s > 0$,

$$\lambda_1((B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}) \leq \lambda_1((A^s \natural_t B^s)^{1/s}). \quad (3.19)$$

Note that for $\alpha, \beta > 0$,

$$((\beta B)^{ts/2}(\alpha A)^{(1-t)s}(\beta B)^{ts/2})^{1/s} = \alpha^{1-t}\beta^t(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}$$

and

$$((\alpha A)^s \natural_t (\beta B)^s)^{1/s} = ((\alpha^s A^s) \natural_t (\beta^s B^s))^{1/s} = \alpha^{(1-t)}\beta^t(A^s \natural_t B^s)^{1/s}.$$

Thus $(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}$ and $A \natural_t B$ have the same order of homogeneity for A, B . Then we may show that

$$\lambda_1((A^s \natural_t B^s)^{1/s}) \leq 1 \Rightarrow \lambda_1((B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}) \leq 1. \quad (3.20)$$

Let $C(s) := A^{-s} \sharp B^s$. Because $\lambda_1((A^s \natural_t B^s)^{1/s}) \leq 1$, which means that

$$A^s \natural_t B^s = C^t(s)A^s C^t(s) \leq I,$$

we know [6, p.114]

$$A^s \leq C^{-2t}(s). \quad (3.21)$$

Applying the Riccati equation to $C(s) := A^{-s} \sharp B^s$, we have from [7, p.11] and (3.21)

$$B^s = C(s)A^s C(s) \leq C^{2(1-t)}(s). \quad (3.22)$$

Since $t \in [0, 1]$ and $1-t \in [0, 1]$, it follows from (3.21), (3.22), and [6, p.115] that

$$A^{(1-t)s} \leq C^{-2t(1-t)}(s) \leq B^{-ts}.$$

It amounts to $B^{ts/2}A^{(1-t)s}B^{ts/2} \leq I$, which means that $\lambda_1(B^{ts/2}A^{(1-t)s}B^{ts/2}) \leq 1$. Thus we have

$$\lambda_1((B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}) = \lambda_1^{1/s}(B^{ts/2}A^{(1-t)s}B^{ts/2}) \leq 1,$$

that is, (3.20) is proved and thus the proof of (3.23) is completed.

Note that when $s = 1$, it is natural to have

$$B^{t/2}A^{1-t}B^{t/2} \prec_{\log} A \natural_t B.$$

By Theorem 3.4, we have (3.18) since \prec_{\log} is transitive. \square

Theorem 3.7. Let $A, B \in \mathbb{P}_n$. For each chosen $t \in [0, 1]$, let $0 < s \leq \min\{1/t, 1/(1-t)\}$. We have

$$(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s} \prec_{\log} A \natural_t B. \quad (3.23)$$

We omit the proof because it is almost identical to the proof of Theorem 3.6. By replacing $C(s) := A^{-s} \sharp B^s$ by $C := A^{-1} \sharp B$ in the above proof, (3.21) should be $A \leq C^{-2t}$ and (3.22) should be $B = CAC \leq C^{2(1-t)}$. As $0 < s \leq \min\{1/t, 1/(1-t)\}$ for each chosen $t \in [0, 1]$, we know $ts, (1-t)s \in [0, 1]$ and thus

$$A^{(1-t)s} \leq C^{-2st(1-t)} \leq B^{-ts}.$$

So we can derive Theorem 3.7 that can compare $(B^{ts/2}A^{(1-t)s}B^{ts/2})^{1/s}$ and $A \natural_t B$ in which the range of s is specified.

Remark 3.8. We remark that the condition $0 < s \leq \min\{1/t, 1/(1-t)\}$ for each chosen t in Theorem 3.7 is more restrictive than the condition $0 < s$ in Theorem 3.4. It is easy to see $\min\{1/t, 1/(1-t)\} \leq 2$ for $t \in [0, 1]$. The following example shows that the upper bound $\min\{1/t, 1/(1-t)\}$ for s is needed. Suppose that $t = 1/2$. Then $\min\{1/t, 1/(1-t)\} = 2$. Now choose $s = 2.1$ and

$$A = \begin{bmatrix} 563.2198 & 77.6893 \\ 77.6893 & 71.7683 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 40.7285 & -25.1376 \\ -25.1376 & 44.0770 \end{bmatrix}.$$

By MATLAB computation, we have

$$(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s} = \begin{bmatrix} 135.6328 & -25.3588 \\ -25.3588 & 51.3716 \end{bmatrix} \quad \text{and} \quad A\sharp_t B = \begin{bmatrix} 139.2433 & -16.7122 \\ -16.7122 & 47.4272 \end{bmatrix},$$

where the spectrum of $(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s}$ is $\{142.6760, 44.3285\}$ and the spectrum of $A\sharp_t B$ is $\{142.1906, 44.4798\}$. Thus $(B^{ts/2} A^{(1-t)s} B^{ts/2})^{1/s} \not\prec_{\log} A\sharp_t B$.

Remark 3.9. For fixed $t \in [0, 1]$, Araki's result [4] and [3, Theorem A] asserts that

$$(B^{tp/2} A^{(1-t)p} B^{tp/2})^{1/p} \prec_{\log} (B^{tq/2} A^{(1-t)q} B^{tq/2})^{1/q}, \quad 0 < p \leq q.$$

According to (3.23), $A\sharp_t B$ is an upper bound for the set of positive definite matrices $(B^{tp/2} A^{(1-t)p} B^{tp/2})^{1/p}$ with respect to \prec_{\log} for $0 < p \leq \min\{1/t, 1/(1-t)\}$.

From (3.15) and (3.17), we have

$$e^{(1-t)A+tB} \prec_{\log} (e^{tpB/2} e^{(1-t)pA} e^{tpB/2})^{1/p} \prec_{\log} (e^{pA}\sharp_t e^{pB})^{1/p}, \quad (3.24)$$

for $A, B \in \mathbb{H}_n$, $t \in [0, 1]$ and $p > 0$. As log majorization implies weak majorization (see [2, p.42] and [27, p.168]), we have the following corollary.

Corollary 3.10. If $A, B \in \mathbb{H}_n$ and $t \in [0, 1]$, then for every $p > 0$

$$\text{tr}(e^{(1-t)A+tB}) \leq \text{tr}(e^{pA}\sharp_t e^{pB})^{1/p}.$$

Moreover, $\text{tr}(e^{pA}\sharp_t e^{pB})^{1/p}$ decreases to $\text{tr}(e^{(1-t)A+tB})$ as $p \searrow 0$.

In the above corollary, $p \searrow 0$ means that $p \in \mathbb{R}$, as a variable, decreases to 0.

4. LIMITS OF t -SPECTRAL MEAN

Hiai and Petz [13, Lemma 3.3] determined the limit of t -geometric mean when p tends to 0:

$$\lim_{p \rightarrow 0} (e^{pA}\sharp_t e^{pB})^{1/p} = e^{(1-t)A+tB}. \quad (4.1)$$

Furthermore, $\| (e^{pA}\sharp_t e^{pB})^{1/p} \|$ is increasing for any unitarily invariant norm $\| \cdot \|$, and thus $\| (e^{pA}\sharp_t e^{pB})^{1/p} \|$ increases to $\| e^{(1-t)A+tB} \|$ as $p \searrow 0$. Ando and Hiai [3] proved that

$$(e^{pA}\sharp_t e^{pB})^{1/p} \prec_{\log} e^{(1-t)A+tB}, \quad p > 0.$$

It means that $e^{(1-t)A+tB}$ is an upper bound for $(e^{pA}\sharp_t e^{pB})^{1/p}$ for all $p \geq 0$, with respect to \prec_{\log} . From Theorem 3.2, we have

$$(A\sharp_t B^p)^{1/p} \prec_{\log} (A^q\sharp_t B^q)^{1/q}, \quad 0 < q \leq p.$$

So we may write

$$(e^{pA} \sharp_t e^{pB})^{1/p} \nearrow_{\prec_{\log}} e^{(1-t)A+tB}, \quad \text{as } p \searrow 0. \quad (4.2)$$

Here the notation $\nearrow_{\prec_{\log}}$ means increasing with respect to \prec_{\log} . On the other hand, recall that from (3.24)

$$e^{(1-t)A+tB} \prec_{\log} (e^{pA} \sharp_t e^{pB})^{1/p}, \quad p > 0,$$

that is, $e^{(1-t)A+tB}$ is a lower bound of $(e^{pA} \sharp_t e^{pB})^{1/p}$ for all $p > 0$. So it would be natural to ask if formulas similar to (4.1) and (4.2) hold for t -spectral mean. The answer is affirmative and is given in the following theorem. The limit in Theorem 4.1 was proved in [1] by differentiation method. Here, we provide another proof. Given two functions $f, g : \mathbb{N} \rightarrow \mathbb{R}^+$ from the set \mathbb{N} of natural numbers to the set \mathbb{R}^+ of positive real numbers, the *little-o* notation $f(n) = o(g(n))$ means that $g(n)$ grows much faster than $f(n)$ intuitively. Rigorously, it means that for all $\varepsilon > 0$, there exists some $k \in \mathbb{N}$ such that $0 \leq f(n) < \varepsilon g(n)$ for all $n \geq k$. Thus $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$.

Theorem 4.1. If $A, B \in \mathbb{H}_n$ and $t \in [0, 1]$, then

$$\lim_{p \rightarrow 0} (e^{pA} \sharp_t e^{pB})^{1/p} = e^{(1-t)A+tB}. \quad (4.3)$$

Moreover,

$$(e^{pA} \sharp_t e^{pB})^{1/p} \searrow_{\prec_{\log}} e^{(1-t)A+tB}, \quad \text{as } p \searrow 0. \quad (4.4)$$

Proof. Suppose $p > 0$. Let $w := -p < 0$. Note that

$$\lim_{w \rightarrow 0^-} (e^{wA} \sharp_t e^{wB})^{1/w} = \lim_{p \rightarrow 0^+} ((e^{pA})^{-1} \sharp_t (e^{pB})^{-1})^{-1/p} = \lim_{p \rightarrow 0^+} (e^{pA} \sharp_t e^{pB})^{1/p},$$

where the last equality follows from $(A \sharp_t B)^{-1} = A^{-1} \sharp_t B^{-1}$. So it suffices to prove

$$e^{(1-t)A+tB} = \lim_{p \rightarrow 0^+} (e^{pA} \sharp_t e^{pB})^{1/p}.$$

Now we consider $p \rightarrow 0^+$, for $p \in (0, 1)$, write $p = (m + s)^{-1}$ and

$$X(p) = e^{pA} \sharp_t e^{pB} \quad \text{and} \quad Y(p) = e^{p[(1-t)A+tB]},$$

where $m = m(p) \in \mathbb{N}$ and $s = s(p) \in [0, 1)$. Let $\|\cdot\|$ be the spectral norm, that is, $\|X\|$ is the largest singular value of $X \in \mathbb{C}_{n \times n}$. Then

$$\|Y(p)\| \leq e^{p[(1-t)\|A\|+t\|B\|]}. \quad (4.5)$$

Since $o(p)/p \rightarrow 0$ as $p \rightarrow 0^+$, we have

$$\begin{aligned} e^{pA} \sharp_t e^{pB} &= e^{pA/2} \left\{ \left[\sum_{k=0}^{\infty} \frac{1}{k!} \left(-\frac{pA}{2} \right)^k \right] \left[\sum_{k=0}^{\infty} \frac{(pB)^k}{k!} \right] \left[\sum_{k=0}^{\infty} \frac{1}{k!} \left(-\frac{pA}{2} \right)^k \right] \right\}^t e^{pA/2} \\ &= e^{pA/2} [I + p(B - A) + o(p)]^t e^{pA/2} \\ &= \left[I + \frac{pA}{2} + o(p) \right] [I + pt(B - A) + o(p)] \left[I + \frac{pA}{2} + o(p) \right] \\ &= I + p[(1-t)A + tB] + o(p). \end{aligned}$$

Thus

$$e^{pA} \sharp_t e^{pB} = I + \frac{p}{2}(A + B) + o(p).$$

Applying it on $X(p)$ yields

$$\begin{aligned} X(p) &= (e^{-pA} \sharp_t e^{pB})^t e^{pA} (e^{-pA} \sharp_t e^{pB})^t \\ &= [I + \frac{p}{2}(-A + B) + o(p)]^t [I + pA + o(p)] [I + \frac{p}{2}(-A + B) + o(p)]^t \\ &= [I + \frac{tp}{2}(-A + B) + o(p)] [I + pA + o(p)] [I + \frac{tp}{2}(-A + B) + o(p)]. \end{aligned}$$

Since $Y(p) = e^{p[(1-t)A+tB]}$, we have

$$\|Y(p)^{1/p} - Y(p)^m\| \rightarrow 0 \quad \text{when } p \rightarrow 0^+. \quad (4.6)$$

Write $Y(p) = I + p[(1-t)A + tB] + o(p)$, so we have

$$X(p) - Y(p) = o(p).$$

By Theorem 3.4 (also see Kubo and Ando [20]), we have

$$\|e^{pA} \sharp_t e^{pB}\| \leq \|Y(p)\| \leq e^{p[(1-t)\|A\|+t\|B\|]}.$$

Then we obtain

$$\begin{aligned} \|X(p)\| &= \|(e^{-pA} \sharp_t e^{pB})^t e^{pA} (e^{-pA} \sharp_t e^{pB})^t\| \\ &\leq \|e^{-pA} \sharp_t e^{pB}\|^t \|e^{pA}\| \|e^{-pA} \sharp_t e^{pB}\|^t \\ &\leq e^{p[(1-t)\|A\|+t\|B\|]} e^{p\|A\|} e^{p[(1-t)\|A\|+t\|B\|]} \\ &= e^{p[(3-2t)\|A\|+2t\|B\|]}. \end{aligned}$$

Hence

$$\begin{aligned} \|X(p)^{1/p} - X(p)^m\| &\leq \|X(p)\|^m \|X(p)^s - I\| \\ &\leq e^{(3-2t)\|A\|+2t\|B\|} \|X(p)^s - I\| \rightarrow 0 \quad \text{when } p \rightarrow 0^+ \quad (4.7) \end{aligned}$$

because $e^{(3-2t)\|A\|+2t\|B\|}$ is bounded and $X(p) \rightarrow I$ as $p \rightarrow 0^+$, while $s \in [0, 1)$.

We also have

$$\begin{aligned} &\|X(p)^m - Y(p)^m\| \\ &\leq m \|X(p) - Y(p)\| (\max\{\|X(p)\|, \|Y(p)\|\})^{m-1} \\ &= m \|X(p) - Y(p)\| \|X(p)\|^{m-1} \\ &\leq (m+s) \|X(p) - Y(p)\| e^{p(m-1)\xi} \quad \text{where } \xi := (3-2t)\|A\| + 2t\|B\| \geq 0 \\ &= \frac{1}{p} \|X(p) - Y(p)\| e^{\frac{m-1}{m+s}\xi} \quad \text{as } p = \frac{1}{m+s} \\ &\leq \frac{1}{p} \|X(p) - Y(p)\| e^\xi \rightarrow 0 \quad \text{when } p \rightarrow 0^+ \quad (4.8) \end{aligned}$$

because $X(p) - Y(p) = o(p)$. Hence by (4.6), (4.7), and (4.8), we have

$$\begin{aligned} & \|e^{pA} \natural_t e^{pB} - e^{(1-t)A+tB}\| \\ &= \|X(p)^{1/p} - Y(p)^{1/p}\| \\ &\leq \|X(p)^{1/p} - X(p)^m\| + \|X(p)^m - Y(p)^m\| + \|Y(p)^{1/p} - Y(p)^m\| \rightarrow 0 \quad \text{as } p \rightarrow 0^+. \end{aligned}$$

The proof of (4.3) is completed.

From (3.6), we have

$$(e^{pA} \natural_t e^{pB})^{1/p} \prec_{\log} (e^{qA} \natural_t e^{qB})^{1/q}, \quad 0 < p \leq q,$$

that is, $p \rightarrow (e^{pA} \natural_t e^{pB})^{1/p}$ is a log majorization increasing function on $(0, \infty)$. Together with (3.24)

$$e^{(1-t)A+tB} \prec_{\log} (e^{pA} \natural_t e^{pB})^{1/p}, \quad p > 0,$$

(4.4) is proved. \square

Corollary 4.2. For $p > 0$, $\|\| (e^{pA} \natural_t e^{pB})^{1/p} \|\|$ decreases to $\|\| e^{(1-t)A+tB} \|\|$ as $p \searrow 0$ for any unitarily invariant norm $\|\| \cdot \|\|$.

In view of Remark 3.9 one may ask whether

$$\lim_{p \rightarrow 0} (B^{pt/2} A^{p(1-t)} B^{pt/2})^{1/p}$$

exists or not when $A, B > 0$, or equivalently,

$$\lim_{p \rightarrow 0} (e^{ptB/2} e^{p(1-t)A} e^{ptB/2})^{1/p}$$

exists or not when $A, B \in \mathbb{H}_n$. The next theorem tells us that the answer is affirmative. Note that

$$e^{ptB/2} e^{p(1-t)A} e^{ptB/2} = I + p[(1-t)A + tB] + o(p),$$

and

$$\|e^{ptB/2} e^{p(1-t)A} e^{ptB/2}\| \leq \|e^{pA} \natural_t e^{pB}\|.$$

We can derive the following theorem by an almost identical proof of Theorem 4.1 by replacing $e^{pA} \natural_t e^{pB}$ with $e^{ptB/2} e^{p(1-t)A} e^{ptB/2}$ and Remark 3.9.

Theorem 4.3. If $A, B \in \mathbb{H}_n$ and $t \in [0, 1]$, then

$$\lim_{p \rightarrow 0} (e^{ptB/2} e^{p(1-t)A} e^{ptB/2})^{1/p} = e^{(1-t)A+tB}.$$

Moreover, for $p > 0$

$$(e^{ptB/2} e^{p(1-t)A} e^{ptB/2})^{1/p} \searrow \prec_{\log} e^{(1-t)A+tB}, \quad \text{as } p \searrow 0.$$

Corollary 4.4. For $p > 0$, $\|\| (e^{ptB/2} e^{p(1-t)A} e^{ptB/2})^{1/p} \|\|$ decreases to $\|\| e^{(1-t)A+tB} \|\|$ as $p \searrow 0$ for any unitarily invariant norm $\|\| \cdot \|\|$.

Remark 4.5. Recently Audenaert and Hiai [5] considered the convergence of the sequences $\{(A^{p/2} B^p A^{p/2})^{1/p}\}_{p \in \mathbb{N}}$ and $\{(A^{\sharp} B^p)^{2/q}\}_{p \in \mathbb{N}}$, where $A, B \geq 0$. They proved that

$$\lim_{p \rightarrow \infty} (A^{p/2} B^p A^{p/2})^{1/p}$$

exists but its explicit form is not known. They also showed that

$$\lim_{p \rightarrow \infty} (A^p \sharp B^p)^{2/q}$$

exists when A, B are 2×2 but the general case is unsettled. We do not know whether the sequence $\{(e^{pA} \sharp_{\mathfrak{k}} e^{pB})^{1/p}\}_{p \in \mathbb{N}}$ converges or not and it would be interesting to know the answer.

5. KOSTANT'S PRE-ORDER AND SYMMETRIC SPACES

We first refer to [12, 18] for the notation on symmetric spaces here. Let G be a non-compact connected semisimple Lie group with Lie algebra \mathfrak{g} . Let $\Theta: G \rightarrow G$ be a Cartan involution of G , and let K be the fixed point set of Θ , which is an analytic subgroup of G . Let $\theta = d\Theta$ be the differential map of Θ . Then $\theta: \mathfrak{g} \rightarrow \mathfrak{g}$ is a Cartan involution and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ is a *Cartan decomposition*, where \mathfrak{k} is the eigenspace of θ corresponding to the eigenvalue 1 (and also the Lie algebra of K) and \mathfrak{p} is the eigenspace of θ corresponding to the eigenvalue -1 (and also an $\text{Ad } K$ -invariant subspace of \mathfrak{g} complementary to \mathfrak{k}). The Killing form B on \mathfrak{g} is negative definite on \mathfrak{k} and positive definite on \mathfrak{p} , and the bilinear form B_θ defined by

$$B_\theta(X, Y) = -B(X, \theta Y), \quad X, Y \in \mathfrak{g}$$

is an inner product on \mathfrak{g} . For each $X \in \mathfrak{g}$, let $e^X = \exp X$ be the exponential of X . Let $P = \{e^X : X \in \mathfrak{p}\}$. The map $\mathfrak{p} \times K \rightarrow G$, defined by $(X, k) \mapsto e^X k$, is a diffeomorphism. So each $g \in G$ can be uniquely written as

$$g = pk \tag{5.1}$$

with $p = p(g) \in P$ and $k = k(g) \in K$. The decomposition $G = PK$ is called a *Cartan decomposition of G* .

Let $*$: $G \rightarrow G$ be the diffeomorphism defined by $*(g) = g^* = \Theta(g^{-1})$. Because K is the fixed point set of Θ and $\exp_{\mathfrak{g}}: \mathfrak{p} \rightarrow P$ is bijective, we see that $p^* = p$ for all $p \in P$ and $k^* = k^{-1}$ for all $k \in K$. By the Cartan decomposition (5.1), we have for all $g \in G$

$$p(g) = (gg^*)^{1/2}. \tag{5.2}$$

An element $X \in \mathfrak{g}$ is called *real semisimple* (resp., *nilpotent*) if $\text{ad } X$ is diagonalizable over \mathbb{R} (resp., nilpotent). An element $g \in G$ is called *hyperbolic* (resp., *unipotent*) if $g = \exp X$ for some real semisimple (resp., nilpotent) $X \in \mathfrak{g}$; in either case X is unique and we write $X = \log g$. An element $g \in G$ is called *elliptic* if $\text{Ad } g$ is diagonalizable over \mathbb{C} with eigenvalues of modulus 1. According to [19, Proposition 2.1], each $g \in G$ can be uniquely written as

$$g = eh u, \tag{5.3}$$

where e is elliptic, h is hyperbolic, u is unipotent, and the three elements e, h and u commute. The decomposition (5.3) is called the *complete multiplicative Jordan decomposition*, abbreviated as CMJD.

The Weyl group W of $(\mathfrak{g}, \mathfrak{a})$ acts simply transitively on \mathfrak{a} (and also on A through the exponential map $\exp: \mathfrak{a} \rightarrow A$). For any real semisimple $X \in \mathfrak{g}$, let $W(X)$ denote the set

of elements in \mathfrak{a} that are conjugate to X , that is,

$$W(X) = \text{Ad } G(X) \cap \mathfrak{a}.$$

It is known from [19, Proposition 2.4] that $W(X)$ is a single W -orbit in \mathfrak{a} . Let $\text{conv } W(X)$ be the convex hull in \mathfrak{a} generated by $W(X)$. For each $g \in G$, define

$$A(g) = \exp \text{conv } W(\log h(g)),$$

where $h(g)$ is the hyperbolic component of g in its CMJD.

The Kostant's pre-order \prec_G on G is defined (see [19, p.426]) by setting \prec_G if

$$A(f) \subset A(g).$$

This pre-order induces a partial order on the conjugacy classes of G . It is known from [19, Theorem 3.1] that this pre-order \prec_G does not depend on the choice of \mathfrak{a} .

Example 5.1. If $G = \text{SL}_n(\mathbb{C})$, then $K = \text{SU}(n)$, the special unitary group and P is the space of positive definite matrices of determinant 1. See [12, p.430-431] for the CMJD of $\text{SL}_n(\mathbb{C})$, which comes from the additive Jordan decomposition. The Kostant's pre-order $\prec_{\text{SL}_n(\mathbb{C})}$ is \prec_{\log} .

Let

$$\pi : G \rightarrow G/K, \quad g \mapsto gK$$

be the natural projection. Then \mathfrak{p} may be identified with the tangent space $T_o(G/K)$ of G/K at the origin $o = eK$ via $d\pi$. Thus any $\text{Ad } K$ -invariant inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{p} induces a unique G -invariant Riemannian metric on G/K [12, p.208-209], that is, a Riemannian metric invariant under the natural action of G on G/K given by

$$(g, xK) \mapsto gxK.$$

Since G is semisimple, the Killing form B on \mathfrak{g} is nondegenerate. If B is negative definite on \mathfrak{k} and positive definite on \mathfrak{p} , then the symmetric space G/K is said to be of noncompact type.

The map $G \rightarrow P$, $g \mapsto gg^*$, is onto. Because for any $g \in G$, it maps gK to a single point gg^* , it follows that the map

$$\psi : G/K \rightarrow P, \quad gK \mapsto gg^* \tag{5.4}$$

is a bijection. It is in fact a diffeomorphism by the Cartan decomposition $G = PK$. Via ψ , P may be identified with G/K , and so may be regarded as a symmetric space of noncompact type. Note that for $p \in P$, $\psi^{-1}(p) = p^{1/2}K$, and G acts on P by

$$(g, p) \mapsto gpg^*. \tag{5.5}$$

Let $G = PK$ be the Cartan decomposition of G . The map $p \mapsto p^{1/2}K$ identifies P with G/K as a symmetric space of noncompact type. See [30, p.349-350] for the example $G = \text{SL}_n(\mathbb{R})$ (the special linear group over \mathbb{R}), $K = \text{SO}(n)$ (the special orthogonal group), P is the set of real positive definite matrices of determinant 1.

The t -geometric mean of $p, q \in P$ was defined by Liao, Liu and Tam [24]:

$$p \sharp_t q = p^{1/2} (p^{-1/2} q p^{-1/2})^t p^{1/2}, \quad 0 \leq t \leq 1. \tag{5.6}$$

It is the unique geodesic in P from p (at $t = 0$) to q (at $t = 1$). When $t = 1/2$, we abbreviate $p\#_{1/2}q$ as $p\#q$. Similarly, the t -spectral mean of $p, q \in P$ was defined by Gan, Liu, and Tam [10]:

$$p\#_tq = (p^{-1}\#q)^t p (p^{-1}\#q)^t, \quad 0 \leq t \leq 1.$$

One may interpret $p\#_tq$ as the outcome of the action of $(p^{-1}\#q)^t$ on p in view of the G -action (5.5) on P . Theorem 3.2 and Theorem 3.4 were extended in [24, Theorem 3.6 and Theorem 3.5, respectively] to P :

Theorem 5.2. (Liao, Liu, and Tam 2014) Let $p, q \in P$ and $t \in [0, 1]$. Then

$$p^r \#_t q^r \prec_G (p \#_t q)^r, \quad r \geq 1, \quad (5.7)$$

$$(p \#_t q)^r \prec_G p^r \#_t q^r, \quad 0 < r \leq 1, \quad (5.8)$$

$$(p^r \#_t q^r)^{1/r} \prec_G (p^s \#_t q^s)^{1/s}, \quad 0 < s \leq r. \quad (5.9)$$

The inequality (5.9) means that given $p, q \in P$, the function $s \mapsto (p^s \#_t q^s)^{1/s}$ is monotonic decreasing in the open interval $(0, \infty)$.

Theorem 5.3. (Liao, Liu, and Tam 2014) Let $p, q \in P$ and $t \in [0, 1]$. If $0 < r \leq 1$, we have

$$\begin{aligned} p \#_t q &\prec_G e^{(1-t) \log p + t \log q} \\ &\prec_G (q^{tr/2} p^{(1-t)r} q^{tr/2})^{1/r} \end{aligned}$$

Theorem 3.5 was extended to P [10]:

Theorem 5.4. (Gan, Liu, and Tam 2021) Let $p, q \in P$ and $t \in [0, 1]$. Then

$$p\#_tq \prec_G p\#_tq. \quad (5.10)$$

Remark 5.5. There is nice geometry hidden in (5.7), (5.8), and (5.9). For instance, let us illustrate (5.7): $p^r \#_t q^r \prec_G (p \#_t q)^r$, $r \geq 1$. From (5.6), $p^r = e\#_r p$ and $q^r = e\#_r q$, where e denotes the identity of P . So (5.7) can be rewritten as

$$(e\#_r p) \#_t (e\#_r q) \prec_G e\#_r (p \#_t q), \quad r \geq 1. \quad (5.11)$$

Consider the geodesic triangle $\Delta(e, p, q)$ determined by the three points $e, p, q \in P$, abbreviated as Δ . Recall that $e\#_\mu p$, $\mu \in [0, 1]$, is the geodesic emanating from e with end point p and clearly $e\#_r p$ is the point on the geodesic corresponding to time $\mu = r$. Similarly, $e\#_r q$ is the point on the geodesic $e\#_\mu q$, $\mu \in [0, 1]$, emanating from e with end point q , corresponding to time $\mu = r$. Now

- (1) $p^r \#_t q^r = (e\#_r p) \#_t (e\#_r q)$ is the point on the geodesic $(e\#_r p) \#_\nu (e\#_r q)$, $\nu \in [0, 1]$, emanating from $e\#_r p$ with end point $e\#_r q$, corresponding to time $\nu = t$.
- (2) $(p \#_t q)^r = e\#_r (p \#_t q)$ is the point on the geodesic $e\#_\xi (p \#_t q)$, $\xi \in [0, 1]$, emanating from e with end point $p \#_t q$, corresponding to time $\xi = r$.

See the following figures. Figure 1 illustrates that the Kostant's pre-order \prec_G relating the different points $p^r \#_t q^r$ and $(p \#_t q)^r$ in (5.7), or more precisely its equivalent form (5.11), reflects the hyperbolic geometry of P . Figure 2 depicts a hypothetical Euclidean space in which the two points $p^r \#_t q^r$ and $(p \#_t q)^r$ would be identical as the space is flat.




FIGURE 1. Hyperbolic space




FIGURE 2. Euclidean space

Using the technique in the proof of [24, Theorem 3.5], we can extend Theorems 2.3, 3.3, 3.6, 3.7, 4.1, and 4.3 in the context of symmetric spaces P of noncompact type as follows.

Theorem 5.6. Let $p, q \in P$ and $t \in [0, 1]$. Then $p \#_t q$ is P -conjugate to $(q \natural_t p)^{1/2} k_t (p \natural_t q)^{1/2}$ for some $k_t \in K$.

Theorem 5.7. For every $p, q \in P$ and $t \in [0, 1]$,

$$(p \natural_t q)^r <_G p^r \natural_t q^r, \quad r \geq 1,$$

or equivalently,

$$\begin{aligned} p^r \sharp_t q^r &\prec_G (p \sharp_t q)^r, \quad 0 < r \leq 1, \\ (p^s \sharp_t q^s)^{1/s} &\prec_G (p^r \sharp_t q^r)^{1/r}, \quad 0 < s \leq r. \end{aligned}$$

Theorem 5.8. Let $p, q \in P$. For $t \in [0, 1]$ and $s > 0$,

$$(q^{ts/2} p^{(1-t)s} q^{ts/2})^{1/s} \prec_G (p^s \sharp_t q^s)^{1/s}.$$

Moreover,

$$p \sharp_t q \prec_G e^{(1-t)\log p + t\log q} \prec_G q^{t/2} p^{(1-t)} q^{t/2} \prec_G p \sharp_t q, \quad 0 \leq t \leq 1.$$

Theorem 5.9. Let $p, q \in P$. For each chosen $t \in [0, 1]$, let $0 < s \leq \min\{1/t, 1/(1-t)\}$. We have

$$(q^{ts/2} p^{(1-t)s} q^{ts/2})^{1/s} \prec_G p \sharp_t q.$$

Theorem 5.10. For any $H, K \in \mathfrak{p}$ and $t \in [0, 1]$,

$$\lim_{p \rightarrow 0} (e^{pH} \sharp_t e^{pK})^{1/p} = e^{(1-t)H + tK}.$$

Moreover,

$$(e^{pH} \sharp_t e^{pK})^{1/p} \searrow_{\prec_G} e^{(1-t)H + tK} \quad \text{as } p \searrow 0.$$

Theorem 5.11. For any $H, K \in \mathfrak{p}$ and $t \in [0, 1]$,

$$\lim_{p \rightarrow 0} (e^{ptK/2} e^{p(1-t)H} e^{ptK/2})^{1/p} = e^{(1-t)H + tK}.$$

Moreover,

$$(e^{ptK/2} e^{p(1-t)H} e^{ptK/2})^{1/p} \searrow_{\prec_G} e^{(1-t)H + tK} \quad \text{as } p \searrow 0.$$

6. SOME REMARKS ON LÖWNER ORDER AND t -SPECTRAL MEAN

The t -geometric mean $A \sharp_t B$ may be viewed as a function $\sharp_t : \mathbb{P}_n \times \mathbb{P}_n \rightarrow \mathbb{P}_n$ for any given $t \in [0, 1]$:

$$(A, B) \mapsto A \sharp_t B = A^{1/2} (A^{-1/2} B A^{-1/2})^t A^{1/2}.$$

It is clearly a jointly continuous function. Ando and Hiai [3, p.118] mentioned that t -geometric mean is jointly (or coordinate-wisely) monotone with respect to the Löwner order.

Theorem 6.1 (Ando and Hiai 1994). Let $A, B, C, D \in \mathbb{P}_n$. If $A \geq C, B \geq D$ and $t \in [0, 1]$, then

$$A \sharp_t B \geq C \sharp_t D. \tag{6.1}$$

An example was given by Dinh and Tam [8, p.778] to show that the Löwner order in (6.1) cannot be replaced by \prec_{\log} . The following example shows that the relation similar to Theorem 6.1 is not true for the t -spectral mean.

Remark 6.2. Suppose $t = 1/3$, $A = \begin{bmatrix} 36.4987 & -34.0028 \\ -34.0028 & 39.8198 \end{bmatrix}$,

$$B_1 = \begin{bmatrix} 6.8259 & -11.0027 \\ -11.0027 & 33.6773 \end{bmatrix} \quad \text{and} \quad B_2 = \begin{bmatrix} 2.5166 & -0.2222 \\ -0.2222 & 3.4253 \end{bmatrix}.$$

It is easy to check $B_1 \geq B_2$. Then

$$A \sharp_t B_1 = \begin{bmatrix} 21.5984 & -24.0515 \\ -24.0515 & 36.6270 \end{bmatrix} \quad \text{and} \quad A \sharp_t B_2 = \begin{bmatrix} 13.4040 & -10.9429 \\ -10.9429 & 15.7328 \end{bmatrix},$$

where the eigenvalues of $A \sharp_t B_1 - A \sharp_t B_2$ are -0.0213 and 29.1098 . Thus $A \sharp_t B_1 - A \sharp_t B_2$ is not positive semidefinite.

Remark 6.3. In 2012, Lim [25] named the operator monotone property (6.1) as Löwner-Heinz inequality. The readers should be alerted that in the literature Löwner-Heinz inequality is sometimes used (for example [31, p.2-3]) for the profound result of Löwner:

$$A \geq B \geq 0 \Rightarrow A^r \geq B^r, \quad 0 \leq r \leq 1. \quad (6.2)$$

Löwner [26] first obtained the result in his 1934 seminal paper and Heinz [11] gave an alternative proof in 1951. Since then, various proofs of Löwner-Heinz inequality have been given by different authors. See the historical notes of Bhatia [6, p.149-150]. That being said, it is true that (6.2) follows from (6.1): set $A = C = I$, (6.1) becomes $B^t = I \sharp_t B \geq I \sharp_t D = D^t$ when $B \geq D$ and $0 \leq t \leq 1$. In other words, Löwner-Heinz inequality can be written in the context of r -geometric mean:

$$A \geq B \geq 0 \Rightarrow I \sharp_r A \geq I \sharp_r B, \quad 0 \leq r \leq 1.$$

Remark 6.4. It is known that [6, Theorem IX.2.6]

$$A, B \geq 0 \Rightarrow \lambda_1(A^s B^s) \leq \lambda_1(AB)^s, \quad 0 \leq s \leq 1. \quad (6.3)$$

Both AB and $A^s B^s$ are diagonalizable with nonnegative eigenvalues so log majorization applies to them according to Remark 3.1. Applying the compound matrix argument on (6.3), we have

$$A, B \geq 0 \Rightarrow A^s B^s \prec_{\log} (AB)^s, \quad 0 \leq s \leq 1. \quad (6.4)$$

Since $A^s B^s$ is similar to $A^{s/2} B^s A^{s/2}$, (6.4) can be rewritten as

$$A, B \geq 0 \Rightarrow A^{s/2} B^s A^{s/2} \prec_{\log} (A^{1/2} B^{1/2} A^{1/2})^s, \quad 0 \leq s \leq 1. \quad (6.5)$$

Let us confine ourselves in \mathbb{P}_n . The advantage of the form in (6.5) is that all elements are now in \mathbb{P}_n . The group $\text{GL}_n(\mathbb{C})$ acts on $\mathbb{P}_n \cong \text{GL}_n(\mathbb{C})/\text{U}(n)$ via (5.5). So (6.5) can be interpreted in the context of group action on \mathbb{P}_n and the s -geometric mean since

$$(A^{s/2}, B^s) \mapsto A^{s/2} B^s A^{s/2} \text{ (group action)}$$

and

$$B^s = I \sharp_s B, \quad (A^{1/2} B^{1/2} A^{1/2})^s = I \sharp_s A^{1/2} B^{1/2} A^{1/2}.$$

Motivated by Figure 1, the interested readers may draw a picture to visualize the geometry associated with (6.5).

Acknowledgement We are thankful to the anonymous referee for carefully reading of our paper and for giving constructive suggestions that helped us to greatly improve the paper.

REFERENCES

- [1] E. Ahn, S. Kim, and Y. Lim. An extended Lie-Trotter formula and its applications. *Linear Algebra Appl.*, 427(2-3):190–196, 2007.
- [2] T. Ando. Majorizations and inequalities in matrix theory. *Linear Algebra Appl.*, 199:17–67, 1994.
- [3] T. Ando and F. Hiai. Log majorization and complementary Golden-Thompson type inequalities. *Linear Algebra Appl.*, 197/198:113–131, 1994. Second Conference of the International Linear Algebra Society (ILAS) (Lisbon, 1992).
- [4] H. Araki. On an inequality of Lieb and Thirring. *Lett. Math. Phys.*, 19(2):167–170, 1990.
- [5] K. M. R. Audenaert and F. Hiai. Reciprocal Lie-Trotter formula. *Linear Multilinear Algebra*, 64(6):1220–1235, 2016.
- [6] R. Bhatia. *Matrix analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.
- [7] R. Bhatia. *Positive definite matrices*. Princeton Series in Applied Mathematics. Princeton University Press, Princeton, NJ, 2007.
- [8] T. H. Dinh, S. Ahsani, and T.-Y. Tam. Geometry and inequalities of geometric mean. *Czechoslovak Math. J.*, 66(141)(3):777–792, 2016.
- [9] M. Fiedler and V. Pták. A new positive definite geometric mean of two positive definite matrices. *Linear Algebra Appl.*, 251:1–20, 1997.
- [10] L. Gan, X. Liu, and T.-Y. Tam. On two geometric means and sum of adjoint orbits. *Linear Algebra Appl.*, 631:156–173, 2021.
- [11] E. Heinz. Beiträge zur Störungstheorie der Spektralzerlegung. *Math. Ann.*, 123:415–438, 1951.
- [12] S. Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.
- [13] F. Hiai and D. Petz. The Golden-Thompson trace inequality is complemented. *Linear Algebra Appl.*, 181:153–185, 1993.
- [14] A. Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *Amer. J. Math.*, 76:620–630, 1954.
- [15] H. Kim and Y. Lim. An extended matrix exponential formula. *J. Math. Inequal.*, 1(3):443–447, 2007.
- [16] S. Kim. Operator inequalities and gyrolines of the weighted geometric means. *Math. Inequal. Appl.*, 24(2):491–514, 2021.
- [17] S. Kim and H. Lee. Relative operator entropy related with the spectral geometric mean. *Anal. Math. Phys.*, 5(3):233–240, 2015.
- [18] A. W. Knap. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, second edition, 2002.
- [19] B. Kostant. On convexity, the Weyl group and the Iwasawa decomposition. *Ann. Sci. École Norm. Sup. (4)*, 6:413–455 (1974), 1973.
- [20] F. Kubo and T. Ando. Means of positive linear operators. *Math. Ann.*, 246(3):205–224, 1979/80.
- [21] J. Lawson and Y. Lim. Means on dyadic symmetric sets and polar decompositions. *Abh. Math. Sem. Univ. Hamburg*, 74:135–150, 2004.
- [22] H. Lee and Y. Lim. Metric and spectral geometric means on symmetric cones. *Kyungpook Math. J.*, 47(1):133–150, 2007.
- [23] L. Li, L. Molnár, and L. Wang. On preservers related to the spectral geometric mean. *Linear Algebra Appl.*, 610:647–672, 2021.
- [24] M. Liao, X. Liu, and T.-Y. Tam. A geometric mean for symmetric spaces of noncompact type. *J. Lie Theory*, 24(3):725–736, 2014.

- [25] Y. Lim. Factorizations and geometric means of positive definite matrices. *Linear Algebra Appl.*, 437(9):2159–2172, 2012.
- [26] K. Löwner. Über monotone Matrixfunktionen. *Math. Z.*, 38(1):177–216, 1934.
- [27] A. W. Marshall, I. Olkin, and B. C. Arnold. *Inequalities: theory of majorization and its applications*. Springer Series in Statistics. Springer, New York, second edition, 2011.
- [28] W. Pusz and S. L. Woronowicz. Functional calculus for sesquilinear forms and the purification map. *Rep. Mathematical Phys.*, 8(2):159–170, 1975.
- [29] R. Rado. An inequality. *J. London Math. Soc.*, 27:1–6, 1952.
- [30] A. Terras. *Harmonic analysis on symmetric spaces—higher rank spaces, positive definite matrix space and generalizations*. Springer, New York, second edition, 2016.
- [31] X. Zhan. *Matrix inequalities*, volume 1790 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEVADA, RENO, RENO, NV
89557-0084, USA

Email address: `lgan@unr.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEVADA, RENO, RENO, NV
89557-0084, USA

Email address: `ttam@unr.edu`