

TRANSVERSE SPECTRAL INSTABILITY IN GENERALIZED KADOMTSEV-PETVIASHVILI EQUATION

BHAVNA, ATUL KUMAR, AND ASHISH KUMAR PANDEY

ABSTRACT. We study transverse stability and instability of one-dimensional small-amplitude periodic traveling waves of a generalized Kadomtsev-Petviashvili equation. We show transverse instability in the KP-II equation with respect to two-dimensional perturbations, which are periodic or square-integrable in the direction of the propagation of the underlying one-dimensional wave and of finite or short wavelength in the transverse direction. We also obtain transverse instability results in KP-BO, KP-ILW, and KP-Whitham equations. Moreover, assuming the spectral stability of one-dimensional wave with respect to one-dimensional square-integrable periodic perturbations, we obtain transverse stability results in KP-fKdV, KP-ILW, and KP-Whitham equations.

1. INTRODUCTION

We propose the generalized Kadomtsev-Petviashvili (gKP) equation

$$(1.1) \quad (u_t + \mathcal{M}u_x - uu_x)_x + \sigma u_{yy} = 0,$$

in which $u(x, y, t)$ depends upon the spatial variables $x, y \in \mathbb{R}$, and the temporal variable $t \in \mathbb{R}$, \mathcal{M} is a multiplier operator given by the symbol $m(k)$ as

$$(1.2) \quad \widehat{\mathcal{M}f}(k) = m(k)\widehat{f}(k),$$

and σ is equal to either 1 or -1 . We make following assumptions on $m(k)$.

Hypothesis 1.1. *The multiplier symbol $m(k)$ in (1.2) should satisfy the following.*

H1. m is real valued, even and without loss of generality, $m(0) = 1$,

H2. $C_1 k^\alpha \leq m(k) \leq C_2 k^\alpha$, $k \gg 1$, $\alpha \geq -1$ and for some $C_1, C_2 > 0$,

H3. m is strictly monotonic for $k > 0$.

Hypotheses H1 and H2 are essential for the proof of the existence of periodic traveling waves, which we discuss later in this section, while H3 is required for stability analysis done in Sections 3 and 4.

Models. The gKP equation (1.1) is a generalization to the Kadomtsev-Petviashvili (KP) equation [KP70]

$$(1.3) \quad (u_t - u_{xxx} - uu_x)_x + \sigma u_{yy} = 0,$$

where $\mathcal{M} = -\partial_x^2$. The KP equation is a natural extension to two spatial dimensions of the well-known Korteweg-de Vries (KdV) equation

$$(1.4) \quad u_t - u_{xxx} - uu_x = 0.$$

Date: April 9, 2024.

The KP equation (1.3) with $\sigma = -1$ (negative dispersion) is called the KP-II equation, whereas the one with $\sigma = 1$ (positive dispersion) is called the KP-I equation. Along the same lines, the gKP equation can be thought of as an extension to two spatial dimensions of the equation

$$(1.5) \quad u_t + \mathcal{M}u_x - uu_x = 0.$$

The gKP equation in the form (1.1) first appears in [Sau95] where the existence and properties of its localized solitary waves were studied. For different values of \mathcal{M} or $m(k)$, (1.5) reduces to various well-known equations like $m(k) = 1 + |k|^\beta$ ($\beta > 1$) is the Fractional KdV (fKdV) equation, $m(k) = 1 + |k|$ is the Benjamin-Ono (BO) equation, $m(k) = k \coth k$ is the Intermediate Long wave (ILW) equation and $m(k) = \sqrt{\tanh k/k}$ is the Whitham equation. For $m(k) = 1 + |k|^\beta$, $1 + |k|$, $k \coth k$, and $\sqrt{\tanh k/k}$, we name the gKP equation as KP-fKdV, KP-BO, KP-ILW, and KP-Whitham, respectively. We add -I or -II to the name if $\sigma = 1$ or -1 , respectively.

Dispersion relation. Assuming a plane-wave solution of the form

$$u(x, t) = e^{i(kx - \omega t + \ell y)}$$

for the linear part

$$(u_t + \mathcal{M}u_x)_x + \sigma u_{yy} = 0$$

of the gKP equation (1.1), we arrive at the dispersion relation given by the phase velocity

$$(1.6) \quad v_p(k) = \frac{\omega}{k} = m(k) + \sigma \frac{\ell^2}{k^2}.$$

From (1.6), we observe that phase velocity v_p is monotonic if $\sigma = 1$ and m is decreasing or $\sigma = -1$ and m is increasing, while in other two combinations it changes its behavior along local extrema.

Small amplitude periodic traveling waves. Seeking y -independent traveling wave solution of (1.1) of the form $u(x, y, t) = U(x - ct)$, where $c \in \mathbb{R}$ is the speed of propagation, then U satisfies the following

$$(-cU' + \mathcal{M}U' - UU')' = 0.$$

Integrating this, we get

$$\mathcal{M}U = cU + \frac{U^2}{2} + b + dx,$$

in which b and d are arbitrary constants. Since we are interested in periodic solutions, we can set $d = 0$ and the equation becomes

$$(1.7) \quad \mathcal{M}U = cU + \frac{U^2}{2} + b.$$

Let U be a $2\pi/k$ -periodic function in x . Then, $w(z) := U(x)$ with $z := kx$ is a 2π -periodic function in z . For each $k > 0$, a family of small amplitude 2π -periodic and smooth solutions

$w(k, a, b)(z)$ exists at $c = c(k, a, b)$, see [HJ15, Proposition 2.2] for more details. Moreover,

$$(1.8) \quad \begin{cases} w(k, a, b)(z) = (1 - m(k))b + a \cos z + a^2(A_0 + A_2 \cos 2z) + a^3 A_3 \cos 3z + O(a^4 + ab + b^2), \\ c(k, a, b) = m(k) - (1 - m(k))b + a^2 c_2 + O(a^4 + ab + b^2), \end{cases}$$

where

$$(1.9) \quad A_0 = \frac{1}{4(1 - m(k))}, \quad A_2 = \frac{1}{4(m(2k) - m(k))}, \quad A_3 = \frac{A_2}{2(m(3k) - m(k))} \text{ and } c_2 = -A_0 - \frac{A_2}{2}.$$

Hypotheses 1.1 H1 and H2 are used to prove the existence of $w(k, a, b)(z)$ and $c(k, a, b)$, see [HJ15, Proposition 2.2] for the proof. Starting now, we denote $w(k, a, b)$ and $c(k, a, b)$ as w and c respectively.

Transverse instability. Clearly, a solution of (1.5) is a y -independent solution of the gKP equation (1.1). The stability (or instability) of such a one-dimensional solution of (1.1) with respect to perturbations which are two-dimensional is generally termed as *transverse stability (or instability)*.^{*} The transverse instability of solitary waves of the KdV in the KP equation was first conducted by Kadomtsev, and Petviashvili [KP70], where it was found that such solutions are stable to transverse perturbations in the case of negative dispersion ($\sigma = -1$), while they are unstable to long-wavelength transverse perturbations in the case of positive dispersion ($\sigma = 1$) even though they are stable in the corresponding one-dimensional problem. Johnson and Zumbrun [JZ10] have studied transverse instability of periodic waves for the KP-gKdV equation, with respect to periodic perturbations in the direction of propagation and of long wavelength in the transverse direction. They have constructed an orientation index by comparing the low and high-frequency behavior of the periodic Evans functions. Mariana Haragus [Har11] also studied the transverse stability of KP-KdV equations, but the author restricted to the case of small periodic waves and considered transverse stability for more general perturbations for the KP-KdV equation. Transverse instability of periodic waves of KP-I and Schrödinger equations have been studied in [HSS12]. Transverse instability of solitary wave solutions of various water-wave models have also been explored by several authors, see [GHS01, PS04, RT09, RT11].

In this article, we study transverse spectral stability of y -independent solution $u(x, y, t) = w(k(x - ct))$, where w and c are given in (1.8), of (1.1) with respect to two-dimensional perturbations which are either periodic or non-periodic in the x -direction and always periodic in the y -direction. If the perturbation is periodic in the x -direction then it is co-periodic with the solution. If the perturbation is non-periodic in the x -direction then it is square-integrable on the whole real line. The periodic nature of the perturbation in the y -direction is classified into two categories: short or finite wavelength and long-wavelength perturbations. Table 1 represents results obtained in this article for different combinations of nature of the two-dimensional perturbation in x - and y -directions. Notably, we have obtained transverse

^{*}The definition of transverse stability can be different in different articles depending on what is the nature of underlying stability analysis, for example, orbital or spectral stability.

instability results for small-amplitude periodic traveling waves of KP-II, KP-BO-I, KP-BO-II, KP-ILW-I, KP-ILW-II, KP-Whitham-I, and KP-Whitham-II equations. Also, we have obtained transverse stability results for these models conditioned on the spectral stability of periodic traveling waves with respect to one-dimensional perturbations.

In Section 2, we linearize the equation and formulate the problem. In Sections 3 and 4, we provide transverse instability analysis for different cases mentioned in Table 1. We discuss our main results and applications of these results for KP-fKdV, KP-BO, KP-ILW, and KP-Whitham equations in Section 5.

		x -direction	
y -direction		Periodic	Non-periodic
	Short or Finite wavelength	$\sigma = 1$ and m decreasing $\sigma = -1$ and m increasing	For all combinations of σ and m
	Long wavelength	$\sigma = 1$ and m increasing $\sigma = -1$ and m decreasing	For no combinations of σ and m

TABLE 1. The combination of value of σ and monotonic nature of m which provide transverse instability for given nature of perturbations in x - and y -directions.

Notations. The following notations are going to be used throughout the article. Here, $L^2(\mathbb{R})$ denotes the set of real or complex-valued, Lebesgue measurable functions over \mathbb{R} such that

$$\|f\|_{L^2(\mathbb{R})} = \left(\int_{\mathbb{R}} |f(x)|^2 dx \right)^{1/2} < +\infty$$

and, $L^2(\mathbb{T})$ denote the space of 2π -periodic, measurable, real or complex-valued functions over \mathbb{R} such that

$$\|f\|_{L^2(\mathbb{T})} = \left(\frac{1}{2\pi} \int_0^{2\pi} |f(x)|^2 dx \right)^{1/2} < +\infty.$$

The space $C_b(\mathbb{R})$ consists of all bounded continuous functions on \mathbb{R} , normed with

$$\|f\| = \sup_{x \in \mathbb{R}} |f(x)|.$$

For $s \in \mathbb{R}$, let $H^s(\mathbb{R})$ consists of tempered distributions such that

$$\|f\|_{H^s(\mathbb{R})} = \left(\int_{\mathbb{R}} (1 + |t|^2)^s |\hat{f}(t)|^2 dt \right)^{\frac{1}{2}} < +\infty$$

and

$$H^s(\mathbb{T}) = \{f \in H^s(\mathbb{R}) : f \text{ is } 2\pi\text{-periodic}\}.$$

We define $L^2(\mathbb{T})$ -inner product as

$$(1.10) \quad \langle f, g \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(z) \overline{g(z)} dz = \sum_{n \in \mathbb{Z}} \hat{f}_n \overline{\hat{g}_n},$$

where \hat{f}_n are Fourier coefficients of the function f defined by

$$\hat{f}_n = \frac{1}{2\pi} \int_0^{2\pi} f(z) e^{inz} dz.$$

The value of σ and monotonic nature of the symbol m in (1.2) appear numerous times, and we use the shorthand notation in Table 2 for them.

		m	
		Increasing	Decreasing
σ	1	$(1, \uparrow)$	$(1, \downarrow)$
	-1	$(-1, \uparrow)$	$(-1, \downarrow)$

TABLE 2. Notations for different values of σ and monotonic nature of m .

2. LINEARIZATION

Linearizing the gKP equation (1.1) about its one-dimensional periodic traveling wave w in (1.8) and using change of variables, abusing notation, $x \rightarrow kz$, $t \rightarrow k^2t$, and $y \rightarrow y$, we arrive at

$$(2.1) \quad v_{tz} - cv_{zz} + \mathcal{M}_k v_{zz} - (wv)_{zz} + \sigma v_{yy} = 0,$$

For $v(z, y, t) = e^{\lambda t + i\ell y} V(z)$, we obtain

$$\lambda V_z - cV_{zz} + \mathcal{M}_k V_{zz} - (wV)_{zz} - \sigma \ell^2 V = 0.$$

which can be rewritten as

$$(2.2) \quad \mathcal{T}_a(\lambda, \ell)V := (\lambda \partial_z - \partial_z^2(c - \mathcal{M}_k + w) - \sigma \ell^2)V = 0$$

We assume that $2\pi/k$ -periodic traveling wave solution $u(x, y, t) = w(k(x - ct))$ of (1.1) is a stable solution of the one-dimensional equation (1.5) where w and c are as in (1.8). We then say that the periodic wave w in (1.8) is transversely spectrally stable with respect to two-dimensional periodic perturbations (resp. non-periodic (localized or bounded perturbations)) if the gKP operator $\mathcal{T}_a(\lambda, \ell)$ acting in $L^2(\mathbb{T})$ (resp. $L^2(\mathbb{R})$ or $C_b(\mathbb{R})$) with domain $H^{\alpha+2}(\mathbb{T})$ (resp. $H^{\alpha+2}(\mathbb{R})$ or $C_b^{\alpha+2}(\mathbb{R})$), where α is in Hypothesis 1.1 H2, is invertible, for any $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$ and any $\ell \neq 0$.

Depending on the space in which we are studying the invertibility of $\mathcal{T}_a(\lambda, \ell)$, we split our study into periodic ($L^2(\mathbb{T})$) and non-periodic perturbations ($L^2(\mathbb{R})$ or $C_b(\mathbb{R})$). Also, depending upon the values of ℓ we distinguish two different regimes: long-wavelength transverse perturbations, when $|\ell| \ll 1$ and short or finite wavelength transverse perturbations, otherwise.

3. PERIODIC PERTURBATIONS

In this section, we study transverse stability with respect to two-dimensional perturbations, which are co-periodic in the direction of the propagation of the wave. Therefore, we check if the operator $\mathcal{T}_a(\lambda, \ell)$ acting in $L^2(\mathbb{T})$ is invertible, for any $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$ and any $\ell \neq 0$. We reformulate the invertibility problem for this particular case.

Proposition 3.1. *The following statements are equivalent:*

- (1) $\mathcal{T}_a(\lambda, \ell)$ acting in $L^2(\mathbb{T})$ with domain $H^{\alpha+2}(\mathbb{T})$ is not invertible.
- (2) The restriction of $\mathcal{T}_a(\lambda, \ell)$ to the subspace $L_0^2(\mathbb{T})$ of $L^2(\mathbb{T})$ is not invertible, where

$$L_0^2(\mathbb{T}) = \left\{ f \in L^2(\mathbb{T}) : \int_0^{2\pi} f(z) dz = 0 \right\}.$$

- (3) λ belongs to the spectrum of the operator $\mathcal{A}_a(\ell)$ acting in $L_0^2(\mathbb{T})$ with domain $H^{\alpha+1}(\mathbb{T}) \cap L_0^2(\mathbb{T})$, where

$$\mathcal{A}_a(\ell) = \partial_z(c - \mathcal{M}_k + w) + \sigma\ell^2\partial_z^{-1}$$

We refer to [Har11, Lemma 4.1, Corollary 4.2] for a detailed proof in a similar situation. Proposition 3.1 reduces the invertibility problem of $\mathcal{T}_a(\lambda, \ell)$ to the study of the spectrum of $\mathcal{A}_a(\ell)$ acting on $L_0^2(\mathbb{T})$ with domain $H^{\alpha+1}(\mathbb{T}) \cap L_0^2(\mathbb{T})$. The operator $\mathcal{A}_a(\ell)$ acting on $L_0^2(\mathbb{T})$ has a compact resolvent so that its spectrum consists of isolated eigenvalues with finite multiplicity. In addition, the spectrum of $\mathcal{A}_a(\ell)$ is symmetric with respect to both the real and imaginary axes.

A straightforward calculation reveals that

$$(3.1) \quad \mathcal{A}_0(\ell)e^{inz} = i\omega_{n,\ell}e^{inz} \quad \text{for all } n \in \mathbb{Z} \setminus \{0\}$$

where

$$(3.2) \quad \omega_{n,\ell} = n(m(k) - m(kn)) - \frac{\sigma\ell^2}{n}.$$

Consequently, $L_0^2(\mathbb{T})$ -spectrum of $\mathcal{A}_0(\ell)$ consists of purely imaginary eigenvalues of finite multiplicity. Since

$$\|\mathcal{A}_a(\ell) - \mathcal{A}_0(\ell)\| = O(|a|)$$

as $a \rightarrow 0$ uniformly in the operator norm. A standard perturbation argument then guarantees the spectrum of $\mathcal{A}_a(\ell)$ and $\mathcal{A}_0(\ell)$ will stay close for $|a|$ small. Recalling that the spectrum of $\mathcal{A}_a(\ell)$ is symmetric with respect to the imaginary axis, it follows then that for $|a|$ small when eigenvalues of $\mathcal{A}_a(\ell)$ bifurcate from the imaginary axis they must bifurcate in pairs resulting from collisions of eigenvalues of $\mathcal{A}_0(\ell)$ on the imaginary axis.

The linear operator $\mathcal{A}_a(\ell)$ can be decomposed as

$$\mathcal{A}_a(\ell) = J\mathcal{L}_a(\ell)$$

where

$$J = \partial_z \quad \text{and} \quad \mathcal{L}_a(\ell) = c - \mathcal{M}_k + w + \sigma\ell^2\partial_z^{-2}$$

The operator J is skew-adjoint whereas the operator $\mathcal{L}_a(\ell)$ is self-adjoint. The *Krein signature* κ_n of an eigenvalue $i\omega_{n,\ell}$ of $\mathcal{A}_0(\ell)$ is defined as

$$(3.3) \quad \kappa_n = \text{sgn}(\langle \mathcal{L}_0(\ell)e^{inz}, e^{inz} \rangle) = \text{sgn} \left(m(k) - m(kn) - \frac{\sigma \ell^2}{n^2} \right), \quad n \in \mathbb{Z} \setminus \{0\}$$

where sgn is the signum function which determines the sign of a real number. A pair of eigenvalues leave imaginary axis after collision only if their Krein signatures κ_n are opposite. We have the following lemma.

Lemma 3.2. *For any $|a|$ sufficiently small, there exists a $\ell_a > 0$ such that for all $|\ell| > \ell_a$, the spectrum of $\mathcal{A}_a(\ell)$ is purely imaginary if (σ, m) is $(1, \uparrow)$ or $(-1, \downarrow)$, where these notations are explained in Table 2.*

Proof. For $(\sigma, m) = (1, \uparrow)$, κ_n is negative for all $n \in \mathbb{Z} \setminus \{0\}$ and for $(\sigma, m) = (-1, \downarrow)$, κ_n is positive for all $n \in \mathbb{Z} \setminus \{0\}$ for all $k > 0$. Therefore, Krein signatures of all eigenvalues remain same in both cases implying that eigenvalues will not bifurcate from the imaginary axis even if there is a collision away from the origin for $|a|$ sufficiently small. The collision at the origin may possibly lead to bifurcation away from the imaginary axis for sufficiently small ℓ (in fact, this is actually the case, see Lemma 3.5). Therefore, there exists an ℓ_a depending on a such that for all $|\ell| > \ell_a$, the spectrum of $\mathcal{A}_a(\ell)$ is purely imaginary. \square

It follows from Lemma 3.2 that the only collision that may lead to instability for $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$ is the collision at the origin between $\omega_{1,0}$ and $\omega_{-1,0}$. Since this collision takes place at $\ell = 0$, the perturbation analysis will take place in the regime $|\ell| \ll 1$. In other words, the underlying transverse perturbations are of long wavelength. The other regime is of finite and short-wavelength perturbations. We split our further analysis into these two regimes.

3.1. Finite and short-wavelength transverse perturbations. We start the analysis of the spectrum of $\mathcal{A}_a(\ell)$ with the values of ℓ away from the origin, $|\ell| \geq \ell_0$, for some $\ell_0 > 0$, i.e., finite and short wavelength transverse perturbations. Using Lemma 3.2, there are no collisions of eigenvalues that may lead to instability for $\ell > 0$ if $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$. Therefore, we restrict our attention to two other cases, $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$.

Let eigenvalues $i\omega_{p,\ell}$ and $i\omega_{q,\ell}$, $p \neq q$, collide at $\ell = \ell_c > 0$. From (3.3), Krein signatures κ_p and κ_q are opposite at $\ell = \ell_c$ when $pq < 0$, i.e. p and q should be of opposite parity. A direct calculation shows that if $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$ then $i\omega_{p,\ell}$ and $i\omega_{-q,\ell}$ collide when

$$\ell^2 = \ell_{p,q}^2 = \frac{\sigma pq}{p+q} (p(m(k) - m(kp)) + q(m(k) - m(kq))) > 0.$$

for all $p, q \in \mathbb{N}$ except $(p, -q) = (1, -1)$.

Let Δ denotes the distance between indices p and $-q$ of colliding eigenvalues. For $\Delta = 1$ and 2 there are no pairs of eigenvalues which can lead to instability. For $\Delta = 3$, there are two such pairs of colliding eigenvalues, $\{\omega_{-1,\ell}, \omega_{2,\ell}\}$ and $\{\omega_{1,\ell}, \omega_{-2,\ell}\}$ which can lead to instability. In the following lemma, we prove that both these pairs lead to instability.

Lemma 3.3. *Let $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$ and $\ell_c^2 = \frac{4\sigma}{3}(m(k) - m(2k))$. For any a sufficiently small, there exists $\varepsilon_a = -\sigma a^2 A_2 + \frac{2\sqrt{2}}{3}|a|^3 A_3 + O(a^4) > 0$ such that the spectrum of $\mathcal{A}_a(\ell)$ have a pair of complex eigenvalues with opposite nonzero real parts for all ℓ satisfying $|\ell^2 - \ell_c^2| < \varepsilon_a$.*

Proof. We will prove the lemma by analyzing the collisions for $\Delta = 3$. Let for some $n \in \mathbb{Z}$, we have

$$(3.4) \quad 0 \neq \omega_{n, \ell_c} = \omega_{n+3, \ell_c} = \omega \text{ (say)}$$

for some $\ell^2 = \ell_c^2 > 0$. Therefore, $i\omega$ is an eigenvalue of $\mathcal{A}_0(\ell_c)$ of multiplicity two with an orthonormal basis of eigenfunctions $\{e^{inz}, e^{i(n+3)z}\}$. For $|\ell - \ell_c|$ and $|a|$ sufficiently small, let $\lambda_{n, a, \ell}$ and $\lambda_{n+3, a, \ell}$ be eigenvalues of $\mathcal{A}_a(\ell)$ bifurcating from $i\omega$ with an orthonormal basis of eigenfunctions $\{\phi_{n, a, \ell}(z), \phi_{n+3, a, \ell}(z)\}$. Note that $\lambda_{n, 0, \ell_c} = \lambda_{n+3, 0, \ell_c} = i\omega$ with $\phi_{n, 0, \ell_c}(z) = e^{inz}$ and $\phi_{n+3, 0, \ell_c}(z) = e^{i(n+3)z}$. Let

$$(3.5) \quad \lambda_{n, a, \ell} = i\omega + i\mu_{n, a, \ell} \quad \text{and} \quad \lambda_{n+3, a, \ell} = i\omega + i\mu_{n+3, a, \ell}$$

We are interested in the location of $\mu_{n, a, \ell}$ and $\mu_{n+3, a, \ell}$ for $|\ell - \ell_c|$ and $|a|$ sufficiently small.

We start with the following expansions of eigenfunctions [CDT21]

$$(3.6) \quad \phi_{a, n, \ell}(z) = e^{inz} + a\phi_{n, 1} + a^2\phi_{n, 2} + a^3\phi_{n, 3} + O(a^4),$$

$$(3.7) \quad \phi_{a, n+3, \ell}(z) = e^{i(n+3)z} + a\phi_{n+3, 1} + a^2\phi_{n+3, 2} + a^3\phi_{n+3, 3} + O(a^4).$$

We use orthonormality of $\phi_{a, n, \ell}(z)$ and $\phi_{a, n+3, \ell}(z)$ to find that

$$\phi_{n, 1} = \phi_{n, 2} = \phi_{n, 3} = \phi_{n+3, 1} = \phi_{n+3, 2} = \phi_{n+3, 3} = 0.$$

Using expansions of w and c in (1.8), we expand $\mathcal{A}_a(\ell)$ in a as


$$(3.8) \quad \mathcal{A}_a(\ell) = \mathcal{A}_0(\ell) + a\partial_z(\cos z) + a^2\partial_z\left(-\frac{A_2}{2} + A_2 \cos 2z\right) + a^3\partial_z(A_3 \cos 3z) + O(a^4)$$

Now, to trace the bifurcation of the eigenvalues from the point of the collision on the imaginary axis for $|\ell - \ell_c|$ and $|a|$ sufficiently small, we compute the action of $\mathcal{A}_a(\ell)$ and identity operators on the extended eigenspace $\{\phi_{a, n, \ell}(z), \phi_{a, n+3, \ell}(z)\}$ and arrive at

$$B_a(\ell) = \begin{pmatrix} i\omega - ia^2\frac{A_2}{2}n - \frac{i\sigma\varepsilon}{n} & ia^3\frac{A_3}{2}(n+3) \\ ia^3\frac{A_3}{2}n & i\omega - ia^2\frac{A_2}{2}(n+3) - \frac{i\sigma\varepsilon}{n+3} \end{pmatrix} + O(a^4).$$

where $\varepsilon = \ell^2 - \ell_c^2$ and \mathbf{I}_a , 2×2 identity matrix, respectively. To locate μ , we compute

$$(3.9) \quad \det(\mathcal{B}_a(\ell) - (i\omega + i\mu)\mathcal{I}_a) = 0$$

FIGURE 1. Collision of eigenvalues at $\ell = \ell_c$ for $k = 1$.

and arrive at a quadratic in μ

$$\begin{aligned} \mu^2 + \mu \left(\sigma \varepsilon \left(\frac{1}{n} + \frac{1}{n+3} \right) + \frac{a^2 A_2}{2} ((n+3) + n) \right) + \frac{a^4 A_2^2 n(n+3)}{4} \\ + \frac{a^2 A_2 \sigma \varepsilon}{2} \left(\frac{n}{n+3} + \frac{n+3}{n} \right) + \frac{\sigma^2 \varepsilon^2}{n(n+3)} - \frac{a^6 A_3^2 n(n+3)}{4} + O(a^8) = 0. \end{aligned}$$

A direct computation shows that the discriminant of the above quadratic is

$$\text{disc}_a(\varepsilon) = \left(\frac{-3\varepsilon}{n(n+3)} + \frac{3\sigma a^2 A_2}{2} \right)^2 + a^6 A_3^2 n(n+3) + O(a^8)$$

The two collisions corresponding to $\Delta = 3$ corresponds to $n = -2$ and $n = -1$. For both these choices

$$\ell_c^2 = \ell_{1,2}^2 = \ell_{2,1}^2 = \frac{4\sigma}{3}(m(k) - m(2k)).$$

For any a sufficiently small if we choose ℓ such that

$$|\ell^2 - \ell_c^2| < -\sigma a^2 A_2 + \frac{2\sqrt{2}}{3}|a|^3 A_3 + O(a^4)$$

then $\text{disc}_a(\varepsilon) < 0$ and it leads to μ to have non-zero imaginary part and therefore, $\mathcal{A}_a(\ell)$ has two eigenvalues with non-zero real parts, hence the lemma. \square

Figure 1 depicts a collision of pair of eigenvalues in KP-II and KP-Whitham-I equations which lead to instability according to Lemma 3.3.

3.2. Long wavelength transverse perturbations. In all four cases $(\sigma, m) = (1, \uparrow), (1, \downarrow), (-1, \uparrow),$ and $(-1, \downarrow)$, there is a collision at the origin of eigenvalues $i\omega_{-1,\ell}$ and $i\omega_{1,\ell}$ at $\ell = 0$. Since m is monotonic for $k > 0$, the remaining eigenvalues at $\ell = 0$ are all simple, purely imaginary, and located outside the open ball $B(0; |m(k) - m(2k)|)$. The perturbation analysis to locate the bifurcation of these eigenvalues for small ℓ and a will correspond to long wavelength transverse perturbations. The following lemma ensures that for sufficiently small ℓ and a , bifurcating eigenvalues from the origin are separated from the rest of the spectrum by a non-zero distance.

Lemma 3.4. *The following properties hold, for any ℓ and a sufficiently small.*

(1) *The spectrum of $\mathcal{A}_a(\ell)$ decomposes as*

$$\text{spec}_0(\mathcal{A}_a(\ell)) \cup \text{spec}_1(\mathcal{A}_a(\ell)),$$

with

$$\text{spec}_0(\mathcal{A}_a(\ell)) \subset B(0; R/3), \quad \text{spec}_1(\mathcal{A}_a(\ell)) \subset \mathbb{C} \setminus \overline{B(0; R/2)}$$

where $R = |m(k) - m(2k)|$.

(2) *The spectral projection $\Pi_a(\ell)$ associated with $\text{spec}_0(\mathcal{A}_a(\ell))$ satisfies $\|\Pi_a(\ell) - \Pi_0(0)\| = O(\ell^2 + |a|)$.*

(3) *The spectral subspace $\mathcal{X}_a(\ell) = \Pi_a(\ell)(L_0^2(0, 2\pi))$ is two dimensional.*

The proof of these properties is similar to [Har11, Lemma 4.7]. In the following lemma, we show that for sufficiently small ℓ and a , two eigenvalues in $\text{spec}_0(\mathcal{A}_a(\ell))$ leave imaginary axis if $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$ but remain on imaginary axis if $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$.

Lemma 3.5. *Assume ℓ and a are sufficiently small. For $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$, there exists $\ell_a^2 = (A_2 - 2A_0)\sigma a^2 + O(a^4) > 0$, such that*

(1) *for any $\ell^2 \geq \ell_a^2$, the spectrum of $\mathcal{A}_a(\ell)$ is purely imaginary.*

(2) *for any $\ell^2 < \ell_a^2$, the spectrum of $\mathcal{A}_a(\ell)$ is purely imaginary, except for a pair of simple real eigenvalues with opposite signs.*

For $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$, the spectrum of $\mathcal{A}_a(\ell)$ is purely imaginary.

Proof. Consider the decomposition of the spectrum of $\mathcal{A}_a(\ell)$ in Lemma 3.4. The eigenvalues in $\text{spec}_0(\mathcal{A}_a(\ell))$ are the eigenvalues of the restriction of $\mathcal{A}_a(\ell)$ to the two-dimensional spectral subspace $\mathcal{X}_a(\ell)$. We determine the location of these eigenvalues by computing successively a basis of $\mathcal{X}_a(\ell)$, the 2×2 matrix representing the action of $\mathcal{A}_a(\ell)$ on this basis, and the eigenvalues of this matrix. Note that for $a = 0$, $\mathcal{X}_0(\ell)$ is spanned by $\{\cos z, \sin z\}$. Moreover, a direct calculation shows that zero is an $L_0^2(\mathbb{T})$ -eigenvalue of $\mathcal{A}_a(0)$ of multiplicity two with eigenfunctions $(\partial_b c)(\partial_a w) - (\partial_a c)(\partial_b w)(z; k, a, 0)$ and $\partial_z w(z; k, a, 0)$. We use expansions of w and c in (1.8) to calculate expansion of a basis $\{\phi_1, \phi_2\}$ for $\mathcal{X}_a(\ell)$ for small a and ℓ as

$$\begin{aligned} \phi_1(z) &:= \frac{1}{(1 - m(k))} ((\partial_b c)(\partial_a w) - (\partial_a c)(\partial_b w))(z; k, a, 0) \\ (3.10) \quad &:= \cos z + 2aA_2 \cos 2z + 3a^2A_3 \cos 2z + O(a^3), \end{aligned}$$

$$(3.11) \quad \phi_2(z) := -\frac{1}{a} \partial_z w(z; k, a, 0) = \sin z + 2aA_2 \sin 2z + 3a^2A_3 \sin 3z + O(a^3).$$

We use expansion of $\mathcal{A}_a(\ell)$ in (3.8) to find actions of $\mathcal{A}_a(\ell)$ and identity operator on $\{\phi_1, \phi_2\}$ as

$$\mathbf{B}_a(\ell) = \begin{pmatrix} 0 & \frac{\sigma \ell^2 + (A_0 - A_2)a^2}{1 + 4a^2A_2^2} \\ \frac{-\sigma \ell^2 - A_0a^2}{1 + 4a^2A_2^2} & 0 \end{pmatrix} + O(a(\ell^2 + a^2)).$$

and

$$(3.12) \quad \mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + O(a^4)$$

for $|a|$ and $|\ell|$ sufficiently small. To locate the two eigenvalues bifurcating from the origin, we examine the characteristic equation $\det(\mathbf{B}_a(\ell) - \lambda \mathbf{I}) = 0$, which leads to

$$(1 + 4a^2 A_2^2) \lambda^2 + (2A_0 - A_2) a^2 \sigma \ell^2 + \ell^4 \sigma^2 + O(a^2 \ell^2 (\ell^2 + a^2)) = 0.$$

From which we conclude that

$$\lambda^2 = - \left(\frac{\ell^2}{(1 + 4a^2 A_2^2)^2} \right) (\ell^2 - (A_2 - 2A_0) \sigma a^2) + O(a^2 \ell^2 (\ell^2 + a^2)).$$

For $\ell = a = 0$, we get zero as a double eigenvalue, which agrees with our calculation. For $(\sigma, m) = (1, \downarrow)$ and $(-1, \uparrow)$, we obtain two purely imaginary eigenvalues for all ℓ and a sufficiently small. For $(\sigma, m) = (1, \uparrow)$ and $(-1, \downarrow)$, we obtain two purely imaginary eigenvalues when $\ell^2 \geq \ell_a^2$, and real eigenvalues, with opposite signs when $\ell^2 < \ell_a^2$ where

$$\ell_a^2 = (A_2 - 2A_0) \sigma a^2 + O(a^4).$$

Now, we study $\text{spec}_1(\mathcal{A}_a(\ell))$ for ℓ and a sufficiently small. For $n \in \mathbb{Z} \setminus \{-1, 0, 1\}$, as ℓ is sufficiently small, κ_n in (3.3) depends upon behavior of $m(k)$, which is monotonic for all $k > 0$. Hence, k_n is negative for all n and σ when m is monotonically increasing, and k_n is positive for all n and σ when m is monotonically decreasing. This implies that even if eigenvalues in $\text{spec}_1(\mathcal{A}_0(\ell))$ collide, they remain on the imaginary axis and for all ℓ and a sufficiently small $\text{spec}_1(\mathcal{A}_a(\ell))$ is a subset of the imaginary axis. This proves the lemma. \square

4. NON-PERIODIC PERTURBATIONS

In this section, we will study two-dimensional perturbations which are non-periodic (localized or bounded) in the direction of the propagation of the wave. For non-periodic perturbations, we study the invertibility of $\mathcal{T}_a(\lambda, \ell)$ in (2.2) acting in $L^2(\mathbb{R})$ or $C_b(\mathbb{R})$ (with domain $H^{\alpha+2}(\mathbb{R})$ or $C_b^{\alpha+2}(\mathbb{R})$), for $\lambda \in \mathbb{C}$, $\Re(\lambda) > 0$, and $\ell \in \mathbb{R}$, $\ell \neq 0$. Since coefficients of $\mathcal{T}_a(\lambda, \ell)$ are periodic functions, using Floquet Theory, all solutions of (2.2) in $L^2(\mathbb{R})$ or $C_b(\mathbb{R})$ are of the form $V(z) = e^{i\xi z} \tilde{V}(z)$ where $\xi \in (-\frac{1}{2}, \frac{1}{2}]$ is the Floquet exponent and \tilde{V} is a 2π -periodic function, see [Har08] for a similar situation. This replaces the study of invertibility of the operator $\mathcal{T}_a(\lambda, \ell)$ in $L^2(\mathbb{R})$ or $C_b(\mathbb{R})$ by the study of invertibility of a family of Bloch operators in $L^2(\mathbb{T})$ parameterized by the Floquet exponent ξ . We present the precise reformulation in the following lemma.

Lemma 4.1. *The linear operator $\mathcal{T}_a(\lambda, \ell)$ is invertible in $L^2(\mathbb{R})$ if and only if the linear operators*

$$\mathcal{T}_{a,\xi}(\lambda, \ell) = \lambda(\partial_z + i\xi) - (\partial_z + i\xi)^2(c - \mathcal{M}_k + w) - \sigma \ell^2,$$

acting in $L^2(\mathbb{T})$ with domain $H^{\alpha+2}(\mathbb{T})$ are invertible for all $\xi \in (-\frac{1}{2}, \frac{1}{2}]$.

We refer to [Har11, Lemma 5.1] for a detailed proof in the similar situation. The fact that the operators $\mathcal{T}_{a,\xi}(\lambda, \ell)$ act in $L^2(\mathbb{T})$ implies that these operators have only point spectrum. Note that $\xi = 0$ corresponds to the periodic perturbations which we have already investigated, so now we would restrict ourselves to the case of $\xi \neq 0$. The operator $\partial_z + i\xi$ is invertible in $L^2(\mathbb{T})$. Using this, we have the following result.

Lemma 4.2. *The operator $\mathcal{T}_{a,\xi}(\lambda, \ell)$ is not invertible in $L^2(\mathbb{T})$ for some $\lambda \in \mathbb{C}$ and $\xi \neq 0$ if and only if $\lambda \in \text{spec}(\mathcal{A}_a(\ell, \xi))$, $L^2(\mathbb{T})$ -spectrum of the operator,*

$$\mathcal{A}_a(\ell, \xi) := (\partial_z + i\xi)(c - \mathcal{M}_k + w) + \sigma\ell^2(\partial_z + i\xi)^{-1}.$$

Note that the operator $(\partial_z + i\xi)^{-1}$ becomes singular, as $\xi \rightarrow 0$, so replacing the study of the invertibility of $\mathcal{T}_{a,\xi}(\lambda, \ell)$ by the study of the spectrum of $\mathcal{A}_a(\ell, \xi)$ is not suitable for small ξ . To avoid this, we only study the spectrum of $\mathcal{A}_a(\ell, \xi)$ for $|\xi| > \epsilon > 0$. Also, for $\xi \in (-\frac{1}{2}, \frac{1}{2}]$ and $\xi \neq 0$, since the spectrum $\text{spec}(\mathcal{A}_a(\ell, \xi))$ is symmetric with respect to the imaginary axis, and $\text{spec}(\mathcal{A}_a(\ell, \xi)) = \text{spec}(-\mathcal{A}_a(\ell, -\xi))$, we can restrict our study to $\xi \in (0, \frac{1}{2}]$.

We will study the $L^2(\mathbb{T})$ -spectra of linear operators $\mathcal{A}_a(\ell, \xi)$ for $|a|$ sufficiently small. It is straightforward to establish the estimate

$$\|\mathcal{A}_a(\ell, \xi) - \mathcal{A}_0(\ell, \xi)\| = O(|a|)$$

as $a \rightarrow 0$ uniformly for $\xi \in (0, \frac{1}{2}]$ in the operator norm. Therefore, In order to locate the spectrum of $\mathcal{A}_a(\ell, \xi)$, we need to determine the spectrum of $\mathcal{A}_0(\ell, \xi)$. A simple calculation yields that

$$\mathcal{A}_0(\ell, \xi)e^{inz} = i\omega_{n,\ell,\xi}e^{inz}, \quad n \in \mathbb{Z}$$

where

$$\omega_{n,\ell,\xi} = (n + \xi)(m(k) - m(k(n + \xi))) - \frac{\sigma\ell^2}{n + \xi}.$$

As in the previous section, the linear operator $\mathcal{A}_a(\ell, \xi)$ can be decomposed as

$$\mathcal{A}_a(\ell) = J_\xi \mathcal{L}_a(\ell, \xi)$$

where

$$J_\xi = \partial_z + i\xi \quad \text{and} \quad \mathcal{L}_a(\ell, \xi) = c - \mathcal{M}_k + w + \sigma\ell^2(\partial_z + i\xi)^{-2}$$

The operator J_ξ is skew-adjoint, whereas the operator $\mathcal{L}_a(\ell)$ is self-adjoint. As defined in (3.3), the Krein signature, $\kappa_{n,\xi}$ of an eigenvalue $i\omega_{n,\ell,\xi}$ in $\text{spec}(\mathcal{A}_0(\ell, \xi))$ is

$$(4.1) \quad \kappa_{n,\xi} = \text{sgn} \left(m(k) - m(k(n + \xi)) - \frac{\sigma\ell^2}{(n + \xi)^2} \right), \quad n \in \mathbb{Z}.$$

Therefore, a necessary condition for bifurcation of colliding eigenvalues, $i\omega_{p,\ell,\xi}$ and $i\omega_{q,\ell,\xi}$, from the imaginary axis is $(p + \xi_0)(q + \xi_0) < 0$, where ξ_0 is the value of the floquet exponent where eigenvalues $i\omega_{p,\ell,\xi}$ and $i\omega_{q,\ell,\xi}$ collide. As in the previous section, we split the analysis in this section in finite and short wavelength, and long wavelength transverse perturbations.

4.1. Finite and short-wavelength transverse perturbations. We start the analysis of the spectrum of $\mathcal{A}_a(\ell, \xi)$ with the values of ℓ away from the origin, $|\ell| \geq \ell_0$, for some $\ell_0 > 0$. We further split the analysis into two cases depending on the value of σ and nature of m .

4.1.1. $\boxed{(\sigma, m) = (1, \uparrow) \text{ or } (-1, \downarrow)}$. It is straightforward to verify that for $(\sigma, m) = (1, \uparrow)$ and $(-1, \downarrow)$, $\kappa_{n,\xi}$ is negative for all $n \in \mathbb{Z} \setminus \{-1, 0\}$ and positive for all $n \in \mathbb{Z} \setminus \{-1, 0\}$, respectively, for all $k > 0$ and $\xi \in (0, \frac{1}{2}]$. However, Krein signatures of $i\omega_{-1,\ell,\xi}$ and $i\omega_{0,\ell,\xi}$ can be positive as well as negative depending upon the values of k, ℓ and ξ . Therefore, collision of $i\omega_{-1,\ell,\xi}$ and $i\omega_{0,\ell,\xi}$ with each other or with any other eigenvalue may lead to instability. Another straightforward calculation reveals that eigenvalues $i\omega_{-1,\ell,\xi}$ and $i\omega_{0,\ell,\xi}$ collide when

$$(4.2) \quad \ell^2 = \ell_c^2 = \sigma\xi(1-\xi)[(1-\xi)(m(k) - m(k(1-\xi))) + \xi(m(k) - m(k\xi))] > 0$$

for all $\xi \in (0, \frac{1}{2}]$ while there are no collisions between pairs $\{i\omega_{0,\ell,\xi}, i\omega_{-n,\ell,\xi}\}$, $n \in \mathbb{N} \setminus \{1\}$ and $\{i\omega_{-1,\ell,\xi}, i\omega_{m,\ell,\xi}\}$, $m \in \mathbb{N}$. Therefore, we are left with only one pair which may bifurcate into potentially unstable eigenvalues which is $\{\omega_{-1,\ell,\xi}, \omega_{0,\ell,\xi}\}$. We further perform perturbation calculations to obtain the following result.

Lemma 4.3. *Assume that $\xi \in (0, \frac{1}{2}]$ and consider ℓ_c^2 given in (4.2). For $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$ and $|a|$ sufficiently small, there exists $\varepsilon_a(\xi) = \xi^{3/2}(1-\xi)^{3/2}|a| + O(a^2) > 0$ such that*

- (1) *for $|\ell^2 - \ell_c^2| \geq \varepsilon_a(\xi)$, the spectrum of $\mathcal{A}_a(\ell, \xi)$ is purely imaginary.*
- (2) *for $|\ell^2 - \ell_c^2| < \varepsilon_a(\xi)$, the spectrum of $\mathcal{A}_a(\ell, \xi)$ is purely imaginary, except for a pair of complex eigenvalues with opposite nonzero real parts.*

Proof. There exists a curve $\ell = \ell_c$ given in (4.2) along which

$$\omega(\ell_c, \xi) := \omega_{-1,\ell_c,\xi} = \omega_{0,\ell_c,\xi}.$$

Furthermore,

$$(4.3) \quad \phi_{0,-1}(z) = e^{-iz} \quad \text{and} \quad \phi_{0,0}(z) = 1$$

forms the corresponding eigenspace for $\mathcal{A}_0(\ell_c, \xi)$ associated with the two eigenvalues. Let

$$(4.4) \quad i\omega(\ell_c, \xi) + i\mu_{a,\ell,-1} \quad \text{and} \quad i\omega(\ell_c, \xi) + i\mu_{a,\ell,0}$$

be the eigenvalues of $\mathcal{A}_a(\ell_c, \xi)$ bifurcating from $i\omega_{-1,\ell_c,\xi}$ and $i\omega_{0,\ell_c,\xi}$ respectively for $|a|$ and $|\ell - \ell_c|$ small. Let $\{\phi_{a,\ell,-1}(z), \phi_{a,\ell,0}(z)\}$ be the extended eigenspace associated with two bifurcating eigenvalues. Following [CDT21], we can take,

$$(4.5) \quad \phi_{a,-1,\ell}(z) = e^{-iz} + a\phi_{-1} + O(a^2),$$

$$(4.6) \quad \phi_{a,0,\ell}(z) = 1 + a\phi_0 + O(a^2)$$

Using constraint of orthonormality on the eigenfunctions $\phi_{a,\ell,-1}$ and $\phi_{a,\ell,0}$, we obtain

$$\phi_{-1} = \phi_0 = 0.$$

To locate eigenvalues, we calculate matrix representations of $\mathcal{A}_a(\ell_c, \xi)$ and identity operators on $\{\phi_{a,-1,\ell}(z), \phi_{a,0,\ell}(z)\}$ for $|a|$ and $|\ell - \ell_c|$ small and find that

$$\mathbf{B}_a(\ell, \xi) = \begin{pmatrix} i\omega(\ell_c, \xi) - i\frac{\sigma\varepsilon}{\xi-1} & \frac{i}{2}a\xi \\ \frac{i}{2}(\xi-1)a & i\omega(\ell_c, \xi) - i\frac{\sigma\varepsilon}{\xi} \end{pmatrix} + O(|a|(|\varepsilon| + |a|)),$$

where $\varepsilon = \ell^2 - \ell_c^2$ and \mathbf{I}_a is the 2×2 identity matrix. Solving the characteristic equation $\det(\mathbf{B}_a(\ell, \xi) - \lambda \mathbf{I}_a) = 0$ for λ of the form

$$\lambda = i\omega(\ell_c, \xi) + i\mu,$$




FIGURE 2. Collision of eigenvalues at $\ell = \ell_c$ for $k = 1$ and $\xi = 0.25$.

leads to the polynomial equation

$$\mu^2 + \mu\sigma\varepsilon \left(\frac{1}{\xi-1} + \frac{1}{\xi} + O(a^2) \right) - \frac{1}{4}\xi(\xi-1)a^2 + \frac{\varepsilon^2}{\xi(\xi-1)} + O(a^2(|\varepsilon| + a^2)) = 0.$$

A direct computation shows that the discriminant of this polynomial is

$$\text{disc}_a(\varepsilon, \xi) = \frac{\varepsilon^2}{\xi^2(\xi-1)^2} + \xi(\xi-1)a^2 + O(a^2(|\varepsilon| + a^2)).$$

For any a sufficiently small there exists

$$\varepsilon_a(\xi) = \xi^{3/2}(1-\xi)^{3/2}|a| + O(a^2) > 0$$

such that the two eigenvalues of $\mathcal{A}_a(\ell, \xi)$ are purely imaginary when $|\ell^2 - \ell_c^2| \geq \varepsilon_a(\xi)$ and complex with opposite nonzero real parts when $|\ell^2 - \ell_c^2| < \varepsilon_a(\xi)$, which proves the lemma. \square

Figure 2 depicts a collision of pair of eigenvalues in KP-ILW-I and KP-Whitham-II equations which lead to instability according to Lemma 4.3.

4.1.2. $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$. In contrast to the Section 4.1.1, for $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$, there are infinitely many pairs of eigenvalues which collide with each other. A pair $\{\omega_{p,\ell,\xi}, \omega_{-q,\ell,\xi}\}$ collide with each other at

$$\ell^2 = \ell_{p,q}^2 = \frac{\sigma(p+\xi)(q-\xi)}{p+q}((p+\xi)(m(k) - m(k(p+\xi))) + (q-\xi)(m(k) - m(k(q-\xi))))$$

for all $\xi \in (0, 1/2]$, and for all $p \in \mathbb{N} \cup \{0\}$, $q \in \mathbb{N}$ except $(p, q) = (0, 1)$. A Krein signature analysis tells that for each pair of colliding eigenvalues; there are intervals of ℓ where they have opposite Krein signatures. Therefore, all such collisions may lead to instability. An additional analysis is required in order to detect the bifurcating eigenvalues which indeed leave the imaginary axis and become unstable. In what follows, we examine two pairs $\{\omega_{-1,\ell,\xi}, \omega_{1,\ell,\xi}\}$ and $\{\omega_{0,\ell,\xi}, \omega_{-2,\ell,\xi}\}$ whose indices differ by two. We have the following result.

Lemma 4.4. Assume that $\xi \in (0, 1/2]$ and set

$$\ell_c^2 = \frac{\sigma(n+2+\xi)(n+\xi)}{2}((n+\xi)(m(k) - m(k(n+\xi))) - (n+2+\xi)(m(k) - m(k(n+2+\xi)))).$$

For $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$, $|a|$ sufficiently small, and $n \in \{-1, -2\}$, there exists $\varepsilon_a = \sigma a^2 A_2(n + \xi)(n + 2 + \xi) + O(a^3)$ such that the spectrum of $\mathcal{A}_a(\ell, \xi)$ have a pair of complex eigenvalues with opposite nonzero real parts for $|\ell^2 - \ell_c^2| < \varepsilon_a$.

Proof. We will prove this result by analyzing collisions whose indices differ by two. Let $i\omega_{n,\ell,\xi}$ and $i\omega_{n+2,\ell,\xi}$ be such two eigenvalues for some $n \in \mathbb{Z}$. Assume that these eigenvalues collide at $\ell = \ell_c$, that is

$$(4.7) \quad 0 \neq \omega_{n,\ell_c} = \omega_{n+2,\ell_c} = \omega \text{ (say)}.$$

Furthermore,

$$(4.8) \quad \phi_{0,n}(z) = e^{inz} \quad \text{and} \quad \phi_{0,n+2}(z) = e^{i(n+2)z}$$

form a basis for the corresponding eigenspace for $\mathcal{A}_0(\ell_c, \xi)$ generated by the two eigenvalues. Let

$$(4.9) \quad i\omega + i\mu_{a,n} \quad \text{and} \quad i\omega + i\mu_{a,n+2}$$

be the eigenvalues of $\mathcal{A}_a(\ell_c, \xi)$ bifurcating from $i\omega_{n,\ell_c,\xi}$ and $i\omega_{n+2,\ell_c,\xi}$ respectively, for $|a|$ small. Let $\{\phi_{a,n}(z), \phi_{a,n+2}(z)\}$ be a orthonormal basis for the corresponding eigenspace. We assume the following expansions

$$(4.10) \quad \phi_{a,n,\ell}(z) = e^{inz} + a\phi_{n,1} + a^2\phi_{n,2} + O(a^3),$$

$$(4.11) \quad \phi_{a,n+2,\ell}(z) = e^{i(n+2)z} + a\phi_{n+2,1} + a^2\phi_{n+2,2} + O(a^3).$$

We use orthonormality of $\phi_{a,n,\ell}$ and $\phi_{a,n+2,\ell}$ to find that

$$\phi_{n,1} = \phi_{n,2} = \phi_{n+2,1} = \phi_{n+2,2} = 0.$$

Next, we compute the action of $\mathcal{A}_a(\ell, \xi)$ and identity operators on the extended eigenspace $\{\phi_{a,n,\ell}(z), \phi_{a,n+2,\ell}(z)\}$ for $|\ell - \ell_c|$ and $|a|$ small. We arrive at

$$B_a(\ell) = \begin{pmatrix} i\omega - ia^2 \frac{A_2}{2}(n + \xi) - \frac{i\sigma\varepsilon}{n + \xi} & ia^2 \frac{A_2}{2}(n + 2 + \xi) \\ ia^2 \frac{A_2}{2}(n + \xi) & i\omega - ia^2 \frac{A_2}{2}(n + 2 + \xi) - \frac{i\sigma\varepsilon}{n + 2 + \xi} \end{pmatrix} + O(a^4).$$

where $\varepsilon = |\ell^2 - \ell_c^2|$ and \mathbf{I}_a , the 2×2 identity matrix, respectively. Solving the characteristic equation $|\mathcal{B}_a(\ell) - (i\omega + i\mu)\mathcal{I}_a| = 0$ leads to

$$\begin{aligned} |\mathcal{B}_a(\ell) - (i\omega + i\mu)\mathcal{I}_a| &= \mu^2 + \mu \left(\sigma\varepsilon \left(\frac{1}{n + \xi} + \frac{1}{n + 2 + \xi} \right) + \frac{a^2 A_2}{2}((n + 2 + \xi) + (n + \xi)) \right) \\ &\quad + \frac{a^2 A_2 \sigma\varepsilon}{2} \left(\frac{n + \xi}{n + 2 + \xi} + \frac{n + 2 + \xi}{n + \xi} \right) + \frac{\sigma^2 \varepsilon^2}{(n + \xi)(n + 2 + \xi)} + O(a^4) = 0. \end{aligned}$$

A direct computation shows that the discriminant of this quadratic in μ is

$$\text{disc}_a(\varepsilon) = \left(\sigma\varepsilon \left(\frac{1}{n + \xi} - \frac{1}{n + 2 + \xi} \right) + \frac{a^2 A_2}{2}((n + 2 + \xi) + (n + \xi)) \right)^2 - \frac{4\sigma\varepsilon a^2 A_2}{n + \xi}.$$




FIGURE 3. Collision of eigenvalues at $\ell = \ell_c$ for $k = 1$ and $\xi = 0.25$.

As a is sufficiently small, we can rewrite $\text{disc}_a(\varepsilon)$ as

$$\text{disc}_a(\varepsilon) = \frac{4\varepsilon}{(n + \xi)^2(n + 2 + \xi)^2}(\varepsilon - \sigma a^2 A_2(n + \xi)(n + 2 + \xi)) + O(a^3).$$

If we choose ℓ such that

$$(4.12) \quad |\ell^2 - \ell_c^2| < \sigma a^2 A_2(n + \xi)(n + 2 + \xi) + O(a^3)$$

then $\text{disc}_a(\varepsilon) < 0$ and it leads to μ to have non-zero imaginary part implying the instability of small amplitude periodic travelling waves (1.8) of (1.1). Note that there are only two such collisions of opposite Krein signatures corresponding to $n = -2$ and $n = -1$ and for both these values of n , LHS in (4.12) is positive. Hence the lemma. \square

Figure 3 depicts a collision of pair of eigenvalues in KP-Whitham-I and KP-II equations which lead to instability according to Lemma 4.4.

4.2. Long wavelength transverse Perturbations. We now consider the spectrum of $\mathcal{A}_a(\ell, \xi)$ for $|\ell|$ small. Recall that ξ is away from the origin, and we have taken $|\xi| > \epsilon$ for some small but fixed $\epsilon > 0$. Because of this, the collision at the origin and collisions away from the origin are well separated for $\mathcal{A}_0(\ell, \xi)$. This separation persists for small $|a|$ using perturbation arguments. More precisely, we have the following lemma.

Lemma 4.5. *For any ℓ and a sufficiently small, the spectrum of $\mathcal{A}_a(\ell, \xi)$ decomposes as*

$$\text{spec}(\mathcal{A}_a(\ell, \xi)) = \text{spec}_0(\mathcal{A}_a(\ell, \xi)) \cup \text{spec}_1(\mathcal{A}_a(\ell, \xi)),$$

with

$$\text{dist}(\text{spec}_0(\mathcal{A}_a(\ell, \xi)), \text{spec}_1(\mathcal{A}_a(\ell, \xi))) \geq |(m(k) - m(k(1 + \epsilon)))| > 0.$$

Let $\ell^* > 0$ be the smallest positive value of ℓ for which a collision of eigenvalues of $\mathcal{A}_0(\ell, \xi)$ takes place. Note that such an ℓ^* exists because the collision at the origin takes place only for $\ell = 0$, and other collisions are well separated from this. Now, for any ℓ with $|\ell| < \ell^*$, there are no collisions between the eigenvalues of $\mathcal{A}_0(\ell, \xi)$ since $|\xi| > \epsilon > 0$. This persists for small values of a using perturbation arguments, and we have the following lemma.

Lemma 4.6. *Assume that $\xi \in (0, \frac{1}{2}]$. There exists $\ell^* > 0$ and $a^* > 0$ such that the spectrum of $\mathcal{A}_a(\ell, \xi)$ is purely imaginary, for any ℓ and a satisfying $|\ell| < \ell^*$ and $|a| < a^*$.*

5. APPLICATIONS

We summarize the transverse stability and instability results of previous sections in the following theorems.

Theorem 5.1. *Assume that small amplitude periodic traveling waves (1.8) of (1.1) are spectrally stable in $L^2(\mathbb{T})$ as a solution of the corresponding y -independent one-dimensional equation. Then, for any a sufficiently small, $k > 0$, and m satisfying Hypotheses 1.1, periodic traveling waves (1.8) of (1.1) are transversely stable with respect to two-dimensional perturbations which are periodic in the direction of propagation of the wave and of*

- (1) *finite or short wavelength in the transverse direction if σ and m in (1.1) satisfy $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$.*
- (2) *long wavelength in the transverse direction if σ and m in (1.1) satisfy $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$.*

Theorem 5.2. *For any a sufficiently small, $k > 0$, and m satisfying Hypotheses 1.1, periodic traveling waves (1.8) of (1.1) are transversely unstable with respect to two-dimensional perturbations which are*

- (1) *periodic in the direction of propagation and of finite or short wavelength in the transverse direction if σ and m in (1.1) satisfy $(\sigma, m) = (1, \downarrow)$ or $(-1, \uparrow)$.*
- (2) *periodic in the direction of propagation of the wave and of long wavelength in the transverse direction if σ and m in (1.1) satisfy $(\sigma, m) = (1, \uparrow)$ or $(-1, \downarrow)$.*
- (3) *non-periodic (localized or bounded) in the direction of propagation of the wave and of finite or short wavelength in the transverse direction for all values of σ and m in (1.1).*

We discuss implications of Theorem 5.2 on KP-fKdV, KP-BO, KP-ILW, and KP-Whitham equations.

5.1. KP-fKdV and KP-BO Equations. The KP-fKdV equation is obtained from (1.1) by taking

$$m(k) = 1 + |k|^\beta, \quad \beta > 1/2$$

The symbol $m(k)$ clearly satisfies Hypotheses 1.1 $H1$, $H2$ ($\alpha = \beta$, $C_1 = 1$, and $C_2 = 2$), and $H3$ (m is strictly increasing for $k > 0$). The two-parameter family of periodic solutions can be obtained from (1.8) and (1.9) by replacing $m(k)$ with $1 + |k|^\beta$. We obtain KP-BO equation from KP-fKdV equation when $\beta = 1$. We obtain following stability and instability result for the KP-fKdV equation using Theorems 5.1 and 5.2.

Corollary 5.3. *Assume that small-amplitude periodic traveling waves (1.8) of the KP-fKdV equation are spectrally stable in $L^2(\mathbb{T})$ as a solution of the fKdV equation. For any a sufficiently small and $k > 0$,*

- (1) *periodic traveling waves (1.8) of the KP-fKdV-I equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*
- (2) *periodic traveling waves (1.8) of the KP-fKdV-II equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of long wavelength in the transverse direction.*

Corollary 5.4. *For any a sufficiently small and $k > 0$,*

- (1) *periodic traveling waves (1.8) of KP-fKdV-I equation are transversely unstable with respect to two-dimensional perturbations, which are*
 - (a) *periodic in the direction of propagation of the wave and of long-wavelength in the transverse direction, and*
 - (b) *non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*
- (2) *periodic traveling waves (1.8) of the KP-fKdV-II equation are transversely unstable with respect to two-dimensional perturbations, which are periodic or non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*

For $\beta = 1$, KP-fKdV reduces to KP-BO equation, Corollaries 5.3 and 5.4 hold true for the KP-BO equation as well. For $\beta = 2$, KP-fKdV equation reduces to the KP equation (1.3). As results in [HK08, BD09] show that all periodic traveling waves of the KdV equation are spectrally stable in $L^2(\mathbb{T})$, from Corollary 5.3, small-amplitude periodic traveling waves (1.8) of KP-I (and KP-II resp.) are transversely stable with respect to two-dimensional perturbations which are periodic in the direction of propagation of the wave and of finite or short-wavelength (and long-wavelength resp.) in the transverse direction. These stability results agree with results in [Har11]. The transverse instability results obtained here for KP-I in Corollary 5.4 agrees with [Har11]. From Corollary 5.4, we obtain the following transverse instability result for the KP-II equation.

Corollary 5.5. *Small amplitude periodic traveling waves (1.8) of the KP-II equation are transversely unstable with respect to two-dimensional perturbations, which are periodic or non-periodic (localized or bounded) in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*

5.2. KP-ILW Equation. The KP-ILW equation is obtained from (1.1) by taking,

$$m(k) = k \coth k$$

The symbol $m(k)$ satisfies Hypotheses 1.1 H1, H2 ($\alpha = 2$, $C_1 = 1$, and $C_2 = 2$), and H3 (m is strictly increasing for $k > 0$). The two-parameter family of periodic solutions can be obtained from (1.8) and (1.9) by replacing $m(k)$ with $k \coth k$. By using Theorem 5.1 and 5.2, we obtain following corollaries.

Corollary 5.6. *Assume that small-amplitude periodic traveling waves (1.8) of the KP-ILW equation are spectrally stable in $L^2(\mathbb{T})$ as a solution of the ILW equation. For any a sufficiently small and $k > 0$,*

- (1) periodic traveling waves (1.8) of KP-ILW-I equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.
- (2) periodic traveling waves (1.8) of KP-ILW-II equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of long wavelength in the transverse direction.

Corollary 5.7. *For any a sufficiently small and $k > 0$,*

- (1) periodic traveling waves (1.8) of KP-ILW-I equation are transversely unstable with respect to two-dimensional perturbations, which are
 - (a) periodic in the direction of propagation of the wave and of long-wavelength in the transverse direction, and
 - (b) non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.
- (2) periodic traveling waves (1.8) of the KP-ILW-II equation are transversely unstable with respect to two-dimensional perturbations, which are periodic or non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.

5.3. KP-Whitham Equation. The KP-Whitham equation is obtained from (1.1) by taking,

$$m(k) = \sqrt{\frac{\tanh k}{k}}$$

The symbol $m(k)$ satisfies Hypotheses 1.1 H1, H2 with $\alpha = -\frac{1}{2}$, $C_1 = 1$, and $C_2 = 2$, and H3 as m is strictly decreasing for $k > 0$. The two-parameter family of periodic solutions can be obtained from (1.8) and (1.9) by replacing $m(k)$ with $\sqrt{\frac{\tanh k}{k}}$. Since $m(k)$ is monotonically decreasing, by using Theorem 5.1 and 5.2, we obtain following results.

Corollary 5.8. *Assume that small-amplitude periodic traveling waves (1.8) of the KP-Whitham equation are spectrally stable in $L^2(\mathbb{T})$ as a solution of the Whitham equation. For any a sufficiently small and $k > 0$,*

- (1) periodic traveling waves (1.8) of the KP-Whitham-I equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of long wavelength in the transverse direction.
- (2) periodic traveling waves (1.8) of the KP-Whitham-II equation are transversely stable with respect to two-dimensional perturbations, which are periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.

Corollary 5.9. *For any a sufficiently small and $k > 0$,*

- (1) periodic traveling waves (1.8) of KP-Whitham-I equation are transversely unstable with respect to two-dimensional perturbations, which are
 - (a) periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction, and

- (b) *non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*
- (2) *periodic traveling waves (1.8) of the KP-Whitham-II equation are transversely unstable with respect to two-dimensional perturbations, which are*
 - (a) *periodic in the direction of propagation of the wave and of long wavelength in the transverse direction, and*
 - (b) *non-periodic in the direction of propagation of the wave and of finite or short wavelength in the transverse direction.*

ACKNOWLEDGEMENT

Bhavna and AKP are supported by the Science and Engineering Research Board (SERB), Department of Science and Technology (DST), Government of India under grant SRG/2019/000741. Bhavna is also supported by Junior Research Fellowships (JRF) by University Grant Commission (UGC), Government of India. AK is supported by JRF by Council of Scientific and Industrial Research (CSIR), Government of India.

REFERENCES

- [BD09] Nate Bottman and Bernard Deconinck, *KdV cnoidal waves are spectrally stable*, Discrete and Continuous Dynamical Systems **25** (2009), no. 4.
- [CDT21] Ryan Creedon, Bernard Deconinck, and Olga Trichtchenko, *High-Frequency Instabilities of the Kawahara Equation: A Perturbative Approach*.
- [GHS01] Mark D. Groves, Mariana Haragus, and Shu-Ming Sun, *Transverse instability of gravity-capillary line solitary water waves*, C. R. Acad. Sci. Paris Sér. I Math. **333** (2001), no. 5, 421–426. MR 1859230
- [Har08] Mariana Haragus, *Stability of periodic waves for the generalized BBM equation*, Rev. Roumaine Maths. Pures Appl. **53** (2008), 445–463.
- [Har11] ———, *Transverse Spectral Stability of Small Periodic Traveling Waves for the KP Equation*, Studies in Applied Mathematics **126** (2011), no. 2.
- [HJ15] Vera Mikiyoung Hur and Mathew A. Johnson, *Modulational Instability in the Whitham Equation for Water Waves*, Studies in Applied Mathematics **134** (2015), no. 1.
- [HK08] Mariana Haragus and Todd Kapitula, *On the spectra of periodic waves for infinite-dimensional Hamiltonian systems*, Physica D: Nonlinear Phenomena **237** (2008), no. 20.
- [HSS12] Sevdzhan Hakkaev, Milena Stanislavova, and Atanas Stefanov, *Transverse instability for periodic waves of KP-I and Schrödinger equations*, Indiana Univ. Math. J. **61** (2012), no. 2, 461–492. MR 3043584
- [JZ10] Mathew A. Johnson and Kevin Zumbrun, *Transverse instability of periodic traveling waves in the generalized Kadomtsev-Petviashvili equation*, SIAM Journal on Mathematical Analysis **42** (2010), no. 6.
- [KP70] B.B. Kadomtsev and V.I. Petviashvili, *On the stability of solitary waves in weakly dispersive media*, Soviet Physics Doklady **192** (1970), 753–756.
- [PS04] Robert L. Pego and S. M. Sun, *On the transverse linear instability of solitary water waves with large surface tension*, Proc. Roy. Soc. Edinburgh Sect. A **134** (2004), no. 4, 733–752. MR 2079803
- [RT09] F. Rousset and N. Tzvetkov, *Transverse nonlinear instability for two-dimensional dispersive models*, Ann. Inst. H. Poincaré Anal. Non Linéaire **26** (2009), no. 2, 477–496. MR 2504040
- [RT11] Frederic Rousset and Nikolay Tzvetkov, *Transverse instability of the line solitary water-waves*, Invent. Math. **184** (2011), no. 2, 257–388. MR 2793858

- [Sau95] Jean Claude Saut, *Recent results on the generalized Kadomtsev-Petviashvili equations*, Acta Applicandae Mathematicae **39** (1995), no. 1-3.

DEPARTMENT OF MATHEMATICS, IIIT DELHI, INDIA 110020

Email address: bhavnai@iiitd.ac.in, atulk@iiitd.ac.in, ashish.pandey@iiitd.ac.in