A GLOBAL SPACE-TIME ESTIMATE FOR DISPERSIVE OPERATORS THROUGH ITS LOCAL ESTIMATE

CHU-HEE CHO, YOUNGWOO KOH, AND JUNGJIN LEE

ABSTRACT. We will show that a local space-time estimate implies a global space-time estimate for dispersive operators. In order for this implication we consider a Littlewood-Paley type square function estimate for dispersive operators in a time variable and a generalization of Tao's epsilon removal lemma in mixed norms. By applying this implication to the fractional Schrödinger equation in \mathbb{R}^{2+1} we obtain the sharp global space-time estimates with optimal regularity from the previous known local ones.

1. INTODUCTION

Let us consider a Cauchy problem of a dispersive equation in \mathbb{R}^{n+1}

$$\begin{cases} i\partial_t u + \Phi(D)u = 0, \\ u(0) = f, \end{cases}$$

$$(1.1)$$

where $\Phi(D)$ is the corresponding Fourier multiplier to the function Φ . We assume that $\Phi \in C^{\infty}(\mathbb{R}^n \setminus \{0\})$ is a real-valued function satisfying the following conditions:

Condition 1.1.

- $|\nabla \Phi(\xi)| \neq 0$ for all $\xi \neq 0$.
- There is a constant $\mu \ge 1$ such that $\mu^{-1} \le |\Phi(\xi)| \le \mu$ for any ξ with $|\xi| = 1$.
- There is a constant $m \ge 1$ such that $\Phi(\lambda \xi) = \lambda^m \Phi(\xi)$ for all $\lambda > 0$ and all $\xi \ne 0$.
- The Hessian $H_{\Phi}(\xi)$ of Φ has rank at least 1 for all $\xi \neq 0$.

The solution u to (1.1) becomes the Schrödinger operator $e^{-it\Delta}f$ if $\Phi(\xi) = |\xi|^2$ and the wave operator $e^{it\sqrt{-\Delta}}f$ if $\Phi(\xi) = |\xi|$. When $\Phi(\xi) = |\xi|^m$ for m > 1, the solution is called the fractional Schrödinger operator $e^{it(\sqrt{-\Delta})^{m/2}}f$.

Let $e^{it\Phi(D)}f$ denote the solution to (1.1). Our interest is to find suitable pairs (q, r) which satisfy the global space-time estimate

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{n};L^{r}_{t}(\mathbb{R}))} \le C\|f\|_{\dot{H}^{s}(\mathbb{R}^{n})},$$
(1.2)

where $\dot{H}^{s}(\mathbb{R}^{n})$ denotes the homogeneous L^{2} Sobolev space of order s. By scaling invariance the regularity s = s(r, q) should be defined as

$$s = n(\frac{1}{2} - \frac{1}{q}) - \frac{m}{r}.$$
(1.3)

This problem for $\mu = 1$ has been studied by many researchers. For the Schrödinger operator, Planchon [14] conjectured that the estimate (1.2) is valid if and only if $r \ge 2$ and $\frac{n+1}{q} + \frac{1}{r} \le \frac{n}{2}$. Kenig-Ponce-Vega [10] showed the conjecture is true for n = 1. In higher dimensions $n \ge 2$

Date: November 5, 2021.

²⁰²⁰ Mathematics Subject Classification. 35A24, 35L10, 35Q41, 35S35.

Key words and phrases. Schrödinger equation, dispersive operator, global space-time estimate.

C. Cho is supported by NRF grant no.2020R1I1A1A01072942 (Republic of Korea). Y. Koh is supported by the research grant of the Kongju National University in 2021. J. Lee is supported by NRF grant no.2020R1I1A1A01062209 (Republic of Korea).

it was proven by Vega [21] that (1.2) holds for $q \geq \frac{2(n+2)}{n}$ and $\frac{n+1}{q} + \frac{1}{r} \leq \frac{n}{2}$. When n = 2Rogers [15] showed it for $2 \leq r < \infty$, $q > \frac{16}{5}$ and $\frac{3}{q} + \frac{1}{r} < 1$, and later the excluded endline $\frac{3}{q} + \frac{1}{r} = 1$ was obtained by Lee–Rogers–Vargas [11]. When $n \geq 3$, Lee–Rogers–Vargas [11] improved the previous known result to $r \geq 2$, $q > \frac{2(n+3)}{n+1}$ and $\frac{n+1}{q} + \frac{1}{r} = \frac{n}{2}$. Recently it is shown by Du–Kim–Wang–Zhang [6] that the estimate (1.2) with $r = \infty$, that is, the maximal estimate fails for $n \geq 3$. For a case of the wave operator it is known that (1.2) holds for (r,q) pairs such $2 \leq r \leq q$, $q \neq \infty$ and $\frac{1}{r} + \frac{n-1}{2q} \leq \frac{n-1}{4}$ (see [8,9,13,18]). Particularly, when $r = \infty$, Rogers–Villarroya [16] showed that (1.2) with regularity $s > n(\frac{1}{2} - \frac{1}{q}) - \frac{1}{r}$ is valid for $q \geq \frac{2(n+1)}{n-1}$. For the fractional Schrödinger operator the known range of (r,q) for which the estimates hold is that $2 \leq r \leq q$, $q \neq \infty$ and $\frac{n}{2q} + \frac{1}{r} \leq \frac{n}{4}$ (see [1,2,4,12,20]).

The case of $\mu > 1$ has an interesting in its own right. The solution u is formally written as

$$u(t,x) = e^{it\Phi(D)}f(x) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(x\cdot\xi + t\Phi(\xi))}\hat{f}(\xi)d\xi.$$

From this form we see that the space-time Fourier transform of u is supported in the surface $S = \{(\xi, \Phi(\xi))\}$. It is known that the operator u is related to the curvature of S such as the sign of Gaussian curvature and the number of nonvanishing principle curvature. The Schrödinger operator corresponds to a paraboloid which has a positive Gaussian curvature, and the wave operator corresponds to a cone whose Gaussian curvature is zero. We are also interested in operators corresponding to a surface with negative Gaussian curvature. When $\mu > 1$ there is a surface with negative Gaussian curvature. For instance, the surface $\{(\xi_1, \xi_2, \xi_1^4 + 2\xi_1^3\xi_2 - 2\xi_1\xi_2^3 + \xi_2^4)\}$ has negative Gaussian curvature on a neighborhood of the point (1, 0, 1).

In this paper we will establish a local-to-global approach as follows.

Theorem 1.2. Let $\mathbb{I} = (0,1)$ be a unit interval and $\mathbb{B} = B(0,1)$ a unit ball in \mathbb{R}^n . Let $q_0, r_0 \in [2,\infty)$, s(r,q) defined as (1.3) and Φ satisfy Condition 1.1. Suppose that the local estimate

$$\|e^{it\Phi(D)}f\|_{L^{q_0}_x(\mathbb{B};L^{r_0}_t(\mathbb{I}))} \le C_{\epsilon}\|f\|_{H^{s(r_0,q_0)+\epsilon}(\mathbb{R}^n)}$$
(1.4)

holds for all $\epsilon > 0$. Then for any $q > q_0$ and $r > r_0$, the global estimate

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{n};L^{r}_{t}(\mathbb{R}))} \leq C\|f\|_{\dot{H}^{s(r,q)}(\mathbb{R}^{n})}$$
(1.5)

holds, where $H^{s}(\mathbb{R}^{n})$ denotes the inhomogeneous L^{2} -Sobolev space of order s and $\dot{H}^{s}(\mathbb{R}^{n})$ denotes homogeneous one.

The maximal estimate, which is (1.4) with $r_0 = \infty$, is related to pointwise convergence problems. When n = 2 it was proven that the maximal estimates with m > 1 and $\mu = 1$ are valid for $q_0 = 3$ and $s > \frac{1}{3}$ (see [3,5]). By interpolating with a Strichartz estimate

$$\|e^{it\Phi(D)}f\|_{L^6_x(\mathbb{R}^2;L^2_t(\mathbb{R}))} \le \|e^{it\Phi(D)}f\|_{L^2_t(\mathbb{R};L^6_x(\mathbb{R}^2))} \le C\|f\|_{\dot{H}^{(2-m)/4}(\mathbb{R}^2)}$$

we have (1.5) for the line $\frac{3}{q} + \frac{1}{r} = 1$ with $r \ge 2$. By Theorem 1.2, we can obtain the following global space-time estimates which is the Planchon conjecture for n = 2 except the endline.

Corollary 1.3. Let m > 1 and $\mu = 1$. For $2 \le r < \infty$ and $\frac{3}{q} + \frac{1}{r} < 1$, the global estimate

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{2};L^{r}_{t}(\mathbb{R}))} \leq C\|f\|_{\dot{H}^{1-\frac{2}{q}-\frac{m}{r}}(\mathbb{R}^{2})}$$

Notation. Throughout this paper let C > 0 denote various constants that vary from line to line, which possibly depend on n, q, r, m and μ . We use $A \leq B$ to denote $A \leq CB$, and if $A \leq B$ and $B \leq A$ we denote by $A \sim B$.

2. Proof of Theorem 1.2

In this section we prove Theorem 1.2 by using two propositions. In subsection 2.1 we consider a Littlewood–Paley type inequality by which the initial data f can be assumed to be Fourier supported in $\{1/2 \leq |\xi| \leq 2\}$. In subsection 2.2 we prove a mixed norm version of Tao's ε removable lemma by which the global estimates with a compact Fourier support are reduced to a local ones. In subsection 2.3 we show the two propositions imply Theorem 1.2.

2.1. A Littlewood-Paley type inequality. We discuss a Littlewood-Paley type inequality for the operator $e^{it\Phi(D)}$ in a time variable.

Let a cut-off function $\phi \in C_0^{\infty}([\frac{1}{2}, 2])$ satisfy $\sum_{k \in \mathbb{Z}} \phi(2^{-k}x) = 1$. We define Littlewood-Paley projection operators P_k and $\widetilde{P_k}$ by

$$\widehat{P_k f}(\xi) = \phi(2^{-k}|\xi|)\widehat{f}(\xi)$$
 and $\widetilde{\widetilde{P_k f}}(\tau) = \phi(2^{-mk}|\tau|)\widehat{f}(\tau)$

for $\xi \in \mathbb{R}^n$ and $\tau \in \mathbb{R}$, respectively.

Lemma 2.1. Suppose that Φ satisfies Condition 1.1. Then for $1 < r < \infty$,

$$\left\| e^{it\Phi(D)} f(x) \right\|_{L_{t}^{r}(\mathbb{R})} \leq C_{m,\mu} \left\| \left(\sum_{\substack{j,k \in \mathbb{Z}: \\ |k-j| \leq \frac{\log 2\mu}{m} + 2}} |\widetilde{P_{j}} e^{it\Phi(D)} P_{k}f(x)|^{2} \right)^{1/2} \right\|_{L_{t}^{r}(\mathbb{R})}$$

for all functions f and all $x \in \mathbb{R}^n$.

Proof. For simplicity,

$$F(t) := e^{it\Phi(D)}f(x)$$
 and $F_k(t) := e^{it\Phi(D)}P_kf(x).$

Since the projection operators are linear, we have an identity

$$F(t) = \sum_{j \in \mathbb{Z}} \widetilde{P_j} F(t) = \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} \widetilde{P_j} F_k(t).$$

For any test function $\psi \in C_0^{\infty}([-2,2])$ with $\psi = 1$ in [-1,1], the Fourier transform \widehat{f} of f is defined by

$$\widehat{f}(\tau) = \lim_{R \to \infty} \frac{1}{2\pi} \int_{\mathbb{R}} e^{it\tau} \psi\left(\frac{t}{R}\right) f(t) dt$$

in the distributional sense.

We claim that $P_i F_k(t) = 0$ if

$$|k-j| > \frac{\log_2 \mu}{m} + 2.$$
 (2.1)

Indeed, using the above definition of the Fourier transform we can write

$$\widehat{\widetilde{P_j}F_k}(\tau) = \frac{1}{(2\pi)^{n+1}} \phi\Big(\frac{|\tau|}{2^{mj}}\Big) \lim_{R \to \infty} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \bigg(\int_{\mathbb{R}} e^{it\tau} e^{it\Phi(\xi)} \psi\Big(\frac{t}{R}\Big) dt \bigg) \phi\Big(\frac{|\xi|}{2^k}\Big) \widehat{f}(\xi) d\xi$$

In the right side of the above equation, we see that the range of (τ, ξ) is contained in

$$2^{m(j-1)} \le |\tau| \le 2^{m(j+1)}$$
 and $2^{(k-1)} \le |\xi| \le 2^{(k+1)}$.

From Condition 1.1 we have a bound

$$\mu^{-1}2^{m(k-1)} \le |\Phi(\xi)| \le \mu 2^{m(k+1)}$$

Then it follows that for k and j satisfying (2.1),

$$|\tau + \Phi(\xi)| > 0.$$

By the integration by parts it implies that there exists a constant $C_0 > 0$ such that

$$\Big|\int_{\mathbb{R}} e^{it\tau} e^{it\Phi(\xi)} \psi\left(\frac{t}{R}\right) dt \Big| \le \frac{1}{C_0 R}$$

From this estimate and the Lebesgue dominated convergence theorem we obtain $\widehat{\widetilde{P_j}F_k} = 0$, which implies the claim.

By the claim, the Littlewood-Paley theory and the Cauchy-Schwarz inequality,

- . /

$$\begin{aligned} \left\| e^{it\Phi(D)} f(x) \right\|_{L_t^r(\mathbb{R})} &= \left\| \sum_{j \in \mathbb{Z}} \widetilde{P_j} \Big(\sum_{k \in \mathbb{Z}} F_k(\cdot, x) \Big) \right\|_{L_t^r(\mathbb{R})} \\ &\leq C \left\| \Big(\sum_{j \in \mathbb{Z}} \Big| \sum_{k \in \mathbb{Z}: |k-j| \le \frac{\log_2 \mu}{m} + 2} \widetilde{P_j} F_k(\cdot, x) \Big|^2 \Big)^{1/2} \right\|_{L_t^r(\mathbb{R})} \\ &\leq C_{m,\mu} \left\| \Big(\sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}: |k-j| \le \frac{\log_2 \mu}{m} + 2} |\widetilde{P_j} F_k(\cdot, x)|^2 \Big)^{1/2} \right\|_{L_t^r(\mathbb{R})}. \end{aligned}$$

This is the desired inequality.

Using the above lemma we can have the following proposition.

...

Proposition 2.2. Let $2 \le q, r < \infty$. Suppose that Φ satisfies Condition 1.1. If the estimate

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{n};L^{r}_{t}(\mathbb{R}))} \leq C\|f\|_{L^{2}(\mathbb{R}^{n})}$$
(2.2)

holds for all f with supp $\hat{f} \subset \{1/2 \leq |\xi| \leq 2\}$, then the estimate

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{n};L^{r}_{t}(\mathbb{R}))} \leq C_{m,\mu}\|f\|_{\dot{H}^{\frac{n}{2}-\frac{n}{q}-\frac{m}{r}}(\mathbb{R}^{n})}$$

holds for all f.

Proof. The Minkowski inequality and Lemma 2.1 allow that

$$\left\| e^{it\Phi(D)} f \right\|_{L^{q}_{x}(\mathbb{R}^{n}; L^{r}_{t}(\mathbb{R}))} \leq C_{m,\mu} \left\| \left(\sum_{|k-j| \leq \frac{\log_{2}\mu}{m} + 2} \left\| \widetilde{P_{j}} \left(e^{it\Phi(D)} P_{k} f \right) \right\|_{L^{r}_{t}(\mathbb{R})}^{2} \right)^{1/2} \right\|_{L^{q}_{x}(\mathbb{R}^{n})}.$$

Since $\widetilde{P_j}$ is bounded in L^p , it is bounded by

$$C_{m,\mu} \left\| \left(\sum_{k \in \mathbb{Z}} \left\| e^{it\Phi(D)} P_k f \right\|_{L^r_t(\mathbb{R})}^2 \right)^{1/2} \right\|_{L^q_x(\mathbb{R}^n)}$$

By the Minkowski inequality we thus have

$$\left\| e^{it\Phi(D)} f \right\|_{L^{q}_{x}(\mathbb{R}^{n}; L^{r}_{t}(\mathbb{R}))} \leq C_{m,\mu} \left(\sum_{k \in \mathbb{Z}} \left\| e^{it\Phi(D)} P_{k} f \right\|_{L^{q}_{x}(\mathbb{R}^{n}; L^{r}_{t}(\mathbb{R}))}^{2} \right)^{1/2}.$$

Apply (2.2) to the right side of the above estimate after parabolic rescaling. Then we obtain

$$\|e^{it\Phi(D)}f\|_{L^{q}_{x}(\mathbb{R}^{n};L^{r}_{t}(\mathbb{R}))} \leq C_{m,\mu} \left(\sum_{k\in\mathbb{Z}} 2^{2k(\frac{n}{2}-\frac{n}{q}-\frac{m}{r})} \|P_{k}f\|_{2}^{2}\right)^{1/2}$$
$$= C_{m,\mu}\|f\|_{\dot{H}^{\frac{n}{2}-\frac{n}{q}-\frac{m}{r}}(\mathbb{R}^{n})}.$$

2.2. Local-to-global arguments. We will show that the global estimate (2.2) is obtained from its local estimate. Adopting the arguments in [19], we consider the dual estimate of (2.2).

Let $S = \{(\xi, \Phi(\xi)) \in \mathbb{R}^n \times \mathbb{R} : 1/2 \le |x| \le 2\}$ be a compact hypersurface with the induced (singular) Lebesgue measure $d\sigma$. We define the Fourier restriction operator \mathfrak{R} for a compact surface S by the restriction of \hat{f} to S, i.e.,

$$\Re f = \tilde{f}|_S$$

Its adjoint operator $\Re^* f = \widehat{fd\sigma}$ can be viewed as $e^{it\Phi(D)}\hat{g}$, where the Fourier transform $\hat{g}(\xi)$ of g corresponds to $f(\xi, \Phi(\xi))$.

Let $\rho > 0$ be the decay of $\widehat{d\sigma}$, i.e.,

$$|\widehat{d\sigma}(x)| \lesssim (1+|x|)^{-\rho}, \qquad x \in \mathbb{R}^{n+1}.$$
(2.3)

It is known that ρ is determined by the number of nonzero principal curvatures of the surface S, which is equal to the rank of the Hessian H_{Φ} . Specifically, if H_{Φ} has rank at least k then

$$\rho = k/2$$

see [17, subsection 5.8, VIII]. From Condition 1.1 we have $k \ge 1$.

When a function f has a compact Fourier support, the $\widehat{f}d\sigma$ decays away from the support of \widehat{f} because of the decay of $\widehat{d\sigma}$. Thus if f and g are compactly Fourier supported and their supports are far away from each other then the interaction between $\widehat{f}d\sigma$ and $\widehat{g}d\sigma$ is negligible.

Definition 2.3. A finite collection $\{Q(z_i, R)\}_{i=1}^N$ of balls in \mathbb{R}^{n+1} with radius R > 0 is called (N, R)-sparse if the centers $\{z_i\}$ are $(NR)^{\gamma}$ -separated where $\gamma := n/\rho \ (\geq 2)$.

From the definition of (N, R)-sparse we have a kind of orthogonality as follows. Let ϕ be a radial Schwartz function which is positive on the ball B(0, 3/2) and $\phi = 1$ on the unit ball B(0, 1) and whose Fourier transform is supported on the ball B(0, 2/3).

Lemma 2.4 ([19, in the proof of Lemma 3.2]). Let $\{Q(z_i, R)\}_{i=1}^N$ be a (N, R)-sparse collection and $\phi_i(z) = \phi(R^{-1}(z-z_i))$ for $i = 1, \dots, N$. Then there is a constant C independent of N such that

$$\left\|\sum_{i=1}^{N} f_i * \hat{\phi}_i \right\|_2 \le C R^{1/2} \left(\sum_{i=1}^{N} \|f_i\|_2^2\right)^{1/2}$$
(2.4)

for all $f_i \in L^2(\mathbb{R}^{n+1})$.

A proof of the above lemma is given in Appendix.

Let $\mathbb{I}_R = (0, R)$ denote an *R*-interval and \mathbb{B}_R the ball of radius *R* centered at the origin in \mathbb{R}^n . Using Lemma 2.4 we have an intermediate result.

Proposition 2.5. Let R > 0 and $1 < q, r \leq 2$. Suppose that there is a constant A(R) such that

$$\|\Re(\chi_{\mathbb{I}_R \times \mathbb{B}_R} f)\|_{L^2(d\sigma)} \le A(R) \|f\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}$$

$$(2.5)$$

for all $f \in L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))$. Then for any (N, R)-sparse collection $\{Q(z_i, R)\}_{i=1}^N$ there is a constant C independent of N such that

$$\|\Re f\|_{L^2(d\sigma)} \le CA(R) \|f\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}$$

$$\tag{2.6}$$

for all f supported in $\cup_{i=1}^{N} Q(z_i, R)$.

Proof. Let $f_i = f \chi_{Q(z_i,R)}$. Then,

$$\Re f_i = \hat{f}_i \big|_S = \widehat{f_i \phi_i} \big|_S = (\hat{f}_i * \hat{\phi}_i) \big|_S,$$

where $\phi_i(z)$ is defined as in Lemma 2.4. Since $\hat{\phi}_i$ is supported on the ball $B(0, \frac{2}{3R})$, we may restrict the support of \hat{f}_i to a O(1/R)-neighborhood of the surface S and write

$$\Re f_i = \left(\hat{f}_i \Big|_{\mathcal{N}_{1/R}(S)} * \hat{\phi}_i \right) \Big|_S$$

where $\mathcal{N}_{1/R}(S)$ is a O(1/R)-neighborhood of the surface S. Let $\mathfrak{\tilde{R}}$ be another restriction operator defined by $\mathfrak{\tilde{R}}f = \hat{f}|_{\mathcal{N}_{1/R}(S)}$. If f is supported in $\bigcup_{i=1}^{N} Q(z_i, R)$, we write

$$\Re f = \sum_{i=1}^{N} (\tilde{\Re} f_i * \hat{\phi}_i) \big|_S.$$

By Lemma 2.4,

$$\|\Re f\|_{L^2(d\sigma)} \le CR^{1/2} \Big(\sum_{i=1}^N \|\tilde{\Re}f_i\|_{L^2(\mathcal{N}_{1/R}(S))}^2\Big)^{1/2}.$$

Since the estimate (2.5) is translation invariant, by a slice argument we have

$$\|\tilde{\mathfrak{R}}f_i\|_{L^2(\mathcal{N}_{1/R}(S))} \le CR^{-1/2}A(R)\|f_i\|_{L^q_x(\mathbb{R}^n;L^r_t(\mathbb{R}))}.$$

By combining the previous two estimates,

$$\|\Re f\|_{L^2(d\sigma)} \le CA(R) \Big(\sum_{i=1}^N \|f_i\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}^2\Big)^{1/2}$$

If $1 \leq r \leq q \leq 2$ then by $\ell^r \subset \ell^q \subset \ell^2$,

$$\begin{split} \left(\sum_{i=1}^{N} \|f_i\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}^2\right)^{1/2} &\leq \left(\sum_{i=1}^{N} \|f_i\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}^q\right)^{1/q} \\ &= \left(\int_{\mathbb{R}^n} \sum_{i=1}^{N} \|f_i\|_{L^r_t(\mathbb{R})}^q dx\right)^{1/q} \\ &\leq \left(\int_{\mathbb{R}^n} \left(\sum_{i=1}^{N} \|f_i\|_{L^r_t(\mathbb{R})}^r\right)^{q/r} dx\right)^{1/q} \\ &= \|f\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}. \end{split}$$

If $1 \le q \le r \le 2$ one can use the embedding $\ell^r \subset \ell^2$ and the Minkowski inequality to get

$$\left(\sum_{i=1}^{N} \|f_i\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}^2\right)^{1/2} \le \left(\sum_{i=1}^{N} \|f_i\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}^r\right)^{1/r}$$
$$\le \left(\int_{\mathbb{R}^n} \left(\sum_{i=1}^{N} \|f_i\|_{L^r_t(\mathbb{R})}^r\right)^{q/r} dx\right)^{1/q}$$
$$= \|f\|_{L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))}.$$

Therefore we have (2.6).

We now extend the (N, R)-sparse sets to the whole space. For this we need the following decomposition lemma.

Lemma 2.6 ([19]). Let E be a subset in \mathbb{R}^n with |E| > 1. Suppose that E is a finite union of finitely overlapping cubes of side-length $c \sim 1$. Then for each $K \in \mathbb{N}$, there are subsets E_1, E_2, \cdots, E_K of E with

$$E = \bigcup_{k=1}^{K} E_k$$

such that each E_k has $O(|E|^{1/K})$ number of $(O(|E|), |E|^{O(\gamma^{k-1})})$ -sparse collections

$$\mathbf{S}_1, \mathbf{S}_2, \cdots, \mathbf{S}_{O(|E|^{1/K})}$$

of which the union $\mathbf{S}_1 \cup \mathbf{S}_2 \cup \cdots \cup \mathbf{S}_{O(|E|^{1/K})}$ is a covering of E_k .

This lemma is a precise version of Lemma 3.3 in [19]. A detailed proof can be found in Appendix.

Using the above lemma we have the following proposition.

Proposition 2.7. Let $1 < q_0, r_0 < \infty$. Suppose that for any $\epsilon > 0$ and any (N, R)-sparse collection $\{Q(z_i, R)\}_{i=1}^N$ in \mathbb{R}^{n+1} , the estimate

$$\|\Re f\|_{L^2(d\sigma)} \le C_{\epsilon} R^{\epsilon} \|f\|_{L^{q_0}_x(\mathbb{R}^n; L^{r_0}_t(\mathbb{R}))}$$

$$\tag{2.7}$$

holds for all f supported in $\bigcup_{i=1}^{N} Q(z_i, R)$. Then for any $1 \le q < q_0$ and $1 \le r < r_0$, the estimate $\|\Re f\|_{L^2(d\sigma)} \le C \|f\|_{L^q(\mathbb{R}^n; L^r_t(\mathbb{R}))}$

holds for all $f \in L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))$.

Proof. By interpolation (see [7]), it suffices to show that for $1 \le q < q_0$ and $1 \le r < r_0$, the restricted type estimate

 $\|\Re\chi_E\|_{L^2(d\sigma)} \le C \|\chi_E\|_{L^q(\mathbb{R}^n;L^r(\mathbb{R}))}$

for all subset E in \mathbb{R}^{n+1} . We may assume |E| > 1, otherwise the estimate is trivial. Since the set S is compact, χ_E can be replaced with $\chi_E * \varphi$, where φ is a bump function supported on a cube of sidelength $c \sim 1$ such that $\hat{\varphi}$ is positive on S. Thus, we may further assume that E is the union of c-cubes.

We denote by $\operatorname{proj}(E)$ the projection of E onto the x-plane. For each grid point $x \in c \mathbb{Z}^n \cap \operatorname{proj}(E)$, we define E_x to be the union of c-cubes in E that intersect $\mathbb{R} \times \{x\}$. Let E^j be the union of E_x which satisfies

 $2^{j-1} <$ the number of *c*-cubes contained in $E_x \leq 2^{j+1}$

for $j \in \mathbb{N}$, (see Figure 1). Then,

$$E = \bigcup_{j \ge 1} E^j.$$

By using Lemma 2.6 with

$$K := \frac{\log(1/\epsilon)}{2\log\gamma} + 1,$$

the E^j is decomposed into E_k^j 's which are covered by $O(|E^j|^{1/K})$ number of $(O(|E^j|), |E^j|^{C\gamma^{k-1}}))$ -sparse collections. We apply (2.7) to these sparse collections and obtain

$$\|\Re\chi_{E_k^j}\|_{L^2(d\sigma)} \le C_{\epsilon} |E^j|^{1/K} (|E^j|^{C\gamma^{k-1}})^{\epsilon} \|\chi_{E_k^j}\|_{L^{q_0}_x(\mathbb{R}^n; L^{r_0}_t(\mathbb{R}))}.$$

Summing over k, we have

$$\begin{aligned} \|\Re\chi_{E^{j}}\|_{L^{2}(d\sigma)} &\leq \sum_{k=1}^{K} \|\Re\chi_{E_{k}^{j}}\|_{L^{2}(d\sigma)} \\ &\leq C_{\epsilon} |E^{j}|^{1/K} (|E^{j}|^{C\gamma^{K-1}})^{\epsilon} \|\chi_{E^{j}}\|_{L_{x}^{q_{0}}(\mathbb{R}^{n};L_{t}^{r_{0}}(\mathbb{R}))} \end{aligned}$$




FIGURE 1. The sets E, projE, E_x and E^j in the proof of Proposition 2.7.

where K is absorbed into C_{ϵ} . Since $|E^j| \leq 2^{j+1} |\operatorname{proj}(E^j)|$, we have

$$\|\Re\chi_{E^j}\|_{L^2(d\sigma)} \le C_{\epsilon} 2^{j(\frac{1}{r_0} + \delta(\epsilon))} |\operatorname{proj}(E^j)|^{\frac{1}{q_0} + \delta(\epsilon)},$$

where

$$\delta(\epsilon) := \frac{1}{K} + C\gamma^{K-1}\epsilon.$$

Since $\lim_{\epsilon \to 0} \delta(\epsilon) = 0$, we can take $\epsilon > 0$ such that

$$0 < \delta(\epsilon) + \epsilon \le \min\left(\frac{1}{q} - \frac{1}{q_0}, \frac{1}{r} - \frac{1}{r_0}\right).$$

Thus,

$$\begin{aligned} \|\Re\chi_E\|_{L^2(d\sigma)} &\leq \sum_{j\geq 1} \|\Re\chi_{E^j}\|_{L^2(d\sigma)} \\ &\leq C_\epsilon \sum_{j\geq 1} 2^{j(\frac{1}{r_0}+\delta(\epsilon))} |\operatorname{proj}(E^j)|^{\frac{1}{q_0}+\delta(\epsilon)} \\ &\leq C \sum_{j\geq 1} 2^{-\epsilon j} 2^{\frac{1}{r}j} |\operatorname{proj}(E^j)|^{\frac{1}{q}} \\ &\leq C \sum_{j\geq 1} 2^{-\epsilon j} \|\chi_E\|_{L^q_x(\mathbb{R}^n;L^r_t(\mathbb{R}))} \\ &\leq C \|\chi_E\|_{L^q_x(\mathbb{R}^n;L^r_t(\mathbb{R}))}. \end{aligned}$$

Combining Proposition 2.5 and Proposition 2.7 we obtain an extension of Tao's epsilon removal lemma as follows.

Proposition 2.8. Let $1 < q_0, r_0 \leq 2$. Suppose that

$$\|\Re(\chi_{\mathbb{I}_R\times\mathbb{B}_R}f)\|_{L^2(d\sigma)} \le C_{\epsilon}R^{\epsilon}\|f\|_{L^{q_0}_x(\mathbb{R}^n;L^{r_0}_t(\mathbb{R}))}$$

for all $f \in L^q_x(\mathbb{R}^n; L^r_t(\mathbb{R}))$.

Now we are ready to prove Theorem 1.2. The theorem follows from Proposition 2.2 and Proposition 2.8 as follows.

2.3. **Proof of Theorem 1.2.** Let P_0 be the Littlewood-Paley projection operator as in subsection 2.1. By rescaling $x \mapsto 2^{-k}x$ and $t \mapsto 2^{-mk}t$, the estimate (1.4) implies

$$\|e^{it\Phi(D)}P_0f\|_{L^{q_0}_x(\mathbb{B}_R;L^{r_0}_t(\mathbb{I}_{R^m}))} \le C_{\epsilon}2^{k\epsilon}\|P_0f\|_{L^2(\mathbb{R}^n)}$$

for all $k \ge 1$ and $\epsilon > 0$. Since $m \ge 1$, we have

$$\|e^{it\Phi(D)}P_0f\|_{L^{q_0}_x(\mathbb{B}_R;L^{r_0}_t(\mathbb{I}_R))} \le C_{\epsilon}2^{k\epsilon}\|P_0f\|_{L^2(\mathbb{R}^n)}.$$

By Proposition 2.8 and duality,

$$\|e^{it\Phi(D)}P_0f\|_{L^q_x(\mathbb{R}^n;L^r_t(\mathbb{R}))} \le C\|P_0f\|_{L^2(\mathbb{R}^n)}$$

By Proposition 2.2, we obtain the desired estimate.

3. Appendix

3.1. Proof of Lemma 2.4. We divide the left side of (2.4) into two parts

$$\|\sum_{i=1}^{N} f_{i} * \hat{\varphi}_{i}|_{S}\|_{2}^{2} = \sum_{i} \|f_{i} * \hat{\varphi}_{i}|_{S}\|_{2}^{2} + \sum_{i \neq j} \int f_{i} * \hat{\varphi}_{i} \overline{f_{j} * \hat{\varphi}_{j}} d\sigma.$$

By a basic restriction estimate we have $||f_i * \hat{\varphi}_i|_S ||_2 \lesssim R^{1/2} ||f_i||_2$. Thus,

$$\sum_{i=1}^{N} \|f_i * \hat{\varphi}_i\|_S \|_2^2 \lesssim R \sum_{i=1}^{N} \|f_i\|_2^2.$$
(3.1)

By Parseval's identity,

$$\int f_i * \hat{\varphi}_i \overline{f_j * \hat{\varphi}_j} d\sigma = \int \overline{\check{f}_j \varphi_j} ((\check{f}_i \varphi_i) * \widehat{d\sigma}),$$

where the *`*denotes the inverse Fourier transform. It is bounded by

$$\left(\sup_{z,w} |\varphi_j^{1/2}(z)\varphi_i^{1/2}(w)\widehat{d\sigma}(z-w)|\right) \|\check{f}_i\varphi_i^{1/2}\|_1 \|\check{f}_j\varphi_j^{1/2}\|_1.$$

By the Cauchy-Schwarz inequality and Plancherel's theorem,

$$\|\check{f}_i\varphi_i^{1/2}\|_1 \lesssim R^{(n+1)/2}\|f_i\|_2.$$

By (2.3),

$$\sup_{z,w} |\varphi_j^{1/2}(z)\varphi_i^{1/2}(w)\widehat{d\sigma}(z-w)| \lesssim |z_i - z_j - 2R|^{-\rho}$$

Since $|z_i - z_j| \ge (NR)^{\gamma}$ and $\gamma \ge 2$, we have that $|z_i - z_j - 2R|$ is comparable to $|z_i - z_j|$. Thus,

$$\sup_{z,w} |\varphi_j^{1/2}(z)\varphi_i^{1/2}(w)\widehat{d\sigma}(z-w)| \lesssim |z_i - z_j|^{-\rho}.$$

Combining these estimates we have

$$\sum_{i \neq j} \int f_i * \hat{\varphi}_i \overline{f_j * \hat{\varphi}_j} d\sigma \lesssim R^{n+1} \sum_{i=1}^N \sum_{j=1}^N |z_i - z_j|^{-\rho} ||f_i||_2 ||f_j||_2$$
$$\lesssim R^{n+1} N \max_{i,j} |z_i - z_j|^{-\rho} \sum_{i=1}^N ||f_i||_2^2.$$

Since $|z_i - z_j| \ge (NR)^{\gamma} \ge N^{\frac{1}{\rho}} R^{\frac{n}{\rho}}$, it follows that

$$\sum_{i \neq j} \int f_i * \hat{\varphi}_i \overline{f_j * \hat{\varphi}_j} d\sigma \lesssim R \sum_{i=1}^N \|f_i\|_2^2$$

From the above estimate and (3.1) we obtain (2.4).

3.2. **Proof of Lemma 2.6.** Fix $K \in \mathbb{N}$. We define $R_0 = 1$ and R_k for $k = 1, 2, \dots, K$ recursively by

$$R_k = |E|^{\gamma} R_{k-1}^{\gamma}. \tag{3.2}$$

From this definition we have $R_k = |E|^{\frac{\gamma^{k+1}-\gamma}{\gamma-1}}$. Let $E_0 = \emptyset$. We define E_k for $k = 1, 2, \dots, K$ to be the set of all $x \in E \setminus \bigcup_{j=0,1,2,\dots,k-1} E_j$ such that

$$|E \cap B(x, R_k)| \le |E|^{k/K}.$$
(3.3)

Then, $E = \bigcup_{k=1}^{K} E_k$. From this construction it follows that for $x \in E_k$, $k = 2, 3, \dots, K$,

$$|E \cap B(x, R_{k-1})| > |E|^{(k-1)/K}.$$
(3.4)

We cover E_k with finitely overlapping R_k -balls $\mathbf{C}_{E_k} := \{B_i = B(x_i, R_k) : x_i \in E_k\}$. Since E is a finite union of cubes of side-length $c \sim 1$, it is obvious that $\#\mathbf{C}_{E_k} \leq |E|$. For each $B_i \in \mathbf{C}_{E_k}$ we cover $E_k \cap B_i$ with finitely overlapping R_{k-1} -balls $\mathbf{C}_{E_k \cap B_i} := \{B'_{ij} = B'(y_j, R_{k-1}) : y_j \in E_k \cap B_i\}$, that is,

$$E_k \cap B_i = \bigcup_{B'_{ij} \in \mathbf{C}_{E_k \cap B_i}} E_k \cap B'_{ij}$$

Since $((E \setminus E_k) \cap B'_{ij}) \subset ((E \setminus E_k) \cap B_i)$ for all j, we have

$$(E_k \cap B_i) \cup ((E \setminus E_k) \cap B_i) \supset \bigcup_{B'_{ij} \in \mathbf{C}_{E_k \cap B_i}} (E_k \cap B'_{ij}) \cup ((E \setminus E_k) \cap B'_{ij}),$$

thus

$$E \cap B_i \supset \bigcup_{B'_{ij} \in \mathbf{C}_{E_k \cap B_i}} E \cap B'_{ij}.$$

By finitely overlapping,

$$\#\mathbf{C}_{E_k\cap B_i} \lesssim \max_{B'_{ij}\in\mathbf{C}_{E_k\cap B_i}} \frac{|E\cap B_i|}{|E\cap B'_{ij}|}.$$

By (3.3) and (3.4) the above is bounded by $C|E|^{1/K}$, and we have $\#\mathbf{C}_{E_k\cap B_i} \leq C|E|^{1/K}$ for all *i*. Thus,

$$E_k \subset \bigcup_{i=1}^{O(|E|)} \bigcup_{j=1}^{O(|E|^{1/K})} B'_{ij}.$$

We choose $O(R_k)$ -separated balls $\{B'_{ij(i)}\}_{i=1}^{O(|E|)}$. Then it becomes a $(O(|E|), R_{k-1})$ -sparse collection because of (3.2). Since $R_{k-1} = |E|^{O(\gamma^{k-1})}$ and every $B_i \in \mathbf{C}_{E_k}$ has the covering

 $\mathbf{C}_{E_k \cap B_i}$ of cardinality $O(|E|^{1/K})$, there are $O(|E|^{1/K})$ number of $(O(|E|), |E|^{O(\gamma^{k-1})})$ -sparse collections $\mathbf{S}_1, \mathbf{S}_2, \cdots, \mathbf{S}_{O(|E|^{1/K})}$ such that

$$E_k \subset \bigcup_{j=1}^{O(|E|^{1/K})} \bigcup_{B' \in \mathbf{S}_j} B'.$$

References

- [1] T. Cazenave, Semilinear Schrödinger Equations, Vol. 10, AMS, 2003.
- [2] Y. Cho, G. Hwang, S. Kwon, and S. Lee, Well-posedness and ill-posedness for the cubic fractional Schrödinger equations, Discrete Contin. Dyn. Syst. 35 (2015), 2863–2880.
- [3] C. Cho and H. Ko, A note on maximal estimates of generalized Schrödinger equation, arXiv:1809.03246.
- [4] Y. Cho, T. Ozawa, and S. Xia, Remarks on some dispersive estimates, Commun. Pure Appl. Anal. 10 (2011), 1121–1128.
- [5] X. Du, L. Guth, and X. Li, A sharp Schrödinger maximal estimate in ℝ², Ann. Math. 186 (2017), no. 2, 607–640.
- [6] X. Du, J. Kim, H. Wang, and R. Zhang, Lower bounds for estimates of the Schrodinger maximal function, Math. Res. Lett., to appear. 27 (2020), no. 3, 687–692.
- [7] D. L. Fernandez, Lorentz spaces, with mixed norms, J. Funct. Anal. 25 (1977), no. 2, 128–146.
- [8] J. Ginibre and G. Velo, Generalized Strichartz inequalities for the wave equation, J. Funct. Anal. 133 (1995), 50–68.
- [9] M. Keel and T. Tao, Endpoint Strichartz estimates, Amer. J. Math. 120 (1999), no. 5, 955–980.
- [10] C. E. Kenig, G. Ponce, and L. Vega, Oscillatory integrals and regularity of dispersive equations, Indiana Univ. Math. J. 40 (1991), no. 1, 33–69.
- [11] S. Lee, K. M. Rogers, and A. Vargas, An endpoint space-time estimate for the Schrödinger equation, Adv. Math. 226 (2011), no. 5, 4266–4285.
- [12] B. Pausader, Global well-posedness for energy critical fourth-order Schrödinger equations in the radial case, Dyn. Partial Differ. Equ. 4 (2007), 197–225.
- [13] H. Pecher, Dispersive estimates and the 2D cubic NLS Nonlinear small data scattering for the wave and Klein-Gordon equation, Math. Z. 185 (1984), no. 2, 261–270.
- [14] F. Planchon, Dispersive estimates and the 2D cubic NLS equation, J. Anal. Math. 86 (2002), 319–334.
- [15] K. M. Rogers, Strichartz estimates via the Schrödinger maximal operator, Math. Ann. 343 (2), no. 3, 603–622.
- [16] K. M. Rogers and P. Villarroya, Sharp estimates for maximal operators associated to the wave equation, Ark. Mat. 46 (2008), no. 1, 143–151.
- [17] E. M. Stein, Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, Vol. 3, Princeton University Press, 1993.
- [18] R.S. Strichartz, Restriction of Fourier transform to quadratic surfaces and decay of solutions of wave equations, Duke Math. J. 44 (1977), 705–774.
- [19] T. Tao, The Bochner-Riesz conjecture implies the restriction conjecture, Duke Math. J. 96 (1999), no. 2, 363–375.
- [20] _____, Nonlinear Dispersive Equations: Local and Global Analysis, Vol. 106, AMS, 2006.
- [21] L. Vega, El multiplicador de Schrödinger. La funcion maximal y los operadores de restricción, 1998.

DEPARTMENT OF MATHEMATICAL SCIENCES AND RIM, SEOUL NATIONAL UNIVERSITY, SEOUL 151–747, RE-PUBLIC OF KOREA

Email address: akilus@snu.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION, KONGJU NATIONAL UNIVERSITY, KONGJU 32588, REPUBLIC OF KOREA

Email address: ywkoh@kongju.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, SCHOOL OF NATURAL SCIENCE, ULSAN NATIONAL INSTITUTE OF SCIENCE AND TECHNOLOGY, UNIST-GIL 50, ULSAN 44919, REPUBLIC OF KOREA

Email address: jungjinlee@unist.ac.kr