Self-normalizing Path Integrals

Ivan M. Burbano*1 and Francisco Calderón^{†2}

¹Department of Physics, University of California, 366 Physics North MC 7300, Berkeley, CA 94720-7300, USA ²Department of Philosophy, University of Michigan, 435 South State Street 2215 Angell Hall, Ann Arbor, MI 48109-1003, USA

September 3, 2021

Abstract

We address the problem of computing the overall normalization constant of path integrals using zeta-function regularization techniques. In particular, we study a phenomenon we called "self-normalization," in which the ambiguity of the integral measure, which would typically need to be renormalized, resolves itself. Hawking had already detected this phenomenon in the context of Gaussian integrals. However, our approach extends Hawking's work for the cases in which the space of fields is not a vector space but instead has another structure which we call a "linear foliation." After describing the general framework, we work out examples in one (the transition amplitudes and partition functions for the harmonic oscillator and the particle on a circle in the presence of a magnetic field) and two (the partition functions for the massive and compact bosons on the torus and the cylinder) spacetime dimensions in a detailed fashion. One of the applications of our results, explicitly shown in the examples, is the computation of the overall normalization of path integrals that do not self-normalize. That is usually done in the literature using different comparison methods involving additional assumptions on the nature of this constant. Our method recovers the normalization without the need for those extra assumptions.

Contents

| 1 | Introduction | 2 |
|---|--------------|---|
| | | |

5

2 Gaussian Path Integrals

^{*}ivan_burbano@berkeley.edu

[†]fcalder@umich.edu

| 3 | Linear Foliations | | | | | |
|---|-----------------------------------|--------|-----------------------|----|--|--|
| 4 | Examples in 1D | | | | | |
| | 4.1 | Harmo | onic Oscillator | 10 | | |
| | | 4.1.1 | Partition Function | 11 | | |
| | | 4.1.2 | Transition Amplitudes | 12 | | |
| | 4.2 Particle on the Circle | | | | | |
| | | 4.2.1 | Partition Function | | | |
| | | 4.2.2 | Transition Amplitudes | 18 | | |
| 5 | Examples in 2D | | | | | |
| | 5.1 | Free B | Boson | 19 | | |
| | | 5.1.1 | On the Torus | 20 | | |
| | | 5.1.2 | On the Cylinder | 21 | | |
| | 5.2 | Compa | oact Boson | | | |
| | | 5.2.1 | On the Torus | | | |
| | | 5.2.2 | On the Cylinder | 25 | | |
| 6 | Conclusions and Open Problems 2 | | | | | |
| 7 | Acknowledgments | | | | | |
| A | Useful Values of Zeta Functions 2 | | | | | |

1 Introduction

The path integral is by now a ubiquitous tool in both physics and mathematics. At a superficial level, it is simple to give an intuitive account of their meaning. Consider some N-dimensional manifold $\mathcal E$ equipped with a volume form called the integral measure. In a coordinate system ϕ , said form can be written as $\mathrm{d}^N \phi \, e^{-\frac{1}{\hbar} S(\phi)}$ for some function S on $\mathcal E$ called the action. Then, we know how to make sense of integrals of the form

$$\int_{\mathcal{E}} d^N \phi \, e^{-\frac{1}{\hbar}S(\phi)} \mathcal{O}(\phi), \tag{1}$$

for functions \mathcal{O} on \mathcal{E} known as insertions. These, in a certain sense, sum up the values of \mathcal{O} along \mathcal{E} by weighing them against the volume form. Of course, the precise meaning of the integral depends on what said form is. For example, if the form describes the infinitesimal size around a point of \mathcal{E} , then the integral corresponds to the volume enclosed by the graph of \mathcal{O} . On the other hand, if the form corresponds to some probability measure along \mathcal{E} , then the integral yields the expectation value of \mathcal{O} .

Path integrals attempt to generalize these to the situation where \mathcal{E} is a space of fields on some manifold M. These are usually taken to be sections of some fiber bundle on M. In

particular, in path integrals \mathcal{E} is infinite-dimensional, and the standard theory of Lebesgue integration on finite-dimensional manifolds breaks down.

Before we delve into the difficulties behind defining such integrals, let us briefly mention some of their applications for the reader that may be just starting to get acquainted with them. The most obvious is in statistical field theory, where physical systems consist of fields on a spatial manifold. In this case, the path integral measure describes the ensemble's probability measure, and the path integral yields expectation values of the observables of the theory.

As discovered by Feynman [1], path integrals also play an important role in quantum mechanics. In this case, the quantum evolution of a system of fields is obtained by considering all the possible trajectories it could traverse. Each such trajectory contributes an infinitesimal amplitude, which has to be summed up to obtain the full amplitude of the quantum process. In this case, M corresponds to the Lorentzian spacetime manifold of the theory, and the path integral measure describes the infinitesimal contributions of each trajectory.

In quantum mechanics, there is an intimate relationship between the evolution of a system and its thermal properties, as formalized, for example, by Tomita-Takesaki theory and its relation to KMS states [2]. In some spacetimes, this relation allows one to compute the thermal properties of a quantum system by performing path integrals using Euclidean versions of the spacetime manifold [3]. These integrals are often of the same form as those encountered in systems of statistical field theory. This relationship has led to an important array of maps relating the thermal properties of quantum theories in d-1 spatial dimensions to the thermal properties of classical theories in d spatial dimensions [4]. Analogous modifications of the spacetime manifold have provided expressions for other important properties of a quantum system, such as out-of-time-order correlators and out-of-equilibrium quantities within the Schwinger-Keldysh formalism [5].

Path integrals have also played an important role in pure mathematics. For example, they have found applications in geometry because the path integral contains information of the manifold M and the structure on M required to define the path integral measure. One of the earliest uses of this idea was developed by Witten [6], who realized that if one uses an integration measure that only depends on the topology of M, the integral should yield topological invariants. This method has been used, for example, to compute knot invariants. It is also a key part of the applications of string theory to mathematics [7].

We hope that the reader is now convinced that path integrals are an important and versatile tool in physics and mathematics. However, their applications are likely to extend far beyond these fields. For example, path integrals have already become an important tool in economics [8]. Still, to this day, no one understands how to give a precise definition of a path integral in full generality. The standard measure-theoretic tools have only found limited success in this task [9]. In fact, it is believed that a satisfactory definition of a path integral will require truly novel mathematics that go beyond the current techniques of measure theory.

One of the additional inputs required to define the path integral is the idea of renormalization. In the currently available methods, if one attempts to compute the path integral using some measure coming from a classical theory, one often runs into undesired infinities. These must be resolved by choosing additional quantum corrections to the path integral method, thus appealing to renormalization techniques. From the quantum point of view, this is expected, for it is clear that no quantization map assigns unequivocally to each classical theory some preferred quantum theory that underlies it [10]. These corrections to the measure are usually expressed as \hbar -dependent corrections to the action and $\mathcal{D}\phi$, the infinite-dimensional analog of $\mathrm{d}^N\phi$.

Several steps have been taken towards giving a mathematically complete theory of path integrals that incorporate this phenomenon. For example, the work of Costello and collaborators [11, 12, 13] has given a mathematical framework for the computation of the perturbative aspects of these integrals, using Feynman diagrammatics. In this framework, a theory is defined by a collection of effective actions that can describe physics up to a particular scale. The renormalization group equation then relates physics at different scales. The absence of anomalies in the quantum path integral measure obtained through these actions is then expressed using the Batalin-Vilkovisky master equation.

In this paper, we will, however, take other complementary and somewhat orthogonal steps towards the definition of a path integral measure. We will consider path integrals corresponding only to free field theories. The simplicity of these theories places them among the few where we can currently make non-perturbative calculations, making them a perfect case study for computing the overall normalization of their path integrals. This issue is commonly disregarded in the literature, given that some of the initial applications of the path integral formalism, particularly in particle physics in conjunction with the LSZ formalism, were independent of this normalization [14]. This has led, at times, to the incorrect belief that the path integral formulation is not powerful enough to compute said normalizations. This is, however, not true. As discussed by Hawking [15], within the path integral formalism, this overall normalization constitutes one of the corrections required in order to achieve a consistent quantization. Accordingly, like the rest of the renormalization program, fixing these requires the introduction of renormalization conditions. Moreover, generic calculations in quantum mechanical systems do require knowledge of these normalizations, as exhibited, for example, by the Schwinger-Keldysh formalism.

This paper came about from the study of the book *Mirror Symmetry* [16]. Several elementary applications of the path integral formalism are exhibited in it, many of which will be revisited in sections 4 and 5. They do not mention the renormalization conditions required for the overall normalization of the integrals. However, their final results somehow end up with the correct normalizations. We will call this phenomenon "self-normalization." In comparison, the application of the same techniques for other examples does not yield the correct overall normalization [17]. We then embarked on the task of characterizing in which examples the phenomenon of self-normalization is found.

As we later found out, Hawking had already given a framework to understand this phenomenon by carefully tracking the renormalization constants involved. He gave a generic condition that explained when a path integral self-normalizes which works for free theories based on a space of fields equipped with a vector structure. In this work, we generalize said

framework to more general spaces of fields. Namely, we relax the linear structure from that of a vector space to what we call a "linear foliation." Heuristically, spaces \mathcal{E} that admit such a linear foliation can be split into leafs that are affine. Accordingly, the path integrals over each leaf can be computed using the standard methods. Moreover, we require that the leafs are bundled together by a finite-dimensional manifold, called the "stem." The key realization then is that the integral over the stem is also affected by quantum corrections. This procedure provides a new self-normalization method where neither the integrals over the leafs nor the integral over the stem self-normalize, but the overall integral does.

In section 2, we give a self-contained review on the computation of free path integrals whose spaces of fields have a vector space structure. We introduce zeta-function regularization there, which will play a major role throughout this paper. We follow Hawking's heuristic treatment and review his self-normalization method. In section 3, we extend this framework to field spaces that admit a linear foliation. We continue using heuristic arguments in order to motivate the precise definition of a linear foliation. We then explain how zeta-function regularization can also be used to define a path integral measure on these foliations. Carefully tracking the associated renormalization constants allows us to generalize Hawking's self-normalization method to these cases. Section 4 then gives concrete examples of these ideas in the case of one-dimensional field theories. The examples treated are the computation of the partition functions and transition amplitudes of the harmonic oscillator and the particle on a circle in the presence of a magnetic field. Section 5 then treats the analogs of these examples in the two-dimensional case. We explore massive and compact bosons on the torus and the cylinder.

For our treatment of the zeta-function regularization techniques necessary to regularize the path integrals studied in this paper, we follow Elizalde; notably, [18]. This work allowed us to do the calculations more rigorously than what can be commonly found in the literature, which sometimes contains erroneous formulas. Among these are $\prod_k \lambda_k \mu_k = \prod_k \lambda_k \prod_k \mu_k$, which obviates the possibility of a multiplicative anomaly. Another one is $\prod_{k \in \mathbb{Z}} \mu = 1$, which is not only not true, even within the realm of zeta-function regularization, but would also trivialize this whole paper.

We want to finish this introduction by noting that we have tried to keep the prerequisites at a minimum and the exposition self-contained. In particular, although we use the language of quantum mechanics throughout to give some physical intuition, very few prerequisites in quantum or statistical physics are required. We do expect the reader to be comfortable with certain notions in linear algebra and differential geometry. However, we have tried our best to stir away from considerations of an analytic nature, making the paper fully algebraic in essence. Therefore, we hope this work will gently introduce an important part of the path integral toolbox to the more mathematically inclined reader.

2 Gaussian Path Integrals

We will now give a self-contained review of the main computation scheme as described in [15]. Let \mathcal{E} be a **vector space** of fields on a spacetime M denoted by ϕ . We will equip \mathcal{E}

with a bilinear form $\langle \cdot, \cdot \rangle$ and a differential operator Q which is self-adjoint with respect to the bilinear form. We will first consider path integrals of the Gaussian form

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)},\tag{2}$$

for actions $S(\phi)$ which, although classically of the form $\frac{1}{2}\langle\phi,Q\phi\rangle$, may require further quantum corrections. This basic structure appears for example in the theory of free Euclidean fields [see e. g. 11, Definition 7.0.1]. The scheme depends crucially on the assumption that Q admits a countable enumeration of a complete set of eigenfields $\{\phi_n \in \mathcal{E} \mid n \in I\}$ which are orthogonal with respect to the bilinear form. We will denote the eigenvalues by λ_n

$$Q\phi_n = \lambda_n \phi_n. \tag{3}$$

Notice that, generically, this will not be a Hamel basis for \mathcal{E} . In fact, since we will not need to equip \mathcal{E} with a topology, we will not ponder on whether it is a Schauder basis or not. Completeness in this setting means that this set contains a basis for every eigenspace of Q.

With this set-up, we can now imagine how the computation would be performed if the set of eigenfields happened to behave as a Schauder basis, i. e., if we were able to express every $\phi \in \mathcal{E}$ as a linear combination of the list above

$$\phi = \sum_{n \in I} c_n \phi_n. \tag{4}$$

We will not worry whether the expression above is well-defined since we only need it to motivate the computation scheme. Then the quadratic form in the integrand would take the form

$$\langle \phi, Q\phi \rangle = \sum_{n \in I} \lambda_n c_n^2. \tag{5}$$

We are then compelled to try to make sense of the integration measure as a measure induced by the bilinear form

$$\mathcal{D}\phi = \prod_{n \in I} \mathrm{d}c_n \,. \tag{6}$$

As far as the quantum corrections to the classical action, the quadratic nature of our ansatz guarantees that no complicated divergences are arising from Feynman diagrams. Accordingly, the quantum corrections appear as constant (or vacuum [14]) counterterms to the action [19]

$$S(\phi) = \frac{1}{2} \langle \phi, Q\phi \rangle + \hbar V. \tag{7}$$

Equivalently, we can incorporate said counterterm as a correction to the measure by redefining

$$\mathcal{D}\phi = \left(\prod_{n \in I} dc_n\right) e^{-V}.$$
 (8)

We can then attempt to make sense of this measure by distributing this extra exponential as a rescaling of the factors. For this we introduce a renormalization constant μ_n for each mode¹

$$\mathcal{D}\phi = \prod_{n \in I} \frac{\mathrm{d}c_n}{\sqrt{2\pi\hbar\mu_n}}.$$
 (9)

These renormalization constants are then related to V by

$$V = \frac{1}{2} \sum_{n \in I} \ln(2\pi\hbar\mu_n) \tag{10}$$

In particular, the simplest choice is to distribute the correction to the measure uniformly so that there is a single renormalization constant $\mu_n = \mu$. We will prefer this latter point of view in which we think of the required renormalization procedure as a correction to the measure instead of a correction to the action.

The choice of setting all $\mu_n = \mu$ makes (10) has a natural interpretation from the point of view of Gaussian integration. Indeed, comparison with the finite-dimensional Gaussian integral

$$\int d^{D} \left(\frac{x}{\sqrt{2\pi\hbar\mu}} \right) e^{-\frac{1}{2\hbar}x \cdot Qx} = \det(\mu Q)^{-1/2}, \tag{11}$$

suggests we set

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{2\hbar}\langle\phi,Q\phi\rangle} = \int \prod_{n\in I} d\left(\frac{c_n}{\sqrt{2\pi\hbar\mu}}\right) e^{-\frac{1}{2\hbar}\sum_{n\in I}\lambda_n c_n^2} = \det(\mu Q)^{-1/2}. \tag{12}$$

We have now motivated the reduction of the computation of path integrals of the form (2) to that of defining the determinant of μQ . Here is where zeta-function regularization comes in. We first try to make sense of the determinant as an infinite product

$$\det(\mu Q) = \prod_{n \in I} \mu \lambda_n. \tag{13}$$

In order to define the latter, we will consider the zeta function associated to Q

$$\zeta_Q(s) := \sum_{n \in I} \lambda_n^{-s}. \tag{14}$$

Under certain conditions [20], this is well defined for Re s big enough and can be analytically extended to a meromorphic function on the whole complex plane \mathbb{C} . In particular, the formal computation

$$\zeta_Q'(s) = -\sum_{n \in I} \lambda_n^{-s} \ln \lambda_n, \tag{15}$$

¹In [15] Hawking uses a different normalization for the regularization constants. We prefer this one since it simplifies the formulas below.

suggests we define

$$\det(Q) := e^{-\zeta_Q'(0)}. (16)$$

Given that $\zeta_{\mu Q}(s) = \mu^{-s} \zeta_Q(s)$, the same logic yields

$$\det(\mu Q) = e^{-\zeta'_{\mu Q}(0)} = \mu^{\zeta_Q(0)} e^{-\zeta'_Q(0)} = \mu^{\zeta_Q(0)} \det(Q). \tag{17}$$

Along with (12), this gives a definition for our initial path integral (2).

In general, the result of the path integral using the scheme above depends on μ . One must then device external consistency conditions, known as renormalization conditions, to fix the value of μ . A way of doing this would be to compare the result with that obtained from operator methods. However, the path integral method truly shines when a computation through operator methods is not feasible.

Another way of doing this would be to find a limit of the parameters in which the path integral is already known. Then, assuming μ is independent of the parameters used to achieve this limit, one can use the known result to fix its value. We will call this the **comparison method**. Since the μ -dependent part of (17) is always an overall normalization, this renormalization condition is usually interpreted via the claim that zeta-function regularization only computes the quotient between the determinant one is interested in and the one that is known. A useful set of limits are those in which the path integral corresponds to a theory with a simple operator formulation that can be used to fix μ [21]. Another particularly useful limit is to consider the integral in an infinitesimal interval of time. For example, when the path integral corresponds to a transition amplitude, one expects that it will reduce to a delta function in this limit [19]. In particular, this limit does not require knowing the operator formulation of a related theory.

Another method that does not require operator methods is to demand that these path integrals compose correctly. Here one compares the path integral on a spacetime to the path integrals obtained by cutting said spacetime. We will call these **cut-paste conditions**. This procedure is useful, for example, when the original path integral and their cut versions are of the same type. One can then invoke consistency to fix the value of μ [21]. This method is particularly powerful because it does not require any underlying assumptions on the behavior of μ in a particular limit of the theory.

We have yet to consider the possibility that the path integral is independent of μ . In this case, one does not need a renormalization condition. From (17) one sees that, for the path integrals we have considered so far, this happens precisely when $\zeta_Q(0) = 0$. Only after checking this are we allowed to set $\mu = 1$ consistently throughout the computation. We say that these path integrals **self-normalize** since no renormalization condition is required. A particularly useful consequence of such integrals is that one can then cut them and use the cut-paste renormalization condition discussed above to renormalize other path integrals.

3 Linear Foliations

In applications to physics, one often finds that the path integrals do not appear in the Gaussian form (2). For example, they may have additional insertions required to compute

correlation functions or interactions appearing as non-quadratic terms in the action. There is a well-developed theory designed to handle these cases in perturbation theory, often using Feynman diagrams. Some modern mathematical accounts of these methods can be found in [11, 12, 13, 22]. Although there are still several unanswered questions regarding these deformations of the Gaussian path integral, particularly regarding their non-perturbative aspects, we will not focus on them in this paper. Instead, we will look at path integrals of the form

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)},\tag{18}$$

where the action S is still quadratic in the fields, but \mathcal{E} is no longer a vector space.

There are many reasons why these path integrals arise in physics. In our examples, there will be two main ones. On the one hand, the path integrals may be defined on spacetimes with boundaries. In these, we need to prescribe boundary conditions, most of which spoil the linearity of the fields. On the other, the fields themselves may take values on topologically non-trivial spaces. Then the non-linearity of these target spaces induces non-linearity in the field content of the integral.

In order to deal with these integrals, we will restrict to spaces \mathcal{E} that can be written as

$$\mathcal{E} \cong \mathcal{J} \times \mathcal{F},\tag{19}$$

where \mathcal{J} is a finite-dimensional manifold and \mathcal{F} is a vector space of quantum fluctuations. We will call this decomposition a **linear foliation** of \mathcal{E} . We will call \mathcal{J} the **stem** of the foliation. Since \mathcal{F} is a vector space, the stem can be identified as a submanifold of \mathcal{E} via $\mathcal{J} \cong \mathcal{J} \times \{0\}$. Sets of the form $\{p\} \times \mathcal{F}$ will be called **leafs** and will be assumed to be affine spaces modelled on \mathcal{F} .

We would then like to use this decomposition to induce a measure on \mathcal{E}

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)} = \int_{\mathcal{J}} \mathrm{d}s \int_{\{s\} \times \mathcal{F}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}. \tag{20}$$

In order to understand this, we will begin by studying the behavior of the action on each leaf. Here is where the affine structure comes in. Let us fix $s \in \mathcal{J}$ and a field $\phi_s \in \{s\} \times \mathcal{F}$. We can then write every $\phi \in \{s\} \times \mathcal{F}$ in the form $\phi = \phi_s + \tilde{\phi}$ for some unique fluctuation $\tilde{\phi} \in \mathcal{F}$ around ϕ_s . Using this decomposition we can expand in a Taylor series

$$S(\phi) = S(\phi_0) + \int dx \left. \frac{\delta S}{\delta \phi(x)} \right|_{\phi_s} \tilde{\phi}(x) + \int dx \, dy \, \tilde{\phi}(x) \left. \frac{\delta^2 S}{\delta \phi(x) \delta \phi(y)} \right|_{\phi_s} \tilde{\phi}(y), \tag{21}$$

called the **semi-classical expansion**, which in this case terminates at the quadratic term given that the action S was quadratic to begin with. We will then attempt to choose ϕ_s such that the linear term vanishes. For variations vanishing on the boundaries of the spacetime M, this means we need to choose a ϕ_s that satisfies the equations of motion. This will be called an **instanton** of the leaf. In fact, in practice, we will usually begin by choosing a set of instantons $\phi_s \in \mathcal{E}$ for the theory. The space \mathcal{J} will then be chosen as the space of

such instantons. We then choose a vector space of fluctuations \mathcal{F} such that the affine spaces $\phi_s + \mathcal{F}$ yield the leafs for an admissible foliation (19). The vanishing of the linear part of the action also imposes non-trivial conditions on the boundary behaviour of the fields on \mathcal{F} .

Having eliminated the linear part, we equip \mathcal{F} with a differential operator Q and a bilinear form $\langle \cdot, \cdot \rangle$, so that

$$S(\phi) = S_s + \frac{1}{2} \left\langle \tilde{\phi}, Q\tilde{\phi} \right\rangle, \tag{22}$$

where $S_s := S(\phi_s)$ is the action of the instanton. The measure is then completely fixed by choosing some vector space \mathcal{V} which extends \mathcal{F} , the bilinear form, and the differential operator. This space has to be small enough so that the condition on the self-adjointness and the spectrum of Q are satisfied on \mathcal{V} . However, it has to be big enough so that the connected components of the foliation, which are completely determined by the connected components of the stem, are subsets of affine subspaces modeled after \mathcal{V} . Then, the measure (6) on \mathcal{V} induces a measure on the foliation \mathcal{E} . The net result of these choices is that we can write

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)} = \int_{\mathcal{J}} \mathrm{d}s \, e^{-\frac{1}{\hbar}S_s} \int_{\mathcal{F}} \mathcal{D}\tilde{\phi} \, e^{-\frac{1}{2\hbar}\left\langle \tilde{\phi}, Q\tilde{\phi} \right\rangle} = \int_{\mathcal{J}} \mathrm{d}s \, e^{-\frac{1}{\hbar}S_s} \det(\mu Q)^{-1/2}. \tag{23}$$

Here, we assume that the integration measure in field space is translation invariant $\mathcal{D}\phi = \mathcal{D}\tilde{\phi}$. These path integrals admit a new self-normalization method. Namely, since the measure on \mathcal{V} depends on μ , and this is used to induce a measure on the whole foliation $\mathcal{J} \times \mathcal{F}$, both contributions to (23) depend on μ . Then, even if the Gaussian integral does not self-normalize by itself, it can be compensated by the stem integral. This happens precisely when the μ -dependence of the stem integral is

$$\int_{\mathcal{I}} ds \, e^{-\frac{1}{\hbar}S_s} \propto \mu^{\frac{1}{2}\zeta_Q(0)}. \tag{24}$$

4 Examples in 1D

We will now exhibit the formalism above by considering examples in one and two dimensions. We will begin with the one-dimensional case since it already exhibits the key points of the formalism, and the computations are much simpler. Quantum field theory in 1D is also known as quantum mechanics. In it, the fields are simply the dynamical variables that are a function of time. In particular, the role of spacetime is played solely by time.

4.1 Harmonic Oscillator

We begin with the important example of the quantum harmonic oscillator. To follow the conventions above, we will stick to the study of the quantum statistics of this system. Therefore, fields are defined over one-dimensional Euclidean time M. They are given by maps

 $x \in C^{\infty}(M)$, which assign to every $\tau \in M$ the position of the oscillator $x(\tau)$. The space of all such fields forms a vector space with bilinear form

$$\langle x, y \rangle = \int d\tau \, xy,$$
 (25)

and a differential operator

$$Q = -\frac{\mathrm{d}^2}{\mathrm{d}\tau^2} + \omega^2 \tag{26}$$

The action in this case takes the form

$$S(x) = \int_{M} d\tau \left(\frac{1}{2}\dot{x}^{2} + \frac{1}{2}\omega^{2}x^{2}\right), \tag{27}$$

with ω the angular frequency of the oscillator. Since the mass term would mutiply the whole action, we will absorb it into \hbar for simplicity.

4.1.1 Partition Function

Let us begin by exploring the partition function of this system. This will be an example of a path integral that self-normalizes using the Hawking mechanism in which $\zeta_Q(0) = 0$. This example should be compared with the calculation in [16], in which they assume from the beginning that μ can be set to 1.

The partition function of the quantum harmonic oscillator at inverse temperature β is given by the path integral (18) with $M = S^1_{\hbar\beta}$, the circle of circumference $\hbar\beta$, and $\mathcal{E} = C^{\infty}(M)$. We will denote this by

$$\bigcirc = \int_{\mathcal{E}} \mathcal{D}x \, e^{-\frac{1}{\hbar}S(x)}. \tag{28}$$

 \mathcal{E} is already a vector space and we can immediately integrate by parts the action (27) to obtain $S(x) = \frac{1}{2}\langle x, Qx \rangle$. Since M has no boundary, there are no boundary terms to worry about. Accordingly, this is a Gaussian integral.

The solutions to the eigenvalue problem

$$\left(-\frac{\mathrm{d}^2}{\mathrm{d}\tau^2} + \omega^2\right)x = \lambda x\tag{29}$$

are generated by the orthonormal set

$$x_n(\tau) = \frac{1}{\sqrt{\hbar\beta}} e^{\frac{2\pi i n}{\hbar\beta}\tau}, \quad n \in \mathbb{Z}, \tag{30}$$

with eigenvalues

$$\lambda_n = \frac{4\pi^2 n^2}{\hbar^2 \beta^2} + \omega^2. \tag{31}$$

Then, for each² $n \in \mathbb{N}$, the eigenvalue λ_n has multiplicity 2 while λ_0 has multiplicity 1.

We can now compute the zeta function associated to Q. For this we will need the Epstein-Hurwitz zeta function and its limits described in the appendix A. By splitting the integers into positive, negative (which contribute the same as the positive) and 0, we obtain

$$\zeta_Q(s) = \omega^{-2s} + 2\sum_{n=1}^{\infty} \left(\frac{4\pi^2 n^2}{\hbar^2 \beta^2} + \omega^2\right)^{-s}
= \omega^{-2s} + 2\left(\frac{4\pi^2}{\hbar^2 \beta^2}\right)^{-s} \zeta_{EH}\left(s; \frac{\hbar^2 \beta^2 \omega^2}{4\pi^2}\right).$$
(32)

Using the result (135), we have $\zeta_Q(0) = 1 + 2(-1/2) = 0$ and the path integral self-normalizes via the method of section 2. We can further compute using (135) and (140)

$$\zeta_Q'(0) = -2\ln(\omega) - 2\ln\left(\frac{4\pi^2}{\hbar^2\beta^2}\right)\zeta_{EH}\left(0; \frac{\hbar^2\beta^2\omega^2}{4\pi^2}\right) + 2\frac{\mathrm{d}}{\mathrm{d}s}\zeta_{EH}\left(s; \frac{\hbar^2\beta^2\omega^2}{4\pi^2}\right)\Big|_{s=0}
= \ln\left(\frac{4\pi^2}{\hbar^2\beta^2\omega^2}\right) - 2\ln\left(\frac{4\pi}{\hbar\beta\omega}\sinh(\hbar\beta\omega/2)\right) = -\ln\left((2\sinh(\hbar\beta\omega/2))^2\right).$$
(33)

We then conclude

$$\det(\mu Q) = \det(Q) = (2\sinh(\hbar\beta\omega/2))^2, \tag{34}$$

and

$$\bigcirc = \int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)} = \det(\mu Q)^{-1/2} = \frac{1}{2\sinh(\hbar\beta\omega/2)}.$$
 (35)

4.1.2 Transition Amplitudes

We will now examine the transition amplitudes of this system. More precisely, since we are working in the Euclidean formalism, these truly correspond to the matrix elements of the Gibbs state at inverse temperature β . They are, however, related to transition amplitudes via Wick rotation. This will be an example of an integral that does not self-normalize. However, by using the partition function, which did self-normalize, and cut-paste renormalization conditions, we will be able to fix the value of μ .

In this example we want to compute (18) for $M = [\tau_i, \tau_f]$, an interval of size $\tau_f - \tau_i = \hbar \beta$, and

$$\mathcal{E} = \{ x \in C^{\infty}(M) \mid x(\tau_i) = x_i \text{ and } x(\tau_f) = x_f \},$$
(36)

for some fixed boundary conditions $x_i, x_f \in \mathbb{R}$. We will denote this path integral by

²In this paper, the convention is that N does not contain zero.

In particular, unless $x_i = x_f = 0$, \mathcal{E} is not a vector space. It is however an affine subspace of $C^{\infty}(M)$ modelled after the space of fluctuations

$$\mathcal{F} = \{ x \in C^{\infty}(M) \mid x(\tau_i) = x(\tau_f) = 0 \}.$$
 (38)

For any $x_0 \in \mathcal{E}$ we obtain a foliation with a trivial stem

$$\{x_0\} \times \mathcal{F} \to \mathcal{E} = x_0 + \mathcal{F}. \tag{39}$$

Of course, we will choose x_0 to be the unique instanton determined by the boundary conditions and the equations of motion $\ddot{x} = \omega^2 x$. It is given by

$$x_0 = \frac{\sinh(\omega(\tau - \tau_i))}{\sinh(\omega(\tau_f - \tau_i))} x_f + \frac{\sinh(\omega(\tau_f - \tau))}{\sinh(\omega(\tau_f - \tau_i))} x_i, \tag{40}$$

and has an action which can be easily computed using integration by parts and the equations of motion

$$S_0 = \left[\frac{1}{2}x\dot{x}\right]_{\tau_i}^{\tau_f} = \frac{\omega}{2}\left(\left(x_f^2 + x_i^2\right)\coth(\hbar\omega\beta) - 2x_f x_i \operatorname{csch}(\hbar\omega\beta)\right). \tag{41}$$

The boundary conditions on \mathcal{F} guarantee that the action separates into the form (22)

$$S(x_0 + \tilde{x}) = S_0 + \frac{1}{2} \langle \tilde{x}, Q\tilde{x} \rangle, \qquad (42)$$

where the operator Q is still given by (26) but must now be considered as an operator on $\mathcal{F} \subseteq C^{\infty}([\tau_i, \tau_f])$ as opposed to an operator on $C^{\infty}(S^1_{\hbar\beta})$. Accordingly, the eigenvalues of the operator are different. They are now given by the sines with periods divisible by $2\hbar\beta$

$$\lambda_n = \frac{\pi^2 n^2}{\hbar^2 \beta^2} + \omega^2, \quad \frac{1}{\sqrt{\hbar \beta}} \sin\left(\frac{\pi n}{\hbar \beta} (\tau - \tau_i)\right), \quad n \in \mathbb{N}.$$
 (43)

 \mathcal{F} also inherits the bilinear product from $C^{\infty}([\tau_i, \tau_f])$ and Q happens to be self-adjoint on \mathcal{F} . In view of the trivial stem, we can just take $\mathcal{V} = \mathcal{F}$.

In order to compute the determinant, we consider the zeta function

$$\zeta_Q(s) = \left(\frac{\pi^2}{\hbar^2 \beta^2}\right)^{-s} \zeta_{EH}\left(s; \frac{\hbar^2 \beta^2 \omega^2}{\pi^2}\right). \tag{44}$$

We then have using (135) that $\zeta_Q(0) = -1/2 \neq 0$, and the path integral does not self-normalize. Further, we have using (140)

$$\zeta_{Q}'(0) = -\ln\left(\frac{\pi^{2}}{\hbar^{2}\beta^{2}}\right)\zeta_{EH}\left(0; \frac{\hbar^{2}\beta^{2}\omega^{2}}{\pi^{2}}\right) + \frac{\mathrm{d}}{\mathrm{d}s}\zeta_{EH}\left(s; \frac{\hbar^{2}\beta^{2}\omega^{2}}{\pi^{2}}\right)\Big|_{s=0}
= \ln\left(\frac{\pi}{\hbar\beta}\right) - \ln\left(\frac{\pi}{\hbar\beta\omega}2\sinh(\hbar\beta\omega)\right) = -\ln\left(\frac{1}{\omega}2\sinh(\hbar\beta\omega)\right).$$
(45)

The determinant is then

$$\det(\mu Q) = \mu^{-\frac{1}{2}} \frac{2\sinh(\hbar\beta\omega)}{\omega}.$$
 (46)

We then conclude that the path integral is

$$\int_{x_i}^{x_f} = \mu^{1/4} \sqrt{\frac{\omega}{2\sinh(\hbar\beta\omega)}} e^{-\frac{\omega}{2\hbar} \left(\left(x_f^2 + x_i^2 \right) \coth(\hbar\omega\beta) - 2x_f x_i \operatorname{csch}(\hbar\omega\beta) \right)}$$
(47)

In order to finish this computation, we need to employ a renormalization condition that will allow us to determine μ . We will use the cut-paste condition that joining the two ends of a line should yield the circle

$$\frac{1}{2\sinh(\hbar\beta\omega/2)} = \bigcirc = \int dx \int_{x}^{x} = \mu^{1/4} \sqrt{\frac{\omega}{2\sinh(\hbar\beta\omega)}} \int dx \, e^{-\frac{\omega}{\hbar}(\coth(\hbar\omega\beta) - \operatorname{csch}(\hbar\omega\beta))x^{2}}$$

$$= \mu^{1/4} \sqrt{\frac{\hbar\pi}{4}} \frac{1}{\sinh(\hbar\beta\omega/2)}.$$
(48)

This fixes

$$\mu = \frac{1}{\pi^2 \hbar^2}.\tag{49}$$

This is of course, not the only cut-paste procedure one could perform. For example, one could realize the circle as the union of two lines of half the length

$$\frac{1}{2\sinh(\hbar\beta\omega/2)} = \bigcirc = \int dx_f dx_i \left[\begin{matrix} x_f \\ \vdots \\ x_i \end{matrix} \right] \frac{x_i}{\hbar\beta/2} \left[\begin{matrix} x_i \\ \vdots \\ x_f \end{matrix} \right] \frac{\hbar\beta/2}{x_f}$$

$$= \sqrt{\mu} \frac{\omega}{2\sinh(\hbar\beta\omega/2)} \int dx_f dx_i e^{-\frac{2}{\hbar}S_0}.$$
(50)

This method has the advantage that the prefactors of $2\sinh(\hbar\beta\omega/2)$ now cancel exactly. The remaining part of the equation constitutes an expression for the renormalization constant in terms of a finite-dimensional integral concerning the instanton action

$$\frac{1}{\omega\sqrt{\mu}} = \int dx_f dx_i e^{-\frac{m\omega}{\hbar} \left(\left(x_f^2 + x_i^2 \right) \coth(\hbar\omega\beta/2) - 2x_f x_i \operatorname{csch}(\hbar\omega\beta/2) \right)} \\
= \frac{\pi\hbar}{\omega} \det \begin{pmatrix} \coth(\hbar\omega\beta/2) & -\operatorname{csch}(\hbar\omega\beta/2) \\ -\operatorname{csch}(\hbar\omega\beta/2) & \coth(\hbar\omega\beta/2) \end{pmatrix}^{-1/2} = \frac{\pi\hbar}{\omega}.$$
(51)

4.2 Particle on the Circle

It is interesting to consider the free limit $\omega \to 0$ of the quantum harmonic oscillator. While the transition amplitude (47) is well defined in this limit, the partition function (35) diverges.

This is not an artifact of the path integral formalism: since the orbits of the free particle are not compact, this result is expected. Another interesting way of seeing this phenomenon is by considering a free particle on a compact space, such as a circle $S^1_{2\pi R}$ of radius R. The free particle should then be recovered in the limit $R \to \infty$. Moreover, since both theories share a common limit, the comparison method suggests that the renormalization constant will be the same.

For the particle on the circle, our space of fields is made out of maps $C^{\infty}(M, S^1_{2\pi R})$ assigning to every Euclidean time $\tau \in M$ a position for the particle $x(\tau)$ on the circle. Unlike the harmonic oscillator, this space does not carry a vector space structure since $S^1_{2\pi R}$ is not a vector space. The action is given by

$$S(x) = \int_{M} d\tau \left(\frac{1}{2} \dot{x}^2 + i \frac{\mathcal{B}}{2R} \dot{x} \right). \tag{52}$$

where \mathcal{B} is a magnetic parameter of the system.³ Note that the electromagnetic term has a factor of i since it stays invariant under Wick rotation.

4.2.1 Partition Function

Following the harmonic oscillator example, we will begin by studying the partition function of our theory. For this we need to compute (18)

$$\bigcirc \equiv \int_{\mathcal{E}} \mathcal{D}x \, e^{-\frac{1}{\hbar}S(x)}. \tag{53}$$

for $M = S_{h\beta}^1$ and $\mathcal{E} = C^{\infty}(M, S_{2\pi R}^1)$. In particular, \mathcal{E} is no longer a vector space. This will be our first example in which a non-trivial foliation appears. Moreover, the path integral will self-normalize through the mechanism explained in section 3. Another treatment of this computation can be found in [16] (with the assumption again that μ can be set to 1 from the start).

In order to find the leaf structure of the theory, we will replace $C^{\infty}(M, S_{\hbar\beta}^1)$ by a vector space \mathcal{W} at the cost of introducing a discrete gauge symmetry. This will be a space of fields defined on the spacetime $\tilde{M} := [0, \hbar\beta]$. We define \mathcal{W} to be the set of fields $x \in C^{\infty}(\tilde{M})$ such that $x(\hbar\beta) = x(0) + 2\pi Rn$, for some $n \in \mathbb{Z}$, and $x^{(k)}(0) = x^{(k)}(\hbar\beta)$ for all derivatives $k \in \mathbb{N}$. The first condition guarantees that the fields are periodic when viewed as taking values in $S_{2\pi R}^1 \cong \mathbb{R}/2\pi R\mathbb{Z}$. Therefore, all such fields descend to functions $S_{\hbar\beta}^1 \to S_{2\pi R}^1$. The n in this condition is called the **winding number** of x. The second condition on the derivatives ensures that the induced functions on the circle are smooth. However, \mathcal{W} is not equivalent to \mathcal{E} . Instead, using the identification $S_{2\pi R}^1 \cong \mathbb{R}/2\pi R\mathbb{Z}$, we obtain a surjective map $\mathcal{W} \to \mathcal{E}$. There is now an ambiguity because for every field $x \in \mathcal{W}$, both x and $x + 2\pi R$ correspond to the same physical field in \mathcal{E} . Accordingly, we now have a discrete gauge group $2\pi R\mathbb{Z}$ and the identification

$$\mathcal{E} \cong \mathcal{W}_{2\pi R\mathbb{Z}}.\tag{54}$$

³For a solenoid of radius r and magnetic field B, piercing transversely the circle in which the particle of charge e and mass m moves, we have $\mathcal{B} = eBr^2/m$.

The action of the theory lifts to W. In [23] it is suggested that we can use the method of Faddeev-Popov to fix the gauge symmetry and express our path integral as an integral over W. However, this path integral is simple enough that we will be able to fix the gauge using more direct methods. The procedure will be first to foliate W using a stem that is transverse to the gauge orbits and then gauge-fix the foliation by restricting the stem.

In order to do this, we will guide ourselves by studying the instantons of the action (52) on W. The magnetic term is topological and therefore does not contribute to the equations of motion $\ddot{x} = 0$. Accordingly, all instantons are of the form

$$x_{n,x_i} = \frac{2\pi Rn}{\hbar\beta}\tau + x_i. \tag{55}$$

They are classified by their winding number n and their initial position x_i . We would then like to choose a vector space of fluctuations $\mathcal{F} \subseteq \mathcal{W}$ such that we obtain a foliation

$$\mathbb{Z} \times \mathbb{R} \times \mathcal{F} \mapsto \mathcal{W}
(n, x_i, \tilde{x}) \mapsto x_{n, x_i} + \tilde{x}.$$
(56)

The power of these types of foliations lies in the fact that they can easily be gauge-fixed to a foliation of \mathcal{E} . Indeed, we can eliminate the $2\pi R\mathbb{Z}$ ambiguity by restricting the set of allowed x_i 's to an interval $I \subseteq \mathbb{R}$ of width $2\pi R$

$$\mathbb{Z} \times I \times \mathcal{F} \to \mathcal{E}. \tag{57}$$

In particular, if we can find an appropriate space of fluctuations, the stem is $\mathcal{J} := \mathbb{Z} \times I$.

One possible choice for \mathcal{F} is to take the space of fields $x \in \mathcal{W}$ of winding number 0 with initial position 0. This would yield a foliation whose leafs have fields with the associated instanton's winding number and initial position. However, this foliation is unnatural from the point of view of our initial space of fields since it heavily depends on two marked points that we decided to call 0. This will become clear in the fact that the path integral obtained via this scheme will not self-normalize. A more sensible and geometrically-motivated choice is obtained by setting \mathcal{F} to be the set of all $x \in \mathcal{W}$ with winding number 0 and average 0. To see that this gives an appropriate foliation, we need to note that every $x \in \mathcal{W}$ can be uniquely written as $x = x_{n,x_i} + \tilde{x}$ with n the winding number of x,

$$x_i := \frac{1}{\hbar \beta} \int_0^{\hbar \beta} d\tau \left(x - \frac{2\pi Rn}{\hbar \beta} \tau \right), \tag{58}$$

and $\tilde{x} := x - x_{n,x_i}$, which happens to lie in \mathcal{F} .

While the magnetic term did not contribute to the equations of motion, it does contribute to the action of the instantons

$$S(x_{n,x_i}) = \frac{2\pi^2 R^2 n^2}{\hbar \beta} + i\pi \mathcal{B}n =: S_n.$$
 (59)

Then the action becomes

$$S(x_{n,x_i} + \tilde{x}) = S_n + \frac{1}{2} \langle \tilde{x}, Q\tilde{x} \rangle.$$
 (60)

The bilinear form and differential operator on \mathcal{F} are inherited from (25) and the $\omega \to 0$ limit of (26) on $\mathcal{V} := C^{\infty}(S^1_{\hbar\beta})$. Indeed, \mathcal{F} can be identified as the subspace of fields in \mathcal{V} with 0 average. This also explains why we have no boundary terms since $S^1_{\hbar\beta}$ has no boundary. Moreover, the connected components of the foliation are

$$\{n\} \times I \times \mathcal{F} \subseteq x_{n,0} + \mathcal{V}.$$
 (61)

The $\omega \to 0$ limit of the treatment of Q in section 4.1.1 yields the measure on \mathcal{V} . This, in turn, induces a measure on \mathcal{F} . To be explicit, on $\{n\} \times I \times \mathcal{F}$ we have coordinates a_k defined by

$$x = \frac{2\pi Rn}{\hbar \beta} \tau + \sum_{k \in \mathbb{Z}} \frac{a_k}{\sqrt{\hbar \beta}} e^{\frac{2\pi i k}{\hbar \beta} \tau}, \tag{62}$$

and the measure by

$$\mathcal{D}x = d\left(\frac{a_0}{\sqrt{2\pi\hbar\mu}}\right) \prod_{k \in \mathbb{Z}} d\left(\frac{a_k}{\sqrt{2\pi\hbar\mu}}\right). \tag{63}$$

We are now ready to compute the path integrals. Over fluctuations, the integral is the same as (34) although we have to omit the zero mode a_0 . Although we could repeat the computation directly, it is easier to just take advantage of our previous result.

$$\det(\mu Q) = \lim_{\omega \to 0} \frac{\left(2\sinh(\hbar\beta\omega/2)\right)^2}{\mu\omega^2} = \frac{\hbar^2\beta^2}{\mu},\tag{64}$$

which depends on μ . We are left with computing the integral over the stem. In terms of x_i the integral measure is

$$d\left(\frac{a_0}{\sqrt{2\pi\hbar\mu}}\right) = \sqrt{\frac{\beta}{2\pi\mu}} dx_i \tag{65}$$

We then have

$$\int_{\mathcal{J}} ds \, e^{-\frac{1}{\hbar}S_s} = \sum_{n \in \mathbb{Z}} e^{-\frac{1}{\hbar}S_n} \sqrt{\frac{\beta}{2\pi\mu}} \int_{I} dx_i = \mu^{-1/2} \sqrt{2\pi\beta R^2} \sum_{n \in \mathbb{Z}} e^{-\frac{1}{\hbar}S_n}. \tag{66}$$

We conclude that

$$\bigcirc = \sqrt{\frac{2\pi R^2}{\hbar^2 \beta}} \sum_{n \in \mathbb{Z}} \exp\left(-\frac{2\pi^2 R^2 n^2}{\hbar^2 \beta} - \frac{i\pi \mathcal{B}n}{\hbar}\right). \tag{67}$$

In particular, the new contribution from the zero mode coming from the integral over \mathcal{J} cancels the μ -dependence of the integral over fluctuations. This thus constitutes an example of the self-normalization method introduced in section 3.

4.2.2 Transition Amplitudes

We will now study the transition amplitudes of this theory. They are given by (18)

$$\int_{x_i}^{x_f} \equiv \int_{\mathcal{E}} \mathcal{D}x \, e^{-\frac{1}{\hbar}S(x)}, \tag{68}$$

with $M = [\tau_i, \tau_f]$, an interval of Euclidean time with length $\hbar \beta$, and

$$\mathcal{E} = \left\{ x \in C^{\infty}(M, S^{1}_{2\pi R}) \mid x(\tau_i) = x_i \quad \text{and} \quad x(\tau_f) = x_f \right\}$$
 (69)

Again, this is not a vector space for very similar reasons as in the partition function computation. Accordingly, we will use the same technique to find an appropriate linear foliation. On the one hand, the leaf structure of the theory will turn out to be simpler. On the other, this path integral will not self-normalize. This will be another instance of how the cut-paste method can be used to renormalize a path integral. Another account of this computation can be found in [17] (where the overall normalization is fixed by using the comparison method).

Let us define W to be the set of fields $x \in C^{\infty}(M)$ such that $x(\tau_i) = x_i + 2\pi Rm$ and $x(\tau_f) = x_f + 2\pi Rn$ for some $m, n \in \mathbb{Z}$. This is not a vector space but it is affine inside of $C^{\infty}(M)$. In here we have chosen some representatives $x_i, x_f \in S^1_{2\pi R} = \mathbb{R}/2\pi R\mathbb{Z}$ in \mathbb{R} . This has the same discrete gauge symmetry

$$\mathcal{E} \cong \mathcal{W}_{2\pi R \mathbb{Z}}.\tag{70}$$

The instantons of (52) are of the form

$$x_{m,n} = \frac{x_f - x_i + 2\pi R(n-m)}{\hbar \beta} (\tau - \tau_i) + x_i + 2\pi Rm, \tag{71}$$

and we will choose the space of fluctuations to be the same as in section 4.1.2

$$\mathcal{F} := \{ x \in C^{\infty}(M) \mid x(\tau_i) = x(\tau_f) = 0 \}.$$
 (72)

This leaves us with leafs

$$x_{m,n} + \mathcal{F} =$$

$$\left\{ x \in C^{\infty}(M) \mid x(\tau_i) = x_i + 2\pi Rm \quad \text{and} \quad x(\tau_f) = x_f + 2\pi Rn \right\}$$

$$(73)$$

that induce the foliation

$$\mathbb{Z} \times \mathbb{Z} \times \mathcal{F} \to \mathcal{W}
(m, n, \tilde{x}) \mapsto x_{m,n} + \tilde{x}.$$
(74)

This foliation can be gauge-fixed to yield a foliation of the space of physical fields $\{m\} \times \mathbb{Z} \times \mathcal{F} \to \mathcal{E}$. In particular, $\mathcal{J} = \mathbb{Z}$ and the connected components $\{m\} \times \{n\} \times \mathcal{F}$ are trivially affine spaces modelled after $\mathcal{V} := \mathcal{F}$.

We are now ready to compute the path integral. The integral over fluctuations is given by a determinant that is simply the $\omega \to 0$ limit of (46)

$$\det(\mu Q) = \frac{2\hbar\beta}{\sqrt{\mu}}.\tag{75}$$

The instantons, on the other hand, have an action

$$S_n := S(x_{0,n}) = \frac{(x_f - x_i + 2\pi Rn)^2}{2\hbar\beta} + i\frac{\mathcal{B}(x_f - x_i + 2\pi Rn)}{2R}.$$
 (76)

Then the path integral is

$$\int_{x_i}^{x_f} = \mu^{1/4} \frac{1}{\sqrt{2\hbar\beta}} \sum_{n \in \mathbb{Z}} e^{-\frac{1}{\hbar}S_n}$$

$$\tag{77}$$

This path integral does not self-normalize. However, it can be renormalized using the cut-paste method on the partition function (67). This is particularly simple in this case since on $x_i = x_f$ the instanton actions (76) and (59) coincide. In particular, they are independent of the endpoints. We thus obtain that

$$\sqrt{\frac{2\pi R^2}{\hbar^2 \beta}} \sum_{n \in \mathbb{Z}} e^{-\frac{1}{\hbar} S_n} = \bigcirc = \int_I dx \int_x^x = \mu^{1/4} \frac{2\pi R}{\sqrt{2\hbar \beta}} \sum_{n \in \mathbb{Z}} e^{-\frac{1}{\hbar} S_n}. \tag{78}$$

We conclude that μ is still given by (49), as suggested by the comparison method.

5 Examples in 2D

We will now move on to the analogs of the examples of section 4 in 2D field theory. For this section, we prefer a field-theoretic notation. The analogy, however, is most clear when one thinks of the examples below as part of a string theory [24]. Quite surprisingly, we will see that all of the examples considered will self-normalize either via the method of section 2 or section 3.

5.1 Free Boson

We will consider fields ϕ in the vector space $C^{\infty}(M)$ for some 2D spacetime M with Euclidean metric g. The action of our theory will be that of a free scalar field of mass m

$$S(\phi) = \int_{M} d^{2}x \sqrt{g} \left(\frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi + \frac{1}{2} m^{2} \phi^{2} \right). \tag{79}$$

This vector space is equipped with the bilinear form

$$\langle \phi, \psi \rangle = \int_{M} d^{2}x \sqrt{g} \,\phi \psi,$$
 (80)

and the differential operator

$$Q = -\Delta + m^2 \tag{81}$$

for the Laplacian

$$\Delta \phi = \frac{1}{\sqrt{g}} \partial_{\mu} (\sqrt{g} \partial^{\mu} \phi). \tag{82}$$

5.1.1 On the Torus

We can generalize the discussion of section 4.1.1 using the action (79) when M is some closed manifold (of arbitrary dimension). In this case, we are interested in computing the path integral (18)

$$\left(\right) \equiv \int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}, \tag{83}$$

with $\mathcal{E} = C^{\infty}(M)$. This is already a vector space. The action can be put into the form $S = \frac{1}{2} \langle \phi, Q\phi \rangle$ via integration by parts. The boundary term vanishes given that M is closed.⁴ We then conclude that the path integral is Gaussian

$$\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)} = \det(\mu Q)^{-1/2}.\tag{84}$$

In other words, the computation of these path integrals reduces to the computation of $\zeta_Q(0)$ and $\zeta_Q'(0)$ up to finding a suitable renormalization condition for μ .

We will consider the case where M is given by a torus $M = S_{2\pi r_1}^1 \times S_{2\pi r_2}^1$ built as a quotient of the complex plane $\mathbb C$ with the standard metric $g = \mathrm{d} x \otimes \mathrm{d} x + \mathrm{d} y \otimes \mathrm{d} y$ in Cartesian coordinates z = x + iy. We will consider the horizontal translation by $2\pi r_2$ and by a complex number of the form $\tau = w + i\hbar\beta$, with $|\tau| = 2\pi r_1$ and $\beta \neq 0$,

$$M = \mathbb{C}_{\tau \mathbb{Z} \times 2\pi r_2 \mathbb{Z}}.$$
 (85)

We can introduce new coordinates σ^{μ} , defined by $x + iy = z = \sigma^1 + \frac{\tau}{|\tau|}\sigma^0$, in which the boundary conditions imposed by the quotient are simpler

$$\sigma^0 \sim \sigma^0 + 2\pi r_1, \quad \sigma^1 \sim \sigma^1 + 2\pi r_2 \tag{86}$$

In these coordinates the metric tensor is constant but not diagonal

$$g = g_{\mu\nu} \,\mathrm{d}\sigma^{\mu} \otimes \mathrm{d}\sigma^{\nu} = \mathrm{d}\sigma^{0} \otimes \mathrm{d}\sigma^{0} + \frac{w}{2\pi r_{1}} (\mathrm{d}\sigma^{0} \otimes \mathrm{d}\sigma^{1} + \mathrm{d}\sigma^{1} \otimes \mathrm{d}\sigma^{0}) + \mathrm{d}\sigma^{1} \otimes \mathrm{d}\sigma^{1}. \tag{87}$$

For example, it is simple to compute in these coordinates the area of the torus

$$A = \int d^2x \sqrt{g} = \frac{\hbar \beta}{2\pi r_1} \int_0^{2\pi r_1} d\sigma^0 \int_0^{2\pi r_2} d\sigma^1 = 2\pi r_2 \hbar \beta.$$
 (88)

⁴This is most easily seen by rewritting (79) using differential forms [see e. g. 25, Example 2.28] and the expression for the Laplacian $d^n x \sqrt{g} \Delta = d \star d$.

The eigenvalue equation of Q is

$$(-\Delta + m^2)\phi = \lambda\phi. \tag{89}$$

The solutions of this are plane waves. In coordinates in which the metric tensor is constant, like our standard Cartesian coordinates or the σ coordinates, they are of the form $e^{ip_{\mu}x^{\mu}}$. The momentum is quantized by the boundary conditions, as is most easily seen in the σ coordinates

$$\phi_{nk} = A^{-1/2} e^{i\left(\frac{n}{r_1}\sigma^0 + \frac{k}{r_2}\sigma^1\right)}, \quad n, k \in \mathbb{Z}, \tag{90}$$

with eigenvalues

$$\lambda_{nk} = g^{\mu\nu} p_{\mu} p_{\nu} + m^2 = \frac{4\pi^2 r_1^2}{\hbar^2 \beta^2} \left(\frac{n^2}{r_1^2} + \frac{k^2}{r_2^2} - 2 \frac{w}{2\pi r_1} \frac{n}{r_1} \frac{k}{r_2} \right) + m^2.$$
 (91)

We can now consider the zeta function of Q. This can be expressed in terms of the two-dimensional inhomogeneous Epstein zeta function (141)

$$\zeta_Q(s) = m^{-2s} + F\left(s; \frac{4\pi^2}{\hbar^2 \beta^2} \frac{r_1^2}{r_2^2}, -\frac{4\pi^2}{\hbar^2 \beta^2} \frac{w}{\pi r_2}, \frac{4\pi^2}{\hbar^2 \beta^2}; m^2\right). \tag{92}$$

In particular, using (144) we obtain $\zeta_Q(0) = 1 - 1 = 0$ and we conclude the path integral self-normalizes via the method of section 2. The full path integral can then be computed by taking the derivative at s = 0. Using the results obtained in [18] in the context of the Wheeler-De Witt equation, one can give a rapidly convergent series expansion for it. Since we have already shown the path integral self-normalizes, we will not cite the full expression here. However, we note that the path integral will diverge to $+\infty$ in the massless limit since the term $-2\ln(m)$ of the zero-mode contribution to $\zeta_Q'(0)$ does. This behavior is analogous to that of the integral considered in section 4.1.1.

5.1.2 On the Cylinder

We now consider the theory on a cylinder $M = [y_i, y_f] \times S_{2\pi r}^1$, obtained by cutting the torus of the previous section along a circle of constant σ^0 . We will consider the case w = 0 so that the standard coordinates and the σ coordinates agree. In particular, $y_f - y_i = \hbar \beta = 2\pi r_1$ and $r = r_2$. This computation should be compared against that of section 4.1.2. The steps of the computation are analogous but, unlike in the 1D case, this path integral will self-normalize with the method of section 2. Accordingly, this will be our first path integral on a non-closed spacetime that will self-normalize.

The boundary conditions are determined by $\phi_i, \phi_f \in C^{\infty}(S_{2\pi r}^1)$ on $\Sigma_i := \{y_i\} \times S_{2\pi r}^1$ and $\Sigma_f := \{y_f\} \times S_{2\pi r}^1$. We are then interested in computing the path integral (18)

$$\underbrace{\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}}_{\phi_i} \equiv \int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}, \tag{93}$$

with

$$\mathcal{E} := \left\{ \phi \in C^{\infty}(M) \mid \phi|_{\Sigma_i} = \phi_i \quad \text{and} \quad \phi|_{\Sigma_f} = \phi_f \right\}. \tag{94}$$

This is not a vector space unless $\phi_i = \phi_f = 0$. However, it is an affine subspace of $C^{\infty}(M)$ modelled after

$$\mathcal{F} = \left\{ \phi \in C^{\infty}(M) \mid \phi|_{\Sigma_i} = \phi|_{\Sigma_f} = 0 \right\}. \tag{95}$$

We have a simple foliation

$$\{\phi_0\} \times \mathcal{F} \to \mathcal{E} = \phi_0 + \mathcal{F},$$
 (96)

for any ϕ_0 in \mathcal{E} . Choosing ϕ_0 to be an instanton,⁵ and letting S_0 be its instanton action, the periodicity along the circle and the boundary conditions of the space of fluctuations guarantee the action separates as in (22)

$$S(\phi_0 + \tilde{\phi}) = S_0 + \frac{1}{2} \left\langle \tilde{\phi}, Q\tilde{\phi} \right\rangle. \tag{97}$$

The eigenfunctions of Q are given now by

$$\phi_{nk} = \sqrt{2}A^{-1/2}\sin\left(\frac{\pi n}{\hbar\beta}(y - y_i)\right)e^{i\frac{k}{r}x}, \quad n \in \mathbb{N}, \quad k \in \mathbb{Z},$$
(98)

corresponding to the eigenvalues

$$\lambda_{nk} = \frac{\pi^2 n^2}{\hbar^2 \beta^2} + \frac{k^2}{r^2} + m^2. \tag{99}$$

The corresponding zeta function can then be expressed in terms of the Epstein-Hurwitz zeta function (134) and the inhomogeneous Epstein zeta function (141) by using the partition

$$\mathbb{Z}^2 \setminus \{(0,0)\} = (\mathbb{N} \times \mathbb{Z}) \cup (-\mathbb{N} \times \mathbb{Z}) \cup (\{0\} \times \mathbb{N}) \cup (\{0\} \times -\mathbb{N}). \tag{100}$$

The left hand side can be identified with the indices of the inhomogeneous Epstein zeta function. The first two terms of the right hand side correspond those of our zeta function of interest since we took w = 0. The last two terms of the right hand side correspond to the indices of Epstein-Hurwitz zeta functions. This yields

$$\zeta_Q(s) = \frac{1}{2} F\left(s; \frac{1}{r^2}, 0, \frac{\pi^2}{\hbar^2 \beta^2}; m^2\right) - r^{2s} \zeta_{EH}(s, m^2 r^2). \tag{101}$$

Then (135) and (144) give us that

$$\zeta_Q(0) = \frac{1}{2}(-1) - \left(-\frac{1}{2}\right) = 0.$$
 (102)

Accordingly, the determinant self-normalizes with the method of section 2. This implies that the whole path integral self-normalizes since the stem integral is trivial.

⁵The existence of such an instanton can be easily established by expanding in a Fourier series along the circle to reduce the equations of motion to a series of ordinary differential equations.

5.2 Compact Boson

Now we will consider the case of a massless boson whose associated fields take values in a compact space. This will be the 2D analog of the particle in the circle treated in section 4.2. Here our fields will be elements of spaces of the form $C^{\infty}(M, S^1_{\hbar\beta})$ for some two-dimensional spacetime M with a metric g. The action will be given by the massless version of (79)

$$S(\phi) = \int_{M} d^{2}x \sqrt{g} \, \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi. \tag{103}$$

5.2.1 On the Torus

We will mimic section 5.1.1 and take M as in (85). Refer to that section for comments on the coordinatization and the metric structure we are interested in for the torus. The computation that follows should be compared to the one done in section 4.2.1. Since most of the steps are simple generalizations of the ones found there, we will not cover them in as much detail. A standard reference for this calculation is [26], but our method clarifies some of the steps found there.

In this section, we are interested in computing the path integral (18)

for $\mathcal{E}:=C^{\infty}(M,S^1_{2\pi R})$. As in section 4.2.1, we will find convenient to introduce a redundant space of fields \mathcal{W} . This will be the space of fields $\phi\in C^{\infty}(\tilde{M})$, where $\tilde{M}=[0,2\pi r_1]\times[0,2\pi r_2]$, satisfying the following gluing properties. On the one hand, we need to impose a combatibility condition between the left $\Sigma_L=[0,2\pi r_1]\times\{0\}$ and right $\Sigma_R=[0,2\pi r_1]\times\{2\pi r_2\}$ boundaries. Namely, we need to demand that $\phi|_{\Sigma_R}=\phi|_{\Sigma_L}+2\pi Rn$ for some $n\in\mathbb{Z}$ called the horizontal winding number. This winding number is uniform throughout these boundaries due to the continuity of the field. A similar condition has to be imposed on the initial $\Sigma_i=\{0\}\times[0,2\pi r_2]$ and final boundaries $\Sigma_f=\{2\pi r_1\}\times[0,2\pi r_2]$. We thus have $\phi_{\Sigma_f}=\phi_{\Sigma_i}+2\pi Rk$ for some $k\in\mathbb{Z}$ called the vertical winding number. The conditions so far guarantee that the fields in \mathcal{W} can be reinterpreted as maps $M\to S^1_{2\pi R}$. The rest of the conditions are now simply that $\partial_I\phi|_{\Sigma_R}=\partial_I\phi|_{\Sigma_L}$ and $\partial_I\phi|_{\Sigma_f}=\partial_I\phi|_{\Sigma_i}$, for any multiindex $I\neq 0$. This ensures that the resulting maps $M\to S^1_{2\pi R}$ are in fact smooth. Of course, having lifted the values of our fields from $S^1_{2\pi R}$ to \mathbb{R} we have incurred once again in the ambiguity that both ϕ and $\phi+2\pi R$ correspond to the same field. In other words,

$$\mathcal{E} \cong \mathcal{W}_{2\pi R \mathbb{Z}}.\tag{105}$$

We will once again find first a foliation of $\mathcal W$ which we will then gauge-fix to $\mathcal E$. The instantons in $\mathcal W$ are of the form

$$\phi_{nk\varphi}(\sigma^0, \sigma^1) = n \frac{R}{r_1} \sigma^0 + k \frac{R}{r_2} \sigma^1 + \varphi, \quad n, k \in \mathbb{Z}, \quad \varphi \in \mathbb{R}.$$
 (106)

We would then like to find a space of fluctuations \mathcal{F} that yields an appropriate foliation

$$\mathbb{Z}^2 \times \mathbb{R} \times \mathcal{F} \to \mathcal{W}
\left(n, k, \varphi, \tilde{\phi}\right) \mapsto \phi_{nk\varphi} + \tilde{\phi}.$$
(107)

As before, the advantage of these foliations is that they can immediately be gauge-fixed to find a foliation $\mathbb{Z}^2 \times I \times \mathcal{F} \to \mathcal{E}$ by choosing an interval I of length $2\pi R$. In other words, the stem would be $\mathcal{J} = \mathbb{Z}^2 \times I$. In fact, as before, we can choose \mathcal{F} to be the set of fields in \mathcal{W} with winding numbers 0 (or, equivalently, the fields in $\mathcal{V} := C^{\infty}(M)$) whose average is 0.

We define

$$S_{nk} := S(\phi_{nk\varphi}) = \frac{1}{2} A R^2 \frac{4\pi^2 r_1^2}{\hbar^2 \beta^2} \left(\frac{n^2}{r_1^2} + \frac{k^2}{r_2^2} - 2 \frac{w}{2\pi r_1} \frac{n}{r_1} \frac{k}{r_2} \right). \tag{108}$$

Then, since the fluctuations are truly functions on M and the latter does not have a boundary, we can integrate by parts to obtain the separation (22)

$$S(\phi_{nk\varphi} + \tilde{\phi}) = S_{nk} + \frac{1}{2} \left\langle \tilde{\phi}, Q\tilde{\phi} \right\rangle. \tag{109}$$

We have once again borrowed the inner product and differential operator on \mathcal{V} from (80) and (81).

The connected components of this foliation are of the form $\{n\} \times \{k\} \times I \times \mathcal{F}$. These are all subsets of the affine spaces $\phi_{n,k,0} + \mathcal{V}$ modelled on \mathcal{V} . Accordingly, the measure on \mathcal{V} , which happens to be the $m \to 0$ limit of the measure on the space of fields of section 5.1.1, induces a measure on our foliation. To be precise, on each connected component we have coordinates a_{pq} given by

$$\phi = \phi_{nk0} + \sum_{p,q \in \mathbb{Z}} a_{pq} \phi_{pq} \tag{110}$$

and the measure

$$\mathcal{D}\phi = d\left(\frac{a_{00}}{\sqrt{2\pi\hbar\mu}}\right) \prod_{p,q\in\mathbb{Z}} d\left(\frac{a_{pq}}{\sqrt{2\pi\hbar\mu}}\right). \tag{111}$$

Then the measure induced on the stem is obtained by identifying

$$\varphi = a_{00}\phi_{00} = A^{-1/2}a_{00},\tag{112}$$

i. e.

$$d\left(\frac{a_{00}}{\sqrt{2\pi\hbar\mu}}\right) = \sqrt{\frac{A}{2\pi\hbar\mu}}d\varphi. \tag{113}$$

The measure on the leafs is obtained by omitting the zero mode a_{00} .

The zeta function for Q is then obtained by ommitting the zero mode from the $m \to 0$ limit of (92)

$$\zeta_Q(s) = F\left(s; \frac{4\pi^2}{\hbar^2 \beta^2} \frac{r_1^2}{r_2^2}, -\frac{4\pi^2}{\hbar^2 \beta^2} \frac{w}{\pi r_2}, \frac{4\pi^2}{\hbar^2 \beta^2}; 0\right). \tag{114}$$

Using equation (144), $\zeta_Q(0) = -1$. On the other hand, the stem integral is

$$\int_{\mathcal{J}} ds \, e^{-\frac{1}{\hbar}S_s} = \sum_{n,k\in\mathbb{Z}} e^{-\frac{1}{\hbar}S_{nk}} \sqrt{\frac{A}{2\pi\hbar\mu}} \int_{I} d\varphi \propto \mu^{-1/2} = \mu^{\frac{1}{2}\zeta_Q(0)}. \tag{115}$$

We conclude this path integral self-normalizes via the method of section 3. In fact, we can compute the full path integral (23) using the derivative of the zeta function⁶ (147). The derivative of the zeta function is

$$\zeta_Q'(0) = -2\ln\left(\hbar\beta \left|\eta\left(\frac{\tau}{2\pi r_2}\right)\right|^2\right),\tag{116}$$

which yields the path integral

$$\left(\begin{array}{c} \left(\begin{array}{c} \left(\begin{array}{c} \left(\begin{array}{c} \left(\frac{1}{2\pi R}\right) \right) \\ \left(\frac{1}{2\pi R}\right) \end{array}\right) & \sum_{n,k\in\mathbb{Z}} e^{-\frac{1}{\hbar}S_{nk}} \\ = \frac{2\pi R}{\sqrt{2\pi\hbar}} \sqrt{\frac{2\pi r_2}{\hbar\beta}} \left| \eta \left(\frac{\tau}{2\pi r_2}\right) \right|^{-2} \sum_{n,k\in\mathbb{Z}} e^{-\frac{1}{\hbar}S_{nk}}.
\end{array} \right) (117)$$

5.2.2 On the Cylinder

Let us now consider $M = [y_i, y_f] \times S^1_{2\pi r}$ to be the cylinder explored in section 5.1.2. Boundary conditions are determined by initial and final fields $\phi_i, \phi_f \in C^{\infty}(S^1_{2\pi r}, S^1_{2\pi R})$. We then want to compute the path integral (18)

$$\underbrace{\int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}}_{\phi_{\delta}} \equiv \int_{\mathcal{E}} \mathcal{D}\phi \, e^{-\frac{1}{\hbar}S(\phi)}, \tag{118}$$

with

$$\mathcal{E} = \left\{ \phi \in C^{\infty}(M, S^{1}_{2\pi R}) \mid \phi|_{\Sigma_{i}} = \phi_{i} \quad \text{and} \quad \phi|_{\Sigma_{f}} = \phi_{f} \right\}.$$
 (119)

This will be analogous to the computation in section 4.2.2.

There is an important subtlety appearing in this case that is new to this example. Note that given a circle $\Sigma = \{y\} \times S^1_{2\pi R} \subseteq M$ and a field $\phi \in C^{\infty}(M, S^1_{2\pi r})$, the field $\phi|_{\Sigma}$ defines an element of $C^{\infty}(S^1_{2\pi r}, S^1_{2\pi R})$. Accordingly, it has a winding number. The discussion in section 5.2.1 shows that smoothness demands that this winding number is independent of the circle

⁶In here instead of using $|\eta(\tau)|$ we have used $|\overline{\eta(\tau)}|$. It is clear they both coincide and taking the complex conjugate of η amounts to changing the sign of the real part of its argument.

 Σ chosen. In that section, we called it the horizontal winding number. Therefore, \mathcal{E} is empty unless both ϕ_i and ϕ_f have the same winding number. Thus, in order to continue employing our techniques, from now on, we will then assume that ϕ_i and ϕ_f have the same winding number $n \in \mathbb{Z}$. In particular, all fields in \mathcal{E} have the horizontal winding number n.

Following the procedure of sections 4.2.1, 4.2.2 and 5.2.1, we want to replace this by a redundant space of fields \mathcal{W} on $\tilde{M} = [y_i, y_f] \times [0, 2\pi r]$ valued in \mathbb{R} . This will require us to lift the initial and final fields ϕ_i and ϕ_f to maps in $C^{\infty}([0, 2\pi r])$, and any lifts will do. We then define \mathcal{W} to be the set of fields $\phi \in C^{\infty}(\tilde{M})$ satisfying the following boundary conditions. On the one hand, we must have the periodicity conditions given by the horizontal winding mode $\phi|_{\Sigma_R} = \phi|_{\Sigma_L} + 2\pi Rn$. These ensure that the fields can be reinterpreted as maps $M \to S^1_{2\pi R}$ with the correct winding number. Moreover, we demand that $\partial_I \phi|_{\Sigma_L} = \partial_I \phi|_{\Sigma_R}$ for all multiindices $I \neq 0$. This ensures that that the maps are smooth. Finally, the boundary conditions take the form $\phi|_{\Sigma_i} = \phi_i + 2\pi Rm_i$ and $\phi|_{\Sigma_f} = \phi_f + 2\pi Rm_f$ for some $m_i, m_f \in \mathbb{Z}$. The resulting space of fields exhibits the same ambiguity

$$\mathcal{E} = \frac{\mathcal{W}}{2\pi R \mathbb{Z}}.\tag{120}$$

In order to find the instantons we first note that all $\phi \in \mathcal{W}$ can be written in the form

$$\phi = n \frac{R}{r_2} x + \frac{1}{\sqrt{2\pi r_2}} \sum_{k \in \mathbb{Z}} c_k e^{i\frac{k}{r_2}x}, \tag{121}$$

for some functions $c_k \in C^{\infty}([y_i, y_f])$. The instanton equation is then

$$0 = \Delta \phi = \frac{1}{\sqrt{2\pi r_2}} \sum_{k \in \mathbb{Z}} \left(\frac{\mathrm{d}^2 c_k}{\mathrm{d} y^2} - \frac{k^2}{r_2^2} c_k \right) e^{i\frac{k}{r_2} x}. \tag{122}$$

Thus, the instanton equations determine ordinary differential equations for the c_k . These equations were studied in section 4.1.2. The initial conditions for these are given by the expansions

$$\phi_i = n \frac{R}{r_2} x + \frac{1}{\sqrt{2\pi r_2}} \sum_{k \in \mathbb{Z}} \phi_{i,k} e^{i\frac{k}{r_2}x}.$$
 (123)

The final conditions are given by the similar expansion for ϕ_f . We conclude that the instantons are of the form

$$\phi_{m_{i}m_{f}} = \varphi_{i} + 2\pi R m_{i} + n \frac{R}{r} x + \frac{\varphi_{f} - \varphi_{i} + 2\pi R (m_{f} - m_{i})}{\hbar \beta} (y - y_{i})$$

$$+ \sum_{k \in \mathbb{Z} \setminus \{0\}} \left(\frac{\sinh(k(y - y_{i})/r)}{\sinh(k\hbar \beta/r)} \phi_{f,k} + \frac{\sinh(k(y_{f} - y)/r)}{\sinh(k\hbar \beta/r)} \phi_{i,k} \right) e^{i\frac{k}{r}x}, \tag{124}$$

where $\varphi_i := \phi_{i,0}/\sqrt{2\pi r}$ and $\varphi_f := \phi_{f,0}/\sqrt{2\pi r}$. It is simple to see that the instanton action is only a function of $m_f - m_i$

$$S(\phi_{m_i, m_f}) =: S_{m_f - m_i}. \tag{125}$$

We would now like to find a space of fluctuations \mathcal{F} that yield a foliation

$$\mathbb{Z}^2 \times \mathcal{F} \to \mathcal{W}$$

$$(m_i, m_f, \tilde{\phi}) \mapsto \phi = \phi_{m_i m_f} + \tilde{\phi}.$$
(126)

This kind of foliations can be gauge-fixed by fixing m_i

$$\{m_i\} \times \mathbb{Z} \times \mathcal{F} \to \mathcal{E}.$$
 (127)

We will choose \mathcal{F} to be the set of fields $\tilde{\phi} \in C^{\infty}(\tilde{M})$ satisfying the following conditions. On the one hand $\partial_I \tilde{\phi}|_{\Sigma_R} = \partial_I \tilde{\phi}|_{\Sigma_R}$ for any multiindex I. In other words, they can be reinterpreted as maps in $C^{\infty}(M)$. We will also require that they vanish at the boundaries $\tilde{\phi}|_{\Sigma_i} = \tilde{\phi}|_{\Sigma_f} = 0$. Since the connected components of the foliation are of the form $\{m_i\} \times \{m_f\} \times \mathcal{F}$, we can take $\mathcal{V} = \mathcal{F}$.

The fluctuations can all be written in the form

$$\tilde{\phi} = \sum_{k \in \mathbb{Z}} \phi_k e^{i\frac{k}{r}x},\tag{128}$$

for some functions $\phi_k \in C^{\infty}([y_i, y_f])$ that vanish at the end points. The eigenvalue equation $-\Delta \tilde{\phi} = \lambda \tilde{\phi}$ then takes the form

$$\frac{\mathrm{d}^2 \phi_k}{\mathrm{d}y^2} - \frac{k^2}{r^2} \phi_k + \lambda \phi_k = 0 \tag{129}$$

such that $\phi_k(\tau_i) = \phi_k(\tau_f) = 0 \ \forall k \in \mathbb{Z} \setminus \{0\}$. The eigenfunctions and their eigenvalues are then

$$\phi_{lk} \propto \sin\left(\frac{l\pi}{\hbar\beta}\right) e^{i\frac{k}{r}\sigma}, \quad \lambda_{lk} = \frac{l^2\pi^2}{\hbar^2\beta^2} + \frac{k^2}{r^2}, \quad l \in \mathbb{N}, \quad k \in \mathbb{Z}.$$
(130)

Repeating the discussion leading up to (101), we conclude the zeta function of the operator is

$$\zeta_Q(s) = \frac{1}{2} F\left(s; \frac{1}{r^2}, 0, \frac{\pi^2}{\hbar^2 \beta^2}; 0\right) + r^{2s} \zeta(2s)$$
(131)

We then conclude that $\zeta_Q(0) = 0$. Since the stem is trivial, this implies that the path integral self-normalizes. We can also compute its derivative using (147)

$$\zeta_Q'(0) = -\ln\left(2\hbar\beta \left|\eta\left(i\frac{\hbar\beta}{\pi r}\right)\right|^2\right) + 2\ln(r)\zeta(0) + 2\zeta'(0) = -\ln\left(2A\left|\eta\left(i\frac{\hbar\beta}{\pi r}\right)\right|^2\right)$$
(132)

This yields the result

$$\frac{\phi_f}{\phi_i} = (2A)^{-1/2} \left| \eta \left(i \frac{\hbar \beta}{\pi r} \right) \right|^{-1} \sum_{m \in \mathbb{Z}} e^{-\frac{1}{\hbar} S_m}, \tag{133}$$

6 Conclusions and Open Problems

This paper addressed the problem of computing the overall normalization constant of path integrals using zeta-function regularization techniques. In particular, we studied the phenomenon we called "self-normalization," in which the ambiguity of the integral measure, which would typically need to be renormalized, resolves itself. Hawking studied the conditions under which this phenomenon happens using zeta-function regularization for the well-known Gaussian integrals. However, this only works when the space of fields has a vector space structure. We devised an extension for the cases in which we do not have a vector space but instead another structure which we called a "linear foliation." The core idea behind this structure is that there is a stem encoding the information associated with the classical equations of motion, on which leafs are attached, which harbor the description of quantum fluctuations. They were modeled to embed these foliations into vector spaces suitable to the more standard techniques. We considered examples in one and two spacetime dimensions for which, even though the transition amplitudes or partition functions were not given by Gaussian integrals, were amenable to calculations by decomposing the space of fields into a linear foliation. Some of these cases self-normalized because the ambiguity of the integral along the stem was compensated by the one in the leafs. Interestingly, all the 2D examples we considered (the massive and compact bosons on a cylinder and a torus) exhibited this behavior.

One of the applications of our results is the computation of the overall normalization of path integrals that do not self-normalize. This computation is usually done in the literature using different comparison methods, which involve additional assumptions on the nature of this constant. However, an alternative method is to use cut-paste methods on self-normalizing path integrals to recover the normalization of these non-self-normalizing ones. This method avoids the need for those extra assumptions. We showed this explicitly in the examples in which the path integrals did not self-normalize.

Multiple possible paths can stem from our paper. First, we could try to extend our methods to other types of integrals of physical importance, like fermionic path integrals. Another possibility would be to find physical examples in which the foliation are non-trivial vector bundles of infinite rank over a finite-dimensional stem. Second, the fact that all of our path integrals on circles self-normalized while those on the line did not, or that all of our 2D examples did, hint strongly at the possibility that the self-normalization of path integrals is hiding topological information about the spacetime manifold in which the fields are evaluated. Clarifying this connection would be an important next step. Along with this same idea, we surmise that these techniques might shed some light on evaluating the path integrals of topological field theories. Third, we would like to highlight that, if supplemented with Picard-Lefschetz techniques, our methods are likely to be equally applicable to non-Euclidean manifolds, among which Minkowskian spacetimes would be physically relevant. Finally, in a similar way in which the work of Sorkin, Dowker, et al. have led to recovering Hilbert spaces from path integrals, we hypothesize that, with an appropriate interpretation of path integrals in terms of events, one could go beyond the construction of a Hilbert space to that of the operator C*-algebras inspired by the use of insertions in (1). Although this paper merely scratches the surface of the potential applications of linear foliation decomposition in path integrals, we think it might become an important part of the (mathematical) physicist's toolbox and, moreover, it lays the ground for future endeavors relevant to both physicists and mathematicians.

7 Acknowledgments

We would like to thank Colegio San Carlos, in Bogotá, Colombia, where we had our primary (in the case of IMB only one year) and secondary education, and first got interested in physics and mathematics. In particular, we would like to thank Fr. Francis Wehri not only for the profound impact he had in our education but also for the role he played in that of countless other Colombians. Similarly, we would like to thank Andrés F. Reyes-Lega, who later became our supervisor during our time in the Universidad de los Andes. We would also like to thank Petr Hořava for useful comments and suggestions on the manuscript. IMB would like to acknowledge financial support from the Berkeley Graduate Division and the Friends of Warren Hellman Fund.

A Useful Values of Zeta Functions

In this appendix, we recall some facts on zeta functions that we will need throughout the body of the text. If presented without a citation or a derivation, the formulas can be found in [18].

The Epstein-Hurwitz zeta function is

$$\zeta_{EH}(s;p) := \sum_{n=1}^{\infty} (n^2 + p)^{-s}
= -\frac{p^{-s}}{2} + \frac{\sqrt{\pi}\Gamma(s - 1/2)}{2\Gamma(s)} p^{-s+1/2} + \frac{2\pi^s p^{-s/2+1/4}}{\Gamma(s)} \sum_{n=1}^{\infty} n^{s-1/2} K_{s-1/2}(2\pi n \sqrt{p}), \tag{134}$$

where K_{ν} are the modified Bessel functions of the second kind. The second expression defines its analytic continuation to the complex plane. Its limit at s=0 is simple to take since $\Gamma(s) \to \pm \infty$. Therefore, only the first term contributes

$$\zeta_{EH}(0;p) = -1/2. \tag{135}$$

To compute its derivative at s=0 we need more information on the behaviour of the Γ function at s=0, namely [27]

$$\frac{\mathrm{d}}{\mathrm{d}s} \left. \frac{1}{\Gamma(s)} \right|_{s=0} = 1. \tag{136}$$

Therefore, as far as the last two terms are concerned, we need only consider the derivatives that act on the Gamma function in the denominator

$$\left. \frac{\partial}{\partial s} \zeta_{EH}(s;p) \right|_{s=0} = \frac{1}{2} \ln p + \frac{\sqrt{\pi} \Gamma(-1/2)}{2} p^{1/2} + 2p^{1/4} \sum_{n=1}^{\infty} n^{-1/2} K_{-1/2}(2\pi n \sqrt{p})$$
 (137)

This leaves a series that can be computed using

$$K_{-1/2}(z) = K_{1/2}(z) = \sqrt{\frac{\pi}{2}} \frac{e^{-z}}{z^{1/2}}$$
 (138)

and the Taylor expansion of the logarithmic function at 1

$$2p^{1/4} \sum_{n=1}^{\infty} n^{-1/2} K_{-1/2}(2\pi n \sqrt{p}) = \sum_{n=1}^{\infty} \frac{e^{-2\pi n \sqrt{p}}}{n} = -\ln(1 - e^{-2\pi\sqrt{p}})$$
 (139)

Using that $\Gamma(-1/2) = -2\sqrt{\pi}$, we obtain

$$\frac{\partial}{\partial s} \zeta_{EH}(s; p) \bigg|_{s=0} = \frac{1}{2} \ln p - \pi \sqrt{p} - \ln \left(1 - e^{-2\pi\sqrt{p}} \right) = -\ln \left(\frac{1}{\sqrt{p}} \left(e^{\pi\sqrt{p}} - e^{-\pi\sqrt{p}} \right) \right)
= -\ln \left(p^{-1/2} 2 \sinh(\pi\sqrt{p}) \right).$$
(140)

Another important zeta function we will need is the two-dimensional inhomogeneous Epstein zeta function

$$F(s; a, b, c; q) = \sum_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} (am^2 + bmn + cn^2 + q)^{-s}.$$
 (141)

The omission of m = n = 0 is customary in the literature, and we will follow suit. However, in some physical applications, we will need the m = n = 0 term, which means we will have to incorporate a summand of q^{-s} explicitly.

The analytic continuation of this is given in terms of the generalized Chowla-Selberg formula

$$F(s; a, b, c; q) = 2\zeta_{EH}(s, 4aq/\Delta)a^{-s} + \frac{2^{2s}\sqrt{\pi}a^{s-1}}{\Gamma(s)\Delta^{s-1/2}}\Gamma(s - 1/2)\zeta_{EH}(s - 1/2, 4aq/\Delta)$$

$$+ \frac{8(2\pi)^{s}}{\Gamma(s)\Delta^{s-1/2}\sqrt{2a}} \sum_{n=0}^{\infty} n^{s-1/2}\cos(n\pi b/a) \times$$

$$\sum_{d|n} d^{1-2s} \left(\Delta + \frac{4aq}{d^{2}}\right)^{s/2-1/4} K_{s-1/2} \left(\frac{\pi n}{a}\sqrt{\Delta + \frac{4aq}{d^{2}}}\right),$$
(142)

with

$$\Delta = 4ac - b^2. \tag{143}$$

For the same reason as in the Epstein-Hurwitz zeta function, only the first term contributes at s=0

$$F(0; a, b, c; q) = 2(-1/2) = -1.$$
(144)

There is a formula for the derivative of this zeta function which is very useful for numerical applications due to its quick convergence. Moreover, in the case q = 0, we can give a closed

form for it. The reason is that, in this case, we can recast the zeta function in terms of the Eisenstein series $F(s; a, b, c; 0) = (cy)^{-s} E(z, s)$, where

$$E(z,s) := \sum_{(m,n)\in\mathbb{Z}^2\setminus\{(0,0)\}} \frac{y^s}{|mz+n|^{2s}} \quad \text{and} \quad z = x + iy = \frac{b + i\sqrt{\Delta}}{2c}.$$
 (145)

The latter has a simple formula for its derivative at 0 which arises as an alternative Kronecker limit formula [28]

$$\left. \frac{\partial}{\partial s} E(z,s) \right|_{s=0} = -2\ln(2\pi) - \ln(y|\eta(z)|^4). \tag{146}$$

This result and (144) in turn yield a formula for the derivative we are interested in

$$\left. \frac{\partial}{\partial s} F(s; a, b, c; 0) \right|_{s=0} = \ln(cy) - 2\ln(2\pi) - \ln(y|\eta(z)|^4) = -2\ln\left(\frac{2\pi}{\sqrt{c}}|\eta(z)|^2\right). \tag{147}$$

References

- [1] Richard P. Feynman and Paul A. M. Dirac. Feynman's Thesis A New Approach to Quantum Theory. Ed. by Laurie M Brown. WORLD SCIENTIFIC, Aug. 2005. DOI: 10.1142/5852.
- [2] Rudolf Haag. Local Quantum Physics. 2nd ed. Springer, Berlin, Heidelberg, 1996. DOI: 10.1007/978-3-642-61458-3.
- [3] Thomas Hartman. Lectures on Quantum Gravity and Black Holes. 2015. URL: http://www.hartmanhe
- [4] Jean Zinn-Justin. Quantum Field Theory and Critical Phenomena. 5th ed. Oxford University Press, 2021. DOI: 10.1093/oso/9780198834625.001.0001.
- [5] Felix M. Haehl et al. "Classification of Out-of-Time-Order Correlators". In: *SciPost Phys.* 6.1 (Jan. 7, 2019). DOI: 10.21468/SciPostPhys.6.1.001.
- [6] Edward Witten. "Quantum Field Theory and the Jones Polynomial". In: Commun. Math. Phys. 121 (Sept. 1989), pp. 351–399. DOI: 10.1007/BF01217730.
- [7] Mina Aganagic. "String Theory and Math: Why This Marriage May Last. Mathematics and Dualities of Quantum Physics". In: *Bull. Amer. Math. Soc.* 53 (2016), pp. 93–115. DOI: 10.1090/bull/1517.
- [8] Vadim Linetsky. "The Path Integral Approach to Financial Modeling and Options Pricing". In: Comput. Econ. 11 (1997), pp. 129–163. DOI: 10.1023/A:1008658226761.
- [9] James Glimm and Arthur Jaffe. Quantum Physics: A Functional Integral Point of View. 2nd ed. Springer-Verlag New York, 1987. DOI: 10.1007/978-1-4612-4728-9.
- [10] Hilbrand J. Groenewold. "On the Principles of Elementary Quantum Mechanics". In: *Physica* 12.7 (Oct. 1946), pp. 405–460. DOI: 10.1016/S0031-8914(46)80059-4.

- [11] Kevin Costello. Renormalization and Effective Field Theory. Mathematical Surveys and Monographs 170. American Mathematical Society, 2011. URL: https://bookstore.ams.org/sur
- [12] Kevin Costello and Owen Gwilliam. Factorization Algebras in Quantum Field Theory. Vol. 1. New Mathematical Monographs 31. Cambridge University Press, 2016. DOI: 10.1017/9781316678626.
- [13] Kevin Costello and Owen Gwilliam. Factorization Algebras in Quantum Field Theory. Vol. 2. 2020. URL: https://people.math.umass.edu/~gwilliam/factorization2.pdf.
- [14] Bryan Gin ge Chen et al., eds. Quantum Field Theory: Lectures of Sidney Coleman. 2018. DOI: 10.1142/9371.
- [15] Stephen W. Hawking. "Zeta Function Regularization of Path Integrals in Curved Spacetime". In: Commun. Math. Phys. 55 (1977), pp. 133–148. DOI: 10.1007/BF01626516.
- [16] Kentaro Hori et al. *Mirror Symmetry*. Clay Mathematics Monographs 1. American Mathematical Society and Clay Mathematics Institute, 2003.
- [17] Gregory W Moore. Chapter 1 Abstract Group Theory. URL: http://www.physics.rutgers.edu/~gmo
- [18] Emilio Elizalde. "An Extension of the Chowla-Selberg Formula Useful in Quantizing with the Wheeler-De Witt Equation". In: *J. Phys. A: Math. Gen.* 27 (1994), pp. 3775–3785. DOI: 10.1088/0305-4470/27/11/027.
- [19] Fiorenzo Bastianelli and Peter van Nieuwenhuizen. Path Integrals and Anomalies in Curved Space. Cambridge University Press, 2006. DOI: 10.1017/CB09780511535031.
- [20] Emilio Elizalde. Ten Physical Applications of Spectral Zeta Functions. 2nd ed. Lecture Notes in Physics 855. Springer, Berlin, Heidelberg, 2012. DOI: 10.1007/978-3-642-29405-1.
- [21] Yoni BenTov. Schwinger-Keldysh Path Integral for the Quantum Harmonic Oscillator. Feb. 9, 2021. arXiv: 2102.05029 [hep-th].
- [22] Kasia Rejzner. Perturbative Algebraic Quantum Field Theory. Mathematical Physics Studies. Springer, Cham, 2016. DOI: 10.1007/978-3-319-25901-7.
- [23] James Gordon and Gordon W. Semenoff. Schwinger Pair Production: Explicit Localization of the World-Line Instanton. Oct. 9, 2017. arXiv: 1612.05909 [hep-th].
- [24] Joseph Polchinski. "Evaluation of the One Loop String Path Integral". In: Commun. Math. Phys. 104 (1986), pp. 37–47. DOI: 10.1007/BF01210791.
- [25] Daniel Freed. "Classical Field Theory and Supersymmetry". In: Quantum Field Theory, Supersymmetry, and Enumerative Geometry. IAS/Park City Mathematics Series 11. American Mathematical Society and Institute for Advanced Study, Oct. 30, 2006, pp. 61–161. DOI: 10.1090/pcms/011/04.
- [26] Paul Ginsparg. Applied Conformal Field Theory. Nov. 11, 1988. arXiv: hep-th/9108028.
- [27] István Mező and Michael E. Hoffman. "Zeros of the Digamma Function and Its Barnes G-Function Analogue". In: Integral Transforms and Special Functions 28 (2017), pp. 846–858. DOI: 10.1080/10652469.2017.1376193.

[28] Cangxiong Chen. "On Asai's Function Analogous to $\log |\eta(z)|$ ". Doctoral thesis. University of Cambridge, 2015. DOI: 10.17863/CAM.53187.