

Characterizations of weighted core inverse in rings with involution

Tingting Li *

Abstract: R is a unital ring with involution. We investigate the characterizations and representations of weighted core inverse of an element in R by idempotents and units. For example, let $a \in R$ and $e \in R$ be an invertible Hermitian element, $n \geq 1$, then a is e -core invertible if and only if there exists an element (or an idempotent) p such that $(ep)^* = ep$, $pa = 0$ and $a^n + p$ (or $a^n(1-p) + p$) is invertible. As a consequence, let $e, f \in R$ be two invertible Hermitian elements, then a is weighted-EP with respect to (e, f) if and only if there exists an element (or an idempotent) p such that $(ep)^* = ep$, $(fp)^* = fp$, $pa = ap = 0$ and $a^n + p$ (or $a^n(1-p) + p$) is invertible. These results generalize and improve conclusions in [14].

Key words: Weighted core inverse, weighted dual core inverse, weighted EP.

AMS subject classifications: 15A09, 16B99, 46L05.

1 Introduction

Let R be a unital ring. An involution $*$ in R is an anti-isomorphism of degree 2 in R , that is to say, $(a^*)^* = a$, $(a+b)^* = a^* + b^*$ and $(ab)^* = b^*a^*$ for all $a, b \in R$. An element $a \in R$ is Hermitian if and only if $a^* = a$. And $a \in R$ is called an idempotent if $a^2 = a$. A Hermitian idempotent is said to be a projection.

Let $e, f \in R$ be two invertible Hermitian elements. We say that $a \in R$ has a weighted Moore-Penrose inverse with weights e, f if there exists an x satisfying the following four equations (see for example, [1-3]):

$$(1) \ axa = a, \quad (2) \ xax = x, \quad (3e) \ (eax)^* = ax, \quad (4f) \ (fxa)^* = xa.$$

If such an x exists, then it is called a weighted Moore-Penrose inverse of a , it is unique and denoted by $a_{e,f}^\dagger$. If $e = f = 1$, then $a_{e,f}^\dagger = a^\dagger$ is the ordinary Moore-Penrose inverse of a . The sets of all Moore-Penrose invertible elements and weighted Moore-Penrose invertible elements with weights e, f are denoted by the symbols R^\dagger and $R_{e,f}^\dagger$, respectively.

If x satisfies the equations (1) and (3e), then x is called a $\{1, 3e\}$ -inverse of a and denoted by $a^{(1,3e)}$, and a $\{1, 4f\}$ -inverse of a can be similarly defined. The symbols $a\{1, 3e\}$ and $a\{1, 4f\}$ denote the sets of all $\{1, 3e\}$ -inverses and $\{1, 4f\}$ -inverses of a , respectively. An element $a \in R$ is group invertible if there is an $x \in R$ satisfying

$$(1) \ axa = a, \quad (2) \ xax = x, \quad (5) \ ax = xa.$$

Such an x is the group inverse of a , and it is unique if it exists and denoted by $a^\#$. The sets of all group invertible, invertible, $\{1, 3e\}$ -invertible and $\{1, 4f\}$ -invertible elements in R are denoted by the symbols $R^\#, R^{-1}, R^{(1,3e)}$ and $R^{(1,4f)}$, respectively.

Mosić et al. [4] give introduced the definitions of the e -core inverse and the f -dual core inverse, which extended the concepts of the core inverse and the dual core inverse

*Tingting Li (Corresponding author E-mail: littnanjing@163.com): School of Mathematical Sciences, Yangzhou University, Yangzhou, 225002, China.

(see [5–7]). The element $a \in R$ has a weighted core inverse with weight e (or e -core inverse), if there exists $x \in R$ such that

$$axa = a, \quad xR = aR, \quad Rx = Ra^*e.$$

If such an x exists, then it is called a e -core inverse of a , it is unique and denoted by $a^{e,\oplus}$. The element $a \in R$ has a weighted dual core inverse with weight f (or f -dual core inverse), if there exists $x \in R$ such that

$$axa = a, \quad fxR = a^*R, \quad Rx = Ra.$$

If such an x exists, then it is called a f -core inverse of a , it is unique and denoted by $a_{f,\oplus}$. If $e = 1$ (resp., $f = 1$), then $a^{e,\oplus} = a^\oplus$ (resp., $a_{f,\oplus} = a_\oplus$) is the ordinary core inverse (resp., dual core inverse) of a . The sets of all e -core invertible elements and f -dual core invertible elements in R are denoted by the symbols $R^{e,\oplus}$ and $R_{f,\oplus}$, respectively. In addition, they defined that an element $a \in R$ is weighted-EP with respect to (e, f) if $a \in R^\# \cap R_{e,f}^\dagger$ and $a^\# = a_{e,f}^\dagger$. When $e = f = 1$, it is the ordinary EP element.

In this paper, we investigate the characterizations of e -core inverse, f -dual core inverse, right e -core inverse and weighted-EP with respect to (e, f) in a ring with involution. These results will generalize and improve conclusions in [14].

We will also use the following notations: $aR = \{ax \mid x \in R\}$, $Ra = \{xa \mid x \in R\}$, ${}^{\circ}a = \{x \in R \mid xa = 0\}$, $a^{\circ} = \{x \in R \mid ax = 0\}$.

Some auxiliary lemmas and results are presented for the further reference.

Lemma 1.1. [8, Proposition 7] *Let $a \in R$. $a \in R^\#$ if and only if $a = a^2x = ya^2$ for some $x, y \in R$. In this case, $a^\# = yax = y^2a = ax^2$.*

Lemma 1.2. [9, 10] *Let $a, b \in R$. $1 + ab$ is invertible if and only if $1 + ba$ is invertible.*

Lemma 1.3. [11, Proposition 2.1 and 2.2] *Let $a \in R$ and let $e \in R$ be an invertible Hermitian element. We have the following results:*

(i) *a is $\{1, 3e\}$ -invertible if and only if $a \in Ra^*ea$. Moreover, if $a = xa^*ea$ for some $x \in R$, then $x^*e \in a\{1, 3e\}$;*

(ii) *a is $\{1, 4f\}$ -invertible if and only if $a \in af^1a^*R$. Moreover, if $a = af^1a^*y$ for some $y \in R$, then $f^{-1}y^* \in a\{1, 4f\}$.*

Lemma 1.4. [4, Theorem 2.1 and 2.2] *Let $a \in R$ and let $e, f \in R$ be two invertible Hermitian elements. Then*

(I) *The following statements are equivalent:*

(i) *$a \in R^{e,\oplus}$;*

(ii) *there exists $x \in R$ such that*

$$(1) axa = a, \quad (2) xax = x, \quad (3e) (eax)^* = eax, \quad (6) xa^2 = a, \quad (7) ax^2 = x;$$

(iii) *there exists $x \in R$ such that*

$$(3e) (eax)^* = eax, \quad (6) xa^2 = a, \quad (7) ax^2 = x;$$

(iv) *$a \in R^\# \cap R^{(1,3e)}$.*

In this case,

$$a^{e,\oplus} = x = a^\#aa^{(1,3e)}.$$

(II) The following statements are equivalent:

(i) $a \in R_{f, \oplus}$;

(ii) there exists $x \in R$ such that

$$(1) axa = a, (2) xax = x, (4f) (fxa)^* = fxa, (8) a^2x = a, (9) x^2a = x;$$

(iii) there exists $x \in R$ such that

$$(4f) (fxa)^* = fxa, (8) a^2x = a, (9) x^2a = x;$$

(iv) $a \in R^\# \cap R^{(1,4f)}$.

In this case,

$$a_{f, \oplus} = x = a^{(1,4f)}aa^\#.$$

Lemma 1.5. [13] Let $a \in R$ and let $e \in R$ be an invertible Hermitian element, $n \geq 2$. Then the following statements are equivalent:

(i) $a \in Ra^*ea \cap a^nR$;

(ii) $a \in R(a^*)^nea$.

2 Main results

In this section, we present some new equivalent conditions for the existences of core inverses.

Theorem 2.1. Let $a \in R$ and let $e \in R$ be an invertible Hermitian element, $n \geq 2$. We have the following results:

(i) $a \in R^{e, \oplus}$ if and only if $a \in R(a^*)^nea \cap Ra^n$. If $a = s(a^*)^nea$ for some $s \in R$, then $a^{e, \oplus} = a^{n-1}s^*e$;

(ii) $a \in R_{f, \oplus}$ if and only if $a \in af^{-1}(a^*)^nR \cap a^nR$. If $a = af^{-1}(a^*)^nt$ for some $t \in R$, then $a_{f, \oplus} = f^{-1}t^*a^{n-1}$.

Proof. (i). According to Lemma 1.1, it is easy to deduce that $a \in R^\#$ if and only if $a \in a^nR \cap Ra^n$ for $n \geq 2$. From Lemma 1.5 we know that $a \in R(a^*)^nea$ if and only if $a \in Ra^*ea \cap a^nR$, thus

$$\begin{aligned} a \in R^{e, \oplus} &\xleftrightarrow{\text{Lemma 1.4}} a \in R^{(1,3e)} \cap R^\# \\ &\xleftrightarrow{\text{Lemma 1.3}} a \in Ra^*ea \cap R^\# \\ &\iff a \in Ra^*ea \cap a^nR \cap Ra^n \\ &\iff a \in R(a^*)^nea \cap Ra^n. \end{aligned}$$

Next, we give the representation of $a^{e, \oplus}$. Since $a \in R(a^*)^nea \cap Ra^n$, there exists $s \in R$ such that $a = s(a^*)^nea$. By Lemma 1.3, we have

$$(s(a^*)^{n-1})^*e = a^{n-1}s^*e \in a\{1, 3e\}.$$

Using Lemma 1.4, we obtain

$$a^{e, \oplus} = a^\#aa^{(1,3e)} = a^\#a(a^{n-1}s^*e) = a^{n-1}s^*e.$$

(ii). Similarly as (i). □

From the above proof, it is easy to see that Theorem 2.1 is also true in a semigroup. Next we characterize the existence of e -core inverse by idempotents and units.

Theorem 2.2. *Let $a \in R$ and let $e \in R$ be an invertible Hermitian element, $n \geq 1$ is a positive integer. The following statements are equivalent:*

- (i) $a \in R^{e, \oplus}$;
- (ii) there exists a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $u = a^n + p \in R^{-1}$;
- (iii) there exists an element s such that $(es)^* = es$, $sa = 0$, $v = a^n + s \in R^{-1}$;
- (iv) there exists a unique idempotent q such that $(eq)^* = eq$, $qa = 0$, $w = a^n(1 - q) + q \in R^{-1}$;
- (v) there exists an element t such that $(et)^* = et$, $ta = 0$, $z = a^n(1 - t) + t \in R^{-1}$.

In this case, when $n = 1$,

$$a^{e, \oplus} = u^{-1}(1 - p) = v^{-1}av^{-1} = (1 - q)w^{-1} = z^{-1}a(1 - t)z^{-1},$$

when $n \geq 2$,

$$a^{e, \oplus} = a^{n-1}u^{-1} = a^{n-1}v^{-1} = a^{n-1}(1 - q)w^{-1} = a^{n-1}(1 - t)z^{-1}.$$

Proof. (i) \Rightarrow (ii). Set $p = 1 - aa^{e, \oplus}$, we observe first that p is an idempotent satisfying $(ep)^* = ep$ and $pa = 0$. Next we show that $u = a^n + p = a^n + 1 - aa^{e, \oplus} \in R^{-1}$.

When $n = 1$, it is easy to verify

$$(a + 1 - aa^{e, \oplus})(a^{e, \oplus} + 1 - a^{e, \oplus}a) = 1 = (a^{e, \oplus} + 1 - a^{e, \oplus}a)(a + 1 - aa^{e, \oplus}),$$

thus $u = a + 1 - aa^{e, \oplus} \in R^{-1}$.

When $n \geq 2$, we prove the invertibility of u by induction on n . Since $1 + aa^{e, \oplus}(a - 1) = a + 1 - aa^{e, \oplus} \in R^{-1}$, from Lemma 1.2 we obtain that $a^2a^{e, \oplus} + 1 - aa^{e, \oplus} = 1 + (a - 1)aa^{e, \oplus} \in R^{-1}$. Therefore, $a^2 + 1 - aa^{e, \oplus} = (a^2a^{e, \oplus} + 1 - aa^{e, \oplus})(a + 1 - aa^{e, \oplus})$ is invertible. Assume that $n > 2$ and $a^{n-1} + 1 - aa^{e, \oplus} \in R^{-1}$, then $a^n + 1 - aa^{e, \oplus} = (a^2a^{e, \oplus} + 1 - aa^{e, \oplus})(a^{n-1} + 1 - aa^{e, \oplus})$ is invertible.

Finally, we prove the uniqueness of the idempotent p . It is necessary to show that ${}^\circ(a^n) = {}^\circ(1 - p)$. If $xa^n = 0$, then

$$0 = xa^n = x(1 - p)a^n = x(1 - p)(a^n + p).$$

Since $a^n + p \in R^{-1}$, we have $x(1 - p) = 0$. Conversely, if $y(1 - p) = 0$, then $ya^n = y(1 - p)a^n = 0$. Thus ${}^\circ(a^n) = {}^\circ(1 - p)$. Assume that p, p_1 are both idempotents which satisfy the statement (ii), then ${}^\circ(1 - p) = {}^\circ(a^n) = {}^\circ(1 - p_1)$. By $p \in {}^\circ(1 - p) = {}^\circ(1 - p_1)$, we obtain $p = pp_1$. Similarly, we can get $p_1 = p_1p$ from $p_1 \in {}^\circ(1 - p_1) = {}^\circ(1 - p)$. Then

$$ep = (ep)^* = (epp_1)^* = (epe^{-1}ep_1)^* = (ep_1)^*(e^{-1})^*(ep)^* = ep_1e^{-1}ep = ep_1p = ep_1,$$

thus $p = p_1$.

(ii) \Rightarrow (i). Suppose that there exists a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $u = a^n + p \in R^{-1}$.

When $n = 1$, set $x = u^{-1}(1 - p)$. Now we prove that x is the e -core inverse of a . Since $(1 - p)u = a$, $ua^2 = a$ and $pu = p$, we obtain

$$1 - p = au^{-1}, \tag{2.1}$$

$$a = u^{-1}a^2, \tag{2.2}$$

$$(1-p)u^{-1} = u^{-1} - p. \quad (2.3)$$

Therefore,

$$eax = eau^{-1}(1-p) \stackrel{(2.1)}{=} e(1-p)(1-p) = e - ep \text{ is Hermitian,}$$

$$xa^2 = u^{-1}(1-p)a^2 = u^{-1}a^2 \stackrel{(2.2)}{=} a,$$

$$\begin{aligned} ax^2 &= au^{-1}(1-p)u^{-1}(1-p) \stackrel{(2.1)}{=} (1-p)u^{-1}(1-p) \\ &\stackrel{(2.3)}{=} (u^{-1} - p)(1-p) = u^{-1}(1-p) = x. \end{aligned}$$

As a consequence, $a \in R^{e,\oplus}$ with $a^{e,\oplus} = u^{-1}(1-p)$ by Lemma 1.4.

When $n \geq 2$, from $u = a^n + p \in R^{-1}$ and $p^* = (e^{-1}ep)^* = epe^{-1}$, we get $u^* = (a^*)^n + epe^{-1} \in R^{-1}$. Since $ua = a^{n+1}$ and $u^*ea = (a^*)^n ea$,

$$a = u^{-1}a^{n+1} \in Ra^n$$

and

$$a = (u^*e)^{-1}(a^*)^n ea \in R(a^*)^n ea.$$

Thus $a \in R(a^*)^n ea \cap Ra^n$, and then $a \in R^{e,\oplus}$ and $a^{e,\oplus} = a^{n-1}(eu)^{-1}e = a^{n-1}u^{-1}$ by Theorem 2.1.

(ii) \Rightarrow (iii) is trivial.

(iii) \Rightarrow (i). Assume that there exists an element s such that $(es)^* = es$, $sa = 0$, $v = a^n + s \in R^{-1}$.

When $n = 1$. Since $v = a^n + s \in R^{-1}$ and

$$s^* = (e^{-1}es)^* = ese^{-1}, \quad (2.4)$$

we get $v^* = a^* + ese^{-1} \in R^{-1}$. From $va = a^2$ and $v^*ea = a^*ea$ we obtain

$$a = v^{-1}a^2 \in Ra^2 \quad (2.5)$$

and

$$a = (v^*e)^{-1}a^*ea \in Ra^*ea. \quad (2.6)$$

It remains to show that $a \in a^2R$. By Lemma 1.3, the equation (2.6) shows that $a \in R^{(1,3e)}$ with $v^{-1} \in a\{1, 3e\}$. Hence

$$a = aa^{(1,3e)}a = av^{-1}a. \quad (2.7)$$

In addition, $sv = s^2$ leads to $s = s^2v^{-1}$, thus

$$\begin{aligned} es^2v^{-1} &= es = (es)^* = (es^2v^{-1})^* = (v^*)^{-1}s^*es \\ &\stackrel{(2.4)}{=} (v^*)^{-1}ese^{-1}es = (v^*)^{-1}es^2. \end{aligned} \quad (2.8)$$

Moreover,

$$\begin{aligned} esv^{-2} &\stackrel{(2.8)}{=} [(v^*)^{-1}es^2]v^{-2} = (v^*)^{-1}(es^2v^{-1})v^{-1} \\ &\stackrel{(2.8)}{=} (v^*)^{-1}[(v^*)^{-1}es^2]v^{-1} = (v^*)^{-2}es^2v^{-1} \stackrel{(2.8)}{=} (v^*)^{-2}es, \end{aligned} \quad (2.9)$$

thus

$$ev^{-1} = evv^{-2} = e(a+s)v^{-2} = eav^{-2} + esv^{-2} \stackrel{(2.9)}{=} eav^{-2} + (v^*)^{-2}es. \quad (2.10)$$

Therefore,

$$a \stackrel{(2.7)}{=} av^{-1}a = ae^{-1}(ev^{-1})a \stackrel{(2.10)}{=} ae^{-1}[eav^{-2} + (v^*)^{-2}es]a = a^2v^{-2}a \in a^2R. \quad (2.11)$$

According to Lemma 1.1, equations (2.5) and (2.11) show that $a \in R^\#$ with $a^\# = v^{-2}a$. And from (2.6) and Lemma 1.4, $a \in R^{e, \oplus}$ with

$$a^{e, \oplus} = a^\#aa^{(1,3e)} = v^{-2}aav^{-1} \stackrel{(2.5)}{=} v^{-1}av^{-1}.$$

When $n \geq 2$, from $v = a^n + s \in R^{-1}$ and $s^* = ese^{-1}$, we get $v^* = (a^*)^n + ese^{-1} \in R^{-1}$. Since $va = a^{n+1}$ and $v^*ea = (a^*)^n ea$,

$$a = v^{-1}a^{n+1} \in Ra^n$$

and

$$a = (v^*e)^{-1}(a^*)^n ea \in R(a^*)^n ea.$$

Thus $a \in R(a^*)^n ea \cap Ra^n$, and then $a \in R^{e, \oplus}$ and $a^{e, \oplus} = a^{n-1}(ev)^{-1}e = a^{n-1}v^{-1}$ by Theorem 2.1.

(i) \Rightarrow (iv). Set $q = 1 - aa^{e, \oplus}$, then q is an idempotent satisfying $(eq)^* = eq$ and $qa = 0$. Now we prove that $w = a^n(1 - q) + q = a^{n+1}a^{e, \oplus} + 1 - aa^{e, \oplus} \in R^{-1}$.

When $n = 1$, from the proof of (i) \Rightarrow (ii) we know that $1 + aa^{e, \oplus}(a-1) = a + 1 - aa^{e, \oplus} \in R^{-1}$, thus $w = a^2a^{e, \oplus} + 1 - aa^{e, \oplus} = 1 + (a-1)aa^{e, \oplus} \in R^{-1}$ by the Lemma 1.2.

When $n \geq 2$, from the proof of (i) \Rightarrow (ii) we know that $1 + aa^{e, \oplus}(a^n - 1) = a^n + 1 - aa^{e, \oplus} \in R^{-1}$, thus $w = a^{n+1}a^{e, \oplus} + 1 - aa^{e, \oplus} = 1 + (a^n - 1)aa^{e, \oplus} \in R^{-1}$ by the Lemma 1.2.

In the end, we prove the uniqueness of the idempotent q . It is necessary to show that ${}^\circ(a^n) = {}^\circ(1 - q)$. If $xa^n = 0$, then

$$0 = xa^n(1 - q) = x(1 - q)[a^n(1 - q) + q].$$

From $a^n(1 - q) + q \in R^{-1}$ we have $x(1 - q) = 0$. Conversely, if $y(1 - q) = 0$, then $ya^n = y(1 - q)a^n = 0$. Hence ${}^\circ(a^n) = {}^\circ(1 - q)$. Suppose that q, q_1 are both idempotents which satisfy the statement (iv), then ${}^\circ(1 - q) = {}^\circ(a^n) = {}^\circ(1 - q_1)$. From $q \in {}^\circ(1 - q) = {}^\circ(1 - q_1)$ we obtain $q = qq_1$. Similarly, $q \in {}^\circ(1 - q_1) = {}^\circ(1 - q)$ leads to $q_1 = q_1q$. Then

$$eq = (eq)^* = (eqq_1)^* = (eqe^{-1}eq_1)^* = (eq_1)^*(e^{-1})^*(eq)^* = eq_1e^{-1}eq = eq_1q = eq_1,$$

thus $q = q_1$.

(iv) \Rightarrow (i). Assume that there exists a unique idempotent q such that $(eq)^* = eq$, $qa = 0$, $w = a^n(1 - q) + q \in R^{-1}$.

When $n = 1$, let $x = (1 - q)w^{-1}$. Now we show that x is the e -core inverse of a . Since $(1 - q)w = a(1 - q)$ and $wa^2 = a$, we get

$$1 - q = a(1 - q)w^{-1}, \quad (2.12)$$

$$a = w^{-1}a^2. \quad (2.13)$$

Therefore,

$$eax = ea(1 - q)w^{-1} \stackrel{(2.12)}{=} e(1 - q) = e - eq \text{ is Hermitian,}$$

$$xa^2 = (1 - q)w^{-1}a^2 \stackrel{(2.13)}{=} (1 - q)a = a,$$

$$ax^2 = a(1 - q)w^{-1}(1 - q)w^{-1} \stackrel{(2.12)}{=} (1 - q)w^{-1} = x.$$

As a consequence, $a \in R^{e, \oplus}$ with $a^{e, \oplus} = (1 - q)w^{-1}$ by Lemma 1.4.

When $n \geq 2$, from $w = a^n(1 - q) + q \in R^{-1}$ and $q^* = (e^{-1}eq)^* = eqe^{-1}$, we get $w^* = (1 - p)^*(a^*)^n + eqe^{-1} \in R^{-1}$. Since $wa = a^{n+1}$ and $w^*ea = (1 - q)^*(a^*)^n ea$,

$$a = w^{-1}a^{n+1} \in Ra^n$$

and

$$a = (w^*e)^{-1}(1 - q)^*(a^*)^n ea \in R(a^*)^n ea.$$

Therefore, $a \in R(a^*)^n ea \cap Ra^n$, and then $a \in R^{e, \oplus}$ and $a^{e, \oplus} = a^{n-1}(1 - q)(ew)^{-1}e = a^{n-1}(1 - q)w^{-1}$ by Theorem 2.1.

(iv) \Rightarrow (v) is trivial.

(v) \Rightarrow (i). Assume that there exists an element t such that $(et)^* = et$, $ta = 0$, $z = a^n(1 - t) + t \in R^{-1}$.

When $n = 1$. Since $z = a^n(1 - t) + t \in R^{-1}$ and

$$t^* = (e^{-1}et)^* = ete^{-1}, \quad (2.14)$$

we get $z^* = (1 - t)^*a^* + ese^{-1} \in R^{-1}$. From $za = a^2$ and $z^*ea = (1 - t)^*a^*ea$ we obtain

$$a = z^{-1}a^2 \in Ra^2 \quad (2.15)$$

and

$$a = (z^*e)^{-1}(1 - t)^*a^*ea \in Ra^*ea. \quad (2.16)$$

It remains to show that $a \in a^2R$. By Lemma 1.3, the equation (2.16) shows that $a \in R^{(1, 3e)}$ with $(1 - t)z^{-1} \in a\{1, 3e\}$. Hence

$$a = aa^{(1, 3e)}a = a(1 - t)z^{-1}a. \quad (2.17)$$

In addition, $tz = t^2$ leads to $t = t^2z^{-1}$, thus

$$\begin{aligned} et^2z^{-1} &= et = (et)^* = (et^2z^{-1})^* = (z^*)^{-1}t^*et \\ &\stackrel{(2.14)}{=} (z^*)^{-1}ete^{-1}et = (z^*)^{-1}et^2. \end{aligned} \quad (2.18)$$

Moreover,

$$\begin{aligned} etz^{-2} &\stackrel{(2.18)}{=} [(z^*)^{-1}et^2]z^{-2} = (z^*)^{-1}(et^2z^{-1})z^{-1} \\ &\stackrel{(2.18)}{=} (z^*)^{-1}[(z^*)^{-1}et^2]z^{-1} = (z^*)^{-2}et^2z^{-1} \stackrel{(2.18)}{=} (z^*)^{-2}et, \end{aligned} \quad (2.19)$$

thus

$$\begin{aligned} ez^{-1} &= ezz^{-2} = e[a(1-t) + t]z^{-2} = ea(1-t)z^{-2} + etz^{-2} \\ &\stackrel{(2.19)}{=} ea(1-t)z^{-2} + (z^*)^{-2}et. \end{aligned} \quad (2.20)$$

Therefore,

$$\begin{aligned} a &\stackrel{(2.17)}{=} a(1-t)z^{-1}a = a(1-t)e^{-1}(ez^{-1})a \\ &\stackrel{(2.20)}{=} a(1-t)e^{-1}[ea(1-t)z^{-2} + (z^*)^{-2}et]a \\ &= a^2(1-t)z^{-2}a \in a^2R. \end{aligned} \quad (2.21)$$

According to Lemma 1.1, equations (2.15) and (2.21) show that $a \in R^\#$ with $a^\# = z^{-2}a$. And from (2.16) and Lemma 1.4 deducing that $a \in R^{e, \oplus}$ with

$$a^{e, \oplus} = a^\#aa^{(1,3e)} = z^{-2}aa(1-t)z^{-1} \stackrel{(2.15)}{=} z^{-1}a(1-t)z^{-1}.$$

When $n \geq 2$, from $z = a^n(1-t) + t \in R^{-1}$ and $t^* = ete^{-1}$ we get $z^* = (1-t)^*(a^*)^n + ete^{-1} \in R^{-1}$. Since $za = a^{n+1}$ and $z^*ea = (1-t)^*(a^*)^nea$,

$$a = z^{-1}a^{n+1} \in Ra^n$$

and

$$a = (z^*e)^{-1}(1-t)^*(a^*)^nea \in R(a^*)^nea.$$

Thus $a \in R(a^*)^nea \cap Ra^n$, and then $a \in R^{e, \oplus}$ and $a^{e, \oplus} = a^{n-1}(1-t)(ev)^{-1}e = a^{n-1}(1-t)v^{-1}$ by Theorem 2.1. \square

When $e = 1$, we get the following result, which improves Theorem 3.3 and Theorem 3.4 in [14].

Corollary 2.3. *Let $a \in R$, $n \geq 1$ is a positive integer. The following statements are equivalent:*

- (i) $a \in R^{\oplus}$;
- (ii) there exists a unique projection p such that $pa = 0$, $u = a^n + p \in R^{-1}$;
- (iii) there exists a Hermitian element s such that $sa = 0$, $v = a^n + s \in R^{-1}$;
- (iv) there exists a unique projection q such that $qa = 0$, $w = a^n(1-q) + q \in R^{-1}$;

(v) there exists a Hermitian element t such that $ta = 0$, $z = a^n(1 - t) + t \in R^{-1}$.
In this case, when $n = 1$,

$$a^{\oplus} = u^{-1}(1 - p) = v^{-1}av^{-1} = (1 - q)w^{-1} = z^{-1}a(1 - t)z^{-1},$$

when $n \geq 2$,

$$a^{\oplus} = a^{n-1}u^{-1} = a^{n-1}v^{-1} = a^{n-1}(1 - q)w^{-1} = a^{n-1}(1 - t)z^{-1}.$$

There is an analogous result for the f -dual core inverse of $a \in R$, we omit its proof.

Theorem 2.4. Let $a \in R$ and let $f \in R$ be an invertible Hermitian element, $n \geq 1$ is a positive integer. The following statements are equivalent:

- (i) $a \in R_{f, \oplus}$;
 - (ii) there exists a unique idempotent p such that $(fp)^* = fp$, $ap = 0$, $u = a^n + p \in R^{-1}$;
 - (iii) there exists an element s such that $(fs)^* = fs$, $as = 0$, $v = a^n + s \in R^{-1}$;
 - (iv) there exists a unique idempotent q such that $(fq)^* = fq$, $aq = 0$, $w = (1 - q)a^n + q \in R^{-1}$;
 - (v) there exists an element t such that $(ft)^* = ft$, $at = 0$, $z = (1 - t)a^n + t \in R^{-1}$.
- In this case, when $n = 1$,

$$a_{f, \oplus} = (1 - p)u^{-1} = v^{-1}av^{-1} = w^{-1}(1 - q) = z^{-1}(1 - t)az^{-1},$$

when $n \geq 2$,

$$a_{f, \oplus} = u^{-1}a^{n-1} = v^{-1}a^{n-1} = (1 - q)a^{n-1}w^{-1} = (1 - t)a^{n-1}z^{-1}.$$

When $f = 1$, we get the following result.

Corollary 2.5. Let $a \in R$, $n \geq 1$ is a positive integer. The following statements are equivalent:

- (i) $a \in R_{\oplus}$;
- (ii) there exists a unique projection p such that $ap = 0$, $u = a^n + p \in R^{-1}$;
- (iii) there exists a Hermitian element s such that $as = 0$, $v = a^n + s \in R^{-1}$;
- (iv) there exists a unique projection q such that $aq = 0$, $w = (1 - q)a^n + q \in R^{-1}$;
- (v) there exists a Hermitian element t such that $at = 0$, $z = (1 - t)a^n + t \in R^{-1}$.

In this case, when $n = 1$,

$$a_{\oplus} = (1 - p)u^{-1} = v^{-1}av^{-1} = w^{-1}(1 - q) = z^{-1}(1 - t)az^{-1},$$

when $n \geq 2$,

$$a_{\oplus} = u^{-1}a^{n-1} = v^{-1}a^{n-1} = (1 - q)a^{n-1}w^{-1} = (1 - t)a^{n-1}z^{-1}.$$

In the derivation of (i) from (ii) of Theorem 2.2, when $n = 1$, $a^{\oplus} = u^{-1}(1 - p)$. From the equation (2.1) (i.e., $1 - p = au^{-1}$) we know that $e(1 - p) = eau^{-1}$ is Hermitian, thus

$$\begin{aligned} a^{\oplus} &= u^{-1}(1 - p) = u^{-1}au^{-1} = u^{-1}e^{-1}eau^{-1} = u^{-1}e^{-1}(eau^{-1})^* \\ &= (u^*eu)^{-1}a^*e = (a^*ea + ep)^{-1}a^*e. \end{aligned}$$

Through this expression, we naturally want to know whether a is e -core invertible or not when there is a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $a^*ea + ep \in R^{-1}$. However, it is not true. Here is a counterexample.

Example 2.6. Let R be an infinite matrix ring over complex field whose rows and columns are both finite, let conjugate transpose be the involution and $a = \sum_{i=1}^{\infty} e_{i+1,i}$. Then $a^*a = 1$, $aa^* = \sum_{i=2}^{\infty} e_{i,i}$. Set $e = 1$, $p = 0$, then p is an idempotent satisfying $(ep)^* = ep$, $pa = 0$ and $a^*ea + ep = 1 \in R^{-1}$. But a is not group invertible, thus a is not e -core invertible.

The above counterexample shows that even if there is a unique idempotent p such that $(ep)^* = ep$, $pa = 0$ and $a^*ea + ep \in R^{-1}$, a is not necessary to be e -core invertible in general rings. However, it is true when we take R as a Dedekind-finite ring which satisfies the property that $ab = 1$ implies $ba = 1$ for any $a, b \in R$. Let R_r^{-1} and R_l^{-1} denote all right invertible and left invertible elements in R , respectively. See the following result.

Theorem 2.7. Let R be a Dedekind-finite ring and $a \in R$, $e \in R$ is an invertible Hermitian element. Then the following statements are equivalent:

- (i) $a \in R^{e,\oplus}$;
 - (ii) there exists a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $a^*ea + ep \in R^{-1}$;
 - (iii) there exists a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $a^*ea + ep \in R_r^{-1}$;
 - (iv) there exists a unique idempotent p such that $(ep)^* = ep$, $pa = 0$, $a^*ea + ep \in R_l^{-1}$.
- In this case,

$$a^{e,\oplus} = (a^*ea + ep)^{-1}a^*e.$$

Proof. Since $a^*ea + ep$ is Hermitian, $a^*ea + ep$ is one-sided invertible if and only if it is invertible, hence the statements (ii), (iii) and (iv) are equivalent. It is sufficient to show that (i) \Leftrightarrow (ii).

(i) \Rightarrow (ii). Suppose that $a \in R^{e,\oplus}$ and let $p = 1 - aa^{e,\oplus}$, then $a + p \in R^{-1}$ by Theorem 2.2. Moreover, $e(a + p) = ea + ep \in R^{-1}$, so $(ea + ep)^* = a^*e + ep \in R^{-1}$. Therefore $a^*ea + ep = (a^*e + ep)(a + p)$ is invertible.

(ii) \Rightarrow (i). Let $u = a + p$, $u^*eu = a^*ea + ep \in R^{-1}$. As R is a Dedekind-finite ring, thus $u \in R^{-1}$, which guarantees $a \in R^{e,\oplus}$ and $a^{e,\oplus} = (a^*ea + ep)^{-1}a^*e$ by Theorem 2.2. \square

There is a dual result for the f -dual core inverse.

Theorem 2.8. Let R be a Dedekind-finite ring and $a \in R$, $f \in R$ is an invertible Hermitian element. Then the following statements are equivalent:

- (i) $a \in R^{e,\oplus}$;
 - (ii) there exists a unique idempotent q such that $(fq)^* = fq$, $aq = 0$, $af^{-1}a^* + qf^{-1} \in R^{-1}$;
 - (iii) there exists a unique idempotent q such that $(fq)^* = fq$, $aq = 0$, $af^{-1}a^* + eqf^{-1} \in R_r^{-1}$;
 - (iv) there exists a unique idempotent q such that $(fq)^* = fq$, $aq = 0$, $af^{-1}a^* + qf^{-1} \in R_l^{-1}$.
- In this case,

$$a^{e,\oplus} = f^{-1}a^*(af^{-1}a^* + qf^{-1})^{-1}.$$

Let $a \in R$ and $e, f \in R$ be two invertible Hermitian elements. Zhu and Wang [11] mentioned that $a \in R^{e,\oplus} \cap R_{f,\oplus}$ if and only if $a \in R^\# \cap R_{e,f}^\dagger$. Mosić et al. [4] mentioned that $a \in R$ is weighted-EP with respect to (e, f) if and only if $a \in R^{e,\oplus} \cap R_{f,\oplus}$ and $a^{e,\oplus} = a_{f,\oplus}$. By Theorem 2.2 and Theorem 2.4, we obtain the following theorem.

Theorem 2.9. *Let $a \in R$ and $e, f \in R$ be two invertible Hermitian elements, $n \geq 1$ is a positive integer. The following conditions are equivalent:*

- (i) $a \in R$ is weighted-EP with respect to (e, f) ;
- (ii) there exists a unique idempotent p such that $(ep)^* = ep$, $(fp)^* = fp$, $ap = pa = 0$, $a^n + p \in R^{-1}$;
- (iii) there exists an element s such that $(es)^* = es$, $(fs)^* = fs$, $as = sa = 0$, $a^n + s \in R^{-1}$.

Proof. (i) \Rightarrow (ii). Suppose a is weighted-EP with respect to (e, f) , so $a \in R_{e,f}^\dagger \cap R^\#$ and $a_{e,f}^\dagger = a^\#$. Let $p = 1 - a^\#a = 1 - a_{e,f}^\dagger a$, it is easily seen that p is an idempotent satisfying $(ep)^* = ep$, $(fp)^* = fp$, $pa = ap = 0$. Since

$$(a^n + 1 - a^\#a)((a^\#)^n + 1 - a^\#a) = 1 = ((a^\#)^n + 1 - a^\#a)(a^n + 1 - a^\#a),$$

thus $a^n + 1 - a^\#a$ is invertible.

(ii) \Rightarrow (iii) is trivial.

(iii) \Rightarrow (i). According to Theorem 2.2 and Theorem 2.4, $a \in R^{e,\oplus} \cap R_{f,\oplus}$. It remains to show that $a^{e,\oplus} = a_{f,\oplus}$. Write $v = a^n + s$. When $n = 1$, we have $a^{e,\oplus} = v^{-1}av^{-1} = a_{f,\oplus}$. When $n \geq 2$, $a^{e,\oplus} = a^{n-1}v^{-1}$, $a_{f,\oplus} = v^{-1}a^{n-1}$. From $pa = ap = 0$, we obtain $av = va$, thus $av^{-1} = v^{-1}a$. Therefore, $a^{e,\oplus} = a^{n-1}v^{-1} = v^{-1}a^{n-1} = a_{f,\oplus}$. As a consequence, a is weighted-EP with respect to (e, f) . \square

ACKNOWLEDGMENTS

This research is supported by the National Natural Science Foundation of China (No.11871145); NSF of Jiangsu Province (BK20200944), Natural Science Foundation of Jiangsu Higher Education Institutions of China (20KJB110001); the QingLan Project of Jiangsu Province.

References

- [1] Bhaskara Rao, K. P. S. (2002). Generalized Inverses over Commutative Rings. London: Taylor and Francis, Ltd.
- [2] Wei, Y. (2014). Generalized inverse of matrices. In: Hogben, L., 2nd ed. Boca Raton: CRC Press.
- [3] Mosić, D., Djordjević, D. S. (2011). Weighted-EP elements in C^* -algebras. Electron. J. Linear Algebra 22:912-930.
- [4] Mosić, D., Deng, C.Y., Ma, H.F.(2018). On a weighted core inverse in a ring with involution. Comm. Algebra. 46: 2332-2345.
- [5] Baksalary O.M.(2010). Trenkler G. Core inverse of matrices. Linear Multilinear Algebra. 58: 681-697.
- [6] Rakić D.S., Dinčić N.Č., Djordjević D.S.(2014). Group, Moore-Penrose, core and dual core inverse in rings with involution. Linear Algebra Appl. 463: 115-133.
- [7] Xu S.Z., Chen J.L., Zhang X.X.(2017). New characterizations for core and dual core inverses in rings with involution. Front. Math. China. 12(1): 231-246.

- [8] Hartwig R.E.(1976). Block generalized inverses. Arch. Retional Mech. Anal. 61: 197-251.
- [9] Barnes B.A.(1998). Common operator properties of the linear operators RS and SR. Proc. Amer. Math. Soc. 126: 1055-1061.
- [10] Cvetković-Ilić D.S.(2007). Harte R.E. On the algebraic closure in rings. Proc. Amer. Math. Soc. 135: 3547-3582.
- [11] Zhu H. H., Wang Q. W. (2021). Weighted Moore-Penrose inverses and weighted core inverses in rings with involution. Chin. Ann. Math. Ser. B. 2021(42)(4): 613-624.
- [12] Wang L., Mosić D. (2021). The one-sided inverse along two elements in rings. Linear Multilinear Algebra. 69(13): 2410-2422.
- [13] Wang L., Mosić D., Gao Y.F.(2019). One sided core inverse and the related generalized inverse. Comm. Algebra. 47(11):4749-4762.
- [14] Li, T.T., Chen, J.L.(2018). Characterizations of core and dual core inverses in rings with involution. Linear Multilinear Algebra. 66: 717-730.