

Single-photon steering

L M Arévalo Aguilar^{a)}

Facultad de Ciencias Físico Matemáticas, Benemérita Universidad Autónoma de Puebla. 18 Sur y Avenida San Claudio, Col. San Manuel, C.P: 72520, Puebla, Pue., Mexico

(Dated: 19 January 2022)

According to Schrödinger, the laws of quantum mechanics *obliges us to admit that by suitable measurement taken on one of the two system only*¹ the state of the other system can not only be determined but steered too. That is, it conveys the potential ability to steer the state of another physical system without interacting with it by implementing independent measurements, this nonlocal phenomenon was named *steering*. On the other hand, the first proposals about the nonlocality of a single photon focus on showing the Bell nonlocality by using a single photon path entanglement. This path entanglement of a single photon was also used for analyzing and experimentally produce the steering of a single photon. However, these established facts have been recently called into question suggesting that single-photon entanglement is not non-local. In this letter, we show that by incorporating and manipulating the internal degrees of freedom of the photon, together with the external path, it is easy to demonstrate the nonlocal effect of steering of a single photon's state. In this sense, the experimental set-up that we propose differs from the one reported in the quantum optics literature which only uses, to the best of our knowledge, the path entanglement of photons for showing the steering phenomenon, i.e. here we exploit the entanglement between the internal and external degrees of freedom of the photon to show this nonlocal effect. The introduction of the photon's internal degrees of freedom in the experimental set-up gives us new insight, advantages and possibilities to control the nonlocal character of the single-photon entangled state.

Keywords: quantum mechanics, steering, single-photon entanglement, quantum non-locality

^{a)}Electronic mail: larevalo@fcfm.buap.mx

I. INTRODUCTION

The steering of quantum states was pointed out by Schrödinger when he was analyzing the *disconcerting fact* that *the particular choice of observation* affects the wavefunction of the other system even if the measurement was made on one system only², the previous situation arises from the Einstein-Podolsky-Rosen analysis³. This investigation was pursued in a second paper where he established the steering phenomenon as a *more complete* control, *with the indirect measurement*, of a second system distantly situated¹. Additionally, in 2007 steering received an operational definition in quantum information theory as the Alice's task to convince another distrustful party that she could share an entangled state⁴. Additionally, there it was demonstrated that steering is a quantum correlation that is stronger than non separability but weaker than Bell nonlocality⁴, it shows that local hidden state models are unable to characterize the conditional states at one party. Besides, it was shown that steering together with the uncertainty principle underlies nonlocality⁵, and that steering is unveiled by measurements of observables that are not jointly measurable^{6,7}. Consequently, nowadays we can say that the nonlocal steering phenomenon is a confirmed fact.

It is worth remembering that the first time when the wave particle duality was proposed was using light, this was given in the Einstein-Planck relations $p = \hbar k$ and $E = \hbar \omega$. The second time was when d'Broglie proposed that such equations should be used also for matter. Similarly, the first time that nonlocal quantum correlations was proposed for a single quantum system was the proposal that single-photon entanglement really possesses nonlocal properties. This was theoretically proved first by Tan, Walls and Collet (TWC)⁸, see also⁹, and experimentally proved, free from the detection loophole, by Guerreiro et al.¹⁰, where it was clearly demonstrated that single-photon entanglement generates single-photon steering, see also¹¹.

However, in recent works this possibility has been denied¹³⁻¹⁹. The perspective employed was conceived using as example the entanglement between the internal degrees of freedom of individual photons¹³. Nevertheless, it is not always possible to generalize from a particular photon's state to all entangled states and scenarios. Besides, the nonlocal effects of the single-particle entanglement might be better understood by means of an intelligible experiment, where single-photon steering could be implemented by using a photon's state that is equivalent to the one analyzed in the single-particle steering case¹², where the steering

of a single-particle traversing the Stern-Gerlach experiment (SGE) was proposed. Thereby, the experiment shown below could be thought as an experimental set-up which is easier to implement than the SGE and additionally it could indirectly corroborate the steering in the SGE because the entangled states are strongly similar. To attain this, the spreading (over large distances) of the entangled wavefunction of the single-photon is employed. Particularly, it will be clear to notice that the nonlocality of the single-photon entangled state allows the photon to "know" the kind of detector that Alice is using to steer Bob's state.

In short, some authors claim that the single-particle entanglement lacks nonlocality¹³: *single-particle entanglement refers to correlations of two different degrees of freedom of an individual particle; this situation cannot lead to nonlocal correlations*, see also¹⁴. A similar assertion has been made by other different authors^{15–18} taking as main sources the references¹³ and¹⁹. For example, in a recent paper the authors claim that entangled states between *two different degrees of freedom of one and the same object* are not *non-local*¹⁵ (these authors allude to the state¹⁵ $|\psi\rangle = (|x_1\rangle|\uparrow\rangle + |x_2\rangle|\downarrow\rangle)/\sqrt{2}$ experimentally created by Wineland et al.²⁰, this state, *is hence prototypic for 'local entanglement' of states which cannot be separated spatially*); this point of view is shared by highly influential journals stating that single-particle entanglement is local only^{16–18}, i.e. lacking nonlocal effects; additionally see a recent paper that also endorses the claim that single-particle entanglement just serves to text noncontextual reality models²¹.

Hence, the previous paragraphs strongly suggest that nonlocal correlations can only occur in entangled states comprising two quantum systems. Additionally, it may result counter-intuitive to name a phenomenon *steering*, which implies nonlocal correlations, when it comes from a single-particle. However, the fact is that this effect was theoretically proved¹² and in the case of single-photon steering it was experimentally proved also¹⁰.

By restricting oneself to the particular kind of entangled states considered in reference¹³, i.e. bipartite entanglement between the polarization and the orbital angular momentum (OAM) of photons, it could lead us to conclude that single-particle entanglement is not nonlocal. However, this assertion based on a particular kind of bipartite entangled states could not be generalized to all entangled states or all scenarios. More important, if we remain embracing the idea that single-particle entangled states do not possess non-local correlations, then many of the non-local task such as quantum key distribution might not be investigated or even implemented by using these resources.

In recent years world leading quantum groups have made a great effort to experimentally demonstrate single-photon steering¹². For example, Fuwa et al. demonstrated the violation of a steering inequality employing a single-photon entangled state by implementing homodyne detection²², additionally see also^{11,23–25}; and Guerreiro et al. have proven the violation of a new steering inequality (employing a single-photon entangled state) free of detection loopholes using as the set of assemblage the eigenstates of the displaced operator¹⁰. Besides, recently it was theoretically shown that single-particle entanglement between an internal DoF and an external DoF produces steering in atoms crossing the famous Stern-Gerlach experiment¹². These experiments^{10,11,22–25} together with the theoretical proposals given in references^{12,26} cast doubts on whether the assertion that *single-particle entanglement cannot produce nonlocal quantum correlations* is a valid generalization.

In this letter, we show that the nonlocal effect of single-photon steering could be produced by using an alternative experimental scheme with a state that is similar to the one that was proposed in the single-particle steering case¹²; that is to say, the entanglement between an internal degree of freedom of the photon and its external path. In particular, from this scheme it is easy to see that the nonlocality of the single-photon entangled state allows Alice to steer Bob's state depending on the kind of measurement apparatuses that she employs. The states that we use in this letter differ from the state used in the first proposal of single photon nonlocality by TWC⁸ who use as an initial state $|1\rangle_u |0\rangle_v$, which is transformed after the first beam splitter as: $|1\rangle_{b_1} |0\rangle_{b_2} + |0\rangle_{b_1} |1\rangle_{b_2}$; and the state used by Hardy⁹ who uses as an initial state $(|1\rangle_s + |0\rangle_s) |0\rangle_t$, which is transformed after a first beam splitter as: $q |0\rangle_{u_1} |0\rangle_{u_2} + \frac{ir}{\sqrt{2}} |1\rangle_{u_1} |0\rangle_{u_2} + \frac{r}{\sqrt{2}} |0\rangle_{u_1} |1\rangle_{u_2}$, due to the fact that they use the entanglement in the photon path only. In this work, alternatively, instead of using the single-photon path entanglement only, we devise a proposal which take into account the internal degree of freedom of the photon and we show that the steering of the single-photon is clearly and effortlessly conceived. On the other hand, in recent interesting works by Das et al.^{27,28}, who also employ the entangled path state $|1\rangle_{b_1} |0\rangle_{b_2} + |0\rangle_{b_1} |1\rangle_{b_2}$, it is argued that the experimental set-up (i.e. the measurement parts) used by TWC does not reliable test the violation of the Bell inequality and that a local hidden variable model could reproduce the thought nonlocal events considered by TWC; besides, these authors claim that the initial Hardy's state and his proposed experimental set-up could reliable produce nonlocal effects²⁷. Essentially, the works by TWC, Hardy and Das et al. rely on the path entanglement of the single photon

(although they differ on the measurement set-up), whereas the proposal carried out in this letter relies on the entanglement between an internal degree of freedom of the photon with the potential paths that the photon follows. Notice that what Das et al. demonstrated is that the experimental set-up together with the kind of Bell inequality that was used by TWC do not allow a test of the violation of the CHCH-Bell inequality due to both: i) the assumption that the total intensity given by $I_j(\lambda) = I_{c_j}(\theta_j, \lambda) + I_{d_j}(\theta_j, \lambda)$ is independent of the local value of θ_j , and ii) the use of a constant amplitude of the weak local field used in the homodyne measurement, see²⁷ for details. Besides, it is worth of mention that the works carried out in references^{10,22} are variants of the TWC analysis.


On the other hand, in references^{12,29} it was proved that the entanglement between the internal and external degrees of freedom of a particle violates the CHSH-Bell inequality; hence, as the state that we employ in this letter is a similar entangled state between the internal and external degrees of freedom it would also violate such inequality. Hence, we will exploit the entanglement between the internal and external degrees of freedom to show the steering of a single photon state; this approach, to the best of our knowledge, has not been previously proposed yet.

II. THE SCHEME

The scheme that is used to help the explanation of the single-photon steering is shown in Fig. (1). It takes as base the schemes given in references^{12,30} and it has as a main goal to prepare the state given by the following equation:

$$|\psi\rangle_{\text{photon}} = \frac{1}{\sqrt{2}} (|a\rangle |H\rangle + |b\rangle |V\rangle), \quad (1)$$

where $|a\rangle$ and $|b\rangle$ are the probable paths that the photon could follow, towards the cities of New York or Puebla; $|H\rangle$ and $|V\rangle$ are the vertical and horizontal polarizations of the photon, respectively. Usually, in quantum optics $|a\rangle = |0\rangle |1\rangle$ and $|b\rangle = |1\rangle |0\rangle$. Eq. (1) represents a single-photon entangled state where the dichotomic entangled variables are the orthogonal polarizations $|H\rangle$ and $|V\rangle$ and the spatial modes $|a\rangle$ and $|b\rangle$ —as the phase produced by the reflection in the Beam splitter could be compensated by a linear optical component we do not write it down—. The preparation of the single-photon entangled state given by Eq. (1) is as follows: First, a single photon is generated by using a postselection method where a



$$|\psi\rangle_{\text{photon}} = \frac{1}{\sqrt{2}} \left(|a\rangle |H\rangle + |b\rangle |V\rangle \right)$$

FIG. 1. A single-photon steering experiment, where Alice is able to steer Bob's state at a spacelike distance between the cities of New York and Puebla.

type-II BBO crystal is pumped creating two orthogonal polarized photons propagating in the same direction, then a Polarized Beam Splitter, PBS-1 in the Fig. (1), divert the vertical polarized photon towards the trigger detector T. Hence, the trigger signal indicates that a single photon with horizontal polarization was created. Second, the half-wave plate (HWP) rotates the polarization of the horizontal polarized photon to 45° just before the second PBS, then the latter beam splitter (PBS) produces the entangled state given by Eq. (1).

The state given by Eq. (1) is an entangled state between the internal (polarization) and external (path) DoFs of the photon, it is equivalent to the following entangled state given in references^{31,32}:

$$|\psi(t)\rangle = C_0 M(x, y) \left(|\psi_+\rangle |\uparrow_z\rangle + |\psi_-\rangle |\downarrow_z\rangle \right), \quad (2)$$

where

$$\langle z | \psi_\pm \rangle = e^{\frac{\mp i t \mu_c}{\hbar} (B_0 + b z)} \exp \left[\frac{-1}{4(\sigma_0^2 + i \hbar / 2m)} \left(z \pm \frac{t^2 \mu_c b}{2m} \right)^2 \right], \quad (3)$$

$$C_0 = e^{\frac{-i t^3 \mu_c^2 b^2}{6 m \hbar}} \frac{1}{\sqrt{2}} \left[\frac{\sigma_0}{(2\pi)^{1/2}} \right]^{3/2} \left(\sigma_0^2 + \frac{i \hbar t}{2m} \right)^{-3/2}, \quad (4)$$

and $M(x, y) = e^{-\sigma_0^2 k_y^2} e^{\frac{4y\sigma_0^2 k_y}{4(\sigma_0^2 + i\hbar/2m)}} \exp \left[\frac{-(x^2 + y^2 - 4\sigma_0^4 k_y^2)}{4(\sigma_0^2 + i\hbar/2m)} \right]$. This state is generated in the famous Stern-Gerlach experiment, and it represents an entangled state between the internal and external DoFs of a single particle. It is worthy of mention that it was already demonstrated that the entangled state given by this equation violates the Bell's Clauser–Horne–Shimony–Holt (CHSH-Bell) kind inequality²⁹.

The entangled state of the single-photon given by Eq. (1) resembles the Einstein's boxes with entangled states given in reference¹², and it allows Alice to steer Bob's states. To perceive this, notice that if Alice decided to measure the polarization of her system then the following two possibilities would occur:

1. If Alice got as result that her photon has a vertical polarization, the wave function would collapse towards $|b\rangle |V\rangle$, with the photon at New York city. That is, as $|b\rangle = |1\rangle |0\rangle$, Alice gets the $|1\rangle$ state and Bob gets the $|0\rangle$ state.
2. However, if Alice detected nothing, the state vector would collapse towards $|a\rangle |H\rangle$ with the photon at Puebla city. That is, as $|a\rangle = |0\rangle |1\rangle$, Alice gets the $|0\rangle$ state and Bob gets the $|1\rangle$ state.

The effect of "detect nothing" could be explained in terms of the Einstein's boxes, where a party could collapse the state of another party by open a box and finding nothing, see reference¹². Also, notice that in the previous consideration Alice is measuring the polarization of the photon; however, if she measured the presence of the photon¹¹ instead she would obtain similar states, but at the cost of destroying the photon.

On the other hand, if Alice decided to measure in different bases, for example in the bases $\frac{1}{\sqrt{2}}(|V\rangle \pm |H\rangle)$ or in the left and right circular polarization, in the former case the following possibilities arise:

1. If Alice got the polarization $|V\rangle + |H\rangle$, the spatial part of the state wavefunction would collapse towards $|a\rangle + |b\rangle$, i.e. a superposition of "being" in New York and Puebla at the same time. Remember that $|a\rangle + |b\rangle = |0\rangle |1\rangle + |1\rangle |0\rangle$; that is to say, what Bob gets is an entangled photon state, i.e. the local Bob's quantum state is $\hat{\rho}_B = Tr_A\{\hat{\rho}_{AB}\} = |0\rangle \langle 0| + |1\rangle \langle 1|$.
2. If Alice got the polarization $|V\rangle - |H\rangle$, the spatial part of the wavefunction would collapse towards $|a\rangle - |b\rangle$, i.e. a superposition of "being" in New York and Puebla at

the same time but with a different relative phase than the previous case. Please take into account that $|a\rangle - |b\rangle = |0\rangle|1\rangle - |1\rangle|0\rangle$; that is to say, what Bob gets is an entangled photon state, i.e. the local Bob's quantum state is $\hat{\rho}_B = Tr_A\{\hat{\rho}_{AB}\} = |0\rangle\langle 0| - |1\rangle\langle 1|$. similar results (but with different phases) would be achieved if Alice decide to measure in the left and right circular polarization bases.

Therefore, by choosing the measurement bases Alice is allowed to steer Bob's state at two space-like separate places like New York city and Puebla city. As it is well known, steering a quantum state means that a nonlocal resource is being used, since steering is a nonlocal property that lies between Bell nonlocality and entanglement^{4,33-37}. It is worthy of mention that the measurements could be made at two different places, i.e. New York city or Puebla city. Hence, if Bob also make measurements, then Bob and Alice would obtain similar or different results and process depending on whether or not they use generalized or projective measurements³⁸. Therefore, in the single-photon entangled state that is customized adapted in the above scheme, the non-local features arises from independent measurements at two faraway places.

Additionally, is worth of noting that if we replaced the HWP of Fig. (1) by a q plate^{39,40}, the q plate would produce a single-particle entangled state between the left and right circular polarization and the orbital angular momentum, $|\psi\rangle = \frac{1}{\sqrt{2}}(|L\rangle_\pi |-2\rangle_l + |R\rangle_\pi |2\rangle_l)$ ³⁹, of the kind mentioned in reference¹³, where $|R\rangle_\pi$ and $|L\rangle_\pi$ are the right and left circular polarization $|2\rangle_l$ and $|-2\rangle_l$ are the angular momentum sates. Hence, by writing this state in terms of the horizontal and vertical polarization we can ascertain the kind of state that is produced after the photon traverse the q plate. Then, the state before the second PBS is $|\psi\rangle = \frac{1}{2} \left[|H\rangle_\pi (|+2\rangle_l + |-2\rangle_l) + i |V\rangle_\pi (|+2\rangle_l - |-2\rangle_l) \right]$

Therefore, after traversing the second PBS the entangled state evolves towards: $|\psi\rangle = \frac{1}{2} \left[|a\rangle |H\rangle_\pi (|+2\rangle_l + |-2\rangle_l) + i |b\rangle |V\rangle_\pi (|+2\rangle_l - |-2\rangle_l) \right]$. Thus, this produces a similar situation that the one reported above which allowed Alice to steer Bob's state by means of suitable measurements. But in this case, it is an entangled state between three DoFs of the photon. Hence, Alice is able to measure in any bases of the three sub-spaces: i) the path, ii) the polarization, or iii) the orbital angular momentum (OAM); this scenario gives her more advantages than in the bipartite case. Notice that two DoFs of this state are the internal polarization and the OAM, hence by measuring in any of these two bases Alice may steer Bob's state. Of course, if Alice used a bipartite entangled state comprising the polarization

and the OAM only, she would not be able to steer Bob's state, as the authors of reference¹³ rightly imply. One of the drawbacks of reference¹³ is perhaps that the authors generalize this single situation for disproving the existence of single-photon nonlocal effects. However, as it is well known it is not always valid to generalize from a single setting, especially when one try to prove nonlocality with two internal degrees of freedom; on the other hand, by means of the scheme given above we could easily explain the phenomena of single-photon steering.

We can draw some conclusions from our discussion: as it is well known, quantum correlations are significantly different from the classical ones; hence, we agree that classical states cannot generate true quantum entanglement, thus the use of the name *classical entanglement* for alluding to classical non-separable states could lead to confusion. On the other hand, quantum single-particle entanglement could legitimately address many issues of quantum theory regarding the applications and understanding of fundamental questions on nonlocal quantum correlations and quantum mechanics. In this letter, we have proposed a different experimental set-up to show nonlocal effects in a single photon. In this new alternative set-up, it was shown that by manipulating the internal and external degrees of freedom of the single photon, the steering effect could be achieved. Particularly, the nonlocal features of the single-photon was addressed by using both i) a single internal and a single external degrees of freedom and ii) two internal degrees of freedom together with an external degree of freedom. These configurations differ from the ones reported in literature to show the nonlocality of the photon which use, as far as we know, the path entanglement of photons only. Hence, if one take into account the conclusions reached in references^{27,28} (that is, that the correlations measured by the TWC's experimental set-up are explained by a local hidden variable model), then the experimental set-up proposed here, could be a good alternative to demonstrate the nonlocality of a single photon. Additionally, it is worth mentioning that this scenario differs from the set-up shown in reference²⁶ where two external degree of freedom are entangled with an internal degree of freedom. These two kinds of entanglement represent two different experimental situations which produce different physical results; for example, the possibility of considering two additional observers in the case of reference²⁶ (i.e two external degrees of freedom entangled with an internal degree of freedom) is not possible in the photon case used in this letter where two internal degrees of freedom are entangled with an external degree of freedom. Finally, an additional difference is that in the case of

the photon, the vacuum state $|0\rangle$ plays a significant role (for example it could allow quantum tomography) that is not present in the situation that was addressed in reference¹². In conclusion, the nonlocal steering phenomenon could be facilitated by allowing entanglement between internal and external degrees of freedom in a single photon.

ACKNOWLEDGMENTS

We thank Alba Julita Chiyopa Robledo for the proffreading of the manuscript. There was not any support of any scientific agency. We thank Vicerrectoría de Investigación y Estudios de Postgrado from BUAP for providing us with computer equipment to carry out our academic and research duties.

REFERENCES

- ¹E. Schrödinger, Probability relations between separated systems. Mathematical Proceedings of the Cambridge Philosophical Society, **32**, 446-452 (1936).
- ²E. Schrödinger, Discussion of probability relations between separated systems. Mathematical Proceedings of the Cambridge Philosophical Society **31**, 555-563 (1936).
- ³A. Einstein, B. Podolsky, and N. Rosen, Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?, Phys. Rev. **47**, 777 (1935).
- ⁴H. M. Wiseman, S. J. Jones, and A. C. Doherty, Steering, entanglement, nonlocality, and the Einstein-Podolsky-Rosen Paradox, Phys. Rev. Lett. **98**, 140402 (2007).
- ⁵R. Ramanathan, D. Goyeneche, S. Muhammad, P. Mironowicz, M. Grünfeld, M. Bourennane, and P. Horodecki, Steering is an essential feature of non- locality in quantum theory, Nature Communications 9, 4244 (2018).
- ⁶Marco Túlio Quintino, Tamás Vértesi, and Nicolas Brunner, Joint Measurability, Einstein-Podolsky-Rosen Steering, and Bell Nonlocality, Phys. Rev. Lett. **113**, 160402 (2016).
- ⁷Roope Uola, Costantino Budroni, Otfried Gühne, and Juha-Pekka Pellonpää. One-to-One Mapping between Steering and Joint Measurability Problems Phys. Rev. Lett. **115**, 230402 (2015).
- ⁸S. M. Tan, D. F. Walls, and M. J. Collett, Nonlocality of a single photon, Phys. Rev. Lett. **66**, 252 (1991).

- ⁹L. Hardy, Nonlocality of a single photon revisited, *Phys. Rev. Lett.* **73**, 252 (1994).
- ¹⁰T. Guerreiro, F. Monteiro, A. Martin, J. B. Brask, T. Vértesi, B. Korzh, M. Caloz, F. Brussi  res, V. B. Verma, A. E. Lita, R. P. Mirin, S. W. Nam, F. Marsilli, M. D. Shaw, N. Gisin, N. Brunner, H. Zbinden, and R. T. Thew, Demonstration of Einstein-Podolsky-Rosen Steering Using Single-Photon Path Entanglement and Displacement-Based Detection, *Phys. Rev. Lett.* **117**, 070404 (2016).
- ¹¹Su-Yong Lee, Jiyong Park, Jaewan Kim, and Changsuk Noh, Single-photon quantum nonlocality: Violation of the Clauser-Horne-Shimony-Holt inequality using feasible measurement setups, *Phys. Rev. A* **95**, 012134 (2017).
- ¹²L. M. Ar  valo Aguilar, Nonlocal single particle steering generated through single particle entanglement, *Sci Rep* **11**, 6744 (2021).
- ¹³E. Karimi and R. W. Boyd, Classical Entanglement?, *Science* **350**, 1172 (2015).
- ¹⁴D. Paneru, E. Cohen, R. Fickler, R. W. Boyd and E. Karimi, Entanglement: quantum or Classical?, *Rep. Prog. Phys.* **83**, 064001 (2020).
- ¹⁵N Korolkova and G Leuchs, Quantum correlations in separable multi-mode states and in classically entangled light, *Rep. Prog. Phys.* **82**, 056001 (2019).
- ¹⁶A. Forbes, A. Aiello, B. Ndagano, Classically Entangled Light, *Progress in Optics* **64**, 99-153 (2019).
- ¹⁷T. Konrad and A. Forbes, Quantum mechanics and classical light, *Contemporary Physics*, **60**, 1-22, (2019).
- ¹⁸S. Azzini S. Mazzucchi V. Moretti, D. Pastorello, and L. Pavesi, Single-Particle Entanglement, *Adv. Quantum Technol.* , 2000014 (2020).
- ¹⁹R. Spreeuw, A classical analogy of entanglement *Found. Phys.* **28**, 361 (1998).
- ²⁰Monroe C, Meekhof D, King B and Wineland D, A ‘Schr  dinger cat’ superposition state of atom, *Science* **272**, 1131 (1996).
- ²¹Feiran Wang *et al*, Experimental Demonstration of a Quantum Controlled-SWAP Gate with Multiple Degrees of Freedom of a Single Photon, arXiv:2011.02581v3 [quant-ph], published in *Quantum Sci. Technol.* **6**, 035005 (2021). We have access only to the arXiv version.
- ²²M. Fuwa, S. Takeda, M. Zwi  rz, H. M. Wiseman, and A. Furusawa , Experimental proof of nonlocal wavefunction collapse for a single particle using homodyne measurements. *Nat Commun* **6**, 6665 (2015).

- ²³Francesco Garrisi, Micol Previde Massara, Alberto Zambianchi, Matteo Galli, Daniele Bajoni, Alberto Rimini, and Oreste Nicosini, Experimental test of the collapse time of a delocalized photon state, *Sci Rep* **9**, 11897 (2019).
- ²⁴Thiago Guerreiro, Bruno Sanguinetti, Hugo Zbinden, Nicolas Gisin, and Antoine Suarez, Single-photon space-like antibunching, *Phys. Lett. A* **376**, 2174–2177 (2012).
- ²⁵R. E. George, L. M. Robledo, O. J. E. Maroney, M. S. Blok, H. Bernien, M. L. Markham, D. J. Twitchen, J. J. L. Morton, G. Andrew D. Briggs, and Ronald Hanson, Opening up three quantum boxes causes classically undetectable wavefunction collapse, *PNAS*, **110** (10), 3777-3781 (2013).
- ²⁶E Benítez Rodríguez, E Piceno Martínez, and L M Arévalo Aguilar, Single-particle steering and nonlocality: The consecutive Stern-Gerlach Experiments, *Phys. Rev. A* **103**, 042217 (2021).
- ²⁷T. Das, M. Karczewski, A. Mandarino, M. Markiewicz, B. Woloncewicz and M. Żukowski, Can single photon of two spatially separated modes lead to a violation of Bell inequality via weak-field homodyne measurements?, *New Jour. Phys.* **23**, 073042 (2021).
- ²⁸T. Das, M. Karczewski, A. Mandarino, M. Markiewicz, B. Woloncewicz and M. Żukowski, No-go for device independent protocols with Tan-Walls-Collet of a single photon, arXiv:2102.03254v1 [quant-ph].
- ²⁹A. E. Piceno Martínez, E. Benítez Rodríguez, J. A. Mendoza Fierro, M. M. Méndez Otero, and L. M. Arévalo Aguilar, Quantum Nonlocality and Quantum Correlations in the Stern–Gerlach Experiment, *Entropy* **20**, 299 (2018).
- ³⁰Y-H Kim, Single-photon two-qubit entangled state: preparation and measurement, *Phys. Rev. A* **67**, 040301 (2003).
- ³¹E. Benítez Rodríguez, L.M. Arévalo Aguilar, and E.Piceno Martínez, A full quantum analysis of the Stern- Gerlach experiment using the evolution operator method: Analysing current issues in teaching quantum mechanics, *Eur. J. Phys.* **38**, 025403 (2017).
- ³²E. Benítez Rodríguez, L.M. Arévalo Aguilar, and E.Piceno Martínez, Corrigendum: ‘A full quantum analysis of the Stern–Gerlach experiment using the evolution operator method: Analysing current issues in teaching quantum mechanics’, *Eur. J. Phys.* **38**, 069501 (2017).
- ³³S. J. Jones, H. M. Wiseman, and A. C. Doherty, Entanglement, Einstein-Podolsky-Rosen correlations, Bell nonlocality, and steering, *Phys. Rev. A* **76**, 052116 (2007).

- ³⁴E. G. Cavalcanti, S. j. Jones, H. M. Wiseman, and M. D. Reid, Experimental criteria for steering and the Einstein-Podolsky-Rosen paradox, *Phys. Rev. A* **80**, 032112 (2009).
- ³⁵R. Uola, A. C. S. Costa, H. C. Nguyen, and O. Güne, Quantum steering, *Rev. Mod. Phys.* **92**, 015001 (2020).
- ³⁶N. Brunner, D. Cavalcanti, S. Pironio, V. Scarini, and S. Wehner, Bell nonlocality, *Rev. Mod. Phys.* **86**, 419 (2014).
- ³⁷H. Cao and Z. Guo, Characterizing Bell nonlocality and EPR steering, *Sci. China Phys. Mech. Astron.* **62**, 30311 (2019).
- ³⁸Aw, C., Dall’Arno, M. and Scarani, V. Knowledge by direct measurement versus inference from steering. *Quantum Stud. Math. Found.* **7**, 247–254 (2020).
- ³⁹E. Nagali, F. Sciarrino, F. De Martini, L. Marrucci, B. Piccirillo, E. Karimi, and Enrico Santamato, Quantum Information Transfer from Spin to Orbital Angular Momentum of Photons, *Phys. Rev. Lett.* **103**, 013601 (2009).
- ⁴⁰L. Marrucci, C. Manzo, and D. Paparo, Optical Spin-to-Orbital Angular Momentum conversion in inhomogeneous anisotropic media, *Phys. Rev. Lett.* **96**, 163905 (2006).