Evidence for an Algebra of G_2 Instantons

Michele Del Zotto[†], Jihwan Oh[‡], and Yehao Zhou^{*}

[†] Mathematics Institute, Uppsala University, Box 480, SE-75106 Uppsala, Sweden

[†] Department of Physics and Astronomy, Uppsala University, Box 516, SE-75120 Uppsala, Sweden

[‡] Mathematical Institute, University of Oxford, Woodstock Road, Oxford, OX2 6GG, United Kingdom

* Perimeter Institute for Theoretical Physics,31 Caroline St. N., Waterloo, ON N2L 2Y5, Canada

Abstract

In this short note, we present some evidence towards the existence of an algebra of BPS G_2 instantons. These are instantonic configurations that govern the partition functions of 7d SYM theories on local G_2 holonomy manifolds \mathcal{X} . To shed light on such structure, we begin investigating the relation with parent 4d $\mathcal{N} = 1$ theories obtained by geometric engineering M-theory on \mathcal{X} . The main point of this paper is to substantiate the following dream: the holomorphic sector of such theories on multi-centered Taub-NUT spaces gives rise to an algebra whose characters organise the G_2 instanton partition function. As a first step towards this program we argue by string duality that a multitude of geometries \mathcal{X} exist that are dual to well-known 4d SCFTs arising from D3 branes probes of CY cones: all these models are amenable to an analysis along the lines suggested by Dijkgraaf, Gukov, Neitzke and Vafa in the context of topological M-theory. Moreover, we discuss an interesting relation to Costello's twisted M-theory, which arises at local patches, and is a key ingredient in identifying the relevant algebras.

September 2021

Contents

1.	Introduction	2
2.	 M-theory and G₂ instantons 2.1. 7d gauge theory and geometry	6 6 7
3.	 \$\mathcal{N} = 1\$ SCFTs, \$G_2\$ manifolds, and dimers 3.1. Lightning review of dimers and mirrors 3.2. From IIA to M-theory 	11 12 13
4.	Towards the algebra of G_2 instantons4.1. Evidence from 4d/7d correspondence	 14 14 16 19 20 21 25
Α.	A brief review of the twisted M-theory background	25
B.	The D2 matrix model from string quantization B.1. Supersymmetry	28 28 31 37
C.	Formal description of the moduli spacesC.1. Moduli space as a subspace of a Nakajima quiver variety	40 44 44 45 49 54
D.	Connectedness of $\mathcal{M}^{N,K}(M_1, M_2)$	55

1. Introduction

The study of string compactifications on backgrounds with G_2 holonomy has received a recent renewal of interest in the physics literature [1–15] due to the discovery of infinitely many novel examples of such varieties. We can group the two kinds of novel G_2 back-

grounds in two broad classes: on the one hand we have the compact ones, which arise from twisted connected sums of asymptotically cylindrical Calabi Yau 3-folds [16–19], on the other hand we have non-compact local models that are obtained either from specific circle fibrations over asymptotic Calabi-Yau cones [20] or as asymptotically conical G_2 spaces in their own right [21].

Our paper builds on the remark that many of these non-compact cases might admit equivariant-like twists as M-theory backgrounds [22-24] — see also [25, 26]. In analogy with what happens for Calabi-Yau 3-folds [22, 27, 28] it is natural to expect these twisted backgrounds are related to suitable generalisations of topological string partition functions, the so called topological M-theories [29, 30], in terms of equivariant Donaldson-Thomas (DT) theories [22].¹ From this perspective in particular, it is natural to expect an interpretation of topological M-theory in terms of a generating function of Witten indices of supersymmetric quantum mechanical sigma models with target suitable supersymmetric versions of the G_2 instanton moduli spaces of a given geometry.

This short note is motivated by the question of understanding the structures governing the enumerative geometry of local G_2 manifolds [35], as well as their physical applications in the context of (equivariantly) twisted M-theory [22,24,36,37] and stringy correspondences [28,38].

Geometric engineering techniques in M-theory can be exploited to associate to a given local G_2 manifold \mathcal{X} a four-dimensional theory with $\mathcal{N} = 1$ supersymmetry which we denote $\mathcal{T}_{\mathcal{X}}$ in this paper. Whenever $\mathcal{T}_{\mathcal{X}}$ has a conserved $U(1)_R$ symmetry one can consider twisting this theory on any Kähler four-manifold, \mathcal{M} , by identifying the $U(1)_R$ symmetry with a U(1) subgroup of its U(2) holonomy [39–42] – see also [43,44]. The resulting theory is not topological in this case, rather it has a dependence on the holomorphic structure of \mathcal{M} . Often it is possible to compute the corresponding partition function $Z_{\mathcal{T}}(\mathcal{M})$, for example by means of localization methods. Then, we have a natural correspondence

$$Z_{\mathcal{T}_{\mathcal{X}}}(\mathcal{M}) = \mathcal{Z}_M(\mathcal{M} \times \mathcal{X}) \tag{1.1}$$

between an M-theory partition function on a background with topology $\mathcal{M} \times \mathcal{X}$ and the twisted 4d $\mathcal{N} = 1$ partition function on the background \mathcal{M} of the field theory $\mathcal{T}_{\mathcal{X}}$. This poses the question of constructing G_2 manifolds corresponding to theories $\mathcal{T}_{\mathcal{X}}$ with an unbroken $U(1)_R$ symmetry. Thanks to string duality, we argue in this paper that an infinite class of such geometries exist, giving rise to local G_2 manifolds that geometrically engineer 4d $\mathcal{N} = 1$ SCFTs.²

In this paper we are interested in the case when either \mathcal{M} is a multicentered Taub-NUT

¹ See also [31–34] for other attempts at a topological string or topological M-theory on G_2 manifolds.

² We warmly thank Iñaki García Etxebarria for suggesting to exploit this duality to us, as well as explaining several of his results that are crucial to argue for the stability of these backgrounds in M-theory.

space with k centers, which we denote \mathbf{TN}_k , or \mathcal{M} is an ALE space of type A_k .³ Recall that the theory obtained from M-theory on \mathbf{TN}_k is a 7d U(k) maximally supersymmetric gauge theory. Similarly, the theory obtained from M-theory on the \mathbf{ALE}_k space is a 7d SU(k) maximally supersymmetric gauge theory.⁴ Therefore, the backgrounds of the form $\mathbf{TN}_k \times \mathcal{X}$ and backgrounds $\mathbf{ALE}_k \times \mathcal{X}$ are such that they admit a distinct interpretation in geometric engineering, as 7d partition functions for these gauge theories, namely

$$Z_{U(k)}^{7d}(\mathcal{X}) = \mathcal{Z}_M(\mathbf{TN}_k \times \mathcal{X}) \quad \text{and} \quad Z_{SU(k)}^{7d}(\mathcal{X}) = \mathcal{Z}_M(\mathbf{ALE}_k \times \mathcal{X}) \quad (1.2)$$

Whenever the G_2 manifolds have compact associative 3-cycles, we argue that the 7d partition functions on the LHS of these identities receive contributions from G_2 instantons for the groups U(k) and SU(k) respectively, which therefore gives an interpretation in terms of enumerative geometry of local G_2 manifolds of the partition functions we are interested in. Moreover, if the manifolds \mathcal{X} admit compact co-associative four-cycles, we expect the parition functions can receive contributions also from 7d monopoles.

From the relation with the M-theory partition function, combining (1.1) with (1.2) we expect a 4d/7d correspondence of the schematic form [38]

which extends the results of [28] to this example, thus giving rise to a dictionary between holomorphically twisted partition functions of 4d $\mathcal{N} = 1$ SCFTs on multi-centered Taub-NUT spaces and the U(k) G_2 -instanton partition function. Whenever \mathcal{X} admits a U(1)isometry, the k = 1 case of this correspondence (in particular, the bottom line in equation (1.3)) coincides with the definition of topological M-theory as given in [30], in terms of a generating function of D2-D6 boundstates arising from exploiting the U(1) fibration of \mathcal{X} as an M-theory circle to reduce to IIA. In this paper we will discuss how this perspective can be exploited to begin unraveling the properties of these partition functions in a simple example. In future papers of this series we will address more complicated geometries and constructions. The main message of this paper is that building on the 4d/7d correspondence, the algebra of operators which govern the holomorphic sector of 4d $\mathcal{N} = 1$ theory is such that the resulting G_2 instanton partition function can be interpreted as a character.

This paper is organised as follows. In section 2 we argue that the partition function

³ We refer our readers to the papers [45] and [28] for a through review of the properties of these spaces and their relations. Of course our results can be generalised to include other ALE spaces, as well as discrete C_3 form fluxes in M-theory to give rise to G_2 instantons in all other simple gauge groups.

⁴ In this paper we are neglecting all the subtleties related to the choices of global structures for these models [46–48]. We plan to revisit our analysis in the near future to take the latter into account.

of the 7d U(K) gauge theory, in presence of compact associative cycles, localises on G_2 instantons. This gives rise to a definition of the partition function $Z_{U(K)}^{7d}(\mathcal{X})$ as a generating function for some sort of equivariant volumes of G_2 -instanton moduli spaces. In section 3 we discuss a string duality that allows to construct infinitely many novel examples of G_2 manifolds with an interpretation as 4d $\mathcal{N} = 1$ SCFTs that will play a role in the construction of the correspondence and give rise to the details of the corresponding dictionary as suggested by geometric engineering. The resulting local G_2 geometries can be understood in terms of fibrations of Taub-NUT spaces over a collection of intersecting associatives. For this class of geometries a strategy to reconstruct the full G_2 instanton partition function is presented, in terms of the 4d side of the correspondence in section 4. Since the G_2 instantons are engineered by M2 branes wrapping these loci, this gives a way of determining their contributions in terms of a matrix model which can also be uplifted to an index for an associated supersymmetric quantum mechanics which we derive by a chain of string dualities.

Exploiting such quantum mechanics it is the key towards understanding the algebra of G_2 instantons for the example we consider, in particular, zooming to the intersection point of a pair of associatives, one has a local model that can be traced back to a wellknown orbifold of the Bryant-Salamon metric for the G_2 cone over the three-dimensional complex projective plane \mathbb{CP}^3 [49], that were discussed in [50, 51]. This is a universal building block for the G_2 instanton partition functions that can be understood in terms of the twisted M-theory à la Costello [23] (see also [25, 26]), along the lines of [52].

Indeed, our search for the algebra of G_2 instantons was initiated by the fascinating observation by Costello⁵: if we instead consider twisted M-theory on $\mathbb{R} \times \mathbb{C} \times TN_K$ not the G_2 cone, one can argue that the Donaldson-Thomas partition function on $\mathbb{C} \times TN_K$ forms a character of an algebra of operators of twisted M-theory. In this paper, we will consider twisted M-theory on the G_2 cone over \mathbb{CP}^3 and compute the G_2 instanton partition, which is presumably a part of the G_2 Donaldson-Thomas partition function. Along the way, we will be able to see that the analogous statement to that of Costello can be made.

As a final comment, let us remark that the main aim of this paper is to pose this problem: here we show that the study of the G_2 instanton parition functions have interesting interconnections with various aspects of geometric engineering and M-theory, which already unveils interesting algebraic structures: this is our evidence for the existence of an algebra of G_2 instantons.

For the benefit of our readers, the vast majority of the technical aspects of the discussion are confined in the Appendix of the manuscript.

⁵We learned this from Kevin Costello's lecture "Cohomological hall algebras from string and M theory" in Perimeter Institute, which can be found in https://pirsa.org/19020061.

2. M-theory and G_2 instantons

In this section we discuss our dictionary between the 7d G_2 instanton partition function and the underlying M-theory geometry. We stress that in this paper, as in [28], the explicit expression of the equivariantly twisted M-theory background is not yet known explicitly for the cases of interest. The non-twisted version of these backgrounds are well known to exist in M-theory, and have been discussed recently in [53, 54]: these are 11 dimensional spaces with topology $\mathcal{M} \times \mathcal{X}$ where \mathcal{M} is a HyperKähler space and \mathcal{X} is a G_2 holonomy space. The backgrounds are supersymmetric and preserve 2 supercharges. Strong evidence towards the existence of similar equivariantly twisted structures comes from the supergravity limit [22, 24–26], whenever the spaces \mathcal{M} and \mathcal{X} have sufficiently large isometry groups.

2.1. 7d gauge theory and geometry

It is well known that M-theory on $\mathbf{TN}_k \times \mathbf{R}^7$ gives rise to a U(k) gauge theory along \mathbf{R}^7 . The space of normalisable forms on \mathbf{TN}_k can be parametrised by k elements,

$$\omega_0, \omega_\alpha \qquad \alpha = 1, \dots, k-1 \tag{2.1}$$

where ω_0 is such that $\int \omega_0 \wedge \omega_\alpha = 0$ for all α and $\int \omega_\alpha \wedge \omega_\beta = C_{\alpha,\beta}$, the Cartan matrix of A_{k-1} type. Consequently, the reduction of the C_3 field of M-theory on such a basis gives rise to 7d vectors:

$$C_3 = A^{(0)} \wedge \omega_0 + \sum_{\alpha} A^{(\alpha)} \wedge \omega_{\alpha}$$
(2.2)

where $A^{(0)}$ is a U(1) gauge potential, while the remaining $A^{(\alpha)}$ are associated to the Cartan of SU(k). In the limit in which the Taub-NUT radius of \mathbf{TN}_k is sent to infinite size, the form ω_0 stops being normalisable and one is left with an \mathbf{ALE}_k metric, which captures only the dynamics of a 7d SU(k) gauge theory. In the moduli space of these geometries we have k-1 compact \mathbb{P}^1_{α} whose volumes correspond to the vevs in the Cartan of SU(k) for the scalars in the corresponding 7d vectormultiplets

$$\langle \phi_{\alpha} \rangle = \operatorname{vol} \mathbb{P}^{1}_{\alpha} \qquad \int_{\mathbb{P}^{1}_{\alpha}} \omega_{\beta} = \delta_{\alpha\beta} \,.$$
 (2.3)

M2 branes wrapping such 2-cycles give rise to the corresponding W-bosons. Sending these volumes to zero we have an enhancement of symmetry and the gauge group becomes non-Abelian.

Besides these degrees of freedom, the resulting 7d gauge theory has the following non-perturbative solitonic degrees of freedom:

• Instantons with worldvolumes of codimension four that are parametrized by M2 branes strechced along **R**⁷;

- Monopoles with worldvolumes of codimension three that are parametrized by M5 branes wrapped around P¹_α;
- Domain walls with worldvolumes of codimension six that are parametrized by M5 branes strechced along **R**⁷.

In this paper, we are interested in understanding some features of the partition function of this theory coupled to a rigid supersymmetric background. In general, one expects to receive contributions to these partition functions from BPS solitons that depend on the properties of the localising BPS background.

2.2. Localising on G₂ instantons

We are interested in the parition function for 7d SYM on a non-compact space with G_2 holonomy \mathcal{X} , suitably regularized to avoid the potential IR divergence with a choice of boundary condition at infinity. Several results about this theory on curved spaces have been obtained recently, see e.g. [55–57], working à la Festuccia-Seiberg [58], however a detailed study of the case of pure G_2 holonomy is complicated by the fact that in this latter case one has only a single parallel spinor, and therefore the localising locus is not reduced to a combination of fixed points arising from a Reeb-like structure. Nevertheless, twisted versions of this theory are well-known to exist and lead to an interesting cohomological field theory depending on the G_2 structure of \mathcal{X} — see [59–61]. In particular, it is expected that these theories receive contributions from analogues of G_2 instanton configurations [62–65].

The main aim of this section is indeed to argue heuristically that the 7d instanton partition function is governed by a suitable supersymmetrization of the topological term

$$\varkappa(F) = \int_{\mathcal{X}} \left(\varphi_3 \wedge ch_2(F) \right) \tag{2.4}$$

where φ_3 is the parallel G_2 form, which is closed and co-closed, i.e. it satisfies

$$d\varphi_3 = 0 \qquad d * \varphi_3 = 0. \tag{2.5}$$

This ensures that $\varkappa(F)$ is a characteristic class. The action for 7d SYM is obtained by dimensional reduction from the 10d SYM Lagrangian (see e.g. [66])

$$\mathcal{L} = -\frac{1}{4} F_{MN} F^{MN} - \frac{i}{2} \overline{\Psi} \Gamma^M D_M \Psi$$
(2.6)

where Ψ is a Majorana-Weyl 10d spinor in the Adjoint of the gauge group. The supersymmetry transformations are

$$\delta A_M = \frac{i}{2} \overline{\epsilon} \Gamma_M \Psi \qquad \qquad \delta \Psi = -\frac{1}{4} F_{MN} \Gamma^{MN} \epsilon \qquad (2.7)$$

The resulting theory has an Adjoint scalar triplet of $SU(2)_R$ symmetry, L_a , that arises from the components of A_M in the directions 789. The corresponding supersymmetric action can be found in [67]. Here we consider the partition function of the 7d U(k) gauge theory on \mathcal{X} , which can be written schematically as

$$Z_{U(K)}^{7d}(\mathcal{X}) = \int \mathcal{D}\Phi e^{-\int_{\mathcal{X}} S[\Phi]},$$
(2.8)

where we collectively denote the fields in the 7d $\mathcal{N} = 1$ SYM as Φ , $S[\Phi]$ is the action of the 7d SYM on the G_2 manifold \mathcal{X} , and of course

$$S[\Phi] \supseteq S_{YM}(F) = \frac{1}{g^2} \int_{\mathcal{X}} \operatorname{tr}_{\mathfrak{g}} F_A \wedge *F_A$$
(2.9)

where F_A is the 7d U(k) curvature with potential A. The trace is taken in the the normalisation for which a single instanton configuration of SU(2) has unit instanton number, and g is the gauge coupling constant of the 7d theory. We proceed by arguing that the corresponding action is dominated by G_2 instantons in the saddle point approximation, following [68]. We present our argument on the Yang-Mills side (2.9).

First of all it is necessary to remind our readers that two forms on a G_2 manifold are naturally organised according the representation theory of G_2 as follows [69] (see also the nice review [70])

$$\wedge^2 T^* \mathcal{X} = \wedge^2_{\mathbf{7}} \oplus \wedge^2_{\mathbf{14}}, \qquad \qquad H^2(\mathcal{X}) = H^2_{\mathbf{7}}(\mathcal{X}) \oplus H^2_{\mathbf{14}}(\mathcal{X}), \qquad (2.10)$$

where the splitting is orthogonal with respect to the canonical pairing

$$\langle \eta, \xi \rangle_{\mathcal{X}} = \int_{\mathcal{X}} \eta \wedge *\xi$$
 (2.11)

induced by the Hodge star of the G_2 manifold. We call π_7 and π_{14} the corresponding projectors.⁶ Then, of course

$$S_{YM}(F) = S_{YM}(F_7) + S_{YM}(F_{14})$$
(2.12)

There is a natural G_2 -equivariant linear map, which is compatible with such a splitting, meaning that [71]

$$T: \wedge^2 T^* \mathcal{X} \to \wedge^2 T^* \mathcal{X} \qquad T(\eta) \equiv *(\eta \wedge \varphi_3) = 2\pi_7(\eta) - \pi_{14}(\eta).$$
(2.13)

⁶ Sometimes we use the short hand $\eta_7 = \pi_7 \eta$ and $\eta_{14} = \pi_{14} \eta$.

It is easy to see that

$$\int_{\mathcal{X}} \eta \wedge \eta \wedge \varphi_3 = 2 \langle \pi_7(\eta), \pi_7(\eta) \rangle_{\mathcal{X}} - \langle \pi_{\mathbf{14}}(\eta), \pi_{\mathbf{14}}(\eta) \rangle_{\mathcal{X}}$$
(2.14)

follows from the definition (2.11) and the property (2.13) by orthogonality.

Therefore, the topological term in (2.4) is such that

$$\alpha \varkappa(F) = \operatorname{tr}_{\mathfrak{g}} \int_{\mathcal{X}} F_A \wedge F_A \wedge \varphi_3 = 2 \operatorname{tr}_{\mathfrak{g}} \langle \pi_{\mathbf{7}}(F_A), \pi_{\mathbf{7}}(F_A) \rangle_{\mathcal{X}} - \operatorname{tr}_{\mathfrak{g}} \langle \pi_{\mathbf{14}}(F_A), \pi_{\mathbf{14}}(F_A) \rangle_{\mathcal{X}}$$
$$= g^2 \Big(2S_{YM}(F_{\mathbf{7}}) - S_{YM}(F_{\mathbf{14}}) \Big)$$
(2.15)

Here α is a constant to keep track of the correct normalisation for the second Chern class. This gives rise to the following identities:

$$S_{YM}(F) = \frac{1}{2} \Big(3S_{YM}(F_{14}) + \frac{\alpha}{g^2} \varkappa(F) \Big) = 3S_{YM}(F_7) - \frac{\alpha}{g^2} \varkappa(F)$$
(2.16)

Since $S_{YM}(F)$ has to be positive, then only two possibilities are allowed, either $F_7 = 0$ and $\varkappa(F) < 0$, or $F_{14} = 0$ and $\varkappa(F) > 0$, which are the G_2 analogues of the self-dual and anti-self dual equations. For compact G_2 manifolds, F_7 is always zero and for this reason here we will also focus on those instanton configurations. The topological term in equation (2.4) is then saturated by G_2 instanton configurations that are supported on an associative 3-cycle σ_3 and have a non-trivial Chern class in the corresponding Poincaré dual four-form. By definition, these G_2 instantons are solutions to the equation

$$*F_A = -F_A \wedge \varphi_3. \tag{2.17}$$

It is interesting to remark that G_2 instanton equations are rigid for compact G_2 spaces, meaning that the corresponding moduli spaces have virtual dimension zero, and the corresponding moduli space is a collection of points [72, 73], however for non-compact manifolds the moduli problem becomes, of course richer, and more interesting moduli spaces are allowed [74]. Of course, for 7d SYM field configurations the relevant moduli spaces will be enhanced by the contribution from the other fields in the 7d SYM Lagrangian.

Remark. In general, \mathcal{X} can have calibrated associative 3-cycles as well as calibrated co-associative 4-cycles. For the examples that we are considering in this paper, however, we always have a phase of the geometry where there are no compact co-associative 4-cycles, so we will assume it to be the case for the time being. We will comment on the phases with compact coassociative 4-cycles below.

Let us denote by $\sigma_{3,i}$ a basis of $H_3(\mathcal{X}, \mathbb{Z})$ and by abuse of notation also their calibrated representatives. The complexified volumes will give rise to complex scalar moduli for M- theory, which we denote

$$q_i = \int_{\sigma_{3,i}} \varphi_3 \qquad \qquad \varphi_3 = \varphi_3 + \frac{i}{2\pi} C_3. \tag{2.18}$$

Here, of course, C_3 is the M-theory 3-form. A G_2 instanton configuration for U(k) with instanton number κ_i along $\sigma_{3,i}$ is engineered in this setting by a stack of Euclidean M2 branes wrapping $\sigma_{3,i}$ for κ_i times. The contribution from such a G_2 instanton to the parition function is proportional to

$$Z^k_{\kappa_i}(\sigma_{3,i})Q^{\kappa_i}_i,\tag{2.19}$$

where $Q_i^{\kappa_i} = e^{-q_i \kappa_i}$ and $Z_{\kappa_i}^k(\sigma_{3,i})$ is the partition function of κ_i M2 branes wrapping the associative 3-cycle $\sigma_{3,i} \subset \mathcal{X}$ in the background of $\mathbf{TN}_k \times \mathcal{X}$. Whenever we have multiwrapping, this gives rise to further contributions proportional to the same power of Q_i . In general, the coefficient multiplying Q_i^N is of the from

$$\sum_{\substack{k_i, m_i \text{ such that} \\ N = \kappa_i m_i}} (Z_{\kappa_i}^k(\sigma_{3,i}))^{m_i} .$$
(2.20)

Whenever the instanton is supported on $\sigma_3 = \sum_{i=1}^{b_3(\mathcal{X})} m_i \sigma_{3,i}$ with instanton numbers κ_i on each $\sigma_{3,i}$, its contribution to the partition function receives a weight

$$Q_{\sigma_3}^{\kappa} = \prod_{i=1}^{b_3(\mathcal{X})} Q_i^{m_i \kappa_i}.$$
(2.21)

Therefore, schematically, for examples without co-associatives we can write

$$\mathcal{Z}_M = \sum_{\sigma \in H_3(\mathcal{X})} \sum_{\kappa} Z_{\kappa}^k(\sigma) Q_{\sigma}^{\kappa}.$$
(2.22)

Remark. The M-theory partition function could in principle receive contributions also from M5 branes wrapping $\mathbb{P}^1_{\alpha} \times S_{4,a}$ where $S_{4,a}$ is a compact co-associative 4-cycle. These contributions will be relevant for slight generalisations of our backgrounds, which are associated to a different twist of the 4d $\mathcal{N} = 1$ theory, in particular, the vevs of such co-associatives will break the $U(1)_R$ symmetry and so the corresponding backgrounds would no longer be twisted holomorphically. A construction of such partition functions on the 4d side of the 4d/7d correspondence can be achieved by exploiting rigid curved supersymmetry and non-minimal $\mathcal{N} = 1$ supergravity coupled to the corresponding Ferrara-Zumino \mathcal{F} multiplet. Since the latter reduces to the \mathcal{R} multiplet whenever we are switching off the offending vevs, we believe that such partition functions are the most natural extension to the partition functions we consider in this note. Based on the above analysis then, naively one expects that

$$\widehat{Z}_{U(k)}^{7D}(\mathcal{X}) = \frac{Z_{U(k)}^{7D}(\mathcal{X})}{Z_{U(k)}^{(0)}(\mathcal{X})} = \sum_{\sigma_3 \in H_3(X)} \sum_{\kappa \ge 0} Q_{\sigma_3}^{\kappa} \int_{\mathcal{M}_{\sigma_3,\kappa}^{U(k)}} \mathbf{1},$$
(2.23)

where:

- $Z_{U(k)}^{(0)}(\mathcal{X})$ is a perturbative contribution;
- $\mathcal{M}_{\sigma_3,\kappa}^{U(k)}$ is the moduli space of a BPS version of the U(k) G_2 instanton, with instanton numbers κ_i over the collection of normal bundles in the associative 3-cycles in the support of the vector m_i associated to $\sigma_3 = \sum_i m_i \sigma_{3,i}$. Each G_2 instanton configuration contributes to the partition function with a suitably regularized volume of such moduli space.⁷

Comparison between (2.23) and (2.19) leads to identify

$$Z_{\kappa_i}^k(\sigma_{3,i}) = \int_{\mathcal{M}_{\kappa_i,\sigma_{3,i}}^{U(k)}} \mathbf{1} \,. \tag{2.24}$$

Building on the idea of topological M-theory [30], for the class of examples that we will consider in this paper we claim that the matrix model computing $Z_{\kappa_i}^k(\sigma_{3,i})$ can be obtained by considering bound states of D2 and D6 branes in the type IIA frame. For the special example we consider in section 4 below, we also comment on an interesting duality to IIB where the same configuration is related to a 3d $\mathcal{N} = 2$ theory with impurities (a close cousin of the theories studied in [77,78]), which reminds also of the discussions in [53,54].⁸

3. $\mathcal{N} = 1$ SCFTs, G_2 manifolds, and dimers

In this section we give a simple argument for the existence of infinitely many examples of $\mathcal{T}_{\mathcal{X}}$ that have a $U(1)_R$ isometry.

⁷ A possible regularization arise in the equivariant setting, where we expect that parameters for each instanton contribution are associated to a local Ω deformation [75,76] of the SO(4) rotation of $\mathcal{N}_{\sigma_{3,i}}$. The explicit form of the glueing rules among such parameters into global ones can be read off by the existence of the global G_2 structure.

⁸ This last connection is very interesting: the theory with impurities considered in these references is associated to an M5 wrapping a co-associative, precisely of the type that would contribute to our parition function in presence of compact 4-cycles. We believe this perspective could be the key to model certain associative transitions, which are the G_2 manifold uplift of the conifold transitions in IIA [50, 79].



Figure 1: Some well-known examples of quivers from dimers.

3.1. Lightning review of dimers and mirrors

It is well known that probing singularities at the tip of Calabi-Yau cones **X** with a stack of N D3 branes in type IIB one obtains 4d $\mathcal{N} = 1$ SCFTs, which we label $\mathcal{T}_{\mathbf{X},N}$.⁹ In particular, when **X** is a local toric CY threefold, the corresponding theories on the stack of N D3 branes have a 4d $\mathcal{N} = 1$ Lagrangian description that is straightforward to compute, being encoded in a dimer model. In general, many of these theories can be captured by a quiver diagram (also in the non-toric cases) that can be read off directly from the structure of the singular orbifold geometry exploiting standard techniques that we are not going to review here. Some examples of such quivers can be found in Figure 1.

A well-known feature of these geometries is that the resulting theories can be represented in Type IIA by exploiting mirror symmetry. By mirror symmetry, the IIB theory on the toric threefold \mathbf{X} is equivalent to IIA on the mirror 3-CY \mathbf{X}^{\vee} . The D3 brane stack in IIB is mapped to a configuration of wrapped *D*6 branes that have support on special Lagrangian 3-cycles in the mirror geometry \mathbf{X}^{\vee} , which can be understood, e.g. in terms of the SYZ fibration [86]. We denote a basis of such cycles $C_i \in H_3(\mathbf{X}^{\vee}, \mathbb{Z})$. For the mirrors of the toric cases we are interested in, the topology of these 3-cycles is either an S^3 or an $S^1 \times S^2$.

On the IIB side, the D3 brane fractions at the singularity **X** in a number b_3 of fractional D3 branes [87]. These are in one-to-one correspondence with the cycles C_i in the mirror dual. The original D3 brane is a bound state of fractional 3-branes, consisting of N_i copies of the *i*-th 3-brane. This is mapped in the mirror to N_i D6 branes wrapping the 3-cycle C_i . These D6 branes give rise to gauge groups $SU(N_i)$ on the 4d theories:¹⁰ if C_i is an S^3 we obtain an $\mathcal{N} = 1$ vectormultiplet, if instead $C_i \simeq S^1 \times S^2$ the resulting degrees of freedom are those of an $\mathcal{N} = 2$ vectormultiplet. The intersection pairing

$$B_{ij} = \mathcal{C}_i \cdot \mathcal{C}_j \tag{3.1}$$

⁹ There is a multitude of references about this subject and therefore it is hard to properly cite them, some references there were key for the preparation of this section are [80–85].

¹⁰ The corresponding U(1) are expected to decouple in the IR.

in the mirror geometry corresponds to an adjacency matrix of the quiver theory describing the D3 brane worldvolume. In particular, if a pair of wrapped cycles C_i, C_j are intersecting, then the corresponding wrapped D6 branes are intersecting, in which case we obtain extra massless matter from open strings ending on them. Generically, we will have configurations in which the 3-cycles are S^3 's that intersect over a collection of $|B_{ij}|$ points. In correspondence to each point with $B_{ij} > 0$ we obtain an $\mathcal{N} = 1$ chiral multiplet in a bifundamental $(\overline{\mathbf{N}}_i, \mathbf{N}_j)$. In cases when the 3-cycles overlap along an S^1 the resulting contribution is an $\mathcal{N} = 2$ hypermultiplet, which however gives $B_{ij} = 0$. For most of the toric models, the topology of the C_i is always that of S^3 and one has an $\mathcal{N} = 1$ theory. In particular, in this case, the $N_i > 0$ must satisfy

$$B_{ij}N_j = 0, (3.2)$$

which guarantees anomaly cancellation on the 4d $\mathcal{N} = 1$ side. Whenever the quiver has loops, the resulting 4d $\mathcal{N} = 1$ theory can also have an interesting tree-level superpotential, which, in favourable circumstances, is also determined using geometry. For the toric cases $N_i = N$ for all *i*.

3.2. From IIA to M-theory

The G_2 geometries that we will consider are captured by the IIA geometry of \mathbf{X}^{\vee} by duality.¹¹ One has to uplift the configuration of intersecting D6 branes over the collection of special Lagrangian 3-cycles C_i to a hyperKähler fibration on a collection of associative cycles $\sigma_{3,i} \simeq C_i$, which gives rise to a G_2 manifold that is a U(1) fibration over \mathbf{X}^{\vee} . The fibration is such that the U(1) bundle degenerates along the cycles $\sigma_{3,i}$ that correspond to the special lagrangian 3-spheres C_i that are wrapped by N_i D6 branes, giving rise to a local $\mathbb{C}^2/\mathbb{Z}_{N_i}$ singularities with locus $\sigma_{3,i}$. The singularities enhance to $\mathbb{C}^2/\mathbb{Z}_{N_i+N_j+1}$ at the intersection where N_i D6s meet with N_j D6s, similar to what happens in other geometric engineering settings [88]. In this context, it is natural to conjecture that the condition for the absence of anomalies (3.2) is equivalent to the condition that the corresponding φ_3 form is not only closed but also co-closed. For toric examples, we have that $N_i = N$, where N is the number of D3-branes probing \mathbf{X} . Therefore, we label this class of G_2 manifolds $\mathcal{X}_{\mathbf{X}^{\vee},N}$. From the perspective of geometric engineering, we have that

$$\mathcal{T}_{\mathbf{X},N} = \mathcal{T}_{\mathcal{X}_{\mathbf{X}^{\vee},N}}.$$
(3.3)

In the toric examples, the mirror geometries are known explicitly. One starts from the

¹¹ It would be interesting to investigate generalising slightly the construction of Foscolo, Haskins, and Nördstrom [20] to include orbifold singularities. We believe many of the geometries in this section are orbifolds of the FHN metrics.

toric diagram of \mathbf{X} and obtains the mirror geometry

$$\mathbf{X}^{\vee} = \{uv = W(X, Y)\} \subset \mathbf{C}^2 \times (\mathbf{C}^*)^2, \tag{3.4}$$

where W(X, Y) is the Newton polynomial obtained from the toric diagram by assigning to the internal dots with coordinates (n, m) the monomial $X^n Y^m$, and X and Y are \mathbb{C}^* variables [89]. We refer our readers to the beautiful detailed discussion in [85] for more aspects of these geometries.

An application of the fact that these models have an explicit description in terms of a toric quiver, is that it is possible to use these gauge theories to argue for the lack of contributions to the superpotential from euclidean M2 branes wrapping the associative 3-cycles $\sigma_{3,i}$. This follows because for all toric dimer models the quiver associated to the corresponding geometry is necessarily such that $N_f = \ell N_c$ where $\ell \geq 2$. Since each gauge node can be treated independently from the perspective of the generation of a nonperturbative superpotential in gauge theory, this is precisely the condition that there are as many incoming as there are outgoing arrows from each node, which is a condition that every toric quivers satisfy. Then there is no non-perturbative superpotential generated [90]: in all these cases the instanton generates no superpotential, but rather higher Fterms [91].¹² The higher F-terms involve lots of fermion insertions and therefore most likely will be irrelevant deformations from the perspective of the SCFT. This guarantees that these geometries are not receiving non-perturbative superpotential contributions from the associative 3-cycles, and therefore evade the main issue in geometric engineering of 4d $\mathcal{N} = 1$ SCFTs, namely that non-perturbative contributions to the superpotential could spoil geometric intuition giving rise to a quantum moduli space that differs from the classical geometrical one — see e.g. [92] for a recent discussion of this effect in the context of F-theory geometric engineering of 4d $\mathcal{N} = 1$ SCFTs.

This concludes our brief overview of these geometries. We stress that there are many applications of this geometric setting, which uplift to G_2 , for instance, the geometrization of dualities by Ooguri and Vafa [80]. We will return to the study of more details of these geometries in future papers in this series.

4. Towards the algebra of G_2 instantons

4.1. Evidence from 4d/7d correspondence

In the previous section we have given an argument to provide infinitely many examples of local G_2 geometries that give rise to systems with an unbroken $U(1)_R$ symmetry, and therefore can be exploited to construct 4d holomorphically twisted \mathbf{TN}_k partition

¹² We thank Iñaki García Etxebarria for an important clarifying discussion about this point.

functions.¹³ We can use this fact to our advantage while proceeding in our investigation of the structure governing the BPS G_2 instantons on the stringy side for the corresponding G_2 manifolds $\mathcal{X}_{\mathbf{X}^{\vee},N}$.

The main feature of all the local models of G_2 geometries we introduced in the previous section is that those are dual to collections of N D6 branes wrapping compact special Lagrangians in the mirrors of toric geometries $\mathcal{X}_{\mathbf{X}^{\vee}}$. We are interested in a configurations of Euclidean D2 branes, wrapping the same special Lagrangians. From geometry it is clear that the most interesting contributions will arise from intersecting loci (other loci would be closer to ordinary instantons). Our task therefore is the following: (1) exploiting branes, identify the contribution of intersecting loci to our putative BPS G_2 instanton partition function; (2) establish gluing rules corresponding to the quiver diagram associated to the corresponding geometry. This is in a sense the analogue of the topological vertex [93, 94] for this special class of G_2 manifolds.

This task is seemingly impossible, however some recent progresses in the context of twisted M-theory can be exploited to our advantage. The main point is that the twisted M-theory background by Costello [24] provides a local model of the form¹⁴

$$\mathbf{TN}_k \times \mathbf{R}^3 \times \mathbf{TN}_{N_i} \tag{4.1}$$

where the twist is holomorphic on \mathbf{TN}_k , and topological on $\mathbf{R}^3 \times \mathbf{TN}_{N_i}$. Such a geometry can be identified with a local patch in the structure of any of the geometries discussed in section 3: it is enough to identify the \mathbf{R}^3 in equation (4.1) with a local neighborhood of one of the associatives 3-cycles $\sigma_{3,i}$. The fact that the twist is holomorphic on \mathbf{TN}_k guarantees that this twisted M-theory construction contributes a subalgebra of the cohomological field theory associated to the holomorphic twist of one of the 4d $\mathcal{N} = 1$ models $\mathcal{T}_{\mathcal{X}_{\mathbf{X}^{\vee},N}}$. More precisely, the contribution from a subsector like (4.1) is encoded in an algebra that we schematically denote $\mathcal{A}_{N_i}^k$. It is natural to conjecture that

$$\mathcal{A}_{N_i}^k \simeq U_{\epsilon_1}(\mathcal{O}_{\epsilon_2}(\mathbf{TN}_k) \otimes \mathfrak{gl}_{N_i}) \tag{4.2}$$

which is the quantum deformation with parameter ϵ_1 of the universal enveloping algebra of the algebra of deformed holomorphic functions on \mathbf{TN}_k tensored by \mathfrak{gl}_{N_i} .

For the class of examples in this note, whenever we have a pair of intersecting D6 branes in the background of \mathbf{TN}_k , we expect to obtain a more general algebra that has two such factors

$$\mathcal{A}_{N_i}^k \otimes_{B_{ij}} \mathcal{A}_{N_j}^k, \tag{4.3}$$

where the notation $\otimes_{B_{ij}}$ indicates that there are some further relations which arise from

¹³ It would be interesting to explicitly construct the holomorphic partition function of these models on multicentered Taub-NUTs.

 $^{^{14}}$ We refer our readers to appendix A for a quick review.

the number of points of intersection among the associatives, and from gauging, and hence the corresponding system is not just a tensor product (although being close to one).

Based on the structure of the geometries $\mathcal{X}_{\mathbf{X}^{\vee},N}$ it is natural to expect that the factors $\mathcal{A}_{N_i}^k \otimes_{B_{ij}} \mathcal{A}_{N_j}^k$ fully determine the algebra of the associated quiver theory. Therefore, by 4d/7d correspondence we claim that for all these examples, there is an algebraic structure underlying the G_2 instanton partition function, which we schematically denote $\mathcal{A}_k(\mathcal{X}_{\mathbf{X}^{\vee},N})$. This is the algebra of G_2 instantons, meaning that the corresponding 7d G_2 instanton partition function is organized according to characters of $\mathcal{A}_k(\mathcal{X}_{\mathbf{X}^{\vee},N})$. The task of finding a precise definition of such an algebra is extremely hard, and in this section we build some evidence towards its existence exploiting string dualities, which allows to capture some of the features of its building blocks. From the perspective of the 4d $\mathcal{N} = 1$ theory, we expect the $\mathcal{A}_k(\mathcal{X}_{\mathbf{X}^{\vee},N})$ to be computable in terms of the structure of the complexes associated to the differential arising from the holomorphic twist on \mathbf{TN}_k . We will devote to this computation another paper in this series.

Exhibiting the details of the gluing rule goes beyond the scope of this short note. In what follows we proceed with a consistency check of (4.3): we consider the k = 1 case, and we prove that indeed equation (4.2) is verified, by considering a local model for the D6 brane intersection.

4.2. Building blocks from intersecting D6 branes

As a local model for the intersection between two D6 branes stacks, we can exploit a famous class of asymptotcally conical G_2 spaces, which are metric cones on complex weighted projective spaces $\mathbf{WCP}^3_{N,N,K,K}$

$$\mathcal{X}_{N,K} \equiv \mathcal{C}(\mathbf{WCP}^3_{N,N,K,K}), \tag{4.4}$$

here the case N = K = 1 coincides with the Bryant-Salamon G_2 cone over the threedimensional complex projective plane \mathbb{CP}^3 [49], while the other cases are natural generalization of the Bryant-Salamon construction [50, 51]. The physics of these models is understood in terms of a single chiral $U(N) \times U(K)$ bifundamental multiplet [50, 51]:

$$\mathcal{T}_{\mathcal{X}_{N,K}} = \left\{ \begin{array}{c} 4d \ \mathcal{N} = 1 \ \text{chiral} \\ U(N) \times U(K) \ \text{bifundamental} \end{array} \right\} \,. \tag{4.5}$$

It is well-known that the G_2 manifolds $\mathcal{X}_{N,K}$ are dual to a IIA configuration of N D6 branes intersecting with K D6 branes at angles. We choose a frame of reference such that the stack of N D6 branes is located along the directions 0, 1, 2, 3, 4, 5, 6 and the stack of K D6 branes is along the directions 0, 1, 2, 3 and it wraps a plane $\mathbf{L}_{\theta} \simeq \mathbf{R}^3$ which extends in the directions 4, 5, 6, 7, 8, 9, such that each of its components form an angle θ_i with the directions 4, 5, 6 respectively for i = 1, 2, 3.¹⁵ The M-theory background $\mathbf{TN}_k \times \mathcal{X}_{N,K}$ is therefore dual to the following type II_A background

	0	1	2	3	4	5	6	7	8	9		
$N \ \mathrm{D6}_1$	•	•	٠	٠	٠	٠	٠					
$M_1 \text{ D}2_1$					٠	•	٠					()
$K \text{ D6}_2$	•	•	•	•			$\mathbf{L}_{oldsymbol{ heta}}$					(4
$M_2 \ \mathrm{D2}_2$							$\mathbf{L}_{oldsymbol{ heta}}$					
\mathbf{TN}_L					•	•	•	•	•	•		

where we are marking the directions that are filled by the corresponding object. In the Type IIA frame the M2 branes wrapping the corresponding associative cycles are dual in this context to two stacks of D2 branes, one is parallel to D6₁ and is wrapping the 456 directions, the other is parallel to D6₂ and it is wrapping $\mathbf{L}_{\boldsymbol{\theta}}$.

From the 4d/7d correspondence we expect that the algebra $\mathcal{A}_N^L \otimes_{B_{12}=1} \mathcal{A}_K^L$ organises the generating function of some suitably regularised volumes of moduli spaces encoded in a quiver matrix model supported at the common intersection of these two stacks of D2 branes. These moduli spaces control the contributions of the M2 branes to the 7d partition function on the G_2 manifold.

Here, we are interested in probing the conjecture with an explicit computation. therefore, we consider the case L = 1 and we send the Taub-NUT radius to infinity, thus focusing on the topological M-theory limit of our parition function. The resulting algebra is the L = 0 version of the G_2 instanton algebra. In the absence of the Taub-NUT background it is relatively straightforward to derive the quiver which captures the D2 brane boundstates from the building blocks we described above, namely we will have two distinct ADHM-like quivers, each corresponding to one of the two factors of the algebra \mathcal{A}_N^0 and \mathcal{A}_K^0 , arising from the D2-D6 systems. On top of that from the intersection of the two D2 branes, we expect to obtain a single chiral bifundamental contributing to the quiver. As a result we obtain the quiver in figure 2.¹⁶

¹⁵In other words,

$$\mathbf{L}_{\boldsymbol{\theta}} = \begin{cases} x_4 = \tan \theta_1 x_5 \\ x_6 = \tan \theta_2 x_7 \\ x_8 = \tan \theta_3 x_9 \end{cases} \qquad \boldsymbol{\theta} = (\theta_1, \theta_2, \theta_3)$$
(4.6)

¹⁶ We review the relevant computation using string quantization in appendix B, and here we only report on the main result.



Figure 2: The quiver for the matrix model related to the $\mathcal{A}_N^0 \otimes_{B_{12}=1} \mathcal{A}_K^0$ algebra.

Increasing B_{12} , the quiver is essentially the same, only we obtain a collection S^{ℓ} of chiral bifundamental fields, at the intersections, which would carry an index $\ell = 1, ..., B_{12}$. Here, we will focus on the case $\ell = 1$, which corresponds to the orbifolds of the Bryant-Salamon G_2 cones that we are using as local models.

In appendix B, we discuss a proposal for the corresponding action for the matrix model supported at the intersection of the two D2 branes. From the D-term of the associated quiver description, we see that the two ADHM quivers satisfy their own complex moment map equations:

$$[X, Y] + IJ = 0,$$

$$[\tilde{X}, \tilde{Y}] + \tilde{I}\tilde{J} = 0,$$

(4.8)

and we also have a modified real moment map equation, which includes the contribution from the chiral bifundamental field ${\cal S}$

$$\begin{bmatrix} X, X^{\dagger} \end{bmatrix} + \begin{bmatrix} Y, Y^{\dagger} \end{bmatrix} + II^{\dagger} - JJ^{\dagger} - S^{\dagger}S = \xi \cdot \mathbf{I}_{M_1 \times M_1}, \begin{bmatrix} \tilde{X}, \tilde{X}^{\dagger} \end{bmatrix} + \begin{bmatrix} \tilde{Y}, \tilde{Y}^{\dagger} \end{bmatrix} + \tilde{I}\tilde{I}^{\dagger} - \tilde{J}\tilde{J}^{\dagger} + SS^{\dagger} = \xi \cdot \mathbf{I}_{M_2 \times M_2},$$
(4.9)

where ξ is an FI term. This is the starting point for a more formal analysis for the structures of the corresponding moduli spaces, which we describe in appendix C.

In Appendix C, we compute the equivariant K-theory index of the corresponding moduli space. Here, we briefly summarize the result, relegating the details to the appendix. The first step is to utilize a well-known fact, namely that the virtual tangent bundle of the moduli space encoded in the quiver diagram is captured by the cohomology of a specific complex:

$$\operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2}) \xrightarrow{\sigma} \operatorname{End}(\mathcal{V}_{1})^{\oplus 2} \oplus \operatorname{End}(\mathcal{V}_{2})^{\oplus 2} \oplus \mathcal{V}_{1}^{\oplus N} \oplus \mathcal{V}_{1}^{* \oplus N} \oplus \mathcal{V}_{2}^{\oplus K} \oplus \mathcal{V}_{2}^{* \oplus N}$$

$$\oplus \operatorname{Hom}(\mathcal{V}_{1}, \mathcal{V}_{2}) \oplus \operatorname{Hom}(\mathcal{V}_{2}, \mathcal{V}_{1}) \xrightarrow{d\mu_{\mathbb{C}}} \operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2}).$$

$$(4.10)$$

Here \mathcal{V}_1 , \mathcal{V}_2 are universal bundles of rank M_1 , M_2 , σ is related to the stability condition or real D-term equation, and μ is complex moment map equation. Our goal is to compute the Euler characteristics of the virtual tangent bundle of this moduli space¹⁷. To do so, it is necessary to find a suitable compactification of the moduli space, which typically is achieved by means of equivariance. We can achieve this by turning on five-torus $\mathbf{T}'_1 \times \mathbf{T}'_2 \times \mathbf{T}_3$ acting on $\mathcal{M}^{N,K}(M_1, M_2)$, defined by

$$(r_1, r_2) \in \mathbf{T}'_1 : (X, Y, I, J) \mapsto (r_1 X, r_2 Y, I, r_1 r_2 J),$$

$$(s_1, s_2) \in \mathbf{T}'_2 : (\tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}) \mapsto (s_1 \tilde{X}, s_2 \tilde{Y}, \tilde{I}, s_1 s_2 \tilde{J}),$$

$$c \in \mathbf{T}_3 : S \mapsto cS.$$

$$(4.11)$$

We show in the appendix that the torus fixed points of the moduli space are in one-to-one correspondence with M_1+2 copies of Young diagrams, see Proposition 5. Using this fact, we are able to compute the equivariant K-theoretic index of the moduli space of the G_2 instantons in Appendix C.5. Especially, the equivariant K-theory index nicely factorizes into two pieces in the limit $r_1r_2 \rightarrow 1$, $s_1s_2 \rightarrow 1$ for the equivariant parameters of the torus action. We show this in Proposition 8. We conjecture that each piece is captured by the Witten index that we will discuss in §4.3. Finally, in Appendix C.6 we explicitly show that in a certain limit of the equivariant parameters the equivariant K-theory index admits a structure of Verma modules of $\mathcal{A}_1^1 \otimes_1 \mathcal{A}_1^1$. This series of propositions is a strong evidence toward the existence of an algebra of G_2 instantons.

In sum, we claim that the contributions from the equivariant volumes of the G_2 instantons to the 7d partition function are captured by the equivariant K-theory index described above, which we denote $\mathcal{I}^1_{M_1,M_2}(N,K)$. This gives rise to the following explicit contribution to the partition function

$$\widehat{Z}_{U(1)}^{7D}(\mathcal{X}_{1,1}) = \sum_{M_1,M_2} Q_1^{M_1 1} Q_2^{M_2 1} \mathcal{I}_{M_1,M_2}^1(1,1) \,. \tag{4.12}$$

The main outcome of such analysis is that indeed, the resulting partition function is organised as a character of $\mathcal{A}_1^1 \otimes_1 \mathcal{A}_1^1$, which is a first consistency check for the ideas discussed here.

4.3. *G*₂ instantons from an SQM Witten index

In this section, we present an alternative way to compute the G_2 instanton partition function as a Witten index [95]. We can achieve this by performing T-duality along two directions on the $D2_1/D6_1/D2_2/D6_2$ brane system. The following table summarizes the resulting D-brane configuration $D0_1/D4_1/D4_2/D8_2$.

¹⁷This Euler character is interpreted physically in terms of a Witten index.

	0	1	2	3	4	5	6	7	8	9
Background					A	В	В	А	В	В
$(D2_1 \rightarrow)D0$					+					
$(D6_1 \rightarrow)D4_1$	+	+	+	+	+					
$(D2_2 \rightarrow)D4_2$					_	+	+		+	+
$(D6_2 \rightarrow)D8$	+	+	+	+	_	+	+		+	+

Table 1: Brane configuration after the T-duality. A and B denote angles and B-field. For instance, D4₂ and D8 branes extend over an oblique line in 47 plane.

Note that we T-dualize two of the directions previously occupied by a stack of $D6_1(D2_1)$ branes, which are forming angles with a stack of $D6_2(D2_2)$ branes. As we explained in appendix B, this T-duality converts D-branes at angle configuration to B-field backgrounds along 5689 directions. It does not modify the result of the string quantization, since the open string boundary condition do not change under the T-duality. In other words, we may simply promote the matrix model field content we discussed in the previous section to a 1d $\mathcal{N} = 4$ quiver supersymmetric quantum mechanics.

4.3.1. Supersymmetric quantum mechanics

strings	$\mathcal{N} = 4$ multiplets	fields	$\mathrm{SU}(2)_{-} \times \mathrm{SU}(2)_{+} \times \mathrm{SU}(2)_{-}^{R} \times \mathrm{SU}(2)_{+}^{R}$	$U(M_1) \times U(N)$			
		gauge field	(1, 1, 1, 1)				
	vector	scalar	(1, 1, 1, 1)				
		fermions	$({f 1},{f 2},{f 1},{f 2})$				
	Fermi	fermions	(2, 1, 2, 1)	(Adi 1)			
D0-D0	twisted hyper	scalars	$({f 1},{f 1},{f 2},{f 2})$	(Auj,1)			
	twisted hyper	fermions	$({f 1},{f 2},{f 2},{f 1})$				
	hypor	scalars	$({f 2},{f 2},{f 1},{f 1})$				
	nyper	fermions	(2, 1, 1, 2)				
	hypor	scalars	$({f 1},{f 2},{f 1},{f 1})$				
D0-D4 ₁	nyper	fermions	(1, 1, 1, 2)	$(\mathbf{M}_1, \mathbf{N})$			
	Fermi	fermions	(1, 1, 2, 1)				

Let us first collect the quantum mechanical fields that are related to D0 and $D4_1$ branes in table 2.

Table 2: The field content of the D0-D4 quantum mechanics. Note that 1d twisted multiplets does not have a matrix model counterpart. Indeed, the field decouples from the computation of the index.

They are the standard ADHM mechanical fields [87, 96–98] (see also [99, 100]). In the table, we organize the fields by their representation under $SO(4) = SU(2)_{-} \times SU(2)_{+}$ and $SO(4)_{R} = SU(2)_{-}^{R} \times SU(2)_{+}^{R}$, where SO(4) acts on 0123 directions and $SO(4)_{R}$ acts on 5689 directions.

Next, the fields that are not related to the usual ADHM quiver are from D0-D4₂ strings and D0-D8 strings, which are T-dual to D2₁-D2₂ strings and D2₁-D6₂ strings. However, we have seen in appendix B that the D2₁-D6₂(D0-D8) are massive and decouple in the IR. Hence, we only need to discuss the D0-D4₂ strings. Their T-dual is summarized in Table 3.

strings	$\mathcal{N} = 4$ multiplets	fields	$SU(2) \times SU(2)_+ \times SU(2)^R \times SU(2)_+^R$	$U(M_1) \times U(M_2)$
$D0_{-}D4_{-}$	twisted hyper	scalar	$({f 1},{f 1},{f 1},{f 2})$	$(\mathbf{M}, \mathbf{M}_{2})$
D0-D42	twisted hyper	fermions	$({f 1},{f 2},{f 1},{f 1})$	(111,1112)
	Fermi	fermions	(1, 1, 2, 1)	

Table 3: The quantum mechanical fields from the D0-D4₂ strings. These will introduce local operators that we can integrate out to form 1-loop determinants— see (4.16).



Figure 3: A subquiver of the whole quiver that is captured by the Witten index.

The quantum mechanical fields collected in Table 2 and Table 3 are fields that appear in a subquiver of the entire quiver for the system: compare figure 3 with figure 2. This is the same quiver that was needed for our analysis in appendix C — see figure Figure 6 there. We showed in Lemma 3 that G_2 instanton moduli space $\mathcal{M}^{N,K}(M_1, M_2)$ is an entire space of a trivial fibration over a single ADHM moduli space $\mathcal{M}^N(M_1)$ with the fiber being the moduli space described by Figure 3. Here we will mimic that construction from the perspective of a Witten index computation.

4.3.2. Witten index of SQM

Here we are interested in computing the following Witten index for our quiver in figure 3

$$\mathcal{I}^{M_{1},N} = Tr_{\mathcal{H}_{QM}} \left[(-1)^{F} e^{\beta \{Q,Q^{\dagger}\}} t^{2(J_{+}+J_{R})} u^{2J_{-}} v^{2J'_{R}} w_{i}^{2\Pi_{i}} x_{l}^{\Pi'_{l}} \right].$$
(4.13)

Here F, J_{\pm} , J_R , J'_R , Π_i , and Π'_l are the Fermion number and charges under the Cartan generators of $SU(2)_{\pm}$, $SU(2)^R_+$, $SU(2)^R_-$, U(N), $U(M_2)$. \mathcal{H}_{QM} is the Hilbert space of the

quantum mechanics.

The Witten index can be computed by supersymmetric localization of the quantum mechanical path integral. As it has now become a standard technique, we will simply take formulas from the original references and direct the reader to [99]. The difference between the original set-up and our set-up is the presence of $D0_1 - D4_2$ strings, which are T-dual of $D2_1 - D2_2$ strings. Since the other extra fields are all massive, the computation reduces to

$$\mathcal{I}^{M_1,N} = \int [d\phi_I] Z^{M_1}_{D0-D0}(\phi_I, \epsilon_+, \epsilon_-, \tilde{\epsilon}) Z^{M_1}_{D0-D4_1}(\phi_i, \epsilon_+, \epsilon_-, \tilde{\epsilon}, a_l) Z^{M_1}_{D0-D4_2}(\phi_I, \epsilon_+, \epsilon_-, m_l),$$
(4.14)

where ϕ_i are Cartan elements of $U(M_1)$, ϵ_{\pm} , $\tilde{\epsilon}$ are Cartan elements of $SU(2)_{\pm}$, $SU(2)_{-}^R$, and a_i , m_l are Cartan elements of U(N) and $U(M_2)$.

 $Z_{D0-D0}^{M_1} Z_{D0-D4_1}^{M_1}$ is given in [99, 101]:

$$Z_{D0-D0}^{M_1} Z_{D0-D4_1}^{M_1} = \prod_{I,J=1}^{M_1} \frac{\sinh'\frac{\phi_I - \phi_J}{2} \sinh\frac{\phi_I - \phi_J + 2\epsilon_+}{2}}{\sinh\frac{\phi_I - \phi_J + 2\epsilon_1}{2} \sinh\frac{\phi_I - \phi_J + 2\epsilon_2}{2}} \prod_{I,J=1}^{M_1} \frac{\sinh\frac{\phi_{IJ} \pm \tilde{\epsilon} - \epsilon_-}{2}}{\sinh\frac{\phi_{IJ} \pm \tilde{\epsilon} - \epsilon_+}{2}} \prod_{I=1}^{M_1} \prod_{i=1}^N \frac{\sinh\frac{\phi_I - a_i \pm \tilde{\epsilon}}{2}}{2\sinh\frac{\phi_I - a_i \pm \epsilon_+}{2}}$$
(4.15)

where the prime on sinh indicates that sinh(x) is omitted when x = 0.

We can read $Z_{D0-D4_2}^{M_1}$ from [101, (3.6)]. The difference here is that we do not have massless $D4_1 - D4_2$ strings. The chiral multiplet S maps to the twisted hypermultiplet in SQM.¹⁸ The corresponding 1-loop determinant is

$$Z_{D0-D4_2}^{M_1} = \prod_{I=1}^{M_1} \prod_{l=1}^{M_2} \frac{\sinh \frac{\phi_I - m_l \pm \epsilon_-}{2}}{\sinh \frac{\phi_I - m_l \pm \epsilon_+}{2}}.$$
(4.16)

After substituting the ingredients (4.15), (4.16) in (4.14), we can use the JK prescription [102, 103], which guides what poles to choose when one evaluates multidimensional complex integrals. Let us explain the contour of the integral (4.14) following [101]. We can classify the poles selected by the JK prescription in two types:

$$\mathcal{P}^{1}: \phi_{I} - a_{i} + \epsilon_{+} = 0, \quad \phi_{I} - \phi_{J} \pm \epsilon_{-} + \epsilon_{+} = 0,$$

$$\mathcal{P}^{2}: \phi_{I} - m_{l} - \epsilon_{+} = 0, \quad \phi_{I} - \phi_{J} \pm \tilde{\epsilon} - \epsilon_{+} = 0.$$
(4.17)

When evaluating the rank $M_1 \phi_I$ integral, one needs to choose M_1 poles out of (4.17) and evaluate the residue integral. In other words, since we have two sets, if we can pick M poles in \mathcal{P}^1 , we need to pick the rest $M_1 - M$ poles in \mathcal{P}^2 . It was shown in [101] that each of the first and second set of poles is classified by the N-colored Young diagrams $Y = \{Y_1, \cdots, Y_N\}$ with total size M and M_2 -colored¹⁹ Young diagrams \tilde{Y} with total size

Here by twisted hypermultiplet we mean a multiplet whose scalar transforms non trivially under $SU(2)^{\check{R}}_{\pm}$ but trivially under $SU(2)_{\pm}$. ¹⁹The case for $M_2 = 1$ was discussed in [101], but the statement can be extended for general M_2 . For

 $M_1 - M$, respectively. Each of the contributions can be naturally understood as the D0 bound states on the $D4_1$ branes and on the $D4_2$ branes. With this information, we can organize the result of the integral as the following double summation over two sets of Young diagrams:

$$\mathcal{I}^{M_1,N} = \sum_{Y} \sum_{\tilde{Y}} f_{Y,\tilde{Y}}(\epsilon_+, \epsilon_-, \tilde{\epsilon}, a_i, m_l)$$
(4.18)

It is assumed in the above formula that we take $e^{\tilde{\epsilon}} \to 0$ limit to decouple the twisted multiplet contribution, as we commented in the caption of Table 2. This is a valid limit, as discussed in [101].

Compared to [101], we do not have a Fermi multiplet, which would come from $D4_1 - D4_2$ strings. The corresponding 1-loop determinant is

$$Z_{D4_1-D4_2} = \prod_{i=1}^{N} \prod_{j=1}^{M_2} 2\sinh\frac{a_i - m_j}{2}.$$
(4.19)

Since $D4_1 - D4_2$ strings are not related to D0 branes(i.e. (4.19) does not depend on ϕ_I), this 1-loop determinant does not affect the ϕ_I integral. Hence, the structure of our integral (4.18) is the same as that of [101].

Computing $\mathcal{I}^{M_1,N}$ for each M_1 , we can form a generating series by weighting each $\mathcal{I}^{M_1,N}$ with the instanton counting parameter Q_1 with power M_1 :

$$1 + \sum_{M_1=1} Q_1^{M_1} \mathcal{I}^{M_1,N}.$$
(4.20)

This gives a part of the G_2 instanton partition function that is associated to the fiber of the bundle

$$\pi: \mathcal{M}^{N,K}(M_1, M_2) \to \mathcal{M}^N(M_1).$$
(4.21)

We naturally conjecture that (4.20) is equivalent to the half (C.46) of the factorization formula in Proposition 8, since both of them are labeled by the same information: two sets of Young diagrams Y and \tilde{Y} .

Similarly, one can do the same computation for the second saddle of the G_2 instanton partition function that is represented by the following subquiver:

instance, quiver gauge theory with higher rank gauge nodes was discussed in [104].



Figure 4:

This is done by first choosing a coordinate frame where we find $D2_2$, $D6_2$ branes aligning in the 4,5,6 coordinate axis and on the other hand $D2_1$, $D6_2$ branes forming angles with 4,5,6 axis. Performing the same T-duality, $D2_2$ brane stack into D0' brane stack. The new configuration is as follows

	0	1	2	3	4	5	6	7	8	9
Background					A	В	В	А	В	В
$(D2_1 \rightarrow)D4'_2$					_	+	+		+	+
$(D6_1 \rightarrow)D8'$	+	+	+	+	_	+	+		+	+
$(D2_2 \rightarrow)D0'$					+					
$(D6_2 \rightarrow)D4_1'$	+	+	+	+	+					

Table 4: Brane configuration after the T-duality. A and B denote angles and B-field. For instance, D4'₂ and D8' branes extend over a line in 47 plane.

Doing the exactly same computation as above, we can form a generating series by weighting each $\mathcal{I}^{M_2,K}$ with the instanton counting parameter Q_2 .

$$1 + \sum_{M_2=1} Q_2^{M_2} \mathcal{I}^{M_2,K}.$$
(4.22)

This gives a part of the G_2 instanton partition function that is associated to the fiber of the bundle

$$\pi': \mathcal{M}^{N,K}(M_1, M_2) \to \mathcal{M}^K(M_2).$$
(4.23)

Similarly to the other half, we expect that (4.22) is equivalent to the other half of the factorization formula in Proposition 8, since both of them are labeled by the same information: two sets of Young diagrams Y and \tilde{Y} .

4.4. A relation with 3d $\mathcal{N} = 2$ theories with impurities

We conclude this section with a remark about a relation between our setup and $3d \mathcal{N} = 4$ systems with supersymmetry breaking impurities. Exploiting the Taub-NUT radius to perform a T-duality with IIB, starting from equation (4.7), we obtain the following setup:



From this T-dual frame, it is interesting to notice a connection between our computation and 3d $\mathcal{N} = 4$ theories with supersymmetry breaking impurities, which is somehow reminiscent of the analysis in [53, 77, 78]. Indeed, the brane setup involving D3₁, D5₁, and NS5 is the famous Hanany-Witten brane engineering for 3d $\mathcal{N} = 4$ systems [105]. On top of that, we have the injection of the D3₂ and D5₂ stacks, which can be described as impurities localised at points from the perspective of the corresponding 3d $\mathcal{N} = 4$ theories. The partition functions of M2 branes contributing to the 7d instanton partition function should have an interpretation also in terms of this kind of systems. In a sense, these would be captured by 3d $\mathcal{N} = 4 S^3$ or $S^1 \times S^2$ parition function with the insertion of impurity operators.

Acknowledgements

We thank Christopher Beem, Kevin Costello, Iñaki García Etxebarria, Dominic Joyce, Raeez Lorgat, Maxim Zabzine for useful discussion and email correspondences. We are especially grateful to Kevin Costello for asking a fascinating question that led us to initiate this project. The work of MDZ and RM has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 851931). Research of JO was supported by ERC Grant numbers 864828 and 682608. Research at the Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development & Innovation.

A. A brief review of the twisted M-theory background

Twisted M-theory is eleven-dimensional supergravity background that is specified by three data (g, C, Ψ) : the metric, M-theory 3-form, and bosonic ghost for a certain component of local supersymmetry. The metric is in general a product metric of a hyperKahler 4-manifold \mathcal{M}_4^{HK} and a G_2 7-manifold $\mathcal{M}_7^{G_2}$. In the original work [24], the geometry was given by

$$\underbrace{\mathbb{C}_z \times \mathbb{C}_w}_{\mathcal{M}_4^{HK}} \times \underbrace{\mathbb{R}_t \times \mathbb{C}_1 \times \mathrm{TN}_K}_{\mathcal{M}_7^{G_2}}.$$
(A.1)

The generalized Killing spinor Ψ imposes a topological and holomorphic twist on \mathcal{M}_4^{HK} and $\mathcal{M}_7^{G_2}$ respectively. More precisely, we turn a nonzero bosonic ghost Ψ [23] for the local supersymmetry of the supergravity. We determine Ψ in the same way we get the scalar supercharge in a twisted QFT [25]. Start with an 11d Killing spinor Ψ_{11} and decompose it into 7d and 4d components. Each of the component spinors decomposes into representations of G_2 and $SU(2)_- \times U(1)_+$, where $U(1)_+$ is the Cartan of $SU(2)_+$.

$$\Psi_{11} = \Psi_7 \otimes \Psi_4 = (\mathbf{1}_{G_2} \oplus 7_{G_2}) \otimes (1_{-1} \oplus 1_{+1} \oplus 2_0).$$
 (A.2)

We keep a non zero bosonic ghost for the scalar $\Psi = (\mathbf{1}_{\mathbf{G}_2} \otimes \mathbf{1}_{-1})$. There are Ψ_m and $\Psi_{\dot{\alpha}}$ such that

$$\{\Psi, \Psi_m\} = P_m, \quad \{\Psi, \Psi_{\dot{\alpha}}\} = P_{-\dot{\alpha}}.$$
(A.3)

This implies that the dynamics in $\mathcal{M}_7^{G_2}$ direction becomes topological and that in \mathcal{M}_4^{HK} becomes holomorphic.

We can deform Ψ into Ψ_{ϵ} [24] such that

$$\Psi_{\epsilon}^2 = \epsilon \vec{\mathcal{V}},\tag{A.4}$$

where $\vec{\mathcal{V}}$ is a Killing vector field on $\mathcal{M}_7^{G_2}$. With this deformation, we can incorporate an Omega deformation Ω_{ϵ_1} on \mathbb{C}_1 and $\Omega_{\epsilon_2} \times \Omega_{\epsilon_3}$ on $TN_K \sim (\mathbb{C}_2 \times \mathbb{C}_3)/\mathbb{Z}_K$. ϵ_i are deformation parameters and from now on we will denote Ω_{ϵ_i} deformed \mathbb{C}_i plane as \mathbb{C}_{ϵ_i} .

Lastly, the M-theory 3-form is

$$C = \epsilon_2 V^d d\bar{z} d\bar{w},\tag{A.5}$$

where V^d is a linear dual 1-form of the vector field V, which generates a rotation along the Taub-NUT circle S_{TN}^1 .

As a result of the Omega deformation, we can expect that a localization happens in the twisted M-theory. We can explicitly see this by going to the type IIA frame, by reducing along S_{TN}^1 . The Taub-NUT geometry converts into K D6 branes and the closed string states decouple from the Q-closed spectrum as discussed in [24]. Moreover, due to Ω_{ϵ_1} , the 7d SYM living on K D6 branes localizes to the 5d U(K) topological holomorphic Chern-Simons theory [106]:

$$\int_{\mathbb{C}_z \times \mathbb{C}_w \times \mathbb{R}_t} dz \wedge dw \wedge \left(A dA + \frac{2}{3} A \star_{\epsilon_2} A \star_{\epsilon_2} A \right).$$
(A.6)

Here, \star_{ϵ_2} is a wedge product deformed by the non-commutative B-field $B = \epsilon_2 d\bar{z} d\bar{w}$, which descends from the M-theory 3-form. This is a typical Moyal product, which is defined between two holomorphic functions as

$$f \star_{\epsilon_2} g = fg + \epsilon_2 \frac{1}{2} \epsilon_{ij} \frac{\partial}{\partial_{z_i}} f \frac{\partial}{\partial_{z_j}} g + \cdots$$
 (A.7)

Another important difference with the usual 5d Chern-Simons action is that the 5d gauge field has only three components:

$$A = A_t dt + A_{\bar{z}} d\bar{z} + A_{\bar{w}} d\bar{w}, \tag{A.8}$$

due to the presence of the holomorphic top-form dzdw in the action.

Due to the non-commutativity, the naive gauge symmetry algebra \mathfrak{gl}_K is modified to $\mathfrak{g} = \operatorname{Diff}_{\epsilon_2} \mathbb{C}_z \otimes \mathfrak{gl}_K$. Without the quantum deformation, i.e. $\epsilon_1 = 0$, the symmetry algebra $A_{\text{classical}} = A_{\epsilon_1=0,\epsilon_2}$ of the 5d CS theory is generated by the Fourier modes of the ghost c[m,n] for the gauge symmetry, together with the BRST differential δ . As a graded associative algebra, $A_{\text{classical}}$ is isomorphic to $C^*(\mathfrak{g})$, which is the Chevalley-Eilenberg algebra of cochains on the Lie algebra \mathfrak{g} .

One of the major achievements of [36] was to use Kozsul duality $C^*(\mathfrak{g}) = U(\mathfrak{g})$ and identify the 5d CS algebra of operators with the protected operator algebra $\mathcal{A}_K^{(N)}$ of NM2 branes(on $\mathbb{R}_t \times \mathbb{C}_{\epsilon_1}$) probing the twisted M-theory background in the large N limit. Moreover, there is a surjective map s [36]

$$s: U_{\epsilon_1}(\mathfrak{gl}(K) \otimes \operatorname{Diff}_{\epsilon_2}(\mathbb{C})) \to \mathcal{A}_K^{(N)},$$
 (A.9)

which is compatible with a sequence of surjective maps $s_i : \mathcal{A}_K^{(i+1)} \to \mathcal{A}_K^{(i)}$. Moreover, the intersection of the kernels of $U_{\epsilon_1}(\mathfrak{gl}(K) \otimes \operatorname{Diff}_{\epsilon_2}(\mathbb{C})) \to \mathcal{A}_K^{(N)}$ for all N is zero. In this sense, one can define $U_{\epsilon_1}(\mathfrak{gl}(K) \otimes \operatorname{Diff}_{\epsilon_2}(\mathbb{C}))$ as a *large* N *limit* of $\mathcal{A}_K^{(N)}$, and denote it by \mathcal{A}_K by dropping the superscript (∞) .

Given the abstract description of $\mathcal{A}_{K}^{(N)}$ and its relation with the 5d CS algebra of operators, let us provide the explicit description of $\mathcal{A}_{K}^{(N)}$. One convenient UV descriptions of the M2 branes worldvolume theory is 3d $\mathcal{N} = 4 U(N)$ gauge theory with 1 adjoint hypermultiplet(with scalars X, Y) and K fundamental hypermultiplets(with scalars I, J) [107]. Due to the topological twist applied to $\mathbb{R}_t \times \mathbb{C}_{\epsilon_1} \subset \mathcal{M}_7^{G_2}$, we pass to a particular Q-cohomology that only captures the Higgs branch chiral ring. This consists of gaugeinvariant operators made of X, Y, I, J, divided by the ideal generated by the F-term relation,

$$X_c^a Y_b^c - X_b^c Y_c^a + I_b J^a = \epsilon_2 \delta_b^a, \tag{A.10}$$

where we only presented gauge indices a, b, c, d, suppressing flavor indices.

 Ω_{ϵ_1} deforms the chiral ring into an algebra \mathcal{A}_K [108–110] by making Poisson brackets

into commutators

$$[X_b^a, Y_d^c] = \epsilon_1 \delta_d^a \delta_b^c, \quad [J^b, I_a] = \epsilon_1 \delta_a^b, \tag{A.11}$$

and the theory localizes on one-dimensional topological quantum mechanics [111, 112]

$$\frac{1}{\epsilon_1} \int_{\mathbb{R}_t} \text{Tr}[\epsilon_2 A_t + X D_t Y + J D_t I] dt.$$
(A.12)

Note that ϵ_1 is a deformation parameter, and ϵ_2 is an FI parameter of the 3d $\mathcal{N} = 4$ gauge theory.

The algebra $\mathcal{A}_{K}^{(N)}$ is generated by

$$t_{m,n} = \frac{1}{\epsilon_1} \mathrm{STr} X^m Y^n, \tag{A.13}$$

where $STr[\bullet]$ means to take a trace of a symmetrization of a polynomial \bullet . The generating relations of the algebra was proposed in [113], proved in [114], and used in [113,115] to compute a large set of correlation functions in 3d $\mathcal{N} = 4$ ADHM gauge theories.

B. The D2 matrix model from string quantization

In this appendix we review the computation that leads to the equation describing the moduli space of vacua for the matrix model supported at the intersection of the two D2 brane stacks in our local model arising from the $\mathcal{X}_{N,K}$ geometry. We will use the language of 3d supersymmetry because the multiplets we obtain organise naturally accordingly. However, these are all fields supported at a point (the point of intersection of the two D2 brane stacks).

B.1. Supersymmetry

We can determine the supersymmetry preserved by the intersecting D6 branes [116] by solving the following equations of 10d spinor (ϵ_L, ϵ_R) equation $(\epsilon_L$ and ϵ_R are 10d Majorana-Weyl spinors with different chiralities).

$$\epsilon_L = \Gamma^{0123456} \epsilon_R,$$

$$\epsilon_L = \Gamma^{01234'5'6'} \epsilon_R,$$
(B.1)

where 456, 4'5'6' denote the directions along two stacks of D6 branes in the 456789 direction and $\Gamma^{i_1...i_n}$ denote the fully anti-symmetrized products of Gamma matrices $\Gamma^{i_1}, \ldots, \Gamma^{i_n}$.

We need to remember that there is a B-field turned on 0123 directions. Let us parametrize it as

$$B_{01} = -B_{10} = b_1, \quad B_{23} = -B_{32} = b_2. \tag{B.2}$$

and set $b_a = \tan \pi v_a$. Hence,

$$-\frac{1}{2} < v_a < \frac{1}{2}.$$
 (B.3)

By [117], the Gamma matrix equation above changes to

$$\epsilon_L = \mathcal{B}_{0123} \Gamma^{0123456} \epsilon_R,$$

$$\epsilon_L = \mathcal{B}_{0123} \Gamma^{01234'5'6'} \epsilon_R,$$
(B.4)

where

$$\mathcal{B}_{0123} = \exp \pi \left((v_1 + 1/2) \Gamma^0 \Gamma^1 + (v_2 + 1/2) \Gamma^2 \Gamma^3 \right)$$
(B.5)

For now, this change will not affect the calculation, since both of (B.1) and (B.4) reduce into

$$\Gamma^{456}\epsilon_R = \Gamma^{4'5'6'}\epsilon_R,\tag{B.6}$$

where we used $(\Gamma^{ij})^2 = 1$. Therefore, it only constrains the 6d part ϵ_{6d} of the 10d spinor $\epsilon_R = \epsilon_{10d} = \epsilon_{4d} \otimes \epsilon_{6d}$ whose eigenvalues under the action of

$$\Gamma^{01}, \quad \Gamma^{23}, \quad \Gamma^{47}, \quad \Gamma^{58}, \quad \Gamma^{69}.$$
 (B.7)

is $(\pm, \pm, \pm, \pm, \pm)$. Those are raised and lowered by the action of

$$\Gamma^{0\pm} = \frac{1}{2} (\pm \Gamma^0 + \Gamma^1), \ \Gamma^{1\pm} = \frac{1}{2} (\Gamma^2 \pm i\Gamma^3),$$

$$\Gamma^{2\pm} = \frac{1}{2} (\Gamma^4 \pm i\Gamma^7), \ \Gamma^{3\pm} = \frac{1}{2} (\Gamma^5 \pm i\Gamma^8), \ \Gamma^{4\pm} = \frac{1}{2} (\Gamma^6 \pm i\Gamma^9).$$
(B.8)

Let \mathbb{R} be the SO(6) rotation that takes the first stack of D6 branes to the second stack of D6 branes.

$$\mathbb{R} = \operatorname{diag}(e^{i\theta_1}, e^{-i\theta_1}, e^{i\theta_2}, e^{-i\theta_3}, e^{i\theta_3}, e^{-i\theta_3}).$$
(B.9)

Then we have

$$\Gamma^{0123456} = \mathbb{R}\left(\Gamma^{01234'5'6'}\right)\mathbb{R}^{-1}.\tag{B.10}$$

The solution for (B.6) only exists when \mathbb{R} belongs to SU(3) subgroup of the SO(6) that rotates 456789. The SU(3) condition is

$$\theta_1 \pm \theta_2 \pm \theta_3 = 0 \mod 2\pi. \tag{B.11}$$

There are 4 spinors $\epsilon_{10d} = \epsilon_{4d} \otimes \epsilon_{6d}$ such that $\epsilon_{4d} = (\pm, \pm)$, ϵ_{6d} with one choice of signs out of 8 possible choices (\pm, \pm, \pm) . In other words, a 4d theory along 0123 directions preserves a chiral $\mathcal{N} = 1$ supersymmetry.

Now, introduce $D2_1$ branes along 456 directions. This gives another spinor equation:

$$\epsilon_L = \Gamma^{456} \epsilon_R. \tag{B.12}$$

Coupling with the first line of (B.4), we get

$$\mathcal{B}_{0123}\Gamma^{0123456}\epsilon_R = \Gamma^{456}\epsilon_R \tag{B.13}$$

Decomposing $E_R = \epsilon_{4d} \otimes \epsilon_{6d}$, we get

$$(\mathcal{B}_{0123}\Gamma^{0123}\epsilon_{4d})\otimes(\Gamma^{456}\epsilon_{6d})=\epsilon_{4d}\otimes(\Gamma^{456}\epsilon_{6d})$$
(B.14)

Hence, it reduces to

$$\mathcal{B}_{0123}\Gamma^{0123}\epsilon_{4d} = -\epsilon_{4d} \tag{B.15}$$

It has a solution only if

$$\exp \pi(\pm i(v_1 + 1/2) \pm i(v_2 + 1/2)) = -1.$$
(B.16)

In other words, if v_1 , v_2 satisfy

$$v_1 = \pm v_2, \tag{B.17}$$

we have

$$\Gamma^{0123}\epsilon_{4d} = \epsilon_{4d}.\tag{B.18}$$

Note, however, by [117] only $v_1 = -v_2$ is allowed.²⁰ In any case, out of 4 spinors $\epsilon_{4d} = (\pm, \pm)$, only 2 spinors survive: $\epsilon_{4d} = (+, +), (-, -)$; the number of the preserved supersymmetry is independent of B as shown in [118].

Similarly, consider $D2_2$ branes along 4'5'6' directions. This gives

$$\epsilon_L = \Gamma^{4'5'6'} \epsilon_R. \tag{B.19}$$

Coupling with the second line of (B.1), we get (B.15). Therefore, out of 4 spinors $\epsilon_{4d} = (\pm, \pm)$, only 2 spinors survive: $\epsilon_{4d} = (+, +), (-, -)$. We are allowed have both $D2_1$ and $D2_2$ branes and preserve the same amount of supersymmetry, since both lead to the same equation (B.15).

Therefore, we conclude that D2/D6/D6' configuration preserves 2 supercharges. In D2 brane worldvolume, this is $3d \mathcal{N} = 1$ supersymmetry.

Before doing the sting quantization, let us review supermultiplets of 3d $\mathcal{N} = 1$ SUSY [119]. First, $\mathcal{N} = 1$ vector multiplet \mathcal{V} consists of a gauge field A_{μ} and a Majorana fermion λ_{α}

$$\mathcal{V} = (A_{\mu}, \lambda_{\alpha}), \text{ where } \mu = 1, 2, 3, \ \alpha = +, -.$$
 (B.20)

Second, $\mathcal{N} = 1$ scalar multiplet Φ consists of a real scalar ϕ , a Majorana fermion ψ_{α} ,

$$\star B = \epsilon_2 \star (d\bar{z}_1 \wedge d\bar{z}_2) = \epsilon_2 \star [(dx_1 - idx_3) \wedge (dx_2 - idx_4)] = -\epsilon_2 d\bar{z}_1 \wedge d\bar{z}_2 = -B$$

²⁰Our B-field is anti-self-dual, which is consistent with $v_1 + v_2 = 0$, as shown in [117].

and an auxiliary field D

$$\Phi = (\phi, \psi_{\alpha}, D). \tag{B.21}$$

B.2. String quantization

We will follow [120] to account for the B-field background– recall we have turned B-field on TN_1^{21} . To quantize open strings, we need to mode-expand worldsheet bosons Z and fermions Ψ that respect two types of boundary conditions at two boundaries. In the presence of the B-field, the equation of motion is given by [121]

$$\partial_{++}Z = e^{-2\pi i\nu}\partial_{--}Z, \quad \Psi^+ = e^{2\pi i\nu}\Psi^- \text{ at } \sigma = 0,$$

$$\partial_{++}Z = e^{-2\pi i\mu}\partial_{--}Z, \quad \Psi^+ = e^{2\pi i\mu}\Psi^- \text{ at } \sigma = \pi,$$

(B.22)

Solving the equations above, we see the worldsheet boson Z and R-sector fermion Ψ have modes $\mathbb{Z} + \theta$ with $\theta = \mu - \nu$. The NS-sector fermions have modes $\mathbb{Z} + \epsilon$ with $\epsilon = \theta + \frac{1}{2}$. We will denote the Neumann, Dirichlet, Twisted(by the B-field effect) boundary conditions as **N**, **D**, **T**.

To arrange the spectra by their energy, we need to find a zero-point energy. The NS sector zero-point energy is

$$E_0^{NS} = \frac{1}{8} - \frac{1}{2} ||\theta| - \frac{1}{2}|.$$
(B.23)

The first excited state has energy $E_1^{NS} = E_0^{NS} + |\epsilon|$ if $-\frac{1}{2} \le \epsilon \le \frac{1}{2}$ and $E_1^{NS} = E_0^{NS} + |1-\epsilon|$ if $\frac{1}{2} < \epsilon < \frac{3}{2}$. The R sector zero-point energy $E_0^R = 0$. The first excited state energy is $E_1^R = |\theta|$ for $0 \le |\theta| \le \frac{1}{2}$ and $1 - |\theta|$ for $\frac{1}{2} \le |\theta| \le 1$.

Let us recall the brane configuration from equation (4.7). We can quantize strings that stretch between two stacks of D-branes. Start with $D2_1 - D2_1$ strings. Although we have determined the minimal amount of supersymmetry that the brane configuration preserves is 2, we will later see that the genuine $3d \mathcal{N} = 1$ multiplet is massive and decoupled in the low energy. Therefore, we will supplement the discussion with the 3d $\mathcal{N} = 2$ representation. Moreover, we will sometimes find helpful to reorganize the $\mathcal{N} = 2$ supermultiplets into the $3d \mathcal{N} = 4$ supermultiplets, even though we do not have such an enhanced supersymmetry. There will be just one genuine $3d \mathcal{N} = 2$ massless multiplet after all.

D2-D2 string NS sector: We know $E_0^{NS} = -\frac{1}{2}$, so to get the massless states, we need to act with NS fermion oscillators $d_1^{\mu\dagger}$. For $\mu = 4, 5, 6$, which are **NN** directions, $d_1^{\mu\dagger}$ gives rise to three states, which consist of the 3d G gauge field A_{μ} . For other d's, which are **TT** and **DD**, they give rise to 7 scalars ϕ_a valued in $adj(\mathfrak{g})$ of G.

²¹Note that the D6-branes at angles configuration is equivalent to D0-D6 brane system with B-field on ND directions [117].

D2-D2 string R sector: R-sector vacuum energy is always zero. Since we have 10 NN+DD directions, we can use all 10 zero modes to act on the R-sector vacuum state and form 32 dimensional ground state. The GSO projection projects out half of 32 and the remaining 16 fermionic states pair up with the bosons determined from the NS sector to complete the 3d $\mathcal{N} = 1$ supermultiplet.

Therefore, we see

$$D2_1 - D2_1$$
 strings $\equiv \mathcal{V}$ and $7 \Phi_a$ (B.24)

One can split 7 Φ_a into { Φ_7, Φ_8, Φ_9 } and { $\Phi_0, \Phi_1, \Phi_2, \Phi_3$ }. In the 3d $\mathcal{N} = 2$ notation, one of the scalars in the first set can be identified with the real scalar σ and the other two scalars in the first set can be thought of as the complex scalar φ . On the other hand, one can split the 4 scalars in the second set into 2 complex scalars. In the presence of $\mathcal{N} = 2$ SUSY, each of them forms $\mathcal{N} = 2$ chiral multiplets X, Y.

D2-D6 string NS sector: Let us denote two complex directions in 4 ND directions as *a* and *b*. We first need to derive NS zero point energy. Recall B-field is only turned on the 4 ND directions. Hence, the zero point energy in these directions is

$$\frac{1}{8} - \frac{1}{2}|v_a| + \frac{1}{8} - \frac{1}{2}|v_b| = \frac{1}{4} - \frac{1}{2}(|v_a| + |v_b|).$$
(B.25)

Other directions do not have twisted boundary conditions by the B-field. Taking two directions out of three D2 world volume directions as the lightcone, we can compute the zero point energy for the **NN** and **DD** directions:

$$\frac{1}{8} - \frac{1}{4} + \frac{1}{8} - \frac{1}{4} = -\frac{1}{4}.$$
(B.26)

Summing (B.25) and (B.26), we get

$$E_0^{NS} = -\frac{1}{2}(|v_a| + |v_b|). \tag{B.27}$$

By applying a suitable oscillator that raises the energy by $|v_a|$ and $|v_b|$, we get 4 states with energies $\frac{1}{2}(\pm v_a \pm v_b)$. 2 out of 4 states are projected out by the GSO projection and the remaining ones have energy $\pm \frac{1}{2}(v_a - v_b)$. Those are two complex scalars transforming in (N, M_1) of $U(N) \times U(M_1)$ of the D6 and D2 gauge groups with masses

$$m^2 = \pm \frac{1}{2}(v_a - v_b).$$
 (B.28)

One of the two is tachyonic, which needs further treatment later. Since for 3d $\mathcal{N} = 1$ the elementary multiplet is a real scalar multiplet, it indicates that there are 4 scalars Φ^{I} .

D2-D6 string R sector: As before, the zero point energy of R-sector is zero. For

this case, we have 6 zero modes from 6 NN+DD directions. Therefore, we have $8 = 2^3$ dimensional ground states. We can label those as $|\alpha, A, \dot{A}\rangle$, where α is 3d spinor index, A, \dot{A} are the spinor index of $SU(2) \times SU(2) = SO(4)$ that rotates the transverse direction of D2 inside D6 worldvolume. Then, these fermionic massless fields make a supersymmetric completion of the bosons obtained from the NS sector analysis.

Therefore, we get 4 scalar multiplets:

$$D2_1 - D6_1 \text{ strings } \equiv 4 \Phi^I. \tag{B.29}$$

In the presence of 3d $\mathcal{N} = 2$ supersymmetry, one can recombine the 4 real scalars into 2 complex scalars I, J. Each of them is a part of the 3d $\mathcal{N} = 2$ fundamental chiral multiplet.

Equivalence between B-field and Angle configuration

It is helpful to recognize the equivalence between B-field background and D-branes at angles [122] to unify the background in a single frame before we do $D2_1 - D6_2$ quantization, whose ND directions involve both B-field background and the angle background.

B-field background along 2n directions is T-dual to the D-branes at angles along 2n direction. For instance, our $D6_1/D6_2$ at angles configuration is T-dual to D0/D6 brane system with B-field on 6 ND directions.

To see this, let us look at the open string boundary condition on 456789 directions in the original $D6_1/D6_2$ system.

$$\partial_n X^{4,5,6} = \partial_\tau X^{7,8,9} = 0,$$

$$\partial_n \left[\cos\varphi_{ij} X^i + \sin\varphi_{ij} X^j \right] = 0,$$

$$\partial_\tau \left[-\sin\varphi_{ij} X^i + \cos\varphi_{ij} X^j \right] = 0,$$

(B.30)

where $i \in \{4, 5, 6\}$ and $j \in \{7, 8, 9\}$.

T-dualizing along X^4 , X^5 , X^6 modifies the equations as

$$\partial_{\tau} X^{4,5,6,7,8,9} = 0,$$

$$\tan \varphi_{ij} \partial_n X^j + \partial_{\tau} X^i = 0,$$

$$\tan \varphi_{ij} \partial_n X^i - \partial_{\tau} X^j = 0.$$

(B.31)

These boundary conditions are exactly that of D0 branes bounded on D6 branes with B-field on 6 ND directions. The explicit dictionary is the following:

$$B_{ij} = -B_{ji} = \tan\varphi_{ij} \tag{B.32}$$

Recall that we had a useful parametrization for the B-field to utilize in the string quant-

ization

$$B_{i,i+3} = b_{i-2} = \tan \pi v_{i-2}. \tag{B.33}$$

Hence, we can identify

$$\varphi_{i,i+3}/\pi = v_{i-2}.\tag{B.34}$$

 $D2_1-D6_2$, $D2_2-D6_1$ strings NS sector: First of all, this configuration preserves 2 supersymmetries, which are the minimal amount among all configuration. We can see this from the following Gamma matrix exercise:

$$\Gamma^{4'5'6'}\epsilon_l = \epsilon_r, \quad \Gamma^{0123456}\epsilon_l = \epsilon_r \implies \Gamma^{0123456}\epsilon_l = \Gamma^{4'5'6'}\epsilon_l. \tag{B.35}$$

Decompose $\epsilon_l = (\epsilon_l)_4 \otimes (\epsilon_l)_6$ then the above becomes

$$\left(\Gamma^{0123}(\epsilon_l)_4\right) \otimes \left(\Gamma^{456}(\epsilon_l)_6\right) = (\epsilon_l)_4 \otimes \left(\Gamma^{4'5'6'}(\epsilon_l)_6\right) \Rightarrow \begin{cases} \Gamma^{0123}(\epsilon_l)_4 = (\epsilon_l)_4 \\ \Gamma^{456}(\epsilon_l)_6 = \Gamma^{4'5'6'}(\epsilon_l)_6. \end{cases} (B.36)$$

We know from [116] the second equation gives one solution and the first equation gives two solutions for $(\epsilon_l)_4$: (+, +) or (-, -).

Let us resume the $D2_1 - D6_2$ quantization. We T-dualize 4, 5, 6 directions to convert the D branes intersecting with angles into transversely intersecting D branes with Bfield background. We will parametrize the B-field background by v_2 , v_3 , v_4 . Now, the $D2_1 - D6_2$ configuration becomes $D(-1)_1 - D9_2$: D-instantons probing D9 brane worldvolume with 4 ND directions with the original B-field and the remaining 6 ND directions with the new B-field. In this unified background, we compute the NS zero point energy. First, start with the 0123 direction where the B-field is applied:

$$\frac{1}{8} - \frac{1}{2}|v_0| + \frac{1}{8} - \frac{1}{8}|v_1| = \frac{1}{4} - \frac{1}{2}(|v_0| + |v_1|).$$
(B.37)

Second, for 456789 directions with the new B-field, we have

$$\frac{1}{8} - \frac{1}{2}|v_2| + \frac{1}{8} - \frac{1}{8}|v_3| + \frac{1}{8} - \frac{1}{8}|v_4| = \frac{3}{8} - \frac{1}{2}(|v_2| + |v_3| + |v_4|).$$
(B.38)

Summing them up, we have

$$E_0^{NS} = \frac{5}{8} - \frac{1}{2}(|v_0| + |v_1| + |v_2| + |v_3| + |v_4|).$$
(B.39)

We produce the excited states by acting NS fermion oscillators that increase energy by $|v_i|$ on the NS vacuum state. First 32 states have energies

$$\frac{5}{8} + \frac{1}{2}(\pm v_0 \pm v_1 \pm v_2 \pm v_3 \pm v_4).$$
(B.40)

Recall that there is a list of constraints:

$$v_0 + v_1 = 0,$$

$$\pi(v_2 \pm v_3 \pm v_4) = 0 \mod 2\pi,$$

$$-\frac{1}{2} < v_i < \frac{1}{2},$$

(B.41)

where each of the conditions comes from (B.17), (B.11), (B.3). Moreover, by the construction of the twisted M-theory, we have a small non-commutativity parameter

$$v_0 = -v_1 = \epsilon_2 \ll 1. \tag{B.42}$$

As for the second constraint in (B.41), we have a freedom to fix $v_2 = v_3 = v_4 = \frac{2}{3}$. To be consistent with the third line, we shift v_2 , v_3 , v_4 by the period of the tangent function, 1:

$$v_2 = v_3 = v_4 = \frac{2}{3} - 1 = -\frac{1}{3}.$$
 (B.43)

Hence, (B.40) becomes

$$\frac{5}{8} + \frac{1}{2}(\pm\epsilon_2 \pm \epsilon_2 \pm \frac{1}{3} \pm \frac{1}{3} \pm \frac{1}{3} \pm \frac{1}{3}).$$
(B.44)

Notice that only the minimal energy configuration can potentially be negative or zero energy depending on the value of ϵ_2 being greater or equal to $\frac{1}{8}$, since

$$E_0 = \frac{5}{8} + \frac{1}{2}(-\epsilon_2 - \epsilon_2 - \frac{1}{3} - \frac{1}{3} - \frac{1}{3}) = \frac{1}{8} - \epsilon_2.$$
(B.45)

From now, we will denote 32 states by the signs in (B.44), e.g. (\pm, \pm, \pm, \pm) .

Now, let us consider GSO projection

$$\Gamma_{GSO} = \frac{1}{2} (1 + (-1)^{F_{NS}}). \tag{B.46}$$

We assign $F_{NS} = 1$ to the NS vacuum state(the minimal energy configuration). In the above convention, it corresponds to the state (-, -, -, -, -). Then, all states with even number of '+' are projected out, since their $(-1)^{F_{NS}} = -1$. Therefore, $|E_0\rangle$ is projected out and we do not have any massless or tachyonic state in the spectrum. In other words, in the low energy limit that we are interested in, there is no interesting state coming from the NS sector of $D2_1 - D6_2$ or $D2_2 - D6_1$ strings.

Still, there are massive scalars. The lightest modes are from 0123 directions, due to the relative strength of the B-field. After the GSO projection, we get 2 bosons.

 $D2_1-D6_2$ and $D2_2-D6_1$ strings R sector: Since two D-branes are fully transverse to each other, there are no NN or DD directions in this configuration. Hence, there are no fermionic zero modes. Since R-sector zero point energy is zero, there is still one zero energy state, the R-sector vacuum. However, the GSO projection projects this state out. Hence, we do not obtain any massless fermionic state.

Still, there are massive fermions that pair up with the massive scalars determined from the NS sector to complete the two lightest massive supermultiplet.

 $D6_1$ - $D6_2$ strings: We already know that they give rise to the 4d $\mathcal{N} = 1$ bi-fundamental chiral multiplets at the 4d intersection [116]. Since we are only interested in the 3d worldvolume theory of either $D2_1$ or $D2_2$ branes, we treat the D6 branes as a heavy background. Hence, $D6_1 - D6_2$ strings will not play any important role in our story.

 $D2_1$ - $D2_2$ strings: We can T-dualize the $D6_1 - D6_2$ configuration in the 4 directions of intersection and arrive at $D2_1 - D2_2$ configuration. Since T-duality does not change the amount of preserved supersymmetry, the $D2_1 - D2_2$ string quantization will yield the dimensional reduction of the 4d $\mathcal{N} = 1$ bifundamental chiral multiplet determined above and live at the intersection of $D2_1$ and $D2_2$. It smears out the 3d worldvolume of both $D2_1$ and $D2_2$ branes. Hence, we will treat it as $3d \mathcal{N} = 2$ bifundamental chiral multiplet S, which consists of two $\mathcal{N} = 1$ scalar multiplets S_1 and S_2 . As we will see later, S will provide an edge between two conventional ADHM quivers and make our moduli space more interesting.

Summary

Let us summarize the result of the string quantization of our brane system in the following

 $D2_{1} - D2_{1} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ vector multiplets } \mathcal{V}_{2} \text{ and 7 scalar multiplets } \Phi_{a}$ $D2_{2} - D2_{2} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ vector multiplets } \mathcal{V}_{2} \text{ and 7 scalar multiplets } \Phi_{a}^{\prime}$ $D2_{1} - D2_{2} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ bifundamental scalar multiplets } S_{1}, S_{2}$ $D2_{1} - D6_{1} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ 4 scalars multiplets } \Phi^{I}$ $D2_{2} - D6_{2} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ 4 scalars multiplets } \Phi^{\prime I}$ $D2_{1} - D6_{2} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ massive scalar multiplet}$ $D2_{2} - D6_{2} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ massive scalar multiplet}$ $D2_{2} - D6_{1} \text{ strings} \equiv 3d \ \mathcal{N} = 1 \text{ massive scalar multiplet}$ $D6_{1} - D6_{1} \text{ strings} \equiv 7d \ U(N) \text{ vector multiplet}$ $D6_{2} - D6_{2} \text{ strings} \equiv 7d \ U(K) \text{ vector multiplet}$ $D6_{1} - D6_{2} \text{ strings} \equiv 4d \ \mathcal{N} = 1 \text{ bifundamental chiral multiplet}$ (B.47)

For our purpose that will be described in the next subsection, we can focus on the strings that are associated with the 3d massless fields: $D2_i - D2_i$, $D2_i - D6_i$, $D2_1 - D2_2$ strings. Considering only the massless states, we get 3d $\mathcal{N} = 2$ theory on the D2 branes.

B.3. Deriving the equations for the moduli space

Given the field content of the 3d $\mathcal{N} = 2$ theory on each of the D2 brane stacks, we can proceed to study the matrix model supported at the point of intersection. It is useful first to recall our situation. We have converted the G_2 geometry and the G_2 instantons into the D6 brane and D2 brane stacks. Hence, the relevant moduli space \mathcal{M} is the moduli space of the intersecting D2 brane stacks fluctuating inside the intersecting D6 brane stacks. The directions of the fluctuation are 0123. Therefore, we need to focus on the quantum fields that arise from the quanta of the strings that correspond to 0123 directions. We have denoted those fields as $\Phi_{0,1,2,3}$, $\Phi^{0,1,2,3}$, $\tilde{\Phi}_{0,1,2,3}$, $\tilde{\Phi}^{0,1,2,3}$. We also need to include the chiral bifundamental S that is supported at the intersection of the D2 stacks.

In terms of 3d $\mathcal{N} = 2$ superfields, \mathcal{M} is parametrized by the scalar components of

$$X = \Phi_1 + i\Phi_2, \ Y = \Phi_3 + i\Phi_4, \ I = \Phi^1 + i\Phi^2, \ J = \Phi^3 + i\Phi^4,$$

$$\tilde{X} = \tilde{\Phi}_1 + i\tilde{\Phi}_2, \ \tilde{Y} = \tilde{\Phi}_3 + i\tilde{\Phi}_4, \ \tilde{I} = \tilde{\Phi}^1 + i\tilde{\Phi}^2, \ \tilde{J} = \tilde{\Phi}^3 + i\tilde{\Phi}^4,$$

$$S = S_1 + iS_2.$$
(B.48)

The moduli space is given by the zeros of the scalar potential, which is supposed to be

$$V = \frac{g^2}{2} \operatorname{Tr} D^2 + \dots, \qquad (B.49)$$

where D is the auxiliary field in the off-shell multiplets. The zeros of V is obtained by setting D = 0. Our goal is to derive a set of D-term equations in the 3d $\mathcal{N} = 2$ system that fully specifies the moduli space. To do that, the first step is to figure out all auxiliary fields in the supermultiplets.

Let us start from $D2_1 - D2_1$ strings:

$$\mathcal{V} = (A_{\mu}, \lambda_{\alpha}),$$

$$\Phi_{7} = (\phi_{7}, \psi_{7,\alpha}, D_{7}), \quad \Phi_{8} = (\phi_{8}, \psi_{8,\alpha}, D_{8}), \quad \Phi_{9} = (\phi_{9}, \psi_{9,\alpha}, D_{9}),$$

$$\Phi_{0} = (\phi_{0}, \psi_{0,\alpha}, D_{0}), \quad \Phi_{1} = (\phi_{1}, \psi_{1,\alpha}, D_{1}), \quad \Phi_{2} = (\phi_{2}, \psi_{2,\alpha}, D_{2}), \quad \Phi_{3} = (\phi_{3}, \psi_{3,\alpha}, D_{3}).$$

(B.50)

We will think of ϕ_7 as the real scalar σ and (ϕ_8, ϕ_9) as the complex scalar φ in 3d $\mathcal{N} = 2$ notation. We find a triplet (D_7, D_8, D_9) of the auxiliary D's. Combine them into D_{ab} , where a, b are SU(2) indices. In the last line, we find 4 scalars $\phi_0, \phi_1, \phi_2, \phi_3$. In $\mathcal{N} = 4$ notation, we can combine $X = \phi_0 + i\phi_1, Y = \phi_2 + i\phi_3$ and collect them into a fictitious²² SU(2) doublet $\mathcal{Q} = (X, Y)$. There is a coupling between \mathcal{Q}_a and D_{ab} :

$$\operatorname{Tr} \, \mathcal{Q}^a D^b_a \mathcal{Q}_b. \tag{B.51}$$

²²Even though there is no such SU(2) symmetry in the 3d $\mathcal{N} = 2$ system, this formal bookkeeping tool enables us to present the material in a compact way.

Also, there is a quartic term of D:

$$\mathrm{Tr} \ |D_a|^2. \tag{B.52}$$

From $D2_1 - D6_1$ strings, we got

$$\Phi^{0} = (\phi^{0}, \psi^{0}_{\alpha}, G^{0}), \quad \Phi^{1} = (\phi^{1}, \psi^{1}_{\alpha}, G^{1}), \quad \Phi^{2} = (\phi^{2}, \psi^{2}_{\alpha}, G^{2}), \quad \Phi^{3} = (\phi^{3}, \psi^{3}_{\alpha}, G^{3}), \tag{B.53}$$

where we distinguish the auxiliary fields from those of $D2_1 - D2_1$ strings by denoting them as G, not D. In $\mathcal{N} = 4$ notation, we can combine $I = \phi^0 + i\phi^1$, $J = \phi^2 + i\phi^3$ and further form a fictitious SU(2) doublet $\mathcal{I} = (I, J)$. It also couples to D_{ab} :

$$\operatorname{Tr} \mathcal{I}^a D^b_a \mathcal{I}_b. \tag{B.54}$$

We can do the similar analysis for $D2_1-D2_2$ strings as they also form 3d $\mathcal{N} = 2$ chiral multiplets. Let us combine two real scalar superfields into a complex chiral multiplet $S = S_1 + iS_2$ and treat it as a component of the fictitious SU(2) doublet $\mathcal{S} = (S, T)$. Since there is no T to pair with, one should treat this expression formally. Then, one can write down, at least formally, the relevant Lagrangian:

$$(\mathcal{S}^a)^I_i (D^b_a)^i_j (\mathcal{S}_b)^j_J, \tag{B.55}$$

where i, j are $U(M_1)$ gauge index for M_1 D2 branes on 456 direction and I, J are $U(M_2)$ gauge index for M_2 D2 branes on 4'5'6' direction.

Finally, as our gauge group contains U(1) as a subgroup, there is an FI term:

$$\operatorname{Tr} \xi^b_a D^a_b. \tag{B.56}$$

Collecting the D-related Lagrangian terms, we have the following matrix model action

$$\operatorname{Tr}\left[|D_a|^2 + \mathcal{Q}^b D_b^a \mathcal{Q}_a + \mathcal{I}^b D_b^a \mathcal{I}_a + \xi_a^b D_b^a + \operatorname{Tr} \mathcal{S}^a D_a^b \mathcal{S}_b\right], \qquad (B.57)$$

which gives the equation of motion of $D_7 = D$

$$D = [X, X^{\dagger}] + [Y, Y^{\dagger}] + II^{\dagger} - JJ^{\dagger} - S^{\dagger}S - \xi \cdot I_{M_1 \times M_1} = 0.$$
 (B.58)

The equation of motion of $D_8 + iD_9 = D_{\mathbb{C}}$ is

$$D_{\mathbb{C}} = [X, Y] + IJ = 0.$$
 (B.59)

We notice that there is no S in this equation, due to the absence of the $\mathcal{N} = 4$ completion of S. In other words, T is zero in the formal expression (B.55).

We can do the same for the second quiver. Collecting all the relevant D-terms and

solving the EOM for $\tilde{D}_7 = \tilde{D}$, we get

$$\tilde{D} = \left[\tilde{X}, \tilde{X}^{\dagger}\right] + \left[\tilde{Y}, \tilde{Y}^{\dagger}\right] + \tilde{I}\tilde{I}^{\dagger} - \tilde{J}\tilde{J}^{\dagger} + SS^{\dagger} - \xi \cdot \mathbf{I}_{M_2 \times M_2} = 0.$$
(B.60)

The equation of motion of $\tilde{D}_8 + i\tilde{D}_9 = \tilde{D}_{\mathbb{C}}$ is

$$\tilde{D}_{\mathbb{C}} = \left[\tilde{X}, \tilde{Y}\right] + \tilde{I}\tilde{J} = 0.$$
(B.61)

Hence, the moduli space $\mathcal{M}_{M_1,M_2}^{N,K}$ is the $U(M_1) \otimes U(M_2)$ quotient of the space of solutions of (B.58), (B.59), (B.60), (B.61), where the groups $U(M_1) \otimes U(M_2)$ act as

$$(X, Y, I, J, S) \mapsto \left(g^{-1}Xg, g^{-1}Yg, g^{-1}I, Jg, g^{-1}S\right), \quad \text{where } g \in U(M_1),$$

$$\left(\tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S\right) \mapsto \left(\tilde{g}^{-1}\tilde{X}\tilde{g}, \tilde{g}^{-1}\tilde{Y}\tilde{g}, \tilde{g}^{-1}\tilde{I}, \tilde{J}\tilde{g}, \tilde{g}^{-1}S\right), \quad \text{where } \tilde{g} \in U(M_2)$$
(B.62)

The FI parameter ξ and the non-commutativity parameter ϵ_2

There is a relation between the FI parameter ξ and the non-commutativity parameter ϵ_2 . We have observed in the $D2_1 - D6_1$ string quantization a tachyon with its mass squared given by

$$m^2 = -(v_1 - v_2). \tag{B.63}$$

We can resolve this potential instability and supersymmetry breaking issue by looking at the induced mass term in the 3d Lagrangian [120]:

$$-\frac{g^2}{2} \operatorname{Tr} \left(-\xi I I^{\dagger} + \xi J^{\dagger} J\right). \tag{B.64}$$

Mass squared of I is $\xi \frac{g^2}{2}$ and that of J is $-\xi \frac{g^2}{2}$. Hence, identifying this information with (B.63), we get:

$$v_1 - v_2 = -\xi,$$
 (B.65)

Because $v_1 = -v_2 = \epsilon_2$ in our background, the FI parameter is related to the noncommutativity parameter ϵ_2 :

$$\xi \sim \epsilon_2. \tag{B.66}$$

This condition makes sure the low energy action to be supersymmetric. For the detailed discussion on the tachyon condensation and restoration of the supersymmetry, see [120, 123].

C. Formal description of the moduli spaces

C.1. Moduli space as a subspace of a Nakajima quiver variety

If we add an arrow T from \mathbb{C}^{M_2} to \mathbb{C}^{M_1} , see Figure 5, then it is the double ²³ of the quiver with same nodes but arrows are $X, I, \tilde{X}, \tilde{I}, S$.



Figure 5:

Let us denote the corresponding Nakajima quiver variety of Figure 5 by $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$. Apparently, $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ is non-empty and is smooth of dimension $2M_1M_2 + 2KM_2 + 2NM_1$. Recall that $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ is defined as the space of solutions to the equations

$$[X, Y] + IJ + TS = 0,$$

$$\begin{bmatrix} \tilde{X}, \tilde{Y} \end{bmatrix} + \tilde{I}\tilde{J} - ST = 0,$$

$$[X, X^{\dagger}] + [Y, Y^{\dagger}] + II^{\dagger} - JJ^{\dagger} - S^{\dagger}S + TT^{\dagger} = \xi \cdot \mathbf{I}_{M_{1} \times M_{1}},$$

$$[\tilde{X}, \tilde{X}^{\dagger}] + [\tilde{Y}, \tilde{Y}^{\dagger}] + \tilde{I}\tilde{I}^{\dagger} - \tilde{J}\tilde{J}^{\dagger} + SS^{\dagger} - T^{\dagger}T = \xi \cdot \mathbf{I}_{M_{2} \times M_{2}},$$

(C.1)

modulo the action of $U(M_1) \times U(M_2)$. The first two equations of (C.1) says that the complex moment map $\mu_{\mathbb{C}}(X, Y, I, J, \tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S, T) = 0$ and the last two equations of (C.1) says that the real moment map $\mu_{\mathbb{R}}(X, Y, I, J, \tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S, T) = (\xi \cdot I_{M_1 \times M_1}, \xi \cdot I_{M_2 \times M_2})$.

On the moduli space $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$, there is a universal rank M_1 bundle denoted by \mathcal{V}_1 , and a universal rank M_2 bundle denoted by \mathcal{V}_2 , and a universal map $\mathcal{T}: \mathcal{V}_2 \to \mathcal{V}_1$.

Assume that $\xi \neq 0$, and denote by $\widetilde{M}^{N,K}(M_1, M_2)$ the space of solutions to (C.1), then it is well-known that $U(M_1) \times U(M_2)$ acts on $\widetilde{M}^{N,K}(M_1, M_2)$ freely, and moreover the pullback of the vanishing locus of \mathcal{T} is exactly the locus in $\widetilde{M}^{N,K}(M_1, M_2)$ where T in the equations (C.1) is zero. So we have

²³For a quiver Q, we define its double \overline{Q} to be the quiver with the same nodes as Q and for every arrow of Q we add an arrow of reverse direction.

Lemma 1. Assume that $\xi \neq 0$, then $\mathcal{M}^{N,K}(M_1, M_2)$ embeds into $\widetilde{M}^{N,K}(M_1, M_2)$ as the vanishing locus of \mathcal{T} .

Recall that $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ has algebro-geometric description [124]. Let's assume that $\xi > 0$, then $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ is the space of **stable** solutions to the equations

$$[X, Y] + IJ + TS = 0,$$

$$[\tilde{X}, \tilde{Y}] + \tilde{I}\tilde{J} - ST = 0,$$

(C.2)

modulo the action of $\operatorname{GL}_{M_1} \times \operatorname{GL}_{M_2}$. Here "stable" means that the only subspace of $\mathbb{C}^{M_1} \oplus \mathbb{C}^{M_2}$, which contains the image of I and \tilde{I} and is invariant under the actions of $X, Y, \tilde{X}, \tilde{Y}, S, T$ is $\mathbb{C}^{M_1} \oplus \mathbb{C}^{M_2}$ itself. This has the following obvious consequences:

Lemma 2. Assume that $\xi > 0$, then we have an isomorphism:

$$\mathcal{M}^{N,K}(M_1, M_2) = \mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}/\text{GL}_{M_1} \times \text{GL}_{M_2}, \qquad (C.3)$$

where "stable" means the locus inside $\mu_{\mathbb{C}}^{-1}(0)$ such that $\{X, Y, I, J, \tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S\}$ satisfies that

$$\mathbb{C}\langle X, Y \rangle \operatorname{Im}(I) = \mathbb{C}^{M_1}, \ \mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle (\operatorname{Im}(\tilde{I}) + \operatorname{Im}(S)) = \mathbb{C}^{M_2}.$$

Remark 1. If we assume $\xi < 0$ instead, then the stability condition in the above lemma becomes

* The only subspace of $\mathbb{C}^{M_1} \oplus \mathbb{C}^{M_2}$, which is annihilated by $J \oplus \tilde{J}$ and invariant under the actions of $X, Y, \tilde{X}, \tilde{Y}, S, T$ is 0.

Remark 2. Lemma 2 says that

$$(\mu_{\mathbb{C}} \times \mu_{\mathbb{R}})^{-1}(0, \xi \cdot \mathbf{I}_{M_1 \times M_1}, \xi \cdot \mathbf{I}_{M_2 \times M_2})/U(M_1) \times U(M_2) = \mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}/\mathrm{GL}_{M_1} \times \mathrm{GL}_{M_2},$$

and the following technical result [124, Proposition 3.5] provide a geometric interpretation of this equality:

• Let x be a point in $\mu_{\mathbb{C}}^{-1}(0)$, then x is stable if and only if the $\operatorname{GL}_{M_1} \times \operatorname{GL}_{M_2}$ orbit through x intersects with $\mu_{\mathbb{R}}^{-1}(\xi \cdot I_{M_1 \times M_1}, \xi \cdot I_{M_2 \times M_2})$.

Namely, if we consider the union of $\operatorname{GL}_{M_1} \times \operatorname{GL}_{M_2}$ orbits of $(\mu_{\mathbb{C}} \times \mu_{\mathbb{R}})^{-1}(0, \xi \cdot I_{M_1 \times M_1}, \xi \cdot I_{M_2 \times M_2})$, then this set is exactly $\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}$, and it is known that $\operatorname{GL}_{M_1} \times \operatorname{GL}_{M_2}$ acts on $\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}$ freely, so we have

$$\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}/\text{GL}_{M_1} \times \text{GL}_{M_2}$$

$$= (\text{GL}_{M_1} \times \text{GL}_{M_2}) \cdot (\mu_{\mathbb{C}} \times \mu_{\mathbb{R}})^{-1} (0, \xi \cdot \text{I}_{M_1 \times M_1}, \xi \cdot \text{I}_{M_2 \times M_2})/\text{GL}_{M_1} \times \text{GL}_{M_2} \qquad (C.4)$$

$$= (\mu_{\mathbb{C}} \times \mu_{\mathbb{R}})^{-1} (0, \xi \cdot \text{I}_{M_1 \times M_1}, \xi \cdot \text{I}_{M_2 \times M_2})/U(M_1) \times U(M_2).$$

Notice that when $\xi > 0$, the stability condition implies that the subset of quiver data (X, Y, I, J) is stable as an individual ADHM quiver. This shows that:

Lemma 3. Assume that $\xi > 0$, then there is a map $\pi : \mathcal{M}^{N,K}(M_1, M_2) \to \mathcal{M}^N(M_1)$, where $\mathcal{M}^N(M_1)$ is the moduli space of torsion-free sheaves on \mathbb{P}^2 framed at \mathbb{P}^1_{∞} with rank N and second Chern number M_1 . Moreover, π is a locally trivial fibration with fibers isomorphic to the quiver variety associated to the quiver diagram.



Figure 6:

The stability condition is that $\mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle (\operatorname{Im}(\tilde{I}) + \operatorname{Im}(S)) = \mathbb{C}^{M_2}$.

Proof. π is defined by $(X, Y, I, J, \tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S) \mapsto (X, Y, I, J)$. On the other hand, if we use the universal bundle \mathcal{V}_1 on $\mathcal{M}^N(M_1)$ to form a fibration of quiver variety associated to Figure 6 with stability condition that $\mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle (\operatorname{Im}(\tilde{I}) + \operatorname{Im}(S)) = \mathbb{C}^{M_2}$, where \mathbb{C}^{M_1} is replaced by the bundle \mathcal{V}_1 , then the total space is exactly the moduli space of stable solutions to equations (4.8).

Proposition 1. Assume that $\xi > 0$ and K = 0, then $\pi : \mathcal{M}^{N,0}(M_1, M_2) \to \mathcal{M}^N(M_1)$ is a locally trivial fibration with fibers isomorphic to $\operatorname{Quot}^{M_2}(\mathcal{O}_{\mathbb{C}^2}^{M_1})^{-24}$.

Proof. In the quiver 6, if we set K = 0, then the F-term equation is $[\tilde{X}, \tilde{Y}] = 0$, and the stability condition reads that $\mathbb{C}[\tilde{X}, \tilde{Y}] \operatorname{Im}(S) = \mathbb{C}^{M_2}$, this is exactly the definition of moduli space of quotient sheaf of $\mathcal{O}_{\mathbb{C}^2}^{M_1}$ such that the \mathbb{C} -dimension of the sheaf is M_2 . Note that \mathbb{C}^{M_2} is the linear space of quotient sheaf and \tilde{X}, \tilde{Y} is the coordinate functions of \mathbb{C}^2 .

In general $\mathcal{M}^{N,K}(M_1, M_2)$ is highly singular, but the following proposition shows that the singularity of $\mathcal{M}^{N,K}(M_1, M_2)$ is not too bad under some mild assumptions.

²⁴Quot^{M_2}($\mathcal{O}_{\mathbb{C}^2}^{M_1}$) is the moduli space of quotient sheaf of $\mathcal{O}_{\mathbb{C}^2}^{M_1}$ such that the \mathbb{C} -dimension of the sheaf is M_2 .

Proposition 2. Assume that $\xi \neq 0$ and $NK \neq 0$, then the embedding $\mathcal{M}^{N,K}(M_1, M_2) \hookrightarrow \widetilde{M}^{N,K}(M_1, M_2)$ is regular of codimension M_1M_2 . Moreover, locally the set of matrix elements of \mathcal{T} is a regular sequence of length M_1M_2 ²⁵.

Proof. It suffices to prove that $\mathcal{M}^{N,K}(M_1, M_2)$ is local complete intersection ²⁶ (l.c.i) of pure dimension ²⁷ $M_1M_2 + 2M_1N + 2M_2K$, then it automatically follows that locally the set of matrix elements of \mathcal{T} is a regular sequence of length M_1M_2 . To this end, we need to show that $\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}$ is l.c.i of pure dimension $M_1M_2 + 2M_1N + 2M_2K + M_1^2 + M_2^2$, since the quotient map $\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}} \to \mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}/\text{GL}_{M_2} \times \text{GL}_{M_1} = \mathcal{M}^{N,K}(M_1, M_2)$ is a principal $\text{GL}_{M_2} \times \text{GL}_{M_1}$ bundle and l.c.i property descends to a smooth morphism.

We claim that the map

 $\mu_{\mathbb{C}}: \mathbf{M_1} \times \mathbf{M_1}^* \times \mathrm{End}(\mathbf{M_1})^2 \times \mathbf{M_2} \times \mathbf{M_2}^* \times \mathrm{End}(\mathbf{M_2})^2 \times \mathrm{Hom}(\mathbf{M_1}, \mathbf{M_2}) \to \mathfrak{gl}_{M_1} \times \mathfrak{gl}_{M_2}$

is flat ²⁸. Note that the component Hom($\mathbf{M_1}, \mathbf{M_2}$) where S takes value in, plays no role in the map $\mu_{\mathbb{C}}$, so $\mu_{\mathbb{C}}$ factors through the product of moment maps

$$\mu_{\mathbb{C}}^{N} \times \mu_{\mathbb{C}}^{K} : \left(\mathbf{M}_{1} \times \mathbf{M}_{1}^{*} \times \operatorname{End}(\mathbf{M}_{1})^{2}\right) \times \left(\mathbf{M}_{2} \times \mathbf{M}_{2}^{*} \times \operatorname{End}(\mathbf{M}_{2})^{2}\right) \to \mathfrak{gl}_{M_{1}} \times \mathfrak{gl}_{M_{2}}.$$

And it is known that $\mu_{\mathbb{C}}^{N} : \mathbf{M}_{\mathbf{1}} \times \mathbf{M}_{\mathbf{1}}^{*} \times \operatorname{End}(\mathbf{M}_{\mathbf{1}})^{2} \to \mathfrak{gl}_{M_{1}}$ is flat, using Crawley-Boevey's criterion on the flatness of moment map [125, Theorem 1.1] ²⁹. Hence both $\mu_{\mathbb{C}}^{N}$ and $\mu_{\mathbb{C}}^{K}$ are flat and we deduce that $\mu_{\mathbb{C}}$ is flat.

Since $\{0\} \hookrightarrow \mathfrak{gl}_{M_1} \times \mathfrak{gl}_{M_2}$ is a regular embedding, the flatness of $\mu_{\mathbb{C}}$ implies that $\mu_{\mathbb{C}}^{-1}(0)$ embeds into the ambient space regularly, and it has dimension

$$\dim \mathbf{M_1} \times \mathbf{M_1}^* \times \operatorname{End}(\mathbf{M_1})^2 \times \mathbf{M_2} \times \mathbf{M_2}^* \times \operatorname{End}(\mathbf{M_2})^2 \times \operatorname{Hom}(\mathbf{M_1}, \mathbf{M_2}) - \dim \mathfrak{gl}_{M_1} \times \mathfrak{gl}_{M_2}$$
$$= M_1 M_2 + 2M_1 N + 2M_2 K + M_1^2 + M_2^2.$$
(C.5)
This shows that $u^{-1}(0)$ is l.c.i of of pure dimension $M_1 M_2 + 2M_1 N + 2M_1 K + M_2^2 + M_2^2$

This shows that $\mu_{\mathbb{C}}^{-1}(0)$ is l.c.i of of pure dimension $M_1M_2 + 2M_1N + 2M_2K + M_1^2 + M_2^2$, and obviously its open subset $\mu_{\mathbb{C}}^{-1}(0)^{\text{stable}}$ has the same property.

In the proof of the proposition, we show that

• Assume that $\xi \neq 0$ and $NK \neq 0$, then every irreducible component of $\mathcal{M}^{N,K}(M_1, M_2)$ has dimension $M_1M_2 + 2M_1N + 2M_2K$. In particular, the virtual dimension is the actual dimension.

²⁵A regular sequence of length n in a commutative ring \mathcal{A} is a set of elements $a_1, \dots, a_n \in \mathcal{A}$ such that a_{i+1} does not have zero-divisor in the quotient ring $\mathcal{A}/(a_1, \dots, a_i)\mathcal{A}$, for all i ranging from 0 to n-1 (set $a_0 = 0$). The key point is that, if \mathcal{A} is smooth, then a sequence $a_1, \dots, a_n \in \mathcal{A}$ is regular if and only if dim Spec $\mathcal{A} - \dim \text{Spec}\mathcal{A}/(a_1, \dots, a_i)\mathcal{A} = n$.

²⁶Local complete intersection means locally embeds into a smooth ambient space and is cut out by a regular sequence of equations.

 $^{^{27}}$ Pure dimension means all irreducible components have the same dimension.

 $^{^{28}{\}rm Flatness}$ means pullback of short exact sequence of sheaves is a short exact sequence.

²⁹We use the equivalence between (1) and (4) in [125, Theorem 1.1], and easy computation shows that (4) holds.

C.2. Virtual tangent bundle

Assume that $\xi \neq 0$ in this subsection. It is well known that the tangent bundle of $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ is the cohomology of the complex

$$\operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2}) \xrightarrow{\sigma} \operatorname{End}(\mathcal{V}_{1})^{\oplus 2} \oplus \operatorname{End}(\mathcal{V}_{2})^{\oplus 2} \oplus \mathcal{V}_{1}^{\oplus N} \oplus \mathcal{V}_{1}^{*\oplus N} \oplus \mathcal{V}_{2}^{\oplus K} \oplus \mathcal{V}_{2}^{*\oplus N} \\
\oplus \operatorname{Hom}(\mathcal{V}_{1}, \mathcal{V}_{2}) \oplus \operatorname{Hom}(\mathcal{V}_{2}, \mathcal{V}_{1}) \xrightarrow{d\mu_{\mathbb{C}}} \operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2})$$
(C.6)

where the middle term has cohomology degree zero. $d\mu_{\mathbb{C}}$ is the differential of the moment map, and explicitly it is written as

$$(x, y, \tilde{x}, \tilde{y}, i, j, \tilde{i}, \tilde{j}, s, t) \mapsto ([X, y] + [x, Y] + iJ + Ij + sT, [\tilde{X}, \tilde{y}] + [\tilde{x}, \tilde{Y}] + \tilde{i}\tilde{J} + \tilde{I}\tilde{j} - St).$$
(C.7)

 σ descends from the action of $\mathfrak{gl}_{M_1} \times \mathfrak{gl}_{M_2}$, and explicitly it is written as

$$(u,v) \mapsto ([u,X],[Y,u],[v,\tilde{X}],[\tilde{Y},v],uI,-Ju,v\tilde{I},-\tilde{J}v,vS-Su,uT-Tv).$$
(C.8)

It is easy to verify that $d\mu_{\mathbb{C}} \circ \sigma = 0$. Note that σ is injective and $d\mu_{\mathbb{C}}$ is surjective by stability. Our moduli space $\mathcal{M}^{N,K}(M_1, M_2)$ embeds into $\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)$ regularly, and is locally defined by M_1M_2 equations coming from matrix elements of $\mathcal{T} \in \text{Hom}(\mathcal{V}_2, \mathcal{V}_1)$, so $\mathcal{M}^{N,K}(M_1, M_2)$ has a *perfect obstruction theory* $T\widetilde{\mathcal{M}}^{N,K}(M_1, M_2)|_{\mathcal{M}^{N,K}(M_1, M_2)} \to$ $\text{Hom}(\mathcal{V}_2, \mathcal{V}_1)$, and this complex is quasi-isomorphic to the tangent complex $\mathbb{T}\mathcal{M}^{N,K}(M_1, M_2)$. This obstruction theory can be written as a complex of tautological bundles:

$$\operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2}) \xrightarrow{\sigma} \operatorname{End}(\mathcal{V}_{1})^{\oplus 2} \oplus \operatorname{End}(\mathcal{V}_{2})^{\oplus 2} \oplus \mathcal{V}_{1}^{\oplus N} \oplus \mathcal{V}_{1}^{* \oplus N} \oplus \mathcal{V}_{2}^{\oplus K} \oplus \mathcal{V}_{2}^{* \oplus N} \\ \oplus \operatorname{Hom}(\mathcal{V}_{1}, \mathcal{V}_{2}) \xrightarrow{d\mu_{\mathbb{C}}} \operatorname{End}(\mathcal{V}_{1}) \oplus \operatorname{End}(\mathcal{V}_{2})$$
(C.9)

The difference between (C.9) and (C.6) is that $\operatorname{Hom}(\mathcal{V}_2, \mathcal{V}_1)$ drops out and chain maps are restricted to the rest of components. In (C.9), σ is injective by stability condition, but $d\mu_{\mathbb{C}}$ is not necessarily surjective. We shall show that $d\mu_{\mathbb{C}}$ fails to be surjective at some points on $\mathcal{M}^{N,K}(M_1, M_2)$.

C.3. Some subvarieties of moduli space

We assume that $\xi > 0$ in this subsection. Let us investigate some subvarieties of $\mathcal{M}^{N,K}(M_1, M_2)$. We start with an open subvariety denoted by $\mathcal{M}^{N,K}(M_1, M_2)$, and it is defined by the open condition ³⁰ that

$$\mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle \operatorname{Im}(\tilde{I}) = \mathbb{C}^{M_2}.$$

 $^{^{30}\}mathrm{This}$ means that the set of point which satisfy this condition is open.

Note that under this condition, two gauge nodes satisfy their own stability condition when considered as individual ADHM quivers. So we have a projection

$$p: \mathring{\mathcal{M}}^{N,K}(M_1, M_2) \to \mathcal{M}^N(M_1) \times \mathcal{M}(M_2, K)$$
(C.10)

and in fact the projection map is a vector bundle where fibers are maps between \mathcal{V}_1 and \mathcal{V}_2 , where \mathcal{V}_1 and \mathcal{V}_2 are universal bundles on $\mathcal{M}^N(M_1)$ and $\mathcal{M}(M_2, K)$ respectively. In other words, we have an isomorphism

$$\mathcal{\dot{M}}^{N,K}(M_1, M_2) = \operatorname{Hom}(\mathcal{V}_1, \mathcal{V}_2)$$
(C.11)

Let us also consider the closed subvariety denoted by $\mathcal{M}_{\tilde{J}=0}^{N,K}(M_1, M_2)$, and it is defined by the closed condition ³¹ that

$$\tilde{J} = 0.$$

The same argument of Lemma 3 shows that $\mathcal{M}_{\tilde{J}=0}^{N,K}(M_1, M_2)$ is a locally trivial fibration over $\mathcal{M}^N(M_1)$ with fibers isomorphic to $\operatorname{Quot}^{M_2}(\mathcal{O}_{\mathbb{C}^2}^{K+M_1})$.

Proposition 3. Assume that $\xi > 0$ and K = 1, then $\mathcal{M}_{\tilde{J}=0}^{N,1}(M_1, M_2)$ is the closure of $\mathcal{M}^{N,1}(M_1, M_2)$.

Proof. If K = 1, then the stability condition $\mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle \operatorname{Im}(\tilde{I}) = \mathbb{C}^{M_2}$ together with the equation $[\tilde{X}, \tilde{Y}] + \tilde{I}\tilde{J} = 0$ implies that $\tilde{J} = 0$, so $\mathcal{M}^{N,1}(M_1, M_2)$ is a subvariety of $\mathcal{M}_{\tilde{J}=0}^{N,1}(M_1, M_2)$. Moreover, we prove in the appendix that $\mathcal{M}_{\tilde{J}=0}^{N,1}(M_1, M_2)$ is irreducible (see Proposition 10), thus $\mathcal{M}_{\tilde{J}=0}^{N,1}(M_1, M_2)$ is the closure of $\mathcal{M}^{N,1}$.

C.4. Torus fixed points

We first introduce some notations. Define the action of tori $\mathbf{T}_i, i \in \{1, 2, 3\}$ as follows:

$$a \in \mathbf{T}_{1} : (X, Y) \mapsto (aX, a^{-1}Y),$$

$$b \in \mathbf{T}_{2} : (\tilde{X}, \tilde{Y}) \mapsto (b\tilde{X}, b^{-1}\tilde{Y}),$$

$$c \in \mathbf{T}_{3} : S \mapsto cS,$$

(C.12)

and denote by $\mathbf{T}_I, I \subset \{1, 2, 3\}$ the torus $\prod_{i \in I} \mathbf{T}_i$.

The next proposition follows from the same argument as the proof of [126, Proposition 2.3.1]

Proposition 4. Assume that $\xi > 0$, then the \mathbf{T}_3 fixed points have a disjoint union decomposition:

$$\mathcal{M}^{N,K}(M_1, M_2)^{\mathbf{T}_3} = \bigsqcup_{M_2^{(1)} + M_2^{(2)} = M_2} \mathcal{M}^{N,0}(M_1, M_2^{(1)}) \times \mathcal{M}^{0,K}(0, M_2^{(2)}).$$
(C.13)

 $^{^{31}\}mathrm{This}$ means that the set of point which satisfy this condition is closed.

Remark 3. As we have mentioned in the Proposition 1, $\mathcal{M}^{N,0}(M_1, M_2^{(1)})$ is a locally trivial fibration over the ADHM moduli space $\mathcal{M}^N(M_1)$ with fibers isomorphic to $\operatorname{Quot}^{M_2^{(1)}}(\mathcal{O}_{\mathbb{C}^2}^{M_1})$. And by definition, $\mathcal{M}_i^{0,K}(0, M_2^{(2)})$ is nothing but the ADHM moduli space $\mathcal{M}(M_2^{(2)}, K)$.

Corollary 1. Assume that $\xi \neq 0$ and N = K = 1, then $\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{\{1,2,3\}}}$ is proper (compact).

Proof. We only prove for the case $\xi > 0$, the case $\xi < 0$ is similar. It is enough to show that:

- (1) $\mathcal{M}(M_1, 1)^{\mathbf{T}_1}$ is proper;
- (2) $\operatorname{Quot}^{M_2^{(1)}}(\mathcal{O}_{\mathbb{C}^2}^{M_1})^{\mathbf{T}_2}$ is proper.

Note that if we can show (1) then the same argument shows that $\mathcal{M}(M_2^{(2)}, 1)^{\mathbf{T}_2}$ is proper as well.

Notice that (1) is a special case of (2), since $\mathcal{M}(M_1, 1) = \operatorname{Quot}^{M_1}(\mathcal{O}_{\mathbb{C}^2})$ and both of the torus act on \mathbb{C}^2 by $(x, y) \mapsto (rx, r^{-1}y)$, where (x, y) is a coordinate system on \mathbb{C}^2 . Let us omit the subscripts and show that $\operatorname{Quot}^M(\mathcal{O}_{\mathbb{C}^2}^L)^{\mathbf{T}}$ is proper. Recall that the Hilbert-Chow map ${}^{32} h : \operatorname{Quot}^M(\mathcal{O}_{\mathbb{C}^2}^L)^{\mathbf{T}} \to \operatorname{Sym}^M\mathbb{C}^2$ is proper and **T**-equivariant, so $\operatorname{Quot}^M(\mathcal{O}_{\mathbb{C}^2}^L)^{\mathbf{T}} \subset h^{-1}((\operatorname{Sym}^M\mathbb{C}^2)^{\mathbf{T}})$. Since $(\operatorname{Sym}^M\mathbb{C}^2)^{\mathbf{T}}$ consists of a single point 0, $\operatorname{Quot}^M(\mathcal{O}_{\mathbb{C}^2}^L)^{\mathbf{T}}$ is a closed subset of $h^{-1}(0)$, which is proper. This concludes the proof. \Box

Remark 4. For general N, K, the fixed point set $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbf{T}_{\{1,2,3\}}}$ can be noncompact. Note that $\mathcal{M}^N(M_1)^{\mathbf{T}_1}$ is the moduli space of \mathbf{T}_1 -weighted representations of ADHM quiver of gauge rank M_1 and flavour rank N, so the gauge node \mathbb{C}^{M_1} decomposes into \mathbf{T}_1 -weight space, and flavour node \mathbb{C}^N has \mathbf{T}_1 -weight zero, and X, Y, I, Jare \mathbf{T}_1 -equivariant. A connected component of $\mathcal{M}^N(M_1)^{\mathbf{T}_1}$ corresponds to a weight decomposition

$$\mathbb{C}^{M_1} = \bigoplus_{i=0}^{M_1-1} \mathbb{C} \cdot e_i,$$

where e_i has weight i, X maps e_i to $\lambda_i e_{i-1}$ and Y maps e_i to $\mu_i e_{i+1}$ (we set $\lambda_i = 0$ if $i \notin \{1, \dots, M_1 - 1\}$ and set $\mu_i = 0$ if $i \notin \{0, \dots, M_1 - 2\}$), and I, J map between $\mathbb{C} \cdot e_0$ and the framing node \mathbb{C}^N . It can be easily deduced from the stability condition that

$$\mu_i \neq 0, \ \forall i \in \{0, \cdots, M_1 - 2\}.$$

It follows from the equation [X, Y] + IJ = 0 that

$$\lambda_i \mu_{i-1} = \lambda_{i+1} \mu_i, \ \forall i \ge 1.$$

Combine the above two facts with the equation $\lambda_{M_1} = 0$, and we deduce that $\lambda_i = 0, \forall i$. In the moduli space of quiver representations, we can scale each node, so we can assume

³²For the definition of Hilbert-Chow map, see for example [127, Section 7.1].

that $\mu_i = 1$ for $i = 0, \dots, M_1 - 2$. The remaining moduli parameters are $I : \mathbb{C}^N \to \mathbb{C} \cdot e_0$ and $J : \mathbb{C} \cdot e_0 \to \mathbb{C}^N$ satisfying IJ = 0 and the stability condition that $I \neq 0$, and it is well-known that this moduli space is $T^* \mathbb{P}^{N-1}$, the cotangent bundle of \mathbb{P}^{N-1} . Hence we see that $\mathcal{M}^N(M_1)^{\mathbf{T}_1}$ contains a connected component isomorphic to $T^* \mathbb{P}^{N-1}$, which is non-compact.

A peculiar fact about K = 1 is that $\mathcal{M}^{0,1}(0, M_2^{(2)}) = \mathcal{M}^{0,1}_{\tilde{J}=1}(0, M_2^{(2)})$ (this is a consequence of stability condition together with the F-term equation), so we can actually turn on a larger torus action, namely the two dimensional torus acting on both \tilde{X} and \tilde{Y} . This torus will be denoted by \mathbf{T}'_2 . Similarly, if N = 1 then we have a two dimensional torus acting on X and Y.

More generally, we have a five dimensional torus $\mathbf{T}'_1 \times \mathbf{T}'_2 \times \mathbf{T}_3$ acting on $\mathcal{M}^{N,K}(M_1, M_2)$, which extends the action of $\mathbf{T}_{\{1,2,3\}}$. The action is defined by

$$(a_1, a_2) \in \mathbf{T}'_1 : (X, Y, I, J) \mapsto (a_1 X, a_2 Y, I, a_1 a_2 J), (b_1, b_2) \in \mathbf{T}'_2 : (\tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}) \mapsto (b_1 \tilde{X}, b_2 \tilde{Y}, \tilde{I}, b_1 b_2 \tilde{J}).$$
(C.14)

Denote this torus by \mathbf{T}_{edge} , then we have the following:

Proposition 5. Under the above assumptions, $\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{edge}}$ is disjoint union of points ³³, points one-to-one correspond to M_1+2 copies of Young diagrams $Y^{(1)}, Y_0^{(2)}, Y_1^{(2)}, \cdots, Y_{M_1}^{(2)}$, where $|Y^{(1)}| = M_1$ and $\sum_{i=0}^{M_1} |Y_i^{(2)}| = M_2$. Moreover, the generating function for the number of \mathbf{T}_{edge} -fixed points can be written as

$$\sum_{M_1,M_2} \# \left(\mathcal{M}^{1,1}(M_1,M_2)^{\mathbf{T}_{edge}} \right) x^{M_1} y^{M_2} = \frac{1}{\left(\frac{x}{(y;y)_{\infty}}; \frac{x}{(y;y)_{\infty}} \right)_{\infty} (y;y)_{\infty}},$$
(C.15)

where $(a; y)_{\infty} = \prod_{k=0}^{\infty} (1 - ay^k)$ is the Pochhammer symbol.

Proof. Recall that the edge torus fixed points of the ADHM quiver variety $\mathcal{M}^{1,0}(M,0)$ is disjoint union of points which are one-to-one correspond to Young diagrams Y with |Y| = M. More precisely, a fixed point $p \in \mathcal{M}^{1,0}(M,0)^{\mathbf{T}'_1}$ corresponds to a unique \mathbf{T}'_1 -equivariant quiver representation of which \mathbb{C}^M decomposes into \mathbf{T}'_1 weight spaces

$$\mathbb{C}^M = \bigoplus_{(i,j)\in Y} \mathbb{C} \cdot e_{i,j},$$

where $(i, j) \in Y$ is the (i, j) box in the Young diagram Y and $e_{i,j}$ is a vector of weight (-i, -j) under the \mathbf{T}'_1 action. We set $e_{i,j} = 0$ if $(i, j) \notin Y$, then X maps $e_{i,j}$ to $e_{i+1,j}$ and Y maps $e_{i,j}$ to $e_{i,j+1}$ and I maps the flavour vector to $e_{0,0}$ and all other arrows are zero.

³³In the scheme-theoretical sense, i.e. there is no infinitesimal deformation of those points inside the T_{edge} -fixed points loci.

Back to our situation, we have

$$\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{edge}} = \bigsqcup_{M_2^{(1)} + M_2^{(2)} = M_2} \mathcal{M}^{1,0}(M_1, M_2^{(1)})^{\mathbf{T}_1' \times \mathbf{T}_2'} \times \mathcal{M}^{0,1}(0, M_2^{(2)})^{\mathbf{T}_2'}.$$

From the discussion above, we see that $\mathcal{M}^{0,1}(0, M_2^{(2)})^{\mathbf{T}'_2}$ is disjoint union of points labelled by Young diagrams with $M_2^{(2)}$ boxes, and there is a map $\mathcal{M}^{1,0}(M_1, M_2^{(1)})^{\mathbf{T}'_1 \times \mathbf{T}'_2} \to \mathcal{M}^{1,0}(M_1, 0)^{\mathbf{T}'_1}$ and the latter is also disjoint union of points labelled by Young diagrams with M_1 boxes. Let us fix Young diagrams $Y^{(1)}$ and $Y_0^{(2)}$ such that $|Y^{(1)}| = M_1$ and $|Y_0^{(2)}| = M_2^{(2)}$. Then $Y^{(1)}$ corresponds to a point p in $\mathcal{M}^{1,0}(M_1, 0)^{\mathbf{T}'_1}$, and a point in the preimage of p in $\mathcal{M}^{1,0}(M_1, M_2^{(1)})^{\mathbf{T}'_1 \times \mathbf{T}'_2}$ corresponds to a $\mathbf{T}'_1 \times \mathbf{T}'_2$ -equivariant quiver representation of which \mathbb{C}^M decomposes into $\mathbf{T}'_1 \times \mathbf{T}'_2$ weight spaces

$$\mathbb{C}^M = \bigoplus_{(i,j)\in Y^{(1)}} \mathbb{C} \cdot e_{i,j},$$

where $(i, j) \in Y$ is the (i, j) box in the Young diagram $Y^{(1)}$ and $e_{i,j}$ is a vector of weight (-i, -j) under the \mathbf{T}'_1 action and of weight zero under the \mathbf{T}'_2 action. Next we decompose $\mathbb{C}^{M_2^{(1)}}$ into \mathbf{T}'_1 weight spaces:

$$\mathbb{C}^{M_2^{(1)}} = \bigoplus_{(i,j)\in Y^{(1)}} V_{i,j},$$

where $V_{i,j}$ has \mathbf{T}'_1 weight (-i, -j). By equivariance, S maps $e_{i,j}$ to $V_{i,j}$ and \tilde{X}, \tilde{Y} maps $V_{i,j}$ to itself, and by stability, we have $\mathbb{C}\langle \tilde{X}, \tilde{Y} \rangle S(e_{i,j}) = V_{i,j}$, this shows that the restriction of \tilde{X}, \tilde{Y} to $V_{i,j}$ is a stable ADHM quiver representation, where the flavour vector is $e_{i,j}$. In other word, we show that the preimage of p in $\mathcal{M}^{1,0}(M_1, M_2^{(1)})^{\mathbf{T}'_1 \times \mathbf{T}'_2}$ is isomorphic to

$$\bigsqcup_{\substack{N_1, \cdots, N_{M_1} \ge 0 \\ N_1 + \cdots + N_{M_1} = M_2^{(1)}}} \mathcal{M}^{0,1}(0, N_1)^{\mathbf{T}'_2} \times \cdots \times \mathcal{M}^{0,1}(0, N_{M_1})^{\mathbf{T}'_2}.$$

Each product factor $\mathcal{M}^{0,1}(0, N_i)^{\mathbf{T}'_2}$ is disjoint union of points which are one-to-one correspond to Young diagrams $Y_i^{(2)}$ with N_i boxes. Hence we prove that $\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{edge}}$ is disjoint union of points which are one-to-one correspond to $M_1 + 2$ copies of Young diagrams $Y^{(1)}, Y^{(2)}_0, Y^{(2)}_1, \cdots, Y^{(2)}_{M_1}$, where $|Y^{(1)}| = M_1$ and $\sum_{i=0}^{M_1} |Y_i^{(2)}| = M_2$.

To enumerate fixed points, notice that

$$\sum_{M_1} \left(\mathcal{M}^{1,0}(M_1,0)^{\mathbf{T}_{edge}} \right) x^{M_1} = \sum_{M_1} p(M_1) x^{M_1} = \prod_{i\geq 1}^{\infty} \frac{1}{1-x^i} = \frac{1}{(x;x)}, \quad (C.16)$$

where p(n) is Euler's partition function counting the number of partitions of n in to

positive integers. Thus we have

$$\begin{split} &\sum_{M_1,M_2} \left(\mathcal{M}^{1,1}(M_1,M_2)^{\mathbf{T}'_1} \right) x^{M_1} y^{M_2} \\ &= \left(\sum_{M_1} \left(\mathcal{M}^{1,0}(M_1,0)^{\mathbf{T}'_1} \right) \sum_{N_1,\cdots,N_{M_1} \ge 0} \prod_{i=1}^{M_1} \left(\mathcal{M}^{0,1}(0,N_i)^{\mathbf{T}'_2} \right) x^{M_1} y^{N_1 + \dots + N_{M_1}} \right) \\ &\times \left(\sum_{M_2^{(2)}} \left(\mathcal{M}^{0,1}(0,M_2^{(2)})^{\mathbf{T}'_2} \right) y^{M_2^{(2)}} \right) \\ &= \left(\sum_{M_1} p(M_1) \left(\frac{x}{(y;y)} \right)^{M_1} \right) \times \left(\sum_{M_2^{(2)}} p(M_2^{(2)}) y^{M_2^{(2)}} \right) \\ &= \frac{1}{\left(\frac{x}{(y;y)_{\infty}}; \frac{x}{(y;y)_{\infty}} \right)_{\infty} (y;y)_{\infty}}. \end{split}$$

In the next subsection, we compute the equivariant K-theory index of $\mathcal{M}^{1,1}(M_1, M_2)$ using \mathbf{T}_{edge} localization. This is a weighted version of (C.15), where the \mathbb{C} coefficients will be replaced by $K_{\mathbf{T}_{\text{edge}}}(\text{pt})_{\text{loc}}$ coefficients, the localized ring of characters of \mathbf{T}_{edge} .

Proposition 6. Assume that $\xi \neq 0$, then $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbf{T}_{edge}}$ is proper (compact).

Proof. We only need to show that $\mathcal{M}^{N}(M_{1})^{\mathbf{T}'_{1}}$ is proper. $\mathcal{M}^{N}(M_{1})^{\mathbf{T}'_{1}}$ is the moduli of stable representations of ADHM quivers where the gauge node is a \mathbf{T}'_{1} representation, and the flavour node is a trivial \mathbf{T}'_{1} representation, and (X, Y, I, J) are equivariant maps. The stability condition $\mathbb{C}\langle X, Y, \rangle \mathrm{Im}(I) = \mathbb{C}^{M_{1}}$ implies that the \mathbf{T}'_{1} weights of $\mathbb{C}^{M_{1}}$ are positive. Since J has weight (1, 1) and \mathbb{C}^{N} has weight zero, J must be zero. This shows that $\mathcal{M}^{N}(M_{1})^{\mathbf{T}'_{1}} = \mathrm{Quot}^{M_{1}}(\mathcal{O}^{N}_{\mathbb{C}^{2}})^{\mathbf{T}'_{1}}$, and the latter is proper as we have shown in the proof of Corollary 1.

C.5. Equivariant K-theory index

Following [22], the equivariant K-theory index of the moduli space $\mathcal{M}^{1,1}(M_1, M_2)$ is defined by

$$\chi(\mathcal{M}^{1,1}(M_1, M_2), \mathcal{O}^{\mathrm{vir}}) \tag{C.17}$$

where \mathcal{O}^{vir} is the virtual structure sheaf of $\mathcal{M}^{1,1}(M_1, M_2)$. Since our moduli space $\mathcal{M}^{1,1}(M_1, M_2)$ is l.c.i by Proposition 2, the virtual structure sheaf is the actual structure sheaf, i.e. $\mathcal{O}^{\text{vir}} = \mathcal{O}$. The equivariant K-theory partition function is defined by

$$Z_{1,1} = \sum_{M_1,M_2} \chi(\mathcal{M}^{1,1}(M_1,M_2),\mathcal{O}^{\text{vir}}) x^{M_1} y^{M_2} = \sum_{M_1,M_2} \chi(\mathcal{M}^{1,1}(M_1,M_2)) x^{M_1} y^{M_2} \quad (C.18)$$

Using equivariant localization, the \mathbf{T}_{edge} -equivariant K-theory index of $\mathcal{M}^{1,1}(M_1, M_2)$ can be written in terms of \mathbf{T}_{edge} -fixed points:

$$\chi(\mathcal{M}^{1,1}(M_1, M_2), \mathcal{O}) = \chi\left(\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{edge}}, \left(S^{\bullet}(T^{vir})^*\right)|_{\mathcal{M}^{1,1}(M_1, M_2)^{\mathbf{T}_{edge}}}\right)$$
(C.19)

where T^{vir} is the virtual tangent bundle (C.9).

Let us denote the equivariant parameters of \mathbf{T}'_1 by r_1, r_2 , and denote the equivariant parameters of \mathbf{T}'_2 by s_1, s_2 , and that of \mathbf{T}_3 by t. Then the \mathbf{T}_{edge} -equivariant K-theory class of the virtual tangent bundle (C.9) can be written as

$$\mathcal{V}_1 + \mathcal{V}_1^* r_1 r_2 - \mathcal{V}_1 \mathcal{V}_1^* (1 - r_1)(1 - r_2) + \mathcal{V}_2 + \mathcal{V}_2^* s_1 s_2 - \mathcal{V}_2 \mathcal{V}_2^* (1 - s_1)(1 - s_2) + t \mathcal{V}_2 \mathcal{V}_1^*$$
(C.20)

where \mathcal{V}_1 and \mathcal{V}_2 are universal bundles on $\mathcal{M}^{1,1}(M_1, M_2)$ of rank M_1 and M_2 respectively, note that they are equivariant under the \mathbf{T}_{edge} action. At a fixed point labelled by

$$(Y^{(1)}, Y_0^{(2)}, Y_1^{(2)}, \cdots, Y_{M_1}^{(2)})$$

the fibers of \mathcal{V}_1 and \mathcal{V}_2 are representations of \mathbf{T}_{edge} , and their \mathbf{T}_{edge} characters can be described as following. Let us introduce the notation

$$w\left(Y^{(1)}\right) = \sum_{(i,j)\in Y^{(1)}} r_1^{-i} r_2^{-j}, \ w\left(Y^{(2)}_{\alpha}\right) = \sum_{(i,j)\in Y^{(2)}_{\alpha}} s_1^{-i} s_2^{-j}, \alpha \in \{0,\cdots,M_1\}.$$
(C.21)

From the description of the \mathbf{T}_{edge} fixed points, we know that $Y_1^{(2)}, \dots, Y_{M_1}^{(2)}$ are one-to-one correspond to boxes (i, j) in the diagram $Y^{(1)}$, so we use the notation $Y_{(i,j)}^{(2)}$ to indicate their relationship to $Y^{(1)}$. Then the characters of \mathcal{V}_1 and \mathcal{V}_2 are

$$\mathcal{V}_1 = w\left(Y^{(1)}\right), \ \mathcal{V}_2 = w\left(Y_0^{(2)}\right) + t^{-1} \sum_{(i,j)\in Y^{(1)}} r_1^{-i} r_2^{-j} w\left(Y_{(i,j)}^{(2)}\right)$$
(C.22)

For example, if $\Phi = \mathcal{O}$, then the \mathbf{T}_{edge} -equivariant K-theory index is

$$\chi(\mathcal{M}^{1,1}(M_1, M_2), \mathcal{O}) = \sum_{Y^{(1)}, Y_0^{(2)}, Y_{(i,j)}^{(2)}} S^{\bullet}(T^{\mathrm{vir}})^*$$
(C.23)

where T^{vir} is given by (C.20), and $S^{\bullet}(T^{\text{vir}})$ is the plethystic exponential of the virtual character T^{vir} , and the * operator maps (r_1, r_2, s_1, s_2, t) to $(r_1^{-1}, r_2^{-1}, s_1^{-1}, s_2^{-1}, t^{-1})$.

Example

Let us compute the example $M_1 = M_2 = 1$ explicitly. In this case there is only one

possible $Y^{(1)}$, which is a single box at (0,0). There are two \mathbf{T}_{edge} -fixed points:

$$Y_0^{(2)} = \Box, Y_1^{(2)} = \emptyset, \text{ and} Y_0^{(2)} = \emptyset, Y_1^{(2)} = \Box,$$
(C.24)

and the virtual tangent characters at these fixed points are

$$T^{\text{vir}} = r_1 + r_2 + s_1 + s_2 + t, \text{ and}$$

$$T^{\text{vir}} = r_1 + r_2 + s_1 + s_2 + (t-1)s_1s_2 + t^{-1},$$
(C.25)

respectively. Plug these into (C.23) we get

$$\chi(\mathcal{M}^{1,1}(1,1),\mathcal{O}) = \left(\frac{1}{(1-r_1)(1-r_2)(1-s_1)(1-s_2)} \left(\frac{1}{1-t} + \frac{1-s_1s_2}{(1-ts_1s_2)(1-t^{-1})}\right)\right)^*$$
$$= \frac{1}{(1-r_1^{-1})(1-r_2^{-1})(1-s_1^{-1})(1-s_2^{-1})} \left(\frac{1}{1-t^{-1}} + \frac{1-s_1^{-1}s_2^{-1}}{(1-t^{-1}s_1^{-1}s_2^{-1})(1-t)}\right)$$
$$= \frac{r_1r_2s_1^2s_2^2t}{(1-r_1)(1-r_2)(1-s_1)(1-s_2)(ts_1s_2-1)}$$
(C.26)

The limit $s_1 s_2 \to 1$

In the example that we computed above, we notice that when $s_1s_2 \to 1$, the contribution from the fixed point $Y_0^{(2)} = \emptyset, Y_1^{(2)} = \Box$ vanishes. This is not a coincidence:

Proposition 7. In the limit $s_1s_2 \rightarrow 1$, we have

$$\chi(\mathcal{M}^{1,1}(M_1, M_2), \mathcal{O})_{s_1 s_2 \to 1} = \chi(\mathring{\mathcal{M}}^{1,1}(M_1, M_2), \mathcal{O})_{s_1 s_2 \to 1}.$$
 (C.27)

In other words, the fixed points outside of $\mathring{\mathcal{M}}^{1,1}(M_1, M_2)$ do not contribute in this limit.

Proof. Suppose that we have a fixed point labelled by M_1+2 Young diagrams $Y^{(1)}, Y^{(2)}_0, Y^{(2)}_{(i,j)}$, where (i, j) takes value in coordinate of box in $Y^{(1)}$. We need to show that $S^{\bullet}(T^{\text{vir}})$ vanishes in the limit $s_1s_2 \to 1$ when there is an $(i, j) \in Y^{(1)}$ such that $Y^{(2)}_{(i,j)}$ is non-empty. We claim that it is enough to show that the virtual character $T^{\text{vir}}_{s_1s_2\to 0}$ has negative constant term. To prove the claim, we notice that constant term in $T^{\text{vir}}_{s_1s_2\to 0}$ comes from

$$\sum_{k} a_k (s_1 s_2)^k - \sum_{l} b_l (s_1 s_2)^k$$

in T^{vir} , where a_k, b_l are positive integers. After taking the plethystic exponential, this term becomes

$$\frac{\prod_l (1 - s_1^l s_2^l)^{b_l}}{\prod_k (1 - s_1^k s_2^k)^{a_k}}$$

and this becomes zero in the limit $s_1s_2 \to 1$ if $\sum_l b_l > \sum_k a_k$, i.e. $T_{s_1s_2\to 0}^{\text{vir}}$ has negative

constant term.

Therefore, it remains to compute the constant term in $T_{s_1s_2\to 0}^{\text{vir}}$. Let us introduce some notations. Define

$$\mathcal{V}_{2}^{(0)} = w\left(Y_{0}^{(0)}\right), \ \mathcal{V}_{2}^{(1)} = \sum_{(i,j)\in Y^{(1)}} r_{1}^{-i} r_{2}^{-j} w\left(Y_{(i,j)}^{(2)}\right), \tag{C.28}$$

so $\mathcal{V}_2 = \mathcal{V}_2^{(0)} + t^{-1} \mathcal{V}_2^{(1)}$. For every $V \in K_{\mathbb{C}^2}(\mathrm{pt}) = \mathbb{C}[u_1, u_1^{-1}, u_2, u_2^{-1}]$, define

$$T(V) = V + V^* - VV^*(1 - u_1)(1 - u_2)$$
(C.29)

For example, let \mathbb{C}^2 be \mathbf{T}'_1 then $u_1 = r_1, u_2 = r_2$, and $T(\mathcal{V}_1) = \mathcal{V}_1 + \mathcal{V}_1^* r_1 r_2 - \mathcal{V}_1 \mathcal{V}_1^* (1 - r_1)(1 - r_2)$. Using this notation, we expand (C.20) as following:

$$T^{\text{vir}} = T(\mathcal{V}_1) + T(\mathcal{V}_2^{(0)}) + t^{-1}\mathcal{V}_2^{(1)} + ts_1s_2\mathcal{V}_2^{(1)} - \mathcal{V}_2^{(1)}\mathcal{V}_2^{(1)*}(1-s_1)(1-s_2) - t\mathcal{V}_2^{(0)}\mathcal{V}_2^{(1)*}(1-s_1)(1-s_2) - t^{-1}\mathcal{V}_2^{(0)*}\mathcal{V}_2^{(1)}(1-s_1)(1-s_2) + t\mathcal{V}_2^{(0)}\mathcal{V}_1^* + \mathcal{V}_2^{(1)}\mathcal{V}_1^*.$$
(C.30)

Recall that if V is the weight space of a Young diagram Y, then [128, Lemma 6]

$$T(V) = \sum_{\Box \in Y} u_1^{-l(\Box)} u_2^{a(\Box)+1} + \sum_{\Box \in Y} u_1^{l(\Box)+1} u_2^{-a(\Box)},$$
(C.31)

where $a(\Box)$ and $l(\Box)$ are arm-length and leg-length respectively:

$$a(\Box) = \#\{j' > j | (i,j') \in Y\}, \ l(\Box) = \#\{i' > i | (i',j) \in Y\}.$$
 (C.32)

In particular, we see that constant terms in $T(\mathcal{V}_1)_{s_1s_2\to 0}$ and $T(\mathcal{V}_2^{(0)})_{s_1s_2\to 0}$ are zero. Next, we simply drop all the other terms in (C.30) which involve t or t^{-1} because they can not be constant in the limit $s_1s_2 \to 0$. Therefore, we only need to compute the constant term in

$$\mathcal{V}_2^{(1)}\mathcal{V}_1^* - \mathcal{V}_2^{(1)}\mathcal{V}_2^{(1)*}(1-s_1)(1-s_2) \tag{C.33}$$

in the limit $s_1s_2 \to 0$. Let us take \mathbf{T}'_1 -invariant first, this amounts to take those terms in (C.33) which do not involve r_1 or r_2 , and the result is

$$\sum_{(i,j)\in Y^{(1)}} w\left(Y_{(i,j)}^{(2)}\right) - \sum_{(i,j)\in Y^{(1)}} w\left(Y_{(i,j)}^{(2)}\right) w\left(Y_{(i,j)}^{(2)}\right)^* (1-s_1)(1-s_2).$$
(C.34)

Notice that this can be rewritten as

$$\sum_{(i,j)\in Y^{(1)}} T\left(w\left(Y_{(i,j)}^{(2)}\right)\right) - \sum_{(i,j)\in Y^{(1)}} w\left(Y_{(i,j)}^{(2)}\right)^* s_1 s_2,\tag{C.35}$$

and each individual $T\left(w\left(Y_{(i,j)}^{(2)}\right)\right)$ has zero constant term in the limit $s_1s_2 \to 0$, as can

be seen from (C.31). Thus the constant term of $T_{s_1s_2\to 1}^{\text{vir}}$ equals to the constant term of

$$\sum_{(i,j)\in Y^{(1)}} w \left(Y^{(2)}_{(i,j)}\right)^*_{s_1s_2\to 1},\tag{C.36}$$

which is minus the number of non-empty Young diagrams in $Y_{(i,j)}^{(2)}$. In particular, it is negative if one of $Y_{(i,j)}^{(2)}$ is non-empty. This concludes the proof of proposition.

The index of the open subset $\mathring{\mathcal{M}}^{1,1}(M_1, M_2)$

We have seen in (C.11) that $\mathcal{\dot{M}}^{1,1}(M_1, M_2)$ is the vector bundle Hom $(\mathcal{V}_1, \mathcal{V}_2)$, so its ring of functions is

$$\bigoplus_{k\geq 0} t^k S^k(\mathcal{V}_2 \otimes \mathcal{V}_1^*) \tag{C.37}$$

as a \mathbf{T}_{edge} -equivariant sheaf on $\mathcal{M}(M_1, 1) \times \mathcal{M}(M_2, 1)$. It is easy to see that

$$S^{k}(\mathcal{V}_{2} \otimes \mathcal{V}_{1}^{*}) = \bigoplus_{|\underline{\lambda}|=k} S^{\underline{\lambda}}(\mathcal{V}_{2}) \otimes S^{\underline{\lambda}}(\mathcal{V}_{1}^{*}),$$
(C.38)

where $\underline{\lambda}$ is a partition of k (given by a Young diagram) and $S^{\underline{\lambda}}(\mathcal{V}_2)$ is the irreducible representation of $GL(\mathcal{V}_2)$ defined by

$$\operatorname{Hom}_{S_k}(R_{\underline{\lambda}}, \mathcal{V}_2^{\otimes k}). \tag{C.39}$$

Here $R_{\underline{\lambda}}$ is the irreducible representation of permutation group S_k with the Young diagram $\underline{\lambda}$. For example,

$$S^{(k)}(\mathcal{V}_2) = S^k(\mathcal{V}_2), \ S^{(1,1,\dots,1)}(\mathcal{V}_2) = \wedge^k(\mathcal{V}_2).$$
 (C.40)

Using the decomposition (C.38), we can write the \mathbf{T}_{edge} -equivariant K-theory index of $\mathcal{\dot{M}}^{1,1}(M_1, M_2)$ as

$$\chi(\mathring{\mathcal{M}}^{1,1}(M_1, M_2), \mathcal{O}) = \sum_{\underline{\lambda}} t^{|\underline{\lambda}|} \chi(\mathcal{M}(M_1, 1), S^{\underline{\lambda}}(\mathcal{V}_1^*)) \chi(\mathcal{M}(M_2, 1), S^{\underline{\lambda}}(\mathcal{V}_2)), \quad (C.41)$$

where the sum is for all Young diagrams $\underline{\lambda}$. Note that

$$\chi(\mathcal{M}(M_1, 1), S^{\underline{\lambda}}(\mathcal{V}_1^*)) = r_1^{-2M_1} r_2^{-2M_1} \chi(\mathcal{M}(M_1, 1), S^{\underline{\lambda}}(\mathcal{V}_1))^*.$$
(C.42)

Therefore, we have

$$\chi(\mathring{\mathcal{M}}^{1,1}(M_1, M_2), \mathcal{O}) = r_1^{-2M_1} r_2^{-2M_1} \sum_{\underline{\lambda}} t^{|\underline{\lambda}|} \chi(\mathcal{M}(M_1, 1), S^{\underline{\lambda}}(\mathcal{V}_1))^* \chi(\mathcal{M}(M_2, 1), S^{\underline{\lambda}}(\mathcal{V}_2)).$$
(C.43)

Recall that the equivariant K-theory partition function of $\mathcal{M}^{1,1}$ is defined as

$$Z_{1,1} = \sum_{M_1,M_2} \chi(\mathcal{M}^{1,1}(M_1,M_2),\mathcal{O}) x^{M_1} y^{M_2}.$$
 (C.44)

Combine (C.43) with Propistion 7, we then have

Proposition 8. In the limit $r_1r_2 \rightarrow 1, s_1s_2 \rightarrow 1$, the equivariant K-theory partition function of $\mathcal{M}^{1,1}$ factorizes as

$$Z_{1,1} = \sum_{\underline{\lambda}} t^{|\underline{\lambda}|} \mathcal{Z}_{1}^{\underline{\lambda}}(r_{1}, r_{2}, x)^{*}_{r_{1}r_{2} \to 1} \mathcal{Z}_{1}^{\underline{\lambda}}(s_{1}, s_{2}, y)_{s_{1}s_{2} \to 1}$$
(C.45)

where $\mathcal{Z}_1^{\underline{\lambda}}(r_1, r_2, x)$ is the equivariant K-theory partition function for ADHM quiver:

$$\mathcal{Z}_{1}^{\underline{\lambda}}(r_{1}, r_{2}, x) = \sum_{M} \chi(\mathcal{M}(M, 1), S^{\underline{\lambda}}(\mathcal{V})) x^{M}.$$
 (C.46)

Remark 5. Consider the ADHM quiver with framing rank one and gauge rank M_1 , and an extra framing node with rank M_2 , see Figure 3, then it is easy to see that the corresponding quiver variety is the universal bundle \mathcal{V} on the ADHM quiver without extra framing. Note that there is an action of $\mathbf{T}'_1 \times \mathbf{T}_3$ on the quiver variety and the equivariant K-theory index for this variety is

$$\sum_{\underline{\lambda}} \chi(\mathcal{M}(M,1), S^{\underline{\lambda}}(\mathcal{V})) t^{|\underline{\lambda}|}.$$
(C.47)

This is exactly the coefficient in the summation (C.46).

For general N, K, we know that $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbf{T}_{edge}}$ is compact, so \mathbf{T}_{edge} -equivariant K-theory index of $\mathcal{M}^{N,K}(M_1, M_2)$ is well-defined. To compute this index, we turn on the maximal torus of the flavour symmtry, i.e. consider the diagonal action of $\mathbf{T}_{\text{framing}} = \mathbb{C}^{\times N} \times \mathbb{C}^{\times N}$ on the framing vector space, and then take the $\mathbf{T}_{edge} \times \mathbf{T}_{\text{framing}}$ fixed points of $\mathcal{M}^{N,K}(M_1, M_2)$ and this will be disjoint union of points (scheme-theoretically), one could follow the same argument in the computation of N = K = 1 case and compute the $\mathbf{T}_{edge} \times \mathbf{T}_{\text{framing}}$ -equivariant K-theory index of $\mathcal{M}^{N,K}(M_1, M_2)$ and then take $\mathbf{T}_{\text{framing}}$ equivariant parameters to 1 and this would be the \mathbf{T}_{edge} -equivariant K-theory index of $\mathcal{M}^{N,K}(M_1, M_2)$.

C.6. Connection to the 5d Chern-Simons algebra of operators

Notice that in the $t \to 0, r_1r_2 \to 1, s_1s_2 \to 1$ limit, the equivariant K-theory partition function of $\mathcal{M}^{1,1}$ equals

$$\sum_{M,N} \chi(\mathcal{M}(M,1),\mathcal{O})\chi(\mathcal{M}(N,1),\mathcal{O})x^M y^N,$$
(C.48)

i.e. it is a product of equivariant K-theory partition functions of ADHM quivers:

$$\lim_{t \to 0} Z_{1,1} = Z_{\text{ADHM}}(r_1, r_2, x)_{r_1 r_2 \to 1} Z_{\text{ADHM}}(s_1, s_2, y)_{s_1 s_2 \to 1}.$$
 (C.49)

Since the equivariant K-theory of ADHM quiver varieties admits a structure of Verma modules of the affine quantum group $U_q(\hat{\mathfrak{gl}}_1)$, we see that t = 0 part of $Z_{1,1}$ is a character of a Verma module for $U_q(\hat{\mathfrak{gl}}_1) \otimes U_q(\hat{\mathfrak{gl}}_1)$. After passing from K-theory to cohomology, part of $Z_{1,1}$ is a character of a Verma module for the affine Yangian of $\mathfrak{gl}_1 \oplus \mathfrak{gl}_1$, which is exactly the algebra of gauge invariant local observables of the 5d holomorphic Chern-Simons theory with gauge group $U(1) \times U(1)$.

D. Connectedness of $\mathcal{M}^{N,K}(M_1, M_2)$

In this appendix, we show that

Proposition 9. $\mathcal{M}^{N,K}(M_1, M_2)$ is connected.

First of all, if $\xi = 0$, then the moduli space is the affine quotient of the solutions to ADHM equations (4.8) by $\operatorname{GL}_N \times \operatorname{GL}_K$. Consider the action of \mathbb{C}^{\times} by

$$(X, Y, I, J, \tilde{X}, \tilde{Y}, \tilde{I}, \tilde{J}, S) \mapsto (rX, rY, rI, rJ, r\tilde{X}, r\tilde{Y}, r\tilde{I}, r\tilde{J}, rS),$$
(D.1)

then it contracts the $\mathcal{M}_{\xi=0}^{N,K}(M_1, M_2)$ to a single point $(0, 0, \dots, 0)$, thus $\mathcal{M}_{\xi=0}^{N,K}(M_1, M_2)$ is connected.

Next, we show that $\mathcal{M}^{N,K}(M_1, M_2)$ is connected when $\xi > 0$, and the $\xi < 0$ case is similar. Consider the \mathbb{C}^{\times} action (D.1), its fixed points in $\mathcal{M}^{N,K}(M_1, M_2)$ correspondes to quiver representations such that \mathbb{C}^{M_1} and \mathbb{C}^{M_2} are \mathbb{Z} -graded and all arrows in the quiver increase the grade by one. From the stability condition, $\mathbb{C}\langle \tilde{X}, \tilde{Y}\rangle(\operatorname{Im}(\tilde{I}) + \operatorname{Im}(S)) = \mathbb{C}^{M_2}$ we deduce that \mathbb{C}^{M_2} must be positively graded. Since the framing vector space has grade zero, we conclude that $\tilde{J} = 0$, i.e. $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbb{C}^{\times}} \subset \mathcal{M}^{N,K}_{\tilde{J}=0}(M_1, M_2)$. Note that the natural projection $q : \mathcal{M}^{N,K}(M_1, M_2) \to \mathcal{M}^{N,K}_{\xi=0}(M_1, M_2)$ defined by

Note that the natural projection $q : \mathcal{M}^{N,K}(M_1, M_2) \to \mathcal{M}_{\xi=0}^{N,K}(M_1, M_2)$ defined by semisimplification is projective and equivariant under the \mathbb{C}^{\times} action. We claim that $\forall x \in \mathcal{M}^{N,K}(M_1, M_2)$, its limit under the \mathbb{C}^{\times} action $\lim_{r\to 0} r(x)$ exists. This follows from the fact that $\lim_{r\to 0} r(q(x))$ exists and the valuative criterion for a proper morphism. Since $\forall s \in \mathbb{C}^{\times}$, we have

$$s\left(\lim_{r \to 0} r(x)\right) = \lim_{r \to 0} sr(x) = \lim_{r' \to 0} r'(x),$$
(D.2)

we see that the limit $\lim_{r\to 0} r(x) \in \mathcal{M}^{N,K}(M_1, M_2)^{\mathbb{C}^{\times}}$, thus every point in $\mathcal{M}^{N,K}(M_1, M_2)$ is connected to some point in $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbb{C}^{\times}}$. As there is an inclusion $\mathcal{M}^{N,K}(M_1, M_2)^{\mathbb{C}^{\times}} \subset \mathcal{M}^{N,K}_{\tilde{J}=0}(M_1, M_2)$, we only need to show that $\mathcal{M}^{N,K}_{\tilde{J}=0}(M_1, M_2)$ is connected. In fact, we prove a stronger result **Proposition 10.** Assume that $\xi > 0$, the $\mathcal{M}_{\tilde{J}=0}^{N,K}(M_1, M_2)$ is irreducible of dimension $2NM_1 + M_2(M_1 + K + 1)$.

Proof. The same argument in Lemma 3 shows that there is locally trivial fibration

$$\pi: \mathcal{M}_{\tilde{J}=0}^{N,K}(M_1,M_2) \to \mathcal{M}^N(M_1)$$

with fibers isomorphic to $\operatorname{Quot}^{M_2}(\mathcal{O}_{\mathbb{C}^2}^{M_1+K})$. It is known that the Nakajima quiver variety $\mathcal{M}^N(M_1)$ is irreducible³⁴ of dimension $2NM_1$, so we only need to show that $\operatorname{Quot}^{M_2}(\mathcal{O}_{\mathbb{C}^2}^{M_1+K})$ is irreducible of dimension $M_2(M_1+K+1)$, and this is proven in the next lemma.

Lemma 4. Let S be a smooth connected algebraic surface and V is a vector bundle of rank N on S, then the Quot scheme $\text{Quot}^{M}(V)$ is irreducible of dimension (N+1)M.

Proof. Consider the Hilbert-Chow map $h : \operatorname{Quot}^{M}(\mathcal{V}) \to \operatorname{Sym}^{M}\mathcal{S}$. Its fiber over a point

$$(\underbrace{x_1,\cdots,x_1}_{L_1},\underbrace{x_2,\cdots,x_2}_{L_2},\cdots,\underbrace{x_n,\cdots,x_n}_{L_n}) \in \operatorname{Sym}^M \mathcal{S}, \ M = \sum_{i=1}^n L_i$$
(D.3)

is isomorphic to

$$\prod_{i=1}^{n} \operatorname{Quot}^{L_{i}}(\mathbb{C}\llbracket x, y \rrbracket^{N})$$
(D.4)

where $\operatorname{Quot}^{L_i}(\mathbb{C}\llbracket x, y \rrbracket^N)$ is the formal Quot scheme defined by the moduli space of quotient module of $\mathbb{C}\llbracket x, y \rrbracket^N$ of \mathbb{C} -dimension L_i on the ring $\mathbb{C}\llbracket x, y \rrbracket$. We claim that $\operatorname{Quot}^{L_i}(\mathbb{C}\llbracket x, y \rrbracket^N)$ is irreducible. The irreducibility of $\operatorname{Quot}^M(\mathcal{V})$ follows from this claim and the irreducibility of $\operatorname{Sym}^M \mathcal{S}$.

To prove this claim, we notice that $\operatorname{Quot}^{L_i}(\mathbb{C}[\![x,y]\!]^N)$ has the form $U/\operatorname{GL}_{L_i}$ where U is an open subvariety of the following variety

$$\{X, Y \in \mathfrak{gl}_{L_i} | [X, Y] = 0, \text{ X and Y are nilpotent} \} \times \operatorname{Hom}(\mathbb{C}^N, \mathbb{C}^{L_i}).$$
(D.5)

The first factor in (D.5) is irreducible [130], and the second factor is obviously irreducible, so U is an irreducible variety, thus $\operatorname{Quot}^{L_i}(\mathbb{C}[\![x, y]\!]^N)$ is irreducible.

Finally, the dimension of $\operatorname{Quot}^{M}(\mathcal{V})$ can be computed using the open locus $h^{-1}(V)$, where $V \subset \operatorname{Sym}^{M} \mathcal{S}$ is the open subset of points (x_{1}, \dots, x_{M}) where x_{i} are distinct. Restrict to $h^{-1}(V)$, h is a locally trivial fibration with fibers isomorphic to product of M copies of \mathbb{P}^{N-1} , so

dim Quot^M(
$$\mathcal{V}$$
) = dim $h^{-1}(V) = (N-1)M + 2M = (N+1)M$ (D.6)

³⁴The Kirwan map $K_{\mathrm{GL}_{M_1}}(\mathrm{pt}) \to K(\mathcal{M}^N(M_1))$ is surjective by [129], so every K-theory class on $\mathcal{M}^N(M_1)$ must have constant rank, then it follows that $\mathcal{M}^N(M_1)$ is connected thus it is irreducible.

References

- J. Halverson and D. R. Morrison, "The landscape of M-theory compactifications on seven-manifolds with G₂ holonomy," *JHEP* 04 (2015) 047, arXiv:1412.4123 [hep-th].
- J. Halverson and D. R. Morrison, "On gauge enhancement and singular limits in G₂ compactifications of M-theory," JHEP 04 (2016) 100, arXiv:1507.05965
 [hep-th].
- [3] A. P. Braun, "Tops as building blocks for G₂ manifolds," *JHEP* 10 (2017) 083, arXiv:1602.03521 [hep-th].
- [4] T. C. da C. Guio, H. Jockers, A. Klemm, and H.-Y. Yeh, "Effective Action from M-Theory on Twisted Connected Sum G₂-Manifolds," *Commun. Math. Phys.* 359 no. 2, (2018) 535-601, arXiv:1702.05435 [hep-th].
- [5] A. P. Braun and M. Del Zotto, "Mirror Symmetry for G₂-Manifolds: Twisted Connected Sums and Dual Tops," JHEP 05 (2017) 080, arXiv:1701.05202 [hep-th].
- [6] A. P. Braun and S. Schäfer-Nameki, "Compact, Singular G₂-Holonomy Manifolds and M/Heterotic/F-Theory Duality," JHEP 04 (2018) 126, arXiv:1708.07215 [hep-th].
- [7] A. P. Braun and M. Del Zotto, "Towards Generalized Mirror Symmetry for Twisted Connected Sum G₂ Manifolds," JHEP 03 (2018) 082, arXiv:1712.06571 [hep-th].
- [8] A. P. Braun, M. Del Zotto, J. Halverson, M. Larfors, D. R. Morrison, and S. Schäfer-Nameki, "Infinitely many M2-instanton corrections to M-theory on G₂-manifolds," *JHEP* 09 (2018) 077, arXiv:1803.02343 [hep-th].
- B. S. Acharya, A. P. Braun, E. E. Svanes, and R. Valandro, "Counting associatives in compact G₂ orbifolds," JHEP 03 (2019) 138, arXiv:1812.04008 [hep-th].
- [10] A. P. Braun, S. Cizel, M. Hübner, and S. Schäfer-Nameki, "Higgs bundles for M-theory on G₂-manifolds," JHEP 03 (2019) 199, arXiv:1812.06072 [hep-th].
- [11] A. P. Braun, S. Majumder, and A. Otto, "On Mirror Maps for Manifolds of Exceptional Holonomy," JHEP 10 (2019) 204, arXiv:1905.01474 [hep-th].

- [12] A. P. Braun, "M-Theory and Orientifolds," JHEP 09 (2020) 065, arXiv:1912.06072 [hep-th].
- [13] R. Barbosa, M. Cvetič, J. J. Heckman, C. Lawrie, E. Torres, and G. Zoccarato, "T-branes and G₂ backgrounds," *Phys. Rev. D* 101 no. 2, (2020) 026015, arXiv:1906.02212 [hep-th].
- B. S. Acharya, L. Foscolo, M. Najjar, and E. E. Svanes, "New G₂-conifolds in M-theory and their field theory interpretation," *JHEP* 05 (2021) 250, arXiv:2011.06998 [hep-th].
- [15] B. S. Acharya, A. Kinsella, and D. R. Morrison, "Non-Perturbative Heterotic Duals of M-Theory on G₂ Orbifolds," arXiv:2106.03886 [hep-th].
- [16] A. Kovalev, "Twisted connected sums and special Riemannian holonomy," J. Reine Angew. Math. 565 (2003) 125–160.
- [17] A. Kovalev and J. Nordström, "Asymptotically cylindrical 7-manifolds of holonomy g2 with applications to compact irreducible g2-manifolds," Annals of Global Analysis and Geometry 38 no. 3, (2010) 221-257, arXiv:0907.0497
 [math.DG].
- [18] A. Corti, M. Haskins, J. Nordström, and T. Pacini, "Asymptotically cylindrical Calabi-Yau 3-folds from weak Fano 3-folds," *Geom. Topol.* 17 no. 4, (2013) 1955–2059.
- [19] A. Corti, M. Haskins, J. Nordström, and T. Pacini, "G₂-manifolds and associative submanifolds via semi-Fano 3-folds," *Duke Math. J.* 164 no. 10, (2015) 1971–2092, arXiv:1207.4470 [math.DG].
- [20] L. Foscolo, M. Haskins, and J. Nordström, "Complete non-compact G2-manifolds from asymptotically conical Calabi-Yau 3-folds," arXiv:1709.04904 [math.DG].
- [21] L. Foscolo, M. Haskins, and J. Nordström, "Infinitely many new families of complete cohomogeneity one G_2-manifolds: G_2 analogues of the Taub-NUT and Eguchi-Hanson spaces," arXiv:1805.02612 [math.DG].
- [22] N. Nekrasov and A. Okounkov, "Membranes and Sheaves," arXiv:1404.2323 [math.AG].
- [23] K. Costello and S. Li, "Twisted supergravity and its quantization," arXiv:1606.00365 [hep-th].
- [24] K. Costello, "M-theory in the Omega-background and 5-dimensional non-commutative gauge theory," arXiv:1610.04144 [hep-th].

- [25] R. Eager and F. Hahner, "Maximally twisted eleven-dimensional supergravity," arXiv:2106.15640 [hep-th].
- [26] I. Saberi and B. R. Williams, "Twisting pure spinor superfields, with applications to supergravity," arXiv:2106.15639 [math-ph].
- [27] A. Iqbal, N. Nekrasov, A. Okounkov, and C. Vafa, "Quantum foam and topological strings," JHEP 04 (2008) 011, arXiv:hep-th/0312022.
- [28] M. Del Zotto, N. Nekrasov, N. Piazzalunga, and M. Zabzine, "Playing with the index of M-theory," arXiv:2103.10271 [hep-th].
- [29] R. Dijkgraaf, C. Vafa, and E. Verlinde, "M-theory and a topological string duality," arXiv:hep-th/0602087.
- [30] R. Dijkgraaf, S. Gukov, A. Neitzke, and C. Vafa, "Topological M-theory as unification of form theories of gravity," *Adv. Theor. Math. Phys.* 9 no. 4, (2005) 603-665, arXiv:hep-th/0411073.
- [31] J. de Boer, A. Naqvi, and A. Shomer, "The Topological G(2) string," Adv. Theor. Math. Phys. 12 no. 2, (2008) 243-318, arXiv:hep-th/0506211.
- [32] G. Bonelli and M. Zabzine, "From current algebras for p-branes to topological M-theory," JHEP 09 (2005) 015, arXiv:hep-th/0507051.
- [33] G. Bonelli, A. Tanzini, and M. Zabzine, "On topological M-theory," Adv. Theor. Math. Phys. 10 no. 2, (2006) 239–260, arXiv:hep-th/0509175.
- [34] G. Bonelli, A. Tanzini, and M. Zabzine, "Computing Amplitudes in topological M-theory," JHEP 03 (2007) 023, arXiv:hep-th/0611327.
- [35] S. K. Donaldson and R. P. Thomas, "Gauge theory in higher dimensions," in Conference on Geometric Issues in Foundations of Science in honor of Sir Roger Penrose's 65th Birthday, pp. 31–47. 6, 1996.
- [36] K. Costello, "Holography and Koszul duality: the example of the M2 brane," arXiv:1705.02500 [hep-th].
- [37] D. Gaiotto and J. Oh, "Aspects of Ω-deformed M-theory," arXiv:1907.06495 [hep-th].
- [38] M. Del Zotto, "Geometric Engineering and Correspondences; to appear,".
- [39] A. Johansen, "Twisting of N = 1 SUSY gauge theories and heterotic topological theories," Int. J. Mod. Phys. A 10 (1995) 4325-4358, arXiv:hep-th/9403017.

- [40] E. Witten, "Supersymmetric Yang-Mills theory on a four manifold," J. Math. Phys. 35 (1994) 5101-5135, arXiv:hep-th/9403195.
- [41] A. Johansen, "Infinite conformal algebras in supersymmetric theories on four manifolds," Nucl. Phys. B 436 (1995) 291-341, arXiv:hep-th/9407109.
- [42] A. Johansen, "Holomorphic currents and duality in N = 1 supersymmetric theories," JHEP 12 (2003) 032, arXiv:hep-th/0309125.
- [43] C. Closset, T. T. Dumitrescu, G. Festuccia, and Z. Komargodski, "The Geometry of Supersymmetric Partition Functions," JHEP 01 (2014) 124, arXiv:1309.5876 [hep-th].
- [44] C. Closset, T. T. Dumitrescu, G. Festuccia, and Z. Komargodski, "From Rigid Supersymmetry to Twisted Holomorphic Theories," *Phys. Rev. D* 90 no. 8, (2014) 085006, arXiv:1407.2598 [hep-th].
- [45] E. Witten, "Branes, Instantons, And Taub-NUT Spaces," JHEP 06 (2009) 067, arXiv:0902.0948 [hep-th].
- [46] I. n. García Etxebarria, B. Heidenreich, and D. Regalado, "IIB flux non-commutativity and the global structure of field theories," *JHEP* 10 (2019) 169, arXiv:1908.08027 [hep-th].
- [47] F. Albertini, M. Del Zotto, I. n. García Etxebarria, and S. S. Hosseini, "Higher Form Symmetries and M-theory," *JHEP* 12 (2020) 203, arXiv:2005.12831 [hep-th].
- [48] D. R. Morrison, S. Schafer-Nameki, and B. Willett, "Higher-Form Symmetries in 5d," JHEP 09 (2020) 024, arXiv:2005.12296 [hep-th].
- [49] R. L. Bryant and S. M. Salamon, "On the construction of some complete metrics with exceptional holonomy," *Duke Mathematical Journal* 58 no. 3, (1989) 829 – 850.
- [50] M. Atiyah and E. Witten, "M theory dynamics on a manifold of G(2) holonomy," Adv. Theor. Math. Phys. 6 (2003) 1-106, arXiv:hep-th/0107177.
- [51] B. S. Acharya and E. Witten, "Chiral fermions from manifolds of G(2) holonomy," arXiv:hep-th/0109152.
- [52] J. Oh and Y. Zhou, "A domain wall in twisted M-theory," arXiv:2105.09537 [hep-th].
- [53] M. Dedushenko, S. Gukov, and P. Putrov, "Vertex algebras and 4-manifold invariants," in *Nigel Hitchin's 70th Birthday Conference*, vol. 1, pp. 249–318. 5, 2017. arXiv:1705.01645 [hep-th].

- [54] S. Cecotti, C. Gerig, and C. Vafa, "G₂ holonomy, Taubes' construction of Seiberg-Witten invariants and superconducting vortices," *JHEP* 04 (2020) 038, arXiv:1909.10453 [hep-th].
- [55] J. A. Minahan and M. Zabzine, "Gauge theories with 16 supersymmetries on spheres," JHEP 03 (2015) 155, arXiv:1502.07154 [hep-th].
- [56] D. Prins, "Supersymmetric Gauge Theory on Curved 7-Branes," Fortsch. Phys. 67 no. 5, (2019) 1900009, arXiv:1812.05349 [hep-th].
- [57] N. Iakovidis, J. Qiu, A. Rocén, and M. Zabzine, "7D supersymmetric Yang-Mills on hypertoric 3-Sasakian manifolds," JHEP 06 (2020) 026, arXiv:2003.12461 [hep-th].
- [58] G. Festuccia and N. Seiberg, "Rigid Supersymmetric Theories in Curved Superspace," JHEP 06 (2011) 114, arXiv:1105.0689 [hep-th].
- [59] B. S. Acharya, M. O'Loughlin, and B. J. Spence, "Higher dimensional analogs of Donaldson-Witten theory," *Nucl. Phys. B* 503 (1997) 657-674, arXiv:hep-th/9705138.
- [60] L. Baulieu, H. Kanno, and I. M. Singer, "Cohomological Yang-Mills theory in eight-dimensions," in APCTP Winter School on Dualities of Gauge and String Theories, pp. 365–373. 4, 1997. arXiv:hep-th/9705127.
- [61] L. Baulieu, H. Kanno, and I. M. Singer, "Special quantum field theories in eight-dimensions and other dimensions," *Commun. Math. Phys.* **194** (1998) 149–175, arXiv:hep-th/9704167.
- [62] S. Fubini and H. Nicolai, "The Octonionic Instanton," Phys. Lett. B 155 (1985) 369–372.
- [63] A. D. Popov, "Spherically symmetric solutions of Yang-Mills equations in D = 7 for arbitrary gauge group," *Europhys. Lett.* 17 (1992) 23–26.
- [64] T. A. Ivanova and A. D. Popov, "Selfdual Yang-Mills fields in d = 7, 8, octonions and Ward equations," *Lett. Math. Phys.* 24 (1992) 85–92.
- [65] D. Harland, T. A. Ivanova, O. Lechtenfeld, and A. D. Popov, "Yang-Mills flows on nearly Kahler manifolds and G(2)-instantons," *Commun. Math. Phys.* 300 (2010) 185–204, arXiv:0909.2730 [hep-th].
- [66] M. B. Green, J. H. Schwarz, and E. Witten, *SUPERSTRING THEORY. VOL. 2:* LOOP AMPLITUDES, ANOMALIES AND PHENOMENOLOGY. ., 7, 1988.

- [67] P. K. Townsend and P. van Nieuwenhuizen, "GAUGED SEVEN-DIMENSIONAL SUPERGRAVITY," Phys. Lett. B 125 (1983) 41–46.
- [68] G. Tian, "Gauge theory and calibrated geometry. 1.," Annals Math. 151 (2000) 193-268, arXiv:math/0010015.
- [69] D. Joyce, Compact Manifolds with Special Holonomy. Oxford mathematical monographs. Oxford University Press, 2000.
- [70] B. S. Acharya and S. Gukov, "M theory and singularities of exceptional holonomy manifolds," *Phys. Rept.* **392** (2004) 121–189, arXiv:hep-th/0409191.
- [71] R. Bryant, Metrics with holonomy G2 or spin (7). Hirzebruch F., Schwermer J., Suter S. (eds) Arbeitstagung Bonn 1984. Lecture Notes in Mathematics, vol 1111. Springer, Berlin, Heidelberg., 1985.
- [72] A. Doan, "Monopoles and Fueter sections on three-manifolds," Ph.D Thesis (2019).
- [73] T. Walpuski, "Gauge theory on G_2 -manifolds," Ph.D Thesis (2013).
- [74] J. Driscoll, "Deformations of asymptotically conical g_2 -instantons," 2021.
- [75] A. Lossev, N. Nekrasov, and S. L. Shatashvili, "Testing Seiberg-Witten solution," NATO Sci. Ser. C 520 (1999) 359–372, arXiv:hep-th/9801061.
- [76] N. A. Nekrasov, "Seiberg-Witten prepotential from instanton counting," Adv. Theor. Math. Phys. 7 no. 5, (2003) 831-864, arXiv:hep-th/0206161.
- [77] S. A. Cherkis, "Moduli Spaces of Instantons on the Taub-NUT Space," Commun. Math. Phys. 290 (2009) 719-736, arXiv:0805.1245 [hep-th].
- [78] S. A. Cherkis, C. O'Hara, and C. Saemann, "Super Yang-Mills Theory with Impurity Walls and Instanton Moduli Spaces," *Phys. Rev. D* 83 (2011) 126009, arXiv:1103.0042 [hep-th].
- [79] S. Kachru and J. McGreevy, "M theory on manifolds of G(2) holonomy and type IIA orientifolds," JHEP 06 (2001) 027, arXiv:hep-th/0103223.
- [80] H. Ooguri and C. Vafa, "Geometry of N=1 dualities in four-dimensions," Nucl. Phys. B 500 (1997) 62-74, arXiv:hep-th/9702180.
- [81] A. E. Lawrence, N. Nekrasov, and C. Vafa, "On conformal field theories in four-dimensions," Nucl. Phys. B 533 (1998) 199-209, arXiv:hep-th/9803015.

- [82] I. R. Klebanov and E. Witten, "Superconformal field theory on three-branes at a Calabi-Yau singularity," Nucl. Phys. B 536 (1998) 199–218, arXiv:hep-th/9807080.
- [83] C. Beasley, B. R. Greene, C. I. Lazaroiu, and M. R. Plesser, "D3-branes on partial resolutions of Abelian quotient singularities of Calabi-Yau threefolds," *Nucl. Phys. B* 566 (2000) 599–640, arXiv:hep-th/9907186.
- [84] A. Hanany and A. Iqbal, "Quiver theories from D6 branes via mirror symmetry," JHEP 04 (2002) 009, arXiv:hep-th/0108137.
- [85] B. Feng, Y.-H. He, K. D. Kennaway, and C. Vafa, "Dimer models from mirror symmetry and quivering amoebae," Adv. Theor. Math. Phys. 12 no. 3, (2008) 489–545, arXiv:hep-th/0511287.
- [86] A. Strominger, S.-T. Yau, and E. Zaslow, "Mirror symmetry is T duality," Nucl. Phys. B479 (1996) 243–259, arXiv:hep-th/9606040.
- [87] M. R. Douglas and G. W. Moore, "D-branes, quivers, and ALE instantons," arXiv:hep-th/9603167.
- [88] S. H. Katz and C. Vafa, "Matter from geometry," Nucl. Phys. B 497 (1997) 146-154, arXiv:hep-th/9606086.
- [89] K. Hori and C. Vafa, "Mirror symmetry," arXiv:hep-th/0002222.
- [90] K. A. Intriligator, R. G. Leigh, and N. Seiberg, "Exact superpotentials in four-dimensions," Phys. Rev. D 50 (1994) 1092-1104, arXiv:hep-th/9403198.
- [91] C. Beasley and E. Witten, "New instanton effects in supersymmetric QCD," JHEP 01 (2005) 056, arXiv:hep-th/0409149.
- [92] F. Apruzzi, J. J. Heckman, D. R. Morrison, and L. Tizzano, "4D Gauge Theories with Conformal Matter," JHEP 09 (2018) 088, arXiv:1803.00582 [hep-th].
- [93] M. Aganagic, A. Klemm, M. Marino, and C. Vafa, "The Topological vertex," Commun. Math. Phys. 254 (2005) 425–478, arXiv:hep-th/0305132.
- [94] A. Iqbal, C. Kozcaz, and C. Vafa, "The Refined topological vertex," JHEP 10 (2009) 069, arXiv:hep-th/0701156.
- [95] E. Witten, "Constraints on Supersymmetry Breaking," Nucl. Phys. B 202 (1982) 253.
- [96] E. Witten, "Sigma models and the ADHM construction of instantons," J. Geom. Phys. 15 (1995) 215-226, arXiv:hep-th/9410052.

- [97] M. R. Douglas, "Branes within branes," NATO Sci. Ser. C 520 (1999) 267-275, arXiv:hep-th/9512077.
- [98] M. R. Douglas, "Gauge fields and D-branes," J. Geom. Phys. 28 (1998) 255-262, arXiv:hep-th/9604198.
- [99] C. Hwang, J. Kim, S. Kim, and J. Park, "General instanton counting and 5d SCFT," JHEP 07 (2015) 063, arXiv:1406.6793 [hep-th]. [Addendum: JHEP 04, 094 (2016)].
- [100] K. Hori, H. Kim, and P. Yi, "Witten Index and Wall Crossing," JHEP 01 (2015) 124, arXiv:1407.2567 [hep-th].
- [101] H.-C. Kim, "Line defects and 5d instanton partition functions," JHEP 03 (2016) 199, arXiv:1601.06841 [hep-th].
- [102] L. Jeffrey and F. Kirwan, "Localization for nonabelian group actions," arXiv:9307001 [alg-geom].
- [103] F. Benini, R. Eager, K. Hori, and Y. Tachikawa, "Elliptic Genera of 2d N = 2 Gauge Theories," Commun. Math. Phys. 333 no. 3, (2015) 1241-1286, arXiv:1308.4896 [hep-th].
- [104] N. Haouzi and C. Kozçaz, "Supersymmetric Wilson Loops, Instantons, and Deformed W-Algebras," arXiv:1907.03838 [hep-th].
- [105] A. Hanany and E. Witten, "Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics," Nucl. Phys. B 492 (1997) 152–190, arXiv:hep-th/9611230.
- [106] K. Costello and J. Yagi, "Unification of integrability in supersymmetric gauge theories," arXiv:1810.01970 [hep-th].
- [107] D. Bashkirov and A. Kapustin, "Supersymmetry enhancement by monopole operators," JHEP 05 (2011) 015, arXiv:1007.4861 [hep-th].
- [108] J. Yagi, "Ω-deformation and quantization," JHEP 08 (2014) 112, arXiv:1405.6714 [hep-th].
- [109] M. Bullimore, T. Dimofte, D. Gaiotto, and J. Hilburn, "Boundaries, Mirror Symmetry, and Symplectic Duality in 3d N = 4 Gauge Theory," JHEP 10 (2016) 108, arXiv:1603.08382 [hep-th].
- [110] M. Bullimore, T. Dimofte, D. Gaiotto, J. Hilburn, and H.-C. Kim, "Vortices and Vermas," Adv. Theor. Math. Phys. 22 (2018) 803-917, arXiv:1609.04406 [hep-th].

- [111] M. Mezei, S. S. Pufu, and Y. Wang, "A 2d/1d Holographic Duality," arXiv:1703.08749 [hep-th].
- [112] M. Dedushenko, S. S. Pufu, and R. Yacoby, "A one-dimensional theory for Higgs branch operators," JHEP 03 (2018) 138, arXiv:1610.00740 [hep-th].
- [113] D. Gaiotto and J. Abajian, "Twisted M2 brane holography and sphere correlation functions," arXiv:2004.13810 [hep-th].
- [114] J. Oh and Y. Zhou, "Feynman diagrams and Ω-deformed M-theory," SciPost Phys. 10 no. 2, (2021) 029, arXiv:2002.07343 [hep-th].
- [115] Y. Hatsuda and T. Okazaki, "Fermi-gas correlators of ADHM theory and triality symmetry," arXiv:2107.01924 [hep-th].
- [116] M. Berkooz, M. R. Douglas, and R. G. Leigh, "Branes intersecting at angles," *Nucl. Phys. B* 480 (1996) 265–278, arXiv:hep-th/9606139.
- [117] E. Witten, "BPS Bound states of D0 D6 and D0 D8 systems in a B field," JHEP 04 (2002) 012, arXiv:hep-th/0012054.
- [118] N. Seiberg and E. Witten, "String theory and noncommutative geometry," JHEP 09 (1999) 032, arXiv:hep-th/9908142.
- [119] S. J. Gates, M. T. Grisaru, M. Rocek, and W. Siegel, Superspace Or One Thousand and One Lessons in Supersymmetry, vol. 58 of Frontiers in Physics. ., 1983. arXiv:hep-th/0108200.
- [120] N. Nekrasov and N. S. Prabhakar, "Spiked Instantons from Intersecting D-branes," Nucl. Phys. B 914 (2017) 257–300, arXiv:1611.03478 [hep-th].
- [121] A. Abouelsaood, C. G. Callan, Jr., C. R. Nappi, and S. A. Yost, "Open Strings in Background Gauge Fields," Nucl. Phys. B 280 (1987) 599–624.
- [122] V. Balasubramanian and R. G. Leigh, "D-branes, moduli and supersymmetry," *Phys. Rev. D* 55 (1997) 6415–6422, arXiv:hep-th/9611165.
- [123] E. Pomoni, W. Yan, and X. Zhang, "Tetrahedron instantons," arXiv:2106.11611 [hep-th].
- [124] H. Nakajima, "Instantons on ALE Spaces, Quiver Varieties, and Kac-Moody Algebras," Duke Mathematical Journal 76 (1994) 365?416.
- [125] W. Crawley-Boevey, "Geometry of the Moment Map for Representations of Quivers," Compositio Mathematica 126 (2001) 257?93.
- [126] D. Maulik and A. Okounkov, "Quantum groups and quantum cohomology," 2018.

- [127] Fantechi, Gottsche, Illusie, Kleiman, Nitsure, and Vistoli, Fundamental Algebraic Geometry: Grothendieck?s FGA Explained. American Mathematical Society., 2005.
- [128] E. Carlsson and A. Okounkov, "Exts and vertex operators," Duke Mathematical Journal 161 no. 9, (Jun, 2012).
- [129] K. McGerty and T. Nevins, "Kirwan surjectivity for quiver varieties," *Inventiones mathematicae* 212 no. 1, (Nov, 2017) 161?187.
- [130] V. Baranovsky, "The Variety of Pairs of Commuting Nilpotent Matrices is Irreducible," *Transformation Groups* 6 (2001) 3–8.