Classical gauge principle - From field theories to classical mechanics

B. F. Rizzuti^{*1} and G. F. Vasconcelos Jr.^{†2}

¹Departamento de Física, Universidade Federal de Juiz de Fora, MG, Brazil ²Instituto de Matemática e Estatística, Universidade de São Paulo, SP, Brazil

Abstract

In this paper we discuss how the gauge principle can be applied to classical-mechanics models with finite degrees of freedom. The local invariance of a model is understood as its invariance under the action of a matrix Lie group of transformations parametrized by arbitrary functions. It is formally presented how this property can be introduced in such systems, followed by modern applications. Furthermore, Lagrangians describing classical-mechanics systems with local invariance are separated in equivalence classes according to their local structures.

1 Introduction

The importance of local invariance in field theoretical models is unquestionable. To illustrate that, we could use, for instance, the gauge principle. It states that a Lagrangian field model invariant under the action of a Lie group with global parameters can be reformulated if we raise the parameters to local ones. That is the basic ingredient R. Utiyama [1] has used to generalize the seminal previous paper by Yang and Mills [2]. The introduction of local parameters into the game implies the insertion of a covariant derivative, as well as a connection, in order to keep the desired invariance. In turn, it leads the initial model to interact with a gauge field. This is the very structure one finds in the unified Standard Model [3]. In this case, the gauge group is given by the unitary product $SU(3) \times SU(2) \times U(1)$, setting together fermions, electroweak boson, gluon and Higgs fields.

The introduction of the gauge principle dates back to 1918, when Hermann Weyl presented his ideas of building a unified field theory, which gave rise to the notion of gauge invariance [4]. Although his ideas were mathematically solid, they did not sound so physically appealing. In effect, his proposal predicted that if one transported two identical measuring rods, from points p_1 to

^{*}brunorizzuti@ice.ufjf.br

[†]guilherme.vasconcelos@ime.usp.br

 p_2 , along different paths which cross regions with distinct electromagnetic fields, generally they would no longer have the same length if measured on the point of arrival. Nonetheless, with the ascendance of quantum mechanics, his ideas were no longer physically implausible. They were actually suitable to link electromagnetism and matter fields and thus played a central role in the development of such areas. It is also worth mentioning that his seminal work still reverberates and provides the necessary background for attempts to relate gravity and electromagnetism [5].

Weyl's proposal was based on the following. In accordance with the theory of General Relativity, spacetime is described by a 4-dimensional manifold Mendowed with a Lorentz metric \tilde{g} . A key aspect of such modeling is that it presupposes the choice of unit length. However, if different observers agree to measure lengths at different points with different units, then not only one Lorentz metric \tilde{g} is being used but a whole set of them. In turn, this set is a class of equivalence \mathcal{G} of metrics g such that $g = e^{\lambda}\tilde{g}$, with λ being an arbitrary smooth function on M. Moreover, a Weyl structure on M is a map $F : \mathcal{G} \to \Lambda^1(M)$, where $\Lambda^1(M)$ is the space of 1-forms on M, defined by $F(e^{\lambda}g) = F(g) - d\lambda$, whilst a manifold with a Weyl structure is called a Weyl manifold [6]. One of the key aspects of this geometrical approach is that Weyl structures allow us to compare lengths of vectors at different points of M through the transport of scalar products, associated to the metrics in \mathcal{G} , along curves over the manifold [7].

Furthermore, such theory enables us to understand the gauge invariance principle in a full geometrical perspective, see [6] for more details. In particular, it reveals how the commonly seen "covariant derivative" $D_{\mu} = \partial_{\mu} + iA_{\mu}$ appears in the context of field theories. Nevertheless, such procedure to define Weyl structures is not the only one, as it can be seen in [8].

The central role played by local invariance is not only restricted to (quantum) field models. Therefore, it is natural to step back and seek its implications on systems with finite degrees of freedom. In effect, there is a long list of relevant models where local or gauge invariance is as important as described previously. For example, the dynamics of any classical-mechanics system can be formulated with a local structure [9,10]. We have only to take into account the corresponding reparametrization-invariant formulation. In this case, reparametrization invariance (RI) can be seen as a gauge symmetry [11]. A mathematical oriented perspective on all that can be seen in [12].

RI has even deeper implications though. It may lead to a common arrow of time and non-negativity of mass [13]. Broader consequences, such as an underlying origin to classical long range interactions (electrodynamics and gravitation) could also be supported by reparametrization-invariant systems [14]. Gauge theory for finite-dimensional dynamical systems has also been explored, running from quantum mechanics to information theory, passing through chemistry and rigid body dynamics [15].

In a more philosophical approach, in the realm of Newtonian physics, one could eliminate the notion of an absolute space by gauging translations and rotations [16]. While this is restricted to the Galileo group, we extend it in

this paper by considering the gauge principle directly applied to arbitrary Lagrangian systems with finite degrees of freedom. We will name it *Classical Gauge Principle*. The prescription is pretty much the same: a Lagrangian model inert under the representation of a Lie group parametrized by global parameters is reformulated by the swap of such parameters to local ones, that is, smooth functions of the evolution variable. Just as before, it implies the introduction of a covariant derivative (and the corresponding connection) into the original model.

The paper is divided as follows. In Section 2 we present a seminal and illustrative example which can be rewritten in order to present local invariance, according to the gauge principle. This example is a key application of such principle and, in turn, the consequences of its usage are then enumerated. In Section 3 we develop the method of introducing local invariance in classical-mechanics models and present two modern examples in which this technique may be applied. In Section 4 we formalize the concept of equivalent classes of Lagrangians with the same local structure. Furthermore, leveraging on a more pragmatic view, two examples are given to elucidate such approach. Finally, Section 5 is left for the conclusions.

2 Free Dirac Lagrangian

Let us consider an underlying example of a model endowed with local invariance. We start with the Lagrangian of a free massive spinor field $\psi = \psi(x^{\mu})$ and write

$$\mathcal{L} = \bar{\psi}(i\gamma^{\mu}\partial_{\mu} - m)\psi. \tag{1}$$

Here, $\bar{\psi} = \psi^{\dagger} \gamma^{0}$, whereas γ^{μ} are the Dirac-gamma matrices and $\gamma^{\mu} \gamma^{\nu} + \gamma^{\nu} \gamma^{\mu} = 2\eta^{\mu\nu}$, with $\eta^{\mu\nu} = \text{diag}(1, -1, -1, -1)$. In turn, \mathcal{L} is invariant under the action of the symmetry group U(1), since the transformations $\psi \mapsto \psi' = e^{i\alpha}\psi$ and $\bar{\psi} \mapsto \bar{\psi}' = e^{-i\alpha}\bar{\psi}$, where α is a real number, do not alter the Lagrangian,

$$\mathcal{L} \mapsto \mathcal{L}' = e^{-i\alpha} \bar{\psi} (i\gamma^{\mu} \partial_{\mu} - m) e^{i\alpha} \psi = \mathcal{L}.$$
⁽²⁾

One may now ask if this Lagrangian would still be invariant under transformations where α is an arbitrary smooth function. Naturally, it would no longer be invariant under the action of the same transformations, due to the derivative $\partial_{\mu}\psi$ term. However, local invariance can be achieved in this scenario. In order to enforce such property, we introduce the vector field A_{μ} in \mathcal{L} and define

$$D_{\mu} \coloneqq \partial_{\mu} - iA_{\mu}. \tag{3}$$

We also fix the transformation of A_{μ} with

$$A_{\mu} \mapsto A'_{\mu} = A_{\mu} + \partial_{\mu} \alpha. \tag{4}$$

This way, we write

$$\mathcal{L} = \bar{\psi}(i\gamma^{\mu}D_{\mu} - m)\psi, \tag{5}$$

whereas $D_{\mu}\psi$ has a simple transformation law,

$$D_{\mu}\psi \mapsto D'_{\mu}\psi' = e^{i\alpha(x)}D_{\mu}\psi. \tag{6}$$

Therefore, local invariance for \mathcal{L} has been established, as previously claimed. Such procedure is the standard way in order to obtain the interaction of the initial free field with the vector A_{μ} , commonly called the gauge field.

Let us make a two elucidating observations.

- 1. Models with local invariance are constructed according to this recipe, known as the gauge principle. Namely, a field model which is invariant under a Lie group of global transformations may be reformulated to be invariant under the same group, although with local parameters. In order to do so, an interaction with a vector field is introduced, whilst the components of this field, named gauge field, are but the components of a 1-form which defines the connection map on a Weyl manifold [6].
- 2. Lagrangian models with local invariance present first class constraints through the corresponding hamiltonization procedure [17, 18]. Theories of this type possess spurious degrees of freedom, with ambiguous evolution. The physical sector may be consistently characterized by gauge invariant variables.

In our illustrative example, A_{μ} is the vector potential. Since it changes under the local transformation, see (4), A_{μ} has no physical interpretation. The possible observables of the model are combinations that are not affected by the local transformations; for instance, the integral $I = \oint \vec{A} \cdot d\vec{l}$ of the spatial part of A_{μ} over any closed contour is invariant under (4). It manifests itself through the Aharonov-Bohm effect [19]. The electromagnetic tensor, represented by $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$, is another example of local invariant which is an observable.

At this point, a natural question that rises is if the gauge principle for field models can be transposed to systems with finite degrees of freedom. The answer is yes and the Classical Gauge Principle will be discussed in details in the next section, followed by applications in both classical and relativistic mechanics.

3 Introducing local invariance in classical-mechanics models

To begin with, let us consider a configuration space parametrized by x^a , $a = 1, \ldots, n$ and a mechanical model defined on it. The corresponding dynamics is governed by a Lagrangian $L(x^a, \dot{x}^a)$, where $\dot{x}^a := \frac{dx^a}{d\tau}$ and τ is an evolution parameter. As usual, the equations of motion are obtained from a least action principle applied to the action functional

$$S = \int d\tau L(x^a, \dot{x}^a). \tag{7}$$

We also impose that L is invariant under the faithful representation of a matrix Lie group \mathcal{G} , whose elements are denoted by G, with respective entries $G^a_{\ b}$ in a field \mathbb{F} . The linear representation is defined by

$$x^a \to x'^a \coloneqq G^a_{\ b} x^b. \tag{8}$$

Although it appears to be a restriction to consider only $n \times n$ -matrix groups directly realized onto the underlying (*n*-dimensional) configuration space, every case of interest to this work possess this very structure. We could go even further and realize that there are many standard examples where this particular representation plays a central role, for instance SO(3) as the rotation group in space or even SO(1,3) describing the Lorentz transformations in spacetime and so on [20].

The corresponding Lie algebra is denoted by

$$\mathfrak{g} = \{\xi^i \Gamma_i \mid \xi^i \in \mathbb{F}, \ [\Gamma_i, \Gamma_j] = c_{ij}^{\ k} \Gamma_k\}, \tag{9}$$

that is, \mathfrak{g} is generated by $\Gamma_i, i = 1, \ldots, n$ and $c_{ij}^{\ k}$ are the structure constants. Any $G \in \mathcal{G}$ sufficiently close to the identity of the group may be written as

$$G = e^{\xi^* \Gamma_i},\tag{10}$$

with $G^a_b = (e^{\xi^i \Gamma_i})^a_b$. It will be sufficient to this work to write the group elements only in first order in the exponential map,

$$G^a_{\ b} \approx \delta^a_{\ b} + \xi^i (\Gamma_i)^a_{\ b}. \tag{11}$$

According to the initial assumption,

$$L(x^{a}, \dot{x}^{a}) \xrightarrow{G} L'(x'^{a}, \dot{x}'^{a}) = L(G^{a}_{\ b}x^{b}, G^{a}_{\ b}\dot{x}^{b}) = L(x^{a}, \dot{x}^{a}).$$
(12)

Now, we may relax this first assumption to include not only constant parameters ξ^i , but also smooth functions

$$\xi^{i}: I \subset \mathbb{R} \to \mathbb{F}, \ \forall \ i \in \{1, \dots, n\}$$

$$\tag{13}$$

$$\tau \mapsto \xi^i(\tau),\tag{14}$$

where I is an open interval of the real line. We denote both the Lie algebra and group so obtained by \mathfrak{g}_{τ} and G_{τ} . It means they are local in the sense that their elements are parametrized by arbitrary smooth functions of τ . Clearly, due to the velocity \dot{x}^a in L, the latter is no longer invariant under the G-action,

$$\frac{dx^a}{d\tau} \to \frac{dx'^a}{d\tau} = \frac{d}{d\tau} (\delta^a_{\ b} + \xi^i (\Gamma_i)^a_{\ b}) x^b \tag{15}$$

$$= G^a{}_b \dot{x}^b + \dot{\xi}^i (\Gamma_i)^a{}_b x^b.$$
⁽¹⁶⁾

We would like to maintain its invariance even under the action of the local group G_{τ} . Hence, we proceed exactly in the same manner one introduces covariant derivatives through the gauge principle in field theories. In our cause, though, we substitute the time derivative for

$$\delta^a_{\ b} \frac{d}{d\tau} \to D^a_{\ b} \coloneqq \delta^a_{\ b} \left(\frac{d}{d\tau} - g\right),\tag{17}$$

where g is a new variable, analogous to the gauge field. The Lagrangian remains invariant under G_{τ} , provided g changes according to

$$\delta^a_{\ b}g \to \delta^a_{\ b}g' = \delta^a_{\ b}g + \dot{\xi}^i (\Gamma_i)^a_{\ b}. \tag{18}$$

In fact, the pathological term comes from the velocity, see (16). However, it is ruled out by the transformation provided above in (18),

$$D^{a}_{\ b}x^{b} \to D'^{a}_{\ b}x'^{b} = G^{a}_{\ b}D^{b}_{\ c}x^{c}.$$
 (19)

In turn, returning back to L,

$$L(x^{a}, D^{a}{}_{b}x^{b}) \to L(x'^{a}, D'^{a}{}_{b}x'^{b}) = L(G^{a}{}_{b}x^{b}, G^{a}{}_{b}D^{b}{}_{c}x^{c})$$
$$= L(x^{a}, D^{a}{}_{b}x^{b}),$$
(20)

where the equality comes from (12).

As claimed previously, we have imposed local invariance to the action we are working with. In the next subsections, we apply the procedure developed here to concrete models, revealing the power of the method while explaining a possible geometrical and algebraic origin to leading proposals, such as the free relativistic particle, as well as more sophisticated examples, such as doubly special relativity (DSR) models and spinning particles.

3.1 Free particle and DSR proposal

Let us consider as a first example a particle model described by the Lagrangian

$$L = \frac{1}{2}m\eta(v,v),\tag{21}$$

defined on (\mathbb{R}^5, η) , a 5-dimensional spacetime endowed with a pseudo-metric η and parametrized by $\{x^A\}$. *m* is a constant that can be interpreted as the mass of the particle. We shall denote $v^A = \dot{x}^A$ and write $\eta_{AB} = \text{diag}(+1, -1, -1, -1, -1)$. We also allow ourselves to multiply position and velocity vectors by complex numbers and impose

$$\eta(\alpha v_1, \beta v_2) = (\alpha^* \beta) \eta(v_1, v_2), \ \forall \ \alpha, \beta \in \mathbb{C}.$$
(22)

With these considerations, we take the representation of the group $U(1) = \{e^{i\gamma}; \gamma \in \mathbb{R}\}$ on (\mathbb{R}^5, η) given by

$$x^A \to x'^A = e^{i\gamma} x^A \Rightarrow v'^A = e^{i\gamma} v^A, \tag{23}$$

which gives us

$$L \to L' = \frac{1}{2}m\eta(v',v') = \frac{1}{2}m\eta(e^{i\gamma}v,e^{i\gamma}v) = \frac{1}{2}m\eta(v,v) = L.$$
 (24)

According to the classical gauge principle, we raise U(1) to the status of a local group, meaning that its elements are now parametrized by smooth functions $\gamma: I \text{ (open)} \subset \mathbb{R} \to \mathbb{R}$. As explained, L is no longer invariant under $x'^{\mu} = e^{i\gamma(\tau)}x^{\mu}$. In accordance with what we have exposed previously, we substitute

$$\frac{d}{d\tau} \to D \coloneqq \frac{d}{d\tau} - i\gamma. \tag{25}$$

This way, in local coordinates the Lagrangian reads

$$L = \frac{1}{2}m\eta_{AB}Dx^{A}Dx^{B} = \frac{1}{2}m\eta_{AB}(\dot{x}^{A} - i\gamma x^{A})(\dot{x}^{B} - i\gamma x^{B}),$$
(26)

which is now invariant under $U(1)_{\tau} = \{e^{i\gamma(\tau)}; \gamma: I \to \mathbb{R}\}$. If we redefine $i\gamma = g$, we obtain exactly the same action proposed in [21]. Due to the local invariance, the Lagrangian presents first class constraints through the Dirac's hamiltonization algorithm [22]. The physical sector of the model describes a free massive particle and the superior bounded speed is obtained in a particular gauge choice. It is also possible to show that the model leads to the Magueijo-Smolin doubly special relativity (MS DSR) proposal [23].

We conclude this section with a brief comment concerning the transformation law given in (18). It implies that the gauge variable g changes as a total derivative,

$$\Delta(\delta^a_{\ b}g) \coloneqq \delta^a_{\ b}(g'-g) = \frac{d}{d\tau} \left(\xi^i(\tau)(\Gamma_i)^a_{\ b}\right). \tag{27}$$

In this case, we could change L in (21) for

$$L \to L' = L + \xi g = \frac{1}{2} m \eta D x^A D x^B - \xi g, \qquad (28)$$

where ξ is a constant that couples the gauge variable g to the dynamical sector of the model. In turn, it can be shown that these Lagrangians are equivalent and possess the same local structure (this notion shall be clarified in Section 4). Moreover, (28) is the Lagrangian described in the work [24], in which it was shown that the MS DSR model is a particular gauge of a free relativistic particle Lagrangian model. In this particular case, ξ is related to the new invariant scale, presented within the model from the beginning.

3.2 Classical spinning particle

Let us discuss another example of imposing local invariance to a classical model. Contrary to our previous example, the covariant derivative encodes the interaction of the model with an external field. Moreover, the local parameters gain clear physical interpretation, as we shall see. On the configuration space parametrized by $\{v^i, g, \varphi\}, i \in \{1, 2, 3\}$, we define the following Lagrangian,

$$L_{spin} = \frac{1}{2g} (\dot{v}^i)^2 + g \frac{b^2}{2a^2} + \frac{1}{\varphi} [(v^i)^2 - a^2].$$
(29)

We are using the short hand notation $(v^i)^2 = \delta_{ij} v^i v^j$, as well as $\dot{v}^i = \frac{dv^i}{d\tau}$. On one hand *a* is an arbitrary constant, while $b^2 = 3\hbar^2/4$.

This model is particularly important to describe internal degrees of freedom of classical spinning particles. In fact, when set together with space-time dynamical variables, we find a Lagrangian whose equations of motion produce the Pauli equation through the canonical quantization method. All the details may be seen in [25].

We shall focus on (29). It is clear that this Lagrangian has manifest SO(3) global invariance,

$$v^{i} \to v^{\prime i} = R^{i}{}_{j}v^{j}; \quad \forall \ R \in SO(3).$$

$$(30)$$

Identifying SO(3) as a matrix Lie group we may write

$$R^i{}_j = \left(e^{\xi^\kappa \Gamma_k}\right)^i{}_j,\tag{31}$$

where $\xi^k \Gamma_k$ is an element of the Lie algebra $\mathfrak{so}(3)$ and ξ^k are interpreted as rotation angles. According to the Classical Gauge Principle, we promote the constant parameters ξ^k to arbitrary smooth functions $\xi^k(\tau)$. In order o maintain the invariance of L_{spin} , we introduce the covariant derivative

$$\delta^{i}{}_{j}\frac{d}{d\tau} \to \delta^{i}{}_{j}\frac{d}{d\tau} + \dot{\xi}^{k}(\Gamma_{k})^{i}{}_{j}.$$
(32)

In this case $\dot{\xi}^k$ may be interpreted as an angular velocity. We redefine it according to

$$\dot{\xi}^k = -\frac{e}{m}B^k,\tag{33}$$

which provides the angular velocity of a charged massive particle in terms of an external magnetic field.

Together with this redefinition, there is an explicit form for the generators Γ ,

$$(\Gamma_k)^i{}_j = \varepsilon_{ikj},\tag{34}$$

where ε_{ijk} is the Levi-Civita symbol. Thus, (29) may be rewritten as

$$L_{spin} = \frac{1}{2g} (\dot{v}^j - \frac{e}{m} \varepsilon_{jik} v^i B^k)^2 + g \frac{b^2}{2a^2} + \frac{1}{\varphi} [(v^i)^2 - a^2].$$
(35)

Our construction provides the correct interaction between an external magnetic field and the spinning degrees of freedom. To verify this, we may look at a possible observable of the theory, given by the angular momentum associated to the v^i -variables,

$$J^i \coloneqq \varepsilon^{ijk} v_j \pi_k, \tag{36}$$

where $\pi_k = \frac{\partial L}{\partial \dot{v}^k}$. The Lagrangian equations of motion imply

$$\dot{J}^{i} = \frac{e}{m} \varepsilon^{ijk} J_j B_k, \tag{37}$$

which is precisely the precession expected from a spinning particle under the effect of an exterior magnetic field.

4 Local structures

Our examples are constructed with a peculiar freedom one has to write out Lagrangian models. Even when two Lagrangians differ, they may possess the same local structure, in the sense of being invariant under (almost) the same gauge transformations. Let us formalize these ideas that unite Lagrangians in equivalence classes. Our background shall be a configuration space parametrized by $\{q^A\}, A \in \{1, ..., N\}$. We write collectively

$$q^A = (x^a, y^\alpha, \dots). \tag{38}$$

That is, the space can be split in different sectors. We may allow Lagrangians to depend only on a particular subset of variables, say, $L(x^a, \dot{x}^a)$. Let \mathscr{L} be the set of all possible Lagrangians of the form $L = L(q^A, \dot{q}^A)$. Now, consider transformations $q^A(\tau) \to q'^A(\tau)$ parametrized by arbitrary functions $\epsilon^{\alpha}(\tau)$; $\alpha \in$ $\{1, ..., n\}$. Their infinitesimal form is given by

$$q^A \to q'^A = q^A + \sum_{k=0}^n \mathscr{R}^A{}_{(k)\alpha}(q^A, \dot{q}^A, ...; \tau) \frac{d^k \epsilon^\alpha(\tau)}{d\tau^k}, \tag{39}$$

where the functions $\mathscr{R}^{A}_{(k)\alpha}$ are the corresponding generators. From here on, we write $\delta q^{A} := q'^{A} - q^{A}$ for short.

Let L_1 and L_2 be two Lagrangians in \mathscr{L} invariant under gauge transformations $\delta_1 q^A$ and $\delta_2 q^A$, respectively. On $\mathscr{L} \times \mathscr{L}$, we set the following relation

$$L_1 \sim L_2 \Leftrightarrow \delta_1 L_1 = \delta_2 L_2 + \frac{dF}{d\tau} \tag{40}$$

whenever there exists such a function F. Clearly, ~ is symmetric, reflexive and transitive, and as so, an equivalence relation. The corresponding equivalence classes will be designated by $[L] \in \mathscr{L}/\sim$ and for $L_1, L_2 \in [L]$, we say that the Lagrangians possess the same *local structure*.

Let us elucidate how this division of ${\mathscr L}$ in classes works with our previous examples.

Example 1. Let us consider the already presented Lagrangians,

$$L_1 = \frac{m}{2} \eta_{AB} D x^A D x^B, \tag{41}$$

$$L_2 = \frac{m}{2} \eta_{AB} D x^A D x^B - \xi g, \qquad (42)$$

see Section 3.

The two Lagrangians are invariant under the same transformations $\delta = \delta_1 = \delta_2$, given by

$$\delta x^{A} = \frac{1}{2} \dot{\alpha} x^{A} - \alpha \dot{x}^{A}$$

$$\delta g = \frac{1}{2} \ddot{\alpha} - \dot{\alpha} g - \alpha \dot{g}$$
(43)

where α is an arbitrary function of the evolution parameter τ . Clearly, $L_1 \neq L_2$, although a straightforward calculation leads to

$$\delta_2 L_2 = \delta_1 L_1 + \left(\xi \alpha g - \frac{\xi \alpha}{2}\right)^2 \Rightarrow L_1 \sim L_2, \tag{44}$$

that is, L_1 and L_2 bear the same local structure.

Example 2. The second example is related to the spinning particle also presented in Sec. 3. In the first place, we take a Lagrangian defined on the configuration space $\{v^i, g\}$ given by

$$L_1 = \frac{1}{2g} (\dot{v}^i)^2 \tag{45}$$

while the second one is defined on an extended space $\{v^i, g, \varphi\}$ and is given by (29). In this case, the local symmetries are given by

$$\begin{aligned}
\delta_1 v^i &= \alpha \dot{v}^i, & \delta_2 v^i &= \alpha \dot{v}^i, \\
\delta_1 g &= (\alpha g)^{\cdot}, & \delta_2 g &= (\alpha g)^{\cdot}, \\
\delta_2 \varphi &= \alpha \dot{\varphi} - \dot{\alpha} \varphi.
\end{aligned}$$
(46)

 α is an arbitrary function of the evolution parameter $\tau.$ The local transformations lead to

$$\delta_2 L_2 = \delta_1 L_1 + \left(\frac{\alpha g b^2}{2a^2} + \frac{\alpha}{\varphi} [(v^i)^2 - a^2]\right) \Rightarrow L_1 \sim L_2.$$

$$\tag{47}$$

This example shows an interesting feature of models with the same local structure. We can impose extra desirable constraints to a particular Lagrangian, enlarging the configuration space, maintaining the different Lagrangians so generated in the same class though. In this particular case, L_1 is not restricted to the sphere $(v^i)^2 = a^2$ as is L_2 . This constraint was introduced in L_2 by the variable φ .

5 Conclusions

This work discusses models with finite degrees of freedom that present local symmetries. Our new results are listed below.

i. We have transposed the gauge principle from field theories to mechanical models. Although there are related works exhibited in the introduction, this is, up to our knowledge, the first attempt to do so with the generality hereby presented. In a few words, we start from a Lagrangian model invariant under

a Lie group parametrized by constant factors. Then the group is raised to a local one and we require that the model stays invariant. With this prescription, the dynamical sector of the initial Lagrangian is now described by a covariant derivative, once a connection naturally appears throughout the method.

ii. Our construction was applied to different current models such as DSR proposals and classical spinning particles. Our prescription provides an original algebraic and geometric origin to the corresponding local invariance imposed, while giving insightful interpretations of relevant physical quantities that naturally emerge in the previously presented models.

iii. Finally, and still concerning the local symmetries, we have shown how the set of Lagrangians is partitioned in equivalence classes. The corresponding equivalence relation was defined in (40). We say that two Lagrangians posses the same local structure when they differ by a total derivative term under local transformations. This composition of the set \mathscr{L} of Lagrangians was particularly important to our own examples. In the first case (regarding DSR proposals), we could deform the initial Lagrangian to a new one, preserving the local structure. In the new framework, the invariant scale ξ presented in DSR models appears as a coupling constant to the gauge variable. The second example (concerning spinning particles) shows how we can add extra variables to the configuration space, enlarging the initial configuration space while still conserving the local structure. For that case, the last term added guaranteed a desirable constraint.

Furthermore, this work may shine some light on the importance of formalizing key aspects of classical mechanics, such as the gauge principle for models with finite degrees of freedom. Even though a more complete description of mechanics is accomplished with quantum field theory, approaches like the one presented here play a central role in the finding and understanding of the physical sector of important models formulated with extra degrees of freedom, thus justifying their formalization.

Acknowledgement

This work was supported by Programa Institucional de Bolsas de Iniciação Científica - XXIX PIBIC/CNPq/UFJF - 2020/2021, project number ID47862.

GFVJr is currently a CNPq fellow - Brazil.

BFR would also like to thank G. Caetano for indicating valuable references concerning the analogy between non-Abelian gauge theories and general relativity.

References

- R. Utiyama, "Invariant theoretical interpretation of interaction," *Phys. Rev.*, vol. 101, pp. 1597–1607, Mar 1956.
- [2] C. N. Yang and R. L. Mills, "Conservation of isotopic spin and isotopic gauge invariance," *Phys. Rev.*, vol. 96, pp. 191–195, Oct 1954.

- [3] J. Baez and J. Huerta, "The algebra of grand unified theories," Bulletin of the American Mathematical Society, vol. 47, pp. 483–552, Mar. 2010.
- [4] H. Weyl, "Gravitation und elektrizität," Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften, pp. 465–478, 1918.
- [5] J. Lindgren and J. Liukkonen, "Maxwell's equations from spacetime geometry and the role of weyl curvature," *Journal of Physics: Conference Series*, vol. 1956, p. 012017, July 2021.
- [6] G. B. Folland, *Quantum field theory : a tourist guide for mathematicians*. American Mathematical Society, 2008.
- [7] G. B. Folland, "Weyl manifolds," *Journal of Differential Geometry*, vol. 4, pp. 145–153, 1970.
- [8] G. S. Hall, "Weyl manifolds and connections," *Journal of Mathematical Physics*, vol. 33, pp. 2633–2638, July 1992.
- [9] F. Debbasch, "Classical mechanics and gauge theories," American Journal of Physics, vol. 61, pp. 1131–1133, Dec. 1993.
- [10] A. A. Deriglazov and B. F. Rizzuti, "Reparametrization-invariant formulation of classical mechanics and the schrödinger equation," *American Jour*nal of Physics, vol. 79, pp. 882–885, Aug. 2011.
- [11] G. G. Fulop, D. M. Gitman, and I. V. Tyutin, "Reparametrization invariance as gauge symmetry," *International Journal of Theoretical Physics*, vol. 38, pp. 1941–1968, 1999.
- [12] E. Massa, E. Pagani, and P. Lorenzoni, "On the gauge structure of classical mechanics," *Transport Theory and Statistical Physics*, vol. 29, pp. 69–91, Jan. 2000.
- [13] V. G. Gueorguiev and A. Maeder, "Reparametrization Invariance and Some of the Key Properties of Physical Systems," *Symmetry*, vol. 13, p. 522, Mar. 2021.
- [14] V. G. Gueorguiev and A. Maeder, "Geometric Justification of the Fundamental Interaction Fields for the Classical Long-Range Forces," *Symmetry*, vol. 13, p. 379, Feb. 2021.
- [15] P. Gurfil, "Gauge theory for finite-dimensional dynamical systems," Chaos: An Interdisciplinary Journal of Nonlinear Science, vol. 17, p. 023107, June 2007.
- [16] R. Ferraro, "Relational mechanics as a gauge theory," *General Relativity and Gravitation*, vol. 48, p. 23, Jan. 2016.

- [17] A. A. Deriglazov and B. F. Rizzuti, "Generalization of the extended lagrangian formalism on a field theory and applications," *Phys. Rev. D*, vol. 83, p. 125011, Jun 2011.
- [18] B. F. Rizzuti, G. F. V. Júnior, and M. A. Resende, "To square root the lagrangian or not: an underlying geometrical analysis on classical and relativistic mechanical models," *Open Access Journal of Mathematical and Theoretical Physics*, vol. 3, pp. 8–15, 2020.
- [19] Y. Aharonov and D. Bohm, "Significance of electromagnetic potentials in the quantum theory," *Phys. Rev.*, vol. 115, pp. 485–491, Aug 1959.
- [20] A. N. Rocha, B. F. Rizzuti, and D. S. Mota, "Transformações de galileu e de lorentz: um estudo via teoria de grupos," *Rev. Bras. Ensino Física*, vol. 35, pp. 4304–9, 2013.
- [21] B. F. Rizzuti and A. A. Deriglazov, "Five-dimensional mechanics as the starting point for the magueijo-smolin doubly special relativity," *Physics Letters B*, vol. 702, no. 2, pp. 173–176, 2011.
- [22] P. A. M. Dirac, Lectures on quantum mechanics. Mineola, NY: Dover Publications, 2001.
- [23] J. Magueijo and L. Smolin, "Lorentz Invariance with an Invariant Energy Scale," *Physical Review Letters*, vol. 88, p. 190403, Apr. 2002.
- [24] B. F. Rizzuti and G. F. Vasconcelos, "5-dimensional so(1,4)-invariant action as an origin to the magueijo-smolin doubly special relativity proposal," *Physics Letters B*, vol. 817, p. 136334, 2021.
- [25] A. A. Deriglazov, "From noncommutative sphere to nonrelativistic spin," Symmetry, Integrability and Geometry: Methods and Applications, vol. 6, p. 016, 2010.