Ergodic convergence rates for time-changed symmetric Lévy processes in dimension one

Tao Wang*

School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education Beijing 100875, China

Abstract

We obtain the lower bounds for ergodic convergence rates, including spectral gaps and convergence rates in strong ergodicity, for time-changed symmetric Lévy processes, by using harmonic function and reversible measure. As direct applications, explicit sufficient conditions for exponential and strong ergodicity are given. Some examples are also presented.

Keywords and phrases: Lévy process; spectral gap; convergence rate in strong ergodicity; harmonic function; time change.

Mathematics Subject classification(2020): 60G51 47A75 35P15

1 Main results and examples

Ergodicity for Lévy-type processes is an important topic in study of Markov processes. In general, criteria are obtained by using Lyapunov functions (cf. [16] for general onedimensional Lévy-type operators, [17] for Lévy-driven SDEs, and [4] for time-changed symmetric stable processes), or coupling methods (see [10] for Lévy-driven SDEs).

Recently, [18] obtains the criteria for strong and exponential ergodicity of one-dimensional time-changed symmetric stable processes; the lower bounds for ergodic convergence rates, including spectral gaps and convergence rates in strong ergodicity are also estimated in [18]. Different from Lyapunov criteria and coupling methods, the main idea in [18] is to estimate the Green function for $\mathbb{R} \setminus \{0\}$ by assuming that the process is pointwise-recurrent.

While [18] deals with the classical α -stable processes for $\alpha \in (1, 2)$, this will exclude some significant pointwise-recurrent Lévy processes such as the diffusion operator with stable jump: $L = a(x)(c_1\Delta + c_2\Delta^{\alpha/2})$, where $\alpha \in (1, 2)$, a is a positive function such that a^{-1} is Lebesgue integrable, c_1 and c_2 are two constants, Δ is the Laplacian operator and $\Delta^{\alpha/2}$ is the fractional Laplacian.

The aim of this paper is to study the ergodic convergence rates for general pointwiserecurrent time-changed symmetric Lévy processes. To this end, we first recall some basic definitions.

^{*}E-mail: wang_tao@mail.bnu.edu.cn.

Let $X = (X_t)_{t \ge 0}$ be a one-dimensional symmetric Lévy process with Lévy measure ν . The characteristic exponent of X is defined as

$$\psi(\xi) := -\log \mathbb{E}_0 e^{i\xi X_1} = \int_{\mathbb{R}} (1 - \cos \xi x) \nu(\mathrm{d}x) + \sigma^2 \xi^2, \quad \xi, \sigma \in \mathbb{R}.$$
(1.1)

The generator \mathcal{A} is a one-dimensional Lévy operator given by

$$\mathcal{A}u(x) := -\int \psi(\xi)\widehat{u}(\xi)e^{ix\xi}\mathrm{d}\xi, \quad u \in C_0^\infty(\mathbb{R}),$$
(1.2)

where $C_0^{\infty}(\mathbb{R})$ is the space of compactly supported smooth functions in \mathbb{R} and $\hat{u}(\xi)$ is the Fourier transform of u. The corresponding regular Dirichlet form $(\mathscr{E}, \mathcal{F})$ is given by

$$\mathscr{E}(u,w) = \int_{\mathbb{R}} \frac{1}{2} \sigma^2 u'(x) w'(x) \mathrm{d}x + \iint_{\mathbb{R} \times \mathbb{R} \setminus \text{diag}} (u(x+h) - u(x)) (w(x+h) - w(x)) \nu(\mathrm{d}h) \mathrm{d}x,$$
(1.3)

for $u, w \in \mathcal{F} := \{f \in L^2(dx) : \mathscr{E}(f, f) < \infty\}$ (see [1, Example 3.13] for more details). Let $p_t(y - x) := p_t(x, y)$ be the transition density of this process. Define

$$H(x) = \int_0^\infty (p_s(0) - p_s(x)) ds = \frac{1}{\pi} \int_0^\infty (1 - \cos xs) \frac{1}{\psi(s)} ds.$$
 (1.4)

Obviously, H is an even function: H(x) = H(-x). Let P_t^0 be the semigroup of the process killed upon hitting the origin, i.e.,

$$P_t^0(x,A) = \mathbb{P}_x[X_t \in A, t < \tau_0], \quad \forall x \in \mathbb{R}, \ \forall A \in \mathscr{B}(\mathbb{R}),$$
(1.5)

where $\tau_0 := \inf\{t > 0 : X_t = 0\}$. By [19, Theorem 1.1], H(x) is the harmonic function for P_t^0 , i.e., for any $x \neq 0$, $P_t^0 H(x) = H(x)$.

Now we consider the time-changed symmetric Lévy processes. Let a be a positive and locally bounded measurable function on \mathbb{R} with $a(x)^{-1}$ integrable. Define $\mu(dx) := a(x)^{-1}dx$, $A_t := \int_0^t 1/a(X_s)ds$, and the time-changed Lévy process $Y_t = X_{\tau_t}$, where

$$\tau_t = \inf \{ s > 0 : A_s > t \}.$$
(1.6)

Then, the generator of time-changed process Y is just the operator $L = a\mathcal{A}$ (for more details about this paragraph, see Section 2).

Let $\mu(f) := \int_{\mathbb{R}} f d\mu$ and $||f||_{L^2(\mu)} := \sqrt{\mu(f^2)}$. We say that Y is (L²-)exponentially ergodic, if there exist non-negative function $C(x) < \infty$ and $\lambda_1 > 0$, such that

$$\|P_t f - \mu(f)\|_{L^2(\mu)} \leqslant e^{-\lambda_1 t} \|f - \mu(f)\|_{L^2(\mu)}.$$
(1.7)

The optimal convergence rate λ_1 in (1.7) (i.e., the L²-spectral gap) is defined by

$$\lambda_1 = \inf\{\mathscr{E}(f, f) : f \in \mathscr{F}, \mu(f^2) = 1, \mu(f) = 0\}.$$
(1.8)

Let $\|\eta\|_{\text{Var}} := \sup_{|f| \leq 1} |\eta(f)|$ be the total variation of a signed measure η . We say that Y is **strongly ergodic**, if there exist constants $C < \infty$ and $\kappa > 0$, such that

$$\sup_{x \in \mathbb{R}} \|P_t(x, \cdot) - \mu\|_{\operatorname{Var}} \leqslant C \mathrm{e}^{-\kappa t}$$

The optimal convergence rate (see [11] for more details)

$$\kappa = -\lim_{t \to \infty} \frac{1}{t} \log \sup_{x \in E} \|P_t(x, \cdot) - \mu\|_{\operatorname{Var}} = -\lim_{t \to \infty} \frac{1}{t} \log \|P_t - \mu\|_{\infty \to \infty}.$$

The following theorem gives the explicit sufficient conditions for exponential ergodicity and strong ergodicity. The explicit lower bounds for ergodic convergence rates are also obtained.

Theorem 1.1. Assume that $\mu(\mathbb{R}) < \infty$. Consider the following conditions:

(A1) $\int_0^\infty (q + \psi(x))^{-1} dx < \infty$, for any q > 0; (A2) $\int_0^1 (\psi(x))^{-1} dx = \infty$; (A3) $\psi(t)/t \to \infty$, as $t \to \infty$. (1) If (A1)-(A3) hold, and

$$\delta := \sup_{x} H(x)\mu\big((-|x|,|x|)^c\big) < \infty, \tag{1.9}$$

then Y is exponentially ergodic and the L^2 -spectral gap

$$\lambda_1 \geqslant \frac{1}{8\delta}.$$

(2) If (A1)-(A2) hold, and

$$I := \int_{\mathbb{R}} a(x)^{-1} H(|x|) dx < \infty,$$

then Y is strongly ergodic and

$$\kappa \geqslant \frac{1}{2I} > 0.$$

Remark 1.2. (1) For symmetric Lévy process, condition (A1) means that the process is not compound Poisson and the origin is regular for itself (see [9, Section 2]). Condition (A2) means that the process is recurrent (see [19, Section 3.2]). If (A1) and (A2) hold, then X is pointwise recurrent.

(2) (A3) indicates that H is differentiable (see the proof of Theorem 1.1).

(3) Note that a time change does not change the recurrence (cf. [13, Corollary 4.3.7]). Therefore, under the conditions (A1)–(A2), Y is also pointwise recurrent, so Y is Lebesgue irreducible (see [14, Page 42] for the definition). Thus, by [14, Theorem 4.1.1 and Theorem 4.2.1], if $\mu(\mathbb{R}) < \infty$, then Y is ergodic.

(4) Let $\psi(x) = |x|^{\alpha}$, $\alpha \in (1, 2)$. Then Y is a time-changed symmetric α -stable process. It is well known that the process is pointwise recurrent; by [19, Example 1.1], the harmonic function $H(x) = \omega_{\alpha} |x|^{\alpha-1}/2$, where $\omega_{\alpha} = -(\cos(\pi \alpha/2)\Gamma(\alpha))^{-1} > 0$. Then we have Y is exponentially ergodic if

$$\delta_1 := \sup_{x} |x|^{\alpha - 1} \mu((-|x|, |x|)^c) < \infty, \tag{1.10}$$

and

$$\lambda_1 \geqslant \frac{1}{4\omega_\alpha \delta_1}$$

Y is strongly ergodic if

$$I_1 = \int_{\mathbb{R}} \sigma(x)^{-\alpha} |x|^{\alpha - 1} \mathrm{d}x < \infty, \qquad (1.11)$$

and

$$\kappa \geqslant \frac{1}{\omega_{\alpha} I_1} > 0.$$

This case is introduced in [18]. In fact, the conditions (1.10) and (1.11) are sufficient and necessary (see [18] for more details).

Next, we discuss a class of extended α -stable processes, which is introduced in [9]. In general, the cases mean that there exists $\alpha > 0$ such that $\psi(\theta)/\theta^{\alpha}$ is comparable to a non-decreasing function on $(0, \infty)$.

We say that ψ satisfies the global weak lower scaling condition, if there exist $\delta > 0$ and $\beta \in (0, 1]$, such that for $\lambda \ge 1$ and $\theta > 0$,

$$\psi(\lambda\theta) \geqslant \beta\lambda^{\delta}\psi(\theta).$$

In short, we write $\psi \in WLSC(\delta, \beta)$ (see [9] for more details). Applying this condition, we have the following result which is a direct corollary by using Theorem 1.1 and [9, Lemma 2.14]:

Corollary 1.3. Let $\psi^*(x) = \sup_{|u| \leq x} \psi(u)$, $x \geq 0$. Assume that there exists a constant c > 0, such that $\psi \geq c\psi^*$, and $\psi \in WLSC(\delta, \beta)$ ($\delta > 1$). If

$$\sup_{x} \frac{\mu\left((-|x|,|x|)^{c}\right)}{|x|\psi(1/x)} < \infty,$$

then Y is exponentially ergodic and the spectral gap

$$\lambda_1 \ge \frac{\pi(\delta - 1)\beta^2}{10} \inf_x \frac{|x|\psi^*(1/|x|)}{\mu((-|x|, |x|)^c)},$$

If

$$\int_{\mathbb{R}} \frac{1}{|x|a(x)\psi(1/x)|} \mathrm{d}x < \infty,$$

then Y is strongly ergodic and

$$\kappa \ge \frac{\pi\beta^2(\delta-1)}{20\int_{\mathbb{R}} \left(|x|a(x)\psi^*(1/|x|) \right)^{-1} \mathrm{d}x}$$

Now we return to the diffusion operator with stable jump.

Example 1.4. Let $\psi(x) = c_1 x^2 + c_2 |x|^{\alpha}$, where c_1 , c_2 are two constants and $\alpha \in (1, 2)$. Then $L = a(x)(c_1\Delta + c_2\Delta^{\alpha/2})$. Denote by Y the corresponding process with generator L. By (1.4), the harmonic function for $c_1\Delta + c_2\Delta^{\alpha/2}$ is

$$H(x) = \frac{1}{\pi} \int_0^\infty \frac{1 - \cos xs}{c_1 s^2 + c_2 s^\alpha} ds$$

Obviously, (A1)-(A3) hold, $\psi^*(x) = \psi(x) = c_1 x^2 + c_2 |x|^{\alpha}$, and $\psi \in \text{WLSC}(\alpha, 1)$. According to [9, (12)], we have

$$H(x) \leq \frac{10}{\pi(\alpha - 1)} \frac{1}{c_1 |x|^{-1} + c_2 |x|^{1 - \alpha}}.$$

Combining it with Corollary 1.3, if

$$\sup_{x} |x|^{\alpha - 1} \mu((-|x|, |x|)^{c}) < \infty,$$

then Y is exponentially ergodic, and the spectral gap

$$\lambda_1 \ge \frac{\pi(\alpha - 1)}{80} \inf_x \frac{c_1 |x|^{-1} + c_2 |x|^{1 - \alpha}}{\mu((-|x|, |x|)^c)};$$

if

$$\int_{\mathbb{R}} \sigma(x)^{-\alpha} |x|^{\alpha - 1} dx < \infty,$$

then Y is strongly ergodic, and the convergence rate in strong ergodicity

$$\kappa \ge \frac{\pi(\alpha - 1)}{20 \int_{\mathbb{R}} (a(x))^{-1} (c_1 |x|^{-1} + c_2 |x|^{1-\alpha})^{-1} dx}$$

Remark 1.5. Note that if $c_1 = 0$ (resp., $c_2 = 0$), then the result is reduced to the timechanged symmetric stable process (resp., time-changed Brownian motion).

Example 1.6. Let $\psi(x) = x^2 + |x|$. The process X associated with $\Delta + \Delta^{1/2}$ is a sum of the Cauchy process and independent Brownian motion. Obviously, **(A1)**–**(A3)** hold. By [9, Lemma 2.14],

$$H(x) \leq \frac{10}{\pi} \int_{1/|x|}^{\infty} \frac{\mathrm{d}r}{r+r^2} = \frac{10}{\pi} \log(1+|x|).$$

Therefore, if

$$\delta_2 := \sup_x \log(1+|x|)\mu((-|x|,|x|)^c) < \infty,$$

then Y is exponentially ergodic, and the spectral gap

$$\lambda_1 \geqslant \frac{\pi}{80\delta_2}.$$

If

$$I_2 := \int_{\mathbb{R}} a(x)^{-1} \log(1+|x|) dx < \infty,$$

then Y is strongly ergodic and

$$\kappa \geqslant \frac{\pi}{20I_2}.$$

2 Time change and Green potential

Let $X = (X_t)_{t \ge 0}$ be a one-dimensional symmetric Lévy process with Lévy measure ν , transition density $p_t(x, y) = p_t(y - x)$, and characteristic exponent ψ given by (1.1).

Recalling that a is a positive and locally bounded measurable function on \mathbb{R} with $a(x)^{-1}$ is Lebesgue integrable, $A_t = \int_0^t 1/a(X_s)ds$ is the positive continuous additive functional and Y is the time-changed Lévy process defined as (1.6). The Revuz measure μ of A_t with respect to dx, is given by (cf. [7])

$$\mu(f) = \lim_{t \to \infty} \frac{1}{t} \int_{\mathbb{R}} \mathbb{E}_x \left[\int_0^t f(X_s) dA_s \right] dx.$$

Since dx is the invariant measure of X, for nonnegative bounded function f, we have

$$\mu(f) = \lim_{t \to \infty} \frac{1}{t} \int_{\mathbb{R}} \mathbb{E}_x \left[\int_0^t f(X_s) a(X_s)^{-1} \mathrm{d}s \right] \mathrm{d}x = \lim_{t \to \infty} \frac{1}{t} \int_0^t \int_{\mathbb{R}} P_s(fa^{-1})(x) \mathrm{d}x \mathrm{d}s$$
$$= \lim_{t \to \infty} \frac{1}{t} \int_0^t \int_{\mathbb{R}} (fa^{-1})(x) \mathrm{d}x \mathrm{d}s = \int_{\mathbb{R}} (fa^{-1})(x) \mathrm{d}x,$$
(2.1)

thus the Revuz measure $\mu(dx) = a(x)^{-1}dx$. Combining this fact and [5, Theorem 5.2.2, Theorem 5.2.8 and Corollary 5.2.12], similar to [4, Page 2807], we know that Y is μ -symmetric and its Dirichlet form $(\widehat{\mathscr{E}}, \widehat{\mathcal{F}})$ is given by

$$\widehat{\mathscr{E}}(f,g) = \mathscr{E}(f,g), \quad f,g \in \widehat{\mathcal{F}} := \mathcal{F}_e \cap L^2(\mu), \tag{2.2}$$

where \mathcal{F}_e is the extended Dirichlet space of $(\mathscr{E}, \mathcal{F})$, i.e., the family of functions u satisfy that there exists an \mathscr{E} -Cauchy sequence $\{u_n\} \subset \mathcal{F}$ such that for a.e. x, $\lim_{n\to\infty} u_n = u$ in $L^2(\mathrm{d}x)$ and $\mathscr{E}(u, u) = \lim_{n\to\infty} \mathscr{E}(u_n, u_n)$. Therefore, the L^2 infinitesimal generator of Yis $\mathcal{L} = a\mathcal{A}$.

By a similar argument to [1, Corollary 4.2], we can also prove that the extended infinitesimal generator $\widetilde{\mathcal{L}}$ (see [15, Definition 2.1]), is also $a\mathcal{A}$.

Recalling that P_t^0 is the killed semigroup of X given by (1.5). Define the *Green* potential measure $G_X^0(x, A)$ for P_t^0 by

$$G_X^0(x,A) = \int_0^\infty P_t^0(x,A) \mathrm{d}t, \quad \forall x \in \mathbb{R}, \ \forall A \in \mathscr{B}(\mathbb{R}).$$

Denote by $P_t^{0,Y}$ the semigroup of Y killed upon hitting the origin, i.e.

$$P_t^{0,Y}(x,A) = \mathbb{P}_x[Y_t \in A, t < \tau_0^Y],$$

where $\tau_0^Y = \inf\{t > 0 : Y_t \neq 0\}$. Let $G_Y^0(x, A)$ be the Green measure of Y killed upon 0:

$$G_Y^0(x,A) = \int_0^\infty P_t^{0,Y}(x,A) \mathrm{d}t, \quad \forall x \in \mathbb{R}, \ \forall A \in \mathscr{B}(\mathbb{R}).$$

Similar to [18, (14)], for the time-changed process Y,

$$G_Y^0 f(x) = \int_{\mathbb{R}} G_X^0(x, y) a(y)^{-1} \mathrm{d}y.$$
(2.3)

3 Proofs of main results

To finish the proof of Theorem 1.1, we need to consider the first Dirichlet eigenvalue

$$\lambda_0 = \inf\{\mathscr{E}(f,f) : f \in \widehat{\mathcal{F}}, \mu(f^2) = 1 \text{ and } f(0) = 0\},$$
(3.1)

which will play a crucial role in the proof of Theorem 1.1.

First, we introduce the following dual variational inequality, which is mainly motivated by the dual variational formulas for one-dimensional diffusion processes (see [3, Theorem 6.1]).

Lemma 3.1. Let G_Y^0 be the Green operator of Y killed upon $\{0\}$, C(B) be the space of all continuous functions on a measurable set $B \subset \mathbb{R}$. Denote by

$$\mathscr{H} = \{ f : f(0) = 0, f \in C(\mathbb{R} \setminus \{0\}) \}$$

$$\widetilde{\mathscr{H}} = \{ f : f(0) = 0, \text{ there exists } x_0 > 0 \text{ such that } f = f(\cdot \land x_0 \lor (-x_0)), f \in C(-x_0, x_0) \}.$$

Then

$$\inf_{f \in \widetilde{\mathscr{H}}} \sup_{x \in \mathbb{R} \setminus \{0\}} \frac{f(x)}{G_Y^0 f(x)} \ge \lambda_0 \ge \sup_{f \in \mathscr{H}} \inf_{x \in \mathbb{R} \setminus \{0\}} \frac{f(x)}{G_Y^0 f(x)}$$

Proof. First we proof the upper bound. Note that for any $g \in \widetilde{\mathscr{H}}$, there exists $x_0 > 0$ such that $g(-x_0) \leq g \leq g(x_0)$. Therefore, $g \in L^2(\pi)$. By a similar argument to the proof of [18, Theorem 2] (see [18, Page 12]), we have $G_Y^0 g \in \widehat{\mathcal{F}}$, and

$$\mathscr{E}(G_Y^0 g, G_Y^0 g) = \int g G_Y^0 g \mathrm{d}\pi$$

By the definition (3.1),

$$\lambda_0 \leqslant \frac{\mathscr{E}(G_Y^0 g, G_Y^0 g)}{\pi((G_Y^0 g)^2)} \leqslant \sup_{x \in \mathbb{R} \setminus \{0\}} \frac{g(x)}{G_Y^0 g(x)}$$

Now we get the upper bound by the arbitrariness of $g \in \widetilde{\mathscr{H}}$.

Next, denote by

$$\lambda_0^{(n)} = \inf \{ \mathscr{E}(f, f) : \mu(f^2) = 1, f |_{(-\infty, -n) \cup (n, \infty)} = 0 \}.$$

According to [18, Lemma 11],

$$\lim_{n \to \infty} \lambda_0^{(n)} = \lambda_0.$$

By a similar argument to the proof of [18, Lemma 7], for regular set $B_n := (-n, 0) \cup (0, n)$ (see [6, Page 68] for the definition of regular set),

$$\lambda_0^{(n)} \ge \sup_{f \in C_b(B_n)} \inf_{x \in B_n} \frac{f(x)}{G_Y^{B_n} f(x)},$$

where $G_Y^{B_n}$ is the Green operator defined as $G_Y^{B_n}(x, A) := \int_0^\infty \mathbb{P}_x[Y_t \in A, t < \tau_{B_n}^Y] dt, \ \tau_{B_n}^Y$ is the exit time from B_n : $\tau_{B_n}^Y := \inf\{t \ge 0 : Y_t \notin B_n\}.$

Next, since $G_Y^{B_n} \varphi \leq G_Y^{\{0\}^c} \varphi$ and $f \in C(\mathbb{R} \setminus \{0\})$ is bounded on B_n , we have that for any $f \in \mathscr{H}$,

$$\lambda_0 = \lim_{n \to \infty} \lambda_0^{(n)} \ge \lim_{n \to \infty} \inf_{x \in B_n} \frac{f(x)}{G_Y^0 f(x)} \ge \inf_{x \in \mathbb{R} \setminus \{0\}} \frac{f(x)}{G_Y^0 f(x)},\tag{3.2}$$

thus we obtain the lower bound.

The following explicit estimates for lower and upper bounds of λ_0 is similar to [18, Theorem 2], and can be obtained directly by Lemma 3.1.

Theorem 3.2. Let δ be defined by (1.9) and

$$\delta^+ = \sup_{x>0} H(x)\mu((x,\infty)), \quad \delta^- = \sup_{x<0} H(x)\mu((-\infty,-x)).$$

Then

$$\frac{1}{\delta_{+}} + \frac{1}{\delta_{-}} \geqslant \lambda_0 \geqslant \frac{1}{8\delta}.$$
(3.3)

Proof. First, we prove the upper bound. According to Lemma 3.1, for any $g \in \widetilde{\mathscr{H}}$,

$$\lambda_0 \leqslant \sup_{x \in \mathbb{R} \setminus \{0\}} \frac{g(x)}{G_Y^0 g(x)} \leqslant \sup_{x>0} \frac{g(x)}{G_Y^0 g(x)} + \sup_{x<0} \frac{g(x)}{G_Y^0 g(x)}.$$
(3.4)

Note that by [9, Proposition 2.4], for xy > 0,

$$G_Y^0(x,y) \ge H(|x| \land |y|),$$

so we get that for x > 0,

$$G_Y^0 g(x) \ge \int_0^\infty H(x \wedge y) g(y) a(y)^{-1} \mathrm{d}y \ge H(x) \int_x^\infty g(y) a(y)^{-1} \mathrm{d}y.$$
 (3.5)

Since $g \in \widetilde{\mathscr{H}}$, there exists $x_0 > 0$ such that $g(x) = g(x \wedge x_0 \vee (-x_0))$. Now by choosing $g(x) = H(x \wedge x_0 \vee (-x_0))$, we obtain that for x > 0,

$$\frac{G_Y^0 g(x)}{g(x)} = \frac{G_Y^0 g(x \wedge x_0)}{g(x \wedge x_0)} \ge H(x_0) \int_{x_0}^{\infty} a(y)^{-1} \mathrm{d}y.$$
(3.6)

Similarly, for x < 0,

$$\frac{G_Y^0 g(x)}{g(x)} = \frac{G_Y^0 g(x \vee (-x_0))}{g(x \vee (-x_0))} \ge H(-x_0) \int_{-\infty}^{-x_0} a(y)^{-1} \mathrm{d}y.$$
(3.7)

Therefore, by combining (3.4), (3.6), and (3.7), we obtain the estimate for upper bound.

Next, we consider the lower bound. Let G^0 be the Green operator of Y killed upon $\mathbb{R} \setminus \{0\}$. According to [9, Proposition 2.3 and 2.4],

$$G_X^0(x,y) = H(x) + H(y) - H(y-x) \le 2(H(x) \land H(y)).$$
(3.8)

By the property of time change and (3.8), for any f with $G_Y^0|f| < \infty$,

$$G_Y^0 f(x) \leqslant \int_{\mathbb{R}} 2(H(y) \wedge H(x)) f(y) a(y)^{-1} \mathrm{d}y.$$
(3.9)

Since $\psi(t)/t \to \infty$ as $t \to \infty$, by Dirichlet criterion, $\int_0^\infty t \sin(xt)\psi(t)^{-1}dt$ is integrable. Therefore, *H* is differentiable, and

$$H'(x) = \frac{1}{\pi} \int_0^\infty \frac{t \sin xt}{\psi(t)} dt < \infty.$$

Recalling that $p_t(x) = (2\pi)^{-1} \int_{\mathbb{R}} e^{-t\psi(\xi) - i\xi x} d\xi$ and $\psi(\xi) = \psi(-\xi)$. Since

$$\int_{\mathbb{R}} e^{-t\psi(\xi)} d\xi = p_t(0) = p_t(x, x) < \infty,$$

 $e^{-t\psi} \in L^1(dx)$. By [8, Theorem 1.1], for $|\eta| = r$,

$$-\frac{1}{2\pi r}\frac{\mathrm{d}}{\mathrm{d}r}p_t(r) = \int_{\mathbb{R}^3} \mathrm{e}^{-t\psi(|x|)} \mathrm{e}^{-2\pi i x \cdot \eta} \mathrm{d}x = \frac{1}{2\pi}q_t(\eta),$$

where $q_t(\eta)$ is the transition density of a 3-dimensional Lévy process with characteristic exponent $\Psi(\eta) = \psi(|\eta|)$. Therefore, $q_t(\eta) \ge 0$, $p_t(x)$ is non-increasing for x, i.e. it is unimodal (see [9]). Thus by [9], H(x) is non-increasing on $(0, \infty)$ and non-decreasing on $(-\infty, 0)$, so we have

$$G_Y^0 f(x) \leq 2 \left(\int_{\mathbb{R} \setminus (-|x|, |x|)} H(x) f(y) \mu(dy) + \int_{-|x|}^{|x|} H(y) f(y) \mu(dy) \right)$$

= $2 \int_0^{|x|} H'(z) \left(\int_{\mathbb{R} \setminus (-z, z)} f(y) \mu(dy) \right) dz.$ (3.10)

Choose $f(x) = \sqrt{H(x)}$. By using integration by parts, for any y > 0,

$$\int_{\mathbb{R}\setminus(-y,y)} \sqrt{H(z)} \mu(dz) \leqslant \sqrt{H(y)} \mu((-y,y)^c) + \int_y^\infty \frac{H'(z)\mu((-z,z)^c)}{2\sqrt{H(z)}} dz.$$

Since $\delta < \infty$, then

$$\int_{\mathbb{R}\setminus(-y,y)}\sqrt{H(z)}\mu(dz) \leqslant \frac{\delta}{\sqrt{H(y)}} + \frac{\delta}{2}\int_{y}^{\infty}H(z)^{-3/2}dH(z) = \frac{2\delta}{\sqrt{H(y)}}.$$

Thus

$$\frac{G_Y^0\sqrt{H}}{\sqrt{H}}(x) \leqslant \frac{2}{\sqrt{H(x)}} \int_0^{|x|} H'(z) \frac{2\delta}{\sqrt{H(z)}} dz = 8\delta.$$

Now by Lemma 3.1 and letting $f = \sqrt{H}$, we obtain that if $\delta < \infty$, then

$$\lambda_0 \ge \inf_{x \neq 0} \frac{\sqrt{H(x)}}{G_Y^0 \sqrt{H(x)}} \ge \frac{1}{8\delta}.$$

Proof of Theorem 1.1.

(1) By [2, Proposition 3.2], $\lambda_1 \ge \lambda_0$. Hence

$$\lambda_1 \geqslant \lambda_0 \geqslant \frac{1}{8\delta},$$

and Y is exponential ergodicity.

(2) Specially, by choosing $f \equiv 1$ in (3.9), we have

$$M_{0} := \sup_{x} \mathbb{E}_{x} \tau_{0}^{Y} = \sup_{x} G_{Y}^{0} 1(x) \leq \int_{\mathbb{R}} 2H(y)a(y)^{-1}dy < \infty,$$

thus by [12, Theorem 1.2(R2)], $\kappa \ge M_0^{-1} \ge (2I)^{-1} > 0$.

Proof of Corollary 1.3.

First, according to [9, Lemma 2.14], $H(x) \approx (|x|\psi(1/x))^{-1}$, thus by using Theorem 1.1, we obtain the exponential ergodicity and strong ergodicity.

By [9, Lemma 2.14 and (12)],

$$H(x) \leq \frac{10}{\pi\beta} \int_{1/x}^{\infty} \frac{1}{\psi^*(s)} ds \leq \frac{10}{\pi\beta^2(\alpha - 1)x\psi^*(1/x)}, \text{ for } x > 0.$$

Then the estimates for λ_1 and κ follow from Theorem 1.1.

Acknowledgements

The author thanks Prof. Yong-Hua Mao and Prof. Jian Wang for valuable conversations of this paper. The author would also like to thank Mr. Zhi-Feng Wei for his helpful suggestions. This work is supported in part by the National Key Research and Development Program of China (2020YFA0712900), the National Natural Science Foundation of China (Grant No.11771047), and the project from the Ministry of Education in China.

References

- B. Bottcher, R. Schilling, and J. Wang. Lévy-type processes: construction, approximation and sample path properties. From: Lévy matters III, volume 2099 of Lecture Notes in Mathematics, with a short biography of Paul Lévy by Jean Jacod. Berlin: Springer, 2013.
- [2] M. F. Chen. Explicit bounds for the first eigenvalues. Sci. in China Ser. A, 43:1051– 1059, 2000.
- [3] M.F. Chen. *Eigenvalues, inequalities, and ergodic theory*. London: Springer, 2004.
- [4] Z. Q. Chen and J. Wang. Ergodicity for time-changed symmetric stable processes. Stoch. Proc. Appl., 124(9):2799–2823, 2014.
- [5] Z.Q. Chen and M. Fukushima. Symmetric Markov processes, time change, and boundary theory. Princeton: Princeton Univ. Press, 2012.

- [6] K.L. Chung. Doubly-Feller process with multiplicative functional. In Seminar on Stochastic Processes, Birkhäuser., 12:63–78, 1986.
- [7] P. J. Fitzsimmons and R. K.Getoor. Revuz measures and time changes. Math. Z., 199(2):233-256, 1988.
- [8] L. Grafakos and G. Teschl. On Fourier transforms of radial functions and distributions. J. Fourier Anal. Appl, 19(1):167–179, 2013.
- [9] T. Grzywny and M.Ryznar. Hitting times of points and intervals for symmetric Lévy processes. *Potential Anal.*, 46:739–777, 2017.
- [10] D. Luo and J. Wang. Refined basic couplings and Wasserstein-type distances for SDEs with Levy noises. Stoch. Proc. Appl., 129:3129—-3173, 2019.
- [11] Y.H. Mao. Convergence rates in strong ergodicity for Markov processes. Stoch. Proc. Appl., 116:1964–1976, 2006.
- [12] Y.H. Mao and T. Wang. Convergence rates in strong ergodicity by hitting times and L²-exponential convergence rates. see ArXiv 2102.07069.
- [13] Y. Oshima. Semi-Dirichlet forms and Markov processes. Berlin: De Gruyter Studies in Mathematics., 2013.
- [14] G. Da Prato and J. Zabczyk. Ergodicity for infinite-dimensional systems. Cambridge: Cambridge University Press, 1996.
- [15] Y. Shiozawa and M. Takeda. Variational formula for Dirichlet forms and estimates of principal eigenvalues for symmetric α -stable processes. *Potent. Anal.*, 23:135–151, 2005.
- [16] J. Wang. Criteria for ergodicity of lévy type operators in dimension one. Stoch. Proc. Appl., 118:1909–1928, 2008.
- [17] J. Wang. Exponential ergodicity and strong ergodicity for SDEs driven by symmetric α -stable processes. Appl. Math. Lett., 26:654–658, 2013.
- [18] T. Wang. Exponential and strong ergodicity for one-dimensional symmetric stable jump diffusions. See ArXiv: 2104.07947, 2021.
- [19] K. Yano. Excursions away from a regular point for one-dimensional symmetric Lévy processes without Gaussian part. *Potential Anal.*, 32(4):305–341, 2010.