

\mathcal{O} -OPERATORS ON LIE ∞ -ALGEBRAS WITH RESPECT TO LIE ∞ -ACTIONS

RAQUEL CASEIRO AND JOANA NUNES DA COSTA

ABSTRACT. We define \mathcal{O} -operators on a Lie ∞ -algebra E with respect to an action of E on another Lie ∞ -algebra and we characterize them as Maurer-Cartan elements of a certain Lie ∞ -algebra obtained by Voronov's higher derived brackets construction. The Lie ∞ -algebra that controls the deformation of \mathcal{O} -operators with respect to a fixed action is determined.

INTRODUCTION

The first instance of Rota-Baxter operator appeared in the context of associative algebras in 1960, in a paper by Baxter [1], as a tool to study fluctuation theory in probability. Since then, these operators were widely used in many branches of mathematics and mathematical physics.

Almost forty years later, Kupersmidt [4] introduced \mathcal{O} -operators on Lie algebras as a kind of generalization of classical r -matrices, thus opening a broad application of \mathcal{O} -operators to integrable systems. Given a Lie algebra $(E, [\cdot, \cdot])$ and a representation Φ of E on a vector space V , an \mathcal{O} -operator on E with respect to Φ is a linear map $T : V \rightarrow E$ such that $[T(x), T(y)] = T(\Phi(T(x))(y) - \Phi(T(y))(x))$. When Φ is the adjoint representation of E , T is a Rota-Baxter operator (of weight zero). \mathcal{O} -operators are also called relative Rota-Baxter operators or generalized Rota-Baxter operators.

In recent years Rota-Baxter and \mathcal{O} -operators, in different algebraic and geometric settings, have deserved a great interest by mathematical and physical communities.

In [9], a homotopy version of \mathcal{O} -operators on symmetric graded Lie algebras was introduced. This was the first step towards the definition of an \mathcal{O} -operator on a Lie ∞ -algebra with respect to a representation on a graded vector space that was given in [6]. The current paper also deals with \mathcal{O} -operators on Lie ∞ -algebras, but with a different approach which uses Lie ∞ -actions instead of representations of Lie ∞ -algebras. Our definition is therefore different from the one given in [6] but there is a relationship between them.

There are two equivalent definitions of Lie ∞ -algebra structure on a graded vector space E , both given by collections of n -ary brackets which are either symmetric or skew-symmetric, depending on the definition we are considering, and must satisfy a kind of generalized Jacobi identities. One goes from one to the other by shifting the degree of E and applying a *décalage* isomorphism. We use the definition in its symmetric version, where the brackets have degree $+1$. Equivalently, this structure can be defined by a degree $+1$ coderivation M_E of $\tilde{S}(E)$, the reduced symmetric algebra of E , such that the commutator $[M_E, M_E]_c$ vanishes.

Representations of Lie ∞ -algebras on graded vector spaces were introduced in [7]. In [6], the authors consider a representation Φ of a Lie ∞ -algebra E on a graded

2020 *Mathematics Subject Classification.* 17B10, 17B40, 17B70, 55P43.

Key words and phrases. Lie ∞ -algebra, \mathcal{O} -operator, Maurer Cartan element.

The authors are partially supported by the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES.

vector space V and define an \mathcal{O} -operator (homotopy relative Rota-Baxter operator) on E with respect to Φ as a degree zero element T of $\text{Hom}(\bar{S}(V), E)$ satisfying a family of suitable identities. Inspired by the notion of an action of a Lie ∞ -algebra on a graded manifold [8], we define an action of a Lie ∞ -algebra (E, M_E) on a Lie ∞ -algebra (V, M_V) as a Lie ∞ -morphism Φ between E and $\text{Coder}(\bar{S}(V))[1]$, the symmetric DGLA of coderivations of $\bar{S}(V)$. An \mathcal{O} -operator on E with respect to the action Φ is a comorphism between $\bar{S}(V)$ and $\bar{S}(E)$ that intertwines the coderivation M_E and a degree +1 coderivation of $\bar{S}(V)$ built from M_V and Φ , which turns out to be a Lie ∞ -algebra structure on V too.

As we said before, the two \mathcal{O} -operator definitions, ours and the one in [6], are different. However, since there is a close connection between Lie ∞ -actions and representations of Lie ∞ -algebras, the two definitions can be related. On the one hand, any representation of (E, M_E) on a complex (V, d) can be seen as a Lie ∞ -action of (E, M_E) on (V, D) , with D the coderivation given by the differential d , and for this very “simple” Lie ∞ -algebra structure on V our \mathcal{O} -operator definition recovers the one given in [6]. On the other hand, any action Φ of (E, M_E) on (V, M_V) yields a representation ρ on the graded vector space $\bar{S}(V)$ and an \mathcal{O} -operator with respect to the action Φ is not the same as an \mathcal{O} -operator with respect to the representation ρ . However, there is a way to relate the two concepts.

A well-known Voronov’s construction [10] defines a Lie ∞ -algebra structure on an abelian Lie subalgebra \mathfrak{h} of $\text{Coder}(\bar{S}(E \oplus V))$ and we show that \mathcal{O} -operators with respect to the action Φ are Maurer-Cartan elements of \mathfrak{h} .

In general, deformations of structures and morphisms are governed by DGLA’s or, more generally, by Lie ∞ -algebras. We do not intend to deeply study deformations of \mathcal{O} -operators on Lie ∞ -algebras with respect to Lie ∞ -actions. Still, we prove that deformations of an \mathcal{O} -operator are controlled by the twisting of a Lie ∞ -algebra, constructed out of a graded Lie subalgebra of $\text{Coder}(\bar{S}(E \oplus V))$.

The paper is organized in four sections. In Section 1 we collect some basic results on graded vector spaces, graded symmetric algebras and Lie ∞ -algebras that will be needed along the paper. In Section 2, after recalling the definition of a representation of a Lie ∞ -algebra on a complex (V, d) [7], we introduce the notion of action of a Lie ∞ -algebra on another Lie ∞ -algebra (Lie ∞ -action) and we prove that a Lie ∞ -action of E on V induces a Lie ∞ -algebra structure on $E \oplus V$. We pay special attention to the adjoint action of a Lie ∞ -algebra. In Section 3 we introduce the main notion of the paper – \mathcal{O} -operator on a Lie ∞ -algebra E with respect to an action of E on another Lie ∞ -algebra, and we give the explicit relation between these operators and \mathcal{O} -operators on E with respect to a representation on a graded vector space introduced in [6]. Given an \mathcal{O} -operator T on E with respect to a Lie ∞ -action Φ on V , we show that V inherits a new Lie ∞ -algebra structure given by a degree +1 coderivation which is the sum of the initial one on V with a degree +1 coderivation obtained out of Φ and T . We prove that symmetric and invertible comorphisms $T : \bar{S}(E^*) \rightarrow S(E)$ are \mathcal{O} -operators with respect to the coadjoint action if and only if a certain element of $\bar{S}(E^*)$, which is defined using the inverse of T , is a cocycle for the Lie ∞ -algebra cohomology of E . Section 3 ends with the characterization of \mathcal{O} -operators as Maurer-Cartan elements of a Lie ∞ -algebra obtained by Voronov’s higher derived brackets construction. The main result in Section 4 shows that Maurer-Cartan elements of a graded Lie subalgebra of $\text{Coder}(\bar{S}(E \oplus V))$ encode a Lie ∞ -algebra on E and an action of E on V . Moreover, we obtain the Lie ∞ -algebra that controls the deformation of \mathcal{O} -operators with respect to a fixed action.

1. LIE ∞ -ALGEBRAS

We begin by reviewing some concepts about graded vector spaces, graded symmetric algebras and Lie ∞ -algebras.

1.1. Graded vector spaces and graded symmetric algebras. We will work with \mathbb{Z} -graded vector spaces with finite dimension over a field $\mathbb{K} = \mathbb{R}$ or \mathbb{C} .

Let $E = \bigoplus_{i \in \mathbb{Z}} E_i$ be a finite dimensional graded vector space. We call E_i the homogeneous component of E of degree i . An element x of E_i is said to be homogeneous with degree $|x| = i$. For each $k \in \mathbb{Z}$, one may shift all the degrees by k and obtain a new grading on E . This new graded vector space is denoted by $E[k]$ and is defined by $E[k]_i = E_{i+k}$.

A morphism $\Phi : E \rightarrow V$ between two graded vector spaces is a degree preserving linear map, i.e. a collection of linear maps $\Phi_i : E_i \rightarrow V_i$, $i \in \mathbb{Z}$. We call $\Phi : E \rightarrow V$ a (homogeneous) morphism of degree k , for some $k \in \mathbb{Z}$, and we write $|\Phi| = k$, if it is a morphism between E and $V[k]$. This way we have a natural grading in the vector space of linear maps between graded vector spaces:

$$\text{Hom}(E, V) = \bigoplus_{i \in \mathbb{Z}} \text{Hom}_i(E, V).$$

In particular, $\text{Hom}(E, E) = \text{End}(E) = \bigoplus_{i \in \mathbb{Z}} \text{End}_i(E)$.

The dual E^* of E is naturally a graded vector space whose component of degree i is, for all $i \in \mathbb{Z}$, the dual $(E_{-i})^*$ of E_{-i} . In equation: $(E^*)_i = (E_{-i})^*$.

Given two graded vector spaces E and V , their direct sum $E \oplus V$ is a vector space with grading

$$(E \oplus V)_i = E_i \oplus V_i$$

and their usual tensor product comes equipped with the grading

$$(E \otimes V)_i = \bigoplus_{j+k=i} E_j \otimes V_k.$$

We will adopt the Koszul sign convention, for homogeneous linear maps $f : E \rightarrow V$ and $g : F \rightarrow W$ the tensor product $f \otimes g : E \otimes F \rightarrow V \otimes W$ is the morphism of degree $|f| + |g|$ given by

$$(f \otimes g)(x \otimes y) = (-1)^{|x||g|} f(x) \otimes g(y),$$

for all homogeneous $x \in E$ and $y \in F$.

For each $k \in \mathbb{N}_0$, let $T^k(E) = \bigotimes^k E$, with $T^0(E) = \mathbb{K}$, and let $T(E) = \bigoplus_k T^k(E)$ be the tensor algebra over E . The **graded symmetric algebra over E** is the quotient

$$S(E) = T(E) / \langle x \otimes y - (-1)^{|x||y|} y \otimes x \rangle.$$

The symmetric algebra $S(E) = \bigoplus_{k \geq 0} S^k(E)$ is a graded commutative algebra, whose product we denote by \odot . For $x = x_1 \odot \dots \odot x_k \in S^k(E)$, we set $|x| = \sum_{i=1}^k |x_i|$.

For $n \geq 1$, let S_n be the permutation group of order n . For any homogeneous elements $x_1, \dots, x_n \in E$ and $\sigma \in S_n$, the Koszul sign is the element in $\{-1, 1\}$ defined by

$$x_{\sigma(1)} \odot \dots \odot x_{\sigma(n)} = \epsilon(\sigma) x_1 \odot \dots \odot x_n.$$

As usual, writing $\epsilon(\sigma)$ is an abuse of notation because the Koszul sign also depends on the x_i .

An element σ of S_n is called an $(i, n-i)$ -unshuffle if $\sigma(1) < \dots < \sigma(i)$ and $\sigma(i+1) < \dots < \sigma(n)$. The set of $(i, n-i)$ -unshuffles is denoted by $Sh(i, n-i)$. Similarly, $Sh(k_1, \dots, k_j)$ is the set of (k_1, \dots, k_j) -unshuffles, i.e., elements of S_n with $k_1 + \dots + k_j = n$ such that the order is preserved within each block of length k_i , $1 \leq i \leq j$.

The reduced symmetric algebra $\bar{S}(E) = \oplus_{k \geq 1} S^k(E)$ has a natural coassociative and cocommutative coalgebra structure given by the coproduct $\Delta : \bar{S}(E) \rightarrow \bar{S}(E) \otimes \bar{S}(E)$,

$$\Delta(x) = 0, \quad x \in E;$$

$$\Delta(x_1 \odot \dots \odot x_n) = \sum_{i=1}^{n-1} \sum_{\sigma \in Sh(i, n-i)} \epsilon(\sigma) (x_{\sigma(1)} \odot \dots \odot x_{\sigma(i)}) \otimes (x_{\sigma(i+1)} \odot \dots \odot x_{\sigma(n)}),$$

for $x_1, \dots, x_n \in E$.

We will mainly use Sweedler notation: given $x \in \bar{S}(E)$,

$$\Delta^{(1)}(x) = \Delta(x) = x_{(1)} \otimes x_{(2)},$$

and the coassociativity yields

$$\Delta^{(n)}(x) = (\text{id} \otimes \Delta^{(n-1)})\Delta(x) = x_{(1)} \otimes \dots \otimes x_{(n+1)}, \quad n \geq 2.$$

Notice that

$$\Delta^{(n)}(x) = 0, \quad x \in S^{\leq n}(E).$$

The cocommutativity of the coproduct is expressed, for homogeneous elements of $\bar{S}(E)$, as

$$x_{(1)} \otimes x_{(2)} = (-1)^{|x_{(1)}||x_{(2)}|} x_{(2)} \otimes x_{(1)}.$$

Let V be another graded vector space. A linear map $f : \bar{S}(E) \rightarrow V$ is given by a collection of maps $f_k : S^k(E) \rightarrow V$, $k \geq 1$, and is usually denoted by $f = \sum_k f_k$.

Remark 1.1. Every linear map $f : S^k(E) \rightarrow V$ corresponds to a graded symmetric k -linear map $f \in \text{Hom}(\otimes^k E, V)$ through the quotient map $p_k : \otimes^k E \rightarrow S^k(E)$ i.e., $f \equiv f \circ p_k$. In the sequel, we shall often write

$$f(x_1 \odot \dots \odot x_k) = f(x_1, \dots, x_k), \quad x_i \in E.$$

A **coalgebra morphism** (or **comorphism**) between the coalgebras $(\bar{S}(E), \Delta_E)$ and $(\bar{S}(V), \Delta_V)$ is a morphism $F : \bar{S}(E) \rightarrow \bar{S}(V)$ of graded vector spaces such that

$$(F \otimes F) \circ \Delta_E = \Delta_V \circ F.$$

There is a one-to-one correspondence between coalgebra morphisms $F : \bar{S}(E) \rightarrow \bar{S}(V)$ and degree preserving linear maps $f : \bar{S}(E) \rightarrow V$. Each f determines F by

$$F(x) = \sum_{k \geq 1} \frac{1}{k!} f(x_{(1)}) \odot \dots \odot f(x_{(k)}), \quad x \in \bar{S}(E),$$

and $f = p_V \circ F$, with $p_V : \bar{S}(V) \rightarrow V$ the projection map.

A degree k **coderivation** of $\bar{S}(E)$, for some $k \in \mathbb{Z}$, is a linear map $Q : \bar{S}(E) \rightarrow \bar{S}(E)$ of degree k such that

$$\Delta \circ Q = (Q \otimes \text{id} + \text{id} \otimes Q) \circ \Delta.$$

We also have a one to one correspondence between coderivations of $\bar{S}(E)$ and linear maps $q = \sum_i q_i : \bar{S}(E) \rightarrow E$:

Proposition 1.2. *Let E be a graded vector space and $p_E : \bar{S}(E) \rightarrow E$ the projection map. For every linear map $q = \sum_i q_i : \bar{S}(E) \rightarrow E$, the linear map $Q : \bar{S}(E) \rightarrow \bar{S}(E)$ given by*

$$(1) \quad Q(x_1 \odot \dots \odot x_n) = \sum_{i=1}^n \sum_{\sigma \in Sh(i, n-i)} \epsilon(\sigma) q_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}) \odot x_{\sigma(i+1)} \odot \dots \odot x_{\sigma(n)},$$

is the unique coderivation of $\bar{S}(E)$ such that $p_E \circ Q = q$.

In Sweedler notation, Equation (1) is written as:

$$Q(x) = q(x_{(1)}) \odot x_{(2)} + q(x), \quad x \in \bar{S}(E).$$

When E is a finite dimensional graded vector space, we may identify $S(E^*)$ with $(SE)^*$. Koszul sign conventions yield, for each homogeneous elements $f, g \in E^*$,

$$(f \odot g)(x \odot y) = (-1)^{|x||g|} f(x) g(y) + f(y) g(x), \quad x, y \in E.$$

1.2. Lie ∞ -algebras. We briefly recall the definition of Lie ∞ -algebra [5], some basic examples and related concepts.

We will consider the symmetric approach to Lie ∞ -algebras.

Definition 1.3. A **symmetric Lie ∞ -algebra** (or a **Lie[1] ∞ -algebra**) is a graded vector space $E = \oplus_{i \in \mathbb{Z}} E_i$ together with a family of degree +1 linear maps $l_k : S^k(E) \rightarrow E$, $k \geq 1$, satisfying

$$(2) \quad \sum_{i+j=n+1} \sum_{\sigma \in Sh(i, j-1)} \epsilon(\sigma) l_j(l_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)}, \dots, x_{\sigma(n)}) = 0,$$

for all $n \in \mathbb{N}$ and all homogeneous elements $x_1, \dots, x_n \in E$.

The *décalage* isomorphism [10] establishes a one to one correspondence between skew-symmetric Lie ∞ -algebra structures $\{l'_k\}_{k \in \mathbb{N}}$ on E and symmetric Lie ∞ -algebra structures $\{l_k\}_{k \in \mathbb{N}}$ on $E[1]$:

$$l_k(x_1, \dots, x_k) = (-1)^{(k-1)|x_1| + (k-2)|x_2| + \dots + |x_{k-1}|} l'_k(x_1, \dots, x_k).$$

In the sequel, we frequently write Lie ∞ -algebra, omitting the term symmetric.

Example 1.4 (Symmetric graded Lie algebra). A symmetric graded Lie algebra is a symmetric Lie ∞ -algebra $E = \oplus_{i \in \mathbb{Z}} E_i$ such that $l_n = 0$ for $n \neq 2$. Then the degree 0 bilinear map on $E[-1]$ defined by

$$(3) \quad \llbracket x, y \rrbracket := (-1)^i l_2(x, y), \quad \text{for all } x \in E_i, y \in E_j,$$

is a graded Lie bracket. In particular, if $E = E_{-1}$ is concentrated on degree -1 , we get a Lie algebra structure.

Example 1.5 (Symmetric DGLA algebra). A symmetric differential graded Lie algebra (DGLA) is a symmetric Lie ∞ -algebra $E = \oplus_{i \in \mathbb{Z}} E_i$ such that $l_n = 0$ for $n \neq 1$ and $n \neq 2$.

Then, from (2), we have that $d := l_1$ is a degree +1 linear map $d : E \rightarrow E$ squaring zero and satisfies the following compatibility condition with the bracket $[\cdot, \cdot] := l_2(\cdot, \cdot)$:

$$\begin{cases} d[x, y] + [d(x), y] + (-1)^{|x|} [x, d(y)] = 0, \\ [[x, y], z] + (-1)^{|y||z|} [[x, z], y] + (-1)^{|x|} [x, [y, z]] = 0, \end{cases}$$

Applying the *décalage* isomorphism, $(E[-1], d, \llbracket \cdot, \cdot \rrbracket)$ is a (skew-symmetric) DGLA, with $\llbracket \cdot, \cdot \rrbracket$ given by (3).

Example 1.6. Let $(E = \oplus_{i \in \mathbb{Z}} E_i, d)$ be a cochain complex. Then $\text{End}(E)[1]$ has a natural symmetric DGLA structure with $l_1 = \partial$, $l_2 = [\cdot, \cdot]$ given by:

$$\begin{cases} \partial \phi = -d \circ \phi + (-1)^{|\phi|+1} \phi \circ d, \\ [\phi, \psi] = (-1)^{|\phi|+1} (\phi \circ \psi - (-1)^{(|\phi|+1)(|\psi|+1)} \psi \circ \phi), \end{cases}$$

for ϕ, ψ homogeneous elements of $\text{End}(E)[1]$. In other words, $\partial \phi = -[d, \phi]_c$ and $[\phi, \psi] = (-1)^{\deg(\phi)} [\phi, \psi]_c$, with $[\cdot, \cdot]_c$ the graded commutator on $\text{End}(E)$ and $\deg(\phi)$ the degree of ϕ in $\text{End}(E)$.

The symmetric Lie bracket $[\cdot, \cdot]$ on $\text{End}(\bar{S}(E))[1]$ preserves $\text{Coder}(\bar{S}(E))[1]$, the space of coderivations of $\bar{S}(E)$, so that $(\text{Coder}(\bar{S}(E))[1], \partial, [\cdot, \cdot])$ is a symmetric DGLA.

The isomorphism between $\text{Hom}(\bar{S}(E), E)$ and $\text{Coder}(\bar{S}(E))$ given by Proposition 1.2, induces a Lie bracket on $\text{Hom}(\bar{S}(E), E)$ known as the Richardson-Nijenhuis bracket:

$$[f, g]_{RN}(x) = f(G(x)) - (-1)^{|f||g|}g(F(x)), \quad x \in \bar{S}(E),$$

for each $f, g \in \text{Hom}(\bar{S}(E), E)$, where F and G denote the coderivations defined by f and g , respectively. In other words, $[F, G]_c$ is the (unique) coderivation of $\bar{S}(E)$ determined by $[f, g]_{RN} \in \text{Hom}(\bar{S}(E), E)$.

Degree +1 elements $l := \sum_k l_k$ of $\text{Hom}(\bar{S}(E), E)$ satisfying $[l, l]_{RN} = 0$ define a Lie ∞ -algebra structure on E . This way we have an alternative definition of Lie ∞ -algebra [5]:

Proposition 1.7. *A Lie ∞ -algebra is a graded vector space E equipped with a degree +1 coderivation M_E of $\bar{S}(E)$ such that*

$$[M_E, M_E]_c = 2M_E^2 = 0.$$

The dual of the coderivation M_E yields a differential d_* on $\bar{S}(E^*)$. The **cohomology of the Lie ∞ -algebra** $(E, M_E \equiv \{l_k\}_{k \in \mathbb{N}})$ is the cohomology defined by the differential d_* .

A **Maurer-Cartan element** of a Lie ∞ -algebra $(E, \{l_k\}_{k \in \mathbb{N}})$ is a degree zero element z of E such that

$$(4) \quad \sum_{k \geq 1} \frac{1}{k!} l_k(z, \dots, z) = 0.$$

The set of Maurer-Cartan elements of E is denoted by $\text{MC}(E)$. Let z be a Maurer-Cartan element of $(E, \{l_k\}_{k \in \mathbb{N}})$ and set, for $k \geq 1$,

$$(5) \quad l_k^z(x_1, \dots, x_k) := \sum_{i \geq 0} \frac{1}{i!} l_{k+i}(z, \dots, z, x_1, \dots, x_k).$$

Then, $(E, \{l_k^z\}_{k \in \mathbb{N}})$ is a Lie ∞ -algebra, called the *twisting of E by z* [3]. For filtered, or even weakly filtered Lie ∞ -algebras, the convergence of the infinite sums defining Maurer-Cartan elements and twisted Lie ∞ -algebras (Equations (4) and (5)) is guaranteed (see [3, 2, 6]).

For a symmetric graded Lie algebra (E, l_2) , the twisting by $z \in \text{MC}(E)$ is the symmetric DGLA $(E, l_1^z = l_2(z, \cdot), l_2^z = l_2)$.

1.3. Lie ∞ -morphisms. A morphism of Lie ∞ -algebras is a morphism between symmetric coalgebras that is compatible with the Lie ∞ -structures.

Definition 1.8. Let $(E, \{l_k\}_{k \in \mathbb{N}})$ and $(V, \{m_k\}_{k \in \mathbb{N}})$ be Lie ∞ -algebras. A **Lie ∞ -morphism** $\Phi : E \rightarrow V$ is given by a collection of degree zero linear maps:

$$\Phi_k : S^k(E) \rightarrow V, \quad k \geq 1,$$

such that, for each $n \geq 1$,

$$\begin{aligned}
 (6) \quad & \sum_{\substack{k+l=n \\ \sigma \in Sh(k,l) \\ l \geq 0, k \geq 1}} \varepsilon(\sigma) \Phi_{1+l} \left(l_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), x_{\sigma(k+1)}, \dots, x_{\sigma(n)} \right) \\
 &= \sum_{\substack{k_1+\dots+k_j=n \\ \sigma \in Sh(k_1, \dots, k_j)}} \frac{\varepsilon(\sigma)}{j!} m_j \left(\Phi_{k_1}(x_{\sigma(1)}, \dots, x_{\sigma(k_1)}), \Phi_{k_2}(x_{\sigma(k_1+1)}, \dots, x_{\sigma(k_1+k_2)}), \dots, \right. \\
 & \quad \left. \Phi_{k_j}(x_{\sigma(k_1+\dots+k_{j-1}+1)}, \dots, x_{\sigma(n)}) \right),
 \end{aligned}$$

If $\Phi_k = 0$ for $k \neq 1$, then Φ is called a **strict Lie ∞-morphism**.

A **curved Lie ∞-morphism** $E \rightarrow V$, with V a weakly filtered Lie ∞-algebra, is a degree zero linear map $\Phi : S(E) \rightarrow V$ satisfying, for $n \geq 0$, an adapted version of (6) where the indexes k_1, \dots, k_j on the right hand side of the equation run from 0 to n . The zero component $\Phi_0 : \mathbb{R} \rightarrow V_0$ gives rise to an element $\Phi_0(1) \in V_0$, which by abuse of notation we denote by Φ_0 . The curved adaptation of (6), for $n = 0$, then reads $0 = \sum_{j \geq 1} \frac{1}{j!} m_j(\Phi_0, \dots, \Phi_0)$. In other words, Φ_0 is a Maurer Cartan element of V [8].

Considering the coalgebra morphism $\Phi : \bar{S}(E) \rightarrow \bar{S}(V)$ defined by the collection of degree zero linear maps

$$\Phi_k : S^k(E) \rightarrow V, \quad k \geq 1,$$

we see that Equation (6) is equivalent to Φ preserving the Lie ∞-algebra structures:

$$\Phi \circ M_E = M_V \circ \Phi.$$

2. REPRESENTATIONS OF LIE ∞-ALGEBRAS

A complex (V, d) induces a natural symmetric DGLA structure in $\text{End}(V)[1]$, see Example 1.6.

Definition 2.1. A **representation** of a Lie ∞-algebra $(E, \{l_k\}_{k \in \mathbb{N}})$ on a complex (V, d) is a Lie ∞-morphism

$$\Phi : (E, \{l_k\}_{k \in \mathbb{N}}) \rightarrow (\text{End}(V)[1], \partial, [\cdot, \cdot]),$$

i.e., $\Phi \circ M_E = M_{\text{End}(V)[1]} \circ \Phi$, where M_E is the coderivation determined by $\sum_k l_k$ and $M_{\text{End}(V)[1]}$ is the coderivation determined by $\partial + [\cdot, \cdot]$.

Equivalently, a representation of E is defined by a collection of degree +1 maps

$$\Phi_k : S^k(E) \rightarrow \text{End}(V), \quad k \geq 1,$$

such that, for each $n \geq 1$,

$$\begin{aligned}
 (7) \quad & \sum_{\substack{i=1 \\ \sigma \in Sh(i, n-i)}}^n \varepsilon(\sigma) \Phi_{n-i+1} \left(l_i(x_{\sigma(1)}, \dots, x_{\sigma(i)}), x_{\sigma(i+1)}, \dots, x_{\sigma(n)} \right) = \\
 &= \partial \Phi_n(x_1, \dots, x_n) + \frac{1}{2} \sum_{\substack{j=1 \\ \sigma \in Sh(j, n-j)}}^{n-1} \varepsilon(\sigma) \left[\Phi_j(x_{\sigma(1)}, \dots, x_{\sigma(j)}), \Phi_{n-j}(x_{\sigma(j+1)}, \dots, x_{\sigma(n)}) \right].
 \end{aligned}$$

Remark 2.2. A representation on a complex (V, d) can be seen as a curved Lie ∞ -morphism $\Phi : E \rightarrow \text{End}(V)[1]$, with $\Phi = \sum_{k \geq 0} \Phi_k$ and $\Phi_0 = d$. In fact, the first term on the right hand-side of Equations (7) is given by

$$\partial \Phi_n(x_1, \dots, x_n) = [\Phi_0, \Phi_n(x_1, \dots, x_n)],$$

and we have a curved Lie ∞ -morphism

$$\Phi : (E, \{l_k\}_{k \in \mathbb{N}}) \rightarrow (\text{End}(V)[1], [\cdot, \cdot])$$

between the Lie ∞ -algebra E and the symmetric graded Lie algebra $(\text{End}(V)[1], [\cdot, \cdot])$ (see [8], Lemma 2.5). This is why sometimes a representation of a Lie ∞ -algebra E on a complex (V, d) is called a representation on the graded vector space V (compatible with the differential d of V).

Any representation $\Phi : E \rightarrow \text{End}(V)[1]$ of a Lie ∞ -algebra E on a complex (V, d) has a dual one. Let

$$* : \text{End}(V) \rightarrow \text{End}(V^*)$$

be the Lie ∞ -morphism given by

$$(8) \quad \langle f^*(\alpha), v \rangle = -(-1)^{|\alpha||f|} \langle \alpha, f(v) \rangle, \quad f \in \text{End}(V), \alpha \in V^*, v \in V.$$

The **dual representation** $^*\Phi : E \rightarrow \text{End}(V^*)[1]$ is obtained by composition of Φ with this Lie ∞ -morphism. It is a representation on the complex (V^*, d^*) and is given by

$$(9) \quad \langle ^*\Phi(e)(\alpha), v \rangle = -(-1)^{(|e|+1)|\alpha|} \langle \alpha, \Phi(e)(v) \rangle, \quad e \in \bar{S}(E), \alpha \in V^*, v \in V.$$

Remark 2.3. Given a representation $\Phi : E \rightarrow \text{End}(V)[1]$ on a complex (V, d) , defined by the collection of degree +1 linear maps $\Phi_k : S^k(E) \rightarrow \text{End}(V)$, $k \geq 1$, one may consider the collection of degree +1 maps $\phi_k : S^k(E) \otimes V \rightarrow V$, $k \geq 0$, where $\phi_0 = d : V \rightarrow V$ and $\phi_k(x, v) = (\Phi_k(x))(v)$, $k \geq 1$.

The embedding $\bar{S}(E) \oplus (S(E) \otimes V) \hookrightarrow \bar{S}(E \oplus V)$, provides a collection of maps

$$\tilde{\Phi}_k : S^k(E \oplus V) \rightarrow E \oplus V, \quad k \geq 1,$$

given by

$$\begin{aligned} & \tilde{\Phi}_k((x_1, v_1), \dots, (x_k, v_k)) \\ &= \left(l_k(x_1, \dots, x_k), \sum_{i=1}^k (-1)^{|x_i|(|x_{i+1}| + \dots + |x_k|)} \phi_{k-1}(x_1, \dots, \widehat{x_i}, \dots, x_k, v_i) \right), \end{aligned}$$

and we may express Equations (7) as

$$(10) \quad \tilde{\Phi}_\bullet \left(\tilde{\Phi}_\bullet(x_{(1)}) \odot x_{(2)} \right) + \tilde{\Phi}_1 \tilde{\Phi}_\bullet(x) = 0, \quad x \in \bar{S}(E \oplus V).$$

Equation (10) means that $\tilde{\Phi}$ equips $E \oplus V$ with a Lie ∞ -algebra structure.

Now suppose the graded vector space V has a Lie ∞ -algebra structure $\{m_k\}_{k \in \mathbb{N}}$ given by a coderivation M_V of $\bar{S}(V)$. By the construction in Example 1.6, the coderivation M_V of $\bar{S}(V)$ defines a symmetric DGLA structure in $\text{Coder}(\bar{S}(V))[1]$:

$$\partial_{M_V} Q = -M_V \circ Q + (-1)^{\deg(Q)} Q \circ M_V,$$

$$[Q, P] = (-1)^{\deg(Q)} \left(Q \circ P - (-1)^{\deg(Q) \deg(P)} P \circ Q \right),$$

where $\deg(Q)$ and $\deg(P)$ are the degrees of Q and P in $\text{Coder}(\bar{S}(V))$.

Generalizing the notion of an action of a graded Lie algebra on another graded Lie algebra, we have the following definition of an action of a Lie ∞ -algebra on another Lie ∞ -algebra:

Definition 2.4. An **action of the Lie ∞ -algebra** $(E, M_E \equiv \{l_k\}_{k \in \mathbb{N}})$ on the Lie ∞ -algebra $(V, M_V \equiv \{m_k\}_{k \in \mathbb{N}})$, or a **Lie ∞ -action** of E on V , is a Lie ∞ -morphism

$$\Phi : (E, \{l_k\}_{k \in \mathbb{N}}) \rightarrow (\text{Coder}(\tilde{S}(V))[1], \partial_{M_V}, [\cdot, \cdot]).$$

Remark 2.5. Being a Lie ∞ -morphism, an action

$$\Phi : E \rightarrow \text{Coder}(\tilde{S}(V))[1]$$

is univocally defined by a collection of degree +1 linear maps

$$\Phi_k : S^k(E) \rightarrow \text{Coder}(\tilde{S}(V)), \quad k \geq 1.$$

By the isomorphism provided in Proposition 1.2, and since each $\Phi_k(x)$, $x \in S^k(E)$, is a coderivation of $\tilde{S}(V)$, we see that an action is completely defined by a collection of linear maps

$$(11) \quad \Phi_{k,i} : S^k(E) \otimes S^i(V) \rightarrow V, \quad i, k \geq 1.$$

We will denote the coderivation $\Phi_k(x)$ simply by Φ_x .

Remark 2.6. If we define $\Phi_0 := M_V$, then an action is equivalent to a curved Lie ∞ -morphism between E and the graded Lie algebra $\text{Coder}(\tilde{S}(V))$ (compatible with the Lie ∞ -structure in V) [8]. In this case, $\Phi = \sum_{k \geq 0} \Phi_k$ is called a **curved Lie ∞ -action**.

There is a close relationship between representations and actions on Lie ∞ -algebras.

First notice that each linear map $\ell : V \rightarrow V$ induces a (co)derivation of $\tilde{S}(V)$. Hence we may see $\text{End}(V)[1]$ as a Lie ∞ -subalgebra of $\text{Coder}(\tilde{S}(V))[1]$. Therefore, given a representation $\Phi : E \rightarrow \text{End}(V)[1]$ of the Lie ∞ -algebra E on the complex (V, d) , we have a natural action of E on the Lie ∞ -algebra (V, M_V) , where M_V is the coderivation defined by the map $d : V \rightarrow V$. In this case, we say **the action is induced by a representation**.

Moreover, for each action $\Phi : E \rightarrow \text{Coder}(\tilde{S}(V))[1]$ of E on the Lie ∞ -algebra $(V, M_V \equiv \{m_k\}_{k \in \mathbb{N}})$, we have a representation of E on V given by the collection of maps $\Phi_{k,1} : S^k(E) \otimes V \rightarrow V$, $k \geq 1$, or equivalently, $\Phi_{k,1} \equiv \rho_k : S^k(E) \rightarrow \text{End}(V)$, $k \geq 1$. The morphism $\rho = \sum_k \rho_k$ is a representation of the Lie ∞ -algebra E on the complex $(V, d = m_1)$, called the **linear representation defined by Φ** .

Finally one should notice that, given a Lie ∞ -algebra (V, M_V) , the graded vector space $\text{Coder}(\tilde{S}(V))[1]$ is a Lie ∞ -subalgebra of $\text{End}(\tilde{S}(V))[1]$. Therefore, any action $\Phi : E \rightarrow \text{Coder}(\tilde{S}(V))[1]$ of the Lie ∞ -algebra E on (V, M_V) yields a representation of E on the graded vector space $\tilde{S}(V)$. We call it the **representation induced by the action Φ** . The coderivation M_V defines a (co)derivation of $\tilde{S}(\tilde{S}(V))$ and the representation is compatible with this (co)derivation.

Remark 2.7. In [8], the authors define an action of a finite dimensional Lie ∞ -algebra E on a graded manifold \mathcal{M} as a Lie ∞ -morphism $\Phi : E \rightarrow \mathfrak{X}(\mathcal{M})[1]$. As the authors point out, when \mathcal{M} is the graded manifold defined by a finite dimensional Lie ∞ -algebra, we have an action of a Lie ∞ -algebra on another Lie ∞ -algebra. The definition presented here is a particular case of theirs because we are only considering coderivations of $\tilde{S}(V)$, i.e. coderivations of $S(V)$ vanishing on the field $S^0(E)$. This restrictive case reduces to the usually Lie algebra action on another Lie algebra (and its semi-direct product) while the definition given in [8], gives rise to general Lie algebra extensions. For our purpose, this definition is more adequate.

Next, with the identification $S^n(E \oplus V) \simeq \oplus_{k=0}^n S^{n-k}(E) \otimes S^k(V)$, we see that the action Φ determines a coderivation of $\tilde{S}(E \oplus V)$. Together with M_E and M_V we have a Lie ∞ -algebra structure on $E \oplus V$. Next proposition can be deduced from [8].

Proposition 2.8. *Let $(E, M_E \equiv \{l_k\}_{k \in \mathbb{N}})$ and $(V, M_V \equiv \{m_k\}_{k \in \mathbb{N}})$ be Lie ∞ -algebras. An action*

$$\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$$

defines a Lie ∞ -algebra structure on $E \oplus V$.

Proof. We consider the brackets $\{l_n\}_{n \in \mathbb{N}}$ on $E \oplus V$ given by:

$$\begin{aligned} l_n(x_1, \dots, x_n) &= l_n(x_1, \dots, x_n), \quad x_i \in E \\ l_n(v_1, \dots, v_n) &= m_n(v_1, \dots, v_n), \quad v_i \in V \\ l_{k+n}(x_1, \dots, x_k, v_1, \dots, v_n) &= \Phi_{k,n}(x_1, \dots, x_k, v_1, \dots, v_n), \end{aligned}$$

with $\Phi_{k,n} : S^k(E) \otimes S^n(V) \rightarrow V$ the collection of linear maps defining Φ (see Remark 2.5).

The collection of linear maps $\Phi_{k,n}$ defines a coderivation of $\bar{S}(E \oplus V)$,

$$\Upsilon : \bar{S}(E \oplus V) \rightarrow (\bar{S}(E) \otimes \bar{S}(V)) \oplus \bar{S}(V) \subset \bar{S}(E \oplus V)$$

related to the action Φ by

$$\Upsilon(x \otimes v) = \Phi_x(v), \quad x \in E, v \in \bar{S}(V)$$

and

$$\Upsilon(x \otimes v) = \Phi_x(v) + (-1)^{|x(1)|} x_{(1)} \otimes \Phi_{x_{(2)}}(v), \quad x \in S^{\geq 2}(E), v \in \bar{S}(V).$$

The degree +1 coderivation of $\bar{S}(E \oplus V)$ determined by $\{l_n\}_{n \in \mathbb{N}}$ is

$$M_{E \oplus V} = M_E + \Upsilon + M_V.$$

Let us prove that $M_{E \oplus V}^2 = 0$. For $x \in \bar{S}(E)$ and $v \in \bar{S}(V)$,

$$M_{E \oplus V}^2(x) = M_E^2(x) = 0 \quad \text{and} \quad M_{E \oplus V}^2(v) = M_V^2(v) = 0$$

while, for mixed terms, we have

$$M_{E \oplus V}(x \otimes v) = M_E(x) \otimes v + (-1)^{|x|} x \otimes M_V(v) + (-1)^{|x(1)|} x_{(1)} \otimes \Phi_{x_{(2)}}(v) + \Phi_x(v)$$

and

$$l(M_{E \oplus V}(x \otimes v)) = (\Phi_{M_E(x)})_{\bullet}(v) + (-1)^{|x|} (\Phi_x)_{\bullet}(M_V(v)) + (-1)^{|x(1)|} (\Phi_{x_{(1)}})_{\bullet}(\Phi_{x_{(2)}}(v)) + m_{\bullet}(\Phi_x(v)).$$

Since Φ is a Lie ∞ -morphism, we have

$$\Phi_{M_E(x)} = -M_V \circ \Phi_x - (-1)^{|x|} \Phi_x \circ M_V + \frac{1}{2} [\Phi_{x_{(1)}}, \Phi_{x_{(2)}}],$$

which implies $M_{E \oplus V}^2 = 0$. □

The Lie ∞ -algebra structure in $E \oplus V$ presented in Remark 2.3, is a particular case of Proposition 2.8, with $M_V = d$.

Adjoint representation and adjoint action. An important example of a representation is given by a Lie ∞ -algebra structure.

Let $(E, M_E \equiv \{l_k\}_{k \in \mathbb{N}})$ be a Lie ∞ -algebra; thus (E, l_1) is a complex. The collection of degree +1 maps

$$\begin{aligned} \text{ad}_k : S^k(E) &\rightarrow \text{End}(E) \\ x_1 \odot \dots \odot x_k &\mapsto \text{ad}_{x_1 \odot \dots \odot x_k} := l_{k+1}(x_1, \dots, x_k, -), \quad k \geq 1, \end{aligned}$$

satisfies Equations (7). (Note that Equations (7) are equivalent to Equations (2)). So, this collection of maps defines a representation $\text{ad} = \sum_k \text{ad}_k$ of the Lie ∞ -algebra E on (E, l_1) .

Definition 2.9. The representation ad is called the **adjoint representation** of the Lie ∞ -algebra $(E, M_E \equiv \{l_k\}_{k \in \mathbb{N}})$.

Moreover, notice that for each $x \in S^i(E)$, $i \geq 1$, we may consider the degree $|x| + 1$ coderivation ad_x^D of $\bar{S}(E)$ defined by the family of linear maps

$$\begin{aligned} (\text{ad}_x)_k : S^k(E) &\rightarrow E \\ e &\mapsto l_{i+k}(x, e), \quad k \geq 1. \end{aligned}$$

So, we have a collection of degree +1 linear maps

$$(12) \quad \begin{aligned} \text{ad}_i : S^i(E) &\rightarrow \text{Coder}(\bar{S}(E)) \\ x &\mapsto \text{ad}_x^D \end{aligned}, \quad i \geq 1,$$

and we set $\mathbf{ad} = \sum_i \text{ad}_i$.

Proposition 2.10. *The collection of degree +1 linear maps given by (12) defines a Lie ∞ -morphism*

$$\mathbf{ad} : (E, \{l_k\}_{k \in \mathbb{N}}) \rightarrow (\text{Coder}(\bar{S}(E))[1], \partial_{M_E}, [\cdot, \cdot])$$

from the Lie ∞ -algebra E to the symmetric DGLA $\text{Coder}(\bar{S}(E))[1]$.

Proof. For each $x \in S^i(E)$, let $\text{ad}_x = \sum_k (\text{ad}_x)_k$ and set $l = \sum_k l_k$.

If $x \in \oplus_{i \geq 2} S^i(E)$ and $e \in \bar{S}(E)$, we have

$$\begin{aligned} M_E(x \odot e) &= M_E(x) \odot e + (-1)^{|x|} x \odot M_E(e) + (-1)^{|e||x(2)|} l(x_{(1)}, e) \odot x_{(2)} \\ &\quad + l(x, e_{(1)}) \odot e_{(2)} + (-1)^{|e(1)||x(2)|} l(x_{(1)}, e_{(1)}) \odot x_{(2)} \odot e_{(2)} + l(x, e) \end{aligned}$$

and so,

$$\begin{aligned} \text{ad}_x(M_E(e)) &= l(x, M_E(e)) \\ &= (-1)^{|x|} \underbrace{l(M_E(x \odot e))}_{=0 \text{ by (2)}} - (-1)^{|x|} l(M_E(x), e) - (-1)^{|x|} l(\text{ad}_x^D(e)) \\ &\quad - (-1)^{|x(1)|+|x(1)||x(2)|} l(x_{(2)}, \text{ad}_{x_{(1)}}^D(e)) \\ &= (-(-1)^{|x|} \text{ad}_{M_E(x)} - (-1)^{|x|} l \circ \text{ad}_x^D - (-1)^{|x(2)|} \text{ad}_{x_{(1)}} \circ \text{ad}_{x_{(2)}}^D)(e), \end{aligned}$$

which is equivalent to

$$\text{ad}_{M_E(x)} = -l \circ \text{ad}_x^D - (-1)^{|x|} \text{ad}_x \circ M_E - (-1)^{|x(2)|} \text{ad}_{x_{(1)}} \circ \text{ad}_{x_{(2)}}^D$$

or to

$$(13) \quad \text{ad}_{M_E(x)} = -[l, \text{ad}_x]_{RN} - \frac{1}{2}(-1)^{|x(1)|} [\text{ad}_{x_{(1)}}, \text{ad}_{x_{(2)}}]_{RN}.$$

Note that the coderivation defined by the second member of (13) is

$$[M_E, \text{ad}_x^D] + \frac{1}{2}[\text{ad}_{x_{(1)}}^D, \text{ad}_{x_{(2)}}^D] = \partial_{M_E}(\text{ad}_x^D) + \frac{1}{2}[\text{ad}_{x_{(1)}}^D, \text{ad}_{x_{(2)}}^D].$$

If $x \in E$, a similar computation gives

$$(14) \quad \text{ad}_{l_1(x)} = -l \circ \text{ad}_x^D - (-1)^{|x|} \text{ad}_x \circ M_E = -[l, \text{ad}_x]_{RN}.$$

Equations (13) and (14) mean that the map $\mathbf{ad} : E \rightarrow \text{Coder}(\bar{S}(E))[1]$ is a Lie ∞ -morphism. \square

Definition 2.11. The linear map $\mathbf{ad} : E \rightarrow \text{Coder}(\bar{S}(E))[1]$ is an action of the Lie ∞ -algebra E on itself, called the **adjoint action of E** .

3. O-OPERATORS ON A LIE ∞ -ALGEBRA

In this section we define \mathcal{O} -operators on a Lie ∞ -algebra E with respect to an action of E on a Lie ∞ -algebra V . This is the main notion of the paper.

3.1. \mathcal{O} -operators with respect to a Lie ∞ -action. Let $(E, M_E \equiv \{l_k\}_{k \geq 1})$ and $(V, M_V \equiv \{m_k\}_{k \geq 1})$ be Lie ∞ -algebras and $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ a Lie ∞ -action of E on V . Remember we are using Sweedler's notation: for each $v \in \bar{S}(V)$,

$$\Delta(v) = v_{(1)} \otimes v_{(2)}$$

and

$$\Delta^2(v) = (\text{id} \otimes \Delta)\Delta(v) = (\Delta \otimes \text{id})\Delta(v) = v_{(1)} \otimes v_{(2)} \otimes v_{(3)}.$$

Each degree zero linear map $T : \bar{S}(V) \rightarrow \bar{S}(E)$ defines a degree +1 linear map $\Phi^T : \bar{S}(V) \rightarrow \bar{S}(V)$ given by

$$\begin{aligned} \Phi^T(v) &= 0, \quad v \in V, \\ \Phi^T(v) &= \Phi_{T(v_{(1)})} v_{(2)}, \quad v \in S^{\geq 2}(V). \end{aligned}$$

Lemma 3.1. *The linear map $\Phi^T : \bar{S}(V) \rightarrow \bar{S}(V)$ is a degree +1 coderivation of $\bar{S}(V)$ and is defined by the collection of linear maps $\sum \Phi_{\bullet, \bullet}(T \otimes \text{id})\Delta$.*

Proof. For the linear map $\Phi^T : \bar{S}(V) \rightarrow \bar{S}(V)$ to be a coderivation it must satisfy:

$$\Delta\Phi^T(v) = (\Phi^T \otimes \text{id} + \text{id} \otimes \Phi^T)\Delta(v), \quad v \in \bar{S}(V).$$

This equation is trivially satisfied for $v \in V$.

For each $v = v_1 \odot v_2 \in S^2(V)$ we have $\Phi^T(v) \in V$ and consequently, $\Delta\Phi^T(v) = 0$. On the other hand, since $\Phi^T|_V = 0$, we see that

$$(\Phi^T \otimes \text{id} + \text{id} \otimes \Phi^T)\Delta(v) = 0$$

and the equation is satisfied in $S^2(V)$.

Now let $v \in S^{\geq 3}(V)$, then

$$\begin{aligned} \Delta\Phi^T(v) &= \Delta\Phi_{T(v_{(1)})}v_{(2)} \\ &= (\Phi_{T(v_{(1)})} \otimes \text{id} + \text{id} \otimes \Phi_{T(v_{(1)})})\Delta(v_{(2)}). \end{aligned}$$

The coassociativity of Δ ensures that

$$\begin{aligned} \Delta\Phi^T(v) &= \Phi_{T(v_{(1)})}v_{(2)} \otimes v_{(3)} + (-1)^{(|v_{(1)}|+1)|v_{(2)}|}v_{(2)} \otimes \Phi_{T(v_{(1)})}v_{(3)} \\ &= \Phi_{T(v_{(1)})}v_{(2)} \otimes v_{(3)} + (-1)^{|v_{(1)}|}v_{(1)} \otimes \Phi_{T(v_{(2)})}v_{(3)} \\ &= (\Phi^T \otimes \text{id} + \text{id} \otimes \Phi^T)\Delta(v). \end{aligned}$$

□

Definition 3.2. Let $(E, M_E \equiv \{l_k\}_{k \geq 1})$ and $(V, M_V \equiv \{m_k\}_{k \geq 1})$ be Lie ∞ -algebras and $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ an action. An \mathcal{O} -operator on E with respect to the action Φ is a (degree 0) morphism of coalgebras $T : \bar{S}(V) \rightarrow \bar{S}(E)$ such that

$$(15) \quad M_E \circ T = T \circ (\Phi^T + M_V).$$

Definition 3.3. A **Rota Baxter operator (of weight 1)** on a Lie ∞ -algebra $(E, M_E \equiv \{l_k\}_{k \geq 1})$ is an \mathcal{O} -operator with respect to the adjoint action.

An \mathcal{O} -operator $T : \bar{S}(V) \rightarrow \bar{S}(E)$ with respect to an action $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ of $(E, M_E \equiv \{l_k\}_{k \geq 1})$ on $(V, M_V \equiv \{m_k\}_{k \geq 1})$ is defined by a linear map $t = \sum_i t_i : \bar{S}(V) \rightarrow E$ satisfying:

- (i) $l_1(t_1(v)) = t_1(m_1(v)), \quad v \in V$
- (ii) $l(T(v)) = t\left(\Phi_{T(v_{(1)})}v_{(2)} + m(v_{(1)}) \odot v_{(2)}\right), \quad v \in \oplus_{i \geq 2} S^i(V).$

In particular, the \mathcal{O} -operator T is a comorphism i.e., for each $v \in S^n(V)$, $n \geq 1$,

$$T(v) = \sum_{k_1 + \dots + k_r = n} \frac{1}{r!} t_{k_1}(v_{(k_1)}) \odot \dots \odot t_{k_r}(v_{(k_r)}),$$

so, detailing (ii) for $v = v_1 \odot v_2$, we get

$$\begin{aligned} l_1(t_2(v_1, v_2)) + l_2(t_1(v_1), t_1(v_2)) &= t_1(\Phi_{t_1(v_1)}v_2 + (-1)^{|v_1||v_2|}\Phi_{t_1(v_2)}v_1 + m_2(v_1, v_2)) \\ &\quad + t_2(m_1(v_1, v_2)) + (-1)^{|v_1|}t_2(v_1, m_1(v_2)). \end{aligned}$$

Generally, for every $v = v_1 \odot \dots \odot v_n \in S^n(V)$, $n \geq 3$, we have

$$\begin{aligned} &\sum_{\substack{k_1 + \dots + k_i = n \\ \sigma \in Sh(k_1, \dots, k_i)}} \frac{\epsilon(\sigma)}{i!} l_i \left(t_{k_1}(v_{\sigma(1)}), \dots, v_{\sigma(k_1)}, \dots, t_{k_i}(v_{\sigma(k_1 + \dots + k_{i-1} + 1)}), \dots, v_{\sigma(n)} \right) \\ &= \sum_{\substack{k_1 + \dots + k_{i+2} = n \\ \sigma \in Sh(k_1, \dots, k_{i+2})}} \frac{\epsilon(\sigma)}{i!} t_{1+k_{i+2}} \left(\Phi_{i, k_{i+1}} \left(t_{k_1}(v_{\sigma(1)}), \dots, v_{\sigma(k_1)} \right) \odot \dots \odot t_{k_i}(v_{\sigma(k_1 + \dots + k_{i-1} + 1)}), \dots, v_{\sigma(k_1 + \dots + k_i)} \right), \\ &\quad v_{\sigma(k_1 + \dots + k_{i+1})} \odot \dots \odot v_{\sigma(k_1 + \dots + k_{i+1} + 1)}, v_{\sigma(k_1 + \dots + k_{i+1} + 1)} \odot \dots \odot v_{\sigma(n)} \Big) \\ &\quad + \sum_{i=1}^n \sum_{\sigma \in Sh(i, n-i)} \epsilon(\sigma) t_{n-i+1} (m_i(v_{\sigma(1)}), \dots, v_{\sigma(i)}, v_{\sigma(i+1)}, \dots, v_{\sigma(n)}). \end{aligned}$$

Remark 3.4. When $M_V = 0$ we are considering V simply as a graded vector space, with no Lie ∞ -algebra attached and an \mathcal{O} -operator must satisfy

$$M_E \circ T = T \circ \Phi^T.$$

In this case, the terms of above equations involving the brackets m_i on V vanish.

Remark 3.5. When $(E, [\cdot, \cdot]_E)$ and $(V, [\cdot, \cdot]_V)$ are Lie algebras, for degree reasons, a morphism $T = t_1$ must be a strict morphism. Moreover, our definition coincides with the usual definition of \mathcal{O} -operator (of weight 1) between Lie algebras [4] :

$$[t_1(v), t_1(w)]_E = t_1(\Phi_{t_1(v)}w - \Phi_{t_1(w)}v + [v, w]_V), \quad v, w \in V.$$

Remark 3.6. When (V, d) is just a complex and the action $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ is induced by a representation $\rho : E \rightarrow \text{End}(V)[1]$ we have that $\Phi(x)$ is the (co)derivation defined by $\rho(x)$. In this case, \mathcal{O} -operators with respect to Φ coincide with \mathcal{O} -operators with respect to ρ (or relative Rota Baxter operators) given in [6].

In [6] the authors define \mathcal{O} -operators with respect to representations of Lie ∞ -algebras. Any action induces a representation and \mathcal{O} -operators with respect to an action are related with \mathcal{O} -operators of with respect to the induced representation. We shall see that this relation is given by the comorphism

$$I = \sum_{n \geq 1} i_n : \bar{S}(V) \rightarrow \bar{S}(\bar{S}(V)),$$

defined by the family of inclusion maps $i_n : S^n(V) \hookrightarrow \bar{S}(V)$, $n \geq 1$.

Notice that any coderivation D of $\bar{S}(V)$ induces a (co)derivation D^d of $\bar{S}(\bar{S}(V))$. The comorphism I preserves these coderivations:

Lemma 3.7. *Let V be a graded vector space and D a coderivation of $\bar{S}(V)$. The map $I : \bar{S}(V) \rightarrow \bar{S}(\bar{S}(V))$ satisfies*

$$I \circ D = D^d \circ I.$$

Proof. We will denote by \cdot the symmetric product in $\bar{S}(\bar{S}(V))$, to distinguish from the symmetric product \odot in $\bar{S}(V)$.

Let $v \in S^n(V)$, $n \geq 1$, and denote by $\{m_k\}_{k \geq 1}$ the family of linear maps defining the coderivation D . For $v \in V$, we immediately have $D^d \circ I(v) = D \circ I(v) = I \circ D(v)$. For $v \in S^{\geq 2}(V)$ we have

$$\begin{aligned} D^d \circ I(v) &= D^d \left(\sum_{k=1}^n \frac{1}{k!} v_{(1)} \cdot \dots \cdot v_{(k)} \right) \\ &= \sum_{k=1}^n \frac{1}{k!} \left(D(v_{(1)}) \cdot v_{(2)} \cdot \dots \cdot v_{(k)} + \dots + (-1)^{|D|(|v_{(1)}| + \dots + |v_{(k-1)}|)} v_{(1)} \cdot \dots \cdot v_{(k-1)} \cdot D(v_{(k)}) \right) \\ &= \sum_{k=1}^n \frac{1}{(k-1)!} D(v_{(1)}) \cdot v_{(2)} \cdot \dots \cdot v_{(k)} \\ &= D(v_{(1)}) \cdot I(v_{(2)}). \end{aligned}$$

On the other hand

$$\begin{aligned} I \circ D(v) &= I(m_{\bullet}(v_{(1)}) \odot v_{(2)}) \\ &= m_{\bullet}(v_{(1)}) \cdot I(v_{(2)}) + (m_{\bullet}(v_{(1)}) \odot v_{(2)}) \cdot I(v_{(3)}) \\ &= D(v_{(1)}) \cdot I(v_{(2)}) \end{aligned}$$

and the result follows. \square

Remark 3.8. In particular, if D defines a Lie ∞ -algebra structure on V , then D^d defines a Lie ∞ -algebra structure on $\bar{S}(V)$ and I is a Lie ∞ -morphism.

Proposition 3.9. *Let $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ be an action of the Lie ∞ -algebra $(E, M_E \equiv \{l_k\}_{k \geq 1})$ on the Lie ∞ -algebra $(V, M_V \equiv \{m_k\}_{k \geq 1})$ and $\tilde{T} : \bar{S}(\bar{S}(V)) \rightarrow E$ be an \mathcal{O} -operator with respect to the induced representation $\rho : E \rightarrow \text{End}(\bar{S}(V))[1]$. Then $T = \tilde{T} \circ I$ is an \mathcal{O} -operator with respect to the action Φ .*

Proof. For each $x \in \bar{S}(E)$, let us denote by

$$\Phi_x^d := \Phi(x)^d = \rho(x)^d,$$

the (co)derivation of $\bar{S}(\bar{S}(V))$ defined by $\rho(x)$.

Let \tilde{T} be an \mathcal{O} -operator with respect to the induced representation. This means that

$$M_E \circ \tilde{T}(w) = \tilde{T} \left(\Phi_{\tilde{T}(w_{(1)})}^d w_{(2)} + M_V^d(w) \right), \quad w \in \bar{S}(\bar{S}(V)).$$

Then, for each $w = I(v)$, $v \in \bar{S}(V)$, we have:

$$M_E \circ \tilde{T}(I(v)) = \tilde{T} \left(\Phi_{\tilde{T}(I(v)_{(1)})}^d I(v)_{(2)} + M_V^d \circ I(v) \right).$$

Using the fact that I is a comorphism and Lemma 3.7, we rewrite last equation as

$$\begin{aligned} M_E \circ T(v) &= \tilde{T} \left(\Phi_{\tilde{T}(I(v)_{(1)})}^d I(v)_{(2)} + M_V^d \circ I(v) \right) \\ &= \tilde{T} \left(I \circ \Phi_{T(v_{(1)})} v_{(2)} + I \circ M_V(v) \right) \\ &= T \left(\Phi_{T(v_{(1)})} v_{(2)} + M_V(v) \right). \end{aligned}$$

Taking into account this equation and that T is a comorphism, because is the composition of two comorphisms, the result follows. \square

Proposition 3.10. *Let T be an \mathcal{O} -operator on $(E, M_E \equiv \{l_k\}_{k \geq 1})$ with respect to a Lie ∞ -action $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ on $(V, M_V \equiv \{m_k\}_{k \geq 1})$. Then, V has a new Lie ∞ -algebra structure*

$$M_{V^T} = \Phi^T + M_V$$

and $T : (V, M_{V^T}) \rightarrow (E, M_E)$ is a Lie ∞ -morphism.

Proof. By Lemma 3.1 we know Φ^T is a degree +1 coderivation of $\bar{S}(V)$ hence so is M_{V^T} .

Since Φ is an action, so that $\Phi \circ M_E = M_{\text{Coder}(\bar{S}(V))[1]} \circ \Phi$, and T is a comorphism, we have, for each $v \in \bar{S}(V)$,

$$(16) \quad \begin{aligned} \Phi_{M_E T(v_{(1)})} v_{(2)} &= -M_V \Phi_{T(v_{(1)})} v_{(2)} - (-1)^{|v_{(1)}|} \Phi_{T(v_{(1)})} M_V(v_{(2)}) \\ &\quad + (-1)^{|v_{(1)}|+1} \Phi_{T(v_{(1)})} \Phi_{T(v_{(2)})} v_{(3)}. \end{aligned}$$

On the other hand, T is an \mathcal{O} -operator:

$$M_E \circ T(v) = T \circ \Phi_{T(v_{(1)})} v_{(2)} + T \circ M_V(v)$$

and this yields

$$(17) \quad \Phi_{M_E T(v_{(1)})} v_{(2)} = \Phi_{T \Phi_{T(v_{(1)})} v_{(2)}} v_{(3)} + \Phi_{T M_V(v_{(1)})} v_{(2)}.$$

Moreover, due to the fact that both Φ^T and M_V are coderivations and $M_V^2 = 0$, we have

$$\begin{aligned} M_{V^T}^2(v) &= (\Phi^T)^2(v) + \Phi^T \circ M_V(v) + M_V \circ \Phi^T(v) \\ &= \Phi_{T(\Phi_{T(v_{(1)})} v_{(2)})} v_{(3)} + (-1)^{|v_{(1)}|} \Phi_{T(v_{(1)})} \Phi_{T(v_{(2)})} v_{(3)} \\ &\quad + \Phi_{T M_V(v_{(1)})} v_{(2)} + (-1)^{|v_{(1)}|} \Phi_{T(v_{(1)})} M_V(v_{(2)}) + M_V(\Phi_{T(v_{(1)})} v_{(2)}). \end{aligned}$$

Taking into account Equations (16) and (17) we conclude $M_{V^T}^2 = 0$. Therefore, M_{V^T} defines a Lie ∞ -algebra structure on V and Equation (15) means that $T : \bar{S}(V) \rightarrow \bar{S}(E)$ is a Lie ∞ -morphism between the Lie ∞ -algebras (V, M_{V^T}) and (E, M_E) . \square

The brackets of the Lie ∞ -algebra structure on V defined by the coderivation M_{V^T} are given by

$$m_1^T(v) = m_1(v)$$

and, for $n \geq 2$,

$$m_n^T(v_1, \dots, v_n) = m_n(v_1, \dots, v_n) + \sum_{\substack{k_1 + \dots + k_i = j \\ 1 \leq j \leq n-1}} \sum_{\sigma \in Sh(k_1, \dots, k_i, n-j)} \epsilon(\sigma) \frac{1}{n!}$$

$$\Phi_{i, n-j} (t_{k_1}(v_{\sigma(1)}, \dots, v_{\sigma(k_1)}) \odot \dots \odot t_{k_i}(v_{\sigma(k_1 + \dots + k_{i-1} + 1)}, \dots, v_{\sigma(j)}), v_{\sigma(j+1)} \odot \dots \odot v_{\sigma(n)}),$$

with $\Phi_{i, n-j}$, $i \geq 1$, the linear maps determined by the action Φ (see (11)).

\mathcal{O} -operators for the coadjoint representation. Let $(E, M_E \equiv \{l_k\}_{k \geq 1})$ be a finite dimensional Lie ∞ -algebra. Next, we consider the dual of the adjoint representation of E (see (9)), called the coadjoint representation.

Definition 3.11. The **coadjoint representation** of E , $\text{ad}^* : E \rightarrow \text{End}(E^*)[1]$, is defined by

$$\langle \text{ad}_x^*(\alpha), v \rangle = -(-1)^{|\alpha|(|x|+1)} \langle \alpha, \text{ad}_x v \rangle, \quad v \in E, x \in \bar{S}(E), \alpha \in E^*.$$

Notice that E^* is equipped with the differential l_1^* (see (8)).

An \mathcal{O} -operator on E with respect to the coadjoint representation $\text{ad}^* : E \rightarrow \text{End}(E^*)[1]$ is a coalgebra morphism $T : \bar{S}(E^*) \rightarrow \bar{S}(E)$ given by a collection of maps $t = \sum_i t_i : \bar{S}(E^*) \rightarrow E$ satisfying

$$(18) \quad \begin{aligned} l(T(\alpha)) &= \sum_{\substack{1 \leq i \leq n-1 \\ \sigma \in Sh(i, n-i)}} \varepsilon(\sigma) t_{n-i+1}(\text{ad}_{T(\alpha_{\sigma(1)} \odot \dots \odot \alpha_{\sigma(i)})}^* \alpha_{\sigma(i+1)}, \alpha_{\sigma(i+2)}, \dots, \alpha_{\sigma(n)}) \\ &\quad + \sum_{i=1}^n (-1)^{|\alpha_1| + \dots + |\alpha_{i-1}|} t_n(\alpha_1, \dots, l_1^* \alpha_i, \dots, \alpha_n), \end{aligned}$$

for all $\alpha = \alpha_1 \odot \dots \odot \alpha_n \in S^n(E^*)$, $n \geq 1$.

We say that T is **symmetric** if

$$\langle \beta, t_n(\alpha_1, \dots, \alpha_n) \rangle = (-1)^{|\alpha| |\beta| + |\alpha_n| (|\alpha_1| + \dots + |\alpha_{n-1}|)} \langle \alpha_n, t_n(\alpha_1, \dots, \alpha_{n-1}, \beta) \rangle,$$

for all $\alpha_1, \dots, \alpha_n, \beta \in E^*$ and $n \geq 1$.

When T is invertible, its inverse $T^{-1} : \bar{S}(E) \rightarrow \bar{S}(E^*)$, given by $t^{-1} = \sum_n t_n^{-1}$, is also symmetric:

$$\langle t_n^{-1}(x_1, \dots, x_n), y \rangle = (-1)^{|y| |x_n|} \langle t_n^{-1}(x_1, \dots, x_{n-1}, y), x_n \rangle,$$

for every $x_1, \dots, x_n, y \in E$, $n \geq 1$.

One should notice that t_n^{-1} is **not** the inverse map of t_n . It simply denotes the n -component of the inverse T^{-1} of T .

For each $n \geq 1$, let $\omega^{(n)} \in \otimes^n E^*$ be defined by $\omega^{(1)} = 0$ and

$$\langle \omega^{(n)}, x_1 \otimes \dots \otimes x_n \rangle = \langle t_{n-1}^{-1}(x_1, \dots, x_{n-1}), x_n \rangle, \quad x_1, \dots, x_n \in E.$$

The symmetry of T^{-1} guarantees that $\omega = \sum_{n \geq 1} \omega^{(n)}$ is an element of $\bar{S}(E^*)$.

Proposition 3.12. *Let $T : \bar{S}(E^*) \rightarrow \bar{S}(E)$ be an invertible symmetric comorphism. The linear map T is an \mathcal{O} -operator with respect to the coadjoint representation if and only if $\omega \in \oplus_{n \geq 2} S^n(E^*)$, given by*

$$\langle \omega, x_1 \odot \dots \odot x_{k+1} \rangle = \langle t_k^{-1}(x_1, \dots, x_k), x_{k+1} \rangle, \quad x_1, \dots, x_{k+1} \in E, \quad k \geq 1,$$

is a cocycle for the Lie ∞ -algebra cohomology.

Proof. When T is invertible, Equation (18) is equivalent to equations

$$t_1^{-1} l_1(x) = l_1^* t_1^{-1}(x), \quad x \in E,$$

and

$$t^{-1} M_E(x) = \text{ad}_{x_{(1)}}^* t^{-1}(x_{(2)}) + l_1^* t_n^{-1}(x), \quad x \in S^n(E), n \geq 2.$$

Let $x = x_1 \odot \dots \odot x_n \in S^n(E)$, $n \geq 1$, and $y \in E$, such that $|y| = |x| + 1$. We have:

$$\begin{aligned} \langle \omega, M_E(x \odot y) \rangle &= \langle t^{-1}(M_E(x)), y \rangle + (-1)^{|x|} \langle t^{-1}(x_1, \dots, x_n), l_1(y) \rangle \\ &\quad + (-1)^{|x_{(1)}|} \langle t^{-1}(x_{(1)}), \text{ad}_{x_{(2)}} y \rangle \\ &= \langle t^{-1}(M_E(x)), y \rangle - \langle l_1^* t^{-1}(x), y \rangle - \langle \text{ad}_{x_{(1)}}^* t^{-1}(x_{(2)}), y \rangle \end{aligned}$$

and the result follows. \square

3.2. \mathcal{O} -operators as Maurer-Cartan elements. Let $(E, M_E \equiv \{l_k\}_{k \geq 1})$ and $(V, M_V \equiv \{m_k\}_{k \geq 1})$ be Lie ∞ -algebras.

The graded vector space of linear maps between $\bar{S}(V)$ and E will be denoted by $\mathfrak{h} := \text{Hom}(\bar{S}(V), E)$. It can be identified with the space of coalgebra morphisms between $\bar{S}(V)$ and $\bar{S}(E)$. On the other hand, since

$$S^n(E \oplus V) \simeq \bigoplus_{k=0}^n (S^{n-k}(E) \otimes S^k(V)), \quad n \geq 1,$$

the space \mathfrak{h} can be seen as a subspace of $\text{Coder}(\bar{S}(E \oplus V))$, the space of coderivations of $\bar{S}(E \oplus V)$. Its elements define coderivations that only act on elements of $\bar{S}(V)$, they are $S(E)$ -linear.

The space $S(E \oplus V)$ has a natural $S(E)$ -bimodule structure. With the above identification we have:

$$e \cdot (x \otimes v) = (e \odot x) \otimes v = (-1)^{|e|(|x|+|v|)}(x \otimes v) \cdot e,$$

for $e \in S(E)$, $x \otimes v \in S(E \oplus V) \simeq S(E) \otimes S(V)$.

Let $t : \bar{S}(V) \rightarrow E$ be an element of \mathfrak{h} defined by the collection of maps $t_k : S^k(V) \rightarrow E$, $k \geq 1$. Let us denote by $T : \bar{S}(V) \rightarrow \bar{S}(E)$ the coalgebra morphism and by \mathfrak{t} the coderivation of $\bar{S}(E \oplus V)$ defined by t . Notice that

$$\mathfrak{t}(v) = t_1(v), \quad v \in V$$

and

$$\mathfrak{t}(v) = t(v_{(1)}) \otimes v_{(2)} + \mathfrak{t}(v), \quad v \in S^{\geq 2}(V).$$

and also, for $x \in \bar{S}(E)$,

$$\mathfrak{t}(x \otimes v) = (-1)^{|x||\mathfrak{t}|} x \cdot \mathfrak{t}(v), \quad v \in \bar{S}(V).$$

Proposition 3.13. *The space \mathfrak{h} is an abelian Lie subalgebra of $\text{Coder}(\bar{S}(E \oplus V))$.*

Proof. Let $t = \sum_i t_i : \bar{S}(V) \rightarrow E$ and $w = \sum_i w_i : \bar{S}(V) \rightarrow E$ be elements of \mathfrak{h} . Denote by \mathfrak{t} and \mathfrak{w} the coderivations of $\bar{S}(E \oplus V)$ defined by t and w , respectively.

Let $v \in \bar{S}(V)$. The Lie bracket of \mathfrak{t} and \mathfrak{w} is given by:

$$\begin{aligned} [\mathfrak{t}, \mathfrak{w}]_c(v) &= \mathfrak{t} \circ (w(v_{(1)}) \otimes v_{(2)}) - (-1)^{|\mathfrak{t}||\mathfrak{w}|} \mathfrak{w} \circ (t(v_{(1)}) \otimes v_{(2)}) \\ &= (-1)^{|\mathfrak{t}|(|w|+|v_{(1)}|)} w(v_{(1)}) \cdot \mathfrak{t}(v_{(2)}) - (-1)^{|\mathfrak{t}||w|} (-1)^{|w|(|\mathfrak{t}|+|v_{(1)}|)} t(v_{(1)}) \cdot \mathfrak{w}(v_{(2)}) \\ &= \left((-1)^{|\mathfrak{t}|(|w|+|v_{(1)}|)} w(v_{(1)}) \cdot t(v_{(2)}) - (-1)^{|\mathfrak{t}||w|} (-1)^{|w|(|\mathfrak{t}|+|v_{(1)}|)} t(v_{(1)}) \cdot w(v_{(2)}) \right) \otimes v_{(3)} \\ &\quad + (-1)^{|\mathfrak{t}|(|w|+|v_{(1)}|)} w(v_{(1)}) \cdot \mathfrak{t}(v_{(2)}) - (-1)^{|\mathfrak{t}||w|} (-1)^{|w|(|\mathfrak{t}|+|v_{(1)}|)} t(v_{(1)}) \cdot w(v_{(2)}) \\ &= \left((-1)^{|\mathfrak{t}|(|w|+|v_{(1)}|)} w(v_{(1)}) \odot t(v_{(2)}) - (-1)^{|\mathfrak{t}|(|w|+|v_{(2)}|)+|v_{(1)}||v_{(2)}|} w(v_{(2)}) \odot t(v_{(1)}) \right) \otimes v_{(3)} \\ &\quad + (-1)^{|\mathfrak{t}|(|w|+|v_{(1)}|)} w(v_{(1)}) \odot t(v_{(2)}) - (-1)^{|\mathfrak{t}|(|w|+|v_{(2)}|)+|v_{(1)}||v_{(2)}|} w(v_{(2)}) \odot t(v_{(1)}), \end{aligned}$$

where we used the fact that \mathfrak{t} and \mathfrak{w} are $\bar{S}(E)$ -linear. Because of cocommutativity of the coproduct, the last expression vanishes. \square

Now, let $\Phi : E \rightarrow \text{Coder}(\bar{S}(V))[1]$ be an action of the Lie ∞ -algebra E on the Lie ∞ -algebra V . By Proposition 2.8, Φ induces a coderivation Υ of $\bar{S}(E \oplus V)$ and $M_{E \oplus V} = M_E + \Upsilon + M_V$ is a Lie ∞ -algebra structure on $E \oplus V$. Let $\mathcal{P} : \text{Coder}(\bar{S}(E \oplus V)) \rightarrow \mathfrak{h}$ be the projection onto \mathfrak{h} .

Then we have:

Proposition 3.14. *The quadruple $(\text{Coder}(\bar{S}(E \oplus V)), \mathfrak{h}, \mathcal{P}, M_{E \oplus V})$ is a V -data and \mathfrak{h} has a Lie ∞ -algebra structure.*

Proof. We already know that $\text{Coder}(\bar{S}(E \oplus V))$, equipped with the commutator, is a graded Lie algebra and \mathfrak{h} is an abelian Lie subalgebra.

Let $p : \bar{S}(E \oplus V) \rightarrow E$ be the projection and $i : \bar{S}(V) \rightarrow \bar{S}(E \oplus V)$ the inclusion.

Notice that, for each $Q \in \text{Coder}(\bar{S}(E \oplus V))$ we have $\mathcal{P}(Q) = p \circ Q \circ i$ so

$$\ker \mathcal{P} = \{Q \in \text{Coder}(\bar{S}(E \oplus V)) : Q \circ i \text{ is a coderivation of } \bar{S}(V)\}$$

is clearly a Lie subalgebra of $\text{Coder}(\bar{S}(E \oplus V))$:

$$\begin{aligned} \mathcal{P}([Q, P]_c) &= p \circ [Q, P]_c \circ i = p \circ QP \circ i - (-1)^{|Q||P|} p \circ PQ \circ i \\ &= p \circ Q \circ i \circ P \circ i - (-1)^{|Q||P|} p \circ P \circ i \circ Q \circ i = 0, \quad P, Q \in \ker \mathcal{P}. \end{aligned}$$

Moreover

$$M_{E \oplus V} \circ i = M_V, \text{ so } M_{E \oplus V} \in (\ker \mathcal{P})_1$$

and, since $M_{E \oplus V}$ defines a Lie ∞ -structure in $E \oplus V$, we have:

$$[M_{E \oplus V}, M_{E \oplus V}]_c = 0.$$

Voronov's construction [10] guarantees that \mathfrak{h} inherits a (symmetric) Lie ∞ -structure given by:

$$\partial_k(t_1, \dots, t_k) = \mathcal{P}([\dots [M_{E \oplus V}, t_1]_{RN} \dots]_{RN}, t_k]_{RN}), \quad t_1, \dots, t_k \in \mathfrak{h}, k \geq 1.$$

□

Remark 3.15. A similar proof as in [6] shows that, with the above structure, \mathfrak{h} is a filtered Lie ∞ -algebra.

Lemma 3.16. *Let $t : \bar{S}(V) \rightarrow E$ be a degree zero element of \mathfrak{h} . For each $v \in \bar{S}(V)$,*

$$\partial_1 t(v) = l_1 t(v) - t \circ M_V(v)$$

and

$$\partial_k(t, \dots, t)(v) = l_k(t(v_{(1)}), \dots, t(v_{(k)})) - k t \left(\Phi_{t(v_{(1)}) \odot \dots \odot t(v_{(k-1)})} v_{(k)} \right), \quad k \geq 2.$$

Proof. Let $p : \bar{S}(E \oplus V) \rightarrow E$ be the projection map and \mathfrak{t} the coderivation of $\bar{S}(E \oplus V)$ defined by t . Notice that

$$\begin{aligned} p \circ \mathfrak{t} &= t \\ p \circ \mathfrak{t}^k &= 0, \quad k \geq 2. \end{aligned}$$

Consequently, for $k = 1$ we have

$$\partial_1 t(v) = p \circ M_{E \oplus V} \circ \mathfrak{t}(v) - p \circ \mathfrak{t} \circ M_V(v) = l_1 t(v) - t \circ M_V(v), \quad v \in \bar{S}(V)$$

and, for $k \geq 2$,

$$\begin{aligned} \partial_k(t, \dots, t) &= p \circ M_{E \oplus V} \circ \mathfrak{t}^k - k p \circ \mathfrak{t} \circ M_{E \oplus V} \circ \mathfrak{t}^{k-1} \\ &= l \circ \mathfrak{t}^k - k t \circ M_{E \oplus V} \circ \mathfrak{t}^{k-1} \end{aligned}$$

and the result follows. □

Remark 3.17. Notice that $\partial_k(t, \dots, t)(v) = 0$, for $v \in S^{<k}(V)$, as a consequence of $\mathfrak{t}^k(v) = 0$ and $\Phi \circ \mathfrak{t}^{k-1}(v) \in \bar{S}(E)$.

Next proposition realizes \mathcal{O} -operators as Maurer-Cartan elements of this Lie ∞ -algebra \mathfrak{h} .

Proposition 3.18. *\mathcal{O} -operators on E with respect to an action Φ are Maurer-Cartan elements of \mathfrak{h} .*

Proof. Let $t : \bar{S}(V) \rightarrow E$ be a degree zero element of \mathfrak{h} . Maurer-Cartan equation yields

$$\partial_1 t + \frac{1}{2} \partial_2(t, t) + \cdots + \frac{1}{k!} \partial_k(t, \dots, t) + \dots = 0.$$

Using Lemma 3.16 we have, for each $v \in S^k(V)$,

$$\begin{aligned} \partial_1 t(v) + \frac{1}{2} \partial_2(t, t)(v) + \cdots + \frac{1}{k!} \partial_k(t, \dots, t)(v) &= \\ &= l_1 t(v) - t \circ M_V(v) \\ &+ \frac{1}{2} l_2(t(v_{(1)}), t(v_{(2)})) - t \left(\Phi_{t(v_{(1)})} v_{(2)} \right) + \cdots + \\ &+ \frac{1}{k!} l_k(t(v_{(1)}), \dots, t(v_{(k)})) - \frac{1}{(k-1)!} t \left(\Phi_{t(v_{(1)}) \odot \dots \odot t(v_{(k-1)})} v_{(k)} \right). \end{aligned}$$

Let $T : \bar{S}(V) \rightarrow \bar{S}(E)$ be the morphism of coalgebras defined by $t : \bar{S}(V) \rightarrow E$. Maurer-Cartan equation can be written as

$$l \circ T(v) - t \circ M_V(v) - t \Phi_{T(v_{(1)})} v_{(2)} = 0,$$

which is equivalent to T being an \mathcal{O} -operator (see Equation (15)). \square

4. DEFORMATION OF \mathcal{O} -OPERATORS

We prove that each Maurer-Cartan element of a special graded Lie subalgebra of $\text{Coder}(\bar{S}(E \oplus V))$ encodes a Lie ∞ -algebra structure on E and a curved Lie ∞ -action of E on V . We study deformations of \mathcal{O} -operators.

4.1. Maurer-Cartan elements of $\text{Coder}(\bar{S}(E \oplus V))$. Let E and V be two graded vector spaces and consider the graded Lie algebra $\mathfrak{L} := (\text{Coder}(\bar{S}(E \oplus V)), [\cdot, \cdot]_c)$. Since $\bar{S}(E \oplus V) \simeq \bar{S}(E) \oplus (\bar{S}(E) \otimes \bar{S}(V)) \oplus \bar{S}(V)$, the space $M := \text{Coder}(\bar{S}(E))$ of coderivations of $\bar{S}(E)$ can be seen as a graded Lie subalgebra of \mathfrak{L} . Also, the space R of coderivations defined by linear maps of the space $\text{Hom}((\bar{S}(E) \otimes \bar{S}(V)) \oplus \bar{S}(V), V)$ can be embedded in \mathfrak{L} . We will use the identifications $M \equiv \text{Hom}(\bar{S}(E), E)$ and $R \equiv \text{Hom}((\bar{S}(E) \otimes \bar{S}(V)) \oplus \bar{S}(V), V)$. Given $\rho \in R$, we will denote by ρ_0 the restriction of the linear map ρ to $\bar{S}(V)$ and by ρ_x the linear map obtained by restriction of ρ to $\{x\} \otimes \bar{S}(V)$, with $x \in \bar{S}(E)$. We set $\mathfrak{L}' := M \oplus R$.

Proposition 4.1. *The space \mathfrak{L}' is a graded Lie subalgebra of $\mathfrak{L} = \text{Coder}(\bar{S}(E \oplus V))$.*

Proof. Given $m \oplus \rho, m' \oplus \rho' \in \mathfrak{L}'$, let us see that

$$[m \oplus \rho, m' \oplus \rho']_{RN} = [m, m']_{RN} \oplus ([m, \rho']_{RN} + [\rho, m']_{RN} + [\rho, \rho']_{RN})$$

is an element of \mathfrak{L}' . It is obvious that $[m, m']_{RN} \in \text{Hom}(\bar{S}(E), E)$. Consider m^D and ρ^D the coderivations of $\bar{S}(E \oplus V)$ defined by the morphisms m and ρ , respectively. For $x \in \bar{S}(E)$ and $v \in \bar{S}(V)$ we have,

$$\begin{aligned} [m, \rho']_{RN}(x) &= [m, \rho']_{RN}(v) = 0 \\ [m, \rho']_{RN}(x \otimes v) &= \left(m \circ \rho'^D - (-1)^{|m||\rho'|} \rho' \circ m^D \right) (x \odot v) \\ &= -(-1)^{|m||\rho'|} \rho'_{m^D(x)}(v) \in V \end{aligned}$$

and

$$\begin{aligned}
[\rho, \rho']_{RN}(x) &= 0 \\
[\rho, \rho']_{RN}(v) &= \rho \circ \rho'^D(v) - (-1)^{|\rho||\rho'|} \rho' \circ \rho^D(v) \in V \\
[\rho, \rho']_{RN}(x \otimes v) &= \underbrace{(-1)^{|x||\rho'|} \rho_x(\rho_0'^D(v)) + (-1)^{|x_{(1)}||\rho'|} \rho_{x_{(1)}}(\rho_{x_{(2)}}'^D(v)) + \rho_0(\rho_x'^D(v))}_{\in V} \\
&\quad - (-1)^{|\rho||\rho'|} \underbrace{\left((-1)^{|x||\rho|} \rho'_x(\rho_0^D(v)) + (-1)^{|x_{(1)}||\rho|} \rho'_{x_{(1)}}(\rho_{x_{(2)}}^D(v)) + \rho'_0(\rho_x^D(v)) \right)}_{\in V},
\end{aligned}$$

which proves that $[m, \rho']_{RN} + [\rho, m']_{RN} + [\rho, \rho']_{RN} \in \text{Hom}((\bar{S}(E) \otimes \bar{S}(V)) \oplus \bar{S}(V), V)$. \square

Next theorem shows that an element $m \oplus \rho \in \mathfrak{L}'$ which is a Maurer-Cartan of $\mathfrak{L} = \text{Coder}(\bar{S}(E \oplus V))$ encodes a Lie ∞ -algebra structure on E and an action of E on the Lie ∞ -algebra V .

Theorem 4.2. *Let E and V be two graded vector spaces and $m \oplus \rho \in \mathfrak{L}' = M \oplus R$. Then, $m \oplus \rho$ is a Maurer-Cartan element of \mathfrak{L}' if and only if m^D defines a Lie ∞ -structure on E and ρ is a curved Lie ∞ -action of E on V .*

Proof. We have

$$(19) \quad [m \oplus \rho, m \oplus \rho]_{RN} = 0 \Leftrightarrow \begin{cases} [m, m]_{RN} = 0 \\ 2[m, \rho]_{RN} + [\rho, \rho]_{RN} = 0. \end{cases}$$

Similar computations to those in the proof of Proposition 4.1 give, for all $v \in \bar{S}(V)$ and $x \in \bar{S}(E)$,

$$\begin{aligned}
&\begin{cases} (2[m, \rho]_{RN} + [\rho, \rho]_{RN})(v) = 0 \\ (2[m, \rho]_{RN} + [\rho, \rho]_{RN})(x \otimes v) = 0 \end{cases} \\
&\Leftrightarrow \begin{cases} \rho_0 \circ \rho_0^D(v) = 0 \\ \rho_{m^D(x)}(v) = \left(-[\rho_0, \rho_x]_{RN} - \frac{(-1)^{|x_{(1)}|}}{2} [\rho_{x_{(1)}}, \rho_{x_{(2)}}]_{RN} \right)(v). \end{cases}
\end{aligned}$$

Since $m \oplus \rho$ is a degree +1 element of \mathfrak{L}' , the right hand-side of (19) means that m^D defines a Lie ∞ -algebra structure on E and $\rho = \sum_{k \geq 0} \rho_k$ is a curved Lie ∞ -action of E on V . Notice that $\rho_0^D : \bar{S}(V) \rightarrow \bar{S}(V)$ equips V with a Lie ∞ -structure.

Reciprocally, if (E, m^D) is a Lie ∞ -algebra and ρ is a curved Lie ∞ -action of E on V , the degree +1 element $m \oplus \rho$ of \mathfrak{L}' is a Maurer-Cartan element of \mathfrak{L}' . \square

Next proposition gives the Lie ∞ -algebra that controls the deformations of the actions of E on V [3].

Proposition 4.3. *Let $m \oplus \rho$ be a Maurer-Cartan element of \mathfrak{L}' and $m' \oplus \rho'$ a degree +1 element of \mathfrak{L}' . Then, $m \oplus \rho + m' \oplus \rho'$ is a Maurer-Cartan element of \mathfrak{L}' if and only if $m' \oplus \rho'$ is a Maurer-Cartan element of $\mathfrak{L}'^{m \oplus \rho}$. Here, $\mathfrak{L}'^{m \oplus \rho}$ denotes the DGLA which is the twisting of \mathfrak{L}' by $m \oplus \rho$.*

4.2. Deformation of \mathcal{O} -operators. Let \mathfrak{h} be the abelian Lie subalgebra of $\mathfrak{L} = \text{Coder}(\bar{S}(E \oplus V))$ considered in Proposition 3.13 and $\mathcal{P} : \mathfrak{L} \rightarrow \mathfrak{h}$ the projection onto \mathfrak{h} . Let $\Delta \in \mathfrak{L}'$ be a Maurer-Cartan element of \mathfrak{L} . Then, $(\mathfrak{L}, \mathfrak{h}, \mathcal{P}, \Delta)$ is a V -data and \mathfrak{h} has a Lie ∞ -structure given by the brackets:

$$\partial_k(a_1, \dots, a_k) = \mathcal{P}([\dots[\Delta, a_1]_{RN} \dots]_{RN}, a_k]_{RN}), \quad k \geq 1.$$

We denote by \mathfrak{h}_Δ the Lie ∞ -algebra \mathfrak{h} equipped with the above structure.

The V -data $(\mathfrak{L}, \mathfrak{h}, \mathcal{P}, \Delta)$ also provides a Lie ∞ -algebra structure on $\mathfrak{L}[1] \oplus \mathfrak{h}$, that we denote by $(\mathfrak{L}[1] \oplus \mathfrak{h})_\Delta$, with brackets [10]:

$$(20) \quad \begin{cases} q_1^\Delta((x, a_1)) = (-[\Delta, x]_{RN}, \mathcal{P}(x + [\Delta, a_1]_{RN})) \\ q_2^\Delta(x, x') = (-1)^{\deg(x)} [x, x']_{RN} \\ q_k^\Delta(x, a_1, \dots, a_{k-1}) = \mathcal{P}([\dots [x, a_1]_{RN}, a_2]_{RN} \dots a_{k-1}]_{RN}), \quad k \geq 2, \\ q_k^\Delta(a_1, \dots, a_k) = \partial_k(a_1, \dots, a_k), \quad k \geq 1, \end{cases}$$

$x, x' \in \mathfrak{L}[1]$ and $a_1, \dots, a_{k-1} \in \mathfrak{h}$. Here, $\deg(x)$ is the degree of x in \mathfrak{L} .

Moreover, since \mathfrak{L}' is a Lie subalgebra of \mathfrak{L} satisfying $[\Delta, \mathfrak{L}'] \subset \mathfrak{L}'$, the brackets $\{q_k^\Delta\}_{k \in \mathbb{N}}$ restricted to $\mathfrak{L}'[1] \oplus \mathfrak{h}$ define a Lie ∞ -algebra structure on $\mathfrak{L}'[1] \oplus \mathfrak{h}$, that we denote by $(\mathfrak{L}'[1] \oplus \mathfrak{h})_\Delta$. Notice that the restrictions of the brackets $\{q_k^\Delta\}$ to $\mathfrak{L}'[1] \oplus \mathfrak{h}$ are given by the same expressions as in (20) except for $k = 1$:

$$q_1^\Delta((x, a_1)) = (-[\Delta, x]_{RN}, \mathcal{P}([\Delta, a_1]_{RN})) = (-[\Delta, x]_{RN}, \partial_1(a_1)),$$

because $\mathcal{P}(\mathfrak{L}') = 0$. Of course, \mathfrak{h}_Δ is a Lie ∞ -subalgebra of $(\mathfrak{L}'[1] \oplus \mathfrak{h})_\Delta$.

Remark 4.4. The brackets (20) that define the Lie ∞ -algebra structure of $(\mathfrak{L}[1] \oplus \mathfrak{h})_\Delta$ coincide with those of \mathfrak{h}_Δ for $x = x' = 0$. So, an easy computation yields

$$t \in \text{MC}(\mathfrak{h}_\Delta) \Leftrightarrow (0, t) \in \text{MC}(\mathfrak{L}'[1] \oplus \mathfrak{h})_\Delta.$$

Theorem 3 in [2] yields:

Proposition 4.5. *Consider the V -data $(\mathfrak{L}, \mathfrak{h}, \mathcal{P}, \Delta)$, with $\Delta \in \text{MC}(\mathfrak{L}')$ and let t be a degree zero element of \mathfrak{h} . Then,*

$$t \in \text{MC}(\mathfrak{h}_\Delta) \Leftrightarrow (\Delta, t) \in \text{MC}(\mathfrak{L}[1] \oplus \mathfrak{h})_\Delta.$$

Recall that, given an element $t \in \mathfrak{h} = \text{Hom}(\bar{S}(V), E)$, the corresponding morphism of coalgebras $T : \bar{S}(V) \rightarrow \bar{S}(E)$ is an \mathcal{O} -operator if and only if t is a Maurer-Cartan element of \mathfrak{h}_Δ (Proposition 3.18). Moreover, given a Maurer-Cartan element $m \oplus \rho$ of \mathfrak{L}' , we know from Theorem 4.2 that (E, m^D) is a Lie ∞ -algebra and ρ is a curved Lie ∞ -action of E on V . So, an \mathcal{O} -operator can be seen as a Maurer-Cartan element of the Lie ∞ -algebra $(\mathfrak{L}'[1] \oplus \mathfrak{h})_\Delta$:

Proposition 4.6. *Let E and V be two graded vector spaces. Consider a morphism of coalgebras $T : \bar{S}(V) \rightarrow \bar{S}(E)$ defined by $t \in \text{Hom}(\bar{S}(V), E)$, and the V -data $(\mathfrak{L}, \mathfrak{h}, \mathcal{P}, \Delta)$, with $\Delta := m \oplus \rho \in \text{MC}(\mathfrak{L}')$. Then, T is an \mathcal{O} -operator on E with respect to the curved Lie ∞ -action ρ if and only if (Δ, t) is a Maurer-Cartan element of $(\mathfrak{L}'[1] \oplus \mathfrak{h})_\Delta$.*

Corollary 4.7. *If T is an \mathcal{O} -operator on the Lie ∞ -algebra (E, m^D) with respect to the curved Lie ∞ -action ρ of E on V , then $((\mathfrak{L}'[1] \oplus \mathfrak{h})_{m \oplus \rho})^{(m \oplus \rho, t)}$ is a Lie ∞ -algebra.*

As a consequence of Theorem 3 in [2], we obtain the Lie ∞ -algebra that controls the deformation of \mathcal{O} -operators on E with respect to a fixed curved Lie ∞ -action on V :

Corollary 4.8. *Let E and V be two graded vector spaces and consider the V -data $(\mathfrak{L}, \mathfrak{h}, \mathcal{P}, \Delta := m \oplus \rho)$. Let T be an \mathcal{O} -operator on (E, m^D) with respect to the curved Lie ∞ -action ρ and $T' : \bar{S}(V) \rightarrow \bar{S}(E)$ a (degree zero) morphism of coalgebras defined by $t' \in \text{Hom}(\bar{S}(V), E)$. Then, $T + T'$ is an \mathcal{O} -operator on E*

with respect to the curved Lie ∞ -action ρ if and only if (Δ, t') is a Maurer-Cartan element of $(\mathfrak{L}'[1] \oplus \mathfrak{h})_{\Delta}^{(\Delta, t)}$.

Proof. Let $t \in \mathfrak{h}$ be the morphism defined by T . Then [2],

$$(\Delta, t + t') \in \text{MC}(\mathfrak{L}'[1] \oplus \mathfrak{h})_{\Delta} \Leftrightarrow (\Delta, t') \in \text{MC}(\mathfrak{L}'[1] \oplus \mathfrak{h})_{\Delta}^{(\Delta, t)}.$$

□

REFERENCES

- [1] Baxter, G.: An analytic problem whose solution follows from a simple algebraic identity. *Pacific J. Math.* (1960) **10**, no. 2, 731–742.
- [2] Frégier, Y. and Zambon, M.: Simultaneous deformations of algebras and morphisms via derived brackets. *Journal of Pure and Applied Algebra* (2015) **219** 5344–5362.
- [3] Getzler, E.: Lie theory for nilpotent L_{∞} -algebras. *Ann. Math.* (2009) **170**, no. 2, 271–301.
- [4] Kupersmidt, B.A.: What a Classical r -Matrix Really Is. *J. Nonlinear Math. Phys.* (1999) **6**, no. 4, 448–488.
- [5] Lada, T. and Stasheff, J.: Introduction to SH Lie algebras for physicists. *Internat. J. Theoret. Phys.* (1993) **32**, no. 7, 1087–1103.
- [6] Lazarev, A.; Sheng, Y. and Tang, R.: Deformations and homotopy theory of relative Rota-Baxter Lie algebras. *Comm. Math. Phys.* (2021) **383**, 595–631.
- [7] Lada, T. and Markl, M.: Strongly homotopy Lie algebras. *Comm. Algebra* (1995) **23**, no. 6, 2147–2161.
- [8] Mehta, R. and Zambon M.: L_{∞} -algebra actions. *Differential Geometry and its Applications* (2012) **30**, 576–587.
- [9] Tang, R., Bai, C., Guo, L., Sheng, Y.: Homotopy Rota-Baxter operators, homotopy O -operators and homotopy post-Lie algebras. *arXiv:1907.13504*.
- [10] Voronov, T.: Higher derived brackets and homotopy algebras. *J. Pure Appl. Algebra* (2005) **202**, no. 1-3, 133–153.

UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, APARTADO 3008, EC SANTA CRUZ, 3001-501 COIMBRA, PORTUGAL
Email address: raquel@mat.uc.pt

UNIVERSITY OF COIMBRA, CMUC, DEPARTMENT OF MATHEMATICS, APARTADO 3008, EC SANTA CRUZ, 3001-501 COIMBRA, PORTUGAL
Email address: jmcosta@mat.uc.pt