

POISSON METRICS AND HIGGS BUNDLES OVER NONCOMPACT KÄHLER MANIFOLDS

DI WU AND XI ZHANG

ABSTRACT. In this paper, we study the existence of Poisson metrics on flat vector bundles over noncompact Riemannian manifolds and discuss related consequence, specially on the applications in Higgs bundles, towards generalizing Corlette-Donaldson-Hitchin-Simpson's nonabelian Hodge correspondence to noncompact Kähler manifolds setting.

1. INTRODUCTION

1.1. Background. Let (E, D) be a vector bundle over a Riemannian manifold (M, g) , we say (E, D) is simple if it admits no proper D -invariant sub-bundle and (E, D) is semi-simple¹ if it is a direct sum of (E_i, D_i) and each of them is simple. The connection D is called irreducible if it admits no nontrivial covariantly constant section of $\text{End}(E)$. Given any metric H on E , it splits the connection as

$$(1.1) \quad D = D_H + \psi_H,$$

where D_H is a connection preserving H and $\psi_H \in \Omega^1(\text{End}(E))$ is self-adjoint with respect to H . We say H is harmonic if it satisfies

$$(1.2) \quad D_H^* \psi_H = 0,$$

where D_H^* is the adjoint of D_H . The notion of harmonic map originated from Eells-Sampson [17] and was generalized to the notion of harmonic metric by Corlette [8]. The motivation for this name may be explained as follows. By the Riemann-Hilbert correspondence, suppose $\rho_D : \pi_1(M) \rightarrow GL(r)$ be the representation corresponding to a flat bundle (E, D) , then any metric H on E induces a ρ_D -equivariant map f_H from \tilde{M} to $GL(r)/U(r)$, where \tilde{M} is the universal of M . Then we can see that H being harmonic is equivalent to f_H being a harmonic map or df_H being a harmonic 1-form with values in $f_H^{-1}T(GL(r)/U(r))$.

The study of harmonic metrics is a fundamental problem which remains an active branch of current research in differential geometry, it builds a bridge between geometry analysis and many other areas including group representations, nonabelian Hodge theory. Let (M, g) be a compact Riemannian manifold, the main existence result is the following

Theorem 1.1 (Donaldson [14], Corlette [8]). *A flat bundle admits a harmonic metric if and only if it arises from a semi-simple representation of the fundamental group $\pi_1(M)$.*

2020 *Mathematics Subject Classification.* 53C07, 53C21.

Key words and phrases. Flat bundle, Noncompact manifold, Analytically stability, Poisson metric, Higgs bundle, Nonabelian Hodge correspondence.

The research was partially supported by the project "Analysis and Geometry on Bundle" of Ministry of Science and Technology of the People's Republic of China, No. SQ2020YFA070080. The authors are partially supported by NSF in China No. 11625106, 11801535 and 11721101.

¹In literature, one sometimes says that complete reducible or reductive.

If (M, g) happens to be a compact Kähler manifold, the significance of this theorem lies in the construction of Higgs bundle. In fact, the pair $(D_H^{0,1}, \psi_H^{1,0})$ will determine a poly-stable Higgs structure, provided H being harmonic. For this reason, sometimes we then say that D satisfies the Hitchin's self-duality equation [19] and we call D a Hitchin connection. Combing this with the Donaldson-Uhlenbeck-Yau theorem([13, 44]) for Higgs bundles [19, 39], one obtains Corlette-Donaldson-Hitchin-Simpson's nonabelian Hodge correspondence:

Theorem 1.2 ([8, 14, 19, 39]). *On a compact Kähler manifold, there is an one to one correspondence between the category of seme-simple flat bundles and the category of poly-stable Higgs bundles with vanishing Chern numbers.*

This identification can be traced back to Narasimhan-Seshadri [34] and has yielded several fascinating topological results, specially on the fundamental groups of compact Kähler manifolds, see [10, 41] for more details. In [21, 22], Jost-Zuo studied the representations of fundamental groups for quasi-projective varieties by using the theory of harmonic metrics on quasi-projective varieties. Besides that, there are also some partial existence results for harmonic metrics on noncompact manifolds. Simpson [40] obtained the existence of harmonic metrics on punctured Riemannian surface in his study of nonabelian Hodge theory over noncompact curves. From the view point of equivariant harmonic maps, Li [26] considered the case that $M = \overline{M} - D$ equipped with a Poincaré-like metric g , where \overline{M} is a compact complex manifold and D be a divisor with normal crossings in \overline{M} . Recently, Mochizuki makes a significant process towards the generations of the Kobayashi-Hitchin correspondence and the nonabelian Hodge correspondence, the existence of pluriharmonic metrics for good filtered flat bundles over smooth projective varieties play an important role, see [29–32]. There are many interesting and important literature concerning the existence of harmonic metrics and related topics, see [3, 6, 9, 12, 20, 25, 35, 48] and the references therein.

Flat bundles occur in great abundance since it corresponds local systems and representations of fundamental groups. It was observed by Loftin [27] that for an affine manifold M , any flat bundle (E, D) over M corresponds a holomorphic bundle \tilde{E} over the tangent bundle TM . And then the classical Donaldson-Uhlenbeck-Yau theorem can be generalized to the affine case, see [5, 27, 38]. Recently, Collins-Jacob-Yau [7] found that the dimension reduction of the Hermitian-Yang-Mills equation on \tilde{E} is given by the Poisson metric equation on (E, D) . Here we say H is a Poisson metric if

$$(1.3) \quad D_H^* \psi_H = c_H \text{id}_E,$$

where c_H is a function on M . Motivated by this and the understanding of stable vector bundles on K3 surface in the large complex structure limit, Collins-Jacob-Yau [7] focused on the research of Poisson metrics on flat vector bundles. Under some suitable assumptions, they related the existence of Poisson metrics on flat bundles with parabolic structures over punctured Riemann surfaces to certain notion of slope stability.

1.2. Main results. In the first part of this paper, we shall attack the existence of Poisson metrics on flat bundles over noncompact Riemannian manifolds of arbitrary dimension. Below we state our first theorem, which can be interpreted as a noncompact version of the result of Donaldson and Corlette.

Theorem 1.3. *Let (M, g) be a Riemannian manifold(not necessarily compact) satisfying the Assumptions 2.1 and either Assumption 2.2 or complete with finite volume, E be a vector bundle over M equipped with a connection D and a metric K such that $|D_K^* \psi_K| \leq C\phi$ for a constant C . Then it admits a unique Poisson metric H such that $\det h = 1$, $|h| \in L^\infty$ and*

$|Dh| \in L^2$ for $h = K^{-1}H$, provided D being flat and K -analytically stable. Conversely, if D is irreducible, it admits such a Poisson metric only if D is K -analytically stable.

For the existence statement, we actually only need the Assumption 2.1 and the condition $|(D_K^*\psi_K)^\perp| \leq C\phi$, where $(D_K^*\psi_K)^\perp$ is the trace-free part of $D_K^*\psi_K$, see Proposition 3.8 for details. The Assumption 2.1 is introduced by Mochizuki [33] for Kähler manifolds and the volume can be infinite. The notion of analytically stability in Theorem 1.3 will be introduced in Section 2.2. If M is compact, we know that D being analytically stable(poly-stable) if and only if it is simple(semi-simple), while the correct substitution should be replaced by analytically stability in noncompact setting. Moreover, if (M, g) be a Kähler manifold and D being flat, the definition coincides with that in Simpson's paper [40], see Remark 2.1.

The second statement in Theorem 1.3 indicates that D is analytically stable, not only with respect to the Poisson metric, but also the background metric K . It may be mentioned that it is facilitated by the feature of the stability of vector bundles, while the issue is more complicated in the setting of holomorphic bundles. We will use Mochizuki's exhaustion method in [33], where he proved the existence of exhaustion function f . Denote by M_s the set where $f \leq s$ and we firstly consider the parabolic heat flow (3.1) on M_s , which can be regard as the vector bundle analogy of Eells-Sampson's flow for harmonic maps or Donaldson's heat flow [13, 15]. For arbitrary D , we show the existence and uniqueness of the long-time solution to (3.1), also the Dirichlet boundary problems for harmonic metric and Poisson metric are uniquely solvable, see Proposition 3.3 and Proposition 3.4. Next we observe in Proposition 3.5 that the crucial identity (3.21), based on which several core a priori estimates are established. Once the zeroth estimate of a sequence of the solutions to the Dirichlet boundary problem is obtained, the evolved metrics are completely controlled and will converge to a Poisson metric. Finally, we prove the uniqueness and the second statement by comparing the analytically stabilities with respect to different reference fiber metrics.

We mention that Corlette [8] studied the following heat flow

$$(1.4) \quad \frac{\partial D_t}{\partial t} = -D_t D_{t,K}^* \psi_{D_t,K},$$

on flat complex vector bundles (E, D_0) over compact Riemannian manifolds, where D_t is a time-dependent family of connections in the orbit of D_0 . Corlette proved the convergence of D_t at $t = \infty$ with the property $D_{\infty,K} \psi_{D_{\infty,K}} = 0$ and if (E, D_0) is simple, D_{∞} must lie in the complex gauge orbit of D_0 . Corlette's proof relies on Uhlenbeck's deep compactness theorem [43] and cannot be generalized to noncompact setting directly. Even in compact case, here the discussion is different from that in [8] and we bypass Uhlenbeck's compactness theorem.

As a corollary of (2.19) and Theorem 1.3, we have

Corollary 1.1. *Under the same assumptions in Theorem 1.3 and $\text{tr } D_K^* \psi_K = 0$, there exist a unique harmonic metric H such that $\det h = 1$, $|h| \in L^\infty$ and $|Dh| \in L^2$ for $h = K^{-1}H$, provided D being flat and K -analytically stable. Conversely, D is K -analytically stable if it is irreducible and admits such a harmonic metric.*

By (2.22), we know that

$$(1.5) \quad \text{tr } D_H^* \psi_H - \text{tr } D_K^* \psi_K = -\frac{1}{2} \text{tr } D_K^* (h^{-1} \delta_K h) = \frac{1}{2} \Delta \log \det h,$$

where $h = K^{-1}H$ and $\delta_K = D_K - \psi_K$. So by solving a scalar Poisson equation(Theorem 4.3 in [45]) and employing Theorem 1.3, it is easy to conclude

Corollary 1.2. *Let (M, g) be a Riemannian manifold satisfying the Assumption 2.1 and Assumption 2.3 with $\phi = 1$, E be a vector bundle over M equipped with a connection D and a metric K such that $|D_K^* \psi_K| \in L^\infty$, $|\psi_K| \in L^2$. Then it admits a unique harmonic metric H such that $\det h = 1$, $|h| \in L^\infty$ and $|Dh| \in L^2$ for $h = K^{-1}H$, provided D being flat and K -analytically stable. Conversely, D is K -analytically stable if it is irreducible and admits such a harmonic metric.*

There is a remark on the situation that there is no stability(simple) assumption. For an arbitrary connection on the vector bundle over a compact Riemannian manifold, there is no obstruction to the existence of L^∞ -approximate solutions of (1.2), see Proposition 3.2.

In the second part of this paper, we shall discuss related applications and consequences after Theorem 1.3. Recall a Higgs bundle $(E, \bar{\partial}_E, \theta)$ over a complex manifold X consists of a holomorphic vector bundle $(E, \bar{\partial}_E)$ over X together with holomorphic section $\theta \in \Omega^{1,0}(\text{End}(E))$ such that $\theta \wedge \theta = 0$. For a Hermitian metric K , we consider the Hitchin-Simpson connection

$$(1.6) \quad D_{\bar{\partial}_E, \theta, K} = \partial_{\theta, K} + \bar{\partial}_{E, \theta}, \quad \partial_{\theta, K} = \partial_K + \theta^* K, \quad \bar{\partial}_{E, \theta} = \bar{\partial}_E + \theta,$$

where $D_{\bar{\partial}_E, K} = \partial_K + \bar{\partial}_E$ is the Chern connection. If the curvature $F_{\bar{\partial}_E, \theta, K} = D_{\bar{\partial}_E, \theta, K}^2$ vanishes, we will call $(\bar{\partial}_E, \theta, K)$ a Higgs flat structure on the underlying bundle E . In particular, if the Chern curvature $F_{\bar{\partial}_E, K} = D_{\bar{\partial}_E, K}^2$ vanishes (that is, $\theta = 0$), we then say $(\bar{\partial}_E, K)$ a flat holomorphic structure.

As an application of Theorem 1.3, we have

Theorem 1.4. *Let (X, ω) be a Kähler manifold satisfying the Assumption 2.1 and (E, D) be a K -analytically stable flat bundle over X with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C , then*

- (1) *If (X, ω) is a complex curve, there is a Higgs flat structure on E .*
- (2) *If (X, ω) is complete with bounded Ricci curvature from below and $|\psi_K| \in L^2$, there is a Higgs flat structure on E .*
- (3) *If (X, ω) is complete with nonnegative Ricci curvature and $|\psi_K| \in L^2$, there is a flat holomorphic structure on E .*

Especially, the third statement of the conclusions in Theorem 1.4 indicates there is a metric compatible flat connection on E . Finding metric compatible flat connections may be of interest in its own right. By Corlette's work in [8], it is easy to see the same conclusion also holds for a flat bundle over a compact Riemannian manifold with nonnegative Ricci curvature and so it follows that the Euler class $e(E)$ of E is zero.

In order to prove Theorem 1.4, we need to construct Higgs structures by employing Theorem 1.3, see Proposition 4.3 and Proposition 4.4, which is a key step to establish the Corlette-Donaldson-Hitchin-Simpson's correspondence. If X is compact and H being harmonic, this is equivalent to determine whether H being pluriharmonic, the issue does not appear and it is essentially due to Siu-Sampson [37, 42] and Corlette [8], see also [41]. If X is noncompact, see [22] for case that the complement of a divisor in a compact Kähler manifold.

Secondly, we will focus on generalizing the Corlette-Donaldson-Hitchin-Simpson's correspondence to noncompact Kähler manifolds setting. Following Simpson [39], if $(\bar{\partial}_E, \theta)$ is a Higgs structure on E , the K -analytically degree is defined by

$$(1.7) \quad \deg_\omega(E, K) = \int_X \sqrt{-1} \text{tr } \Lambda_\omega F_{\bar{\partial}_E, \theta, K} \, \text{dvol}_\omega.$$

If $\mathcal{V} \subseteq \mathcal{O}_X(E)$ is a sub-sheaf, then it can be seen as a sub-bundle V outside a singular set $\Sigma_{\mathcal{V}}$ of codimension at least two. Let π_K denotes the projection onto V and K restricts to

a metric on V so that we can define $\deg_\omega(\mathcal{V}, K)$ by integrating outside $\Sigma_{\mathcal{V}}$. We have the Chern-Weil formula written in terms of $\Lambda_\omega F_{\bar{\partial}_E, \theta, K}$,

$$(1.8) \quad \deg_\omega(\mathcal{V}, K) = \int_{X \setminus \Sigma_{\mathcal{V}}} \left(\sqrt{-1} \operatorname{tr}(\pi_K \circ \Lambda_\omega F_{\bar{\partial}_E, \theta, K}) - |\bar{\partial}_E, \theta \pi_K|_K^2 \right) \operatorname{dvol}_\omega.$$

$(\bar{\partial}_E, \theta)$ is called K -analytically stable if for any proper sub-Higgs sheaf $\mathcal{V} \subset \mathcal{O}_X(E)$, it holds

$$(1.9) \quad \frac{\deg_\omega(\mathcal{V}, K)}{\operatorname{rank}(\mathcal{V})} < \frac{\deg_\omega(E, K)}{\operatorname{rank}(E)}.$$

From now on, we fix a Hermitian metric K on the vector bundle E , write

- $\mathcal{M}_{Flat, K}$: the set of isomorphic classes of K -analytically stable and irreducible flat structures such that $|(D_K^* \psi_K)^\perp| \in L^1$.
- $\mathcal{M}_{Higgs, K}$: the set of isomorphic classes of K -analytically stable and irreducible Higgs structures with $c_1(E, K) = 0$ and $|\Lambda_\omega F_{\bar{\partial}_E, \theta, K}| \in L^1$.
- $\mathcal{M}_{Flat, K, \phi}$: the set of isomorphic classes of K -analytically stable and irreducible flat structures such that $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C .
- $\mathcal{M}_{Higgs, K, \phi}$: the set of isomorphic classes of K -analytically stable and irreducible Higgs structures with $c_1(E, K) = 0$ and $|\Lambda_\omega F_{\bar{\partial}_E, \theta, K}| \leq C\phi$ for a constant C .

In the above, isomorphic means that they are in the the same \mathcal{G} -orbit, where \mathcal{G} denotes the set of L^∞ -sections of $\operatorname{Aut}(E)$. We say a Higgs structure $(\bar{\partial}_E, \theta)$ is irreducible if $D_{\bar{\partial}_E, \theta, H}$ is irreducible, for any Hermitian-Einstein metric H with $\det h = 1$ and $|h| \in L^\infty$ for $h = K^{-1}H$.

Our third theorem can be stated as the follows.

Theorem 1.5. *Assume (X, ω) be a noncompact complex curve satisfying the Assumption 2.1 and either Assumption 2.2 or complete with finite volume, then we have*

- (1) *There is map from $\mathcal{M}_{Flat, K, \phi}$ to $\mathcal{M}_{Higgs, K}$.*
- (2) *There is map from $\mathcal{M}_{Higgs, K, \phi}$ to $\mathcal{M}_{Flat, K}$.*

Notice that the complex plane \mathbb{C} and compact Riemann surfaces punctured finite points satisfy the assumptions in Theorem 1.5. Moreover, the objects are analytically stable with respect to the same metric K . For higher dimensional case, under some extra technical assumptions(see [33, 39]), the direction that from K -analytically stable and irreducible Higgs structures to K -analytically stable and irreducible flat structures can be finished by the results of Simpson [39], Mochizuki [33] and the discussion as that in (2) of Theorem 1.5. Conversely, start with a K -analytically stable and irreducible flat structure, using Proposition 4.4(or Proposition 4.5) instead, we can obtain an irreducible Higgs structure. In the process, the analytic estimates in Theorem 1.5 remain valid, but currently it is not clear that when the resulting Higgs structure is stable with respect to K . We will further discuss this problem in the future.

Theorem 1.5 is established based on the existence of Poisson metrics(Theorem 1.3) and Hermitian-Einstein metrics([33, 39]), while analytic estimates and the comparison of different analytically stabilities play the significant roles. If X is compact, Theorem 1.5 and the corresponding result for high dimensional case reduce to the classical nonabelian Hodge correspondence on compact Kähler manifolds. This correspondence has been generalised to parabolic and noncompact setting, specially on the case of a projective variety equipped with normal crossing divisors, in a series of works including those of Simpson [40] and Biquard-Boalch [2] for the tame and wild case in complex dimension one. And the higher dimensional case can be found in [1, 29, 31, 32].

1.3. Organization. The rest of this paper is organized as follows. In Section 2, we discuss the assumptions on the base manifolds, introduce the notation of stability of vector bundles, and present some useful calculations. In Section 3, we solve the Dirichlet boundary problems for harmonic metric equation and Poisson metric equation, then give the proof of Theorem 1.3. In Section 4, we firstly investigate the uniqueness of Poisson metrics and then prove Theorem 1.4, Theorem 1.5. As a byproduct, also a vanishing theorem of Kamber-Tondeur classes is obtained.

2. PRELIMINARIES

2.1. Assumptions on the base spaces. Assume (M, g) be a Riemannian manifold, we list the following assumptions that needed in this paper.

Assumption 2.1. *There is a function $\phi : M \rightarrow \mathbb{R}_{\geq 0}$ with $\phi \in L^1$, such that if f is a nonnegative bounded function on M satisfying $\Delta f \geq -B\phi$ for a positive constant B in distribution sense, we have*

$$(2.1) \quad \sup_M f \leq C(B)(1 + \int_M f \phi \, d\text{vol}_g).$$

Furthermore, if f satisfies $\Delta f \geq 0$, we have $\Delta f = 0$.

Assumption 2.2. (M, g) admits an exhaustion function $\rho : M \rightarrow \mathbb{R}_{\geq 0}$ with $|\Delta \rho| \in L^1$.

Assumption 2.3. (M, g) admits an exhaustion function $\rho : M \rightarrow \mathbb{R}_{\geq 0}$ with bounded $\Delta \rho$.

The above assumptions are introduced for Kähler manifolds in [33] and [39].

Example 2.1. *Compact Riemannian manifolds satisfy Assumptions 2.1, 2.2 and 2.3.*

Example 2.2. *Let M be a Zariski open subset of a compact Kähler manifold $(\overline{X}, \overline{g})$, g is the restriction of \overline{g} , then it satisfies Assumptions 2.1, 2.2 and 2.3.*

Example 2.3. *Consider $(\mathbb{R}^2, g_{\mathbb{R}^2})$ with the Euclidean metric $g_{\mathbb{R}^2}$ and $\phi_{\mathbb{R}^2} = (1 + |r|^2)^{-1-\delta}$, $\delta > 0$, then it has infinite volume and satisfies Assumptions 2.1. Moreover, $(\mathbb{R}^2, \phi_{\mathbb{R}^2} g_{\mathbb{R}^2})$ satisfies Assumptions 2.1 and has finite volume.*

Example 2.4. *Any product manifold $(\mathbb{R}^2, g_{\mathbb{R}^2}) \times (N, g_N)$ has infinite volume and satisfies Assumptions 2.1, where (N, g_N) is a compact Riemannian manifold.*

2.2. Stability of vector bundles. Assume (E, D) be a vector bundle over a Riemannian manifold (M, g) , equipped with a metric K on E .

Definition 2.1. *The K -analytically degree of (E, D) is defined by*

$$(2.2) \quad \deg_g(E, D, K) = - \int_M \text{tr } D_K^* \psi_K \, d\text{vol}_g.$$

For any D -invariant sub-bundle $S \subseteq E$, we define

$$(2.3) \quad \deg_g(S, D, K) = - \int_M \text{tr } D_{K_S}^* \psi_{K_S} \, d\text{vol}_g,$$

where K_S is the restricted metric on S . Then (E, D) is said to be K -analytically stable if for any nontrivial D -invariant sub-bundle S , it holds that

$$(2.4) \quad \frac{\deg_g(S, D, K)}{\text{rank}(S)} < \frac{\deg_g(E, D, K)}{\text{rank}(E)}.$$

Remark 2.1. If (M, ω) be a Kähler manifold and D is a flat connection, we have

$$(2.5) \quad \deg_g(E, D, K) = \deg_\omega(E, D^{0,1}, K) = -\deg_\omega(E, \delta_K^{0,1}, K),$$

where $\delta_K^{0,1} = D_K^{0,1} - \psi_K^{0,1}$, $\deg_\omega(E, D^{0,1}, K)$, $\deg_\omega(E, \delta_K^{0,1}, K)$ are the K -analytically degree of the holomorphic bundles $(E, D^{0,1})$ and $(E, \delta_K^{0,1})$ respectively.

For any D -invariant sub-bundle S , we have the Chern-Weil formula written in terms of $D_K^* \psi_K$,

$$(2.6) \quad \deg_g(S, D, K) = - \int_M \left(\text{tr}(\pi_K \circ D_K^* \psi_K) + \frac{1}{2} |D\pi_K|_K^2 \right) \text{dvol}_g,$$

where π_K is the orthogonal projection onto S .

Proposition 2.1. If a vector bundle (E, D) admits a Poisson metric K , it is an orthogonal direct sum of K -analytically stable bundles.

Proof. For any nontrivial D -invariant sub-bundle $S \subset E$, the Chern-Weil formula yields

$$(2.7) \quad \begin{aligned} \frac{\deg_g(S, D, K)}{\text{rank}(S)} &= \frac{- \int_M \left(\text{tr}(\pi_K \circ D_K^* \psi_K) + \frac{1}{2} |D\pi_K|_K^2 \right) \text{dvol}_g}{\text{rank}(S)} \\ &= \frac{- \int_M \left(\frac{\text{rank}(S)}{\text{rank}(E)} \text{tr} D_K^* \psi_K + \frac{1}{2} |D\pi_K|_K^2 \right) \text{dvol}_g}{\text{rank}(S)} \\ &\leq \frac{\deg_g(E, D, K)}{\text{rank}(E)}. \end{aligned}$$

If the equality holds, we obtain $D\pi_K = 0$. Let S^\perp denote the orthogonal complement of S in E with respect to K , then it is also a D -invariant sub-bundle such that $(E, D) \cong (S, D_S) \oplus (S^\perp, D_{S^\perp})$, where D_S and D_{S^\perp} are the induced connections. Moreover, the induced metrics K_S and K_{S^\perp} both are Poisson metrics and we complete the proof by an easy induction. \square

2.3. Some useful calculations. We present some important calculations, it is emphasized that we are not working under the local flat frame and hence most of them need not the assumption that the underlying connection is flat.

Given a vector bundle E over a Riemannian manifold (M, g) , the exterior differential operator $D : \Omega^p(E) \rightarrow \Omega^{p+1}(E)$ relative to a connection is given by

$$(2.8) \quad \begin{aligned} D\omega(e_0, \dots, e_p) &= \sum_{k=0}^p (-1)^k D_{e_k}(\omega(e_0, e_1, \dots, \hat{e}_k, \dots, e_p)) \\ &\quad + \sum_{k < l} (-1)^{k+l} \omega([e_k, e_l], e_1, \dots, \hat{e}_k, \dots, \hat{e}_l, \dots, e_p). \end{aligned}$$

Since the Levi-Civita connection on TM is torsion-free, we also have

$$(2.9) \quad D\omega(e_0, \dots, e_p) = \sum_{k=0}^p (-1)^k (\tilde{\nabla}_{e_k} \omega)(e_0, \dots, \hat{e}_k, \dots, e_p),$$

where $\tilde{\nabla}$ is the induced connection on $\Omega^*(E)$.

For a metric K on E , we define the point-wise inner product

$$(2.10) \quad K(\omega, \theta)(x) = \sum_{i_1 < \dots < i_p} (\omega(e_{i_1}, \dots, e_{i_p}), \theta(e_{i_1}, \dots, e_{i_p}))_K,$$

where $\omega, \theta \in \Omega^p(E)$ and $\{e_i\}_{i=1}^{\dim M}$ is the orthogonal unit basis of $T_x M$. Then the decomposition (1.1) is achieved by choosing ψ_K such that

$$(2.11) \quad K(\psi_K(X), Y) = \frac{1}{2}(K(DX, Y) + K(X, DY) - dK(X, Y)),$$

for any $X, Y \in \Omega^0(E)$. With respect to the Riemannian structures of E and TM , the co-differential operator $D_K^* : \Omega^p(E) \rightarrow \Omega^{p-1}(E)$ is characterized via the formula

$$(2.12) \quad \int_M K(D_K \omega, \theta) \, \text{dvol}_g = \int_M K(\omega, D_K^* \theta) \, \text{dvol}_g,$$

for any $\omega \in \Omega^{p-1}(E), \theta \in \Omega^p(E)$ and either ω or θ is compactly supported. Then it holds

$$(2.13) \quad D_K^* \theta(e_1, \dots, e_{p-1}) = - \sum_i (\nabla_{K, e_i} \theta)(e_i, e_1, \dots, e_{p-1}),$$

where ∇_K is the connection on $\Omega^*(E)$, induced by the Levi-Civita connection ∇ and D_K .

If H be another metric on E , we define the positive endomorphism $h = K^{-1}H$ by setting $H(\cdot, \cdot) = K(h(\cdot), \cdot)$. A straightforward calculation, using (2.11), shows that

$$(2.14) \quad D_H = D_K + \frac{1}{2}h^{-1}\delta_K h = \frac{1}{2}(D_K + h^{-1} \circ D_K \circ h + \psi_K - h^{-1} \circ \psi_K \circ h),$$

$$(2.15) \quad \psi_H = \psi_K - \frac{1}{2}h^{-1}\delta_K h = \frac{1}{2}(\psi_K + h^{-1} \circ \psi_K \circ h + D_K - h^{-1} \circ D_K \circ h),$$

$$(2.16) \quad \delta_H = \delta_K + h^{-1}\delta_K h = h^{-1} \circ \delta_K \circ h,$$

where $\delta_K = D_K - \psi_K$.

Lemma 2.1. *If $H(t)$ is a one-parameter family of fiber metrics on the vector bundle (E, D) , then we have*

$$(2.17) \quad \frac{\partial D_{H(t)}}{\partial t} = \frac{1}{2}D_{H(t)}(h^{-1}(t)\frac{\partial h(t)}{\partial t}) - \frac{1}{2}[\psi_H(t), h^{-1}(t)\frac{\partial h(t)}{\partial t}],$$

$$(2.18) \quad \frac{\partial \psi_{H(t)}}{\partial t} = -\frac{1}{2}D_{H(t)}(h^{-1}(t)\frac{\partial h(t)}{\partial t}) + \frac{1}{2}[\psi_H(t), h^{-1}(t)\frac{\partial h(t)}{\partial t}],$$

$$(2.19) \quad \begin{aligned} \frac{\partial D_{H(t)}^* \psi_{H(t)}}{\partial t} &= -\frac{1}{2}D_{H(t)}^* D_{H(t)}(h^{-1}(t)\frac{\partial h(t)}{\partial t}) + \frac{1}{2}[D_{H(t)}^* \psi_{H(t)}, h^{-1}(t)\frac{\partial h(t)}{\partial t}] \\ &\quad - \frac{1}{2}g^{ij}[\psi_{H(t)}(\frac{\partial}{\partial x^i}), [\psi_{H(t)}(\frac{\partial}{\partial x^j}), h^{-1}(t)\frac{\partial h(t)}{\partial t}]], \end{aligned}$$

where $h(t) = K^{-1}H(t)$.

Proof. It suffice to prove the (2.17) and (2.19). By making use of (2.14), (2.15), we deduce

$$\begin{aligned}
(2.20) \quad \frac{\partial D_H}{\partial t} &= \frac{1}{2} \frac{\partial}{\partial t} (h^{-1} D_K h - h^{-1} \circ \psi_K \circ h) \\
&= -\frac{1}{2} h^{-1} \frac{\partial h}{\partial t} h^{-1} D_K h + \frac{1}{2} h^{-1} D_K \frac{\partial h}{\partial t} + \frac{1}{2} h^{-1} \frac{\partial h}{\partial t} h^{-1} \psi_K h - \frac{1}{2} h^{-1} \psi_K \frac{\partial h}{\partial t} \\
&= -\frac{1}{2} h^{-1} \frac{\partial h}{\partial t} h^{-1} D_K h + \frac{1}{2} h^{-1} D_K \frac{\partial h}{\partial t} - \frac{1}{2} [\psi_H, h^{-1} \frac{\partial h}{\partial t}] \\
&\quad + \frac{1}{2} h^{-1} \frac{\partial h}{\partial t} (\psi_H - \psi_K - D_K + h^{-1} \circ D_K \circ h) \\
&\quad - \frac{1}{2} (\psi_H - \psi_K - D_K + h^{-1} \circ D_K \circ h) h^{-1} \frac{\partial h}{\partial t} \\
&= \frac{1}{2} D_H (h^{-1} \frac{\partial h}{\partial t}) - \frac{1}{2} [\psi_H, h^{-1} \frac{\partial h}{\partial t}] - \frac{1}{2} h^{-1} \frac{\partial h}{\partial t} h^{-1} D_K h \\
&\quad + \frac{1}{2} h^{-1} D_K \frac{\partial h}{\partial t} + \frac{1}{2} [h^{-1} \frac{\partial h}{\partial t}, h^{-1} \circ D_K \circ h] \\
&= \frac{1}{2} D_H (h^{-1} \frac{\partial h}{\partial t}) - \frac{1}{2} [\psi_H, h^{-1} \frac{\partial h}{\partial t}].
\end{aligned}$$

On the other hand, let's choose the local normal coordinate (x^1, \dots, x^n) at the considered point, using (2.17) and (2.18), we have

$$\begin{aligned}
(2.21) \quad \frac{\partial D_H^* \psi_H}{\partial t} &= -\frac{\partial}{\partial t} [D_{H, \frac{\partial}{\partial x^i}}, \psi_H(\frac{\partial}{\partial x^i})] \\
&= -[\frac{\partial}{\partial t} D_{H, \frac{\partial}{\partial x^i}}, \psi_H(\frac{\partial}{\partial x^i})] - [D_{H, \frac{\partial}{\partial x^i}}, \frac{\partial}{\partial t} \psi_H(\frac{\partial}{\partial x^i})] \\
&= -\frac{1}{2} [D_{H, \frac{\partial}{\partial x^i}} (h^{-1} \frac{\partial h}{\partial t}) - [\psi_H(\frac{\partial}{\partial x^i}), h^{-1} \frac{\partial h}{\partial t}], \psi_H(\frac{\partial}{\partial x^i})] \\
&\quad + \frac{1}{2} [D_{H, \frac{\partial}{\partial x^i}}, D_{H, \frac{\partial}{\partial x^i}} (h^{-1} \frac{\partial h}{\partial t}) - [\psi_H(\frac{\partial}{\partial x^i}), h^{-1} \frac{\partial h}{\partial t}]] \\
&= \frac{1}{2} D_{H, \frac{\partial}{\partial x^i}} D_{H, \frac{\partial}{\partial x^i}} (h^{-1} \frac{\partial h}{\partial t}) - \frac{1}{2} [\psi_H(\frac{\partial}{\partial x^i}), [\psi_H(\frac{\partial}{\partial x^i}), h^{-1} \frac{\partial h}{\partial t}]] \\
&\quad - \frac{1}{2} [D_{H, \frac{\partial}{\partial x^i}} (h^{-1} \frac{\partial h}{\partial t}), \psi_H(\frac{\partial}{\partial x^i})] - \frac{1}{2} [D_{H, \frac{\partial}{\partial x^i}}, [\psi_H(\frac{\partial}{\partial x^i}), h^{-1} \frac{\partial h}{\partial t}]] \\
&= -\frac{1}{2} D_H^* D_H (h^{-1} \frac{\partial h}{\partial t}) + \frac{1}{2} [D_H^* \psi_H, h^{-1} \frac{\partial h}{\partial t}] \\
&\quad - \frac{1}{2} [\psi_H(\frac{\partial}{\partial x^i}), [\psi_H(\frac{\partial}{\partial x^i}), h^{-1} \frac{\partial h}{\partial t}]].
\end{aligned}$$

□

Under the local coordinate (x^1, \dots, x^n) , by using (2.14) and (2.15), it follows

$$\begin{aligned}
(2.22) \quad D_H^* \psi_H &= D_K^* \psi_K - \frac{1}{2} D_K^* (h^{-1} \delta_K h) - \frac{1}{2} g^{ij} [h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, \psi_K(\frac{\partial}{\partial x^j})] \\
&= D_K^* \psi_K + \frac{1}{2} g^{ij} D_{\frac{\partial}{\partial x^i}} (h^{-1} \delta_{K, \frac{\partial}{\partial x^j}} h) - \frac{1}{2} g^{ij} \Gamma_{ij}^k h^{-1} \delta_{K, \frac{\partial}{\partial x^k}} h.
\end{aligned}$$

From now on, the norms below are taken with respect to K unless indicated explicitly. Based on (2.22), we prove the following lemma.

Lemma 2.2. *Let H and K be two fiber metrics on the vector bundle (E, D) , it holds*

$$(2.23) \quad (D_H^* \psi_H - D_K^* \psi_K, h) = \frac{1}{2} \Delta \operatorname{tr} h - \frac{1}{2} |h^{-\frac{1}{2}} \delta_K h|^2,$$

$$(2.24) \quad (D_K^* \psi_K - D_H^* \psi_H, h^{-1})_H = \frac{1}{2} \Delta \operatorname{tr} h^{-1} - \frac{1}{2} |h^{\frac{1}{2}} \delta_H h^{-1}|_H^2,$$

where $h = K^{-1}H$.

Proof. We only proof (2.23) and (2.24) follows from (2.23). Choose the local normal coordinate (x^1, \dots, x^n) at the considered point, it follows

$$(2.25) \quad \begin{aligned} (D_H^* \psi_H - D_K^* \psi_K, h) &= -\frac{1}{2} (D_K^* (h^{-1} \delta_K h) + [h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, \psi_K(\frac{\partial}{\partial x^i})], h) \\ &= \frac{1}{2} (D_{K, \frac{\partial}{\partial x^i}} (h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h), h) - \frac{1}{2} ([h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, \psi_K(\frac{\partial}{\partial x^i})], h) \\ &= \frac{1}{2} \frac{\partial}{\partial x^i} (h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, h) - \frac{1}{2} (h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, D_{K, \frac{\partial}{\partial x^i}} h) \\ &\quad + \frac{1}{2} (h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, [\psi_K(\frac{\partial}{\partial x^i}), h]) \\ &= \frac{1}{2} \frac{\partial}{\partial x^i} \operatorname{tr} \delta_{K, \frac{\partial}{\partial x^i}} h - \frac{1}{2} (h^{-1} \delta_{K, \frac{\partial}{\partial x^i}} h, \delta_{K, \frac{\partial}{\partial x^i}} h) \\ &= \frac{1}{2} \Delta \operatorname{tr} h - \frac{1}{2} |h^{-\frac{1}{2}} \delta_K h|^2. \end{aligned}$$

□

Following [13], the Donaldson's distance on the space of metrics is defined by

$$(2.26) \quad \sigma(K, H) = \operatorname{tr}(K^{-1}H) + \operatorname{tr}(H^{-1}K) - 2 \operatorname{rank}(E).$$

It is obvious that $\sigma(H, K) \geq 0$ with equality if and only if $H = K$.

By Lemma 2.2, we get

Corollary 2.1. *Let H and K be two harmonic metrics on the vector bundle (E, D) , then we have*

$$(2.27) \quad \Delta \sigma(H, K) = |h^{-\frac{1}{2}} \delta_K h|^2 + |h^{\frac{1}{2}} \delta_H h^{-1}|_H^2,$$

where $h = K^{-1}H$. In particular, if (E, D) is simple and (M, g) satisfies the Assumption 2.1, the mutually bounded harmonic metrics are unique up to scaling.

Now we consider the evolution equation

$$(2.28) \quad H^{-1}(t) \frac{\partial H(t)}{\partial t} = 2D_{H(t)}^* \psi_{H(t)},$$

a direct calculation yields

Lemma 2.3. *Let (E, D) be a vector bundle over (M, g) , assume $H(t)$ and $K(t)$ be two solutions of the heat flow (2.28), then we have*

$$(2.29) \quad \left(\frac{\partial}{\partial t} - \Delta \right) \sigma(H(t), K(t)) = -|\tilde{h}(t)^{-\frac{1}{2}} \delta_{K(t)} \tilde{h}(t)|_{K(t)}^2 - |\tilde{h}^{\frac{1}{2}}(t) \delta_{H(t)} \tilde{h}^{-1}(t)|_{H(t)}^2,$$

where $\tilde{h}(t) = K^{-1}(t)H(t)$.

Lemma 2.4. *Let (E, D) be a vector bundle over (M, g) , along the heat flow (2.28) it holds*

$$(2.30) \quad \frac{d}{dt} \|\psi_H\|_{H, L^2}^2 = -2 \|D_H^* \psi_H\|_{H, 2}^2 - 2 \int_M d\eta_H,$$

$$(2.31) \quad \frac{d}{dt} \|D_H^* \psi_H\|_{H, L^2}^2 = -2 \|D_H D_H^* \psi_H\|_{H, L^2}^2 - 2 \|[\psi_H, D_H^* \psi_H]\|_{H, L^2}^2 + 2 \int_M d\tilde{\eta}_H,$$

$$(2.32) \quad \left(\frac{\partial}{\partial t} - \Delta\right) |D_H^* \psi_H|_H^2 = -2 |D_H D_H^* \psi_H|_H^2 - 2 |[\psi_H, D_H^* \psi_H]|_H^2,$$

where $\eta_H = H(D_H^* \psi_H, * \psi_H)$, $\tilde{\eta}_H = H(D_H^* \psi_H, * D_H D_H^* \psi_H)$. Furthermore, if D is a flat connecton, we have

$$(2.33) \quad \left(\frac{\partial}{\partial t} - \Delta\right) |\psi_H|_H^2 = -|[\psi_H, \psi_H]|_H^2 - 2(\psi_H \circ \text{Ric}, \psi_H)_H - 2 |\nabla_H \psi_H|_H^2,$$

where Ric is the Ricci transformation of (M, g) .

Proof. Firstly, applying the equality (2.18) yields

$$(2.34) \quad \begin{aligned} \frac{d}{dt} \|\psi_H\|_{H, L^2}^2 &= 2 \int_M \left(-\frac{1}{2} D_H (H^{-1} \frac{\partial H}{\partial t}) + \frac{1}{2} [\psi_H, H^{-1} \frac{\partial H}{\partial t}], \psi_H\right)_H \text{dvol}_g \\ &= -2 \int_M (D_H D_H^* \psi_H, \psi_H)_H \text{dvol}_g \\ &= -2 \|D_H^* \psi_H\|_{H, 2}^2 - 2 \int_M d\eta_H. \end{aligned}$$

Next, (2.19) indicates that

$$(2.35) \quad \frac{\partial D_H^* \psi_H}{\partial t} = -D_H^* D_H D_H^* \psi_H - g^{ij} [\psi_H(\frac{\partial}{\partial x^i}), [\psi_H(\frac{\partial}{\partial x^j}), D_H^* \psi_H]],$$

as it is easy to check $D_H^* \psi_H$ is self-adjoint with respect to H , it then follows

$$(2.36) \quad \begin{aligned} \frac{d}{dt} \|D_H^* \psi_H\|_{H, L^2}^2 &= -2 \int_M (D_H^* D_H D_H^* \psi_H, D_H^* \psi_H)_H \text{dvol}_g \\ &\quad - 2 \int_M g^{ij} ([\psi_H(\frac{\partial}{\partial x^i}), [\psi_H(\frac{\partial}{\partial x^j}), D_H^* \psi_H]], D_H^* \psi_H)_H \text{dvol}_g \\ &= -2 \|D_H D_H^* \psi_H\|_{H, L^2}^2 - 2 \|[\psi_H, D_H^* \psi_H]\|_{H, L^2}^2 + 2 \int_M d\tilde{\eta}_H, \end{aligned}$$

and

$$(2.37) \quad \begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) |D_H^* \psi_H|_H^2 &= -2 (D_H^* D_H D_H^* \psi_H, D_H^* \psi_H)_H \\ &\quad - 2 g^{ij} ([\psi_H(\frac{\partial}{\partial x^i}), [\psi_H(\frac{\partial}{\partial x^j}), D_H^* \psi_H]], D_H^* \psi_H)_H \\ &\quad + 2 (D_H^* D_H D_H^* \psi_H, D_H^* \psi_H)_H - 2 |D_H D_H^* \psi_H|_H^2 \\ &= -2 |D_H D_H^* \psi_H|_H^2 - 2 |[\psi_H, D_H^* \psi_H]|_H^2. \end{aligned}$$

On the other hand, we have

$$(2.38) \quad \begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) |\psi_H|_H^2 &= 2 \left(-\frac{1}{2} D_H (H^{-1} \frac{\partial H}{\partial t}) + \frac{1}{2} [\psi_H, H^{-1} \frac{\partial H}{\partial t}], \psi_H\right)_H \\ &\quad + 2 (\nabla_H^* \nabla_H \psi_H, \psi_H)_H - 2 |\nabla_H \psi_H|_H^2 \\ &= -2 (D_H D_H^* \psi_H - \nabla_H^* \nabla_H \psi_H, \psi_H)_H - 2 |\nabla_H \psi_H|_H^2, \end{aligned}$$

Now using the flatness of D : $D_H\psi_H = 0$ and $D_H^2 = -\frac{1}{2}[\psi_H, \psi_H]$, it is equal to

$$\begin{aligned}
(2.39) \quad & 2g^{ij}(D_H^2(\cdot, \frac{\partial}{\partial x^i})(\psi_H(\frac{\partial}{\partial x^j})) - \psi_H(R_\nabla(\cdot, \frac{\partial}{\partial x^i})(\frac{\partial}{\partial x^j})), \psi_H)_H - 2|\nabla_H\psi_H|_H^2 \\
& = -(g^{ij}[[\psi_H, \psi_H(\frac{\partial}{\partial x^i})], \psi_H(\frac{\partial}{\partial x^j})] + 2\psi_H \circ \text{Ric}, \psi_H)_H - 2|\nabla_H\psi_H|_H^2 \\
& = -|\psi_H, \psi_H|_H^2 - 2(\psi_H \circ \text{Ric}, \psi_H)_H - 2|\nabla_H\psi_H|_H^2.
\end{aligned}$$

□

3. ANALYTICALLY STABILITY AND EXISTENCE OF POISSON METRICS

Given a vector bundle (E, D) over a compact Riemannian manifold (X, g) , consider the following heat flow

$$(3.1) \quad \begin{cases} H^{-1}\frac{\partial H}{\partial t} = 2D_H^*\psi_H, \\ H(0) = K. \end{cases}$$

And if X has nonempty smooth boundary ∂X , for any given compatible data \tilde{H} , we impose the boundary condition $H(t)|_{\partial X} = \tilde{H}$ in the above system.

With the calculations in the last sections in hand and following [13], we have

Proposition 3.1. *Let (E, D) be a vector bundle over a compact Riemannian manifold (X, g) (with possibly nonempty boundary), then heat flow (3.1) admits unique solution defined on $[0, \infty)$.*

Proposition 3.2. *Assume (E, D) be a vector bundle over a compact Riemannian manifold (M, g) , then for any $\epsilon > 0$, there exists a fiber metric H_ϵ such that $\|D_{H_\epsilon}\psi_{H_\epsilon}\|_{L^\infty} \leq \epsilon$.*

Proof. Let $H = H(t)$ be the solution to (3.1), defined on $[0, +\infty)$. Since (3.1) is the negative gradient flow of the energy $\|\psi_H\|_{H, L^2}$, by (2.31) we know $\|D_H^*\psi_H\|_{H, L^2}^2 \rightarrow 0$, as t goes infinity. Then the proof can be finished by using (2.32) and parabolic Moser's iteration. □

Proposition 3.3. *Let (E, D) be a vector bundle over a compact Riemannian manifold (X, g) with smooth boundary ∂X . Then for any given data K , there exists a unique harmonic metric H on (E, D) such that $H = K$ on ∂X .*

Proof. Due to Corollary 2.1, it remain to prove the existence statement. By the standard elliptic equation theory, let u be the solution to the following Dirichlet problem:

$$(3.2) \quad \Delta u = -|D_K^*\psi_K|, u|_{\partial X} = 0.$$

Proposition 3.1 ensures the existence of long-time solution $H(t)$ of (3.1) with boundary condition $H(t)|_{\partial X} = K$. Set

$$(3.3) \quad v(x, t) = \int_0^t |D_{H(s)}^*\psi_{H(s)}|_{H(s)} ds - u,$$

by (2.32) it is easy to see

$$(3.4) \quad \begin{cases} (\frac{\partial}{\partial t} - \Delta)v(x, t) \leq 0, \\ v(x, 0) = -u(x), \\ v(x, t)|_{\partial X} = 0. \end{cases}$$

Then the maximum principle gives

$$(3.5) \quad \int_0^t |D_{H(s)}^*\psi_{H(s)}|_{H(s)} ds \leq \max_X u.$$

Now for any $t_1 \leq t \leq t_2$ and $\tilde{h}(t) = H^{-1}(t_1)H(t)$, $\hat{h}(t) = H^{-1}(t)H(t_1)$, it is straightforward to check

$$(3.6) \quad \left| \frac{\partial}{\partial t} \log \operatorname{tr} \tilde{h}(t) \right| \leq 2 |D_{H(t)}^* \psi_{H(t)}|_{H(t)}, \left| \frac{\partial}{\partial t} \log \operatorname{tr} \hat{h}(t) \right| \leq 2 |D_{H(t)}^* \psi_{H(t)}|_{H(t)}.$$

A simple integration yields

$$(3.7) \quad \sigma(H(t_1), H(t_2)) \leq 2 \operatorname{rank}(E) (e^{2 \int_{t_1}^{t_2} |D_{H(s)}^* \psi_{H(s)}|_{H(s)} ds} - 1).$$

So by combining (3.5) and (3.7), we know that $H(t)$ converge in C^0 -topology to H_∞ as $t \rightarrow \infty$. Using (3.5) and arguing as Lemma 3.3 in [47], the elliptic regularity theory implies that there exists a subsequence $H(t_i)$ converging to H_∞ in C^∞ -topology and H_∞ is a harmonic metric satisfying the desired boundary condition. \square

Proposition 3.4. *Let (E, D) be a vector bundle over a compact Riemannian manifold (X, g) with smooth boundary ∂X . Then for any given data K , there exists a unique Poisson metric H on (E, D) such that $H = K$ on ∂X and $\det H = \det K$.*

Proof. Due to Proposition 3.3, there exists a unique harmonic metric \tilde{H} such that $\tilde{H} = K$ on ∂X . Set $H = \tilde{H}e^f$, it follows

$$(3.8) \quad D_H^* \psi_H = -\frac{1}{2} D_{\tilde{H}}^* (df \otimes \operatorname{id}_E) = \frac{\operatorname{tr} D_{\tilde{H}}^* \psi_{\tilde{H}}}{\operatorname{rank}(E)} \operatorname{id}_E.$$

Now we set $f = \frac{\log \det(\tilde{H}^{-1}K)}{\operatorname{rank}(E)}$, then we have $H = K$ on ∂X and $\det H = \det K$. \square

On each M_s , the following Dirichlet problem is solvable,

$$(3.9) \quad \begin{cases} D_{H_s}^* \psi_{H_s} = c_s \operatorname{id}_E, \\ H|_{\partial M_s} = K, \\ \det h_s = 1, h_s = K^{-1} H_s. \end{cases}$$

Due to the elementary inequality $\log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}) \geq \log(\det h_s)^{\frac{1}{r}} = 0$ and the boundary condition, we have

$$(3.10) \quad \frac{\partial \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)})}{\partial \vec{n}} \geq 0,$$

where \vec{n} is the inward unit normal vector field at ∂M_s . We extend $\log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)})$ to the function $\widetilde{\log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)})}$ on whole M by setting 0 outside M_s . For any non-negative compactly supported function φ , the Green's formula yields

$$(3.11) \quad \begin{aligned} \int_M \widetilde{\log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)})} \Delta \varphi \, d\operatorname{vol}_g &= \int_{M_s} \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}) \Delta \varphi \, d\operatorname{vol}_g \\ &= \int_{M_s} \varphi \Delta \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}) \, d\operatorname{vol}_g \\ &\quad + \int_{\partial M_s} \varphi \frac{\partial \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)})}{\partial \vec{n}} \, d\operatorname{vol}_{\partial M_s} \\ &\quad - \int_{\partial M_s} \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}) \frac{\partial \varphi}{\partial \vec{n}} \, d\operatorname{vol}_{\partial M_s} \\ &\geq \int_{M_s} \varphi \Delta \log(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}) \, d\operatorname{vol}_g. \end{aligned}$$

Now by Lemma 2.2, we see

$$\begin{aligned}
(3.12) \quad \Delta \log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right) &= \frac{2(D_{H_s}^* \psi_{H_s} - D_K^* \psi_K, h_s) + |h_s^{-\frac{1}{2}} \delta_K h_s|^2}{\operatorname{tr} h_s} - \frac{|\operatorname{tr} \delta_K h_s|^2}{(\operatorname{tr} h_s)^2} \\
&\geq \frac{2(D_{H_s}^* \psi_{H_s} - D_K^* \psi_K, h_s)}{\operatorname{tr} h_s} \\
&= \frac{-2((D_K^* \psi_K)^\perp, h_s)}{\operatorname{tr} h_s} + \frac{2((\operatorname{tr} D_{H_s}^* \psi_{H_s} - \operatorname{tr} D_K^* \psi_K) \operatorname{id}_E, h_s)}{\operatorname{rank}(E) \operatorname{tr} h_s} \\
&\geq -2|(D_K^* \psi_K)^\perp| + \frac{\Delta \log \det h_s}{\operatorname{rank}(E)} \\
&\geq -2C\phi,
\end{aligned}$$

where we have used the following simple inequality

$$(3.13) \quad \frac{|\operatorname{tr} \delta_K h_s|^2}{\operatorname{tr} h_s} \leq |h_s^{-\frac{1}{2}} \delta_K h_s|^2.$$

And therefore

$$(3.14) \quad \int_M \widetilde{\log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right)} \Delta \varphi \, d\operatorname{vol}_g \geq -2C \int_{M_s} \varphi \phi \, d\operatorname{vol}_g \geq -2C \int_M \varphi \phi \, d\operatorname{vol}_g.$$

That is, $\widetilde{\Delta \log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right)} \geq -2C\phi$ as a distribution. So by Assumption 2.1 we get

$$(3.15) \quad \max_{M_s} \log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right) \leq C_1(1 + \int_{M_s} \log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right) \phi \, d\operatorname{vol}_g),$$

where C_1 depends on C . Combing this with $\det h_s = 1$ yields

$$(3.16) \quad \max_{M_s} |\log h_s| \leq C_2(1 + \int_{M_s} |\log h_s| \phi \, d\operatorname{vol}_g),$$

for a constant C_2 depending only on C_1 and $\operatorname{rank}(E)$, where we have used that

$$(3.17) \quad \log\left(\frac{\operatorname{tr} h_s}{\operatorname{rank}(E)}\right) \leq |\log h_s| \leq \operatorname{rank}(E)^{\frac{3}{2}} \log \operatorname{tr} h_s.$$

Our goal is to show the quantity

$$(3.18) \quad \int_{M_s} |\log h_s| \phi \, d\operatorname{vol}_g$$

is uniformly bounded, under the assumption that D is K -analytically stable.

For further consideration, we recall some notations. Given $\rho \in C^\infty(\mathbb{R}, \mathbb{R})$, $\Theta \in C^\infty(\mathbb{R} \times \mathbb{R}, \mathbb{R})$, $\chi \in \Omega^\bullet(\operatorname{End}(E))$ and a self-adjoint bundle endomorphism σ , we define $\rho[\sigma]$ and $\Theta[\sigma](\chi)$ as follows. For every $x \in M$, we set

$$(3.19) \quad \rho[\sigma] = \rho(\lambda_\alpha) e^\alpha \otimes e_\alpha, \Theta[\sigma](\chi) = \Theta(\lambda_\alpha, \lambda_\beta) \chi_\alpha^\beta e^\alpha \otimes e_\beta,$$

where $\{e_\alpha\}_{\alpha=1}^{\operatorname{rank}(E)}$ is an orthogonal basis with respect to K , such that

$$(3.20) \quad \sigma(e_\alpha) = \lambda_\alpha e_\alpha, \chi = \chi_\alpha^\beta e^\alpha \otimes e_\beta.$$

Next we prove the following proposition.

Proposition 3.5. *Let (E, D) be a vector bundle over a compact Riemannian manifold (X, g) with empty boundary, and H, K be two fiber metrics with $s = \log(K^{-1}H)$ and $s|_{\partial X} = 0$. Then we have*

$$(3.21) \quad \int_X (D_H^* \psi_H - D_K^* \psi_K, s) \, \text{dvol}_g = -\frac{1}{2} \int_X (\Theta[s](Ds), Ds) \, \text{dvol}_g,$$

in the sense of (3.19), where

$$(3.22) \quad \Theta(x, y) = \begin{cases} \frac{e^{y-x}-1}{y-x}, & x \neq y, \\ 1, & x = y. \end{cases}$$

Proof. Consider the local normal coordinate at the considered point and for $h = K^{-1}H$, noting that $(h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, s) = \text{tr}(s\delta_{K, \frac{\partial}{\partial x^i}} s)$, through computing, we have

$$(3.23) \quad \begin{aligned} (D_H^* \psi_H - D_K^* \psi_K, s) &= -\frac{1}{2} (D_K^* (h^{-1}\delta_K h), s) - \frac{1}{2} ([h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, \psi_K(\frac{\partial}{\partial x^i})], s) \\ &= \frac{1}{2} (D_{K, \frac{\partial}{\partial x^i}} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h), s) + \frac{1}{2} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, [\psi_K(\frac{\partial}{\partial x^i}), s]) \\ &= \frac{1}{2} \frac{\partial}{\partial x^i} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, s) - \frac{1}{2} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, D_{K, \frac{\partial}{\partial x^i}} s) \\ &\quad + \frac{1}{2} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, [\psi_K(\frac{\partial}{\partial x^i}), s]) \\ &= \frac{1}{2} \frac{\partial}{\partial x^i} \text{tr}(s\delta_{K, \frac{\partial}{\partial x^i}} s) - \frac{1}{2} (h^{-1}\delta_{K, \frac{\partial}{\partial x^i}} h, \delta_{K, \frac{\partial}{\partial x^i}} s) \\ &= \frac{1}{2} \frac{\partial}{\partial x^i} \text{tr}(s\frac{\partial}{\partial x^i} s) - \frac{1}{2} (h^{-1}\delta_K h, \delta_K s). \end{aligned}$$

On the other hand, according to Section 7.4 in [28], there is an open dense subset $W \subset X$ and for every $x \in W$, we may choose an orthogonal basis $\{e_\alpha\}_{\alpha=1}^{\text{rank}(E)}$ for E with respect to K , defined over a neighborhood of x , such that

$$(3.24) \quad h = \sum_{\alpha=1}^{\text{rank}(E)} e^{\lambda_\alpha} e_\alpha \otimes e^\alpha, \quad s = \sum_{\alpha=1}^{\text{rank}(E)} \lambda_\alpha e_\alpha \otimes e^\alpha.$$

We set $De_\alpha = A_\alpha^\beta e_\beta$, then it follows

$$(3.25) \quad Dh \circ h^{-1} = \Theta[s](Ds) = \sum_{\alpha=1}^{\text{rank}(E)} d\lambda_\alpha e_\alpha \otimes e^\alpha + \sum_{\alpha \neq \beta} (1 - e^{\lambda_\alpha - \lambda_\beta}) A_\beta^\alpha e_\alpha \otimes e^\beta,$$

and therefore

$$(3.26) \quad (h^{-1}\delta_K h, \delta_K s) = (Ds, Dh \circ h^{-1}) = (\Theta[s](Ds), Ds).$$

Finally, by (3.23) and (3.26), we have

$$(3.27) \quad \begin{aligned} \int_X (D_H^* \psi_H - D_K^* \psi_K, s) \, \text{dvol}_g &= \frac{1}{4} \int_X d(*d|s|^2) - \frac{1}{2} \int_X (\Theta[s](Ds), Ds) \, \text{dvol}_g \\ &= -\frac{1}{2} \int_X (\Theta[s](Ds), Ds) \, \text{dvol}_g. \end{aligned}$$

□

We shall prove (3.18) by contradictory argument and the method is a combination of Proposition 3.5 and Simpson's trick. Otherwise, we may assume that there exists a sequence $s \rightarrow \infty$, such that

$$(3.28) \quad l_s = \int_{M_s} |\log h_s| \phi \, d\text{vol}_g \rightarrow +\infty.$$

Let's set $u_s = \log h_s / l_s$ with $\text{tr } u_s = 0$ and $\int_{M_s} |u_s| \phi \, d\text{vol}_g = 1$. By (3.16) we see

$$(3.29) \quad \sup_{M_s} |u_s| \leq \frac{C_2}{l_s} (1 + l_s) \leq C_3,$$

for a constant C_3 doesn't depend on s . Using Proposition 3.5 and the fact that H_s is a Poisson metric, one has

$$(3.30) \quad \begin{aligned} \int_{M_s} l_s (\Theta[l_s u_s](Du_s), Du_s) \, d\text{vol}_g &= 2 \int_{M_s} (D_K^* \psi_K - D_{H_s}^* \psi_{H_s}, u_s) \, d\text{vol}_g \\ &= 2 \int_{M_s} ((D_K^* \psi_K)^\perp, u_s) \, d\text{vol}_g. \end{aligned}$$

Notice that

$$(3.31) \quad l\Theta(lx, ly) \rightarrow \begin{cases} \frac{1}{x-y}, & x > y, \\ +\infty, & x \leq y, \end{cases}$$

increases monotonically as $l \rightarrow +\infty$. It holds that

$$(3.32) \quad \int_{M_s} (\rho[u_s](Du_s), Du_s) \, d\text{vol}_g \leq 2 \int_{M_s} ((D_K^* \psi_K)^\perp, u_s) \, d\text{vol}_g.$$

for any $\rho : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ with $\rho(x, y) < \frac{1}{x-y}$ whenever $x > y$ and s large enough.

Due to the zeroth order estimate of u_s , and $|(D_K^* \psi_K)^\perp| \leq C\phi$, by taking ρ small enough, it follows that Du_s is bounded in L^2 -norm on any compact subset of M . We may assume that $u_s \rightarrow u_\infty$ weakly in $L^2_{1,\text{loc}}$ -topology with $\|u_\infty\|_{L^\infty} \leq C_3$. Since $\phi \in L^1$, for any $\epsilon > 0$ there exists a s_ϵ such that for any $s \geq s_\epsilon$, it holds

$$(3.33) \quad \int_{M-M_s} \phi \, d\text{vol}_g \leq \epsilon.$$

Hence for $s_\epsilon \leq s_1 \leq s_2$, (3.29) indicates

$$(3.34) \quad 1 - C_3\epsilon \leq \int_{M_{s_2}} |u_{s_2}| \phi \, d\text{vol}_g - \int_{M_{s_2}-M_{s_1}} |u_{s_2}| \phi \, d\text{vol}_g = \int_{M_{s_1}} |u_{s_2}| \phi \, d\text{vol}_g \leq 1.$$

Fixing s_1 and by the weak compactness of $\{u_s\}$ in $L^2_{1,\text{loc}}$ -topology, we get

$$(3.35) \quad 1 - C_3\epsilon \leq \int_{M_{s_1}} |u_\infty| \phi \, d\text{vol}_g \leq 1,$$

and then taking $s_1 \rightarrow \infty$ and $\epsilon \rightarrow 0$ yields $\int_M |u_\infty| \phi \, d\text{vol}_g = 1$, which implies $u_\infty \neq 0$.

Now it follows from (3.29), (3.32), (3.33) and $|(D_K^* \psi_K)^\perp| \leq C\phi$ that

$$\begin{aligned}
(3.36) \quad \int_{M_{s_1}} (\rho[u_{s_2}](Du_{s_2}), Du_{s_2}) \, \text{dvol}_g &\leq \int_{M_{s_2}} (\rho[u_{s_2}](Du_{s_2}), Du_{s_2}) \, \text{dvol}_g \\
&\leq 2 \left(\int_{M_{s_2}-M_{s_1}} + \int_{M_{s_1}} \right) ((D_K^* \psi_K)^\perp, u_{s_2}) \, \text{dvol}_g \\
&\leq 2CC_3\epsilon + 2 \int_{M_{s_1}} ((D_K^* \psi_K)^\perp, u_{s_2}) \, \text{dvol}_g.
\end{aligned}$$

Let's take $s_2 \rightarrow \infty, s_1 \rightarrow \infty$ and $\epsilon \rightarrow 0$ successively, it yields

$$(3.37) \quad \int_M (\rho[u_\infty](Du_\infty), Du_\infty) \, \text{dvol}_g \leq 2 \int_M ((D_K^* \psi_K)^\perp, u_\infty) \, \text{dvol}_g \leq 2CC_3 \int_M \phi \, \text{dvol}_g.$$

As long as (3.37) is established, by using Simpson's discussion in [39](p886-888) and Loftin's observation in [27], it's easy to conclude (3.18) and therefore we obtain

Proposition 3.6. *If (M, g) satisfies the Assumption 2.1 and the flat bundle (E, D) is K -analytically stable with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C , it holds*

$$(3.38) \quad \max_{M_s} |\log h_s| \leq \tilde{C}_1,$$

for a constant \tilde{C}_1 that doesn't depend on s .

Using Lemma 2.2 again shows

$$\begin{aligned}
(3.39) \quad |h_s^{-\frac{1}{2}} \delta_K h_s|_{L^2(M_s)}^2 &= \int_{M_s} (\Delta \operatorname{tr} h_s - 2(D_{H_s}^* \psi_{H_s} - D_K^* \psi_K, h_s)) \, \text{dvol}_g \\
&= \int_{M_s} \left(\Delta \operatorname{tr} h_s + 2((D_K^* \psi_K)^\perp, h_s) \right) \, \text{dvol}_g \\
&= - \int_{\partial M_s} \frac{\partial \operatorname{tr} h_s}{\partial \vec{n}} \, \text{dvol}_{\partial M_s} + 2 \int_{M_s} ((D_K^* \psi_K)^\perp, h_s) \, \text{dvol}_g \\
&\leq 2 \int_{M_s} ((D_K^* \psi_K)^\perp, h_s) \, \text{dvol}_g \\
&\leq CC_4 \int_M \phi \, \text{dvol}_g,
\end{aligned}$$

where C_4 depends only on \tilde{C}_1 and $\operatorname{rank}(E)$. So we obtain

Proposition 3.7. *If (M, g) satisfies the Assumption 2.1 and the flat bundle (E, D) is K -analytically stable with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C , we have*

$$(3.40) \quad \|Dh_s\|_{L^2(M_s)} \leq \tilde{C}_2 \|\phi\|_{L^1}^{\frac{1}{2}},$$

for a constant \tilde{C}_2 that depends only on C, \tilde{C}_1 and $\operatorname{rank}(E)$.

Proposition 3.8. *If (M, g) satisfies the Assumption 2.1 and the flat bundle (E, D) is K -analytically stable with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C , then there exists a Poisson metric H such that $\det h = 1$, $|h| \in L^\infty$ and $|Dh| \in L^2$ for $h = K^{-1}H$.*

Proof. Using the flatness of D , we deduce

$$\begin{aligned}
\Delta|\psi_{H_s}|_{H_s}^2 &\geq -2g^{ij}(D_{H_s}^2(\cdot, \frac{\partial}{\partial x^i})(\psi_{H_s}(\frac{\partial}{\partial x^j})), \psi_{H_s})_{H_s} \\
&\quad + 2g^{ij}(\psi_{H_s}(R_{\nabla}(\cdot, \frac{\partial}{\partial x^i})(\frac{\partial}{\partial x^j})), \psi_{H_s})_{H_s} - 2(D_{H_s}D_{H_s}^*\psi_{H_s}, \psi_{H_s})_{H_s} \\
&= |[\psi_{H_s}, \psi_{H_s}]|_{H_s}^2 + 2(\psi_{H_s} \circ \text{Ric}, \psi_{H_s}) - 2(D_{H_s}D_{H_s}^*\psi_{H_s}, \psi_{H_s})_{H_s} \\
(3.41) \quad &\geq -C_5|\psi_{H_s}|_{H_s}^2 - \frac{2}{\text{rank}(E)}(d \text{tr } D_{H_s}^*\psi_{H_s} \otimes \text{id}_E, \psi_{H_s})_{H_s} \\
&= -C_5|\psi_{H_s}|_{H_s}^2 - \frac{2}{\text{rank}(E)}(d \text{tr } D_K^*\psi_K \otimes \text{id}_E, \psi_{H_s})_{H_s} \\
&\geq -C_6|\psi_{H_s}|_{H_s}^2 - C_7,
\end{aligned}$$

where C_6 depends only on the lower bounded of the Ricci curvature at the considered point, C_7 depends only on $|d \text{tr } D_K^*\psi_K|$ and $\text{rank}(E)$.

On the other hand, (2.15) implies there exists two constants C_8, C_9 that depend only on \tilde{C}_1 and $\text{rank}(E)$, such that

$$(3.42) \quad |\psi_{H_s}|_{H_s}^2 \leq C_8(|\psi_K|^2 + |Dh_s|^2),$$

$$(3.43) \quad |Dh_s|^2 \leq C_9(|\psi_K|^2 + |\psi_{H_s}|_{H_s}^2).$$

So combining (3.42), (3.43) with Proposition 3.7, (3.41) and the Theorem 9.20 in [18], we get the uniform local boundedness of $|Dh_s|$. Now on each M_s , by the Poisson metric equation, we have

$$(3.44) \quad D_K^*(h_s^{-1}\delta_K h_s) = 2(D_K^*\psi_K)^\perp - g^{ij}[h_s^{-1}\delta_{K, \frac{\partial}{\partial x^i}}h_s, \psi_K(\frac{\partial}{\partial x^j})],$$

the standard bootstrapping procedure implies the uniform local higher order estimates of $\{h_s\}$ and by diagonal subsequence argument, H_s converge to a metric H_∞ on the whole M in C_{loc}^∞ -topology. It obvious that H_∞ is a Poisson metric on (E, D) such that $\det h = 1$, $|h| \in L^\infty$ and $|Dh| \in L^2$. \square

Proof of Theorem 1.3. The existence is given by Proposition 3.8 and the uniqueness will be proved in Proposition 4.1. Conversely, we assume D is irreducible and H be a Poisson metric which shares the same properties with H_∞ , we shall demonstrate that it is also analytically stable with respect to K . For any D -invariant sub-bundle $S \subset E$, K and H restrict to the metrics K_S and H_S on S . Taking the orthogonal complement of S in E with respect to K , we have the orthogonal decomposition $E = S \oplus S^\perp$ and the projection π_K onto S .

If $|D\pi_K| \notin L^2$, by (2.6) and $|D_K^*\psi_K| \in L^1$, we know

$$(3.45) \quad \deg_g(S, D, K) = -\infty.$$

If $|D\pi_K| \in L^2$, we know $\deg_g(S, D, K)$ is finite. Moreover, we have

$$(3.46) \quad h_S = \pi_K \circ h \circ \pi_K,$$

$$(3.47) \quad D_S h_S = \pi_K \circ Dh \circ \pi_K + D\pi_K \circ (\text{id}_E - \pi_K) \circ h \circ \pi_K,$$

where $h_S = K_S^{-1}H_S$ and D_S is the induced connection on S . It then follows

$$(3.48) \quad \int_M |\text{tr}(h_S^{-1}\delta_{D_S, K_S}h_S)|^2 \text{dvol}_g \leq \int_M |h_S^{-1}|^2 |D_S h_S|^2 \text{dvol}_g < \infty,$$

where $\delta_{D_S, K_S} = D_{S, K} - \psi_{D_S, K_S}$. So Simpson's Lemma 5.2 in [39] (or Yau's Lemma in [46]) implies

$$(3.49) \quad \lim_{j \rightarrow \infty} \int_{M_j} \frac{1}{2} d(*d \log \det h_S) = \lim_{j \rightarrow \infty} \int_{M_j} \frac{1}{2} d(*\operatorname{tr}(h_S^{-1} \delta_{D_S, K_S} h_S)) = 0,$$

where each M_j is an exhaustion subset. It then follows from the convergence theorem that

$$(3.50) \quad \begin{aligned} \lim_{j \rightarrow \infty} \int_{M_j} \operatorname{tr} D_{H_S}^* \psi_{H_S} \, \mathrm{dvol}_g &= \lim_{j \rightarrow \infty} \int_{M_j} \operatorname{tr} D_{K_S}^* \psi_{K_S} \, \mathrm{dvol}_g \\ &+ \lim_{j \rightarrow \infty} \int_{M_j} \frac{1}{2} \Delta \log \det h_S \, \mathrm{dvol}_g \\ &= -\deg_g(S, D, K), \end{aligned}$$

and hence

$$(3.51) \quad \lim_{j \rightarrow \infty} \int_{M_j} \operatorname{tr} D_{H_S}^* \psi_{H_S} \, \mathrm{dvol}_g = \int_M \operatorname{tr} D_{H_S}^* \psi_{H_S} \, \mathrm{dvol}_g = -\deg_g(S, D, H).$$

In summary, we conclude either

$$(3.52) \quad \deg_g(S, D, K) = -\infty$$

or

$$(3.53) \quad \deg_g(S, D, K) = \deg_g(S, D, H).$$

On the other hand, it follows from Proposition 2.1 that E is H -analytically stable and hence the proof is completed. \square

4. SOME APPLICATIONS AND CONSEQUENCES

4.1. Uniqueness of Poisson metrics. Firstly, we get the following uniqueness result, which implies the limiting Poisson metric in Theorem 1.3 is unique.

Proposition 4.1. *Let (E, D) be a K -analytically stable bundle over a Riemannian manifold (M, g) satisfying the Assumptions 2.1 and either Assumption 2.2 or complete with finite volume. Suppose $\operatorname{tr} D_K^* \psi_K \in L^1$ and \tilde{h} be a Poisson metric such that $\det h = 1, |h| \in L^\infty$ and $|Dh| \in L^2$, where $h = K^{-1}H$. If \tilde{H} is another Poisson metric which is mutually bounded with H and $\det H = \det \tilde{H}$, we have $H = \tilde{H}$.*

Proof. Let $\tilde{h} = H^{-1}\tilde{H}$ and then we have

$$(4.1) \quad \begin{aligned} \Delta \operatorname{tr} \tilde{h} &= 2(D_{\tilde{H}}^* \psi_{\tilde{H}} - D_H^* \psi_H, \tilde{h})_H + |\tilde{h}^{-\frac{1}{2}} \delta_H \tilde{h}|_H^2 \\ &= \frac{(\Delta \log \det \tilde{h} \operatorname{id}_E, \tilde{h})_H}{\operatorname{rank}(E)} + |\tilde{h}^{-\frac{1}{2}} \delta_H \tilde{h}|_H^2 \\ &= |\tilde{h}^{-\frac{1}{2}} \delta_H \tilde{h}|_H^2. \end{aligned}$$

So Assumption 2.1 implies \tilde{h} is D -parallel and set $E = \bigoplus_{j=1}^m E_j$ be the eigendecomposition of \tilde{h} , then this decomposition is orthogonal with respect H and \tilde{H} . Moreover, we have $\tilde{H}|_{E_j} = c_j H|_{E_j}$ for some constants c_j and

$$(4.2) \quad -\int_M \operatorname{tr} D_H^* \psi_H \, \mathrm{dvol}_g = -\sum_{j=1}^m \int_M \operatorname{tr} D_{H|_{E_j}}^* \psi_{H|_{E_j}} \, \mathrm{dvol}_g.$$

Then running the same argument in the proof of Theorem 1.3 shows that D being H -analytically stable. As a consequence,

$$(4.3) \quad \begin{aligned} \frac{\deg(E, D, H)}{\text{rank}(E)} &= - \sum_{j=1}^m \frac{\int_M \text{tr } D_{H|E_j}^* \psi_{H|E_j} \, \text{dvol}_g}{\text{rank}(E_j)} \frac{\text{rank}(E_j)}{\text{rank}(E)} \\ &\leq \frac{\deg(E, D, H)}{\text{rank}(E)}, \end{aligned}$$

the strict inequality occurs if and only if $m > 1$ and therefore we complete the proof. \square

If D being flat, the following is a flat bundle analogy of Mochizuki's uniqueness result on Hermitian-Einstein metrics in [33].

Proposition 4.2. *Let (E, D) be a K -analytically stable flat bundle over a Riemannian manifold (M, g) satisfying Assumption 2.1 and there is a positive exhaustion function ρ such that $|d \log \rho| \in L^1$. Assume that $|\psi_K| \in L^2$ and H_1 be a Poisson metric such that $|h_1|, |h_1^{-1}| \in L^\infty$ and $|Dh_1| \in L^2$, where $h_1 = K^{-1}H_1$. If H_2 is another Poisson metric which is mutually bounded with H_1 and $\det H_1 = \det H_2$, then $H_1 = H_2$.*

Proof. Similar, set $E = \bigoplus_{j=1}^m E_j$ be the eigendecomposition of $h_{12} = H_1^{-1}H_2$ and $H_1|_{E_j} = c_j H_2|_{E_j}$ for some constants c_j . Let π_j denotes the projection onto E_j and $h_i = K^{-1}H_i$, then π_j are bounded with respect to K , as K and H_i are mutually bounded.

Now (2.16) indicates $\delta_K \pi_j = \delta_{H_1} \pi_j - [h_1^{-1} \delta_K h_1, \pi_j] = -[h_1^{-1} \delta_K h_1, \pi_j]$, this implies $\delta_K \pi_j$ and $D\pi_j^{*K}$ are L^2 . We consider the fiber metric H determined by the direct sum of $K|_{E_j}$, then H and K are mutually bounded. By definition, we have $h := K^{-1}H = \sum_{j=1}^m \pi_j^{*K} \circ \pi_j$ and it follows

$$(4.4) \quad h^{-1} Dh = h^{-1} \sum_{j=1}^m D\pi_j^{*K} \circ \pi_j, \quad h^{-1} \delta_K h = h^{-1} \sum_{j=1}^m \pi_j^{*K} \circ \delta_K \pi_j,$$

which implies $h^{-1} Dh$ and $h^{-1} \delta_K h$ are L^2 .

Next we compute $D(h^{-1} \delta_K h) = -h^{-1} Dh h^{-1} \delta_K h + h^{-1} D(\delta_K h)$ and

$$(4.5) \quad \begin{aligned} D(\delta_K h) &= D\left(\sum_{j=1}^m \pi_j^{*K} \circ \delta_K \pi_j\right) \\ &= \sum_{j=1}^m \left(D\pi_j^{*K} \circ \delta_K \pi_j + \pi_j^{*K} \circ D(\delta_K \pi_j)\right) \\ &= \sum_{j=1}^m \left(D\pi_j^{*K} \circ \delta_K \pi_j + 4\pi_j^{*K} \circ [D_K^2, \pi_j]\right) \\ &= \sum_{j=1}^m \left(D\pi_j^{*K} \circ \delta_K \pi_j - 2\pi_j^{*K} \circ [[\psi_K, \psi_K], \pi_j]\right). \end{aligned}$$

where we have use the flatness of D . As a consequence, we find $\Delta \log \det h$ is L^1 .

Finally we set $\chi_N = \eta(\frac{\rho}{N})$, where η is a nonnegative function with $\eta(t) = 0$ if $t \geq 2$ and $\eta(t) = 1$ if $t \leq 1$. We then deduce

$$\begin{aligned}
\int_M \chi_N \Delta \log \det h \, d\text{vol}_g &= - \int_M \nabla \chi_N \cdot \nabla \log \det h \, d\text{vol}_g \\
&= - \int_{\{N \leq \rho \leq 2N\}} \frac{\eta'(\frac{\rho}{N})}{N} \nabla \rho \cdot \nabla \log \det h \, d\text{vol}_g \\
(4.6) \quad &\leq \int_{\{N \leq \rho \leq 2N\}} \frac{2|\eta'(\frac{\rho}{N})|}{\rho} |\nabla \rho| |\nabla \log \det h| \, d\text{vol}_g \\
&= \int_{\{N \leq \rho \leq 2N\}} 2|\eta'(\frac{\rho}{N})| |\nabla \log \rho| |\text{tr}(h^{-1} \delta_K h)| \, d\text{vol}_g.
\end{aligned}$$

Combing this and using the fact that $\Delta \log \det h$ lies in L^1 , it follows

$$\begin{aligned}
- \int_M \text{tr} D_K^* \psi_K \, d\text{vol}_g &= - \int_M \text{tr} D_H^* \psi_H \, d\text{vol}_g + \frac{1}{2} \int_M \Delta \log \det h \, d\text{vol}_g \\
(4.7) \quad &= - \sum_{j=1}^m \int_M \text{tr} D_{K|E_j}^* \psi_{K|E_j} \, d\text{vol}_g.
\end{aligned}$$

Since D is K -analytically stable, we obtain $m = 1$ and the proof is completed. \square

4.2. Applications to the nonabelian Hodge correspondence.

Proposition 4.3. *Let (X, ω) be a noncompact complex curve satisfying the Assumption 2.1, and (E, D) be a K -analytically stable flat bundle over X with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C , then there exists a Higgs structure on E .*

Proof. According to Theorem 1.3, there is a Poisson metric H on (E, D) and we have

$$(4.8) \quad D_H \psi_H^\perp = D_H(\psi_H^{\perp, 1, 0} + \psi_H^{\perp, 0, 1}) = 0,$$

$$(4.9) \quad D_H^* \psi_H^\perp = \sqrt{-1} * D_H(\psi_H^{\perp, 1, 0} - \psi_H^{\perp, 0, 1}) = 0,$$

where ψ_H^\perp is the trace-free part of ψ_H . Then we easily get $D_H^{0,1} \psi_H^{\perp, 1, 0} = 0$ and the pair $(D_H^{0,1}, \psi_H^{\perp, 1, 0})$ gives rise a Higgs structure on E . \square

For higher dimensional case, we have

Proposition 4.4. *Let (X, ω) be a complete Kähler manifold with bounded Ricci curvature from below and satisfying the Assumption 2.1. Assume (E, D) be a K -analytically stable flat bundle over X with $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C and $|\psi_K| \in L^2$. Then there exists a Higgs structure on E .*

Proof. Due to Theorem 1.3, there is a Poisson metric H and we compute

$$\begin{aligned}
(4.10) \quad \Delta|\psi_H^\perp|_H^2 &= -2(\nabla_H^* \nabla_H \psi_H^\perp, \psi_H^\perp)_H + 2|\nabla_H \psi_H^\perp|_H^2 \\
&= -2g^{ij}(D_H^2(\cdot, \frac{\partial}{\partial x^i})(\psi_H^\perp(\frac{\partial}{\partial x^j})), \psi_H^\perp)_H \\
&\quad + 2g^{ij}(\psi_H^\perp(R_\nabla(\cdot, \frac{\partial}{\partial x^i})(\frac{\partial}{\partial x^j})), \psi_H^\perp)_H + 2|\nabla_H \psi_H^\perp|_H^2 \\
&= g^{ij}([\psi_H, \psi_H(\frac{\partial}{\partial x^i})], \psi_H^\perp(\frac{\partial}{\partial x^j})) + 2\psi_H^\perp \circ \text{Ric}, \psi_H^\perp)_H + 2|\nabla_H \psi_H^\perp|_H^2 \\
&= |[\psi_H^\perp, \psi_H^\perp]|_H^2 + 2(\psi_H^\perp \circ \text{Ric}, \psi_H^\perp)_H + 2|\nabla_H \psi_H^\perp|_H^2 \\
&\geq -C_*|\psi_H^\perp|_H^2 + 2|\nabla_H \psi_H^\perp|_H^2,
\end{aligned}$$

where $-C_*$ the lower bound of Ricci curvature. Let R be any positive constant and we fix point x_0 , choose a cut-off function η satisfying

$$(4.11) \quad \begin{cases} \eta(x) = 1, x \in B_{x_0}(R), \\ \eta(x) = 0, x \in X \setminus B_{x_0}(2R), \\ 0 \leq \eta \leq 1, |\nabla \eta| \leq \frac{C_1}{R}, \end{cases}$$

where C_1 is a positive constant and $B_{x_0}(R)$ is the geodesics ball centered at x_0 with radius R . Then we have

$$\begin{aligned}
(4.12) \quad 2 \int_X \eta^2 |\nabla_H \psi_H^\perp|_H^2 \, d\text{vol}_\omega &\leq \int_X \eta^2 (\Delta|\psi_H^\perp|_H^2 + C_*|\psi_H^\perp|_H^2) \, d\text{vol}_\omega \\
&\leq \int_X 4\eta |\nabla \eta| |\psi_H^\perp|_H |\nabla_H \psi_H^\perp|_H \, d\text{vol}_\omega + C_* \int_X \eta^2 |\psi_H^\perp|_H^2 \, d\text{vol}_\omega \\
&\leq \int_X \eta^2 |\nabla_H \psi_H^\perp|_H^2 \, d\text{vol}_\omega + \int_X (C_*\eta^2 + 4|\nabla \eta|^2) |\psi_H^\perp|_H^2 \, d\text{vol}_\omega \\
&\leq \int_X \eta^2 |\nabla_H \psi_H^\perp|_H^2 \, d\text{vol}_\omega + \int_X (C_*\eta^2 + \frac{4C_1^2}{R^2}) |\psi_H^\perp|_H^2 \, d\text{vol}_\omega.
\end{aligned}$$

And it also holds

$$(4.13) \quad \int_X |\psi_H^\perp|_H^2 \, d\text{vol}_\omega = \int_X |\psi_K^\perp - \frac{1}{2}h^{-1}Dh + h^{-1}[\psi_K^\perp, h]|_H^2 \, d\text{vol}_\omega < \infty,$$

where $h = K^{-1}H$. So by letting R goes infinity, we see

$$(4.14) \quad \int_X |\nabla_H \psi_H^\perp|_H^2 \, d\text{vol}_\omega \leq C_* \int_X |\psi_H^\perp|_H^2 \, d\text{vol}_\omega < \infty.$$

Now let's consider the pseudo-curvature operator $G_H = (D_H^{0,1} + \psi_H^{1,0})^2$. Notice the flatness of D implies

$$(4.15) \quad D_H^{1,0}\psi_H^{0,1} = -D_H^{0,1}\psi_H^{1,0}, D_H^{1,0}D_H^{1,0} = -\frac{1}{2}[\psi_H^{1,0}, \psi_H^{1,0}], D_H^{0,1}D_H^{0,1} = -\frac{1}{2}[\psi_H^{0,1}, \psi_H^{0,1}],$$

and it follows that

$$\begin{aligned}
(4.16) \quad G_H^* &= \left((D_H^{0,1})^2 + D_H^{0,1}\psi_H^{1,0} + \psi_H^{1,0} \wedge \psi_H^{1,0} \right)^* \\
&= -D_H^{1,0}D_H^{1,0} + D_H^{1,0}\psi_H^{0,1} - \psi_H^{0,1} \wedge \psi_H^{0,1} \\
&= G_H^{2,0} - G_H^{1,1} + G_H^{0,2}.
\end{aligned}$$

By Kähler identity we know $\sqrt{-1}\Lambda_\omega G_H^\perp = \frac{1}{2}(D_H^* \psi_H)^\perp = 0$ and

$$(4.17) \quad \text{tr}(G_H^\perp \wedge G_H^\perp) \wedge \frac{\omega^{n-2}}{(n-2)!} = \text{tr}\left(G_H^\perp \wedge *(G_H^\perp)^*\right) = |G_H^\perp|_H^2 \text{dvol}_\omega,$$

where n is the complex dimension of X . On the other hand, a direct calculation shows

$$(4.18) \quad \begin{aligned} \text{tr}(G_H^\perp \wedge G_H^\perp) \wedge \omega^{n-2} &= -2 \text{tr}(\psi_H^{0,1} \wedge \psi_H^{0,1} \wedge \psi_H^{1,0} \wedge \psi_H^{1,0}) \wedge \omega^{n-2} \\ &\quad - \text{tr}(D_H^{0,1} \psi_H^{1,0} \wedge D_H^{1,0} \psi_H^{0,1}) \wedge \omega^{n-2} \\ &\quad - \frac{\text{tr}(D_H^{0,1} \psi_H^{1,0}) \wedge \text{tr}(D_H^{1,0} \psi_H^{0,1})}{\text{rank}(E)} \wedge \omega^{n-2} \\ &= -\bar{\partial} \left(\text{tr}(\psi_H^{1,0} \wedge D_H^{1,0} \psi_H^{0,1}) \wedge \omega^{n-2} \right) \\ &\quad + \bar{\partial} \left(\frac{\text{tr} \psi_H^{1,0} \wedge \text{tr}(D_H^{1,0} \psi_H^{0,1})}{\text{rank}(E)} \wedge \omega^{n-2} \right) \\ &= -\bar{\partial} \left(\text{tr}(\psi_H^{1,0} \wedge D_H^{1,0} \psi_H^\perp) \wedge \omega^{n-2} \right). \end{aligned}$$

where we have used (4.15), $D_H^{0,1} \psi_H^{0,1} = 0$ and $[D_H^{1,0}, D_H^{0,1}] = -[\psi_H^{1,0}, \psi_H^{0,1}]$.

Now as it holds

$$(4.19) \quad D_H^{1,0} \psi_H^\perp = \sum_{i=1}^n dz^i \wedge \nabla_{H, \frac{\partial}{\partial z^i}} \psi_H^\perp,$$

it follows from (4.14) that

$$(4.20) \quad \int_X |\text{tr}(\psi_H^{1,0} \wedge D_H^{1,0} \psi_H^\perp) \wedge \omega^{n-2}| \text{dvol}_\omega < \infty.$$

Therefore, by (4.17), (4.18), (4.20) and Yau's Lemma in [46], we conclude G_H^\perp vanishes identically and the pair $(D_H^{0,1}, \psi_H^{\perp,1,0})$ is a Higgs structure on E . \square

Proof of Theorem 1.4. Proposition 4.3 and Proposition 4.4 indicate $(D_H^{0,1}, \psi_H^{\perp,1,0})$ determines a Higgs structure on E . Moreover, we see the corresponding Hitchin-Simpson curvature

$$(4.21) \quad F_{D_H^{0,1}, \psi_H^{\perp,1,0}, H} = D_H^2 + D_H \psi_H^\perp + \frac{1}{2}[\psi_H^\perp, \psi_H^\perp] = 0,$$

where we have used the flatness of D . Hence we complete of proof of (1) and (2). For (3), if (X, ω) has nonnegative Ricci curvature, we have for any $\epsilon > 0$ that

$$(4.22) \quad \begin{aligned} \Delta\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{1}{2}} &= \frac{1}{4} \frac{\Delta|\psi_H^\perp|_H^2}{\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{1}{2}}} - \frac{1}{4} \frac{\frac{1}{2}d|\psi_H^\perp|_H^2|^2}{\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{3}{2}}} \\ &\geq \frac{1}{4} \frac{2|\nabla_H \psi_H^\perp|_H^2}{\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{1}{2}}} - \frac{1}{4} \frac{|\psi_H^\perp|_H^2 |\nabla_H \psi_H^\perp|_H^2}{\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{3}{2}}} \\ &= \frac{1}{2} \frac{|\nabla_H \psi_H^\perp|_H^2}{\left(\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon\right)^{\frac{1}{2}}} \left(1 - \frac{\frac{1}{2}|\psi_H^\perp|_H^2}{\frac{1}{2}|\psi_H^\perp|_H^2 + \epsilon}\right) \\ &\geq 0, \end{aligned}$$

and by letting ϵ goes zero, it follows that $|\psi_H^\perp|_H$ is subharmonic. On the other hand, as $\|\psi_H^\perp\|_{H, L^2} < \infty$, Yau's Liouville theorem in [46] applies and we conclude $|\psi_H^\perp|_H$ is constant. But the volume of (X, ω) is infinite, so we know $\psi_H^\perp = 0$ and then (3) follows easily. \square

In a similar way, using Corollary 1.2 instead, we obtain the following proposition, which means the harmonic metric produced there is pluriharmonic.

Proposition 4.5. *Let (X, ω) be a Kähler manifold satisfying the Assumptions 2.1, 2.3 with $\phi = 1$ and (E, D) be a K -analytically stable flat bundle over X such that $|D_K^* \psi_K| \in L^\infty$ and $|\psi_K| \in L^2$. Assume either $\dim_{\mathbb{C}} X = 1$ or (X, ω) being complete with bounded Ricci curvature from below, then (E, D) comes from a Higgs bundle.*

Suppose (E, D) be a flat bundle coming from a Higgs bundle $(E, \bar{\partial}_E, \theta)$ via a harmonic metric H , then we call (E, D, H) is a harmonic bundle. Denote by $H_{DR}^*(X, E)$ and $H_{Dol}^*(X, E)$ be the set of L^∞ -elements in the cohomology groups of the complexes $(\Omega^*(E), D)$ and $(\Omega^*(E), \bar{\partial}_{E, \theta})$ respectively.

Lemma 4.1. *Assume (X, ω) be a Kähler manifold satisfying Assumption 2.1.*

- (1) *If (E, D) be a flat bundle and H is a harmonic metric, then any D -parallel L^∞ -section is also $D_H^{0,1} + \psi_H^{1,0}$ -parallel.*
- (2) *If $(E, \bar{\partial}_E, \theta)$ be a Higgs bundle and H is a Hermitian-Einstein metric such that the $\sqrt{-1}\Lambda_\omega F_{\bar{\partial}_E, \theta, H} = 0$, then any $\bar{\partial}_{E, \theta}$ -parallel L^∞ -section is also $D_{\bar{\partial}_E, \theta, H}$ -parallel.*
- (3) *If (E, D, H) be a harmonic bundle, then $H_{DR}^0(X, E) = H_{Dol}^0(X, E)$.*

Proof. The statement (3) follows from (1) and (2), Firstly suppose f is D -parallel, we have

$$\begin{aligned}
\Delta|f|_H^2 &= \sqrt{-1}\Lambda_\omega(D_H^{1,0}D_H^{0,1}f, f)_H + \sqrt{-1}\Lambda_\omega(D_H^{1,0}f, D_H^{1,0}f)_H \\
&\quad - \sqrt{-1}\Lambda_\omega(D_H^{0,1}f, D_H^{0,1}f)_H + \sqrt{-1}\Lambda_\omega(f, D_H^{0,1}D_H^{1,0}f)_H \\
(4.23) \quad &= |D_H^{0,1}f|_H^2 + |\psi_H^{1,0}f|_H^2 - \sqrt{-1}\Lambda_\omega(D_H^{1,0}\psi_H^{0,1}f, f)_H - \sqrt{-1}\Lambda_\omega(f, D_H^{0,1}\psi_H^{1,0}f)_H \\
&= |D_H^{0,1}f|_H^2 + |\psi_H^{1,0}f|_H^2 + \sqrt{-1}\Lambda_\omega(\psi_H^{0,1}D_H^{1,0}f, f)_H + \sqrt{-1}\Lambda_\omega(f, \psi_H^{1,0}D_H^{0,1}f)_H \\
&= |D_H^{0,1}f|_H^2 + |\psi_H^{1,0}f|_H^2 - \sqrt{-1}\Lambda_\omega(D_H^{1,0}f, \psi_H^{1,0}f)_H + \sqrt{-1}\Lambda_\omega(\psi_H^{0,1}f, D_H^{0,1}f)_H \\
&= 2|(D_H^{0,1} + \psi_H^{1,0})f|_H^2.
\end{aligned}$$

By Assumption 2.1 we have f is $D_H^{0,1} + \psi_H^{1,0}$ -parallel.

On the other hand, if f is $\bar{\partial}_{E, \theta}$ -parallel, we deduce

$$\begin{aligned}
\Delta|f|_H^2 &= \sqrt{-1}\Lambda_\omega(\partial_H f, \partial_H f)_H + \sqrt{-1}\Lambda_\omega(f, \bar{\partial}_E \partial_H f)_H \\
(4.24) \quad &= |\partial_H f|_H^2 + \sqrt{-1}\Lambda_\omega(f, F_{\bar{\partial}_E, \theta, H} f)_H - \sqrt{-1}\Lambda_\omega(f, [\theta, \theta^{*H}]f)_H \\
&= |\partial_{\theta, H} f|_H^2,
\end{aligned}$$

and then we see f is $D_{\bar{\partial}_E, \theta, H}$ -parallel. □

Given a vector bundle E over a complex manifold (X, ω) , equipped with a Hermitian metric H and a connection D . The contraction of pseudo-curvature operator can be written as

$$\begin{aligned}
\sqrt{-1}\Lambda_\omega G_H &= \frac{\sqrt{-1}}{4}\Lambda_\omega[D^{0,1} + \delta_H^{0,1}, D^{1,0} - \delta_H^{1,0}] \\
(4.25) \quad &= \frac{\sqrt{-1}}{4}\Lambda_\omega \left(D^2 - (\delta_H^{1,0} + D^{0,1})^2 + (D^{1,0} + \delta_H^{0,1})^2 - \delta_H^2 \right),
\end{aligned}$$

If D is flat, we see

$$\begin{aligned}
\sqrt{-1}\Lambda_\omega G_H &= \frac{\sqrt{-1}}{4}\Lambda_\omega \left(-(\delta_H^{1,0} + D^{0,1})^2 + (D^{1,0} + \delta_H^{0,1})^2 \right) \\
(4.26) \quad &= \frac{\sqrt{-1}}{4}\Lambda_\omega (-[D^{0,1}, \delta_K^{1,0} + h^{-1}\delta_K^{1,0}h] + [\delta_K^{0,1} + h^{-1}\delta_K^{0,1}h, D^{1,0}]) \\
&= \sqrt{-1}\Lambda_\omega G_K + \frac{\sqrt{-1}}{4}\Lambda_\omega D(h^{-1}D_K^c h),
\end{aligned}$$

where $h = K^{-1}H$ and $D_K^c = \delta_K^{0,1} - \delta_K^{1,0}$.

Finally, we are in the position prove Theorem 1.5.

Proof of Theorem 1.5. Given $[D] \in \mathcal{M}_{Flat,K,\phi}$, by Theorem 1.3, we yield a Poisson metric H with $\det h = 1, |h| \in L^\infty, \|Dh\|_{L^2} \leq C\|\phi\|_{L^1}^{\frac{1}{2}}$ for a positive constant C and $h = K^{-1}H$. Then Proposition 4.3 implies $(\bar{\partial}_E, \theta) = (D_H^{0,1}, \psi_H^{\perp,1,0})$ determines an irreducible Higgs structure with $c_1(E, H) = c_1(E, K) = 0$. Moreover, it can be seen H is harmonic with respect to $D_H + \psi_H^\perp$ and due to Lemma 4.1, the assignment $[D] \mapsto [(\bar{\partial}_E, \theta)]$ is well-defined. Then we compute

$$\begin{aligned}
\sqrt{-1}\Lambda_\omega F_{\bar{\partial}_E, \theta, K} &= \sqrt{-1}\Lambda_\omega F_{\bar{\partial}_E, \theta, H} + \sqrt{-1}\Lambda_\omega \bar{\partial}_{E, \theta}(h\partial_{\theta, H}h^{-1}) \\
&= \sqrt{-1}\Lambda_\omega D_{\bar{\partial}_E, \theta, H}(h\partial_{\theta, H}h^{-1}) - \sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\partial_{\theta, H}h^{-1}) \\
&= -\sqrt{-1}\Lambda_\omega D_{\bar{\partial}_E, \theta, H}(hD_{\bar{\partial}_E, \theta, H}^c h^{-1}) + \sqrt{-1}\Lambda_\omega D_{\bar{\partial}_E, \theta, H}(h\bar{\partial}_{E, \theta}h^{-1}) \\
&\quad - \sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\partial_{\theta, H}h^{-1}) \\
(4.27) \quad &= 4\sqrt{-1}\Lambda_\omega G_H^\perp - 4\sqrt{-1}\Lambda_\omega G_K^\perp + \sqrt{-1}\Lambda_\omega D_{\bar{\partial}_E, \theta, H}(h\bar{\partial}_{E, \theta}h^{-1}) \\
&\quad - \sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\partial_{\theta, H}h^{-1}) \\
&= -2(D_K^* \psi_K)^\perp + \sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\bar{\partial}_{E, \theta}h^{-1}) \\
&\quad + \sqrt{-1}\Lambda_\omega \bar{\partial}_{E, \theta}(h\bar{\partial}_{E, \theta}h^{-1}) - \sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\partial_{\theta, H}h^{-1}),
\end{aligned}$$

and

$$\begin{aligned}
\sqrt{-1}\Lambda_\omega \partial_{\theta, H}(h\bar{\partial}_{E, \theta}h^{-1}) &= \sqrt{-1}\Lambda_\omega (\partial_{\theta, H}h \wedge \bar{\partial}_{E, \theta}h^{-1}) + \sqrt{-1}\Lambda_\omega (\bar{\partial}_{E, \theta}h \wedge \partial_{\theta, H}h^{-1}) \\
(4.28) \quad &\quad - \sqrt{-1}\Lambda_\omega (\bar{\partial}_{E, \theta}h \wedge \partial_{\theta, H}h^{-1}) - \sqrt{-1}\Lambda_\omega (h\bar{\partial}_{E, \theta}\partial_{\theta, H}h^{-1}) \\
&= \sqrt{-1}\Lambda_\omega (\partial_{\theta, H}h \wedge \bar{\partial}_{E, \theta}h^{-1}) + \sqrt{-1}\Lambda_\omega (\bar{\partial}_{E, \theta}h \wedge \partial_{\theta, H}h^{-1}) \\
&\quad - \sqrt{-1}\Lambda_\omega F_{\bar{\partial}_E, \theta, K}.
\end{aligned}$$

where we have used (4.26). We obtain from above two equalities that

$$\begin{aligned}
\sqrt{-1}\Lambda_\omega F_{\bar{\partial}_E, \theta, K} &= -(D_K^* \psi_K)^\perp + \frac{\sqrt{-1}}{2}\Lambda_\omega (\partial_{\theta, H}h \wedge \bar{\partial}_{E, \theta}h^{-1}) \\
(4.29) \quad &\quad + \frac{\sqrt{-1}}{2}\Lambda_\omega (\bar{\partial}_{E, \theta}h \wedge \partial_{\theta, H}h^{-1}) + \frac{\sqrt{-1}}{2}\bar{\partial}_{E, \theta}(h\bar{\partial}_{E, \theta}h^{-1}) \\
&\quad - \frac{\sqrt{-1}}{2}\Lambda_\omega \partial_{\theta, H}(h\partial_{\theta, H}h^{-1}),
\end{aligned}$$

from which we conclude $|\Lambda_\omega F_{\bar{\partial}_E, \theta, K}| \in L^1$.

For any θ -invariant sub-holomorphic bundle $V \subset E$, we have the orthogonal decomposition $E = V \oplus V^\perp$ with respect to K and the projection π_K onto V . We denote by K_V, H_V the

metrics on V induced by K and H . If $|\bar{\partial}_{E,\theta}\pi_K| \notin L^2$, by Chern-Weil formula (1.8) and $|\Lambda_\omega F_{\bar{\partial}_{E,\theta},K}| \in L^1$, we know

$$(4.30) \quad \deg_\omega(V, K) = -\infty.$$

If $|\bar{\partial}_{E,\theta}\pi_K| \in L^2$, the same reason indicates $\deg_\omega(V, K)$ is finite. It also holds

$$(4.31) \quad h_V = \pi_K \circ h \circ \pi_K,$$

$$(4.32) \quad \bar{\partial}_{V,\theta_V} h_V = \pi_K \circ \bar{\partial}_{E,\theta} h \circ \pi_K + \bar{\partial}_{E,\theta}\pi_K \circ (\text{id}_E - \pi_K) \circ h \circ \pi_K,$$

where $h_V = K_V^{-1} H_V$ and $\bar{\partial}_{V,\theta_V}$ is the induced Higgs structure on V . Therefore

$$(4.33) \quad \|\text{tr}(h_V^{-1} \partial_{\theta_V, K_V} h_V)\|_{L^2} \leq \|h_V^{-1}\|_{L^\infty} \|\bar{\partial}_{V,\theta_V} h_V\|_{L^2} < \infty.$$

So Simpson's Lemma 5.2 in [39] (or Yau's Lemma in [46]) applies,

$$(4.34) \quad \lim_{j \rightarrow \infty} \int_{X_j} \sqrt{-1} \bar{\partial} \text{tr}(h_V^{-1} \partial_{\theta_V, K_V} h_V) = 0,$$

where $\{X_j\}$ is a sequence of exhaustion subsets. And

$$(4.35) \quad \begin{aligned} \lim_{j \rightarrow \infty} \int_{X_j} \sqrt{-1} \text{tr} F_{\bar{\partial}_V, \theta_V, H_V} &= \lim_{j \rightarrow \infty} \int_{X_j} \sqrt{-1} \text{tr} F_{\bar{\partial}_V, \theta_V, K_V} \\ &+ \lim_{j \rightarrow \infty} \int_{X_j} \sqrt{-1} \text{tr}(\bar{\partial}_{V,\theta_V}(h_V^{-1} \partial_{\theta_V, K} h_V)) \\ &= \deg_\omega(V, K), \end{aligned}$$

then we have

$$(4.36) \quad \lim_{j \rightarrow \infty} \int_{X_j} \sqrt{-1} \text{tr} F_{\bar{\partial}_V, \theta_V, H_V} = \int_X \sqrt{-1} \text{tr} F_{\bar{\partial}_V, \theta_V, H_V} = \deg_\omega(V, H).$$

Since $(\bar{\partial}_E, \theta)$ is H -analytically stable, we conclude by (4.30), (4.35) and (4.36) that $(\bar{\partial}_E, \theta)$ is K -analytically stable.

Next assume \tilde{H} be another Hermitian-Einstein metric such that $\det \tilde{h} = 1$ and $|\tilde{h}| \in L^\infty$ for $\tilde{h} = K^{-1} \tilde{H}$. Set $\hat{h} = H^{-1} \tilde{H}$, we have $\sqrt{-1} \Lambda_\omega \partial \bar{\partial} \text{tr} \hat{h} = |\hat{h}^{-\frac{1}{2}} \partial_{\theta, H} \hat{h}|_H^2$ and the Assumption 2.1 implies $\partial_{\theta, H} \hat{h} = 0$. We let $E = \bigoplus_{j=1}^m E_j$ be the eigendecomposition of \hat{h} , which is orthogonal

with respect to H and \tilde{H} . Moreover, it holds $\bar{\partial}_{E,\theta}\pi_j = 0$ for the projection π_j onto E_j . But $(\bar{\partial}_E, \theta)$ is H -analytically stable, we know $m = 1$, $H = \tilde{H}$ and therefore $[(\bar{\partial}_E, \theta)] \in \mathcal{M}_{Higgs, K}$.

Conversely, given $[(\bar{\partial}_E, \theta)] \in \mathcal{M}_{Higgs, K, \phi}$, by Mochizuki's work [33] we know that there exists a Hermitian-Einstein metric H with $\det h = 1$, $|h| \in L^\infty$ and $\|\bar{\partial}_{E,\theta} h\|_{L^2} \leq C \|\phi\|_{L^1}^{\frac{1}{2}}$ for a positive constant C , where $h = K^{-1} H$. Moreover, the Hitchin-Simpson connection $D_{\bar{\partial}_E, \theta, H}$ is flat and irreducible. Due to Lemma 4.1, the map $[(\bar{\partial}_E, \theta)] \mapsto [D_{\bar{\partial}_E, \theta, H}]$ is well-defined. Next as H is harmonic with respect to $D = D_{\bar{\partial}_E, \theta, H}$, a similar discussion as that in (4.29)

shows

$$\begin{aligned}
(4.37) \quad D_K^* \psi_K &= -\sqrt{-1} \Lambda_\omega F_{\bar{\partial}_{E,\theta},K} + \frac{\sqrt{-1}}{2} \Lambda_\omega (\partial_{\theta,H} h \wedge \bar{\partial}_{E,\theta} h^{-1}) \\
&+ \frac{\sqrt{-1}}{2} \Lambda_\omega (\bar{\partial}_{E,\theta} h \wedge \partial_{\theta,H} h^{-1}) + \frac{\sqrt{-1}}{2} \bar{\partial}_{E,\theta} (h \bar{\partial}_{E,\theta} h^{-1}) \\
&- \frac{\sqrt{-1}}{2} \Lambda_\omega \partial_{\theta,H} (h \partial_{\theta,H} h^{-1}).
\end{aligned}$$

and therefore $|(D_K^* \psi_K)^\perp| = |D_K^* \psi_K| \in L^1$. Then proceeding the argument in the proof of the Theorem 1.3, we conclude $D_{\bar{\partial}_{E,\theta},H}$ is analytically stable with respect to the background metric K and hence $[D_{\bar{\partial}_{E,\theta},H}] \in \mathcal{M}_{Flat,K}$. \square

4.3. Vanishing theorems of characteristic classes. Associated with a vector bundle (E, D) , up to some normalization coefficients, the Kamber-Tondeur classes are defined by

$$(4.38) \quad \alpha_{2k+1}(E, D) = [\text{tr } \psi_H^{2k+1}] \in H_{DR}^{2k+1}(M, \mathbb{R}),$$

and the class is independent of the choice of fiber metrics, see [4, 11, 16, 23, 35] for the related study of these characteristic classes.

By Proposition 4.4, we know $\psi_H^{1,0} \wedge \psi_H^{1,0} = 0$ and following [24], for any $k \geq 1$ we deduce

$$\begin{aligned}
(4.39) \quad \text{tr } \psi_H^{2k+1} &= \text{tr} \left((\psi_H^{1,0} \wedge \psi_H^{0,1})^{2k} \wedge \psi_H^{1,0} \right) + \text{tr} \left((\psi_H^{0,1} \wedge \psi_H^{1,0})^{2k} \wedge \psi_H^{0,1} \right) \\
&= \text{tr} \left(\psi_H^{1,0} \wedge (\psi_H^{1,0} \wedge \psi_H^{0,1})^{2k} \right) + \text{tr} \left(\psi_H^{0,1} \wedge (\psi_H^{0,1} \wedge \psi_H^{1,0})^{2k} \right) \\
&= 0.
\end{aligned}$$

So we conclude the following vanishing theorem.

Theorem 4.1. *Let (X, ω) be a complete Kähler manifold with bounded Ricci curvature from below and satisfying the Assumptions 2.1. Assume (E, D) be a K -analytically stable flat bundle over X such that $|(D_K^* \psi_K)^\perp| \leq C\phi$ for a constant C and $|\psi_K| \in L^2$. Then the characteristic classes $\alpha_{2k+1}(E, D)$ vanish for all $k \geq 1$.*

We mention that in the compact setting, this result was firstly proved in [36].

REFERENCES

- [1] O. Biquard, Fibrés de Higgs et connexions intégrables: le cas logarithmique (diviseur lisse). Ann Sci École Norm Sup (4), 1997, **30**(1): 41-96.
- [2] O. Biquard and P. Boalch, Wild non-abelian Hodge theory on curves. Compos Math, 2004, **140**(1): 179-204.
- [3] O. Biquard, O. García-Prada and I. Mundet i Riera, Parabolic Higgs bundles and representations of the fundamental group of a punctured surface into a real group. Adv Math, 2020, **372**: 107305.
- [4] J. M. Bismut and J. Lott, Flat vector bundles, direct images and higher real analytic torsion. J Amer Math Soc, 1995, **8**(2): 291-363.
- [5] I. Biswas, J. Loftin and M. Stemmler, Flat bundles on affine manifolds. Arab J Math (Springer), 2013, **2**(2): 159-175.
- [6] I. Biswas and H. Kasuya, Higgs bundles and flat connections over compact Sasakian manifolds. Commun Math Phys, 2021, **385**(1): 267-290.
- [7] T. C. Collins, A. Jacob and S. T. Yau, Poisson metrics on flat vector bundles over non-compact curves. Comm Anal Geom, 2019, **27**(3): 529-597.

- [8] K. Corlette, Flat G-bundles with canonical metrics. *J. Differential Geom*, 1988, **28**(3): 361-382.
- [9] K. Corlette, Archimedean superrigidity and hyperbolic geometry. *Ann of Math* (2), 1992, **135**(1): 165-182.
- [10] K. Corlette, Nonabelian Hodge theory, In: *Differential geometry: geometry in mathematical physics and related topics* (Los Angeles, CA, 1990), 125-144. *Proc. Sympos. Pure Math.*, 54, Part 2, Amer. Math. Soc., Providence, RI, 1993.
- [11] J. Daniel, On some characteristic classes of flat bundles in complex geometry. *Ann Inst Fourier (Grenoble)*, 2019, **69**(2019): 729-751.
- [12] G. Daskalopoulos, C. Mese and G. Wilkin, Higgs bundles over cell complexes and representations of finitely presented groups. *Pacific J Math*, 2018, **296**(1): 31-55.
- [13] S. K. Donaldson, Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. *Proc London Math Soc* (3), 1985, **50**(1): 1-26.
- [14] S. K. Donaldson, Twisted harmonic maps and the self-duality equations. *Proc London Math Soc* (3), 1987, **55**(1): 127-131.
- [15] S. K. Donaldson, Boundary value problems for Yang-Mills fields. *J. Geom. Phys*, 1992, **8**(1-4): 89-122.
- [16] J. Dupont, Simplicial de Rham cohomology and characteristic classes of flat bundles. *Topology*, 1976, **15**(3): 233-245.
- [17] J. Eells and J. H. Sampson, Harmonic mappings of Riemannian manifolds. *Amer J Math*, 1964, **86**: 109-160.
- [18] D. Gilbarg and N. S. Trudinger, Elliptic partial differential equations of second order. Reprint of the 1998 edition. *Classics in Mathematics*. Springer-Verlag, Berlin, 2001. xiv+517 pp. ISBN: 3-540-41160-7.
- [19] N. J. Hitchin, The self-duality equations on a Riemann surface. *Proc London Math Soc* (3), 1987, **55**(1): 59-126.
- [20] J. Jost and S. T. Yau, Harmonic maps and group representations, In: *Differential Geometry*, 241-259. *Pitman Monogr Surveys Pure Appl Math.*, 52, Longman Sci Tech., Harlow, 1991.
- [21] J. Jost and K. Zuo, Harmonic maps and $Sl(r, \mathbb{C})$ -representations of fundamental groups of quasiprojective manifolds. *J Algebraic Geom*, 1996, **5**(1): 77-106.
- [22] J. Jost and K. Zuo, Harmonic maps of infinite energy and rigidity results for representations of fundamental groups of quasiprojective varieties. *J. Differential Geom*, 1997, **47**(3): 469-503.
- [23] F. Kamber and P. Tondeur, Characteristic invariants of foliated bundles. *Manuscripta Math*, 1974, **11**: 51-89.
- [24] E. Korman, Characteristic classes of Higgs bundles and Reznikov's theorem. *Manuscripta Math*, 2017, **152**(3-4): 433-442.
- [25] F. Labourie, Existence d'applications harmoniques tordues à valeurs dans les variétés à courbure négative. *Proc Amer Math Soc*, 1991, **111**(3): 877-882.
- [26] J. Y. Li, Hitchin's self-duality equations on complete Riemannian manifolds. *Math Ann*, 1996, **306**(3): 419-428.
- [27] J. Loftin, Affine Hermitian-Einstein metrics. *Asian J Math*, 2009, **13**(1): 101-130.
- [28] M. Lübke and A. Teleman, *The Kobayashi-Hitchin Correspondence*. World Scientific Publishing Co., Inc., River Edge, NJ, 1995. x+254 pp. ISBN: 981-02-2168-1.
- [29] T. Mochizuki, Kobayashi-Hitchin correspondence for tame harmonic bundles and an application. *Astérisque* No. 309(2006), viii+117 pp. ISBN: 978-2-85629-226-6.

- [30] T. Mochizuki, Asymptotic behaviour of tame harmonic bundles and an application to pure twistor D -modules I. Mem. Amer. Math. Soc. 185(2007), no. 869, xii+324 pp.
- [31] T. Mochizuki, Kobayashi-Hitchin correspondence for tame harmonic bundles II. Geom Topol, 2009, **13**(1): 359-455.
- [32] T. Mochizuki, Wild harmonic bundles and wild pure twistor D -modules. Astérisque No. 340(2011), x+607 pp. ISBN: 978-2-85629-332-4
- [33] T. Mochizuki, Kobayashi-Hitchin correspondence for analytically stable bundles. Trans Amer Math Soc, 2020, **373**(1): 551-596.
- [34] M. S. Narasimhan and C. S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface. Ann Math (2), 1965, **82**: 540-567.
- [35] C. P. Pan, C. J. Zhang and X. Zhang, Projectively flat bundles and semi-stable Higgs bundles. arXiv: 1911.03593, 2019.
- [36] A. Reznikov, All regulators of flat bundles are torsion. Ann Math (2), 1995, **141**(2): 373-386.
- [37] J. H. Sampson, Applications of harmonic maps to Kähler geometry, In: Complex differential geometry and nonlinear differential equations(Brunswick, Maine, 1984), 125-134. Contemp. Math., 49, Amer. Math. Soc., Providence, RI, 1986.
- [38] Z. H. Shen, C. J. Zhang and X. Zhang, Flat Higgs bundles over non-compact affine Gauduchon manifolds. arXiv: 1909.12577, 2019.
- [39] C. T. Simpson, Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization. J Amer Math Soc, 1988, **1**(4): 867-918.
- [40] C. T. Simpson, Harmonic bundles on noncompact curves. J Amer Math Soc, 1990, **3**(3): 713-770.
- [41] C. T. Simpson, Higgs bundles and local systems. Inst Hautes Études Sci Publ Math, 1992, No. 75: 5-95.
- [42] Y. T. Siu, The complex-analyticity of harmonic maps and strong rigidity of complex Kähler manifolds. Ann of Math (2), 1980, **112**(2): 73-111.
- [43] K. Uhlenbeck, Connections with L^p bounds on curvature. Comm Math Phys, 1982, **83**(1): 31-42.
- [44] K. Uhlenbeck and S. T. Yau, On the existence of Hermitian-Yang-Mills connections in stable vector bundles. Comm Pure Appl Math, 1986, **39**(S): 257-293.
- [45] Y. Wang and X. Zhang, Twisted holomorphic chains and vortex equations over non-compact Kähler manifolds. J Math Anal Appl, 2011, **373**(1): 179-202.
- [46] S. T. Yau, Some function-theoretic properties of complete Riemannian manifold and their applications to geometry. Indiana Univ Math J, 1976, **25**(7): 659-670.
- [47] X. Zhang, Hermitian-Einstein metrics on holomorphic vector bundles over Hermitian manifolds. J Geom Phys, 2005, **53**(3): 315-335.
- [48] X. Zhang, The limit of the harmonic flow on flat complex vector bundle. arXiv: 2101.07443, 2021.

DI WU, SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI 230026, PEOPLE'S REPUBLIC OF CHINA

Email address: wudi123@mail.ustc.edu.cn

XI ZHANG, SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI 230026, PEOPLE'S REPUBLIC OF CHINA

Email address: mathzx@ustc.edu.cn