

VARIATIONAL BIHAMILTONIAN COHOMOLOGIES AND INTEGRABLE HIERARCHIES II: VIRASORO SYMMETRIES

SI-QI LIU, ZHE WANG, YOUJIN ZHANG

ABSTRACT. We prove that for any tau-symmetric bihamiltonian deformation of the tau-cover of the Principal Hierarchy associated with a semisimple Frobenius manifold, the deformed tau-cover admits an infinite set of Virasoro symmetries.

CONTENTS

1. Introduction	1
2. Super tau-covers of bihamiltonian integrable hierarchies	3
2.1. Bihamiltonian structures on the infinite jet space	3
2.2. Frobenius manifolds and super tau-covers of the Principal Hierarchies	7
2.3. Super extensions of bihamiltonian integrable hierarchies	11
2.4. Deformation of the super tau-covers	18
3. Deformation of Virasoro symmetries: formulation	22
3.1. Variational bihamiltonian cohomologies	22
3.2. Virasoro symmetries of the Principal Hierarchies	24
3.3. Formulation of the deformation problem	26
3.4. Example: one-dimensional Frobenius manifold	29
4. Deformation of Virasoro symmetries: existence and uniqueness	32
4.1. Locality conditions	32
4.2. Closedness conditions	38
4.3. Vanishing of the genus one obstruction	42
4.4. Lifting to the tau-covers	45
5. Conclusion	52
References	54

1. INTRODUCTION

This is the second one of the series of papers devoted to the study of the deformations of the Virasoro symmetries of bihamiltonian integrable hierarchies. In the first one [20], we developed a cohomology theory on the space of differential forms of the infinite jet space

Date: July 17, 2022.

Key words and phrases. Virasoro symmetries, Super tau-covers, Variational Bihamiltonian Cohomologies, Integrable Hierarchies.

of a super manifold for a given bihamiltonian structure of the hydrodynamic type, and we call such a cohomology theory the *variational bihamiltonian cohomology*. It can be viewed as a generalization of the bihamiltonian cohomology introduced in [12], and it provides a suitable tool for us to study deformations of Virasoro symmetries of bihamiltonian integrable hierarchies.

The purpose of the present paper is to prove the following theorem.

Main Theorem. *For the Principal Hierarchy associated with a semisimple Frobenius manifold and any of its tau-symmetric bihamiltonian deformations, there exists a unique deformation of its Virasoro symmetries such that they are symmetries of the deformed integrable hierarchy. Moreover, the action of the Virasoro symmetries on the tau-function Z of the deformed integrable hierarchy can be represented in the form*

$$(1.1) \quad \frac{\partial Z}{\partial s_m} = L_m Z + O_m Z, \quad m \geq -1,$$

where L_m are the Virasoro operators constructed in [11] and O_m are some differential polynomials, and the flows $\frac{\partial}{\partial s_m}$ satisfy the Virasoro commutation relations

$$\left[\frac{\partial}{\partial s_k}, \frac{\partial}{\partial s_l} \right] = (l - k) \frac{\partial}{\partial s_{k+l}}, \quad k, l \geq -1.$$

Let us briefly explain the basic idea for proving this theorem. Consider the following system of evolutionary PDEs with time variable t and spacial variable x :

$$(1.2) \quad \frac{\partial u^i}{\partial t} = A_j^i(u) u_x^j + \varepsilon (B_j^i(u) u_{xx}^j + C_{jk}^i(u) u_x^j u_x^k) + \cdots, \quad i = 1, \dots, n,$$

We assume that this system is bihamiltonian with respect to the bihamiltonian structure (P_0, P_1) whose leading term is semisimple. We can associate with it a super extension by introducing odd unknown functions θ_i for $i = 1, \dots, n$, and by adding odd flows $\frac{\partial}{\partial \tau_0}$ and $\frac{\partial}{\partial \tau_1}$ which correspond respectively to the Hamiltonian structure P_0 and P_1 (see [19] and Sect. 2.3 given below for details). Thus the super extension of (1.2) consists of the flows $\frac{\partial}{\partial t}$, $\frac{\partial}{\partial \tau_0}$ and $\frac{\partial}{\partial \tau_1}$ for the unknown functions u^i and θ_i , and the fact that the system (1.2) is bihamiltonian with respect to (P_0, P_1) is equivalent to the following commutation relations:

$$\left[\frac{\partial}{\partial t}, \frac{\partial}{\partial \tau_i} \right] = 0, \quad \left[\frac{\partial}{\partial \tau_i}, \frac{\partial}{\partial \tau_j} \right] = 0, \quad i, j = 0, 1.$$

Example 1.1. *Consider the following Korteweg-de Vries (KdV) equation:*

$$\frac{\partial u}{\partial t} = uu_x + \frac{\varepsilon^2}{12} u_{xxx}.$$

It admits a bihamiltonian structure given by the following Poisson brackets:

$$\begin{aligned} \{u(x), u(y)\}_0 &= \delta'(x - y), \\ \{u(x), u(y)\}_1 &= u(x) \delta'(x - y) + \frac{u_x}{2} \delta(x - y) + \frac{\varepsilon^2}{8} \delta'''(x - y). \end{aligned}$$

We introduce an odd unknown function θ , and construct the following super extension of the KdV equation:

$$(1.3) \quad \frac{\partial u}{\partial t} = uu_x + \frac{\varepsilon^2}{12}u_{xxx}, \quad \frac{\partial \theta}{\partial t} = u\theta_x + \frac{\varepsilon^2}{12}\theta_{xxx},$$

$$(1.4) \quad \frac{\partial u}{\partial \tau_0} = \theta_x, \quad \frac{\partial u}{\partial \tau_1} = v\theta_x + \frac{1}{2}v_x\theta + \frac{\varepsilon^2}{8}\theta_{xxx},$$

$$(1.5) \quad \frac{\partial \theta}{\partial \tau_0} = 0, \quad \frac{\partial \theta}{\partial \tau_1} = \frac{1}{2}\theta\theta_x.$$

It is easy to check directly that the flows in the extended system mutually commute.

Remark 1.1. The flow (1.3) also appeared in [1].

According to the theory of bihamiltonian cohomology given in [7], we know that if there is another system of evolutionary PDEs given by the flow $\frac{\partial}{\partial t}$ satisfying the commutation relation

$$(1.6) \quad \left[\frac{\partial}{\partial t}, \frac{\partial}{\partial \tau_0} \right] = \left[\frac{\partial}{\partial t}, \frac{\partial}{\partial \tau_1} \right] = 0,$$

then it gives a symmetry of the system (1.2), i.e.,

$$\left[\frac{\partial}{\partial t}, \frac{\partial}{\partial t} \right] = 0.$$

We note that the above results is for a flow $\frac{\partial}{\partial t}$ given by differential polynomials. In order to use the above results to consider symmetries of more general forms such as Virasoro symmetries, we introduce the notion of super tau-cover and develop the theory of variational bihamiltonian cohomology [20]. Then we are able to consider the commutation relation between the Virasoro symmetries $\frac{\partial}{\partial s_m}$ and the odd flows $\frac{\partial}{\partial \tau_0}$ and $\frac{\partial}{\partial \tau_1}$. By applying the results of variational bihamiltonian cohomology proved in [20], we manage to prove the main theorem.

This paper is organized as follows. In Sect. 2, we construct the super tau-cover for a given tau-symmetric bihamiltonian deformation of the Principal Hierarchy associated with a semisimple Frobenius manifold. This construction builds a bridge which relates the Virasoro symmetries to the bihamiltonian structures. In Sect. 3, we explain how the deformation problem of the Virasoro symmetries can be solved via the theory of the variational bihamiltonian cohomology. Sect. 4 is devoted to the proof of the main theorem. Finally some concluding remarks are made in Sect. 5.

2. SUPER TAU-COVERS OF BIHAMILTONIAN INTEGRABLE HIERARCHIES

2.1. Bihamiltonian structures on the infinite jet space. Let us start by recalling the basic constructions of bihamiltonian structures as local functionals on infinite jet spaces. One may refer to [17] for a detailed introduction to this topic.

Let M be a smooth manifold of dimension n and \hat{M} be the super manifold of dimension $(n|n)$ obtained by reversing the parity of the fibers of the cotangent bundle of M . In another word, if we choose a local canonical coordinate system $(u^1, \dots, u^n; \theta_1, \dots, \theta_n)$

on T^*M , then \hat{M} can be described locally by the same chart while regarding the fiber coordinates as odd variables:

$$\theta_i \theta_j + \theta_j \theta_i = 0, \quad i, j = 1, \dots, n.$$

We call that the odd coordinates θ_i are dual to u^i . The transition functions between two local trivializations $(u^1, \dots, u^n; \theta_1, \dots, \theta_n)$ and $(w^1, \dots, w^n; \phi_1, \dots, \phi_n)$ are given by the same formula as those of the cotangent bundle:

$$\phi_\alpha = \frac{\partial u^\beta}{\partial w^\alpha} \theta_\beta, \quad \alpha = 1, \dots, n.$$

Here and henceforth, summation of the repeated upper and lower Greek indices is assumed.

Denote by $J^\infty(\hat{M})$ the infinite jet bundle of \hat{M} . It is a fiber bundle over \hat{M} with fiber \mathbb{R}^∞ . If we choose a local chart $(u^1, \dots, u^n; \theta_1, \dots, \theta_n)$ of \hat{M} , a trivialization can be realized by choosing the fiber coordinate being $(u^{\alpha,s}; \theta_\alpha^s)$ for $\alpha = 1, \dots, n$ and $s \geq 1$. The transition functions between different charts are given by the chain rule

$$\begin{aligned} w^{\alpha,1} &= \frac{\partial w^\alpha}{\partial u^\beta} u^{\beta,1}, \quad w^{\alpha,2} = \frac{\partial w^\alpha}{\partial u^\beta} u^{\beta,2} + \frac{\partial^2 w^\alpha}{\partial u^\beta \partial u^\gamma} u^{\beta,1} u^{\gamma,1}, \quad \dots \\ \phi_\alpha^1 &= \frac{\partial u^\beta}{\partial w^\alpha} \theta_\beta^1 + \frac{\partial^2 u^\beta}{\partial w^\gamma \partial w^\alpha} \frac{\partial w^\gamma}{\partial u^\lambda} u^{\lambda,1} \theta_\beta, \quad \dots \end{aligned}$$

Denote by $\hat{\mathcal{A}}$ the ring of differential polynomials, locally it is given by

$$\hat{\mathcal{A}} = C^\infty(u)[[u^{i,s+1}, \theta_i^s \mid i = 1, \dots, n; s \geq 0]].$$

It is graded with respect to the super degree \deg_θ defined by

$$\deg_\theta u^{\alpha,s} = 0, \quad \deg_\theta \theta_\alpha^s = 1; \quad \alpha = 1, \dots, n, \quad s \geq 0.$$

Here and henceforth we use the notation $u^{\alpha,0} = u^\alpha$, $\theta_\alpha^0 = \theta_\alpha$. The set of homogeneous elements with super degree p is denoted by $\hat{\mathcal{A}}^p$.

Introduce a global vector field ∂_x on $J^\infty(\hat{M})$ which is locally described by

$$\partial_x = \sum_{s \geq 0} u^{\alpha,s+1} \frac{\partial}{\partial u^{\alpha,s}} + \theta_\alpha^{s+1} \frac{\partial}{\partial \theta_\alpha^s}.$$

Therefore we see that $u^{\alpha,s} = \partial_x^s u^\alpha$ and $\theta_\alpha^s = \partial_x^s \theta_\alpha$. Hence we can grade the ring $\hat{\mathcal{A}}$ with respected to the differential degree \deg_{∂_x} defined by

$$\deg_{\partial_x} u^{\alpha,s} = s, \quad \deg_{\partial_x} \theta_\alpha^s = s; \quad \alpha = 1, \dots, n, \quad s \geq 0.$$

We use the notation $\hat{\mathcal{A}}_d$ to denote the set of homogeneous elements with differential degree d , and $\hat{\mathcal{A}}_d^p = \hat{\mathcal{A}}^p \cap \hat{\mathcal{A}}_d$.

Using the vector field ∂_x , one can construct the space $\hat{\mathcal{F}}$ of local functionals via the quotient $\hat{\mathcal{F}} := \hat{\mathcal{A}} / \partial_x \hat{\mathcal{A}}$. Since the vector field ∂_x is homogeneous with respect to both the super degree and the differential degree, the quotient space $\hat{\mathcal{F}}$ admits natural gradations induced from $\hat{\mathcal{A}}$ and we will use the notation $\hat{\mathcal{F}}^p$, $\hat{\mathcal{F}}_d$ and $\hat{\mathcal{F}}_d^p$ to denote the subspace of homogeneous elements. For any element $f \in \hat{\mathcal{A}}$, we will use $\int f \in \hat{\mathcal{F}}$ to denote its image of the natural projection $\hat{\mathcal{A}} \rightarrow \hat{\mathcal{F}}$.

For a differential polynomial $f \in \hat{\mathcal{A}}$, one may define the variational derivatives by

$$\frac{\delta f}{\delta u^\alpha} = \sum_{s \geq 0} (-\partial_x)^s \frac{\partial f}{\partial u^{\alpha, s}}, \quad \frac{\delta f}{\delta \theta_\alpha} = \sum_{s \geq 0} (-\partial_x)^s \frac{\partial f}{\partial \theta_\alpha^s}.$$

It is easy to verify that the variational derivatives annihilate the elements in $\partial_x \hat{\mathcal{A}}$, hence the variational derivatives are also well-defined on the quotient space $\hat{\mathcal{F}}$. For any $F \in \hat{\mathcal{F}}$, we have

$$\frac{\delta F}{\delta u^\alpha} = \sum_{s \geq 0} (-\partial_x)^s \frac{\partial f}{\partial u^{\alpha, s}}, \quad \frac{\delta F}{\delta \theta_\alpha} = \sum_{s \geq 0} (-\partial_x)^s \frac{\partial f}{\partial \theta_\alpha^s},$$

with $f \in \hat{\mathcal{A}}$ being an arbitrary lift of F such that $F = \int f$. With the help of the notion of the variational derivatives, one can define the so-called Schouten-Nijenhuis bracket, which is a bilinear map $[-, -] : \hat{\mathcal{F}} \times \hat{\mathcal{F}} \rightarrow \hat{\mathcal{F}}$ given by

$$[P, Q] = \int \left(\frac{\delta P}{\delta \theta_\alpha} \frac{\delta Q}{\delta u^\alpha} + (-1)^p \frac{\delta P}{\delta u^\alpha} \frac{\delta Q}{\delta \theta_\alpha} \right), \quad P \in \hat{\mathcal{F}}^p, Q \in \hat{\mathcal{F}}^q.$$

This bracket satisfies the graded commuting relation

$$[P, Q] = (-q)^{pq} [Q, P], \quad P \in \hat{\mathcal{F}}^p, Q \in \hat{\mathcal{F}}^q,$$

and the graded Jacobi identity:

$$(-1)^{rp} [[P, Q], R] + (-1)^{pq} [[Q, R], P] + (-1)^{qr} [[R, P], Q] = 0, \quad P \in \hat{\mathcal{F}}^p, Q \in \hat{\mathcal{F}}^q, R \in \hat{\mathcal{F}}^r.$$

Any local functional $P \in \hat{\mathcal{F}}^p$ gives rise to a graded derivation $D_P \in \text{Der}(\hat{\mathcal{A}})^{p-1}$ by

$$(2.1) \quad D_P = \sum_{s \geq 0} \partial_x^s \left(\frac{\delta P}{\delta \theta_\alpha} \right) \frac{\partial}{\partial u^{\alpha, s}} + (-1)^p \partial_x^s \left(\frac{\delta P}{\delta u^\alpha} \right) \frac{\partial}{\partial \theta_\alpha^s}.$$

Here the space $\text{Der}(\hat{\mathcal{A}})^p$ ($p \in \mathbb{Z}$) is the space of linear maps $D : \hat{\mathcal{A}}^q \rightarrow \hat{\mathcal{A}}^{q+p}$ satisfying the graded Leibniz rule:

$$D(fg) = D(f)g + (-1)^{kp} f D(g), \quad f \in \hat{\mathcal{A}}^k, g \in \hat{\mathcal{A}}.$$

and we denote $\text{Der}(\hat{\mathcal{A}}) = \bigoplus_{p \in \mathbb{Z}} \text{Der}(\hat{\mathcal{A}})^p$, which is a graded Lie algebra with the graded commutator

$$[D_1, D_2] = D_1 \circ D_2 - (-1)^{kl} D_2 \circ D_1, \quad D_1 \in \text{Der}(\hat{\mathcal{A}})^k, D_2 \in \text{Der}(\hat{\mathcal{A}})^l.$$

$\text{Der}(\hat{\mathcal{A}})$ is also graded by the differential degree

$$\text{Der}(\hat{\mathcal{A}})_d = \{D \in \text{Der}(\hat{\mathcal{A}}) \mid D(\hat{\mathcal{A}}_k) \subseteq \hat{\mathcal{A}}_{k+d}\}.$$

For $P \in \hat{\mathcal{F}}^p$ and $Q \in \hat{\mathcal{F}}^q$, the derivation defined in (2.1) satisfies the identity

$$(2.2) \quad (-1)^{p-1} D_{[P, Q]} = [D_P, D_Q].$$

The following identities are useful:

$$(2.3) \quad \frac{\delta}{\delta u^\alpha} [P, Q] = D_P \left(\frac{\delta Q}{\delta u^\alpha} \right) + (-1)^{pq} D_Q \left(\frac{\delta P}{\delta u^\alpha} \right);$$

$$(2.4) \quad (-1)^{p-1} \frac{\delta}{\delta \theta_\alpha} [P, Q] = D_P \left(\frac{\delta Q}{\delta \theta_\alpha} \right) - (-1)^{(p-1)(q-1)} D_Q \left(\frac{\delta P}{\delta \theta_\alpha} \right).$$

A Hamiltonian structure is defined as a local functional $P \in \hat{\mathcal{F}}^2$ such that $[P, P] = 0$. We can associate a matrix valued differential operator $\mathcal{P} = (\mathcal{P}^{\alpha\beta})$ with $\mathcal{P}^{\alpha\beta} = \sum_{s \geq 0} \mathcal{P}_s^{\alpha\beta} \partial_x^s$ to a bivector $P \in \hat{\mathcal{F}}^2$. Here $\mathcal{P}_s^{\alpha\beta} \in \hat{\mathcal{A}}$ is defined by

$$\frac{\delta P}{\delta \theta_\alpha} = \sum_{s \geq 0} \mathcal{P}_s^{\alpha\beta} \theta_\beta^s, \quad \alpha = 1, \dots, n.$$

If P is a Hamiltonian structure, then we call \mathcal{P} the Hamiltonian operator of P .

Theorem 2.1 ([10]). *Let $P \in \hat{\mathcal{F}}_1^2$ whose associated differential operator is given by*

$$\mathcal{P}^{\alpha\beta} = g^{\alpha\beta}(u) \partial_x + \Gamma_\gamma^{\alpha\beta}(u) u^{\gamma,1}, \quad \det(g^{\alpha\beta}) \neq 0.$$

Then P is a Hamiltonian structure if and only if $g = (g_{\alpha\beta}) = (g^{\alpha\beta})^{-1}$ defines a flat (pseudo-)Riemannian metric on M and the Christoffel symbols of the Levi-Civita connection of g are given by $\Gamma_{\alpha\beta}^\gamma = -g_{\alpha\lambda} \Gamma_\beta^{\lambda\gamma}$.

The Hamiltonian structure P satisfying the conditions of the above theorem is called of hydrodynamic type. It follows from the above theorem that for each Hamiltonian structure P of hydrodynamic type, there exists a local coordinate system $(v^\alpha; \sigma_\alpha)$ on \hat{M} such that

$$P = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1,$$

where $\eta^{\alpha\beta}$ is a constant non-degenerate matrix. The coordinates v^α and σ_α are called flat coordinates of P .

A bihamiltonian structure (P_0, P_1) is a pair of Hamiltonian structure satisfying an additional compatibility condition $[P_0, P_1] = 0$. Assume that the bihamiltonian structure is of hydrodynamic type, then according to Theorem 2.1, we have two flat contravariant metrics $g_0^{\alpha\beta}$ and $g_1^{\alpha\beta}$. We say that this bihamiltonian structure is semisimple if the roots of the characteristic equation

$$\det(g_1^{\alpha\beta} - \lambda g_0^{\alpha\beta}) = 0$$

are distinct and not constant. In this case, the roots $\lambda^1(u), \dots, \lambda^n(u)$ can serve as local coordinates of M and it is called the canonical coordinate of the semisimple bihamiltonian structure. It is proved in [14] that in canonical coordinate,

$$P_0 = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} f^i(\lambda) \theta_i \theta_j^1 + A^{ij} \theta_i \theta_j), \quad P_1 = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} g^i(\lambda) \theta_i \theta_j^1 + B^{ij} \theta_i \theta_j),$$

where f^i are non-vanishing functions, $g^i = \lambda^i f^i$ and the functions A^{ij} and B^{ij} are given by

$$(2.5) \quad A^{ij} = \frac{1}{2} \left(\frac{f^i}{f^j} \frac{\partial f^j}{\partial \lambda^i} \lambda^{j,1} - \frac{f^j}{f^i} \frac{\partial f^i}{\partial \lambda^j} \lambda^{i,1} \right), \quad B^{ij} = \frac{1}{2} \left(\frac{g^i}{f^j} \frac{\partial f^j}{\partial \lambda^i} \lambda^{j,1} - \frac{g^j}{f^i} \frac{\partial f^i}{\partial \lambda^j} \lambda^{i,1} \right).$$

Here by abusing the notation, we still use θ_i to denote the fiber coordinates of \hat{M} dual to λ^i . We also call λ^i and θ_i the canonical coordinates of (P_0, P_1) .

From now on, for a semisimple hydrodynamic bihamiltonian structure (P_0, P_1) , we will use $(v^\alpha; \sigma_\alpha)$ to denote the flat coordinate of P_0 such that

$$P_0 = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1, \quad P_1 = \frac{1}{2} \int g^{\alpha\beta}(v) \sigma_\alpha \sigma_\beta^1 + \Gamma_\gamma^{\alpha\beta}(v) v^{\gamma,1} \sigma_\alpha \sigma_\beta,$$

and we will use $(u^i; \theta_i)$ to denote the canonical coordinate for (P_0, P_1) such that

$$(2.6) \quad P_0 = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} f^i(u) \theta_i \theta_j^1 + A^{ij} \theta_i \theta_j), \quad P_1 = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} u^i f^i(u) \theta_i \theta_j^1 + B^{ij} \theta_i \theta_j).$$

Note that we will not adopt Einstein summation for Latin indices.

In terms the notations introduced above, a system of evolutionary PDEs

$$\frac{\partial u^\alpha}{\partial t} = X^\alpha, \quad X^\alpha \in \hat{\mathcal{A}}^0$$

can be represented as a local functional $X = \int X^\alpha \theta_\alpha$, and it is said to be bihamiltonian if there exists a bihamiltonian structure (P_0, P_1) and two Hamiltonians $G, H \in \hat{\mathcal{F}}^0$ such that

$$X = -[G, P_0] = -[H, P_1].$$

Example 2.1. *The KdV equation*

$$(2.7) \quad \frac{\partial u}{\partial t} = uu_x + \frac{\varepsilon^2}{12} u_{xxx}$$

can be represented by $X = \int (uu_x + \frac{\varepsilon^2}{12} u_{xxx}) \theta$. Its bihamiltonian structure is given by

$$P_0 = \frac{1}{2} \int \theta \theta_x, \quad P_1 = \frac{1}{2} \int u \theta \theta^1 + \frac{\varepsilon^2}{8} \theta \theta^3.$$

The two Hamiltonians with respect to the bihamiltonian structure are given by

$$X = - \left[\int \frac{u^3}{6} - \frac{\varepsilon^2}{24} u_x^2, P_0 \right] = - \left[\int \frac{u^2}{3}, P_1 \right].$$

2.2. Frobenius manifolds and super tau-covers of the Principal Hierarchies. In this section, we will recall some basic facts of Frobenius manifolds and the construction of the associated Principal Hierarchies following the work [4, 5, 6, 12]. Then we recall the construction of the super tau-covers of the Principal Hierarchies given in [19].

The notion of Frobenius manifold is a geometric description of genus zero 2D topological field theories. An n -dimensional Frobenius manifold M can be locally described by a solution $F(v^1, \dots, v^n)$ of the following Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) associativity equation [2, 24]:

$$(2.8) \quad \partial_\alpha \partial_\beta \partial_\lambda F \eta^{\lambda\mu} \partial_\mu \partial_\gamma \partial_\delta F = \partial_\delta \partial_\beta \partial_\lambda F \eta^{\lambda\mu} \partial_\mu \partial_\gamma \partial_\alpha F.$$

Here $\partial_\alpha = \frac{\partial}{\partial v^\alpha}$ and we require $(\eta_{\alpha\beta}) := (\partial_1 \partial_\alpha \partial_\beta F)$ is a constant non-degenerate matrix with inverse $(\eta^{\alpha\beta})$. The function $F(v^1, \dots, v^n)$ is called the potential of M and it defines a Frobenius algebra structure on TM :

$$\langle \partial_\alpha, \partial_\beta \rangle = \eta_{\alpha\beta}, \quad \partial_\alpha \cdot \partial_\beta = c_{\alpha\beta}^\gamma \partial_\gamma,$$

where the functions $c_{\alpha\beta}^\gamma$ are defined by

$$c_{\alpha\beta}^\gamma = \eta^{\gamma\lambda} c_{\lambda\alpha\beta}, \quad c_{\lambda\alpha\beta} = \partial_\lambda \partial_\alpha \partial_\beta F.$$

The potential F is required to be quasi-homogeneous in the sense that there exists a vector field

$$E = \sum_{\alpha=1}^n \left(\left(1 - \frac{d}{2} - \mu_\alpha \right) v^\alpha + r_\alpha \right) \partial_\alpha,$$

called the Euler vector field, such that

$$E(F) = (3-d)F + \frac{1}{2} A_{\alpha\beta} v^\alpha v^\beta + B_\alpha v^\alpha + C.$$

Here the diagonal matrix $\mu = \text{diag}(\mu_1, \dots, \mu_n)$ is part of the isomonodromy data of M and it is assumed that $\mu_1 = -d/2$ and $r_1 = 0$. The matrix μ satisfies the property

$$(2.9) \quad (\mu_\alpha + \mu_\beta) \eta_{\alpha\beta} = 0, \quad \forall \alpha, \beta.$$

An important property of Frobenius manifolds is that the affine connection

$$\tilde{\nabla}_X Y = \nabla_X Y + z X \cdot Y, \quad \forall X, Y \in \Gamma(TM), \quad z \in \mathbb{C}$$

is flat for arbitrary z , here ∇ is the Levi-Civita connection of the flat metric $(\eta_{\alpha\beta})$. It can be extended to be a flat connection on $M \times \mathbb{C}^*$ by viewing z as the coordinate on \mathbb{C}^* and defining

$$\tilde{\nabla}_{\partial_z} X = \partial_z X + E \cdot X - \frac{1}{z} \mu X, \quad \tilde{\nabla}_{\partial_z} \partial_z = \tilde{\nabla}_X \partial_z = 0.$$

The connection $\tilde{\nabla}$ is called the deformed flat connection or the Dubrovin connection. For such a flat connection, one can find a system of flat coordinates of the form

$$(\tilde{v}^1(v, z), \dots, \tilde{v}^n(v, z)) = (h_1(v, z), \dots, h_n(v, z)) z^\mu z^R.$$

Here R is a constant matrix. The constant matrices η , μ and R form the monodromy data of M at $z = 0$. The matrix R can be decomposed into a finite sum $R = R_1 + \dots + R_m$, and they satisfy the condition

$$(2.10) \quad [\mu, R_k] = k R_k, \quad \eta_{\alpha\gamma} (R_k)^\gamma_\beta = (-1)^{k+1} \eta_{\beta\gamma} (R_k)^\gamma_\alpha.$$

The functions $h_\alpha(v, z)$ are analytic at $z = 0$ and has the expansion $h_\alpha(v, z) = \sum_{p \geq 0} h_{\alpha,p}(v) z^p$. The coefficients $h_{\alpha,p}$ satisfy the following recursion condition:

$$h_{\alpha,0} = \eta_{\alpha\beta} v^\beta, \quad \partial_\beta \partial_\gamma h_{\alpha,p+1} = c_{\beta\gamma}^\lambda \partial_\lambda h_{\alpha,p}, \quad p \geq 0.$$

They also satisfy the following quasi-homogeneous condition

$$E(\partial_\beta h_{\alpha,p}) = (p + \mu_\alpha + \mu_\beta) \partial_\beta h_{\alpha,p} + \sum_{k=1}^p (R_k)^\gamma_\alpha \partial_\beta h_{\gamma,p-k},$$

and the following normalization condition

$$\langle \nabla h_\alpha(v, z), \nabla h_\beta(v, -z) \rangle = \eta_{\alpha\beta}.$$

A choice of the functions $h_{\alpha,p}$ satisfying all the conditions above is called a calibration of M and a Frobenius manifold M is called calibrated if such a choice is fixed. In what follows, we assume that all the Frobenius manifolds are calibrated.

The Principal Hierarchy associated with a Frobenius manifold M is a bihamiltonian integrable hierarchy of hydrodynamic type. Denote by σ_α the odd variables dual to the flat coordinates v^α , then the Principal Hierarchy can be described by the local functionals $X_{\alpha,p} \in \hat{\mathcal{F}}^1$ of the form

$$(2.11) \quad X_{\alpha,p} = \int \eta^{\lambda\gamma} \partial_x (\partial_\gamma h_{\alpha,p+1}) \sigma_\lambda, \quad \alpha = 1, \dots, n, \quad p \geq 0,$$

or equivalently, we can represent the hierarchy as follows:

$$\frac{\partial v^\lambda}{\partial t^{\alpha,p}} = \eta^{\lambda\gamma} \partial_x (\partial_\gamma h_{\alpha,p+1}).$$

Define two local functionals

$$(2.12) \quad P_0 = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1, \quad P_1 = \frac{1}{2} \int g^{\alpha\beta} \sigma_\alpha \sigma_\beta^1 + \Gamma_\gamma^{\alpha\beta} v^{\gamma,1} \sigma_\alpha \sigma_\beta,$$

where the functions $g^{\alpha\beta}$ and $\Gamma_\gamma^{\alpha\beta}$ are given by

$$g^{\alpha\beta} = E^\varepsilon c_\varepsilon^{\alpha\beta}, \quad \Gamma_\gamma^{\alpha\beta} = \left(\frac{1}{2} - \mu_\beta \right) c_\gamma^{\alpha\beta}; \quad c_\gamma^{\alpha\beta} = \eta^{\alpha\lambda} c_{\lambda\gamma}^\beta.$$

Then we have the following theorem.

Theorem 2.2 ([5]). *Let M be a Frobenius manifold, then*

- (1) *The local functionals P_0, P_1 defined in (2.12) form a bihamiltonian structure which is exact in the sense that*

$$P_0 = [Z, P_1], \quad Z = \int \sigma_1.$$

- (2) *The Principal Hierarchy $X_{\alpha,p}$ associated with M is bihamiltonian with respect to the bihamiltonian structure (P_0, P_1) and*

$$X_{\alpha,p} = -[H_{\alpha,p}, P_0], \quad H_{\alpha,p} = \int h_{\alpha,p+1}.$$

- (3) *The following bihamiltonian recursion relation holds true:*

$$[H_{\alpha,p-1}, P_1] = \left(p + \frac{1}{2} + \mu_\alpha \right) [H_{\alpha,p}, P_0] + \sum_{k=1}^p (R_k)_\alpha^\gamma [H_{\gamma,p-k}, P_0], \quad p \geq 0.$$

Another important property satisfied by the Principal Hierarchy is that it is tau-symmetric. Let functions $\Omega_{\alpha,p;\beta,q}$ for $\alpha, \beta = 1, \dots, n$ and $p, q \geq 0$ be defined by the generating function

$$\sum_{p \geq 0, q \geq 0} \Omega_{\alpha,p;\beta,q}(v) z_1^p z_2^q = \frac{\langle \nabla h_\alpha(v, z_1), \nabla h_\beta(v, z_2) \rangle - \eta_{\alpha\beta}}{z_1 + z_2}.$$

They have the following properties [5]:

$$\begin{aligned} \Omega_{\alpha,p;1,0} &= h_{\alpha,p}, & \Omega_{\alpha,p;\beta,0} &= \partial_\beta h_{\alpha,p+1}; \\ \Omega_{\alpha,p;\beta,q} &= \Omega_{\beta,q;\alpha,p}, & \frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial t^{\lambda,k}} &= \frac{\partial \Omega_{\lambda,k;\beta,q}}{\partial t^{\alpha,p}}. \end{aligned}$$

It follows from these identities that one can extend the Principal Hierarchy by introducing another family of unknown functions $f_{\alpha,p}$ satisfying the following equations:

$$(2.13) \quad \frac{\partial f_{\alpha,p}}{\partial t^{\beta,q}} = \Omega_{\alpha,p;\beta,q}, \quad \frac{\partial v^\alpha}{\partial t^{\beta,q}} = \eta^{\alpha\lambda} \partial_x \Omega_{\lambda,0;\beta,q}.$$

The system (2.13) is called the tau-cover of the Principal Hierarchy.

In order to study the relations between the bihamiltonian structures and the Virasoro symmetries, the notion of super tau-covers was introduced in [19]. Let us briefly recall the construction of the super tau-covers since it provides the main motivation of our work presented in the next subsection. We first introduce a family of odd unknown functions $\sigma_{\alpha,k}^s$ for $s, k \geq 0$ with $\sigma_{\alpha,0}^s = \sigma_\alpha^s$. In what follows we will also use $\sigma_{\alpha,k}$ to denote $\sigma_{\alpha,k}^0$. We extend the action of ∂_x to include these new odd variables as follows:

$$\partial_x = \sum_{s \geq 0} v^{\alpha,s+1} \frac{\partial}{\partial v^{\alpha,s}} + \sum_{k,s \geq 0} \sigma_{\alpha,k}^{s+1} \frac{\partial}{\partial \sigma_{\alpha,k}^s}.$$

These odd variables are required to satisfy the following bihamiltonian recursion relation:

$$(2.14) \quad \eta^{\alpha\beta} \sigma_{\beta,k+1}^1 = g^{\alpha\beta} \sigma_{\beta,k}^1 + \Gamma_\gamma^{\alpha\beta} v^{\gamma,1} \sigma_{\beta,k}, \quad \alpha = 1, \dots, n; k \geq 0.$$

We also introduce a family of odd flows $\frac{\partial}{\partial \tau_m}$ for $m \geq 0$. The first two odd flows are determined by the bihamiltonian structure (P_0, P_1) :

$$\frac{\partial v^\alpha}{\partial \tau_i} = \frac{\delta P_i}{\delta \sigma_\alpha}, \quad \frac{\partial \sigma_\alpha}{\partial \tau_i} = \frac{\delta P_i}{\delta v^\alpha}, \quad i = 0, 1.$$

Note that $\frac{\partial}{\partial \tau_i} = D_{P_i}$ for $i = 0, 1$, where D_{P_i} is defined by (2.1). The actions of $\frac{\partial}{\partial \tau_i}$ can be extended to all the other odd variables $\sigma_{\alpha,k}$ such that the flows $\frac{\partial}{\partial \tau_i}$ are compatible with the recursion relations (2.14). Furthermore we can define infinitely many odd flows $\frac{\partial}{\partial \tau_m}$ for $m \geq 2$ which can be viewed as flows corresponding to certain non-local Hamiltonian structures.

We have the following theorem.

Theorem 2.3 ([19]). *We have the following mutually commuting flows associated with any given Frobenius manifold M :*

$$\begin{aligned} \frac{\partial v^\alpha}{\partial t^{\beta,p}} &= \eta^{\alpha\gamma} (\partial_\lambda \partial_\gamma h_{\beta,p+1}) v^{\lambda,1}, & \frac{\partial \sigma_{\alpha,k}}{\partial t^{\beta,p}} &= \eta^{\gamma\varepsilon} (\partial_\alpha \partial_\varepsilon h_{\beta,p+1}) \sigma_{\gamma,k}^1, \\ \frac{\partial v^\alpha}{\partial \tau_m} &= \eta^{\alpha\beta} \sigma_{\beta,m}^1, & \frac{\partial \sigma_{\alpha,k}}{\partial \tau_m} &= -\frac{\partial \sigma_{\alpha,m}}{\partial \tau_k} = \Gamma_\alpha^{\gamma\beta} \sum_{i=0}^{m-k-1} \sigma_{\beta,k+i} \sigma_{\gamma,m-i-1}^1, \quad 0 \leq k \leq m. \end{aligned}$$

where $\alpha, \beta = 1, \dots, n$, and $m, p \geq 0$. These flows are well-defined in the sense that they are compatible with the recursion relations (2.14).

The system described in Theorem 2.3 is a super extension of the Principal Hierarchy, since the reduction obtained by setting all the odd variables to be zero yields the original Principal Hierarchy. The super extension of the tau-cover (2.13) can be constructed by introducing another family of odd variables $\Phi_{\alpha,p}^m$ for $p, m \geq 0$ and we call it the super tau-cover of the Principal Hierarchy. It is given in the following theorem.

Theorem 2.4 ([19]). *The mutually commuting flows*

$$\begin{aligned}\frac{\partial f_{\alpha,p}}{\partial t^{\beta,q}} &= \Omega_{\alpha,p;\beta,q}, \\ \frac{\partial f_{\alpha,p}}{\partial \tau_m} &= \Phi_{\alpha,p}^m, \\ \frac{\partial \Phi_{\alpha,p}^m}{\partial t^{\beta,q}} &= \frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial \tau_m}, \\ \frac{\partial \Phi_{\alpha,p}^m}{\partial \tau_k} &= \Delta_{\alpha,p}^{k,m},\end{aligned}$$

together with the ones presented in Theorem 2.3, give the super tau-cover of the Principal Hierarchy associated with M , where $\Delta_{\alpha,p}^{k,m}$ are defined by the formula

$$\Delta_{\alpha,p}^{k,m} = -\Delta_{\alpha,p}^{m,k} = \eta^{\gamma\lambda} \partial_\lambda h_{\alpha,p} \Gamma_\gamma^{\delta\mu} \left(\sum_{i=0}^{k-m-1} \sigma_{\mu,m+i} \sigma_{\delta,k-i-1}^1 \right), \quad k \geq m.$$

It was shown in [19] that the odd variables $\Phi_{\alpha,p}^m$ satisfy the recursion relations

$$-\left(\frac{2p-1}{2} + \mu_\alpha\right) \Phi_{\alpha,p}^m = \left(\frac{1}{2} + \mu_\lambda\right) \eta^{\lambda\varepsilon} (\partial_\lambda h_{\alpha,p}) \sigma_{\varepsilon,m} + \sum_{k=1}^p (R_k)_\alpha^\xi \Phi_{\xi,p-k}^m - \Phi_{\alpha,p-1}^{m+1}$$

with the initial condition $\Phi_{\alpha,0}^m = \sigma_{\alpha,m}$. So when the diagonal matrix μ of a Frobenius manifold M satisfies the condition $\frac{1-2k}{2} \notin \text{Spec}(\mu)$ for any $k = 1, 2, \dots$, all the variables $\Phi_{\alpha,p}^m$ are linear combinations of $\sigma_{\varepsilon,k}$ with coefficients being smooth functions of v^1, \dots, v^n .

For an arbitrary tau-symmetric bihamiltonian deformation of the Principal Hierarchy associated with a semisimple Frobenius manifold, we are to construct in the remaining part of this section its super extension and super tau-cover by generalizing the constructions given in Theorem 2.3 and Theorem 2.4.

2.3. Super extensions of bihamiltonian integrable hierarchies. In this subsection, we construct a super extension for a given bihamiltonian integrable hierarchy with hydrodynamic limit.

We fix an n -dimensional smooth manifold M and a semisimple hydrodynamic bihamiltonian structure $(P_0^{[0]}, P_1^{[0]})$ defined on $J^\infty(\hat{M})$. Let us choose $(w^\alpha; \phi_\alpha)$ as local coordinates on \hat{M} . Recall that a Miura type transformation is a choice of n differential polynomials $\tilde{w}^1, \dots, \tilde{w}^n \in \hat{\mathcal{A}}^0$ such that

$$\det \left(\frac{\partial \tilde{w}_0^\alpha}{\partial w^\beta} \right) \neq 0,$$

where \tilde{w}_0^α is the differential degree zero component of \tilde{w}^α . By defining $\tilde{w}^{\alpha,s} = \partial_x^s \tilde{w}^\alpha$, it is easy to see that we can represent any differential polynomial in $w^{\alpha,s}$ by a differential polynomial in $\tilde{w}^{\alpha,s}$. Therefore a Miura type transformation can be viewed as a special type of coordinate transformation on $J^\infty(M)$. The extension of the Miura type transformations on $J^\infty(\hat{M})$ is given by the following theorem [22].

Theorem 2.5 ([22]). *A Miura type transformation induces a change of coordinates from $(w^{\alpha,s}; \phi_\alpha^s)$ to $(\tilde{w}^{\alpha,s}; \tilde{\phi}_\alpha^s)$ given by*

$$\phi_\alpha^s = \partial_x^s \sum_{t \geq 0} (-\partial_x)^t \left(\frac{\partial \tilde{w}^\beta}{\partial w^{\alpha,t}} \tilde{\phi}_\beta \right).$$

Now let (P_0, P_1) be any given deformation of $(P_0^{[0]}, P_1^{[0]})$. Denote by \mathcal{P}_0 and \mathcal{P}_1 the Hamiltonian operators of P_0 and P_1 under the coordinate $(w^{\alpha,s}; \phi_\alpha^s)$. We introduce another family of odd variables $\phi_{\alpha,m}^s$ for $m \geq 0$ and extend the vector field ∂_x to the following one:

$$\partial_x = \sum_{s \geq 0} w^{\alpha,s+1} \frac{\partial}{\partial w^{\alpha,s}} + \sum_{k,s \geq 0} \phi_{\alpha,k}^{s+1} \frac{\partial}{\partial \phi_{\alpha,k}^s}.$$

In what follows we also use the notations $\phi_{\alpha,0}^s = \phi_\alpha^s$ and $\phi_{\alpha,m}^0 = \phi_{\alpha,m}$. Inspired by (2.14), we require that these new odd variables satisfy the recursion relations

$$(2.15) \quad \mathcal{P}_0^{\alpha\beta} \phi_{\beta,m+1} = \mathcal{P}_1^{\alpha\beta} \phi_{\beta,m}, \quad m \geq 0.$$

We first show that (2.15) is well defined in the sense that it is invariant under Miura type transformations.

Proposition 2.6. *The Miura type transformation from $(w^{\alpha,s}; \phi_\alpha^s)$ to $(\tilde{w}^{\alpha,s}; \tilde{\phi}_\alpha^s)$ induces a transformation for the new odd variables $\phi_{\alpha,m}^s$ given by*

$$(2.16) \quad \phi_{\alpha,m}^s = \partial_x^s \sum_{t \geq 0} (-\partial_x)^t \left(\frac{\partial \tilde{w}^\beta}{\partial w^{\alpha,t}} \tilde{\phi}_{\beta,m} \right), \quad m \geq 1$$

such that the recursion relation (2.15) is invariant.

Proof. Denote by \mathcal{P}_i and $\tilde{\mathcal{P}}_i$ the Hamiltonian operator of P_i in the coordinates $(w^{\alpha,s}; \phi_\alpha^s)$ and $(\tilde{w}^{\alpha,s}; \tilde{\phi}_\alpha^s)$ respectively for $i = 0, 1$. Then it is well known that

$$\tilde{\mathcal{P}}_i^{\alpha\beta} = \sum_{s \geq 0} \frac{\partial \tilde{w}^\alpha}{\partial w^{\lambda,s}} \partial_x^s \circ \mathcal{P}_i^{\lambda\varepsilon} \circ \sum_{s \geq 0} (-\partial_x)^s \circ \frac{\partial \tilde{w}^\beta}{\partial w^{\varepsilon,s}}.$$

Therefore by using the relation (2.15), it is easy to see that:

$$\begin{aligned} \tilde{\mathcal{P}}_0^{\alpha\beta} \tilde{\phi}_{\beta,m+1} &= \sum_{s \geq 0} \frac{\partial \tilde{w}^\alpha}{\partial w^{\lambda,s}} \partial_x^s \circ \mathcal{P}_0^{\lambda\varepsilon} \circ \sum_{s \geq 0} (-\partial_x)^s \circ \frac{\partial \tilde{w}^\beta}{\partial w^{\varepsilon,s}} (\tilde{\phi}_{\beta,m+1}) \\ &= \sum_{s \geq 0} \frac{\partial \tilde{w}^\alpha}{\partial w^{\lambda,s}} \partial_x^s \circ \mathcal{P}_0^{\lambda\varepsilon} \phi_{\varepsilon,m+1} \\ &= \sum_{s \geq 0} \frac{\partial \tilde{w}^\alpha}{\partial w^{\lambda,s}} \partial_x^s \circ \mathcal{P}_1^{\lambda\varepsilon} \phi_{\varepsilon,m} \\ &= \sum_{s \geq 0} \frac{\partial \tilde{w}^\alpha}{\partial w^{\lambda,s}} \partial_x^s \circ \mathcal{P}_1^{\lambda\varepsilon} \circ \sum_{s \geq 0} (-\partial_x)^s \circ \frac{\partial \tilde{w}^\beta}{\partial w^{\varepsilon,s}} (\tilde{\phi}_{\beta,m}) \\ &= \tilde{\mathcal{P}}_1^{\alpha\beta} \tilde{\phi}_{\beta,m}. \end{aligned}$$

Thus we see that the recursion relations (2.15) are preserved under the change of variables (2.16). The proposition is proved. \square

By using the theory of bihamiltonian cohomology [7] for $(P_0^{[0]}, P_1^{[0]})$, we can choose a coordinate system $(v^{\alpha,s}; \sigma_\alpha^s)$ such that

$$P_0 = P_0^{[0]} = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1,$$

and the $\hat{\mathcal{F}}_2^2$ component of P_1 vanishes. From now on, we will always use $(v^{\alpha,s}; \sigma_\alpha^s)$ to denote the coordinate system described above. We use the notation $\hat{\mathcal{A}}^+$ to denote the extension of $\hat{\mathcal{A}}$ by including the odd variables $\sigma_{\alpha,m}^s$ for $m \geq 1$ satisfying the recursion relations

$$(2.17) \quad \eta^{\alpha\beta} \sigma_{\beta,m+1}^1 = \mathcal{P}_1^{\alpha\beta} \sigma_{\beta,m}, \quad m \geq 0.$$

As before we use the notation $\sigma_{\alpha,0}^s = \sigma_\alpha^s$ and $\sigma_{\alpha,m}^0 = \sigma_{\alpha,m}$. We will still use ∂_x to denote the vector field on $\hat{\mathcal{A}}^+$ given by

$$\partial_x = \sum_{s \geq 0} v^{\alpha,s+1} \frac{\partial}{\partial v^{\alpha,s}} + \sum_{s,m \geq 0} \sigma_{\alpha,m}^{s+1} \frac{\partial}{\partial \sigma_{\alpha,m}^s}.$$

For any element $f \in \hat{\mathcal{A}}^+$, we say f is *local* if it can be represented by an element in $\hat{\mathcal{A}}$ and we say f is *non-local* if it is not local. Note that on the space $\hat{\mathcal{A}}^+$, the differential degree is not well-defined but the super degree is still well defined by setting the super degree of $\sigma_{\alpha,m}^s$ being 1. We will use $\hat{\mathcal{A}}^{+,p}$ to denote the set of homogeneous elements with super degree p .

Example 2.2. Consider the following bihamiltonian structure of the KdV equation (2.7):

$$\mathcal{P}_0 = \partial_x, \quad \mathcal{P}_1 = v \partial_x + \frac{1}{2} v_x + \frac{\varepsilon^2}{8} \partial_x^3.$$

We introduce odd variables σ_m^s for $s, m \geq 0$ such that they satisfy the recursion relations

$$\sigma_{m+1}^1 = v \sigma_m^1 + \frac{1}{2} v_x \sigma_m + \frac{\varepsilon^2}{8} \sigma_m^3, \quad m \geq 0.$$

Then the ring $\hat{\mathcal{A}}^+$ is given by the quotient

$$\hat{\mathcal{A}}^+ = C^\infty(v)[[v^{(s+1)}, \sigma_m^s \mid m, s \geq 0]]/J,$$

where J is the differential ideal generated by

$$v \sigma_m^1 + \frac{1}{2} v_x \sigma_m + \frac{\varepsilon^2}{8} \sigma_m^3 - \sigma_{m+1}^1, \quad m \geq 0.$$

Then we see that σ_1^1 is local but σ_2^1 is non-local.

Definition 2.1. For $k, l \geq 0$, we define the shift operators

$$T_k : \hat{\mathcal{A}}^1 \rightarrow \hat{\mathcal{A}}^{+,1}, \quad T_{k,l} : \hat{\mathcal{A}}^2 \rightarrow \hat{\mathcal{A}}^{+,2}$$

to be the linear operators given by

$$T_k(f \sigma_{\alpha,0}^s) = f \sigma_{\alpha,k}^s, \quad f \in \hat{\mathcal{A}}^0,$$

$$T_{k,l} = -T_{l,k}; \quad T_{k,l}(f\sigma_{\alpha,0}^t\sigma_{\beta,0}^s) = f \sum_{i=0}^{l-k-1} \sigma_{\alpha,k+i}^t \sigma_{\beta,l-i-1}^s, \quad k \leq l, \quad f \in \hat{\mathcal{A}}^0.$$

In particular, $T_{k,k} = 0$.

The following lemmas are obvious from the definition.

Lemma 2.7. *The shift operators T_k and $T_{k,l}$ commute with ∂_x .*

Lemma 2.8. *The shift operators T_k and $T_{k,l}$ are globally defined, i.e. they are invariant under Miura type transformations.*

Example 2.3. *Using the shift operator, the recursion relation (2.17) can be represented by the following formula*

$$(2.18) \quad T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha,0}} = T_m \frac{\delta P_1}{\delta \sigma_{\alpha,0}}, \quad m \geq 0.$$

Example 2.4. *The recursion relation (2.17) can also be represented by the following formulae:*

$$T_{m+1}(D_{P_0}f) = T_m(D_{P_1}f), \quad m \geq 0, \quad f \in \hat{\mathcal{A}}^0,$$

here D_{P_i} are the derivations defined in (2.1). Indeed, when $f = v^\alpha$ we recover the relation (2.18); for general $f \in \hat{\mathcal{A}}^0$, by definition (2.1), we have

$$T_{m+1}(D_{P_0}f) = T_{m+1} \sum_{s \geq 0} \frac{\partial f}{\partial v^{\alpha,s}} \partial_x^s \frac{\delta P_0}{\delta \sigma_{\alpha,0}} = T_m \sum_{s \geq 0} \frac{\partial f}{\partial v^{\alpha,s}} \partial_x^s \frac{\delta P_1}{\delta \sigma_{\alpha,0}} = T_m(D_{P_1}f).$$

With the help of the shift operator, we can generalize the construction given in the previous subsection. We first introduce the following notation.

Definition 2.2. *We define a family of odd derivations $\frac{\partial}{\partial \tau_m}$ on $\hat{\mathcal{A}}^+$ by*

$$(2.19) \quad \frac{\partial v^\alpha}{\partial \tau_m} = T_m \frac{\delta P_0}{\delta \sigma_{\alpha,0}}, \quad \frac{\partial \sigma_{\alpha,k}}{\partial \tau_m} = T_{k,m} \frac{\delta P_1}{\delta v^\alpha}, \quad \left[\frac{\partial}{\partial \tau_m}, \partial_x \right] = 0.$$

We need to check that that this definition is well-defined, i.e., it is compatible with the recursion relations (2.18).

Lemma 2.9. *The following identity holds true for any $X \in \hat{\mathcal{A}}^1$ and $m, k \geq 0$:*

$$\frac{\partial}{\partial \tau_k} T_m(X) = T_{m,k}(D_{P_1}(X)) - T_{m+1,k}(D_{P_0}(X)).$$

Proof. Since all the operators are linear, we may assume $X = f\sigma_{\beta,0}^l$ for some $f \in \hat{\mathcal{A}}^0$ and $l \geq 0$. We first assume that $k \geq m$. The case $k = m$ can be easily verified as follows:

$$\frac{\partial}{\partial \tau_m}(f\sigma_{\beta,m}^l) = T_m(D_{P_0}f)\sigma_{\beta,m}^l = T_{m,m+1}(D_{P_0}(f\sigma_{\beta,0}^l)).$$

Here we use the fact that $P_0 = P_0^{[0]}$ and $D_{P_0}\sigma_{\beta,0} = 0$. Now we assume $k \geq m+1$, then by using the definition of the shift operators and the recursion relations (2.18) we obtain

the following identities:

$$\begin{aligned}
 \frac{\partial}{\partial \tau_k} (f \sigma_{\beta, m}^l) &= T_k (D_{P_0} f) \sigma_{\beta, m}^l + T_{m, k} (f D_{P_1} \sigma_{\beta, 0}^l) \\
 &= T_{k-1} (D_{P_1} f) \sigma_{\beta, m}^l + T_{m, k} (f D_{P_1} \sigma_{\beta, 0}^l) \\
 &= T_{m, k} ((D_{P_1} f) \sigma_{\beta, 0}^l) - \sum_{i=0}^{k-m-2} T_{m+i} (D_{P_1} f) \sigma_{\beta, k-i-1}^l + T_{m, k} (f D_{P_1} \sigma_{\beta, 0}^l) \\
 &= T_{m, k} (D_{P_1} (f \sigma_{\beta, 0}^l)) - \sum_{i=0}^{k-m-2} T_{m+i+1} (D_{P_0} f) \sigma_{\beta, k-i-1}^l \\
 &= T_{m, k} (D_{P_1} (f \sigma_{\beta, 0}^l)) - T_{m+1, k} (D_{P_0} (f \sigma_{\beta, 0}^l)).
 \end{aligned}$$

The case $k < m$ is proved in exactly the same way. The lemma is proved. \square

Proposition 2.10. *The flows $\frac{\partial}{\partial \tau_k}$ are compatible with the recursion relations (2.18), i.e.,*

$$\frac{\partial}{\partial \tau_k} T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} = \frac{\partial}{\partial \tau_k} T_m \frac{\delta P_1}{\delta \sigma_{\alpha, 0}}, \quad m, k \geq 0.$$

Proof. Using the fact that

$$P_0 = P_0^{[0]} = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_{\alpha} \sigma_{\beta}^1,$$

it is easy to obtain the following identities:

$$\frac{\partial}{\partial \tau_k} T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} = \frac{\partial}{\partial \tau_k} \eta^{\alpha\beta} \sigma_{\beta, m+1}^1 = T_{m+1, k} \left(\eta^{\alpha\beta} \partial_x \frac{\delta P_1}{\delta v^{\beta}} \right) = T_{m+1, k} \left(D_{P_1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} \right).$$

Since $[P_0, P_1] = 0$, it follows from the identity (2.4) that

$$\frac{\partial}{\partial \tau_k} T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} = -T_{m+1, k} \left(D_{P_0} \frac{\delta P_1}{\delta \sigma_{\alpha, 0}} \right).$$

Thus by using $[P_1, P_1] = 0$ and Lemma 2.9 we finish the proof of the proposition. \square

Proposition 2.11. *The odd flows $\frac{\partial}{\partial \tau_m}$ mutually commute, i.e.,*

$$\left[\frac{\partial}{\partial \tau_m}, \frac{\partial}{\partial \tau_k} \right] = 0, \quad m, k \geq 0.$$

Proof. By the definition of the flows $\frac{\partial}{\partial \tau_m}$, it is easy to see that

$$\left[\frac{\partial}{\partial \tau_m}, \frac{\partial}{\partial \tau_k} \right] v^{\alpha} = \eta^{\alpha\beta} \partial_x T_{k, m} \frac{\delta P_1}{\delta v^{\beta}} + \eta^{\alpha\beta} \partial_x T_{m, k} \frac{\delta P_1}{\delta v^{\beta}} = 0.$$

To show the commutation relation

$$\left[\frac{\partial}{\partial \tau_l}, \frac{\partial}{\partial \tau_k} \right] \sigma_{\alpha, m} = 0,$$

it suffices to verify the case $m = 0$ due to the recursion relations (2.17). By using the trivial relation

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial \tau_0} \right] \sigma_{\alpha, 0} = 0,$$

the recursion relations (2.17), and by induction on k , we arrive at

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial \tau_0} \right] \sigma_{\alpha, k} = 0, \quad k \geq 0.$$

This commutation relation is equivalent to

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial \tau_k} \right] \sigma_{\alpha, 0} = 0$$

due to the definition of the odd flows. Using induction again we arrive at the identity

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial \tau_k} \right] \sigma_{\alpha, l} = 0$$

for any $l \geq 0$. Therefore we have

$$\frac{\partial}{\partial \tau_k} \frac{\partial \sigma_{\alpha, 0}}{\partial \tau_l} = - \frac{\partial}{\partial \tau_k} \frac{\partial \sigma_{\alpha, l}}{\partial \tau_0} = \frac{\partial}{\partial \tau_0} \frac{\partial \sigma_{\alpha, l}}{\partial \tau_k}.$$

It follows from the definition of the odd flows that the right hand side is anti-symmetric with respect to the indices k, l , hence we prove that

$$\left[\frac{\partial}{\partial \tau_l}, \frac{\partial}{\partial \tau_k} \right] \sigma_{\alpha, 0} = 0.$$

The proposition is proved. \square

Now let $X_i \in \hat{\mathcal{F}}^1$, $i \in I$ be a family of bihamiltonian vector fields with respect to an index set I , i.e., each X_i satisfies the equations $[X_i, P_0] = [X_i, P_1] = 0$. Recall that the family $\{X_i\}$ corresponds to a bihamiltonian integrable hierarchy given by

$$(2.20) \quad \frac{\partial v^\alpha}{\partial t_i} = \frac{\delta X_i}{\delta \sigma_{\alpha, 0}}, \quad i \in I.$$

In what follows we will extend this integrable hierarchy such that it becomes a system of mutually commuting vector fields on $\hat{\mathcal{A}}^+$.

Definition 2.3. For a bihamiltonian vector field $X \in \hat{\mathcal{F}}^1$ we associate it with the following system of PDEs on $\hat{\mathcal{A}}^+$:

$$\frac{\partial v^\alpha}{\partial t_X} = D_X v^\alpha, \quad \frac{\partial \sigma_{\alpha, m}}{\partial t_X} = T_m D_X \sigma_{\alpha, 0}, \quad \left[\frac{\partial}{\partial t_X}, \partial_x \right] = 0.$$

It is called the super extended flow given by X .

Proposition 2.12. The super extended flow $\frac{\partial}{\partial t_X}$ given by a bihamiltonian vector field X is compatible with the recursion relations (2.18), i.e.,

$$(2.21) \quad \frac{\partial}{\partial t_X} T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} = \frac{\partial}{\partial t_X} T_m \frac{\delta P_1}{\delta \sigma_{\alpha, 0}}.$$

Proof. By definition of the flow $\frac{\partial}{\partial t_X}$, it is easy to see that

$$\frac{\partial}{\partial t_X} T_{m+1} \frac{\delta P_0}{\delta \sigma_{\alpha, 0}} = T_{m+1} D_X \frac{\delta P_0}{\delta \sigma_{\alpha, 0}}, \quad \frac{\partial}{\partial t_X} T_m \frac{\delta P_1}{\delta \sigma_{\alpha, 0}} = T_m D_X \frac{\delta P_1}{\delta \sigma_{\alpha, 0}}.$$

Using the fact that $[X, P_0] = [X, P_1] = 0$ and the identity (2.4), we see that (2.21) is equivalent to the following identity

$$T_{m+1}D_{P_0}\frac{\delta X}{\delta\sigma_{\alpha,0}} = T_mD_{P_1}\frac{\delta X}{\delta\sigma_{\alpha,0}},$$

which is equivalent to the recursion relations (2.18) according to Example 2.4. The proposition is proved. \square

Proposition 2.13. *Let X and Y be two bihamiltonian vector fields, then their super extended flows commute.*

Proof. From the theory of the bihamiltonian cohomology [7] we know that $[X, Y] = 0$, hence it follows from (2.2) that

$$\left[\frac{\partial}{\partial t_X}, \frac{\partial}{\partial t_Y}\right]v^\alpha = 0, \quad \left[\frac{\partial}{\partial t_X}, \frac{\partial}{\partial t_Y}\right]\sigma_{\alpha,0} = 0.$$

Using the definition of the super extended flow, it is easy to see that

$$\left[\frac{\partial}{\partial t_X}, \frac{\partial}{\partial t_Y}\right]\sigma_{\alpha,m} = T_m\left[\frac{\partial}{\partial t_X}, \frac{\partial}{\partial t_Y}\right]\sigma_{\alpha,0} = 0.$$

The proposition is proved. \square

Now let us prove that the super extended flow given by a bihamiltonian vector field commutes with the odd flows $\frac{\partial}{\partial\tau_m}$.

Lemma 2.14. *For any $\mathcal{D} \in \text{Der}(\hat{\mathcal{A}})^0$ satisfying the condition $[\mathcal{D}, \partial_x] = 0$, we extend its action to $\hat{\mathcal{A}}^+$ by setting*

$$\mathcal{D}\sigma_{\alpha,m} = T_m\mathcal{D}\sigma_{\alpha,0}, \quad m \geq 0.$$

Then the following identities hold true:

$$T_k \circ (\mathcal{D}|_{\hat{\mathcal{A}}^1}) = (\mathcal{D}|_{\hat{\mathcal{A}}^{1,1}}) \circ T_k; \quad T_{k,l} \circ (\mathcal{D}|_{\hat{\mathcal{A}}^2}) = (\mathcal{D}|_{\hat{\mathcal{A}}^{2,2}}) \circ T_{k,l}, \quad k, l \geq 0.$$

Proof. The first identity is obvious from the definition $\mathcal{D}\sigma_{\alpha,k} = T_k\mathcal{D}\sigma_{\alpha,0}$. The second one is also easy to verify by using the definition of the shift operator $T_{k,l}$. The lemma is proved. \square

Proposition 2.15. *The odd flows $\frac{\partial}{\partial\tau_m}$ commute with the super extended flow given by a bihamiltonian vector field X .*

Proof. Using Lemma 2.14 and the definition of the odd flows, it is easy to see that

$$\frac{\partial}{\partial t_X} \frac{\partial v^\alpha}{\partial \tau_m} = \frac{\partial}{\partial t_X} T_m \frac{\delta P_0}{\delta \sigma_{\alpha,0}} = T_m D_X \left(\frac{\delta P_0}{\delta \sigma_{\alpha,0}} \right), \quad \frac{\partial}{\partial \tau_m} \frac{\partial v^\alpha}{\partial t_X} = T_m D_{P_0} \left(\frac{\partial v^\alpha}{\partial t_X} \right).$$

Therefore it follows from $[X, P_0] = 0$ and the identity (2.4) that

$$\left[\frac{\partial}{\partial \tau_m}, \frac{\partial}{\partial t_X}\right]v^\alpha = 0.$$

Similarly by using Lemma 2.14 again we have

$$\frac{\partial}{\partial t_X} \frac{\partial \sigma_{\alpha,k}}{\partial \tau_m} = \frac{\partial}{\partial t_X} T_{k,m} \frac{\delta P_1}{\delta v^\alpha} = T_{k,m} D_X \left(\frac{\delta P_1}{\delta v^\alpha} \right).$$

On the other hand, by using Lemma 2.9 we obtain

$$\frac{\partial}{\partial \tau_m} \frac{\partial \sigma_{\alpha,k}}{\partial t_X} = -\frac{\partial}{\partial \tau_m} T_k \left(D_X \left(\frac{\delta X}{\delta v^\alpha} \right) \right) = -T_{k,m} \left(D_{P_1} \left(\frac{\delta X}{\delta v^\alpha} \right) \right).$$

Here we use the fact that $[P_0, X] = 0$ and $\frac{\delta P_0}{\delta v^\alpha} = 0$. Now by using (2.3) and $[X, P_1] = 0$, it is easy to conclude that

$$\left[\frac{\partial}{\partial \tau_m}, \frac{\partial}{\partial t_X} \right] \sigma_{\alpha,k} = 0.$$

The proposition is proved. \square

Let us summarize the constructions given in this subsection in the following theorem.

Theorem 2.16. *Let (P_0, P_1) be a bihamiltonian structure with semisimple hydrodynamic leading terms and $\{X_i\}$ be a family of bihamiltonian vector fields. Then we have the following super extended integrable hierarchy:*

$$\begin{aligned} \frac{\partial v^\alpha}{\partial t_i} &= D_{X_i} v^\alpha, & \frac{\partial \sigma_{\alpha,m}}{\partial t_i} &= T_m D_{X_i} \sigma_{\alpha,0}; \\ \frac{\partial v^\alpha}{\partial \tau_m} &= T_m \frac{\delta P_0}{\delta \sigma_{\alpha,0}}, & \frac{\partial \sigma_{\alpha,k}}{\partial \tau_m} &= T_{k,m} \frac{\delta P_1}{\delta v^\alpha}. \end{aligned}$$

The flows in this super extended hierarchy mutually commute.

2.4. Deformation of the super tau-covers. In this section, we will construct super tau-covers for tau-symmetric bihamiltonian deformations of the Principal Hierarchy associates with a semisimple Frobenius manifold. Let us first recall how to construct the deformations of the tau-cover (2.13) of the Principal Hierarchy following [9].

We fix a semisimple Frobenius manifold M and use $(P_0^{[0]}, P_1^{[0]})$ to denote the bihamiltonian structure (2.12). The two-point functions in the tau-cover (2.13) will be denoted by $\Omega_{\alpha,p;\beta,q}^{[0]}$ and similarly we denote $h_{\alpha,p}^{[0]} = \Omega_{\alpha,p;1,0}^{[0]}$. Let (P_0, P_1) be a deformation of $(P_0^{[0]}, P_1^{[0]})$, then it determines a unique deformation of the Principal Hierarchy according to [7, 23]. By using the results proved in [13], we know that a bihamiltonian deformation of the Principal Hierarchy is tau-symmetric if and only if the central invariants of the deformation of the bihamiltonian structure are constants. In such a case, after an appropriate Miura type transformation the Hamiltonian structure P_0 can be represented in the form

$$P_0 = P_0^{[0]} = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1,$$

and P_1 has no $\hat{\mathcal{F}}_2^2$ components. Moreover, we can also require that the condition of exactness of the bihamiltonian structure is preserved [9, 13]:

$$P_0 = [Z, P_1], \quad Z = \int \sigma_1.$$

In what follows, we will always assume that P_0 and Z take the above forms. We still use the same notation $X_{\alpha,p} \in \hat{\mathcal{F}}^1$, as we have already used in (2.11), to denote the unique deformed flows of the Principal Hierarchy and we will also use $\frac{\partial}{\partial t^{\alpha,p}}$ to denote the vector

field $D_{X_{\alpha,p}}$. Let $H_{\alpha,p} \in \hat{\mathcal{F}}^0$ be the unique deformation of the Hamiltonian of the Principal Hierarchy such that

$$(2.22) \quad X_{\alpha,p} = -[H_{\alpha,p}, P_0], \quad \alpha = 1, \dots, n, \quad p \geq 0,$$

and

$$(2.23) \quad H_{\alpha,-1} := \int \eta_{\alpha\beta} v^\beta.$$

Let us define the following differential polynomials:

$$(2.24) \quad h_{\alpha,p} = D_Z H_{\alpha,p}, \quad \alpha = 1, \dots, n, \quad p \geq 0.$$

Note we use an index convention that is different from the one used in [9].

Proposition 2.17 ([9]). *We have the following results:*

- (1) $D_{X_{1,0}} = \partial_x$.
- (2) *The functionals $H_{\alpha,p}$ defined in (2.22), (2.23) and the differential polynomials defined in (2.24) satisfy the relations*

$$H_{\alpha,p} = \int h_{\alpha,p+1}, \quad p \geq -1.$$

We also have the following properties for the Hamiltonians and two-point functions of the deformed Principal Hierarchy.

Proposition 2.18. *The following bihamiltonian recursion relations hold true:*

$$(2.25) \quad [H_{\alpha,p-1}, P_1] = \left(p + \frac{1}{2} + \mu_\alpha\right) [H_{\alpha,p}, P_0] + \sum_{k=1}^p (R_k)_\alpha^\gamma [H_{\gamma,p-k}, P_0], \quad p \geq 0.$$

Proof. Denote the following by $Y_{\alpha,p} \in \hat{\mathcal{F}}^1$ for $p \geq 0$:

$$Y_{\alpha,p} = [H_{\alpha,p-1}, P_1] - \left(p + \frac{1}{2} + \mu_\alpha\right) [H_{\alpha,p}, P_0] - \sum_{k=1}^p (R_k)_\alpha^\gamma [H_{\gamma,p-k}, P_0].$$

According to Theorem 2.2, we know that $Y_{\alpha,p} \in \hat{\mathcal{F}}_{\geq 2}^1$. Since the flows $X_{\alpha,p} = -[H_{\alpha,p}, P_0]$ is bihamiltonian, we easily conclude that $Y_{\alpha,p}$ is a bihamiltonian vector field. Therefore by using the results of bihamiltonian cohomology given in [7], we arrive at $Y_{\alpha,p} = 0$. \square

Theorem 2.19 ([9]). *There exist differential polynomials $\Omega_{\alpha,p;\beta,q}$ such that they are deformations of $\Omega_{\alpha,p;\beta,q}^{[0]}$, and satisfy the following properties:*

- (1) $\partial_x \Omega_{\alpha,p;\beta,q} = \frac{\partial h_{\beta,q}}{\partial t^{\alpha,p}}$.
- (2) $\Omega_{\alpha,p;\beta,q} = \Omega_{\beta,q;\alpha,p}$ and $\Omega_{\alpha,p;1,0} = h_{\alpha,p}$.
- (3) $\frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial t^{\lambda,k}} = \frac{\partial \Omega_{\lambda,k;\beta,q}}{\partial t^{\alpha,p}}$.

To construct the tau-cover of the deformed Principal Hierarchy, we introduce the following *normal coordinates* as in [12]:

$$(2.26) \quad w_\alpha = h_{\alpha,0}, \quad w^\alpha = \eta^{\alpha\beta} w_\beta,$$

then we see that the differential polynomials w^α and v^α are related by a Miura type transformation. In particular, it follows from Proposition 2.17 that

$$(2.27) \quad H_{\alpha,-1} = \int w_\alpha = \int \eta_{\alpha\beta} v^\beta.$$

In terms of the normal coordinates, the tau-cover of the deformed Principal Hierarchy can be represented in the form (c.p. (2.13))

$$(2.28) \quad \frac{\partial f_{\alpha,p}}{\partial t^{\beta,q}} = \Omega_{\alpha,p;\beta,q}, \quad \frac{\partial w^\alpha}{\partial t^{\beta,q}} = \eta^{\alpha\lambda} \partial_x \Omega_{\lambda,0;\beta,q}.$$

Let us proceed to construct its super tau-cover. To this end we introduce odd variables $\Phi_{\alpha,p}^m$, as we do for the super tau-cover of the Principal Hierarchy, such that

$$(2.29) \quad \partial_x \Phi_{\alpha,p}^m = \frac{\partial h_{\alpha,p}}{\partial \tau_m}, \quad m \geq 0.$$

Here the odd flows $\frac{\partial}{\partial \tau_m}$ are defined by 2.19. By using Theorem 2.19 we obtain the following identity:

$$\frac{\partial}{\partial t^{\beta,q}} \frac{\partial h_{\alpha,p}}{\partial \tau_m} = \partial_x \frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial \tau_m}.$$

Therefore we conclude that the following definitions of the evolutions of the odd variables $\Phi_{\alpha,p}^m$ along the flows $\frac{\partial}{\partial t^{\beta,q}}$ are compatible with (2.29):

$$\frac{\partial \Phi_{\alpha,p}^m}{\partial t^{\beta,q}} = \frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial \tau_m}.$$

To define the evolutions of $\Phi_{\alpha,p}^m$ along the odd flows $\frac{\partial}{\partial \tau_k}$, we need the following lemma.

Lemma 2.20. *There exist differential polynomials $F_{\alpha,p} \in \hat{\mathcal{A}}^2$ such that*

$$\frac{\partial}{\partial \tau_k} \frac{\partial h_{\alpha,p}}{\partial \tau_m} = T_{m,k} \partial_x F_{\alpha,p}, \quad \alpha = 1, \dots, n; \quad p, m, k \geq 0.$$

Proof. By using the definition (2.1) of D_{P_i} and the fact that the vector fields $X_{\alpha,p}$ are bihamiltonian, it is easy to see that

$$\int D_{P_1} D_{P_0} h_{\alpha,p} = \int [P_1, [P_0, H_{\alpha,p-1}]] = 0.$$

Therefore there exists $F_{\alpha,p} \in \hat{\mathcal{A}}^2$ such that

$$\frac{\partial}{\partial \tau_1} \frac{\partial h_{\alpha,p}}{\partial \tau_0} = \partial_x F_{\alpha,p}.$$

Now from Lemma 2.9 it follows that

$$\frac{\partial}{\partial \tau_k} \frac{\partial h_{\alpha,p}}{\partial \tau_m} = T_{m,k} \partial_x F_{\alpha,p}.$$

The lemma is proved. \square

Now we are ready to construct the following super tau-cover of the deformed Principal Hierarchy:

Theorem 2.21. *Let M be a semisimple Frobenius manifold and (P_0, P_1) be a deformation of the bihamiltonian structure (2.12) with constant central invariants. Then we have the following deformation of the super tau-cover of the Principal Hierarchy:*

$$\begin{aligned}\frac{\partial f_{\alpha,p}}{\partial t^{\beta,q}} &= \Omega_{\alpha,p;\beta,q}, \\ \frac{\partial f_{\alpha,p}}{\partial \tau_m} &= \Phi_{\alpha,p}^m, \\ \frac{\partial \Phi_{\alpha,p}^m}{\partial t^{\beta,q}} &= \frac{\partial \Omega_{\alpha,p;\beta,q}}{\partial \tau_m}, \\ \frac{\partial \Phi_{\alpha,p}^m}{\partial \tau_k} &= T_{m,k} F_{\alpha,p}.\end{aligned}$$

Recall that when the diagonal matrix μ of a Frobenius manifold M satisfies the condition $\frac{1-2k}{2} \notin \text{Spec}(\mu)$ for any $k = 1, 2, \dots$, the odd variables $\Phi_{\alpha,p}^m$ for the super tau-cover of the Principal Hierarchy are redundant since they can be represented by elements in $\hat{\mathcal{A}}^+$. We want to show that the deformed super tau-cover have the same property which will play an important role in our consideration of the deformation of the Virasoro symmetries.

We start with the definition of generalized shift operators.

Definition 2.4. *We define the shift operators \hat{T}_k for $k \geq 0$ to be linear operators $\hat{\mathcal{A}}^{+,1} \rightarrow \hat{\mathcal{A}}^{+,1}$ such that*

$$\hat{T}_k(f\sigma_{\alpha,m}^l) = f\sigma_{\alpha,m+k}^l, \quad f \in \hat{\mathcal{A}}^0, \quad m \geq 0.$$

The following lemma is easy to verify.

Lemma 2.22. *The operators \hat{T}_k commute with ∂_x and are compatible with the recursion relations (2.17).*

Lemma 2.23. *If $\Phi_{\alpha,p}^0$ can be represented by an element in $\hat{\mathcal{A}}^{+,1}$, then so does $\Phi_{\alpha,p}^m$ and $\Phi_{\alpha,p}^m = \hat{T}_m \Phi_{\alpha,p}^0$ for $m \geq 1$.*

Proof. By using the relation (2.29), it is easy to see that

$$\partial_x \hat{T}_m \Phi_{\alpha,p}^0 = \hat{T}_m \frac{\partial h_{\alpha,p}}{\partial \tau_0} = \frac{\partial h_{\alpha,p}}{\partial \tau_m}.$$

The lemma is proved. □

We will use the notation $\Phi_{\alpha,p}^m \in \hat{\mathcal{A}}^+$ to mean that $\Phi_{\alpha,p}^m$ can be represented by an element in $\hat{\mathcal{A}}^+$.

Proposition 2.24. *We have $\Phi_{\alpha,0}^m \in \hat{\mathcal{A}}^+$ and*

$$(2.30) \quad \left(\prod_{k=1}^p \left(k - \frac{1}{2} + \mu_\alpha \right) \right) \Phi_{\alpha,p}^m \in \hat{\mathcal{A}}^+, \quad p \geq 1.$$

Proof. It follows from Lemma 2.23 that it is enough to prove this lemma for $m = 0$. For $\Phi_{\alpha,0}^0$, it is easy to see from (2.27) that there exist differential polynomials $g_\alpha \in \hat{\mathcal{A}}_{\geq 1}^0$ such that

$$(2.31) \quad h_{\alpha,0} = \eta_{\alpha\beta} v^\beta + \partial_x g_\alpha.$$

Therefore we arrive at

$$(2.32) \quad \Phi_{\alpha,0}^0 = \sigma_{\alpha,0} + \frac{\partial g_\alpha}{\partial \tau_0} \in \hat{\mathcal{A}}.$$

We proceed to consider $\Phi_{\alpha,p}^0$ for $p \geq 1$. By using Proposition 2.17 we can rewrite (2.25) as follows:

$$(2.33) \quad \left[\int h_{\alpha,p}, P_1 \right] = \left(p + \frac{1}{2} + \mu_\alpha \right) \left[\int h_{\alpha,p+1}, P_0 \right] + \sum_{k=1}^p (R_k)_\alpha^\gamma \left[\int h_{\gamma,p-k+1}, P_0 \right].$$

Take $p = 0$ in (2.33) and rewrite it in the form

$$\left(\frac{1}{2} + \mu_\alpha \right) \int \frac{\partial h_{\alpha,1}}{\partial \tau_0} = \int \frac{\partial h_{\alpha,0}}{\partial \tau_1},$$

which means that there exists a differential polynomial $p_{\alpha,1} \in \hat{\mathcal{A}}^1$ such that

$$\left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial h_{\alpha,1}}{\partial \tau_0} = \frac{\partial h_{\alpha,0}}{\partial \tau_1} + \partial_x p_{\alpha,1}.$$

Therefore we have

$$\left(\frac{1}{2} + \mu_\alpha \right) \Phi_{\alpha,1}^0 = \Phi_{\alpha,0}^1 + p_{\alpha,1} \in \hat{\mathcal{A}}^+.$$

For general $p \geq 1$, we can prove (2.30) by using (2.33) and induction on p . The proposition is proved. \square

3. DEFORMATION OF VIRASORO SYMMETRIES: FORMULATION

In this section, we first recall the theory of variational bihamiltonian cohomology given in [20] and then explain how to use it to study Virasoro symmetries of the deformed Principal Hierarchies. We also use the example of the deformation of the Riemann hierarchy to illustrate our approach to the study of Virasoro symmetries.

3.1. Variational bihamiltonian cohomologies. In [20], we established a cohomology theory on the space $\text{Der}^\partial(\hat{\mathcal{A}})$ consisting of derivations on $\hat{\mathcal{A}}$ that commute with ∂_x . This theory provides us suitable tools to study Virasoro symmetries of deformations of the Principal Hierarchies. We recall the basic definitions and results in this subsection.

Let us define the space $\text{Der}^\partial(\hat{\mathcal{A}})$ by

$$\text{Der}^\partial(\hat{\mathcal{A}}) = \{X \in \text{Der}(\hat{\mathcal{A}}) \mid [X, \partial_x] = 0\}.$$

We grade the space $\text{Der}^\partial(\hat{\mathcal{A}})$ by the differential gradation and the super gradation, and use the notation $\text{Der}^\partial(\hat{\mathcal{A}})_d^p$ to denote the subspace of homogeneous elements with super degree p and differential degree d .

Lemma 3.1. $\text{Der}^\partial(\hat{\mathcal{A}})_d^p = 0$ for $p \leq -2$ or $d < 0$.

Proof. Let us choose a local coordinate system (w^α, ϕ_α) on \hat{M} . Assume $X \in \text{Der}^\partial(\hat{\mathcal{A}})$ with super degree $p \leq -2$ or $d < 0$. Then by definition this means

$$X(w^\alpha) = X(\phi_\alpha) = 0.$$

Since $[X, \partial_x] = 0$, we immediately see that $X(w^{\alpha, s}) = 0$ and $X(\phi_\alpha^s) = 0$ for $s \geq 0$. Hence $X = 0$ and the lemma is proved. \square

Let $P^{[0]}$ be a Hamiltonian structure of hydrodynamic type and $(P_0^{[0]}, P_1^{[0]})$ be a semisimple bihamiltonian structure of hydrodynamic type. Then by using (2.2) we have a complex $(\text{Der}^\partial(\hat{\mathcal{A}}), D_{P^{[0]}})$ and a double complex $(\text{Der}^\partial(\hat{\mathcal{A}}), D_{P_0^{[0]}}, D_{P_1^{[0]}})$. We define the following cohomology groups:

$$(3.1) \quad H_d^p(\text{Der}^\partial(\hat{\mathcal{A}}), P^{[0]}) = \frac{\text{Der}^\partial(\hat{\mathcal{A}})_d^p \cap \ker D_{P^{[0]}}}{\text{Der}^\partial(\hat{\mathcal{A}})_d^p \cap \text{Im } D_{P^{[0]}}}, \quad p, d \geq 0,$$

$$(3.2) \quad BH_d^p(\text{Der}^\partial(\hat{\mathcal{A}}), P_0^{[0]}, P_1^{[0]}) = \frac{\text{Der}^\partial(\hat{\mathcal{A}})_d^p \cap \ker D_{P_0^{[0]}} \cap \ker D_{P_1^{[0]}}}{\text{Der}^\partial(\hat{\mathcal{A}})_d^p \cap \text{Im } D_{P_0^{[0]}} D_{P_1^{[0]}}}, \quad p, d \geq 0.$$

Note that the spaces $\text{Der}^\partial(\hat{\mathcal{A}})_d^{-1} \neq 0$ for $d \geq 0$ and they must be taken into account while computing the cohomology groups. For example, the space $H_d^0(\text{Der}^\partial(\hat{\mathcal{A}}), P^{[0]})$ is given by

$$H_d^0(\text{Der}^\partial(\hat{\mathcal{A}}), P^{[0]}) = \frac{\ker(D_{P^{[0]}} : \text{Der}^\partial(\hat{\mathcal{A}})_d^0 \rightarrow \text{Der}^\partial(\hat{\mathcal{A}})_{d+1}^1)}{\text{Im}(D_{P^{[0]}} : \text{Der}^\partial(\hat{\mathcal{A}})_{d-1}^{-1} \rightarrow \text{Der}^\partial(\hat{\mathcal{A}})_d^0)}.$$

By using the canonical symplectic structure on \hat{M} , we can identify the space $\text{Der}^\partial(\hat{\mathcal{A}})$ with the space $\bar{\Omega}$ of local functionals of variational 1-forms, and the vector fields $D_{P^{[0]}}$, $D_{P_0^{[0]}}$ and $D_{P_1^{[0]}}$ induce Lie derivatives on $\bar{\Omega}$. This is the reason why we call the above cohomology groups the variational cohomology groups. In [20], the cohomology groups (3.1) and (3.2) are computed by converting them to the cohomology groups on the space $\bar{\Omega}$. The details of the computation of these cohomology groups are not used in the present paper, so we omit them and refer the readers to [20]. The following result plays an essential role in the present paper.

Theorem 3.2 ([20]). *We have the following results on the cohomology groups (3.1) and (3.2):*

- (1) $H_d^p(\text{Der}^\partial(\hat{\mathcal{A}}), P^{[0]}) = 0$ for $p \geq 0, d > 0$.
- (2) $BH_{\geq 2}^0(\text{Der}^\partial(\hat{\mathcal{A}}), P_0^{[0]}, P_1^{[0]}) = 0$.
- (3) $BH_{\geq 4}^1(\text{Der}^\partial(\hat{\mathcal{A}}), P_0^{[0]}, P_1^{[0]}) = 0$.
- (4) $BH_3^1(\text{Der}^\partial(\hat{\mathcal{A}}), P_0^{[0]}, P_1^{[0]}) \cong \oplus_{i=1}^n C^\infty(\mathbb{R})$, and if we denote the action of a cocycle $X \in \text{Der}^\partial(\hat{\mathcal{A}})_3^1$ on the i -th canonical coordinate u^i by

$$X(u^i) = \sum_{j=1}^n \sum_{k=0}^3 X_{i,j}^k \theta_j^{3-k}, \quad X_{i,j}^k \in \hat{\mathcal{A}}_k^0,$$

then the cohomology class $[X]$ is determined by the following functions:

$$c_1 = \frac{X_{1,1}^0}{(f^1)^2}, \quad c_2 = \frac{X_{2,2}^0}{(f^2)^2}, \quad \dots, \quad c_n = \frac{X_{n,n}^0}{(f^n)^2}.$$

Here each function c_i depends only on the i -th canonical coordinate u^i , and f^i is the function defined in (2.6).

3.2. Virasoro symmetries of the Principal Hierarchies. In this subsection, we recall the construction of Virasoro symmetries of the super tau-cover of the Principal Hierarchy following [19]. In [11], a family of infinitely many symmetries $\frac{\partial}{\partial s_m^{even}}$ for $m \geq -1$ of the tau-cover of the Principal Hierarchy associated with a Frobenius manifold M was constructed. This family of symmetries are called the Virasoro symmetries due to the property

$$\left[\frac{\partial}{\partial s_k^{even}}, \frac{\partial}{\partial s_l^{even}} \right] = (l - k) \frac{\partial}{\partial s_{k+l}^{even}}, \quad k, l \geq -1.$$

These symmetries can be represented by a family of quadratic differential operators L_m^{even} of the form

$$L_m^{even} = a_m^{\alpha,p;\beta,q} \frac{\partial^2}{\partial t^{\alpha,p} \partial t^{\beta,q}} + b_{m;\alpha,p}^{\beta,q} t^{\alpha,p} \frac{\partial}{\partial t^{\beta,q}} + c_{m;\alpha,p;\beta,q} t^{\alpha,p} t^{\beta,q} + \frac{1}{4} \delta_{m,0} \text{tr} \left(\frac{1}{4} - \mu^2 \right),$$

where $a_m^{\alpha,p;\beta,q}$, $b_{m;\alpha,p}^{\beta,q}$, $c_{m;\alpha,p;\beta,q}$ are some constants determined from the monodromy data of M and one may refer to [11] for details. These operators satisfy the Virasoro commutation relation

$$[L_k^{even}, L_l^{even}] = (k - l) L_{l+k}^{even}.$$

In this paper, we only need the explicit expressions for L_{-1}^{even} and L_2^{even} which are given by

$$(3.3) \quad L_{-1}^{even} = \frac{1}{2} \eta_{\alpha\beta} t^{\alpha,0} t^{\beta,0} + \sum_{p \geq 1} t^{\alpha,p} \frac{\partial}{\partial t^{\alpha,p-1}},$$

$$(3.4) \quad L_2^{even} = a^{\alpha\beta} \frac{\partial^2}{\partial t^{\alpha,1} \partial t^{\beta,0}} + b^{\alpha\beta} \frac{\partial^2}{\partial t^{\alpha,0} \partial t^{\beta,0}} + \mathcal{L}_2^{even} + c_{2;\alpha,p;\beta,q} t^{\alpha,p} t^{\beta,q},$$

where the constants have the expressions

$$(3.5) \quad a^{\alpha\beta} = \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\beta \right) \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{3}{2} + \mu_\alpha \right),$$

$$(3.6) \quad b^{\alpha\beta} = \frac{1}{2} \eta^{\beta\gamma} (R_1)_\gamma^\alpha \left(\frac{1}{4} + 3\mu_\beta - 3\mu_\beta^2 \right),$$

and the operator \mathcal{L}_2^{even} is given by

$$\begin{aligned}
 (3.7) \quad & \sum_{p \geq 0} \left(p + \frac{1}{2} + \mu_\alpha \right) \left(p + \frac{3}{2} + \mu_\alpha \right) \left(p + \frac{5}{2} + \mu_\alpha \right) t^{\alpha,p} \frac{\partial}{\partial t^{\alpha,p+2}} \\
 & + \sum_{p \geq 0} \sum_{1 \leq r \leq p+2} \left(3 \left(p + \frac{1}{2} + \mu_\alpha \right)^2 + 6 \left(p + \frac{1}{2} + \mu_\alpha \right) + 2 \right) (R_r)_\alpha^\beta t^{\alpha,p} \frac{\partial}{\partial t^{\beta,p-r+2}} \\
 & + \sum_{p \geq 0} \sum_{2 \leq r \leq p+2} \left(3p + \frac{9}{2} + 3\mu_\alpha \right) (R_{r,2})_\alpha^\beta t^{\alpha,p} \frac{\partial}{\partial t^{\beta,p-r+2}} \\
 & + \sum_{p \geq 1} \sum_{3 \leq r \leq p+2} (R_{r,3})_\alpha^\beta t^{\alpha,p} \frac{\partial}{\partial t^{\beta,p-r+2}}.
 \end{aligned}$$

The explicit expressions for the matrices $R_{k,l}$ and constants $c_{2;\alpha,p;\beta,q}$ are not used in this paper, so we omit them.

We have the following theorem for the Virasoro symmetries of the tau-cover of the Principal Hierarchy.

Theorem 3.3 ([11]). *Let us define the following time-dependent flows for $m \geq -1$:*

$$\begin{aligned}
 \frac{\partial f_{\lambda,k}}{\partial s_m^{even}} &= \frac{\partial}{\partial t^{\lambda,k}} \left(\sum a_m^{\alpha,p;\beta,q} f_{\alpha,p} f_{\beta,q} + \sum b_{m;\alpha,p}^{\beta,q} t^{\alpha,p} f_{\beta,q} + \sum c_{m;\alpha,p;\beta,q} t^{\alpha,p} t^{\beta,q} \right), \\
 \frac{\partial v^\lambda}{\partial s_m^{even}} &= \eta^{\lambda\gamma} \frac{\partial}{\partial t^{1,0}} \frac{\partial f_{\gamma,0}}{\partial s_m^{even}}.
 \end{aligned}$$

Then the following commutation relations hold true:

$$\left[\frac{\partial}{\partial s_k^{even}}, \frac{\partial}{\partial t^{\alpha,p}} \right] = 0, \quad \left[\frac{\partial}{\partial s_k^{even}}, \frac{\partial}{\partial s_m^{even}} \right] = (m-k) \frac{\partial}{\partial s_{k+m}^{even}}, \quad k, m \geq -1.$$

We also have the following theorem for the Virasoro symmetries of the super tau-cover of the Principal Hierarchy.

Theorem 3.4 ([19]). *Let us denote*

$$\frac{\partial}{\partial s_m^{odd}} = \sum_{p \geq 0} (p + c_0) \tau_p \frac{\partial}{\partial \tau_{p+m}}, \quad m \geq -1,$$

where c_0 is an arbitrary constant. Then the following flows are symmetries of the super tau-cover of the Principal Hierarchy associated with a Frobenius manifold:

$$\begin{aligned}
 \frac{\partial f_{\alpha,p}}{\partial s_m} &= \frac{\partial f_{\alpha,p}}{\partial s_m^{even}} + \frac{\partial f_{\alpha,p}}{\partial s_m^{odd}}, \quad \frac{\partial \Phi_{\alpha,p}^n}{\partial s_m} = \frac{\partial}{\partial \tau_n} \left(\frac{\partial f_{\alpha,p}}{\partial s_m} \right), \\
 \frac{\partial v^\alpha}{\partial s_m} &= \frac{\partial v^\alpha}{\partial s_m^{even}} + \frac{\partial v^\alpha}{\partial s_m^{odd}}, \quad \frac{\partial \sigma_{\alpha,p}}{\partial s_m} = \frac{\partial}{\partial \tau_p} \left(\frac{\partial f_{\alpha,0}}{\partial s_m} \right).
 \end{aligned}$$

Moreover, these flows satisfy the commutation relation

$$\left[\frac{\partial}{\partial s_k}, \frac{\partial}{\partial s_m} \right] = (m-k) \frac{\partial}{\partial s_{k+m}}, \quad k, m \geq -1.$$

3.3. Formulation of the deformation problem. In this subsection, we first state the main problem of this paper, then explain the motivation and strategy of our proof.

From now on, we fix a semisimple Frobenius manifold M of dimension n and let (P_0, P_1) be a deformation of the bihamiltonian structure (2.12) with constant central invariants. Then (P_0, P_1) determines a unique deformation of the Principal Hierarchy associated with M . After a suitable Miura type transformation we may assume

$$P_0 = \frac{1}{2} \int \eta^{\alpha\beta} \sigma_\alpha \sigma_\beta^1 = [Z, P_1], \quad Z = \int \sigma_1,$$

and P_1 has no $\hat{\mathcal{F}}_2^2$ components. We also introduce odd variables $\sigma_{\alpha,m}$ for $m \geq 0$ as we explained in Sect. 2.3.

Our problem can be stated as follows: to find a unique derivation $X \in \text{Der}^\partial(\hat{\mathcal{A}})_{\geq 0}^0$ such that the flow defined by

$$(3.8) \quad \begin{aligned} \frac{\partial v_\lambda}{\partial s_2} = & a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\ & + X v_\lambda + \mathcal{L}_2 v_\lambda, \end{aligned}$$

$$(3.9) \quad \begin{aligned} \frac{\partial \sigma_{\lambda,0}}{\partial s_2} = & a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\ & + X \sigma_{\lambda,0} + M_\lambda^\zeta \sigma_{\zeta,2} + N_\lambda^\zeta \sigma_{\zeta,1} + \mathcal{L}_2 \sigma_{\lambda,0}, \end{aligned}$$

satisfies the conditions

$$(3.10) \quad \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial \tau_0} \right] = \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial \tau_1} \right] = 0,$$

and we require that the leading term of X is determined by Virasoro symmetry of the super tau-cover of the Principal Hierarchy. Here the operator \mathcal{L}_2 is given by

$$\mathcal{L}_2 = \mathcal{L}_2^{\text{even}} + \sum_{p \geq 0} (p + c_0) \tau_p \frac{\partial}{\partial \tau_{p+2}},$$

and $M_\lambda^\zeta, N_\lambda^\zeta \in \hat{\mathcal{A}}^0$ are some differential polynomials whose definitions will be given later.

If we find such a derivation X , then we can prove that the flow $\frac{\partial}{\partial s_2}$ is a symmetry of the deformed Principal Hierarchy, i.e.,

$$\left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] = 0, \quad \alpha = 1, \dots, n, \quad p \geq 0.$$

To prove this fact, we only need to show that

$$(3.11) \quad \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] \in \text{Der}(\hat{\mathcal{A}})_{\geq 2}^0,$$

since this commutator is a cocycle in $(\text{Der}^\partial(\hat{\mathcal{A}})_{\geq 2}^0, D_{P_0}, D_{P_1})$, and by using the property $BH_{\geq 2}^0(\text{Der}^\partial(\hat{\mathcal{A}}), D_{P_0^{[0]}}, D_{P_1^{[0]}}) = 0$ it is easy to conclude that this commutator vanishes. Note that a priori we have

$$\left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] v_\lambda \in \hat{\mathcal{A}}, \quad \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] \sigma_{\lambda,0} \in \hat{\mathcal{A}}^+,$$

so the condition (3.11) is actually a *locality condition*, i.e., it is equivalent to

$$(3.12) \quad \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] v_\lambda \in \hat{\mathcal{A}}, \quad \left[\frac{\partial}{\partial s_2}, \frac{\partial}{\partial t^{\alpha,p}} \right] \sigma_{\lambda,0} \in \hat{\mathcal{A}},$$

which is the condition we actually need to check.

Let us explain how to find a unique $X \in \text{Der}^\partial(\hat{\mathcal{A}})$ such that the conditions (3.10) hold true. To this end, we first rewrite the conditions (3.10) into the equations for X as follows:

$$(3.13) \quad \left[\frac{\partial}{\partial \tau_0}, X \right] = I_0, \quad \left[\frac{\partial}{\partial \tau_1}, X \right] = I_1,$$

where I_0 and I_1 are some derivations that will be given later. The uniqueness of X is a consequence of the equations (3.13) and the fact that $BH_{\geq 2}^0(\text{Der}^\partial(\hat{\mathcal{A}})) = 0$, since the leading term of X is fixed. We will prove the existence of X by using the following steps.

Step 1. To check the *locality condition* (3.12) and to prove that

$$(3.14) \quad I_0, I_1 \in \text{Der}(\hat{\mathcal{A}}).$$

Step 2. To check the *closedness condition*

$$(3.15) \quad \left[\frac{\partial}{\partial \tau_0}, I_0 \right] = 0.$$

When Step 2 is finished, we can find a derivation $X^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})$ by using the property $H_{>0}^1(\text{Der}^\partial(\hat{\mathcal{A}})) = 0$ such that

$$I_0 = \left[\frac{\partial}{\partial \tau_0}, X^\circ \right],$$

and the leading term of X° is given by the Virasoro symmetry of the super tau-cover of the Principal Hierarchy. We define $\mathcal{C} = X - X^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})_{\geq 2}^0$, then the equations (3.13) for X are transformed to the following equations for \mathcal{C} :

$$(3.16) \quad \left[\frac{\partial}{\partial \tau_0}, \mathcal{C} \right] = 0, \quad \left[\frac{\partial}{\partial \tau_1}, \mathcal{C} \right] = I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right].$$

Step 3. To check the closedness conditions

$$(3.17) \quad \left[\frac{\partial}{\partial \tau_0}, I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right] \right] = 0, \quad \left[\frac{\partial}{\partial \tau_1}, I_1 \right] = 0.$$

Step 4. To check that the differential degree 3 component of the derivation $I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right]$ vanishes in the cohomology group $BH_3^1(\text{Der}^\partial(\hat{\mathcal{A}}))$.

We call the above fact the *vanishing of the genus one obstruction* for the following reason. By using the first equation in (3.16) and the vanishing of $H_{\geq 2}^0(\text{Der}^\partial(\hat{\mathcal{A}}))$, we see that there exists a unique $\mathcal{T} \in \text{Der}^\partial(\hat{\mathcal{A}})_{\geq 1}^{-1}$ such that

$$\mathcal{C} = \left[\frac{\partial}{\partial \tau_0}, \mathcal{T} \right].$$

The derivation \mathcal{T} must also satisfy the second equation in (3.16)

$$(3.18) \quad \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, \mathcal{T} \right] \right] = I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right].$$

If the differential degree 3 component of the derivation $I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right]$ does not vanish in the cohomology group $BH_3^1(\text{Der}^\partial(\hat{A}))$, then such \mathcal{T} does not exist.

However if the genus one obstruction vanishes, the derivation \mathcal{T} exists upto differential degree 1. Then by using $BH_{\geq 4}^1(\text{Der}^\partial(\hat{A})) = 0$ and the closedness conditions (3.17), we can solve \mathcal{T} from (3.18) degree by degree and hence we can solve X such that (3.10) holds true.

Step 5. To lift the symmetry $\frac{\partial v_\lambda}{\partial s_2}$ to a symmetry of the tau-cover of the deformed Principal Hierarchy, and to define all the other flows $\frac{\partial}{\partial s_m}$ for $m \geq 0$ for the Virasoro symmetries. Note that the symmetry $\frac{\partial}{\partial s_{-1}}$ is constructed in [9]. We remark that we can also lift the symmetry (3.8) and (3.9) to a symmetry of the super tau-cover of the deformed Principal Hierarchy, but it is not necessary for the consideration for our problem.

In the remaining part of this subsection, we explain how the equations (3.8) and (3.9) are derived from the equation (1.1) of the main theorem. Let \mathcal{Z} be a tau-function of the tau-cover (2.28) of the deformed Principal Hierarchy, i.e.,

$$(3.19) \quad f_{\alpha,p} = \frac{\partial \log \mathcal{Z}}{\partial t^{\alpha,p}}, \quad w^\alpha = \eta^{\alpha\beta} \frac{\partial^2 \log \mathcal{Z}}{\partial t^{1,0} \partial t^{\beta,0}}$$

give a solution of the tau-cover (2.28). Our goal is to find a symmetry of the following form

$$(3.20) \quad \frac{\partial \mathcal{Z}}{\partial s_2} = L_2^{\text{even}} \mathcal{Z} + O_2 \mathcal{Z},$$

where L_2^{even} is the operator (3.4) and O_2 is a differential polynomial. If we assume that this is indeed a symmetry, then by using (3.19) we obtain the flow

$$\begin{aligned} \frac{\partial f_{\lambda,k}}{\partial s_2} &= a^{\alpha\beta} \left(f_{\alpha,1} \Omega_{\beta,0;\lambda,k} + f_{\beta,0} \Omega_{\alpha,1;\lambda,k} + \frac{\partial \Omega_{\alpha,1;\beta,0}}{\partial t^{\lambda,k}} \right) \\ &\quad + b^{\alpha\beta} \left(f_{\alpha,0} \Omega_{\beta,0;\lambda,k} + f_{\beta,0} \Omega_{\alpha,0;\lambda,k} + \frac{\partial \Omega_{\alpha,0;\beta,0}}{\partial t^{\lambda,k}} \right) \\ &\quad + b_{2;\lambda,k}^{\beta,q} f_{\beta,q} + c_{2;\lambda,k;\beta,q} t^{\beta,q} + c_{2;\beta,q;\lambda,k} t^{\beta,q} + \frac{\partial O_2}{\partial t^{\lambda,k}} + \mathcal{L}_2^{\text{even}} f_{\lambda,k}; \\ \frac{\partial w_\lambda}{\partial s_2} &= a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial w_\lambda}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial w_\lambda}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial w_\lambda}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial w_\lambda}{\partial t^{\beta,0}} \right) \\ &\quad + W_\lambda + \mathcal{L}_2^{\text{even}} w_\lambda, \end{aligned}$$

here W_λ are some differential polynomials. Now recall that v_λ and w_ζ are related by a Miura type transformation, hence by using the equation

$$\frac{\partial v_\lambda}{\partial s_2} = \sum_{s \geq 0} \partial_x^s \frac{\partial w_\zeta}{\partial s_2} \frac{\partial v_\lambda}{\partial w_\zeta^{(s)}}$$

we know that there exist differential polynomials $X_\lambda^0 \in \hat{\mathcal{A}}^0$ such that

$$(3.21) \quad \frac{\partial v_\lambda}{\partial s_2} = a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) + X_\lambda^0 + \mathcal{L}_2^{even} v_\lambda.$$

To write down the evolutions of the odd variables, we must replace \mathcal{L}_2^{even} by \mathcal{L}_2 . By using (2.32) it is easy to see that

$$\frac{\partial \mathbf{f}_0}{\partial \tau_0} = \mathbf{A} \sigma_0,$$

where $\mathbf{f}_0 = (f_{1,0}, \dots, f_{n,0})^T$, $\sigma_0 = (\sigma_{1,0}, \dots, \sigma_{n,0})^T$ and \mathbf{A} is a matrix of differential operator of the form

$$\mathbf{A} = \sum_{g \geq 0} \sum_{k=0}^{2g} \mathbf{A}_{g,k} \partial_x^k, \quad \mathbf{A}_{g,k} \in M_n(\hat{\mathcal{A}}_{2g-k}^0).$$

Note that $\mathbf{A}_{0,0}$ is the identity matrix, therefore \mathbf{A} is invertible as a differential operator, i.e., there exists $\mathbf{B} = \sum_{g \geq 0} \sum_{k=0}^{2g} \mathbf{B}_{g,k} \partial_x^k$ such that $\mathbf{A}\mathbf{B} = \mathbf{B}\mathbf{A} = \mathbf{I}$, and in particular $\mathbf{B}_{0,0}$ is the identity matrix. Thus we can represent the odd variable $\sigma_{\lambda,0}$ in the form

$$\sigma_{\lambda,0} = \frac{\partial f_{\lambda,0}}{\partial \tau_0} + \sum_{g \geq 1} \sum_{k=0}^{2g} \mathbf{B}_{g,k} \partial_x^k \frac{\partial f_{\lambda,0}}{\partial \tau_0}.$$

This identity leads us to the following definition of the evolutions of the odd variables $\sigma_{\lambda,0}$ along the flow $\frac{\partial}{\partial s_2}$:

$$\frac{\partial \sigma_{\lambda,0}}{\partial s_2} = \frac{\partial}{\partial \tau_0} \frac{\partial f_{\lambda,0}}{\partial s_2} + \sum_{g \geq 1} \sum_{k=0}^{2g} \frac{\partial}{\partial s_2} \left(\mathbf{B}_{g,k} \partial_x^k \right) \frac{\partial f_{\lambda,0}}{\partial \tau_0} + \sum_{g \geq 1} \sum_{k=0}^{2g} \mathbf{B}_{g,k} \partial_x^k \frac{\partial}{\partial \tau_0} \frac{\partial f_{\lambda,0}}{\partial s_2}.$$

By using Proposition 2.24 and Lemma 2.23, we can see that the odd variables $\Phi_{\alpha,p}^m$ appearing in $\frac{\partial}{\partial \tau_0} \frac{\partial f_{\lambda,0}}{\partial s_2}$ can be represented by elements of $\hat{\mathcal{A}}^+$. So there exist differential polynomials $M_\lambda^\zeta, N_\lambda^\zeta \in \hat{\mathcal{A}}^0$ and $X_\lambda^1 \in \hat{\mathcal{A}}^1$ such that

$$\begin{aligned} \frac{\partial \sigma_{\lambda,0}}{\partial s_2} &= a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\ &\quad + X_\lambda^1 + M_\lambda^\zeta \sigma_{\zeta,2} + N_\lambda^\zeta \sigma_{\zeta,1} + \mathcal{L}_2 \sigma_{\lambda,0}. \end{aligned}$$

Finally we define a derivation $X \in \text{Der}^\partial(\hat{\mathcal{A}})^0$ such that $Xv_\lambda = X_\lambda^0$, where X_λ^0 is the differential polynomial introduced in (3.21), and $X\sigma_{\lambda,0} = X_\lambda^1$. Therefore our problem of finding such a derivation X is a necessary condition of the main theorem.

3.4. Example: one-dimensional Frobenius manifold. In this subsection we present an example to illustrate how the general framework described in the previous subsection works. We consider the one-dimensional Frobenius manifold M , it has the following potential and Euler vector field:

$$F = \frac{1}{6} v^3, \quad E = v \partial_v.$$

Due to the dimension reason, we will omit the Greek indices, for example, we will use $v^{(s)}$ and σ_m^s instead of $v^{1,s}$ and $\sigma_{1,m}^s$. The Principal Hierarchy associated with M is the Riemann hierarchy

$$\frac{\partial v}{\partial t_p} = \frac{v^p}{p!} v_x, \quad p \geq 0,$$

whose bihamiltonian structure is given by

$$P_0^{[0]} = \frac{1}{2} \int \sigma_0 \sigma_0^1, \quad P_1^{[0]} = \frac{1}{2} \int v \sigma_0 \sigma_0^1.$$

It is proved in [7, 21] that every deformation (P_0, P_1) of $(P_0^{[0]}, P_1^{[0]})$ with a constant central invariant is equivalent to the bihamiltonian structure given by

$$P_0 = \frac{1}{2} \int \sigma_0 \sigma_0^1, \quad P_1 = \frac{1}{2} \int v \sigma_0 \sigma_0^1 + \varepsilon^2 c \sigma_0 \sigma_0^3$$

via a certain Miura type transformation. Here the dispersion parameter ε is added for clearness, and the central invariant of (P_0, P_1) is $\frac{c}{3}$. In particular, when $c = \frac{1}{8}$ the corresponding deformed Riemann Hierarchy is the KdV hierarchy that controls the 2D gravity [16, 25].

We introduce the odd variables σ_m satisfying the recursion relations

$$\sigma_{m+1}^1 = v \sigma_m^1 + \frac{1}{2} v_x \sigma_m + \varepsilon^2 c \sigma_m^3, \quad m \geq 0.$$

The following useful results can be obtained directly:

$$\begin{aligned} h_1 &= \frac{v^2}{2} + \frac{2}{3} \varepsilon^2 c v_{xx}, \quad h_2 = \frac{v^3}{6} + \varepsilon^2 c \left(\frac{1}{3} v_x^2 + \frac{2}{3} v v_{xx} \right) + \frac{4}{15} \varepsilon^4 c^2 v^{(4)}; \\ \frac{\partial v}{\partial t_1} &= v v_x + \frac{2}{3} \varepsilon^2 c v^{(3)}, \quad \frac{\partial \sigma_0}{\partial t_1} = v \sigma_0^1 + \frac{2}{3} \varepsilon^2 c \sigma_0^3; \\ \frac{\partial v}{\partial t_2} &= \frac{1}{2} v^2 v_x + \varepsilon^2 c \left(\frac{4}{3} v_x v_{xx} + \frac{2}{3} v v^{(3)} \right) + \frac{4}{15} \varepsilon^4 c^2 v^{(5)}, \\ \frac{\partial \sigma_0}{\partial t_2} &= \frac{1}{2} v^2 \sigma_0^1 + \varepsilon^2 c \left(\frac{2}{3} v_{xx} \sigma_0^1 + \frac{2}{3} v_x \sigma_0^2 + \frac{2}{3} v \sigma_0^3 \right) + \frac{4}{15} \varepsilon^4 c^2 \sigma_0^5; \\ \varepsilon \frac{\partial f_0}{\partial \tau_0} &= \sigma_0, \quad \varepsilon \frac{\partial f_1}{\partial \tau_0} = 2 \sigma_1 - v \sigma_0 - \frac{4}{3} \varepsilon^2 c \sigma_0^2; \\ \varepsilon \frac{\partial f_2}{\partial \tau_0} &= \frac{4}{3} \sigma_2 - \frac{2}{3} v \sigma_1 - \frac{1}{6} v^2 \sigma_0 - \varepsilon^2 c \left(\frac{2}{3} v_{xx} \sigma_0 + \frac{4}{3} v_x \sigma_0^2 + \frac{4}{3} v \sigma_0^3 \right) - \frac{16}{15} \varepsilon^4 c^2 \sigma_0^4. \end{aligned}$$

We first note that

$$L_2 = \frac{3}{8} \varepsilon^2 \frac{\partial^2}{\partial t_1 \partial t_0} + \mathcal{L}_2, \quad \mathcal{L}_2 = \sum_{p \geq 0} \frac{\Gamma(\frac{7}{2} + p)}{\Gamma(\frac{1}{2} + p)} t_p \frac{\partial}{\partial t_{p+2}} + (p + c_0) \tau_p \frac{\partial}{\partial \tau_{p+2}}.$$

Then the equations (3.8) and (3.9) for this example have the following form:

$$\begin{aligned}\frac{\partial v}{\partial s_2} &= \frac{3}{8}\varepsilon \left(v_x f_1 + \frac{\partial v}{\partial t_1} f_0 \right) + Xv + \mathcal{L}_2 v, \\ \frac{\partial \sigma_0}{\partial s_2} &= \frac{3}{8}\varepsilon \left(\sigma_0^1 f_1 + \frac{\partial \sigma_0}{\partial t_1} f_0 \right) + \left(\frac{5}{2} + c_0 \right) \sigma_2 - \frac{v}{2} \sigma_1 + X\sigma_0 + \mathcal{L}_2 \sigma_0.\end{aligned}$$

We are to find the derivation $X \in \text{Der}^\partial(\hat{\mathcal{A}})^0$ such that the flow $\frac{\partial}{\partial s_2}$ commutes with $\frac{\partial}{\partial \tau_0}$ and $\frac{\partial}{\partial \tau_1}$. These conditions yield the following equations for X :

$$\left[\frac{\partial}{\partial \tau_0}, X \right] = I_0, \quad \left[\frac{\partial}{\partial \tau_1}, X \right] = I_1,$$

where the derivations I_0 and I_1 are defined by

$$(3.22) \quad I_0 v = vv_x \sigma_0 + \frac{7}{2} v^2 \sigma_0^1 + \varepsilon^2 c \left(v^{(3)} \sigma_0 + \frac{13}{2} v_{xx} \sigma_0^1 + 8v_x \sigma_0^2 + 6v \sigma_0^3 \right) + 3\varepsilon^4 c^2 \sigma_0^5,$$

$$(3.23) \quad I_0 \sigma_0 = v \sigma_0 \sigma_0^1 - \varepsilon^2 c \left(\frac{1}{2} \sigma_0^1 \sigma_0^2 - \sigma_0 \sigma_0^3 \right),$$

$$\begin{aligned}(3.24) \quad I_1 v &= \frac{5}{4} v^2 v_x \sigma_0 + \frac{5}{2} v^3 \sigma_0^1 + \varepsilon^2 c \left(\frac{7}{2} v_x v_{xx} \sigma_0 + 2v v^{(3)} \sigma_0 + \frac{45}{4} v_x^2 \sigma_0^1 + \frac{31}{2} v v_{xx} \sigma_0^1 \right) \\ &\quad + \varepsilon^2 c \left(26v v_x \sigma_0^2 + \frac{19}{2} v^2 \sigma_0^3 \right) + \varepsilon^4 c^2 \left(v^{(5)} \sigma_0 + \frac{17}{2} v^{(4)} \sigma_0^1 + \frac{45}{2} v^{(3)} \sigma_0^2 \right) \\ &\quad + \varepsilon^4 c^2 \left(\frac{59}{2} v_{xx} \sigma_0^3 + \frac{43}{2} v_x \sigma_0^4 + 9v \sigma_0^5 \right) + 3\varepsilon^6 c^3 \sigma_0^7,\end{aligned}$$

$$\begin{aligned}(3.25) \quad I_1 \sigma_0 &= \frac{5}{4} v^2 \sigma_0 \sigma_0^1 + \varepsilon^2 c \left(\frac{5}{2} v_{xx} \sigma_0 \sigma_0^1 + \frac{5}{2} v_x \sigma_0 \sigma_0^2 - \frac{1}{2} v \sigma_0^1 \sigma_0^2 + 2v \sigma_0 \sigma_0^3 \right) \\ &\quad - \varepsilon^4 c^2 \left(\frac{1}{2} \sigma_0^1 \sigma_0^4 - \sigma_0 \sigma_0^5 \right).\end{aligned}$$

It follows from (3.22) and (3.23) that we can choose $X^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})$ such that $[\frac{\partial}{\partial \tau_0}, X^\circ] = I_0$, whose actions on v and σ_0 are given by

$$\begin{aligned}X^\circ v &= v^3 + \varepsilon^2 c \left(\frac{5}{4} v_x^2 + 3v v_{xx} \right) + \varepsilon^4 c^2 v^{(4)}, \\ X^\circ \sigma_0 &= -\frac{1}{2} v^2 \sigma_0 - \varepsilon^2 c \left(v_{xx} \sigma_0 + \frac{5}{2} v_x \sigma_0^1 + 3v \sigma_0^2 \right) - 2\varepsilon^4 c^2 \sigma_0^4.\end{aligned}$$

Then according to the general discussion given in the previous subsection, the derivation $\mathcal{C} = X - X^\circ$ satisfies the equations

$$\left[\frac{\partial}{\partial \tau_0}, \mathcal{C} \right] = 0, \quad \left[\frac{\partial}{\partial \tau_1}, \mathcal{C} \right] = I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ \right].$$

Finally we can solve the above equations and obtain the unique derivation \mathcal{C} that is defined by

$$\begin{aligned}\mathcal{C}v &= \varepsilon^2 c (3v_x^2 + 3vv_{xx}) + 2\varepsilon^4 c^2 v^{(4)}, \\ \mathcal{C}\sigma_0 &= \varepsilon^2 c (3v_x\sigma_0^1 + v\sigma_0^2) + 2\varepsilon^4 c^2 \sigma_0^4.\end{aligned}$$

By forgetting all the odd variables we obtain the following symmetry for the deformed Riemann hierarchy:

$$\frac{\partial v}{\partial s_2} = \frac{3}{8}\varepsilon \left(v_x f_1 + \frac{\partial v}{\partial t_1} f_0 \right) + v^3 + \varepsilon^2 c \left(\frac{17}{4}v_x^2 + 6vv_{xx} \right) + 3\varepsilon^4 c^2 v^{(4)} + \mathcal{L}_2^{even} v.$$

It is easy to check that the action of this symmetry on the tau function \mathcal{Z} of the deformed Riemann hierarchy can be represented by

$$(3.26) \quad \frac{\partial \mathcal{Z}}{\partial s_2} = L_2^{even} \mathcal{Z} + \left(3c - \frac{3}{8} \right) \left(\frac{v^2}{2} + \frac{2}{3}\varepsilon^2 c v_{xx} \right) \mathcal{Z}.$$

In particular, when $c = \frac{1}{8}$, this symmetry is given by a linear action on \mathcal{Z} .

4. DEFORMATION OF VIRASORO SYMMETRIES: EXISTENCE AND UNIQUENESS

In this section, we present details of the proof of the main theorem following the framework described in Sect. 3.3.

4.1. Locality conditions. We start by verifying the locality conditions (3.14).

Let us first find the differential polynomials M_λ^ζ and N_λ^ζ in (3.9) to ensure that I_0 is local. By using the fact (2.32) and the equations (3.8) and (3.9), we arrive at

$$(4.1) \quad \left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial s_2} \right] v_\lambda = c_0 \sigma_{\lambda,2}^1 + a^{\alpha\beta} \frac{\partial f_{\alpha,1}}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} - \partial_x \left(M_\lambda^\zeta \sigma_{\zeta,2} + N_\lambda^\zeta \sigma_{\zeta,1} \right) + loc.$$

Here and henceforth we will use *loc* to denote the local terms, i.e., terms belonging to $\hat{\mathcal{A}}$. We need to find M_λ^ζ and N_λ^ζ such that the right hand side of (4.1) is local.

Lemma 4.1. *The following identities hold true:*

$$\frac{\partial}{\partial \sigma_{\beta,0}} \frac{\delta P_1}{\delta \sigma_{\lambda,0}} = \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial v^\lambda}{\partial t^{\alpha,0}}, \quad \frac{\partial}{\partial \sigma_{\beta,0}} \frac{\delta P_1}{\delta v^\lambda} = \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}}.$$

Proof. By using the identity (2.25) and the fact that $H_{\alpha,-1}$ are Casimirs of P_0 , we obtain the identity

$$[P_1, H_{\alpha,-1}] = \left(\frac{1}{2} + \mu_\alpha \right) [P_0, H_{\alpha,0}].$$

By taking the variational derivatives of both sides of the identity above and by using (2.3) and (2.4), it is easy to prove the desired identities. The lemma is proved. \square

Lemma 4.2. *We have the following relation:*

$$(4.2) \quad \sigma_{\lambda,2}^1 = \left(\frac{1}{2} + \mu_\varepsilon \right) \eta^{\zeta\varepsilon} \frac{\partial v_\lambda}{\partial t^{\varepsilon,0}} \sigma_{\zeta,1} + loc.$$

Proof. Let us denote by \mathcal{P}_1 the Hamiltonian operator of P_1 and represent it in the form

$$\mathcal{P}_1 = \sum_{k \geq 0} \mathcal{P}_{1,k} \partial_x^k, \quad \mathcal{P}_{1,k} \in M_n(\hat{\mathcal{A}}).$$

Then according to the recursion relation

$$\eta^{\gamma\lambda} \sigma_{\lambda,2}^1 = \mathcal{P}_1^{\gamma\lambda} \sigma_{\lambda,1} = \sum_{k \geq 0} \mathcal{P}_{1,k}^{\gamma\lambda} \sigma_{\lambda,1}^k,$$

it is easy to see that

$$\sigma_{\lambda,2}^1 = \eta_{\lambda\gamma} \mathcal{P}_{1,0}^{\gamma\zeta} \sigma_{\zeta,1} + loc.$$

Now it follows from the definition of the Hamiltonian operator that

$$\mathcal{P}_{1,0}^{\gamma\zeta} = \frac{\partial}{\partial \sigma_{\zeta,0}} \frac{\delta P_1}{\delta \sigma_{\gamma,0}}.$$

Therefore by using Lemma 4.1 we arrive at the identity (4.2) and the lemma is proved. \square

Proposition 4.3. *There exist unique differential polynomials M_λ^ζ and N_λ^ζ such that (4.1) is local. More explicitly, we have*

$$M_\lambda^\zeta = \left(\frac{5}{2} + \mu_\lambda + c_0 \right) \delta_\lambda^\zeta, \quad \partial_x N_\lambda^\zeta = \left(\frac{1}{2} + \mu_\alpha \right) \eta^{\alpha\zeta} (\mu_\zeta - \mu_\lambda - 1) \frac{\partial v_\lambda}{\partial t^{\alpha,0}},$$

here c_0 is the arbitrary constant appearing in the operator \mathcal{L}_2 .

Proof. To ensure the vanishing of the coefficient of $\sigma_{\zeta,2}$ in the right hand side of (4.1), M_λ^ζ should be a constant, so it is determined by the leading terms of the right hand sides of (3.8) and (3.9), which are fixed by the Virasoro symmetry $\frac{\partial}{\partial s_2}$ of the super tau-cover of the Principal Hierarchy. Hence we obtain

$$M_\lambda^\zeta = \left(\frac{5}{2} + \mu_\lambda + c_0 \right) \delta_\lambda^\zeta.$$

By using Proposition 2.24 we have

$$(4.3) \quad \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial f_{\alpha,1}}{\partial \tau_0} = \sigma_{\alpha,1} + loc.$$

Then by using this equation, the recursion relations (2.17) and the identity (4.2), we see that the vanishing of the coefficient of $\sigma_{\zeta,1}$ in the right hand side of (4.1) gives the equation

$$(4.4) \quad \partial_x N_\lambda^\zeta = \left(\frac{1}{2} + \mu_\alpha \right) \eta^{\alpha\zeta} (\mu_\zeta - \mu_\lambda - 1) \frac{\partial v_\lambda}{\partial t^{\alpha,0}}.$$

Both sides of (4.4) are total x -derivatives, so we can integrate (4.4) to obtain N_λ^ζ upto a constant, which has differential degree zero. Since the differential degree zero part of N_λ^ζ is fixed by genus zero Virasoro symmetry, N_λ^ζ is uniquely determined. The proposition is proved. \square

Now by a direct computation of the condition

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial s_2} \right] = 0,$$

we obtain the following explicit expression for I_0 which is defined by (3.13):

$$(4.5) \quad I_0 v_\lambda = a^{\alpha\beta} \left(f'_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + f'_{\alpha,1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) - a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\ + b^{\alpha\beta} \left(f'_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + f'_{\alpha,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\ + \left(\frac{5}{2} + \mu_\lambda \right) \sigma_{\lambda,2}^1 + W \sigma_{\lambda,0} + \partial_x \left(N_\lambda^\zeta \sigma_{\zeta,1} \right),$$

$$(4.6) \quad I_0 \sigma_{\lambda,0} = -a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\ + \left(\frac{5}{2} + \mu_\lambda \right) \frac{\partial \sigma_{\lambda,0}}{\partial \tau_2} - \frac{\partial}{\partial \tau_0} \left(N_\lambda^\zeta \sigma_{\zeta,1} \right),$$

here and henceforth we will use W to denote the coefficient of $t^{1,0}$ of the operator \mathcal{L}_2 , more explicitly, we have

$$W = \left(\frac{1}{2} + \mu_1 \right) \left(\frac{3}{2} + \mu_1 \right) \left(\frac{5}{2} + \mu_1 \right) \frac{\partial}{\partial t^{1,2}} \\ + \sum_{k=1}^2 \left(3 \left(\frac{1}{2} + \mu_1 \right)^2 + 6 \left(\frac{1}{2} + \mu_1 \right) + 2 \right) (R_k)_1^\beta \frac{\partial}{\partial t^{\beta,2-k}} \\ + \left(\frac{9}{2} + 3\mu_1 \right) (R_{2,2})_1^\beta \frac{\partial}{\partial t^{\beta,0}}.$$

Proposition 4.4. *The derivation I_0 given by (4.5), (4.6) is local.*

Proof. The locality of (4.5) is equivalent to the definition of N_λ^ζ and we only need to check the locality of (4.6). By using the definition of the odd flows we know that

$$\frac{\partial \sigma_{\lambda,0}}{\partial \tau_2} = T_{0,2} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} = \sum_{s \geq 0} \sigma_{\beta,1}^s \frac{\partial}{\partial \sigma_{\beta,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1}.$$

Therefore it follows from Lemma 4.1 that

$$\frac{\partial \sigma_{\lambda,0}}{\partial \tau_2} = \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + loc.$$

Then the locality of (4.6) is proved by using (4.3) and the following obvious fact:

$$\frac{\partial N_\lambda^\zeta}{\partial \tau_0} = \left(\frac{1}{2} + \mu_\alpha \right) \eta^{\alpha\zeta} (\mu_\zeta - \mu_\lambda - 1) \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}}.$$

The proposition is proved. \square

Next let us prove the locality of I_1 . Similar to the expression (4.5) and (4.6), we can write down the explicit expression for I_1 , which can be found in the next subsection. However the locality of I_1 can not be derived from this expression directly, so we turn to prove the following equivalent conditions:

$$(4.7) \quad \left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] v_\lambda \in \hat{\mathcal{A}}, \quad \left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] \sigma_{\lambda,0} \in \hat{\mathcal{A}}.$$

By a direct computation, we have

$$(4.8) \quad \begin{aligned} \left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] v_\lambda &= a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\ &\quad + b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) + (1+c_0) \frac{\partial v_\lambda}{\partial \tau_3} \\ &\quad - \sum_{s \geq 0} \left(\left(\frac{5}{2} + c_0 + \mu_\zeta \right) \sigma_{\zeta,2}^s + \partial_x^s (N_\zeta^\varepsilon \sigma_{\zeta,1}) \right) \frac{\partial}{\partial \sigma_{\zeta,0}^s} \frac{\partial v_\lambda}{\partial \tau_1} + loc. \end{aligned}$$

Lemma 4.5. *There exists a unique differential polynomial Z_α^β such that*

$$\left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial f_{\alpha,1}}{\partial \tau_1} = \sigma_{\alpha,2} + Z_\alpha^\beta \sigma_{\beta,1} + loc,$$

where Z_α^β satisfies the following equation

$$(4.9) \quad \partial_x Z_\alpha^\beta = - \left(\frac{1}{2} + \mu_\varepsilon \right) \eta^{\beta\varepsilon} \frac{\partial v_\alpha}{\partial t^{\varepsilon,0}}.$$

Proof. The existence and uniqueness of Z_α^β can be obtained from Proposition 2.24, hence we only need to derive (4.9), which can be obtained by using the identity (4.2) and the fact that $\partial_x (\sigma_{\alpha,2} + Z_\alpha^\beta \sigma_{\beta,1})$ is local. The lemma is proved. \square

The following equation can be derived from the definition of the odd flows and the recursion relation (2.17):

$$\frac{\partial v_\lambda}{\partial \tau_3} = \sum_{s \geq 0} \sigma_{\zeta,2}^s \frac{\partial}{\partial \sigma_{\zeta,0}^s} \frac{\partial v_\lambda}{\partial \tau_1}.$$

This equation together with Lemma 4.5 and the identity (4.2) enables us to rewrite (4.8) in the form

$$\left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] v_\lambda = U_\lambda^\zeta \sigma_{\zeta,2} + V_\lambda^\zeta \sigma_{\zeta,1} + loc,$$

where the differential polynomials U_λ^ζ and V_λ^ζ are given by

$$\begin{aligned} U_\lambda^\zeta &= \eta^{\zeta\beta} \left(\frac{1}{2} + \mu_\beta \right) \left(\frac{3}{2} + \mu_\zeta \right) \frac{\partial v_\lambda}{\partial t^{\beta,0}} - \left(\frac{3}{2} + \mu_\zeta \right) \frac{\partial}{\partial \sigma_{\zeta,0}} \frac{\partial v_\lambda}{\partial \tau_1}, \\ V_\lambda^\zeta &= a^{\alpha\zeta} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + (b^{\alpha\zeta} + b^{\zeta\alpha}) \frac{\partial v_\lambda}{\partial t^{\alpha,0}} - \sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial v_\lambda}{\partial \tau_1}. \end{aligned}$$

The vanishing of U_λ^ζ is a consequence of Lemma 4.1 and the vanishing of V_λ^ζ is due to the following lemma.

Lemma 4.6. *We have $V_\lambda^\zeta = 0$, that is*

$$\sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial v_\lambda}{\partial \tau_1} = a^{\alpha\zeta} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + (b^{\alpha\zeta} + b^{\zeta\alpha}) \frac{\partial v_\lambda}{\partial t^{\alpha,0}}.$$

Proof. Consider the functional

$$F^\zeta = \eta^{\alpha\zeta} \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{1}{2} + \mu_\zeta \right) \int h_{\alpha,1},$$

then by using (4.4) and (4.9), we can check that there exist constants C_λ^ζ such that

$$N_\lambda^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\lambda^\zeta = \frac{\delta F^\zeta}{\delta v_\lambda} + C_\lambda^\zeta.$$

Then by using (2.4), we obtain

$$\sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial v_\lambda}{\partial \tau_1} = -\frac{\delta}{\delta \sigma_{\lambda,0}} [F^\zeta, P_1] + C_\gamma^\zeta \frac{\partial}{\partial \sigma_{\gamma,0}} \frac{\partial v_\lambda}{\partial \tau_1}.$$

Using the bihamiltonian recursion relations (2.25) and Lemma 4.1 we see that the right hand side of the above equation is a linear combination of the flows $\frac{\partial v_\lambda}{\partial t^{\alpha,p}}$, hence so is V_λ^ζ . The genus zero Virasoro symmetry implies that $V_\lambda^\zeta \in \hat{\mathcal{A}}_{\geq 2}^0$, so by using the theory of bihamiltonian cohomology [7], we know that V_λ^ζ must vanish. The lemma is proved. \square

We have verified the first relation in (4.7) and let us proceed to prove the second one. We have

$$\begin{aligned} (4.10) \quad \left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] \sigma_{\lambda,0} &= a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\ &\quad + b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\ &\quad + \left(\frac{5}{2} + c_0 + \mu_\lambda \right) \frac{\partial \sigma_{\lambda,2}}{\partial \tau_1} + \frac{\partial N_\lambda^\zeta}{\partial \tau_1} \sigma_{\zeta,1} + (1 + c_0) \frac{\partial \sigma_{\lambda,0}}{\partial \tau_3} \\ &\quad - \sum_{s \geq 0} \left(\left(\frac{5}{2} + c_0 + \mu_\zeta \right) \sigma_{\zeta,2}^s + \partial_x^s (N_\zeta^\varepsilon \sigma_{\zeta,1}) \right) \frac{\partial}{\partial \sigma_{\zeta,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} + loc. \end{aligned}$$

By using the equations

$$\frac{\partial \sigma_{\lambda,0}}{\partial \tau_3} = \sum_{s \geq 0} \sigma_{\gamma,2}^s \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} + \frac{\partial \sigma_{\lambda,1}}{\partial \tau_2}, \quad \frac{\partial \sigma_{\lambda,1}}{\partial t^{\alpha,p}} = \frac{\partial}{\partial \tau_1} \frac{\delta h_{\alpha,p+1}}{\delta v^\lambda},$$

$$\frac{\partial \sigma_{\lambda,1}}{\partial \tau_2} = \sigma_{\zeta,1} \left(\frac{1}{2} + \mu_\varepsilon \right) \eta^{\varepsilon\zeta} \frac{\partial}{\partial \tau_1} \frac{\delta h_{\varepsilon,1}}{\delta v^\lambda} + loc$$

we obtain that

$$\left[\frac{\partial}{\partial \tau_1}, \frac{\partial}{\partial s_2} \right] \sigma_{\lambda,0} = \sigma_{\zeta,2} \tilde{U}_\lambda^\zeta + \sigma_{\zeta,1} \tilde{V}_\lambda^\zeta + loc,$$

where the differential polynomials $\tilde{U}_\lambda^\zeta, \tilde{V}_\lambda^\zeta$ are given by

$$\begin{aligned}\tilde{U}_\lambda^\zeta &= \eta^{\zeta\beta} \left(\frac{1}{2} + \mu_\beta \right) \left(\frac{3}{2} + \mu_\zeta \right) \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} - \left(\frac{3}{2} + \mu_\zeta \right) \frac{\partial}{\partial \sigma_{\zeta,0}} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1}, \\ \tilde{V}_\lambda^\zeta &= a_{\alpha\zeta} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + (b_{\alpha\zeta} + b_{\zeta\alpha}) \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} - \frac{\partial}{\partial \tau_1} \left(N_\lambda^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\lambda^\zeta \right) \\ &\quad - \sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1}.\end{aligned}$$

These differential polynomials actually vanish, the proof of which is similar to the one for the vanishing of U_λ^ζ and V_λ^ζ given above. Hence the locality condition (3.14) is verified.

Finally let us consider the locality condition (3.12). The first condition

$$\left[\frac{\partial}{\partial t^{\delta,j}}, \frac{\partial}{\partial s_2} \right] v_\lambda \in \hat{\mathcal{A}}$$

is obvious. On the other hand, we have

$$\begin{aligned}(4.11) \quad \left[\frac{\partial}{\partial t^{\delta,j}}, \frac{\partial}{\partial s_2} \right] \sigma_{\lambda,0} &= \left(\frac{5}{2} + c_0 + \mu_\lambda \right) \frac{\partial \sigma_{\lambda,2}}{\partial t^{\delta,j}} + \frac{\partial N_\lambda^\zeta}{\partial t^{\delta,j}} \sigma_{\zeta,1} \\ &\quad - \sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta \sigma_{\zeta,1} + \left(\frac{5}{2} + c_0 + \mu_\gamma \right) \sigma_{\gamma,2} \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\delta,j}} + loc.\end{aligned}$$

Proposition 4.7. *The right hand side of the equation (4.11) is local.*

Proof. By using the following identity proved in [22]

$$\frac{\partial}{\partial \sigma_{\beta,0}} \frac{\delta}{\delta v^\alpha} = \frac{\delta}{\delta v^\alpha} \frac{\delta}{\delta \sigma_{\beta,0}}$$

we immediately obtain that

$$\frac{\partial}{\partial \sigma_{\beta,0}} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,p}} = \frac{\partial}{\partial \sigma_{\beta,0}} \frac{\delta X_{\alpha,p}}{\delta v^\lambda} = 0.$$

This means that the flows $\frac{\partial \sigma_{\lambda,2}}{\partial t^{\delta,j}}$ can be written as

$$\frac{\partial \sigma_{\lambda,2}}{\partial t^{\delta,j}} = T_2 \frac{\partial \sigma_{\lambda,0}}{\partial t^{\delta,j}} = \sum_{s \geq 0} \sigma_{\beta,2}^s \frac{\partial}{\partial \sigma_{\beta,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\delta,j}} = B_\lambda^\zeta \sigma_{\zeta,1} + loc,$$

where B_λ^ζ are certain differential polynomials. By using the identity (4.2), we know that

$$B_\lambda^\zeta = - \frac{\partial Z_\lambda^\zeta}{\partial t^{\delta,j}}.$$

By using (4.2) again, we can represent the right hand side of (4.11) in the form

$$V_\lambda^\zeta \sigma_{\zeta,1} + loc,$$

where $V_{\delta,j;\lambda}^\zeta$ is the following differential polynomial:

$$\begin{aligned} V_{\delta,j;\lambda}^\zeta &= \frac{\partial}{\partial t^{\delta,j}} \left(N_\lambda^\zeta - \left(\frac{5}{2} + c_0 + \mu_\gamma \right) Z_\lambda^\zeta \right) \\ &\quad - \sum_{s \geq 0} \partial_x^s \left(N_\gamma^\zeta - \left(\frac{5}{2} + c_0 + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\delta,j}}. \end{aligned}$$

Define the following functional

$$F^\zeta = \eta^{\alpha\zeta} \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{3}{2} + c_0 + \mu_\zeta \right) \int h_{\alpha,1},$$

then by applying (2.3) we have the following identity:

$$V_{\delta,j;\lambda}^\zeta = \frac{\delta}{\delta v^\lambda} [F^\zeta, X_{\delta,j}],$$

here $X_{\delta,j}$ is the vector field defined in (2.22). Since the functional F^ζ is a conserved quantity of the deformed Principal hierarchy, hence $[F^\zeta, X_{\delta,j}] = 0$. The proposition is proved. \square

4.2. Closedness conditions. In this subsection we prove the closedness condition (3.15) and (3.17). The verification of (3.15) is straightforward by using the explicit expression (4.5) and (4.6), so we omit the details here.

Let us proceed to prove the closedness condition (3.17). We fix a choice of $X^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})$ such that X° satisfies

$$\left[\frac{\partial}{\partial \tau_0}, X^\circ \right] = I_0$$

and that the differential degree zero part of X° is given by genus zero Virasoro symmetry.

We first write down the explicit expression for I_1 defined in (3.13). Define a derivation $\hat{I}_1 \in \text{Der}(\hat{\mathcal{A}})^0$ by the formulae $\hat{I}_1 v_\gamma = \hat{I}_1 \sigma_{\gamma,0} = 0$ and

$$\begin{aligned} \hat{I}_1 v_\gamma^{(n)} &= n \partial_x^{n-1} W(v_\gamma) + a^{\alpha\beta} \partial_x^n \left(f_{\beta,0} \frac{\partial v_\gamma}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial v_\gamma}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \partial_x^n \left(f_{\beta,0} \frac{\partial v_\gamma}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial v_\gamma}{\partial t^{\beta,0}} \right) \\ &\quad - a^{\alpha\beta} \left(f_{\beta,0} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\alpha,1}} + f_{\alpha,1} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(f_{\beta,0} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\alpha,0}} + f_{\alpha,0} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\beta,0}} \right), \\ \hat{I}_1 \sigma_{\gamma,0}^n &= n \partial_x^{n-1} W(\sigma_{\gamma,0}) + a^{\alpha\beta} \partial_x^n \left(f_{\beta,0} \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \partial_x^n \left(f_{\beta,0} \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) \\ &\quad - a^{\alpha\beta} \left(f_{\beta,0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(f_{\beta,0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) \\ &\quad + \left(\frac{3}{2} + \mu_\gamma \right) (\sigma_{\gamma,2}^n + \partial_x^n Z_\gamma^\varepsilon \sigma_{\varepsilon,1}) + \partial_x^n (N_\gamma^\varepsilon \sigma_{\varepsilon,1}) - \partial_x^n N_\gamma^\varepsilon \sigma_{\varepsilon,1}, \end{aligned}$$

here $n \geq 1$. Note that this derivation does NOT commute with ∂_x . It is easy to see that \hat{I}_1 is indeed local. By a careful computation, we obtain the following expression for I_1 :

$$\begin{aligned} I_1 v_\lambda &= \hat{I}_1 \left(\frac{\partial v_\lambda}{\partial \tau_1} \right) + A^\alpha \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + B^\alpha \frac{\partial v_\lambda}{\partial t^{\alpha,0}}, \\ I_1 \sigma_{\lambda,0} &= \hat{I}_1 \left(\frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} \right) + A^\alpha \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + B^\alpha \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} \\ &\quad + \left(\frac{3}{2} + \mu_\lambda \right) \left(\frac{\partial \sigma_{\lambda,1}}{\partial \tau_2} - \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial \sigma_{\lambda,1}}{\partial t^{\alpha,0}} \right), \end{aligned}$$

where A^α and B^α are local differential polynomials given by

$$\begin{aligned} A^\alpha &= -a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} - \sigma_{\beta,1} \right), \\ B^\alpha &= -\eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{3}{2} + \mu_\beta \right) \left(\left(\frac{1}{2} + \mu_\beta \right) \frac{\partial f_{\beta,1}}{\partial \tau_1} - \sigma_{\beta,2} - Z_\beta^\varepsilon \sigma_{\varepsilon,1} \right) \\ &\quad - (b^{\alpha\beta} + b^{\beta\alpha}) \left(\frac{\partial f_{\beta,0}}{\partial \tau_1} - \sigma_{\beta,1} \right). \end{aligned}$$

We start by proving the first identity in (3.17), which can also be written as

$$\left[\frac{\partial}{\partial \tau_0}, I_1 \right] + \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, X^\circ \right] \right] = 0.$$

The following lemmas are obtained by lengthy but straightforward computations.

Lemma 4.8. *Let $Q \in \hat{\mathcal{A}}$ be any local differential polynomial, then we have*

$$\begin{aligned} \left[\hat{I}_1, \partial_x \right] Q &= W(Q) + a_{\alpha\beta} \left(f'_{\beta,0} \frac{\partial Q}{\partial t^{\alpha,1}} + f'_{\alpha,1} \frac{\partial Q}{\partial t^{\beta,0}} \right) + b_{\alpha\beta} \left(f'_{\beta,0} \frac{\partial Q}{\partial t^{\alpha,0}} + f'_{\alpha,0} \frac{\partial Q}{\partial t^{\beta,0}} \right) \\ &\quad + \sum_{n \geq 0} \sigma_{\zeta,1}^1 \partial_x^n \left(Y_\gamma^\zeta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\zeta \right) \frac{\partial}{\partial \sigma_{\gamma,0}^n} Q \\ &\quad + \left(\frac{3}{2} + \mu_\gamma \right) \partial_x (\sigma_{\gamma,2} + Z_\gamma^\varepsilon \sigma_{\varepsilon,1}) \frac{\partial}{\partial \sigma_{\gamma,0}} Q. \end{aligned}$$

Lemma 4.9. *The following identities hold true:*

$$\begin{aligned} \left[\frac{\partial}{\partial \tau_0}, I_1 \right] v_\lambda &= \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, \hat{I}_1 \right] \right] v_\lambda + \frac{\partial A^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + \frac{\partial B^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}}, \\ \left[\frac{\partial}{\partial \tau_0}, I_1 \right] \sigma_{\lambda,0} &= \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, \hat{I}_1 \right] \right] \sigma_{\lambda,0} + \frac{\partial A^\alpha}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + \frac{\partial B^\alpha}{\partial \tau_0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} \\ &\quad + \left(\frac{3}{2} + \mu_\lambda \right) \left(\frac{\partial \sigma_{\lambda,1}}{\partial \tau_2} - \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial \sigma_{\lambda,1}}{\partial t^{\alpha,0}} \right). \end{aligned}$$

Lemma 4.10. *We have the following decomposition:*

$$\left[\frac{\partial}{\partial \tau_0}, \hat{I}_1 + X^\circ \right] = D + \frac{\partial}{\partial \tau_2},$$

where D is a derivation $\hat{\mathcal{A}} \rightarrow \hat{\mathcal{A}}^+$ given by the formulae

$$\begin{aligned}
& Dv_\gamma^{(n)} \\
&= -a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_0} \partial_x^n \frac{\partial v_\gamma}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \partial_x^n \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_0} \partial_x^n \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\
&\quad + \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{1}{2} + \mu_\beta \right) \partial_x^n \frac{\partial v_\gamma}{\partial t^{\alpha,0}}, \\
& D\sigma_{\gamma,0}^n \\
&= -a^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,1}} + \frac{\partial f_{\alpha,1}}{\partial \tau_0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) - b^{\alpha\beta} \left(\frac{\partial f_{\beta,0}}{\partial \tau_0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,0}} + \frac{\partial f_{\alpha,0}}{\partial \tau_0} \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\beta,0}} \right) \\
&\quad + \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \left(\frac{1}{2} + \mu_\beta \right) \partial_x^n \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,0}} + \frac{\partial \sigma_{\beta,0}}{\partial \tau_1} \partial_x^n \left(N_\gamma^\beta - \left(\frac{3}{2} + \mu_\gamma \right) Z_\gamma^\beta \right) \\
&\quad - \delta_{n,0} \left(\frac{3}{2} + \mu_\gamma \right) \frac{\partial}{\partial \tau_0} (\sigma_{\gamma,2} + Z_\gamma^\varepsilon \sigma_{\varepsilon,1}).
\end{aligned}$$

Proposition 4.11. *The first identity of the closedness condition (3.17) holds true, i.e.,*

$$\left[\frac{\partial}{\partial \tau_0}, I_1 \right] + \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, X^\circ \right] \right] = 0.$$

Proof. It follows from Lemma 4.9 that

$$\begin{aligned}
& \left[\frac{\partial}{\partial \tau_0}, I_1 \right] v_\lambda + \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, X^\circ \right] \right] v_\lambda \\
&= \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, \hat{I}_1 + X^\circ \right] \right] v_\lambda + \frac{\partial A^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + \frac{\partial B^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}}.
\end{aligned}$$

Due to Lemma 4.10 we only need to verify the following identity:

$$\left[\frac{\partial}{\partial \tau_1}, D \right] v_\lambda + \frac{\partial A^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + \frac{\partial B^\alpha}{\partial \tau_0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} = 0,$$

which can be checked directly from the definition of D . Similarly we can prove that

$$\left[\frac{\partial}{\partial \tau_0}, I_1 \right] \sigma_{\lambda,0} + \left[\frac{\partial}{\partial \tau_1}, \left[\frac{\partial}{\partial \tau_0}, X^\circ \right] \right] \sigma_{\lambda,0} = 0.$$

The proposition is proved. \square

It remains to prove the second identity of (3.17).

Lemma 4.12. *The following identities hold true:*

$$\begin{aligned} \left[I_1, \frac{\partial}{\partial \tau_1} \right] v_\lambda &= \sum_{s \geq 0} G_s^\gamma \frac{\partial}{\partial v_\gamma^{(s)}} \frac{\partial v_\lambda}{\partial \tau_1} + F_s^\gamma \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial v_\lambda}{\partial \tau_1} \\ &\quad - A^\alpha \frac{\partial}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} - B^\alpha \frac{\partial}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + \frac{\partial B^\alpha}{\partial \tau_1} \frac{\partial v_\lambda}{\partial t^{\alpha,0}}, \\ \left[I_1, \frac{\partial}{\partial \tau_1} \right] \sigma_{\lambda,0} &= \sum_{s \geq 0} G_s^\gamma \frac{\partial}{\partial v_\gamma^{(s)}} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} + F_s^\gamma \frac{\partial}{\partial \sigma_{\gamma,0}^s} \frac{\partial \sigma_{\lambda,0}}{\partial \tau_1} \\ &\quad - A^\alpha \frac{\partial}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} - B^\alpha \frac{\partial}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + \frac{\partial B^\alpha}{\partial \tau_1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}}, \end{aligned}$$

where G_s^γ and F_s^γ are differential polynomials defined by

$$G_s^\gamma = \left[\frac{\partial}{\partial \tau_1}, \hat{I}_1 \right] v_\gamma^{(s)} + \partial_x^s (I_1 v_\gamma), \quad F_s^\gamma = \left[\frac{\partial}{\partial \tau_1}, \hat{I}_1 \right] \sigma_{\gamma,0}^s + \partial_x^s (I_1 \sigma_{\gamma,0}).$$

Lemma 4.13. *The differential polynomials G_s^γ and F_s^γ defined in Lemma 4.12 have the following expressions:*

$$\begin{aligned} G_s^\gamma &= A^\alpha \partial_x^s \frac{\partial v_\gamma}{\partial t^{\alpha,1}} + B^\alpha \partial_x^s \frac{\partial v_\gamma}{\partial t^{\alpha,0}}, \\ F_s^\gamma &= A^\alpha \partial_x^s \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,1}} + B^\alpha \partial_x^s \frac{\partial \sigma_{\gamma,0}}{\partial t^{\alpha,0}} \\ &\quad + \delta_{s,0} \left(\frac{3}{2} + \mu_\gamma \right) \left(\frac{\partial \sigma_{\gamma,1}}{\partial \tau_2} - \sigma_{\beta,1} \eta^{\alpha\beta} \left(\frac{1}{2} + \mu_\alpha \right) \frac{\partial \sigma_{\gamma,1}}{\partial t^{\alpha,0}} \right). \end{aligned}$$

Proof. By using the definition of G_s^γ we can obtain the identity

$$G_{s+1}^\gamma - \partial_x G_s^\gamma = \frac{\partial}{\partial \tau_1} \left(\left[\hat{I}_1, \partial_x \right] v_\gamma^{(s)} \right) + \left[\partial_x, \hat{I}_1 \right] \frac{\partial v_\gamma^{(s)}}{\partial \tau_1}.$$

Therefore it follows from Lemma 4.8 that

$$G_{s+1}^\gamma - \partial_x G_s^\gamma = -\partial_x A^\alpha \partial_x^s \frac{\partial v_\gamma}{\partial t^{\alpha,1}} - \partial_x B^\alpha \partial_x^s \frac{\partial v_\gamma}{\partial t^{\alpha,0}}.$$

Hence G_s^γ can be solved recursively starting from the initial condition

$$G_0^\gamma = I_1 v_\gamma - \hat{I}_1 \frac{\partial v_\gamma}{\partial \tau_1} = A^\alpha \frac{\partial v_\gamma}{\partial t^{\alpha,1}} + B^\alpha \frac{\partial v_\gamma}{\partial t^{\alpha,0}}.$$

The differential polynomials F_s^γ can be computed similarly. The lemma is proved. \square

Proposition 4.14. *The second identity of the closedness condition (3.17) holds true, i.e.,*

$$\left[\frac{\partial}{\partial \tau_1}, I_1 \right] = 0.$$

Proof. The proof of the proposition is straightforward by combining the results of Lemma 4.12 and Lemma 4.13. \square

4.3. Vanishing of the genus one obstruction. In this subsection, we will work with the canonical coordinates of the Frobenius manifold. Let us start by recalling some useful formulae related to the canonical coordinates. For the details one may refer to the work [5, 12].

We denote by u^1, \dots, u^n the local canonical coordinates of a semisimple Frobenius manifold M and denote by $(u^i; \theta_i)$ the corresponding coordinates of \hat{M} . Under this system of local coordinates, the bihamiltonian structure (2.12) takes the form (2.6), i.e.,

$$P_0^{[0]} = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} f^i \theta_i \theta_j^1 + A^{ij} \theta_i \theta_j), \quad P_1^{[0]} = \frac{1}{2} \int \sum_{i,j=1}^n (\delta_{i,j} u^i f^i \theta_i \theta_j^1 + B^{ij} \theta_i \theta_j).$$

Introduce functions

$$(4.12) \quad \psi_{i1} = \frac{1}{\sqrt{f^i}},$$

where the sign of the square root can be arbitrarily chosen, and define

$$\psi_{i\alpha} = \frac{1}{\psi_{i1}} \frac{\partial v_\alpha}{\partial u^i}, \quad \gamma_{ij} = \frac{1}{\psi_{j1}} \frac{\partial \psi_{i1}}{\partial u^j}, \quad i \neq j.$$

Then it is proved in [5] that

$$(4.13) \quad \frac{\partial v_\alpha}{\partial u^i} = \psi_{i1} \psi_{i\alpha}, \quad \frac{\partial u^i}{\partial v^\alpha} = \frac{\psi_{i\alpha}}{\psi_{i1}},$$

$$(4.14) \quad \frac{\partial \psi_{i\alpha}}{\partial u^k} = \gamma_{ik} \psi_{k\alpha}, \quad i \neq k; \quad \frac{\partial \psi_{i\alpha}}{\partial u^i} = - \sum_{k \neq i} \gamma_{ik} \psi_{k\alpha}.$$

Let $V_{ij} = \mu_\alpha \eta^{\alpha\beta} \psi_{i\alpha} \psi_{j\beta}$, then we have the identity

$$(4.15) \quad \gamma_{ij} (u^j - u^i) = V_{ij}.$$

Now we are going to describe the deformation problem given in Sect. 3.3 in terms of the canonical coordinates. Introduce the odd variables $\theta_{i,m}$ by the recursion relations (2.15), then it follows from the relation (2.16) that the equations (3.8) and (3.9) can be represented in the form

$$(4.16) \quad \frac{\partial u^i}{\partial s_2} = a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial u^i}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial u^i}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial u^i}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial u^i}{\partial t^{\beta,0}} \right) + Xu^i + \mathcal{L}_2 u^i,$$

$$(4.17) \quad \frac{\partial \theta_{i,0}}{\partial s_2} = a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \theta_{i,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial \theta_{i,0}}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \theta_{i,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial \theta_{i,0}}{\partial t^{\beta,0}} \right) + X\theta_{i,0} + \sum_j A_i^j \theta_{j,2} + B_i^j \theta_{j,1} + \mathcal{L}_2 \theta_{i,0}.$$

In the canonical coordinates, the Hamiltonian operator \mathcal{P}_1 of P_1 has the form

$$\mathcal{P}_1^{ij} = u^i f^i \delta_{ij} \partial_x + \frac{1}{2} \partial_x (u^i f^i) \delta_{ij} + B^{ij} + Q^{ij} \partial_x^3 + \dots$$

where B^{ij} is defined in (2.5) and Q^{ij} is given by the following formula (c.f., e.g., [13])

$$(4.18) \quad Q^{ij} = 3c_i (f^i)^2 \delta_{ij} + \frac{1}{2} (u^i - u^j) (f^j \partial_j f^i c_i - f^i \partial_i f^j c_j),$$

here c_i is the i -th central invariant. Let us also denote the evolutions of θ_i along the flows $\frac{\partial}{\partial t^{\alpha,p}}$ by the following equations:

$$\frac{\partial \theta_i}{\partial t^{\alpha,p}} = T_{\alpha,p}^i \theta_i^1 + \cdots + K_{\alpha,p;j}^i \theta_j^3 + \cdots, \quad T_{\alpha,p} \in \hat{\mathcal{A}}_0^0, \quad K_{\alpha,p;j}^i \in \hat{\mathcal{A}}_3^0,$$

here and henceforth we omit the terms that do not contribute to the relevant computations given later. Note that the coefficients of θ_j^1 of the leading term of $\frac{\partial \theta_i}{\partial t^{\alpha,p}}$ are zero for $j \neq i$, this is due to the fact that the leading term of the flow $\frac{\partial}{\partial t^{\alpha,p}}$ is diagonal, one may refer to [9] for details.

Let us turn to the proof of the vanishing of the genus one obstruction. Due to the closedness condition (3.15), we can choose a derivation X° such that when we take $X = X^\circ$ in (4.16) and (4.17) we have

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial s_2} \right] = 0.$$

We require that the leading term of X° is given by the genus zero Virasoro symmetry, hence it follows from the result of [11] that the actions of X° on u^i and θ_i take the form

$$\begin{aligned} X^\circ u^i &= (u^i)^3 + \sum_j L_j^i u^{j,2} + \cdots, \quad L_j^i \in \hat{\mathcal{A}}_0^0, \\ X^\circ \theta_i &= \sum_j M_i^j \theta_j + \sum_j J_i^j \theta_j^2 + \cdots, \quad M_i^j, J_i^j \in \hat{\mathcal{A}}_0^0. \end{aligned}$$

Lemma 4.15. *We have the following identity:*

$$\begin{aligned} J_i^i - L_i^i &= \frac{1}{f^i} 2c_0 (u^i)^2 Q^{ii} - \sum_j \frac{A_i^j}{f^j} (u^i + u^j) Q^{ij} - \sum_j \frac{B_i^j}{f^j} Q^{ij} \\ &\quad - b_{2;1,0}^{\beta,q} K_{\beta,q;i}^i - a_2^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 K_{\alpha,p;i}^i + (f'_{\alpha,p})_0 K_{\beta,q;i}^i \right), \end{aligned}$$

where $b_{2;\alpha,p}^{\beta,q}$ and $a_2^{\alpha,p;\beta,q}$ are the constants that appear in the operator \mathcal{L}_2 , and $(f'_{\alpha,p})_0$ denotes the differential degree zero component of $f'_{\alpha,p}$.

Proof. We can prove this lemma by computing the coefficient of θ_i^3 of differential degree 3 component of the left hand side of the equation

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial s_2} \right] u^i = 0.$$

The lemma is proved. □

Remark 4.1. *Note that c_0 is the arbitrary constant that appears in the operator \mathcal{L}_2 , and c_1, \dots, c_n are the central invariants.*

According to the discussion given in Sect. 3.3, we need to compute the cohomology class of the differential degree 3 component of the derivation $\left[\frac{\partial}{\partial \tau_1}, \mathcal{C}\right] = I_1 - \left[\frac{\partial}{\partial \tau_1}, X^\circ\right]$. Due to Theorem 3.2, it suffices to prove that the coefficient of θ_i^3 of the differential degree 3 component of

$$I_1 u^i - \left[\frac{\partial}{\partial \tau_1}, X^\circ\right] u^i$$

vanishes. By using Lemma 4.15 and a straightforward computation, we obtain the following lemma.

Lemma 4.16. *The coefficient of θ_i^3 of the differential degree 3 component of $I_1 u^i - \left[\frac{\partial}{\partial \tau_1}, X^\circ\right] u^i$ reads*

$$(4.19) \quad \sum_j Q^{ij} (M_j^i + A_j^i (u^i)^2 + B_j^i u^i) + E^3 Q^{ii} - (6 + c_0) (u^i)^2 Q^{ii} \\ + 3Q^{ii} b_{2;1,0}^{\beta,q} T_{\beta,q}^i + 3Q^{ii} a_2^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 T_{\alpha,p}^i + (f'_{\alpha,p})_0 T_{\beta,q}^i \right),$$

here E^3 is the cubic of the Euler vector field, which is given by

$$E^3 = \sum_i (u^i)^3 \frac{\partial}{\partial u^i}.$$

Lemma 4.17. *We have the identities*

$$M_i^i + A_i^i (u^i)^2 + B_i^i u^i \\ = (3 + c_0) (u^i)^2 - \frac{1}{f^i} E^3 f^i - b_{2;1,0}^{\beta,q} T_{\beta,q}^i - a_2^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 T_{\alpha,p}^i + (f'_{\alpha,p})_0 T_{\beta,q}^i \right),$$

and $M_i^j + A_i^j (u^j)^2 + B_i^j u^j = 0$ for $i \neq j$.

Proof. We can prove this lemma by computing the coefficient of θ_j^1 of the differential degree 1 component of the left hand side of the equation

$$\left[\frac{\partial}{\partial \tau_0}, \frac{\partial}{\partial s_2}\right] u^i = 0.$$

The lemma is proved. \square

In order to prove the vanishing of the genus one obstruction, we need to verify that the expression (4.19) vanishes. To this end, we only need to check, due to Lemma 4.17 and the expression (4.18) for Q^{ii} , the following identity:

$$E^3 f^i + 2f^i \left(b_{2;1,0}^{\beta,q} T_{\beta,q}^i + a_2^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 T_{\alpha,p}^i + (f'_{\alpha,p})_0 T_{\beta,q}^i \right) \right) = 3(u^i)^2 f^i.$$

Proposition 4.18. *For $m \geq -1$, we have*

$$E^{m+1} f^i + 2f^i \left(b_{m;1,0}^{\beta,q} T_{\beta,q}^i + a_m^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 T_{\alpha,p}^i + (f'_{\alpha,p})_0 T_{\beta,q}^i \right) \right) = (1 + m) (u^i)^m f^i,$$

here E^{m+1} is the $(m+1)$ -th power of the Euler vector field E which is given by

$$E^{m+1} = \sum_i (u^i)^{m+1} \frac{\partial}{\partial u^i}.$$

Proof. Let us consider the following generating functions:

$$\begin{aligned}
 \sum_{m \geq -1} \frac{1}{\lambda^{m+2}} E^{m+1} f^i &= \sum_j \frac{1}{\lambda - u^j} \frac{\partial f^i}{\partial u^j}; \\
 \sum_{m \geq -1} \frac{1}{\lambda^{m+2}} (1+m)(u^i)^m f^i &= \frac{f^i}{(\lambda - u^i)^2}; \\
 \sum_{m \geq -1} \frac{1}{\lambda^{m+2}} \left(b_{m;1,0}^{\beta,q} T_{\beta,q}^i + a_m^{\alpha,p;\beta,q} \left((f'_{\beta,q})_0 T_{\alpha,p}^i + (f'_{\alpha,p})_0 T_{\beta,q}^i \right) \right) \\
 &= \frac{1}{2} \frac{1}{(\lambda - u^i)^2} + \sum_{j \neq i} \frac{\psi_{j1}}{\psi_{i1}} \frac{V_{ij}}{(\lambda - u^i)(\lambda - u^j)},
 \end{aligned}$$

where the last generating function was computed in [12], one may refer to Lemma 3.10.18 and the proof of Theorem 3.10.29 of [12] for details. Then the proposition is proved by using (4.12), (4.14) and (4.15). \square

Finally we have the following theorem.

Theorem 4.19. *There exists a unique $X \in \text{Der}^0(\hat{\mathcal{A}})^0$ such that the following flow defines a symmetry of the deformed Principal Hierarchy which is a deformation of the genus zero Virasoro symmetry $\frac{\partial}{\partial s_2}$ of the Principal Hierarchy:*

$$\begin{aligned}
 (4.20) \quad \frac{\partial v_\lambda}{\partial s_2} &= a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial v_\lambda}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial v_\lambda}{\partial t^{\beta,0}} \right) \\
 &\quad + X v_\lambda + \mathcal{L}_2 v_\lambda,
 \end{aligned}$$

$$\begin{aligned}
 (4.21) \quad \frac{\partial \sigma_{\lambda,0}}{\partial s_2} &= a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial \sigma_{\lambda,0}}{\partial t^{\beta,0}} \right) \\
 &\quad + X \sigma_{\lambda,0} + \left(\frac{5}{2} + c_0 + \mu_\lambda \right) \sigma_{\lambda,2} + N_\lambda^\zeta \sigma_{\zeta,1} + \mathcal{L}_2 \sigma_{\lambda,0},
 \end{aligned}$$

where $N_\lambda^\zeta \in \hat{\mathcal{A}}_{\geq 0}^0$ is the differential polynomial described in Lemma 4.3.

4.4. Lifting to the tau-covers. In order to lift the symmetry (4.20) to the tau-cover of the deformed Principal Hierarchy, we first need to rewrite (4.20) in terms of the normal coordinates w^1, \dots, w^n of M . We start by proving the following lemmas.

Lemma 4.20. *Let $g_\lambda \in \hat{\mathcal{A}}_{\geq 1}^0$ be the differential polynomials given in (2.31) which satisfy the identities*

$$h_{\lambda,0} = v_\lambda + \partial_x g_\lambda, \quad \lambda = 1, \dots, n.$$

Then we have:

- (1) $g_1 = 0$.
- (2) For any $\alpha = 1, \dots, n$ and $p \geq 0$,

$$\Omega_{\alpha,p;\lambda,0} = \frac{\delta h_{\alpha,p+1}}{\delta v^\lambda} + \frac{\partial g_\lambda}{\partial t^{\alpha,p}}.$$

Proof. Due to Proposition 2.17, we have $D_{X_{1,0}} = \partial_x$ and $X_{1,0} = -[H_{1,0}, P_0]$, from which it follows that

$$v_\alpha = \frac{\delta H_{1,0}}{\delta v^\alpha}.$$

By taking $\alpha = 1$, we obtain the first property by using the definition (2.24). The second one is obvious due to Theorem 2.19. The lemma is proved. \square

Lemma 4.21. *There exists a derivation $X^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})$ such that its leading term is given by the Virasoro symmetry $\frac{\partial}{\partial s_2}$ of the Principal Hierarchy, and it satisfies the equations $\left[\frac{\partial}{\partial \tau_0}, X^\circ\right] = I_0$ and*

$$(4.22) \quad \begin{aligned} X^\circ h_{\lambda,0} + \hat{I}_1 h_{\lambda,0} = & a^{\alpha\beta} \left(f'_{\alpha,1} \Omega_{\beta,0;\lambda,0} + f'_{\beta,0} \Omega_{\alpha,1;\lambda,0} + \frac{\partial^2 \Omega_{\alpha,1;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} \right) \\ & + b^{\alpha\beta} \left(f'_{\alpha,0} \Omega_{\beta,0;\lambda,0} + f'_{\beta,0} \Omega_{\alpha,0;\lambda,0} + \frac{\partial^2 \Omega_{\alpha,0;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} \right) \\ & + b_{2;1,0}^{\beta,q} \Omega_{\beta,q;\lambda,0} + b_{2;\lambda,0}^{\beta,q} \Omega_{\beta,q;1,0} + 2c_{2;\lambda,0;1,0}. \end{aligned}$$

Proof. We first find a particular solution \tilde{X}° of the equation $\left[\frac{\partial}{\partial \tau_0}, X\right] = I_0$, then we modify it by a solution $\tilde{\mathcal{C}}$ of the homogeneous equation $\left[\frac{\partial}{\partial \tau_0}, \mathcal{C}\right] = 0$ such that $X^\circ := \tilde{X}^\circ + \tilde{\mathcal{C}}$ satisfies (4.22).

Let us define $\tilde{X}^\circ \in \text{Der}^\partial(\hat{\mathcal{A}})$ as follows:

$$\begin{aligned} \tilde{X}^\circ v_\lambda = & a^{\alpha\beta} \left(f'_{\alpha,1} \frac{\delta h_{\beta,1}}{\delta v^\lambda} + f'_{\beta,0} \frac{\delta h_{\alpha,2}}{\delta v^\lambda} \right) + b^{\alpha\beta} \left(f'_{\alpha,0} \frac{\delta h_{\beta,1}}{\delta v^\lambda} + f'_{\beta,0} \frac{\delta h_{\alpha,1}}{\delta v^\lambda} \right) \\ & + b_{2;1,0}^{\beta,q} \frac{\delta h_{\beta,q+1}}{\delta v^\lambda} + b_{2;\lambda,0}^{\beta,q} \frac{\delta h_{\beta,q+1}}{\delta v^1} + 2c_{2;\lambda,0;1,0}; \\ \tilde{X}^\circ \sigma_{\lambda,0} = & a^{\alpha\beta} \left(\frac{\partial f_{\alpha,1}}{\partial \tau_0} \frac{\delta h_{\beta,1}}{\delta v^\lambda} + \frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\delta h_{\alpha,2}}{\delta v^\lambda} \right) + b^{\alpha\beta} \left(\frac{\partial f_{\alpha,0}}{\partial \tau_0} \frac{\delta h_{\beta,1}}{\delta v^\lambda} + \frac{\partial f_{\beta,0}}{\partial \tau_0} \frac{\delta h_{\alpha,1}}{\delta v^\lambda} \right) \\ & + b_{2;\lambda,0}^{\beta,q} \frac{\partial f_{\beta,q}}{\partial \tau_0} - \left(\frac{5}{2} + \mu_\lambda \right) \sigma_{\lambda,2} - N_\lambda^\zeta \sigma_{\zeta,1}. \end{aligned}$$

Firstly, from the definition of N_λ^ζ it follows that \tilde{X}° is indeed local. We also note that the leading terms of $\tilde{X}^\circ v_\lambda$ and $\tilde{X}^\circ \sigma_{\lambda,0}$ coincide with the local terms of the Virasoro symmetry $\frac{\partial v_\lambda}{\partial s_2}$ and $\frac{\partial \sigma_{\lambda,0}}{\partial s_2}$ of the Principal Hierarchy. By a direct computation, it is easy to check that $\left[\frac{\partial}{\partial \tau_0}, \tilde{X}^\circ\right] = I_0$.

Next we want to determine $\tilde{\mathcal{C}} \in \text{Der}^\partial(\hat{\mathcal{A}})$ such that $X^\circ = \tilde{X}^\circ + \tilde{\mathcal{C}}$ satisfies (4.22). By using the definition of $\tilde{X}^\circ v_\lambda$ and Lemma 4.8, 4.20, it is straightforward to show that

$$(4.23) \quad \tilde{\mathcal{C}}(h_{\lambda,0}) = a^{\alpha\beta} \frac{\partial^2 \Omega_{\alpha,1;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} + b^{\alpha\beta} \frac{\partial^2 \Omega_{\alpha,0;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} - \partial_x \hat{I}_1(g_\lambda) - \partial_x \tilde{X}^\circ(g_\lambda),$$

which uniquely determines the actions $\tilde{\mathcal{C}} v_\lambda$.

Finally we need to check such $\tilde{\mathcal{C}}$ satisfies $\left[\frac{\partial}{\partial\tau_0}, \tilde{\mathcal{C}}\right] = 0$. By using the next lemma, we know that it suffices to show that $\int \tilde{\mathcal{C}}v_\lambda = 0$, which is obvious from (4.23) and (2.31). The lemma is proved. \square

Lemma 4.22. *Let U_1, \dots, U_n be differential polynomials with $U_\lambda \in \hat{\mathcal{A}}_{\geq 2}^0$. Then there exists $\mathcal{C} \in \text{Der}^\partial(\hat{\mathcal{A}})_{\geq 1}^0$ such that $\left[\frac{\partial}{\partial\tau_0}, \mathcal{C}\right] = 0$ and $\mathcal{C}v_\lambda = U_\lambda$ if and only if $\int U_\lambda = 0$.*

Proof. Let $\mathcal{C} \in \text{Der}^\partial(\hat{\mathcal{A}})^0$ be a derivation such that $\left[\frac{\partial}{\partial\tau_0}, \mathcal{C}\right] = 0$. Then by using the triviality of the variational Hamiltonian cohomology $H_{\geq 1}^0(\text{Der}^\partial(\hat{\mathcal{A}}))$, we know the existence of a certain $\mathcal{K} \in \text{Der}^\partial(\hat{\mathcal{A}})^{-1}$ such that $\left[\frac{\partial}{\partial\tau_0}, \mathcal{K}\right] = \mathcal{C}$. Let us denote $\mathcal{K}\sigma_{\lambda,0} = V_\lambda \in \hat{\mathcal{A}}^0$. Then it is easy to see that

$$\mathcal{C}v_\lambda = \left[\frac{\partial}{\partial\tau_0}, \mathcal{K}\right]v_\lambda = \mathcal{K}\sigma_{\lambda,0}^1 = \partial_x V_\lambda.$$

Therefore we have $\int U_\lambda = \int \partial_x V_\lambda = 0$.

Conversely, if $U_\lambda = \partial_x V_\lambda$ for some $V_\lambda \in \hat{\mathcal{A}}^0$, we can define a derivation $\mathcal{C} \in \text{Der}^\partial(\hat{\mathcal{A}})^0$ by

$$\mathcal{C}v_\lambda = \partial_x V_\lambda, \quad \mathcal{C}\sigma_{\lambda,0} = \frac{\partial V_\lambda}{\partial\tau_0},$$

then it is easy to check that $\left[\frac{\partial}{\partial\tau_0}, \mathcal{C}\right] = 0$ and the lemma is proved. \square

Now let us denote $\mathcal{C} = X - X^\circ$, where the derivation X is described in Theorem 4.19 and X° satisfies (4.22). Then we can rewrite the Virasoro symmetry in terms of the normal coordinates as follows:

$$\begin{aligned} \frac{\partial w_\lambda}{\partial s_2} &= a^{\alpha\beta} \left(f_{\beta,0} \frac{\partial w_\lambda}{\partial t^{\alpha,1}} + f_{\alpha,1} \frac{\partial w_\lambda}{\partial t^{\beta,0}} \right) + b^{\alpha\beta} \left(f_{\beta,0} \frac{\partial w_\lambda}{\partial t^{\alpha,0}} + f_{\alpha,0} \frac{\partial w_\lambda}{\partial t^{\beta,0}} \right) \\ &\quad + a^{\alpha\beta} \left(f'_{\alpha,1} \Omega_{\beta,0;\lambda,0} + f'_{\beta,0} \Omega_{\alpha,1;\lambda,0} + \frac{\partial^2 \Omega_{\alpha,1;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} \right) \\ &\quad + b^{\alpha\beta} \left(f'_{\alpha,0} \Omega_{\beta,0;\lambda,0} + f'_{\beta,0} \Omega_{\alpha,0;\lambda,0} + \frac{\partial^2 \Omega_{\alpha,0;\beta,0}}{\partial t^{\lambda,0} \partial t^{1,0}} \right) \\ &\quad + b_{2;1,0}^{\beta,q} \Omega_{\beta,q;\lambda,0} + b_{2;\lambda,0}^{\beta,q} \Omega_{\beta,q;1,0} + 2c_{2;\lambda,0;1,0} + \mathcal{C}(h_{\lambda,0}) + \mathcal{L}_2 w_\lambda. \end{aligned}$$

According to Lemma 4.22, there exists $Q_\lambda \in \hat{\mathcal{A}}_{\geq 1}^0$ such that $\mathcal{C}(h_{\lambda,0}) = \partial_x Q_\lambda$. In particular, the fact that

$$\frac{\partial}{\partial t^{\gamma,0}} \frac{\partial w_\lambda}{\partial s_2} = \frac{\partial}{\partial t^{\lambda,0}} \frac{\partial w_\gamma}{\partial s_2}$$

gives the equation

$$\frac{\partial Q_\lambda}{\partial t^{\gamma,0}} = \frac{\partial Q_\gamma}{\partial t^{\lambda,0}},$$

which means that $\int Q_1$ is a conserved quantity of the flows $\frac{\partial}{\partial t^{\lambda,0}}$. If the dimension of the Frobenius manifold $\dim M \geq 2$, then it follows from Lemma 4.12 and Theorem A.2 of [9] that there exists $Q \in \hat{\mathcal{A}}_{\geq 0}^0$ such that $Q_\lambda = \frac{\partial Q}{\partial t^{\lambda,0}}$.

Theorem 4.23. *Let \mathcal{Z} be a tau-function of the tau-cover (2.28) of the deformed Principal Hierarchy. Then there exists a differential polynomial $O_2 \in \hat{\mathcal{A}}_{\geq 0}^0$ which yields a symmetry of the tau-cover given by*

$$\frac{\partial \mathcal{Z}}{\partial s_2} = L_2^{even} \mathcal{Z} + O_2 \mathcal{Z},$$

where the evolutions of $f_{\alpha,p}$ and w_λ along the flow $\frac{\partial}{\partial s_2}$ are defined by

$$\frac{\partial f_{\alpha,p}}{\partial s_2} = \frac{\partial}{\partial t^{\alpha,p}} \frac{\partial \log Z}{\partial s_2}, \quad \frac{\partial w_\lambda}{\partial s_2} = \frac{\partial}{\partial t^{1,0}} \frac{\partial f_{\lambda,0}}{\partial s_2}.$$

Proof. When $\dim M \geq 2$ we take $O_2 = Q$ which is just given above. When $\dim M = 1$, O_2 is given in (3.26). The theorem is proved. \square

Now let us proceed to determine all other Virasoro symmetries $\frac{\partial}{\partial s_m}$ of the tau-cover of the deformed Principal Hierarchy. In what follows we will denote by $\mathcal{F} = \log \mathcal{Z}$ and denote

$$\begin{aligned} \mathcal{G}_m &= a_m^{\alpha,p;\beta,q} \left(\frac{\partial \mathcal{F}}{\partial t^{\alpha,p}} \frac{\partial \mathcal{F}}{\partial t^{\beta,q}} + \frac{\partial^2 \mathcal{F}}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) + c_{m;\alpha,p;\beta,q} t^{\alpha,p} t^{\beta,q}, \\ \mathcal{L}_m^{even} &= b_{m;\alpha,p}^{\beta,q} t^{\alpha,p} \frac{\partial}{\partial t^{\beta,q}}, \quad m \geq -1. \end{aligned}$$

It is proved in [9] that

$$\frac{\partial \mathcal{F}}{\partial s_{-1}} = \mathcal{G}_{-1} + \mathcal{L}_{-1}^{even} \mathcal{F}$$

induces a symmetry of the tau-cover of the deformed Principal Hierarchy. On the other hand, by using Theorem 4.23 we know that

$$\frac{\partial \mathcal{F}}{\partial s_2} = \mathcal{G}_2 + O_2 + \mathcal{L}_2^{even} \mathcal{F}$$

also induces a symmetry of the tau-cover of the deformed Principal Hierarchy. From the commutation relation

$$[L_{-1}^{even}, L_2^{even}] = -3L_1^{even},$$

it follows that

$$\left[\frac{\partial}{\partial s_{-1}}, \frac{\partial}{\partial s_2} \right] \mathcal{F} = 3(\mathcal{G}_1 + \mathcal{L}_1^{even} \mathcal{F}) + \frac{\partial O_2}{\partial s_{-1}} - \mathcal{L}_{-1}^{even} O_2.$$

Since $\frac{\partial O_2}{\partial s_{-1}} - \mathcal{L}_{-1}^{even} O_2$ is a differential polynomial, we can define

$$\frac{\partial \mathcal{F}}{\partial s_1} = \frac{1}{3} \left[\frac{\partial}{\partial s_{-1}}, \frac{\partial}{\partial s_2} \right] \mathcal{F} = \mathcal{G}_1 + O_1 + \mathcal{L}_1^{even} \mathcal{F}, \quad O_1 = \frac{1}{3} \left(\frac{\partial O_2}{\partial s_{-1}} - \mathcal{L}_{-1}^{even} O_2 \right).$$

Then it is obvious that $\frac{\partial \mathcal{F}}{\partial s_1}$ induces a symmetry of the tau-cover of the deformed Principal Hierarchy. Similarly we define the symmetry

$$\frac{\partial \mathcal{F}}{\partial s_0} = \frac{1}{2} \left[\frac{\partial}{\partial s_{-1}}, \frac{\partial}{\partial s_1} \right] \mathcal{F} = \mathcal{G}_0 + O_0 + \mathcal{L}_0^{even} \mathcal{F}, \quad O_0 = \frac{1}{2} \left(\frac{\partial O_1}{\partial s_{-1}} - \mathcal{L}_{-1}^{even} O_1 \right).$$

Now let us define $\frac{\partial \mathcal{F}}{\partial s_m}$ for $m \geq 3$ recursively in the following way. Assume we have defined

$$\frac{\partial \mathcal{F}}{\partial s_m} = \mathcal{G}_m + O_m + \mathcal{L}_m^{\text{even}} \mathcal{F}$$

such that it induces a symmetry of the tau-cover of the deformed Principal Hierarchy for $m \geq 2$. Then we have

$$\frac{\partial}{\partial s_1} \frac{\partial \mathcal{F}}{\partial s_m} = \frac{\partial \mathcal{G}_m}{\partial s_1} + \frac{\partial O_m}{\partial s_1} + \mathcal{L}_m^{\text{even}} (\mathcal{G}_1 + O_1 + \mathcal{L}_1^{\text{even}} \mathcal{F}).$$

It follows from the definition of \mathcal{G}_m that

$$\begin{aligned} \frac{\partial \mathcal{G}_m}{\partial s_1} &= \frac{\partial}{\partial s_1} \left(a_m^{\alpha,p;\beta,q} \left(\frac{\partial \mathcal{F}}{\partial t^{\alpha,p}} \frac{\partial \mathcal{F}}{\partial t^{\beta,q}} + \frac{\partial^2 \mathcal{F}}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) + c_{m;\alpha,p;\beta,q} t^{\alpha,p} t^{\beta,q} \right) \\ &= a_m^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial}{\partial t^{\beta,q}} (\mathcal{G}_1 + O_1 + \mathcal{L}_1^{\text{even}} \mathcal{F}) + f_{\beta,q} \frac{\partial}{\partial t^{\alpha,p}} (\mathcal{G}_1 + O_1 + \mathcal{L}_1^{\text{even}} \mathcal{F}) \right) \\ &\quad + a_m^{\alpha,p;\beta,q} \frac{\partial^2}{\partial t^{\alpha,p} \partial t^{\beta,q}} (\mathcal{G}_1 + O_1 + \mathcal{L}_1^{\text{even}} \mathcal{F}) \\ &= a_m^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial O_1}{\partial t^{\beta,q}} + f_{\beta,q} \frac{\partial O_1}{\partial t^{\alpha,p}} + \frac{\partial^2 O_1}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) + \dots, \end{aligned}$$

here \dots stands for remaining terms that do not contain O_1 . In a similar way we can compute $\frac{\partial}{\partial s_m} \frac{\partial \mathcal{F}}{\partial s_1}$. By using the commutation relation

$$[L_1^{\text{even}}, L_m^{\text{even}}] = (1-m)L_{m+1}^{\text{even}}$$

we obtain the following result:

$$\begin{aligned} \left[\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_m} \right] \mathcal{F} &= (m-1) (\mathcal{G}_{m+1} + \mathcal{L}_{m+1}^{\text{even}} \mathcal{F}) \\ &\quad + \frac{\partial O_m}{\partial s_1} - a_1^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial O_m}{\partial t^{\beta,q}} + f_{\beta,q} \frac{\partial O_m}{\partial t^{\alpha,p}} + \frac{\partial^2 O_m}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) - \mathcal{L}_1^{\text{even}} O_m \\ &\quad - \frac{\partial O_1}{\partial s_m} + a_m^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial O_1}{\partial t^{\beta,q}} + f_{\beta,q} \frac{\partial O_1}{\partial t^{\alpha,p}} + \frac{\partial^2 O_1}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) + \mathcal{L}_m^{\text{even}} O_1. \end{aligned}$$

Therefore we obtain the symmetry $\frac{\partial \mathcal{F}}{\partial s_{m+1}}$ by defining

$$\frac{\partial \mathcal{F}}{\partial s_{m+1}} = \frac{1}{m-1} \left[\frac{\partial}{\partial s_1}, \frac{\partial}{\partial s_m} \right] \mathcal{F} = \mathcal{G}_{m+1} + O_{m+1} + \mathcal{L}_{m+1}^{\text{even}} \mathcal{F},$$

where O_{m+1} is the differential polynomial given by

$$\begin{aligned} O_{m+1} &= \frac{1}{m-1} \left(\frac{\partial O_m}{\partial s_1} - a_1^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial O_m}{\partial t^{\beta,q}} + f_{\beta,q} \frac{\partial O_m}{\partial t^{\alpha,p}} + \frac{\partial^2 O_m}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) - \mathcal{L}_1^{\text{even}} O_m \right) \\ &\quad - \frac{1}{m-1} \left(\frac{\partial O_1}{\partial s_m} - a_m^{\alpha,p;\beta,q} \left(f_{\alpha,p} \frac{\partial O_1}{\partial t^{\beta,q}} + f_{\beta,q} \frac{\partial O_1}{\partial t^{\alpha,p}} + \frac{\partial^2 O_1}{\partial t^{\alpha,p} \partial t^{\beta,q}} \right) - \mathcal{L}_m^{\text{even}} O_1 \right). \end{aligned}$$

Thus we have defined recursively an infinite set of symmetries of the tau-cover of the deformed Principal Hierarchy. Their actions on \mathcal{F} can be represented by

$$\frac{\partial \mathcal{F}}{\partial s_m} = \mathcal{G}_m + O_m + \mathcal{L}_m^{\text{even}} \mathcal{F}.$$

Next we show that we can further adjust O_m by adding certain constants such that these symmetries satisfy the Virasoro commutation relation

$$\left[\frac{\partial}{\partial s_k}, \frac{\partial}{\partial s_l} \right] = (l - k) \frac{\partial}{\partial s_{k+l}}, \quad k, l \geq -1.$$

Lemma 4.24. *There is a unique choice of constants κ_m for $m \geq -1$ such that the flows*

$$\frac{\partial \mathcal{F}}{\partial s_m} = \mathcal{G}_m + O_m + \kappa_m + \mathcal{L}_m^{\text{even}} \mathcal{F}$$

satisfy the Virasoro commutation relation

$$(4.24) \quad \left[\frac{\partial}{\partial s_k}, \frac{\partial}{\partial s_l} \right] = (l - k) \frac{\partial}{\partial s_{k+l}}, \quad k, l \geq -1.$$

Proof. Let us first fix an arbitrary choice of O_m and denote

$$\frac{\partial \mathcal{F}}{\partial \hat{s}_m} = \mathcal{G}_m + O_m + \mathcal{L}_m^{\text{even}} \mathcal{F}.$$

Then we obtain the differential polynomials \tilde{O}_{k+l} such that

$$\left[\frac{\partial}{\partial \hat{s}_k}, \frac{\partial}{\partial \hat{s}_l} \right] \mathcal{F} = (l - k) \left(\mathcal{G}_{l+k} + \tilde{O}_{l+k} + \mathcal{L}_{l+k}^{\text{even}} \mathcal{F} \right).$$

But both $\frac{\partial}{\partial \hat{s}_{l+k}}$ and $\left[\frac{\partial}{\partial \hat{s}_k}, \frac{\partial}{\partial \hat{s}_l} \right]$ are symmetries of the tau-cover of the deformed Principal Hierarchy, so we conclude that

$$\frac{\partial \mathcal{F}}{\partial s} := \frac{\partial \mathcal{F}}{\partial \hat{s}_{k+l}} - \frac{1}{l - k} \left[\frac{\partial}{\partial \hat{s}_k}, \frac{\partial}{\partial \hat{s}_l} \right] \mathcal{F} = O_{k+l} - \tilde{O}_{k+l}$$

is also a symmetry of the tau-cover of the deformed Principal Hierarchy. The action of this symmetry on the normal coordinates has the expression

$$\frac{\partial w_\lambda}{\partial s} = \frac{\partial^2}{\partial t^{\lambda,0} \partial t^{1,0}} \left(O_{k+l} - \tilde{O}_{k+l} \right).$$

Thus $\frac{\partial w_\lambda}{\partial s} \in \hat{\mathcal{A}}_{\geq 2}$, and therefore such a symmetry must vanish due to the result of the bihamiltonian cohomology [7]. Hence we conclude that

$$O_{k+l} - \tilde{O}_{k+l} = c_{k,l}$$

for some constant $c_{k,l}$, and this means that

$$(4.25) \quad \left[\frac{\partial}{\partial \hat{s}_k}, \frac{\partial}{\partial \hat{s}_l} \right] = (l - k) \frac{\partial}{\partial \hat{s}_{k+l}} + c_{k,l}, \quad k, l \geq -1.$$

Let us denote by \mathfrak{W}_1 the Lie algebra of formal vector fields on a line, which is an infinite dimensional Lie algebra with a basis

$$e_m = z^{m+1} \frac{d}{dz}, \quad m \geq -1.$$

Then the relation (4.25) implies that the Lie algebra $\left\{ \frac{\partial}{\partial \hat{s}_m} \right\}$ defines a central extension of \mathfrak{W}_1 . It is computed in [15] that $H^2(\mathfrak{W}_1, \mathbb{R}) = 0$ and hence every central extension is

trivial. Therefore we can modify each O_m by adding an appropriate constant κ_m such that the modified flows

$$\frac{\partial \mathcal{F}}{\partial s_m} = \mathcal{G}_m + O_m + \kappa_m + \mathcal{L}_m^{\text{even}} \mathcal{F}$$

satisfy the commutation relations (4.24). Moreover, the choice of κ_m is unique since $H^1(\mathfrak{W}_1, \mathbb{R}) = 0$ (see [15]). The lemma is proved. \square

Thus we have proved the following theorem.

Theorem 4.25. *For every tau-symmetric bihamiltonian deformation of the Principal Hierarchy associated with a semisimple Frobenius manifold, the deformed integrable hierarchy possesses an infinite set of Virasoro symmetries. The actions of these symmetries on the tau function \mathcal{Z} are represented by*

$$\frac{\partial \mathcal{Z}}{\partial s_m} = L_m^{\text{even}} \mathcal{Z} + O_m \mathcal{Z}, \quad m \geq -1,$$

where $O_m \in \hat{\mathcal{A}}$ are certain differential polynomials, and the flows $\frac{\partial}{\partial s_m}$ satisfy the commutation relations

$$\left[\frac{\partial}{\partial s_k}, \frac{\partial}{\partial s_l} \right] = (l - k) \frac{\partial}{\partial s_{k+l}}, \quad k, l \geq -1.$$

Example 4.1. *Let M be the 2-dimensional Frobenius manifold defined on the orbit space of the Weyl group of type B_2 . Its potential and Euler vector field are given by*

$$F = \frac{1}{2}v^2u + \frac{4}{15}u^5, \quad E = v\partial_v + \frac{1}{2}u\partial_u.$$

Here $v = v^1$ and $u = v^2$ are the flat coordinates of M . We denote by σ_1 and σ_2 the dual coordinates of the fiber of \hat{M} , then the bihamiltonian structure $(P_0^{[0]}, P_1^{[0]})$ associated with M is given by

$$P_0^{[0]} = \frac{1}{2} \int \sigma_1 \sigma_2^1 + \sigma_2 \sigma_1^1, \quad P_1^{[0]} = \frac{1}{2} \int 8u^3 \sigma_1 \sigma_1^1 + \frac{1}{2} u \sigma_2 \sigma_2^1 + 2v \sigma_1 \sigma_2^1 - \frac{1}{2} v_x \sigma_1 \sigma_2.$$

Let us first write down the Virasoro symmetry $\frac{\partial}{\partial s_1}$ of the tau-cover of the Principal Hierarchy associated with M . The Virasoro operator L_1^{even} has the expression

$$L_1^{\text{even}} = \frac{3}{16} \frac{\partial^2}{\partial t^{1,0} \partial t^{2,0}} + \mathcal{L}_1^{\text{even}},$$

where

$$\mathcal{L}_1^{\text{even}} = \sum_{p \geq 0} \left(p + \frac{1}{4} \right) \left(p + \frac{5}{4} \right) t^{1,p} \frac{\partial}{\partial t^{1,p+1}} + \left(p + \frac{3}{4} \right) \left(p + \frac{7}{4} \right) t^{2,p} \frac{\partial}{\partial t^{2,p+1}}.$$

Then the action of $\frac{\partial}{\partial s_1}$ on the genus zero free energy $\mathcal{F}^{[0]}$ of the tau-cover of the Principal Hierarchy is given by

$$\frac{\partial \mathcal{F}^{[0]}}{\partial s_1} = \frac{3}{16} f_{1,0} f_{2,0} + \mathcal{L}_1^{\text{even}} \mathcal{F}^{[0]}.$$

Consider the bihamiltonian structure (P_0, P_1) of the Drinfel'd-Sokolov hierarchy [3] associated with the untwisted affine Kac-Moody algebra $B_2^{(1)}$. After performing a suitable

Miura type transformation we have $P_0 = P_0^{[0]}$, and the Hamiltonian operator \mathcal{P}_1 of P_1 has the expression

$$\mathcal{P}_1 = \begin{pmatrix} 8u^3\partial_x + 12u^2u_x & v\partial_x + \frac{1}{4}v_x \\ v\partial_x + \frac{3}{4}v_x & \frac{1}{2}u\partial_x + \frac{1}{4}u_x \end{pmatrix} + \varepsilon^2 \begin{pmatrix} D_1 & D_2 \\ D_3 & D_4 \end{pmatrix} + O(\varepsilon^4),$$

where the differential operators D_i are given by $D_2 = u\partial_x^3 + \frac{3}{4}u_x\partial_x^2$, $D_4 = \frac{5}{8}\partial_x^3$ and

$$D_1 = 14u^2\partial_x^3 + 42uu_x\partial_x^2 + \left(20u_x^2 + 16uu_{xx} + \frac{1}{2}v_{xx}\right)\partial_x + 12u_xu_{xx} + 6uu^{(3)} + \frac{1}{4}v^{(3)},$$

$$D_3 = u\partial_x^3 + \frac{9}{4}u_x\partial_x^2 + \frac{3}{2}u_{xx}\partial_x + \frac{1}{4}u^{(3)}.$$

The bihamiltonian structure (P_0, P_1) is a deformation of $(P_0^{[0]}, P_1^{[0]})$ with central invariants $c_1 = \frac{1}{6}$, $c_2 = \frac{1}{12}$ (see [8, 21]), and it determines a unique deformation of the Principal Hierarchy associated with M . We can find the Virasoro symmetry $\frac{\partial}{\partial s_1}$ of the tau-cover of the deformed Principal Hierarchy by using the results developed in the present paper. It turns out that the action of $\frac{\partial}{\partial s_1}$ on the tau-function \mathcal{Z} can be represented by

$$(4.26) \quad \frac{\partial \mathcal{Z}}{\partial s_1} = L_1^{\text{even}} \mathcal{Z} + \left(\frac{1}{2}u^2 + \frac{1}{4}v + \frac{1}{4}\varepsilon^2 u_{xx} \right) \mathcal{Z}.$$

A similar result is also given in the Example 5.5 of [26] by using the Kac-Moody-Virasoro algebra.

5. CONCLUSION

In the present paper, we prove the existence of an infinite set of Virasoro symmetries for a given tau-symmetric bihamiltonian deformation of the Principal Hierarchy associated with a semisimple Frobenius manifold. These symmetries can be represented in terms of the tau-function Z of the integrable hierarchy in the form

$$(5.1) \quad \frac{\partial Z}{\partial s_m} = L_m Z + O_m Z, \quad m \geq -1.$$

Note that the differential polynomials O_m depend on the choice of the representative, in the equivalence class of Miura type transformations, of the deformations of the bihamiltonian structure of hydrodynamic type $(P_0^{[0]}, P_1^{[0]})$. It is proved in [9] that, for two different choices of representatives (P_0, P_1) and $(\tilde{P}_0, \tilde{P}_1)$, the corresponding normal coordinates

$$w^\alpha = \eta^{\alpha\beta} \frac{\partial^2 \log Z}{\partial t^{1,0} \partial t^{\beta,0}}, \quad \tilde{w}^\alpha = \eta^{\alpha\beta} \frac{\partial^2 \log \tilde{Z}}{\partial t^{1,0} \partial t^{\beta,0}}, \quad \alpha = 1, \dots, n$$

of the deformed Principal Hierarchy are related by a Miura type transformation

$$\tilde{w}^\alpha = w^\alpha + \eta^{\alpha\beta} \frac{\partial^2 G}{\partial t^{1,0} \partial t^{\beta,0}},$$

where $G \in \hat{\mathcal{A}}$ is a differential polynomial, and the tau-functions are related by the equation

$$(5.2) \quad \tilde{Z} = \exp(G)Z.$$

Conversely, any differential polynomial $G \in \hat{\mathcal{A}}$ defines a Miura type transformation for the deformed bihamiltonian structure and the integrable hierarchy in the manner described above.

After a Miura type transformation induced from (5.2), the Virasoro symmetries (5.1) are transformed to the form

$$\frac{\partial \tilde{Z}}{\partial s_m} = L_m \tilde{Z} + \tilde{O}_m \tilde{Z}, \quad m \geq -1,$$

where the differential polynomials \tilde{O}_m can be computed from O_m and G .

We are going to study the problem of linearization of Virasoro symmetries in subsequent work, i.e., to study whether it is possible to find a suitable differential polynomial G such that all the functions \tilde{O}_m vanish.

Let us exam the possibility of linearizing the Virasoro symmetries given in the example of Sect. 3.4 for the one-dimensional Frobenius manifold. We want to find a certain Miura type transformation given by (5.2) which linearizes the Virasoro symmetry (3.26) and leaves the expression of the Virasoro symmetry

$$\frac{\partial Z}{\partial s_{-1}} = L_{-1}^{even} Z$$

unchanged. It follows from these requirements that the differential degree zero component G_0 of G must satisfy the equations

$$\frac{\partial G_0}{\partial v} = 0, \quad v^3 \frac{\partial G_0}{\partial v} = - \left(3c - \frac{3}{8} \right) \frac{v^2}{2},$$

which do not possess any solution unless $c = \frac{1}{8}$. When $c = \frac{1}{8}$, the linearized Virasoro symmetries for this example are well known [25], and the central invariant of the corresponding deformed bihamiltonian structure is $\frac{1}{3}c = \frac{1}{24}$.

We can do a similar computation for Example 4.1. We want to find a Miura type transformation given by (5.2) to linearize the Virasoro symmetry (4.26) and to preserve the expression of the Virasoro symmetry

$$\frac{\partial Z}{\partial s_{-1}} = L_{-1}^{even} Z.$$

Then the differential degree zero component G_0 of G must satisfy the equations

$$\frac{\partial G_0}{\partial v} = 0, \quad uv \frac{\partial G_0}{\partial u} = - \left(\frac{1}{2}u^2 + \frac{1}{4}v \right),$$

which have no solution. Therefore the Virasoro symmetries given by the bihamiltonian structure (P_0, P_1) in this example cannot be linearized.

In general we have the following conjecture.

Conjecture 5.1. *The Virasoro symmetries for a given tau-symmetric bihamiltonian deformation of the Principal Hierarchy associated with a semisimple Frobenius manifold is linearizable if and only if the central invariants of the corresponding deformed bihamiltonian structure are all equal to $\frac{1}{24}$.*

We will study this conjecture and give a proof of it in the paper [18].

REFERENCES

- [1] BECKER, K., AND BECKER, M. Non-perturbative solution of the super-Virasoro constraints. *Modern Phys. Lett. A* 8, 13 (1993), 1205–1214.
- [2] DIJKGRAAF, R., VERLINDE, H., AND VERLINDE, E. Topological strings in $d < 1$. *Nuclear Phys. B* 352, 1 (1991), 59–86.
- [3] DRINFEL'D, V. G., AND SOKOLOV, V. V. Lie algebras and equations of Korteweg-de Vries type. *J. Soviet Math.* 30, 2 (1985), 1975–2036.
- [4] DUBROVIN, B. Integrable systems and classification of 2-dimensional topological field theories. In *Integrable systems*. Springer, 1993, pp. 313–359.
- [5] DUBROVIN, B. Geometry of 2D topological field theories. In *Integrable systems and quantum groups*. Springer, 1996, pp. 120–348.
- [6] DUBROVIN, B. Painlevé transcendents in two-dimensional topological field theory. In *The Painlevé property*. Springer, 1999, pp. 287–412.
- [7] DUBROVIN, B., LIU, S.-Q., AND ZHANG, Y. On Hamiltonian perturbations of hyperbolic systems of conservation laws I: Quasi-Triviality of bi-Hamiltonian perturbations. *Comm. Pure Appl. Math.* 59, 4 (2006), 559–615.
- [8] DUBROVIN, B., LIU, S.-Q., AND ZHANG, Y. Frobenius manifolds and central invariants for the Drinfeld-Sokolov biHamiltonian structures. *Adv. Math.* 219, 3 (2008), 780–837.
- [9] DUBROVIN, B., LIU, S.-Q., AND ZHANG, Y. Bihamiltonian cohomologies and integrable hierarchies II: the tau structures. *Comm. Math. Phys.* 361, 2 (2018), 467–524.
- [10] DUBROVIN, B., AND NOVIKOV, S. Hamiltonian Formalism of One-Dimensional Systems of Hydrodynamic Type, and the Bogolyubov-Whitman Averaging Method. *Sov. Math. Dokl.* (1983), 665–669.
- [11] DUBROVIN, B., AND ZHANG, Y. Frobenius manifolds and Virasoro constraints. *Selecta Math.* 5, 4 (1999), 423–466.
- [12] DUBROVIN, B., AND ZHANG, Y. Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants. *arXiv:math/0108160v1 [math.DG]* (2001).
- [13] FALQUI, G., AND LORENZONI, P. Exact poisson pencils, τ -structures and topological hierarchies. *Phys. D* 241, 23-24 (2012), 2178–2187.
- [14] FERAPONTOV, E. Compatible Poisson brackets of hydrodynamic type. *J. Phys. A* 34, 11 (2001), 2377.
- [15] GELFAND, I. M., AND FUKS, D. B. Cohomology of Lie algebras of formal vector fields. *Izv. Akad. Nauk SSSR* 34, 2 (1970), 322–337.
- [16] KONTSEVICH, M. Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.* 147, 1 (1992), 1–23.
- [17] LIU, S.-Q. Lecture notes on bihamiltonian structures and their central invariants. In *B-Model Gromov-Witten Theory*. Springer, 2018, pp. 573–625.
- [18] LIU, S.-Q., WANG, Z., AND ZHANG, Y. Linearization of Virasoro symmetries associated with semisimple Frobenius manifolds. *In preparation*.
- [19] LIU, S.-Q., WANG, Z., AND ZHANG, Y. Super tau-covers of bihamiltonian integrable hierarchies. *arXiv:2009.01143 [math.DG]* (2020).
- [20] LIU, S.-Q., WANG, Z., AND ZHANG, Y. Variational bihamiltonian cohomologies and integrable hierarchies I: foundations. *arXiv:2106.13038 [math.DG]* (2021).
- [21] LIU, S.-Q., AND ZHANG, Y. Deformations of semisimple bihamiltonian structures of hydrodynamic type. *J. Geom. Phys.* 54, 4 (2005), 427–453.
- [22] LIU, S.-Q., AND ZHANG, Y. Jacobi structures of evolutionary partial differential equations. *Adv. Math.* 227, 1 (2011), 73–130.
- [23] LIU, S.-Q., AND ZHANG, Y. Bihamiltonian cohomologies and integrable hierarchies I: a special case. *Comm. Math. Phys.* 324, 3 (2013), 897–935.
- [24] WITTEN, E. On the structure of the topological phase of two-dimensional gravity. *Nuclear Phys. B* 340, 2-3 (1990), 281–332.
- [25] WITTEN, E. Two-dimensional gravity and intersection theory on moduli space. *Surv. Differ. Geom.* 1, 1 (1990), 243–310.

- [26] Wu, C.-Z. Tau functions and Virasoro symmetries for Drinfeld–Sokolov hierarchies. *Adv. Math.* 306 (2017), 603–652.

Si-Qi Liu,
 Department of Mathematical Sciences, Tsinghua University
 Beijing 100084, P.R. China
 liusq@tsinghua.edu.cn

Zhe Wang,
 Department of Mathematical Sciences, Tsinghua University
 Beijing 100084, P.R. China
 zhe-wang17@mails.tsinghua.edu.cn

Youjin Zhang,
 Department of Mathematical Sciences, Tsinghua University
 Beijing 100084, P.R. China
 youjin@tsinghua.edu.cn