

THE STRUCTURE OF CONNECTED (GRADED) HOPF ALGEBRAS REVISITED

C.-C. LI, G.-S. ZHOU

ABSTRACT. Let H be a connected graded Hopf algebra over a field of characteristic zero and K an arbitrary graded Hopf subalgebra of H . We show that there is a family of homogeneous elements of H and a total order on the index set that satisfy several desirable conditions, which reveal some interesting connections between H and K . As one of its consequences, we see that H is a graded iterated Hopf Ore extension of K of derivation type provided that H is of finite Gelfand-Kirillov dimension. The main tool of this work is Lyndon words, along the idea developed by Lu, Shen and the second-named author in [24].

INTRODUCTION

Throughout the paper, we work over a fixed field denoted by k which is of characteristic zero. All vector spaces, algebras, coalgebras and unadorned tensors are over k .

A *connected Hopf algebra* is a Hopf algebra with coradical of dimension one. Though the condition of being connected is quite restrictive, it has rich examples from diverse fields of mathematics. Two well-known classes of examples are the universal enveloping algebras of Lie algebras and the coordinate rings of unipotent algebraic groups, which are cocommutative and commutative respectively. Another celebrated class of examples appear in combinatorics, including the Hopf algebras of symmetric functions, of quasi-symmetric functions and of permutations [10, 1, 2]. These Hopf algebras tend to have bases naturally parametrized by combinatorial objects (such as partitions, compositions, permutations, etc.), and they are always of infinite Gelfand-Kirillov dimension (GK dimension, for short). Recently, connected Hopf algebras of GK dimensions up to 4 are completely classified by Wang, Zhang and Zhuang [27, 22], when the base field is algebraically closed. It is a part of the longstanding project of classifying Hopf algebras that satisfy some finiteness conditions. Some new interesting connected Hopf algebras occur in the classification results.

The motivation of this paper is to understand the structure of connected Hopf algebras. We do it by considering the subclass of *connected graded Hopf algebras*, which are Hopf algebras H equipped with a grading $H = \bigoplus_{n \geq 0} H_n$ such that H is simultaneously a graded algebra and a graded coalgebra, the antipode preserves the given grading and moreover H_0 is one dimensional. It is shown in [24, Theorem A] that every connected graded Hopf algebra has *PBW generators* which satisfy some desirable conditions. By PBW generators of an algebra A we mean a family of elements of A indexed by a totally ordered set such that the nondecreasing product of these indexed elements form a linear basis of A . In this paper, we prove a relative version of [24, Theorem A] as follows.

2010 *Mathematics Subject Classification.* 16T05, 68R15, 16P90, 16S15, 16W50.

Key words and phrases. connected Hopf algebra, Hopf Ore extension, Lyndon word, Gelfand-Kirillov dimension.

Theorem A. (Theorem 3.1) Let H be a connected graded Hopf algebra and K a graded Hopf subalgebra. Then there is an indexed family $(z_\gamma)_{\gamma \in \Gamma}$ of homogeneous elements of H and a total order \leq on Γ such that

- (1) $\Delta(z_\gamma) \in 1 \otimes z_\gamma + z_\gamma \otimes 1 + (H_+^{<\gamma} \otimes H_+^{<\gamma})_{\deg(z_\gamma)}$ for every index $\gamma \in \Gamma$, where $H^{<\gamma}$ is the subalgebra of H generated by K and $\{z_\delta \mid \delta \in \Gamma, \delta < \gamma\}$;
- (2) $[z_\gamma, H^{<\gamma}] \subseteq H^{<\gamma}$ for every index $\gamma \in \Gamma$;
- (3) $\{z_{\gamma_1} \cdots z_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n\}$ is a homogeneous basis of H as a left graded K -module as well as a right graded K -module;
- (4) $\text{GKdim } H = \text{GKdim } K + \#(\Gamma)$, where $\#(\Gamma)$ is the number of elements of Γ .

The above theorem has some interesting consequences. For example, it follows that every commutative connected (graded) Hopf algebra is a (graded) polynomial algebra over any of its (graded) Hopf subalgebras in some family of (graded) variables, see Corollary 3.2 and Corollary 3.5. As special cases, some classical results are reconfirmed: (1) every commutative affine connected Hopf algebra is a polynomial algebra [4, Theorem 0.1]; (2) the shuffle algebras are graded polynomial algebras [19, Theorem 3.1.1]; (3) the Hopf algebra of quasi-symmetric functions is a graded polynomial algebra over the Hopf algebra of symmetric functions [15, Corollary 2.2]. We expect more applications of Theorem A for the study of group schemes and combinatorial Hopf algebras.

Another interesting consequence of Theorem A concerns Hopf Ore extensions. This line of research begun in [18] and has been made great progress recently [5, 11, 6, 24]. Recall that an n -step iterated Hopf Ore extension (n -step IHOE, for short) of a Hopf algebra R is a Hopf algebra H which contains R as a Hopf subalgebra and there is a chain of Hopf subalgebras $R = H^0 \subset H^1 \subset \dots \subset H^n = H$ with each of the extensions $H^i \subset H^{i+1}$ being an Ore extension as algebras (Definition 3.6). According to [24, Theorem B], connected graded Hopf algebras of GK dimension d are all graded d -step IHOEs' of the base field. In this paper, we generalize [24, Theorem B] as follows.

Theorem B. (Theorem 3.7) Let H be a connected graded Hopf algebra of GK dimension d and K a graded Hopf subalgebra of GK dimension e . Then H is a graded $(d - e)$ -step IHOE of K of derivation type.

Note that connected Hopf algebras of finite GK dimension are not necessarily IHOEs' of the base field. However, the known counterexamples are all the universal enveloping algebras of certain Lie algebras [5, Example 3.1 (iv)]. By the classification results obtained in [27, 22], connected Hopf algebras of GK dimensions up to 4 are all IHOEs' of the universal enveloping algebra of its primitive space. In view of the above theorem, it is natural to ask the following question.

Question C. Is every connected Hopf algebra of finite GK dimension can be obtained as an IHOE of the universal enveloping algebra of its primitive space?

Now let us sketch the proof of Theorem A. First we choose a set X of homogeneous generators for H such that $X' := X \cap K$ generates K , and then fix a well order on X satisfying certain conditions related to X' . Denote by I (resp. J) the ideal of defining relations on X (resp. X') for H (resp. K). In addition,

let \mathcal{N}_I (resp. \mathcal{N}_J) be the set of I -irreducible (resp. J -irreducible) Lyndon words on X (resp. X'). By the combinatorial properties of Lyndon words and the standard bracketing on words, one may construct from X a new family of homogeneous generators $(z_\gamma)_{\gamma \in \mathcal{N}_I}$ for H . The coalgebra structure of H assures that this new family of generators satisfy some desirable conditions (Proposition 2.4). Note that several key parts of this observation was already obtained in [24, Proposition 2.4], which is not easy to prove and has its own idea trace back to [12, 21]. The decisive new observation for this work is the equality $\mathcal{N}_J = \mathcal{N}_I \cap \langle X' \rangle$, due to the feature of Lyndon words (Lemma 2.7). Combine the above-mentioned two observations as well as some others, we conclude that the family $(z_\gamma)_{\gamma \in \Gamma}$, where $\Gamma := \mathcal{N}_I \setminus \mathcal{N}_J$, together with the restriction of the lexicographic order on Γ satisfy the requirements.

The paper is organized as follows. In section 1, we recall the definition and basic facts of Lyndon words. In section 2, we study the structure of connected graded algebras that admit pseudo-comultiplications (Definition 2.1) by taking advantage of the excellent combinatorial properties of Lyndon words. This section is the technical part of the paper and as a matter of fact, Theorem A is a direct consequence of Theorem 2.2. The final section is devoted to study the structure of connected (graded) Hopf algebras. In particular, the above two main theorems are proved there.

1. PRELIMINARIES

In this section, we introduce some basic notations that will be used in the sequel, recall the definition of Lyndon words and present some of their well-known properties for reader's convenience. The materials of Lyndon words are mostly extracted or modified from [7, 12, 14, 24].

Throughout this section, X stands for a well-ordered alphabet. If $X = \{x_1, \dots, x_\theta\}$ for some positive integer θ , then X is tacitly equipped with the natural order $x_1 < \dots < x_\theta$. We denote by $\langle X \rangle$ the set of all words on X . The empty word is denoted by 1. The *length* of a word u is denoted by $|u|$.

Let u, v be two words on X . We call v a *factor* of u if there exist words w_1, w_2 on X such that $w_1 v w_2 = u$. If w_1 (resp. w_2) can be chosen to be the empty word then v is called a *prefix* (resp. *suffix*) of u . A factor (resp. prefix, resp. suffix) v of u is called *proper* if $v \neq u$. The *lexicographic order* on $\langle X \rangle$, denoted by \leq_{lex} , is defined as follows. For words $u, v \in \langle X \rangle$,

$$u <_{\text{lex}} v \iff \begin{cases} v \text{ is a proper prefix of } u, & \text{or} \\ u = rxs, v = ryt, & \text{with } x, y \in X; x < y; r, s, t \in \langle X \rangle. \end{cases}$$

Clearly, it is a total order compatible with the concatenation of words from left but not from right. For example if $x, y \in X$ with $x > y$, one has $x >_{\text{lex}} x^2$ but $xy <_{\text{lex}} x^2y$.

Lemma 1.1. *For any integer $n \geq 1$, the restriction of \leq_{lex} on the set of all words on X of degree less than or equal to n is a well order.*

Proof. It suffices to see every nonincreasing sequence $u_1 \geq_{\text{lex}} u_2 \geq_{\text{lex}} \dots$ of words of degree $\leq n$ stabilizes eventually. We prove it by induction on n . If $u_r \neq 1$ for some r then $u_s \neq 1$ for $s \geq r$, so we may assume first that all u_r are nonempty. Since \leq_{lex} coincides with \leq on X , the result is true for $n = 1$. For $n \geq 2$, consider the decompositions $u_r = v_r w_r$ for $r \geq 1$, where $v_r \in X$. Clearly,

$$v_1 \geq v_2 \geq v_3 \geq \dots$$

by the definition of \leq_{lex} . Since \leq is a well order on X , there exists $p \geq 1$ such that $v_p = v_{p+1} = \cdots$. So we get a nonincreasing sequence $w_p \geq_{\text{lex}} w_{p+1} \geq_{\text{lex}} \cdots$ of words of degree $\leq n-1$. By induction, there exists $q \geq p$ such that $w_q = w_{q+1} = \cdots$, hence $u_q = u_{q+1} = \cdots$. \square

Definition 1.2. A word $u \in \langle X \rangle$ is called *Lyndon* if u is nonempty and $u >_{\text{lex}} wv$ for every factorization $u = vw$ with v, w nonempty. The set of all Lyndon words on X is denoted by $\mathbb{L} = \mathbb{L}(X)$.

For a word u of length ≥ 2 , define $u_R \in \langle X \rangle$ to be the lexicographically largest proper suffix of u and define $u_L \in \langle X \rangle$ by the decomposition $u = u_L u_R$. The pair of words

$$\text{Sh}(u) := (u_L, u_R)$$

is called the *Shirshov factorization* of u . As an example, $\text{Sh}(x_2^2 x_1 x_2 x_1) = (x_2^2 x_1, x_2 x_1)$.

Lyndon words share many excellent combinatorial properties, we collect below some of them for reader's interests. Due to the different favor of the definition of the lexicographic order in literatures, the statements that presented may be adjusted accordingly from the given references.

Proposition 1.3.

- (L1) ([14, Proposition 5.1.2], [12, Lemma 2]) *A nonempty word u is Lyndon if and only if it is lexicographically greater than any one of its proper nonempty suffixes.*
- (L2) ([14, Proposition 5.1.3] & [7, Lemma 4.3 (3)]) *Let $u = w_1 w_2$, $v = w_2 w_3$ be Lyndon words. If $u >_{\text{lex}} v$ (which holds in priori when $w_2 \neq 1$), then $w_1 w_2 w_3$ is a Lyndon word.*
- (L3) ([14, Proposition 5.1.3]) *Let u be word of length ≥ 2 . Then u is a Lyndon word if and only if u_L and u_R are both Lyndon words and $u_L >_{\text{lex}} u_R$.*
- (L4) ([14, Proposition 5.1.4]) *Let u, v be Lyndon words such that $u >_{\text{lex}} v$. Then $\text{Sh}(uv) = (u, v)$ if and only if either u is a letter or u is of length ≥ 2 with $u_R \leq_{\text{lex}} v$.*
- (L5) ([14, Theorem 5.1.5]) *Every nonempty word u can be written uniquely as a (lexicographically) nondecreasing product $u = u_1 u_2 \cdots u_r$ of Lyndon words.*
- (L6) ([7, Lemma 4.5]) *Let $u = u_1 u_2 \cdots u_r$ be a nondecreasing product of Lyndon words. If v is a Lyndon factor of u then v is a factor of some u_i .* \square

Let $k\langle X \rangle$ be the free algebra on X , which has all words on X as a linear basis. Elements of $k\langle X \rangle$ are frequently called (noncommutative) polynomials on X (with coefficients in k). In the sequel, let us assume that the free algebra $k\langle X \rangle$ is connected graded with each letter homogeneous of positive degree. The degree of a homogeneous polynomial f is denoted by $\deg(f)$.

We need some elementary notions from (noncommutative) Groebner basis theory. The *graded lex order* on $\langle X \rangle$, denoted by \leq_{glex} , is defined as follows. For words u, v ,

$$u <_{\text{glex}} v \iff \begin{cases} \deg(u) < \deg(v), & \text{or} \\ \deg(u) = \deg(v) \text{ and } u <_{\text{lex}} v. \end{cases}$$

Clearly, it is an *admissible order* on $\langle X \rangle$, i.e. a well order on $\langle X \rangle$ that compatible with the concatenation of words from both sides. The *leading word* of a nonzero polynomial f , denoted by $\text{LW}(f)$, is the

largest word that occurs in f (with respect to \leq_{glex}). The coefficient of $\text{LW}(f)$ in f is called the *leading coefficient* of f . For any set $F \subseteq k\langle X \rangle$, we write $\text{LW}(F) := \{ \text{LW}(f) \mid f \in F, f \neq 0 \}$.

Let I be an ideal of $k\langle X \rangle$. A word on X is called *I -reducible* if it is the leading word of some polynomials in I . A word that is not I -reducible is called *I -irreducible*. An *obstruction* of I (with respect to \leq_{glex}) is an I -reducible word whose proper factors are all I -irreducible. We denote by \mathcal{N}_I the set of all I -irreducible Lyndon words, by \mathcal{O}_I the set of all obstructions of I , and by \mathcal{B}_I the set of all lexicographically nondecreasing product of Lyndon words in \mathcal{N}_I .

Lemma 1.4. *Let I be an ideal of $k\langle X \rangle$. The set of all I -irreducible words is included in \mathcal{B}_I , and these two sets are equal if and only if \mathcal{O}_I consists of Lyndon words.*

Proof. It is an easy consequence of Proposition 1.3 (L5, L6) and the fact that every I -reducible word has a factor in \mathcal{O}_I . \square

2. PSEUDO-COMULTIPLICATIONS ON CONNECTED GRADED ALGEBRAS

The primary interest of this work is the structure of connected (graded) Hopf algebras. The techniques that employed or developed in this paper are aimed to deal with it. In our process, it turns out that the coassociativity of the comultiplication plays no role in some key observations. In this section, we study the structure of connected graded algebras A which admit algebra homomorphisms $A \rightarrow A \otimes A$ that share some features of the comultiplication of connected graded Hopf algebras. One of the main tools is the Lyndon words reviewed in the previous section.

Recall that a *connected graded algebra* is an algebra A on which a decomposition $A = \bigoplus_{n \geq 0} A_n$ is specified such that $A_0 = k 1_A$ and $A_i \cdot A_j \subseteq A_{i+j}$ for $i, j \geq 0$. We write $A_+ := \bigoplus_{n \geq 1} A_n$ in the sequel.

Definition 2.1. Let $A = \bigoplus_{n \geq 0} A_n$ be a connected graded algebra. A *pseudo-comultiplication* on A is a homomorphism of graded algebras $\Delta : A \rightarrow A \otimes A$ such that

$$\Delta(a) \in 1 \otimes a + a \otimes 1 + (A_+ \otimes A_+)_{\deg(a)}$$

for every homogeneous element a of positive degree. Clearly, to check the above condition it suffices to show it for some family of homogeneous generators of A .

The whole section is devoted to prove the following result.

Theorem 2.2. *Let A be a connected graded algebra and $B \subseteq A$ a graded subalgebra. Let $\Delta : A \rightarrow A \otimes A$ be a pseudo-comultiplication on A such that $\Delta(B) \subseteq B \otimes B$. Then there is an indexed family $(z_\gamma)_{\gamma \in \Gamma}$ of homogeneous elements of A of positive degree and a total order \leq on Γ such that*

- (1) $\Delta(z_\gamma) \in 1 \otimes z_\gamma + z_\gamma \otimes 1 + (A_+^{<\gamma} \otimes A_+^{<\gamma})_{\deg(z_\gamma)}$ for every index $\gamma \in \Gamma$, where $A^{<\gamma}$ is the subalgebra of A generated by B and $\{ z_\delta \mid \delta \in \Gamma, \delta < \gamma \}$;
- (2) $[z_\gamma, A^{<\gamma}] \subseteq A^{<\gamma}$ for every index $\gamma \in \Gamma$;
- (3) $\{ z_{\gamma_1} \cdots z_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n \}$ is a homogeneous basis of A as a left graded B -module as well as a right graded B -module;
- (4) $\text{GKdim } A = \text{GKdim } B + \#(\Gamma)$, where $\#(\Gamma)$ is the number of elements of Γ .

The proof of this theorem will be addressed after some preparatory results.

Remark 2.3. The totally ordered set (Γ, \leq) appeared in Theorem 2.2 may have no smallest element. Let A be the universal enveloping algebra of the Lie algebra with basis $\{e_n\}_{n \geq 1}$ and Lie bracket $[e_i, e_j] = (j-i)e_{i+j}$. Then A is a connected graded algebra with e_n of degree n . Take B to be the trivial subalgebra $k \cdot 1_A$. Then the family $(e_n)_{n \in \Gamma}$ with $\Gamma = \{1, 2, 3, \dots\}$ and the total order \leq on Γ given by $1 > 2 > 3 > \dots$ satisfy the requirements. Clearly, in this case, (Γ, \leq) has no smallest element.

Let us proceed to show Theorem 2.2. We continue to use the notations and conventions employed in Section 1. So X stands for a well-ordered alphabet and the free algebra $k\langle X \rangle$ is connected graded with letters homogeneous of positive degrees. In addition to the Lyndon words, another brick to construct the required family of generators in Theorem 2.2 is the so-called *standard bracketing* on $\langle X \rangle$, which is the map $[-] : \langle X \rangle \rightarrow k\langle X \rangle$ defined as follows. First set $[1] = 1$ and $[x] := x$ for $x \in X$; and then for words u of length ≥ 2 , inductively set

$$[u] = \begin{cases} [[u_L], [u_R]], & u \text{ is Lyndon,} \\ [u_L][u_R], & u \text{ is not Lyndon.} \end{cases}$$

Here, $[-, -] : k\langle X \rangle \times k\langle X \rangle \rightarrow k\langle X \rangle$ stands for the commutator operation on $k\langle X \rangle$ defined by $[f, g] = fg - gf$. Note that $[u_1 u_2 \cdots u_n] = [u_1][u_2] \cdots [u_n]$ for Lyndon words $u_1 \leq_{\text{lex}} u_2 \leq_{\text{lex}} \cdots \leq_{\text{lex}} u_n$.

For a word w , we denote by $k\langle X|I \rangle^{<w}$ (resp. $k\langle X|I \rangle^{\leq w}$) the subalgebra of $k\langle X \rangle$ generated by

$$\{ [u] \mid u \in \mathcal{N}_I, u <_{\text{lex}} w \} \quad \left(\text{resp. } \{ [u] \mid u \in \mathcal{N}_I, u \leq_{\text{lex}} w \} \right).$$

Note that they are all graded subalgebras of $k\langle X \rangle$.

Proposition 2.4. *Let I be a homogeneous proper ideal of $k\langle X \rangle$ and $A := k\langle X \rangle/I$. Assume that there exists a pseudo-comultiplication Δ on $k\langle X \rangle$ such that $\Delta(I) \subseteq k\langle X \rangle \otimes I + I \otimes k\langle X \rangle$.*

- (1) \mathcal{B}_I equals to the set of I -irreducible words, and \mathcal{O}_I consists of Lyndon words.
- (2) $[v] \in k\langle X|I \rangle_{\deg(v)}^{<v} + I$ for every I -reducible Lyndon words v .
- (3) $\Delta([v]) \in 1 \otimes [v] + [v] \otimes 1 + ((k\langle X|I \rangle_+^{<v} + I) \otimes (k\langle X|I \rangle_+^{<v} + I))_{\deg(v)}$ for every $v \in \mathcal{N}_I$.
- (4) $[u][v] - [v][u] \in k\langle X|I \rangle_{\deg(uv)}^{\leq uv} + I$ for every pair $u, v \in \mathcal{N}_I$ with $u >_{\text{lex}} v$.
- (5) $\{ [w] + I \mid w \in \mathcal{B}_I \}$ is a homogeneous basis of A .
- (6) $\text{GKdim } A = \#(\mathcal{N}_I)$, where $\#(\mathcal{N}_I)$ is the number of elements of \mathcal{N}_I .

Proof. Parts (2), (4) and (5) are obtained in [24, Proposition 2.4], whose proof is a bit of complicate. By Part (5) and the fact that $\text{LW}([u]) = u$ for any word u on X , it is not hard to see that $\{ [w] + I \mid w \in \mathcal{B}_I \}$ is a basis of A . By Lemma 1.4, Part (1) follows. By [24, Proposition 2.3], one has

$$\Delta([v]) \in 1 \otimes [v] + [v] \otimes 1 + (k\langle X \rangle_+^{<v} \otimes k\langle X \rangle_+^{<v})_{\deg(v)},$$

where $k\langle X \rangle_+^{<v}$ is the subalgebra of $k\langle X \rangle$ generated by $\{ [u] \mid u <_{\text{lex}} v, u \in \mathbb{L} \}$. Note that

$$k\langle X|I \rangle_+^{<v} + I = k\langle X \rangle_+^{<v} + I$$

by Part (2), one then obtains Part (3). Finally we show Part (6). For any finite subset Ξ of \mathcal{N}_I , let A_Ξ be the subalgebra of A generated by $\{[v] + I \mid v \in \Xi\}$. For any integer n , let $d_\Xi(n)$ be the number of nondecreasing sequences (v_1, \dots, v_p) in Ξ such that $\deg(v_1 \cdots v_p) = n$. Then

$$\sum_{n \geq 0} \dim(A_\Xi)_n t^n \geq \sum_{n \geq 0} d_\Xi(n) t^n = \prod_{v \in \Xi} (1 - t^{\deg(v)})^{-1}$$

by Part (5) and a simple combinatorial argument, where the inequality means that the difference of the series has no negative coefficients. Consequently,

$$(*) \quad \text{GKdim}(A_\Xi) = \limsup_{n \rightarrow \infty} \log_n \sum_{i \leq n} \dim(A_\Xi)_i \geq \limsup_{n \rightarrow \infty} \log_n \sum_{i \leq n} d_\Xi(i) = \#(\Xi).$$

Here, the first equality is by [13, Lemma 6.1 (b)] and the second equality is by [3, Proposition 2.21]. It follows that if \mathcal{N}_I is infinite then $\text{GKdim}(A) \geq \text{GKdim}(A_\Xi) \geq \#(\Xi)$ for any finite subset Ξ of \mathcal{N}_I and hence $\text{GKdim}(A) = \infty$. If \mathcal{N}_I is finite then $A = A_{\mathcal{N}_I}$ and the inequality in (*) becomes an equality for $\Xi = \mathcal{N}_I$ by Part (5), whence $\text{GKdim}(A) = \text{GKdim}(A_{\mathcal{N}_I}) = \#(\mathcal{N}_I)$. \square

Remark 2.5. Connected graded algebras of the form $A = k\langle X \rangle / I$ with \mathcal{O}_I consists of Lyndon words are studied in [8, 23, 26], where some important homological and algebraic invariants of A are described in terms of \mathcal{N}_I . One interesting source of such algebras are Artin-Schelter regular algebras of global dimension ≤ 5 that satisfy some extra conditions [25, Theorem 8.1]. Proposition 2.4 provides us with another natural source of such algebras, including all connected graded Hopf algebras.

Lemma 2.6. *Let I be a homogeneous ideal of $k\langle X \rangle$ and $A := k\langle X \rangle / I$. Assume that there exists a pseudo-comultiplication Δ on $k\langle X \rangle$ such that $\Delta(I) \subseteq k\langle X \rangle \otimes I + I \otimes k\langle X \rangle$. Then for every sequence (w_1, \dots, w_m) of Lyndon words in \mathcal{N}_I , the coset $[w_1] \cdots [w_m] + I$ in A is a linear combination of cosets of the form $[v_1] \cdots [v_p] + I$ with $v_1, \dots, v_p \in \mathcal{N}_I$ satisfying that*

$$v_1 \leq_{\text{lex}} \cdots \leq_{\text{lex}} v_p \leq_{\text{lex}} \max\{w_1, \dots, w_m\} \quad \text{and} \quad \deg(v_1 \cdots v_p) = \deg(w_1 \cdots w_m),$$

where $\max\{w_1, \dots, w_m\}$ is the lexicographically maximal word among w_1, \dots, w_m .

Proof. Let $\mathcal{N}_I^{(\infty)}$ be the set of finite sequences in \mathcal{N}_I , including the empty sequence. Let \leq be the partial order on $\mathcal{N}_I^{(\infty)}$ defined as follows. For $a = (a_1, \dots, a_m)$ and $b = (b_1, \dots, b_n)$ in $\mathcal{N}_I^{(\infty)}$, $a < b$ if and only if $\sum_{i=1}^m \deg(a_i) = \sum_{j=1}^n \deg(b_j)$ but there exists an integer $p \leq \min\{m, n\}$ such that $a_i = b_i$ for $i < p$ and $a_p <_{\text{lex}} b_p$. Clearly, \leq is compatible with the left and right concatenation of finite sequences. In addition, \leq satisfies the descending chain condition by Lemma 1.1. Now we proceed to show the result by induction on (w_1, \dots, w_m) with respect to \leq . If (w_1, \dots, w_m) is nondecreasing, there is nothing to prove. Otherwise, assume $w_i >_{\text{lex}} w_{i+1}$ for some i . By Proposition 2.4 (4),

$$\begin{aligned} [w_1] \cdots [w_m] &\in [w_1] \cdots [w_{i-1}][w_{i+1}][w_i][w_{i+2}] \cdots [w_m] \\ &\quad + \sum_{(c_1, \dots, c_s)} \lambda_{(c_1, \dots, c_s)}^{w_i, w_{i+1}} [w_1] \cdots [w_{i-1}][c_1] \cdots [c_s][w_{i+2}] \cdots [w_m] + I, \end{aligned}$$

where (c_1, \dots, c_s) runs over finite sequences in \mathcal{N}_I such that

$$\deg(c_1 \cdots c_s) = \deg(w_i w_{i+1}) \quad \text{and} \quad \max\{c_1, \dots, c_s\} \leq_{\text{lex}} w_i w_{i+1} <_{\text{lex}} \max\{w_1, \dots, w_m\}.$$

It follows that $\max\{w_1, \dots, w_{i-1}, c_1, \dots, c_s, w_{i+2}, \dots, w_m\} \leq_{\text{lex}} \max\{w_1, \dots, w_m\}$ and

$$(w_1, \dots, w_{i-1}, c_1, \dots, c_s, w_{i+2}, \dots, w_m) < (w_1, \dots, w_m).$$

Also we have $(w_1, \dots, w_{i-1}, w_{i+1}, w_i, w_{i+2}, \dots, w_m) < (w_1, \dots, w_m)$. By the induction hypothesis, $[w_1] \cdots [w_m] + I$ has the required decomposition. \square

The following result is a crucial observation for the proof of Theorem 2.2. First let us recall more terminologies from standard Groebner basis theory. A set $F \subseteq k\langle X \rangle$ is called *reduced* (with respect to \leq_{glex}) if $0 \notin F$ and for every $f \in F$, the leading coefficient of f is 1 and all words occur in f has no factor in $\text{LW}(F \setminus \{f\})$. For an ideal I of $k\langle X \rangle$, a *Groebner basis* of I (with respect to \leq_{glex}) is a subset $\mathcal{G} \subseteq I$ such that every I -reducible word has a factor in $\text{LW}(\mathcal{G})$. It is well-known that every ideal admits a unique reduced Groebner basis [20, Proposition 14]. Note that all these definitions and facts as mentioned are workable with respect to any admissible orders on $\langle X \rangle$.

Lemma 2.7. *Let I be a homogeneous proper ideal of $k\langle X \rangle$, X' a subset of X and $J = I \cap k\langle X' \rangle$. Assume that $x_1 < x_2$ for every $x_1 \in X'$ and $x_2 \in X \setminus X'$, and assume there is a pseudo-comultiplication Δ on $k\langle X \rangle$ such that $\Delta(k\langle X' \rangle) \subseteq k\langle X' \rangle \otimes k\langle X' \rangle$ and $\Delta(I) \subseteq k\langle X \rangle \otimes I + I \otimes k\langle X \rangle$. Then*

$$\mathcal{N}_J = \mathcal{N}_I \cap \langle X' \rangle, \quad \mathcal{O}_J = \mathcal{O}_I \cap \langle X' \rangle \quad \text{and} \quad \mathcal{G}_J = \mathcal{G}_I \cap k\langle X' \rangle,$$

where \mathcal{G}_I (resp. \mathcal{G}_J) is the reduced Groebner basis of I (resp. J) with respect to \leq_{glex} .

Proof. First we show $\mathcal{N}_J = \mathcal{N}_I \cap \langle X' \rangle$. Let v be a Lyndon word on X' . Clearly, if v is J -reducible then it is I -reducible. Conversely, assume v is I -reducible. Since the biggest letter in a Lyndon word must occur in the left-most place, Lyndon words on X that lexicographically smaller than v are necessarily Lyndon words on X' . It is then clear that v is J -reducible by Proposition 2.4 (2).

Next we show $\mathcal{O}_J = \mathcal{O}_I \cap \langle X' \rangle$. By Proposition 2.4 (1), \mathcal{O}_I and \mathcal{O}_J consist of Lyndon words. Note that a word on X (resp. X') is I -irreducible (resp. J -irreducible) if and only if its Lyndon factors are all so. For any Lyndon word u on X' ,

$$\begin{aligned} u \in \mathcal{O}_J &\iff u \text{ is } J\text{-irreducible and its proper Lyndon factors are all } J\text{-irreducible} \\ &\iff u \text{ is } I\text{-irreducible and its proper Lyndon factors are all } I\text{-irreducible} \\ &\iff u \in \mathcal{O}_I. \end{aligned}$$

Here, the first and the third equivalences are clear, and the second equivalence is by $\mathcal{N}_J = \mathcal{N}_I \cap \langle X' \rangle$ shown as above. The desired equality follows directly.

Finally, we show $\mathcal{G}_J = \mathcal{G}_I \cap k\langle X' \rangle$. Since $\text{LW}(\mathcal{G}_I) = \mathcal{O}_I$ and $\text{LW}(\mathcal{G}_J) = \mathcal{O}_J$ by standard Groebner basis theory, they both consist of Lyndon words and $\text{LW}(\mathcal{G}_J) = \text{LW}(\mathcal{G}_I) \cap \langle X' \rangle$. For any word $u \in \text{LW}(\mathcal{G}_J)$, let f_u (resp. g_u) be the unique element of \mathcal{G}_J (resp. \mathcal{G}_I) with leading word u . By the definition of \mathcal{G}_J , all words occur in $f_u - u$ have no factor in $\text{LW}(\mathcal{G}_J)$, and hence they have no factor in $\text{LW}(\mathcal{G}_I)$. Thus $f_u - u$ is a linear combination of I -irreducible words. Consequently, $f_u = g_u$ for any $u \in \text{LW}(\mathcal{G}_J)$ and the desired equality follows directly. \square

Remark 2.8. For a proper ideal I of $k\langle X \rangle$ and a subset $X' \subset X$, knowing a generating set G for I , it is not necessarily true that $G' = G \cap k\langle X' \rangle$ is a generating set for $J = I \cap k\langle X' \rangle$. To make it happen, a well-known choice of G is an arbitrary Groebner basis of I with respect to an arbitrary admissible order σ on $\langle X \rangle$ that eliminates $X \setminus X'$ in the sense of [20, Definition 17]. It is because that G' is a Groebner basis of J with respect to the restriction of σ to $\langle X' \rangle$ (see [20, Proposition 19]). Nevertheless, Lemma 2.7 provides more choices of G when I satisfy specific conditions.

Now we are ready to prove Theorem 2.2.

Proof of Theorem 2.2. By choosing a set of homogeneous generators of A of positive degree which contains a generating set of B , we may fix a set X of graded variables, a decomposition $X = X' \cup X''$, and a homogeneous ideal I of $k\langle X \rangle$ such that $A = k\langle X \rangle / I$ as graded algebras, the cosets of variables in X' form a generating set of B and the cosets of variables in X'' lie in $A \setminus B$. Further, equip X with a well order \leq as follows. First fix a well order \leq'_r on $X'_r := \{x \in X' \mid \deg(x) = r\}$ and a well order \leq''_r on $X''_r := \{x \in X'' \mid \deg(x) = r\}$ for each integer $r \geq 1$; then for $x_1, x_2 \in X$,

$$x_1 \leq x_2 \iff \begin{cases} x_1 \in X' \text{ and } x_2 \in X'', & \text{or} \\ x_1, x_2 \in X' \text{ and } \begin{cases} \deg(x_1) < \deg(x_2), & \text{or} \\ \deg(x_1) = \deg(x_2) \text{ and } x_1 \leq'_{\deg(x_1)} x_2, \end{cases} & \text{or} \\ x_1, x_2 \in X'' \text{ and } \begin{cases} \deg(x_1) < \deg(x_2), & \text{or} \\ \deg(x_1) = \deg(x_2) \text{ and } x_1 \leq''_{\deg(x_1)} x_2, \end{cases} \end{cases}$$

In addition, one may readily lift Δ to a triangular comultiplication $\tilde{\Delta} : k\langle X \rangle \rightarrow k\langle X \rangle \otimes k\langle X \rangle$ such that

$$\tilde{\Delta}(k\langle X' \rangle) \subseteq k\langle X' \rangle \otimes k\langle X' \rangle \quad \text{and} \quad \tilde{\Delta}(I) \subseteq k\langle X \rangle \otimes I + I \otimes k\langle X \rangle.$$

Clearly, the canonical projection from $k\langle X \rangle$ to A induces a graded algebra isomorphism

$$k\langle X' \rangle / J \cong B, \quad \text{where } J = I \cap k\langle X' \rangle.$$

Let $\Gamma := \mathcal{N}_I \setminus \mathcal{N}_J$ and define \leq to be the restriction of \leq_{lex} on Γ . For every $\gamma \in \Gamma$, let

$$z_\gamma := [\gamma] + I.$$

Since $\mathcal{N}_J = \mathcal{N}_I \cap \langle X' \rangle$ by Lemma 2.7, the left-most letter of every Lyndon word in Γ must belong to X'' . Therefore, every Lyndon word in \mathcal{N}_J is lexicographically smaller than that in Γ . Note that B is generated by $\{[\nu] + I \mid \nu \in \mathcal{N}_J\}$, Parts (1) and (2) follow by Proposition 2.4 (3, 4). Compare the basis of A and B that constructed from \mathcal{N}_I and \mathcal{N}_J respectively as in Proposition 2.4 (5), it is easy to see that A is a free left B -module with a basis as the given set in Part (3). Since $\text{GKdim } A = \#(\mathcal{N}_I)$ and $\text{GKdim } B = \#(\mathcal{N}_J)$ by Proposition 2.4 (6), Part (4) follows directly.

Next we show A is spanned by the given set in Part (3) as a right B -module. For $m \geq 1$, let Γ_m be the set of all nondecreasing finite sequences $(\gamma_1, \dots, \gamma_q)$ in Γ with $\sum_{i=1}^q \deg(z_{\gamma_i}) \leq m$. Clearly,

$$A_m = \bigoplus_{(\gamma_1, \dots, \gamma_q) \in \Gamma_m} A_m^{(\gamma_1, \dots, \gamma_q)}, \quad \text{where } A_m^{(\gamma_1, \dots, \gamma_q)} := B_{m - \sum_{i=1}^q \deg(z_{\gamma_i})} \cdot (z_{\gamma_1} \cdots z_{\gamma_q}).$$

Let \leq_m be the total order on Γ_m defined as follows. For $\bar{\gamma} = (\gamma_1, \dots, \gamma_s)$ and $\bar{\delta} = (\delta_1, \dots, \delta_t)$ in Γ_m , $\bar{\gamma} <_m \bar{\delta}$ if and only if $\bar{\gamma}$ is a proper right part of $\bar{\delta}$ or there exists an integer $l \leq \min\{m, n\}$ such that $\gamma_{m-i} = \delta_{n-i}$ for $i < l$ and $\gamma_{m-l} < \delta_{n-l}$. By a similar argument as that in Lemma 1.1, it is not hard to see \leq_m is a well-order on Γ_m with the empty sequence \emptyset as the smallest element. Note that

$$A_m^\emptyset = B_m.$$

For any sequence $(\gamma_1, \dots, \gamma_s, \overbrace{\gamma, \dots, \gamma}^p) \in \Gamma_m$, where $\gamma_s < \gamma$ and $p \geq 1$, one has by Part (2) that

$$\lambda \cdot (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) \in (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) \cdot \lambda + \sum_{i=0}^{p-1} A^{<\gamma} \cdot z_\gamma^i, \quad \lambda \in B.$$

Since $A^{<\gamma}$ is spanned by

$$(2.1) \quad \{z_{\gamma_1} \cdots z_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n < \gamma\}$$

as a graded left B -module by Lemma 2.6, it follows that

$$A_m^{(\gamma_1, \dots, \gamma_s, \overbrace{\gamma, \dots, \gamma}^p)} \subseteq (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) \cdot B + \sum_{(\delta_1, \dots, \delta_t, \overbrace{\gamma, \dots, \gamma}^i)} A_m^{(\delta_1, \dots, \delta_t, \overbrace{\gamma, \dots, \gamma}^i)},$$

where $(\delta_1, \dots, \delta_t, \overbrace{\gamma, \dots, \gamma}^i)$ runs over elements in Γ_m with $i < p$ and $\delta_t < \gamma$. Note that

$$(\delta_1, \dots, \delta_t, \overbrace{\gamma, \dots, \gamma}^i) <_m (\gamma_1, \dots, \gamma_s, \overbrace{\gamma, \dots, \gamma}^p),$$

so by induction on Γ_m with respect to \leq_m , $A_m^{(\gamma_1, \dots, \gamma_q)}$ is contained in the span of the given set in Part (3) as a right B -module for all $(\gamma_1, \dots, \gamma_q) \in \Gamma_m$, and therefore $A = \sum_{m \geq 0} A_m$ is too.

Finally, we show the given set in Part (3) is right linearly independent over B . Assume not, then

$$\sum_{p=0}^m \sum_{(\gamma_1, \dots, \gamma_s)} (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) \cdot \lambda_{(\gamma_1, \dots, \gamma_s)}^{(p)} = 0, \quad \lambda_{(\gamma_1, \dots, \gamma_s)}^{(p)} \in B,$$

for some $\gamma \in \Gamma$ and $m \geq 1$, where $(\gamma_1, \dots, \gamma_s)$ runs over all nondecreasing finite sequences in $\{\delta \in \Gamma \mid \delta < \gamma\}$ and $\lambda_{(\gamma_1, \dots, \gamma_s)}^{(m)} \neq 0$ for some index $(\gamma_1, \dots, \gamma_s)$. Note that

$$(z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) \cdot \lambda_{(\gamma_1, \dots, \gamma_s)}^{(p)} \in \lambda_{(\gamma_1, \dots, \gamma_s)}^{(p)} \cdot (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^p) + \sum_{i=0}^{p-1} A^{<\gamma} \cdot z_\gamma^i,$$

so

$$0 \in \sum_{(\gamma_1, \dots, \gamma_s)} \lambda_{(\gamma_1, \dots, \gamma_s)}^{(m)} \cdot (z_{\gamma_1} \cdots z_{\gamma_s} z_\gamma^m) + \sum_{i=0}^{m-1} A^{<\gamma} \cdot z_\gamma^i.$$

Since the set (2.1) spans $A^{<\gamma}$ as a left B -module and the given set in Part (3) is left B -linearly independent, $\lambda_{(\gamma_1, \dots, \gamma_s)}^{(m)} = 0$ for all $(\gamma_1, \dots, \gamma_s)$, a contradiction. This completes the proof of Part (3). \square

3. APPLICATIONS TO CONNECTED (GRADED) HOPF ALGEBRAS

In this section we study the structure of connected (graded) Hopf algebras.

Let us begin by recalling some notations and definitions on Hopf algebras. For a general Hopf algebra H , the usual notations Δ_H , ε_H and S_H are employed to denote the comultiplication, counit and antipode of H respectively. The *coradical* of a Hopf algebra H is defined to be the sum of all simple subcoalgebras of H . It is denoted by $H_{(0)}$. Also, the *coradical filtration* of H ([16, Section 5.2]) is denoted by $\{H_{(n)}\}_{n \geq 0}$. Note that the notations for coradical and coradical filtration we used differ from that of [16]. For an element $a \in H$, let $\rho_H(a)$ be the smallest number n such that $a \in H_{(n)}$. A Hopf algebra is called *connected* if its coradical is one-dimensional.

By a *graded Hopf algebra* we mean a Hopf algebra H equipped with a grading $H = \bigoplus_{n \geq 0} H_n$ such that H is both a graded algebra and a graded coalgebra, and the antipode preserves the given grading. Such a graded Hopf algebra is called *connected* if $H_0 = k$. It is easy to show that, when H is a connected graded Hopf algebra, one has $\ker \varepsilon_H = H_+ := \bigoplus_{n \geq 1} H_n$ and

$$\Delta_H(x) \in x \otimes 1 + 1 \otimes x + (H_+ \otimes H_+)_n, \quad x \in H_n.$$

So the comultiplication of a connected graded Hopf algebra is a pseudo-comultiplication. Clearly connected graded Hopf algebras are connected Hopf algebras. For a connected Hopf algebra H , the coradical filtration $\{H_{(n)}\}_{n \geq 0}$ is a Hopf algebra filtration [16, p. 62], whence the associated graded space $\text{gr}_c(H) := \bigoplus_{n \geq 0} H_{(n)}/H_{(n-1)}$ is a connected graded Hopf algebra in the natural way.

Theorem 3.1. *Let H be a connected graded Hopf algebra and K a graded Hopf subalgebra. Then there is an indexed family $(z_\gamma)_{\gamma \in \Gamma}$ of homogeneous elements of H and a total order \leq on Γ such that*

- (1) $\Delta(z_\gamma) \in 1 \otimes z_\gamma + z_\gamma \otimes 1 + (H_+^{<\gamma} \otimes H_+^{<\gamma})_{\deg(z_\gamma)}$ for every index $\gamma \in \Gamma$, where $H^{<\gamma}$ is the subalgebra of H generated by K and $\{z_\delta \mid \delta \in \Gamma, \delta < \gamma\}$;
- (2) $[z_\gamma, H^{<\gamma}] \subseteq H^{<\gamma}$ for every index $\gamma \in \Gamma$;
- (3) $\{z_{\gamma_1} \cdots z_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n\}$ is a homogeneous basis of H as a left graded K -module as well as a right graded K -module;
- (4) $\text{GKdim } H = \text{GKdim } K + \#\Gamma$, where $\#\Gamma$ is the number of elements of Γ .

Proof. It is a direct consequence of Theorem 2.2. □

Corollary 3.2. *Let H be a commutative connected graded Hopf algebra and K a graded Hopf subalgebra. Then H is isomorphic as a graded algebra to the graded polynomial algebra over K in some family of graded variables.*

Proof. It is a direct consequence of Theorem 3.1 (3). □

Corollary 3.3. *Let H be a commutative Hopf algebra which is connected graded as an algebra. Let K be a Hopf subalgebra as well as a graded subalgebra. If $(H_+)^2 \cap K = (K_+)^2$ then H is isomorphic as a graded algebra to the graded polynomial algebra over K in some family of graded variables.*

Proof. Let $\mathfrak{m} := H_+ = \sum_{n \geq 1} H_n$ and $\mathfrak{n} := K_+ = \sum_{n \geq 1} K_n$. By the argument of [4, Lemma 2.1 (2)], to see the result we may assume $\mathfrak{m} = \ker(\varepsilon_H)$, in which case $\mathfrak{n} = \ker(\varepsilon_K)$. By [9, Lemma 3.3],

$A = \bigoplus_{i \geq 0} \mathfrak{m}^i / \mathfrak{m}^{i+1}$ and $B = \bigoplus_{i \geq 0} \mathfrak{n}^i / \mathfrak{n}^{i+1}$ are connected graded Hopf algebras with comultiplication and counit defined in [9, Lemma 3.2]. Since A (resp. B) is generated by $A_1 = \mathfrak{m} / \mathfrak{m}^2$ (resp. $B_1 = \mathfrak{n} / \mathfrak{n}^2$) and commutative, Theorem 3.1 (3) tells us that A (resp. B) is freely generated as a commutative algebra on any choice of bases for $A_1 = \mathfrak{m} / \mathfrak{m}^2$ (resp. $B_1 = \mathfrak{n} / \mathfrak{n}^2$). Assume $\mathfrak{m}^2 \cap K = \mathfrak{n}^2$. Then the canonical map $B_1 \rightarrow A_1$ is injective, so the canonical map $B \rightarrow A$ are injective homomorphism of graded Hopf algebras. Therefore, we may consider B as a graded Hopf subalgebra of A . Note that A is actually \mathbb{Z}^2 -graded as an algebra with $A_{(i,j)} = (\mathfrak{m}^i / \mathfrak{m}^{i+1})_j$, and B a \mathbb{Z}^2 -graded subalgebra. Choose an indexed family of homogeneous elements $\{a_\gamma\}_{\gamma \in \Gamma}$ of \mathfrak{m} such that the family of cosets $\{\overline{a_\gamma}\}_{\gamma \in \Gamma}$, where $\overline{a_\gamma} := a_\gamma + \mathfrak{m}^2$, forms a basis of a complement of B_1 in A_1 . Fix a total order \leq on Γ . Then the set

$$\{\overline{a_{\gamma_1}} \cdots \overline{a_{\gamma_n}} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n\}$$

is a homogeneous basis of A as a \mathbb{Z}^2 -graded B -module. By a standard application of filtered-graded method and the observation that $(\mathfrak{m}^i)_r = 0$ for integers $0 \leq r < i$, it is not hard to conclude that the set

$$\{a_{\gamma_1} \cdots a_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n\}$$

forms a homogeneous basis of H as a graded K -module. Since H is commutative, H is a graded polynomial algebra over K in some family of graded variables. \square

Let H be a connected Hopf algebra and K a Hopf subalgebra. By a *filter basis* of H as a left (resp. right) K -module we mean a basis $(a_i)_{i \in I}$ of H as a left (resp. right) K -module such that

$$H_{(n)} = \sum_{i \in I} K_{(n-\rho_H(a_i))} \cdot a_i \quad (\text{resp. } H_{(n)} = \sum_{i \in I} a_i \cdot K_{(n-\rho_H(a_i))}), \quad n \geq 0.$$

Corollary 3.4. *Let H be a connected Hopf algebra and K a Hopf subalgebra. Then there is an indexed family $(x_\gamma)_{\gamma \in \Gamma}$ of elements of $\ker(\varepsilon_H)$ and a total order \leq on Γ such that $\text{GKdim } H = \text{GKdim } K + \#\Gamma$, where $\#\Gamma$ is the number of elements of Γ , and $\{x_{\gamma_1} \cdots x_{\gamma_n} \mid n \geq 0, \gamma_1, \dots, \gamma_n \in \Gamma, \gamma_1 \leq \dots \leq \gamma_n\}$ is a filter basis of H as a left K -module as well as a right K -module.*

Proof. By [16, Lemma 5.2.12], we may consider $\text{gr}_c(K)$ as a Hopf subalgebra of $\text{gr}_c(H)$ in the natural way. Choose an indexed family $\{z_\gamma\}_{\gamma \in \Gamma}$ of homogeneous elements of $\text{gr}_c(H)$ of positive degrees and a total order \leq on Γ as in Theorem 3.1 for the pair $(\text{gr}_c(H), \text{gr}_c(K))$. Then, for each index γ with $\deg(z_\gamma) = n_\gamma$, pick an element $x_\gamma \in H_{(n_\gamma)}$ such that $x_\gamma + H_{(n_\gamma-1)} = z_\gamma$ in $\text{gr}_c(H)$. Replacing x_γ by $x_\gamma - \varepsilon_H(x_\gamma)$, one may assume $x_\gamma \in \ker(\varepsilon_H)$. Note that $\rho_H(x_\gamma) = n_\gamma$ and $\rho_H(x_{\gamma_1} \cdots x_{\gamma_n}) = \rho_H(x_{\gamma_1}) + \dots + \rho_H(x_{\gamma_n})$. By [17, Lemma I.5.1 (3)], the first two parts follow directly. In addition, by [24, Proposition 3.4] and Theorem 3.1 (4), one has $\text{GKdim } H = \text{GKdim } \text{gr}_c(H) = \text{GKdim } \text{gr}_c(K) + \#\Gamma = \text{GKdim } K + \#\Gamma$. \square

Corollary 3.5. *Let H be a commutative connected Hopf algebra and K a Hopf subalgebra. Then H is isomorphic as an algebra to the polynomial algebra over K in some family of variables.*

Proof. It is a direct consequence of Corollary 3.4 (1). \square

Now we turn to consider the structure of connected (graded) Hopf algebras of finite GK dimension. First we recall the definition of Hopf Ore extensions and iterated Hopf Ore extensions.

Definition 3.6. Let R be a Hopf algebra and $n \geq 2$.

- (1) A Hopf algebra H is called a *Hopf Ore extension (HOE, for short)* of R if it contains R as a Hopf subalgebra and is an Ore extension of R as algebras, that is $H = R[X; \sigma, \delta]$ for some algebra automorphism σ of R and σ -derivation δ of R . We call the HOE H is of *derivation type* if σ can be chosen to be the identity map of R .
- (2) A Hopf algebra H is called an *(n -step) iterated Hopf Ore extension ((n -step) IHOE, for short)* of R if it contains R as a Hopf subalgebra and there is a chain of Hopf subalgebras

$$R = H^0 \subset H^1 \subset \cdots \subset H^n = H$$

such that H^i is an Ore extension of H^{i-1} as algebras for $i = 1, \dots, n$. We call the IHOE H is of *derivation type* if all successive HOEs' can be chosen to be of derivation type .

By convention, a 0-step IHOE of R is just R itself; a 1-step IHOE of R is just a HOE of R . One similarly define the concept of *graded HOE* and *graded n -step IHOE* for graded Hopf algebras.

Theorem 3.7. *Let H be a connected graded Hopf algebra of GK dimension d and K a graded Hopf subalgebra of GK dimension e . Then H is a graded $(d - e)$ -step IHOE of K of derivation type.*

Proof. Let $l = d - e$. Fix an indexed family $\{z_\gamma\}_{\gamma \in \Gamma}$ of homogeneous elements of H of positive degrees and a total order \leq on Γ as in Theorem 3.1. Since $\#\Gamma = l$, we may assume $\Gamma = \{1, \dots, l\}$ equipped with the natural order. Let $H^0 = K$; let H^i be the subalgebra of H generated by K and z_1, \dots, z_i for $i = 1, \dots, l$. Clearly, H^i are all clearly Hopf subalgebras of H by Theorem 3.1 (1) and induction on i . Also, one may readily conclude by Theorem 3.1 (2, 3) and induction on i that the set

$$\{1, z_i, z_i^2, \dots\}$$

is a homogeneous basis of H^i as a graded left H^{i-1} -module for $i = 1, \dots, d$. In addition, according to 3.1 (2), one may define a graded map $\delta_i : H^{i-1} \rightarrow H^{i-1}$ of degree $\deg(z_i)$, $i = 1, \dots, d$, by

$$f \mapsto z_i \cdot f - f \cdot z_i, \quad f \in H^{i-1}.$$

It is easy to check that δ_i is a derivation of H^{i-1} and $H^i = H^{i-1}[z_i; \delta_i]$. □

Corollary 3.8. *Let H be a connected graded Hopf algebra of GK dimension d . Then H is a $(d - e)$ -step graded IHOE of the universal enveloping algebra of $P(H)$ of derivation type, where $P(H)$ is the primitive space of H and e is the dimension of $P(H)$.*

Proof. Note that the universal enveloping algebra of $P(H)$ is canonically a graded Hopf subalgebra of H and of GK dimension e . The result then follows directly by Theorem 3.7. □

Proposition 3.9. *Let H be a connected Hopf algebra of GK dimension d and K a Hopf subalgebra of GK dimension $d - 1$. Let r be the smallest number such that $H_{(r)} \neq K_{(r)}$. Assume that K is generated by $K_{(r-1)}$ as an algebra. Then H is a Hopf Ore extension of K .*

Proof. By Corollary 3.4, there's an element $x \in H$ satisfying the following conditions:

- (1) $\{1, x, x^2, \dots\}$ forms a basis of H as a left K -module;

- (2) $\{1, x, x^2, \dots\}$ forms a basis of H as a right K -module;
 (3) $H_{(n)} = \sum_{i \geq 0} K_{(n-\rho_H(x) \cdot i)} \cdot x^i = \sum_{i \geq 0} x^i \cdot K_{(n-\rho_H(x) \cdot i)}$ for $n \geq 0$.

Then clearly, $\rho_H(x) = r$,

$$x \cdot K_{(r-1)} \subseteq K_{(2r-1)} = K_{(r-1)} \cdot x + K_{(2r-1)} \quad \text{and} \quad K_{(r-1)} \cdot x \subseteq K_{(2r-1)} = x \cdot K_{(r-1)} + K_{(2r-1)}.$$

Since K is generated by $K_{(r-1)}$, it follows that

$$x \cdot K \subseteq K \cdot x + K \quad \text{and} \quad K \cdot x \subseteq x \cdot K + K.$$

By (1) and $x \cdot K \subseteq K \cdot x + K$, one obtains that $H = K[x; \sigma, \delta]$ for some endomorphism σ of K and σ -derivation δ of K . It remains to show σ is an automorphism. If $\sigma(a) = 0$ then $x \cdot a = \delta(a)$, and hence $a = 0$ by (2). So σ is injective. Since $K \cdot x \subseteq x \cdot K + K$, for any $a \in K$ one has

$$a \cdot x = x \cdot a_1 + a_2 = \sigma(a_1) \cdot x + \delta(a_1) + a_2$$

for some $a_1, a_2 \in K$. Then $a = \sigma(a_1)$ by (1). So σ is surjective. \square

Acknowledgments. The second-named author would like to thank Houyi Yu for helpful discussions, particularly for calling his attention to [15]. G.-S. Zhou is supported by the NSFC (Grant Nos. 11971141 & 11871186) and K. C. Wong Magna Fund in Ningbo University.

REFERENCES

- [1] M. Aguiar, N. Bergeron, F. Sottile, *Combinatorial Hopf algebras and generalized Dehn-Sommerville relations*, *Compositio Math.* **142** (2006), 1-30.
- [2] M. Aguiar, F. Sottile, *Structure of the Malvenuto-Reutenauer Hopf algebras of permutations*, *Adv. Math.* **191** (2005), 225-275.
- [3] M. Artin, J. Tate, M. Van den Bergh, *Modules over regular algebras of dimension 3*, *Invent. Math.* **106** (1991), no. 2, 335-388.
- [4] K. A. Brown, P. Gilmartin, J. J. Zhang, *Connected (graded) Hopf algebras*, *Trans. Amer. Math. Soc.* **372** (2019), 3283-3317.
- [5] K. A. Brown, S. O'Hagan, J. J. Zhang, G. Zhuang, *Connected Hopf algebras and iterated Ore extensions*, *J. Pure Appl. Alg.* **219** (2015), 2405-2433.
- [6] K. A. Brown, J. J. Zhang, *Iterated Hopf Ore extensions in positive characteristic*, Preprint, arXiv: 2004.14702 (2020). To appear in *J. Noncommut. Geom.*
- [7] T. Gateva-Ivanova, G. Fløystad, *Monomial algebras defined by Lyndon words*, *J. Alg.* **403** (2014), 470-496.
- [8] T. Gateva-Ivanova, *Algebras defined by Lyndon words and Artin-Schelter regularity*, Preprint, arXiv:1905.11281 (2019).
- [9] K. R. Goodearl, J. J. Zhang, *Noetherian Hopf algebra domains of Gelfand-Kirillov dimension two*, *J. Alg.* **324** (2010), 3131-3168.
- [10] D. Grinberg, V. Reiner, *Hopf algebras in combinatorics*, Preprint, arXiv: 1409.8356v7 (2020).
- [11] H. Huang, *Hopf Ore extensions*, *Alg. Represent. Theory* **23** (2020), 1477-1486.
- [12] V. K. Kharchenko, *A quantum analogue of Poincaré-Birkhoff-Witt theorem*, *Alg. Log.*, **38** (1999), 259-276.
- [13] G. R. Krause, T. H. Lenagan, *Growth of algebras and Gelfand-Kirillov dimension*, revised edition, *Graduate Studies in Mathematics*, Vol **22**, American Mathematical Society, Rhode Island, (1999).

- [14] M. Lothaire, *Algebraic combinatorics on words*, Encyclopedia in mathematics and its applications, **17** GC Rota, Editor Addison-Wesley Publishing Company, (1983).
- [15] C. Malvenuto, C. Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, J. Alg. **177** (1995), 967-982.
- [16] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics, **82**, American Mathematics Society, Providence, RI, (1993).
- [17] C. Năstăsescu, F. Van Oystaeyen, *Graded and filtered rings and modules*, Lect. Notes in Math. vol. 758, Springer Verlag (1979), Berlin.
- [18] A. N. Panov, *Ore extensions of Hopf algebras*, Mat. Zametki **74** (2003), 425-434.
- [19] D. E. Radford, *A natural ring basis for the Shuffle algebra and an application to group schemes*, J. Alg. **58**, (1979), 432-454
- [20] P. Nordbeck, *Cononical bases for algebraic computations*, Phd Thesis, Lund University, (2001).
- [21] S. Ufer, *PBW bases for a class of braided Hopf algebras*, J. Alg. **280** (2004), 84-119.
- [22] D.-G. Wang, J. J. Zhang, G. Zhuang, *Connected Hopf algebras of Gelfand-Kirillov dimension four*, Trans. Amer. Math. Soc. **367** (2015), 5597-5632.
- [23] G.-S. Zhou, *An application of Lyndon words in associative algebras*, (Chinese) Appl. Math. J. Chinese Univ. Ser. A **30** (2015), 245-252.
- [24] G.-S. Zhou, Y. Shen, D.-M. Lu, *The structure of connected (graded) Hopf algebras*, Adv. Math. **372** (2020), 107292.
- [25] G.-S. Zhou, D.-M. Lu, *Artin-Schelter regular algebras of dimension five with two generators*, J. Pure Appl. Alg. **218** (2014), 937-961.
- [26] G.-S. Zhou, D.-M. Lu, *Lyndon words for Artin-Schelter regular algebras*, Preprint, arXiv:1403.0385 (2014).
- [27] G. Zhuang, *Properties of pointed and connected Hopf algebras of finite Gelfand-Kirillov dimension*, J. Lond. Math. Soc. **87** (2013), 877-898.

Li

School of Mathematics and Statistics, Ningbo University, Ningbo 315211, China

E-mail: thesilencesea@gmail.com

Zhou

School of Mathematics and Statistics, Ningbo University, Ningbo 315211, China

E-mail: zhouguisong@nbu.edu.cn