

Incentives for Collective Innovation*

Gregorio Curello[†]

February 26, 2022

Abstract

Innovators with shared interests face a dynamic collective-action problem with two special features: uncertainty about the returns to (research) effort, and an exploration-exploitation trade-off in which exploitation becomes more attractive as discoveries accumulate. I study this problem in a general model with long-lived firms that exert effort to obtain improvements of random size in a technology they share. The firms' flow payoffs grow as the technology improves, but so does the marginal cost of effort. In (the unique symmetric Markov) equilibrium, small discoveries may hurt all firms simultaneously as they drastically reduce effort. Allowing each firm to discard the innovations it obtains (after observing their size) yields uniformly higher effort and welfare in equilibrium. If firms may instead conceal innovations from the others for a period of time, under linear payoffs, there exists an equilibrium in which improvements are kept secret until they reach a cutoff, and effort stops after disclosure. Although concealment is inefficient due to forgone benefits and the risk of redundancy, firms may be better off compared to the baseline equilibrium with forced disclosure.

1 Introduction

Technological innovation frequently has the features of a dynamic collective-action problem. Firms in the same 'network' gradually expand a stock of shared knowledge, concerning e.g. a production process, or a technology to be sold as part of a product.¹ Moreover,

*I thank Péter Eső, John Quah, Sven Rady, Meg Meyer, Francesc Dilmé, George Georgiadis, Johannes Hörner, Ludvig Sinander, Matteo Escudé, Mathijs Janssen, Claudia Herresthal, Alessio Piccolo, and the seminar participants at Oxford, Bonn, and Aalto universities for helpful comments.

[†]Institute for Microeconomics, University of Bonn. E-mail: gcurello@uni-bonn.de.

1. This occurs within R&D partnerships as well as, less formally, among firms with shared interests. See Powell and Giannella (2010) for a survey on networks of innovators and their 'collective' output, and Pénin (2007) for one on the free revealing of innovations. Recent empirical studies of knowledge spillovers among innovators include Akcigit et al. (2018) and Kerr (2008). Free revealing is characteristic of 'user' innovation as well: Harhoff and Lakhani (2016) is a recent analysis.

progress eventually decreases the incentive to contribute further.² This occurs in the biotechnology, semiconductor, and software industries, among others.³

However, collective innovation differs from traditional ‘games’ of public-good provision in at least two respects. First, the magnitude of discoveries is uncertain, as the value of new technologies is often difficult to predict. Second, firms face a resource trade-off between exploiting an existent technology and exploring new possibilities.⁴ In particular, as a technology improves, the opportunity cost of refining it rises.

In this paper, I study this problem in a general model that captures these key features. Long-lived identical agents exert hidden effort to induce increments in the stock of a public good. Increments have random size (their frequency being increasing with effort). Moreover, agents’ flow payoffs are a general function of their effort and of the current stock, and the marginal cost of effort increases (weakly) with the stock. In the main interpretation of the model, agents are firms and the public good is a technology that they share. The stock of the good corresponds to the current quality of the technology, and an increment in the stock is an innovation. Firms invest in R&D to obtain innovations, as well as in other activities (e.g. production, marketing, or product development). The latter induce a private flow of profits that grows as the technology improves.

I characterise the social-welfare benchmark (Theorem 1) and the unique symmetric Markov perfect equilibrium with the stock as the state variable (Theorem 2). In equilibrium, continuation payoffs may *fall* after an innovation (i.e. an increment in the stock), even though they necessarily increase in the welfare benchmark. This is because opponents exert less effort as the stock grows, delaying future innovations, and this may offset the benefit of a moderately larger stock. I derive a sufficient condition for innovations to be detrimental in this way with strictly positive probability (Proposition 1). Essentially, it suffices that the payoff function take a multiplicative form, and that innovations be sufficiently rare relative to the curvature-to-slope ratio of the cost of effort. In particular, if the condition is met, raising the value of innovations (either by altering how the stock

2. According to Powell and Giannella (2010): ‘Once knowledge accumulates to the stage that tangible outcomes are possible, private interests may take hold and commercialise particular streams of technology that emerge from collective invention [...] as technological uncertainty recedes, firms develop private R&D and focus on their own specific applications. Reliance on collective invention [...] wanes.’ Decreasing returns are ubiquitous in the literature on dynamic games of public-good provision, from Fershtman and Nitzan (1991) to Battaglini, Nunnari, and Palfrey (2014) (see Section 1.1 for a review).

3. See e.g. Powell, Koput, and Smith-Doerr (1996) for biotechnologies, and Chesbrough (2003) and Lim (2009) for the semiconductor industry. Von Hippel (1987) finds evidence of this phenomenon among steel producers. See the citations in footnote 1 for further evidence, and Bessen and Nuvolari (2016) for historical examples.

4. There is empirical evidence that financial constraints hold back innovation activities (e.g. Hottenrott and Peters 2012), and that the latter often rely on internal funds. Thus, firms face a trade-off between investing in innovation and other costly activities, such as production or marketing. Moreover, R&D resources must be split between the improvement of public technologies (which benefits all firms using them) and e.g. product development (which only benefits the firm engaging in it). A similar resource trade-off is omnipresent in games of strategic experimentation, a prominent way to model firms’ R&D activities (see Section 1.1 for a review).

enters the payoff function, or improving the distribution of their size) cannot restore monotonicity.

Because innovations have adverse incentive effects, agents who obtain them may wish to delay their disclosure and adoption. To explore this idea, I first enrich the model by allowing agents to *discard* innovations, after observing their size. While this is a strong assumption, it leads to sharp results.⁵ Namely, the ensuing game admits a unique symmetric Markov perfect equilibrium in which, at any given stock, both effort and continuation payoffs are higher than in the equilibrium of the baseline game with forced disclosure (Theorem 3). Moreover, if innovations are guaranteed to be beneficial when disclosure is compulsory, disposal does not occur even if allowed, and the equilibria coincide. Otherwise, the ex-ante payoffs in the equilibrium with disposal are strictly higher. Thus, in spite of the fact that discarding innovations is clearly inefficient, allowing the agents to do so enhances equilibrium welfare.

In reality, a firm that does not wish to disclose an innovation is more likely to conceal it than to discard it, and may disclose a refined version at a more profitable time. To check the robustness of the social benefits of selective disclosure, I enrich the baseline game by allowing any agent who obtains an innovation to conceal it from other agents, forgoing the larger payoff flow. At any later time, possibly after having obtained and concealed further innovations, the agent may disclose any portion of the *total* increment, increasing her payoffs and the public stock. Moreover, undisclosed increments obtained by different agents are ‘perfect substitutes’, so that concealment leads to redundant innovations.

If payoffs are linear, no symmetric Markov perfect equilibrium with full disclosure exists in this environment (Lemma 3). Given the richness of the setting, this raises the question of whether a tractable equilibrium exists. I show that this is the case. Namely, if payoffs have the aforementioned structure, there exists a symmetric perfect Bayesian equilibrium in which a single innovation is disclosed (the first to exceed a common, time-varying cutoff) and, after this occurs, no effort is exerted (Theorem 4). Although the agents’ payoff flow increases only after the disclosure, and all innovations except one are effectively redundant, ex-ante payoffs may exceed those of the equilibrium with forced disclosure. More specifically, this occurs as long as the number of agents is sufficiently large, and either the initial stock is sufficiently low, or innovations are sufficiently rare. Thus, even though concealing innovations is clearly inefficient, doing so is incentive-compatible and may improve equilibrium welfare.

The rest of the paper is organised as follows. I summarise the relevant literature in Section 1.1. Section 2 contains a description of the model and a discussion of the assumptions. In Section 3, I present the social-welfare benchmark. In Section 4, I characterise

5. Nothing would change if agents could instead conceal innovations, provided they cannot covertly refine the improvements that they hide. This is justified if innovations are aimed at solving a given problem (so that any two are perfect substitutes), and a new problem is tackled after a solution is adopted. See Section 5.1 for details.

the unique symmetric Markov perfect equilibrium. In Section 5, I analyse the game with disposal. In Section 6, I analyse the game with concealment.

1.1 Literature review

This paper belongs to the literature on dynamic games of public-good provision. The baseline model is closely related to Fershtman and Nitzan (1991), Marx and Matthews (2000), and Battaglini, Nunnari, and Palfrey (2014), as agents are long-lived and gradually add to the stock of the public good, and the returns to production decrease as the stock grows.⁶ I contribute to this literature in two ways. First, I impose weaker restrictions on the payoff function and, in particular, allow the production cost to depend on the current stock. An important insight is that, although a higher initial stock increases (ex-ante) payoffs in the social-welfare benchmark, it may lead to lower equilibrium payoffs, due to its adverse effect on incentives. Second, by allowing the stock to make discrete, randomly-sized jumps, I show that agents may have an incentive to discard or conceal increments in the stock,⁷ and I analyse the impact of allowing the agents to do so.

Another strand of the literature on dynamic public-good games assumes that agents are rewarded only once aggregate effort reaches a given threshold.⁸ Thus, even though free-riding remains a central concern, the incentive to exert effort *grows* with the total amount exerted to date. While this is not the case in the baseline model that I analyse, I show that a similar structure arises *in equilibrium* if agents are allowed to conceal the innovations that they obtain (Section 6). Namely, agents are rewarded as soon as the total increment obtained by one of them reaches a common threshold, and each agent exerts more effort as the size of her increment grows. However, in contrast to the literature, increments obtained by different agents are perfect substitutes.⁹

This paper is also related to the literature on strategic experimentation, even though it does not itself feature experimentation.¹⁰ Indeed, games of experimentation may be

6. In Bliss and Nalebuff (1984) and Gradstein (1992), a single unit of the good is valuable, and long-lived agents have private information about their heterogeneous costs of production. In Lockwood and Thomas (2002) and Matthews (2013), agents adjust the extent of their co-operation over time, and cannot ever reduce it.

7. To the best of my knowledge, studies on games of experimentation (which I discuss below) are the only ones within the literature on dynamic provision of public goods to feature either a production cost that varies with the stock, or production involving randomly-sized lumps. The (joint) study of these two features is justified since they are *necessary* to obtain the adverse effects just described, as I show in Appendix I.

8. Major contributions include Radner, Myerson, and Maskin (1986), Admati and Perry (1991), Strausz (1999), Compte and Jehiel (2004), Yildirim (2006), and Georgiadis (2015).

9. In Campbell, Ederer, and Spinnewijn (2014), agents obtain ‘successes’ at random times, and may conceal them in order to preserve their partners’ incentive to exert effort. Discarding or concealing innovations in my model is beneficial for the same reason. However, successes obtained by different agents in Campbell, Ederer, and Spinnewijn (2014) are perfect *complements*, whether or not they are concealed. As I show in Appendix I, if innovations have a fixed value in my model (as successes do in theirs), agents do not conceal them in equilibrium, as concealed increments are perfect substitutes.

10. Important contributions to this literature include Bolton and Harris (1999), Keller, Rady, and

viewed as a class of dynamic public-good games, where the ‘stock’ is the public belief about the quality of the risky arm, and ‘producing’ the good amounts to pulling this arm. Moreover, in these games as in my model, production is stochastic and the (net) production cost varies with the stock. An important feature distinguishing games of experimentation from the model I analyse (as well as from the rest of the literature on public-good games) is that payoffs and incentives to produce move jointly: ‘good news’ simultaneously makes agents better off and the risky arm more attractive.¹¹ This feature implies that equilibrium continuation payoffs are increasing in the probability that the risky arm is ‘good’. That is, the aforementioned phenomena that I analyse do not arise.¹²

Finally, the main interpretation of the model links it to the large theoretical literature on innovation. Reinganum (1983), part of the strand concerning industrial organisation, argued that industry leaders are likely to be overtaken by new entrants in the innovation race.¹³ An overtaking phenomenon also features in Jovanovic and Nyarko (1996) and Redding (2002), which belong to the literature on endogenous growth. Overtaking occurs in my model as well, and for similar reasons (Section 4.1). However, it is more severe than in previous analyses in that the ‘leader’ may be ex-ante worse-off than the ‘follower’.¹⁴ In the endogenous-growth models of Lucas and Moll (2014) and Perla and Tonetti (2014), agents face a resource trade-off between using an existing technology and improving it, exactly as in my model. However, innovations are always beneficial in their models, as aggregate technological progress is deterministic.

2 Model and assumptions

In this section, I describe the model and introduce a specific example which will be useful to illustrate the results. The section ends with a discussion of the assumptions.

Time is continuous and indexed by $t \in \mathbb{R}_+$. Each of $n \geq 2$ identical agents, indexed

Cripps (2005), Keller and Rady (2010), Bonatti and Hörner (2011), Klein and Rady (2011), Heidhues, Rady, and Strack (2015), and Keller and Rady (2015, 2020). Moreover, experimentation features in several papers on the economics of innovation, including Garfagnini and Strulovici (2016) and those cited in footnote 12.

11. In contrast, in standard public-good games as well as in my model, an increase in the stock is beneficial but discourages further production.

12. Nevertheless, concealing information plays an important role in models involving strategic experimentation. See e.g. Akcigit and Liu (2015), Halac, Kartik, and Liu (2017), and Bimpikis, Ehsani, and Mostagir (2019).

13. Important contributions to this literature include Brander and Spencer (1983), Spence (1984), Katz (1986), d’Aspremont and Jacquemin (1988), Kamien, Muller, and Zang (1992), Suzumura (1992), and Leahy and Neary (1997).

14. In detail, consider two groups of agents (e.g. two distinct R&D partnerships) playing the equilibrium of the baseline game (Theorem 2). If the first innovation obtained across groups makes the group who obtained it (the ‘leaders’) worse-off, the group of ‘followers’ (which therefore has a lower stock at the time of the innovation) is likely to have a higher stock in the near future. This is because the (symmetric) continuation payoffs of the followers are higher, even though their *flow* payoffs are lower as long as their stock lies below that of the leaders.

by $i \in \{1, \dots, n\}$, exerts costly effort to produce a public good. Let $x_t \geq 0$ denote the stock of the good at time $t \geq 0$. At any time $t \geq 0$, agent i exerts effort $a_t^i \in [0, 1]$ and receives a flow payoff $u(a_t^i, x_t)$, for some $u : [0, 1] \times \mathbb{R}_+ \rightarrow \mathbb{R}$. The payoff is discounted at rate $r > 0$.

The stock x_t takes some initial value $x_0 \geq 0$ and, for $t > 0$, is determined as follows. Agent i produces an increment in x_t with hazard rate λa_t^i , where $\lambda > 0$. Each increment has random size z , drawn from a CDF F with mean $\mu < \infty$ and such that $F(0) = 0$ and

$$F(z) > 0 \quad \text{for all } z > 0. \quad (1)$$

In words, increments of arbitrarily small (but strictly positive) size are possible. The production and the size of increments are independent across agents, time, and from each other. By rescaling the rate λ of arrival of increments by $1/r$, we may assume without loss of generality that $r = 1$. Write \tilde{x}_t for the random variable describing the stock at time $t \geq 0$, and x_t for its realisation.

Assume that u is twice continuously differentiable, and increasing and concave in x .¹⁵ That is, keeping the effort fixed, payoffs increase as the stock grows, but at a decreasing rate. Note that payoffs u may be expressed as

$$u(a, x) = b(x) - c(a, x)$$

where $b(x) := u(0, x)$ are gross benefits, and $c(a, x) := u(0, x) - u(a, x)$ is the opportunity cost of effort. Suppose that $c(a, x)$ is increasing and convex in a , and strictly increasing in a if $x > 0$. Suppose also that $c_1(a, x)$ and $c_{11}(a, x)$ are increasing in x ; that is, the cost of effort becomes steeper and more convex as the stock grows.¹⁶ Suppose also that

$$\lim_{x \rightarrow \infty} \lambda \mu n b'(x) - c_1(a, x) < 0 \quad \text{for all } 0 < a \leq 1. \quad (2)$$

This means that exerting any fixed amount of positive effort becomes inefficient as the stock diverges. In particular, $\lambda \mu n b'(x)$ approximates the marginal social benefit of effort when the stock x is large, whereas $c_1(a, x)$ is its marginal cost.¹⁷

I restrict agents to play Markov strategies, so that effort is pinned down by the current stock. Formally, a *Markov strategy* (for Agent i) is a Lebesgue measurable map $\alpha : \mathbb{R}_+ \rightarrow [0, 1]$ such that $a_t^i = \alpha(x_t)$ for all $t \geq 0$.¹⁸ If agents play a Markov profile $\alpha := (\alpha^i)_{i=1}^n$, the time- t continuation payoff to agent i is a function of the current stock

15. Notions such as ‘increasing’ and ‘concave’ are always meant in the weak sense.

16. The conditions on c are equivalent to the following conditions on u : u is decreasing and concave in a , and strictly decreasing in a if $x > 0$; u_1 and u_{11} are decreasing in x .

17. The (individual) benefit of obtaining one innovation, $\mathbb{E}_F[b(x + \tilde{z})] - b(x)$, is approximately $\mu b'(x)$ when x is large. This is because b is increasing and concave, hence approximately linear for large x .

18. Measurability ensures that the process $(\tilde{x}_t)_{t \geq 0}$ is well-defined for any Markov profile $(\alpha^i)_{i=1}^n$.

x_t , given by

$$v^i(x_t) := \mathbb{E} \left[\int_t^\infty r e^{-r(s-t)} [b(\tilde{x}_s) - c(\sigma^i(\tilde{x}_s), \tilde{x}_s)] ds \right] \quad (3)$$

where the dependence on α is implicit.¹⁹

In the main interpretation of the model, the stock x_t denotes the (quality of a shared) *technology*, and increments in x_t are *innovations*. These terms are used throughout the discussion. Below is a parametrised example which will help illustrate the results.

Example. Suppose that, in each period, agents face a binary decision between using and improving the technology. Improving the technology ($a_t^i = 1$) yields no payoff, and using it ($a_t^i = 0$) yields payoff x_t , where x_t is the current quality of the technology. We may interpret $0 < a_t^i < 1$ as improving the technology with probability a_t^i and using it with probability $1 - a_t^i$. Assuming that agents are expected utility maximisers, their flow payoff is

$$u(a_t^i, x_t) := (1 - a_t^i)x_t.$$

Finally, suppose that the distribution of the size of innovations is exponential; that is, $F(z) := 1 - e^{-z/\mu}$ for some $\mu > 0$.

2.1 Discussion of the assumptions

The literature on dynamic public-good games assumes that payoffs are separable. Formally, they may be expressed as

$$u(a_t^i, x_t) = b(x_t) - c(a_t^i). \quad (4)$$

This model is a generalisation. Namely, the cost of effort becomes steeper and more convex as the stock grows.²⁰ Another standard assumption in the literature is that the marginal benefit of increasing the stock vanishes as the stock grows (formally, $\lim_{x \rightarrow \infty} b'(x) = 0$). Condition (2) is weaker. The added generality encompasses linear payoffs, such as those of the example introduced above. Condition (2) ensures that the symmetric Markov equilibrium is unique, both in the baseline model (Theorem 2) as well as in the game with disposal (Theorem 3).

This model also deviates from a large part of the literature as it features stochastic production. However, condition (1) is only used to obtain a simple characterisation of the

19. The process $(\tilde{x}_t)_{t \geq 0}$ is piecewise constant, and is thus described by $(\tilde{\tau}_m, \tilde{z}_m : 1 \leq m \leq \tilde{m})$, where $\tilde{m} \geq 0$ is the number of jumps and, for any $1 \leq m \leq \tilde{m}$, the m^{th} jump occurs at time $\tilde{\tau}_m$ and has size \tilde{z}_m . The law of $(\tilde{\tau}_m, \tilde{z}_m : 1 \leq m \leq \tilde{m})$ may be described as follows. For each $m \in \{0, 1, \dots\}$, conditional on $\tilde{m} \geq m$ and on $\tilde{y}_m := x_0 + \sum_{l=1}^m \tilde{z}_l$ (where $\tilde{y}_0 := x_0$), $\tilde{m} = m$ a.s. if $\alpha^i(\tilde{y}_m) = 0$ for all i . Otherwise, $\tilde{m} > m$ a.s., $\tilde{\tau}_{m+1}$ and \tilde{z}_{m+1} are independent, $\tilde{\tau}_{m+1}$ is exponentially distributed with mean $[\lambda \sum_{i=1}^n \alpha^i(\tilde{y}_m)]^{-1}$, and $\tilde{z}_{m+1} \sim F$.

20. I show in Appendix I that the assumption is necessary to guarantee that the efficient effort decreases as the stock grows, a central feature in the literature on dynamic public-good games.

fact that innovations are detrimental (Lemma 2, Section 4.1), of the inefficiency of the equilibria (Corollary 2), and of the incentive to conceal innovations in the equilibrium of the baseline game (Lemma 3). All other results go through whether (1) holds or not. In particular, they continue to hold if production is deterministic (formally, if F is degenerate).

The distribution F of the random size of increments is independent of the effort a of the agent producing them. This is a substantial assumption. However, it is without loss of generality if effort is binary; that is, if picking $0 < a < 1$ is viewed as a randomisation between $a = 0$ and $a = 1$.²¹

Finally, all results continue to hold if agents may choose (pure) non-Markov strategies.²²

3 Social-welfare benchmark

In this section, I describe how non-strategic agents should behave in order to maximise aggregate payoffs. The main features of this benchmark are common in dynamic public-good games. In particular, any innovation is beneficial.

Note that, fixing a profile $\alpha = (\alpha^i)_{i=1}^n$, for all $x \geq 0$,

$$v^i(x) = b(x) - c(\alpha^i(x), x) + \lambda \sum_{j=1}^n \alpha^j(x) \{ \mathbb{E}_F[v^i(x + \tilde{z})] - v^i(x) \}. \quad (5)$$

In words, agent i 's continuation payoff is the sum of the current payoff flow $b(x) - c(\alpha^i(x), x)$, and the net expected future benefit. This is given by the rate $\lambda \sum_{j=1}^n \alpha^j(x)$ at which innovations occur, times their expected social value; that is, the difference between the continuation payoff $v^i(x + z)$ after an innovation of size $z > 0$, and the current payoff $v^i(x)$, weighted by the distribution F of z .

21. Note also that allowing for unbounded effort while keeping the rest of the assumptions unchanged necessarily leads to separable payoffs. Indeed, unless (4) holds, $c(a, \hat{x}) > c(a, x)$ for some a and $\hat{x} > x$. Moreover, for any $\hat{a} \geq a$, $b(\hat{x}) - b(x) \geq c(\hat{a}, \hat{x}) - c(\hat{a}, x) \geq (\hat{a} - a)[c(a, \hat{x}) - c(a, x)]$, so that effort must be bounded.

22. Assuming that effort is hidden, the game is well-defined under minimal measurability conditions on strategies, analogous to those imposed in the extension with concealment (Appendix A). Moreover, beliefs play no role since past effort is sunk and agents do not mix. Then, by standard dynamic-programming arguments, welfare is maximised by a profile of Markov strategies (so that Theorem 1 holds), and there exists a Markov best response to any Markov profile (so that Theorem 2 holds).

23. To derive (5) note that, for a small time interval dt , $v^i(x) = [b(x) - c(\alpha^i(x), x)]dt + e^{-dt} \mathbb{E}[v^i(\tilde{x}_{t+dt}) | \tilde{x}_t = x]$ and, up to a first-order approximation, $\mathbb{E}[v^i(\tilde{x}_{t+dt}) | \tilde{x}_t = x] = \lambda \sum_{j=1}^n \alpha^j(x) dt \mathbb{E}_F[v^i(x + \tilde{z})] + (1 - \lambda \sum_{j=1}^n \alpha^j(x) dt) v^i(x)$. Neglecting terms in dt^2 (and dropping the subscript t) yields (5).

Welfare is the average of agents' payoffs, given by

$$w(x) := \frac{1}{n} \sum_{i=1}^n v^i(x).$$

Since agents are identical, the cost of effort $c(a, x)$ is convex in $a \in [0, 1]$, and the rate of arrival of innovations λa is linear in effort, it is efficient for all agents to exert the same amount of effort. Then, from (5), the Bellman equation for the maximal welfare achievable in the game with initial stock $x \geq 0$ is

$$w(x) = \max_{a \in [0, 1]} \{b(x) - c(a, x) + a\lambda n \{\mathbb{E}_F[w(x + \tilde{z})] - w(x)\}\}. \quad (6)$$

Standard dynamic-programming arguments imply that, if (6) admits a (well-behaved) solution w_* then, for all $x_0 \geq 0$, $w_*(x_0)$ is the maximal welfare achievable in the game with initial stock x_0 . Moreover, any profile α of Markov strategies solving (for all i and $x \geq 0$)

$$\alpha^i(x) \in \arg \max_{a \in [0, 1]} \{a\lambda n \{\mathbb{E}_F[w_*(x + \tilde{z})] - w_*(x)\} - c(a, x)\} \quad (7)$$

induces welfare $w_*(x_0)$ for any initial stock $x_0 \geq 0$.

The following result shows that, in order to maximise welfare, agents should exert symmetric effort, and diminish it as the stock grows. Moreover, any increment in x_t makes agents better off. The proof is in Appendix C.²⁴

Theorem 1. *There exists a Markov profile α_* maximising welfare for all initial stocks $x_0 \geq 0$. Effort $\alpha_*^i(x_t)$ is symmetric and decreasing in the stock x_t , and $\alpha_*^i(\tilde{x}_t) \rightarrow 0$ a.s. as $t \rightarrow \infty$. Maximal welfare $w_*(x_0)$ is increasing in x_0 , and $w_*(x_0) - b(x_0)$ is decreasing.*²⁵

The fact that effort α_*^i is decreasing is standard in dynamic public-good games, and is generally due to concave payoffs. In this model, not only are payoffs concave in x_t , but the cost of effort becomes steeper and more convex as x_t grows (formally, $c_1(a, x)$ and $c_{11}(a, x)$ are increasing in x). The latter force adds to the former and does not alter the qualitative features of the social-welfare benchmark. However, as shown in the next section, it plays an important role in the symmetric equilibrium.

Maximal welfare w_* is increasing in the initial stock x_0 because higher x_0 yields a larger payoff flow ($b(x) - c(a, x)$ is increasing in x) without altering the productivity of

24. To prove Theorem 1, I construct the map w_* as a fixed point of a functional P , defined as follows. For an arbitrary welfare function w , $Pw(x_0)$ is the maximal welfare achievable in the game with initial stock x_0 , given that continuation welfare after the first innovation is given by $w(x_0 + z)$, where z is the size of the innovation. I restrict attention to absolutely continuous w 's that lie above b (and below a given linear map \bar{v}), and are increasing, but at a rate lower than that of b . Existence of w_* follows from Tarski's fixed-point theorem. The fact that $w_* - b$ is decreasing in x is used to show that α_* is also decreasing in x .

25. Unless c is strictly convex, (7) may admit more than one solution α . α_* is the pointwise smallest.

effort (the frequency and size of innovations does not depend on x). Moreover, since each agent's time- t continuation payoff when α_* is played equals $w_*(x_t)$, any increment in the stock (that is, any innovation) is beneficial. We shall see that, despite this, welfare need not increase with x_0 in equilibrium, so that innovations may be detrimental.

The fact that $w_* - b$ is decreasing in x_0 , where $b(x_0)$ is the 'myopic' payoff from exerting no effort in the game with initial stock x_0 , is also standard in dynamic public-good games. It follows from the fact that, as the stock grows, the flow of (gross) benefits increases more slowly. We shall see in the next section that this property also holds in equilibrium, and we will use it to derive a sufficient condition for innovations to be detrimental (Section 4.1).

The next result gives a necessary and sufficient condition for effort to cease in finite time. It is proved in Appendix C.

Corollary 1. *If*

$$\lambda n \{\mathbb{E}_F[b(x + \tilde{z})] - b(x)\} \leq c_1(0, x) \quad (8)$$

admits a solution then it admits a smallest solution x_ . In this case, no effort is exerted if $x_0 \geq x_*$ and, if $x_0 < x_*$, effort $\alpha_*(\tilde{x}_t)$ ceases a.s. in finite time, when \tilde{x}_t reaches x_* . If (8) holds with ' $>$ ' for all $x \geq 0$, then effort never ceases and $\tilde{x}_t \rightarrow \infty$ a.s. as $t \rightarrow \infty$.*

If (8) holds with equality at some x then, when the stock equals x , the cost and benefit of exerting a small amount of effort until an innovation occurs, and no effort thereafter, even out. The right-hand side of (8) is the marginal cost of effort when no effort is exerted. The left-hand side is the marginal social value of effort when the size of the stock is x , given that no effort is exerted after an innovation occurs. In particular, the net welfare benefit of an innovation of size z is $n[b(x + z) - b(x)]$.




Figure 1: Effort (left) and welfare (right) in the social-welfare benchmark of the example, as functions of the stock x_t . Parameter values are $n = 4$ and $\lambda = \mu = 1$.

Example. Figure 1 depicts effort and welfare in the social-welfare benchmark of the example introduced in Section 2. Effort α_*^i ceases at $x = \lambda\mu n$. Moreover, since the cost of effort is linear, α_*^i is ‘bang-bang’: $\alpha_*^i(x) = 1$ for $x < \lambda\mu n$, and $\alpha_*^i(x) = 0$ for $x > \lambda\mu n$. Welfare w_* is given by

$$w_*(x) = \begin{cases} \lambda\mu n e^{\frac{x/\mu - \lambda n}{1 + \lambda n}} & \text{if } x \leq \lambda\mu n \\ x & \text{if } x \geq \lambda\mu n. \end{cases} \quad (9)$$

4 Equilibrium

This section is devoted to the unique symmetric Markov perfect equilibrium of the game. Among other things, I show that continuation payoffs may drop after an innovation.

If the opponents of agent i play strategies $\alpha^{-i} := (\alpha^j)_{j \neq i}$, the largest continuation payoff that agent i can achieve, across all strategies, is a (value) function \hat{v}^i of the current stock x_t (where the dependence on α^{-i} is implicit). From (5), \hat{v}^i solves the Bellman equation

$$v^i(x) = \max_{a \in [0,1]} \left\{ b(x) - c(a, x) + \lambda \left[a + \sum_{j \neq i} \alpha^j(x) \right] \{ \mathbb{E}_F[v^i(x + \tilde{z})] - v^i(x) \} \right\}. \quad (10)$$

Moreover, a strategy α^i is a best response for agent i against α^{-i} (for any initial stock x_0) if and only if it solves, for all $x \geq x_0$,

$$\alpha^i(x) \in \arg \max_{a \in [0,1]} \{ a \lambda \{ \mathbb{E}_F[\hat{v}^i(x + \tilde{z})] - \hat{v}^i(x) \} - c(a, x) \}. \quad (11)$$

A profile α of Markov strategies is a *Markov Perfect Equilibrium* (MPE) if each α^i is a best response to α^{-i} , for any initial stock x_0 . The following result characterises the unique symmetric MPE of the game. The proof is in Appendix D and it is outlined below.

Theorem 2. *There exists a unique symmetric MPE α_e . Effort $\alpha_e^i(x_t)$ is continuous and decreasing in the stock x_t , and lies below the benchmark $\alpha_*^i(x_t)$. Moreover, $v_e(x_t) - b(x_t)$ is decreasing in x_t , where $v_e(x_t)$ is the equilibrium continuation payoff given x_t .*

Equilibrium effort $\alpha_e^i(x_t)$ decreases as the stock x_t increases and, moreover, it is inefficiently low in general. This is caused by *intertemporal free riding* (Marx and Matthews 2000): since α_e is decreasing, agents are reluctant to exert effort as this causes their opponents to exert less effort in the future. Thus α_e is inefficient, except for the trivial case in which no effort is optimal. This is the next result, proved in Appendix D.

26. Note that w_* is not differentiable at $x = \lambda\mu n$. In particular, the smooth-pasting property does not hold in this setting, as the stopping boundary is not regular. See Peskir and Shiryaev (2006).

Corollary 2. $v_e(x_0) < w_*(x_0)$ if and only if $\alpha_*^i(x_0) > 0$.²⁷

The following result parallels Corollary 1, and describes the long-run behaviour of the equilibrium effort schedule. I omit its proof as it is essentially the same as that of Corollary 1.

Corollary 3. *If*

$$\lambda\{\mathbb{E}_F[b(x + \tilde{z})] - b(x)\} \leq c_1(0, x) \quad (12)$$

admits a solution then it admits a smallest solution x_e . In this case, no effort is exerted if $x_0 \geq x_e$ and, if $x_0 < x_e$, then effort $\alpha_e^i(\tilde{x}_t)$ ceases a.s. in finite time, when \tilde{x}_t reaches x_e . If (12) holds with ' $>$ ' for all $x \geq 0$, then effort never ceases and $\tilde{x}_t \rightarrow \infty$ a.s. as $t \rightarrow \infty$.

The 'encouragement effect' (Bolton and Harris 1999) does not arise in α_e . That is, effort ceases in α_e precisely when it would cease in the single-agent problem. More specifically, in either setting, effort ceases eventually if and only if (12) admits a solution. If this is the case, effort ceases as the stock reaches the smallest solution to (12) both in α_e and in the single-agent problem.²⁸

The argument for the existence and uniqueness of α_e is similar to the proof in Bolton and Harris (1999) for the analogous result.²⁹ In particular, the argument shows the existence of (symmetric, Markov perfect) equilibria $\underline{\alpha}_e$ and $\bar{\alpha}_e$, inducing payoffs \underline{v}_e and \bar{v}_e such that, for any equilibrium α_e inducing payoffs v_e

$$\underline{\alpha}_e(x) \leq \alpha_e(x) \leq \bar{\alpha}_e(x) \quad \& \quad \underline{v}_e(x) \leq v_e(x) \leq \bar{v}_e(x)$$

for all $x \geq 0$. That is, $\underline{\alpha}_e$ (resp. $\bar{\alpha}_e$) is the equilibrium in which least (resp. most) effort is exerted, and payoffs are lowest (highest), regardless of the value of the stock. This structure arises since, although the effort choices $\alpha^i(x)$ and $\alpha^j(x)$ of distinct agents at the same stock x are strategic substitutes, their choices $\alpha^i(x)$ and $\alpha^j(x')$ at distinct stocks $x' \neq x$ are strategic 'complements'. That is, if $x' > x$, the best response $\alpha^i(x)$ is increasing in $\alpha^j(x')$.³⁰

27. Corollary 2 relies on (1). Indeed, α_e is efficient if the incentive to exert effort is strong initially and is guaranteed to drop suddenly, so that $\alpha_e(x_t)$ and $\alpha_*(x_t)$ drop from 1 to 0 at the same time. Since α_e^i is continuous (Theorem 2), this can only occur if innovations are guaranteed to be sufficiently large, a phenomenon ruled out by (1).

28. In Curello (2021), I show that this does not hinge on the restriction to symmetric equilibria, nor to Markov strategies. Indeed, no (pure-strategy) perfect Bayesian equilibrium features the encouragement effect.

29. Given the difference in the settings, and the generality of Theorem 2, the proof of Theorem 2 is not a straightforward adaptation of that in Bolton and Harris (1999).

30. This can be seen by rephrasing (10) as

$$v^i(x) = \max_{a \in [0,1]} \left\{ \frac{b(x) - c(a, x)}{1 + \lambda[a + \sum_{j \neq i} \alpha^j(x)]} + \frac{\lambda[a + \sum_{j \neq i} \alpha^j(x)]}{1 + \lambda[a + \sum_{j \neq i} \alpha^j(x)]} \mathbb{E}_F[v^i(x + \tilde{z})] \right\}.$$

Equilibrium uniqueness follows from the fact that, in the efficient benchmark, effort vanishes in the long run. To see why, note that $\bar{\alpha}_e(x) \geq \underline{\alpha}_e(x)$ for all $x \geq 0$. Then, (11) implies that

$$\mathbb{E}_F[\bar{v}_e(x + \tilde{z})] - \bar{v}_e(x) \geq \mathbb{E}_F[\underline{v}_e(x + \tilde{z})] - \underline{v}_e(x).$$

That is, the expected value of an innovation is higher when $\bar{\alpha}_e$ is played than when $\underline{\alpha}_e$ is played. Rearranging yields:

$$\mathbb{E}_F[\bar{v}_e(x + \tilde{z}) - \underline{v}_e(x + \tilde{z})] \geq \bar{v}_e(x) - \underline{v}_e(x).$$

Hence, roughly speaking, the distance between the continuation payoffs increases as the stock rises. But, since \bar{v}_e and \underline{v}_e lie between the myopic payoff b and maximal welfare w_* , and w_* approximates b as the stock diverges (because efficient effort vanishes, as shown in Theorem 1), so do \bar{v}_e and \underline{v}_e . Hence $\bar{v}_e = \underline{v}_e$, which implies that the equilibrium is unique.




Figure 2: Effort (left) and payoffs (right) in the symmetric MPE of the example, as functions of the stock x_t . Parameter values are $n = 4$ and $\lambda = \mu = 1$.

Example. Figure 2 depicts effort and welfare in the symmetric MPE of the example introduced in Section 2. Effort ceases at $\lambda\mu$ and is given by

$$\alpha_e^i(x) = \begin{cases} 1 & \text{if } x \leq y_e \\ \frac{1}{n-1} \left[\frac{x}{2\lambda\mu} + \frac{\mu}{x} \left(\frac{\lambda}{2} + 1 \right) - \frac{1}{\lambda} - 1 \right] & \text{if } y_e \leq x \leq \lambda\mu \\ 0 & \text{if } x \geq \lambda\mu \end{cases}$$

Clearly, the maximand a is decreasing in $\alpha^j(x)$ for each $j \neq i$, and increasing in $\mathbb{E}_F[v^i(x + \tilde{z})]$. If some α^j increases pointwise on (x, ∞) , then so does v^i . Hence the best response $\alpha^i(x)$ also increases.

where y_e is such that α_e^i is continuous at y_e .³¹ Payoffs are given by

$$v_e(x) = \begin{cases} ny_e e^{\frac{x-y_e}{\mu(1+\lambda n)}} & \text{if } x \leq y_e \\ \frac{x}{\lambda} \left(\frac{x}{2\mu} - 1 \right) + \mu \left(\frac{\lambda}{2} + 1 \right) & \text{if } y_e \leq x \leq \lambda\mu \\ x & \text{if } x \geq \lambda\mu. \end{cases} \quad (13)$$

Note that v_e is not monotone, so that an innovation may cause payoffs to drop.³² For instance, the first innovation lowers payoffs if it has size 1 and $x_0 = 0$. We shall see that this phenomenon arises under general conditions.

4.1 Detrimental innovations

In this section, I give a sufficient condition for the time- t continuation payoff $v_e(x_t)$ in the symmetric MPE to drop, with strictly positive probability, over the course of the game. Essentially, it is sufficient that the payoff function $b(x) - c(a, x)$ take a multiplicative form, and that innovations be sufficiently rare (λ be small) relative to the curvature-to-slope ratio of the cost of effort. In particular, if the condition is met, raising the value of innovations (either by altering how the stock enters the payoff function, or improving the distribution F of their size) cannot restore monotonicity.

Note first that, in equilibrium, innovations are beneficial on average.

Lemma 1. *For all $x \geq 0$, $\mathbb{E}_F[v_e(x + \tilde{z})] \geq v_e(x)$.*

Proof. Fix $x \geq 0$. Suppose first that $\alpha_e(x) > 0$. Then, from (11),

$$\lambda \{ \mathbb{E}_F[v_e(x + \tilde{z})] - v_e(x) \} \geq c_1(\alpha_e(x), x) \geq 0.$$

It remains to consider the case $\alpha_e(x) = 0$. In this case, since α_e is decreasing (Theorem 2), $\alpha_e(y) = 0$ for all $y \geq x$. Then

$$\mathbb{E}_F[v_e(x + \tilde{z})] = \mathbb{E}_F[b(x + \tilde{z})] \geq b(x) = v_e(x)$$

where the inequality holds since b is increasing. □

Lemma 1 states that, no matter the value of the current stock, each innovation increases the equilibrium continuation payoff *in expectation*; that is, when averaging across its possible sizes. Yet, as Proposition 1 below shows, innovations may be detrimental *ex post*. The next definition formalises this notion. Given a Markov profile α and x_0 , write $(\tilde{x}_t^\alpha : t \geq 0)$ for the stochastic process describing the evolution of the stock when α is played in the game with initial stock x_0 .

31. That is, $y_e = \mu(1 + \lambda n - \sqrt{(n^2 - 1)\lambda^2 + 2(n - 1)\lambda + 1})$. In Figure 2, $y_e \approx 0.31$.

32. Equivalently, decreasing the initial stock x_0 may be beneficial.

Definition 1. *Innovations are detrimental* if $\Pr(t \mapsto v_e(\tilde{x}_t^{\alpha_e}) \text{ is increasing} | x_0) < 1$.

In words, innovations are detrimental whenever, when α_e is played in the game with initial stock x_0 , with some (strictly positive) probability, we can find periods $t < T$ such that $x_t < x_T$ and $v_e(x_t) > v_e(x_T)$. That is, with some probability, the time- t continuation payoff $v_e(x_t)$ is not an increasing function of time. If this occurs, then the payoff $v_e(x_t)$ must drop after an innovation (that is, an increment in x_t). The following result characterises when this occurs. It is proved in Appendix E.

Lemma 2. *Innovations are detrimental if and only if v_e is not increasing on $[x_0, \infty)$.*

Note that the ‘only if’ part is immediate, since the stock x_t increases over time. The ‘if’ part relies on the possibility that arbitrarily small innovations may arise (condition (1), Section 2). Essentially, for any $x'' > x' \geq x_0$, with some probability, the stock will be close to x' at some $t \geq 0$ and close to x'' at some $T > t$. Thus, if $v_e(x') > v_e(x'')$, the payoff will not always increase over time.³³

In order to derive a sufficient condition for innovations to be detrimental, I restrict attention to payoff functions that take a particular multiplicative form.

Definition 2. The payoff function u (introduced in Section 2) is *multiplicative* if it may be expressed as

$$u(a, x) = g(1 - a)h(x) \quad (14)$$

for some $g : [0, 1] \rightarrow \mathbb{R}_+$ and $h : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $g(1) = 1$.³⁴

Note that the example introduced in Section 2 satisfies (14) with linear g and h . Assuming that g and h are twice continuously differentiable, the restrictions on u set out in Section 2 are equivalent to g being strictly increasing and concave and h being increasing, not constant, and concave.

Multiplicative payoffs may be interpreted as follows. Agent i is endowed with a unit of resources in each interval of time $[t, t + dt)$, and devotes a fraction $a_t^i \in [0, 1]$ to the production of the good (or, more concretely, the improvement of a shared technology), and $1 - a_t^i$ to its use. Agent i ’s payoff increases if she devotes more resources to the use of the technology (g increasing), or if the technology improves (h increasing), but both marginal returns are decreasing (g and h concave).

The following result gives a sufficient condition for innovations to be detrimental.

Proposition 1. *Suppose that payoffs are multiplicative and, for some $a \in (0, 1)$,*

$$\lambda < \frac{[g'(1 - a)]^2 + g(1 - a)g''(1 - a)}{g'(1 - a) - 2ag''(1 - a)}, \quad (15)$$

$$g'(1 - a)h(x_0) < \lambda \{ \mathbb{E}_F[h(x_0 + \tilde{z})] - [g(1 - a) + nag'(1 - a)]h(x_0) \}. \quad (16)$$

33. This occurs as long as effort is bounded away from zero on $[x_0, x'']$. The proof of Lemma 2 shows that, if v_e is not increasing, we may pick x' and x'' such that this is the case.

34. The assumption that $g(1) = 1$ is without loss of generality and is used to simplify Proposition 1.

Then, innovations are detrimental. More precisely, there is $x \geq x_0$ such that $\alpha_e(x) = a$ and v_e is not (weakly) increasing in a neighbourhood of x .

Roughly speaking, condition (15) states that innovations are sufficiently rare (that is, their rate of arrival λ is sufficiently low), compared to the ‘slope-to-curvature’ ratio of g at $1 - a$. Indeed, the right-hand side is strictly positive if and only if the map

$$\psi_g : y \mapsto g(y)g'(y)$$

is strictly increasing at $y = 1 - a$.³⁵ Essentially, this means that g grows faster than g' shrinks near $1 - a$.³⁶ Equation (15) guarantees that, if effort takes value a at some stock $x \geq x_0$, the payoff v_e is not increasing near x .

The purpose of (16) is to ensure that agents exert effort a at some stock $x \geq x_0$. More specifically, (16) implies that effort at x_0 must exceed a (which is sufficient as α_e is continuous and vanishes as the stock diverges). Condition (16) states that, if the effort exerted at x_0 did not exceed a , the marginal cost of effort would be strictly lower than its marginal benefit (as the latter exceeds the right-hand side).

Note that, if the hypotheses of Proposition 1 are satisfied, then innovations continue to be detrimental after an increase in (the distribution of) their size, or their value. Formally, improving F in the FOSD sense, or increasing h on (x_0, ∞) , does not affect (15), and increases the term $\mathbb{E}_F[h(x_0 + \tilde{z})]$ in (16), ‘weakening’ the constraint imposed by the latter inequality.

Note also that Proposition 1 yields a sufficient condition for innovations to be detrimental as long as *some* effort is exerted in equilibrium (formally, $\alpha_e(x_0) > 0$). The condition is that (15) hold for $a > 0$ sufficiently small. This is the case for $\lambda > 0$ sufficiently small as long as $[g'(1)]^2 + g''(1) > 0$.³⁷ From (12), some effort is exerted in equilibrium as long as

$$\lambda\{\mathbb{E}[h(x_0 + \tilde{z})] - h(x_0)\} > g'(0)h(x_0). \quad (17)$$

Then, innovations are detrimental if (17) holds and (15) holds for $a > 0$ sufficiently small.

Finally, Proposition 1 yields sufficient conditions for innovations to be detrimental regardless of the number of agents. For example, the conditions described in the last

35. To see why, note that the numerator of the right-hand side of (15) is precisely $\psi'_g(1 - a)$, whereas the denominator is strictly positive since g is concave and strictly increasing.

36. Moreover, if (15) holds then making g steeper or less concave at $1 - a$ preserves (15). Formally,

$$y \mapsto \frac{y^2 - g(1 - a)g''(1 - a)}{y - 2ag''(1 - a)} \quad \& \quad y \mapsto \frac{[g'(1 - a)]^2 - g(1 - a)y}{g'(1 - a) - 2ay}$$

are increasing on $(0, \infty)$ and $(-\infty, 0]$, respectively.

37. For example, if $g(y) = y^\beta$ with $\beta \in (1/2, 1]$, (15) becomes $\lambda < \frac{2\beta-1}{3-2\beta}$ in the limit as a tends to 0.

paragraph are sufficient, as they do not depend on n .³⁸

To gain an insight into why the equilibrium payoff $v_e(x)$ may drop as the stock x increases, totally differentiate the Bellman equation (10) with respect to x to obtain (assuming that (14) holds)

$$\begin{aligned} v'_e(x) = & \underbrace{g(1 - \alpha_e(x))h'(x)}_A + \underbrace{\lambda(n - 1)\alpha'_e(x)\{\mathbb{E}_F[v_e(x + \tilde{z})] - v_e(x)\}}_B \\ & + \underbrace{\lambda n \alpha_e(x)\{\mathbb{E}_F[v'_e(x + \tilde{z})] - v'_e(x)\}}_C. \end{aligned}$$

An increase in the stock increases the flow payoff (A), and alters the net expected future benefits. In particular, the rate of arrival of innovations drops (B) since effort does (Theorem 2), and the benefit of innovations, averaged over their size, changes (C). Under the hypotheses of Proposition 1, B dominates at some x , so that $v'_e(x) < 0$. In words, innovations can be detrimental because, although they increase the payoff flow and provide the basis for future improvements, they raise the value of using the technology, which lowers the incentive to improve it further. Therefore, due to a strategic effect, the positive externality arising from opponents' efforts shrinks. Further innovations are delayed, hampering the growth of the payoff flow.

I show in Appendix I that, for innovations to be detrimental, production must be stochastic *and* payoffs must not be separable. In other words, the co-existence of the two main features distinguishing this model from the literature on public good-games are necessary for payoffs to drop after an increase in the stock.³⁹

Example. Under what conditions is the equilibrium payoff v_e given in (13) increasing? Note that v_e is increasing on $[0, y_e]$ and convex on $[y_e, \infty)$, and $\arg \min_{[y_e, \infty)} v_e = \max\{y_e, \mu\}$. Then, v_e is increasing if and only if $y_e \leq \mu$ or, equivalently,

$$n \leq 1 + \frac{1}{2} \left(\lambda - \frac{1}{\lambda} \right). \quad (18)$$

In particular, if $\lambda < 1 + \sqrt{2}$, v_e is non-monotone regardless of the number of agents $n \geq 2$. This is the case in Figure 2, as $\lambda = 1$.⁴⁰

38. Another sufficient condition is that $h(x_0) = 0$ (which requires $x_0 = 0$) and (15) hold for $a < 1$ sufficiently large. The latter is satisfied if $\lambda > 0$ is small enough, $\lim_{y \downarrow 0} g'(y) < \infty$ and $g(0) = 0$. The requirement that $h(x_0) = 0$ means that the technology has no value initially, so that effort has no cost. The condition $g(0) = 0$ means that devoting no resources to the use of the technology yields no benefit, no matter its value.

39. Specifically, I show that v_e is increasing if the payoff function u may be expressed as in (4) (Proposition 6), and that innovations are not detrimental if the distribution F is degenerate (Corollary 5). In particular, Lemma 2, which relies on (1), fails if F is degenerate.

40. How likely is payoff to drop with the first innovation? Given the parameters of Figure 2, if $x_0 = 0$, the probability that this occurs is approximately 0.34. (13) implies that, across all parameters, the

5 Disposal of innovations

In this section, I analyse the impact of allowing each agent to freely dispose of the innovations that she induces, after observing their size. Among other things, I show that this raises equilibrium effort and improves payoffs relative to the equilibrium with forced revelation.

5.1 Model

Enrich the model as follows. Whenever agent i obtains an increment of size $z > 0$ at time $t \geq 0$, she (immediately) decides whether or not to reduce z to 0, after observing z . In other words, agent i may either ‘reveal’ ($d_t^i = 1$) or ‘dispose’ of the increment ($d_t^i = 0$).

In this setting, a *Markov strategy* is a pair $\sigma := (\alpha, \delta)$ of (Lebesgue measurable) maps $\alpha : \mathbb{R}_+ \rightarrow [0, 1]$ and $\delta : \mathbb{R}_+ \times (0, \infty) \rightarrow \{0, 1\}$ such that $a_t^i = \alpha(x_t)$ for all $t \geq 0$ and, if agent i obtains an innovation of size $z > 0$ at time t , then $d_t^i = \delta(x_t, z)$. If agents play a profile $\sigma := (\sigma^i)_{i=1}^n$, assuming that no agent obtains an innovation at time t , the time- t continuation payoff to agent i is a function v^i of the current stock x_t given by (3), where the law of the process $(\tilde{x}_t)_{t \geq 0}$ is different than in the baseline model of Section 2.⁴¹ Moreover, if agent j obtains an innovation of size $z > 0$ at time t , the time- t payoff to agent i is $v^i(x_t + z)$ if agent j reveals the innovation (i.e. if $\delta^j(x_t, z) = 1$), and $v^i(x_t)$ if agent j discards it.

We recover the baseline model by restricting the agents’ strategies. Namely, by imposing that agents may never dispose of any innovation (so that $d_t^i = 0$ always). I refer to the baseline model (Section 2) as the game *with forced disclosure*, and to this model as the game *with disposal*.

The game with disposal may be viewed as a reduced form of the following richer environment. Each agent possesses a set of concealed increments $Z_t^i \subset (0, \infty)$, initially empty. Whenever agent i obtains an increment of size $z > 0$, agent i chooses $d_t^i \in Z_t^i \cup \{z, 0\}$. If $d_t^i = 0$, the value z is added to the set Z_t^i (and the stock x_t does not grow). If instead $d_t^i \neq 0$, the stock x_t increases by d_t^i and Z_t^1, \dots, Z_t^n are emptied.⁴² Say that agent

probability of this event is maximised if $x_0 = 0$ and n is large. In the limit as n diverges, the first innovation is detrimental if and only if it has size less than 2μ . This occurs with probability $1 - e^{-2} \approx 0.86$.

41. The process $(\tilde{x}_t)_{t \geq 0}$ is piecewise constant, and is thus described by $(\tilde{\tau}_m, \tilde{z}_m : 1 \leq m \leq \tilde{m})_{m=1}^{\tilde{m}}$, where $\tilde{m} \geq 0$ is the number of jumps and, for any $1 \leq m \leq \tilde{m}$, the m^{th} jump occurs at time $\tilde{\tau}_m$ and has size \tilde{z}_m . The law of $(\tilde{\tau}_m, \tilde{z}_m : 1 \leq m \leq \tilde{m})$ may be described as follows. For each $m \in \{0, 1, \dots\}$, conditional on $\tilde{m} \geq m$ and on $\tilde{y}_m := x_0 + \sum_{l=1}^m \tilde{z}_l$ (where $\tilde{y}_0 := x_0$), $\tilde{m} = m$ a.s. if either $\alpha^i(\tilde{y}_m) = 0$ or $\delta^i(\tilde{y}_m, \cdot) = 0$ F -a.s. Otherwise, given independent $(\tilde{\pi}_{i,l}, \tilde{w}_{i,l} : i = 1, \dots, n, l \in \mathbb{N})$ such that, for each i and l , $\tilde{\pi}_{i,l}$ is exponentially-distributed with mean $[\lambda \alpha^i(\tilde{y}_m)]^{-1}$ (with $\tilde{\pi}_{i,l} := \infty$ if $\alpha^i(\tilde{y}_m) = 0$), and $\tilde{w}_{i,l} \sim F$, $\tilde{\tau}_m = \sum_{l=1}^{\tilde{o}} \tilde{\pi}_{i,l}$ and $\tilde{z}_m = \tilde{w}_{i,\tilde{o}}$, where the pair (\tilde{i}, \tilde{o}) minimises $\sum_{l=1}^{\tilde{o}} \tilde{\pi}_{i,l}$ across all $i \in \{1, \dots, n\}$ and $\tilde{o} \in \mathbb{N}$ such that $\delta^i(\tilde{y}_m, \tilde{w}_{i,\tilde{o}}) = 1$.

42. More formally, if $d_t^i = 0$, then $x_{t+} = x_t$, $Z_{t+}^i = Z_t^i \cup \{z\}$, and $Z_{t+}^j = Z_t^j$ for all $j \neq i$. If instead $d_t^i \neq 0$, $x_{t+} = x_t + d_t^i$, and $Z_{t+}^1 = \dots = Z_{t+}^n = \emptyset$. The event that two or more agents obtain an increment in the same period occurs with zero probability and may therefore be neglected.

i *discloses* an innovation (of size d_t^i at time t) if $d_t^i > 0$, and *conceals* otherwise.⁴³ Assume that the arrival and size of increments as well as the sets Z_t^i are private information, and the concealment of innovations is unobserved. In contrast, agents are immediately informed of any innovation that is disclosed, including its size and the identity of the agent revealing it.

This richer environment may be interpreted as follows. At any point in time, firms collectively seek solutions to a given problem. The arrival of an increment corresponds to the discovery of a solution, and the size of the increment is the quality of the solution. Disclosing an increment corresponds to (revealing and) implementing a particular solution. When this occurs, a new problem is chosen and the search for solutions begins anew.⁴⁴ Solutions geared towards the same problem are perfect substitutes; in particular, when one is implemented, the other solutions may be neglected. As will be clear from the analysis in this section, all results continue to hold (with no changes) in this richer model.⁴⁵

5.2 Welfare benchmark

In this section, I argue that the social-welfare benchmark is unaffected by the introduction of disposal. In particular, disposal of any innovation is inefficient.

The argument used to derive (5) in Section 3 implies that, fixing a profile $(\alpha^i, \delta^i)_{i=1}^n$, for all $x \geq 0$,

$$v^i(x) = b(x) - c(\alpha^i(x), x) + \lambda \mathbb{E}_F \left\{ \left[\sum_{j=1}^n \alpha^j(x) \delta^j(x, \tilde{z}) \right] [v^i(x + \tilde{z}) - v^i(x)] \right\}. \quad (19)$$

In contrast to (5), the rate of arrival of (disclosed) innovations $\sum_{j=1}^n \alpha^j(x) \delta^j(x, z)$ varies with their size z .

The social-welfare benchmark (Theorem 1) is unaffected by the introduction of dis-

43. Since time is continuous, the times at which agents can disclose innovations must be restricted in order for the game to be well-defined. Allowing agents to immediately disclose any innovation that they obtain ensures that any strategy of the game with forced disclosure is a strategy in this game. As will be clear from the analysis in this section, allowing the agents to disclose more frequently does not alter the results.

44. In particular, agents may not search for solutions to a future problem before a solution for the current one is adopted. This may be due to contractual restrictions if agents are firms engaged in a formal partnership. This is also the case if firms cannot predict what the future problem will be unless they know which solution will be implemented for the current one, or unless they reveal the solutions they found.

45. More specifically, Theorem 3 continues to hold under the following notion of ‘Markov’ strategy: a pair $\sigma^i := (\alpha^i, \delta^i)$ of (Lebesgue measurable) maps $\alpha^i : \mathbb{R}_+ \rightarrow [0, 1]$ and $\delta^i : \mathbb{R}_+ \times (0, \infty) \rightarrow \{0, 1\}$ such that $a_t^i = \alpha^i(x_t)$ for all $t \geq 0$ and, if agent i obtains an innovation of size $z > 0$ at time t , then $d_t^i = z \delta^i(x_t, z)$. In particular, innovations that are not immediately disclosed are never adopted. In principle, allowing the choice of d_t^i to depend on Z_t^i would enable the opponents of agent i to alter their beliefs about Z_t^i as time passes without a disclosure, and may prompt them to alter their effort over time. This suggests that a stationary equilibrium in which agent i ’s disclosures explicitly condition on Z_t^i is unlikely to exist.

posal. This is because, in the non-strategic setting, the effects of disposal on incentives can be ignored. Moreover, disposal of innovations hinders the growth of the payoff flow, since $b(x) - c(a, x)$ is increasing in x . I state the result for completeness, and omit the proof as it is straightforward.

Proposition 2. *The social-welfare benchmark in the game with disposal is identical to the baseline one. In particular, disposal of any innovation is inefficient.*

5.3 Equilibrium

In this section, I show that the game with disposal admits a unique symmetric Markov perfect equilibrium. Moreover, equilibrium effort and payoffs are higher than in the equilibrium of the game with forced disclosure.

If the opponents of agent i play strategies $\sigma^{-i} := (\alpha^j, \delta^j)_{j \neq i}$, assuming that no agent obtains an innovation at time t , the largest continuation payoff that agent i can achieve, across all strategies, is a (value) function \hat{v}^i of x_t (where the dependence on σ^{-i} is implicit). If agent i obtains an innovation of size z at time t , her value is $\max\{\hat{v}^i(x_t), \hat{v}^i(x_t + z)\}$.⁴⁶ Moreover, from (5), \hat{v}^i solves the Bellman equation

$$\begin{aligned} v^i(x) = \max_{a \in [0,1]} & \left\{ b(x) - c(a, x) + a\lambda \mathbb{E}_F [\max\{v^i(x + \tilde{z}) - v^i(x), 0\}] \right. \\ & \left. + \lambda \mathbb{E}_F \left[\left(\sum_{j \neq i} \alpha^j(x) \delta^j(x, \tilde{z}) \right) [v^i(x + \tilde{z}) - v^i(x)] \right] \right\}, \end{aligned} \quad (20)$$

and a strategy $\sigma^i := (\alpha^i, \delta^i)$ is a best response for agent i against σ^{-i} (for any initial stock x_0) if and only if, for all $x \geq 0$ and $z > 0$,

$$\alpha^i(x) \in \arg \max_{a \in [0,1]} \{a\lambda \mathbb{E}_F [\max\{\hat{v}^i(x + \tilde{z}) - \hat{v}^i(x), 0\}] - c(a, x)\}, \quad (21)$$

$$\hat{v}^i(x + z) > (<) \hat{v}^i(x) \quad \Rightarrow \quad \delta^i(x, z) = 1 \text{ (0)}. \quad (22)$$

As before, a strategy profile σ is a *Markov perfect equilibrium* (MPE) if each σ^i is a best response against σ^{-i} , for every initial stock x_0 .

The following result characterises the unique symmetric MPE of the game. The proof is in Appendix D. The (unique symmetric Markov perfect) equilibrium α_e of the game with forced disclosure is described in Theorem 2. The notion of detrimental innovations, which applies to the equilibrium α_e , is defined in Section 4.1.

Theorem 3. *The game with disposal admits an (essentially) unique symmetric MPE σ_d . In the absence of innovations at time t , and for any stock x_t , effort $\alpha_d^i(x_t)$ and continuation payoffs $v_d(x_t)$ are no lower than their analogues $\alpha_e^i(x_t)$ and $v_e(x_t)$ in the*

46. If, at time t , agent i does not obtain an innovation one of her opponents reveals (discards) an innovation of size z , the value to agent i is $\hat{v}^i(x_t + z)$ ($\hat{v}^i(x_t)$).

equilibrium with forced disclosure. Moreover, ex-ante payoffs $v_d(x_0)$ strictly exceed $v_e(x_0)$ if innovations are detrimental in α_e . Otherwise, no disposal occurs in σ_d and the equilibria coincide.⁴⁷

Allowing agents to dispose of innovations increases equilibrium payoffs, and strictly so unless all innovations are guaranteed to be beneficial in the equilibrium with forced disclosure. This is because, as can be seen from (22), private and public incentives for the disposal of innovations are aligned since σ_d is symmetric. In other words, whenever an agent discards an innovation in equilibrium, her opponents benefit from it. Moreover, the fact that detrimental innovations will be disposed of at any time in the future increases continuation payoffs at all stock values exceeding the current one and, therefore, the current incentive to exert effort. As a result, equilibrium effort is higher in the game with disposal (formally $\alpha_d^i(x) \geq \alpha_e^i(x)$ for all $x \geq 0$). This strengthens the positive externality arising from the public nature of the good, increasing the payoff further.⁴⁸

The properties of α_e described in Theorem 2, Corollary 2 and Corollary 3 are inherited by σ_d . Namely, effort $\alpha_d^i(x_t)$ is continuous and decreasing in the stock x_t , and below the benchmark $\alpha_*^i(x_t)$. The equilibrium σ_d is inefficient unless the initial stock is so large that exerting no effort is efficient (i.e. unless $\alpha_*^i(x_0) = 0$). Assuming that (12) admits a smallest solution y_e , no effort is exerted if $x_0 \geq y_e$ and, if $x_0 < y_e$, effort $\alpha_d^i(\tilde{x}_t)$ ceases a.s. in finite time, when \tilde{x}_t reaches y_e . If (12) admits no solution, effort never ceases and $\tilde{x}_t \rightarrow \infty$ a.s. as $t \rightarrow \infty$. Finally, $v_d(x_t) - b(x_t)$ is decreasing in the current stock x_t .

Example. Figure 3 depicts effort and payoffs in the equilibrium with disposal of the example introduced in Section 2. Assuming that $\lambda \leq 1$, equilibrium effort α_d^i is given by

$$\alpha_d^i(x) = \begin{cases} 1 & \text{if } x \leq y_d \\ \frac{\mu \log \frac{\lambda\mu}{x}}{(n-1)x} & \text{if } y_d < x < \lambda\mu \\ 0 & \text{if } x \geq \lambda\mu. \end{cases}$$

where y_d is such that α_d^i is continuous at y_d .⁴⁹ On $[y_d, \infty)$, the equilibrium disposal policy

47. Although there may exist multiple equilibria, effort α_d^i and the payoff function v_d are uniquely pinned down. Moreover, for any disclosure policy δ^i , the strategy (α_d^i, δ^i) induces a symmetric MPE if and only if (22) holds with $\hat{v}^i = v_d$ (for all $x \geq 0$ and $z > 0$). In words, multiplicity arises whenever agents are indifferent about the disposal of some innovations, and any way of breaking the indifferences yields an equilibrium.

48. Moreover, because detrimental innovations are discarded, equilibrium continuation payoffs are guaranteed to increase over time. However, as can be seen from the example below, the ex-ante equilibrium payoff $v_d(x_0)$ need not be increasing in the initial stock x_0 . That is, reducing the initial stock may still be beneficial.

49. That is, it is the unique solution $x \in (0, \lambda\mu)$ to $(n-1)x = \mu \log \frac{\lambda\mu}{x}$. In Figure 3, $y_d \approx 0.35$.




Figure 3: Effort (left) and payoffs (right) in the symmetric MPE σ_d of the example, as functions of the stock x_t . The dotted lines are effort α_e^i (left) and payoffs v_e (right) in the symmetric MPE of the game with forced disclosure. Parameter values are $n = 4$ and $\lambda = \mu = 1$.

δ_d^i and payoffs v_d are given by

$$\delta_d^i(x, z) = \begin{cases} \mathbb{1}_{z \geq \mu \log \frac{\lambda \mu}{x}} & \text{if } y_d \leq x < \lambda \mu \\ 1 & \text{if } x \geq \lambda \mu, \end{cases} \quad \& \quad v_d(x) = \begin{cases} x + \mu \log \frac{\lambda \mu}{x} & \text{if } y_d \leq x < \lambda \mu \\ x & \text{if } x \geq \lambda \mu. \end{cases}$$

Note that y_d is decreasing in n with $\lim_{n \rightarrow \infty} y_d = 0$. Moreover, although there is no closed-form solution for v_d on $[0, y_d]$, we know that $x \mapsto v_d(x) - x$ is decreasing, so that $v_d(0) \geq v_d(y_d) - y_d = \log[(\lambda \mu)/y_d]$. Then, since $\lim_{n \rightarrow \infty} y_d = 0$,

$$\lim_{n \rightarrow \infty} v_d(0) = \infty.$$

In words, if the initial stock is small and the population is large, the ex-ante payoff in the equilibrium with disposal is arbitrarily large. This is because, since the cost of effort is small, agents exert maximal effort, so that aggregate effort is large. Then, agents produce innovations arbitrarily fast, until one is disclosure. Because of this, the cutoff for the disposal of innovations is high: only large increments are kept. We shall see in Section 6 that a similar phenomenon occurs in the game with concealment.

In contrast, in the equilibrium with forced disclosure, the ex-ante payoff is bounded for small stocks and large populations.⁵¹ This is because, even though the first innovation is produced arbitrarily fast, its size is finite (as it is drawn from F), and effort drops drastically after it occurs.

50. From (22), δ_d^i is uniquely pinned down on $[y_d, \infty) \times (0, \infty)$ except at $(x, \mu \log(\lambda \mu/x))$ for $x \in [y_d, \lambda \mu]$.

51. To see why, note that the cutoff y_e (defined in footnote 31) tends to 0 as n diverges. Then, from (13), $v_e(0) \rightarrow \mu(\lambda/2 + 1)$ as $n \rightarrow \infty$.

6 Concealment of innovations

In this section, I extend the baseline model by allowing agents to conceal the innovations that they induce for a period of time, after observing their size. Among other things, I show that the resulting game admits a relatively simple equilibrium which, under some conditions, improves on the equilibrium α_e with forced disclosure. Throughout, I restrict attention to the payoff function of the example introduced in Section 2.

6.1 Model

As before, x_t denotes the (public) stock of the good and, in any interval of time $[t, t + dt)$, agent i exerts hidden effort $a_t^i \in [0, 1]$ and obtains payoff $(1 - a_t^i)x_t dt$.

For any $t \geq 0$ and $i \in \{1, \dots, n\}$, let $k_t^i \geq x_t$ denote agent i 's *private stock* of the good. At $t = 0$, $k_0^i = x_0$ for each i . In any interval of time $[t, t + dt)$, agent i obtains an increment in k_t^i (instead of x_t) with probability $\lambda a_t^i dt$. The distribution F of the size of increments inherits the properties described in Section 2. Moreover, given $\mathcal{T} \subset [0, \infty)$ such that $\mathcal{T} \cap [0, T]$ is finite for all $T > 0$, each agent i picks a *disclosure* $d_t^i \in [x_t, k_t^i]$ at any time t such that either $t \in \mathcal{T}$, or $x_t = x_0$, or at which agent i obtains a private increment.⁵² If a (non-empty) set of agents $I \subseteq \{1, \dots, n\}$ pick disclosures $(d_t^i)_{i \in I}$ at some time t , the public stock x_t , as well as the private stocks k_t^j of all agents, jump up to $\max\{d_t^i : i \in I\}$ if they laid strictly below it, and are unchanged otherwise.⁵³ In particular, neither x_t nor any k_t^j increase if $d_t^i = x_t$ for all $i \in I$. Whenever agent i picks d_t^i at time t , say that she *conceals fully* if $d_t^i = x_t$, *discloses fully* if $d_t^i = k_t^i > x_t$, and *discloses partially* if $x_t < d_t^i < k_t^i$.

Note that, in any period t at which agent i does not obtain a private increment, and such that $x_t > x_0$ and $t \notin \mathcal{T}$, agent i does *not* pick d_t^i . Thus, each agent is free to disclose at any point in time as long as no disclosure occurred but, after a disclosure occurs, she can only disclose at one of the times in \mathcal{T} , or immediately after having obtained a private increment. This and other assumptions are discussed in the next section.

Assume that agents' effort, private stocks, as well as the arrival and size of innovations are private information, and that the full concealment of innovations is not observed. In other words, if agent i obtains an innovation at time t and conceals it fully, her opponents will not be able to distinguish this event from the event that agent i does not obtain an innovation at time t . In contrast, if $d_t^i > x_t$, the opponents of agent i are immediately informed of the value of d_t^i , as well as of the identity of agent i . However, as they do not observe k_t^i , they cannot tell whether $d_t^i = k_t^i$; that is, they cannot distinguish between partial and full disclosures. I restrict each agent i to play pure strategies that do not condition on the past trajectory $(k_s^i : 0 \leq s \leq t)$ of her own private stock, but merely

52. If agent i obtains a private increment at time t , k_t^i denotes her private stock *after* the increment.

53. Formally, $x_{t+} = \max\{d_t^i : i \in I\}$ and $k_{t+}^j = \max\{k_t^j\} \cup \{d_t^i : i \in I\}$ for all $j \in \{1, \dots, n\}$.

on its current value k_t^i .⁵⁴ Unlike in previous sections, I also allow agents to condition their behaviour on calendar time and on past disclosures. Formal definitions of histories, strategies, and payoffs are in Appendix A.

We recover the game with forced disclosure (Section 2) by restricting the agents' strategies, namely, by imposing that agents always disclose fully. I refer to this model as the *game with concealment*.

Recall that, in the game *with disposal* (Section 5), whenever an agent obtains an innovation, she decides whether to discard or disclose it, after observing its size. Concealing innovations is more profitable than discarding them, as it allows the agent to 'build' on the hidden increment. To see this, consider an agent who obtains innovations of sizes $z' > 0$ and $z'' > 0$ at times $0 < t' < t''$. If, in the game with disposal, the agent discards the first increment and discloses the second, the stock x_t takes value x_0 until time t'' , then jumps up to $x_0 + z''$. In contrast, if the agent conceals the first increment fully, and discloses fully upon obtaining the second increment, the public stock x_t jumps to $x_0 + z' + z''$ at time t'' . In other words, in the latter case, the agent merely delays the disclosure of the first increment until the second occurs.

6.2 Discussion of the assumptions

Concealment of innovations leads to a stark form of redundancy. To see why, suppose that agent 1 obtains a private increment at time t' raising her private stock to $k_{t'}^1$, and conceals it fully, and agent 2 obtains a private increment at time $t'' > t'$, raising her private stock to some value $k_{t''}^2 > k_{t'}^1$, and discloses it fully. In this case, the private stock of agent 1 rises to $k_{t''}^2$ at time t'' , but no further. That is, the innovations obtained by agents 1 and 2 at times t' and t'' , respectively, are perfect substitutes. Since agent 2's innovation is superior, agent 1's innovation is lost at time t'' , when agent 2 discloses – it is as if the increment had not occurred. A more realistic model would feature some degree of complementarity between innovations. In the example above, the disclosure by agent 2 at time t'' would raise $k_{t''}^1$ to a value above $k_{t''}^2$, at least with some probability. However, despite this extreme form of redundancy, concealment may be beneficial in equilibrium (Proposition 4).

Whereas the time of the first disclosure is unrestricted, after the first disclosure, agents can only disclose at specific times, or whenever they obtain a private increment. This restriction is imposed entirely for tractability. It ensures that only finitely many disclosures occur a.s. in any finite period of time. Without it, the game would not be well-defined, as time is continuous. Moreover, imposing no restriction on the time of the first disclosure ensures that the choice of \mathcal{T} does not affect (on path) the equilibrium that

54. Results continue to hold if strategies may condition on $(k_s^i : 0 \leq s \leq t)$. This is ruled out to simplify the definition of beliefs.

I analyse. Finally, allowing agents to disclose whenever they obtain an increment ensures that the equilibrium α_e of the game with forced disclosure is a well-defined strategy profile in this environment.

A simpler model would constrain the agents to either full disclosure or full concealment. Formally, it would require that $d_t^i \in \{x_t, k_t^i\}$. In that case, an agent may have an incentive to disclose fully as evidence to her opponents that she is not concealing more substantial innovations. Doing so would raise the opponents' incentives to exert effort, and benefit the agent. Hence, this is likely to play an important role in equilibrium. I do not analyse this setting as it seems economically less relevant.⁵⁵

6.3 Welfare benchmark

The social-welfare benchmark (Theorem 1) is unaffected by the introduction of concealment. This is because, in the non-strategic setting, the effects of delayed disclosure on incentives can be ignored. Moreover, delayed disclosure hinders the growth of the payoff flow, and may lead to redundant innovations. I state the result for completeness, and omit the proof as it is straightforward.

Proposition 3. *The social-welfare benchmark in the game with concealment is identical to that of the game with forced disclosure. In particular, it is efficient to disclose all innovations immediately.*

6.4 Equilibrium

In this section, I construct a symmetric perfect Bayesian equilibrium of the game with concealment. We shall see in the next section that, under some conditions, ex-ante payoffs in this equilibrium exceed those of the equilibrium α_e with forced disclosure.

A *perfect Bayesian equilibrium* (PBE) is composed of a profile of strategies ρ , and a profile of beliefs β .⁵⁶ The equilibrium α_e with forced disclosure (Theorem 2), coupled with the ‘full-disclosure’ policy, is a natural candidate equilibrium in the game with concealment. I begin by showing that it is a PBE (for some belief profile) if and only if the initial stock is so large that no effort is exerted. Since effort $\alpha_e^i(x_t)$ is decreasing in the current stock x_t , no effort is exerted in equilibrium if and only if $\alpha_e(x_0) = 0$. From Corollary 3, $\alpha_e(x_0) = 0$ if and only if $x_0 \geq \lambda\mu$, where μ is the average size of innovations.

55. Requiring instead that $d_t^i \in \{k_s^i : 0 \leq s \leq t\} \cap [x_t, \infty)$ would have no effect on the results.

56. Definitions of strategies, beliefs, and PBE are in Appendix A. The standard notion of PBE, defined for discrete-time games, applies to this setting. Indeed, since effort is hidden and I restrict attention to pure strategies, agents reach a new information set precisely whenever they either obtain a private increment, or one of their opponents discloses (partially or fully). Given the restrictions on the arrival of private increments, as well as on the periods in which disclosure is possible, agents reach finitely many different information sets within any bounded interval of time. This implies that PBE are well-defined.

Lemma 3. *The effort policy α_e^i , coupled with the ‘full-disclosure’ policy, forms a symmetric PBE of the game with concealment (for some profile of beliefs) if and only if $x_0 \geq \lambda\mu$.*

Lemma 3 is proved in Appendix G. It shows that agents may have an incentive to conceal innovations even if they are not detrimental in equilibrium (in the sense defined in Section 4.1). For instance, if $F(z) = 1 - e^{-z/\mu}$, innovations are detrimental if and only if (18) fails. Yet, full disclosure is incentive-compatible if and only if $x_0 \geq \lambda\mu$. This is because, as noted in Section 5, innovations are detrimental if and only if agents have an incentive to discard them whereas, as argued at the end of Section 6.1, concealment is more profitable than disposal, as it allows agents to ‘build’ on the concealed increments.⁵⁷

The next result describes a symmetric PBE of the game with concealment. The proof is in Appendix G and is outlined below.

Theorem 4. *There exists a symmetric PBE (ρ_c, β_c) such that, in any period t prior to which no disclosure occurred, agent i exerts effort $\alpha_t^c(k_t^i)$ and conceals (resp. discloses) fully if $k_t^i < (\geq) q(t)$, where $\alpha_t^c(k)$ is increasing in k and $q(t) \in [\lambda\mu, \lambda\mu n]$. On the equilibrium path, no effort is exerted after a disclosure occurs.⁵⁸*

The equilibrium (ρ_c, β_c) is such that, if $x_0 \geq \lambda\mu$, then no effort is exerted and the equilibrium coincides with α_e (Lemma 3). The rest of the discussion focuses on the case $x_0 < \lambda\mu$. In this case, agents conceal innovations as long as their private stocks k_t^i lie below the common, time-varying cutoff $q(t)$. As soon as the private stock of some agent i reaches $q(t)$, agent i discloses fully. Moreover, at the time of disclosure, $k_t^i \geq q(t) \geq k_t^j$ for all $j \neq i$, so that each k_t^j rises to k_t^i . In other words, the first innovation disclosed (on the equilibrium path) is the best available at the time of disclosure. After the first disclosure, the opponents of the agent who disclosed exert no effort.⁵⁹ On the equilibrium path, the agent who disclosed does not exert any effort either; from Lemma 3, this is incentive-compatible since the public stock x_t must now lie above $\lambda\mu$, as q is bounded below by $\lambda\mu$.

At any time t prior to the first disclosure, each agent believes that the private stocks of her opponents are i.i.d. with the same distribution (call it G_t). As time passes *without disclosure*, agents need not become more optimistic about their opponents’ private stocks. Formally, G_t need not grow in the FOSD-sense over time. For this reason, the cutoff $q(t)$ need not increase over time either. In particular, the first disclosure need not happen

57. I show in Appendix I that, if either payoffs are separable (that is, they satisfy (4)) or the size of innovations is fixed, then the effort policy α_e^i , coupled with the full-disclosure policy, forms an equilibrium of the game with concealment. In other words, both of the main features distinguishing this model from the literature on public-good games (namely, a trade-off between ‘production’ and ‘use’ of the good, and stochastic production) are necessary to induce agents to not only discard, but also conceal innovations.

58. The cutoff $q(t)$ is lower-semicontinuous, so that the (random) time of disclosure is well-defined.

59. If more than one agent disclose simultaneously, all agents except one among those who disclosed exert no effort. At the end of this section, I describe the off-path beliefs that sustain this drop in effort.

due to an innovation, but can also occur after a drop in $q(t)$, due to growing pessimism about opponents' progress.

Since agent i exerts effort $\alpha_t^c(k_t^i)$ at any time t prior to the first disclosure, her (pre-disclosure) continuation payoff may be expressed as $v_t^c(k_t^i)$ for some function $v^c : \mathbb{R}_+ \times [x_0, \infty) \rightarrow \mathbb{R}$. One can show that, for all $t \geq 0$ and $k \geq x_0$,

$$\alpha_t^c(k) \in \arg \max_{a \in [0,1]} \{a(\lambda \{\mathbb{E}_F[v_t^c(k + \tilde{z})] - v_t^c(k)\} - x_0)\}. \quad (23)$$

This condition is analogous to (11). Increasing effort by a small amount $da > 0$ at time t increases the probability of obtaining a private increment within the time period $[t, t + dt)$ by approximately $\lambda dt da$, and decreases the payoff within this period by $x_0 dt da$. Moreover, the net expected benefit of obtaining a private increment (averaged across the size z of the increment) is $\mathbb{E}_F[v_t^c(k_t^i + \tilde{z})] - v_t^c(k_t^i)$.

Effort $\alpha_t^c(k_t^i)$ is increasing in k_t^i . Intuitively, this is because effort is beneficial ex-post if and only if it leads to a disclosure. The smaller the distance between k_t^i and the cutoff $q(t)$, the larger the impact of effort on the time to a disclosure, and thus, the higher the benefit of exerting effort. Moreover, the cost of effort does not grow as k_t^i grows, since it is determined by the public stock x_t , which is constant at x_0 until the disclosure.

The payoff $v_t^c(k_t^i)$ is increasing in the private stock k_t^i since a higher k_t^i brings agent i closer to a disclosure, but does not weaken her opponents' incentives for effort, as they do not observe k_t^i . As a consequence, when the disclosure occurs, all agents are better off. Indeed, if agent i discloses at time t , then she discloses $k_t^i \geq q(t)$, so that the payoff to any other agent j rises from $v_t^c(k_t^j)$ to $k_t^i = v_t^c(k_t^i)$, as $k_t^j \leq q(t)$.

The equilibrium (ρ_c, β_c) involves punishments for any agent i disclosing a value $d_t^i < q(t)$ at any time t before which no disclosure occurred. If agent i does so, her opponents believe her private stock to be large, and exert no effort and disclose nothing for the remainder of the game, no matter how agent i behaves after time t .⁶⁰

To prove Theorem 4, I obtain a Bellman equation for the payoff function v^c and derive a 'principle of optimality' guaranteeing that candidates α^c and q induce a Bayes' Nash equilibrium. I then construct a sequence of triplets (α^c, q, v^c) satisfying the conditions of the principle of optimality for an increasingly precise approximation of the Bellman equation. I argue that the sequence lies in a suitably compact space and, using Berge's

60. To see why this punishment is sustainable, note that the opponents of agent i expect her to disclose fully as soon as she obtains an innovation, or at the earliest time $t' \in \mathcal{T}$ after t , if one such time exists. If agent i discloses a lower value (at any time after t), or does not disclose at time t' , her opponents will detect another deviation, and continue to believe her private stock to be large. This equilibrium does not satisfy the Intuitive Criterion (Cho and Kreps 1987) as, if k_t^i were indeed large, agent i would be strictly better off if she disclosed k_t^i at time t than a value $d_t^i < q(t)$. An equilibrium that is both symmetric on path and satisfies the Intuitive Criterion is unlikely to exist. As argued in footnote 62, symmetry is likely to require the 'cutoff' disclosure pattern of (ρ_c, β_c) . Moreover, off-path beliefs satisfying the Intuitive Criterion cannot discourage an agent i from disclosing at time t if $q(t)$ is large and $k_t^i > x_0$ is small compared to $q(t)$.

maximum theorem, I construct a triplet (α^c, q, v^c) that induces a Bayes-Nash equilibrium. Finally, I specify off-path behaviour and beliefs inducing a PBE.

6.5 Concealment can be beneficial

In this section, I derive a sufficient condition for the ex-ante payoffs in the equilibrium (ρ_c, β_c) of the game with concealment to exceed those of the equilibrium α_e of the game with forced disclosure. Essentially, assuming that the population is sufficiently large, it is sufficient that either $\lambda \leq 1$ (that is, the frequency of innovations be low), or that the initial stock x_0 be sufficiently small.

I described the equilibrium (ρ_c, β_c) in Theorem 4, and characterised the equilibrium α_e in Theorem 2. The ex-ante payoffs in (ρ_c, β_c) and α_e are $v_0^c(x_0)$ and $v_e(x_0)$, respectively.

Proposition 4. *If n is sufficiently large given x_0 , λ and F and, moreover, $F(\lambda\mu) < 1$ and*

$$x_0 \leq \lambda \mathbb{E}_F[\max\{0, \tilde{z} + x_0 - \lambda\mu\}], \quad (24)$$

then $v_0^c(x_0) \geq v_e(x_0)$.

Proposition 4 is proved in Appendix H. It shows that, even though redundant innovations are likely to be produced in (ρ_c, β_c) , and no benefit can be derived from innovations until a sufficiently large one is obtained, the latter feature can strengthen the effort incentives so much as to improve efficiency over the equilibrium α_e with forced disclosure.

We have seen in Section 6.4 that, if $x_0 \geq \lambda\mu$, then no effort is exerted in either α_e or (ρ_c, β_c) , and the equilibria coincide. Proposition 4 states that, apart from this case, (ρ_c, β_c) improves over α_e in large populations, provided (24) holds.

Note that (24) holds as long as $\lambda \leq 1$ and $x_0 \leq \lambda\mu$ and, assuming that $\lambda > 1$, (24) holds if and only if the initial stock x_0 is sufficiently small.⁶¹ Condition (24) is essentially equivalent to precisely one innovation being disclosed (almost surely) in the equilibrium σ_d of the game with disposal (characterised in Theorem 3). More specifically, the equivalence holds as long as $0 < x_0 < \lambda\mu$ and n is sufficiently large. In other words, (24) rules out the possibility that agents build on each others' innovations in σ_d . Because this never occurs in (ρ_c, β_c) , (24) allows to compare (ex-ante) payoffs in (ρ_c, β_c) and σ_d . This comparison is useful since (24) also implies that innovations are detrimental in α_e , so that payoffs in σ_d exceed those in α_e (Theorem 3).

If either (24) fails, or the number of agents is small, agents may in principle be *worse-off* in the equilibrium with concealment than in the equilibrium with forced disclosure. Among other things, this is because agents cannot build on each others' innovations

61. In detail, (24) is equivalent to $\psi(x_0) \geq 0$, where $\psi(x_0) := \lambda \mathbb{E}_F[\max\{0, \tilde{z} + x_0 - \lambda\mu\}] - x_0$. Moreover, ψ is absolutely continuous on \mathbb{R}_+ with $\psi(\lambda\mu) = 0$ and a.e. derivative $\psi'(x_0) = \lambda[1 - F(\max\{\lambda\mu - x_0, 0\})] - 1$. Then, if $\lambda \leq 1$, ψ is decreasing and hence positive on $[0, \lambda\mu]$. If $\lambda > 1$ then $\psi(0) > 0$ and ψ is convex, so that it crosses 0 from above on $[0, \lambda\mu]$.

in (ρ_c, β_c) . This suggests that the game with concealment admits another, potentially superior PBE with a richer disclosure pattern. However, an equilibrium of this kind is unlikely to be symmetric on path and, as a consequence, could not be compared to α_e in a straightforward manner.⁶²

References

- Admati, A., and M. Perry. 1991. “Joint projects without commitment.” *The Review of Economic Studies* 58 (2): 259–276.
- Akcigit, U., and Q. Liu. 2015. “The role of information in innovation and competition.” *Journal of the European Economic Association* 14 (4): 828–870.
- Akcigit, U., S. Caicedo, E. Miguelez, S. Stantcheva, and V. Sterzi. 2018. *Dancing with the stars: Innovation through interactions*. Working Paper 24466. National Bureau of Economic Research.
- Battaglini, M., S. Nunnari, and T. R. Palfrey. 2014. “Dynamic free riding with irreversible investments.” *American Economic Review* 104 (9): 2858–2871.
- Bessen, J., and A. Nuvolari. 2016. “Knowledge sharing among inventors: some historical perspectives.” Chap. 7 in *Revolutionizing Innovation: Users, Communities and Open Innovation*, 135–157. MIT Press.
- Bimpikis, K., S. Ehsani, and M. Mostagir. 2019. “Designing dynamic contests.” *Operations Research* 67 (2): 339–356.
- Bliss, C., and B. Nalebuff. 1984. “Dragon-slaying and ballroom dancing: The private supply of a public good.” *Journal of Public Economics* 25 (1-2): 1–12.
- Bolton, P., and C. Harris. 1999. “Strategic experimentation.” *Econometrica* 67 (2): 349–374.
- Bonatti, A., and J. Hörner. 2011. “Collaborating.” *American Economic Review* 101 (2): 632–63.

62. To see why symmetry may fail, consider a candidate PBE in which the first disclosure is precisely the first private stock entering some set $(x_0, q_0(t)] \cup [q_1(t), \infty)$, where $q_1(t) > q_0(t) > x_0$. Symmetry requires that, after a first disclosure $d_t^i \leq q_0(t)$ (due to agent i at some time t , say), for any $j \neq i$, the belief of agent i about k_t^j matches the belief of agent j about k_t^i . This constrains disclosure behaviour after an increment which, prior to the first disclosure, raises one’s private stock to a value within $[q_0(t), q_1(t)]$. If full concealment occurs in this case, then agent j would deduce at time t that $k_t^i = d_t^i$ a.s., whereas agent i would believe that $k_t^j > d_t^i$ with strictly positive probability. In contrast, if $q_0(t)$ is disclosed, assuming that $d_t^i = q_0(t)$, agent j would be more optimistic about k_t^i after agent i discloses $d_t^i = q_0(t)$ at time t than agent i is about k_t^j .

- Brander, J. A., and B. J. Spencer. 1983. "Strategic commitment with R&D: the symmetric case." *The Bell Journal of Economics* 14 (1): 225–235.
- Brunk, H. D., G. M. Ewing, and W. R. Utz. 1956. "Some Helly theorems for monotone functions." *Proceedings of the American Mathematical Society* 7 (5): 776–783.
- Campbell, A., F. Ederer, and J. Spinnewijn. 2014. "Delay and deadlines: Freeriding and information revelation in partnerships." *American Economic Journal: Microeconomics* 6 (2): 163–204.
- Chesbrough, H. W. 2003. *Open innovation: The new imperative for creating and profiting from technology*. Harvard Business Press.
- Cho, I.-K., and D. M. Kreps. 1987. "Signaling games and stable equilibria." *The Quarterly Journal of Economics* 102 (2): 179–221.
- Compte, O., and P. Jehiel. 2004. "Gradualism in bargaining and contribution games." *The Review of Economic Studies* 71 (4): 975–1000.
- Curello, G. 2021. "Incentives for Collective Innovation: Omitted material." Working paper, Sep 2021.
- d'Aspremont, C., and A. Jacquemin. 1988. "Cooperative and noncooperative R & D in duopoly with spillovers." *The American Economic Review* 78 (5): 1133–1137.
- Fershtman, C., and S. Nitzan. 1991. "Dynamic voluntary provision of public goods." *European Economic Review* 35 (5): 1057–1067.
- Garfagnini, U., and B. Strulovici. 2016. "Social experimentation with interdependent and expanding technologies." *Review of Economic Studies* 83 (4): 1579–1613.
- Georgiadis, G. 2015. "Projects and team dynamics." *The Review of Economic Studies* 82 (1): 187–218.
- Gradstein, M. 1992. "Time dynamics and incomplete information in the private provision of public goods." *Journal of political Economy* 100 (3): 581–597.
- Halac, M., N. Kartik, and Q. Liu. 2017. "Contests for Experimentation." *Journal of Political Economy* 125 (5): 1523–1569.
- Harhoff, D., and K. R. Lakhani. 2016. *Revolutionizing innovation: Users, communities, and open innovation*. MIT Press.
- Heidhues, P., S. Rady, and P. Strack. 2015. "Strategic experimentation with private payoffs." *Journal of Economic Theory* 159 (1): 531–551.

- Hottenrott, H., and B. Peters. 2012. “Innovative capability and financing constraints for innovation: more money, more innovation?” *Review of Economics and Statistics* 94 (4): 1126–1142.
- Jovanovic, B., and Y. Nyarko. 1996. “Learning by Doing and the Choice of Technology.” *Econometrica* 64 (6): 1299–1310.
- Kamien, M. I., E. Muller, and I. Zang. 1992. “Research joint ventures and R&D cartels.” *The American Economic Review* 82 (5): 1293–1306.
- Katz, M. L. 1986. “An analysis of cooperative research and development.” *The RAND Journal of Economics* 17 (4): 527–543.
- Keller, G., and S. Rady. 2015. “Breakdowns.” *Theoretical Economics* 10 (1): 175–202.
- . 2010. “Strategic experimentation with Poisson bandits.” *Theoretical Economics* 5 (2): 275–311.
- . 2020. “Undiscounted bandit games.” *Games and Economic Behavior* 124 (1): 43–61.
- Keller, G., S. Rady, and M. Cripps. 2005. “Strategic experimentation with exponential bandits.” *Econometrica* 73 (1): 39–68.
- Kerr, W. R. 2008. “Ethnic scientific communities and international technology diffusion.” *The Review of Economics and Statistics* 90 (3): 518–537.
- Klein, N., and S. Rady. 2011. “Negatively correlated bandits.” *The Review of Economic Studies* 78 (2): 693–732.
- Leahy, D., and J. P. Neary. 1997. “Public policy towards R&D in oligopolistic industries.” *The American Economic Review* 87 (4): 642–662.
- Lim, K. 2009. “The many faces of absorptive capacity: spillovers of copper interconnect technology for semiconductor chips.” *Industrial and Corporate Change* 18 (6): 1249–1284.
- Lockwood, B., and J. P. Thomas. 2002. “Gradualism and irreversibility.” *The Review of Economic Studies* 69 (2): 339–356.
- Lucas, R. E., and B. Moll. 2014. “Knowledge growth and the allocation of time.” *Journal of Political Economy* 122 (1): 1–51.
- Marx, L., and S. Matthews. 2000. “Dynamic voluntary contribution to a public project.” *The Review of Economic Studies* 67 (2): 327–358.

- Matthews, S. A. 2013. “Achievable outcomes of dynamic contribution games.” *Theoretical Economics* 8 (2): 365–403.
- Pénin, J. 2007. “Open knowledge disclosure: An overview of the evidence and economic motivations.” *Journal of Economic Surveys* 21 (2): 326–347.
- Perla, J., and C. Tonetti. 2014. “Equilibrium imitation and growth.” *Journal of Political Economy* 122 (1): 52–76.
- Peskir, G., and A. Shiryaev. 2006. *Optimal stopping and free-boundary problems*. Springer.
- Polyanin, A. D., and A. V. Manzhirov. 2008. *Handbook of integral equations*. CRC press.
- Powell, W. W., and E. Giannella. 2010. “Collective invention and inventor networks.” Chap. 13 in *Handbook of the Economics of Innovation*, 1:575–605. North Holland.
- Powell, W. W., K. W. Koput, and L. Smith-Doerr. 1996. “Interorganizational collaboration and the locus of innovation: Networks of learning in biotechnology.” *Administrative science quarterly* 41 (1): 116–145.
- Radner, R., R. Myerson, and E. Maskin. 1986. “An example of a repeated partnership game with discounting and with uniformly inefficient equilibria.” *The Review of Economic Studies* 53 (1): 59–69.
- Redding, S. 2002. “Path dependence, endogenous innovation, and growth.” *International Economic Review* 43 (4): 1215–1248.
- Reinganum, J. F. 1983. “Uncertain innovation and the persistence of monopoly.” *The American Economic Review* 73 (4): 741–748.
- Spence, M. 1984. “Cost reduction, competition, and industry performance.” *Econometrica: Journal of the Econometric Society* 52 (1): 101–121.
- Strausz, R. 1999. “Efficiency in sequential partnerships.” *Journal of Economic Theory* 85 (1): 140–156.
- Suzumura, K. 1992. “Cooperative and Noncooperative R&D in an Oligopoly with Spillovers.” *The American Economic Review* 82 (5): 1307–1320.
- Von Hippel, E. 1987. “Cooperation between rivals: Informal know-how trading.” *Research Policy* 16 (6): 291–302.
- Yildirim, H. 2006. “Getting the ball rolling: Voluntary contributions to a large-scale public project.” *Journal of Public Economic Theory* 8 (4): 503–528.

Appendices

A The game with concealment

This appendix is devoted to the rigorous definition of the game with concealment, described informally in Section 6.1.

Histories. An *event* is a triplet

$$e := (t, z, s)$$

where $t \in \mathbb{R}_+$, $s := (s^i)_{i=1}^n \in \mathbb{R}_+^n$ and $z := (z^i)_{i=1}^n \in \mathbb{R}_+^n$ are such that one of the following holds (i) if $z^i > 0$ then $t \notin T$ and $z^j = s^j = 0$ for each $j \neq i$, and (ii) $t \in \mathcal{T}$ if $z^i = s^i = 0$ for all i . The event e states that, at time t , for each i , obtained an increment of size z^i if $z^i > 0$, and did not obtain an increment otherwise, and that she disclosed the value $d_t^i = x_t + s^i$ if $s^i > 0$, and did not disclose otherwise. Condition (i) follows from the fact that increments occur at unpredictable times and (almost surely) only finitely many of them occur within any given interval of time. Condition (ii) states that the event that no agent obtained a private increment or disclosed is recorded only at times within \mathcal{T} . Let E be the set of all events, viewed as a subset of \mathcal{R}^{2n+1} , and let \mathcal{L} be the Lebesgue σ -algebra on \mathcal{R}^{2n+1} . Let $E_\infty := \bigcup_{m=1}^\infty \mathcal{R}^{(2n+1) \times m}$ and let \mathcal{E}_∞ be the σ -algebra on E_∞ such that $(E_\infty, \mathcal{E}_\infty)$ is the disjoint union of the spaces $(\mathcal{R}^{(2n+1) \times m}, \mathcal{L}^m)_{m=1}^\infty$.

An *omniscient history* is a finite set $\omega \subset E$ such that, if $\omega \neq \emptyset$, then

$$\omega := \{(t_m, z_m, d_m) : m = 1, \dots, M\} \tag{25}$$

for some $M \in \mathbb{N}$ such that

- (i) $(t_m)_{m=0}^M$ is strictly increasing and $\mathcal{T} \cap [0, t_M] \subset (t_m)_{m=0}^M$
- (ii) if $s_m^i > 0$ then, either $t_m \in \mathcal{T}$, or $s_l^j = 0$ for all j and $l < m$, or $z_m^i > 0$.

A (non-empty) omniscient history ω records the events occurred up to time t_M . Condition (i) states that at most one event occurs at each given time, and that an event always occurs at times in \mathcal{T} . Condition (ii) reflects the restriction on disclosures: if agent i discloses at time t_m then either $t_m \in \mathcal{T}$, or $x_{t_m} = 0$ (that is, no agent disclosed in the past), or agent i obtained an increment (of size z_m^i) at time t_m . Let Ω be the set of all omniscient histories, viewed as a subset of E_∞ , and let \mathcal{O} be the sub- σ -algebra derived from \mathcal{E}_∞ . Also let $T, X : \Omega \rightarrow \mathbb{R}_+$ be given by

$$T(\omega) := \max\{0\} \cup \{t : \exists z, s, (t, z, s) \in \omega\} \quad \& \quad X(\omega) := x_0 + \sum_{(t, z, s) \in \omega} \max_i s^i,$$

and $K : \Omega \rightarrow \mathbb{R}_+^n$ be given by

$$K^i(\omega) := \begin{cases} x_0 & \text{if } \omega = \emptyset \\ \max\{X(\omega), K(\omega \setminus \{(t_M, z_M, s_M)\}) + z^i\} & \text{if } \omega \neq \emptyset \text{ is expressed as (25).} \end{cases}$$

The maps T , X , and K keep track of the current time t , the public stock x_t , and the private stocks $(k_t^i)_{i=1}^n$, respectively.

For each $i \in \{1, \dots, n\}$, let $\psi_i : E \rightarrow \mathbb{R}^{2n+1} \cup \{\emptyset\}$ be given by

$$\psi_i(t, z, s) = \begin{cases} \emptyset & \text{if } z^j > 0 = d^j \text{ for some } j \neq i. \\ (t, (0, \dots, 0, z^i, 0, \dots, 0), s) & \text{otherwise.} \end{cases}$$

The map ψ_i describes the information obtained by agent i after an event occurs. Each agent observes disclosures, as well as the increments that she obtains, but does not observe the increments obtained by her opponents. Let $E^i := \psi_i(E) \setminus \{\emptyset\}$.

A *private history* for agent i is a $h \subset E^i$ such that $h = \psi_i(\omega) \setminus \{\emptyset\}$ for some omniscient history $\omega \in \Omega$. That is, a subsets of events observable to agent i that are derived from an omniscient history. Let H^i be the set of all private histories for agent i , viewed as a subset of E_∞ , and let \mathcal{H}^i be the sub- σ -algebra derived from \mathcal{E}_∞ .

Let $\psi_0 : E \cup (\bigcup_{i=1}^n E^i) \rightarrow \mathbb{R}^{2n+1} \cup \{\emptyset\}$ be given by

$$\psi_0(t, z, s) = \begin{cases} \emptyset & \text{if } z^i > 0 = d^i \text{ for some } i. \\ (t, 0, s) & \text{otherwise.} \end{cases}$$

The map ψ_0 describes the information obtained by all agent after an event occurs. Let $E^0 := \psi_0(E) \setminus \{\emptyset\}$. A *public history* is a $h \subset E^0$ such that $h = \psi_0(\gamma) \setminus \{\emptyset\}$ for some $\gamma \in \Omega \cup (\bigcup_{i=1}^n H^i)$. Let H^0 be the set of all public histories, and extend the definitions of T , X and K to $\bigcup_{i=0}^n H^i$ in the natural way.

Strategies. Let A be the quotient space of Lebesgue measurable maps $a : \mathbb{R}_+ \rightarrow [0, 1]$ by the equivalence relation \sim_A such that $a \sim \hat{a}$ if and only if $\int_0^t a_s ds = \int_0^t \hat{a}_s ds$ for all $t > 0$, endowed with the topology of pointwise convergence of the function $t \mapsto \int_0^t a_s ds$. Let \mathcal{A} be the Borel σ -algebra of A . Let D be the space of Lebesgue measurable $d : \mathbb{R}_+^2 \rightarrow [0, 1]$ by the ' $\ell \times F_0$ -a.e. equal' relation, where ℓ denotes the Lebesgue measure, and F_0 is the average between F and the degenerate probability measure that puts all mass at 0. Endow D with the topology of $\ell \times F_0$ -a.e. convergence, and let \mathcal{D} be the Borel σ -algebra of D .

A (*pure*) *strategy* for agent i is a pair $\rho^i := (\xi^i, \chi^i)$ such that

- (i) $\xi^i : (H^i, \mathcal{H}^i) \rightarrow (A, \mathcal{A})$ and $\chi^i : (H^i, \mathcal{H}^i) \rightarrow (D, \mathcal{D})$ are measurable,
- (ii) for all $h \in H^i$ and $t \geq 0$ such that $[\chi^i(h)](t, 0) > 0$, either $X(h) = x_0$ or $T(h) + t \in \mathcal{T}$,

- (iii) $\xi^i(h) = t \mapsto [\xi^i(h')](T(h) - T(h') + t)$ and $\chi^i(h) = (t, z) \mapsto [\chi^i(h')](T(h) - T(h') + t, z)$ for any $h, h' \in H^i$ such that $T(h) \geq T(h')$, $K^i(h) = K^i(h')$ and $\psi_0(h) \setminus \{\emptyset\} = \psi_0(h') \setminus \{\emptyset\}$.

Refer to ξ^i and χ^i as *effort* and *disclosure policies*.

Given $h \in H^i$, ρ^i pins down agent i 's behaviour between $T(h)$ and the earliest time $t \geq T(h)$ at which either agent i obtains an innovation, or one of her opponents discloses, or $[\chi^i(h)](t - T(h), 0) > 0$, where $t = \infty$ if none of these events occur. For all $T(h) < s < t$, agent i exerts effort $[\xi^i(h)](s - T(h))$ at time s . Moreover, if $t < \infty$ and agent i obtains an innovation of size $z > 0$ at time t , agent i picks the disclosure

$$d_t^i = X(h) + [K^i(h) + z - X(h)][\chi^i(h)](t - T(h), z).$$

Finally, if $t < \infty$ and agent i obtains no innovation at time t then agent i discloses $X(h) + [K_i(h) - X(h)][\chi^i(h)](t - T(h), 0) > X(h)$ at time t (where the inequality holds by definition of t).

Condition (ii) states that, if agent i discloses in the absence of an increment at time $T(h) + t$, then either no disclosure occurred in the past, or this time lies in \mathcal{T} . Condition (iii) states that the strategy ρ^i does not condition on the full trajectory, of agent i 's private stock, but only on its current value. Let R^i be the set of all strategies for agent i , and $R := \prod_{i=1}^n R^i$.

Beliefs and information sets. A *belief* for agent i is a probability measure $b := \prod_{j \neq i} b_j$ over \mathbb{R}_+^{n-1} . Let B^i be the set of all beliefs for agent i . Given $h \in H^i$, say that $b := \prod_{j \neq i} b_j \in B^i$ is *admissible* at h if the support of b_j lies above $X(h)$ for all $j \neq i$. In words, we require that the belief of agent i is consistent with the fact that $k_t^j \geq x_t$ for each j and t . Given h , b_j refers to the distribution of k_t^j , where $t = T(\psi_0(h))$ is the time of the last ‘public’ event in h . Let $B^i(h)$ be the set of beliefs for agent i that are admissible at h . An *information set for agent i* is a pair $y = (h, b)$ such that $h \in H^i$ and $b \in B^i(h)$. Let Y^i be the set of all information sets for agent i . A *system of beliefs for agent i* is a map $\beta^i : H^i \rightarrow B^i$ such that $(h, \beta^i(h)) \in Y^i$ for any $h \in H^i$. Given β^i , $h \in H^i$, and $j \neq i$, let $\beta_j^i(h) := b_j$ where $\beta^i(h) = \prod_{l \neq i} b^l$. Let \mathcal{B}^i be the set of systems of beliefs for agent i , for each $i \leq n$, and $\mathcal{B} := \prod_{i=1}^n \mathcal{B}^i$.

Payoffs. Fix a strategy profile $\rho \in R$, i , and an information set $y := (h, b) \in Y^i$. Playing ρ after agent i reaches y induces a random path of play involving a (random) total number $\tilde{m} \in \{0, 1, \dots, \infty\}$ of events. Moreover, the path of play may be expressed as a random sequence $(\tilde{\omega}_m)_{m=0}^{\tilde{m}} \subset \Omega$ where $\psi_i(\tilde{\omega}_0) = h$, $(\psi_j(\tilde{\omega}_0))_{j \neq i} \sim b$ and, for all $m \in \mathbb{N}$, conditional on $\tilde{m} \geq m$, $\tilde{\omega}_m = \tilde{\omega}_{m-1} \cup \{\tilde{e}_m\}$ for some $\tilde{e}_m \in E$. The (*continuation*) *payoff*

to agent i at information set $y := (h, b)$, when the strategy profile ρ is played, is

$$v_\rho^i(y) := \mathbb{E} \left[\sum_{m=0}^{\tilde{m}} \int_{T(\tilde{\omega}_m)}^{T(\tilde{\omega}_{m+1})} r e^{r[T(h)-t]} u([\xi^i(\psi_i(\tilde{\omega}_m))](t - T(\psi_i(\tilde{\omega}_m))), X(\tilde{\omega}_m)) dt \right]$$

where $T(\tilde{\omega}_{\tilde{m}+1}) := \infty$ if $\tilde{m} < \infty$.

Equilibrium. Given $\rho := (\rho^i)_{i=1}^n \in R$ and i , let $\rho^{-i} := (\rho^j)_{j \neq i}$. Given $y \in Y$, ρ^i is a *best response* for agent i against ρ^{-i} at y if

$$\rho^i \in \arg \max_{\hat{\rho}^i \in \rho^i} v_{(\hat{\rho}^i, \rho^{-i})}^i(y).$$

For each i , let $\underline{b}^i \in B^i(\emptyset)$ be the belief assigning probability 1 to (x_0, \dots, x_0) . Given a pair (ρ, β) where $\rho := (\rho^i)_{i=1}^n \in R$ is a profile of strategies and $\beta := (\beta^i)_{i=1}^n \in \mathcal{B}$ is a profile of systems of beliefs, is a *(weak) perfect Bayesian equilibrium* if, for all i , $\beta^i(\emptyset) = \underline{b}^i$ and, for all $h \in H^i$, all of the following hold:

- (a) ρ^i is a best response for agent i against ρ^{-i} at $(h, \beta^i(h))$
- (b) for all j and $h' \in H^j$ such that $T(h) = T(h')$ and $\psi_0(h) = \psi_0(h')$, and any $l \notin \{i, j\}$, $\beta_l^i(h) = \beta_l^j(h')$.
- (c) for $j \neq i$ and $e, \hat{e} \in E$ such that $h \cup \{e\}, h \cup \{\hat{e}\} \in H^i$, $\beta_j^i(h \cup \{e\}) = \beta_j^i(h \cup \{\hat{e}\})$ if $s^j = \hat{s}^j$, where $e = (t, z, s)$ and $\hat{e} = (\hat{t}, \hat{z}, \hat{s})$.
- (d) for all $h' \in H^i$ such that $\Pr(\tilde{\gamma} \neq \emptyset) > 0$ where $\tilde{\gamma} =: \{m \in \{0, \dots, \tilde{m}\} : \psi_i(\tilde{\omega}_m) = h'\}$ and $(\tilde{\omega}_m)_{m=0}^{\tilde{m}}$ is the random path of play (defined above) arising when ρ is played at $(h, \beta^i(h))$, $(\psi_j(\tilde{\omega}_{\min \tilde{\gamma}}) | \tilde{\gamma} \neq \emptyset)_{j \neq i} \sim \beta^i(h')$.

Condition (b) states that the beliefs of agents i and j about agent l 's private information are identical at private histories that contain the same public information. Condition (c) states that, agent i 's beliefs about agent j 's private information are identical at any two private histories that differ only in their latest event, as long as agent j 's role is identical across these two events. Finally, Condition (d) states that, for any private history h' that is reached with strictly positive probability from agent i 's information set $(h, \beta^i(h))$ when ρ is played, the belief $\beta^i(h')$ of agent i at h' is derived from her belief $\beta^i(h)$ at h using Bayes' rule.

B Preliminary results

This section contains some preliminary results used in various proofs. Note that, since b is concave, there are $k, k' \geq 0$ such that $b(x) \leq k + k'x$ for all $x \geq 0$. Let $\bar{v} : \mathbb{R}_+ \rightarrow \mathbb{R}$ be

given by

$$\bar{v}(x) := k + k'(x + \lambda\mu n) \quad (26)$$

and let \mathcal{V} be the set of Lebesgue-measurable $v : \mathbb{R}_+ \rightarrow \mathbb{R}$ such that $b(x) \leq v(x) \leq \bar{v}(x)$. Given $x \geq 0$ and $v \in \mathcal{V}$, let

$$L^e v(x) := \mathbb{E}_F[v(x + \tilde{z})] \quad \& \quad L^d v(x) = \mathbb{E}_F[\max\{v(x), v(x + \tilde{z})\}]. \quad (27)$$

Claim 1. Let $v \in \mathcal{V}$ be such that $v - b$ is decreasing. Then $L^k v - b$ is decreasing for $k \in \{d, e\}$.

Proof. Fix $0 \leq x_1 \leq x_2$ and note that

$$\begin{aligned} L^e v(x_2) - L^e v(x_1) &= \mathbb{E}_F[v(x_2 + \tilde{z}) - v(x_1 + \tilde{z})] \leq \mathbb{E}_F[b(x_2 + \tilde{z}) - b(x_1 + \tilde{z})] \\ &\leq b(x_2) - b(x_1) \end{aligned}$$

where the first inequality holds since $v - b$ is decreasing, and the second since b is concave.⁶³ Moreover,

$$\begin{aligned} L^d v(x_2) - L^d v(x_1) &= \mathbb{E}_F[\max\{v(x_2 + \tilde{z}), v(x_2)\} - \max\{v(x_1 + \tilde{z}), v(x_1)\}] \\ &\leq \mathbb{E}_F[\max\{v(x_2 + \tilde{z}) - v(x_1 + \tilde{z}), v(x_2) - v(x_1)\}] \\ &\leq \mathbb{E}_F[\max\{b(x_2 + \tilde{z}) - b(x_1 + \tilde{z}), b(x_2) - b(x_1)\}] \\ &= b(x_2) - b(x_1). \end{aligned}$$

where the second inequality holds since $v - b$ is decreasing. □

Let $T : [0, 1] \times [0, n] \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ be given by

$$T(a, \alpha, x, l) := \frac{b(x) - c(a, x) + \alpha\lambda l}{1 + \alpha\lambda} \quad (28)$$

and $\bar{T} : [0, 1] \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ be given by $\bar{T}(a, x, l) := T(a, na, x, l)$.

Claim 2. $T(a, \alpha, x, L^k v(x)) \leq \bar{v}(x)$ for all $a, \alpha, x, v \in \mathcal{V}$ and $k \in \{d, e\}$.

Proof. Fix x, v and k and note that $L^k v(x) \leq L^k \bar{v}(x)$, since $v \leq \bar{v}$. Then, fixing a and a' ,

63. In detail, $\mathbb{E}_F[b(x_2 + \tilde{z}) - b(x_1 + \tilde{z})] = \mathbb{E}_F\left[\int_{x_1}^{x_2} b'(x + \tilde{z})dx\right] \leq \mathbb{E}_F\left[\int_{x_1}^{x_2} b'(x)dx\right] = b(x_2) - b(x_1)$.

we have

$$\begin{aligned}
T(a, a', x, L^k v(x)) &= \frac{b(x) - c(a, x) + a' \lambda L^k v(x)}{1 + a' \lambda} \\
&\leq \frac{k + k' x + n \lambda L^k \bar{v}(x)}{1 + n \lambda} \\
&= \frac{k + k' x + n \lambda \{k + k' [x + \mu(1 + \lambda n)]\}}{1 + n \lambda} \\
&= \bar{v}(x).
\end{aligned}$$

where the first inequality holds since $L^k v(x) \leq L^k \bar{v}(x)$ and $b(x) \leq k + k' x \leq L^k \bar{v}(x)$. \square

C Proofs of Theorem 1 and Corollary 1

The proof of Theorem 1 is based on the following ‘principle of optimality’, which I state without proof.

Lemma 4. *If (6) admits a solution $w \in \mathcal{V}$, then $w_* = w$. Moreover, a strategy profile α induces welfare w_* for any $x_0 \geq 0$ if and only if (7) holds.*

Proof of Theorem 1. The proof has several steps.

Step 1. w_* is the unique solution $w \in \mathcal{V}$ to (6), w_* is absolutely continuous and increasing, and $w_* - b$ is decreasing. From Lemma 4, it suffices to exhibit an absolutely continuous and increasing solution $w \in \mathcal{V}$ to (6) such that $w - b$ is decreasing.

Given $w \in \mathcal{V}$, $x \geq 0$, and $a \in [0, 1]$, let

$$Pw(x) := \max_{a \in [0, 1]} \bar{T}(a, x, L^e w(x)) \quad (29)$$

where \bar{T} is defined below (28). Let \mathcal{V}_* be the subset of increasing $w \in \mathcal{V}$ such that $w - b$ is decreasing. Note that $w \in \mathcal{V}$ solves (6) if (and only if) it is a fixed-point of P .⁶⁴ Moreover, elements of \mathcal{V}_* are increasing and absolutely continuous, since b is concave. Then, it suffices to show that P has a fixed point in \mathcal{V}_* . Note that (\mathcal{V}_*, \geq) is a complete lattice,⁶⁵ where \geq is the point-wise order, and P is increasing. Then, from Tarski’s fixed-point theorem, it suffices to show that $Pw \in \mathcal{V}_*$ for all $w \in \mathcal{V}_*$.

64. Indeed, $w = Pw$ if and only if

$$0 = \max_{a \in [0, 1]} L^e w(x) + \frac{b(x) - c(a, x) - L^e w(x)}{1 + a \lambda n} - w(x) \Leftrightarrow 0 = \max_{a \in [0, 1]} a \lambda n L^e w(x) + b(x) - c(a, x) - w(x)(1 + a \lambda n)$$

and the second equation is clearly equivalent to (6).

65. It suffices to show that, for an arbitrary $V \subset \mathcal{V}_*$, the maps $\tilde{w} : x \mapsto \sup_{w \in V} w(x)$ and $\hat{w} : x \mapsto \inf_{w \in V} w(x)$ belong to \mathcal{V}_* . For the former, note that \tilde{w} is Lebesgue-measurable since each $w \in V$ is and, given $x \geq 0$, $b(x) \leq \tilde{w}(x) \leq \bar{w}(x)$ since $b(x) \leq w(x) \leq \bar{w}(x)$ for all $w \in V$. Then, $\tilde{w} \in \mathcal{V}$. Now fix $y \geq x$ and note that, for each $w \in V$, $0 \leq w(y) - w(x) \leq \tilde{w}(y) - w(x)$ and $w(y) - \tilde{w}(x) \leq w(y) - w(x) \leq b(y) - b(x)$. As $w \in V$ is arbitrary, $0 \leq \tilde{w}(y) - \tilde{w}(x) \leq b(y) - b(x)$. As x and y are arbitrary, \tilde{w} is increasing and $\tilde{w} - b$ is decreasing, so that $\tilde{w} \in \mathcal{V}_*$. The argument for \hat{w} is similar.

Fix $w \in \mathcal{V}_*$. Clearly, Pw is Lebesgue-measurable. Moreover, $Pw(x) \geq \bar{T}(0, x, L^e w(x)) = b(x)$ and, from Claim 2, $Pw \leq \bar{v}$. Also, since w is absolutely continuous and u is continuously differentiable, $L^e w$ and therefore Pw are absolutely continuous. Finally, since $w' \geq 0$ then $(L^e w)' \geq 0$ and, since u is increasing in x , $(Pw)' \geq 0$.

It remains to show that $(Pw)' \leq b'$. Note that, for all $x \geq 0$ such that $L^e w$ is differentiable at x , and for $a \in \arg \max_{a' \in [0,1]} \bar{T}(a', x, L^e w(x))$,

$$(Pw)'(x) = \frac{b'(x) - c_2(a, x) + a\lambda n(L^e w)'(x)}{1 + a\lambda n} \leq \frac{b'(x) - c_2(a, x) + a\lambda n b'(x)}{1 + a\lambda n} \leq b'(x)$$

where the equality holds by the envelope theorem, the first inequality follows from Claim 1, and the second holds since $c_2 \geq 0$.

Step 2. w_* is induced by some symmetric, decreasing Markov profile $\alpha_* \in A^n$. Let $\alpha_* \in A^n$ given by

$$\alpha_*^i(x) := \min \arg \max_{a \in [0,1]} a\lambda n \{ \mathbb{E}_F[w_*(x + \tilde{z})] - w_*(x) \} - c(a, x). \quad (30)$$

Since $w_* \in \mathcal{V}$ solves (6), Lemma 4 implies that w_* is induced by α_* . Moreover, α_* is symmetric, so that it remains to show that $\alpha_*^i(x)$ is decreasing. From the argument in footnote 64,

$$\alpha_*^i(x) \in \min \arg \max_{a \in [0,1]} \bar{T}(a, x, L^e w_*(x))$$

for all $x \geq 0$. Note that $\Psi^* : a \mapsto \bar{T}(a, x, L^e w_*(x))$ is differentiable on $(0, 1)$ for all $x \geq 0$, and its derivative has the same sign as

$$\psi^*(a, x) := L^e w_*(x) - [b(x) - c(a, x)] - \left(\frac{1}{n\lambda} + a \right) c_1(a, x). \quad (31)$$

Hence, it suffices to show that ψ^* is decreasing in a for all $x \geq 0$ (so that Ψ^* is quasi-concave) and decreasing in x for all $a \in [0, 1]$. To this end note that, since $w_* \in \mathcal{V}_*$ (from Step 1) and u is twice continuously differentiable, ψ^* is absolutely continuous and, for almost every (a, x) ,

$$\begin{aligned} \psi_1^*(a, x) &= -\left(\frac{1}{n\lambda} + a \right) c_{11}(a, x) \\ \psi_2^*(a, x) &= (L^e w_*)'(x) - [b'(x) - c_2(a, x)] - \left(\frac{1}{n\lambda} + a \right) c_{12}(a, x) \end{aligned}$$

Since ψ^* is absolutely continuous and $c_{11} \geq 0$, then ψ^* is decreasing in a for all $x \geq 0$. It remains to prove that $\psi_2^* \leq 0$. Note that, since c_{12} is increasing in a , ψ_2^* is decreasing in a . Then

$$\psi_2^*(a, x) \leq \psi_2^*(0, x) \leq (L^e w_*)'(x) - b'(x) \leq 0$$

where the second inequality holds since $c_2(0, x) = 0$ and $c_{12} \geq 0$, and the last follows from Claim 1, since $w_* \in \mathcal{V}_*$.

Step 3. $\alpha_*^i(\tilde{x}_t) \rightarrow 0$ as $t \rightarrow \infty$, and $\lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$. Note that $\lim_{t \rightarrow \infty} \tilde{x}_t$ exists (a.s.) since \tilde{x}_t is increasing in t . Since α_*^i is decreasing, it is clear that $\lim_{t \rightarrow \infty} \tilde{x}_t > x$ a.s. for any $x \geq 0$ such that $\alpha_*^i(x) > 0$. Then, $\alpha_*^i(\tilde{x}_t) \rightarrow 0$ as $t \rightarrow \infty$ as long as $\lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$.⁶⁶

To prove that $\lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$, note that

$$\mathbb{E}_F[w_*(x + \tilde{z}) - w_*(x)] \leq \mathbb{E}_F[b(x + \tilde{z}) - b(x)] \leq b'(x)\mu$$

where the first inequality holds since $w_* - b$ is decreasing, and the second follows from the mean-value theorem, since b' is decreasing. Then, for all $a \in (0, 1]$, (2) implies that

$$\lim_{x \rightarrow \infty} \lambda n \{ \mathbb{E}_F[w_*(x + \tilde{z})] - w_*(x) \} - c_1(a, x) < 0$$

so that, by definition of α_*^i , $\lim_{x \rightarrow \infty} \alpha_*^i(x) < a$. As a is arbitrary, $\lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$. \square

Proof of Corollary 1. Suppose first that (8) admits a solution $x \geq 0$. By continuity, it admits a smallest solution $\hat{x} \geq 0$, and (8) holds with equality at \hat{x} . Since $\alpha_*^i(\tilde{x}_t) \rightarrow 0$ a.s. as $t \rightarrow \infty$ (Theorem 1), it suffices to show that $\alpha_*^i(\hat{x}) = 0$ and $\alpha_*^i(x) > 0$ for all $0 \leq x < \hat{x}$.

For the former, note that the left-hand side of (8) is decreasing in x , since b is concave, whereas the right-hand side is increasing. Then, (8) holds with ' \leq ' for all $x \geq \hat{x}$. Hence, b solves (6) at all $x \geq \hat{x}$. Therefore, $w_*(x) = b(x)$ for all $x \geq \hat{x}$, as w_* is the unique solution to (6) in \mathcal{V} (Step 1 of the proof of Theorem 1). Then, from (30), $\alpha_*^i(\hat{x}) = 0$.

For the latter, suppose that $\alpha_*^i(\tilde{x}) = 0$ for some $0 \leq \tilde{x} < \hat{x}$, and seek a contradiction. Then, since α_*^i is decreasing (Theorem 1), $\alpha_*^i(x) = 0$ for all $x \geq \tilde{x}$ and, since α_*^i induces w_* , $w_*(x) = b(x)$ for all $x \geq \tilde{x}$. Since $\tilde{x} < \hat{x}$, (8) holds with '>'. But this contradicts the fact that α_*^i solves (7) at \tilde{x} .

Suppose now that (8) holds with '<' for all $x \geq 0$. Then, from the argument in the previous paragraph, α_*^i is strictly positive, so that effort never ceases. Finally, $\tilde{x}_t \rightarrow \infty$ a.s. as $t \rightarrow \infty$ since $\alpha_*^i(\tilde{x}_t) \rightarrow 0$ a.s. as $t \rightarrow \infty$ (Theorem 1). \square

D Proof of Theorem 2

I derive a general result (Theorem 5), state a 'principle of optimality' (Lemma 5), and then prove Theorem 2. Theorem 5 is used to prove Theorem 3, too.

66. Indeed, if $\alpha_*^i(x) > 0$ for all $x \geq 0$ then $\lim_{t \rightarrow \infty} \tilde{x}_t = \infty$ a.s., so that $\lim_{t \rightarrow \infty} \alpha_*^i(\tilde{x}_t) \stackrel{\text{a.s.}}{=} \lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$. If instead $\alpha_*^i(x) = 0$ for some $x \geq 0$, let $\hat{x} = \inf\{x \geq 0 : \alpha_*^i(x) = 0\}$. By definition of α_*^i , $\alpha_*^i(\hat{x}) = 0$. Then $\Pr(\lim_{t \rightarrow \infty} \alpha_*^i(\tilde{x}_t) = 0) = \Pr(\lim_{t \rightarrow \infty} \tilde{x}_t \geq \hat{x}) = \lim_{m \rightarrow \infty} \Pr(\lim_{t \rightarrow \infty} \tilde{x}_t > \hat{x} - 1/m) = 1$.

Let \mathcal{V}' be the set of $v \in \mathcal{V}$ such that $v \leq w_*$. The set A was defined in Appendix A. For $k \in \{e, d\}$, let \mathcal{V}_k be the set of pairs $(\alpha_k^i, v_k) \in A \times \mathcal{V}'$ that solve the system of functional equations

$$v_k(x) = \max_{a \in [0,1]} [b(x) - c(a, x)] + \lambda [a + (n-1)\alpha_k^i(x)] [L^k v_k(x) - v_k(x)] \quad (32)$$

$$\alpha_k^i(x) \in \arg \max_{a \in [0,1]} a \lambda [L^k v_k(x) - v_k(x)] - c(a, x), \quad (33)$$

where L^k was defined in (27). The next general result is used to prove Theorem 2 (as well as Theorem 3). It is proved at the end of this section. The efficient schedule α_*^i was defined in (30).

Theorem 5. *Given $k \in \{e, d\}$, $\mathcal{V}_k = \{(\alpha_k^i, v_k)\}$. Moreover, α_k^i is continuous and decreasing, and below α_*^i . v_k is continuous and $v_k - b$ is decreasing.*

The following ‘principle of optimality’, stated without proof, is used to prove Theorem 2.

Lemma 5. *A Markov strategy $\alpha^i \in A$ induces a symmetric Markov perfect equilibrium if and only if there is $v \in \mathcal{V}'$ such that v solves (10) with $\alpha^j = \alpha^i$ for all j , and α^i solves (11) with $v^i = v$. Moreover, in this case, the time- t continuation payoff of each agent is $v(x_t)$.*

Proof of Theorem 2. Let α_e^i and v_e satisfy the conditions of Theorem 5, and let α_e be the symmetric profile associated with α_e^i . To prove existence, note that v_e solves (10) with $\alpha^j = \alpha_e^i$ for all j , and α_e^i solves (11) with $v^i = v_e$. Then, from Lemma 5, α_e^i forms a symmetric Markov perfect equilibrium, and v_e is the equilibrium payoff function.

To prove that the equilibrium is unique, fix strategy $\alpha^i \in A$ inducing a symmetric Markov perfect equilibrium, and let v be the associated payoff function. Then, from Lemma 5, $v \in \mathcal{V}'$, v solves (10) with $\alpha^j = \alpha^i$ for all j , and α^i solves (11) with $v^i = v$. Then, $(\alpha^i, v) \in \mathcal{V}_e$. Hence, from Theorem 5, $\alpha^i = \alpha_e^i$ and $v = v_e$.

The other parts of Theorem 2 follow immediately from Theorem 5. \square

Proof of Theorem 5. Fix $k \in \{e, d\}$. The proof has several steps.

Step 1. *There are $(\underline{\alpha}_k^i, \underline{v}_k), (\bar{\alpha}_k^i, \bar{v}_k) \in \mathcal{V}_k$ such that $\underline{\alpha}_k^i \leq \alpha_k^i \leq \bar{\alpha}_k^i \leq \alpha_*^i$ and $\underline{v}_k \leq v_k \leq \bar{v}_k$ for all $(\alpha^i, v) \in \mathcal{V}_k$.*

Note that, for any $x \geq 0$ and $l \geq b(x)$, there is a unique $Q(x, l) \in [0, 1]$ such that

$$Q(x, l) \in \arg \max_{a \in [0,1]} a \lambda \frac{l - [b(x) - c(Q(x, l), x)]}{1 + \lambda n Q(x, l)} - c(a, x). \quad (34)$$

Indeed, the objective is continuously differentiable and concave in a , and its derivative

has the same sign as $\psi(Q(x, l)|x, l)$ where

$$\psi(a|x, l) := l - [b(x) - c(a, x)] - \left(\frac{1}{\lambda} + na\right)c_1(a, x)$$

is decreasing in a , since $c(a, x)$ is convex in a . Given $v \in \mathcal{V}'$, and $x \geq 0$, let

$$P_kv(x) := \bar{T}(Q(x, L^k v(x)), x, L^k v(x))$$

where \bar{T} is defined below (28).

Claim 3. $\bar{T}(a, x, L^k v(x)) \leq P_kv(x)$ for $x \geq 0$, $v \in \mathcal{V}'$, and $0 \leq a \leq Q(x, L^k v(x))$.

Claim 3 is proved at the end of this proof. I prove that P_k maps \mathcal{V}' to itself. Note that

$$L^k v \leq L^k w_* = L^e w_* \quad \text{for all } v \in \mathcal{V}' \quad (35)$$

where the inequality holds since $v \leq w_*$, and the equality since w_* is increasing (Theorem 1). Fix $v \in \mathcal{V}'$. Note that Q and, therefore, P_kv are measurable. Fix $x \geq 0$. Since $Q(x, L^k v(x)) \geq 0$, from Claim 3, $P_kv(x) \geq \bar{T}(0, x, L^k v(x)) = b(x)$. Then,

$$P_kv(x) \leq \bar{T}(Q(x, L^k v(x)), x, L^e w^*(x)) \leq Pw_*(x) = w_*(x)$$

where the first inequality follows from (35) since $\bar{T}(a, x, l)$ is increasing in l , the second follows from (29), and the equality from the proof of Step 1 of the proof of Theorem 1. Since $x \geq 0$ is arbitrary, $P_kv \in \mathcal{V}'$ as desired.

I show that P_k is increasing (in the pointwise order). Fix $v, w \in \mathcal{V}'$ such that $w \leq v$, and $x \geq 0$. Then $L^k w(x) \leq L^k v(x)$. Then $P_kw(x) \leq \bar{T}(Q(x, L^k w(x)), x, L^k v(x))$ as $\bar{T}(a, x, l)$ is increasing in l , and $Q(x, L^k w(x)) \leq Q(x, L^k v(x))$ as $Q(x, l)$ is increasing in l (since $\psi(a|x, l)$ is). Together with Claim 3, the latter implies that $\bar{T}(Q(x, L^k w(x)), x, L^k v(x)) \leq P_kv(x)$. Hence, $P_kw(x) \leq P_kv(x)$ and, since $x \geq 0$ is arbitrary, P_k is increasing.

Note that (\mathcal{V}', \geq) is a complete lattice. Since P_k is increasing and maps \mathcal{V}' to itself, Tarski's fixed-point theorem implies that P_k has smallest and largest fixed points $\underline{v}_k, \bar{v}_k \in \mathcal{V}'$. Let $\underline{\alpha}_k^i(x) := Q(x, L^k \underline{v}_k(x))$ and $\bar{\alpha}_k^i(x) := Q(x, L^k \bar{v}_k(x))$. Note that, for any $(\alpha_k^i, v_k) \in A \times \mathcal{V}'$ and $x \geq 0$, '(32) and (33)' is equivalent to ' $v_k(x) = P_kv_k(x)$ and $\alpha_k^i(x) = Q(x, L^k v_k(x))$ '.⁶⁷ Then $(\underline{\alpha}_k^i, \underline{v}_k), (\bar{\alpha}_k^i, \bar{v}_k) \in \mathcal{V}_k$. Moreover, for any $(\alpha_k^i, v_k) \in \mathcal{V}_k$, v_k is a fixed point of P_k , and $\alpha_k^i(x) = Q(x, L^k v_k(x))$ for all $x \geq 0$. Then, $\underline{v}_k \leq v_l \leq \bar{v}_k$ so that $L^k \underline{v}_k \leq L^k v_k \leq L^k \bar{v}_k$. Since $Q(x, l)$ is increasing in l (as noted above), $\underline{\alpha}_k^i \leq \alpha_k^i \leq \bar{\alpha}_k^i$.

It remains to prove that $\bar{\alpha}_k^i(x) \leq \alpha_k^i(x)$ for all $x \geq 0$. Fix $x \geq 0$ and assume without

67. Indeed, given (33), (32) is equivalent to ' $v_k(x) = b(x) - c(\alpha_k^i(x), x) + n\alpha_k^i(x)[L^k v_k(x) - v_k(x)]$ ', i.e. ' $v_k(x) = \bar{T}(x, \alpha_k^i(x), L^k v_k(x))$ '. Given ' $v_k(x) = \bar{T}(x, \alpha_k^i(x), L^k v_k(x))$ ', (33) is equivalent to '(34) with $\alpha_k^i(x) = Q(x, l)$ and $l = L^k v_k(x)$ ', which is equivalent to ' $\alpha_k^i(x) = Q(x, L^k v_k(x))$ '. Finally, given ' $\alpha_k^i(x) = Q(x, L^k v_k(x))$ ', ' $v_k(x) = \bar{T}(x, \alpha_k^i(x), L^k v_k(x))$ ' is equivalent to ' $v_k(x) = P_kv_k(x)$ '.

loss of generality that $\bar{\alpha}_k^i(x) > 0$ and $\alpha_*^i(x) < 1$. Then, from Step 2 of the proof of Theorem 1, it suffices to show that $\psi^*(\bar{\alpha}_k^i(x), x) > 0$ where ψ^* was defined in (31). Note that

$$\psi^*(\bar{\alpha}_k^i(x), x) > \psi(\bar{\alpha}_k^i(x)|x, L^k \bar{v}_k(x)) \geq 0$$

where the first inequality follows from (35) (as $\bar{v}_k \in \mathcal{V}'$) and the fact that $c_1(\bar{\alpha}_k^i(x), x) > 0$ (as $\bar{\alpha}_k^i(x) > 0$), and the second holds since $\bar{\alpha}_k^i(x) > 0$.

Step 2: \mathcal{V}_k is a singleton. From Step 1, it suffices to show that $(\underline{\alpha}_k^i, \underline{v}_k) = (\bar{\alpha}_k^i, \bar{v}_k)$. By definition of $\underline{\alpha}_k^i$ and $\bar{\alpha}_k^i$, it suffices to show that $\underline{v}_k = \bar{v}_k$. I claim that, for all $x \geq 0$,

$$\bar{v}_k(x) - \underline{v}_k(x) \leq L^k \bar{v}_k(x) - L^k \underline{v}_k(x). \quad (36)$$

Fix x and consider three cases. Suppose first that $\bar{\alpha}_k(x) = 0$. Then, $\underline{\alpha}_k(x) = 0$ since $\bar{\alpha}_k \geq \underline{\alpha}_k$. Then, $\bar{v}_k(x) = b(x) = \underline{v}_k(x)$, so that (36) holds since $\bar{v}_k \geq \underline{v}_k$. Suppose now that $\underline{\alpha}_k(x) = 1$. Then, $\bar{\alpha}_k(x) = 1$ so that, from (32),

$$[L^k \bar{v}_k(x) - \bar{v}_k(x)] - [L^k \underline{v}_k(x) - \underline{v}_k(x)] = \frac{\bar{v}_k(x) - \underline{v}_k(x)}{\lambda n} \geq 0.$$

Then, (36) follows by rearranging. Finally, assume that $\bar{\alpha}_k(x) > 0$ and $\underline{\alpha}_k(x) < 1$. Then, from (33),

$$L^k \bar{v}_k(x) - \bar{v}_k(x) \geq \frac{1}{\lambda} c(\bar{\alpha}_k(x), x) \geq \frac{1}{\lambda} c(\underline{\alpha}_k(x), x) \geq L^k \underline{v}_k(x) - \underline{v}_k(x)$$

where the first inequality holds since $\bar{\alpha}_k(x) > 0$, the second since $\bar{\alpha}_k(x) \geq \underline{\alpha}_k(x)$, and the third since $\underline{\alpha}_k(x) < 1$. Then, (36) follows by rearranging. This proves the claim.

Claim 4. and $\lim_{x \rightarrow \infty} w_*(x) - b(x) = 0$

Claim 4 is proved at the end of this proof. Suppose that $\bar{v}_k(x) - \underline{v}_k(x) > \epsilon$ for some $\epsilon > 0$ and $x \geq 0$, and seek a contradiction. Let

$$\hat{x} := \inf \left\{ x \geq 0 : \sup_{(x, \infty)} \bar{v}_k - \underline{v}_k < \epsilon \right\}$$

and note that $\hat{x} < \infty$ from Claim 4, since $b \leq \underline{v}_k \leq \bar{v}_k \leq w_*$. Consider a sequence $(x_m)_{m \in \mathbb{N}} \subset (0, \hat{x}]$ with $\lim_m x_m = \hat{x}$ and such that $\lim_m \bar{v}_k(x_m) - \underline{v}_k(x_m) \geq \epsilon$. From, (36), it suffices to show that

$$\limsup_{m \rightarrow \infty} L^k \bar{v}_k(x_m) - L^k \underline{v}_k(x_m) < \epsilon. \quad (37)$$

By definition of \hat{x} , $\bar{v}_k(x) - \underline{v}_k(x) < \epsilon$ for all $x > \hat{x}$. Then, if $k = e$, (37) holds since $F(0) = 0$. Hence, assume that $k = d$. Given $\tilde{z} \sim F$, let E_m be the event ' $\bar{v}_d(x_m + \tilde{z}) > \bar{v}_d(x_m)$ ' for all $m \in \mathbb{N}$. If there were a subsequence $(x_{s(m)})_{m \in \mathbb{N}} \subseteq (x_m)_{m \in \mathbb{N}}$ along

which $\lim_m \Pr(E_{s(m)}) = 0$ then, from (32), $\lim_m \bar{v}_d(x_{s(m)}) = b(\hat{x})$. Then, since $\underline{v}_k \geq b$, $\lim_m \bar{v}_d(x_{s(m)}) - \underline{v}_d(x_{s(m)}) = 0$, contradicting the definition of $(x_m)_{m \in \mathbb{N}}$. Hence, by considering an appropriate subsequence, we may assume without loss of generality that $\Pr(E_m) > 0$ for all $m \in \mathbb{N}$, and $\lim_m \Pr(E_m) > 0$.

Fix $m \in \mathbb{N}$ and note that

$$\begin{aligned} L^d \bar{v}_d(x_m) - L^d \underline{v}_d(x_m) &= \mathbb{E}_F[\max\{\bar{v}_d(x_m + \tilde{z}), \bar{v}_d(x_m)\} - \max\{\underline{v}_d(x_m + \tilde{z}), \underline{v}_d(x_m)\}] \\ &\leq \Pr(E_m) \mathbb{E}[\bar{v}_d(x_m + \tilde{z}) - \underline{v}_d(x_m + \tilde{z}) | E_m] \\ &\quad + [1 - \Pr(E_m)] [\bar{v}_d(x_m) - \underline{v}_d(x_m)] \\ \Rightarrow L^d \bar{v}_d(x_m) - L^d \underline{v}_d(x_m) &\leq \mathbb{E}[\bar{v}_d(x_m) - \underline{v}_d(x_m + \tilde{z}) | E_m] \end{aligned}$$

where the last step follows from (36) since $\Pr(E_m) > 0$. Then, taking the limit $m \rightarrow \infty$ yields (37) since $\lim_m \Pr(E_m) > 0$, $F(0) = 0$ and, by definition of \hat{x} , $\sup_{(x, \infty)} \bar{v}_d - \underline{v}_d < \epsilon$ for all $x > \hat{x}$.

Step 3: $v_k - b$ and α_k^i are decreasing. Let \mathcal{V}'' be the set of $v \in \mathcal{V}'$ such that $v - b$ is decreasing. From Steps 1 and 2, it suffices to find a fixed point v of P_k in \mathcal{V}'' such that $Q(x, L^k v(x))$ is decreasing in x .

I claim that $Q(x, L^k v(x))$ is decreasing in x for any $v \in \mathcal{V}''$. To see why, fix $v \in \mathcal{V}''$. From Claim 1, $L^k v(x) - b(x)$ is decreasing. Then, $\psi(a|x, L^k v(x))$ is decreasing in x , since $c_1, c_{12} \geq 0$. Hence $Q(x, L^k v(x))$ is decreasing in x .

In light of the claim, it suffices to show that P_k has a fixed point in \mathcal{V}'' . I claim that P_k maps \mathcal{V}'' to itself. Fix $v \in \mathcal{V}''$. From Step 1, $P_k v \in \mathcal{V}'$. Then, it suffices to show that $P_k v(x_2) - P_k v(x_1) \leq b(x_2) - b(x_1)$ for all $0 \leq x_1 \leq x_2$. Fix x_1 and x_2 , and let $q_i = Q(x_i, L^k v_i(x_i))$ for $i = 1, 2$, so that $q_1 \geq q_2$. Hence

$$P_k v(x_2) - P_k v(x_1) \leq \bar{T}(q_2, x_2, L^k v(x_2)) - \bar{T}(q_2, x_1, L^k v(x_1)) \quad (38)$$

$$\leq \frac{b(x_2) - b(x_1) + \lambda n q_2 [L^k v(x_2) - L^k v(x_1)]}{1 + \lambda n q_2} \quad (39)$$

$$\leq b(x_2) - b(x_1) \quad (40)$$

where (38) follows Claim 3 since $q_2 \leq q_1$, (39) holds since c is increasing in x , and (40) follows from Claim 1. Hence $P_k v \in \mathcal{V}''$, as desired.

Note that \mathcal{V}'' is a complete lattice⁶⁸ and (from Step 1), P_k is increasing on \mathcal{V}'' . Then, from Tarski's fixed-point theorem, P_k admits a fixed point in \mathcal{V}'' , as desired. \square

Proof of Claim 3. Fix x , v , and a , and let $\hat{a} := Q(x, L^k v(x))$. Let $\phi : [0, 1] \rightarrow \mathbb{R}$ be such that

$$\phi(e) := T(e, e + (n-1)\hat{a}, L^k v(x))$$

68. The argument is similar to that of footnote 65.

where T was defined in (28). Note that ϕ is differentiable with

$$\phi'(e) = \lambda \frac{L^k v(x) - [b(x) - c(e, x)] - \left[\frac{1}{\lambda} + e + (n-1)\hat{a}\right] c_1(e, x)}{\{1 + \lambda[e + (n-1)a']\}^2}. \quad (41)$$

Note that ϕ is quasi-concave, and $\phi'(\hat{a})$ has the same sign as $\psi(\hat{a}|x, L^k v(x))$. Then, ϕ is maximised at $e = \hat{a}$. Moreover, $\phi(\hat{a}) = P_k v(x)$ so that $P_k v(x) \geq \phi(a)$.

Then, it suffices to show that $\phi(a) \geq \bar{T}(a, x, L^k v(x))$. Since ϕ is maximised at $e = \hat{a}$ and it is quasi-concave, and $a \leq \hat{a}$, $\phi'(a) \geq 0$. Then, (41) implies that $b(x) - c(a, x) \leq L^k v(x)$, so that

$$\begin{aligned} \phi(a) &= \frac{b(x) - c(a, x)}{1 + \lambda[a + (n-1)\hat{a}]} + \left[1 - \frac{1}{1 + \lambda[a + (n-1)\hat{a}]}\right] L^k v(x) \\ &\geq \frac{b(x) - c(a, x)}{1 + \lambda na} + \left[1 - \frac{1}{1 + \lambda na}\right] L^k v(x) = \bar{T}(a, x, L^k v(x)) \end{aligned}$$

where the inequality holds since $a \leq \hat{a}$. \square

Proof of Claim 4. From Step 1 of the proof of Theorem 2, $w_* - b$ is decreasing. Hence, for any $x \geq 1$, $w_*(x) \leq w_*(1) - b(1) + b(x)$, so that

$$L^e w_*(x) \leq w_*(1) - b(1) + L^e b(x) \leq w_*(1) - b(1) + b(x) + b'(1)\mu$$

where the second inequality holds since b is concave. Hence

$$\begin{aligned} w_*(x) &= \frac{b(x) - c(\alpha_*^i(x), x) + \alpha_*^i(x)\lambda n L^e w_*(x)}{1 + \alpha_*^i(x)\lambda n} \\ &\leq b(x) + \frac{-c(\alpha_*^i(x), x) + \alpha_*^i(x)\lambda n [w_*(1) - b(1) + b'(1)\mu]}{1 + \alpha_*^i(x)\lambda n} \end{aligned}$$

Therefore, since $\lim_{x \rightarrow \infty} \alpha_*^i(x) = 0$ (Step 3 of the proof of Theorem 2),

$$\lim_{x \rightarrow \infty} w_*(x) - b(x) \leq - \lim_{x \rightarrow \infty} c(\alpha_*^i(x), x) \leq 0.$$

Then $\lim_{x \rightarrow \infty} w_*(x) - b(x) = 0$ since, clearly, $w_* \geq b$. \square

E Proofs of Lemma 2 and Proposition 1

Proof of Lemma 2. The ‘only if’ part is immediate, since \tilde{x}_t increases over time. For the ‘if’ part, suppose that v_e is not increasing on $[x_0, \infty)$. Then, there are $x_0 \leq x' < x''$ such that $v_e(x') > v_e(x'')$.

We claim that we may choose x' and x'' such that $\alpha_e(x'') > 0$. Without loss of generality, we may assume that $\alpha_e(x) = 0$ for some $x \geq 0$. Since α_e is continuous, we

may define

$$x_e := \min\{x \geq 0 : \alpha_e(x) = 0\}.$$

Since α_e is decreasing, α_e is zero on $[x_e, \infty)$. Then v_e matches b on $[x_e, \infty)$. In particular, v_e is increasing on $[x_e, \infty)$. Then $x' < x_e$ and we may assume without loss of generality that $x'' \geq x_e$. Then $v_e(x_e) \leq v_e(x'')$. Since v_e is continuous and $v_e(x') > v_e(x'')$, $v_e(x') > v_e(\hat{x}'')$ for $\hat{x}'' \in (x', x_e)$ sufficiently close to x_e . Then, x' and \hat{x}'' satisfy the requirements. This proves the claim.

Since v_e and α_e are continuous, there are neighbourhoods U' and U'' of x' and x'' such that $v_e(y') > v_e(y'')$ and $\alpha_e(y'') > 0$ for all $y' \in U'$ and $y'' \in U''$. Since α_e is decreasing, α_e is strictly positive on $[x_0, \sup U']$. Then, from (1), $\Pr(\exists t \geq 0, \tilde{x}_t^{\alpha_e} \in U' | x_0) > 0$, and $\Pr(\exists T > t, \tilde{x}_t^{\alpha_e} \in U'' | \exists t \geq 0, \tilde{x}_t^{\alpha_e} \in U') > 0$. \square

Proof of Proposition 1. First, I show that $\alpha_e^i(x_0) > a$. Rearranging (16) yields

$$\begin{aligned} g'(1-a)h(x_0) &< \frac{\mathbb{E}_F[h(x_0 + \tilde{z})] - g(1-a)h(x_0)}{1/\lambda + na} \\ &\leq \frac{\mathbb{E}_F[v_e(x_0 + \tilde{z})] - g(1-a)h(x_0)}{1/\lambda + na} = \mathbb{E}[v_e(x_0 + \tilde{z})] - v_e(x_0) \end{aligned}$$

where the second inequality holds since $v_e \geq b$ and $b = h$ (as $g(1) = 1$), and the equality follows from (5). Then, $\alpha_e^i(x_0) > a$ follows from (11), since $c_1(a, x_0) = g'(1-a)h(x_0)$.

From Theorem 2, α_e^i is continuous and $\lim_{x \rightarrow \infty} \alpha_e^i(x) \leq \lim_{x \rightarrow \infty} \alpha_*^i(x) = 0 < a$, where the equality follows from Step 3 of the proof of Theorem 1. Then, by the intermediate value theorem, there is $x_a \in (0, \infty)$ such that $\alpha_e^i(x_a) = a$. It remains to show that v_e is not increasing in a neighbourhood of x_a . Suppose that v_e is increasing in an neighbourhood of x_a , and seek a contradiction. Then, v_e is differentiable a.e. on this neighbourhood.

Since $\alpha_e^i(x_a) \in (0, 1)$, (11) yields the first-order condition

$$\lambda \{\mathbb{E}_F[v_e(x + \tilde{z})] - v_e(x)\} = g'(1 - \alpha_e^i(x))h(x) \quad (42)$$

in a neighbourhood of x_a . Note that, since $v_e - h$ is decreasing (Theorem 2), Claim 1 implies that

$$\mathbb{E}_F[v_e'(x + \tilde{z})] \leq h'(x) \quad (43)$$

in a neighbourhood of x_a . Moreover

$$h'(x) > 0$$

in a neighbourhood of x_a . Indeed, as h' continuous and decreasing, $h'(x_a) = 0$ would imply that h is constant on $[x_a, \infty)$. But then exerting effort at x_a has no benefit, and a strictly positive cost (since $h(x_a) > 0$ as $x_a > 0$, and g is strictly increasing), contradicting

$\alpha_e^i(x_a) > 0$. We consider two cases.

Case 1. g is linear in a neighbourhood of $1 - a$. In this case, since α_e^i is continuous, totally differentiating (42) with respect to x in a neighbourhood of x_a yields

$$v_e'(x) = \mathbb{E}_F[v_e'(x + \tilde{z})] - \frac{g'(1 - \alpha_e^i(x))h'(x)}{\lambda} \leq h'(x) \left(1 - \frac{g'(1 - \alpha_e^i(x))}{\lambda}\right)$$

where the inequality follows from (43). Then, $v_e'(x) < 0$ near x_a since $h'(x) > 0$ and, from (15), $\lambda < g'(1 - a)$.

Case 2. g is not linear in a neighbourhood of $1 - a$. Since α_e^i is decreasing, it is differentiable a.e. Then, since α_e^i is continuous, there exists an x arbitrarily close to x_a such that $g''(1 - \alpha_e^i(x)) < 0$, and α_e^i and v_e are differentiable at x . Totally differentiating (42) with respect to this x yields

$$\begin{aligned} \alpha_e^{i'}(x) &= \frac{\lambda\{\mathbb{E}_F[v_e'(x + \tilde{z})] - v_e'(x)\} - g'(1 - \alpha_e^i(x))h'(x)}{-g''(1 - \alpha_e^i(x))h(x)} \\ &\leq \frac{[\lambda - g'(1 - \alpha_e^i(x))]h'(x)}{-g''(1 - \alpha_e^i(x))h(x)} \end{aligned} \quad (44)$$

where the inequality follows from (43) and the fact that, by hypothesis, $v_e'(x) \geq 0$. Then, by the envelope theorem, differentiating (10) with respect to x yields

$$\begin{aligned} v_e'(x) &= g(1 - \alpha_e^i(x))h'(x) + \lambda n \alpha_e^i(x) \{\mathbb{E}_F[v_e'(x + \tilde{z})] - v_e'(x)\} \\ &\quad + \lambda(n - 1) \alpha_e^{i'}(x) \{\mathbb{E}_F[v_e(x + \tilde{z})] - v_e(x)\} \\ &\leq [g(1 - \alpha_e^i(x)) + \lambda n \alpha_e^i(x)]h'(x) + \lambda(n - 1) \alpha_e^{i'}(x) \{\mathbb{E}_F[v_e(x + \tilde{z})] - v_e(x)\} \\ &= [g(1 - \alpha_e^i(x)) + \lambda n \alpha_e^i(x)]h'(x) + (n - 1) \alpha_e^{i'}(x) g'(1 - \alpha_e^i(x))h(x) \\ &\leq \left\{ g(1 - \alpha_e^i(x)) - \frac{(n - 1)[g'(1 - \alpha_e^i(x))]^2}{-g''(1 - \alpha_e^i(x))} \right. \\ &\quad \left. + \lambda \left[n \alpha_e^i(x) + \frac{(n - 1)g'(1 - \alpha_e^i(x))}{-g''(1 - \alpha_e^i(x))} \right] \right\} h'(x) \end{aligned}$$

where the first inequality follows from (43) and the fact that $v_e'(x) \geq 0$, the second equality from (42), and the second inequality from (44). Then (15) implies that $v_e'(x) < 0$, since $n \geq 2$ and $h'(x) > 0$. \square

F Proof of Theorem 3

I state a ‘principle of optimality’, then prove Theorem 3, relying on Theorem 5.

Lemma 6. *A Markov strategy $\sigma^i := (\alpha^i, \delta^i) \in \Sigma$ induces a Markov perfect equilibrium of the game with disposal if and only if there is $v \in \mathcal{V}'$ such that v solves (20) with $\sigma^j = \sigma^i$*

for all j , α^i solves (21) with $v^i = v$, and δ^i solves (22) with $v^i = v$. Moreover, in this case, the time- t continuation payoff of each agent is $v(x_t)$, assuming no agent obtains an innovation at time t .

Proof of Theorem 3. Let α_d and v_d satisfy the conditions of Theorem 5. To prove existence, let $\delta_d^i \in D$ satisfy (22) with $v^i = v_d$, and let $\sigma_d \in \Sigma^n$ be the symmetric profile induced by $\sigma_d^i := (\alpha_d^i, \delta_d^i)$. Note that v_d solves (20) with $\sigma^j = \sigma^i$ for all j , and α_e^i solves (21) with $v^i = v_e$. Then, by definition of δ_d^i , Lemma 6 implies that σ_d forms a symmetric Markov perfect equilibrium, and $v_d(x_t)$ is the time- t continuation payoff of each agent, assuming no agent obtains an innovation at time t .

To prove that effort α_d^i and payoffs v_d are pinned down in equilibrium, fix a Markov strategy $\sigma^i := (\alpha^i, \delta^i)$ inducing a symmetric Markov perfect equilibrium, and let v be the associated payoff function. Then, from Lemma 6, $v \in \mathcal{V}'$, v solves (20) with $\sigma^j = \sigma^i$ for all j , α^i solves (21) with $v^i = v$, and δ^i solves (22) with $v^i = v$. Then, $(\alpha^i, v) \in \mathcal{V}_d$. Then, from Theorem 5, $\alpha^i = \alpha_d^i$ and $v = v_d$.

It remains to prove that $\alpha_d \geq \alpha_e$ and $v_d \geq v_e$. We begin with $v_d \geq v_e$. From Steps 1 and 2 of the proof of Theorem 5, it suffices to show that the map (P_e, P_d) admits a fixed point in the set \mathcal{W} of pairs $(v, w) \in \mathcal{V}' \times \mathcal{V}'$ such that $v \leq w$. From Step 1, (\mathcal{W}, \geq) is a complete lattice and (P_e, P_d) is increasing. Then, from Tarski's fixed-point theorem, it suffices to show that (P_e, P_d) maps \mathcal{W} to itself.

Fix $(v, w) \in \mathcal{W}$. From Step 1, $(P_e v, P_d w) \in \mathcal{V}' \times \mathcal{V}'$. Moreover, $L^e v \leq L^d w$ since $v \leq w$. Then, $Q(x, L^e v(x)) \leq Q(x, L^d w(x))$, since $Q(x, l)$ is increasing in l (Step 1 of the proof of Theorem 5). Hence, for all $x \geq 0$,

$$\begin{aligned} P_e v(x) &= \bar{T}(Q(x, L^e v(x)), x, L^e v(x)) \leq \bar{T}(Q(x, L^e v(x)), x, L^d w(x)) \\ &\leq \bar{T}(Q(x, L^d w(x)), x, L^d w(x)) = P_d w(x) \end{aligned}$$

where the first inequality holds since $\bar{T}(a, x, l)$ is increasing in l , and the second follows from Claim 3, since $Q(x, L^e v(x)) \leq Q(x, L^d w(x))$. Then, $(P_e v, P_d w) \in \mathcal{W}$, as desired.

It remains to prove that $\alpha_d^i \geq \alpha_e^i$. Note that $L^d v_d \geq L^e v_e$ since $v_d \geq v_e$. Then, result follows since $\alpha_k(x) = Q(x, L^k v_k(x))$ for all $x \geq 0$ and $k \in \{e, d\}$, and $Q(x, l)$ is increasing in l . \square

G Proofs of Lemma 3 and Theorem 4

Proof of Lemma 3. The ‘if’ part is immediate, since concealment has no value if opponents exert no effort. For the ‘only if’ part, fix an initial stock $x_0 < \lambda\mu$. It remains to show that (α_e, ω_0) is not an equilibrium. From (1), there exists $z < \lambda\mu - x_0$ in the support of F .

Suppose that (α_e, ω_0) is played at initial stock $\lambda\mu - z$ and the first innovation, obtained by some agent i at some time t , has size z . Then, agent i discloses fully at time t , and no agent exerts any effort thereafter. This raises the public stock to $\lambda\mu$, so that the time- t continuation payoff (to each agent) is $\lambda\mu$. Because $\lambda\mu$ solves (12), after time t , agent i would also earn $\lambda\mu$ if she exerted maximal effort until she obtained a further innovation, disclosed it fully, and exerted no effort thereafter. Consider the time- t deviation for agent i in which she does precisely this, *but* conceals fully at time t . Conditional on the opponents of agent i obtaining no innovation after time t , agent i 's continuation payoff is again $\lambda\mu$. This is because exerting full effort yields no flow of payoff (i.e. $u(0, x_t) \equiv 0$), so the fact that the public stock x_t equals $\lambda\mu - z$ instead of $\lambda\mu$ after time t has no effect. Because $z > 0$, Corollary 3 implies that $\alpha_e^i(\lambda\mu - z) > 0$. Then, the above deviation is (strictly) profitable, as the opponents of agent i will obtain an innovation before her with strictly positive probability.

By continuity, there are open neighbourhoods X of $\lambda\mu - z$ and Z of z such that, if (α_e, ω_0) is played at x_0 , and an agent obtains an innovation of size within Z at some time t such that $x_t \in X$, agent i has a profitable deviation at time t . From (1), this occurs with strictly positive probability, so that (α_e, ω_0) is not an equilibrium. \square

It remains to prove Theorem 4. I begin with several definitions. Let \mathcal{Q} be the set of $q : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $q(t) \leq \liminf_{s \downarrow t} q(s)$ for all t , $Q_0 := [\lambda\mu, \lambda\mu n]$, and \mathcal{Q}_0 be the set of lower-semicontinuous $q \in \mathcal{Q}$ with image in Q_0 , such that $q(0) = \liminf_{t \downarrow 0} q(t)$.

Let \mathcal{A} be the quotient space of Lebesgue measurable $\alpha : \mathbb{R}_+^2 \rightarrow [0, 1]$ such that $\alpha_t : \mathbb{R}_+ \rightarrow [0, 1]$ is increasing for all $t \geq 0$, with respect to the equivalence relation $\sim_{\mathcal{A}}$, defined by $\alpha \sim_{\mathcal{A}} \alpha'$ if and only if $\int_0^t \alpha_s(k) ds = \int_0^t \alpha'_s(k) ds$ for each $t, k \geq 0$. Given $\alpha \in \mathcal{A}$ and $q \in \mathcal{Q}$, let $R(\alpha, q)$ be the set of symmetric strategy profiles ρ such that, on path, no effort is exerted after a disclosure occurs and, for all i , in any period t prior to which no disclosure occurred, agent i exerts effort $\alpha_t(k_t^i)$, conceals fully if either $k_t^i = x_0$ or $k_t^i < q(t)$, and discloses fully otherwise.

To prove Theorem 4, it suffices to exhibit $\alpha \in \mathcal{A}$ and $q \in \mathcal{Q}_0$ such that some $\rho \in R(\alpha, q)$ forms a PBE. This section is devoted to the following intermediary result.

Proposition 5. *There are $\alpha \in \mathcal{A}$ and $q \in \mathcal{Q}_0$ such that any $\rho \in R(\alpha, q)$ is a Bayes-Nash equilibrium.*

I complete the proof of Theorem 4 in Curello (2021), by constructing a specific $\rho \in R(\alpha, q)$, as well as a system of beliefs, and showing that they form a PBE. The proof of Proposition 5 is in Appendix G.2. It relies on a principle of optimality (Lemma 7 below), and on an ‘approximate’ solution to the Bellman equation underlying this principle, which I construct in Appendix G.1.

I begin with several definitions. For all $x \geq 0$, let $\underline{v}(x)$ be the value of the single-agent

game with initial stock x .⁶⁹ Let V be the space of 1-Lipschitz continuous, increasing and convex maps $v : \mathbb{R}_+ \rightarrow \mathbb{R}$, and let V_0 be the set of $v \in V$ such that $v - \underline{v}$ is decreasing and $v = \underline{v}$ on $[\lambda\mu n, \infty)$. Let \mathcal{V}_0 be the set of maps $v : \mathbb{R}_+ \rightarrow V_0$ that are continuous in the supremum metric. Let E be the quotient space of Lebesgue measurable $a : \mathbb{R}_+ \rightarrow [0, 1]$ with respect to the equivalence relation \sim_E , defined by $a_1 \sim_E a_2$ if and only if $\int_0^t a_1 = \int_0^t a_2$ for all $t \geq 0$.

Given $\delta \in [0, \infty]$ and $a \in E$, let $H(a, \delta)$ be the distribution of the random time π of the first increment obtained by an agent who exerts effort a_s at any $s < \delta$, and no effort after δ (with ' $\pi = \infty$ ' if no increment arises).⁷⁰ Given $\alpha \in \mathcal{A}$ and $q \in \mathcal{Q}$, suppose that the opponents of agent i play some $\rho \in R(\alpha, q)$. Note that, if no disclosure occurred by some time t , assuming that agent i does not disclose after time t , the joint distribution of (τ, κ) , where $t + \tau$ is the time of the first disclosure by any opponent of agent i after time t , and κ is the value disclosed (with ' $\tau = \infty$ ' and $\kappa = 0$ if no opponent ever discloses), is pinned down by t , α , and q . Label it $J(t, \alpha, q)$.

Lemma 7. *Let $\alpha \in \mathcal{A}$, $q \in \mathcal{Q}_0$, and $v \in \mathcal{V}_0$ satisfy, for each $t \geq 0$ and $k \geq x_0$,*

$$v_t(k) := \sup_{\substack{\delta \in [0, \infty] \\ a \in E}} \mathbb{E} \left[x_0 \int_0^{\tilde{T}} e^{-s} (1 - a_s) ds + e^{-\tilde{T}} (\mathbb{1}_{\tilde{\tau}=\tilde{T}} \tilde{\kappa} + \mathbb{1}_{\tilde{\pi}=\tilde{T}} v_{t+\tilde{T}}(k + \tilde{z}) + \mathbb{1}_{\tilde{T}=\delta} \underline{v}(k)) \right] \quad (45)$$

where $(\tilde{z}, (\tilde{\tau}, \tilde{\kappa}), \tilde{\pi}) \sim F \times J(t, \alpha, q) \times H(a, \delta)$ and $\tilde{T} := \tilde{\pi} \wedge \tilde{\tau} \wedge \delta$. Suppose that the objective in (45) is maximised by

$$\delta := \min \{s \geq 0 : k \geq q(t + s)\} \cup \{\infty\} \quad \& \quad a_s := \alpha_{t+s}(k). \quad (46)$$

Then, any $\rho \in R(\alpha, q)$ is a Bayes-Nash equilibrium and, if ρ is played, at any time t prior to the first disclosure, agent i 's continuation payoff is $v_t(k_t^i)$.

The Bellman equation (45) may be understood as follows. At any time t prior to the first disclosure, an agent i with private stock $k_t^i = k$ picks a delay δ after which, if she obtained no increment and none of her opponents disclosed, she discloses fully (where $\delta = \infty$ if agent i does not disclose in this situation), and a schedule a describing her effort until either she obtains an increment, or an opponent discloses, or time $t + \delta$ is reached.⁷¹ Agent i 's value after she discloses fully is $\underline{v}(k)$ as, following such a disclosure, her opponents do not disclose. Moreover, given a realisation (τ, κ) of $(\tilde{\tau}, \tilde{\kappa})$, agent i 's

69. That is, $\underline{v}(x)$ satisfies (9) with $n = 1$.

70. That is, $[H(a, \delta)](\pi) := 1 - e^{-\lambda \int_0^{\pi \wedge \delta} a_s ds}$ for all $\pi \in \mathbb{R}_+$.

71. In (45), we neglect the constraint that agent i cannot disclose prior to obtaining the first increment. This is without loss of generality as any $q \in \mathcal{Q}_0$ is bounded below by $\lambda\mu > x_0$ so that, from (46), $\delta = \infty$ if $k = x_0$.

value after an opponent discloses κ is precisely κ .⁷²

I end this section with a restatement of Proposition 5 (Remark 1 below), in a language that allows to formulate a discrete-time approximation of (45) for any time length $\Delta > 0$. I require several definitions. Given $\Delta > 0$, let $T_\Delta := \{0, \Delta, 2\Delta, \dots\}$, $T'_\Delta := T_\Delta \cup \{\infty\}$, and E_Δ be the set of $a \in E$ that are constant over $[(m-1)\Delta, m\Delta)$ for all $m \in \mathbb{N}$. Also let \mathcal{V} be the set of Lebesgue measurable $v : \mathbb{R}_+^2 \rightarrow \mathbb{R}$ such that $v_t \in V$ for all $t \geq 0$, and

$$Y' := [0, \infty] \times E \quad \& \quad \Theta_0 := \mathbb{R}_+^3 \times \mathcal{A} \times \mathcal{Q} \times \mathcal{V}$$

with typical elements $y := (\delta, a)$ and $\theta := (\Delta, t, k, \alpha, q, v)$, respectively. Let $I : Y' \times \Theta_0 \rightarrow \mathbb{R}$ be given by

$$\begin{aligned} I(y, \theta) := & \mathbb{E} \left\{ x_0 \int_0^{\tilde{T}} e^{-s} (1 - a_s) ds + e^{-\tilde{T}} [\mathbb{1}_{\tilde{\tau} > \tilde{\pi} \leq \delta} v_{t+\tilde{T}}(k + \tilde{z}) \right. \\ & \left. + (1 - \mathbb{1}_{\tilde{\tau} > \tilde{\pi} \leq \delta}) \underline{v}_\Delta((k + \mathbb{1}_{\tilde{\pi} = \tilde{T}} \tilde{z}) \vee \mathbb{1}_{\tilde{\tau} = \tilde{T}} \tilde{\kappa})] \right\} \end{aligned}$$

where $y := (\delta, a)$, $\theta := (\Delta, t, k, \alpha, q, v)$, $\tilde{T} := \tilde{\tau} \wedge \tilde{\pi} \wedge \delta$,

$$(\tilde{\tau}, \tilde{\pi}) := \begin{cases} (\Delta \lceil \tilde{\tau}_0 / \Delta \rceil, \Delta \lceil \tilde{\pi}_0 / \Delta \rceil) & \text{if } \Delta > 0 \\ (\tilde{\tau}_0, \tilde{\pi}_0) & \text{if } \Delta = 0, \end{cases} \quad (47)$$

and $(\tilde{z}, (\tilde{\tau}_0, \tilde{\kappa}), \tilde{\pi}_0) \sim F \times J(t, \alpha, q) \times H(a, \delta)$. Define the correspondence $C : \Theta_0 \rightarrow Y'$ by

$$C(\Delta, t, k, \alpha, q, v) := \begin{cases} T'_\Delta \times E_\Delta & \text{if } \Delta > 0 \\ Y' & \text{if } \Delta = 0, \end{cases}$$

and let $I^* : \Theta_0 \rightarrow \mathbb{R}$ be given by

$$I^*(\theta) := \sup_{y \in C(\theta)} I(y, \theta).$$

Remark 1. In light of Lemma 7, Proposition 5 is equivalent to: ‘There are $\alpha \in \mathcal{A}$, $q \in \mathcal{Q}_0$, and $v \in \mathcal{V}$ such that, for all $t \geq 0$ and $k \geq x_0$,

$$v_t(k) = I^*(\theta) = I((\delta, a), \theta) \quad \text{where } \delta \text{ and } a \text{ are given by (46),} \quad (48)$$

holds with $\theta := (0, t, k, \alpha, q, v)$.’

72. To see why, note that $\kappa \geq q(\tau) \geq \lambda\mu$ where the first inequality holds since $\rho \in R(\alpha, q)$ is played and the second as $q \in \mathcal{Q}_0$. Then, the result holds as long as $k \leq \kappa$. Thus it suffices to show that it is optimal for agent i to disclose (fully) after obtaining an increment (at time t , say) that raises her private stock to a value $k \geq q(t)$. To this end, we may assume without loss that $k = q(t)$. Then result follows since, from (46), the objective in (45) is maximised by $\delta = 0$.

G.1 Approximate Bellman equation

In this section, I define a discrete-time, finite-horizon approximation of the Bellman equation (45) and show that it admits a solution for any time length $\Delta > 0$ and horizon $M \in \mathbb{N}$ (Lemma 9 below). As part of the analysis, I establish bounds for the disclosure cutoff q that is part of the solution, and show that they converge to the desired values as the approximation becomes more accurate (Lemma 8 below).

I begin with several definitions. Let A be the set of increasing $\alpha : \mathbb{R}_+ \rightarrow [0, 1]$. Given $m \in \mathbb{N}$, let

$$Y := \{0, 1\} \times [0, 1] \quad \& \quad \Theta_m := \mathbb{R}_+ \times A^m \times \mathbb{R}_+^m \times V$$

with typical elements $y := (\delta, a)$ and $\theta := (k, \alpha, q, v)$, respectively. Given $a \in [0, 1]$ and $\Delta > 0$, define the CDF $F^{\Delta, a}$ with support within \mathbb{R}_+ by

$$F^{\Delta, a}(\omega) := (1 - e^{-a\Delta})F(\omega) + e^{-a\Delta} \quad \text{for all } \omega \in \mathbb{R}_+.$$

Let $\underline{v}_\Delta : \mathbb{R}_+ \rightarrow \mathbb{R}$ be the unique element of V solving the Bellman equation

$$\underline{v}_\Delta(x) = \max_{a \in [0, 1]} x(1 - e^{-\Delta})(1 - a) + e^{-\Delta} \mathbb{E}_{F^{\Delta, a}}[\underline{v}_\Delta(x + \tilde{\omega})].^{73} \quad (49)$$

Let $I_{\Delta, m} : Y \times \Theta_m \rightarrow \mathbb{R}$ be given by

$$I_{\Delta, m}(y, \theta) = \begin{cases} \underline{v}_\Delta(k) & \text{if } \delta = 0 \\ x_0(1 - e^{-\Delta})(1 - a) \\ \quad + e^{-\Delta} \mathbb{E}[\mathbb{1}_{\tilde{\tau} \leq \Delta} \underline{v}_\Delta(\tilde{\kappa} \vee (k + \tilde{\omega})) + \mathbb{1}_{\tilde{\tau} > \Delta} v(k + \tilde{\omega})] & \text{if } \delta = 1 \end{cases}$$

where $y := (\delta, a)$, $\theta := (k, \alpha, q, v)$, $(\tilde{\omega}, (\tilde{\tau}, \tilde{\kappa})) \sim F^{\Delta, a} \times J((m-1)\Delta, \hat{\alpha}, \hat{q})$, and $\hat{\alpha} \in \mathcal{A}$ and $\hat{q} \in \mathcal{Q}$ are such that $\hat{\alpha}_t = \alpha_l$ and $\hat{q}_t = q_l$ for all $l \in \{1, \dots, m\}$ and $t \in [(l-1)\Delta, l\Delta]$. Also let $I_{\Delta, m}^* : \Theta_m \rightarrow \mathbb{R}$ be given by

$$I_{\Delta, m}^*(\theta) := \max_{y \in Y} I_{\Delta, m}(y, \theta).$$

Next, I define bounds q_Δ^+ and q_Δ^- for the disclosure cutoff q for each $\Delta > 0$, and show that they converge to the desired values as Δ vanishes (Lemma 8 below). Given $\Delta > 0$,

⁷³ We may view \underline{v}_Δ as a discrete-time approximation of the map \underline{v} for time length Δ . In particular, $\underline{v}_\Delta \rightarrow \underline{v}$ uniformly as $\Delta \rightarrow 0$.

let q_Δ^- be the largest $x \geq 0$ such that the objective in (49) is maximised by $a = 1$,⁷⁴ and

$$q_\Delta^+ := \sup \left\{ k \in \mathbb{R}_+ : \arg \max_{a \in [0,1]} I_{\Delta,m}((1,a), (k, \alpha, q, v)) \geq k \geq \max q \text{ for some } m \in \mathbb{N}, \right. \\ \left. \alpha \in A^m, q \in \mathbb{R}_+^m \text{ and } v \in V, \text{ such that } v(l) = l \text{ for all } l \geq k \right\}.$$

Lemma 8. *For all $\Delta > 0$, $q_\Delta^- \leq q_\Delta^+ < \infty$. Moreover*

$$\liminf_{\Delta \rightarrow 0} q_\Delta^- \geq \lambda\mu \quad \& \quad \limsup_{\Delta \rightarrow 0} q_\Delta^+ \leq n\lambda\mu.$$

Lemma 8 is proved in Curello (2021). Given $\Delta > 0$, let $Q_\Delta := [q_\Delta^-, q_\Delta^+]$, and V_Δ be the set of $v \in V$ such that $v - \underline{v}_\Delta$ is decreasing and $v(k) = \underline{v}_\Delta(k)$ for $k \geq q_\Delta^+$. Let \mathcal{Q}_Δ be the set of $q \in \mathcal{Q}$ that have image in Q_Δ and are constant on $[(m-1)\Delta, m\Delta)$ for all $m \in \mathbb{N}$. Similarly, let \mathcal{A}_Δ (\mathcal{V}_Δ) be the set of $\alpha \in \mathcal{A}$ ($v \in \mathcal{V}$) such that, for all $k \geq 0$, $t \mapsto \alpha_t(k)$ ($t \mapsto v_t(k)$) is constant over $[(m-1)\Delta, m\Delta)$ for all $m \in \mathbb{N}$.

Remark 2. Let $\Delta > 0$ and $(\alpha_m, q_m, v_m)_{m \in \mathbb{N}} \subset A \times Q_\Delta \times V_\Delta$ be such that, for all $m \in \mathbb{N}$ and $k \geq x_0$,

$$v_m(k) = I_{\Delta,m}^*(k, \theta_m) = I_{\Delta,m}((\mathbf{1}_{k < q_m}, \alpha_m), \theta_m) \quad (50)$$

where $\theta_m := (k, (\alpha_l)_{l=1}^m, (q_l)_{l=1}^m, v_{m+1})$. Define $(\hat{\alpha}, \hat{q}, \hat{v}) \in \mathcal{A}_\Delta \times \mathcal{Q}_\Delta \times \mathcal{V}_\Delta$ by

$$(\hat{\alpha}_t, \hat{q}_t, \hat{v}_t) = (\alpha_m, q_m, v_m) \quad \text{for all } m \in \mathbb{N} \text{ and } t \in [(m-1)\Delta, m\Delta). \quad (51)$$

Then \hat{v}_t satisfies (48) with $\theta := (\Delta, t, k, \hat{\alpha}, \hat{q}, \hat{v})$ for any $t \in T_\Delta$ and $k \geq x_0$.

Remark 2 is proved in Curello (2021). Given $m \in \mathbb{N}$, $\alpha \in A^m$, $q \in \mathbb{R}_+^m$, let

$$\beta^{\Delta,m}(\alpha, q) := \beta_{m\Delta}(\hat{\alpha}, \hat{q}) \quad (52)$$

where $\hat{\alpha} \in \mathcal{A}$ and $q \in \mathcal{Q}$ are given by $\hat{\alpha}_t = \alpha_l$ and $\hat{q}_t = q_l$ for all $l \in \{1, \dots, m\}$ and $t \in [(l-1)\Delta, l\Delta)$.⁷⁵

Lemma 9. *There are $\Delta_0 > 0$ and $\gamma : (0, \infty) \rightarrow (0, \infty)$ such that, for any $\Delta \in (0, \Delta_0)$ and $M \in \mathbb{N}$, there are $\alpha \in A^M$, $q \in Q_\Delta^M$, and $v \in V_\Delta^{M+1}$ such that, given any $m = 1, \dots, M$, (50) holds for all $k \geq x_0$ and, moreover,*

$$\sup |v_{m+1} - v_m| \vee |G_m - G_{m-1}| \leq \gamma(m\Delta)\Delta, \quad (53)$$

where $G_m := \beta^{\Delta,m}((\alpha_l)_{l=1}^m, (q_l)_{l=1}^m)$ for each $m = 1, \dots, M$, and G_0 is degenerate at x_0 .

74. The objective in (49) is maximised by $a = 1$ for $x = 0$, so that q_Δ^- is well-defined. Moreover, it is easy to show that $a = 0$ is the unique maximiser of (49) for $x > \Delta\lambda\mu/(e^\Delta - 1)$, so that q_Δ^- is finite.

75. The map $\beta^{\Delta,m}$ is well-defined since $\beta_t(\alpha, q) = \beta_t(\alpha', q')$ for all $t > 0$, $\alpha, \alpha' \in \mathcal{A}$, and $q, q' \in \mathcal{Q}$ such that $\alpha = \alpha'$ and $q = q'$ on $[0, t)$.

Lemma 9 is proved in Appendix G.3.

G.2 Proof of Proposition 5

In this section, I show that the approximate solution to the Bellman equation constructed in Lemma 9 lies in a suitably compact space (Remarks 3 to 5), and that the functional underlying the Bellman equation is continuous in the limit as the approximation becomes more accurate (Lemma 10). I end with the proof of Proposition 5.

Endow \mathcal{A} with the topology such that $\alpha^m \rightarrow \alpha$ if and only if $\int_0^t \alpha_s^m(k) ds \rightarrow \int_0^t \alpha_s(k) ds$ for all t and k .

Remark 3. \mathcal{A} is sequentially compact.⁷⁶

Endow \mathcal{Q} with the topology such that $q^m \rightarrow q$ if and only if $\inf_{[t', t'']} q^m \rightarrow \inf_{[t', t'']} q$ for all but countably many $0 < t' < t'' < \infty$.

Remark 4. For any $(\Delta^m)_{m \in \mathbb{N}} \rightarrow 0$ and $(q^m)_{m \in \mathbb{N}} \in \prod_{m \in \mathbb{N}} \mathcal{Q}_{\Delta^m}$, there is $q \in \mathcal{Q}_0$ such that $q^m \rightarrow q$ along a subsequence.

Remark 4 is proved in Curello (2021).

Endow V with the supremum metric, and \mathcal{V} with the topology such that $v^m \rightarrow v$ if and only if v^m converges uniformly to v on $[0, T]$ for all $T > 0$.

Remark 5. Let $(\Delta^m)_{m \in \mathbb{N}} \rightarrow 0$ and $(v^m)_{m \in \mathbb{N}} \in \prod_{m \in \mathbb{N}} \mathcal{V}_{\Delta^m}$ be such that, for all $T > 0$, there is $\gamma > 0$ such that, for all $m \in \mathbb{N}$, v^m is γ -Lipschitz on $T_{\Delta^m} \cap [0, T]$ in the supremum metric for all $m \in \mathbb{N}$, there is $v \in \mathcal{V}_0$ such that $v^m \rightarrow v$ along a subsequence.⁷⁷

Let

$$\Theta := \left(\bigcup_{\Delta > 0} \{\Delta\} \times T_{\Delta} \times \mathbb{R}_+ \times \mathcal{A}_{\Delta} \times \mathcal{Q}_{\Delta} \times \mathcal{V}_{\Delta} \right) \cup (\{0\} \times \mathbb{R}_+^2 \times \mathcal{A} \times \mathcal{Q}_0 \times \mathcal{V}_0).$$

and note that $\Theta \subset \Theta_0$. Endow Θ_0 with the product topology and Θ with the topology such that $\theta^m := (\Delta^m, t^m, k^m, \alpha^m, q^m, v^m)$ converges to $\theta := (\Delta, t, k, \alpha, q, v)$ if and only if either $\theta_m = \theta$ for m large enough, or $\Delta = 0$, $\theta_m \rightarrow \theta$ in Θ_0 , and $\inf_{[t, t']} q^m \rightarrow \inf_{[t, t']} q$ for all but countably many $t' > t$. Endow Y' with the product topology, and E with the topology such that $a \rightarrow a'$ if and only if $\int_0^t a \rightarrow \int_0^t a'$ for all $t \geq 0$.

Lemma 10. *Let $(\theta_m)_{m \in \mathbb{N}} \subset \Theta$ converge to some $\theta \in \Theta$. Suppose that, for all $T > 0$, there is $\gamma > 0$ such that $t \mapsto \beta_t(\alpha^m, q^m)$ and v^m are γ -Lipschitz on $[0, T] \cap T_{\Delta}$ in the supremum metric, where $\theta_m := (\Delta^m, t^m, k_m, \alpha^m, q^m, v^m)$. Then $I^*(\theta_m) \rightarrow I^*(\theta)$ and, for any $(y_m)_{m \in \mathbb{N}} \subset Y'$ converging to some $y \in Y'$, $I(y_m, \theta_m) \rightarrow I(y, \theta)$.*

76. This follows from (a multi-dimensional version of) Helly's selection theorem (for example, Theorem 2 in Brunk, Ewing, and Utz (1956)), together with a diagonalisation argument. This is because, for any $\alpha \in \mathcal{A}$, the map $(t, k) \mapsto \int_0^t \alpha_s(k) ds$ is increasing and bounded by T on $[0, T] \times \mathbb{R}_+$, for each $T > 0$.

77. This is immediate for Helly's selection theorem for arbitrary metric spaces.

Lemma 10 is proved in Curello (2021).

Proof of Proposition 5. To construct α , q and v , fix γ and Δ_0 satisfying the conditions of Lemma 9.

Claim 5. For all $\Delta \in (0, \Delta_0)$, there are $\hat{\alpha} \in \mathcal{A}_\Delta$, $\hat{q} \in \mathcal{Q}_\Delta$, and $\hat{v} \in \mathcal{V}_\Delta$ such that

- (a) (48) holds with $\theta := (\Delta, t, k, \hat{\alpha}, \hat{q}, \hat{v})$ for all $t \in T_\Delta$ and $k \geq x_0$, and
- (b) $s \mapsto \beta_s(\alpha, q)$ and v are $\gamma(T)$ -Lipschitz on $[0, t] \cap T_\Delta$ in the supremum metric, for all $T > 0$.

Proof. Fix Δ . For all $M \in \mathbb{N}$, let $\alpha^M \in A^M$, $q^M \in Q_\Delta^M$, and $v^M \in V_\Delta^{M+1}$ satisfy the conditions of Lemma 9, and label $\rho^M := (\alpha^M, q^M, v^M)$. Note that A , Q_Δ , and V_Δ are compact.⁷⁸ Then, there are strictly increasing sequences $\mathbb{N} \supseteq (s_{0l})_{l \in \mathbb{N}} \supseteq (s_{1l})_{l \in \mathbb{N}} \supseteq (s_{2l})_{l \in \mathbb{N}}, \dots$ such that $(\rho_1^{s_{ml}}, \dots, \rho_m^{s_{ml}})_{l \in \mathbb{N}}$ converges in $(A \times Q_\Delta \times V_\Delta)^m$ for all $m \in \mathbb{N}$. In particular, $(\rho_m^{s_{ml}})_{l \in \mathbb{N}}$ converges in $A \times Q_\Delta \times V_\Delta$ for all $m \in \mathbb{N}$.

Let $(\alpha_m, q_m, v_m)_{m \in \mathbb{N}} \subset A \times Q_\Delta \times V_\Delta$ be given by

$$(\alpha_m, q_m, v_m) := \lim_l \rho_m^{s_{ml}} \quad \text{for all } m \in \mathbb{N},$$

and note that, for all $m \in \mathbb{N}$, (50) holds since $I_{\Delta, m}^*$ and $I_{\Delta, m}$ are continuous,⁷⁹ and (53) holds since β_m^Δ and β_{m+1}^Δ are continuous.

Let $\hat{\alpha} \in \mathcal{A}_\Delta$, $\hat{q} \in \mathcal{Q}_\Delta$ and $\hat{v} \in \mathcal{V}_\Delta$ be given by (51). It remains to show (a) and (b). The former is immediate from Remark 2, while the latter holds as (53) holds for all $m \in \mathbb{N}$. \square

Fix a sequence $(\Delta^m)_{m \in \mathbb{N}} \subset (0, \Delta_0)$ that converges to 0. For all $m \in \mathbb{N}$, let $\alpha^m \in \mathcal{A}_{\Delta^m}$, $q^m \in \mathcal{Q}_{\Delta^m}$ and $v^m \in \mathcal{V}_{\Delta^m}$ satisfy the conditions of Claim 5 with $\Delta := \Delta^m$. From Remark 3, by considering a subsequence of $(\Delta^m)_{m \in \mathbb{N}}$, we may assume without loss of generality that $\alpha^m \rightarrow \alpha$ in \mathcal{A} for some $\alpha \in \mathcal{A}$. Moreover, from Remark 4, by considering a further subsequence we may also assume that, $q^m \rightarrow q$ in \mathcal{Q} for some $q \in \mathcal{Q}_0$. Finally, for all $m \in \mathbb{N}$ and $T > 0$, the variation of v^m on $[0, T]$ is bounded by $\gamma(T + \Delta_0)$, as v^m is $\gamma(T)$ -Lipschitz on $[0, T] \cap T_{\Delta^m}$ (from (b)). Then, from Remark 5, by considering

78. This follows from Helly's selection theorem for A , and from the Arzela-Ascoli theorem for V_Δ .

79. In particular, writing $\bar{\alpha}^m := (\alpha_i)_{i=1}^m$, $\bar{q}^m := (q_i)_{i=1}^m$, $\bar{\alpha}_l^m := (\alpha_i^{s_{il}})_{i=1}^m$ and $\bar{q}_l^m := (q_i^{s_{il}})_{i=1}^m$,

$$I_{\Delta, m}((\mathbb{1}_{k < q_m}, \alpha_m), (k, \bar{\alpha}^m, \bar{q}^m, v_{m+1})) = \lim_l I_{\Delta, m}((\mathbb{1}_{k < q_m^{s_{ml}}}, \alpha_m^{s_{ml}}), (k, \bar{\alpha}_l^m, \bar{q}_l^m, v_{m+1}^{s_{(m+1)l}}))$$

even if $(\mathbb{1}_{k < q_m^{s_{ml}}})_{l \in \mathbb{N}}$ does not converge to $\mathbb{1}_{k < q_m}$, since in this case $k = q_m$ and

$$\begin{aligned} I_{\Delta, m}((\mathbb{1}_{k < q_m}, \alpha_m), (k, \bar{\alpha}^m, \bar{q}^m, v_{m+1})) &= \lim_{k' \downarrow k} I_{\Delta, m}((\mathbb{1}_{k' < q_m}, \alpha_m), (k', \bar{\alpha}^m, \bar{q}^m, v_{m+1})) = I_{\Delta, m}^*(k, \bar{\alpha}^m, \bar{q}^m, v_{m+1}) \\ &= \lim_l I_{\Delta, m}^*(k, \bar{\alpha}_l^m, \bar{q}_l^m, v_{m+1}^{s_{(m+1)l}}) = \lim_l I_{\Delta, m}((\mathbb{1}_{k < q_m^{s_{ml}}}, \alpha_m^{s_{ml}}), (k, \bar{\alpha}_l^m, \bar{q}_l^m, v_{m+1}^{s_{(m+1)l}})). \end{aligned}$$

a further subsequence, we may assume without loss of generality that $v^m \rightarrow v$ in \mathcal{V} for some $v \in \mathcal{V}_0$.

From Remark 1, it remains to prove that (48) holds with $\theta := (0, t, k, \alpha, q, v)$ for all $t \geq 0$ and $k \geq x_0$. Fix t and k and, let a and δ be given by (46). We consider two cases.

Case 1. $\inf_{[t, t']} q^m \rightarrow \inf_{[t, t']} q$ for all but countably $t' > t$. In this case, there exist sequences $(t^m)_{m \in \mathbb{N}} \subset \mathbb{R}_+$ and $(k_m)_{m \in \mathbb{N}} \subset [x_0, \infty)$ such that $t^m \rightarrow t$, $k_m \rightarrow k$, $t^m \in T_{\Delta^m}$ for all $m \in \mathbb{N}$, and $\delta_m \rightarrow \delta$, where

$$\delta_m := \min\{s \geq 0 : k_m \geq q^m(t^m + s)\} \cup \{\infty\} \quad \text{for all } m \in \mathbb{N}.^{80}$$

Let $a^m : s \mapsto \alpha_{t^m+s}^m(k_m)$ and $\theta_m := (\Delta^m, t^m, k_m, \alpha^m, q^m, v^m)$ for all $m \in \mathbb{N}$. Then, from Claim 5,

$$v_{t^m}^m(k_m) = I^*(\theta_m) = I((\delta_m, a^m), \theta_m) \quad \text{for all } m \in \mathbb{N}. \quad (54)$$

Note that $v_{t^m}^m(k) \rightarrow v_t(k)$. Moreover, $a^m \rightarrow a$ in E ,⁸¹ so that $((a^m, \delta_m), \theta_m) \rightarrow ((a, \delta), \theta)$ in $Y' \times \Theta$, by the hypothesis of this case. Then, $v_t(k) = I^*(\theta) = I((\delta, a), \theta)$ from (b) and Lemma 10.

Case 2. General case. We claim that there is a sequence $(t^m)_{m \in \mathbb{N}} \rightarrow t$ such that the hypothesis Case 1 is satisfied for any t in the sequence, and such that $\inf_{[t^m, t']} q \rightarrow \inf_{[t, t']} q$ for all $t' > t$. Since $q^m \rightarrow q$ in \mathcal{Q} , the hypothesis of Case 1 is satisfied for all but countably many t . Moreover, if $t = 0$ then $\inf_{[t^m, t']} q \rightarrow \inf_{[t, t']} q$ for all $t' > t$ for any $(t^m)_{m \in \mathbb{N}} \rightarrow t$ since $q(0) = \liminf_{s \downarrow 0} q(s)$, and if $t > 0$ then $\inf_{[t^m, t']} q \rightarrow \inf_{[t, t']} q$ for all $t' > t$ as long as $(t^m)_{m \in \mathbb{N}}$ is bounded above by t , as q is lower-semicontinuous (since $q \in \mathcal{Q}_0$). The claim follows.

In light of Case 1, $v_{t^m}(k) = I^*(\theta_m) = I((\delta_m, a^m), \theta_m)$ for all $m \in \mathbb{N}$, where $\theta_m :=$

80. If $\delta = 0$, take $t^m := \Delta^m \lfloor t/\Delta^m \rfloor$. Note that $\inf_{[t^l, t^l+2/m)} q^l \leq \inf_{[t, t+1/m)} q^l$ for any $l, m \in \mathbb{N}$ such that $\Delta^l < \frac{1}{m}$. Moreover, $\inf_{[t, t+1/m)} q \leq k$ as $\delta = 0$. Then $\inf_{[t^l, t^l+2/m)} q^l < k + \frac{1}{m}$ for l sufficiently large, as $\lim_l \inf_{[t, t+1/m)} q^l = \inf_{[t, t+1/m)} q$. Choose $(l_m)_{m \in \mathbb{N}} \subseteq \mathbb{N}$ to be strictly increasing and such that $\Delta^{l_m} < \frac{1}{m}$ and $\inf_{[t^{l_m}, t^{l_m}+2/m)} q^{l_m} < k + \frac{1}{m}$ for all $m \in \mathbb{N}$. Define $(k_l)_{l \in \mathbb{N}}$ by $k_l := k + \frac{1}{m}$ for all $m \in \mathbb{N}$ and $l = l_{m-1} + 1, \dots, l_m$, (where $l_0 := 0$). Then, $\lim_l k_l = k$ and $\lim_l \delta_l = \delta$ as $\delta_{l_m} \leq \frac{2}{m}$ for all $m \in \mathbb{N}$.

If $\delta > 0$, take $t^m := \Delta^m \lfloor t/\Delta^m \rfloor$. If $\delta = \infty$, use $k_m := k$. Since $t^m \rightarrow t$, for all $\delta' \in (0, \infty)$, $\lim_m \inf_{[t^m, t^m+\delta')} q^m \geq \lim_m \inf_{[t, t+2\delta')} q^m = \inf_{[t, t+2\delta')} q > k$ where the strict inequality holds as $\delta > 2\delta'$, so that $\liminf_m \delta_m \geq \delta'$. Then, $\lim_m \delta_m = \delta$ as $\delta' \in (0, \infty)$ is arbitrary.

If $\delta \in (0, \infty)$ then $k > x_0$ as $\inf q \geq \lambda\mu > x_0$. Let $m' := \lceil \min\{\frac{\delta}{2}, k-x_0\}^{-1} \rceil$. For any $m \geq m'$ and $l \in \mathbb{N}$ such that $\Delta^l < \delta \wedge \frac{1}{m}$, $\inf_{[t^l, t^l+\delta-2/m)} q^l \geq \inf_{[t, t+\delta-1/m)} q^l$ and $\inf_{[t+\delta, t+\delta+1/m)} q^l \geq \inf_{[t^l, t^l+\delta+1/m)} q^l$. Moreover, $\inf_{[t, t+\delta-1/m)} q > k \geq \inf_{[t+\delta, t+\delta+1/m)} q$ as $\delta \in (0, \infty)$. Then, for l sufficiently large, there is $k' \in (k - \frac{1}{m}, k + \frac{1}{m})$ such that $\inf_{[t^l, t^l+\delta-2/m)} q^l > k' > \inf_{[t^l, t^l+\delta+1/m)} q^l$, as $\lim_l \inf_{[t, t+\delta-1/m)} q^l = \inf_{[t, t+\delta-1/m)} q$ and $\lim_l \inf_{[t+\delta, t+\delta+1/m)} q^l = \inf_{[t+\delta, t+\delta+1/m)} q$. Choose $(l_m)_{m=m'}^\infty \subseteq \mathbb{N}$ to be strictly increasing and such that, for all $m \geq m'$, there is $k'_m \in (k - \frac{1}{m}, k + \frac{1}{m})$ such that $\inf_{[t^{l_m}, t^{l_m}+\delta-2/m)} q^{l_m} > k'_m > \inf_{[t^{l_m}, t^{l_m}+\delta+1/m)} q^{l_m}$. Define $(k_l)_{l \in \mathbb{N}}$ by $k_l := k'_m$ for all $m \geq m'$ and $l = l_{m-1} + 1, \dots, l_m$, (where $l_{m'-1} := 0$). Then $\lim_l k_l = k$ and $\lim_l \delta_l = \delta$ as $|\delta - \delta_{l_m}| \leq \frac{2}{m} + \Delta^{l_m}$ for all $m \geq m'$.

81. It suffices to show that $\int_0^s a_r^m dr \rightarrow \int_0^s a_r dr$ for all $s \geq 0$. Fix s and note that $\int_0^s a_r^m dr = \int_0^s \alpha_{t^m+r}^m(k) dr = \int_0^{t^m+s} \alpha_r^m(k) dr - \int_0^{t^m} \alpha_r^m(k) dr$. Similarly, $\int_0^s a_r dr = \int_0^{t+s} \alpha_r(k) dr - \int_0^t \alpha_r(k) dr$. Then, $a^m \rightarrow a$ since $\int_0^{t^m+s} \alpha_r^m(k) dr \rightarrow \int_0^{t+s} \alpha_r(k) dr$ and $\int_0^{t^m} \alpha_r^m(k) dr \rightarrow \int_0^t \alpha_r(k) dr$, as $t^m \rightarrow t$, α^m and α take values in $[0, 1]$, and $\alpha^m \rightarrow \alpha$.

$(0, t^m, k, \alpha, q, v)$, and δ_m and a^m are given by (46) with t replaced by t^m . Letting $m \rightarrow \infty$ yields $v_t(k) = I^*(\theta) = I((\delta, a), \theta)$ since $\inf_{[t^m, t']} q \rightarrow \inf_{[t, t']} q$ for all $t' > t$, and v is continuous (as $v \in \mathcal{V}_0$). \square

G.3 Proof of Lemma 9

Endow A with the topology of pointwise convergence. I begin with two remarks, proved in Curello (2021).

Remark 6. Given $\Delta > 0$ and $m \in \mathbb{N}$,

$$(\alpha, q, v) \mapsto \left(k \mapsto \arg \max_{a \in [0, 1]} I_{\Delta, m}((1, a), (k, \alpha, q, v)) \right)$$

is a well-defined and continuous map $A^m \times Q_\Delta^m \times V_\Delta \rightarrow A$.

Remark 7. Given $\Delta > 0$ and $m \in \mathbb{N}$,

$$(\alpha, q, v) \mapsto (k \mapsto I_{\Delta, m}^*(k, \alpha, q, v))$$

is a well-defined and continuous map $A^m \times Q_\Delta^m \times V_\Delta \rightarrow V_\Delta$.

Proof of Lemma 9. From Lemma 8, we can choose $\Delta_0 > 0$ such that $\sup_{\Delta \in (0, \Delta_0)} q_\Delta^+ < \infty$. To define γ , note that there exists $R > 0$ such that, for all $m \in \mathbb{N}$, $\Delta \in (0, \Delta_0)$, $a \in [0, 1]$, $k \leq q_\Delta^+$, $\alpha \in A^m$, $q \in Q_\Delta^m$, and $v \in V_\Delta$,

$$\left| \int_{q_m}^\infty \text{ld} G^{n-1}(l) + G^{n-1}(q_m) v(k) - I_{\Delta, m}((1, a), (k, \alpha, q, v)) \right| < R\Delta$$

where G is degenerate with value x_0 if $m = 1$, and $G := \beta^{\Delta, m-1}((\alpha_l)_{l=1}^{m-1}, (q_l)_{l=1}^{m-1})$ otherwise. Moreover, as F is Lipschitz, there is an increasing map $\gamma_0 : \mathbb{R}_+ \rightarrow (0, \infty)$ such that $\beta_t(\alpha, q)$ is $\gamma_0(t)$ -Lipschitz on $[x_0, \infty)$ for all $\alpha \in \mathcal{A}$, $q \in \mathcal{Q}$, and $t \geq 0$. Define $\gamma : (0, \infty) \rightarrow (0, \infty)$ by

$$\gamma(t) := (R \vee \lambda) + 2R(n-1)\gamma_0(t) \sup\{q_\Delta^+ : \Delta \in (0, \Delta_0)\} \cup \{1\}.$$

Now fix $\Delta \in (0, \Delta_0)$ and $M \in \mathbb{N}$. To construct $\alpha \in A^M$, $q \in Q_\Delta^M$, and $v \in V_\Delta^{M+1}$, let $w : A^M \times Q_\Delta^M \rightarrow V_\Delta^{M+1}$ be given by $w_{M+1}(\alpha, q) = \underline{v}_\Delta$, and

$$[w_m(\alpha, q)](k) := I_{\Delta, m}^*(k, \alpha^m, q^m, w_{m+1}(\alpha, q))$$

for each $m \in \{1, \dots, M\}$, where $\alpha^m = (\alpha_l)_{l=1}^m$ and $q^m := (q_l)_{l=1}^m$. Note that w_{M+1} is well-defined and continuous. Then Remark 7 implies that w is well-defined and continuous.

Define the map $\psi : A^M \times Q_\Delta^M \times V_\Delta^{M+1} \rightarrow A^M \times Q_\Delta^M \times V_\Delta^{M+1}$ by

$$\psi(\alpha, q, v) := \left(\left(k \mapsto \arg \max_{a \in [0,1]} I_{\Delta,m}((1, a), (\alpha^m, q^m, w_{m+1}(\alpha, q))) : m = 1, \dots, M \right), \right. \\ \left. ([w_m(\alpha, q)](q_m) : m = 1, \dots, M), w(\alpha, q) \right).$$

Since w is continuous and V is endowed with the topology of uniform convergence, ψ is well-defined and continuous by Remark 6. Then ψ admits a fixed point (α, q, v) by Brower's fixed-point theorem.

Fix $m \in \{1, \dots, M\}$. To prove (50) note that, by definition of ψ , it suffices to prove that $v_m(k) \leq (\geq) \underline{v}_\Delta(k)$ for all $k \geq (\leq) q_m$. This holds since $v_m(q_m) = q_m \leq \underline{v}_\Delta(q_m)$, and $v_m - \underline{v}_\Delta$ is positive and decreasing (as $v_m \in V_\Delta$).

It remains to prove (53). Let $\widehat{G} := G_{m-1}^{n-1}$.

Claim 6. $1 - \widehat{G}(q_m) \leq 2R\Delta(n-1)\gamma_0(m\Delta)$.

Proof. Assume without loss of generality that $G_{m-1}(q_m) < 1$. By definition of γ_0 , G_{m-1} is $\gamma_0((m-1)\Delta)$ -Lipschitz on $[x_0, \infty)$. Then, as γ_0 is increasing, G_{m-1} is $\gamma_0(m\Delta)$ -Lipschitz on $[x_0, \infty)$, so that \widehat{G} is $(n-1)\gamma_0(m\Delta)$ -Lipschitz on $[x_0, \infty)$. Then, \widehat{G} conditioned on the event (q_m, ∞) FOSD-dominates $U[q_m, q_m + 1/\{(n-1)\gamma_0(m\Delta)\}]$. Hence

$$q_m + \frac{1}{2(n-1)\gamma_0(m\Delta)} \leq \frac{\int_{q_m}^\infty l d\widehat{G}}{1 - \widehat{G}(q_m)} \quad \Leftrightarrow \quad 1 - \widehat{G}(q_m) \leq 2(n-1)\gamma_0(m\Delta) \int_{q_m}^\infty l - q_m d\widehat{G}.$$

Then, result follows as

$$\int_{q_m}^\infty l - q_m d\widehat{G} \leq \int_{q_m}^\infty l - q_m d\widehat{G} + \widehat{G}(q_m)[v_{m+1}(q_m) - q_m] \\ = \int_{q_m}^\infty l d\widehat{G} + \widehat{G}(q_m)v_{m+1}(q_m) - I_{\Delta,m}((1, \alpha_m), (q_m, \alpha^m, q^m, v_{m+1})) \leq R\Delta$$

where the first inequality holds since $v_{m+1}(q_m) \geq q_m$ (as $v_{m+1} \in V_\Delta$), and the equality holds because $q_m = I_{\Delta,m}((1, \alpha_m), (q_m, \alpha^m, q^m, v_{m+1}))$. \square

Since $v_{m+1} - \underline{v}_\Delta$ is positive and decreasing and $v_m(k) = \underline{v}_\Delta(k)$ for $k \geq q_m$, $v_{m+1} - v_m$ is positive and decreasing on $[q_m, \infty)$. Hence, we may restrict attention to $k \leq q_m$, so that

$$|v_{m+1}(k) - v_m(k)| = \left| \left(\int_{q_m}^\infty v_{m+1}(k) - l d\widehat{G}(l) \right) + \left(\int_{q_m}^\infty l d\widehat{G}(l) + \widehat{G}(q_m)v_{m+1}(k) - v_m(k) \right) \right| \\ \leq 2R(n-1)\gamma_0(m\Delta)\Delta \sup_{\Delta' \in (0, \Delta_0)} q_{\Delta'}^+ + R\Delta \leq \gamma(m\Delta)\Delta$$

where the inequality follows from Claim 6.

Finally note that, for any $j \neq i$, if agent j plays a strategy $\rho \in R(\hat{\alpha}, \hat{q})$, where $\hat{\alpha} \in \mathcal{A}$ and $\hat{q} \in \mathcal{Q}$ are such that $\hat{\alpha}_t = \alpha_l$ and $\hat{q}_t = q_l$ for all $t \in [(l-1)\Delta, l\Delta)$ and $l = 1, \dots, m$, then the probability that $k_{(m-1)\Delta}^j \leq q_m$ and that k^j is constant over $[(m-1)\Delta, m\Delta)$ is $e^{-\alpha_m \Delta \lambda} G_m(q_m)$. Then

$$\begin{aligned} |G_m - G_{m-1}| &\leq 1 - e^{-\alpha_m \Delta \lambda} G_{m-1}(q_m) \\ &\leq 1 - (1 - \Delta \lambda)[1 - 2R(n-1)\gamma_0(m\Delta)\Delta] \leq \gamma(m\Delta)\Delta \end{aligned}$$

where the second inequality follows from Claim 6. \square

H Proof of Proposition 4

The argument relies on the following result, proved in Curello (2021).

Lemma 11. *For all $t' \geq 0$ and $k \geq x_0$, (23) holds for almost all $t > t'$ such that $k < \inf_{[t', t)} q$.*

Lemma 12. *Suppose that (24) holds. Then $v_d(x_0) \geq \lambda\mu$, and there are $N \in \mathbb{N}$ and $\epsilon > 0$ such that $v_d(x_0) \geq v_e(x_0) + \epsilon$ for any $n \geq N$.*

Proof. I prove that $v_d(x_0) \geq \lambda\mu$. Note that $v_d(x) \geq x$ for all $x \geq 0$, hence assume without loss of generality that $x_0 < \lambda\mu$, so that $\alpha_d(x_0) > 0$. Then $v_d(x_0) \geq \lambda\mu$, for otherwise

$$x_0 \geq \lambda \mathbb{E}_F[\max\{0, v_d(x_0 + \tilde{z}) - v_d(x_0)\}] > \lambda \mathbb{E}_F[\max\{0, x_0 + \tilde{z} - \lambda\mu\}] \geq x_0,$$

where the first inequality follows from (21) as $\alpha_d(x_0) > 0$, the second holds since $F(\lambda\mu) < 1$, and the last is (24).

For the last part, note that there is a unique continuous $\bar{v}_e : (0, \infty) \rightarrow \mathbb{R}$ such that $\bar{v}_e(x) = x$ on $[\lambda\mu, \infty)$ and

$$\lambda\{\mathbb{E}_F[\bar{v}_e(x + \tilde{z})] - \bar{v}_e(x)\} = x \tag{55}$$

on $(0, \lambda\mu]$. Similarly, there is a unique continuous $\bar{v}_d : (0, \infty) \rightarrow \mathbb{R}$ such that $\bar{v}_d(x) = x$ on $[\lambda\mu, \infty)$ and

$$\lambda \mathbb{E}_F[\max\{0, \bar{v}_d(x + \tilde{z}) - \bar{v}_d(x)\}] = x \tag{56}$$

on $(0, \lambda\mu]$. I claim that, for any $\hat{x} \in (0, \lambda\mu)$, there is $N \in \mathbb{N}$ such that $v_e = \bar{v}_e$ and $v_d = \bar{v}_d$ on $[\hat{x}, \infty)$ for $n \geq N$. To prove the claim, fix \hat{x} and note that $\alpha_e^{-1}(0) = \alpha_d^{-1}(0) = [\lambda\mu, \infty)$. In particular, $v_e(x) = v_d(x) = x$ for $x \geq \lambda\mu$. Then, there is $N \in \mathbb{N}$ such that $\alpha_e(\hat{x}) < 1$ and $\alpha_d(\hat{x}) < 1$ for $n \geq N$, since effort is costly (as $\hat{x} > 0$) and the benefit of exerting effort vanishes as the aggregate effort exerted by opponents grows. Since α_e and α_d are

decreasing, $0 < \alpha_e < 1$ and $0 < \alpha_d < 1$ on $[\hat{x}, \lambda\mu)$ so that v_e satisfies (55) and v_d satisfies (56) on $[\hat{x}, \lambda\mu)$. Then, $v_e = \bar{v}_e$ and $v_d = \bar{v}_d$ on $[\hat{x}, \infty)$, as v_e and v_d are continuous.

Case 1. $x_0 > 0$. In light of the claim, it suffices to show that $\bar{v}_d(x_0) > \bar{v}_e(x_0)$. Fix n such that $v_e = \bar{v}_e$ and $v_d = \bar{v}_d$ on $[x_0, \infty)$. Then, $\bar{v}_d(x_0) > \bar{v}_e(x_0)$ unless $v_e(x_0) \geq \lambda\mu$, since $v_e(x_0) \geq \lambda\mu$. Moreover, if $v_e(x_0) > \lambda\mu$ then v_e is non-monotone on $[x_0, \infty)$, as $v_e(x) < \lambda\mu$ for some $x \in (x_0, \lambda\mu)$, by (55). Then, $\bar{v}_d(x_0) > \bar{v}_e(x_0)$ by Theorem 3, as desired.

Case 2. $x_0 = 0$. Note that, as $x \downarrow 0$, $\bar{v}_e(x)$ and $\bar{v}_d(x)$ converge to some $\bar{v}_e(0) \in \mathbb{R}_+$ and $\bar{v}_d(0) \in [0, \infty]$, respectively. Moreover $\bar{v}_d(x) \geq \bar{v}_e(x)$ for all $x \geq 0$, where the inequality is strict for $x > 0$ sufficiently small, by Case 1. Hence, $\bar{v}_d(0) > \bar{v}_e(0)$. Then, result follows since, clearly, $v_e(0) \rightarrow \bar{v}_e(0)$ and $v_d(0) \rightarrow \bar{v}_d(0)$ as n grows. \square

Proof of Proposition 4. The case $x_0 \geq \lambda\mu$ is clear, hence suppose that $x_0 < \lambda\mu$. From Lemma 12, there are $N_1 \in \mathbb{N}$, $T > 0$ and $\epsilon \in (0, 1)$ such that

$$(1 - \epsilon)e^{-T}v_d(x_0) > v_e(x_0)$$

for all $n \geq N_1$. Fix $T' \in [0, T)$ and $\hat{x} > v_e(x_0)e^T/\epsilon$ and note that there are $N_2 \in \mathbb{N}$ and $Q \in \mathbb{R}_+$ such that

$$v_T^c(\hat{x}) - v_T^c(x_0) \leq \hat{x} - \frac{v_e(x_0)e^T}{\epsilon}$$

if $n \geq N_2$, and $q(t) \geq Q$ and $\alpha_t^c(x_0) = 1$ for all $T' \leq t \leq T$. Intuitively, if no disclosure occurs by time T , any agent i assigns high probability to the event that $k_T^j \geq \hat{x}$ for some $j \neq i$, since her numerous opponents exerted maximal effort within the interval $[T', T]$, and the disclosure cutoff q was high within this time interval. Hence, the net benefit to agent i of having $k_T^i = \hat{x}$ instead of $k_T^i = x_0$ is low.

Given $t \in [0, T]$ and $n \in \mathbb{N}$, let $p_{t,n}$ the probability that a disclosure occurs prior to time t when ρ^c is played. Note that there is $N_3 \in \mathbb{N}$ such that, for any $n \geq N_3$, $q \geq Q$ over $[T', T]$ if $p_{T',n} \leq 1 - \epsilon$ and $\alpha_t^c(x_0) = 1$ for all $t \leq T$.

Let $N := \max\{N_1, N_2, N_3\}$. It remains to show that $v_0^c(x_0) \geq v_e(x_0)$ for any $n \geq N$. Fix n and note that, if a value d is disclosed at time t when ρ^c is played, then $d \geq q(t) \geq v_d(x_0)$, and the continuation payoff after the disclosure is d . If no disclosure occurs within $[0, t]$, the continuation payoff to agent i is $v_t^c(k_t^i) \geq v_t^c(x_0)$, where the inequality holds since v_t^c is increasing and $k_t^i \geq x_0$. Then

$$v_t^c(x_0) \geq e^{-t}[p_{t,n}v_d(x_0) + (1 - p_{t,n})v_t^c(x_0)]. \quad (57)$$

Since $v_t^c(x_0) \geq 0$, $v_0^c(x_0) \geq v_e(x_0)$ if $p_{T,n} \geq 1 - \epsilon$. Hence, assume that $p_{T,n} < 1 - \epsilon$.

Suppose that $\alpha_t^c(x_0) = 1$ for a.e. $t \leq T$. Then, from above,

$$v_0^c(x_0) \geq \epsilon e^{-T}v_T^c(x_0) \geq \epsilon e^{-T}\{\hat{x} - [v_T^c(\hat{x}) - v_T^c(x_0)]\} \geq v_e(x_0)$$

where the first inequality follows from (57), and the second holds since $v_T^c(\hat{x}) \geq \hat{x}$.

Finally, suppose that $\alpha_t^c(x_0) < 1$ for a non-nul set of times $t \leq T$. Then, by (23), for some $t \leq T$,

$$v_t^c(x_0) \geq \mathbb{E}_F[v_t^c(x_0 + \tilde{z})] - \frac{x_0}{\lambda} \geq \mathbb{E}_F[\max\{x_0 + \tilde{z}, v_t^c(x_0)\}] - \frac{x_0}{\lambda}$$

where the second inequality holds since v_t^c is increasing and, clearly, $v_t^c(k) \geq k$ for all $k \geq x_0$. Moreover, $\alpha_d(x_0) < 1$ as $x_0 < \lambda\mu$, so that

$$v_d(x_0) \leq \mathbb{E}_F[\max\{x_0 + \tilde{z}, v_d(x_0)\}] - \frac{x_0}{\lambda}$$

by (21), since $v_d(x) = x$ for $x \geq \lambda\mu$ and $v_d(x_0) \geq \lambda\mu$ (Lemma 12). Then, $v_t^c(x_0) \geq v_d(x_0) \geq v_e(x_0)$. \square

Supplemental appendices

I Necessity of assumptions

In this appendix, I argue that various substantial assumptions of the model are necessary to obtain important results. In Appendix I.1 I show that, if payoffs are separable, then innovations are not detrimental in the equilibrium with forced disclosure, and concealing innovations is unprofitable. In Appendix I.2, I show that the same is true for arbitrary payoffs if innovations have fixed size. In Appendix I.3, I show via an example that, if the cost of effort does not become more convex as the stock grows, then it may be efficient to *raise* effort after an increment in the stock.

I.1 Payoffs are not necessarily separable

Recall that continuation payoffs in the symmetric equilibrium with forced disclosure (Theorem 2) are $v_e(x_t)$, where x_t is the current stock.

Proposition 6. *If the cost of effort does not increase with the stock, i.e. $c(a, x) = c(a)$, then v_e is increasing.*

Proof. Recall from the proofs of Theorem 5 and Theorem 2 that v_e is the unique fixed point of P_e in \mathcal{V}'' , that P_e maps \mathcal{V}'' to itself and is increasing, and that \mathcal{V}'' is a complete lattice. Let \mathcal{V}''' be the set of $v \in \mathcal{V}''$ that are increasing. It suffices to show that P_e admits a fixed point in \mathcal{V}''' . Note that \mathcal{V}''' is a complete lattice. Then, since P_e is increasing on \mathcal{V}'' , by Tarski's fixed point theorem, it suffices to show that P_e maps \mathcal{V}''' to itself. Fix $v \in \mathcal{V}'''$. Note that $P_e v \in \mathcal{V}''$, so that it remains to show that $P_e v$ is increasing.

Since v is increasing and $v - b$ is decreasing, $L^e v$ is continuous. Then, $Q(L^e v(x), x)$ and, therefore, $P_e v$, are continuous. Moreover, from Step 3 of the proof of Theorem 5, $Q(x, L^e v(x))$ is decreasing in x . Let $I_a := \{x \geq 0 : Q(x, L^e v(x)) = a\}$ for $a \in \{0, 1\}$ and $I = \mathbb{R}_+ \setminus (I_0 \cup I_1)$. If $Q(x, L^e v(x)) = 0$ for some $x \geq 0$, then I_0 is an interval, and $P_e v = b$ over I_0 , so that it is increasing on I_0 . If $Q(x, L^e v(x)) \in (0, 1)$ for some $x \geq 0$, then I is also an interval. Moreover, for any $x \in I$,

$$\lambda[L^e v(x) - P_e v(x)] = \lambda \frac{L^e v(x) - [b(x) - c(a, x)]}{1 + \lambda n a} = c'(a)$$

where $a = Q(x, L^e v(x))$, the first equality follows by definition of P_e , and the second from (34). Since v is increasing and $Q(x, L^e v(x))$ is decreasing, $P_e v$ is then increasing on I . Finally, if $Q(x, L^e v(x)) = 1$ for some $x \geq 0$, then I_1 is an interval, and

$$P_e v(x) = \frac{b(x) - c(1, x) + \lambda n L^e v(x)}{1 + \lambda n}$$

over I . Then, $P_e v$ is increasing on I_1 since $b(x) - c(1, x)$ and v are increasing. Therefore, $P_e v$ is decreasing since I_0, I_1 and I partition \mathbb{R}_+ and $P_e v$ is continuous. \square

Given Lemma 2, Proposition 6 implies that, if payoffs are separable, then innovations are not detrimental.

In the remainder of this section, I show that agents have no incentive to conceal innovations in equilibrium if payoffs are separable. Remember that α_e is the effort policy in the equilibrium with forced disclosure (Theorem 2), and that ω_0 is the ‘full-disclosure’ policy in the game with concealment. The next result describes when (α_e, ω_0) is an equilibrium in the game with concealment. I omit the proof as it follows from standard dynamic-programming arguments.

Lemma 13. *Given $x_0 \geq 0$, the following are equivalent:*

1. *The strategy profile (α_e, ω_0) forms a PBE of the game with concealment.*
2. *For all $k \geq x \geq x_0$, $\Delta(k, x) \leq 0$ where*

$$\begin{aligned} \Delta(k, x) = & \max_{a \in [0, 1]} b(x) - c(a, x) + a \lambda \{ \mathbb{E}_F[v_e(k + \tilde{z})] - v_e(k) \} \\ & + \alpha_e(x) \lambda (n - 1) \mathbb{E}_F \{ \mathbf{1}_{x + \tilde{z} > k} [v_e(x + \tilde{z}) - v_e(k)] \} - v_e(k). \end{aligned} \quad (58)$$

Assuming that, when (α_e, ω_0) is played, some agent i obtains an innovation at time t raising her stock to $k_t^i > x_t$, $\Delta(k_t^i, x_t)$ has the same sign as the net benefit of the deviation: ‘conceal fully, and exert optimal effort until the next innovation (by any agent), then

disclose fully and revert to (α_e, ω_0) '.⁸² The first three terms are the continuation payoffs from delaying the disclosure, whereas $v_e(k)$ is the payoff from disclosing immediately.

The next result shows that, if the cost of effort does not increase with the stock, then concealment is not profitable in equilibrium.

Corollary 4. *If the cost of effort does not increase with the stock, i.e. $c(a, x) = c(a)$, then the strategy profile (α_e, ω_0) forms a PBE of the game with concealment.*

Proof. Fix $x_0 \geq 0$. From Proposition 13, it suffices to show that $\Delta(k, x) \leq 0$ for all $k \geq x \geq x_0$. Note that $\Delta(x_0, x_0) = 0$ since v_e satisfies (10). Hence, it suffices to show that $\Delta(k, x_0)$ is decreasing in k on $[x_0, \infty)$. From Proposition 6, v_e is increasing. Then, the second expectation in (58) is decreasing in k . Hence, it suffices to show that the map $\psi : k \mapsto \mathbb{E}_F[v_e(k + \tilde{z})] - v_e(k)$ is decreasing.

Let

$$I_1 = \{x \geq 0 : \alpha_e(x) = 1\}, \quad I = \{x \geq 0 : \alpha_e(x) \in (0, 1)\}, \quad I_0 = \{x \geq 0 : \alpha_e(x) = 0\}.$$

From Theorem 2, α_e is continuous and decreasing, so that I_1 , I and I_0 are (possibly empty) intervals. Since ψ is clearly continuous, it suffices to show that it is decreasing on each of I_1 , I and I_0 . On I_0 , $v_e = b$, so ψ is decreasing since b is concave. On I , (11) implies that $\psi(k) = c(\alpha_e(k))$. Then ψ is decreasing since α_e is. On I_1 ,

$$\psi(k) = \frac{\mathbb{E}_F[v_e(k + \tilde{z})] - b(k)}{1 + \lambda n}.$$

Then, ψ is decreasing on I_1 from Claim 1, since $v_e - b$ is decreasing (see Theorem 2). \square

I.2 Innovations have random size

Note that Lemma 1 continues to hold if innovations have fixed size. Then, it is immediate that, in this case, innovations are not detrimental in the equilibrium with forced disclosure.

Corollary 5. *If innovations have fixed size, then innovations are not detrimental.*

The next result shows that, if innovations have fixed size, then concealment is not profitable in equilibrium.

82. Assuming that agent i is indifferent between this and no deviation, her continuation payoff if deviating is the sum of all but the last term $-v_e(k)$, and $v_e(k)$ is her payoff from not deviating. More specifically, if she deviates and exerts optimal effort a , she obtains flow payoff $b(x) - c(a, x)$ until a disclosure and, if she obtains an innovation of size z before her opponents disclose, her continuation payoff increases by $v_e(k + z) - v_e(k)$; if instead one of her opponents discloses an innovation of size z , agent i discloses k immediately afterwards if and only if $x + z < k$.

Corollary 6. *If innovations have fixed size, then the strategy profile (α_e, ω_0) forms a PBE of the game with concealment.*

Proof. Fix $x_0 \geq 0$. From Lemma 13, it suffices to show that $\Delta(k, x) \leq 0$ for all $k \geq x \geq x_0$ such that $k - x = l\mu$ for some $l \in \mathbb{N}$. From Lemma 1, $v(k) \geq v(x + \mu)$. Then, as F is degenerate with value μ ,

$$\Delta(x, k) = \max_{a \in [0, 1]} b(x) - c(a, x) + a\lambda\{\mathbb{E}_F[v_e(k + \tilde{z})] - v_e(k)\} - v_e(k).$$

Then, $\Delta(k, x) \leq 0$ since v_e satisfies (10), $\mathbb{E}_F[v_e(k + \tilde{z})] - v_e(k) \geq 0$ (Lemma 1), and $b(x) - c(a, x)$ is increasing in x . \square

I.3 The cost becomes more convex as the stock grows

I exhibit a parametrised example satisfying all the assumptions of Section 2 except the requirement that $c_{11}(a, x)$ be increasing in x , and show that the efficient effort schedule α_* (described in Theorem 1) is non-monotone in the stock.

Consider

$$u(a, x) = (1 + x)(1 - a) - \frac{a^2}{2}e^{-x}. \quad (59)$$

It is easy to see that u satisfies all the assumptions of the model except the requirement that $u_{112} \geq 0$.⁸³ Let $F(z) = 1 - e^{-z}$ and fix $\lambda n > 1$. Note that $b(x)$ solves (6) precisely over $[x_*, \infty)$, where $x_* = \lambda n - 1$. Then, efficient effort α_* ceases at x_* .

Note also that there exists $\underline{x} \in [0, x_*)$ such that the differential equation

$$\alpha^{i'}(x) = \left[\left(\frac{\alpha^i(x)}{2} + \frac{2}{\lambda n} \right) \alpha^i(x) - \left(1 - \frac{x}{\lambda n} \right) e^x \right] \left(\alpha^i(x) + \frac{1}{\lambda n} \right)^{-1} \quad (60)$$

admits a solution α^i on $(\underline{x}, x_*]$ satisfying $\alpha^i(x_*) = 0$. Moreover, it is clear that we may choose α^i and \underline{x} such that α^i is strictly positive over (\underline{x}, x_*) and either $\alpha^i(\underline{x}) = 0$ or $\underline{x} = 0$. Then, by the Picard–Lindelöf theorem, α^i is uniquely determined on $[\underline{x}, x_*]$. Simulations imply that, for $\lambda n = 2.55$, $\underline{x} = 0$, and α^i is non-monotone and within $(0, 1)$ over $[0, x_*)$ (see Figure 4). Then, simple algebra implies that α_*^i matches α^i on $[0, x_*]$.⁸⁴ In particular, efficient effort is non-monotone.

⁸³. The other assumptions are that u is twice continuously differentiable, increasing and concave in x , convex and strictly decreasing in a , and that $u_{12} \leq 0$.

⁸⁴. To see why, extend α^i to \mathbb{R}_+ by setting $\alpha^i(x) = 0$ for $x > x_*$, and let $w : \mathbb{R}_+ \rightarrow \mathbb{R}$ be given by $w(x) := u(\alpha^i(x), x) + \alpha^i(x)c_1(\alpha^i(x), x)$ for $x \leq x_*$, and $w(x) = 1 + x$ for $x > x_*$. Then, it suffices to show that w and α^i solve (6) and (7). Note that (60) can be rearranged into $\lambda n w'(x) = c_1(\alpha^i(x), x) - \frac{d}{dx}c_1(\alpha^i(x), x)$. Integrating and using the fact that $x_* = \lambda n - 1$ yields

$$w(x) = \int_{\lambda n - 1}^x \frac{c_1(\alpha^i(z), z)}{\lambda n} dz - \frac{c_1(\alpha^i(x), x)}{\lambda n} + 1 - \lambda n$$




Figure 4: Efficient effort given payoff function (59), $F(z) = 1 - e^{-z}$, and $\lambda n = 2.6$.

over $[0, \lambda n - 1]$. Then w solves the Volterra equation of the second kind

$$w(x) + \int_{\lambda n - 1}^x e^{x-y} w(y) dy = (\lambda n + 1) e^{x+1-\lambda n} - \frac{c_1(\alpha^i(x), x)}{\lambda n}$$

over $[0, \lambda n - 1]$ (see e.g. Polyanin and Manzhirov (2008), page 144). Equivalently, $\lambda n \{\mathbb{E}_F[w(x + \tilde{z})] - w(x)\} = c_1(\alpha^i(x), x)$ over $[0, x_*]$. Then, (7) holds and $w(x) = u(\alpha^i(x), x) + \alpha^i(x) \lambda n \{\mathbb{E}_F[w(x + \tilde{z})] - w(x)\}$ for all $x \geq 0$. Hence, (6) holds as well.