

Cohomology of weighted Rota-Baxter Lie algebras and Rota-Baxter paired operators

Apurba Das

*Department of Mathematics and Statistics,
Indian Institute of Technology, Kanpur 208016, Uttar Pradesh, India.
Email: apurbadas348@gmail.com*

Abstract

In this paper, we define representations and cohomology of weighted Rota-Baxter Lie algebras. As applications of cohomology, we study abelian extensions and formal 1-parameter deformations weighted Rota-Baxter Lie algebras. Finally, we consider weighted Rota-Baxter paired operators that induces a weighted Rota-Baxter Lie algebra and a representation of it. We define suitable cohomology for such paired operators that govern deformation.

2020 MSC classification: 17B38, 17B56, 16S80.

Keywords: Weighted Rota-Baxter Lie algebras, Representations, Cohomology, Deformations, Rota-Baxter paired operators.

Contents

1	Introduction	1
2	Representations of weighted Rota-Baxter Lie algebras	3
3	Cohomology of weighted Rota-Baxter Lie algebras	7
4	Applications of cohomology	12
5	Rota-Baxter paired operators	16

1 Introduction

Rota-Baxter operators were first appeared in the work of Baxter in his study of the fluctuation theory [6] and further developed by Rota in combinatorics [25]. Subsequently, Cartier [7] and Atkinson [2] studied further properties of Rota-Baxter operators. In last twenty years, Rota-Baxter operators on associative algebras pay very much attention due to its connection with combinatorics of shuffle algebras [15], Yang-Baxter equations [1], dendriform algebras [1,13], renormalizations in quantum field theory [8], multiple zeta values in number theory [16] and splitting of algebraic operads [3]. See [17] for more details about Rota-Baxter operators on associative algebras. On the other hand, Rota-Baxter operators on Lie algebras was first considered by Kupersmidt in the study of classical r -matrices [19]. They are also connected with pre-Lie algebras [3], integrable systems [26] and combinatorics of Lyndon-Shirshov words [23].

Deformation theory of some algebraic structure goes back to Gerstenhaber [14] for associative algebras and Nijenhuis-Richardson [24] for Lie algebras. Recently, deformation theory has been adapted to Rota-Baxter operators on Lie algebras [27] and subsequently developed in [9] for associative algebras. In both these papers, the authors only considered deformations of Rota-Baxter operators by keeping the underlying algebras intact. In [11,21] the authors dealt with deformations of Rota-Baxter algebras in which they simultaneously deform Rota-Baxter operators and the underlying algebras. Note that all these works are concern about Rota-Baxter operators of weight zero.

Rota-Baxter operators with arbitrary weight (also called weighted Rota-Baxter operators) was considered in [4,5]. They are related with tridendriform algebras [12], post-Lie algebras and modified Yang-Baxter equations [4], weighted infinitesimal bialgebras and weighted Yang-Baxter equations [32], combinatorics of rooted forests [31], among others. Recently, the authors in [18] defined the cohomology of Rota-Baxter operators of weight 1 on Lie algebras and Lie groups. This motivates the present author to study cohomology and deformations of weighted Rota-Baxter operators on both associative and Lie algebras [10]. However, the simultaneous deformations of (associative) algebras and weighted Rota-Baxter operators are considered in a recent preprint of Wang and Zhou [30]. More precisely, they considered weighted Rota-Baxter associative algebras and define cohomology of them with coefficients in a suitable Rota-Baxter bimodule. When considering the cohomology with coefficients in itself, it governs the simultaneous deformations of algebras and weighted Rota-Baxter operators.

Our aim in this paper is to apply the approach of [30] to weighted Rota-Baxter Lie algebras. More precisely, we first consider representations of weighted Rota-Baxter Lie algebras and provide various constructions of representations. Then we define the cohomology of a weighted Rota-Baxter Lie algebra with coefficients in a representation. This cohomology is obtained as a byproduct of the standard Chevalley-Eilenberg cohomology of the underlying Lie algebra and the cohomology of the underlying weighted Rota-Baxter operator. When the weight is zero, our cohomology coincides with the one introduced in [21]. We interpret our second cohomology group as the isomorphism classes of abelian extensions of weighted Rota-Baxter Lie algebras. Then we consider deformations of weighted Rota-Baxter Lie algebras in which we simultaneously deform underlying Lie algebras and weighted Rota-Baxter operators. The infinitesimals of such deformations are 2-cocycles in the cohomology and equivalent deformations produce cohomologous 2-cocycles. Hence they correspond to the same element in the second cohomology group. We also find a sufficient condition for the rigidity of a weighted Rota-Baxter Lie algebra.

Finally, given a Lie algebra and a representation, we introduce Rota-Baxter paired operators that induces a weighted Rota-Baxter Lie algebra and representation of it. The terminology of Rota-Baxter paired operators is motivated by a paper by Zheng, Guo and Zhang [33] where the authors introduced Rota-Baxter paired modules in the associative context. We define a suitable differential graded Lie algebra that characterize Rota-Baxter paired operators as its Maurer-Cartan elements. This suggests us to define the cohomology of Rota-Baxter paired operators that control formal deformations.

The paper is organized as follows. In Section 2 we consider weighted Rota-Baxter Lie algebras and their representations. We also give several new constructions of representations. The cohomology of a weighted Rota-Baxter Lie algebra with coefficients in a representation is defined in Section 3. Applications of such cohomology to abelian extensions and formal 1-parameter deformations are given in Section 4. Finally, in Section 5, we define Rota-Baxter paired operators. We provide the cohomology of such paired operators and consider their deformations.

All vector spaces, (multi)linear maps, wedge products are over a field \mathbf{k} of characteristic 0.

2 Representations of weighted Rota-Baxter Lie algebras

In this section, we consider weighted Rota-Baxter Lie algebras [28] and introduce their representations. We also provide various examples and new constructions. Let $\lambda \in \mathbf{k}$ be a fixed scalar unless specified otherwise.

2.1 Definition. (i) Let \mathfrak{g} be a Lie algebra. A linear map $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ is said to be a λ -weighted Rota-Baxter operator if \mathfrak{T} satisfies

$$[\mathfrak{T}(x), \mathfrak{T}(y)] = \mathfrak{T}([\mathfrak{T}(x), y] + [x, \mathfrak{T}(y)] + \lambda[x, y]), \quad \text{for } x, y \in \mathfrak{g}. \quad (1)$$

(ii) A λ -weighted Rota-Baxter Lie algebra is a pair $(\mathfrak{g}, \mathfrak{T})$ consisting of a Lie algebra \mathfrak{g} together with a λ -weighted Rota-Baxter operator on it.

2.2 Example. (i) For any Lie algebra \mathfrak{g} , the pair $(\mathfrak{g}, \text{id}_{\mathfrak{g}})$ is a (-1) -weighted Rota-Baxter Lie algebra.

(ii) Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Then for any $\mu \in \mathbf{k}$, the pair $(\mathfrak{g}, \mu\mathfrak{T})$ is a $(\mu\lambda)$ -weighted Rota-Baxter Lie algebra.

(iii) Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Then $(\mathfrak{g}, -\lambda\text{id}_{\mathfrak{g}} - \mathfrak{T})$ is so.

(iv) Given a λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ and an automorphism $\psi \in \text{Aut}(\mathfrak{g})$ of the Lie algebra \mathfrak{g} , the pair $(\mathfrak{g}, \psi^{-1} \circ \mathfrak{T} \circ \psi)$ is a λ -weighted Rota-Baxter Lie algebra.

(v) Let \mathfrak{g} be a Lie algebra which splits as the direct sum of two subalgebras \mathfrak{g}_- and \mathfrak{g}_+ . Then $(\mathfrak{g}, \mathfrak{T})$ is a λ -weighted Rota-Baxter Lie algebra, where $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ is given by

$$\mathfrak{T}(x_-, x_+) = (0, -\lambda x_+), \quad \text{for } (x_-, x_+) \in \mathfrak{g}_- \oplus \mathfrak{g}_+ = \mathfrak{g}.$$

(vi) This example generalizes the previous one. Let \mathfrak{g} be a Lie algebra which splits as the direct sum of three subalgebras \mathfrak{g}_- , \mathfrak{g}_0 and \mathfrak{g}_+ in which \mathfrak{g}_- and \mathfrak{g}_+ are both \mathfrak{g}_0 -modules (i.e., representations of \mathfrak{g}_0). If $(\mathfrak{g}_0, \mathfrak{T}_0)$ is a λ -weighted Rota-Baxter Lie algebra then $(\mathfrak{g}, \mathfrak{T})$ is so, where $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ is given by

$$\mathfrak{T}(x_-, x_0, x_+) = (0, \mathfrak{T}_0(x_0), -\lambda x_+), \quad \text{for } (x_-, x_0, x_+) \in \mathfrak{g}.$$

2.3 Definition. Let $(\mathfrak{g}, \mathfrak{T})$ and $(\mathfrak{g}', \mathfrak{T}')$ be two λ -weighted Rota-Baxter Lie algebras. A morphism from $(\mathfrak{g}, \mathfrak{T})$ to $(\mathfrak{g}', \mathfrak{T}')$ is a Lie algebra homomorphism $\phi : \mathfrak{g} \rightarrow \mathfrak{g}'$ satisfying additionally $\phi \circ \mathfrak{T} = \mathfrak{T}' \circ \phi$. It is called an isomorphism if ϕ is so.

Let \mathfrak{g} be a Lie algebra. Recall that a representation of \mathfrak{g} is a vector space \mathcal{V} with a linear map (called the action map) $\rho : \mathfrak{g} \rightarrow \text{End}(\mathcal{V})$ satisfying

$$\rho[x, y] = \rho(x) \circ \rho(y) - \rho(y) \circ \rho(x), \quad \text{for } x, y \in \mathfrak{g}.$$

We denote a representation of \mathfrak{g} as above by (\mathcal{V}, ρ) or simply by \mathcal{V} when no confusion arises. Note that any Lie algebra \mathfrak{g} is a representation of itself with the action map $\rho : \mathfrak{g} \rightarrow \text{End}(\mathfrak{g})$ given by $\rho(x)(y) = [x, y]$, for $x, y \in \mathfrak{g}$. This is called the adjoint representation.

2.4 Definition. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. A representation of it is a pair $(\mathcal{V}, \mathcal{T})$ in which $\mathcal{V} = (\mathcal{V}, \rho)$ is a representation of the Lie algebra \mathfrak{g} and $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$ is a linear map satisfying

$$\rho(\mathcal{T}x)(\mathcal{T}u) = \mathcal{T}(\rho(\mathfrak{T}x)u + \rho(x)(\mathcal{T}u) + \lambda\rho(x)u), \quad \text{for } x \in \mathfrak{g}, u \in \mathcal{V}. \quad (2)$$

2.5 Example. Any λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ is a representation of itself. We call this the adjoint representation.

2.6 Example. Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. Then the pair $(\mathcal{V}, \text{id}_{\mathcal{V}})$ is a representation of the (-1) -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \text{id}_{\mathfrak{g}})$.

2.7 Example. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then for any scalar $\mu \in \mathbf{k}$, the pair $(\mathcal{V}, \mu\mathcal{T})$ is a representation of the $(\mu\lambda)$ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mu\mathfrak{T})$.

2.8 Proposition. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then $(\mathcal{V}, -\lambda\text{id}_{\mathcal{V}} - \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, -\lambda\text{id}_{\mathfrak{g}} - \mathfrak{T})$.

Proof. For any $x \in \mathfrak{g}$ and $u \in \mathcal{V}$, we observe that

$$\begin{aligned} & \rho((-\lambda\text{id}_{\mathfrak{g}} - \mathfrak{T})x)(-\lambda\text{id}_{\mathcal{V}} - \mathcal{T})(u) \\ &= \lambda^2\rho(x)u + \lambda\rho(\mathfrak{T}x)u + \lambda\rho(x)\mathcal{T}u + \rho(\mathfrak{T}x)\mathcal{T}u \\ &= \lambda^2\rho(x)u + \lambda\rho(\mathfrak{T}x)u + \lambda\rho(x)\mathcal{T}u + \mathcal{T}(\rho(\mathfrak{T}x)u + \rho(x)\mathcal{T}u + \lambda\rho(x)u). \end{aligned} \quad (3)$$

On the other hand,

$$\begin{aligned} & (-\lambda\text{id}_{\mathcal{V}} - \mathcal{T})\left(\rho(-\lambda\text{id}_{\mathfrak{g}} - \mathfrak{T}(x))u + \rho(x)(-\lambda\text{id}_{\mathcal{V}} - \mathcal{T})(u) + \lambda\rho(x)u\right) \\ &= \lambda^2\rho(x)u + \lambda\mathcal{T}(\rho(x)u) + \lambda\rho(\mathfrak{T}x)u + \mathcal{T}(\rho(\mathfrak{T}x)u) + \lambda^2\rho(x)u + \lambda\mathcal{T}(\rho(x)u) + \lambda\rho(x)\mathcal{T}u + \mathcal{T}(\rho(x)\mathcal{T}u) \\ & \quad - \lambda^2\rho(x)u - \lambda\mathcal{T}(\rho(x)u). \end{aligned} \quad (4)$$

The expressions in (3) and (4) are same. Hence $(\mathcal{V}, -\lambda\text{id}_{\mathcal{V}} - \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, -\lambda\text{id}_{\mathfrak{g}} - \mathfrak{T})$. \square

2.9 Proposition. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $\{(\mathcal{V}_i, \mathcal{T}_i)\}_{i \in I}$ be a family of representations of it. Then $(\oplus_{i \in I} \mathcal{V}_i, \oplus_{i \in I} \mathcal{T}_i)$ is also a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$.

Proof. Let $\rho_i : \mathfrak{g} \rightarrow \text{End}(\mathcal{V}_i)$ denote the action of the Lie algebra \mathfrak{g} on the representation \mathcal{V}_i . Then it follows that $\rho : \mathfrak{g} \rightarrow \text{End}(\oplus_{i \in I} \mathcal{V}_i)$, $\rho(x)(u_i)_{i \in I} = (\rho_i(x)u_i)_{i \in I}$ is a representation of \mathfrak{g} on $\oplus_{i \in I} \mathcal{V}_i$. Moreover, for any $x \in \mathfrak{g}$ and $(u_i)_{i \in I} \in \oplus_{i \in I} \mathcal{V}_i$, we have

$$\begin{aligned} \rho(\mathfrak{T}x)(\oplus_{i \in I} \mathcal{T}_i)(u_i)_{i \in I} &= (\rho_i(\mathfrak{T}x)\mathcal{T}_i(u_i))_{i \in I} \\ &= (\mathcal{T}_i(\rho_i(\mathfrak{T}x)u_i + \rho_i(x)\mathcal{T}_i(u_i) + \lambda\rho_i(x)u_i))_{i \in I} \\ &= (\oplus_{i \in I} \mathcal{T}_i)\left(\rho(\mathfrak{T}x)(u_i)_{i \in I} + \rho(x)(\oplus_{i \in I} \mathcal{T}_i)(u_i)_{i \in I} + \lambda\rho(x)(u_i)_{i \in I}\right). \end{aligned}$$

Hence the result follows. \square

Let \mathfrak{g} be a Lie algebra and (\mathcal{V}, ρ) be a representation. Then there is a representation of the Lie algebra \mathfrak{g} on the space $\text{End}(\mathcal{V})$ with the action given by

$$\tilde{\rho} : \mathfrak{g} \rightarrow \text{End}(\text{End}(\mathcal{V})), \quad (\tilde{\rho}(x)f)u = -f(\rho(x)u), \quad \text{for } x \in \mathfrak{g}, f \in \text{End}(\mathcal{V}) \text{ and } u \in \mathcal{V}.$$

With this representation, we have the following.

2.10 Proposition. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then $(\text{End}(\mathcal{V}), \tilde{\mathcal{T}})$ is also a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$, where

$$\tilde{\mathcal{T}} : \text{End}(\mathcal{V}) \rightarrow \text{End}(\mathcal{V}), \quad \tilde{\mathcal{T}}(f)(u) = -\lambda f(u) - f(\mathcal{T}(u)), \quad \text{for } u \in \mathcal{V}.$$

Proof. For any $x, y \in \mathfrak{g}$, $f \in \text{End}(\mathcal{V})$ and $u \in \mathcal{V}$,

$$\left(\tilde{\rho}(\mathfrak{T}x) \tilde{T}(f) \right) u = -\tilde{T}(f)(\rho(\mathfrak{T}x)u) = \lambda f(\rho(\mathfrak{T}x)u) + f(\mathcal{T}(\rho(\mathfrak{T}x)u)). \quad (5)$$

On the other hand,

$$\begin{aligned} & \left(\tilde{T}(\tilde{\rho}(\mathfrak{T}x)f + \tilde{\rho}(x)\tilde{T}(f) + \lambda\tilde{\rho}(x)f) \right) u \\ &= -\lambda(\tilde{\rho}(\mathfrak{T}x)f + \tilde{\rho}(x)\tilde{T}(f) + \lambda\tilde{\rho}(x)f)u - (\tilde{\rho}(\mathfrak{T}x)f + \tilde{\rho}(x)\tilde{T}(f) + \lambda\tilde{\rho}(x)f)(\mathcal{T}u) \\ &= \lambda \left(f(\rho(\mathfrak{T}x)u) + \tilde{T}(f)(\rho(x)u) + \lambda f(\rho(x)u) \right) + \left(f(\rho(\mathfrak{T}x)(\mathcal{T}u)) + \tilde{T}(f)(\rho(x)(\mathcal{T}u)) + \lambda f(\rho(x)(\mathcal{T}u)) \right) \\ &= \lambda \left(f(\rho(\mathfrak{T}x)u) - \cancel{\lambda f(\rho(x)u)} - \cancel{f(\mathcal{T}(\rho(x)u))} + \cancel{\lambda f(\rho(x)u)} \right) \\ &\quad + f \left(\mathcal{T}(\rho(\mathfrak{T}x)u) + \cancel{\mathcal{T}(\rho(x)(\mathcal{T}u))} + \cancel{\mathcal{T}(\rho(x)u)} - \cancel{\lambda \rho(x)(\mathcal{T}u)} - \cancel{\mathcal{T}(\rho(x)(\mathcal{T}u))} + \cancel{\lambda \rho(x)(\mathcal{T}u)} \right) \\ &= \lambda f(\rho(\mathfrak{T}x)u) + f(\mathcal{T}(\rho(\mathfrak{T}x)u)). \end{aligned} \quad (6)$$

It follows from (5) and (6) that $(\text{End}(\mathcal{V}), \tilde{T})$ is a representation of $(\mathfrak{g}, \mathfrak{T})$. \square

In the following, we construct the semidirect product in the context of λ -weighted Rota-Baxter Lie algebras.

2.11 Proposition. *Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then $(\mathfrak{g} \oplus \mathcal{V}, \mathfrak{T} \oplus \mathcal{T})$ is a λ -weighted Rota-Baxter Lie algebra, where the Lie bracket on $\mathfrak{g} \oplus \mathcal{V}$ is given by the semidirect product*

$$[(x, u), (y, v)]_{\ltimes} := ([x, y], \rho(x)v - \rho(y)u), \quad \text{for } x, y \in \mathfrak{g}, u, v \in \mathcal{V}. \quad (7)$$

Proof. We have

$$\begin{aligned} & [(\mathfrak{T} \oplus \mathcal{T})(x, u), (\mathfrak{T} \oplus \mathcal{T})(y, v)]_{\ltimes} \\ &= ([\mathfrak{T}(x), \mathfrak{T}(y)], \rho(\mathfrak{T}x)(\mathcal{T}v) - \rho(\mathfrak{T}y)(\mathcal{T}u)) \\ &= (\mathfrak{T}[\mathfrak{T}(x), y], \mathcal{T}(\rho(\mathfrak{T}x)v - \rho(\mathfrak{T}y)u)) + (\mathfrak{T}[x, \mathfrak{T}(y)], \mathcal{T}(\rho(x)(\mathcal{T}v) - \rho(y)(\mathcal{T}u))) \\ &\quad + \lambda (\mathfrak{T}[x, y], \mathcal{T}(\rho(x)v - \rho(y)u)) \\ &= (\mathfrak{T} \oplus \mathcal{T}) \left([(\mathfrak{T} \oplus \mathcal{T})(x, u), (\mathfrak{T} \oplus \mathcal{T})(y, v)]_{\ltimes} + [(x, u), (\mathfrak{T} \oplus \mathcal{T})(y, v)]_{\ltimes} + \lambda [(x, u), (y, v)]_{\ltimes} \right). \end{aligned}$$

This shows that $\mathfrak{T} \oplus \mathcal{T}$ is a λ -weighted Rota-Baxter operator on the semidirect product Lie algebra. Hence the result follows. \square

2.12 Remark. The converse of the above proposition is also true. More precisely, let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Let \mathcal{V} be a vector space and $\rho : \mathfrak{g} \rightarrow \text{End}(\mathcal{V})$, $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$ be two linear maps. Then $(\mathcal{V} = (\mathcal{V}, \rho), \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ if and only if $(\mathfrak{g} \oplus \mathcal{V}, \mathfrak{T} \oplus \mathcal{T})$ is a λ -weighted Rota-Baxter Lie algebra, where $\mathfrak{g} \oplus \mathcal{V}$ is equipped with the bracket (7).

2.13 Remark. Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. Let $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ be a λ -weighted Rota-Baxter operator on \mathfrak{g} which makes $(\mathfrak{g}, \mathfrak{T})$ a λ -weighted Rota-Baxter Lie algebra. Intuitively, it follows from the above proposition that $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$ can be considered as a representation of \mathfrak{T} with respect to the \mathfrak{g} -representation \mathcal{V} . In Section 5, we call the tuple $(\mathfrak{T}, \mathcal{T})$ as λ -weighted Rota-Baxter paired operators.

2.14 Proposition. *Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Then we have the followings.*

(i) The pair $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{T}})$ is a Lie algebra, where

$$[x, y]_{\mathfrak{T}} := [\mathfrak{T}(x), y] + [x, \mathfrak{T}(y)] + \lambda[x, y], \quad \text{for } x, y \in \mathfrak{g}.$$

We denote this Lie algebra by $\mathfrak{g}_{\mathfrak{T}}$.

(ii) The pair $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$ is a λ -weighted Rota-Baxter Lie algebra and the map $\mathfrak{T} : \mathfrak{g}_{\mathfrak{T}} \rightarrow \mathfrak{g}$ is a morphism of λ -weighted Rota-Baxter Lie algebras.

Proof. (i) This is a standard result that $\mathfrak{g}_{\mathfrak{T}} = (\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{T}})$ is a Lie algebra. See for instance [10, Proposition 5.5]. We also observe that

$$\begin{aligned} [\mathfrak{T}(x), \mathfrak{T}(y)]_{\mathfrak{T}} &= [\mathfrak{T}^2(x), \mathfrak{T}(y)] + [\mathfrak{T}(x), \mathfrak{T}^2(y)] + \lambda[\mathfrak{T}(x), \mathfrak{T}(y)] \\ &= \mathfrak{T}([\mathfrak{T}(x), y]_{\mathfrak{T}} + [x, \mathfrak{T}(y)]_{\mathfrak{T}} + \lambda[x, y]_{\mathfrak{T}}) \end{aligned}$$

which shows that \mathfrak{T} is a λ -weighted Rota-Baxter operator on the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$.

(ii) Since \mathfrak{T} is a λ -weighted Rota-Baxter operator on \mathfrak{g} , it follows from (1) that

$$\mathfrak{T}([x, y]_{\mathfrak{T}}) = [\mathfrak{T}(x), \mathfrak{T}(y)], \quad \text{for } x, y \in \mathfrak{g}_{\mathfrak{T}}.$$

This implies that $\mathfrak{T} : \mathfrak{g}_{\mathfrak{T}} \rightarrow \mathfrak{g}$ is a morphism of λ -weighted Rota-Baxter Lie algebras from $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$ to $(\mathfrak{g}, \mathfrak{T})$. \square

2.15 Theorem. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Define a map $\bar{\rho} : \mathfrak{g} \rightarrow \text{End}(\mathcal{V})$ by

$$\bar{\rho}(x)u := \rho(\mathfrak{T}x)u + \rho(x)(\mathcal{T}u) + \lambda\rho(x)u, \quad \text{for } x \in \mathfrak{g}, u \in \mathcal{V}.$$

Then

(i) $\bar{\rho}$ satisfies $\mathcal{T}(\bar{\rho}(x)u) = \rho(\mathfrak{T}(x))\mathcal{T}(u)$,

(ii) $(\bar{\mathcal{V}} = (\mathcal{V}, \bar{\rho}), \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$.

Proof. The part (i) follows from (2). To prove the part (ii) we first observe that

$$\begin{aligned} &\bar{\rho}(x)\bar{\rho}(y)u - \bar{\rho}(y)\bar{\rho}(x)u \\ &= \rho(\mathfrak{T}x)\bar{\rho}(y)u + \rho(x)\mathcal{T}(\bar{\rho}(y)u) + \lambda\rho(x)\bar{\rho}(y)u - \rho(\mathfrak{T}y)\bar{\rho}(x)u + \rho(y)\mathcal{T}(\bar{\rho}(x)u) + \lambda\rho(y)\bar{\rho}(x)u \\ &= \rho(\mathfrak{T}x)\rho(\mathfrak{T}y)u + \rho(\mathfrak{T}x)\rho(y)\mathcal{T}(u) + \lambda\rho(\mathfrak{T}x)\rho(y)u + \rho(x)\rho(\mathfrak{T}y)\mathcal{T}(u) \\ &\quad + \lambda\rho(x)\rho(\mathfrak{T}y)u + \lambda\rho(x)\rho(y)\mathcal{T}(u) + \lambda^2\rho(x)\rho(y)u \\ &\quad - \rho(\mathfrak{T}y)\rho(\mathfrak{T}x)u - \rho(\mathfrak{T}y)\rho(x)\mathcal{T}(u) - \lambda\rho(\mathfrak{T}y)\rho(x)u - \rho(y)\rho(\mathfrak{T}x)\mathcal{T}(u) \\ &\quad - \lambda\rho(y)\rho(\mathfrak{T}x)u - \lambda\rho(y)\rho(x)\mathcal{T}(u) - \lambda^2\rho(y)\rho(x)u \\ &= \rho([\mathfrak{T}x, \mathfrak{T}y])u + \rho([\mathfrak{T}x, y] + [x, \mathfrak{T}y] + \lambda[x, y])\mathcal{T}(u) + \lambda\rho([\mathfrak{T}x, y] + [x, \mathfrak{T}y] + \lambda[x, y])u \\ &= \rho([\mathfrak{T}x, y]_{\mathfrak{T}})u + \rho([x, y]_{\mathfrak{T}})\mathcal{T}(u) + \lambda\rho([x, y]_{\mathfrak{T}})u \\ &= \bar{\rho}([x, y]_{\mathfrak{T}})u. \end{aligned}$$

This shows that $\bar{\mathcal{V}} = (\mathcal{V}, \bar{\rho})$ is a representation of the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$. Moreover, we have

$$\begin{aligned} \bar{\rho}(\mathfrak{T}x)\mathcal{T}(u) &= \rho(\mathfrak{T}^2(x))\mathcal{T}(u) + \rho(\mathfrak{T}x)\mathcal{T}^2(u) + \lambda\rho(\mathfrak{T}x)\mathcal{T}(u) \\ &= \mathcal{T}(\bar{\rho}(\mathfrak{T}x)u + \bar{\rho}(x)\mathcal{T}(u) + \lambda\bar{\rho}(x)u) \quad (\text{from (i)}). \end{aligned}$$

This shows that $(\bar{\mathcal{V}}, \mathcal{T})$ is a representation of $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$. \square

2.16 Remark. When considering the adjoint representation $(\mathfrak{g}, \mathfrak{T})$ of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$, the representation of the above theorem is the adjoint representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$.

In the following, we will prove another relevant result that will be useful in the next section to construct the cohomology of λ -weighted Rota-Baxter Lie algebras.

2.17 Theorem. Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Define a map $\tilde{\rho} : \mathfrak{g} \rightarrow \text{End}(\mathcal{V})$ by

$$\tilde{\rho}(x)u = \rho(\mathfrak{T}(x))u - \mathcal{T}(\rho(x)u), \quad \text{for } x \in \mathfrak{g}, u \in \mathcal{V}.$$

Then $\tilde{\rho}$ defines a representation of the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$ on \mathcal{V} . Moreover, $(\tilde{\mathcal{V}} = (\mathcal{V}, \tilde{\rho}), \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$.

Proof. For any $x, y \in \mathfrak{g}_{\mathfrak{T}}$ and $u \in \mathcal{V}$, we have

$$\begin{aligned} & \tilde{\rho}(x)\tilde{\rho}(y)u - \tilde{\rho}(y)\tilde{\rho}(x)u \\ &= \tilde{\rho}(x)(\rho(\mathfrak{T}y)u - \mathcal{T}(\rho(y)u)) - \tilde{\rho}(y)(\rho(\mathfrak{T}x)u - \mathcal{T}(\rho(x)u)) \\ &= \rho(\mathfrak{T}x)\rho(\mathfrak{T}y)u - \rho(\mathfrak{T}x)\mathcal{T}(\rho(y)u) - \mathcal{T}(\rho(x)\rho(\mathfrak{T}y)u - \rho(x)\mathcal{T}(\rho(y)u)) \\ &\quad - \rho(\mathfrak{T}y)\rho(\mathfrak{T}x)u + \rho(\mathfrak{T}y)\mathcal{T}(\rho(x)u) + \mathcal{T}(\rho(y)\rho(\mathfrak{T}x)u - \rho(y)\mathcal{T}(\rho(x)u)) \\ &= \rho(\mathfrak{T}x)\rho(\mathfrak{T}y)u - \mathcal{T}\left(\rho(\mathfrak{T}x)\rho(y)u + \underline{\rho(x)\mathcal{T}(\rho(y)u)} + \lambda\rho(x)\rho(y)u\right) - \mathcal{T}\left(\rho(x)\rho(\mathfrak{T}y)u - \underline{\rho(x)\mathcal{T}(\rho(y)u)}\right) \\ &\quad - \rho(\mathfrak{T}y)\rho(\mathfrak{T}x)u + \mathcal{T}\left(\rho(\mathfrak{T}y)\rho(x)u + \underline{\rho(y)\mathcal{T}(\rho(x)u)} + \lambda\rho(y)\rho(x)u\right) + \mathcal{T}\left(\rho(y)\rho(\mathfrak{T}x)u - \underline{\rho(y)\mathcal{T}(\rho(x)u)}\right) \\ &= \rho([\mathfrak{T}x, \mathfrak{T}y])u - \mathcal{T}(\rho([\mathfrak{T}x, y] + [x, \mathfrak{T}y] + \lambda[x, y])u) \\ &= \rho(\mathfrak{T}[x, y]_{\mathfrak{T}})u - \mathcal{T}(\rho([x, y]_{\mathfrak{T}})u) \\ &= \tilde{\rho}([x, y]_{\mathfrak{T}})u. \end{aligned}$$

This shows that $\tilde{\mathcal{V}} = (\mathcal{V}, \tilde{\rho})$ defines a representation of the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$. Moreover, we have

$$\begin{aligned} & \tilde{\rho}(\mathfrak{T}x)(\mathcal{T}u) \\ &= \rho(\mathfrak{T}^2(x))(\mathcal{T}u) - \mathcal{T}(\rho(\mathfrak{T}x)(\mathcal{T}u)) \\ &= \mathcal{T}(\rho(\mathfrak{T}^2(x))u + \rho(\mathfrak{T}x)(\mathcal{T}u) + \lambda\rho(\mathfrak{T}x)u) - \mathcal{T}^2(\rho(\mathfrak{T}x)u + \rho(x)(\mathcal{T}u) + \lambda\rho(x)u) \\ &= \mathcal{T}(\rho(\mathfrak{T}^2(x))u - \mathcal{T}(\rho(\mathfrak{T}x)u)) + \mathcal{T}(\rho(\mathfrak{T}x)(\mathcal{T}u) - \mathcal{T}(\rho(x)(\mathcal{T}u))) + \mathcal{T}(\lambda\rho(\mathfrak{T}x)u - \lambda\mathcal{T}(\rho(x)u)) \\ &= \mathcal{T}(\tilde{\rho}(\mathfrak{T}x)u + \tilde{\rho}(x)(\mathcal{T}u) + \lambda\tilde{\rho}(x)u) \end{aligned}$$

which shows that $(\tilde{\mathcal{V}}, \mathcal{T})$ is a representation of $(\mathfrak{g}_{\mathfrak{T}}, \mathfrak{T})$. □

3 Cohomology of weighted Rota-Baxter Lie algebras

In this section, we first recall the Chevalley-Eilenberg cohomology of a Lie algebra with coefficients in a representation. Then we define the cohomology of a weighted Rota-Baxter Lie algebra with coefficients in a representation. This cohomology is obtained as a byproduct of the Chevalley-Eilenberg cohomology of the underlying Lie algebra with the cohomology of the weighted Rota-Baxter operator. Applications of cohomology to abelian extensions and formal 1-parameter deformations are given in the next section.

Let \mathfrak{g} be a Lie algebra and $\mathcal{V} = (\mathcal{V}, \rho)$ be a representation of it. The Chevalley-Eilenberg cohomology of \mathfrak{g} with coefficients in \mathcal{V} is given by the cohomology of the cochain complex $\{C_{\text{CE}}^*(\mathfrak{g}, \mathcal{V}), \delta_{\text{CE}}\}$, where

$C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) = \text{Hom}(\wedge^n \mathfrak{g}, \mathcal{V})$ for $n \geq 0$ and the coboundary map $\delta_{\text{CE}} : C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \rightarrow C_{\text{CE}}^{n+1}(\mathfrak{g}, \mathcal{V})$ given by

$$\begin{aligned} (\delta_{\text{CE}} f)(x_1, \dots, x_{n+1}) &= \sum_{i=1}^{n+1} (-1)^{i+n} \rho(x_i) f(x_1, \dots, \widehat{x}_i, \dots, x_{n+1}) \\ &\quad + \sum_{1 \leq i < j \leq n+1} (-1)^{i+j+n+1} f([x_i, x_j], x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_{n+1}), \end{aligned}$$

for $f \in C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V})$ and $x_1, \dots, x_{n+1} \in \mathfrak{g}$.

Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then we have seen in Theorem 2.17 that $\tilde{\mathcal{V}} = (\mathcal{V}, \tilde{\rho})$ is a representation of the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$. Therefore, one can define the corresponding Chevalley-Eilenberg cohomology. More precisely, for each $n \geq 0$, we define

$$C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) = \text{Hom}(\wedge^n \mathfrak{g}, \mathcal{V})$$

and a coboundary map $\partial_{\text{CE}} : C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \rightarrow C_{\text{CE}}^{n+1}(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}})$ given by

$$\begin{aligned} (\partial_{\text{CE}} f)(x_1, \dots, x_{n+1}) &= \sum_{i=1}^{n+1} (-1)^{i+n} \tilde{\rho}(x_i) f(x_1, \dots, \widehat{x}_i, \dots, x_{n+1}) \\ &\quad + \sum_{1 \leq i < j \leq n+1} (-1)^{i+j+n+1} f([x_i, x_j]_{\mathfrak{T}}, x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_{n+1}) \\ &= \sum_{i=1}^{n+1} (-1)^{i+n} \rho(\mathfrak{T}(x_i)) f(x_1, \dots, \widehat{x}_i, \dots, x_{n+1}) - \sum_{i=1}^{n+1} (-1)^{i+n} \mathcal{T}(\rho(x_i) f(x_1, \dots, \widehat{x}_i, \dots, x_{n+1})) \\ &\quad + \sum_{1 \leq i < j \leq n+1} (-1)^{i+j+n+1} f([\mathfrak{T}(x_i), x_j] + [x_i, \mathfrak{T}(x_j)] + \lambda[x_i, x_j], x_1, \dots, \widehat{x}_i, \dots, \widehat{x}_j, \dots, x_{n+1}). \end{aligned}$$

Then $\{C_{\text{CE}}^*(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}), \partial_{\text{CE}}\}$ is a cochain complex. The corresponding cohomology groups are called the cohomology of \mathfrak{T} with coefficients in the representation \mathcal{T} .

3.1 Remark. When $(\mathcal{V}, \mathcal{T}) = (\mathfrak{g}, \mathfrak{T})$ is the adjoint representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$, one may consider the cohomology of \mathfrak{T} with coefficients in the representation \mathfrak{T} itself. In [10] the author defines the cohomology of a λ -weighted Rota-Baxter operators motivated from their Maurer-Cartan characterizations. It follows that the cohomology of \mathfrak{T} in the sense of [10] is isomorphic to our cohomology of \mathfrak{T} with coefficients in the representation \mathfrak{T} itself.

We will now in a position to define the cohomology of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the representation $(\mathcal{V}, \mathcal{T})$. We first consider the two cochain complexes, namely,

- ◇ the Chevalley-Eilenberg cochain complex $\{C_{\text{CE}}^*(\mathfrak{g}, \mathcal{V}), \delta_{\text{CE}}\}$ defining the cohomology of the Lie algebra with coefficients in the representation \mathcal{V} ,
- ◇ the complex $\{C_{\text{CE}}^*(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}), \partial_{\text{CE}}\}$ defining the cohomology of \mathfrak{T} with coefficients in the representation \mathcal{T} .

The following result is similar to [30, Proposition 5.1].

3.2 Proposition. *The collection of maps $\{\Phi^n : C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \rightarrow C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}})\}_{n \geq 0}$ defined by*

$$\Phi^0 = \text{id}_{\mathcal{V}}, \quad \text{and}$$

$$\Phi^n(f)(x_1, \dots, x_n) = f(\mathfrak{T}(x_1), \dots, \mathfrak{T}(x_n)) - \sum_{k=0}^{n-1} \lambda^{n-k-1} \sum_{i_1 < \dots < i_k} \mathcal{T} \circ f(x_1, \dots, \mathfrak{T}(x_{i_1}), \dots, \mathfrak{T}(x_{i_k}), \dots, x_n)$$

is a morphism of cochain complexes from $\{C_{\text{CE}}^*(\mathfrak{g}, V), \delta_{\text{CE}}\}$ to $\{C_{\text{CE}}^*(\mathfrak{g}_{\mathfrak{T}}, \tilde{V}), \partial_{\text{CE}}\}$, i.e., $\partial_{\text{CE}} \circ \Phi^n = \Phi^{n+1} \circ \delta_{\text{CE}}$, for $n \geq 0$.

Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. For each $n \geq 0$, we define an abelian group $C_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})$ by

$$C_{\text{RB}}^n(\mathfrak{g}, \mathcal{V}) = \begin{cases} C_{\text{CE}}^0(\mathfrak{g}, \mathcal{V}) = \mathcal{V} & \text{if } n = 0 \\ C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \oplus C_{\text{CE}}^{n-1}(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) = \text{Hom}(\wedge^n \mathfrak{g}, \mathcal{V}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}, \mathcal{V}) & \text{if } n \geq 1 \end{cases}$$

and a map $\delta_{\text{RB}} : C_{\text{RB}}^n(\mathfrak{g}, \mathcal{V}) \rightarrow C_{\text{RB}}^{n+1}(\mathfrak{g}, \mathcal{V})$ by

$$\begin{aligned} \delta_{\text{RB}}(v) &= (\delta_{\text{CE}}(v), -v), \quad \text{for } v \in C_{\text{RB}}^0(\mathfrak{g}, \mathcal{V}) = \mathcal{V}, \\ \delta_{\text{RB}}(f, g) &= (\delta_{\text{CE}}(f), -\partial_{\text{CE}}(g) - \Phi^n(f)), \quad \text{for } (f, g) \in C_{\text{RB}}^n(\mathfrak{g}, \mathcal{V}). \end{aligned}$$

Note that

$$\begin{aligned} (\delta_{\text{RB}})^2(v) &= \delta_{\text{RB}}(\delta_{\text{CE}}(v), -v) \\ &= ((\delta_{\text{CE}})^2(v), \partial_{\text{CE}}(v) - \Phi^2 \circ \delta_{\text{CE}}(v)) = 0 \quad (\because \partial_{\text{CE}} \circ \Phi^0 = \Phi^1 \circ \delta_{\text{CE}}) \end{aligned}$$

and

$$\begin{aligned} (\delta_{\text{RB}})^2(f, g) &= \delta_{\text{RB}}(\delta_{\text{CE}}(f), -\partial_{\text{CE}}(g) - \Phi^n(f)) \\ &= ((\delta_{\text{CE}})^2(f), (\partial_{\text{CE}})^2(g) + \partial_{\text{CE}} \circ \Phi^n(f) - \Phi^{n+1} \circ \delta_{\text{CE}}(f)) = 0 \quad (\because \partial_{\text{CE}} \circ \Phi^n = \Phi^{n+1} \circ \delta_{\text{CE}}). \end{aligned}$$

This shows that $\{C_{\text{RB}}^*(\mathfrak{g}, \mathcal{V}), \delta_{\text{RB}}\}$ is a cochain complex. Let $Z_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})$ and $B_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})$ denote the space of n -cocycles and n -coboundaries, respectively. Then we have $B_{\text{RB}}^n(\mathfrak{g}, \mathcal{V}) \subset Z_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})$, for $n \geq 0$. The corresponding quotients

$$H_{\text{RB}}^n(\mathfrak{g}, \mathcal{V}) := \frac{Z_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})}{B_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})}, \quad \text{for } n \geq 0$$

are called the cohomology of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the representation $(\mathcal{V}, \mathcal{T})$.

Observe that there is a short exact sequence of cochain complexes

$$0 \rightarrow C_{\text{CE}}^{*-1}(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \xrightarrow{i} C_{\text{RB}}^*(\mathfrak{g}, \mathcal{V}) \xrightarrow{p} C_{\text{CE}}^*(\mathfrak{g}, \mathcal{V}) \rightarrow 0$$

given by $i(g) = (0, (-1)^{n-1}g)$ and $p(f, g) = f$, for $(f, g) \in C_{\text{RB}}^n(\mathfrak{g}, \mathcal{V})$. This short exact sequence induces the following long exact sequence on cohomology groups

$$0 \rightarrow H_{\text{RB}}^0(\mathfrak{g}, \mathcal{V}) \rightarrow H_{\text{CE}}^0(\mathfrak{g}, \mathcal{V}) \rightarrow H_{\text{CE}}^0(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \rightarrow H_{\text{RB}}^1(\mathfrak{g}, \mathcal{V}) \rightarrow H_{\text{CE}}^1(\mathfrak{g}, \mathcal{V}) \rightarrow H_{\text{CE}}^1(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \rightarrow \cdots$$

3.3 Proposition. *Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then the cohomology of $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the representation $(\mathcal{V}, \mathcal{T})$ is isomorphic to the cohomology of $(\mathfrak{g}, -\text{lid}_{\mathfrak{g}} - \mathfrak{T})$ with coefficients in the representation $(\mathcal{V}, -\text{lid}_{\mathcal{V}} - \mathcal{T})$.*

Proof. Let $\mathfrak{T}' = -\text{lid}_{\mathfrak{g}} - \mathfrak{T}$ and $\mathcal{T}' = -\text{lid}_{\mathcal{V}} - \mathcal{T}$. Then it is easy to see that the Lie algebra $\mathfrak{g}_{\mathfrak{T}'}$ of Proposition 2.14 (i) is negative to the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$. Moreover, the representation $\tilde{\rho}'$ of the Lie algebra $\mathfrak{g}_{\mathfrak{T}'}$ on \mathcal{V} (we denote this representation by $\tilde{\rho}'$) as of Theorem 2.17 is given by the negative of the representation $\tilde{\rho}$ of the Lie algebra $\mathfrak{g}_{\mathfrak{T}}$ on \mathcal{V} . Hence the corresponding Chevalley-Eilenberg differentials

$$\partial_{\text{CE}} : C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \rightarrow C_{\text{CE}}^{n+1}(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \quad \text{and} \quad \partial'_{\text{CE}} : C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}'}, \tilde{\mathcal{V}}') \rightarrow C_{\text{CE}}^{n+1}(\mathfrak{g}_{\mathfrak{T}'}, \tilde{\mathcal{V}}')$$

are related by $\partial'_{\text{CE}} f = -\partial_{\text{CE}} f$. Finally, if $(\Phi')^n : C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \rightarrow C_{\text{CE}}^n(\mathfrak{g}_{\mathfrak{T}'}, \tilde{\mathcal{V}}')$ be the map as of Proposition 3.2 replacing \mathfrak{T} by \mathfrak{T}' and replacing \mathcal{T} by \mathcal{T}' , then we have $(\Phi')^n(f) = (-1)^n \Phi^n(f)$.

For each $n \geq 0$, define an isomorphism of vector spaces

$$\Xi^n : C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \oplus C_{\text{CE}}^{n-1}(\mathfrak{g}_{\mathfrak{T}}, \tilde{\mathcal{V}}) \rightarrow C_{\text{CE}}^n(\mathfrak{g}, \mathcal{V}) \oplus C_{\text{CE}}^{n-1}(\mathfrak{g}_{\mathfrak{T}'}, \tilde{\mathcal{V}}') \quad \text{by} \quad \Xi^n(f, g) = (f, (-1)^{n-1}g).$$

Moreover,

$$\begin{aligned} (\delta'_{\text{RB}} \circ \Xi^n)(f, g) &= \delta'_{\text{RB}}(f, (-1)^{n-1}g) = (\delta_{\text{CE}}f, (-1)^n \partial'_{\text{CE}}g - (\Phi')^n(f)) \\ &= (\delta_{\text{CE}}f, (-1)^{n+1} \partial_{\text{CE}}g - (-1)^n \Phi^n(f)) \\ &= (\Xi^{n+1} \circ \delta_{\text{RB}})(f, g). \end{aligned}$$

This shows that the collection of maps $\{\Xi^n\}_{n \geq 0}$ commute with the respective coboundary maps. Hence they induce isomorphism on cohomology. \square

3.1 H^0 and H^1

Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. An element $v \in \mathcal{V}$ is in $Z_{\text{RB}}^0(\mathfrak{g}, \mathcal{V})$ if and only if $(\delta_{\text{CE}}(v), -v) = 0$. This holds only when $v = 0$. Therefore, it follows from the definition that $H_{\text{RB}}^0(\mathfrak{g}, \mathcal{V}) = 0$.

A pair $(\gamma, v) \in \text{Hom}(\mathfrak{g}, \mathcal{V}) \oplus \mathcal{V}$ is said to be a derivation on the weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the representation $(\mathcal{V}, \mathcal{T})$ if they satisfies

$$\begin{aligned} \gamma([x, y]) &= \rho(x)(\gamma(y)) - \rho(y)(\gamma(x)), \\ \gamma(\mathfrak{T}(x)) - \mathcal{T}(\gamma(x)) &= \mathcal{T}(\rho(x)v) - \rho(\mathfrak{T}(x))v, \quad \text{for } x, y \in \mathfrak{g}. \end{aligned}$$

It follows from the first condition that γ is a derivation on the Lie algebra \mathfrak{g} with coefficients in \mathcal{V} . The second condition says that the obstruction of vanishing $\gamma \circ \mathfrak{T} - \mathcal{T} \circ \gamma$ is measured by the presence of v . We denote the set of all derivations by $\text{Der}(\mathfrak{g}, \mathcal{V})$.

A derivation is said to be inner if it is of the form $(-\delta_{\text{CE}}(v), v)$, for some $v \in \mathcal{V}$. The set of all inner derivations are denoted by $\text{InnDer}(\mathfrak{g}, \mathcal{V})$.

It follows from the definition that $Z_{\text{RB}}^1(\mathfrak{g}, \mathcal{V}) = \text{Der}(\mathfrak{g}, \mathcal{V})$ and $B_{\text{RB}}^1(\mathfrak{g}, \mathcal{V}) = \text{InnDer}(\mathfrak{g}, \mathcal{V})$. Hence we have $H_{\text{RB}}^1(\mathfrak{g}, \mathcal{V}) = \frac{\text{Der}(\mathfrak{g}, \mathcal{V})}{\text{InnDer}(\mathfrak{g}, \mathcal{V})}$, the space of outer derivations.

3.2 Relation with the cohomology of weighted Rota-Baxter associative algebras

In [30] Wang and Zhou defined the cohomology of a weighted Rota-Baxter associative algebra with coefficients in a Rota-Baxter bimodule. In this subsection, we show that our cohomology is related to the cohomology of [30] by suitable skew-symmetrization.

3.4 Definition. (i) A λ -weighted Rota-Baxter associative algebra is a pair $(\mathfrak{A}, \mathfrak{R})$ in which \mathfrak{A} is an associative algebra and $\mathfrak{R} : \mathfrak{A} \rightarrow \mathfrak{A}$ is a linear map satisfying

$$\mathfrak{R}(a) \cdot \mathfrak{R}(b) = \mathfrak{R}(\mathfrak{R}(a) \cdot b + a \cdot \mathfrak{R}(b) + \lambda a \cdot b), \quad \text{for } a, b \in \mathfrak{A}.$$

(ii) Let $(\mathfrak{A}, \mathfrak{R})$ be a λ -weighted Rota-Baxter associative algebra. A Rota-Baxter bimodule over it consists of a pair $(\mathcal{M}, \mathcal{R})$ in which \mathcal{M} is an \mathfrak{A} -bimodule (denote both left and right actions by cdot) and

$\mathcal{R} : \mathcal{M} \rightarrow \mathcal{M}$ is a linear map satisfying for $a \in \mathfrak{A}$, $m \in \mathcal{M}$,

$$\begin{aligned}\mathfrak{R}(a) \cdot \mathcal{R}(m) &= \mathcal{R}(\mathfrak{R}(a) \cdot m + a \cdot \mathcal{R}(m) + \lambda a \cdot m), \\ \mathcal{R}(m) \cdot \mathfrak{R}(a) &= \mathcal{R}(\mathcal{R}(m) \cdot a + m \cdot \mathfrak{R}(a) + \lambda m \cdot a).\end{aligned}$$

3.5 Remark. It follows from the above definition that any λ -weighted Rota-Baxter associative algebra $(\mathfrak{A}, \mathfrak{R})$ is a Rota-Baxter bimodule over itself. This is called the adjoint Rota-Baxter bimodule.

The following result is straightforward.

3.6 Proposition. *Let $(\mathfrak{A}, \mathfrak{R})$ be a λ -weighted Rota-Baxter associative algebra. Then $(\mathfrak{A}_c, \mathfrak{R})$ is a λ -weighted Rota-Baxter Lie algebra, where \mathfrak{A}_c is the vector space \mathfrak{A} with the commutator Lie bracket*

$$[a, b]_c = a \cdot b - b \cdot a, \quad \text{for } a, b \in \mathfrak{A}_c.$$

(This is called the skew-symmetrization). Moreover, if $(\mathcal{M}, \mathcal{R})$ is a Rota-Baxter bimodule over $(\mathfrak{A}, \mathfrak{R})$, then $(\mathcal{M}_c, \mathcal{R})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{A}_c, \mathfrak{R})$, where $\mathcal{M}_c = \mathcal{M}$ as a vector space and the representation of the Lie algebra \mathfrak{A}_c on \mathcal{M}_c is given by $\rho(a)(m) = a \cdot m - m \cdot a$, for $a \in \mathfrak{A}_c$ and $m \in \mathcal{M}_c$.

Let $(\mathfrak{A}, \mathfrak{R})$ be a λ -weighted Rota-Baxter associative algebra. Then it is known that $(\mathfrak{A}, *_{\mathfrak{R}})$ is an associative algebra [12], where

$$a *_{\mathfrak{R}} b = \mathfrak{R}(a) \cdot b + a \cdot \mathfrak{R}(b) + \lambda a \cdot b, \quad \text{for } a, b \in \mathfrak{A}.$$

We denote this associative algebra by $\mathfrak{A}_{\mathfrak{R}}$. Moreover, if $(\mathcal{M}, \mathcal{R})$ is a Rota-Baxter bimodule over the λ -weighted Rota-Baxter associative algebra $(\mathfrak{A}, \mathfrak{R})$, then it has been observed in [30] that \mathcal{M} carries a bimodule structure over the associative algebra $\mathfrak{A}_{\mathfrak{R}}$ with left and right actions

$$a \cdot m = \mathfrak{R}(a) \cdot m - \mathcal{R}(a \cdot m) \quad \text{and} \quad m \cdot a = m \cdot \mathfrak{R}(a) - \mathcal{R}(m \cdot a), \quad \text{for } a \in \mathfrak{A}_{\mathfrak{R}} \text{ and } m \in \mathcal{M}.$$

Denote this $\mathfrak{A}_{\mathfrak{R}}$ -bimodule by $\widetilde{\mathcal{M}}$. To define the cohomology of the λ -weighted Rota-Baxter associative algebra $(\mathfrak{A}, \mathfrak{R})$ with coefficients in the Rota-Baxter bimodule $(\mathcal{M}, \mathcal{R})$, the authors in [30] considered two Hochschild cochain complexes, namely $\{C_{\mathfrak{H}}^*(\mathfrak{A}, \mathcal{M}), \delta_{\mathfrak{H}}\}$ and $\{C_{\mathfrak{H}}^*(\mathfrak{A}_{\mathfrak{R}}, \widetilde{\mathcal{M}}), \partial_{\mathfrak{H}}\}$. The first one is the Hochschild complex of the given algebra \mathfrak{A} with coefficients in the bimodule \mathcal{M} , whereas the second one is the Hochschild complex of the algebra $\mathfrak{A}_{\mathfrak{R}}$ with coefficients in the bimodule $\widetilde{\mathcal{M}}$. They proved that the collection $\{\Psi^n : C_{\mathfrak{H}}^n(\mathfrak{A}, \mathcal{M}) \rightarrow C_{\mathfrak{H}}^n(\mathfrak{A}_{\mathfrak{R}}, \widetilde{\mathcal{M}})\}_{n \geq 0}$ of maps given by

$$\Psi^0 = \text{id}_{\mathcal{M}}, \quad \text{and}$$

$$\Psi^n(f)(a_1, \dots, a_n) = f(\mathfrak{R}(a_1), \dots, \mathfrak{R}(a_n)) - \sum_{k=0}^{n-1} \lambda^{n-k-1} \sum_{i_1 < \dots < i_k} \mathcal{R} \circ f(a_1, \dots, \mathfrak{R}(a_{i_1}), \dots, \mathfrak{R}(a_{i_k}), \dots, a_n)$$

defines a morphism of cochain complexes. This allows the authors to consider the cochain complex $\{C_{\text{RB}}^*(\mathfrak{A}, \mathcal{M}), \delta_{\text{RB}}^{\text{ass}}\}$, where

$$C_{\text{RB}}^0(\mathfrak{A}, \mathcal{M}) = C_{\mathfrak{H}}^0(\mathfrak{A}, \mathcal{M}) = \mathcal{M} \quad \text{and} \quad C_{\text{RB}}^n(\mathfrak{A}, \mathcal{M}) = C_{\mathfrak{H}}^n(\mathfrak{A}, \mathcal{M}) \oplus C_{\mathfrak{H}}^{n-1}(\mathfrak{A}_{\mathfrak{R}}, \widetilde{\mathcal{M}}), \quad \text{for } n \geq 1$$

and $\delta_{\text{RB}}^{\text{ass}} : C_{\text{RB}}^n(\mathfrak{A}, \mathcal{M}) \rightarrow C_{\text{RB}}^{n+1}(\mathfrak{A}, \mathcal{M})$ given by

$$\delta_{\text{RB}}^{\text{ass}} = (\delta_{\mathfrak{H}}(f), \quad -\partial_{\mathfrak{H}}(g) - \Psi^n(f)), \quad \text{for } (f, g) \in C_{\text{RB}}^n(\mathfrak{A}, \mathcal{M}).$$

The corresponding cohomology groups are called the cohomology of the λ -weighted Rota-Baxter associative algebra $(\mathfrak{A}, \mathfrak{R})$ with coefficients in the Rota-Baxter bimodule $(\mathcal{M}, \mathcal{R})$.

To find the connection between the cohomology of a weighted Rota-Baxter associative algebra and the cohomology of the corresponding skew-symmetrized weighted Rota-Baxter Lie algebra, we need the following result.

3.7 Proposition. *Let $(\mathfrak{A}, \mathfrak{R})$ be a λ -weighted Rota-Baxter associative algebra and $(\mathcal{M}, \mathcal{R})$ be a Rota-Baxter bimodule. Then $(\mathfrak{A}_c)_{\mathfrak{R}} = (\mathfrak{A}_{\mathfrak{R}})_c$ as a Lie algebra. Here $(\mathfrak{A}_c)_{\mathfrak{R}}$ is the Lie algebra structure on \mathfrak{A} induced by the λ -weighted Rota-Baxter operator \mathfrak{R} on the Lie algebra \mathfrak{A}_c as in Proposition 2.14 (i). Moreover, the representation of the Lie algebra $(\mathfrak{A}_c)_{\mathfrak{R}}$ on $\widetilde{\mathcal{M}}_c$ and the representation of the Lie algebra $(\mathfrak{A}_{\mathfrak{R}})_c$ on $(\widetilde{\mathcal{M}})_c$ coincide.*

It is known that the standard skew-symmetrization gives rise to a morphism from the Hochschild cochain complex of an associative algebra to the Chevalley-Eilenberg cochain complex of the corresponding skew-symmetrized Lie algebra [20]. Hence the following diagrams commute

$$\begin{array}{ccc} C_H^n(\mathfrak{A}, \mathcal{M}) & \xrightarrow{\delta_H} & C_H^{n+1}(\mathfrak{A}, \mathcal{M}) \\ S_n \downarrow & & \downarrow S_{n+1} \\ C_{CE}^n(\mathfrak{A}_c, \mathcal{M}_c) & \xrightarrow{\delta_{CE}} & C_{CE}^{n+1}(\mathfrak{A}_c, \mathcal{M}_c) \end{array} \quad \begin{array}{ccc} C_H^n(\mathfrak{A}_{\mathfrak{R}}, \widetilde{\mathcal{M}}) & \xrightarrow{\partial_H} & C_H^{n+1}(\mathfrak{A}_{\mathfrak{R}}, \widetilde{\mathcal{M}}) \\ S_n \downarrow & & \downarrow S_{n+1} \\ C_{CE}^n((\mathfrak{A}_{\mathfrak{R}})_c, \widetilde{\mathcal{M}}_c) & \xrightarrow{\partial_{CE}} & C_{CE}^{n+1}((\mathfrak{A}_{\mathfrak{R}})_c, \widetilde{\mathcal{M}}_c). \end{array}$$

Here S_* are the skew-symmetrization maps. As a consequence, we get the following.

3.8 Theorem. *Let $(\mathfrak{A}, \mathfrak{R})$ be a λ -weighted Rota-Baxter associative algebra and $(\mathcal{M}, \mathcal{R})$ be a Rota-Baxter bimodule. Then the collection of maps*

$$\mathbb{S}_n : C_{RB}^n(\mathfrak{A}, \mathcal{M}) \rightarrow C_{RB}^n(\mathfrak{A}_c, \mathcal{M}_c), \quad \text{for } n \geq 0$$

defined by $\mathbb{S}_0 = \text{id}_{\mathcal{M}}$ and $\mathbb{S}_n = (S_n, S_{n-1})$ for $n \geq 1$, induces a morphism from the cohomology of $(\mathfrak{A}, \mathfrak{R})$ with coefficients in the Rota-Baxter bimodule $(\mathcal{M}, \mathcal{R})$ to the cohomology of $(\mathfrak{A}_c, \mathfrak{R})$ with coefficients in the representation $(\mathcal{M}_c, \mathcal{R})$.

Proof. We only need to check that the maps $\{\mathbb{S}_n\}_{n \geq 0}$ commute with corresponding coboundary maps. For $(f, g) \in C_{RB}^n(\mathfrak{A}, \mathcal{M})$,

$$\begin{aligned} (\delta_{RB} \circ \mathbb{S}_n)(f, g) &= \delta_{RB}(S_n f, S_{n-1} g) \\ &= (\delta_{CE} \circ S_n(f), -\partial_{CE} \circ S_{n-1}(g) - (\Phi^n \circ S_n)(f)) \\ &= (S_{n+1} \circ \delta_H(f), -S_n \circ \partial_H(g) - S_n \circ \Psi^n(f)) \quad (\because \Phi^n \circ S_n = S_n \circ \Psi^n) \\ &= \mathbb{S}_{n+1}(\delta_H(f), -\partial_H(g) - \Psi^n(f)) = (\mathbb{S}_{n+1} \circ \delta_{RB}^{ss})(f, g). \end{aligned}$$

Hence the result follows. \square

4 Applications of cohomology

In this section, we study abelian extensions and formal 1-parameter deformations of weighted Rota-Baxter Lie algebras in terms of cohomology.

4.1 Abelian extensions and H^2

Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Let $(\mathcal{V}, \mathcal{T})$ be a pair of a vector space \mathcal{V} and a linear map $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$. Note that $(\mathcal{V}, \mathcal{T})$ can be considered as a λ -weighted Rota-Baxter Lie algebra where the Lie bracket on \mathcal{V} is assumed to be trivial.

4.1 Definition. An abelian extension of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$ is a short exact sequence of morphisms of λ -weighted Rota-Baxter Lie algebras

$$0 \longrightarrow (\mathcal{V}, \mathcal{T}) \xrightarrow{i} (\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}}) \xrightarrow{p} (\mathfrak{g}, \mathfrak{T}) \longrightarrow 0. \quad (8)$$

In this case, we say that $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ is an abelian extension of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$.

4.2 Definition. Let $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ and $(\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}')$ be two abelian extensions of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$. They are said to be isomorphic if there exists an isomorphism $\phi : (\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}}) \rightarrow (\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}')$ of λ -weighted Rota-Baxter Lie algebras which makes the following diagram commutative

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\mathcal{V}, \mathcal{T}) & \xrightarrow{i} & (\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}}) & \xrightarrow{p} & (\mathfrak{g}, \mathfrak{T}) \longrightarrow 0 \\ & & \parallel & & \downarrow \phi & & \parallel \\ 0 & \longrightarrow & (\mathcal{V}, \mathcal{T}) & \xrightarrow{i'} & (\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}') & \xrightarrow{p'} & (\mathfrak{g}, \mathfrak{T}) \longrightarrow 0 \end{array} \quad (9)$$

Let $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ be an abelian extension of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$ as of (8). A section of the map p is a linear map $s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ satisfying $p \circ s = \text{id}_{\mathfrak{g}}$. Note that a section of p always exists.

Let $s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ be a section of the map p . We define a linear map $\rho : \mathfrak{g} \rightarrow \text{End}(\mathcal{V})$ by $\rho(x)(u) := [s(x), i(u)]_{\widehat{\mathfrak{g}}}$, for $x \in \mathfrak{g}$ and $u \in \mathcal{V}$. Then it can be easily check that $\mathcal{V} = (\mathcal{V}, \rho)$ is a representation of the Lie algebra \mathfrak{g} . More generally, $(\mathcal{V}, \mathcal{T})$ is a representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$. One can easily check that this representation doesn't depend on the choice of the section s . We call this as the induced representation on $(\mathcal{V}, \mathcal{T})$.

Suppose $(\mathcal{V}, \mathcal{T})$ is a given representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$. We denote by $\text{Ext}(\mathfrak{g}, \mathcal{V})$ the isomorphism classes of abelian extensions of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$ for which the induced representation on $(\mathcal{V}, \mathcal{T})$ is the prescribed one.

4.3 Theorem. Let $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ be a λ -weighted Rota-Baxter Lie algebra and $(\mathcal{V}, \mathcal{T})$ be a representation of it. Then there is a one-to-one correspondence between $\text{Ext}(\mathfrak{g}, \mathcal{V})$ and the second cohomology group $H_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$.

Proof. Let $(\psi, \chi) \in Z_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$ be a 2-cocycle, i.e., we have $\delta_{\text{CE}}\psi = 0$ and $-\partial_{\text{CE}}(\chi) - \Phi^2(\psi) = 0$. Consider the space $\mathfrak{g} \oplus \mathcal{V}$ with the bracket

$$[(x, u), (y, v)] = ([x, y], \rho(x)v - \rho(y)u + \psi(x, y)).$$

Since $\delta_{\text{CE}}\psi = 0$, it follows that the above bracket makes $\mathfrak{g} \oplus \mathcal{V}$ into a Lie algebra. We denote this Lie algebra by $\widehat{\mathfrak{g}}$. We also define a map $\widehat{\mathfrak{T}} : \widehat{\mathfrak{g}} \rightarrow \widehat{\mathfrak{g}}$ by $\widehat{\mathfrak{T}}(x, u) = (\mathfrak{T}(x), \mathcal{T}(u) + \chi(x))$, for $(x, u) \in \widehat{\mathfrak{g}}$. Since $-\partial_{\text{CE}}(\chi) - \Phi^2(\psi) = 0$, it follows that $\widehat{\mathfrak{T}}$ is a λ -weighted Rota-Baxter operator on the Lie algebra $\widehat{\mathfrak{g}}$. In other words, $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ is a λ -weighted Rota-Baxter Lie algebra. Moreover, the short exact sequence

$$0 \longrightarrow (\mathcal{V}, \mathcal{T}) \xrightarrow{i} (\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}}) \xrightarrow{p} (\mathfrak{g}, \mathfrak{T}) \longrightarrow 0$$

defines an abelian extension of $(\mathfrak{g}, \mathfrak{T})$ by $(\mathcal{V}, \mathcal{T})$, where $i(u) = (0, u)$ and $p(x, u) = x$, for $x \in \mathfrak{g}$ and $u \in \mathcal{V}$. Note that the canonical section $s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$, $s(x) = (x, 0)$ induces $(\mathcal{V}, \mathcal{T})$ with the original representation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$.

Let $(\psi', \chi') \in Z_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$ be another 2-cocycle cohomologous to (ψ, χ) . Then there exists a pair $(\gamma, v) \in \text{Hom}(\mathfrak{g}, \mathcal{V}) \oplus \mathcal{V}$ such that

$$(\psi, \chi) - (\psi', \chi') = (\delta_{\text{CE}}(\gamma), -\partial_{\text{CE}}v - \Phi^1(\gamma)).$$

Consider the map $\phi : \mathfrak{g} \oplus \mathcal{V} \rightarrow \mathfrak{g} \oplus \mathcal{V}$ given by

$$\phi(x, u) = (x, u - \gamma(x) - \partial_{\text{CE}}(v)(x)).$$

Then it can be easily checked that ϕ defines an isomorphism of abelian extensions from $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ to $(\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}')$. Therefore, there is a well-defined map $\Upsilon : H_{\text{RB}}^2(\mathfrak{g}, \mathcal{V}) \rightarrow \text{Hom}(\mathfrak{g}, \mathcal{V})$.

Conversely, let $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ be an abelian extension given by (8). Let $s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ be a section of the map p . We define elements $\psi \in \text{Hom}(\wedge^2 \mathfrak{g}, \mathcal{V})$ and $\chi \in \text{Hom}(\mathfrak{g}, \mathcal{V})$ by

$$\psi(x, y) = [s(x), s(y)]_{\widehat{\mathfrak{g}}} - s[x, y] \quad \text{and} \quad \chi(x) = \widehat{\mathfrak{T}}(s(x)) - s(\mathfrak{T}(x)), \quad \text{for } x, y \in \mathfrak{g}.$$

Then it follows from a straightforward computation that the pair (ψ, χ) defines a 2-cocycle in $Z_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$. Moreover, the corresponding cohomology class in $H_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$ doesn't depend on the choice of the section s .

Let $(\widehat{\mathfrak{g}}, \widehat{\mathfrak{T}})$ and $(\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}')$ be two isomorphic abelian extensions as of (9). Let $s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ be a section of the map p . Then we have

$$p' \circ (\phi \circ s) = p \circ s = \text{id}_{\mathfrak{g}}.$$

This shows that $s' = \phi \circ s : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}'$ is a section of p' . If (ψ', χ') denotes the 2-cocycle in $Z_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$ corresponding to the abelian extension $(\widehat{\mathfrak{g}}', \widehat{\mathfrak{T}}')$ and section $s' : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}'$ of the map p' , then

$$\begin{aligned} \psi'(x, y) &= [s'(x), s'(y)]_{\widehat{\mathfrak{g}}'} - s'[x, y] \\ &= [\phi \circ s(x), \phi \circ s(y)]_{\widehat{\mathfrak{g}}'} - \phi \circ s[x, y] \\ &= \phi([s(x), s(y)]_{\widehat{\mathfrak{g}}} - s[x, y]) \\ &= \phi(\psi(x, y)) = \phi(x, y) \quad (\because \phi|_{\mathcal{V}} = \text{id}_{\mathcal{V}}) \end{aligned}$$

and

$$\begin{aligned} \chi'(x) &= \widehat{\mathfrak{T}}'(s'(x)) - s'(\mathfrak{T}(x)) \\ &= \widehat{\mathfrak{T}}'(\phi \circ s(x)) - \phi \circ s(\mathfrak{T}(x)) \\ &= \phi(\widehat{\mathfrak{T}}(s(x)) - s(\mathfrak{T}(x))) = \phi(\chi(x)) = \chi(x) \quad (\because \phi|_{\mathcal{V}} = \text{id}_{\mathcal{V}}). \end{aligned}$$

This show that (ψ, χ) and (ψ', χ') correspond to the same elemwnt in $H_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$. Therefore, we have a well-defined map $\Omega : \text{Ext}(\mathfrak{g}, \mathcal{V}) \rightarrow H_{\text{RB}}^2(\mathfrak{g}, \mathcal{V})$. Finally, the map Υ and Ω are inverses to each other. This completes the proof. \square

4.2 Formal deformations

In this subsection, we will consider formal deformations of a λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$. Here we will simultaneously deform the Lie bracket on \mathfrak{g} and the λ -weighted Rota-Baxter operator \mathfrak{T} . We find the relation between such deformations and cohomology of $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the adjoint representation.

Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. Let $\mu \in C_{\text{CE}}^2(\mathfrak{g}, \mathfrak{g}) = \text{Hom}(\wedge^2 \mathfrak{g}, \mathfrak{g})$ be the element that corresponds to the Lie bracket on \mathfrak{g} , i.e., $\mu(x, y) = [x, y]$, for $x, y \in \mathfrak{g}$. Consider the space $\mathfrak{g}[[t]]$ of formal power series in t with coefficients from \mathfrak{g} . Then $\mathfrak{g}[[t]]$ is a $\mathbf{k}[[t]]$ -module.

4.4 Definition. A formal 1-parameter deformation of $(\mathfrak{g}, \mathfrak{T})$ consists of a pair (μ_t, \mathfrak{T}_t) of two formal power

series

$$\begin{aligned}\mu_t &= \sum_{i \geq 0} \mu_i t^i, \quad \text{where } \mu_i \in \text{Hom}(\wedge^2 \mathfrak{g}, \mathfrak{g}) \text{ with } \mu_0 = \mu, \\ \mathfrak{T}_t &= \sum_{i \geq 0} \mathfrak{T}_i t^i, \quad \text{where } \mathfrak{T}_i \in \text{Hom}(\mathfrak{g}, \mathfrak{g}) \text{ with } \mathfrak{T}_0 = \mathfrak{T}\end{aligned}$$

such that the $\mathbf{k}[[t]]$ -bilinear map μ_t defines a Lie algebra structure on $\mathfrak{g}[[t]]$ and the $\mathbf{k}[[t]]$ -linear map $\mathfrak{T}_t : \mathfrak{g}[[t]] \rightarrow \mathfrak{g}[[t]]$ is a λ -weighted Rota-Baxter operator. In other words, $(\mathfrak{g}[[t]] = (\mathfrak{g}[[t]], \mu_t), \mathfrak{T}_t)$ is a λ -weighted Rota-Baxter Lie algebra over $\mathbf{k}[[t]]$.

Thus, (μ_t, \mathfrak{T}_t) is a formal 1-parameter deformation of $(\mathfrak{g}, \mathfrak{T})$ if and only if

$$\begin{aligned}\mu_t(x, \mu_t(y, z)) + \mu_t(y, \mu_t(z, x)) + \mu_t(z, \mu_t(x, y)) &= 0, \\ \mu_t(\mathfrak{T}_t(x), \mathfrak{T}_t(y)) &= \mathfrak{T}_t(\mu_t(\mathfrak{T}_t(x), y) + \mu_t(x, \mathfrak{T}_t(y)) + \lambda \mu_t(x, y)),\end{aligned}$$

for $x, y, z \in \mathfrak{g}$. They are equivalent to the following two systems of identities: for $n \geq 0$ and $x, y, z \in \mathfrak{g}$,

$$\sum_{i+j=n} \mu_i(x, \mu_j(y, z)) + \mu_i(y, \mu_j(z, x)) + \mu_i(z, \mu_j(x, y)) = 0, \quad (10)$$

$$\sum_{i+j+k=n} \mu_i(\mathfrak{T}_j(x), \mathfrak{T}_k(y)) = \sum_{i+j+k=n} \mathfrak{T}_i(\mu_j(\mathfrak{T}_k(x), y) + \mu_j(x, \mathfrak{T}_k(y))) + \sum_{i+j=n} \lambda \mathfrak{T}_i(\mu_j(x, y)). \quad (11)$$

4.5 Definition. Two formal deformations (μ_t, \mathfrak{T}_t) and $(\mu'_t, \mathfrak{T}'_t)$ of a λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ are said to be equivalent if there is a formal isomorphism

$$\phi_t = \sum_{i \geq 0} \phi_i t^i : \mathfrak{g}[[t]] \rightarrow \mathfrak{g}[[t]], \quad \text{where } \phi_i \in \text{Hom}(\mathfrak{g}, \mathfrak{g}) \text{ with } \phi_0 = \text{id}_{\mathfrak{g}}$$

such that the $\mathbf{k}[[t]]$ -linear map ϕ_t is a morphism of λ -weighted Rota-Baxter Lie algebras from $(\mathfrak{g}[[t]]' = (\mathfrak{g}[[t]], \mu'_t), \mathfrak{T}'_t)$ to $(\mathfrak{g}[[t]] = (\mathfrak{g}[[t]], \mu_t), \mathfrak{T}_t)$.

Therefore, (μ_t, \mathfrak{T}_t) and $(\mu'_t, \mathfrak{T}'_t)$ are equivalent if the followings are hold:

$$\phi_t(\mu'_t(x, y)) = \mu_t(\phi_t(x), \phi_t(y)) \quad \text{and} \quad \phi_t \circ \mathfrak{T}'_t = \mathfrak{T}_t \circ \phi_t.$$

They can be equivalently expressed by the following system of equations: for $n \geq 0$,

$$\sum_{i+j=n} \phi_i(\mu'_j(x, y)) = \sum_{i+j+k=n} \mu_i(\phi_j(x), \phi_k(x)), \quad (12)$$

$$\sum_{i+j=n} \phi_i \circ \mathfrak{T}'_j(x) = \sum_{i+j=n} \mathfrak{T}_i \circ \phi_j(x). \quad (13)$$

4.6 Theorem. Let (μ_t, \mathfrak{T}_t) be a formal 1-parameter deformation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$. Then $(\mu_1, \mathfrak{T}_1) \in C_{\text{RB}}^2(\mathfrak{g}, \mathfrak{g})$ is a 2-cocycle in the cohomology of $(\mathfrak{g}, \mathfrak{T})$ with coefficients in the adjoint representation. Moreover, the corresponding cohomology class in $H_{\text{RB}}^2(\mathfrak{g}, \mathfrak{g})$ depends only on the equivalence class of the deformation (μ_t, \mathfrak{T}_t) .

Proof. Since (μ_t, \mathfrak{T}_t) is a formal 1-parameter deformation, we get from (10) and (11) for $n = 1$ that

$$[x, \mu_1(y, z)] + \mu_1(x, [y, z]) + [y, \mu_1(z, x)] + \mu_1(y, [z, x]) + [z, \mu_1(x, y)] + \mu_1(z, [x, y]) = 0$$

and

$$\begin{aligned} & \mu_1(\mathfrak{T}(x), \mathfrak{T}(y)) + [\mathfrak{T}_1(x), \mathfrak{T}(y)] + [\mathfrak{T}(x), \mathfrak{T}_1(y)] \\ &= \mathfrak{T}_1([\mathfrak{T}(x), y] + [x, \mathfrak{T}(y)]) + \mathfrak{T}([\mathfrak{T}_1(x), y] + [x, \mathfrak{T}_1(y)] + \mu_1(\mathfrak{T}(x), y) + \mu_1(x, \mathfrak{T}(y))) \\ &+ \lambda \mathfrak{T}(\mu_1(x, y)) + \lambda \mathfrak{T}_1([x, y]). \end{aligned}$$

The first identity is equivalent to $\delta_{\text{CE}}\mu_1 = 0$ while the second identity is equivalent to $-\partial_{\text{CE}}(\mathfrak{T}_1) - \Phi^2(\mu_1) = 0$. This implies that

$$\delta_{\text{RB}}(\mu_1, \mathfrak{T}_1) = (\delta_{\text{CE}}\mu_1, -\partial_{\text{CE}}(\mathfrak{T}_1) - \Phi^2(\mu_1)) = 0.$$

This proves the first part. For the second part, we let (μ_t, \mathfrak{T}_t) and $(\mu'_t, \mathfrak{T}'_t)$ be two equivalent deformations. For $n = 1$, it follows from (12) and (13) that

$$\begin{aligned} (\mu'_1 - \mu_1)(x, y) &= [\phi_1(x), y] + [x, \phi_1(y)] - \phi_1[x, y], \\ (\mathfrak{T}'_1 - \mathfrak{T}_1)(x) &= \mathfrak{T}(\phi_1(x)) - \phi_1(\mathfrak{T}(x)). \end{aligned}$$

Hence $(\mu'_1, \mathfrak{T}'_1) - (\mu_1, \mathfrak{T}_1) = (\delta_{\text{CE}}\phi_1, -\Phi^1(\phi_1)) = \delta_{\text{RB}}(\phi_1, 0)$. This shows that (μ_1, \mathfrak{T}_1) and $(\mu'_1, \mathfrak{T}'_1)$ correspond to the same cohomology class in $H_{\text{RB}}^2(\mathfrak{g}, \mathfrak{g})$. This completes the proof. \square

We will now prove the following interesting result about formal 1-parameter deformations.

4.7 Theorem. *Let $(\mathfrak{g}, \mathfrak{T})$ be a λ -weighted Rota-Baxter Lie algebra. If $H_{\text{RB}}^2(\mathfrak{g}, \mathfrak{g}) = 0$ then any formal 1-parameter deformation of $(\mathfrak{g}, \mathfrak{T})$ is equivalent to the trivial one $(\mu'_t = \mu, \mathfrak{T}'_t = \mathfrak{T})$.*

Proof. Let (μ_t, \mathfrak{T}_t) be a formal 1-parameter deformation of the λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$. It follows from Theorem 4.6 that (μ_1, \mathfrak{T}_1) is a 2-cocycle. Thus, from the hypothesis, there exists $(\psi_1, v) \in \text{Hom}(\mathfrak{g}, \mathcal{V}) \oplus \mathcal{V}$ such that

$$(\mu_1, \mathfrak{T}_1) = \delta_{\text{RB}}(\psi_1, v) = (\delta_{\text{CE}}(\psi_1), -\partial_{\text{CE}}x - \Phi^1(\psi_1)). \quad (14)$$

Let $\phi_1 : \mathfrak{g} \rightarrow \mathfrak{g}$ be the map $\phi_1 = \text{id}_{\mathfrak{g}} + \phi_1 t$. Then

$$(\bar{\mu}_t = \phi_t^{-1} \circ \mu_t \circ (\phi_t \otimes \phi_t), \bar{\mathfrak{T}}_t = \phi_t^{-1} \circ \mathfrak{T}_t \circ \phi_t)$$

is a deformation of $(\mathfrak{g}, \mathfrak{T})$ equivalent to (μ_t, \mathfrak{T}_t) . Using (14), it can be easily check that the linear terms of $\bar{\mu}_t$ and $\bar{\mathfrak{T}}_t$ are vanish. In other words, $\bar{\mu}_t$ and $\bar{\mathfrak{T}}_t$ are of the form

$$\bar{\mu}_t = \mu + \bar{\mu}_2 t^2 + \dots \quad \text{and} \quad \bar{\mathfrak{T}}_t = \mathfrak{T} + \bar{\mathfrak{T}}_2 t^2 + \dots.$$

Hence by repeating this argument, we conclude that (μ_t, \mathfrak{T}_t) is equivalent to $(\mu'_t = \mu, \mathfrak{T}'_t = \mathfrak{T})$. Hence the proof. \square

4.8 Remark. A λ -weighted Rota-Baxter Lie algebra $(\mathfrak{g}, \mathfrak{T})$ is said to be rigid if any deformation of it is equivalent to the trivial one. It follows from the above theorem that the vanishing of the second cohomology group $H_{\text{RB}}^2(\mathfrak{g}, \mathfrak{g})$ is a sufficient condition for the rigidity of $(\mathfrak{g}, \mathfrak{T})$.

5 Rota-Baxter paired operators

Let \mathfrak{g} be a Lie algebra and $\mathcal{V} = (\mathcal{V}, \rho)$ be a representation of it. In this section, we introduce weighted Rota-Baxter paired operators that are related to weighted Rota-Baxter Lie algebras together with their representations. We introduce a differential graded Lie algebra whose Maurer-Cartan elements are weighted

Rota-Baxter paired operators. Using such characterization, we define the cohomology of Rota-Baxter paired operators. Finally, we discuss deformations of weighted Rota-Baxter paired operators in terms of cohomology.

5.1 Definition. Let \mathfrak{g} be a Lie algebra and $\mathcal{V} = (\mathcal{V}, \rho)$ be a representation of it. A pair $(\mathfrak{T}, \mathcal{T})$ of linear maps $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ and $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$ is said to be a λ -weighted Rota-Baxter paired operators if they satisfy the followings: for $x, y \in \mathfrak{g}$ and $u \in \mathcal{V}$,

$$\begin{aligned} [\mathcal{T}(x), \mathfrak{T}(y)] &= \mathfrak{T}([\mathfrak{T}(x), y] + [x, \mathfrak{T}(y)] + \lambda[x, y]), \\ \rho(\mathcal{T}x)(\mathcal{T}u) &= \mathcal{T}(\rho(\mathfrak{T}x)u + \rho(x)(\mathcal{T}u) + \lambda\rho(x)u), \end{aligned}$$

5.2 Remark. Observe that the first condition is equivalent to say that $(\mathfrak{g}, \mathfrak{T})$ is a λ -weighted Rota-Baxter Lie algebra, while the second condition is equivalent to say that $(\mathcal{V}, \mathcal{T})$ is a representation of $(\mathfrak{g}, \mathfrak{T})$. Therefore, λ -weighted Rota-Baxter paired operators are closely related with weighted Rota-Baxter Lie algebras and their representations. Hence examples of Rota-Baxter paired operators follow from the examples of Section 2.

Let \mathfrak{g} be a Lie algebra. Then the direct sum $\mathfrak{g} \oplus \mathfrak{g}$ carries a Lie algebra structure with bracket given by

$$[(x, x'), (y, y')]_{\lambda} = ([x, y], [x, y'] + [x', y] + \lambda[x', y']).$$

We denote this Lie algebra by $(\mathfrak{g} \oplus \mathfrak{g})_{\lambda}$. Moreover, if $\mathcal{V} = (\mathcal{V}, \rho)$ is a representation of the Lie algebra \mathfrak{g} , then $\mathcal{V} \oplus \mathcal{V}$ can be equipped with a representation of the Lie algebra $(\mathfrak{g} \oplus \mathfrak{g})_{\lambda}$ with the action given by

$$\rho_{\lambda}((x, x'))(u, u') = (\rho(x)u, \rho(x)u' + \rho(x')u + \lambda\rho(x')u').$$

Hence the direct sum $\mathfrak{g} \oplus \mathfrak{g} \oplus \mathcal{V} \oplus \mathcal{V}$ carries the semidirect product Lie algebra structure whose bracket is explicitly given by

$$\begin{aligned} &\{[(x, x', u, u'), (y, y', v, v')]\} \\ &= ([x, y], [x, y'] + [x', y] + \lambda[x', y'], \rho(x)v - \rho(y)u, \rho(x)v' + \rho(x')v + \lambda(x')v' - \rho(y)u' - \rho(y')u - \rho(y')u'). \end{aligned}$$

We denote this Lie algebra by $(\mathfrak{g} \oplus \mathfrak{g} \oplus \mathcal{V} \oplus \mathcal{V})_{\lambda}$. The following proposition is straightforward, hence we omit the details.

5.3 Proposition. Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. A pair $(\mathfrak{T}, \mathcal{T})$ of linear maps $\mathfrak{T} : \mathfrak{g} \rightarrow \mathfrak{g}$ and $\mathcal{T} : \mathcal{V} \rightarrow \mathcal{V}$ forms a λ -weighted Rota-Baxter paired operators if and only if

$$Gr_{\mathfrak{T}, \mathcal{T}} := \{(\mathfrak{T}x, x, \mathcal{T}u, u) \mid x \in \mathfrak{g}, u \in \mathcal{V}\} \subset \mathfrak{g} \oplus \mathfrak{g} \oplus \mathcal{V} \oplus \mathcal{V}$$

is a Lie subalgebra of $(\mathfrak{g} \oplus \mathfrak{g} \oplus \mathcal{V} \oplus \mathcal{V})_{\lambda}$.

Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. We denote \mathfrak{g}' and \mathcal{V}' by another copies of \mathfrak{g} and \mathcal{V} , respectively. Consider the vector space $\mathfrak{l} = \mathfrak{g} \oplus \mathfrak{g}' \oplus \mathcal{V} \oplus \mathcal{V}'$ and the Nijenhuis-Richardson bracket on the space $\oplus_{n \geq 1} \text{Hom}(\wedge^n \mathfrak{l}, \mathfrak{l})$ of skew-symmetric multilinear maps on \mathfrak{l} , given by

$$[f, g]_{\text{NR}} := f \diamond g - (-1)^{(m-1)(n-1)} g \diamond f,$$

where

$$(f \diamond g)(l_1, \dots, l_{m+n-1}) = \sum_{\sigma \in \mathbb{S}_{(n, m-1)}} (-1)^{\sigma} f(g(l_{\sigma(1)}, \dots, l_{\sigma(n)}), l_{\sigma(n+1)}, \dots, l_{\sigma(m+n-1)}),$$

for $f \in \text{Hom}(\wedge^m \mathfrak{l}, \mathfrak{l})$ and $g \in \text{Hom}(\wedge^n \mathfrak{l}, \mathfrak{l})$. The Nijenhuis-Richardson bracket $[\cdot, \cdot]_{\text{NR}}$ makes the shifted graded vector space $\oplus_{n \geq 0} \text{Hom}(\wedge^{n+1} \mathfrak{l}, \mathfrak{l})$ into a graded Lie algebra. Moreover, it can be easily checked that the graded subspace $\mathfrak{a} = \oplus_{n \geq 0} (\text{Hom}(\wedge^{n+1} \mathfrak{g}', \mathfrak{g}) \oplus \text{Hom}(\wedge^n \mathfrak{g}' \otimes \mathcal{V}', \mathcal{V}))$ is an abelian subalgebra.

In the subsequent, we will use the following notations. Let

- $\diamond \mu \in \text{Hom}(\wedge^2 \mathfrak{g}, \mathfrak{g})$ and $\mu' \in \text{Hom}(\wedge^2 \mathfrak{g}', \mathfrak{g}')$ denote the multiplications of \mathfrak{g} and \mathfrak{g}' , respectively;
- $\diamond \text{ad}' : \mathfrak{g} \times \mathfrak{g}' \rightarrow \mathfrak{g}'$ be the adjoint representation of \mathfrak{g} on \mathfrak{g}' ;
- $\diamond \rho : \mathfrak{g} \times \mathcal{V} \rightarrow \mathcal{V}$ and $\rho' : \mathfrak{g} \times \mathcal{V}' \rightarrow \mathcal{V}'$ denote the representations of \mathfrak{g} on \mathcal{V} and \mathcal{V}' , respectively;
- $\diamond \varrho : \mathfrak{g}' \times \mathcal{V} \rightarrow \mathcal{V}$ and $\varrho' : \mathfrak{g}' \times \mathcal{V}' \rightarrow \mathcal{V}'$ denote the representations of \mathfrak{g}' on \mathcal{V} and \mathcal{V}' , respectively.

We consider the element

$$\theta = \mu + \text{ad}' + \rho + \rho' + \varrho \in \text{Hom}(\wedge^2 \mathfrak{l}, \mathfrak{l}).$$

It is easy to see that θ is a Maurer-Cartan element in the graded Lie algebra $(\oplus_{n \geq 0} \text{Hom}(\wedge^{n+1} \mathfrak{l}, \mathfrak{l}), [\cdot, \cdot]_{\text{NR}})$. Therefore, it induces a differential $d_\theta := [\theta, -]_{\text{NR}}$ on the graded vector space $\oplus_{n \geq 0} \text{Hom}(\wedge^{n+1} \mathfrak{l}, \mathfrak{l})$. Hence by the derived bracket construction of Voronov [29], the shifted graded space

$$\mathfrak{a}[-1] = \oplus_{n \geq 1} (\text{Hom}(\wedge^n \mathfrak{g}', \mathfrak{g}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}' \otimes \mathcal{V}', \mathcal{V}))$$

inherits a graded Lie algebra structure with bracket given by

$$[\mathbf{P}, \mathbf{Q}] = (-1)^m [d_\theta(\mathbf{P}), \mathbf{Q}]_{\text{NR}}, \quad (15)$$

for $\mathbf{P} = (\mathbf{P}_1, \mathbf{P}_2) \in \text{Hom}(\wedge^m \mathfrak{g}', \mathfrak{g}) \oplus \text{Hom}(\wedge^{m-1} \mathfrak{g}' \otimes \mathcal{V}', \mathcal{V})$ and $\mathbf{Q} = (\mathbf{Q}_1, \mathbf{Q}_2) \in \text{Hom}(\wedge^n \mathfrak{g}', \mathfrak{g}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}' \otimes \mathcal{V}', \mathcal{V})$.

On the other hand, the element $\theta' = -(\mu' + \varrho') \in \text{Hom}(\wedge^2 \mathfrak{l}, \mathfrak{l})$ is also a Maurer-cartan element in the graded Lie algebra $(\oplus_{n \geq 0} \text{Hom}(\wedge^{n+1} \mathfrak{l}, \mathfrak{l}), [\cdot, \cdot]_{\text{NR}})$. Hence it induces a differential $d_{\theta'} = [\theta', -]_{\text{NR}}$ on $\oplus_{n \geq 1} \text{Hom}(\wedge^n \mathfrak{l}, \mathfrak{l})$. The differential $d_{\theta'}$ restricts to a differential \mathfrak{d} on the space $\mathfrak{a}[-1]$. Finally, the elements θ and θ' additionally satisfies $[\theta, \theta'] = 0$ which in turn implies that \mathfrak{d} is a derivation for the bracket $[\cdot, \cdot]$ on $\mathfrak{a}[-1]$. In other words, the triple $(\mathfrak{a}[-1], [\cdot, \cdot], \mathfrak{d})$ is a differential graded Lie algebra.

By replacing \mathfrak{g}' by \mathfrak{g} and replacing \mathcal{V}' by \mathcal{V} , we get the following.

5.4 Theorem. *Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. Then the triple*

$$(\oplus_{n \geq 1} (\text{Hom}(\wedge^n \mathfrak{g}, \mathfrak{g}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g} \otimes \mathcal{V}, \mathcal{V})), [\cdot, \cdot], \mathfrak{d})$$

is a differential graded Lie algebra. A pair $\mathbf{T} = (\mathfrak{T}, \mathcal{T})$ is a Rota-Baxter paired operators if and only if \mathbf{T} is a Maurer-Cartan element in the above differential graded Lie algebra.

Proof. The first part is clear from previous discussions. For any $\mathbf{T} = (\mathfrak{T}, \mathcal{T}) \in \text{Hom}(\mathfrak{g}, \mathfrak{g}) \oplus \text{Hom}(\mathcal{V}, \mathcal{V})$, by expanding (15) similar to [27], we get that

$$\begin{aligned} [\mathbf{T}, \mathbf{T}]_1(x, y) &= 2(\mathfrak{T}([\mathfrak{T}(x), y] + [x, \mathfrak{T}(y)]) - [\mathfrak{T}(x), \mathfrak{T}(y)]), \\ [\mathbf{T}, \mathbf{T}]_2(x, u) &= 2(\mathcal{T}(\rho(\mathfrak{T}(x))u + \rho(x)(\mathcal{T}u)) - \rho(\mathfrak{T}(x))(\mathcal{T}u)), \end{aligned}$$

for $x \in \mathfrak{g}$ and $u \in \mathcal{V}$. On the other hand, we can also check (similar to [10]) that

$$(\mathfrak{d}\mathbf{T})_1(x, y) = \lambda \mathfrak{T}[x, y] \quad \text{and} \quad (\mathfrak{d}\mathbf{T})_2(x, u) = \lambda \mathcal{T}(\rho(x)u).$$

Hence it follows that $\mathfrak{d}\mathbf{T} + \frac{1}{2}[\mathbf{T}, \mathbf{T}] = 0$ if and only if $\mathbf{T} = (\mathfrak{T}, \mathcal{T})$ is a Rota-Baxter paired operators. \square

It follows from the above theorem that a Rota-Baxter paired operators $\mathbf{T} = (\mathfrak{T}, \mathcal{T})$ induces a differential

$$d_{\mathbf{T}} : \text{Hom}(\wedge^n \mathfrak{g}, \mathfrak{g}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g} \otimes \mathcal{V}, \mathcal{V}) \rightarrow \text{Hom}(\wedge^{n+1} \mathfrak{g}, \mathfrak{g}) \oplus \text{Hom}(\wedge^n \mathfrak{g} \otimes \mathcal{V}, \mathcal{V}), \text{ for } n \geq 1,$$

$$d_{\mathbf{T}} = \mathfrak{d} + \llbracket \mathbf{T}, - \rrbracket.$$

We define $C_{\text{RBp}}^n(\mathfrak{g}, \mathcal{V}) = \text{Hom}(\wedge^n \mathfrak{g}, \mathfrak{g}) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g} \otimes \mathcal{V}, \mathcal{V})$, for $n \geq 1$. Then $\{C_{\text{RBp}}^n(\mathfrak{g}, \mathcal{V}), d_{\mathbf{T}}\}$ is a cochain complex. The corresponding cohomology groups are called the cohomology of the Rota-Baxter paired operators $(\mathfrak{T}, \mathcal{T})$.

Note that, for a Rota-Baxter paired operators $\mathbf{T} = (\mathfrak{T}, \mathcal{T})$, the differential $d_{\mathbf{T}}$ makes the tuple

$$(\oplus_{n \geq 1} C_{\text{RBp}}^n(\mathfrak{g}, \mathcal{V}), \llbracket \cdot, \cdot \rrbracket, d_{\mathbf{T}})$$

into a differential graded Lie algebra. In the following, we will show that this differential graded Lie algebra governs the Maurer-Cartan deformations of \mathbf{T} .

5.5 Proposition. *Let \mathfrak{g} be a Lie algebra and \mathcal{V} be a representation of it. Let $\mathbf{T} = (\mathfrak{T}, \mathcal{T})$ be a Rota-Baxter paired operators. Then for a pair $\mathbf{T}' = (\mathfrak{T}', \mathcal{T}')$ of maps $\mathfrak{T}' : \mathfrak{g} \rightarrow \mathfrak{g}$ and $\mathcal{T}' : \mathcal{V} \rightarrow \mathcal{V}$, the sum $\mathbf{T} + \mathbf{T}' = (\mathfrak{T} + \mathfrak{T}', \mathcal{T} + \mathcal{T}')$ is a Rota-Baxter paired operators if and only if \mathbf{T}' is a Maurer-Cartan element in the differential graded Lie algebra $(\oplus_{n \geq 1} C_{\text{RBp}}^n(\mathfrak{g}, \mathcal{V}), \llbracket \cdot, \cdot \rrbracket, d_{\mathbf{T}})$.*

Proof. We observe that

$$\begin{aligned} & \mathfrak{d}(\mathbf{T} + \mathbf{T}') + \frac{1}{2} \llbracket \mathbf{T} + \mathbf{T}', \mathbf{T} + \mathbf{T}' \rrbracket \\ &= \mathfrak{d}\mathbf{T} + \mathfrak{d}\mathbf{T}' + \frac{1}{2} (\llbracket \mathbf{T}, \mathbf{T} \rrbracket + 2\llbracket \mathbf{T}, \mathbf{T}' \rrbracket + \llbracket \mathbf{T}', \mathbf{T}' \rrbracket) \\ &= \mathfrak{d}\mathbf{T}' + \llbracket \mathbf{T}, \mathbf{T}' \rrbracket + \frac{1}{2} \llbracket \mathbf{T}', \mathbf{T}' \rrbracket = d_{\mathbf{T}}(\mathbf{T}') + \frac{1}{2} \llbracket \mathbf{T}', \mathbf{T}' \rrbracket. \end{aligned}$$

Hence the result follows from the Maurer-Cartan characterization of Rota-Baxter paired operators. \square

5.6 Remark. In [10] the present author studied formal 1-parameter deformations of weighted Rota-Baxter operators by keeping the underlying algebra intact. Given a Lie algebra \mathfrak{g} and a representation \mathcal{V} , one could also study formal 1-parameter deformations of Rota-Baxter paired operators by keeping the algebra \mathfrak{g} and the representation \mathcal{V} intact. Then it is easy to see that the above-defined cohomology governs such deformations. More precisely, the infinitesimals lie in the first cohomology group, and the obstruction to extending a finite order deformation lies in the second cohomology group.

Acknowledgements. The research is supported by the fellowship of Indian Institute of Technology (IIT) Kanpur.

References

- [1] M. Aguiar, Pre-Poisson algebras, *Lett. Math. Phys.* 54 (2000), 263-277.
- [2] F. V. Atkinson, Some aspects of Baxter's functional equation, *J. Math. Anal. Appl.* 7 (1963), 1-30.
- [3] C. Bai, O. Bellier, L. Guo and X. Ni, Splitting of operations, Manin products and Rota-Baxter operators, *Int. Math. Res. Not. IMRN* 2013 (2013), 485-524.

- [4] C. Bai, L. Guo and X. Ni, Nonabelian generalized Lax pairs, the classical Yang-Baxter equation and PostLie algebras, *Comm. Math. Phys.* 297 (2010), 553-596.
- [5] C. Bai, L. Guo and X. Ni, Relative Rota-Baxter algebras and tridendriform algebras, *J. Algebra Appl.* 12 (2013) 1350027.
- [6] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* 10 (1960), 731-742.
- [7] P. Cartier, On the structure of free Baxter algebras, *Adv. Math.* 9 (1972) 253-265.
- [8] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* 210 (2000), 249-273.
- [9] A. Das, Deformations of associative Rota-Baxter operators, *J. Algebra* 560 (2020) 144-180.
- [10] A. Das, Cohomology and deformations of weighted Rota-Baxter operators, arXiv:2108.05411
- [11] A. Das and S. K. Mishra, The L_∞ -deformations of associative Rota-Baxter algebras and homotopy Rota-Baxter operators, arXiv:2008.11076
- [12] K. Ebrahimi-Fard, Loday-type algebras and the Rota-Baxter relation, *Lett. Math. Phys.* 61 (2002) 139-147.
- [13] K. Ebrahimi-Fard and L. Guo, Rota-Baxter algebras and dendriform algebras, *J. Pure Appl. Algebra* 212 (2008) 320-339.
- [14] M. Gerstenhaber, On the deformation of rings and algebras, *Ann. of Math.* (2) 79 (1964), 59-103.
- [15] L. Guo and W. Keigher, Baxter algebras and shuffle products, *Adv. Math.* 150 (2000) 117-149.
- [16] L. Guo and B. Zhang, Renormalization of multiple zeta values, *J. Algebra* 319 (2008) 3770-3809.
- [17] L. Guo, An introduction to Rota-Baxter algebra, *Surveys of Modern Mathematics*, 4. International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- [18] J. Jiang, Y. Sheng and C. Zhu, Cohomologies of relative Rota-Baxter operators on Lie groups and Lie algebras, arXiv:2108.02627v1
- [19] B. A. Kupersmidt, What a classical r-matrix really is, *J. Nonlinear Math. Phys.* 6 (1999), no. 4, 448-488.
- [20] J.-L. Loday, Cyclic homology, Grundlehren der Mathematischen Wissenschaften 301, Springer-Verlag, Berlin, 1998. xx+513 pp.
- [21] A. Lazarev, Y. Sheng and R. Tang, Deformations and homotopy theory of relative Rota-Baxter Lie algebras, *Comm. Math. Phys.* 383 (2021) 595-631.
- [22] J. B. Miller, Some properties of Baxter operators, *Acta Math. Acad. Sci. Hungar.* 17 (1966), 387-400.
- [23] J. Qiu and Y. Chen, Gröbner-Shirshov bases for Lie Ω -algebras and free Rota-Baxter Lie algebras, *J. Algebra Appl.* Vol. 16, No. 10 (2017) 1750190.
- [24] A. Nijenhuis and R. Richardson, Deformation of Lie algebra structures, *J. Math. Mech.* 17 (1967), 89-105.

- [25] G.-C. Rota, Baxter algebras and combinatorial identities, I, II, *Bull. Amer. Math. Soc.* 75 (1969), 325-329; *ibid.* 75 1969 330-334.
- [26] M. A. Semenov-Tyan Shanskii, What is a classical r -matrix ? *Funct. Anal. Appl.* 17 (1893) 259-272.
- [27] R. Tang, C. Bai, L. Guo and Y. Sheng, Deformations and their controlling cohomologies of \mathcal{O} -operators, *Comm. Math. Phys.* 368 (2019), no. 2, 665-700.
- [28] R. Tang, C. Bai, L. Guo and Y. Sheng, Homotopy Rota-Baxter operators, homotopy \mathcal{O} -operators and homotopy post-Lie algebras, arXiv:1907.13504
- [29] Th. Voronov, Higher derived brackets and homotopy algebras, *J. Pure Appl. Algebra* 202 (2005), 133-153.
- [30] K. Wang and G. Zhou, Deformations and homotopy theory of Rota-Baxter algebras of any weight, arXiv:2108.06744
- [31] Y. Zhang, D. Chen, X. Gao and Y. Luo, Weighted infinitesimal unitary bialgebras on rooted forests and weighted cocycles, *Pacific J. Math.* 302 (2019), 741-766.
- [32] Y. Zhang, X. Gao and J. Zheng, Weighted infinitesimal unitary bialgebras on matrix algebras and weighted associated Yang-Baxter equations, arXiv:1811.00842
- [33] H. Zheng, L. Guo and L. Zhang, Rota-Baxter paired modules and their constructions from Hopf algebras, *J. Algebra* 559 (2020) 601-624.