

# Representations of the Yangians associated with Lie superalgebras $\mathfrak{osp}(1|2n)$

A. I. Molev

## Abstract

We classify the finite-dimensional irreducible representations of the Yangians associated with the orthosymplectic Lie superalgebras  $\mathfrak{osp}_{1|2n}$  in terms of the Drinfeld polynomials. The arguments rely on the description of the representations in the particular case  $n = 1$  obtained in our previous work.

## 1 Introduction

The finite-dimensional irreducible representations of the Yangian  $Y(\mathfrak{g})$  associated with a simple Lie algebra  $\mathfrak{g}$  were classified by Drinfeld [5]. The arguments rely on the work of Tarasov [9] on the particular case of  $Y(\mathfrak{sl}_2)$ , where the classification was carried over in the language of monodromy matrices within the quantum inverse scattering method; see [7, Sec. 3.3] for a detailed adapted exposition of these results. This description of the representations of the Yangian  $Y(\mathfrak{sl}_2)$ , along with some other low rank cases, should also play an essential role in the classification of the finite-dimensional irreducible representations of the Yangians associated with simple Lie superalgebras. One of these cases was considered in our previous work [8], where the representations of the Yangian  $Y(\mathfrak{osp}_{1|2})$  were described.

These two basic cases turn out to be sufficient to complete the classification in the case of the Yangians associated with the orthosymplectic Lie superalgebras  $\mathfrak{osp}_{1|2n}$ . We prove in this paper that, similar to the classification results of [5], the finite-dimensional irreducible representations of the Yangian  $Y(\mathfrak{osp}_{1|2n})$  are in one-to-one correspondence with the  $n$ -tuples of monic polynomials  $(P_1(u), \dots, P_n(u))$ , and so we call them the *Drinfeld polynomials*.

To describe the results in more detail, recall that the Yangian  $Y(\mathfrak{osp}_{M|2n})$ , as introduced by Arnaudon *et al.* [1], can be considered as a quotient of the extended Yangian  $X(\mathfrak{osp}_{M|2n})$  defined via an *RTT* relation. A standard argument shows that every finite-dimensional irreducible representation of  $X(\mathfrak{osp}_{M|2n})$  is a highest weight representation. It is isomorphic to the irreducible quotient  $L(\lambda(u))$  of the Verma module  $M(\lambda(u))$  associated with an  $(n + 1)$ -tuple  $\lambda(u) = (\lambda_1(u), \dots, \lambda_{n+1}(u))$  of formal series  $\lambda_i(u) \in 1 + u^{-1}\mathbb{C}[[u^{-1}]]$ . The tuple is called the *highest weight* of the representation. The key step in the classification is to find the conditions on the highest weight for the representation  $L(\lambda(u))$  to be finite-dimensional. The required necessary conditions are derived by induction from those for the associated actions of the Yangians

$Y(\mathfrak{gl}_2)$  and  $X(\mathfrak{osp}_{1|2})$  on the respective cyclic spans of the highest vector of  $L(\lambda(u))$ . The sufficiency of these conditions is verified by constructing the *fundamental representations* of the Yangian  $X(\mathfrak{osp}_{M|2n})$ ; cf. [3], [4]. The following is our main result.

**Main Theorem.** *Every finite-dimensional irreducible representation of the algebra  $X(\mathfrak{osp}_{1|2n})$  is isomorphic to  $L(\lambda(u))$  for a certain highest weight  $\lambda(u)$ . The representation  $L(\lambda(u))$  is finite-dimensional if and only if*

$$\frac{\lambda_{i+1}(u)}{\lambda_i(u)} = \frac{P_i(u+1)}{P_i(u)}, \quad i = 1, \dots, n, \quad (1.1)$$

for some monic polynomials  $P_i(u)$  in  $u$ . The finite-dimensional irreducible representations of the Yangian  $Y(\mathfrak{osp}_{1|2n})$  are in a one-to-one correspondence with the  $n$ -tuples of monic polynomials  $(P_1(u), \dots, P_n(u))$ .

## 2 Definitions and preliminaries

For any integer  $n \geq 1$  introduce the involution  $i \mapsto i' = 2n - i + 2$  on the set  $\{1, 2, \dots, 2n+1\}$ . Consider the  $\mathbb{Z}_2$ -graded vector space  $\mathbb{C}^{1|2n}$  over  $\mathbb{C}$  with the basis  $e_1, e_2, \dots, e_{2n+1}$ , where the vectors  $e_i$  and  $e_{i'}$  with  $i = 1, \dots, n$  are odd and the vector  $e_{n+1}$  is even. We set

$$\bar{i} = \begin{cases} 1 & \text{for } i = 1, \dots, n, n', \dots, 1', \\ 0 & \text{for } i = n+1. \end{cases}$$

The endomorphism algebra  $\text{End } \mathbb{C}^{1|2n}$  gets a  $\mathbb{Z}_2$ -gradation with the parity of the matrix unit  $e_{ij}$  found by  $\bar{i} + \bar{j} \pmod{2}$ .

We will consider even square matrices with entries in  $\mathbb{Z}_2$ -graded algebras, their  $(i, j)$  entries will have the parity  $\bar{i} + \bar{j} \pmod{2}$ . The algebra of even matrices over a superalgebra  $\mathcal{A}$  will be identified with the tensor product algebra  $\text{End } \mathbb{C}^{1|2n} \otimes \mathcal{A}$ , so that a matrix  $A = [a_{ij}]$  is regarded as the element

$$A = \sum_{i,j=1}^{2n+1} e_{ij} \otimes a_{ij} (-1)^{\bar{i}\bar{j}+\bar{j}} \in \text{End } \mathbb{C}^{1|2n} \otimes \mathcal{A}.$$

We will use the involutive matrix *super-transposition*  $t$  defined by  $(A^t)_{ij} = A_{j'i'} (-1)^{\bar{i}\bar{j}+\bar{j}} \theta_i \theta_j$ , where we set

$$\theta_i = \begin{cases} 1 & \text{for } i = 1, \dots, n+1, \\ -1 & \text{for } i = n+2, \dots, 2n+1. \end{cases}$$

This super-transposition is associated with the bilinear form on the space  $\mathbb{C}^{1|2n}$  defined by the anti-diagonal matrix  $G = [\delta_{ij}, \theta_i]$ . We will also regard  $t$  as the linear map

$$t : \text{End } \mathbb{C}^{1|2n} \rightarrow \text{End } \mathbb{C}^{1|2n}, \quad e_{ij} \mapsto e_{j'i'} (-1)^{\bar{i}\bar{j}+\bar{i}} \theta_i \theta_j. \quad (2.1)$$

In the case of multiple tensor products of the endomorphism algebras, we will indicate by  $t_a$  the map (2.1) acting on the  $a$ -th copy of  $\text{End } \mathbb{C}^{1|2n}$ .

A standard basis of the general linear Lie superalgebra  $\mathfrak{gl}_{1|2n}$  is formed by elements  $E_{ij}$  of the parity  $\bar{i} + \bar{j} \pmod 2$  for  $1 \leq i, j \leq 2n + 1$  with the commutation relations

$$[E_{ij}, E_{kl}] = \delta_{kj} E_{il} - \delta_{il} E_{kj} (-1)^{(\bar{i}+\bar{j})(\bar{k}+\bar{l})}.$$

We will regard the orthosymplectic Lie superalgebra  $\mathfrak{osp}_{1|2n}$  associated with the bilinear form defined by  $G$  as the subalgebra of  $\mathfrak{gl}_{1|2n}$  spanned by the elements

$$F_{ij} = E_{ij} - E_{j'i'} (-1)^{\bar{i}\bar{j}+\bar{i}} \theta_i \theta_j.$$

Introduce the permutation operator  $P$  by

$$P = \sum_{i,j=1}^{2n+1} e_{ij} \otimes e_{ji} (-1)^{\bar{j}} \in \text{End } \mathbb{C}^{1|2n} \otimes \text{End } \mathbb{C}^{1|2n}$$

and set

$$Q = P^{t_1} = P^{t_2} = \sum_{i,j=1}^{2n+1} e_{ij} \otimes e_{i'j'} (-1)^{\bar{i}\bar{j}} \theta_i \theta_j \in \text{End } \mathbb{C}^{1|2n} \otimes \text{End } \mathbb{C}^{1|2n}.$$

The  $R$ -matrix associated with  $\mathfrak{osp}_{1|2n}$  is the rational function in  $u$  given by

$$R(u) = 1 - \frac{P}{u} + \frac{Q}{u - \kappa}, \quad \kappa = -n - 1/2.$$

This is a super-version of the  $R$ -matrix originally found in [10]. Following [1], we define the *extended Yangian*  $X(\mathfrak{osp}_{1|2n})$  as a  $\mathbb{Z}_2$ -graded algebra with generators  $t_{ij}^{(r)}$  of parity  $\bar{i} + \bar{j} \pmod 2$ , where  $1 \leq i, j \leq 2n + 1$  and  $r = 1, 2, \dots$ , satisfying certain quadratic relations. In order to write them down, introduce the formal series

$$t_{ij}(u) = \delta_{ij} + \sum_{r=1}^{\infty} t_{ij}^{(r)} u^{-r} \in X(\mathfrak{osp}_{1|2n})[[u^{-1}]] \quad (2.2)$$

and combine them into the matrix  $T(u) = [t_{ij}(u)]$  so that

$$T(u) = \sum_{i,j=1}^{2n+1} e_{ij} \otimes t_{ij}(u) (-1)^{\bar{i}\bar{j}+\bar{j}} \in \text{End } \mathbb{C}^{1|2n} \otimes X(\mathfrak{osp}_{1|2n})[[u^{-1}]].$$

Consider the algebra  $\text{End } \mathbb{C}^{1|2n} \otimes \text{End } \mathbb{C}^{1|2n} \otimes X(\mathfrak{osp}_{1|2n})[[u^{-1}]]$  and introduce its elements  $T_1(u)$  and  $T_2(u)$  by

$$T_1(u) = \sum_{i,j=1}^{2n+1} e_{ij} \otimes 1 \otimes t_{ij}(u) (-1)^{\bar{i}\bar{j}+\bar{j}}, \quad T_2(u) = \sum_{i,j=1}^{2n+1} 1 \otimes e_{ij} \otimes t_{ij}(u) (-1)^{\bar{i}\bar{j}+\bar{j}}.$$

The defining relations for the algebra  $X(\mathfrak{osp}_{1|2n})$  take the form of the  $RTT$ -relation

$$R(u - v) T_1(u) T_2(v) = T_2(v) T_1(u) R(u - v). \quad (2.3)$$

As shown in [1], the product  $T(u) T^t(u - \kappa)$  is a scalar matrix with

$$T(u - \kappa) T^t(u) = c(u)1, \quad (2.4)$$

where  $c(u)$  is a series in  $u^{-1}$ . All its coefficients belong to the center  $ZX(\mathfrak{osp}_{1|2n})$  of  $X(\mathfrak{osp}_{1|2n})$  and generate the center.

The Yangian  $Y(\mathfrak{osp}_{1|2n})$  is defined as the subalgebra of  $X(\mathfrak{osp}_{1|2n})$  which consists of the elements stable under the automorphisms

$$t_{ij}(u) \mapsto f(u) t_{ij}(u) \quad (2.5)$$

for all series  $f(u) \in 1 + u^{-1}\mathbb{C}[[u^{-1}]]$ . We have the tensor product decomposition

$$X(\mathfrak{osp}_{1|2n}) = ZX(\mathfrak{osp}_{1|2n}) \otimes Y(\mathfrak{osp}_{1|2n}). \quad (2.6)$$

The Yangian  $Y(\mathfrak{osp}_{1|2n})$  can be equivalently defined as the quotient of  $X(\mathfrak{osp}_{1|2n})$  by the relation

$$T(u - \kappa) T^t(u) = 1.$$

We will also use a more explicit form of the defining relations (2.3) written in terms of the series (2.2) as follows:

$$\begin{aligned} [t_{ij}(u), t_{kl}(v)] = & \frac{1}{u - v} (t_{kj}(u) t_{il}(v) - t_{kj}(v) t_{il}(u)) (-1)^{\bar{i}\bar{j} + \bar{i}\bar{k} + \bar{j}\bar{k}} \\ & - \frac{1}{u - v - \kappa} \left( \delta_{ki'} \sum_{p=1}^{2n+1} t_{pj}(u) t_{p'l}(v) (-1)^{\bar{i} + \bar{i}\bar{j} + \bar{j}\bar{p}} \theta_i \theta_p \right. \\ & \left. - \delta_{lj'} \sum_{p=1}^{2n+1} t_{kp'}(v) t_{ip}(u) (-1)^{\bar{j} + \bar{p} + \bar{i}\bar{k} + \bar{j}\bar{k} + \bar{i}\bar{p}} \theta_j \theta_p \right). \end{aligned} \quad (2.7)$$

For any  $a \in \mathbb{C}$  the mapping

$$t_{ij}(u) \mapsto t_{ij}(u + a) \quad (2.8)$$

defines an automorphism of the algebra  $X(\mathfrak{osp}_{1|2n})$ .

The universal enveloping algebra  $U(\mathfrak{osp}_{1|2n})$  can be regarded as a subalgebra of  $X(\mathfrak{osp}_{1|2n})$  via the embedding

$$F_{ij} \mapsto \frac{1}{2} (t_{ij}^{(1)} - t_{j'i'}^{(1)} (-1)^{\bar{j} + \bar{i}\bar{j}} \theta_i \theta_j) (-1)^{\bar{i}}. \quad (2.9)$$

This fact relies on the Poincaré–Birkhoff–Witt theorem for the orthosymplectic Yangian which was pointed out in [1] and [2]. It states that the associated graded algebra for  $Y(\mathfrak{osp}_{1|2n})$  is isomorphic to  $U(\mathfrak{osp}_{1|2n}[u])$ . A detailed proof of the theorem can be given by extending the arguments of [3, Sec. 3] to the super case with the use of the vector representation recalled below in (3.6).

The extended Yangian  $X(\mathfrak{osp}_{1|2n})$  is a Hopf algebra with the coproduct defined by

$$\Delta : t_{ij}(u) \mapsto \sum_{k=1}^{2n+1} t_{ik}(u) \otimes t_{kj}(u). \quad (2.10)$$

For the image of the series  $c(u)$  we have  $\Delta : c(u) \mapsto c(u) \otimes c(u)$  and so the Yangian  $Y(\mathfrak{osp}_{1|2n})$  inherits the Hopf algebra structure from  $X(\mathfrak{osp}_{1|2n})$ .

### 3 Highest weight representations

We will start by deriving a general reduction property for representations of the extended Yangians  $X(\mathfrak{osp}_{1|2n})$  analogous to [3, Lemma 5.13]. For an  $X(\mathfrak{osp}_{1|2n})$ -module  $V$  set

$$V^+ = \{\eta \in V \mid t_{1j}(u)\eta = 0 \text{ for } j > 1 \text{ and } t_{i1'}(u)\eta = 0 \text{ for } i < 1'\}. \quad (3.1)$$

**Proposition 3.1.** *The subspace  $V^+$  is stable under the action of the operators  $t_{ij}(u)$  subject to  $2 \leq i, j \leq 2n$ . Moreover, the assignment  $\bar{t}_{ij}(u) \mapsto t_{i+1,j+1}(u)$  for  $1 \leq i, j \leq 2n-1$  defines a representation of the algebra  $X(\mathfrak{osp}_{1|2n-2})$  on  $V^+$ , where the  $\bar{t}_{ij}(u)$  denote the respective generating series for  $X(\mathfrak{osp}_{1|2n-2})$ .*

*Proof.* Suppose that  $2 \leq k, l \leq 2n$  and  $j > 1$ . For any  $\eta \in V^+$  apply (2.7) to get

$$t_{1j}(u)t_{kl}(u)\eta = \frac{1}{u-v-\kappa} \delta_{lj'}(-1)^{\bar{j}+\bar{k}+\bar{j}\bar{k}} \theta_j t_{k1'}(v) t_{11}(u)\eta.$$

Another application of (2.7) yields

$$t_{k1'}(v) t_{11}(u)\eta = -[t_{11}(u), t_{k1'}(v)]\eta = \frac{1}{u-v-\kappa} t_{k1'}(v) t_{11}(u)\eta,$$

implying  $t_{1j}(u)t_{kl}(u)\eta = 0$ . A similar calculation shows that  $t_{i1'}(u)t_{kl}(u)\eta = 0$  for  $i < 1'$  thus proving the first part of the proposition.

Now suppose that  $2 \leq i, j, k, l \leq 2n$ . By (2.7) the super-commutator  $[t_{ij}(u), t_{kl}(v)]$  of the operators in  $V^+$  equals

$$\begin{aligned} & \frac{1}{u-v} (t_{kj}(u) t_{il}(v) - t_{kj}(v) t_{il}(u)) (-1)^{\bar{i}\bar{j}+\bar{i}\bar{k}+\bar{j}\bar{k}} \\ & - \frac{1}{u-v-\kappa} \left( \delta_{ki'} \sum_{p=2}^{2n} t_{pj}(u) t_{p'l}(v) (-1)^{\bar{i}+\bar{i}\bar{j}+\bar{j}\bar{p}} \theta_i \theta_p \right. \\ & \quad \left. - \delta_{lj'} \sum_{p=2}^{2n} t_{kp'}(v) t_{ip}(u) (-1)^{\bar{j}+\bar{p}+\bar{i}\bar{k}+\bar{j}\bar{k}+\bar{i}\bar{p}} \theta_j \theta_p \right) \end{aligned}$$

plus the additional terms

$$- \frac{1}{u-v-\kappa} \left( \delta_{ki'} t_{1j}(u) t_{1'l}(v) (-1)^{\bar{i}+\bar{i}\bar{j}+\bar{j}} \theta_i + \delta_{lj'} t_{k1'}(v) t_{i1}(u) (-1)^{\bar{j}+\bar{i}\bar{k}+\bar{j}\bar{k}+\bar{i}} \theta_j \right).$$

To transform these terms, use (2.7) again to get the relations

$$\begin{aligned} t_{1j}(u) t_{1'l}(v) &= \frac{1}{u-v-\kappa-1} \sum_{p=2}^{2n} t_{pj}(u) t_{p'l}(v) (-1)^{\bar{j}+\bar{j}\bar{p}} \theta_p \\ &\quad - \frac{1}{u-v-\kappa-1} \delta_{lj'} t_{1'1'}(v) t_{11}(u) \theta_j \end{aligned}$$

and

$$\begin{aligned} t_{k1'}(v) t_{i1}(u) &= [t_{i1}(u) t_{k1'}(v)](-1)^{\bar{i}+\bar{k}+\bar{i}\bar{k}} = \frac{1}{u-v-\kappa-1} \delta_{ki'} t_{11}(u) t_{1'1'}(v) (-1)^{\bar{i}} \theta_i \\ &\quad - \frac{1}{u-v-\kappa-1} \sum_{p=2}^{2n} t_{kp'}(v) t_{ip}(u) (-1)^{\bar{i}+\bar{p}+\bar{i}\bar{p}} \theta_p. \end{aligned}$$

Now combine the expressions together and observe that the actions of the operators  $t_{11}(u)$  and  $t_{1'1'}(v)$  in  $V^+$  commute. Taking into account the change of the value  $\kappa \mapsto \kappa + 1$  for the algebra  $X(\mathfrak{osp}_{1|2n-2})$ , we find that the formula for the super-commutator  $[t_{ij}(u), t_{kl}(v)]$  agrees with the defining relations of  $X(\mathfrak{osp}_{1|2n-2})$ .  $\square$

*Remark 3.2.* The reduction property of Proposition 3.1 should be related to a super-version of the embedding theorem for the orthogonal and symplectic Yangians proven in [6, Thm 3.1]. The arguments of that paper should apply to the super-case to lead to a Drinfeld-type presentation of the Yangians  $Y(\mathfrak{osp}_{1|2n})$  extending the work [2].  $\square$

A representation  $V$  of the algebra  $X(\mathfrak{osp}_{1|2n})$  is called a *highest weight representation* if there exists a nonzero vector  $\xi \in V$  such that  $V$  is generated by  $\xi$ ,

$$\begin{aligned} t_{ij}(u) \xi &= 0 & \text{for } 1 \leq i < j \leq 2n+1, & \quad \text{and} \\ t_{ii}(u) \xi &= \lambda_i(u) \xi & \text{for } i = 1, \dots, 2n+1, \end{aligned} \quad (3.2)$$

for some formal series

$$\lambda_i(u) \in 1 + u^{-1} \mathbb{C}[[u^{-1}]]. \quad (3.3)$$

The vector  $\xi$  is called the *highest vector* of  $V$ .

**Proposition 3.3.** *The series  $\lambda_i(u)$  associated with a highest weight representation  $V$  satisfy the consistency conditions*

$$\lambda_i(u) \lambda_{i'}(u + n - i + 1/2) = \lambda_{i+1}(u) \lambda_{(i+1)'}(u + n - i + 1/2) \quad (3.4)$$

for  $i = 1, \dots, n$ . Moreover, the coefficients of the series  $c(u)$  act in the representation  $V$  as the multiplications by scalars determined by

$$c(u) \mapsto \lambda_1(u) \lambda_{1'}(u + n + 1/2).$$

*Proof.* To derive the consistency conditions, we will use the induction on  $n$  with the base case  $n = 1$  already considered in [8]. Suppose that  $n \geq 2$  and introduce the subspace  $V^+$  by (3.1). The vector  $\xi$  belongs to  $V^+$ , and applying Proposition 3.1 we find that the cyclic span  $X(\mathfrak{osp}_{1|2n-2}) \xi$  is a highest weight submodule with the highest weight  $(\lambda_2(u), \dots, \lambda_{2'}(u))$ . By the induction hypothesis, this implies conditions (3.4) with  $i = 2, \dots, n$ . Furthermore, using the defining relations (2.7), we get

$$t_{12}(u) t_{1'2'}(v) \xi = \frac{1}{u-v-\kappa} \left( t_{12}(u) t_{1'2'}(v) - \lambda_1(u) \lambda_{1'}(v) + \lambda_2(u) \lambda_{2'}(v) \right) \xi$$

and so

$$(u - v - \kappa - 1) t_{12}(u) t_{1'2'}(v) \xi = \left( -\lambda_1(u) \lambda_{1'}(v) + \lambda_2(u) \lambda_{2'}(v) \right) \xi.$$

Setting  $v = u - \kappa - 1 = u + n - 1/2$  we obtain (3.4) for  $i = 1$ . Finally, the last part of the proposition is obtained by using the expression for  $c(u)$  implied by taking the  $(1', 1')$  entry in the matrix relation (2.4).  $\square$

As Proposition 3.3 shows, the series  $\lambda_i(u)$  in (3.2) with  $i > n + 1$  are uniquely determined by the first  $n + 1$  series. The corresponding  $(n + 1)$ -tuple  $\lambda(u) = (\lambda_1(u), \dots, \lambda_{n+1}(u))$  is called the *highest weight* of  $V$ .

Given an arbitrary  $(n + 1)$ -tuple  $\lambda(u) = (\lambda_1(u), \dots, \lambda_{n+1}(u))$  of formal series of the form (3.3), introduce the series  $\lambda_i(u)$  with  $i = n + 2, \dots, 2n + 1$  to satisfy the consistency conditions (3.4). Define the *Verma module*  $M(\lambda(u))$  as the quotient of the algebra  $X(\mathfrak{osp}_{1|2n})$  by the left ideal generated by all coefficients of the series  $t_{ij}(u)$  with  $1 \leq i < j \leq 2n + 1$ , and  $t_{ii}(u) - \lambda_i(u)$  for  $i = 1, \dots, 2n + 1$ . As in [3, Prop. 5.14], the Poincaré–Birkhoff–Witt theorem for the algebra  $X(\mathfrak{osp}_{1|2n})$  implies that the Verma module  $M(\lambda(u))$  is nonzero, and we denote by  $L(\lambda(u))$  its irreducible quotient.

**Proposition 3.4.** *Every finite-dimensional irreducible representation of the algebra  $X(\mathfrak{osp}_{1|2n})$  is isomorphic to  $L(\lambda(u))$  for a certain highest weight  $\lambda(u) = (\lambda_1(u), \dots, \lambda_{n+1}(u))$ .*

*Proof.* The argument is essentially the same as for the proof of the corresponding counterparts of the property for the Yangians associated with Lie algebras; cf. [3, Thm 5.1], [7, Sec. 3.2]. We outline some key steps.

Suppose that  $V$  is a finite-dimensional irreducible representation of the algebra  $X(\mathfrak{osp}_{1|2n})$  and introduce its subspace  $V^0$  by

$$V^0 = \{\eta \in V \mid t_{ij}(u) \eta = 0, \quad 1 \leq i < j \leq 2n + 1\}.$$

First we note that  $V^0$  is nonzero, which follows by considering the set of weights of  $V$ , regarded as an  $\mathfrak{osp}_{1|2n}$ -module defined via the embedding (2.9). This set is finite and hence contains a maximal weight with respect to the standard partial ordering on the set of weights of  $V$ . A weight vector with this weight belongs to  $V^0$ .

Furthermore, we show that  $V^0$  is stable under the action of all operators  $t_{ii}(u)$ . This follows by straightforward calculations similar to those used in the proof of Proposition 3.1, relying on the defining relations (2.7). In a similar way, we verify that all the operators  $t_{ii}(u)$  with  $i = 1, \dots, 2n + 1$  form a commuting family of operators on  $V^0$ . Hence they have a simultaneous eigenvector  $\xi \in V^0$ . Since the representation  $V$  is irreducible, the submodule  $X(\mathfrak{osp}_{1|2n})\xi$  must coincide with  $V$  thus proving that  $V$  is a highest weight module.

By considering the  $\mathfrak{osp}_{1|2n}$ -weights of  $V$  we can also conclude that the highest vector  $\xi$  of  $V$  is determined uniquely, up to a constant factor.  $\square$

Proposition 3.4 yields the first part of the Main Theorem. Our next step is to show that the conditions in the theorem are necessary for the representation  $L(\lambda(u))$  to be finite-dimensional. So we now suppose that  $\dim L(\lambda(u)) < \infty$  and argue by induction on  $n$ . The conditions (1.1) in the base case  $n = 1$  are implied by the main result of [8]. Suppose further that  $n \geq 2$ .



Recall that the Yangian  $Y(\mathfrak{gl}_n)$  for the general linear Lie algebra  $\mathfrak{gl}_n$  is defined as a unital associative algebra with countably many generators  $t_{ij}^{(1)\circ}, t_{ij}^{(2)\circ}, \dots$  where  $1 \leq i, j \leq n$ , and the defining relations

$$(u - v) [t_{ij}^\circ(u), t_{kl}^\circ(v)] = t_{kj}^\circ(u) t_{il}^\circ(v) - t_{kj}^\circ(v) t_{il}^\circ(u)$$

written in terms of the series

$$t_{ij}^\circ(u) = \delta_{ij} + t_{ij}^{(1)\circ} u^{-1} + t_{ij}^{(2)\circ} u^{-2} + \dots \in Y(\mathfrak{gl}_n)[[u^{-1}]];$$

see [7] for a detailed exposition of the algebraic structure and representation of the Yangians associated with  $\mathfrak{gl}_n$ . The Yangian  $Y(\mathfrak{gl}_n)$  can be regarded as a subalgebra of  $X(\mathfrak{osp}_{1|2n})$  via the embedding

$$Y(\mathfrak{gl}_n) \hookrightarrow X(\mathfrak{osp}_{1|2n}), \quad t_{ij}^\circ(u) \mapsto t_{ij}(-u) \quad \text{for } 1 \leq i, j \leq n. \quad (3.5)$$

The cyclic span  $Y(\mathfrak{gl}_n)\xi \subset L(\lambda(u))$  is a highest weight module over  $Y(\mathfrak{gl}_n)$ . Its highest weight is the  $n$ -tuple  $(\lambda_1(-u), \dots, \lambda_n(-u))$ . Since  $\dim L(\lambda(u)) < \infty$ , the corresponding conditions for finite-dimensional highest weight representations of  $Y(\mathfrak{gl}_n)$  must be satisfied; see [7, Sec. 3.4]. This implies conditions (1.1) of the Main Theorem for  $i = 1, \dots, n - 1$ .

Furthermore, by Proposition 3.1, the subspace  $L(\lambda(u))^+$  is a module over the extended Yangian  $X(\mathfrak{osp}_{1|2n-2})$ . The vector  $\xi$  generates a highest weight  $X(\mathfrak{osp}_{1|2n-2})$ -module with the highest weight  $(\lambda_2(u), \dots, \lambda_{n+1}(u))$ . Since this module is finite-dimensional, conditions (1.1) hold for  $i = 2, \dots, n$  by the induction hypothesis. This completes the proof of the necessity of the conditions.

Now suppose that conditions (1.1) hold and derive that the corresponding module  $L(\lambda(u))$  is finite-dimensional. The  $n$ -tuple of Drinfeld polynomials  $(P_1(u), \dots, P_n(u))$  determines the highest weight  $\lambda(u)$  up to a simultaneous multiplication of all components  $\lambda_i(u)$  by a series  $f(u) \in 1 + u^{-1}\mathbb{C}[[u^{-1}]]$ . This operation corresponds to twisting the action of the algebra  $X(\mathfrak{osp}_{1|2n})$  on  $L(\lambda(u))$  by the automorphism (2.5). Hence, it suffices to prove that a particular module  $L(\lambda(u))$  corresponding to a given set of Drinfeld polynomials is finite-dimensional.

Suppose that  $L(\lambda(u))$  and  $L(\mu(u))$  are the irreducible highest weight modules with the highest weights

$$\lambda(u) = (\lambda_1(u), \dots, \lambda_{n+1}(u)) \quad \text{and} \quad \mu(u) = (\mu_1(u), \dots, \mu_{n+1}(u)).$$

By the coproduct rule (2.10), the cyclic span  $X(\mathfrak{osp}_{1|2n})(\xi \otimes \xi')$  of the tensor product of the respective highest vectors of  $L(\lambda(u))$  and  $L(\mu(u))$  is a highest weight module with the highest weight

$$(\lambda_1(u)\mu_1(u), \dots, \lambda_{n+1}(u)\mu_{n+1}(u)).$$

This observation implies that the cyclic span corresponds to the set of Drinfeld polynomials  $(P_1(u)Q_1(u), \dots, P_n(u)Q_n(u))$ , where the  $P_i(u)$  and  $Q_i(u)$  are the Drinfeld polynomials for  $L(\lambda(u))$  and  $L(\mu(u))$ , respectively. Therefore, we only need to establish the sufficiency of conditions (1.1) for the *fundamental representations* of  $X(\mathfrak{osp}_{1|2n})$  associated with the  $n$ -tuples of



Drinfeld polynomials such that  $P_j(u) = 1$  for all  $j \neq i$  and  $P_i(u) = u + b$  for a certain  $i \in \{1, \dots, n\}$  and  $b \in \mathbb{C}$ ; cf. [4]. Moreover, it is sufficient to take one particular value of  $b \in \mathbb{C}$ ; the general case will then follow by twisting the action of the algebra  $X(\mathfrak{osp}_{1|2n})$  in such representations by automorphisms of the form (2.8).

Consider the vector representation of  $X(\mathfrak{osp}_{1|2n})$  on  $\mathbb{C}^{1|2n}$  defined by

$$t_{ij}(u) \mapsto \delta_{ij} + u^{-1} e_{ij}(-1)^{\bar{i}} - (u + \kappa)^{-1} e_{j'i'}(-1)^{\bar{i}j} \theta_i \theta_j. \quad (3.6)$$

The homomorphism property follows from (2.3) by applying the standard transposition to one copy of  $\text{End } \mathbb{C}^{1|2n}$  in the Yang–Baxter equation satisfied by  $R(u)$ . Now use the coproduct (2.10) and suitable automorphisms (2.8) to equip the tensor product space  $(\mathbb{C}^{1|2n})^{\otimes k}$  with the action of  $X(\mathfrak{osp}_{1|2n})$  by setting

$$t_{ij}(u) \mapsto \sum_{a_1, \dots, a_{k-1}=1}^{2n+1} t_{ia_1}(u) \otimes t_{a_1 a_2}(u-1) \otimes \dots \otimes t_{a_{k-1} j}(u-k+1), \quad (3.7)$$

where the generators act in the respective copies of the vector space  $\mathbb{C}^{1|2n}$  via the rule (3.6). For the values  $k = 1, \dots, n$  introduce the vectors

$$\xi_k = \sum_{\sigma \in \mathfrak{S}_k} \text{sgn } \sigma \cdot e_{\sigma(1)} \otimes \dots \otimes e_{\sigma(k)} \in (\mathbb{C}^{1|2n})^{\otimes k}.$$

Now verify that each vector  $\xi_k$  has the properties

$$t_{ij}(u) \xi_k = 0 \quad \text{for } 1 \leq i < j \leq n+1 \quad (3.8)$$

and

$$t_{ii}(u) \xi_k = \begin{cases} \frac{u-k}{u-k+1} \xi_k & \text{for } i = 1, \dots, k, \\ \xi_k & \text{for } i = k+1, \dots, n+1. \end{cases} \quad (3.9)$$

The expression for the vector  $\xi_k$  involves only tensor products of the basis vectors  $e_i$  with  $i \leq n$ . This implies that for the application of the operators  $t_{ij}(u)$  with  $1 \leq i \leq j \leq n$  to  $\xi_k$  we may restrict the sum in formula (3.7) to the values  $a_p \in \{1, \dots, n\}$ .

By using the embedding (3.5), we may regard the cyclic span  $Y(\mathfrak{gl}_n) \xi_k$  as a  $Y(\mathfrak{gl}_n)$ -module. Moreover, this module is isomorphic to  $A^{(k)}(\mathbb{C}^n)^{\otimes k}$ , where  $A^{(k)}$  is the anti-symmetrization operator. It is well-known that this  $Y(\mathfrak{gl}_n)$ -module is isomorphic to the evaluation module  $L(1, \dots, 1, 0, \dots, 0)$  (with  $k$  ones) twisted by a shift automorphism  $u \mapsto u + k - 1$ ; see e.g. [7, Sec. 6.5]. This yields formulas (3.8) and (3.9) with  $1 \leq i \leq j \leq n$ . They are easily verified directly for the remaining generators.

Formulas (3.9) show that the corresponding set of Drinfeld polynomials for the highest weight module  $X(\mathfrak{osp}_{1|2n}) \xi_k$  has the form  $P_i(u) = 1$  for  $i \neq k$ , while  $P_k(u) = u - k$ . This completes the proof of the second part of the Main Theorem concerning conditions (1.1). The last part is immediate from the decomposition (2.6); cf. [3, Sec. 5.3].

## References

- [1] D. Arnaudon, J. Avan, N. Crampé, L. Frappat, E. Ragoucy, *R-matrix presentation for super-Yangians  $Y(\mathfrak{osp}(m|2n))$* , J. Math. Phys. **44** (2003), 302–308.
- [2] D. Arnaudon, N. Crampé, L. Frappat, E. Ragoucy, *Super Yangian  $Y(\mathfrak{osp}(1|2))$  and the universal R-matrix of its quantum double*, Comm. Math. Phys. **240** (2003), 31–51.
- [3] D. Arnaudon, A. Molev and E. Ragoucy, *On the R-matrix realization of Yangians and their representations*, Annales Henri Poincaré **7** (2006), 1269–1325.
- [4] V. Chari and A. Pressley, *Fundamental representations of Yangians and rational R-matrices*, J. Reine Angew. Math. **417** (1991), 87–128.
- [5] V. G. Drinfeld, *A new realization of Yangians and quantized affine algebras*, Soviet Math. Dokl. **36** (1988), 212–216.
- [6] N. Jing, M. Liu and A. Molev, *Isomorphism between the R-matrix and Drinfeld presentations of Yangian in types B, C and D*, Comm. Math. Phys. **361** (2018), 827–872.
- [7] A. Molev, *Yangians and classical Lie algebras*, Mathematical Surveys and Monographs, 143. AMS, Providence, RI, 2007.
- [8] A. Molev, *Representations of the orthosymplectic Yangian*, [arXiv:2108.10104](https://arxiv.org/abs/2108.10104).
- [9] V. O. Tarasov, *Irreducible monodromy matrices for the R-matrix of the XXZ-model and lattice local quantum Hamiltonians*, Theor. Math. Phys. **63** (1985), 440–454.
- [10] A. B. Zamolodchikov and Al. B. Zamolodchikov, *Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field models*, Ann. Phys. **120** (1979), 253–291.

School of Mathematics and Statistics  
University of Sydney, NSW 2006, Australia  
[alexander.molev@sydney.edu.au](mailto:alexander.molev@sydney.edu.au)