

Asymptotic critical behavior of holographic phase transition at finite topological charge - a potentially new quantum phase transition at finite chemical potential

Toan T. Nguyen*

*Key Laboratory for Multiscale Simulation of Complex Systems,
University of Science, Vietnam National University, Hanoi,
334 Nguyen Trai street, Thanh Xuan district, Hanoi 100000, Vietnam*

Tran Huu Phat

*Vietnam Atomic Energy Commission, 59 Ly Thuong Kiet, Hanoi, Vietnam
(Dated: May 28, 2022)*

We investigate analytically the asymptotic critical behavior at large chemical potential of the conformal field living at the AdS boundary of a four-dimensional spacetime Einstein gravity. The threshold values of the chemical potential for the appearance of condensate states are discrete, equal spacing, with the gap approaches zero logarithmically in the limit $T \rightarrow 0$. Numerical results surprisingly show that, the result apply even for states with low quantum number, as low as for the first or second excited states of the condensate, especially on the liquid side of the black hole van der Waals - like phase transition. We postulate that, at the exact limit $T = 0$ where the gap is zero, all excite states of the condensate are activated above a finite chemical potential, suggesting a new quantum phase transition as a function of the chemical potential.

We start from the model of an Abelian Higgs field and a Maxwell field in the four-dimensional spacetime Einstein gravity. The bulk action is given by,

$$S = \frac{1}{16\pi G_N} \int d^4x \sqrt{-g} \left(R - \frac{16}{L^2} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - |\nabla\Psi - iqA\Psi|^2 - m^2|\Psi|^2 \right), \quad (1)$$

where G_N is the Newton constant. In the uncondensed phase, the solutions to Eq. (1) are the Reissner-Nordstrom black hole (BH).

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_{2,k}, \quad (2)$$

where

$$f(r) = k - \frac{2M}{r} + \frac{Q^2}{r^2} + \frac{r^2}{L^2}. \quad (3)$$

The event horizon r_0 is the largest solution of the equation $f(r_0) = 0$. The BH Hawking temperature T is:

$$T_{\text{Hawking}} = f'(r_0)/4\pi, \quad (4)$$

which is also regarded as the temperature of the conformal field at the AdS boundary.

For $k > 0$, using the entropy $S = \pi r_0^2$, pressure $P = 3/8\pi L^2$, one discovered a total analogy between the small-large BH phase transition and the liquid-gas phase transition of the van der Waals (vdW) theory [1]. In this set up, M becomes the enthalpy and k is interpreted as the measure of a new charge, the topological charge [2]. For $k = 1$, $L < 6$ corresponds to the liquid side, while $L > 6$ corresponds to the gas side of the vdW transition [3].

Using the ansatz, $A_\mu = (\Phi(r), 0, 0, 0)$, $\Psi = \Psi(r)$, after minimizing the action with respect to $\Phi(r)$ and $\Psi(r)$, we

arrive at the equations of motion for the fields:

$$\Psi'' + \left(\frac{2}{r} + \frac{f'}{f} \right) \Psi' + \left(\frac{q^2 \Phi^2}{f^2} + \frac{2}{L^2 f} \right) \Psi = 0, \quad (5)$$

$$\Phi'' + \frac{2}{r} \Phi' - \frac{2q^2 |\Psi|^2}{f} \Phi = 0. \quad (6)$$

At the AdS boundary $r \rightarrow \infty$,

$$\Phi(r \rightarrow \infty) = \mu - \frac{\rho}{r} + \dots \quad (7)$$

$$\Psi(r \rightarrow \infty) = \frac{\Psi^{(1)}}{r} + \frac{\Psi^{(2)}}{r^2} + \dots \quad (8)$$

We will work in the grand canonical ensemble and fix the chemical potential μ of the conformal field living at the boundary, ρ is the corresponding density [4].

At large temperature, the system has only a “trivial” normal solution:

$$\Phi(r) = \mu(1 - r_0/r), \quad (9)$$

$$\Psi(r) = 0. \quad (10)$$

As the temperature decreases below a critical value T_c , the system permits a non-vanishing $\Psi(r)$ condensate solution. Depending on the value of the chemical potential, as the temperature decreases even further, other solutions become permissible, which are identified as the 1st excited state, 2nd excited state, ... of the condensate [5]. This work attempts to deal with critical properties of these states, combining both analytical and numerical methods, especially the asymptotic behavior of critical behaviors of systems at large chemical potential μ .

Without losing generality, we set $Q = q = 1$. One also makes a change of coordinate to $z = r_0/r$ and rescale various quantities to simplify formulas. In the new coordinate, the AdS boundary corresponds to $z = 0$, the BH

event horizon corresponds to $z = 1$. Various quantities are scaled as following:

$$f(z) = \frac{f(r)}{z_h^2} = \frac{k}{z_h^2} - \frac{2M}{z_h}z + z^2 + \frac{1/L^2 z_h^4}{z^2}, \quad (11)$$

$$\Phi(z) = z_h^{-3}\Phi(r = 1/z), \quad (12)$$

$$\Psi(z) = r\Psi(r = 1/z)/z_h^2, \quad (13)$$

here $z_h = 1/r_0$, $T = 4\pi T_{\text{Hawking}}/z_h^3 = -f'(1)$; $\Psi^{(1)}$, $\Psi^{(2)}$, μ , ρ , are scaled accordingly, and are not listed here to avoid crowding. In the rest of the paper, the rescaled version of these quantities is always assumed, unless explicitly stated otherwise.

Notice that, for convenience, the field $\Psi(r)$ is not only rescaled, but also changed. With the new definition, the boundary conditions at the AdS boundary are:

$$\Psi(z \rightarrow 0) = \Psi^{(1)} + \Psi^{(2)}z + \dots, \quad (14)$$

$$\Phi(z \rightarrow 0) = \mu - \rho z + \dots. \quad (15)$$

Because $z = 1$ is the smallest positive solution of $f(z)$, the latter can be rewritten in a more compact form:

$$f(z) = z^{-2}(z-1) [(z-1)(z^2 + 2bz + b) - z^2 T], \quad (16)$$

where $b = 1/L^2 z_h^4$. The equations of motion for dimensionless fields are

$$\Psi'' + \frac{F'}{F}\Psi' + \frac{zF' - 2F + 2b}{z^2 F}\Psi + \frac{\Phi^2}{F^2}\Psi = 0, \quad (17)$$

$$\Phi'' - 2\Psi^2\Phi/F = 0, \quad (18)$$

with $F(z) = z^2 f(z)$.

For condensate solution at low temperature, the expectation value of the operator, $\langle \mathcal{O}_\delta \rangle = \sqrt{2}\Psi^{(\delta)}$ ($\delta = 1$ or 2) is non-zero. Analytical expressions are not available even for simplest forms of the metric $f(z)$ and one typically resorts to numerical solution. In Fig. 1, condensate solutions for $k = 1$, $L = 1$, $\mu = 8$, and $T = 0.1T_c$ are shown as a typical example. As the temperature decreases below the critical temperature $T_c = 1.699$, the condensate ground state is available. Below the critical temperature $T_c = 0.295$, the first excited state is available, the field $\Psi(z)$ has one node (zero) in the interval $(0, 1)$. Finally, below the critical temperature $T_c = 0.0806$, the second excited state is available, $\Psi(z)$ has two nodes.

Nevertheless, in numerical solutions, it is easy to overlook various hidden symmetries in the system. In this paper, we attempt to investigate behavior of the system near criticality using semi-analytical methods, which reveals several interesting features of the system.

The critical temperature T_c

First, let us investigate, given the chemical potential μ , how many solutions are permitted, what are the corresponding critical transition temperature for the ground




FIG. 1. Numerical solutions of equations of motion for $\langle \mathcal{O}_1 \rangle$ condensate at $\mu = 8$, $L = 1$, $k = 1$, $T/T_c = 0.1$ for the field $\Phi(z)$ (left) and $\Psi(z)$ (right) as functions of z . There are totally three different condensate solutions with this set of parameters: ground state solution (red), first excited state solution (green), blue excited state solution (blue).

state, for the 1st excited state, for the second excited state, ... Near criticality, the operator expectation values are small. one does not need to solve the full set of equation of motion to obtain the critical temperature [6]. The analysis of this section provides a check on our formulation by comparing with previous results obtained by directly solving the full system of equations of motion.

Indeed, near criticality $T \approx T_c$, the condensate solution is very small, $\Psi(z) \approx 0$ v \grave{a} $\Phi(z) \approx \mu(1 - z)$. The equation for Ψ becomes:

$$\Psi'' + \frac{F'}{F}\Psi' + \frac{zF' - 2F + 2b}{z^2 F}\Psi + \frac{\mu^2(1 - z)^2}{F^2}\Psi = 0, \quad (19)$$

Equation (19) is to be solved with appropriate boundary conditions depending on the type of condensate. At a given μ , the procedure for numerical calculation of the critical transition temperature T_c is following. Since it is a linear ordinary differential equation, one sets $\Psi(1) = 1$, $\Psi'(1) = (2b - T)/T$. Numerically integrating the equation from $z = 1$ to $z = 0$, one obtains unique boundary values $\Psi(0) = \Psi^{(1)}(\mu, T)$ and $\Psi'(0) = \Psi^{(2)}(\mu, T)$ as functions of μ and T . For $\langle \mathcal{O}_1 \rangle$ condensate, for example, by varying T , every time $\Psi^{(2)}(\mu, T)$ crosses zero, the corresponding T is the critical temperature for the ground state, 1st excited state, ... of the condensate.




FIG. 2. $\Psi'(0)$ as functions of z_h from numerically solving Eq. (19) for $\mu = 0.1$ (left) and 0.6 (right). Note that, there is a one-to-one correspondence between z_h and T . Smaller z_h corresponds to higher temperature T .

As an example, in Fig. 2, we plot $\Psi^{(2)}(\mu, T)$ as function of T for the case $L = 9$, with two different values of

the chemical potential $\mu = 0.1$ and 0.6 respectively. At $\mu = 0.1$, $\Psi^{(2)}(\mu, T)$ never crosses zero. This means that the system is always in the normal state, no phase transition at any finite T . As μ increases to 0.6 , $\Psi^{(2)}(\mu, T)$ crosses zero three times. This means that, at this value of the chemical potential, the system goes through phase transition to the ground state, 1st excited state, 2nd excited state at $z_h = 0.0236, 0.171$, and 0.652 . These values exactly match critical temperatures obtained by directly solving the full system of equations of motion [7].

The threshold chemical potential and its asymptotic behavior

As demonstrated, for small μ , the system permits no condensate solutions. As μ increases beyond a threshold μ_0 , a condensate solution is permitted below certain temperature T_c where the expectation values of the operators are non-zero, $\langle \mathcal{O}_\delta \rangle \neq 0$. This T_c is the critical temperature of the holographic transition. As μ increases further, more solutions are permitted at lower T_c which one identifies as excited states of the condensate. In this section, we would like to ask, at which threshold value of the chemical potential μ_0 , the condensate ground state solution is available; at which μ_1 , the first excited state solution is available,

Clearly, at each threshold μ where a new solution (new condensate state) appears, its critical temperature T_c is zero. As μ increases, T_c increases from zero. Therefore, we seek to investigate the criticality $\Psi(z)$ equation, Eq. (19), in the limit $T \rightarrow 0$. Before going to analytical discussions, let us give typical examples of threshold μ behaviors obtained from numerical solution. Fig. 3 shows




FIG. 3. The chemical potential μ where the ground and first four excited states appears when T decreases, plotted for two cases $L = 1$ (left) and $L = 9$ (right). The threshold μ are limits of these curves at $T \rightarrow 0$.

the threshold μ for five lowest energy states as the temperature $T \rightarrow 0$, for the two cases $L = 1$ and $L = 9$. In both cases, the topological charge $k = 1$. One sees that the threshold μ for ground state, and maybe first few excited states, shows saturation, but for higher states it shows no signs of saturation for T as low as 10^{-10} .

To understand better the threshold behavior, we resort

to analytic discussions. Eq. (19) is rewritten as:

$$-(F\Psi)' - \frac{zF' - 2F + 2b}{z^2}\Psi = \mu^2 \frac{(1-z)^2}{F}\Psi. \quad (20)$$

This is the Sturm–Liouville (S–L) eigenvalue problem. Its spectrum consists of an infinite number of eigenvalues which are real, countable, bounded from below and unbounded from above (more on this property later):

$$\mu_0^2 < \mu_1^2 < \mu_2^2 < \dots \quad \text{with} \quad \lim_{n \rightarrow \infty} \mu_n^2 \rightarrow \infty. \quad (21)$$

The corresponding eigenfunction, $\Psi_n(z)$, has exactly n zeros in the interval $(0, 1)$. Therefore, we easily identify these μ_n as the threshold chemical potential above which the system permits at least n different solutions: the ground state and $n - 1$ excited states. For $\mu < \mu_0$, no non-trivial solutions are available, the system cannot undergo phase transition and the normal state prevails all the way to $T = 0$ limit. The rest of this section concerns with explicit estimates for these eigenvalues.

We introduce new coordinate x and field $\omega(x)$:

$$x = \int_0^z ds (1-s)/F(s), \quad (22)$$

$$\omega(x) = \Psi(z)\sqrt{1-z}, \quad (23)$$

the S–L equation becomes,

$$-\frac{d^2\omega(x)}{dx^2} + V(x)\omega(x) = \mu^2\omega(x), \quad (24)$$

which is exactly the one-dimensional Schrödinger equation for a unit mass particle moving in the “potential”:

$$V(z(x)) = -\frac{z[2(z-1) + 2b - T]F}{(1-z)^2} - \frac{1}{2} \frac{1}{(1-z)^2} \left\{ \frac{3F^2}{2(1-z)^2} + \frac{F_z F}{1-z} \right\}. \quad (25)$$

The Schrödinger equation is to be solved with appropriate boundary conditions. Because $F(z) > 0$ almost everywhere, the relation, Eq. (22), gives one-to-one mapping between x and z . The interval $z \in [0, 1)$ maps into the interval $x \in [0, c)$ where

$$c = \int_0^1 ds (1-s)/F(s), \quad (26)$$

is positive and finite at finite T , and grows logarithmically with $1/T$ as $T \rightarrow 0$. For analytical solution, it is more appropriate to specify boundary conditions at the regular $x = 0$ boundary, requiring only that $\Psi(1)$ is finite (remind that we are working with a linear equation). Thus, $\omega(c) = 0$, a common physical requirement for Schrödinger equations at a singular boundary.

Boundary conditions at $x = 0$ depend on the type of condensate operator one wishes to study, either $\Psi(0) = 1$

and $\Psi'(0) = 0$ for $\langle \mathcal{O}_1 \rangle$ condensate; or $\Psi(0) = 0$ and $\Psi'(0) = 1$ for $\langle \mathcal{O}_2 \rangle$ condensate. These transform into corresponding boundary conditions for the $\omega(x)$ field:

$$\omega(0) = \Psi(0) = 1 \text{ or } 0, \quad (27)$$

$$\omega'(0) = b[\Psi'(0) - \Psi(0)/2] = -b/2 \text{ or } b. \quad (28)$$

for $\langle \mathcal{O}_1 \rangle$ or $\langle \mathcal{O}_2 \rangle$ condensate.

With the familiar Schrödinger form, Eq. (24), we are in a much better position to understand physical behaviors of the system using many well-known results from non-relativistic quantum mechanics. Fig. 4 shows the typical




FIG. 4. The typical potential $V(x)$ at low temperature for $L = 1$ (left) and $L = 9$ (right). At $x \rightarrow 0$, $V(x) = -\frac{3}{4}b^2 + b^2(1 - 2b)x$, which is negative. At $x \rightarrow c$, it is again negative and diverges as $-\frac{1}{4}(x - c)^{-2}$. $V(x)$ is bounded from above.

structure of the potential $V(x)$. From this, we can make many qualitative discussions. The first observation is that, near the singular boundary $x = c$,

$$V(x \rightarrow c) \simeq -\frac{2b - T}{c - x} - \frac{1/4}{(c - x)^2}. \quad (29)$$

The second dominant term near $x = c$ is the famous $-1/x^2$ potential [8]. It is well-known that if the range of x is infinite, the $-\alpha/x^2$ potential is unstable (not bounded from below, no ground state) unless $\alpha < 1/4$ and, when this is the case, all the eigenvalues are positive and the spectrum is continuous. In our case, α is exactly equal to the limiting value $1/4$ (this coincident is not well understood yet). However, we expect the first term to keep the eigenvalue spectrum bounded from below. Additionally, since the range of the potential is finite, the eigenvalue spectrum is discretized regardless.

Because of the exact $1/4$ value, the endpoint $x = c$ is of limit-circle non-oscillatory type. Rigorous mathematical investigation [9] shows that it can be regularized to an equivalent regular S-L equation. Thus, all the general properties of the regular S-L equation are applicable in our case. Most importantly, the spectrum is discrete and countable, unbounded from above and bounded from below, as already shown in Eq. (21). The eigenfunction $\omega_n(x)$ has exactly n zeros in $(0, c)$. Therefore, they correspond to the threshold μ for the ground state, 1st excited state, that we are seeking. The regularization requires introduction of regularization parameters and boundary conditions at $x = c$. However, they are regularization-scheme dependent, is irrelevant to our problem, and do not influence physical quantities.

For the second observation, one sees from Fig. 4 that $V(x)$ is bounded from above, $V(x) \leq V_{\max}$. Therefore, one can obtain analytical expressions in the limit of asymptotically large eigenvalues μ_n . Indeed, at $\mu^2 \gg V_{\max}$, our equation is simply

$$\omega_{xx} + \mu^2 \omega = 0. \quad (30)$$

This is the Schrödinger equation for a free particle in a infinite square-well potential. The general solution is $\omega(x) = Ae^{i\mu x} + Be^{-i\mu x}$. Since the range of x is finite, the wave number is quantized

$$\mu_n = n \frac{\pi}{c} + \text{const.} \quad (31)$$

To the leading order, this spectrum does not depend on the specific potential $V(x)$ or specific boundary conditions. These details only contribute a constant phase shift in the eigenfunctions and is different for different types of condensate.

Remind that c grows logarithmically with $1/T$. As the temperature $T \rightarrow 0$, Eq. (31) shows that the spectrum of eigenvalues with high quantum number n remains discrete, but the gap between μ_n shrinks to an infinitesimal value. This explains non-saturation behaviors observed in Fig. 3. Surprisingly, this asymptotic behavior seems to apply even for relatively low n in those figures, as low as $n \geq 2$. Note that, for $L = 9$ case (the gas side of the vdW BH transition), threshold values μ for the three lowest energy states immediately saturate, the gap between them does not shrink with decreasing T . On the other hand, for $L = 1$ case (the liquid side of the vdW BH transition), the gap between the 1st excited state and the ground state keeps shrinking. This might correlate with the fact that we do not see gap in the conductivity at low temperature for excited states in the gas side of the vdW transition [7].

The consequence of this observation is more profound. We postulate that at the exact limit $T = 0$ where *the gap at large n is zero*, all higher excited states of the condensate are activated beyond a finite chemical potential (approximately μ_1 or μ_2 depending on liquid or gas side of the BH transition), suggesting a potentially new quantum phase transition as function of the chemical potential. Unfortunately, the $T = 0$ limit is non-analytical and more investigations are needed to confirm this.

For the third and final observation, Eq. (29) shows that the range of the singular $-1/x^2$ term is within $1/8b$ of the “origin” $x = c$. As $T \rightarrow 0$, c grows logarithmically, this localized region becomes parametrically small. Therefore, we regard this term as a singular potential that “pins” the wave function to zero at the origin, $\omega(c) = 0$. Away from this parametrically small region, the Coulomb-like $-\alpha/x$ term in Eq. (29) dominates. Its spectrum is well-known [10]:

$$\mu_n^2 = -(2b - T)^2/4n^2.$$

To avoid negative values for μ^2 in the ground state in the original problem, one would need the constant shift $V_{\max} \geq (2b - T)^2/4$. This condition seems to be satisfied for all the set of L, k parameters we used so far, thus there are no negative eigenvalues in our problem. The ground state condensate occurs beyond a positive threshold chemical potential $\mu_0 > 0$. We do not pursue this further in this work, since $[z(x) - 1]^2 V(x)$ is a six-degree polynomial and no perturbative treatment of small parameters seems to be available in general cases.

As one last comment, let us study the system below but close to the critical temperature to obtain the critical exponent. Setting $\Psi_n(z) = \langle \mathcal{O}_n^\delta \rangle z^{\delta-1} \eta_n(z)$, from the equation for $\Phi(z)$ we have

$$\Phi'' = 2\Psi_n^2 \Phi / F = 2 \langle \mathcal{O}_n^\delta \rangle^2 z^{2(\delta-1)} \eta_n^2(z) \Phi / F. \quad (32)$$

Near T_c , $\langle \mathcal{O}_n^\delta \rangle$ is very small, we expand in this parameter:

$$\Phi(z) = \mu(1 - z) + \langle \mathcal{O}_n^\delta \rangle^2 \chi(z), \quad (33)$$

with $\chi(0) = \chi(1) = 0$, $\chi(z) < 0$. Substituting into Eq. (32), keeping only the leading terms, one gets

$$\chi''(z) = \mu(1 - z) 2z^{2(\delta-1)} \eta_n^2(z) / F(z). \quad (34)$$

$$\chi(z) = \chi'(0)z + \mu \int_0^z dz_1 \int_0^{z_1} ds \frac{2s^{2(\delta-1)} \eta_n^2(s)}{F(s)/(1-s)}. \quad (35)$$

The condition $\chi(1) = 0$ gives

$$\chi'(0) = -C_n = \mu \int_0^1 dz' \int_0^{z'} ds \frac{2s^{2(\delta-1)} \eta_n^2(s)}{F(s)/(s-1)} \quad (36)$$

By this definition, the constant C_n is finite and positive. From the limit $\Phi(z \rightarrow 0) = \mu - \rho z$, we have

$$\mu - \rho z = \mu(1 - z) - \langle \mathcal{O}_n^\delta \rangle^2 C_n z, \quad (37)$$

$$\langle \mathcal{O}_n^\delta \rangle \simeq \sqrt{\mu/C_n} [1 - T/T_c]^{1/2}. \quad (38)$$

Here we use the fact that near T_c , $\rho \simeq \mu T_c / T$. Eq. (38) shows that the condensates obey universal mean-field critical exponent 1/2 for all ground and excited states.

In conclusion, by considering holographic phase transition near criticality in the AdS/CFT duality, excited states of the condensate of the conformal field living at the AdS boundary at large quantum number are investigated. The spectrum of threshold chemical potentials of the states are discrete, asymptotically equal spacing, with the gap between consecutive values shrinks to zero logarithmically in the limit $T \rightarrow 0$. Surprisingly, numerical solution shows that this asymptotic behavior is valid for the quantum number n as low as 2 or 3 depending on the nature of the small-large BH vdW-like transition. We postulate that, at the exact limit $T = 0$ where the gap vanished, all excited states are activated beyond a finite chemical potential, suggesting a potentially new quantum phase transition. More investigation of this new transition will be conducted in the near future.

We thank D. T. Son, P. H. Lien, and H. V. Quyet for useful discussions.

* toannt@hus.edu.vn, toannt@vnu.edu.vn

- [1] M. Ammon and J. Erdmenger, *Gauge/gravity duality: Foundations and applications* (Cambridge University Press, 2015).
- [2] D. Kubizňák, R. B. Mann, and M. Teo, *Class. and Quantum Gravity* **34**, 063001 (2017).
- [3] T. H. Phat and T. T. Nguyen, *Eur. Phys. J. C* **81**, 1 (2021).
- [4] C. Herzog, P. Kovtun, and D. Son, *Phys. Rev. D* **79**, 066002 (2009).
- [5] Y.-Q. Wang, T.-T. Hu, Y.-X. Liu, J. Yang, and L. Zhao, *Excited states of holographic superconductors* (2020), arXiv:1910.07734 [hep-th].
- [6] G. Siopsis and J. Therrien, *J. High Energy Phys.* **2010** (5), 1.
- [7] T. H. Phat and T. T. Nguyen (2021), manuscript in preparation.
- [8] A. M. Essin and D. J. Griffiths, *Am. J. Phys.* **74**, 109 (2006).
- [9] A. Zettl, *Sturm-liouville theory*, 121 (American Mathematical Soc., 2010).
- [10] H. N. Yezpez, C. Vargas, and A. S. Brito, *Eur. J. Phys.* **8**, 189 (1987).