

On mass spectra of primordial black holes

Alexander A. Kirillov* ¹ and Sergey G. Rubin† ^{1,2}

¹ National Research Nuclear University MEPhI
(Moscow Engineering Physics Institute),
115409 Kashirskoe shosse 31, Moscow, Russia

² N. I. Lobachevsky Institute of Mathematics and Mechanics,
Kazan Federal University,
420008 Kremlevskaya street 18, Kazan, Russia

Abstract

Evidences for the primordial black holes (PBH) presence in the early Universe renew permanently. New limits on their mass spectrum challenge existing models of PBH formation. One of the known model is based on the closed walls collapse after the inflationary epoch. Its intrinsic feature is multiple production of small mass PBH which might contradict observations in the nearest future. We show that the mechanism of walls collapse can be applied to produce substantially different PBH mass spectra if one takes into account the classical motion of scalar fields together with their quantum fluctuations at the inflationary stage.

1 Introduction

Interest in the primordial black holes (PBHs) is dramatically increasing since the gravitational waves discovery from the black holes mergers [1]. However, PBHs origin and possible formation mechanisms are still a topical issue of the modern astrophysics and cosmology. The first ideas of such mechanisms had been proposed in [2–4] and lately developed in many other works (see reviews and references within [5, 6]). The different PBH spectra are used in papers [7–13] depending on specific needs.

The phase transitions of the first [14–18] and the second order [19, 20] might also underlay a mechanism of the PBH formation. In this paper, we continue elaboration of the model based on the second type phase transitions during the inflationary epoch [19–21]. However, the described model has a flaw. It inevitably leads to a multiple production of small mass PBHs. That problem

*AAKirillov@mephi.ru

†SGRubin@mephi.ru

could not be avoided within the framework of the discussed scenario, and typical mass spectra have the falling form $dN/dM \propto M^{-\alpha}$, $\alpha > 0$ (see review [22]). Such form of spectra could be unfavorable for explaining the observable effects. Moreover, the overproduction of low-mass PBHs could contradict future experiments. The way out discussed in this paper consists of an involving the classical motion of scalar fields together with their quantum fluctuations.

We study the way to circumvent the problem supposing a complicated form of scalar field potential. The latter is used in a variety of inflationary models predicting the potential landscape. Moreover, the inflaton might consist of multiple fields [23]. For instance, the supergravity often produces more than one physical scalar field [24] and predicts nontrivial forms of inflaton potentials [25]. The string theory also predicts the landscape with a large number of vacua, local peaks and saddle points [26, 27]. Such complicated potential can be presented both random and quasi-periodic shape [28] and leads to the multi-field inflation such as multi-stream inflation [29, 30], assisted inflation [31] or multi-field inflation with a random potential [32, 33]. Therefore, other non-inflaton scalar fields might have complicated potential as well.

This paper is organized as follows. In Section 2, we elaborate the way to involve the classical part of scalar fields into the expression for its fluctuations probability. The PBH spectrum depends on an initial position of the scalar field that allow us to adjust the model predictions to future observational data without inserting small parameters. The numerical results are represented in Section 3. Finally, Section 4 concludes the paper.

2 Quantum fluctuations accompanied by classical motion at the inflationary stage

The discussed mechanism of PBHs production requires closed domain walls formation due to the quantum fluctuations of scalar fields at the inflation epoch [19, 20]. Let us take into account both quantum and classical motion of fields. Consider the scalar field Φ of mass m and the standard action

$$S = \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{g_4} \left[R + (\partial\Phi)^2 - m^2\Phi^2 \right]. \quad (1)$$

Here, M_{Pl} is the Planck mass. The scalar field could be the inflaton field as well as a spectator one. The field equation in the de Sitter space is represented as [34]

$$\begin{aligned} \frac{\partial\Phi}{\partial t} - \frac{1}{3H} \left[e^{-2Ht} \Delta\Phi - \frac{\partial V(\Phi)}{\partial\Phi} \right] &= y(\mathbf{x}, t); \\ y(\mathbf{x}, t) &\equiv \left(-\frac{1}{3H} \frac{\partial^2}{\partial t^2} - \frac{\partial}{\partial t} + \frac{1}{3H} e^{-2Ht} \Delta \right) Q(\mathbf{x}, t). \end{aligned} \quad (2)$$

Here, $Q(x, t)$ is the “quick” part of the Fourier field decomposition. This equation was simplified: we have omitted the second time derivative due to a slow

roll approximation and have neglected higher powers of the function $y(\mathbf{x}, t)$. The latter is supposed to be small so that we may find a solution to the equation in the form

$$\Phi = \Phi_{\text{cl}} + \phi. \quad (3)$$

The deterministic part of the classical field Φ_{cl} is governed by the equation

$$\frac{\partial \Phi_{\text{cl}}}{\partial t} - \frac{1}{3H} \left[e^{-2Ht} \Delta \Phi_{\text{cl}} - \frac{\partial V(\Phi_{\text{cl}})}{\partial \Phi_{\text{cl}}} \right] = 0, \quad (4)$$

while its random part ϕ depends strictly on quantum fluctuations according to the linear equation

$$\frac{\partial \phi}{\partial t} - \frac{1}{3H} \left[e^{-2Ht} \Delta \phi - V''(\Phi_{\text{cl}}) \phi \right] = y(\mathbf{x}, t). \quad (5)$$

Here, we consider the limit $\Phi_{\text{cl}} \gg \phi$ which is valid if the random “force” $y(\mathbf{x}, t)$ is small.

Let us denote

$$m^2(t) \equiv V''(\Phi_{\text{cl}}). \quad (6)$$

The parameter m is positive if we are near the bottom of potential and is imaginary if we are near the potential maximum. It is supposed that $m(t)$ varies slowly during inflation.

We are interested in the super horizon scales where the fluctuations do not depend on the space coordinates. The uniform distribution $\Phi = \Phi(t)$ is governed by the more simple equation

$$\frac{\partial \Phi_{\text{cl}}}{\partial t} + \frac{1}{3H} \frac{\partial V(\Phi_{\text{cl}})}{\partial \Phi_{\text{cl}}} = 0, \quad (7)$$

$$\frac{\partial \phi}{\partial t} + \mu(t) \phi = y(t); \quad \mu(t) \equiv \frac{m^2(t)}{3H}, \quad (8)$$

provided that $H(t) = \text{const}$. The correlator of the random function $y(t)$ may be approximated as follows [35]

$$\langle y(t_1) y(t_2) \rangle = D(\mathbf{x}, t_1; \mathbf{x}, t_2) = \frac{H}{4\pi^2} \delta(t_1 - t_2). \quad (9)$$

The delta function in the rhs of this expression indicates that the random function $y(t)$ is distributed according to the Gauss law with the density

$$W(y) = \text{const} \exp \left[-\frac{1}{2\sigma^2} \int y^2(t) dt \right], \quad \sigma = \frac{H^{3/2}}{2\pi}. \quad (10)$$

The probability distribution of the function ϕ is proportional to that of the function $y(t)$ due to their linear relationship (8). It means that the probability to find the specific value $\phi(t)$ inside some small interval is equal to [36]

$$dP(\phi) = \text{const} \mathcal{D}\phi \exp \left[-\frac{1}{2\sigma^2} \int \left[\frac{\partial \phi}{\partial t} + \mu(t) \phi \right]^2 dt \right]. \quad (11)$$

Let's obtain the probability to find a quantum part of the field ϕ_2 at an instant t_2 provided that a value ϕ_1 at an instant t_1 is known. Evidently, we have to integrate over all values of the field inside the interval (t_1, t_2) and come to the expression

$$dP(\phi_2, t_2; \phi_1, t_1) = \text{const} \int_{\phi_1}^{\phi_2} \mathcal{D}\phi \exp \left[-\frac{1}{2\sigma^2} \int_{t_1}^{t_2} \left[\frac{\partial \phi}{\partial t} + \mu(t)\phi \right]^2 dt \right]. \quad (12)$$

The constant factor in this equation is determined by normalization condition

$$\int_{-\infty}^{\infty} \frac{dP(\phi_2, t_2; \phi_1, t_1)}{d\phi_2} d\phi_2 = 1. \quad (13)$$

Functional integral (12) can be calculated in the standard manner by finding an extreme trajectory of the integral in the exponent

$$\ddot{\phi} - \mu^2(t)\phi = 0, \quad (14)$$

where the term $\dot{\mu}$ is neglected due to slow variation of $\mu(t)$. The boundary conditions for (14) are as follows

$$\phi(t_1) = \phi_1; \quad \phi(t_2) = \phi_2. \quad (15)$$

Exact solution to this equation is

$$\phi(t) = A \exp(M(t)) + B \exp(-M(t)); \quad M(t) \equiv \int_{t_1}^t \mu(t') dt'; \quad (16)$$

$$A = \frac{\phi_2 - \phi_1 e^{-M(t_2)}}{2 \sinh(M(t_2))}, \quad B = -\frac{\phi_2 - \phi_1 e^{M(t_2)}}{2 \sinh(M(t_2))}. \quad (17)$$

Notice that $M(t_1) = 0$ by definition.

Substituting this solution into the integral in the exponent of the expression (12) one obtains the desired probability in the saddle point approximation

$$\begin{aligned} dP(\phi_2, t_2; \phi_1, t_1) &= \text{const} \int_{\phi_1}^{\phi_2} d\phi_2 \exp \left[-\frac{2A^2}{\sigma^2} \int_{t_1}^{t_2} \mu^2(t) e^{2M(t)} dt \right] = \\ &= \text{const} \int_{\phi_1}^{\phi_2} d\phi_2 \exp \left[-q^2 \left(\phi_2 - \phi_1 e^{-M(t_2)} \right)^2 \right], \end{aligned} \quad (18)$$

$$q^2 \equiv \frac{1}{2\sigma^2 \sinh^2(M(t_2))} \int_{t_1}^{t_2} \mu^2(t) e^{2M(t)} dt. \quad (19)$$

It describes the probability to find specific value of "quantum" part of the field (3). The "classical" part of the field Φ_{cl} is incorporated into the function

$M(t)$. The probability for the field value Φ (the distribution function f) is easily obtained by substitution $\phi(t) = \Phi(t) - \Phi_{\text{cl}}(t)$ into the formula above.

$$\begin{aligned} f(\Phi_2, t) &= \frac{dP(\Phi_2, t; \Phi_1, t_1)}{d\Phi_2} = \\ &= \sqrt{\frac{q^2}{\pi}} \exp \left[-q^2 \left(\Phi_2 - \Phi_{\text{cl}}(t_2) - (\Phi_1 - \Phi_{\text{cl}}(t_1)) e^{-M(t_2)} \right)^2 \right]. \end{aligned} \quad (20)$$

The limit $m \rightarrow 0$ restores the textbook formula.

The aim of the next section is to demonstrate how the obtained formulas can be applied to a particular scalar field potential. It is assumed that the potential may possess many extremes of different kind. In our consideration, we choose a part of phase space containing two maxima and at least one saddle point.

3 The PBH formation

In this section, we show that the classical motion of fields together with their quantum fluctuations influence the PBH mass spectra. To this end, we have to find the classical trajectory and use the probability (20) derived above.

The fields move between potential local maxima that leads to complicated spectra of fluctuations. The latter are discussed in papers [23, 24]. At the same time, the presence of saddle points is the reason of the closed domain walls formation, see details in [37, 38]. In the following, they could collapse to black holes [20, 22].

Let us consider the model of two real scalar fields with the Lagrangian

$$\mathcal{L} = \frac{1}{2} (\partial_\mu \Phi \partial^\mu \Phi + \partial_\mu X \partial^\mu X) - V(\Phi, X). \quad (21)$$

We choose the potential possessing n peaks and saddle points

$$\begin{aligned} V(\Phi, X) &= \frac{m_\phi^2}{2} \Phi^2 + \frac{m_\chi^2}{2} X^2 + \sum_{i=1}^n \delta V_i(\Phi, X), \\ \delta V_i(\Phi, X) &= \Lambda_i^4 \exp \left(- \left(\frac{\Phi - \phi_i}{\Delta_i} \right)^2 - \left(\frac{X - \chi_i}{\Delta_i} \right)^2 \right). \end{aligned} \quad (22)$$

Here, δV_i describes the i -th local maximum. The global minimum of the potential is located at the point $(\phi_{\text{min}}, \chi_{\text{min}}) = (0, 0)$ with exponentially small errors. Hereinafter, all variables are taken in the Hubble units H where $H \approx 10^{13}$ GeV at the inflationary epoch.

For our estimates, we choose the fields masses $m_\phi = 0.4$ and $m_\chi = 0.5$. For simplicity, we consider the potential with two peaks ($n = 2$) with the coordinates $\phi_1 = -9.0$, $\chi_1 = 3.0$ and $\phi_2 = -1.7$, $\chi_2 = 4.5$. The parameters corresponding to the peaks heights are $\Lambda_1 = 3.0$ and $\Lambda_2 = 1.5$, and the peaks widths are set with




Figure 1: The contour plot of the potential (22) with the parameters $m_1 = 0.4$, $m_2 = 0.5$, $\Lambda_1 = 3.0$, $\Lambda_2 = 1.5$, $\phi_1 = -9.0$, $\chi_1 = 3.0$, $\phi_2 = -1.7$, $\chi_2 = 4.5$, $\Delta_1 = 0.5$, $\Delta_2 = 1.5$ is shown. The red circles illustrate the classical trajectory of the fields Φ , X , and the black cross shows the potential minimum. The initial fields values for (23) are $\phi_{\text{in}} = -8.0$ and $\chi_{\text{in}} = 45.0$.

$\Delta_1 = 0.5$ and $\Delta_2 = 1.5$. The initial fields values are $\phi_{\text{in}} = -8.0$ and $\chi_{\text{in}} = 45.0$. Note, all chosen parameters have the values $\sim \mathcal{O}(1)$.

Following Section 2, the first step consists of finding the classical trajectory $\Phi_{\text{cl}}(t)$, $X_{\text{cl}}(t)$ of the fields Φ , X . Starting from the initial values $(\phi_{\text{in}}, \chi_{\text{in}})$ at the inflation epoch, the scalar fields tend to the potential minimum. The process is described by the classical motion equations

$$\begin{cases} \Phi_{\text{cl},tt} + 3H \Phi_{\text{cl},t} + \frac{\partial V(\Phi_{\text{cl}}, X_{\text{cl}})}{\partial \Phi_{\text{cl}}} = 0, \\ X_{\text{cl},tt} + 3H X_{\text{cl},t} + \frac{\partial V(\Phi_{\text{cl}}, X_{\text{cl}})}{\partial X_{\text{cl}}} = 0. \end{cases} \quad (23)$$

The classical evolution of the fields and the form of the specific potential are shown in Figure 1. At the same time, quantum fluctuations lead to fields “diffusion” during inflation. The probability density f to find the fields Φ or X in any point of the physical space is given by formula (20).

Both Φ and X distributions depend on a classical position of the fields at the instant t . In our estimates, we suppose that the probability function for the quantum parts ϕ or χ of the fields is separated into two independent fluctuation processes

$$f(t, \phi, \chi) = f_\phi(t, \phi) f_\chi(t, \chi) = \frac{dP}{d\phi} \frac{dP}{d\chi}, \quad (24)$$

where distribution functions of each field ϕ, χ can be defined as in (18).

As noted above, the fields should reach a saddle point of potential for domain wall formation. Suppose that some quantum fluctuation crosses a saddle point

and ends up at a point of the area Ω . As was shown in [37, 38], it causes nontrivial field solutions of the system (23) characterized by a nonzero winding number. Such configurations might lead to the domain walls formation after the inflation is finished. Detailed explanation might be found in [37]. The calculation of an exact shape of the area Ω is a separate, quite complicated task, so that we limit ourselves with the following approximation. Let us assume that the area Ω is bordered by two lines $\chi_{\text{cl-sp}}(\Phi)$ and $\chi_{\text{min-sp}}(\Phi)$ in the phase space. The first line connects the classical value at the instant t and the saddle point $(\phi_{\text{sp}}, \chi_{\text{sp}})$

$$\chi_{\text{cl-sp}}(\Phi) = \chi_{\text{sp}} + \frac{\Phi - \phi_{\text{sp}}}{\phi_{\text{cl}}(t) - \phi_{\text{sp}}} (\chi_{\text{cl}}(t) - \phi_{\text{sp}}). \quad (25)$$

The second one connects the vacuum value and the saddle point

$$\chi_{\text{min-sp}}(\Phi) = \chi_{\text{sp}} + \frac{\Phi - \phi_{\text{sp}}}{\phi_{\text{min}} - \phi_{\text{sp}}} (\chi_{\text{min}} - \phi_{\text{sp}}). \quad (26)$$

Thus, the probability for the fields to attain the area Ω where domain walls might form is calculated by integrating (24)

$$P(t) = \iint_{\Omega} f(\phi, \chi, t) dX d\Phi = \int_{\phi_{\text{sp}}}^{+\infty} f_{\phi}(\phi, t) d\Phi \int_{\chi_{\text{cl-sp}}(\Phi)}^{\chi_{\text{min-sp}}(\Phi)} f_{\chi}(\chi, t) dX. \quad (27)$$

Here, both distribution functions f_{ϕ} and f_{χ} are defined in (20) and according to (3) $\phi = \Phi - \Phi_{\text{cl}}$, $\chi = X - X_{\text{cl}}$. The algorithm for calculating the probability (27) discussed above is more accurate than that used in the previous papers.

Now, let us find the mass spectra of primordial black holes. In the considered model, they are formed due to collapse of domain walls. Here, we briefly reproduce the idea, while details may be found in the review [22]. As was shown in [37, 38], domain walls might be formed due to the quantum fluctuations in field models with potential possessing at least one saddle point and a local maximum. The protosoliton is formed if the fields achieve a saddle point (in our model, we have noted this area as Ω). These protosoliton field configurations are quickly expanded during inflation. The final scale of such configuration depends on a e-fold number N . More definitely, the configuration scale is stretched in the factor $\sim e^{N_{\text{inf}} - N}$ to the end of inflation. The soliton is quickly formed after the end of inflation. The total mass of the field configuration is proportional to its area. Evidently, it could collapse into a black hole after the end of inflation [19–21].

The regions number where the fields reach the critical values ϕ_{cr} and χ_{cr} belonging to Ω can be found as

$$n(t) = P(t) e^{3Ht}. \quad (28)$$

Here, the term e^{3Ht} is the number of causally independent regions of the size H^{-1} at the instant t from the beginning of inflation. After the end of inflation

at $t = N_{\text{inf}}H^{-1} = 60H^{-1}$, the size of each region is expanded

$$r_0(t) = H^{-1} \exp(N_{\text{inf}} - Ht) \gg r_{h,0}. \quad (29)$$

Here, $N_{\text{inf}} \approx 60$ is the total e-folds number, and $r_{h,0}$ is the horizon size at the end of inflation. At the radiation stage (RD), each region expands as $\propto \sqrt{\tau}$ while the horizon size is $r_h = H^{-1}(\tau) = 2\tau$. Here, τ is time after the beginning of the RD epoch. After a domain wall goes under the horizon, its collapse begins (the details of the process taking into account detachment from the Hubble flow may be found in [22, 39, 40]). Thus, the maximal size of a domain wall can be written as the function of the instant t

$$r(t) \approx \frac{1}{2} \frac{e^{2(N_{\text{inf}} - Ht)}}{HN_{\text{inf}}}. \quad (30)$$

After eliminating of t from (28) and (30), one can finally get the distribution $n(r)$ of closed walls sizes which can be rearranged into the mass spectrum.

Next, we have to find masses of PBHs. For simplicity, we assume total energy of domain wall converts to a black hole mass during its collapse and neglect nonsphericity of a domain wall and losses caused by gravitational waves. The energy density of a domain wall might be found by a common way. The energy momentum tensor for the Lagrangian (21) is given by

$$T_{\nu}^{\mu} = \sum_{i=1,2} \left(\partial^{\mu} \varphi_i \partial_{\nu} \varphi_i - \frac{1}{2} \partial^{\alpha} \varphi_i \partial_{\alpha} \varphi_i \delta_{\nu}^{\mu} \right) + V \delta_{\nu}^{\mu}, \quad (31)$$

where φ_1, φ_2 correspond to the fields Φ_{cl} and X_{cl} , respectively. Then, the energy density of a domain wall is found to be

$$\varepsilon(x) = T_0^0 = \frac{1}{2} \sum_{i=1,2} ((\partial_t \varphi_i)^2 + (\partial_x \varphi_i)^2) + V. \quad (32)$$

Upon integrating (32) over the all possible values of x (infinite interval), the surface energy density of a domain wall σ may be found. Finally, masses of black holes are $M(r(t)) \simeq 4\pi\sigma r^2(t)$. Taking into account (28) and (30), one can find the mass spectrum of primordial black holes. The PBH mass distribution for the parameters of the Lagrangian (21) is shown in Figure 2. Note, the mass spectrum has the non-power form due to taking into account both the quantum and classical motion of scalar fields and the compound form of the potential leading to nontrivial classical fields trajectory. The obtained spectrum is free from overproduction of light PBHs and, therefore, is much more adaptable to new observational effects. We leave the detailed analysis of these possibilities for future research.

4 Conclusion

In this note, we have shown that the mechanism of PBHs formation in the second order phase transitions of scalar fields might produce a wide variety of




Figure 2: The PBH mass distribution is shown.

the PBH mass spectra. It is expected that observations will help to select an appropriate one. The key point is the classical field motion which was taken into account together with the quantum fluctuations at the inflationary stage. We show here that the probability to find a particular value of scalar field at a space point depends on its classical dynamics. We have derived the appropriate analytical formula and have applied it to obtain one of the PBH mass spectra. The elaborated method is the useful tool to fit an observable spectrum in the near future.

Acknowledgements

This research was funded by the Ministry of Science and Higher Education of the Russian Federation, Project “Fundamental properties of elementary particles and cosmology” № 0723-2020-0041.

References

- [1] B. P. Abbott et al., “Observation of Gravitational Waves from a Binary Black Hole Merger”, [Phys. Rev. Lett. **116**, 061102, 061102 \(2016\)](#), [arXiv:1602.03837 \[gr-qc\]](#).
- [2] Y. B. Zel’dovich and I. D. Novikov, “The Hypothesis of Cores Retarded during Expansion and the Hot Cosmological Model”, [Sov. Astron. J. **10**, 602–603 \(1967\)](#).
- [3] S. Hawking, “Gravitationally collapsed objects of very low mass”, [Mon. Not. R. Astron. Soc. **152**, 75 \(1971\)](#).
- [4] B. J. Carr and S. W. Hawking, “Black holes in the early Universe”, [Mon. Not. R. Astron. Soc. **168**, 399–416 \(1974\)](#).

- [5] M. Y. Khlopov, “Primordial black holes”, *Res. Astron. Astrophys.* **10**, 495–528 (2010), [arXiv:0801.0116](#).
- [6] B. Carr and F. Kühnel, “Primordial Black Holes as Dark Matter: Recent Developments”, *Annu. Rev. Nucl. Part. Sci.* **70**, 355–394 (2020), [arXiv:2006.02838 \[astro-ph.CO\]](#).
- [7] B. J. Carr, “The Primordial black hole mass spectrum”, *Astrophys. J.* **201**, 1–19 (1975).
- [8] A. Dolgov and J. Silk, “Baryon isocurvature fluctuations at small scales and baryonic dark matter”, *Phys. Rev. D* **47**, 4244–4255 (1993).
- [9] Y. Sendouda, S. Nagataki, and K. Sato, “Mass spectrum of primordial black holes from inflationary perturbation in the Randall Sundrum braneworld: a limit on blue spectra”, *J. Cosmol. Astropart. Phys.* **2006**, 003, 003 (2006), [arXiv:0603509 \[astro-ph\]](#).
- [10] S. Clesse and J. García-Bellido, “Massive primordial black holes from hybrid inflation as dark matter and the seeds of galaxies”, *Phys. Rev. D* **92**, 023524 (2015), [arXiv:1501.07565 \[astro-ph.CO\]](#).
- [11] J. Garriga, A. Vilenkin, and J. Zhang, “Black holes and the multiverse”, *J. Cosmol. Astropart. Phys.* **2016**, 064 (2016), [arXiv:1512.01819 \[hep-th\]](#).
- [12] B. Carr and F. Kühnel, “Primordial black holes with multimodal mass spectra”, *Phys. Rev. D* **99**, 103535 (2019), [arXiv:1811.06532 \[astro-ph.CO\]](#).
- [13] J. Liu, Z.-K. Guo, and R.-G. Cai, “Primordial black holes from cosmic domain walls”, *Phys. Rev. D* **101**, 023513 (2020), [arXiv:1908.02662 \[astro-ph.CO\]](#).
- [14] H. Kodama, M. Sasaki, and K. Sato, “Abundance Of Primordial Holes Produced By Cosmological First Order Phase Transition”, *Prog. Theor. Phys.* **68**, 1979 (1982).
- [15] S. W. Hawking, I. G. Moss, and J. M. Stewart, “Bubble Collisions in the Very Early Universe”, *Phys. Rev. D* **26**, 2681 (1982).
- [16] R. V. Konoplich et al., “Black hole production in first order phase transitions in the Universe”, *Sov. Pis'ma Astron. J.* **24**, 1 (1998).
- [17] K. Jedamzik and J. C. Niemeyer, “Primordial black hole formation during first-order phase transitions”, *Phys. Rev. D* **59**, 124014 (1999), [arXiv:astro-ph/9901293 \[astro-ph\]](#).
- [18] R. V. Konoplich et al., “Formation of black holes in first-order phase transitions as a cosmological test of symmetry-breaking mechanisms”, *Phys. of Atom. Nucl.* **62**, 1593–1600 (1999).
- [19] S. G. Rubin, M. Y. Khlopov, and A. S. Sakharov, “Primordial Black Holes from Non-Equilibrium Second Order Phase Transition”, *Grav. Cosmol. S* **6**, 51–58 (2000), [arXiv:hep-ph/0005271](#).
- [20] S. G. Rubin, A. S. Sakharov, and M. Y. Khlopov, “The Formation of Primary Galactic Nuclei during Phase Transitions in the Early Universe”, *J. Exp. Theor. Phys.* **92**, 921–929 (2001), [arXiv:hep-ph/0106187](#).

- [21] M. Y. Khlopov, S. G. Rubin, and A. S. Sakharov, “Strong Primordial Inhomogeneities and Galaxy Formation”, (2002), [arXiv:0202505 \[astro-ph\]](#).
- [22] K. M. Belotsky et al., “Clusters of Primordial Black Holes”, *Eur. Phys. J. C* **79**, 246 (2019), [arXiv:1807.06590 \[astro-ph.CO\]](#).
- [23] D. Wands, “Multiple Field Inflation”, in *Lecture notes in physics*, Vol. 738, edited by M. Lemoine, J. Martin, and P. Peter (Springer, Berlin, 2007), p. 275.
- [24] S. V. Ketov and A. A. Starobinsky, “Inflation and nonminimal scalar-curvature coupling in gravity and supergravity”, *J. Cosmol. Astropart. Phys.* **2012**, 022 (2012), [arXiv:1203.0805 \[hep-th\]](#).
- [25] S. V. Ketov, “Multi-Field versus Single-Field in the Supergravity Models of Inflation and Primordial Black Holes”, *Universe* **7**, 115 (2021).
- [26] L. Susskind, “The Anthropic Landscape of String Theory”, in The davis meeting on cosmic inflation (2003), p. 26, [arXiv:0302219 \[hep-th\]](#).
- [27] J. M. Cline, “Inflation from String Theory”, arXiv e-prints, hep-th/0501179 (2005), [arXiv:0501179 \[hep-th\]](#).
- [28] S. Li, Y. Liu, and Y.-S. Piao, “Inflation in a web”, *Phys. Rev. D* **80**, 123535 (2009), [arXiv:0906.3608 \[hep-th\]](#).
- [29] M. Li and Y. Wang, “Multi-stream inflation”, *J. Cosmol. Astropart. Phys.* **7**, 033, 33 (2009), [arXiv:0903.2123 \[hep-th\]](#).
- [30] F. Duplessis, Y. Wang, and R. Brandenberger, “Multi-stream inflation in a landscape”, *J. Cosmol. Astropart. Phys.* **4**, 012, 12 (2012), [arXiv:1201.0029 \[hep-th\]](#).
- [31] D. Battefeld and T. Battefeld, “Multi-field inflation on the landscape”, *J. Cosmol. Astropart. Phys.* **2009**, 027, 027 (2009), [arXiv:0812.0367 \[hep-th\]](#).
- [32] S.-H. H. Tye, J. Xu, and Y. Zhang, “Multi-field inflation with a random potential”, *J. Cosmol. Astropart. Phys.* **4**, 018, 18 (2009), [arXiv:0812.1944 \[hep-th\]](#).
- [33] J. Frazer and A. R. Liddle, “Multi-field inflation with random potentials: field dimension, feature scale and non-Gaussianity”, *J. Cosmol. Astropart. Phys.* **2012**, 039, 039 (2012), [arXiv:1111.6646 \[astro-ph.CO\]](#).
- [34] M. Y. Khlopov and S. G. Rubin, *Cosmological pattern of microphysics in the inflationary universe* (Kluwer Academic Publishers, P.O. Box 17, 3300 AA Dordrecht, The Netherlands, 2004).
- [35] S.-J. Rey, “Dynamics of inflationary phase transition”, *Nuclear Physics B* **284**, 706–728 (1987).
- [36] R. Feynman, A. Hibbs, and D. Styer, *Quantum mechanics and path integrals*, Dover Books on Physics (Dover Publications, 2010).

- [37] V. A. Gani, A. A. Kirillov, and S. G. Rubin, “Classical transitions with the topological number changing in the early Universe”, *J. Cosmol. Astropart. Phys.* **2018**, 042, 042 (2018), [arXiv:1704.03688 \[hep-th\]](#).
- [38] B. S. Murygin, A. A. Kirillov, and V. V. Nikulin, “Cosmological formation of $(2 + 1)$ -dimensional soliton structures in models possessing potentials with local peaks”, *Physics* **3**, 563–568 (2021).
- [39] M. Y. Khlopov, S. G. Rubin, and A. S. Sakharov, “Primordial structure of massive black hole clusters”, *Astropart. Phys.* **23**, 265–277 (2005), [arXiv:astro-ph/0401532 \[astro-ph\]](#).
- [40] V. I. Dokuchaev, Y. N. Eroshenko, and S. G. Rubin, “Early formation of galaxies induced by clusters of black holes”, *Astron. Rep.* **52**, 779–789 (2008), [arXiv:0801.0885](#).